A LOCAL-GLOBAL PRINCIPLE FOR PARAMETRIZED ∞ -CATEGORIES

HADRIAN HEINE, UNIVERSITY OF OSLO, NORWAY, HADRIAH@MATH.UIO.NO

ABSTRACT. We prove a local-global principle for ∞ -categories over any base ∞ -category \mathbb{C} : we show that any ∞ -category $\mathbb{B} \to \mathbb{C}$ over \mathbb{C} is determined by the following data: the collection of fibers \mathbb{B}_X for X running through the set of equivalence classes of objects of \mathbb{C} endowed with the action of the space of automorphisms $\operatorname{Aut}_X(\mathbb{B})$ on the fiber, the local data, together with a locally cartesian fibration $\mathbb{D} \to \mathbb{C}$ and $\operatorname{Aut}_X(\mathbb{B})$ -linear equivalences $\mathbb{D}_X \simeq \mathcal{P}(\mathbb{B}_X)$ to the ∞ -category of presheaves on \mathbb{B}_X , the gluing data.

As applications we describe the ∞ -category of small ∞ -categories over [1] in terms of the ∞ -category of left fibrations and prove an end formula for mapping spaces of the internal hom of $\infty \operatorname{Cat}_{/[1]}$ and the conditionally existing internal hom of $\infty \operatorname{Cat}_{/[C]}$.

Considering functoriality in \mathcal{C} we obtain as a corollary that the double ∞ -category CORR of correspondences is the pullback of the double ∞ -category PR^L of presentable ∞ -categories along the functor ∞ Cat \rightarrow Pr^L taking presheaves. We deduce that ∞ -categories over any ∞ -category \mathcal{C} are classified by normal lax 2-functors.

CONTENTS

	1.	Introduction	1
	2.	Notation and terminology	5
	3.	Parametrized ∞ -categories of presheaves	6
	4.	A local-global principle	13
	5.	Correspondences as left fibrations	14
	6.	An end formula for mapping spaces of parametrized functor ∞ -categories	16
	7.	Correspondences as profunctors	19
References		ferences	26

1. INTRODUCTION

For every map of spaces $\phi: Y \to X$ we can decompose the space $X \simeq \coprod_{L \in \pi_0(X)} BAut_L(X)$ as the disjoint union of path components $BAut_L(X)$, the deloopings of the loop space at L, to obtain a decomposition of ϕ as

(1)
$$\coprod_{L \in \pi_0(X)} \phi^{-1}(B \operatorname{Aut}_L(X)) \to \coprod_{L \in \pi_0(X)} B \operatorname{Aut}_L(X).$$

In this sense the theory of spaces over X splits in the theories of spaces over any path component $BAut_L(X)$ for $L \in X$, i.e. the theory of spaces with an action of the grouplike automorphism space $Aut_L(X)$.

In ∞ -category theory the situation is very different: by the presence of non-invertible morphisms an ∞ -category \mathcal{C} rarely splits into all full subcategories spanned by a single

object running over all representatives of equivalence classes. On the other hand for a functor $\phi : \mathcal{D} \to X$ to a space there is a similar decomposition (1) like in topology. For an arbitrary functor $\phi : \mathcal{D} \to \mathcal{C}$ of ∞ -categories we can decompose the pullback $\iota(\mathcal{C}) \times_{\mathcal{C}} \mathcal{D} \to \iota(\mathcal{C})$ to the maximal subspace of \mathcal{C} as

$$\coprod_{L \in \pi_0(\iota(\mathfrak{C}))} \phi^{-1}(B\operatorname{Aut}_L(\iota(\mathfrak{C}))) \to \coprod_{L \in \pi_0(\iota(\mathfrak{C}))} B\operatorname{Aut}_L(\iota(\mathfrak{C}))$$

and hope to be able to reconstruct $\phi : \mathcal{D} \to \mathcal{C}$ from this decomposition via coherent gluing data. For $\mathcal{C} = [1]$ for example [17, 8.3.2. Proposition] a functor $\mathcal{M} \to [1]$ may be thought of as ∞ -categories $\mathcal{C} := \mathcal{M}_0, \mathcal{D} := \mathcal{M}_1$ and the collection of mapping spaces $\mathcal{M}(A, B)$ for $A \in \mathcal{C}, B \in \mathcal{D}$, organized to a restricted mapping space functor

$$\mathcal{C}^{\mathrm{op}} \times \mathcal{D} \subset \mathcal{M}^{\mathrm{op}} \times \mathcal{M} \xrightarrow{\mathcal{M}(-,-)} S.$$

The latter corresponds to a functor $\mathcal{D} \to \operatorname{Fun}(\mathcal{C}^{\operatorname{op}}, \mathbb{S})$ and so by the universal property of the ∞ -category of presheaves to a left adjoint functor $\operatorname{Fun}(\mathcal{D}^{\operatorname{op}}, \mathbb{S}) \to \operatorname{Fun}(\mathcal{C}^{\operatorname{op}}, \mathbb{S})$. In this sense a functor $\mathcal{M} \to [1]$ may be decomposed as its fibers over 0 and 1, the local data, and a cartesian fibration $\mathcal{N} \to [1]$ that classifies a left adjoint functor $\mathcal{N}_1 \to \mathcal{N}_0$ equipped with equivalences $\mathcal{N}_i \simeq \operatorname{Fun}(\mathcal{M}_i^{\operatorname{op}}, \mathbb{S})$ for i = 0, 1, the gluing data.

We extend this situation to any base ∞ -category \mathcal{C} . Let $\infty \operatorname{Cat}_{/\mathcal{C}}$ be the $(\infty, 2)$ -category of small ∞ -categories over \mathcal{C} and $\operatorname{LoCART}_{\mathcal{C}}^{\mathrm{L}}$ the $(\infty, 2)$ -category of locally cartesian fibrations over \mathcal{C} whose fibers are presentable and whose fiber transports are left adjoints, and functors over \mathcal{C} that are fiberwise left adjoints. We prove the following reconstruction principle:

Theorem 1.1. (Theorem 4.1) For every small ∞ -category \mathcal{C} there is a pullback square of $(\infty, 2)$ -categories



where $\mathfrak{P}^{\mathfrak{C}}$ sends a functor $\mathfrak{D} \to \mathfrak{C}$ to a locally cartesian fibration $\mathfrak{P}^{\mathfrak{C}}(\mathfrak{D}) \to \mathfrak{C}$ whose fiber over any $X \in \mathfrak{C}$ is the ∞ -category of presheaves on the fiber \mathfrak{D}_X .

For $\mathcal{C} = [2]$ for example Theorem 1.1 guarantees that a functor $\mathcal{M} \to [2]$ can be decomposed as its fibers over 0,1 and 2, the local data, and a locally cartesian fibration $\mathcal{N} \to [2]$ classifying left adjoint functors $\alpha : \mathcal{N}_2 \to \mathcal{N}_1, \beta : \mathcal{N}_1 \to \mathcal{N}_0, \gamma : \mathcal{N}_2 \to \mathcal{N}_0$ and a natural transformation $\beta \alpha \to \gamma$ equipped with equivalences $\mathcal{N}_i \simeq \operatorname{Fun}(\mathcal{M}_i^{\operatorname{op}}, \mathcal{S})$ for i = 0, 1, 2, the gluing data.

Specializing to $\mathcal{C} = [1]$ we obtain a pullback square



where $\Pr^{L} = \operatorname{LoCART}_{[0]}^{L}$ is the $(\infty, 2)$ -category of presentable ∞ -categories and left adjoint functors. We use the latter result to prove the following theorem, where $\mathcal{L} \subset \operatorname{Fun}([1], \infty \operatorname{Cat})$ is the full subcategory of left fibrations and $(-)^{\operatorname{op}} : \infty \operatorname{Cat} \to \infty \operatorname{Cat}$ forms the opposite ∞ -category:

Theorem 1.2. (Theorem 5.2) There is a canonical pullback square of ∞ -categories:



where the right hand vertical functor evaluates at the target.

The induced equivalence from the left upper corner of the square to the pullback sends a functor $\mathcal{M} \to [1]$ to the triple

$$(\mathcal{M}_0, \mathcal{M}_1, \mathcal{M}_0^{\mathrm{op}} \times \mathcal{M}_1 \times_{\mathcal{M}^{\mathrm{op}} \times \mathcal{M}} \mathrm{Tw}(\mathcal{M}) \to \mathcal{M}_0^{\mathrm{op}} \times \mathcal{M}_1),$$

where $\operatorname{Tw}(\mathcal{M}) \to \mathcal{M}^{\operatorname{op}} \times \mathcal{M}$ is the twisted arrow left fibration classifying the mapping space functor of \mathcal{M} .

Theorem 1.2 gives the following important corollaries:

Corollary 1.3. Let \mathcal{C}, \mathcal{D} be small ∞ -categories. The fiber of the functor

 $\infty Cat_{[1]} \rightarrow \infty Cat \times \infty Cat$

over $(\mathfrak{C}, \mathfrak{D})$ is the ∞ -category Fun $(\mathfrak{C}^{\mathrm{op}}, \operatorname{Fun}(\mathfrak{D}, \mathfrak{S}))$.

Let $\text{Cocart}_{[1]} \subset \infty \text{Cat}_{/[1]}$ be the full subcategory of cocartesian fibrations.

Corollary 1.4. Let \mathcal{C}, \mathcal{D} be small ∞ -categories. The fiber of the functor

$$\operatorname{Cocart}_{[1]} \to \infty \operatorname{Cat} \times \infty \operatorname{Cat}$$

over $(\mathcal{C}, \mathcal{D})$ is the ∞ -category $\operatorname{Fun}(\mathcal{C}^{\operatorname{op}}, \mathcal{D}^{\operatorname{op}}) \simeq \operatorname{Fun}(\mathcal{C}, \mathcal{D})^{\operatorname{op}}$.

Gepner-Haugseng-Nikolaus [7, Proposition 5.1.] prove an end formula for mapping spaces of functor ∞ -categories: they prove that for any ∞ -categories \mathcal{C}, \mathcal{D} the space of natural transformations from a functor $F : \mathcal{C} \to \mathcal{D}$ to a functor $G : \mathcal{C} \to \mathcal{D}$ is the end $\int_{X \in \mathcal{C}} \mathcal{D}(F(X), G(X))$. The authors define the latter as the limit of the functor

$$\mathrm{Tw}(\mathcal{C}) \to \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \xrightarrow{\mathrm{F}^{\mathrm{op}} \times \mathrm{G}} \mathcal{D}^{\mathrm{op}} \times \mathcal{D} \xrightarrow{\mathcal{D}(-,-)} S.$$

where $\operatorname{Tw}(\mathcal{C}) \to \mathcal{C}^{\operatorname{op}} \times \mathcal{C}$ is the twisted arrow left fibration whose fiber over any pair (A, B) in \mathcal{C} is the space of maps $A \to B$ in \mathcal{C} . As a corollary of Theorem 1.1 we obtain a global version of [7, Proposition 5.1.] that describes the twisted arrow left fibration of the ∞ -category of functors $\mathcal{C} \to \mathcal{D}$:

Corollary 1.5. Let \mathcal{C}, \mathcal{D} be ∞ -categories. There is a canonical equivalence

 $\operatorname{Tw}(\operatorname{Fun}(\mathcal{C},\mathcal{D})) \simeq \operatorname{Fun}(\mathcal{C},\mathcal{D})^{\operatorname{op}} \times \operatorname{Fun}(\mathcal{C},\mathcal{D}) \times_{\operatorname{Fun}(\operatorname{Tw}(\mathcal{C}),\mathcal{D}^{\operatorname{op}} \times \mathcal{D})} \operatorname{Fun}(\operatorname{Tw}(\mathcal{C}),\operatorname{Tw}(\mathcal{D}))$

over

$$\operatorname{Fun}(\mathcal{C},\mathcal{D})^{\operatorname{op}} \times \operatorname{Fun}(\mathcal{C},\mathcal{D})$$

which induces on the fiber over every functors $F, G : \mathfrak{C} \to \mathfrak{D}$ an equivalence

$$\operatorname{Fun}(\mathcal{C},\mathcal{D})(\mathrm{F},\mathrm{G})\simeq\int_{\mathrm{X}\in\mathcal{C}}\mathcal{D}(\mathrm{F}(\mathrm{X}),\mathrm{G}(\mathrm{X})).$$

Corollary 1.5 desribes the mapping spaces in the ∞ -category of functors, the internal hom of ∞ Cat. In a similar way we describe the mapping spaces in the internal hom of ∞ Cat/[1] (Theorem 6.5) and the conditionally existing internal hom of ∞ Cat/ $_{\mathbb{C}}$ (Corollary 6.6).

Ayala-Francis [2, Theorem 0.8] extend the ∞ -categorical Grothendieck construction for cartesian and cocartesian fibrations to a Grothendieck construction for exponential fibration, where an exponential fibration is a functor ϕ such that taking pullback along ϕ preserves

small colimits. Ayala-Francis prove the existence of a small ∞ -precategory Corr - or flagged ∞ -category in their language - that is the target of the following straightening: for every small ∞ -category \mathbb{C} there is a canonical equivalence between the space of functors $\mathbb{C} \to \mathbb{C}$ orr and the space of exponential fibrations over \mathbb{C} extending the straightening of cocartesian fibrations [19, Theorem 3.2.0.1.], [21], [20], [1]. By the self-duality of exponential fibrations the latter may equivalently be phrased as an equivalence between the space of functors $\mathbb{C}^{\text{op}} \to \mathbb{C}$ orr and the space of exponential fibrations over \mathbb{C} extending the straightening of cartesian fibrations. To determine the ∞ -precategory Corr the authors [2, Theorem 0.4] prove that the (derived version of the) nerve establishes an equivalence between ∞ -precategories and Segal spaces. Having this the authors show that mapping a simplex $[n] \in \Delta$ to the space of exponential fibrations over [n] defines a Segal space representing the ∞ -precategory Corr. We extend the picture of Ayala-Francis [2] via the concept of Segal ∞ -category or double ∞ -category [6, Definition 2.4.3.], which is a category object, i.e. Segal object, in ∞ Cat and so a categorical enhancement of a Segal space.

Removing the non-invertible vertical morphisms from a double ∞ -category gives rise to an $(\infty, 2)$ -precategory that we call the underlying horizontal $(\infty, 2)$ -precategory. Removing the non-invertible horizontal morphisms and 2-morphisms from a double ∞ -category gives rise to an ∞ -category that we call the underlying vertical ∞ -category. The latter process determines a cartesian fibration from the ∞ -category of double ∞ -categories to the ∞ category of ∞ -categories. In other words, we can pullback any double ∞ -category along any functor mapping to the underlying vertical ∞ -category.

We use Theorem 1.1 to prove that mapping a simplex $[n] \in \Delta$ to the ∞ -category of exponential fibrations over [n] determines a double ∞ -category CORR (Theorem 7.19). We have chosen this name to emphasize that CORR is a double-categorical refinement of the ∞ -precategory Corr. This way we answer a question of [2, Question 0.14] asking for a 2-categorical refinement of Corr.

Via Theorem 1.1 we build CORR from a simpler double ∞ -category: the ∞ -category ∞ Cat is cartesian closed and so an $(\infty, 2)$ -category. We prove (Proposition 7.20) that the $(\infty, 2)$ -category ∞ Cat is the underlying horizontal $(\infty, 2)$ -precategory of the double ∞ -category sending a simplex $[n] \in \Delta$ to the full subcategory of ∞ Cat_[n] of cartesian fibrations over [n]. We conclude that the $(\infty, 2)$ -category \Pr^L of presentable ∞ -categories and left adjoint functors, a locally full subcategory of the $(\infty, 2)$ -category ∞ Cat of large ∞ -categories, is the underlying horizontal $(\infty, 2)$ -precategory of a double ∞ -category \Pr^L . We deduce the following corollary from Theorem 1.1 considering functoriality in the base:

Theorem 1.6. (Theorem 7.19) The double ∞ -category CORR is the pullback of the double ∞ -category $\operatorname{PR}^{\operatorname{L}}$ along the functor $\infty \operatorname{Cat} \to \operatorname{Pr}^{\operatorname{L}}$ forming the ∞ -category of presheaves.

Passing to underlying $(\infty, 2)$ -precategories we obtain the following corollary:

Corollary 1.7. The underlying $(\infty, 2)$ -precategory of CORR is the pullback of the $(\infty, 2)$ -category $\operatorname{Pr}^{\mathrm{L}}$ along the map $\iota(\infty\mathsf{Cat}) \to \iota(\operatorname{Pr}^{\mathrm{L}})$ forming the ∞ -category of presheaves.

Corollary 1.7 immediately implies that the Rezk-completion of the underlying $(\infty, 2)$ -precategory of CORR is the full subcategory of the $(\infty, 2)$ -category \Pr^{L} spanned by the ∞ -categories of presheaves (Corollary 7.25).

Theorems 1.1 and 1.6 imply the straightening result of Ayalay-Francis [2, Theorem 0.8] as a corollary and moreover give an enhancement of their result. While Ayala-Francis prove for every small ∞ -category \mathcal{C} an equivalence of spaces between the space of functors $\mathcal{C}^{\text{op}} \to \text{Corr}$ and the space of exponential fibrations over \mathcal{C} , we obtain an equivalence of ∞ -categories inducing the equivalence of Ayala-Francis on maximal subspaces. We prove the following corollary, where N is the (derived version of the) nerve of an ∞ -category:

Corollary 1.8. (Corollary 7.26) Let \mathcal{C} be a small ∞ -category. There is a canonical equivalence of ∞ -categories

$$FUN(N(\mathcal{C}^{op}), CORR) \simeq EXP_{\mathcal{C}}$$

between the ∞ -category of maps of double ∞ -categories $\mathbb{C}^{\mathrm{op}} \to \mathrm{CORR}$ and the full subcategory $\mathrm{EXP}_{\mathbb{C}} \subset \infty \mathrm{Cat}_{/\mathbb{C}}$ spanned by the exponential fibrations over \mathbb{C} .

Corollary 1.8 in different formulation is also claimed without proof in [5, 4.8. Proposition]. Moreover Theorems 1.1 and 1.6 give rise to further generalization:

Corollary 1.9. (Corollary 7.26) Let C be a small ∞ -category. There is a canonical equivalence of ∞ -categories

LaxFun(N(
$$\mathcal{C}^{op}$$
), CORR) $\simeq \infty Cat_{\mathcal{C}}$,

where the left hand side is the ∞ -category of lax normal maps of double ∞ -categories $\mathbb{C}^{\mathrm{op}} \to \mathrm{CORR}$.

2. NOTATION AND TERMINOLOGY

We fix a hierarchy of Grothendieck universes whose objects we call small, large, very large, etc. We call a space small, large, etc. if its set of path components and its homotopy groups are for any choice of base point. We call an ∞ -category small, large, etc. if its maximal subspace and all its mapping spaces are.

We write

- Set for the category of small sets.
- Δ for (a skeleton of) the category of finite, non-empty, partially ordered sets and order preserving maps, whose objects we denote by $[n] = \{0 < ... < n\}$ for $n \ge 0$.
- S for the ∞ -category of small spaces.
- ∞ Cat for the ∞ -category of small ∞ -categories.
- ∞Cat^{cc} for the ∞-category of large ∞-categories with small colimits and small colimits preserving functors.

We often indicate ∞ -categories of large objects by (-), for example we write $\widehat{S}, \infty \widehat{\mathsf{Cat}}$ for the ∞ -categories of large spaces, ∞ -categories.

For any ∞ -category \mathcal{C} containing objects A, B we write

- $\mathcal{C}(A, B)$ for the space of maps $A \to B$ in \mathcal{C} ,
- $\mathcal{C}_{/A}$ for the ∞ -category of objects over A,
- Ho(C) for its homotopy category,
- C[⊲], C[▷] for the ∞-category arising from C by adding an initial, final object, respectively,
- $\iota(\mathcal{C})$ for the maximal subspace in \mathcal{C} .

Note that $Ho(\infty Cat)$ is cartesian closed and for small ∞ -categories \mathcal{C}, \mathcal{D} we write $Fun(\mathcal{C}, \mathcal{D})$ for the internal hom, the ∞ -category of functors $\mathcal{C} \to \mathcal{D}$.

We often call a fully faithful functor an embedding. We call a functor an inclusion if if it induces an embedding on maximal subspaces and on all mapping spaces.

3. Parametrized ∞-categories of presheaves

Our key tool to prove a local global principle for parametrized ∞ -categories, i.e. ∞ -categories over a given base ∞ -category C, is a parametrized version of presheaf ∞ -category that we construct in this section.

Definition 3.1. A right fibration $\mathcal{A} \to \mathcal{B}$ is representable if \mathcal{A} has a final object.

Remark 3.2. A right fibration $\mathcal{A} \to \mathcal{B}$ is representable if and only if $\mathcal{A} \to \mathcal{B}$ classifies a representable presheaf.

Notation 3.3. Let

$$\mathcal{R} \subset \operatorname{Fun}([1], \infty \operatorname{Cat})$$

be the full subcategory of right fibrations and

 $\mathcal{U} \subset \mathcal{R}$

the full subcategory of representable right fibrations.

Lemma 3.4. Evaluation at the target $\rho : \mathbb{R} \to \infty$ Cat is a cocartesian and cartesian fibration that restricts to a cocartesian fibration $\kappa : \mathbb{U} \to \infty$ Cat with the same cocartesian morphisms.

Proof. The functor γ : Fun([1], ∞ Cat) $\rightarrow \infty$ Cat evaluating at the target is a cartesian fibration because ∞ Cat admits pullbacks. Since right fibrations are stable under pullback, γ restricts to a cartesian fibration $\rho: \mathcal{R} \rightarrow \infty$ Cat with the same cartesian morphisms. For any functor $\psi: \mathcal{C} \rightarrow \mathcal{D}$ the fiber transport $\mathcal{R}_{\mathcal{D}} \rightarrow \mathcal{R}_{\mathcal{C}}$ of $\mathcal{R} \rightarrow \infty$ Cat admits a left adjoint $\psi_{!}$ that takes the left Kan extension along the functor $\mathcal{C}^{\mathrm{op}} \rightarrow \mathcal{D}^{\mathrm{op}}$. This guarantees that the cartesian fibration $\rho: \mathcal{R} \rightarrow \infty$ Cat is also a cocartesian fibration. Since $\psi_{!}$ preserves representables, the cocartesian fibration $\rho: \mathcal{R} \rightarrow \infty$ Cat restricts to a cocartesian fibration $\kappa: \mathcal{U} \rightarrow \infty$ Cat with the same cocartesian morphisms.

Definition 3.5. For every cocartesian fibration $\mathcal{D} \to \mathcal{C}$ classifying a functor $\psi : \mathcal{C} \to \infty \mathsf{Cat}$ let $\mathcal{P}^{\mathcal{C}}(\mathcal{D}) \to \mathcal{C}$ be the pullback of $\rho : \mathcal{R} \to \infty \mathsf{Cat}$ along $\psi : \mathcal{C} \to \infty \mathsf{Cat}$

Remark 3.6. Evaluation at the target $\mathcal{R} \to \infty Cat$ is a cocartesian and cartesian fibration by Lemma 3.4. So for every cocartesian fibration $\mathcal{D} \to \mathcal{C}$ whose fibers are small, the functor $\mathcal{P}^{\mathcal{C}}(\mathcal{D}) \to \mathcal{C}$ is a cocartesian and cartesian fibration.

Lemma 3.7. (1) The cocartesian fibration $\kappa : \mathfrak{U} \to \infty$ Cat classifies the identity.

By (1) for every cocartesian fibration $\mathbb{D} \to \mathbb{C}$ classifying a functor $\mathbb{C} \to \infty$ Cat there is a canonical embedding of cocartesian fibrations over \mathbb{C} :

$$\mathcal{D} \simeq \mathcal{C} \times_{\infty \mathsf{Cat}} \mathcal{U} \subset \mathcal{P}^{\mathsf{C}}(\mathcal{D}) = \mathcal{C} \times_{\infty \mathsf{Cat}} \mathcal{R}.$$

(2) The embedding

$$\mathcal{D} \simeq \mathcal{C} \times_{\infty \mathsf{Cat}} \mathcal{U} \subset \mathcal{P}^{\mathfrak{C}}(\mathcal{D}) = \mathcal{C} \times_{\infty \mathsf{Cat}} \mathcal{R} \subset \mathcal{C} \times_{\mathrm{Fun}(\{1\},\infty \mathsf{Cat})} \mathrm{Fun}([1],\infty \mathsf{Cat})$$

corresponds to a natural transformation of functors $\mathbb{D} \to \infty \mathsf{Cat}$ that is classified by the canonical map $\mathbb{D}^{[1]} \to \mathbb{D} \times_{\mathbb{C}} \mathbb{D}$ of cocartesian fibrations over \mathbb{D} .

Proof. (1): Let $\mathcal{V} \to \infty \mathsf{Cat}$ be the cocartesian fibration classifying the identity. We will construct an equivalence $\mathcal{V} \simeq \mathcal{U}$ over $\infty \mathsf{Cat}$ by naturally identifying for any ∞ -category \mathcal{A} the set of equivalence classes of functors $\mathcal{A} \to \mathcal{V}$ over $\infty \mathsf{Cat}$ and $\mathcal{A} \to \mathcal{U}$ over $\infty \mathsf{Cat}$.

A functor $\mathcal{A} \to \mathcal{V}$ over $\infty \mathsf{Cat}$ is classified by a cocartesian fibration $\mathcal{C} \to \mathcal{A}$ equipped with a section of $\mathcal{C} \to \mathcal{A}$. A functor $\mathcal{A} \to \mathcal{U} \subset \operatorname{Fun}([1], \infty \mathsf{Cat})$ over $\infty \mathsf{Cat}$ is classified by a map of cocartesian fibrations $\mathcal{D} \to \mathcal{C}$ over \mathcal{A} that induces on the fiber over any $\mathcal{A} \in \mathcal{A}$ a representable right fibration. A section α of $\mathcal{C} \to \mathcal{A}$ gives rise to a map $\mathcal{A} \times_{\mathcal{C}^{\{1\}}} \mathcal{C}^{\{1\}} \to \mathcal{C}^{\{0\}}$ of cocartesian fibrations over \mathcal{A} that induces on the fiber over any $A \in \mathcal{A}$ the representable right fibration $(\mathcal{C}_A)_{/\alpha(A)} \to \mathcal{C}_A$ and induces on sections the functor $\operatorname{Fun}_{\mathcal{A}}(\mathcal{A}, \mathcal{C})_{/\alpha} \to \operatorname{Fun}_{\mathcal{A}}(\mathcal{A}, \mathcal{C})$, which is an equivalence if and only if α is a final object in $\operatorname{Fun}_{\mathcal{A}}(\mathcal{A}, \mathcal{C})$. This is the case if for any $A \in \mathcal{A}$ the image $\alpha(A)$ is final in \mathcal{C}_A .

Let $\psi : \mathcal{D} \to \mathbb{C}$ be a map of cocartesian fibrations over \mathcal{A} such that for any $A \in \mathcal{A}$ the fiber \mathcal{D}_A has a final object. Then the ∞ -category $\operatorname{Fun}_{\mathcal{A}}(\mathcal{A}, \mathcal{D})$ has a final object β such that for any $A \in \mathcal{A}$ the image $\beta(A)$ is final in \mathcal{D}_A . So we get a section $\psi \circ \beta$ of $\mathbb{C} \to \mathcal{A}$. For $\mathcal{D} = \mathcal{A} \times_{\mathbb{C}^{\{1\}}} \mathbb{C}^{[1]}$ we have $\operatorname{Fun}_{\mathcal{A}}(\mathcal{A}, \mathcal{D}) \simeq \operatorname{Fun}_{\mathcal{A}}(\mathcal{A}, \mathbb{C})_{/\alpha}$ over $\operatorname{Fun}_{\mathcal{A}}(\mathcal{A}, \mathbb{C})$ so that $\psi \circ \beta = \alpha$. In general the induced map $\mathcal{A} \times_{\mathcal{D}^{\{1\}}} \mathcal{D}^{[1]} \to \mathcal{A} \times_{\mathbb{C}^{\{1\}}} \mathbb{C}^{[1]}$ of cocartesian fibrations over \mathcal{A} induces on the fiber over any $A \in \mathcal{A}$ the equivalence $(\mathcal{D}_A)_{/\beta(A)} \simeq (\mathbb{C}_A)_{/\alpha(A)}$ (coming from the fact that $\mathcal{D}_A \to \mathbb{C}_A$ is a right fibration). Hence the induced map $\mathcal{D} \simeq \mathcal{A} \times_{\mathcal{D}^{\{1\}}} \mathcal{D}^{[1]} \to \mathcal{A} \times_{\mathbb{C}^{\{1\}}} \mathbb{C}^{[1]}$ of cocartesian fibrations over \mathcal{A} is an equivalence. So $\mathcal{D} \to \mathbb{C}$ is equivalent over \mathbb{C} to $\mathcal{A} \times_{\mathbb{C}^{\{1\}}} \mathbb{C}^{[1]} \to \mathbb{C}^{\{0\}}$. The proof of (1) shows (2).

Definition 3.8. Let $\mathcal{C} \to \mathcal{A}$ be a functor. The enveloping cocartesian fibration of $\mathcal{C} \to \mathcal{A}$ is the pullback

$$\operatorname{Env}(\mathcal{C}) \coloneqq \operatorname{Fun}([1], \mathcal{A}) \times_{\operatorname{Fun}(\{0\}, \mathcal{A})} \mathcal{C} \to \operatorname{Fun}(\{1\}, \mathcal{A}).$$

The diagonal embedding $\mathcal{A} \to \operatorname{Fun}([1], \mathcal{A})$ induces an embedding $\theta : \mathcal{C} \to \operatorname{Env}(\mathcal{C})$ over \mathcal{A} . The following proposition is [9, Proposition A.2.]:

Proposition 3.9. Let $\mathcal{C} \to \mathcal{A}$ be a functor. Restriction along θ induces for every cocartesian fibration $\mathcal{D} \to \mathcal{A}$ an equivalence

$$\operatorname{Fun}_{\mathcal{A}}^{\operatorname{cocart}}(\operatorname{Env}(\mathcal{C}), \mathcal{D}) \to \operatorname{Fun}_{\mathcal{A}}(\mathcal{C}, \mathcal{D}),$$

where the left hand side is the full subcategory of maps of cocartesian fibrations over A.

Lemma 3.10. Let $\mathcal{C} \to \mathcal{A}$ be a cocartesian fibration between small ∞ -categories. There is a canonical embedding

$$\mathbb{P}(\mathcal{C}) \hookrightarrow \mathcal{P}^{\mathcal{A}}(\operatorname{Env}(\mathcal{C}))$$

over \mathcal{A} that induces on the fiber over every $A \in \mathcal{A}$ the embedding $(\theta_A)_! : \mathcal{P}(\mathcal{C}_A) \subset \mathcal{P}(\operatorname{Env}(\mathcal{C})_A)$.

Proof. By [9, Lemma A.7.] the embedding $\theta : \mathbb{C} \subset \operatorname{Env}(\mathbb{C})$ admits a left adjoint L relative to \mathcal{A} , which is a map of cocartesian fibrations over \mathcal{A} . The relative left adjoint gives rise to a map $L_! : \mathcal{P}^{\mathcal{A}}(\operatorname{Env}(\mathbb{C})) \to \mathcal{P}^{\mathcal{A}}(\mathbb{C})$ of cocartesian fibrations over \mathcal{A} . For every $A \in \mathcal{A}$ the induced localization $L_A : \operatorname{Env}(\mathbb{C})_A \rightleftharpoons \mathbb{C}_A : \theta_A$ induces a localization $(L_!)_A \simeq (L_A)_! :$ $\mathcal{P}(\operatorname{Env}(\mathbb{C})_A) \rightleftharpoons \mathcal{P}^{\mathcal{C}}(\mathbb{C})$ of cocartesian fibrations over \mathcal{A} admits the map $L_! : \mathcal{P}^{\mathcal{A}}(\operatorname{Env}(\mathbb{C})) \to \mathcal{P}^{\mathcal{A}}(\mathbb{C})$ of cocartesian fibrations over \mathcal{A} admits a fully faithful right adjoint relative to \mathcal{A} that induces on the fiber over $A \in \mathcal{A}$ the embedding $(\theta_A)_!$.

Lemma 3.10 motivates the following extension of Definition 3.5.

Definition 3.11. For every functor $\mathcal{C} \to \mathcal{A}$ between small ∞ -categories let

$$\mathcal{P}^{\mathcal{A}}(\mathcal{C}) \subset \mathcal{P}^{\mathcal{A}}(\operatorname{Env}(\mathcal{C}))$$

be the full subcategory spanned by the objects of $\mathcal{P}^{\mathcal{A}}(\operatorname{Env}(\mathcal{C}))_{A} \simeq \mathcal{P}(\operatorname{Env}(\mathcal{C})_{A})$ that belong to the essential image of the embedding $(\theta_{A})_{!} : \mathcal{P}(\mathcal{C}_{A}) \subset \mathcal{P}(\operatorname{Env}(\mathcal{C})_{A})$ for some $A \in \mathcal{A}$.

Remark 3.12. The embedding $\mathcal{C} \subset \operatorname{Env}(\mathcal{C}) \to \mathcal{P}^{\mathcal{A}}(\operatorname{Env}(\mathcal{C}))$ over \mathcal{A} of Lemma 3.7 induces an embedding $\mathcal{C} \to \mathcal{P}^{\mathcal{A}}(\mathcal{C})$ over \mathcal{A} .

Lemma 3.13. Let $\phi : \mathbb{D} \to \mathcal{A}$ be a locally cocartesian fibration and $\mathbb{C} \subset \mathbb{D}$ a full subcategory such that for every $A \in \mathcal{A}$ the induced embedding $\mathbb{C}_A \subset \mathbb{D}_A$ admits a left adjoint L_A . The restriction $\phi' : \mathbb{C} \subset \mathbb{D} \to \mathcal{A}$ is a locally cocartesian fibration. A morphism of \mathbb{C} lying over a morphism $f : A \to B$ of \mathcal{A} is locally ϕ' -cocartesian if and only if it factors as $X \to Y \to Y'$, where $X \to Y$ is ϕ -cocartesian and $Y \to Y'$ is a L_B -local equivalence.

Proof. For every $Z \in \mathcal{C}$ lying over $B \in \mathcal{A}$ the canonical map

$$\mathcal{C}_{B}(Y',Z) \simeq \mathcal{D}_{B}(Y,Z) \rightarrow \{f\} \times_{\mathcal{A}(A,B)} \mathcal{D}(X,Z)$$

is an equivalence.

Proposition 3.14. For every functor $\mathbb{C} \to \mathcal{A}$ of small ∞ -categories the functor $\mathbb{P}^{\mathcal{A}}(\mathbb{C}) \to \mathcal{A}$ is a locally cartesian fibration. For every morphism $A \to B$ of \mathcal{A} the induced functor $\mathbb{P}^{\mathcal{A}}(\mathbb{C})_B \to \mathbb{P}^{\mathcal{A}}(\mathbb{C})_A$ factors as

$$\mathcal{P}(\mathcal{C}_{\mathrm{B}}) \subset \mathcal{P}(\mathrm{Env}(\mathcal{C})_{\mathrm{B}}) \to \mathcal{P}(\mathcal{C}_{\mathrm{A}}),$$

where the last functor is restriction along the functor $\mathcal{C}_A \subset \operatorname{Env}(\mathcal{C})_A \to \operatorname{Env}(\mathcal{C})_B$, and thus is a left adjoint.

Proof. The functor $\mathcal{P}^{\mathcal{A}}(\operatorname{Env}(\mathbb{C})) \to \mathcal{A}$ is a cartesian fibration and for every $A \in \mathcal{A}$ the induced embedding $(\theta_A)_! : \mathcal{P}^{\mathcal{A}}(\mathbb{C})_A \simeq \mathcal{P}(\mathbb{C}_A) \subset \mathcal{P}^{\mathcal{A}}(\operatorname{Env}(\mathbb{C}))_A \simeq \mathcal{P}(\operatorname{Env}(\mathbb{C})_A)$ admits a right adjoint. Thus by Lemma 3.13 the restriction $\mathcal{P}^{\mathcal{A}}(\mathbb{C}) \subset \mathcal{P}^{\mathcal{A}}(\operatorname{Env}(\mathbb{C})) \to \mathcal{A}$ is a locally cartesian fibration.

Notation 3.15. Let $\mathcal{C} \to \mathcal{A}, \mathcal{D} \to \mathcal{B}$ be functors. Let

$$\Theta(\mathcal{C}, \mathcal{D}) \coloneqq \operatorname{Fun}(\mathcal{A}, \mathcal{B}) \times_{\operatorname{Fun}(\mathcal{C}, \mathcal{B})} \operatorname{Fun}(\mathcal{C}, \mathcal{D})$$

and

$$\Theta(\mathcal{C},\mathcal{D})^{\mathrm{L}} \subset \Theta(\mathcal{C},\mathcal{D})$$

the full subcategory spanned by the commutative squares



such that for every $A \in \mathcal{A}$ the induced functor $\mathcal{C}_A \to \mathcal{D}_{\psi(A)}$ on the fiber over A preserves small colimits.

The following universal property of $\mathcal{P}^{\mathcal{A}}(\mathcal{C}) \to \mathcal{A}$ is crucial for our main theorem:

Proposition 3.16. Let $\mathcal{D} \to \mathcal{B}$ be a locally cartesian fibration whose fibers admit small colimits.

(1) Let $\mathfrak{C} \to \mathcal{A}$ be a functor between small ∞ -categories. Restriction along the embedding $\mathfrak{C} \to \mathfrak{P}^{\mathcal{A}}(\mathfrak{C})$ induces an equivalence

$$\theta_{\mathcal{D}}: \Theta(\mathcal{P}^{\mathcal{A}}(\mathcal{C}), \mathcal{D})^{\mathrm{L}} \to \Theta(\mathcal{C}, \mathcal{D}).$$

(2) Let $\mathbb{C} \to \mathcal{A}$ be a functor between ∞ -categories and $\mathbb{P}^{\mathcal{A}}(\mathbb{C})$ the full subcategory of $\widehat{\mathbb{P}}^{\mathcal{A}}(\mathbb{C})$ such that for every $A \in \mathcal{A}$ the fiber $\mathbb{P}^{A}(\mathbb{C})_{A} \subset \widehat{\mathbb{P}}^{\mathcal{A}}(\mathbb{C})_{A} \simeq \widehat{\mathbb{P}}(\mathbb{C}_{A})$ is generated by \mathbb{C}_{A} under small colimits. Restriction along the induced embedding $\mathbb{C} \to \mathbb{P}^{\mathcal{A}}(\mathbb{C}) \subset \widehat{\mathbb{P}}^{\mathcal{A}}(\mathbb{C})$ induces an equivalence

$$\theta_{\mathcal{D}}: \Theta(\mathcal{P}^{\mathcal{A}}(\mathcal{C}), \mathcal{D})^{\mathrm{L}} \to \Theta(\mathcal{C}, \mathcal{D}).$$

Proof. (2) is similar to (1). We prove (1). By the opposite of Lemma 3.17 (applied to a larger universe) there is an embedding $\mathcal{D} \subset \mathcal{E}$ over \mathcal{B} that induces on the fiber over every object of \mathcal{A} a small colimits preserving functor, where $\mathcal{E} \to \mathcal{A}$ is a cocartesian and cartesian fibration whose fibers admit large colimits and are locally large. Since the ∞ -category of presheaves is generated by the representable presheaves under small colimits [8, Lemma 2.10.], the functor $\theta_{\mathcal{D}}$ is the pullback of the functor $\theta_{\mathcal{E}}$. So it is enough to check that $\theta_{\mathcal{E}}$ is an equivalence and we can assume that $\mathcal{D} \to \mathcal{A}$ is a cocartesian and cartesian fibration whose fibers admit large colimits and are locally large.

Since $\mathcal{D} \to \mathcal{B}$ is a cocartesian and cartesian fibration, the functor $\theta_{\mathcal{D}}$ is a map of cocartesian and cartesian fibrations over Fun(\mathcal{A}, \mathcal{B}). Consequently, it is enough to see that $\theta_{\mathcal{D}}$ induces on the fiber over every functor $\psi : \mathcal{A} \to \mathcal{B}$ an equivalence. The functor $\theta_{\mathcal{D}}$ induces on the fiber over ψ the canonical functor

$$\operatorname{Fun}_{\mathcal{A}}^{\operatorname{L}}(\mathcal{P}^{\mathcal{A}}(\mathcal{C}), \mathcal{A} \times_{\mathcal{B}} \mathcal{D}) \to \operatorname{Fun}_{\mathcal{A}}(\mathcal{C}, \mathcal{A} \times_{\mathcal{B}} \mathcal{D}),$$

where the left hand side is the full subcategory of functors over \mathcal{A} that induce on every fiber a small colimits preserving functor. Therefore it is enough to see that for every cocartesian and cartesian fibration $\mathcal{D} \to \mathcal{A}$ whose fibers admit large colimits and are locally large the canonical functor

$$\kappa : \operatorname{Fun}_{\mathcal{A}}^{\operatorname{L}}(\mathcal{P}^{\mathcal{A}}(\mathcal{C}), \mathcal{D}) \to \operatorname{Fun}_{\mathcal{A}}(\mathcal{C}, \mathcal{D})$$

is an equivalence. The functor κ is conservative since for every $A \in \mathcal{A}$ the ∞ -category $\mathcal{P}^{\mathcal{A}}(\mathbb{C})_A \simeq \mathcal{P}(\mathbb{C}_A)$ is generated by \mathcal{C}_A under small colimits. Thus it is enough to check that κ admits a fully faithful left adjoint. By Proposition 3.14 the functor $\mathcal{P}^{\mathcal{A}}(\mathbb{C}) \to \mathcal{A}$ is a locally cartesian fibration. This implies that $\mathcal{P}^{\mathcal{A}}(\mathbb{C})$ is large because \mathcal{A} is large and the fibers of $\mathcal{P}^{\mathcal{A}}(\mathbb{C}) \to \mathcal{A}$ are large. Since $\mathcal{P}^{\mathcal{A}}(\mathbb{C})$ is large and the fibers of the cocartesian fibration $\mathcal{D} \to \mathcal{A}$ admit large colimits preserved by the fiber transports, by [19, Corollary 4.3.2.14.] and [19, Proposition 4.3.2.17.] the induced functor

$$\operatorname{Fun}_{\mathcal{A}}(\mathcal{P}^{\mathcal{A}}(\mathcal{C}), \mathcal{D}) \to \operatorname{Fun}_{\mathcal{A}}(\mathcal{C}, \mathcal{D})$$

admits a fully faithful left adjoint ι taking the left Kan extension relative to \mathcal{A} along the embedding $\mathcal{C} \to \mathcal{P}^{\mathcal{A}}(\mathcal{C})$. We complete the proof by showing that the left Kan extension $F: \mathcal{P}^{\mathcal{A}}(\mathcal{C}) \to \mathcal{D}$ relative to \mathcal{A} of any functor $\mathcal{C} \to \mathcal{D}$ over \mathcal{A} preserves fiberwise small colimits. Because $\mathcal{P}^{\mathcal{A}}(\operatorname{Env}(\mathcal{C}))$ is large, the induced functor

$$\operatorname{Fun}_{\mathcal{A}}(\mathcal{P}^{\mathcal{A}}(\operatorname{Env}(\mathcal{C})), \mathcal{D}) \to \operatorname{Fun}_{\mathcal{A}}(\mathcal{P}^{\mathcal{A}}(\mathcal{C}), \mathcal{D})$$

admits a fully faithful left adjoint β . Since the embedding $\mathcal{P}^{\mathcal{A}}(\mathcal{C}) \subset \mathcal{P}^{\mathcal{A}}(\operatorname{Env}(\mathcal{C}))$ yields on the fiber over every object of \mathcal{A} a left adjoint functor, it is enough to see that $\beta(F)$ preserves fiberwise small colimits. Since $\beta(F) \simeq \beta(\iota(F_{|\mathcal{C}}))$ is the left Kan extension relative to \mathcal{A} of $F_{|\mathcal{C}} : \mathcal{C} \to \mathcal{D}$ along the embedding $\mathcal{C} \subset \operatorname{Env}(\mathcal{C}) \to \mathcal{P}^{\mathcal{A}}(\operatorname{Env}(\mathcal{C}))$ over \mathcal{A} , it is enough to check that the left Kan extension $\overline{H} : \mathcal{P}^{\mathcal{A}}(\operatorname{Env}(\mathcal{C})) \to \mathcal{D}$ relative to \mathcal{A} of any functor $H : \operatorname{Env}(\mathcal{C}) \to \mathcal{D}$ over \mathcal{A} preserves fiberwise small colimits. Because $\operatorname{Env}(\mathcal{C}) \subset \mathcal{P}^{\mathcal{A}}(\operatorname{Env}(\mathcal{C}))$ is a map of cocartesian fibrations over \mathcal{A} , by [19, Proposition 4.3.3.10.] for every $A \in \mathcal{A}$ the induced functor

$$\overline{\mathrm{H}}_{\mathrm{A}}: \mathcal{P}^{\mathcal{A}}(\mathrm{Env}(\mathcal{C}))_{\mathrm{A}} \simeq \mathcal{P}(\mathrm{Env}(\mathcal{C})_{\mathrm{A}}) \to \mathcal{D}_{\mathrm{A}}$$

is the left Kan extension of $H_A : Env(\mathcal{C})_A \to \mathcal{D}_A$, which by [19, Lemma 5.1.5.5.] preserves small colimits.

Lemma 3.17. Let $\mathcal{D} \to \mathcal{A}$ be a locally cocartesian fibration of small ∞ -categories. There is an embedding $\mathcal{D} \subset \mathcal{E}$ over \mathcal{A} that induces on the fiber over every object of \mathcal{A} a small limits

preserving functor, where $\mathcal{E} \to \mathcal{A}$ is a cocartesian and cartesian fibration whose fibers admit small limits and are locally small.

Proof. Since $\mathcal{D} \to \mathcal{A}$ is a locally cocartesian fibration, by [9, Lemma A.7.] the embedding $\mathcal{D} \subset \operatorname{Env}(\mathcal{D})$ induces on the fiber over every object of \mathcal{A} a right adjoint functor. Since $\mathcal{D} \to \mathcal{A}$ is a locally cocartesian fibration of small ∞ -categories, also \mathcal{D} is small. Take $\mathcal{E} := \mathcal{P}^{\mathcal{A}}(\operatorname{Env}(\mathcal{D}))$. The embedding $\mathcal{D} \subset \operatorname{Env}(\mathcal{D}) \to \mathcal{P}^{\mathcal{A}}(\operatorname{Env}(\mathcal{D}))$ induces on every fiber a small limits preserving functor.

Remark 3.18. Proposition 3.16 gives for any functors $\mathcal{A}' \to \mathcal{A}, \mathcal{C} \to \mathcal{A}$ between small ∞ -categories a canonical functor

(2)
$$\mathcal{P}^{\mathcal{A}'}(\mathcal{A}' \times_{\mathcal{A}} \mathcal{C}) \to \mathcal{A}' \times_{\mathcal{A}} \mathcal{P}^{\mathcal{A}}(\mathcal{C})$$

over \mathcal{A}' , which is a map of locally cartesian fibrations over \mathcal{A}' , and induces equivalences on fibers and so is an equivalence.

Lemma 3.19. Let $\mathbb{C} \to \mathcal{A}$ be a locally cartesian fibration between small ∞ -categories. The embedding $\mathbb{C} \subset \mathbb{P}^{\mathcal{A}}(\mathbb{C})$ is a map of locally cartesian fibrations over \mathcal{A} . The functor $\mathbb{C} \to \mathcal{A}$ is a cartesian fibration if and only if the functor $\mathbb{P}^{\mathcal{A}}(\mathbb{C}) \to \mathcal{A}$ is a cartesian fibration.

Proof. We first prove that the embedding $C \subset \mathcal{P}^{\mathcal{A}}(\mathcal{C})$ is a map of locally cartesian fibrations over \mathcal{A} if $\mathcal{C} \to \mathcal{A}$ is a locally cartesian fibration. By Remark 3.18 for every functor $\mathcal{A}' \to \mathcal{A}$ there is a canonical equivalence

$$\mathcal{A}' \times_{\mathcal{A}} \mathcal{P}^{\mathcal{A}}(\mathcal{C}) \simeq \mathcal{P}^{\mathcal{A}'}(\mathcal{A}' \times_{\mathcal{A}} \mathcal{C})$$

over \mathcal{A}' so that we can reduce to the case $\mathcal{A} = [1]$. Let $\mathcal{M} \to [1]$ be a functor and $\mathcal{C} := \mathcal{M}_0, \mathcal{D} := \mathcal{M}_1$ the fibers. The functor $\mathcal{P}(\mathcal{M})^{[1]} \to [1]$ is a cartesian fibration classifying the functor $\phi : \mathcal{P}(\mathcal{D}) \subset \mathcal{P}(\mathcal{M}) \to \mathcal{P}(\mathcal{C})$, where the first functor takes left Kan extension along the embedding $\mathcal{D} \subset \mathcal{M}$ and the second functor restricts along the embedding $\mathcal{C} \subset \mathcal{M}$. If $\mathcal{M} \to [1]$ is a cartesian fibration classifying a functor $G : \mathcal{D} \to \mathcal{C}$, then ϕ is the extension of the functor $\mathcal{D} \xrightarrow{G} \mathcal{C} \subset \mathcal{P}(\mathcal{C})$. So the embedding $\mathcal{M} \subset \mathcal{P}(\mathcal{M})^{[1]}$ is a map of cartesian fibrations over [1].

A locally cartesian fibration ϕ is a cartesian fibration if and only if the collection of ϕ cartesian morphisms is closed under composition. Hence $\mathbb{C} \to \mathcal{A}$ is a cartesian fibration if the functor $\mathcal{P}^{\mathcal{A}}(\mathbb{C}) \to \mathcal{A}$ is a cartesian fibration. Conversely, if $\mathbb{C} \to \mathcal{A}$ is a cartesian fibration, the functor $\mathcal{P}^{\mathcal{A}}(\mathbb{C}) \to \mathcal{A}$ is a cartesian fibration since the fibers of $\mathcal{P}^{\mathcal{A}}(\mathbb{C}) \to \mathcal{A}$ are generated by the representable presheaves under small colimits, and the fiber transports of $\mathcal{P}^{\mathcal{A}}(\mathbb{C}) \to \mathcal{A}$ preserve small colimits.

Lemma 3.20. Let $\mathbb{C} \to \mathbb{D}$ be a map of locally (co)cartesian fibrations over \mathcal{A} . The functor $\mathbb{P}^{\mathcal{A}}(\mathbb{C}) \to \mathbb{P}^{\mathcal{A}}(\mathbb{D})$ is a map of locally (co)cartesian fibrations over \mathcal{A} .

Proof. By Lemma 3.19 for any locally cartesian fibration $\mathcal{C} \to \mathcal{A}$ the functor $\mathcal{P}^{\mathcal{A}}(\mathcal{C}) \to \mathcal{A}$ is a locally cartesian fibration and the embedding $\mathcal{C} \to \mathcal{P}^{\mathcal{A}}(\mathcal{C})$ is a map of locally cartesian fibrations over \mathcal{A} . By Lemma 3.10 and Remark 3.18 for every locally cocartesian fibration $\mathcal{C} \to \mathcal{A}$ the functor $\mathcal{P}^{\mathcal{A}}(\mathcal{C}) \to \mathcal{A}$ is a locally cocartesian and locally cartesian fibration and the embedding $\mathcal{C} \to \mathcal{P}^{\mathcal{A}}(\mathcal{C})$ is a map of locally cocartesian fibrations over \mathcal{A} . Since the fibers of $\mathcal{P}^{\mathcal{A}}(\mathcal{C}) \to \mathcal{A}$ are generated under small colimits by the representables and the fiber transports of the locally (co)cartesian fibration $\mathcal{P}^{\mathcal{A}}(\mathcal{C}) \to \mathcal{A}$ preserve small colimits by Proposition 3.14, the induced functor $\mathcal{P}^{\mathcal{A}}(\mathcal{C}) \to \mathcal{P}^{\mathcal{A}}(\mathcal{D})$ is a map of locally (co)cartesian fibrations over \mathcal{A} . \Box **Definition 3.21.** A functor $\phi : \mathcal{D} \to \mathcal{C}$ of small ∞ -categories is an exponential fibration if the functor $\mathcal{D} \times_{\mathcal{C}} (-) : \infty \mathsf{Cat}_{/\mathcal{C}} \to \infty \mathsf{Cat}_{/\mathcal{C}}$ taking pullback along ϕ preserves small colimits or equivalently admits a right adjoint.

Notation 3.22. For every exponential fibration $\phi : \mathcal{D} \to \mathcal{C}$ let $\operatorname{Fun}^{\mathcal{C}}(\mathcal{D}, -) : \infty \operatorname{Cat}_{/\mathcal{C}} \to \infty \operatorname{Cat}_{/\mathcal{C}}$ be the right adjoint of the functor $\mathcal{D} \times_{\mathcal{C}} (-) : \infty \operatorname{Cat}_{/\mathcal{C}} \to \infty \operatorname{Cat}_{/\mathcal{C}}$.

The next remark is [10, Remark 3.71.]:

Remark 3.23. The pullback of any exponential fibration $\mathcal{D} \to \mathcal{C}$ along any functor $\mathcal{B} \to \mathcal{C}$ is an exponential fibration and for every functor $\mathcal{E} \to \mathcal{C}$ there is a canonical equivalence $\mathcal{B} \times_{\mathcal{C}} \operatorname{Fun}^{\mathcal{C}}(\mathcal{D}, \mathcal{E}) \simeq \operatorname{Fun}^{\mathcal{B}}(\mathcal{B} \times_{\mathcal{C}} \mathcal{D}, \mathcal{B} \times_{\mathcal{C}} \mathcal{E})$ over \mathcal{B} .

Proposition 3.24. A functor $\mathbb{C} \to \mathcal{A}$ of small ∞ -categories is an exponential fibration if and only if the locally cartesian fibration $\mathbb{P}^{\mathcal{A}}(\mathbb{C}) \to \mathcal{A}$ is a cartesian fibration.

Proof. A functor $\mathcal{B} \to \mathcal{A}$ is an exponential fibration (cartesian fibration) if and only if the pullbacks along any functors $[2] \to \mathcal{A}$ are exponential fibrations (cartesian fibrations). So by Remark 3.18 we can reduce to the case that $\mathcal{A} = [2]$. The locally cartesian fibration $\mathcal{P}^{[2]}(\mathcal{C}) \to [2]$ classifies three functors $f : \mathcal{P}(\mathcal{C}_2) \to \mathcal{P}(\mathcal{C}_1), g : \mathcal{P}(\mathcal{C}_1) \to \mathcal{P}(\mathcal{C}_0), h : \mathcal{P}(\mathcal{C}_2) \to \mathcal{P}(\mathcal{C}_0)$ and a natural transformation $\alpha : g \circ f \to h$. By [18, Proposition B.3.2.] a functor $\mathcal{C} \to [2]$ is an exponential fibration if and only if α is an equivalence. By [19, Proposition 2.4.2.8.] the locally cartesian fibration $\mathcal{P}^{\mathcal{A}}(\mathcal{C}) \to \mathcal{A}$ is a cartesian fibration if and only if α is an equivalence.

Corollary 3.25. Every functor $\mathcal{D} \to [1]$ is an exponential fibration. So the ∞ -category $\infty \operatorname{Cat}_{I[1]}$ is cartesian closed.

Corollary 3.26. Let $\phi : \mathcal{A} \to \mathcal{B}$ be an inclusion of ∞ -categories such that for every morphisms $X \to Y, Y \to Z$ we have that $Y \in \mathcal{A}$ if $X, Z \in \mathcal{A}$. Then ϕ is an exponential fibration.

Proof. By Proposition 3.24 the functor ϕ is an exponential fibration if and only if the locally cartesian fibration $\mathcal{P}^{\mathcal{B}}(\mathcal{A}) \to \mathcal{B}$ is a cartesian fibration. By [19, Proposition 2.4.2.8.] the locally cartesian fibration $\mathcal{P}^{\mathcal{B}}(\mathcal{A}) \to \mathcal{B}$ is a cartesian fibration if and only if for every morphisms $\alpha : X \to Y, \beta : Y \to Z$ the induced natural transformation $\theta : \alpha^* \circ \beta^* \to (\beta \circ \alpha)^* : \mathcal{P}(\mathcal{A}_Z) \to \mathcal{P}(\mathcal{A}_X)$ is an equivalence. If X does not belong to \mathcal{A} , the fiber of ϕ over X is empty so that $\mathcal{P}(\mathcal{A}_X)$ is contractible. So there is nothing to show and we assume that X belongs to \mathcal{A} so that $\mathcal{P}(\mathcal{A}_X) \simeq S$. If Z does not belong to \mathcal{A} , the fiber of ϕ over Z is empty so that $\mathcal{P}(\mathcal{A}_X) \cong S$. If Z does not belong to show, too, because α^*, β^* preserve small colimits and so the initial object so that θ is an endomorphism of the initial object in $\mathcal{P}(\mathcal{A}_X) \simeq S$ and so an equivalence. So we can assume that Z belongs to \mathcal{A} . By assumption we find that Y belongs to \mathcal{A} . In this case the small colimits preserving functors $\alpha^*, \beta^* : S \to S$ are the identity on objects and so preserve the final object. This implies that $\alpha^*, \beta^* : S \to S$ are equivalences and θ is an equivalence since the component of θ at the final space is an endomorphism of the final space of the final space is an endomorphism of the final space.

Definition 3.27. A map $[n] \rightarrow [m]$ of Δ is inert if it is of the form $[n] \cong \{i+0, ..., i+n\} \subset [m]$ for some $i \ge 0$.

Corollary 3.28. Every inert map $[n] \rightarrow [m]$ of Δ (viewed as a functor of ∞ -categories) is an exponential fibration.

Corollary 3.29. A locally cartesian fibration of small ∞ -categories is a cartesian fibration if and only if it is an exponential fibration.

Proof. Let $\mathcal{C} \to \mathcal{A}$ be a locally cartesian fibration of small ∞ -categories. By Lemma 3.19 the functor $\mathcal{C} \to \mathcal{A}$ is a cartesian fibration if and only if the functor $\mathcal{P}^{\mathcal{A}}(\mathcal{C}) \to \mathcal{A}$ is a cartesian fibration. By Proposition 3.24 the functor $\mathcal{C} \to \mathcal{A}$ is an exponential fibration if and only if the functor $\mathcal{P}^{\mathcal{A}}(\mathcal{C}) \to \mathcal{A}$ is a cartesian fibration.

Notation 3.30. Let

12

$$LoCART \subset Fun([1], \infty Cat)$$

be the full subcategory of locally cartesian fibrations.

Notation 3.31. Let

$$LoCART^{L} \subset LoCART$$

be the subcategory of locally cartesian fibrations whose target is small and whose fibers are presentable and commutative squares



such that for every $A \in \mathcal{A}$ the induced functor $\mathcal{C}_A \to \mathcal{D}_{\psi(A)}$ on the fiber over A preserves small colimits.

Corollary 3.32. The inclusion of $(\infty, 2)$ -categories

$$LoCART^{L} \subset Fun([1], \infty \widehat{Cat})$$

admits a left adjoint that sends a functor $\mathcal{C} \to \mathcal{A}$ between small ∞ -categories to $\mathcal{P}^{\mathcal{A}}(\mathcal{C}) \to \mathcal{A}$.

Corollary 3.33. Let \mathcal{A} be a small ∞ -category. The inclusion of $(\infty, 2)$ -categories

$$LoCART^{L}_{\mathcal{A}} \subset \infty \widehat{Cat}_{/\mathcal{A}}$$

admits a left adjoint that sends a functor $\mathcal{C} \to \mathcal{A}$, where \mathcal{C} is small, to $\mathcal{P}^{\mathcal{A}}(\mathcal{C}) \to \mathcal{A}$.

In the following we compare Definition 3.5 to another one (Corollary 3.37).

Notation 3.34. Let $\mathcal{L} \subset \operatorname{Fun}([1], \infty \operatorname{Cat})$ be the full subcategory of left fibrations.

Proposition 3.35. There is a canonical equivalence

$$\mathcal{L} \simeq \operatorname{Fun}^{\infty \mathsf{Cat}}(\mathcal{U}, \mathcal{S} \times \infty \mathsf{Cat})$$

over ∞Cat , where $\mathcal{L} \to \infty Cat$ is evaluation at the target.

Proof. Let $\mathcal{C} \to \infty \mathsf{Cat}$ be a functor classifying a cocartesian fibration $\mathcal{B} \to \mathcal{C}$. We prove that for any ∞-category \mathcal{C} functors $\mathcal{C} \to \mathcal{L}$ over ∞ Cat naturally correspond to functors $\mathcal{C} \to \mathsf{Fun}^{\infty\mathsf{Cat}}(\mathcal{U}, \mathbb{S} \times \infty\mathsf{Cat})$ over ∞ Cat . Functors $\mathcal{C} \to \mathsf{Fun}^{\infty\mathsf{Cat}}(\mathcal{U}, \mathbb{S} \times \mathbb{C})$ over ∞ Cat correspond to sections of the functor $\mathsf{Fun}^{\mathcal{C}}(\mathcal{B}, \mathbb{S} \times \mathcal{C}) \to \mathcal{C}$. Such sections correspond to functors $\mathcal{B} \to \mathbb{S} \times \mathcal{C}$ over \mathcal{C} and so to functors $\mathcal{B} \to \mathbb{S}$. A functor $\mathcal{B} \to \mathcal{L} \subset \mathsf{Fun}([1], \infty\mathsf{Cat})$ over ∞ Cat is classified by a map of cocartesian fibrations $\mathcal{A} \to \mathcal{B}$ over \mathcal{C} that is fiberwise a left fibration and so a left fibration by [19, Proposition 2.4.2.4., Proposition 2.4.2.11.]. Such a functor corresponds to a left fibration $\mathcal{A} \to \mathcal{B}$ classifying a functor $\mathcal{B} \to \mathbb{S}$.

Corollary 3.36. There is a canonical equivalence

$$\mathcal{R} \simeq \operatorname{Fun}^{\infty \mathsf{Cat}}(\mathcal{U}^{\operatorname{rev}}, \mathcal{S} \times \infty \mathsf{Cat})$$

over ∞Cat , where $\mathcal{R} \rightarrow \infty Cat$ is evaluation at the target.

Corollary 3.37. Let $\mathcal{C} \to \mathcal{A}$ be a cocartesian fibration. There is a canonical equivalence

 $\mathcal{P}^{\mathcal{A}}(\mathcal{C}) \simeq \operatorname{Fun}^{\mathcal{A}}(\mathcal{C}^{\operatorname{rev}}, \mathcal{S} \times \mathcal{A})$

over \mathcal{A} .

Remark 3.38. In [4, Example 9.9.] the authors give a different construction of a parametrized ∞ -category of presheaves associated to a cocartesian fibration. Corollary 3.37 implies that our construction of parametrized ∞ -category of presheaves associated to a functor $\phi : \mathcal{C} \to \mathcal{A}$ agrees with the one of [4] in case that ϕ is a cocartesian fibration.

4. A local-global principle

Now we are ready to prove the main theorem.

Theorem 4.1. Let \mathcal{A} be an ∞ -category. The canonical commutative square of ∞ -categories

is a pullback square.

Proof. Let $\mathcal{C} \to \mathcal{A}, \mathcal{D} \to \mathcal{A}$ be functors of small ∞ -categories. Square (3) induces on morphism ∞ -categories the commutative square

which identifies with the commutative square

where we use Remark 3.18. The bottom and top functor of square (4) are fully faithful, which guarantees that square (4) is a pullback square. This implies that the functor

$$\theta: \infty \mathsf{Cat}_{/\mathcal{A}} \to \infty \mathsf{Cat}_{/\iota(\mathcal{A})} \times_{\mathrm{LoCART}_{\iota(\mathcal{A})}^{\mathrm{L}}} \mathrm{LoCART}_{\mathcal{A}}^{\mathrm{L}}$$

is fully faithful. So it remains to see that θ is essentially surjective.

Let $\mathcal{B} \to \iota(\mathcal{A})$ be a functor, $\mathcal{D} \to \mathcal{A}$ a locally cartesian fibration whose fibers admit small colimits and whose fiber transports preserve small colimits, and $\alpha : \mathcal{P}^{\iota(\mathcal{A})}(\mathcal{B}) \simeq \iota(\mathcal{A}) \times_{\mathcal{A}} \mathcal{D}$ an equivalence over $\iota(\mathcal{A})$. Let $\mathcal{C} \subset \mathcal{D}$ be the essential image of the Yoneda-embedding $\mathcal{B} \subset \mathcal{P}^{\iota(\mathcal{A})}(\mathcal{B}) \simeq \iota(\mathcal{A}) \times_{\mathcal{A}} \mathcal{D} \subset \mathcal{D}$. Then there is a canonical equivalence $\iota(\mathcal{A}) \times_{\mathcal{A}} \mathcal{C} \simeq \mathcal{B}$ over $\iota(\mathcal{A})$ and the embedding $\mathcal{C} \subset \mathcal{D}$ over \mathcal{A} extends to a canonical functor $\beta : \mathcal{P}^{\mathcal{A}}(\mathcal{C}) \to \mathcal{D}$ over A, whose pullback to $\iota(\mathcal{A})$ is the equivalence $\alpha : \iota(\mathcal{A}) \times_{\mathcal{A}} \mathcal{P}^{\mathcal{A}}(\mathbb{C}) \simeq \mathcal{P}^{\iota(\mathcal{A})}(\mathfrak{B}) \simeq \iota(\mathcal{A}) \times_{\mathcal{A}} \mathcal{D}$. Therefore it is enough to see that β is an equivalence. For this by Remark 3.18 we can reduce to the case that $\mathcal{A} = [1]$ because a functor over \mathcal{A} is an equivalence if all pullbacks along functors $[1] \to \mathcal{A}$ are equivalences. Since β is essentially surjective, we need to check that for any $X \in \mathcal{P}(\mathbb{C}_0), Y \in \mathcal{P}(\mathbb{C}_1)$ the canonical map $\theta_{X,Y} : \mathcal{P}^{[1]}(\mathbb{C})(X,Y) \to \mathcal{D}(\beta(X),\beta(Y))$ induced by β is an equivalence. Let $G : \mathcal{P}(\mathbb{C}_1) \to \mathcal{P}(\mathbb{C}_0)$ be the left adjoint functor classified by the cartesian fibration $\mathcal{P}^{[1]}(\mathbb{C}) \to [1]$ and $G' : \mathcal{D}_1 \to \mathcal{D}_0$ the left adjoint functor classified by the cartesian fibration $\mathcal{D} \to [1]$. The unique cartesian lift $G(Y) \to Y$, where $G(Y) \in \mathcal{P}(\mathbb{C}_0)$, is sent to a morphism $\beta(G(Y)) \to \beta(Y)$ that factors as $\beta(G(Y)) \xrightarrow{\rho_Y} G'(\beta(Y)) \to \beta(Y)$, where $G'(\beta(Y)) \in \mathcal{D}_0$. The map $\theta_{X,Y}$ identifies with the map

$$\mathcal{P}(\mathcal{C}_0)(\mathbf{X}, \mathbf{G}(\mathbf{Y})) \to \mathcal{D}_0(\beta(\mathbf{X}), \beta(\mathbf{G}(\mathbf{Y}))) \to \mathcal{D}_0(\beta(\mathbf{X}), \mathbf{G}'(\beta(\mathbf{Y}))).$$

Since the first map is an equivalence, it is enough to see that ρ_{Y} is an equivalence for every $Y \in \mathcal{P}(\mathcal{C}_{1})$. Since $\beta_{\mathcal{C}}$ is fully faithful, $\theta_{X,Y}$ is an equivalence if $X \in \mathcal{C}_{0}, Y \in \mathcal{C}_{1}$. Because $\mathcal{P}(\mathcal{C}_{0})$ is generated by \mathcal{C}_{0} under small colimits, $\theta_{X,Y}$ is an equivalence for all $X \in \mathcal{P}(\mathcal{C}_{0}), Y \in \mathcal{C}_{1}$. Thus ρ_{Y} is an equivalence for every $Y \in \mathcal{C}_{1}$. Therefore ρ_{Y} is also an equivalence for every $Y \in \mathcal{P}(\mathcal{C}_{1})$ because $\mathcal{P}(\mathcal{C}_{1})$ is generated by \mathcal{C}_{1} under small colimits and the functors G, G' preserve small colimits and ρ_{Y} is natural in Y.

Corollary 4.2. The canonical commutative square

(5)
$$\begin{array}{c} \operatorname{Fun}([1], \infty \operatorname{Cat}) \xrightarrow{\mathcal{P}} \operatorname{LoCART}^{\mathrm{L}} \\ \downarrow \\ \iota^{*}(\operatorname{Fun}([1], \infty \operatorname{Cat})) \xrightarrow{\iota^{*}(\mathcal{P})} \iota^{*}(\operatorname{LoCART}^{\mathrm{L}}) \end{array}$$

of cartesian fibrations over ∞ Cat is a pullback square.

Proof. Square (5) induces on the fiber over every small ∞ -category \mathcal{A} the pullback square (3) of ∞ -categories.

Notation 4.3. Let

- EXP \subset Fun([1], ∞ Cat) be the full subcategory of exponential fibrations,
- CART ⊂ Fun([1],∞Cat) be the full subcategory of cartesian fibrations,
- $CART^{L} \subset LoCART^{L}$ be the full subcategory of cartesian fibrations.

Corollary 4.2 and Proposition 3.24 imply the following corollary:

Corollary 4.4. The commutative square





of cartesian fibrations over ∞ Cat is a pullback square.

5. Correspondences as left fibrations

The next definition is [14, Notation 3.3.]:

Definition 5.1. Let \mathcal{C} be an ∞ -category. The twisted arrow left fibration $\operatorname{Tw}(\mathcal{C}) \to \mathcal{C}^{\operatorname{op}} \times \mathcal{C}$ is the left fibration classifying the mapping space functor $\mathcal{C}^{\operatorname{op}} \times \mathcal{C} \to S$.

Theorem 5.2. There is a canonical pullback square of ∞ -categories



where the right hand vertical functor evaluates at the target. The induced equivalence from the left upper corner of the square to the pullback sends a functor $\mathcal{M} \to [1]$ to the triple

 $(\mathfrak{M}_0, \mathfrak{M}_1, \mathfrak{M}_0^{\mathrm{op}} \times \mathfrak{M}_1 \times_{\mathfrak{M}^{\mathrm{op}} \times \mathfrak{M}} \mathrm{Tw}(\mathfrak{M}) \to \mathfrak{M}_0^{\mathrm{op}} \times \mathfrak{M}_1).$

Proof. By Theorem 4.1 there is a canonical pullback square of ∞ -categories:



Consequently, it is enough to construct an equivalence between the pullbacks

$$\begin{array}{c} \mathcal{A} & \longrightarrow \mathcal{L} \\ \downarrow & & \downarrow \\ \infty \mathsf{Cat} \times \infty \mathsf{Cat} \xrightarrow{(-)^{\mathrm{op}} \times (-)} & \infty \mathsf{Cat} \end{array}$$

and

$$\begin{array}{c} \mathcal{B} & \longrightarrow \operatorname{CART}_{[1]}^{\operatorname{L}} \\ \downarrow & \qquad \downarrow \\ \infty \operatorname{Cat} \times \infty \operatorname{Cat} & \xrightarrow{\mathcal{P} \times \mathcal{P}} & \operatorname{Pr}^{\operatorname{L}} \times \operatorname{Pr}^{\operatorname{L}}. \end{array}$$

We construct this equivalence by giving for every ∞ -category \mathcal{A} a bijection between equivalence classes of objects of the ∞ -categories Fun(\mathcal{A}, \mathcal{A}), Fun(\mathcal{A}, \mathcal{B}) natural in \mathcal{A} . A functor $\mathcal{A} \to \mathcal{B}$ is classified by a pair of cocartesian fibrations $\mathfrak{X} \to \mathcal{A}, \mathcal{Y} \to \mathcal{A}$ and a map of cocartesian fibrations $\mathcal{M} \to \mathcal{A} \times [1]$ over \mathcal{A} that induces on the fiber over every object $A \in \mathcal{A}$ a cartesian fibration $\mathcal{M}_A \to [1]$ classifying a small colimits preserving functor $\mathcal{M}_A^1 \to \mathcal{M}_A^0$ and such that there are equivalences $\mathcal{M}_0 \simeq \mathcal{P}^{\mathcal{A}}(\mathcal{X}), \mathcal{M}_1 \simeq \mathcal{P}^{\mathcal{A}}(\mathcal{Y})$ over \mathcal{A} . By [11, Lemma 2.44.] a functor $\mathcal{W} \to \mathcal{A} \times [1]$ is a map of cocartesian fibrations over \mathcal{A} that induces on the fiber over every object A $\in \mathcal{A}$ a cartesian fibration over [1] if and only if it is a map of cartesian fibrations over [1] that induces on the fiber over every object of [1] a cocartesian fibration over A. Consequently, a functor $\mathcal{A} \to \mathcal{B}$ is likewise classified by a pair of cocartesian fibrations $\mathfrak{X} \to \mathcal{A}, \mathfrak{Y} \to \mathcal{A}$ and a map of cartesian fibrations $\mathcal{M} \to \mathcal{A} \times [1]$ over [1] that induces on the fibers over $0, 1 \in [1]$ the cocartesian fibrations $\mathcal{P}^{\mathcal{A}}(\mathcal{X}), \mathcal{P}^{\mathcal{A}}(\mathcal{Y})$, respectively, and induces on the fiber over every object $A \in \mathcal{A}$ a cartesian fibration $\mathcal{M}_A \rightarrow [1]$ classifying a small colimits preserving functor $\mathcal{M}^1_A \to \mathcal{M}^0_A$. In other words, a functor $\mathcal{A} \to \mathcal{B}$ is classified by a pair of cocartesian fibrations $\mathfrak{X} \to \mathcal{A}, \mathfrak{Y} \to \mathcal{A}$ and a functor $\mathcal{P}^{\mathcal{A}}(\mathfrak{Y}) \to \mathcal{P}^{\mathcal{A}}(\mathfrak{X})$ over \mathcal{A} that induces on the fiber over every object of A a small colimits preserving functor. By Proposition 3.16 the latter is uniquely determined by its restriction, a functor $\mathcal{Y} \to \mathcal{P}^{\mathcal{A}}(\mathcal{X})$ over \mathcal{A} , which corresponds by Proposition 3.37 to a functor $\mathfrak{X}^{rev} \times_{\mathcal{A}} \mathfrak{Y} \to \mathfrak{S}$. The latter is classified by a left fibration $\mathcal{A} \to \mathfrak{X}^{\text{rev}} \times_{\mathcal{A}} \mathcal{Y}$ or equivalently by a map of cocartesian fibrations $\mathcal{A} \to \mathfrak{X}^{\text{rev}} \times_{\mathcal{A}} \mathcal{Y}$ over \mathcal{A} that induces on the fiber over every object of \mathcal{A} a left fibration. So a functor $\mathcal{A} \to \mathcal{B}$

is classified by a pair of cocartesian fibrations $\mathcal{X} \to \mathcal{A}, \mathcal{Y} \to \mathcal{A}$ and a map of cocartesian fibrations $\mathcal{A} \to \mathcal{X}^{\text{rev}} \times_{\mathcal{A}} \mathcal{Y}$ over \mathcal{A} that induces on the fiber over every object of \mathcal{A} a left fibration. The latter is precisely classified by a functor $\mathcal{A} \to \mathcal{A}$.

Remark 5.3. Under the equivalence

(7)
$$\infty \operatorname{Cat}_{/[1]} \simeq (\infty \operatorname{Cat} \times \infty \operatorname{Cat}) \times_{\infty \operatorname{Cat}} \mathcal{L}$$

of Theorem 5.2 the opposite ∞ -category involution on $\infty \operatorname{Cat}_{[1]}$ corresponds to an involution on the right hand pullback induced by the flip action on the product $\infty \operatorname{Cat} \times \infty \operatorname{Cat}$. This gives an easy way to understand the opposite ∞ -category involution on $\infty \operatorname{Cat}_{[1]}$ and is used in [14, Theorem 5.21.] and [15, Theorem 6.27.] to compare different models of dualities. In fact in [14, Proposition 5.23.] equivalence (7) is enhanced to a C₂-equivariant equivalence for the C₂-actions refining the opposite ∞ -category involution and the flip action.

Notation 5.4. Let $\text{Cocart} \subset \text{Fun}([1], \infty \text{Cat})$ be the full subcategory of cocartesian fibrations.

Corollary 5.5. Let \mathcal{C}, \mathcal{D} be small ∞ -categories. There is a canonical equivalence

$$\{(\mathcal{C},\mathcal{D})\}\times_{\infty \mathsf{Cat}\times\infty \mathsf{Cat}} \mathsf{Cat}_{/[1]}\simeq \operatorname{Fun}(\mathcal{C}^{\operatorname{op}},\operatorname{Fun}(\mathcal{D},\mathcal{S}))$$

that sends a functor $\mathbb{M} \to [1]$ and $\mathbb{M}_0 \simeq \mathbb{C}, \mathbb{M}_1 \simeq \mathbb{D}$ to the functor $\mathbb{C}^{\mathrm{op}} \times \mathbb{D} \subset \mathbb{M}^{\mathrm{op}} \times \mathbb{M} \xrightarrow{\mathbb{M}(-,-)} S$. The latter equivalence restricts to an equivalence

 $\{(\mathcal{C}, \mathcal{D})\} \times_{\infty \mathsf{Cat} \times \infty \mathsf{Cat}} \operatorname{Cocart}_{[1]} \simeq \operatorname{Fun}(\mathcal{C}^{\operatorname{op}}, \mathcal{D}^{\operatorname{op}}) \simeq \operatorname{Fun}(\mathcal{C}, \mathcal{D})^{\operatorname{op}}$

that sends a functor $\mathcal{M} \to [1]$ and $\mathcal{M}_0 \simeq \mathcal{C}, \mathcal{M}_1 \simeq \mathcal{D}$ to the functor $\mathcal{C} \to \mathcal{D}$ classified by $\mathcal{M} \to [1]$.

Proof. By Theorem 5.2 there is a canonical equivalence

 $\{(\mathfrak{C}, \mathfrak{D})\} \times_{\infty \mathsf{Cat} \times \infty \mathsf{Cat}} \mathsf{Cat}_{/[1]} \simeq \{\mathfrak{C}^{\mathrm{op}} \times \mathfrak{D}\} \times_{\infty \mathsf{Cat}} \mathcal{L} \simeq \mathrm{Fun}(\mathfrak{C}^{\mathrm{op}} \times \mathfrak{D}, \mathfrak{S}) \simeq \mathrm{Fun}(\mathfrak{C}^{\mathrm{op}}, \mathrm{Fun}(\mathfrak{D}, \mathfrak{S}))$

that sends a functor $\mathcal{M} \to [1]$ and $\mathcal{M}_0 \simeq \mathcal{C}, \mathcal{M}_1 \simeq \mathcal{D}$ to the functor $\mathcal{C}^{\mathrm{op}} \times \mathcal{D} \subset \mathcal{M}^{\mathrm{op}} \times \mathcal{M} \xrightarrow{\mathcal{M}(-,-)} S$. The latter equivalence restricts to an equivalence

 $\{(\mathcal{C},\mathcal{D})\}\times_{\infty \mathsf{Cat}\times\infty \mathsf{Cat}} \operatorname{Cocart}_{[1]} \simeq \operatorname{Fun}(\mathcal{C}^{\operatorname{op}},\mathcal{D}^{\operatorname{op}}) \simeq \operatorname{Fun}(\mathcal{C},\mathcal{D})^{\operatorname{op}}$

that sends a functor $\mathcal{M} \to [1]$ and $\mathcal{M}_0 \simeq \mathcal{C}, \mathcal{M}_1 \simeq \mathcal{D}$ to the functor $\mathcal{C} \to \mathcal{D}$ classified by $\mathcal{M} \to [1]$.

For every cocartesian fibration $\mathcal{M} \to [1]$ and $\mathcal{M}_0 \simeq \mathcal{C}, \mathcal{M}_1 \simeq \mathcal{D}$ classifying a functor $\alpha : \mathcal{C} \to \mathcal{D}$ there is a canonical equivalence between the functor $\mathcal{C}^{\text{op}} \times \mathcal{D} \subset \mathcal{M}^{\text{op}} \times \mathcal{M} \xrightarrow{\mathcal{M}(-,-)} S$ and the functor $\mathcal{C}^{\text{op}} \times \mathcal{D} \xrightarrow{\alpha^{\text{op}} \times \mathcal{D}} \mathcal{D}^{\text{op}} \times \mathcal{D} \xrightarrow{\mathcal{D}(-,-)} S$.

6. An end formula for mapping spaces of parametrized functor ∞-categories

In this section we describe the mapping space functor of the internal hom Fun^[1](\mathcal{M}, \mathcal{N}) between two functors $\mathcal{M} \to [1], \mathcal{N} \to [1]$ (Theorem 6.5). Having this description we relate the mapping spaces of the conditionally existing internal hom Fun^{\mathcal{C}}(\mathcal{M}, \mathcal{N}) between two functors $\mathcal{M} \to \mathcal{C}, \mathcal{N} \to \mathcal{C}$ to the mapping space functors of \mathcal{M}, \mathcal{N} (Corollary 6.6). As a corollary we compute the mapping space functor of the ∞ -category Fun(\mathcal{A}, \mathcal{B}) of functors between two ∞ -categories \mathcal{A}, \mathcal{B} (Corollary 6.9) recovering a global version of a result of [7, Proposition 5.1.].

Lemma 6.1. Let \mathcal{V} be a closed monoidal ∞ -category, $\mathcal{W} \subset \operatorname{Fun}([1], \mathcal{V})$ a full monoidal subcategory such that the base change of any object $X \to Y$ of W along any morphism $Z \to X$ of \mathcal{V} exists and is an object of \mathcal{W} and for every object $X \to Y$ of \mathcal{W} and every object Z of \mathcal{V} the induced morphism $\operatorname{Mor}_{\mathcal{V}}(Z, X) \to \operatorname{Mor}_{\mathcal{V}}(Z, Y)$ in \mathcal{V} is an object of \mathcal{W} . Let $\alpha : \mathcal{A} \to \mathcal{V}$ be a monoidal functor. The pullback $\mathcal{A} \times_{\mathcal{V}} \mathcal{W}$ of evaluation at the target $\mathcal{W} \to \mathcal{V}$ along α , which is a monoidal ∞ -category, is closed. The internal hom of every $(A, f : X \to \alpha(A)), (B, g : Y \to \alpha(B)) \in \mathcal{A} \times_{\mathcal{V}} \mathcal{W}$ is

$$(\operatorname{Mor}_{\mathcal{A}}(A,B), \alpha(\operatorname{Mor}_{\mathcal{A}}(A,B)) \times_{\operatorname{Mor}_{\mathcal{V}}(X,\alpha(B))} \operatorname{Mor}_{\mathcal{V}}(X,Y) \to \alpha(\operatorname{Mor}_{\mathcal{A}}(A,B))) \in \mathcal{A} \times_{\mathcal{V}} \mathcal{W}.$$

Proof. For every $(A, f : X \to \alpha(A)), (B, g : Y \to \alpha(B)) \in \mathcal{A} \times_{\mathcal{V}} \mathcal{W}, (C, h : Z \to \alpha(C)) \in \mathcal{A} \times_{\mathcal{V}} \mathcal{W}$ the canonical morphism

 $(Mor_{\mathcal{A}}(A, B), \alpha(Mor_{\mathcal{A}}(A, B)) \times_{Mor_{\mathcal{V}}(X, \alpha(B))} Mor_{\mathcal{V}}(X, Y) \otimes (A, f : X \to \alpha(A)) =$

 $(\operatorname{Mor}_{\mathcal{A}}(A, B) \otimes A, \alpha(\operatorname{Mor}_{\mathcal{A}}(A, B) \otimes A) \times_{\operatorname{Mor}_{\mathcal{V}}(X, \alpha(B)) \otimes X} \operatorname{Mor}_{\mathcal{V}}(X, Y) \otimes X) \to (B, g : Y \to \alpha(B))$ in $\mathcal{A} \times_{\mathcal{V}} \mathcal{W}$ induces a map

$$\mathcal{A} \times_{\mathcal{V}} \mathcal{W}((\mathbf{A}, \mathbf{f} : \mathbf{X} \to \alpha(\mathbf{A})),$$

$$\begin{split} & (\operatorname{Mor}_{\mathcal{A}}(\operatorname{B},\operatorname{C}), \alpha(\operatorname{Mor}_{\mathcal{A}}(\operatorname{B},\operatorname{C})) \times_{\operatorname{Mor}_{\mathcal{V}}(\operatorname{Y},\alpha(\operatorname{C}))} \operatorname{Mor}_{\mathcal{V}}(\operatorname{Y},\operatorname{Z}) \to \alpha(\operatorname{Mor}_{\mathcal{A}}(\operatorname{B},\operatorname{C})))) \simeq \\ & \mathcal{A} \times_{\mathcal{V}} \mathcal{W}((\operatorname{A},\operatorname{f}:\operatorname{X} \to \alpha(\operatorname{A})) \otimes (\operatorname{B},\operatorname{g}:\operatorname{Y} \to \alpha(\operatorname{B})), (\operatorname{C},\operatorname{h}:\operatorname{Z} \to \alpha(\operatorname{C}))) = \\ & \mathcal{A} \times_{\mathcal{V}} \mathcal{W}((\operatorname{A} \otimes \operatorname{B},\operatorname{f} \otimes \operatorname{g}:\operatorname{X} \otimes \operatorname{Y} \to \alpha(\operatorname{A} \otimes \operatorname{B})), (\operatorname{C},\operatorname{h}:\operatorname{Z} \to \alpha(\operatorname{C}))) \end{split}$$

over $\mathcal{A}(A, Mor_{\mathcal{A}}(B, C)) \simeq \mathcal{A}(A \otimes B, C)$ that induces on the fiber over every morphism θ : $A \rightarrow Mor_{\mathcal{A}}(B, C)$ corresponding to a morphism $\theta' : A \otimes B \rightarrow C$ the canonical equivalence

$$\mathcal{V}_{/\mathrm{Mor}_{\mathcal{V}}(\mathrm{Y},\alpha(\mathrm{C}))}(\mathrm{X},\mathrm{Mor}_{\mathcal{V}}(\mathrm{Y},\mathrm{Z})) \simeq \mathcal{V}_{/\alpha(\mathrm{C})}(\mathrm{X}\otimes\mathrm{Y},\mathrm{Z}).$$

Corollary 6.2. The pullback



in cartesian symmetric monoidal ∞ -categories is closed. The internal hom of every $(\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{C} \rightarrow \mathcal{A}^{\mathrm{op}} \times \mathcal{B}), (\mathcal{A}', \mathcal{B}', \mathcal{C}', \mathcal{C}' \rightarrow \mathcal{A}'^{\mathrm{op}} \times \mathcal{B}')$ is

 $(\operatorname{Fun}(\mathcal{A},\mathcal{A}'),\operatorname{Fun}(\mathcal{B},\mathcal{B}'),\operatorname{Fun}(\mathcal{A},\mathcal{A}')^{\operatorname{op}}\times\operatorname{Fun}(\mathcal{B},\mathcal{B}')\times_{\operatorname{Fun}(\mathcal{C},\mathcal{A}'^{\operatorname{op}}\times\mathcal{B}')}\operatorname{Fun}(\mathcal{C},\mathcal{C}')$

$$\rightarrow$$
 Fun($\mathcal{A}, \mathcal{A}'$)^{op} × Fun($\mathcal{B}, \mathcal{B}'$)).

Notation 6.3. For every functor $\mathcal{M} \to [1]$ let $\widetilde{\mathrm{Tw}}(\mathcal{M})$ be the pullback

$$\mathcal{M}_{0}^{\mathrm{op}} \times \mathcal{M}_{1} \times_{\mathcal{M}^{\mathrm{op}} \times \mathcal{M}} \mathrm{Tw}(\mathcal{M}) \to \mathcal{M}_{0}^{\mathrm{op}} \times \mathcal{M}_{1}.$$

Remark 6.4. Let \mathcal{C} be an ∞ -category. Projection $\mathcal{C} \times [1] \to \mathcal{C}$ induces an equivalence $\widetilde{\mathrm{Tw}}(\mathcal{C} \times [1]) \simeq \mathrm{Tw}(\mathcal{C})$ over $\mathcal{C}^{\mathrm{op}} \times \mathcal{C}$.

Theorem 6.5. Let $\mathcal{M} \to [1], \mathcal{N} \to [1]$ be functors. There is a canonical equivalence

 $\widetilde{\mathrm{Tw}}(\mathrm{Fun}^{[1]}(\mathcal{M},\mathcal{N})) \simeq$

 $\operatorname{Fun}(\mathcal{M}_0,\mathcal{N}_0)^{\operatorname{op}}\times\operatorname{Fun}(\mathcal{M}_1,\mathcal{N}_1)\times_{\operatorname{Fun}(\widetilde{\operatorname{Tw}}(\mathcal{M}),\mathcal{N}_0^{\operatorname{op}}\times\mathcal{N}_1)}\operatorname{Fun}(\widetilde{\operatorname{Tw}}(\mathcal{M}),\widetilde{\operatorname{Tw}}(\mathcal{N}))$

over

$$\operatorname{Fun}(\mathcal{M}_0, \mathcal{N}_0)^{\operatorname{op}} \times \operatorname{Fun}(\mathcal{M}_1, \mathcal{N}_1)$$

that induces on the fiber over any functors $F: \mathcal{M}_0 \to \mathcal{N}_0, G: \mathcal{M}_1 \to \mathcal{N}_1$ an equivalence

$$\operatorname{Fun}^{[1]}(\mathcal{M},\mathcal{N})(\mathrm{F},\mathrm{G}) \simeq \lim(\widetilde{\operatorname{Tw}}(\mathcal{M}) \to \mathcal{M}_0^{\operatorname{op}} \times \mathcal{M}_1 \xrightarrow{\mathrm{F}^{\operatorname{op}} \times \mathrm{G}} \mathcal{N}_0^{\operatorname{op}} \times \mathcal{N}_1 \subset \mathcal{N}^{\operatorname{op}} \times \mathcal{N} \xrightarrow{\mathcal{N}(-,-)} \mathcal{S}).$$

Proof. By Theorem 5.2 there is a canonical equivalence θ between $\infty Cat_{/[1]}$ and the pullback



Consequently, the internal hom of $\infty \operatorname{Cat}_{[1]}$ corresponds under θ to the internal hom of Ω for the cartesian monoidal structures. The description of θ of Theorem 5.2 and the description of the internal hom of Corollary 6.2 imply the result.

Corollary 6.6. Let $\mathcal{A} \to \mathcal{C}, \mathcal{B} \to \mathcal{C}$ be functors, $f : X \to Y$ a morphism in \mathcal{C} and $F : \mathcal{A}_X \to \mathcal{B}_X, G : \mathcal{A}_Y \to \mathcal{B}_Y$ functors. There is a canonical equivalence

$$\{\alpha\} \times_{\mathcal{C}(X,Y)} \operatorname{Fun}^{\mathcal{C}}(\mathcal{A},\mathcal{B})(F,G) \simeq$$

$$\lim(\widetilde{\mathrm{Tw}}([1]\times_{\mathbb{C}}\mathcal{A})\to\mathcal{A}_{\mathrm{X}}^{\mathrm{op}}\times\mathcal{A}_{\mathrm{Y}}\xrightarrow{\mathrm{F}^{\mathrm{op}}\times\mathrm{G}}\mathcal{B}_{\mathrm{X}}^{\mathrm{op}}\times\mathcal{B}_{\mathrm{Y}}\to([1]\times_{\mathbb{C}}\mathcal{B})^{\mathrm{op}}\times([1]\times_{\mathbb{C}}\mathcal{B})\xrightarrow{([1]\times_{\mathbb{C}}\mathcal{B})(-,-)}\mathcal{S}).$$

Proof. There is a canonical equivalence

$$\{\alpha\} \times_{\mathcal{C}(X,Y)} \operatorname{Fun}^{\mathbb{C}}(\mathcal{A}, \mathcal{B})(F, G) \simeq ([1] \times_{\mathbb{C}} \operatorname{Fun}^{\mathbb{C}}(\mathcal{A}, \mathcal{B}))(F, G) \simeq$$
$$\operatorname{Fun}^{[1]}([1] \times_{\mathbb{C}} \mathcal{A}, [1] \times_{\mathbb{C}} \mathcal{B})(F, G) \simeq$$
$$\operatorname{lim}(\widetilde{\operatorname{Tw}}([1] \times_{\mathbb{C}} \mathcal{A}) \to \mathcal{A}_{X}^{\operatorname{op}} \times \mathcal{A}_{Y} \xrightarrow{F^{\operatorname{op}} \times G} \mathcal{B}_{X}^{\operatorname{op}} \times \mathcal{B}_{Y} \to ([1] \times_{\mathbb{C}} \mathcal{B})^{\operatorname{op}} \times ([1] \times_{\mathbb{C}} \mathcal{B}) \xrightarrow{([1] \times_{\mathbb{C}} \mathcal{B})(-,-)} S).$$

Definition 6.7. Let \mathcal{C}, \mathcal{D} be ∞ -categories and $\phi : \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \to \mathcal{D}$ a functor. The end of ϕ , denoted by $\int \phi$ if it exists, is the limit of the functor $\mathrm{Tw}(\mathcal{C}) \xrightarrow{\mathrm{q}} \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \xrightarrow{\phi} \mathcal{D}$.

Remark 6.8. Let $\phi : \mathbb{C}^{\text{op}} \times \mathbb{C} \to \mathbb{S}$ be a functor classified by a left fibration $\mathcal{B} \to \mathbb{C}^{\text{op}} \times \mathbb{C}$. The limit of the functor $\operatorname{Tw}(\mathbb{C}) \xrightarrow{q} \mathbb{C}^{\text{op}} \times \mathbb{C} \xrightarrow{\phi} \mathbb{S}$ is the space

$$\operatorname{Fun}_{\operatorname{Tw}(\mathcal{C})}(\operatorname{Tw}(\mathcal{C}), q^*(\mathcal{B}))$$

Theorem 6.5 implies the following corollary in view of Remark 6.4:

Corollary 6.9. Let \mathcal{C}, \mathcal{D} be ∞ -categories. There is a canonical equivalence

 $\operatorname{Tw}(\operatorname{Fun}(\mathcal{C},\mathcal{D})) \simeq \operatorname{Fun}(\mathcal{C},\mathcal{D})^{\operatorname{op}} \times \operatorname{Fun}(\mathcal{C},\mathcal{D}) \times_{\operatorname{Fun}(\operatorname{Tw}(\mathcal{C}),\mathcal{D}^{\operatorname{op}} \times \mathcal{D})} \operatorname{Fun}(\operatorname{Tw}(\mathcal{C}),\operatorname{Tw}(\mathcal{D}))$ over

 $\operatorname{Fun}(\mathcal{C}, \mathcal{D})^{\operatorname{op}} \times \operatorname{Fun}(\mathcal{C}, \mathcal{D}).$

The latter induces on the fiber over every functors $F, G : \mathcal{C} \to \mathcal{D}$ an equivalence

$$\operatorname{Fun}(\mathfrak{C},\mathfrak{D})(F,G)\simeq\operatorname{Fun}_{\operatorname{Tw}(\mathfrak{C})}(\operatorname{Tw}(\mathfrak{C}),\operatorname{q}^*((F^{\operatorname{op}}\times G)^*(\operatorname{Tw}(\mathfrak{D}))))\simeq\int \mathfrak{D}(F(-),G(-)).$$

7. Correspondences as profunctors

In the following we use the notion of double ∞ -category [6, Definition 2.4.3.], which is an ∞ -category internal to ∞ Cat :

Definition 7.1. A double ∞ -category is a cartesian fibration $\mathcal{B} \to \Delta$ such that for every $[n] \in \Delta$ the induced functor

$$\mathcal{B}_{[n]} \to \mathcal{B}_{[1]} \times_{\mathcal{B}_{[0]}} \dots \times_{\mathcal{B}_{[0]}} \mathcal{B}_{[1]}$$

is an equivalence, where the i-th map $[1] \rightarrow [n]$ sends 0 to i – 1 and 1 to i.

Example 7.2. The functor

$$p:\infty \mathsf{Cat}_{/\Delta}^{\mathrm{cart}} \to \infty \mathsf{Cat}$$

taking the fiber over [0] admits a fully faithful right adjoint that sends an ∞ -category \mathcal{C} to the cartesian fibration $\Delta_{\mathcal{C}} \rightarrow \Delta$ classifying the functor

$$\Delta^{\mathrm{op}} \subset \infty \mathsf{Cat}^{\mathrm{op}} \xrightarrow{\mathrm{Fun}(-,\mathcal{C})} \infty \mathsf{Cat}.$$

The cartesian fibration $\Delta_{\mathfrak{C}} \rightarrow \Delta$ is a double ∞ -category.

Remark 7.3. The functor $\rho : \infty \mathsf{Cat}_{/\Delta}^{cart} \to \infty \mathsf{Cat}$ is a cartesian fibration, where a morphism $\mathcal{B} \to \mathcal{C}$ over Δ is ρ -cartesian if and only if the following square is a pullback square:



This holds since for every cartesian fibration $\mathcal{A} \to \Delta$ the induced commutative square

is equivalent to the commutative square

In particular, the ρ -cartesian lift to a double ∞ -category $\mathcal{C} \to \Delta$ of a functor $\mathcal{A} \to \mathcal{C}_{[0]}$ is the double ∞ -category $\Delta_{\mathcal{A}} \times_{\Delta_{\mathcal{C}_{[0]}}} \mathcal{C}$.

Notation 7.4. Let $\mathcal{A} \to \Delta, \mathcal{B} \to \Delta$ be double ∞ -categories. We write

$$\operatorname{FUN}(\mathcal{A}, \mathcal{B}) \subset \operatorname{Fun}_{\Delta}(\mathcal{A}, \mathcal{B})$$

for the full subcategory spanned by the functors over Δ preserving cocartesian lifts of morphisms of Δ .

Definition 7.5. An $(\infty, 2)$ -precategory is a double ∞ -category $\phi : \mathcal{B} \to \Delta$ such that \mathcal{B}_0 is a space.

Remark 7.6. By [17, 5.6.1. Corollary] there is an equivalence between $(\infty, 2)$ -precategories in the sense of Definition 7.5 and ∞ -precategories enriched in ∞ Cat, which justifies our terminology.

Example 7.7. For every double ∞ -category $\phi : \mathcal{B} \to \Delta$ the pullback

$$\Delta_{\iota(\mathcal{B}_{[0]})} \times_{\Delta_{\mathcal{B}_{[0]}}} \mathcal{B} \to \Delta_{\iota(\mathcal{B}_{[0]})}$$

is an $(\infty, 2)$ -precategory, which we call the underlying $(\infty, 2)$ -precategory of ϕ .

Example 7.8. Viewing spaces as ∞ -categories any Segal space classifies an $(\infty, 2)$ -precategory. For every ∞ -category \mathcal{C} the nerve of \mathcal{C} defined by $N(\mathcal{C}) : \Delta^{\mathrm{op}} \to \mathcal{S}, [n] \mapsto \infty \mathsf{Cat}([n], \mathcal{C})$ is a Segal space and so classifies an $(\infty, 2)$ -precategory.

For the next notation let $\tau := (-)^{\text{op}} : \infty \text{Cat} \to \infty \text{Cat}$ be the opposite ∞ -category involution.

Notation 7.9. Let

- (1) $\infty \text{CAT} \coloneqq \tau^*(\Delta \times_{\infty \text{Cat}} \text{CART}) \to \Delta$,
- (2) $\operatorname{PR}^{\mathrm{L}} \coloneqq \tau^*(\Delta \times_{\infty \widehat{\mathsf{Cat}}} \operatorname{CART}^{\mathrm{L}}) \to \Delta,$
- (3) CORR := $\tau^*(\Delta \times_{\infty Cat} EXP) \to \Delta$,
- (4) CORR_{Lax} := $\tau^*(\Delta \times_{\infty Cat} Fun([1], \infty Cat)) \rightarrow \Delta$.

Construction 7.10. Let $H: K \to \infty$ Cat be a functor that admits a colimit \mathcal{A} . The canonical ∞ Cat-linear functor $Cat_{\infty/\mathcal{A}} \to \lim_{k \in K} Cat_{\infty/H(k)}$ induces on morphism ∞ -categories between two functors $\mathcal{C} \to \mathcal{A}, \mathcal{D} \to \mathcal{A}$ a functor

(8)
$$\theta : \operatorname{Fun}_{\mathcal{A}}(\mathcal{C}, \mathcal{D}) \to \lim_{k \in K} \operatorname{Fun}_{\mathrm{H}(k)}(\mathrm{H}(k) \times_{\mathcal{A}} \mathcal{C}, \mathrm{H}(k) \times_{\mathcal{A}} \mathcal{D}).$$

Lemma 7.11. Let $H : K \to \infty$ Cat be a functor with colimit \mathcal{A} and $\mathcal{C} \to \mathcal{A}, \mathcal{D} \to \mathcal{A}$ functors. If the functor $\mathcal{C} \to \mathcal{A}$ is an exponential fibration, the functor (8) is an equivalence.

Proof. Since $\mathcal{C} \to \mathcal{A}$ is an exponential fibration, the functor $(-) \times_{\mathcal{A}} \mathcal{C} : \operatorname{Cat}_{\infty/\mathcal{A}} \to \operatorname{Cat}_{\infty/\mathcal{C}} \to \operatorname{Cat}_{\infty/\mathcal{A}}$ preserves small colimits. Hence the canonical functor $\operatorname{colim}_{k \in K}(\operatorname{H}(k) \times_{\mathcal{A}} \mathcal{C}) \to \mathcal{C}$ over $\operatorname{colim}(\operatorname{H}) \simeq \mathcal{A}$ is an equivalence. Since the functor $\operatorname{Fun}_{\mathcal{A}}(-, \mathcal{D}) : \operatorname{Cat}_{\infty/\mathcal{A}}^{\operatorname{op}} \to \infty$ Cat preserves limits, the following canonical functor is an equivalence:

$$\operatorname{Fun}_{\mathcal{A}}(\mathcal{C},\mathcal{D}) \to \operatorname{Fun}_{\mathcal{A}}(\operatorname{colim}_{k \in K} \operatorname{H}(k) \times_{\mathcal{A}} \mathcal{C}, \mathcal{D}) \simeq \lim_{k \in K} \operatorname{Fun}_{\operatorname{H}(k)}(\operatorname{H}(k) \times_{\mathcal{A}} \mathcal{C}, \operatorname{H}(k) \times_{\mathcal{A}} \mathcal{D}).$$

For the next corollary we use that for every $n \ge 0$ the category [n] is the colimit in ∞ Cat of the canonical diagram

(9)
$$\{ 0 < 1 \} \coprod_{\{1\}} \{ 1 < 2 \} \coprod_{\{2\}} \dots \coprod_{\{n-1\}} \{ n-1 < n \},$$

in which all maps are inert. Lemma 7.11 gives the following corollary:

Corollary 7.12. Let $n \ge 0$ and $\mathcal{B} \to \mathbb{C}$ a functor over [n] whose pullback along any inert map $[1] \to [n]$ is an equivalence. For every exponential fibration $\mathcal{D} \to [n]$ the induced functor $\operatorname{Fun}_{[n]}(\mathcal{D},\mathcal{B}) \to \operatorname{Fun}_{[n]}(\mathcal{D},\mathbb{C})$

is an equivalence.

Notation 7.13. Let $n \ge 0$ and $\phi : \mathcal{C} \to [n]$ a functor. Pulling back diagram (9) along ϕ we obtain a diagram

$$\{0 < 1\} \times_{[n]} \mathcal{C} \times_{(\{1\}\times_{[n]}\mathcal{C})} \{1 < 2\} \times_{[n]} \mathcal{C} \times_{(\{2\}\times_{[n]}\mathcal{C})} \dots \times_{(\{n-1\}\times_{[n]}\mathcal{C})} \{n-1 < n\} \times_{[n]} \mathcal{C} \times_{(n-1)} \mathcal{C} \times_{(n-1$$

over $\mathfrak{C}.$ We write \mathfrak{C}' for the colimit of the latter diagram.

Remark 7.14. By Corollary 3.28 the inert maps $[1] \cong \{i - 1 < i\} \subset [n]$ for any $i \leq 0$ are exponential fibrations. Hence for every $i \leq 0$ the pullback of the canonical functor $\mathcal{C}' \to \mathcal{C}$ over [n] along the map $[1] \cong \{i - 1 < i\} \subset [n]$ identifies with the equivalence

$$\varnothing \times_{(\{i-1\}\times_{\lceil n\rceil} \mathcal{C})} \{i-1 < i\} \times_{\lceil n\rceil} \mathcal{C} \times_{(\{i\}\times_{\lceil n\rceil} \mathcal{C})} \varnothing \to \{i-1 < i\} \times_{\lceil n\rceil} \mathcal{C}.$$

Corollary 7.12 and Remark 7.14 give the following:

Corollary 7.15. Let $n \ge 0$ and $\mathbb{C} \to [n]$ a functor. For every exponential fibration $\mathcal{D} \to [n]$ the induced functor

$$\operatorname{Fun}_{[n]}(\mathcal{D}, \mathcal{C}') \to \operatorname{Fun}_{[n]}(\mathcal{D}, \mathcal{C})$$

is an equivalence.

We obtain the following proposition:

Proposition 7.16. Let $n \ge 0$. The embedding $EXP_{[n]} \subset \infty Cat_{/[n]}$ admits a right adjoint. A functor $C \to D$ over [n] is a colocal equivalence if and only if it induces an equivalence after pulling back along any inert map $[1] \to [n]$.

Proposition 7.17. The cartesian fibration CART $\rightarrow \infty$ Cat classifies the functor

$$FUN(\tau^*N, \infty CAT) : \infty Cat^{op} \to \infty \widehat{Cat}.$$

Proof. By Yoneda the cartesian fibration $\Delta \times_{\infty Cat} Cart = \tau^* \infty CAT \rightarrow \Delta$ classifies the functor

$$\mathrm{FUN}(\mathrm{N}_{|\Delta},\tau^*\infty\mathrm{CAT})\simeq\mathrm{FUN}(\tau^*\mathrm{N}_{|\Delta},\infty\mathrm{CAT}):\Delta^{\mathrm{op}}\to\infty\overline{\mathsf{Cat}}.$$

Since the full subcategory $\Delta \subset \infty \mathsf{Cat}$ is dense, it is enough to verify that the functor $\infty \mathsf{Cat}^{\operatorname{op}} \to \infty \widehat{\mathsf{Cat}}$ classified by the cartesian fibration CART $\to \infty \mathsf{Cat}$ and the functor $\operatorname{FUN}(\tau^*\mathrm{N}, \infty \operatorname{CAT}) : \infty \operatorname{Cat}^{\operatorname{op}} \to \infty \widehat{\mathsf{Cat}}$ send for every ∞ -category \mathcal{C} the colimit of the functor $\rho : \Delta \times_{\infty \operatorname{Cat}} \infty \operatorname{Cat}_{/\mathcal{C}} \to \Delta \subset \infty \operatorname{Cat}$ to a limit. We first prove the second statement: because the functor $\operatorname{FUN}(-, \infty \operatorname{Cat}) : \operatorname{Seg}(\infty \operatorname{Cat})^{\operatorname{op}} \to \infty \operatorname{Cat}$ preserves small limits, it suffices to show that the embedding $\mathrm{N} : \infty \operatorname{Cat} \to \operatorname{Seg}(\mathcal{S}) \subset \operatorname{Seg}(\infty \operatorname{Cat})$ preserves the colimit of the functor $\Delta \times_{\infty \operatorname{Cat}} \infty \operatorname{Cat}_{/\mathcal{C}} \to \Delta \subset \infty \operatorname{Cat}$. The embedding $\mathrm{N} : \infty \operatorname{Cat} \to \operatorname{Seg}(\mathcal{S})$ preserves this colimit because $\operatorname{Seg}(\mathcal{S})$ is a localization of $\mathcal{P}(\Delta)$ and so for every ∞ -category \mathcal{C} the Segal space $\mathrm{N}(\mathcal{C})$ is the colimit of the functor $\Delta \times_{\operatorname{Seg}(\mathcal{S})} \operatorname{Seg}(\mathcal{S})_{/\mathrm{N}(\mathcal{C})} \to \Delta \subset \infty \operatorname{Cat} \subset \operatorname{Seg}(\mathcal{S})$, which factors as $\mathrm{N}\rho$. The embedding $\operatorname{Seg}(\mathcal{S}) \subset \operatorname{Seg}(\infty \operatorname{Cat})$ preserves small colimits since the embedding $\operatorname{Fun}(\Delta^{\operatorname{op}}, \mathcal{S}) \subset \operatorname{Fun}(\Delta^{\operatorname{op}}, \infty \operatorname{Cat})$ preserve small colimits and Segal equivalences: the generating Segal equivalences for $\operatorname{Fun}(\Delta^{\operatorname{op}}, \infty \operatorname{Cat})$ are the canonical morphisms $\Delta^1 \coprod_{\Delta^0} \ldots \amalg_{\Delta^0} \Delta^1 \to \Delta^n$ for $n \geq 0$, while the generating Segal equivalences for $\operatorname{Fun}(\Delta^{\operatorname{op}}, \infty \operatorname{Cat})$ are the canonical morphisms $\Delta^1 \amalg_{\Delta^0} \ldots \amalg_{\Delta^0} \Delta^1 \times \mathcal{A} \to \Delta^n \times \mathcal{A}$ for $n \geq 0$ and \mathcal{A} a small ∞ -category.

We finish the proof by showing that the cartesian fibration $CART \rightarrow \infty Cat$ classifies a small limits preserving functor $\infty Cat^{op} \rightarrow \infty Cat$. Let $H: K \rightarrow \infty Cat$ be a functor that admits a colimit \mathcal{A} . The canonical ∞Cat -linear functor $\rho: CART_{\mathcal{A}} \rightarrow \lim_{k \in K} CART_{H(k)}$ induces on morphism ∞ -categories between two functors $\mathcal{C} \rightarrow \mathcal{A}, \mathcal{D} \rightarrow \mathcal{A}$ the functor (8). The latter is an equivalence by Lemma 7.11 because by Corollary 3.29 every cartesian fibration is an exponential fibration. The induced map $\iota(\rho)$ is essentially surjective because it identifies with the canonical equivalence

$$\iota(\operatorname{Fun}(\mathcal{A},\infty\mathsf{Cat}^{\operatorname{op}})) \to \lim_{\mathbf{k}\in K} \iota(\operatorname{Fun}(\mathbf{H}(\mathbf{k}),\infty\mathsf{Cat}^{\operatorname{op}})).$$

Corollary 7.18. The cartesian fibration $CART^{L} \to \infty Cat$ classifies the functor $FUN(\tau^*N, PR^{L}) : \infty Cat^{op} \to \infty \widehat{Cat}.$

Corollary 4.4 implies the following theorem:

Theorem 7.19. The cartesian fibrations (1)-(3) of Notation 7.9 are double ∞ -categories and there is a canonical pullback square of double ∞ -categories:



For the next proposition we view ∞ Cat as $(\infty, 2)$ -category via its internal hom:

Proposition 7.20. The $(\infty, 2)$ -category underlying the double ∞ -category ∞ CAT is ∞ Cat viewed as enriched in itself via the internal hom.

Proof. Let ∞ CAT' be the $(\infty, 2)$ -category underlying the double ∞ -category ∞ CAT. Then the space of objects of ∞ CAT' is $\iota(\infty$ CAT_[0]) $\simeq \iota(\infty$ Cat) and for every A, B $\in \infty$ Cat there is a canonical equivalence

$$\operatorname{Mor}_{\infty \operatorname{CAT}'}(A,B) \simeq \{(A,B)\} \times_{\infty \operatorname{CAT}'_{[0]} \times \infty \operatorname{CAT}'_{[0]}} \infty \operatorname{CAT}'_{[1]} \simeq \{(A,B)\} \times_{\infty \operatorname{CAT}_{[0]} \times \infty \operatorname{CAT}_{[0]}} \infty \operatorname{CAT}_{[1]}$$

By the dual of Corollary 5.5 there is a canonical equivalence

$$\{(\mathbf{A},\mathbf{B})\}\times_{\infty \mathrm{CAT}_{[0]}\times\infty \mathrm{CAT}_{[0]}}\infty \mathrm{CAT}_{[1]}\simeq \mathrm{Fun}(\mathbf{A},\mathbf{B}).$$

In particular, for any $A, B, C \in \infty Cat$ there is a canonical equivalence

 $Mor_{\infty CAT'}(A \times B, C) \simeq Mor_{\infty CAT'}(A, Mor_{\infty CAT'}(B, C))$

so that $A \times B$ is the tensor of A and B in the sense of [12, Definition 2.51.] of the ∞ Catenriched ∞ -category ∞ CAT'. Thus ∞ CAT' admits all tensors. By [10, Remark 3.63., Corollary 6.13.] this implies that there is a unique 2-functor $\rho : \infty$ Cat $\rightarrow \infty$ CAT' that preserves tensors and sends the final ∞ -category to the final ∞ -category. By [13, Corollary 4.50.] the 2-functor ρ is an equivalence since the 2-functor $Mor_{\infty CAT'}(*, -) : \infty$ Cat $\rightarrow \infty$ Cat preserves tensors.

Notation 7.21. Let $\infty \widehat{\mathsf{Cat}}$ be the $(\infty, 2)$ -category of large ∞ -categories. Let $\Pr^{\mathsf{L}} \subset \infty \widehat{\mathsf{Cat}}$ be the subcategory of the $(\infty, 2)$ -category $\infty \widehat{\mathsf{Cat}}$ of presentable ∞ -categories and left adjoints.

The following remark gives another interpretation of the $(\infty, 2)$ -category Pr^{L} :

Remark 7.22. By [18, Proposition 4.8.1.15.] the underlying ∞ -category of \Pr^{L} admits a closed symmetric monoidal structure such that the inclusion $j:\Pr^{L} \subset \infty \widehat{\mathsf{Cat}}$ of ∞ -categories refines to a lax symmetric monoidal functor. Being symmetric monoidal closed \Pr^{L} is enriched in itself and so by transfer of enrichment along j also enriched in $\infty \widehat{\mathsf{Cat}}$. The lax symmetric monoidal inclusion j gives rise to an inclusion of $(\infty, 2)$ -categories from \Pr^{L} - endowed with the transfered $\infty \widehat{\mathsf{Cat}}$ -enrichment - to $\infty \widehat{\mathsf{Cat}}$ endowed with the enrichment in itself by the internal hom, and so induces an equivalence from the transfered $\infty \widehat{\mathsf{Cat}}$ -enrichment on \Pr^{L} to the subcategory of the $(\infty, 2)$ -category $\infty \widehat{\mathsf{Cat}}$ of presentable ∞ -categories and left adjoint functors.

Corollary 7.23. The $(\infty, 2)$ -category underlying the double ∞ -category PR^{L} is Pr^{L} .

Corollary 7.24. There is a canonical pullback square of $(\infty, 2)$ -precategories:



Corollary 7.25. The completion of the $(\infty, 2)$ -precategory Corr is the full subcategory of the $(\infty, 2)$ -category Pr^{L} spanned by the ∞ -categories of presheaves.

Corollary 7.26. Let C be a small ∞ -category. There is a canonical equivalence of ∞ -categories

$$FUN(N(\mathcal{C}), CORR) \simeq EXP_{\mathcal{C}^{op}}$$

Proof. By Corollary 4.4 and Proposition 7.18 there is a canonical equivalence

 $\mathrm{FUN}(\mathrm{N}(\mathfrak{C}),\mathrm{CORR})\simeq\mathrm{FUN}(\mathrm{N}(\mathfrak{C}),\Delta_{\infty}\mathsf{Cat})\times_{\mathrm{FUN}(\mathrm{N}(\mathfrak{C}),\Delta_{\mathrm{Pr}^{\mathrm{L}}})}\mathrm{FUN}(\mathrm{N}(\mathfrak{C}),\mathrm{PR}^{\mathrm{L}})\simeq$

$$\begin{aligned} &\operatorname{Fun}(\iota(\mathcal{C}),\infty\mathsf{Cat})\times_{\operatorname{Fun}(\iota(\mathcal{C}),\operatorname{Pr}^{\mathrm{L}})}\operatorname{FUN}(\operatorname{N}(\mathcal{C}),\operatorname{PR}^{\mathrm{L}})\simeq\\ & & & & & \\ & & & & \\ & & & & \\ &$$

In the following we prove a refinement of Corollary 7.26.

Notation 7.27. Let $\phi : \mathcal{A} \to \Delta$ be a cartesian fibration. Let $\mathcal{A}^{\vee} \to \Delta^{\mathrm{op}}$ be the corresponding cocartesian fibration classifying the same functor as ϕ .

Definition 7.28. Let $\mathcal{A} \to \Delta, \mathcal{B} \to \Delta$ be double ∞ -categories.

- A lax map of double ∞ -categories $\mathcal{A} \to \mathcal{B}$ is a functors $\mathcal{A}^{\vee} \to \mathcal{B}^{\vee}$ over Δ^{op} preserving cocartesian lifts of inert morphisms of Δ (Definition 3.27).
- A lax normal map of double ∞-categories A → B is a functors A[∨] → B[∨] over Δ^{op} preserving cocartesian lifts of inert morphisms of Δ and cocartesian lifts of morphisms of the form [n] → [0] for n ≥ 0.

The embedding $\text{CORR} \subset \text{CORR}_{\text{Lax}}$ of cartesian fibrations over Δ induces an embedding $\text{CORR}^{\vee} \subset \text{CORR}_{\text{Lax}}^{\vee}$ of cocartesian fibrations over Δ^{op} . Proposition 7.16 and [18, Proposition 7.3.2.6.] give the following corollary:

Corollary 7.29. The embedding $\text{CORR}^{\vee} \subset \text{CORR}^{\vee}_{\text{Lax}}$ of cocartesian fibrations over Δ^{op} admits a right adjoint relative to Δ^{op} .

Example 7.30. The relative right adjoint $\text{CORR}_{\text{Lax}}^{\vee} \to \text{CORR}^{\vee}$ preserves cocartesian lifts of inert morphisms because the left adjoint does. So the right adjoint determines a lax map $\text{CORR}_{\text{Lax}} \to \text{CORR}$ of double ∞ -categories, which is a lax normal map.

Remark 7.31. Let $\mathcal{A} \to \Delta, \mathcal{B} \to \Delta$ be double ∞ -categories. There is a canonical equivalence $\operatorname{FUN}(\mathcal{A}, \mathcal{B}) \simeq \operatorname{Fun}_{\operatorname{Aop}}^{\operatorname{cocart}}(\mathcal{A}^{\vee}, \mathcal{B}^{\vee})$

to the full subcategory of $\operatorname{Fun}_{\Delta^{\operatorname{op}}}(\mathcal{A}^{\vee}, \mathcal{B}^{\vee})$ spanned by the maps $\mathcal{A}^{\vee} \to \mathcal{B}^{\vee}$ of cocartesian fibrations over $\Delta^{\operatorname{op}}$.

Notation 7.32. Let $\mathcal{A} \to \Delta, \mathcal{B} \to \Delta$ be double ∞ -categories.

• Let

$$\operatorname{LaxFun}(\mathcal{A},\mathcal{B}) \subset \operatorname{Fun}_{\Delta^{\operatorname{op}}}(\mathcal{A}^{\vee},\mathcal{B}^{\vee})$$

the full subcategory spanned by the lax maps of double ∞ -categories $\mathcal{A} \to \mathcal{B}$.

• Let

$$\operatorname{NLaxFun}(\mathcal{A},\mathcal{B}) \subset \operatorname{Fun}_{\Delta^{\operatorname{op}}}(\mathcal{A}^{\vee},\mathcal{B}^{\vee})$$

the full subcategory spanned by the lax normal maps of double ∞ -categories $\mathcal{A} \to \mathcal{B}$.

Notation 7.33. Let Act \subset Fun([1], Δ) be the full subcategory of active morphisms, i.e. order preserving maps $[n] \rightarrow [m]$ preserving the minimum and maximum.

Let $\mathcal{C} \to \Delta$ be a double ∞ -category. By [9, Proposition 3.102.] evaluation at the target

(10)
$$\operatorname{Act}^{\operatorname{op}} \times_{\Delta^{\operatorname{op}}} \mathcal{C}^{\vee} \to \operatorname{Act}^{\operatorname{op}} \to \Delta^{\operatorname{op}}$$

is a cocartesian fibration, where the pullback is formed along evaluation at the source.

Notation 7.34. Let $\mathcal{C} \to \Delta$ be a double ∞ -category. Let $\text{Env}(\mathcal{C}) \to \Delta$ be the cartesian fibration corresponding to the cocartesian fibration (10).

By [9, Proposition 3.102.] the cartesian fibration $\operatorname{Env}(\mathcal{C}) \to \Delta$ is a double ∞ -category and for every double ∞ -category $\mathcal{D} \to \Delta$ the induced functor $\operatorname{FUN}(\operatorname{Env}(\mathcal{C}), \mathcal{D}) \to \operatorname{LaxFun}(\mathcal{C}, \mathcal{D})$ is an equivalence.

Example 7.35. The diagonal embedding $\Delta \subset \text{Act}$ induces an embedding $\mathcal{C}^{\vee} \subset \text{Act}^{\text{op}} \times_{\Delta^{\text{op}}} \mathcal{C}^{\vee}$ over Δ^{op} that determines a lax map of double ∞ -categories $\mathcal{C} \to \text{Env}(\mathcal{C})$.

Remark 7.36. By [9, Lemma 3.103.] the embedding $\mathcal{C}^{\vee} \subset \operatorname{Act}^{\operatorname{op}} \times_{\Delta^{\operatorname{op}}} \mathcal{C}^{\vee}$ over $\Delta^{\operatorname{op}}$ admits a left adjoint relative to $\Delta^{\operatorname{op}}$, which is a map of cocartesian fibrations over $\Delta^{\operatorname{op}}$. So this left adjoint corresponds to a map $\operatorname{Env}(\mathcal{C}) \to \mathcal{C}$ of cartesian fibrations over Δ , in other words to a map of double ∞ -categories.

Remark 7.37. Let $\mathcal{C} \to \Delta$ be a double ∞ -category. There is a canonical equivalence $\operatorname{Env}(\mathcal{C})_{[0]} \simeq \mathcal{C}_{[0]}^{\vee} \simeq \mathcal{C}_{[0]}$ since $\operatorname{Act}_{[0]/}$ is contractible. Consequently, $\operatorname{Env}(\mathcal{C}) \to \Delta$ is an $(\infty, 2)$ -precategory if $\mathcal{C} \to \Delta$ is an $(\infty, 2)$ -precategory.

Remark 7.38. Let $\mathcal{C} \to \Delta$ be a double ∞ -category. Every morphism of $\text{Env}(\mathcal{C})$, i.e. object of $\text{Env}(\mathcal{C})_{[1]}$, lies in the image of the restricted composition functor

 $\mathcal{C}_{[1]} \times_{\mathcal{C}_{[0]}} \ldots \times_{\mathcal{C}_{[0]}} \mathcal{C}_{[1]} \subset \operatorname{Env}(\mathcal{C})_{[1]} \times_{\operatorname{Env}(\mathcal{C})_{[0]}} \ldots \times_{\operatorname{Env}(\mathcal{C})_{[0]}} \operatorname{Env}(\mathcal{C})_{[1]} \to \operatorname{Env}(\mathcal{C})_{[1]}.$

In other words, any horizontal morphism of $\text{Env}(\mathcal{C}) \to \Delta$ is of the form $\alpha_1 \circ \ldots \circ \alpha_n$ for some $n \ge 0$ and composable horizontal morphisms $\alpha_1, \ldots, \alpha_n$ of \mathcal{C} , where \circ is the horizontal composition.

Remark 7.39. Let $\mathcal{C} \to \Delta$ be an $(\infty, 2)$ -category. Then $\operatorname{Env}(\mathcal{C}) \to \Delta$ is an $(\infty, 2)$ -category: by Remark 7.37 the canonical map $\iota(\operatorname{Env}(\mathcal{C})_{[0]}) \to \iota(\operatorname{Env}(\mathcal{C})_{[1]})$ factors as $\iota(\operatorname{Env}(\mathcal{C})_{[0]}) \simeq \iota(\mathcal{C}_{[0]}) \to \iota(\mathcal{C}_{[1]}) \subset \iota(\operatorname{Env}(\mathcal{C})_{[1]})$ and so is an embedding if \mathcal{C} is an $(\infty, 2)$ -category. Hence in view of Remark 7.38 it suffices to observe that for every n, m ≥ 0 and composable horizontal morphisms $\alpha_1, ..., \alpha_n, \beta_1, ..., \beta_m$ of $\mathcal{C} \to \Delta$ such that the horizontal composition $\alpha_1 \circ ... \circ \alpha_n \circ \beta_1 \circ ... \circ \beta_m$ in $\operatorname{Env}(\mathcal{C}) \to \Delta$ is the 0-fold composition, we find that n+m = 0 so that n = m = 0.

Notation 7.40. Let $\phi : \mathcal{C} \to \Delta$ be an ∞ -precategory and \mathcal{W} a set of morphisms of $\mathcal{C}_{[1]}$ corresponding to 2-morphisms of $\mathcal{C} \to \Delta$. For every $n \ge 1$ let \mathcal{W}_n be the set of morphisms of $\mathcal{C}_{[n]}$ whose image under the functor $\mathcal{C}_{[n]} \to \mathcal{C}_{[1]}$ induced by the map $[1] \cong \{i - 1 < i\} \to [n]$ for any $0 \le i < n$ is a morphism of \mathcal{W} . Let \mathcal{W}_0 be the set of all morphisms of $\mathcal{C}_{[0]}$. Let $\overline{\mathcal{W}}$ be the smallest set of morphisms of \mathcal{C} containing all images of morphisms of \mathcal{W}_n in \mathcal{C} for $n \ge 0$.

Let $\phi : \mathbb{C} \to \Delta$ be an ∞ -precategory and \mathcal{W} a set of morphisms of $\mathbb{C}_{[1]}$. Let $\mathbb{C}[\mathcal{W}] := \mathbb{C}[\mathcal{W}]$ be the localization of the ∞ -category \mathbb{C} with respect to the set $\overline{\mathcal{W}}$. The functor $\phi : \mathbb{C} \to \Delta$ induces a functor $\mathbb{C}[\mathcal{W}] \to \Delta$, which by [16, 2.1.4. Proposition] is a cartesian fibration whose

24

fiber over any $[n] \in \Delta$ is the localization $C_{[n]}[\mathcal{W}_n]$. In particular, $C[\mathcal{W}]_{[0]} \simeq C_{[0]}[\mathcal{W}_0] \simeq C_{[0]}$. By the next lemma the cartesian fibration $C[\mathcal{W}] \to \Delta$ is an ∞ -precategory.

Lemma 7.41. Let \mathcal{C} be a space, $\mathcal{A} \to \mathcal{C}, \mathcal{B} \to \mathcal{C}$ functors and \mathcal{W} a set of morphisms of \mathcal{A} and \mathcal{V} a set of morphisms of \mathcal{B} containing all equivalences. The canonical functor θ : $(\mathcal{A} \times_{\mathcal{C}} \mathcal{B})[\mathcal{W} \times_{\mathcal{C}} \mathcal{V}] \to \mathcal{A}[\mathcal{W}] \times_{\mathcal{C}} \mathcal{B}[\mathcal{V}]$ is an equivalence.

Proof. The functor θ is a functor over the space C and so is an equivalence if it induces an equivalence on the fiber over every $Z \in C$. For every $Z \in C$ let W_Z be the set of morphisms of \mathcal{A}_Z whose image in \mathcal{A} belongs to \mathcal{W} . Similarly, we define \mathcal{V}_Z . The functor θ induces on the fiber over Z the canonical functor

$$(\mathcal{A} \times_{\mathfrak{C}} \mathfrak{B})[\mathcal{W} \times_{\mathfrak{C}} \mathcal{V}]_{\mathbb{Z}} \to \mathcal{A}[\mathcal{W}]_{\mathbb{Z}} \times \mathfrak{B}[\mathcal{V}]_{\mathbb{Z}},$$

which by [16, 2.1.4. Proposition] identifies with the canonical functor

$$(\mathcal{A}_{Z} \times \mathcal{B}_{Z})[\mathcal{W}_{Z} \times \mathcal{V}_{Z}] \to \mathcal{A}_{Z}[\mathcal{W}_{Z}] \times \mathcal{B}_{Z}[\mathcal{V}_{Z}].$$

The latter is an equivalence by universal property of the localization using that \mathcal{W}, \mathcal{V} contain all equivalences.

Definition 7.42. Let $\mathcal{C} \to \Delta$ be an ∞ -precategory. Let $\operatorname{NEnv}(\mathcal{C}) \to \Delta$ be the localization of $\operatorname{Env}(\mathcal{C}) \to \Delta$ with respect to the smallest set of $\operatorname{Env}(\mathcal{C})_{[1]}$ containing the morphisms $\operatorname{id}_X^{\operatorname{Env}(\mathcal{C})} \to \operatorname{id}_X^{\mathcal{C}}$ of $\operatorname{Env}(\mathcal{C})_{[1]}$ for $X \in \mathcal{C}_{[0]}$ and all equivalences of $\operatorname{Env}(\mathcal{C})_{[1]}$ and closed under horizontal composition of the ∞ -precategory $\operatorname{Env}(\mathcal{C}) \to \Delta$.

Remark 7.43. The localization $\operatorname{Env}(\mathcal{C})^{\vee} \to \mathcal{C}^{\vee}$ relative to $\Delta^{\operatorname{op}}$ whose left adjoint inverts local equivalences, gives rise to a localization $\operatorname{NEnv}(\mathcal{C})^{\vee} \to \mathcal{C}^{\vee}$ relative to $\Delta^{\operatorname{op}}$. Hence the right adjoint is an embedding $\mathcal{C}^{\vee} \subset \operatorname{NEnv}(\mathcal{C})^{\vee}$ that factors as $\mathcal{C}^{\vee} \to \operatorname{Env}(\mathcal{C})^{\vee} \to \operatorname{NEnv}(\mathcal{C})^{\vee}$ and so corresponds to a lax normal map $\mathcal{C} \to \operatorname{NEnv}(\mathcal{C})$ of double ∞ -categories.

Remark 7.44. By definition the canonical functor $\operatorname{Env}(\mathcal{C}) \to \operatorname{NEnv}(\mathcal{C})$ over Δ induces an equivalence $\operatorname{Env}(\mathcal{C})_{[0]} \simeq \operatorname{NEnv}(\mathcal{C})_{[0]}$. Thus the embedding $\mathcal{C}^{\vee} \subset \operatorname{NEnv}(\mathcal{C})^{\vee}$ over Δ induces an equivalence $\mathcal{C}_{[0]} \simeq \operatorname{NEnv}(\mathcal{C})_{[0]}$.

Remark 7.45. Let $\mathcal{C} \to \Delta$ be an $(\infty, 2)$ -category. Then $\operatorname{NEnv}(\mathcal{C}) \to \Delta$ is an $(\infty, 2)$ -category by a similar argument like in Remark 7.39: by Remark 7.44 the canonical map $\iota(\operatorname{NEnv}(\mathcal{C})_{[0]}) \to \iota(\operatorname{NEnv}(\mathcal{C})_{[1]})$ factors as $\iota(\operatorname{NEnv}(\mathcal{C})_{[0]}) \simeq \iota(\mathcal{C}_{[0]}) \to \iota(\mathcal{C}_{[1]}) \subset \iota(\operatorname{NEnv}(\mathcal{C})_{[1]})$ and so is an embedding if \mathcal{C} is an $(\infty, 2)$ -category. So it suffices to observe that for every $n, m \ge 0$ and composable horizontal morphisms $\alpha_1, ..., \alpha_n, \beta_1, ..., \beta_m$ of $\mathcal{C} \to \Delta$ different from identities such that the horizontal composition $\alpha_1 \circ ... \circ \alpha_n \circ \beta_1 \circ ... \circ \beta_m$ in $\operatorname{NEnv}(\mathcal{C}) \to \Delta$ is the 0-fold composition, we have that n + m = 0 and so n = m = 0.

Remark 7.46. Let $\mathcal{C} \to \Delta, \mathcal{D} \to \Delta$ be double ∞ -categories. By definition the induced functor FUN(NEnv(\mathcal{C}), \mathcal{D}) \to FUN(Env(\mathcal{C}), \mathcal{D}) is fully faithful and the equivalence

$$\mathrm{FUN}(\mathrm{Env}(\mathcal{C}), \mathcal{D}) \simeq \mathrm{LaxFun}(\mathcal{C}, \mathcal{D})$$

restricts to an equivalence

 $\operatorname{FUN}(\operatorname{NEnv}(\mathcal{C}), \mathcal{D}) \to \operatorname{NLaxFun}(\mathcal{C}, \mathcal{D}).$

Proposition 7.47. Let \mathcal{C} be a small $(\infty, 2)$ -category. There is a canonical equivalence

 $\operatorname{LaxFun}(\mathcal{C},\infty \operatorname{CAT}) \simeq \operatorname{LoCART}_{\tau^*\mathcal{C}}.$

Proof. There is a chain of canonical equivalences of ∞ -categories

$$LaxFun(\mathcal{C}, \infty CAT) \simeq FUN(NEnv(\mathcal{C}), \infty CAT) \simeq CART_{\tau^*NEnv(\mathcal{C})} \simeq$$

 $\operatorname{CART}_{\operatorname{NEnv}(\tau^*\mathcal{C})} \simeq \operatorname{LoCART}_{\tau^*\mathcal{C}}.$

The first equivalence holds by the universal property of NEnv(\mathcal{C}) $\rightarrow \Delta$ of Remark 7.46. The second equivalence is straightening of cartesian fibrations. The third equivalence is induced by the canonical equivalence $\tau^* \text{NEnv}(\mathcal{C}) \simeq \text{NEnv}(\tau^* \mathcal{C})$ over Δ , which follows immediately from the construction of Env(\mathcal{C}) $\rightarrow \Delta$. The last equivalence is [3, Theorem B.4.3.].

Corollary 7.48. Let \mathcal{C} be a small ∞ -category. There is a canonical equivalence LaxFun(N(\mathcal{C}), CORR) $\simeq \infty \operatorname{Cat}_{/\mathcal{C}^{\operatorname{op}}}$.

Proof. By Proposition 7.47 (applied to a larger universe) there is a canonical equivalence

 $LaxFun(N(\mathcal{C}), \infty \widehat{CAT}) \simeq Lo \widehat{CART}_{\mathcal{C}^{op}}$

that restricts to an equivalence

$$\operatorname{LaxFun}(\operatorname{N}(\operatorname{\mathcal{C}}),\operatorname{PR}^{\operatorname{L}})\simeq\operatorname{LoCART}^{\operatorname{L}}_{\operatorname{C^{\operatorname{op}}}}.$$

Hence by Corollary 4.4 and Proposition 7.18 we obtain a canonical equivalence

 $\operatorname{LaxFun}(N(\mathcal{C}), \operatorname{CORR}) \simeq \operatorname{LaxFun}(N(\mathcal{C}), \Delta_{\infty\mathsf{Cat}}) \times_{\operatorname{LaxFun}(N(\mathcal{C}), \Delta_{\operatorname{Pr}^L})} \operatorname{LaxFun}(N(\mathcal{C}), \operatorname{PR}^L) \simeq$

$$\begin{aligned} &\operatorname{Fun}(\iota(\mathcal{C}),\infty\mathsf{Cat})\times_{\operatorname{Fun}(\iota(\mathcal{C}),\operatorname{Pr}^{\mathrm{L}})}\operatorname{LaxFun}(\mathrm{N}(\mathcal{C}),\operatorname{PR}^{\mathrm{L}}) \\ & & & & \\ & & & & \\ & & & \\ & & & & \\ & & & \\ & & & & \\$$

References

- [1] Fernando Abellán, Andrea Gagna, and Rune Haugseng. Straightening for lax transformations and adjunctions of $(\infty, 2)$ -categories. arXiv: 2404.03971, 2024.
- [2] David Ayala and John Francis. Fibrations of ∞ -categories. High. Struct., 4(1):168–265, 2020.
- [3] David Ayala, Aaron Mazel-Gee, and Nick Rozenblyum. Stratified noncommutative geometry. Memoirs of the American Mathematical Society, 2019.
- [4] Clark Barwick, Emanuele Dotto, Saul Glasman, Denis Nardin, and Jay Shah. Parametrized higher category theory and higher algebra: Exposé i – elements of parametrized higher category theory.
- [5] Clark Barwick and Jay Shah. Fibrations in ∞ -category theory. arXiv: 1607.04343.
- [6] David Gepner and Rune Haugseng. Enriched ∞-categories via non-symmetric ∞-operads. Adv. Math., 279:575-716, 2015.
- [7] David Gepner, Rune Haugseng, and Thomas Nikolaus. Lax colimits and free fibrations in ∞-categories. Documenta Mathematica, 22, 01 2015.
- [8] Hadrian Heine. A topological model for cellular motivic spectra. arXiv: 1712.00521.
- [9] Hadrian Heine. An equivalence between enriched ∞ -categories and ∞ -categories with weak action. arXiv:2009.02428, 2023.
- [10] Hadrian Heine. An equivalence between enriched ∞-categories and ∞-categories with weak action. Advances in Mathematics, 417:108941, 2023.
- [11] Hadrian Heine. A monadicity theorem for higher algebraic structures. arXiv:1712.00555, 2023.
- [12] Hadrian Heine. The higher algebra of weighted colimits. arXiv: 2406.08925, 2024.
- [13] Hadrian Heine. On bi-enriched ∞-categories. arXiv: 2406.09832, 2024.
- [14] Hadrian Heine, Alejo Lopez-Avila, and Markus Spitzweck. Infinity categories with duality and hermitian multiplicative infinite loop space machines. arXiv: 1610.10043, 2024.
- [15] Hadrian Heine, Markus Spitzweck, and Paula Verdugo. Real k-theory for waldhausen infinity categories with genuine duality. arXiv: 1911.11682, 2021.
- [16] Vladimir Hinich. Dwyer-kan localization revisited. Homology, Homotopy and Applications, 18, 11 2013.
- [17] Vladimir Hinich. Yoneda lemma for enriched ∞ -categories. Advances in Mathematics, 367:107129, 2020.
- [18] Jacob Lurie. Higher Algebra. available at http://www.math.harvard.edu/ lurie/.

- [19] Jacob Lurie. Higher topos theory, volume 170 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 2009.
- [20] Lyne Moser, Nima Rasekh, and Martina Rovelli. An (∞, n) -categorical straightening-unstraightening construction. arXiv: 2307.07259, 2023.
- [21] Joost Nuiten. On straightening for segal spaces. arXiv: 2108.11431, 2023.