

A NOTE ON INVARIANTS OF FOLIATED 3-SPHERE BUNDLES

NILS PRIGGE

ABSTRACT. In this note we prove that $H^*(\mathrm{BSO}(4); \mathbb{Q})$ injects into the group cohomology of $\mathrm{Diff}^+(\mathbb{S}^3)$ with rational coefficients. The proof is based on an idea of Nariman who proved that the monomials in the Euler and Pontrjagin classes are nontrivial in $H^*(\mathrm{BDiff}_+^\delta(\mathbb{S}^{2n-1}); \mathbb{Q})$.

1. INTRODUCTION

Let $\mathrm{BS}\Gamma_d$ be the classifying space of Haefliger structures for codimension d foliations that are transversely oriented and denote by $\nu : \mathrm{BS}\Gamma_d \rightarrow \mathrm{BGL}_d^+(\mathbb{R})$ the map that records the normal bundle of the foliation (see [Hae71] for details). We consider the space of bundle maps $\mathrm{Bun}(TM, \nu^*\gamma_d)$ where γ_d denotes the universal oriented vector bundle over $\mathrm{BGL}_d^+(\mathbb{R})$. It has a $\mathrm{Diff}_+(M)$ -action by precomposition with the differential of a diffeomorphism, and Nariman proved (cf. [Nar17] and [Nar23, Cor. 2.5]) that the map $\mathrm{BDiff}_+^\delta(M) \rightarrow \mathrm{BDiff}_+(M)$ induced by the inclusion of $\mathrm{Diff}_+^\delta(M)$, the group of diffeomorphisms with the discrete topology, factors through an acyclic map

$$\beta : \mathrm{BDiff}_+^\delta(M) \longrightarrow \mathrm{Bun}(TM, \nu^*\gamma_d) // \mathrm{Diff}_+(M).$$

This means in particular that β induces an isomorphism

$$(1.1) \quad H^*(\mathrm{BDiff}_+^\delta(M); \mathbb{Q}) \cong H^*(\mathrm{Bun}(TM, \nu^*\gamma_d) // \mathrm{Diff}_+(M); \mathbb{Q}),$$

and under this isomorphism the map on cohomology induced by $\mathrm{BDiff}_+^\delta(M) \rightarrow \mathrm{BDiff}_+(M)$ agrees with the map induced by the projection

$$p : \mathrm{Bun}(TM, \nu^*\gamma_d) // \mathrm{Diff}_+(M) \rightarrow \mathrm{BDiff}_+(M).$$

If a Lie group G acts smoothly on M , there is a commutative diagram

$$(1.2) \quad \begin{array}{ccc} \mathrm{Bun}(TM, \nu^*\gamma_d) // G & \longrightarrow & \mathrm{Bun}(TM, \nu^*\gamma_d) // \mathrm{Diff}_+(M) \\ \downarrow & & \downarrow \\ \mathrm{BG} & \longrightarrow & \mathrm{BDiff}_+(M) \end{array}$$

and Nariman shows in [Nar23, Sect. 3] that if G is a torus acting freely on M , then $\mathrm{Bun}(TM, \nu^*\gamma_d)$ has a G -fixed point so the left vertical map has a section. Hence, the map on cohomology induced by $\mathrm{BG} \rightarrow \mathrm{BDiff}_+(M)$ factors as

$$H^*(\mathrm{BDiff}_+(M)) \longrightarrow H^*(\mathrm{BDiff}_+^\delta(M)) \longrightarrow H^*(\mathrm{BG}).$$

For an odd sphere one can define classes $e, p_1, \dots, p_{n-1} \in H^*(\mathrm{BDiff}_+(\mathbb{S}^{2n-1}))$ that pull back to the Euler and Pontrjagin classes along the map $\mathrm{BSO}(2n) \rightarrow \mathrm{BDiff}_+(\mathbb{S}^{2n-1})$ induced by the action of $\mathrm{SO}(2n)$ on \mathbb{S}^{2n-1} . If we further restrict to the free S^1 -action, the image of the monomials in the Euler and Pontrjagin classes in the cohomology of BS^1 are nontrivial and it follows that they are nontrivial in $H^*(\mathrm{BDiff}_+^\delta(\mathbb{S}^{2n-1}))$ [Nar23, Thm 1.2]. But this poses the question about injectivity of the map

$$(1.3) \quad \mathbb{Q}[e, p_1, \dots, p_{n-1}] \subset H^*(\mathrm{BDiff}_+(\mathbb{S}^{2n-1}); \mathbb{Q}) \rightarrow H^*(\mathrm{BDiff}_+^\delta(\mathbb{S}^{2n-1}); \mathbb{Q}).$$

The obvious obstacle to making a statement about injectivity of (1.3) is that the Krull dimension of $H^*(BS^1; \mathbb{Q})$ is too small. Instead, we should use the action of the maximal torus $T^n \subset \mathrm{SO}(2n)$ on \mathbb{S}^{2n-1} . However, T^n is not acting freely on \mathbb{S}^{2n-1} so that one does not know whether $\mathrm{Bun}(T\mathbb{S}^{2n-1}, \nu^* \gamma_{2n-1})$ has a T^n -fixed point from Nariman's construction. The main result of this note circumvents the need to construct actual fixed points for $n = 2$.

Theorem 1.4. *The induced map on cohomology*

$$(1.5) \quad H^*(BT^2; \mathbb{Q}) \longrightarrow H^*(\mathrm{Bun}(T\mathbb{S}^3, \nu^* \gamma_3) // T^2; \mathbb{Q})$$

is an injection.

Corollary 1.6. *The subring $\mathbb{Q}[p_1, e] \subset H^*(\mathrm{BDiff}_+(\mathbb{S}^3); \mathbb{Q})$ defined by the Euler and Pontrjagin class injects into $H^*(\mathrm{BDiff}_+^\delta(\mathbb{S}^3); \mathbb{Q})$*

Proof. We know that $\mathbb{Q}[p_1, \dots, p_{n-1}, e] \subset H^*(\mathrm{BDiff}_+(\mathbb{S}^{2n-1}; \mathbb{Q}))$ injects into $H^*(BT^n; \mathbb{Q})$ under the map $BT^n \rightarrow \mathrm{BDiff}_+(\mathbb{S}^{2n-1})$. It then follows from commutativity of (1.2) and Theorem 1.4 that the composite

$$\mathbb{Q}[p_1, e] \rightarrow H^*(\mathrm{BDiff}_+^\delta(\mathbb{S}^3)) \rightarrow H^*(\mathrm{Bun}(T\mathbb{S}^3, \nu^* \gamma_3) // \mathrm{Diff}_+(\mathbb{S}^3)).$$

is an injection, so that the first map is injective as well. \square

Remark 1.7. In a forthcoming paper [Pri] that previously appeared as an appendix to [Nar23] we proved that $H^*(\mathrm{BSO}(4); \mathbb{R})$ injects into the smooth cohomology $H_{\mathrm{sm}}^*(\mathrm{Diff}_+(\mathbb{S}^3); \mathbb{R})$ using the method developed by Haefliger [Hae78]. This statement follows from Corollary 1.6, but the real homotopy theory computation is still interesting as it showcases an interesting relation between p_1^2 and a continuously varying cohomology class as pointed out by Morita (cf. [Nar23]).

The idea for the proof of Theorem 1.4 is quite simple and we learned it from [AP93, Example 3.1.16]. Given a space X with a torus action so that infinitely many distinct subtori $K_i \subset T$ of codimension 1 occur as stabilizers, the projection $p: X//T \rightarrow BT$ induces an injection on cohomology. This is because the map induced by the projection $p: X//K_i \rightarrow BK_i$ is an injection on cohomology (as there is a K_i -fixed point) and from the commutativity of the diagram

$$\begin{array}{ccc} H^*(X//K_i) & \longleftarrow & H^*(X//T) \\ \uparrow & & \uparrow \\ H^*(BK_i) & \longleftarrow & H^*(BT) \end{array}$$

it follows that

$$\ker(H^*(BT) \rightarrow H^*(X//T)) \subset \ker(H^*(BT) \rightarrow H^*(BK_i)),$$

where $\ker(H^*(BT) \rightarrow H^*(BK_i))$ is an ideal generated by a linear polynomial (with integral coefficients) $f_i \in H^2(BT)$. Hence, an element $x \in \ker(H^*(BT) \rightarrow H^*(X//T))$ is divisible by infinitely many distinct linear polynomials f_i as the tori K_i are distinct and therefore $x = 0$.

So the idea of the proof is to construct bundle maps $\tau \in \mathrm{Bun}(T\mathbb{S}^{2n-1}, \nu^* \gamma_{2n-1})$ with prescribed stabilizers, which as it turns out is still difficult to do. However, for $n = 2$ we are able to show that there exists infinitely many tori $K \subset T^2$ with the property that

$$H^*(BK) \hookrightarrow H^*(\mathrm{Bun}(T\mathbb{S}^3, \nu^* \gamma_3) // K).$$

without using actual fixed points of the action, which proves Theorem 1.4.

2. MAIN RESULT

For integers $m, n \in \mathbb{Z}$ we denote by $c_{m,n}: S^1 \rightarrow T^2$ the group homomorphism defined by $c_{m,n}(\lambda) = (\lambda^m, \lambda^n)$. Codimension 1 tori are indexed by integers $(m, n) \in \mathbb{Z}^2 / ((m, n) \sim -(m, n))$ with $\gcd(m, n) = 1$ as the image $K_{m,n} = \text{im}(c_{m,n})$. Theorem 1.4 follows from the following statement which we prove in this section.

Proposition 2.1. *Let $m \in \mathbb{N}$ be odd and $n = m + 2$, then the map*

$$H^*(BK_{m,n}; \mathbb{Q}) \longrightarrow H^*(\text{Bun}(T\mathbb{S}^3, \nu^* \gamma_3) // K_{m,n}; \mathbb{Q})$$

is injective.

We consider the T^2 -action on \mathbb{S}^3 given by $(\lambda_1, \lambda_2) \cdot (z_1, z_2) = (\lambda_1 \cdot z_1, \lambda_2 \cdot z_2)$, and we denote by $(\mathbb{S}^3, K_{m,n})$ the 3-sphere with the restricted $K_{m,n}$ -action. For $m = n = 1$ this corresponds to the usual free S^1 -action on \mathbb{S}^3 which we denote by (\mathbb{S}^3, S^1) .

Lemma 2.2. *The map*

$$\pi_{m,n}: (\mathbb{S}^3, S^1) \rightarrow (\mathbb{S}^3, K_{m,n}), \quad \pi_{m,n}(z_1, z_2) = \frac{(z_1^m, z_2^n)}{\|(z_1^m, z_2^n)\|}$$

is equivariant with respect to $c_{m,n}$.

The pullback $\pi_{m,n}^* T\mathbb{S}^3$ is an S^1 -equivariant vector bundle over (\mathbb{S}^3, S^1) and there is a canonical bundle map $\tilde{\pi}_{m,n}: \pi_{m,n}^* T\mathbb{S}^3 \rightarrow T\mathbb{S}^3$ covering $\pi_{m,n}$ which is equivariant with respect to $c_{m,n}$. Precomposition with $\tilde{\pi}_{m,n}$ defines a map

$$(2.3) \quad \psi: \text{Bun}(T\mathbb{S}^3, \nu^* \gamma_3) \longrightarrow \text{Bun}(\pi_{m,n}^* T\mathbb{S}^3, \nu^* \gamma_3)$$

which is equivariant by construction with respect to $c_{m,n}^{-1}: K_{m,n} \rightarrow S^1$ if the inverse exists (i.e. if $\gcd(m, n) = 1$).

Lemma 2.4. *The map ψ in (2.3) is a rational equivalence.*

Proof. The tangent bundle $T\mathbb{S}^3$ is (non-equivariantly) trivial and hence so is the pullback $\pi_{m,n}^* T\mathbb{S}^3$. This implies that the space of bundle maps is (non-equivariantly) homotopy equivalent to $\text{map}(\mathbb{S}^3, \overline{\text{B}\Gamma}_3)$, where $\overline{\text{B}\Gamma}_d$ denotes the homotopy fibre of $\nu: \text{B}\text{S}\Gamma_d \rightarrow \text{BGL}_d^+(\mathbb{R})$. Since $\overline{\text{B}\Gamma}_3$ is 4-connected [Thu74], these mapping spaces are connected and ψ corresponds to the map of mapping spaces induced by precomposition with $\pi_{m,n}$, which is a rational equivalence since $\pi_{m,n}$ is (this follows directly by inspection of the rational models of mapping spaces, for example cf. [Ber15]). \square

In the following, we denote by $V_k = (\mathbb{C}, \rho_k)$ for $k \in \mathbb{Z}$ the complex S^1 -representation given by $\rho_k(\lambda)(z) = \lambda^k \cdot z$. We obtain T^2 -representations by pulling back along the group homomorphism $\Delta: T^2 \rightarrow S^1$ given by $\Delta(\lambda_1, \lambda_2) = \lambda_2 / \lambda_1$. In the next section we show that $T\mathbb{S}^3$ is T^2 -equivariantly V -trivial for $V = \Delta^* V_1 \oplus \mathbb{R}$, where \mathbb{R} denotes the trivial representation (see Lemma 3.1). This implies that $\pi_{m,n}^* T\mathbb{S}^3$ is $c_{m,n}^* V$ -trivial and we use the following simple criterion to show that it is also \mathbb{R}^3 -trivial.

Lemma 2.5. *Let G be a Lie group acting on X and V, W be two G -representations. There is an isomorphism of equivariant G -vector bundles $X \times V \cong_G X \times W$ if and only if there is a G -equivariant map $X \rightarrow \text{Iso}(V, W)$, where $\text{Iso}(V, W)$ denotes the space of isomorphisms with respect to the conjugation G -action.*

Proposition 2.6. *Let $n = m + 2$, then the pullback $\pi_{m,n}^* T\mathbb{S}^3$ is isomorphic to the trivial S^1 -bundle $\mathbb{S}^3 \times \mathbb{R}^3$, i.e. S^1 acts diagonally on $\mathbb{S}^3 \times \mathbb{R}^3$ and trivially on \mathbb{R}^3 .*

Proof. By Lemma 3.1 the tangent bundle $T\mathbb{S}^3$ is T^2 -equivariantly V -trivial for $V = \mathbb{R} \oplus \Delta^* V_1$, and hence $\pi_{m,n}^* T\mathbb{S}^3 \cong_{S^1} \mathbb{S}^3 \times (\mathbb{R} \oplus c_{m,n}^* \Delta^* V_1)$. The statement follows from Lemma 2.5 if we can construct an S^1 -equivariant map

$$(2.7) \quad \mathbb{S}^3 \longrightarrow \text{Iso}(\mathbb{R}^3, \mathbb{R} \oplus c_{m,n}^* \Delta^* V_1).$$

We identify $\text{Iso}(\mathbb{R}^3, V) \cong \text{GL}_3(\mathbb{R})$ and denote by $D(\lambda) \in \text{SO}(2)$ the rotation corresponding to $\lambda \in S^1$. Then under this identification the S^1 -action is given left multiplication with

$$\begin{pmatrix} 1 & 0 \\ 0 & D(\lambda^{n-m}) \end{pmatrix} \in \text{SO}(3).$$

If $n - m = 2$ then the double cover $f: \mathbb{S}^3 \rightarrow \text{SO}(3) \subset \text{GL}_3(\mathbb{R})$ is S^1 -equivariant with respect to this action by Lemma 3.4 which concludes the proof. \square

Proof of Prop. 2.1. It follows from Proposition 2.6 there is an equivariant homeomorphism $\text{Bun}(\pi_{m,n}^* T\mathbb{S}^3, \nu^* \gamma_3) \approx_{S^1} \text{Bun}(\mathbb{S}^3 \times \mathbb{R}^3, \nu^* \gamma_3)$. The latter has a fixed point by the same argument as in [Nar23, Sect. 3] so that

$$H^*(\text{BS}^1) \rightarrow H^*(\text{Bun}(\pi_{m,n}^* T\mathbb{S}^3, \nu^* \gamma_3) // S^1)$$

is injective. If $m \in \mathbb{N}$ is odd and $n = m + 2$ then $\text{gcd}(m, n) = 1$ and $c_{m,n}: S^1 \rightarrow K_{m,n}$ is an isomorphism, and by Proposition 2.4 we have a commutative diagram

$$(2.8) \quad \begin{array}{ccc} H^*(\text{Bun}(\pi_{m,n}^* T\mathbb{S}^3, \nu^* \gamma_3) // S^1) & \xrightarrow[\cong]{\psi^*} & H^*(\text{Bun}(T\mathbb{S}^3, \nu^* \gamma_3) // K_{m,n}) \\ \uparrow & & \uparrow \\ H^*(\text{BS}^1) & \xrightarrow[\cong]{H^*(c_{m,n}^{-1})} & H^*(\text{BK}_{m,n}) \end{array}$$

which proves the statement. \square

3. TWO FACTS ABOUT \mathbb{S}^3

We prove two elementary statements about the 3-sphere that were used in Proposition 2.6, both relying on the fact that \mathbb{S}^3 can be identified with the group of unit quaternions.

Lemma 3.1. *The tangent bundle $T\mathbb{S}^3$ is T^2 -equivariantly V trivial for $V = \Delta^* V_1 \oplus \mathbb{R}$, i.e. there is an isomorphism of T^2 -equivariant vector bundles $T\mathbb{S}^3 \cong_{T^2} \mathbb{S}^3 \times V$.*

Proof. Choose an inner product on \mathbb{H} with orthonormal basis given by $1, i, j, k$ and consider \mathbb{S}^3 as the set of unit quaternions $\{z_1 + z_2 j \mid z_i \in \mathbb{C}, |z_1|^2 + |z_2|^2 = 1\} \subset \mathbb{H}$ and $T\mathbb{S}^3 = \{(w_1, w_2) \in \mathbb{S}^3 \times \mathbb{H} \mid w_2 \perp w_1\} \subset \mathbb{S}^3 \times \mathbb{H}$. We choose a basis of $T_1 \mathbb{S}^3$ given by $e_1 = (1, i)$, $e_2 = (1, j)$ and $e_3 = (1, k)$ and define a trivialization of $T\mathbb{S}^3$ by

$$\tau: \mathbb{S}^3 \times T_1 \mathbb{S}^3 \rightarrow T\mathbb{S}^3, \quad (p, X_1) \mapsto Dl_p(X_1)$$

where $l_w: \mathbb{S}^3 \rightarrow \mathbb{S}^3$ denotes left multiplication with $w = z_1 + z_2 j \in \mathbb{S}^3$ so that

$$\begin{aligned} \tau(w, e_1) &= (w, z_1 i - z_2 k), \\ \tau(w, e_2) &= (w, -z_2 + z_1 j), \\ \tau(w, e_3) &= (w, z_2 i + z_1 k). \end{aligned}$$

The tangent bundle $T\mathbb{S}^3$ has a left T^2 -action via the differential and one computes that

$$\begin{aligned} D\mu_{(\lambda_1, \lambda_2)} \tau(w, e_1) &= ((\lambda_1, \lambda_2) \cdot w, \lambda_1 z_1 i - \lambda_2 z_2 k), \\ D\mu_{(\lambda_1, \lambda_2)} \tau(w, e_2) &= ((\lambda_1, \lambda_2) \cdot w, -\lambda_1 z_2 + \lambda_2 z_1 j), \\ D\mu_{(\lambda_1, \lambda_2)} \tau(w, e_3) &= ((\lambda_1, \lambda_2) \cdot w, \lambda_1 z_2 i + \lambda_2 z_1 k), \end{aligned}$$

where $\mu_{\lambda_1, \lambda_2}: \mathbb{S}^3 \rightarrow \mathbb{S}^3$ denotes action of $(\lambda_1, \lambda_2) \in T^2$.

We need to compute $\tau^{-1} d\mu_{\lambda_1, \lambda_2} \tau$ in order to understand the induced T^2 -action on $\mathbb{S}^3 \times T_1 S^1$. We see directly that

$$(3.2) \quad D\mu_{(\lambda_1, \lambda_2)} \tau(w, e_1) = \tau((\lambda_1, \lambda_2) \cdot w, e_1),$$

and for the other two cases let $\psi \in [0, 2\pi)$ such that $\lambda_1/\lambda_2 = \cos(\psi) + i \cdot \sin(\psi)$, then a straight forward calculation shows that

$$(3.3) \quad \begin{aligned} D\mu_{(\lambda_1, \lambda_2)} \tau(w, e_2) &= \cos(\psi) \tau((\lambda_1, \lambda_2) \cdot w, e_2) - \sin(\psi) \tau((\lambda_1, \lambda_2) \cdot w, e_3) \\ D\mu_{(\lambda_1, \lambda_2)} \tau(w, e_3) &= \sin(\psi) \tau((\lambda_1, \lambda_2) \cdot w, e_2) + \cos(\psi) \tau((\lambda_1, \lambda_2) \cdot w, e_3). \end{aligned}$$

Finally, observe that the coefficients in (3.3) only depend on (λ_1, λ_2) and not on w , so the induced action on $\mathbb{S}^3 \times T_1 S^1$ is a product action. Hence, $T_1 S^1$ is a T^2 -representation and $f: T_1 S^1 \rightarrow \mathbb{R} \oplus \Delta^* V_1$ defined by $f(e_1) = (1, 0)$, $f(e_2) = (0, 1)$ and $f(e_3) = (0, i)$ is an isomorphism of T^2 -representations. \square

Lemma 3.4. *The double cover group homomorphism $f: \mathbb{S}^3 \rightarrow \text{SO}(3)$ is S^1 -equivariant with respect to the S^1 -action on $\text{SO}(3)$ given by left multiplication with*

$$\begin{pmatrix} 1 & 0 \\ 0 & D(\lambda^2) \end{pmatrix} \in \text{SO}(3).$$

for $\lambda \in S^1$.

Proof. Identify \mathbb{R}^3 with the subspace of pure quaternions $\{xi + yj + zk \mid x, y, z \in \mathbb{R}\} \subset \mathbb{H}$ and $\mathbb{S}^3 \subset \mathbb{H}$ with the unit quaternions, i.e. $z = z_1 + z_2 j \in \mathbb{S}^3$ and $w = ix + z_3 j \in \mathbb{R}^3$ for $x \in \mathbb{R}$ and $z_1, z_2, z_3 \in \mathbb{C}$. We then define

$$f(z)(ix + jy + kz) := z \cdot w \cdot \bar{z},$$

which is a group homomorphism and preserves the inner product on \mathbb{R}^3 . Observe that $zj = j\bar{z}$ for $z \in \mathbb{C}$ and hence for any pure quaternion $w = ix + zj$ and $\lambda \in S^1 \subset \mathbb{C}$ we have that

$$\lambda w \bar{\lambda} = ix + \lambda z j \bar{\lambda} = ix + \lambda z \lambda j = ix + \lambda^2 z j,$$

which corresponds to the rotation by

$$(3.5) \quad \begin{pmatrix} 1 & 0 \\ 0 & D(\lambda^2) \end{pmatrix} \in \text{SO}(3).$$

Finally, since

$$f(\lambda z)(w) = \lambda z \cdot w \cdot \bar{\lambda} = \lambda f(z)(w) \bar{\lambda},$$

this corresponds to the left multiplication of $f(z)$ by (3.5). \square

Acknowledgements. I would like to thank Sam Nariman for introducing me to this aspect of foliation theory and posing many interesting questions in our conversations. I also thank Ronno Das for numerous helpful discussions and reading this note. This research was supported by the Knut and Alice Wallenberg foundation through grant no. 2019.0519.

REFERENCES

- [AP93] C. Allday and V. Puppe. *Cohomological methods in transformation groups*, volume 32 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1993.
- [Ber15] A. Berglund. Rational homotopy theory of mapping spaces via Lie theory for L_∞ -algebras. *Homology Homotopy Appl.*, 17(2):343–369, 2015.
- [Hae71] A. Haefliger. Homotopy and integrability. In *Manifolds–Amsterdam 1970 (Proc. Nuffic Summer School)*, volume Vol. 197 of *Lecture Notes in Math.*, pages 133–163. Springer, Berlin-New York, 1971.
- [Hae78] A. Haefliger. On the Gelfand-Fuks cohomology. *Enseign. Math. (2)*, 24(1-2):143–160, 1978.

- [Nar17] S. Nariman. Stable homology of surface diffeomorphism groups made discrete. *Geom. Topol.*, 21(5):3047–3092, 2017.
- [Nar23] S. Nariman. On invariants of foliated sphere bundles. *arXiv preprint arXiv:2308.16310*, 2023.
- [Pri] N. Prigge. A note on relative Gelfand-Fuks cohomology of spheres, to appear.
- [Thu74] W. Thurston. Foliations and groups of diffeomorphisms. *Bull. Amer. Math. Soc.*, 80:304–307, 1974.

STOCKHOLMS UNIVERSITET, MATEMATISKA INSTITUTIONEN, 106 91 STOCKHOLM
Email address: nils.prigge@math.su.se