Aggregating multiple test results to improve medical decision-making

Lucas Böttcher^{1,*}, Maria R. D'Orsogna^{2,3}, Tom Chou^{3,4}

1 Dept. of Computational Science and Philosophy, Frankfurt School of Finance and Management, Frankfurt am Main, Germany

2 Dept. of Mathematics, California State University at Northridge, Los Angeles, CA, 91330, USA

3 Dept. of Computational Medicine, University of California, Los Angeles, Los Angeles, CA, 90095-1766, USA

4 Dept. of Mathematics, University of California, Los Angeles, Los Angeles, CA, 90095-1555, USA

* l.boettcher@fs.de

Abstract

Gathering observational data for medical decision-making often involves uncertainties arising from both type I (false positive) and type II (false negative) errors. In this work, we develop a statistical model to study how medical decision-making can be improved by repeating diagnostic and screening tests, and aggregating their results. This approach is relevant not only in clinical settings, such as medical imaging, but also in public health, as highlighted by the need for rapid, cost-effective testing methods during the SARS-CoV-2 pandemic. Our model enables the development of testing protocols with an arbitrary number of tests, which can be customized to meet requirements for type I and type II errors. This allows us to adjust sensitivity and specificity according to application-specific needs. Additionally, we derive generalized Rogan–Gladen estimates for estimating disease prevalence, accounting for an arbitrary number of tests with potentially different type I and type II errors. We also provide the corresponding uncertainty quantification.

Author summary

Our work focuses on medical decision-making, particularly on addressing uncertainties associated with screening and diagnostic tests. No test is perfect, so finding a balance between false positives (misidentifying a condition) and false negatives (missing a condition) is crucial in many biomedical applications. Implementing accurate and efficient testing is important not only for individual diagnoses but also for population-wide testing during a pandemic. Since cost-effective and rapid tests are often quite inaccurate, a common goal is to obtain accurate assessments from repeated testing and meaningfully combining their results. However, using the multitude of tests and their different sequences of administration to design effective test protocols is a challenge that requires new statistical tools. In this study, we develop tools for aggregating test results in ways that can be tailored to specific applications by tuning the false positive-false negative ratio. Furthermore, we demonstrate how our method can improve disease prevalence estimates and thus aid in the implementation of effective public health measures. 4 Dept. of Mathematics, University of California, Los Angeles, Los Angeles, CA, 90065-1555, USA
 $^{\circ}$ 1 lost there of the control of the control

Introduction

Administering effective diagnostic and screening tests plays an important role in most biomedical decision-making. Recent advancements in biotechnology have made a wide array of biochemical tests readily available on a large scale. For example, in the case of SARS-CoV-2, a systematic review identified 49 different antigen tests [\[1\]](#page-19-0) which are cost-effective and can provide results in 15–30 minutes. However, their sensitivity (*i.e.*, true positive rate) can be as low as 34.3% in symptomatic patients and 28.6% in asymptomatic patients [\[1\]](#page-19-0). This indicates that some tests correctly identify an infected individual as positive in only about one third of cases, leaving a significant portion of those with the disease undetected. Besides sensitivity, another metric used to assess the accuracy of a test is its specificity (*i.e.*, true negative rate). Highly sensitive tests prioritize identifying individuals with a disease, while highly specific tests prioritize identifying those who do not have the disease. In most cases, sensitivity and specificity are inversely related; both are important when assessing the value of a medical test $[2,3]$.

Given the availability of various tests with differing sensitivities and specificities, how can one repeat tests and integrate results to minimize both type I errors (false positives) and type II errors (false negatives)? Although this is a key question across many different clinical settings, including diabetes testing [\[4,](#page-19-3) [5\]](#page-19-4), medical imaging [\[6–](#page-19-5)[8\]](#page-19-6), prostate cancer testing [\[9\]](#page-19-7), and stool sample analysis in colon cancer testing [\[10,](#page-19-8) [11\]](#page-19-9), our primary focus will be on aggregating results from different tests within the context of SARS-CoV-2 due to the availability of comprehensive studies on properties of the corresponding tests.

The SARS-CoV-2 pandemic emphasized the crucial role of testing in managing the spread of an infectious disease. During the early stages of the pandemic, test shortages were common, causing delays in diagnoses, underreporting of COVID-19 cases, and hindering the effectiveness of public health measures. Due to the intensifying crisis, regulatory agencies expedited the review and approval process of dedicated tests developed by different suppliers in different countries that used different technologies. These tests were often in use at the same time.^{[1](#page-1-0)} Early detection methods relied on genetic sequencing and reverse transcription polymerase chain reaction (RT-PCR) tests to detect viral genetic material. Antibody tests were also introduced to detect the presence of the virus in previously infected individuals who had developed an immune response. Both tests used samples collected from nasopharyngeal swabs and required specialized laboratory equipment and personnel to process them, making the diagnosis of active infections (RT-PCR tests) or of an activated immune response (antibody tests) available only after a few hours or even days. As the pandemic surged, the prioritization of rapid testing methods led to the development of rapid antigen tests, capable of detecting viral proteins and providing results within minutes. Subsequent saliva-based tests offered a less invasive experience compared to those based on nasopharyngeal swabs. Finally, the retreat of the pandemic was accompanied by the introduction of home testing kits. Current research is focused on perfecting new methods, including breathalyzer tests and wastewater monitoring.

Each testing method has its specific advantages and limitations. For example, RT-PCR tests are highly sensitive and specific and can detect even small amounts of viral RNA. However, there may be long delays in obtaining actionable results. Antibody tests may not detect antibodies in the early stages of the infection and are prone to large false-positive results due to cross-reactivity with antibodies from other viruses. Antigen tests are usually less sensitive than RT-PCR tests, but yield the most rapid response. Further variability in sensitivity and specificity arises within each type of testing method due to differences among test manufacturers and periodic modifications to the biochemical protocols, which are made to ensure the detection of any novel viral mutations or variants.

Our collective past experience with the spread of the SARS-CoV-2 virus poses several preparedness challenges to better respond to future pandemics, including how to best allocate scarce resources and enhance testing and classification strategies. The development of appropriate mathematical and computational methods plays a fundamental role in addressing these challenges. For example, one way to stretch resources is to test pooled samples, allowing one to eliminate large numbers of uninfected individuals with a small number of tests. Several mathematical approaches have been developed to study the optimization of both sample pooling and testing [\[13\]](#page-19-10). Different proposed strategies depend on test sensitivity and specificity [\[14\]](#page-19-11), estimated prevalence [\[15,](#page-19-12) [16\]](#page-19-13), disease dynamics [\[17\]](#page-20-0), and available social

¹One distinguishes between two primary categories of SARS-CoV-2 tests: (i) viral tests and (ii) antibody (or serological) tests [\[12\]](#page-19-14). Within the viral test category, there exist two main subclasses: nucleic acid amplification tests (NAATs), such as reverse transcription polymerase chain reaction (RT-PCR) tests that typically detect viral RNA, and antigen tests that detect specific antigen proteins on the surface of the virus. Antibody tests serve to identify antibodies produced as part of the adaptive immune system response. In the context of SARS-CoV-2, antibody tests may target anti-nucleocapsid antibodies, indicative of current or past infection, and anti-spike protein antibodies, generated through infection or vaccination.

contact information [\[18\]](#page-20-1). Other mathematical approaches aimed at improving testing efficiency account for uncertainty in disease prevalence [\[19\]](#page-20-2), indeterminate test results [\[20\]](#page-20-3), time-dependent prevalence and antibody levels [\[21,](#page-20-4) [22\]](#page-20-5), high-dimensional data analysis to improve classification accuracy [\[23\]](#page-20-6), and multiple classes such as vaccinated, previously infected, and unexposed individuals [\[24\]](#page-20-7).

In this paper, we focus on developing mathematical and computational methods that can help improve medical decision-making by repeating tests and aggregating their results. Several related studies have highlighted the potential of this approach [\[25](#page-20-8)[–33\]](#page-20-9), often using different terms such as "all heuristic" [\[30,](#page-20-10) [31\]](#page-20-11), "believe-the-negative rule" [\[34\]](#page-20-12), "conjunctive positivity criterion" [\[27,](#page-20-13) [35,](#page-21-0) [36\]](#page-21-1), and "orthogonal testing" [\[37\]](#page-21-2) to refer to the same protocol where all tests must return a positive result in order to classify an individual as infected. In Boolean algebra, this corresponds to an aggregation using the binary AND operator. Another aggregation method is the "any heuristic" [\[30,](#page-20-10) [31\]](#page-20-11) also termed the "believe-the-positive rule" [\[34\]](#page-20-12) or "disjunctive positivity criterion" [\[27,](#page-20-13) [35,](#page-21-0) [36\]](#page-21-1). In this protocol, all tests must return a negative result in order to classify an individual as not infected. It is thus sufficient for one test to be positive for a positive diagnosis. In Boolean algebra, this aggregation method is represented by the binary OR operator.

The US Food and Drug Administration (FDA) has also recognized the relevance of repeated testing and released an Excel-based calculator to compute properties of two combined tests [\[38\]](#page-21-3). However, most available result aggregation methods, including the FDA calculator, only consider two tests and usually employ very few (between one and three) basic aggregation methods. Nevertheless, there are instances where jurisdictions have implemented testing protocols involving three and four tests, such as in Vienna, Austria [\[39\]](#page-21-4), and Santiago, Chile [\[40\]](#page-21-5). Without appropriate mathematical insight and computational tools, however, it is challenging to analyze the properties of all possible aggregation methods due to the vast number of tests and their combinations. The lack of theoretical understanding often results in the implementation of ad-hoc and suboptimal aggregation protocols, rather than the most efficient ones. In addition to determining the disease status of an individual, combined tests can improve estimates of disease prevalence [\[26,](#page-20-14) [41\]](#page-21-6), which is helpful in infectious-disease surveillance and management [\[36,](#page-21-1) [42](#page-21-7)[–50\]](#page-21-8). In this context, it is also important to develop suitable mathematical tools to compare disease prevalence estimates across jurisdictions, as different public health organizations employ different testing protocols and aggregation methods [\[39,](#page-21-4) [40,](#page-21-5) [51–](#page-22-0)[53\]](#page-22-1).

Here, we combine concepts from biostatistics and Boolean algebra to develop a broadly applicable statistical model that can guide medical decision-making after repeated screening or diagnostic testing. We show how our model enables the development of testing protocols whose overall sensitivity and specificity can be tuned to satisfy application-specific requirements on type I and type II errors. Additionally, we present an algorithm capable of determining the best way to aggregate results from a given set of tests in terms of efficient sensitivity-specificity pairs. Furthermore, we integrate our aggregation approach with population-level prevalence estimation, demonstrating how repeated testing can enhance prevalence monitoring. Specifically, we generalize the Rogan–Gladen prevalence estimate [\[26,](#page-20-14) [41\]](#page-21-6) to account for an arbitrary number of tests, each with potentially different type I and type II errors.

Results

Aggregating two tests

As a starting point, we examine a testing protocol that combines the results of $n = 2$ tests (possibly of different types), denoted by binary random variables Y_1 and Y_2 , where $Y_1, Y_2 \in \{0, 1\}$. The disease status of an individual, classified as either positive $(+)$ or negative $(-)$, is represented by another binary random variable $X \in \{0, 1\}.$

The true positive rates TPRs (or sensitivities) of each of the two (type 1 and type 2) tests are defined as

$$
TPR_1 = Pr(Y_1 = 1 | X = 1) \text{ and } TPR_2 = Pr(Y_2 = 1 | X = 1), \tag{1}
$$

Both input signals are measured at the same time.

The signal on the left input is measured before the signal on the right input.

Fig 1. Parallel and series testing protocols using two tests. Positive $(+)$ and negative $(-)$ test outcomes are combined using the two Boolean functions AND (\Box) and OR (\Box). In parallel testing, both inputs are assessed simultaneously, while in series testing, the left input is examined before the right. Hence, if the initial test in a series protocol yields a negative result with aggregation through an AND gate, the final result will be negative, irrespective of the second input. In series testing with an OR gate, the overall result will be positive if the first test is positive, regardless of the outcome of the second test.

respectively. The corresponding true negative rates TNRs (or specificities) are

$$
TNR_1 = \Pr(Y_1 = 0 \mid X = 0) \text{ and } TNR_2 = \Pr(Y_2 = 0 \mid X = 0), \tag{2}
$$

respectively. These individual-test TPRs and TNRs serve as building blocks for modeling the overall TPR and TNR of a testing and aggregation protocol involving multiple tests.

For $n = 2$ ordered tests, there are $r = 2ⁿ = 2² = 4$ possible sequences of test results (*i.e.*, the inputs are permutations of "+" and "-" of length 2): $(+, +)$, $(+, -)$, $(-, +)$, and $(-, -)$. Given that the final, aggregated output can be either "+" or "−", the number of possible output sequences that one can assign to each of the $r = 4$ input configurations is $2^r = 16$, which is equivalent to the total number of two-input Boolean gates.

In Figure [1](#page-3-0) shows two (out of 16) possible output sequences associated with the aggregation of test results that involve $n = 2$ tests. In this example, aggregation can be processed through an AND gate or an OR gate. Depending on how the individual test results are processed, the output of the chosen aggregation function yields a positive or negative overall result. Notice that the aggregated output sequences of the parallel and series testing protocols shown are the same. The only implicit difference lies in how the input signals are processed (parallel or series). Both AND and OR aggregation functions have been used in SARS-CoV-2 seroprevalence studies (see Table [1\)](#page-9-0) and we will analyze both in this paper.

Aggregating test results using an AND gate produces a positive result if and only if both inputs are positive, corresponding to a "conjunctive positivity criterion" [\[27,](#page-20-13) [35,](#page-21-0) [36\]](#page-21-1). Otherwise the output result is negative. For an OR gate, the aggregate test is negative if and only if both inputs are negative, corresponding to a "disjunctive positivity criterion" [\[27,](#page-20-13) [35,](#page-21-0) [36\]](#page-21-1). In the remaining cases, the OR aggregation method outputs a positive result. The process of aggregation is sometimes referred to as the "all heuristic" [\[30,](#page-20-10) [31\]](#page-20-11) and the "believe-the-negative rule" [\[34\]](#page-20-12) when using AND aggregation. Similarly, OR aggregation is sometimes termed the "any heuristic' [\[30,](#page-20-10) [31\]](#page-20-11) and the "believe-the-positive rule" [\[34\]](#page-20-12).

For the two possible test-administration orderings (parallel and series) and the two aggregation procedures (AND and OR gates), we denote the corresponding cases as series AND, series OR, parallel AND, and parallel OR.

We use the random variable $Z \in \{0,1\}$ to denote the aggregated output and first examine the parallel testing protocol with an AND aggregation function. For $n = 2$ parallel tests, the sensitivity and specificity are

$$
\text{TPR}_{1\wedge 2}^{(p)} = \Pr(Z = 1 \mid X = 1) = \Pr(Y_1 = 1, Y_2 = 1 \mid X = 1) = \text{TPR}_1 \text{TPR}_2 \tag{3}
$$

and

$$
\begin{aligned}\n\text{TNR}_{1\wedge 2}^{(\text{p})} &= \Pr(Z = 0 \mid X = 0) \\
&= 1 - \Pr(Y_1 = 1, Y_2 = 1 \mid X = 0) \\
&= \Pr(Y_1 = 0, Y_2 = 0 \mid X = 0) + \Pr(Y_1 = 0, Y_2 = 1 \mid X = 0) + \Pr(Y_1 = 1, Y_2 = 0 \mid X = 0) \\
&= \text{TNR}_1 \text{TNR}_2 + \text{TNR}_1 (1 - \text{TNR}_2) + \text{TNR}_2 (1 - \text{TNR}_1) \\
&= \text{TNR}_1 + (1 - \text{TNR}_1) \text{TNR}_2 \,,\n\end{aligned} \tag{4}
$$

respectively. In Eqs. [\(3\)](#page-4-0) and [\(4\)](#page-4-1), we assumed that the results of different tests are conditionally independent given the disease status.

Conditioned on a disease state, this independence assumption is reasonable provided the tests do not interfere with each other, chemically or through biases in administration and interpretation, or perturb the patient. For example, the assays used in Elecsys® Anti-SARS-CoV-2 (Elecsys-N) and Elecsys® Anti-SARS-CoV-2 S (Elecsys-S) target antibodies against different proteins of the virus, nucleocapsid and spike proteins, respectively, so that test interactions can be neglected. Repeated tests that involve interpreting images can instead carry correlations due to changing operator bias. This phenomenon has been studied in [\[7\]](#page-19-15).

The derivations presented above are applicable to $n = 2$ parallel tests, where both the first and the second test results must be positive for classifying a sample as positive (an AND gate). If, however, the classification is based on the first or the second result being positive (an OR gate), the sensitivity and specificity are

$$
\begin{aligned} \text{TPR}_{1\vee 2}^{(p)} &= \Pr(Z = 1 \mid X = 1) \\ &= 1 - \Pr(Y_1 = 0, Y_2 = 0 \mid X = 1) \\ &= \Pr(Y_1 = 1, Y_2 = 1 \mid X = 1) + \Pr(Y_1 = 1, Y_2 = 0 \mid X = 1) + \Pr(Y_1 = 0, Y_2 = 1 \mid X = 1) \\ &= \text{TPR}_1 \text{TPR}_2 + \text{TPR}_1 (1 - \text{TPR}_2) + \text{TPR}_2 (1 - \text{TPR}_1) \\ &= \text{TPR}_1 + (1 - \text{TPR}_1) \text{TPR}_2 \end{aligned} \tag{5}
$$

and

$$
TNR_{1\vee 2}^{(p)} = Pr(Z = 0 \mid X = 0) = Pr(Y_1 = 0, Y_2 = 0 \mid X = 0) = TNR_1 TNR_2, \qquad (6)
$$

respectively. Given the assumptions in deriving the AND and OR aggregation protocols, we expect that the true positive rate is lower under AND aggregation (since all tests must be positive for a positive diagnosis) and vice-versa that the true negative rate is lower under OR aggregation (since all tests must be negative for a negative diagnosis). Based on Eqs. (3) – (6) , we obtain

$$
\text{TPR}_{1\vee 2}^{(p)} \ge \text{TPR}_{1\wedge 2}^{(p)} \quad \text{and} \quad \text{TNR}_{1\vee 2}^{(p)} \le \text{TNR}_{1\wedge 2}^{(p)} \tag{7}
$$

for all TPR_i and TNR_i $(i \in \{1, 2\})$.

Instead of administering two tests in parallel, one may also consider series testing in which the second test is administered depending on the outcome of the first test. In contrast to parallel testing with an AND aggregation, the second test in the corresponding sequential testing protocol does not have to be

Fig 2. The ratio of the number of parallel tests to the number of series tests necessary to determine the aggregated output from $n = 2$ tests as a function of prevalence f. Results in panels (a) and (b) are based on AND and OR aggregations of two tests, using Eqs. [\(11\)](#page-6-0) and [\(12\)](#page-6-1), respectively. We consider three different combinations of true positive and true negative rates (solid black lines: $TPR_1 = 0.95$ and $TNR_1 = 0.95$; dashed red lines: $TPR_1 = 0.90$ and $TNR_1 = 0.95$; dash-dotted blue lines: $TPR_1 = 0.95$ and $TNR_1 = 0.90$. The critical values f_c for which the ratios in panel (a) are larger than the ratios in panel (b) are given, respectively, by $f_c = 0.50, 0.47, 0.53$. For $f < f_c$ greater savings are achieved by utilizing the AND-aggregated series tests, compared to the OR-aggregated series test.

performed if the outcome of the first test is negative. The sensitivity and specificity of series testing under AND aggregation are

$$
\text{TPR}_{1\wedge 2}^{(\text{s})} = \text{TPR}_1 \text{TPR}_2 \quad \text{and} \quad \text{TNR}_{1\wedge 2}^{(\text{s})} = \text{TNR}_1 + (1 - \text{TNR}_1)\text{TNR}_2 \,, \tag{8}
$$

respectively. For the corresponding series OR test, we have

$$
TPR_{1\vee 2}^{(s)} = TPR_1 + (1 - TPR_1)TPR_2 \text{ and } TNR_{1\vee 2}^{(s)} = TNR_1TNR_2.
$$
 (9)

Notice that the sensitivities and specificities of the aggregated tests are the same regardless of whether a parallel or sequential aggregation protocol is employed. However, in a sequential protocol, fewer tests need to be administered, making this option more economically viable, especially for rapid antigen tests, characterized by lower sensitivity. For tests with extended processing times, such as enzyme-linked immunosorbent assay (ELISA) and RT-PCR tests, one may still prefer parallel test protocols to avoid substantial delays between the first and second tests.

Mathematically, the TPRs and TNRs of the studied combined testing protocols bound the TPRs and TNRs of the constituent tests according to

$$
\text{TPR}_{1\wedge 2} \le \text{TPR}_{i} \le \text{TPR}_{1\vee 2} \quad \text{and} \quad \text{TNR}_{1\vee 2} \le \text{TNR}_{i} \le \text{TNR}_{1\wedge 2} \quad \text{for} \quad i \in \{1, 2\}. \tag{10}
$$

We will show that this bounding result also holds for $n \geq 3$ tests.

Saving tests with series testing

To fully cover a population comprising N individuals using parallel testing would require $2N$ tests. In contrast, series testing involves administering an initial test to all individuals. In the series AND

aggregation function, a second test is required if and only if the first test yields a positive result. The probability of this event is $fTPR_1 + (1 - f)(1 - TNR_1)$, where $f \in [0, 1]$ is the prevalence, the fraction of the total population carrying a disease. In the series OR aggregation function, a second test is necessary if and only if the first test is negative, and the probability of this event is $f(1 - TPR_1) + (1 - f)TNR_1$. Both series testing protocols achieve the same sensitivity and specificity as the parallel test but with fewer tests, specifically, $N(1 + fTPR_1 + (1 - f)(1 - TNR_1))$ tests for the series AND function and $N(1 + f(1 - TPR_1) + (1 - f)TNR_1)$ tests for the series OR function, instead of 2N

when conducted in parallel.

We assume the number of tests available is such that there are enough of them to cover the entire population N. Since under parallel testing all individuals must be tested at least twice, this assumption implies that the population is at the most half the number of available tests. Given the disease prevalence f, we now compute the ratio of the number of required tests under parallel testing and the corresponding number of sequential tests. The ratios for the AND and OR aggregation methods are

$$
\frac{\text{parallel tests}}{\text{series tests}}\bigg|_{1\wedge 2} = \frac{2}{1 + f\text{TPR}_1 + (1 - f)(1 - \text{TNR}_1)}\tag{11}
$$

and

$$
\frac{\text{parallel tests}}{\text{series tests}}\bigg|_{1\vee 2} = \frac{2}{1 + f(1 - \text{TPR}_1) + (1 - f)\text{TNR}_1},\tag{12}
$$

respectively. Both ratios lie between 1 and 2, indicating that parallel testing always requires more tests than series testing. Besides the ground truth prevalence f, this ratio also depends on the disposition of the first test that determines if a second test is warranted. The first test result in turn, depends on its sensitivity TPR_1 and specificity TNR_1 . Figure [2](#page-5-0) shows these ratios as a function of prevalence f for three different combinations of true positive and true negative rates: (i) $TPR_1 = 0.95$ and $TNR_1 = 0.95$, (ii) $TPR_1 = 0.90$ and $TNR_1 = 0.95$, and (ii) $TPR_1 = 0.95$ and $TNR_1 = 0.90$. When there are no infected individuals in the population (i.e., $f = 0$), the parallel to series ratios are $2/(2 - TNR_1)$ and $2/(1 + \text{TNR}_1)$ for the AND and OR aggregation schemes, respectively. If all N individuals in a population are infected (*i.e.*, $f = 1$), the ratios are $2/(1 + \text{TPR}_1)$ and $2/(2 - \text{TPR}_1)$ for the AND and OR aggregation schemes, respectively.

It is also straightforward to show that for $f < f_c$, where f_c is the critical prevalence defined as

$$
f_{\rm c} = \frac{2TNR_1 - 1}{2(TPR_1 + TNR_1 - 1)},\tag{13}
$$

the number of required AND-aggregated series tests is less than the number of required OR-aggregated series tests. Equivalently, for $f < f_c$, the curve representing the ratio of the required parallel to series tests under the AND protocol given in Eq. [\(11\)](#page-6-0) falls above the corresponding OR protocol curve given in Eq. (12) . The trends observed in the parallel-to-series ratios as a function of prevalence f, shown in Figure [2\(](#page-5-0)a), confirm that AND aggregation yields greater test savings through series testing for prevalences $f < f_c$. In contrast, OR aggregation results in larger savings for $f > f_c$, as illustrated in Figure [2\(](#page-5-0)b).

According to Eq. [\(13\)](#page-6-2), the quantity f_c is meaningful only when TNR₁ \geq 1/2 and when $TPR₁ + TNR₁ > 1$. The latter condition implies that the true positive rate of the first test is greater than its false positive rate $(i.e., 1 - TNR₁)$. A test that satisfies this condition is said to have "discriminatory power" [\[36\]](#page-21-1). Typical values of TPR₁ and TNR₁ yield intermediate values of $f_c \approx 0.5$ as shown in Figure [2.](#page-5-0) Another scenario in which f_c is mathematically meaningful is when TNR₁ \leq 1/2 and $TPR_1 + TNR_1 < 1$. In this case, the trends in Figures $2(a,b)$ are reversed compared to the ones just discussed. This scenario, however, is highly unrealistic, as the first test is misleading since its false positive rate is greater that its true positive rate.

Positive predictive value

Measures such as sensitivity and specificity fail to appropriately take into account the prevalence of a disease f [\[54\]](#page-22-2). In this context, a more appropriate measure is the positive predictive value (PPV), also

Fig 3. Positive predictive value (PPV) and negative predictive value (NPV) as a function of prevalence f. The results that we show in panels (a,c) and (b,d) are based on AND and OR aggregations of $n = 2$ tests, using Eqs.[\(14\)](#page-7-0) and [\(15\)](#page-7-1), respectively. We denote the sensitivities and specificities of the two tests $i \in \{1,2\}$ by TPR_i and TNR_i, respectively. We consider two different combinations of true positive and true negative rates (solid black lines: TPR_i = 0.95 and TNR_i = 0.95; dashed red lines: TPR_i = 0.90 and $TNR_i = 0.90$. As a reference, we also show results for single tests without further aggregation (dash-dotted blue line: $TPR = 0.95$ and $TNR = 0.95$; dash-dot-dotted orange line: $TPR = 0.90$ and TNR = 0.90). These curves are independent of the ordering (parallel or series) method used.

known as precision, defined as

$$
PPV = \frac{fTPR}{fTPR + (1 - f)(1 - TNR)}.
$$
\n(14)

The PPV is the number of true positives divided by the number of positive calls. Similarly, we define the negative predictive (NPV) value as the number of true negatives divided by the number of negative calls,

$$
NPV = \frac{(1 - f)TNR}{(1 - f)TNR + f(1 - TPR)}.
$$
\n
$$
(15)
$$

Here, TPR and TNR represent the overall true positive and true negative rates of the aggregate testing protocol. By defining the utility gain associated with treating a sick individual and the utility loss associated with treating a healthy individual, it is possible to establish a relationship between PPV, NPV, and the treatment threshold. This threshold is the point where the expected treatment gain equals the expected treatment loss [\[36\]](#page-21-1).

Based on Eqs. (14) and (15) , one can show that the PPV is an increasing function of f and that the NPV is a decreasing function of f. These equations also yield $PPV \ge NPV$ when

$$
f \ge \frac{\sqrt{\text{TNR}(1 - \text{TNR})}}{\sqrt{\text{TNR}(1 - \text{TNR})} + \sqrt{\text{TPR}(1 - \text{TPR})}}.
$$
\n(16)

For multiple tests, the PPV and NPV are independent of the test ordering (parallel or series); however, they depend on the TPRs and TNRs of the individual tests in different ways depending whether the AND or the OR aggregation protocol is used. Specifically, we have

$$
PPV_{1\wedge2} \ge PPV_{1\vee2}, \qquad \forall f \tag{17}
$$

if

$$
\frac{\text{TPR}_1 + \text{TPR}_2}{\text{TPR}_1 \text{TPR}_2} \le \frac{(1 - \text{TNR}_1) + (1 - \text{TNR}_2)}{(1 - \text{TNR}_1)(1 - \text{TNR}_2)}.
$$
\n(18)

Similarly, we find

$$
NPV_{1\wedge 2} \le NPV_{1\vee 2}, \qquad \forall f \tag{19}
$$

if

$$
\frac{\text{TNR}_1 + \text{TNR}_2}{\text{TNR}_1 \text{TNR}_2} \le \frac{(1 - \text{TPR}_1) + (1 - \text{TPR}_2)}{(1 - \text{TPR}_1)(1 - \text{TPR}_2)}.
$$
\n(20)

The conditions in Eqs. [\(18\)](#page-8-0) and [\(20\)](#page-8-1) are always satisfied if tests with discriminatory power are used, *i.e.* if $TPR_1 + TNR_1 \geq 1$ and $TPR_2 + TNR_2 \geq 1$.

In Figure [3,](#page-7-2) we show the dependence of PPV and NPV on the prevalence f , using sensitivities and specificities associated with AND and OR aggregations for two tests [see Eqs. $(3)-(6)$ $(3)-(6)$ $(3)-(6)$]. We also include the corresponding PPV and NPV of individual (unaggregated) tests for reference. As can be seen, tests aggregated with the AND function yield substantially higher PPVs compared to those aggregated with an OR function for all f, while the OR aggregation results in notably higher NPVs compared to those obtained with the AND aggregation. Figure [3](#page-7-2) also shows that for low prevalence f the highest PPV and NPV values are obtained under AND aggregation, whereas OR aggregation is best for high prevalence f. For low f, the main source of test error is the false positive rate $1 - \text{TR}$. This term is minimized under the AND aggregation as per Eq. (10) . Similarly, for high f, the primary source of test error is the false negative rate $1 - TPR$, which is minimized under the OR aggregation as per in Eq. [\(10\)](#page-5-1).

So far, we have shown that when tests with discriminatory power are used for diseases with prevalence $f < f_c$, the AND aggregation protocol leads to the greatest reduction in the number of required tests when applied in series. Additionally, the AND aggregation protocol leads to larger PPV values compared to the OR protocol. Conversely, the potential savings under the OR aggregation protocol are smaller, and the NPV is larger than under the AND protocol. Thus, our analysis suggests that for $n = 2$ tests, the most suitable protocol for minimizing test usage and maximizing the PPV estimate in low-prevalence scenarios is the series AND method.

To provide further analytical insight into the properties of repeated tests, we consider aggregation functions involving more than two tests in the next section.

Aggregating more than two tests

In Table [1,](#page-9-0) we list examples of SARS-CoV-2 seroprevalence studies where up to four tests were administered using various combinations of parallel and series ordering with AND and OR

	parallel	series
AND	$n = 2$: Slovenia (nationwide) [51]	$n=2$: Norrbotten County, Sweden [52]
		$n=3$: Vienna, Austria [39]
ОR	$n = 2$: South Africa (three communities) [53]	
	$n = 4$: Santiago, Chile [40]	

Table 1. Examples of parallel and series test protocols that have been used in COVID-19 seroprevalence studies.

aggregation [\[40\]](#page-21-5). For $n = 3$ tests, there are $r = 2^3 = 8$ possible output sequences and $m = 2^r = 2^8 = 256$ possible input-output mappings. For $n = 4$, these numbers increase to $r = 2^4 = 16$ and $m = 2^r = 2^{16} = 65,536$ respectively. Given the large number of possible ways of combining n tests, we will derive sensitivities and specificities for a few select choices and otherwise resort to an algorithmic evaluation of test performances as detailed in the following section.

Equations [\(3\)](#page-4-0)–[\(8\)](#page-5-2) show that for $n = 2$, parallel and series test protocols carry the same sensitivities and specificities. This equivalence remains valid for $n \geq 3$ tests, so for notational simplicity we suppress the "s" and "p" superscripts that distinguish them.

For $n = 3$ tests and an AND aggregation, the sensitivity and specificity are

$$
TPR_{1\wedge2\wedge3} = TPR_1TPR_2TPR_3 \tag{21}
$$

and

$$
\begin{aligned} \text{TNR}_{1 \wedge 2 \wedge 3} &= \text{TNR}_1 + \text{TNR}_2 + \text{TNR}_3 - \text{TNR}_1 \text{TNR}_2 \\ &- \text{TNR}_1 \text{TNR}_3 - \text{TNR}_2 \text{TNR}_3 + \text{TNR}_1 \text{TNR}_2 \text{TNR}_3 \,, \end{aligned} \tag{22}
$$

respectively. Similarly, the sensitivity and specificity of an OR test protocol with $n = 3$ tests are

$$
\begin{aligned} \text{TPR}_{1\vee 2\vee 3} = \text{TPR}_1 + \text{TPR}_2 + \text{TPR}_3 - \text{TPR}_1 \text{TPR}_2 \\ &- \text{TPR}_1 \text{TPR}_3 - \text{TPR}_2 \text{TPR}_3 + \text{TPR}_1 \text{TPR}_2 \text{TPR}_3 \end{aligned} \tag{23}
$$

and

$$
TNR_{1\vee 2\vee 3} = TNR_1 TNR_2 TNR_3. \qquad (24)
$$

The overall sensitivity and specificity of the limiting AND and OR aggregations for general n-tests are

$$
\text{TPR}_{n}^{\text{AND}} = \prod_{i=1}^{n} \text{TPR}_{i}, \quad \text{TNR}_{n}^{\text{AND}} = 1 - \prod_{i=1}^{n} (1 - \text{TNR}_{i}), \tag{25}
$$

and

$$
\text{TPR}_{n}^{\text{OR}} = 1 - \prod_{i=1}^{n} (1 - \text{TPR}_{i}), \quad \text{TNR}_{n}^{\text{OR}} = \prod_{i=1}^{n} \text{TNR}_{i}. \tag{26}
$$

In line with Eq. [\(10\)](#page-5-1), the TPRs and TNRs of the combined testing protocols satisfy

$$
\text{TPR}_{n}^{\text{AND}} \leq \text{TPR}_{i} \leq \text{TPR}_{n}^{\text{OR}} \quad \text{and} \quad \text{TNR}_{n}^{\text{OR}} \leq \text{TNR}_{i} \leq \text{TNR}_{n}^{\text{AND}} \quad \forall i \in \{1, \dots, n\}. \tag{27}
$$

For odd $n \geq 3$, one can also employ a majority aggregation, where at least $(n + 1)/2$ tests have to be positive for the combined test to be positive. The majority aggregation is intermediate relative to the "all" and "any" characteristics of the AND and OR aggregations, respectively. The sensitivity of a majority aggregation of $n = 3$ tests is

$$
\text{TPR}_{\text{M}(1,2,3)} = \text{TPR}_1 \text{TPR}_2 + \text{TPR}_1 \text{TPR}_3 + \text{TPR}_2 \text{TPR}_3 - 2 \text{TPR}_1 \text{TPR}_2 \text{TPR}_3, \tag{28}
$$

and the corresponding specificity is

$$
TNR_{M(1,2,3)} = TNR_1TNR_2 + TNR_1TNR_3 + TNR_2TNR_3 - 2TNR_1TNR_2TNR_3.
$$
 (29)

Fig 4. Receiver operating characteristic (ROC) curves for various combinations of tests and aggregation functions. (a) We consider $n = 2$ tests and two distinct aggregation functions (disks: AND aggregation; triangles: OR aggregation). (b) We consider $n = 3$ tests and the same aggregation functions as in panel (a) along with the majority function represented by inverted triangles. Markers in black, blue, and red correspond to underlying tests $i \in \{1, \ldots, n\}$ with sensitivities (TPR_i) and specificities (TNR_i) set to 0.8, 0.9, and 0.95, respectively. Dashed lines indicate the sensitivities and false positive rates (*i.e.*, 1 − TNR) of the individual isolated tests. Under AND aggregation, both the sensitivities and false positive rates of the aggregated tests are smaller than those of the individual tests. The opposite holds for OR aggregation. When considering $n = 3$ tests, the majority function results in higher sensitivities and smaller false positive rates compared to the individual isolated tests. This function provides a tradeoff between the "all" and "any" characteristics of AND and OR aggregations. The results shown are independent of the ordering (parallel or series) method used.

In Figure [4,](#page-10-0) we show receiver operating characteristic (ROC) curves for various combinations of tests and aggregation functions. In Figure $4(a)$, we present the sensitivities and false positive rates for AND and OR aggregations with $n = 2$ tests. Additionally, in Figure [4\(](#page-10-0)b), we consider AND and OR, and majority aggregation for $n = 3$ tests. Under AND aggregation, the sensitivities and false positive rates of the aggregated tests are lower than those of the individual tests. The opposite holds for OR aggregation. These findings are in agreement with the analytical results in Eq. (27) . Finally, when examining $n = 3$ tests, the majority function yields greater sensitivities and reduced false positive rates compared to the individual isolated tests.

Efficiently combining n tests

For a given set of n tests, what are the best aggregation protocols in terms of sensitivities and specificities of the aggregated tests? As discussed in the prior sections, there exist numerous possibilities to combine individual tests, and the mathematical expressions for aggregated sensitivities and specificities can be quite lengthy. Therefore, we adopt an algorithmic approach to compute ROC curves associated with n tests, each potentially having distinct sensitivities and specificities. In this context, we use the term "efficient test" to denote a test whose underlying sensitivity-specificity pairs are optimized to achieve the highest possible values.

Aggregation and optimization

Algorithm 1 shown below computes the most efficient combination of n conditionally independent tests for given TPR_i and TNR_i of each test $i \in \{1, \ldots, n\}$. The following example illustrates our algorithm. We define $\mathcal{P} = \{P_1, \ldots, P_r\}$ as the set of possible ordered outcomes deriving from the administration of n tests, where $r = 2^n$. For example, for $n = 2$ tests, there are $r = 2^n = 2^2 = 4$ permutations and $P_1 = (+, +), P_2 = (+, -), P_3 = (-, +), \text{ and } P_4 = (-, -).$ Thus, we have

$$
\mathcal{P} = \{ (+, +), (+, -), (-, +), (-, -) \}.
$$
\n(30)

Each of these four outcomes can be mapped to either a positive aggregate diagnosis "+" or a negative one "−". Hence, there are $m = 2^r = 2^4 = 16$ mappings in total. For example, the output sequence $S = (+, -, -, -)$ means that only the input $P_1 = (+, +)$ is mapped to an aggregated "+", and the other permutations $P_2 = (+,-), P_3 = (-,+)$, and $P_4 = (-,-)$ are mapped to "−.". This case corresponds to the AND aggregation protocol. Similarly, $S = (+, +, +, -)$ corresponds to the OR aggregation protocol.

We define the sensitivity $TPR_{\mathcal{S}}$ associated with the output sequence $\mathcal{S} = (S_1, \ldots, S_r)$ in two steps. First, we define TPR_S as the sum over the sensitivities TPR_{S_j} $(j \in \{1, ..., r\})$ associated with elements S_j of S. That is,

$$
TPR_{\mathcal{S}} = \sum_{j=1}^{r} TPR_{S_j}.
$$
\n(31)

Second, we define TPR_{S_j} as follows. If element S_j is "−" (*i.e.*, if the input state P_j gets classified as negative), then we pose $TPR_{S_i} = 0$. Otherwise, if element S_j is "+", we calculate products of TPR_i and $1-\text{TPR}_i$ depending on whether the result from test $i \in \{1,\ldots,n\}$ is positive or negative. That is,

$$
\text{TPR}_{S_j} = \begin{cases} \prod_{i=1}^n \left[\text{TPR}_i \delta_{i,+} + (1 - \text{TPR}_i) \delta_{i,-} \right], & \text{if } S_j \text{ is } +\\ 0, & \text{if } S_j \text{ is } - \end{cases} \tag{32}
$$

where $\delta_{i,+} = 1$ if test i is positive and 0 otherwise. Likewise, $\delta_{i,-} = 1$ if test i is negative and 0 otherwise. For $n = 2$ tests and $S = (+, -, -, -)$, Eq. [\(32\)](#page-11-0) reduces to the AND aggregation TPR_{1∧2} result given in Eqs. [\(3\)](#page-4-0) and [\(8\)](#page-5-2). Similarly, for $n = 2$ tests and $S = (+, +, +, -)$, Eq. [\(32\)](#page-11-0) reduces to the OR aggregation TPR_{1∨2} result given in Eqs. [\(5\)](#page-4-3) and [\(9\)](#page-5-3). We follow the same steps to define the specificity TNR_S of the output sequence $\mathcal S$ so that

$$
TNR_{\mathcal{S}} = \sum_{j=1}^{r} TNR_{S_j}
$$
\n(33)

where

$$
\text{TNR}_{S_j} = \begin{cases} 0, & \text{if } S_j \text{ is } +\\ \prod_{i=1}^n \left[(1 - \text{TNR}_i)\delta_{i,+} + \text{TNR}_i \delta_{i,-} \right], & \text{if } S_j \text{ is } - \end{cases} \tag{34}
$$

For $n = 2$ tests and $S = (+, -, -, -)$, Eq. [\(34\)](#page-11-1) reduces to the AND aggregation TNR_{1∧2} result given in Eqs. [\(4\)](#page-4-1) and [\(8\)](#page-5-2). Similarly, for $n = 2$ tests and $S = (+, +, +, -)$, Eq. [\(34\)](#page-11-1) simplifies to the OR aggregation TNR_{1∨2} result given in Eqs. [\(6\)](#page-4-2) and [\(9\)](#page-5-3).

We identify two limit cases. One is the output sequence $S = (+, +, +, +)$ where all input permutations P_i are mapped to "+" outcomes. In this case, the aggregated sensitivity and specificity are TPR_S = 1 and TNR_S = 0, respectively. The other limit case is the output sequence $S = (-,-,-,-)$ where all input permutations P_i are mapped to "−" outcomes. Here, the aggregated sensitivity and specificity are TPR $_{\mathcal{S}} = 0$ and TNR $_{\mathcal{S}} = 1$, respectively.

Once we have determined all pairs (TPR_S, TNR_S) associated with the $m = 2^r$ test aggregations, we identify the most efficient test combinations, i.e., those combinations where the underlying sensitivity-specificity pairs reach the highest values. This is achieved by employing a convex-hull algorithm, such as Graham scan [\[55\]](#page-22-4) and Quickhull [\[56,](#page-22-5) [57\]](#page-22-6), to determine the ROC frontier in the $(TPR_{\mathcal{S}}, 1 - TNR_{\mathcal{S}})$ space (*i.e.*, true positive-false positive rate space). We summarize all steps of our algorithm in Python pseudocode in Algorithm [1.](#page-12-0)

Algorithm 1 Compute the most efficient combinations of n conditionally independent tests.

	$1:$ Inputs:		
	n , TPRs, TNRs, CONVEXHULL $()$		
	2: Outputs:		
	roc_frontier		
	3. input_permutations \leftarrow list(itertools.product([0, 1], repeat=n))	\triangleright Generate input permutations $\mathcal P$	
	4. input_output_mappings \leftarrow list(itertools.product([0, 1], repeat=2 ⁿ))	\triangleright Generate output sequences S	
	5: TPR_arr \leftarrow []		
	6: TNR_arr \leftarrow []		
	7. for input output map in input output mappings do		
8:	TPR -combined \leftarrow []		
9:	$TNR_{\text{-combined}} \leftarrow$		
10:	for perm, output_value in zip(input_permutations, input_output_map) do		
11:	if output_value then		
12:	TPR_combined.append $(\prod_{i=0}^{n-1}$ (TPRs[i] if perm[i] else 1-TPRs[i])) \triangleright see Eq. (32)		
13:	else		
14:	TNR_combined.append $(\prod_{i=0}^{n-1} (1-TNRs[i])$ if perm[i] else TNRs[i]))	\triangleright see Eq. (34)	
15:	end if		
16:	end for		
17:	$TPR_arr.append(sum(TPR_combined))$	\triangleright Compute aggregated sensitivity using Eq. (31)	
18:	$TNR_arr.append(sum(TNR_combined))$	\triangleright Compute aggregated specificity using Eq. (33)	
	19: end for		
	20: points \leftarrow concatenate(1-TNR_arr, TPR_arr)		
	21: convex_hull \leftarrow CONVEXHULL(points)		
	22: roc_frontier \leftarrow []		
	23: for edge in convex_hull do		
24:	if (points[edge, 1][0] \ge points[edge, 0][0]) and (points[edge, 1][1] \ge points[edge, 0][1]) then		
25:	roc_frontier.append(points[edge])	\triangleright ROC points must satisfy TPR ≥ 1 – TNR	
26:	end if		
	27: end for		
	28: return roc_frontier		

An example with three antigen tests

As an example, we apply Algorithm 1 to the aggregation of $n = 3$ commonly used SARS-CoV-2 antigen tests [\[1\]](#page-19-0). We list their median sensitivities and specificities along with their 95% confidence intervals (CIs) in Table [2.](#page-13-0) In this case there are $r = 2^n = 2^3 = 8$ permutations of test results and $m = 2^r = 256$ possible input-output mappings. The set of permutations is

$$
\mathcal{P} = \{(+,+,+),(+,+,-),(+,-,+),(+,-,-),(-,+,+),(-,-,-),(-,-,-),\}, \qquad (35)
$$

and the corresponding output sequence is $S = (S_1, \ldots, S_r)$, where $r = 8$ and $S_j \in \{+, -\}.$ To make the notation simpler, we introduce the Boolean variable $Y_i \in \{0,1\}$ for each test $i \in \{1,\ldots,n\}$ and map the aggregation method S to its corresponding Boolean expression as shown below.

We begin by using the median sensitivities and specificities of the three tests from Table [2](#page-13-0) as inputs in Algorithm [1](#page-12-0) for various aggregation protocols and use them to derive the corresponding ROC curve shown in Figure [5\(](#page-13-1)a). On this curve, there exist two limit cases: (i) an aggregation method where both sensitivity TPR_S and false positive rate $1 - \text{TNR}_{\mathcal{S}}$ are equal to 0, effectively classifying all input sequences as negative. This corresponds to $S_j = -$ for all $j \in \{1, \ldots, 8\}$; and (ii) an aggregation method with sensitivity $TPR_{\mathcal{S}}$ and false positive rate $1 - TNR_{\mathcal{S}}$ both at 1, resulting in the classification of all input sequences as positive, corresponding to $S_j = +$ for all $j \in \{1, \ldots, 8\}$.

We also include four more aggregation methods on the ROC curve shown in Figure [5\(](#page-13-1)a). The first of these requires that only the last test (i.e., Siemens) be positive, irrespective of the outcomes of the other two. This aggregation method corresponds to $S = (+,-, +,-, +,-, +,-)$ and is denoted Y_3 . It exhibits the smallest possible false positive rate, $1 - \text{TNR}_{\mathcal{S}} = 0$, which is intuitive given that the Siemens test also has the lowest median false positive rate of 0. Its sensitivity is $TPR_{\mathcal{S}} = 68.7\%$.

The subsequent aggregated test result shown on the ROC curve requires both the first and the second tests (i.e., Abbott and Innova), or the last one (i.e., Siemens) to yield positive results. This protocol corresponds to $S = (+, +, +, -, +, -)$ and can be written in Boolean algebra as $(Y_1 \wedge Y_2) \vee Y_3$.

	sensitivity	specificity
Abbott – Panbio COVID-19 Ag	74.8% $(67.6 - 80.8\%)$	99.7% (99.6 – 99.8%)
Innova Medical Group - Innova SARS-CoV-2 Ag	68.1% (47.2 – 83.6\%)	99.0% (98.5 – 99.3%)
Siemens - CLINITEST Rapid COVID-19 Ag	68.7% $(48.0 - 83.8\%)$	100% (98.0 – 100%)

Table 2. Median sensitivities and specificities of three commonly used SARS-CoV-2 antigen tests that are based on studies involving symptomatic patients [\[1\]](#page-19-0). Numbers in parentheses denote 95% CIs.

Using the values listed in Table [2](#page-13-0) and Eqs. [\(32\)](#page-11-0) and [\(34\)](#page-11-1), it can be verified that its sensitivity and false positive rate are TPR_S = 84.6% and $1 - \text{TNR}_S = 3.0 \times 10^{-3}\%$, respectively.

We can improve the aggregated sensitivity by omitting the second test $(i.e., Innova)$, which has the lowest sensitivity at 68.1%; the tradeoff is to accept a slightly higher false positive rate. This protocol yields the next point on the ROC curve. It corresponds to $S = \{+, +, +, +, +, -, +, -\}$ and can be written using an OR aggregation over the first and last tests, *i.e.*, $Y_1 \vee Y_3$. Equations [\(32\)](#page-11-0) and [\(34\)](#page-11-1) yield the sensitivity TPR $_{\mathcal{S}} = 92.1\%$ and the false positive rate $1 - \text{TNR}_{\mathcal{S}} = 0.3\%$.

Finally, the largest sensitivity smaller than 100% is achieved through an OR aggregation over all tests *i.e.*, for $Y_1 \vee Y_2 \vee Y_3$. This corresponds to $S = \{+, +, +, +, +, +, -\}$, an output sequence with sensitivity TPR_S = 97.5% and false positive rate $1 - \text{TNR}_{\mathcal{S}} = 1.3\%$ as per Eqs. [\(32\)](#page-11-0) and [\(34\)](#page-11-1).

For a more detailed comparison between aggregated and individual tests, we show a magnified view of the four non-trivial aggregations in the ROC frontier in Figure [5\(](#page-13-1)b) and include individual tests. In this plot, we incorporate CIs alongside median sensitivities and false positive rates. We generate these CIs from 10^5 samples of beta distributions capturing the 95% CIs of the underlying individual sensitivities and specificities (see Materials and methods for further details). We observe that the two OR protocols, $Y_1 \vee Y_3$ and $Y_1 \vee Y_2 \vee Y_3$, exhibit significantly higher sensitivity compared to each individual test.

Fig 5. ROC curves associated with the aggregation of three antigen tests (Abbot, Innova, and Siemens). The sensitivities and specificities of the $n = 3$ tests are listed in Table [2.](#page-13-0) (a) The ROC curve associated with the aggregation of the three antigen tests as derived from Eqs. [\(32\)](#page-11-0) and [\(34\)](#page-11-1). We use $Y_i \in \{0, 1\}$ to denote the outcome of test $i \in \{1,2,3\}$. (b) A magnified view of the ROC curve without the trivial aggregated tests that classify all samples as either negative or positive. The error bars indicate the 95% CIs that we generated from 10^6 samples of beta distributions capturing the 95% CIs of the underlying individual sensitivities and specificities.

Estimating prevalence

In the preceding sections, we have observed how repeating and aggregating results from diagnostic and screening tests can substantially enhance sensitivity and specificity. This enhancement can contribute to improved infectious-disease surveillance and management $[42-45]$ by providing more accurate estimates \hat{f} of the true prevalence f in a population. The prevalences \hat{f} and f may be time-dependent and stratified, e.g., according to age, $(i.e., \hat{f} \equiv \hat{f}(a_k, t)$ and $\hat{f} \equiv f(a_k, t)$ where a_k is a given age).

Correcting test errors

We distinguish between the measured prevalence $\hat{f}_{\mathcal{S}}^{*}(a_k, t)$ of a disease derived from testing a given sample of the population and using the aggregation method S , and the measured, error-corrected prevalence $f(a_k, t)$, which is instead the resulting estimate of the true disease prevalence $f(a_k, t)$. If we also assume that the selected sample is unbiased and representative of the infection behavior in the entire population, we can identify the estimate $f(a_k, t)$ with the actual prevalence $f(a_k, t)$ and write $\hat{f}(a_k,t) = f(a_k,t)$. For n aggregated tests with output sequence S, the quantities $\hat{f}^*_\mathcal{S}(a_k,t)$ and $\hat{f}(a_k,t)$ are related via

$$
\hat{f}_{\mathcal{S}}^*(a_k, t) = \hat{f}(a_k, t)\text{TPR}_{\mathcal{S}} + (1 - \hat{f}(a_k, t))(1 - \text{TNR}_{\mathcal{S}}),\tag{36}
$$

which yields

$$
\hat{f}(a_k, t) = \frac{\hat{f}_\mathcal{S}^*(a_k, t) + \text{TNR}_\mathcal{S} - 1}{\text{TPR}_\mathcal{S} + \text{TNR}_\mathcal{S} - 1},\tag{37}
$$

a generalized Rogan–Gladen prevalence estimate [\[41\]](#page-21-6) that accounts for the sensitivity and specificity of the aggregated tests with output sequence S. We omit the subscript S in $f(a_k, t)$ since the error-corrected prevalence is an estimate of the true prevalence and should not depend of the method used for aggregating test results. For example, for $n = 2$ tests under AND and OR aggregation and using Eqs. $(3)-(8)$ $(3)-(8)$, we have

$$
\hat{f}(a_k, t) = \frac{\hat{f}_{1 \wedge 2}^*(a_k, t) + \text{TNR}_1 + \text{TNR}_2 - \text{TNR}_1 \text{TNR}_2 - 1}{\text{TPR}_1 \text{TPR}_2 + \text{TNR}_1 + \text{TNR}_2 - \text{TNR}_1 \text{TNR}_2 - 1},
$$
\n(38)

and

$$
\hat{f}(a_k, t) = \frac{\hat{f}_{1\vee 2}^*(a_k, t) + \text{TNR}_1 \text{TNR}_2 - 1}{\text{TPR}_1 + \text{TPR}_2 - \text{TPR}_1 \text{TPR}_2 + \text{TNR}_1 \text{TNR}_2 - 1},
$$
\n(39)

respectively.

In Figure [6,](#page-15-0) we show the measured (uncorrected) prevalences $\hat{f}_{1\wedge 2}^*$ and $\hat{f}_{1\vee 2}^*$ associated with the AND and OR aggregations using Eq. [\(36\)](#page-14-0) and Eqs. [\(3\)](#page-4-0)–[\(8\)](#page-5-2) for the corresponding TPR_S and TNR_S, and using different sensitivities and specificities for $n = 2$ tests. For simplicity, we assume that samples are unbiased and that the measured, error-corrected prevalence \hat{f} can be identified with the true prevalence f.

In line with our findings regarding PPV and NPV and the trends shown in Figure [3,](#page-7-2) we observe in Figure [6](#page-15-0) that $\hat{f}_{1\wedge 2}^*$, the measured prevalence under AND aggregation, deviates only slightly from the true prevalence f, for low true prevalence, whereas under OR aggregation deviations between $\hat{f}_{1\vee2}^*$ and f are small at large true prevalence.

As a real-world example of prevalence correction under aggregated testing, we consider the seroprevalence study from Norrbotten Sweden (May $25 -$ June 5, 2020) [\[52\]](#page-22-3). In this study, two SARS-CoV-2 tests, Abbott SARS-CoV-2 IgG and the Euroimmun Anti-SARS-CoV-2 ELISA (IgG), were administered to an age-stratified population and combined using an AND function. The non age-stratified measured prevalence is estimated at $\hat{f}_{1\wedge 2}^* = 1.9\%$; other details of the testing protocols employed in this study are discussed in the Materials and methods. We use the $\hat{f}^*_{1\wedge 2}$ estimate to calculate the measured, error-corrected prevalence using Eq. [\(38\)](#page-14-1) and the underlying individual test sensitivities and specificities given in the Materials and methods section. We also calculate the corresponding 95% CIs by generating 10^6 samples from beta distributions capturing the measured

Fig 6. Measured prevalence \hat{f}^* as a function of true prevalence f under the assumption that the measured, error-corrected prevalence \hat{f} in Eq. [\(36\)](#page-14-0) can be identified with the true prevalence f. The results that we show in panels (a) and (b) are based on AND and OR aggregations of two tests $i \in \{1, 2\}$, respectively. We consider three different combinations of true positive and true negative rates (solid black lines: TPR_i = 0.95 and TNR_i = 0.95; dashed red lines: TPR_i = 0.90 and TNR_i = 0.95; dash-dotted blue lines: $TPR_i = 0.95$ and $TNR_i = 0.90$). Grey lines indicate measured prevalences associated with individual tests.

prevalence $\hat{f}_{1\wedge 2}^*$. We present the age-stratified measured prevalences $\hat{f}_{1\wedge 2}^*$ and the corresponding measured, error-corrected prevalences \hat{f} for various age groups in Table [3.](#page-16-0)

Equation [\(38\)](#page-14-1) yields a non-stratified measured, error-corrected prevalence $\hat{f} = 2.5\%$ (1.1 – 5.0%), which is higher than the measured prevalence $\hat{f}^*_{1\wedge 2} = 1.9\%$ (0.8 - 3.7%). Because the sensitivity of tests combined using an AND function is lower compared to the sensitivity of the underlying constituent tests, the measured prevalence associated with this aggregation function usually underestimates the true prevalence. Hence, the measured, error-corrected prevalence is substantially larger in this example than the measured one.

An application in fatality and hospitalization monitoring

Prevalence estimates commonly arise in infection fatality and hospitalization ratios, which are useful measures for monitoring outbreak severity. For a given jurisdiction at time t , the infection fatality ratio IFR(a_k , t) of the population of age in the interval $[a_k, a_{k+1})$ is

$$
IFR(a_k, t) = \frac{D(a_k, t)}{f(a_k, t)N(a_k)},
$$
\n
$$
(40)
$$

where $f(a_k, t)$ and $D(a_k, t)$ respectively denote the age-stratified true proportion of infected individuals at time t and the total number of infection-caused fatalities up to time t measured from the start of an outbreak and within the age interval $[a_k, a_{k+1})$. In the above definition, we assume that the overall population $N(a_k)$ of age in the interval $[a_k, a_{k+1})$ is constant in the time horizon of interest. The denominator $f(a_k, t)N(a_k)$ in Eq. [\(40\)](#page-15-1) quantifies the total number of age-stratified infections at time t since the start of an outbreak *(i.e., current and prior infections)*.

The number of infection-caused fatalities, $D(a_k, t)$, may be difficult to infer because of various confounding factors. These factors include variations in protocols for attributing the cause of death, the

age group		measured prevalence $\hat{f}_{1\wedge2}^*$ error-corrected prevalence \hat{f}
$20-29$ years	6.6% $(1.8 - 15.9\%)$	8.8% $(2.4 - 21.6\%)$
$30-64$ years	0.7% $(0.1 - 2.7\%)$	0.9% $(0.1 - 3.3\%)$
$65 - 80$ years	2.1% $(0.3 - 7.3\%)$	2.8% $(0.4 - 9.5\%)$

Table 3. Measured and error-corrected prevalence in Norrbotten, Sweden (May 25 – June 5, 2020) [\[52\]](#page-22-3). The error correction method we employed takes into account the two tests used in the seroprevalence study from Norrbotten: (i) the Abbott SARS-CoV-2 IgG kit and (ii) the Euroimmun Anti-SARS-CoV-2 ELISA (IgG). These tests have been aggregated using an AND function. We calculated the measured, error-corrected prevalence through Eq. (38) and their corresponding 95% CIs by generating 10^6 samples from beta distributions capturing the measured prevalence as well as the underlying individual test sensitivities and specificities. Details of the study are listed in Materials and methods.

existence of co-morbidities [\[58\]](#page-22-7), and delays in reporting. In jurisdictions where underreporting is prevalent, statistics on excess deaths may offer a more accurate assessment of the overall death toll [\[42,](#page-21-7) [45\]](#page-21-9).

Analogous to the IFR, the infection hospitalization ratio $\text{IHR}(a_k, t)$ of the population of age in the interval $[a_k, a_{k+1})$ in a given jurisdiction is

$$
IHR(a_k, t) = \frac{H(a_k, t)}{f(a_k, t)N(a_k)},
$$
\n(41)

where $H(a_k, t)$ is the corresponding total number of age-stratified infection-caused hospitalizations up to time t measured from the start of an outbreak. Because of the time lag between infection and resolution, both the IFR and IHR may underestimate the true burden of an outbreak, especially in the early stages when the number of new cases increases rapidly [\[59\]](#page-22-8). In Table [4,](#page-17-0) we summarize the main variables used in outbreak severity measures.

The true proportion of infections $f(a_k, t)$ used in the denominators of both IFR and IHR is usually difficult to quantify for large populations. We can thus employ prevalence estimates $f(a_k, t)$ as derived in Eq. [\(37\)](#page-14-2) that are usually based on serological testing of random samples of the entire population. Estimated proportions of infections $\hat{f}(a_k, t)$ that have been obtained using serological tests can be assumed to be close to the true proportions $f(a_k, t)$ if antibody waning is negligible and if the population sample is unbiased and representative of the whole population.

We denote the corresponding IFR and IHR estimates by

$$
\widehat{\text{IFR}}(a_k, t) = \frac{D(a_k, t)}{\hat{f}(a_k, t)N(a_k)},
$$
\n(42)

and

$$
\widehat{\text{IHR}}(a_k, t) = \frac{H(a_k, t)}{\hat{f}(a_k, t)N(a_k)},
$$
\n(43)

respectively.

The seroprevalence study from Norrbotten, Sweden (May 25 – June 5, 2020) [\[52\]](#page-22-3) yields an overall measured, error-corrected seroprevalence of $\hat{f} = 2.5\%$ (1.1 – 5.0%). We assume that this prevalence estimate, obtained for a subpopulation aged 20 to 80 years, is reflective of the prevalence in the entire population of 249,614 individuals. Using the total number of 59 fatalities and 242 hospitalizations documented throughout the entire study duration, along with Eqs. [\(42\)](#page-16-1) and [\(43\)](#page-16-2), we obtain IFR $= 0.9\%$ $(0.5 - 2.2\%)$ and IHR = 3.8% (1.9 – 9.1%). These values are lower than the fatality ratio of 1.2% (0.6 – 3.0%) and hospitalization ratio of 5.1% (2.6 – 12.1%) obtained with the uncorrected, measured prevalence $\hat{f}_{1\wedge2}^* = 1.9\%$ (0.8 - 3.7%).

Symbol	Definition	
$N(a_k) \in \mathbb{N}$	population of age in the interval $[a_k, a_{k+1})$ in a given jurisdiction	
$D(a_k, t) \in \mathbb{N}$	total number of infection-caused fatalities of age in the interval $[a_k, a_{k+1})$ in a given jurisdiction at time t (measured from the start of an outbreak)	
$H(a_k, t) \in \mathbb{N}$	total number of infection-caused hospitalizations of age in the interval $[a_k, a_{k+1})$ in a given jurisdiction at time t (measured from the start of an outbreak)	
$f(a_k, t) : [0, 1]$	true proportion of infected individuals of age in the interval $[a_k, a_{k+1})$ at time t in a given jurisdiction	

Table 4. Main variables used in outbreak severity measures. Population, fatality, hospitalization, and prevalence statistics are often reported for N_a age intervals $[a_{k-1}, a_k)$ $(k \in \{1, ..., N_a\})$ with $a_k = a_0 + \sum_{\ell=1}^k \Delta a_\ell$. Here, a_0 is the smallest age value in the data set and Δa_ℓ is the width of the ℓ -th age window. We assume that the population size $N(a_k)$ is constant in the considered time window. The closed interval $[0, 1]$ contains $[0, 1]$, and all numbers in between, and $\mathbb N$ denotes the set of non-negative integers.

Discussion

Repeating and aggregating results from diagnostic and screening tests can significantly enhance overall test performance. While our primary focus has been on aggregating tests within the context of infectious-disease surveillance, similar concepts hold broad clinical applicability, such as in diabetes testing [\[4,](#page-19-3) [5\]](#page-19-4), medical imaging [\[6](#page-19-5)[–8\]](#page-19-6), and cancer screening [\[10,](#page-19-8) [11\]](#page-19-9). The complex clinical conditions are usually probed by tests performing multiclass discrimination, requiring generalizations of the ROC surface and other reduction schemes [\[60\]](#page-22-9).

Starting from the aggregation of two tests, for which there are 16 fundamental two-input Boolean gates, we derived expressions for the sensitivity and specificity of aggregated tests, assuming their conditional independence. We quantified the potential for saving tests when employing series testing compared to parallel testing, without compromising sensitivity and specificity. Additionally, we discussed the strong dependence of the positive predictive value (PPV) (*i.e.*, the ratio of true positives to positive calls) and negative predictive value (NPV) $(i.e.,$ the ratio of true negatives to negative calls) on the employed aggregation mechanism. For example, AND aggregation yields good relative large PPVs and NPVs at low prevalence values, while OR aggregation does so for larger prevalences.

Expressions of sensitivity and specificity for aggregations of more than two tests can also be derived. However, these expressions may become very lengthy. Thus, we developed an algorithm capable of identifying the best way of aggregating results from a given set of tests in terms of efficient sensitivity-specificity pairs (i.e., sensitivity-specificity values that lie on an ROC frontier). We applied this algorithm to three commonly used SARS-CoV-2 tests and demonstrated how their individual sensitivities and specificities can be significantly improved when combined.

Finally, in this work we established a connection between aggregating tests and prevalence estimates in infectious-disease surveillance. Such estimates are pertinent for computing measures like the infection fatality ratio (IFR) and infection hospitalization ratio (IHR).

Although our work addresses various factors related to aggregating tests, there are additional aspects that we have not considered. For instance, certain tests may entail higher costs or varying levels of complications for patients (see, e.g., chapter 7.4 in [\[36\]](#page-21-1)). Other refinements may incorporate test-avoidance, or increasing levels of test-fatigue when multiple tests are to be administered. Incorporating these effects requires formulating appropriate target functions and adjusting our optimization approach. In the context of an ROC curve, a target function that quantifies the utility gain associated with treating a sick individual and the utility loss associated with treating a healthy individual enables the identification of the optimum aggregation approach [\[61\]](#page-22-10). Moreover, although we

have incorporated sensitivity and specificity data for numerous tests in our analysis, it would be beneficial to validate the results of our model through experimental data on aggregated tests.

In addition to the described applications, our work can help inspire aggregation methods in social choice theory and decision-making under uncertainty, where the objective is to effectively combine individual opinions [\[62](#page-22-11)[–66\]](#page-22-12). For instance, it can inform decision-making processes in organizations where decision makers also possess sensitivities and specificities with respect to a given decision task. Furthermore, our paper is closely connected to historical works on fault-tolerant computing by von Neumann [\[67\]](#page-23-1), Moore, and Shannon [\[68–](#page-23-2)[70\]](#page-23-3), who studied how reliable (Boolean) computing elements can be constructed from unreliable components.

Materials and methods

Beta distribution sampler

To calculate CIs associated with aggregated tests and related quantities that depend on multiple factors such as sensitivity, specificity, and prevalence, we employ a Monte Carlo sampling technique. In this work, we consider samples drawn from a beta distribution

$$
\mathbb{P}(x;\alpha,\beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)} x^{\alpha-1} (1-x)^{\beta-1},\tag{44}
$$

where $x \in [0, 1], \alpha, \beta$ are shape parameters, and $\Gamma(\cdot)$ denotes the gamma function. Sensitivities, specificities, and prevalences are quantities with a support of [0, 1], so beta distributions are plausible approximations of their underlying distributions.

We determine shape parameters such that the corresponding distributions capture the median and 95% CIs of the underlying quantities. To do so, we minimize the sum of squared differences between the cumulative distribution at the 2.5%, 50%, and 97.5% quantiles, and the corresponding empirical median and 95% CI values. We carry out this optimization process by employing the fmin function implemented in scipy.optimize in Python. Further implementation details are available at [\[71\]](#page-23-4).

Hospitalization, fatality, and serology data

In the main text, we use data from a seroprevalence study conducted in Norrbotten, Sweden, during weeks 22 and 23 of 2020 (May 25 to June 5) [\[52\]](#page-22-3). We considered the hospitalization, fatality, and seroprevalence data provided in this study to illustrate how errors associated with aggregated tests can be addressed. The study encompassed a population of 182,828 adults aged 20 to 80 years. The age distribution within this population was as follows: 16.2% were aged 20 to 29 years, 57.8% were aged 30 to 64 years, and 25.9% were aged 65 to 80 years. From this population, 500 individuals were randomly selected and contacted, out of which 425 participated in the study. A total of 242 individuals with confirmed infection had been hospitalized since the beginning of the outbreak, and 59 people with confirmed infection had passed away.

The study revealed a population-wide measured prevalence $\hat{f}_{1\wedge 2}^*$ of 1.9% (0.8 – 3.7%). Seroprevalence was assessed using two different assays: (i) Abbott SARS-CoV-2 IgG and (ii) Euroimmun Anti-SARS-CoV-2 ELISA (IgG). The former has a sensitivity and specificity of 83.1% (75.4 – 100%) and 100%, respectively [\[72\]](#page-23-5). The sensitivity and specificity of the latter are 91.1% (80.7 – 96.1%) and 100% $(96.5 - 100\%)$, respectively [\[73\]](#page-23-6).

Every individual who tested positive in Abbott's assay underwent confirmation using Euroimmun's Anti-SARS-CoV-2 ELISA (IgG). This process represents an AND aggregation.

Acknowledgments

The authors thank Stefan Felder and Nana Owusu-Boaitey for helpful comments. We acknowledge financial support from hessian.AI (LB), the ARO through grant W911NF-23-1-0129 (MRD and LB), and

References

- 1. Dinnes J, Sharma P, Berhane S, van Wyk S, Nyaaba N, Domen J, et al. Rapid, point-of-care antigen tests for diagnosis of SARS-CoV-2 infection. Cochrane Database of Systematic Reviews. 2022;7(7).
- 2. Glaros AG, Kline RB. Understanding the accuracy of tests with cutting scores: The sensitivity, specificity, and predictive value model. Journal of Clinical Psychology. 1988;44(6):1013–1023.
- 3. Akobeng AK. Understanding diagnostic tests 1: sensitivity, specificity and predictive values. Acta Paediatrica. 2007;96(3):338–341.
- 4. Brohall G, Behre CJ, Hulthe J, Wikstrand J, Fagerberg B. Prevalence of diabetes and impaired glucose tolerance in 64-year-old Swedish women: experiences of using repeated oral glucose tolerance tests. Diabetes Care. 2006;29(2):363–367.
- 5. Kermani SK, Khatony A, Jalali R, Rezaei M, Abdi A. Accuracy and precision of measured blood sugar values by three glucometers compared to the standard technique. Journal of Clinical and Diagnostic Research. 2017;11(4):OC05.
- 6. Weinstein S, Obuchowski NA, Lieber ML. Clinical evaluation of diagnostic tests. American Journal of Roentgenology. 2005;184(1):14–19.
- 7. Zou KH, Bhagwat JG, Carrino JA. Statistical combination schemes of repeated diagnostic test data. Academic Radiology. 2006;13(5):566–572.
- 8. Brennan PC, Ganesan A, Eckstein MP, Ekpo EU, Tapia K, Mello-Thoms C, et al. Benefits of independent double reading in digital mammography: a theoretical evaluation of all possible pairing methodologies. Academic Radiology. 2019;26(6):717–723.
- 9. Vickers AJ, Cronin AM, Gönen M. A simple decision analytic solution to the comparison of two binary diagnostic tests. Statistics in Medicine. 2013;32:1865–1876.
- 10. Neuhauser D, Lewicki AM. What do we gain from the sixth stool guaiac? New England Journal of Medicine. 1975;293(5):226–228.
- 11. Collins JF, Lieberman DA, Durbin TE, Weiss DG, Group* VACS. Accuracy of screening for fecal occult blood on a single stool sample obtained by digital rectal examination: a comparison with recommended sampling practice. Annals of Internal Medicine. 2005;142(2):81–85.
- 12. Testing for SARS-CoV-2 Infection; Accessed: March 13, 2024. Available from: [https:](https://www.cdc.gov/coronavirus/2019-ncov/hcp/testing-overview.html#TestingInfection) [//www.cdc.gov/coronavirus/2019-ncov/hcp/testing-overview.html#TestingInfection](https://www.cdc.gov/coronavirus/2019-ncov/hcp/testing-overview.html#TestingInfection).
- 13. Daniel EA, Esakialraj L BH, S A, Muthuramalingam K, Karunaianantham R, Karunakaran LP, et al. Pooled Testing Strategies for SARS-CoV-2 diagnosis: A comprehensive review. Diagnostic Microbiology and Infectious Disease. 2021;101(2):115432.
- 14. Wunsch M, Aschemeier D, Heger E, Ehrentraut D, Krüger J, Hufbauer M, et al. Safe and effective pool testing for SARS-CoV-2 detection. Journal of Clinical Virology. 2021;145:105018.
- 15. Mutesa L, Ndishimye P, Butera Y, Souopgui J, Uwineza A, Rutayisire R, et al. A pooled testing strategy for identifying SARS-CoV-2 at low prevalence. Nature. 2020;589:276–280.
- 16. Schenk H, Caf Y, Knabl L, Mayerhofer C, Rauch W. High prevalence group testing in epidemiology with geometrically inspired algorithms. Scientific Reports. 2023;13:18910.
- 17. Cleary B, Hay JA, Blumenstiel B, Harden M, Cipicchio M, Bezney J, et al. Using viral load and epidemic dynamics to optimize pooled testing in resource-constrained settings. Science Translational Medicine. 2021;13:eabf1568.
- 18. Zhang J, Heath LS. Adaptive group testing strategy for infectious diseases using social contact graph partitions. Scientific Reports. 2023;13:12102.
- 19. Patrone PN, Kearsley AJ. Classification under uncertainty: data analysis for diagnostic antibody testing. Mathematical Medicine and Biology: A Journal of the IMA. 2021;38(3):396–416.
- 20. Patrone PN, Bedekar P, Pisanic N, Manabe YC, Thomas DL, Heaney CD, et al. Optimal decision theory for diagnostic testing: Minimizing indeterminate classes with applications to saliva-based SARS-CoV-2 antibody assays. Mathematical Biosciences. 2022;351:108858.
- 21. Bedekar P, Kearsley AJ, Patrone PN. Prevalence estimation and optimal classification methods to account for time dependence in antibody levels. Journal of Theoretical Biology. 2023;559:111375.
- 22. Bedekar P, Luke RA, Kearsley AJ. Prevalence estimation methods for time-dependent antibody kinetics of infected and vaccinated individuals: a graph-theoretic approach; 2024.
- 23. Luke RA, Kearsley AJ, Pisanic N, Manabe YC, Thomas DL, Heaney CD, et al. Modeling in higher dimensions to improve diagnostic testing accuracy: theory and examples for multiplex saliva-based SARS-CoV-2 antibody assays. PLOS ONE. 2023;18(3):e0280823.
- 24. Luke RA, Kearsley AJ, Patrone PN. Optimal classification and generalized prevalence estimates for diagnostic settings with more than two classes. Mathematical Biosciences. 2023;358:108982.
- 25. Cebul RD, Hershey JC, et al. Using multiple tests: series and parallel approaches. Clinics in Laboratory Medicine. 1982;2(4):871–890.
- 26. McClish D, Quade D. Improving estimates of prevalence by repeated testing. Biometrics. 1985; p. 81–89.
- 27. Hershey JC, Cebul RD, Williams SV. Clinical guidelines for using two dichotomous tests. Medical Decision Making. 1986;6(2):68–78.
- 28. Marshall RJ. The predictive value of simple rules for combining two diagnostic tests. Biometrics. 1989; p. 1213–1222.
- 29. Ramdas K, Darzi A, Jain S. 'Test, re-test, re-test': using inaccurate tests to greatly increase the accuracy of COVID-19 testing. Nature Medicine. 2020;26(6):810–811.
- 30. Jain S, Jónasson JO, Pauphilet J, Flower B, Moshe M, Fontana G, et al. A new combination testing methodology to identify accurate and economical point-of-care testing strategies. medRxiv. 2021;.
- 31. Jain S, Jónasson JO, Pauphilet J, Ramdas K. Robust combination testing: methods and application to COVID-19 detection. Management Science. 2023;.
- 32. Salvatore PP, Shah MM, Ford L, Delaney A, Hsu CH, Tate JE, et al. Quantitative comparison of SARS-CoV-2 nucleic acid amplification test and antigen testing algorithms: a decision analysis simulation model. BMC Public Health. 2022;22(1):1–10.
- 33. Perkmann T, Koller T, Perkmann-Nagele N, Ozsvar-Kozma M, Eyre D, Matthews P, et al. Increasing test specificity without impairing sensitivity: lessons learned from SARS-CoV-2 serology. Journal of Clinical Pathology. 2023;76(11):770–777.
- 34. Politser P. Reliability, decision rules, and the value of repeated tests. Medical Decision Making. 1982;2(1):47–69.
- 35. Ament A, Hasman A. Optimal test strategy in the case of two tests and one disease. International Journal of Biomedical Computing. 1993;33(3–4):179–197.
- 36. Felder S, Mayrhofer T. Medical Decision Making: A Health Economic Primer, 3rd Ed. Berlin, Germany: Springer-Verlag; 2022.
- 37. Lau CS, Aw TC. Disease prevalence matters: Challenge for SARS-CoV-2 testing. Antibodies. $2021;10(4):50.$
- 38. Food and Drug Administration calculator for positive predictive value (PPV) and negative predictive value (NPV) for individual tests and combined; Accessed: March 14, 2024. Available from: <https://www.fda.gov/media/137612/download>.
- 39. Breyer MK, Breyer-Kohansal R, Hartl S, Kundi M, Weseslindtner L, Stiasny K, et al. Low SARS-CoV-2 seroprevalence in the Austrian capital after an early governmental lockdown. Scientific Reports. 2021;11(1):10158.
- 40. Vial PA, Gonz´alez C, Apablaza M, Vial C, Lav´ın ME, Araos R, et al. First wave of SARS-CoV-2 in Santiago Chile: Seroprevalence, asymptomatic infection and infection fatality rate. Epidemics. 2022;40:100606.
- 41. Rogan WJ, Gladen B. Estimating prevalence from the results of a screening test. American Journal of Epidemiology. 1978;107(1):71–76.
- 42. Böttcher L, D'Orsogna MR, Chou T. Using excess deaths and testing statistics to determine COVID-19 mortalities. European Journal of Epidemiology. 2021;36(5):545–558.
- 43. Zhang Q. Data science approaches to infectious disease surveillance. Philosophical Transactions of the Royal Society A. 2022;380(2214):20210115.
- 44. Schneider T, Dunbar OR, Wu J, Böttcher L, Burov D, Garbuno-Inigo A, et al. Epidemic management and control through risk-dependent individual contact interventions. PLOS Computational Biology. 2022;18(6):e1010171.
- 45. Böttcher L, D'Orsogna MR, Chou T. A statistical model of COVID-19 testing in populations: effects of sampling bias and testing errors. Philosophical Transactions of the Royal Society A. 2022;380(2214):20210121.
- 46. Meyer MJ, Yan S, Schlageter S, Kraemer JD, Rosenberg ES, Stoto MA. Adjusting COVID-19 seroprevalence survey results to account for test sensitivity and specificity. American Journal of Epidemiology. 2022;191(4):681–688.
- 47. Zhou L, Díaz-Pachón DA, Zhao C, Rao JS, Hössjer O. Correcting prevalence estimation for biased sampling with testing errors. Statistics in Medicine. 2023;42(26):4713–4737.
- 48. Owusu-Boaitey N, Russell TW, Meyerowitz-Katz G, Levin AT, Herrera-Esposito D. Dynamics of SARS-CoV-2 seroassay sensitivity: a systematic review and modelling study. Eurosurveillance. 2023;28(21):2200809.
- 49. Levin AT, Owusu-Boaitey N, Pugh S, Fosdick BK, Zwi AB, Malani A, et al. Assessing the burden of COVID-19 in developing countries: systematic review, meta-analysis and public policy implications. BMJ Global Health. 2022;7(5):e008477.
- 50. Liu K, Bai Z, He D, Lou Y. Getting Jab or Regular Test: Observations from an Impulsive Epidemic COVID-19 Model. Bulletin of Mathematical Biology. 2023;85(10):97.
- 51. Poljak M, Valenčak AO, Štrumbelj E, Vodičar PM, Vehovar V, Rus KR, et al. Seroprevalence of severe acute respiratory syndrome coronavirus 2 in Slovenia: results of two rounds of a nationwide population study on a probability-based sample, challenges and lessons learned. Clinical Microbiology and Infection. 2021;27(7):1039–e1.
- 52. The seroprevalence of COVID-19 antibodies in Norrbotten County, Sweden; Accessed: December 14, 2023. Available from: [https://web.archive.org/web/20200623071108/http://norrbotten.se/sv/](https://web.archive.org/web/20200623071108/http://norrbotten.se/sv/Halsa-och-sjukvard/Smittskydd-i-Norrbotten/Information-om-nya-coronaviruset/The-seroprevalence-of-covid-19-antibodies-in-Norrbotten-County-Sweden/) [Halsa-och-sjukvard/Smittskydd-i-Norrbotten/Information-om-nya-coronaviruset/](https://web.archive.org/web/20200623071108/http://norrbotten.se/sv/Halsa-och-sjukvard/Smittskydd-i-Norrbotten/Information-om-nya-coronaviruset/The-seroprevalence-of-covid-19-antibodies-in-Norrbotten-County-Sweden/) [The-seroprevalence-of-covid-19-antibodies-in-Norrbotten-County-Sweden/](https://web.archive.org/web/20200623071108/http://norrbotten.se/sv/Halsa-och-sjukvard/Smittskydd-i-Norrbotten/Information-om-nya-coronaviruset/The-seroprevalence-of-covid-19-antibodies-in-Norrbotten-County-Sweden/).
- 53. Wolter N, Tempia S, von Gottberg A, Bhiman JN, Walaza S, Kleynhans J, et al. Seroprevalence of Severe Acute Respiratory Syndrome Coronavirus 2 After the Second Wave in South Africa in Human Immunodeficiency Virus–Infected and Uninfected Persons: A Cross-Sectional Household Survey. Clinical Infectious Diseases. 2022;75(1):e57–e68.
- 54. Eisenberg MJ. Accuracy and predictive values in clinical decision-making. Cleveland Clinic Journal of Medicine. 1995;62(5):311–316.
- 55. Graham RL. An Efficient algorithm for determining the convex hull of a finite planar set. Information Processing Letters. 1972;1(4):132–133. doi:10.1016/0020-0190(72)90045-2.
- 56. Greenfield JS. A Proof for a QuickHull Algorithm. Electrical Engineering and Computer Science; 1990.
- 57. Barber CB, Dobkin DP, Huhdanpaa H. The quickhull algorithm for convex hulls. ACM Transactions on Mathematical Software. 1996;22(4):469–483.
- 58. Centers for Disease Control and Prevention. Conditions contributing to COVID-19 deaths, by state and age, provisional 2020-2023; Accessed: March 15, 2024.
- 59. Böttcher L, Xia M, Chou T. Why case fatality ratios can be misleading: individual-and population-based mortality estimates and factors influencing them. Physical Biology. 2020;17(6):065003.
- 60. Allwein EL, Schapire RE, Singer Y. Reducing multiclass to binary: A unifying approach for margin classifiers. Journal of Machine Learning Research. 2000;1:113–141.
- 61. Felder S, Werblow A, Robra BP. A priori risk and optimal test accuracy in prenatal diagnostics. Medical Decision Making. 2003;23(5):406–413.
- 62. Smoke WH, Zajonc RB. Reliable groups of unreliable people. Ann Arbor, Michigan: Office of Naval Research, Research Center for Group Dynamics, The Institute for Social Research, The University of Michigan; 1960. 7.
- 63. Brandt F, Conitzer V, Endriss U, Lang J, Procaccia AD. Handbook of computational social choice. Cambridge, UK: Cambridge University Press; 2016.
- 64. Böttcher L, Kernell G. Examining the limits of the Condorcet Jury Theorem: Tradeoffs in hierarchical information aggregation systems. Collective Intelligence. 2022;1(2):26339137221133401.
- 65. Kurvers RH, Nuzzolese AG, Russo A, Barabucci G, Herzog SM, Trianni V. Automating hybrid collective intelligence in open-ended medical diagnostics. Proceedings of the National Academy of Sciences. 2023;120(34):e2221473120.
- 66. Böttcher L, Klingebiel R. Collective screening of innovation. Organization Science. 2004;
- 67. von Neumann J. Probabilistic logics and the synthesis of reliable organisms from unreliable components. Automata Studies. 1956;34(34):43–98.
- 68. Moore EF, Shannon CE. Reliable circuits using less reliable relays, part I. Journal of the Franklin Institute. 1956;262:191–208.
- 69. Moore EF, Shannon CE. Reliable circuits using less reliable relays, part II. Journal of the Franklin Institute. 1956;262:281–297.
- 70. Kochen M. Extension of Moore–Shannon model for relay circuits. IBM Journal of Research and Development. 1959;3(2):169–186.
- 71. Böttcher L. GitLab repository; 2024. Available from: <https://gitlab.com/ComputationalScience/testing-statistics>.
- 72. Castro MDM, Caicedo I, Ortiz-Rojas HJ, Castillo CM, Medina AG, Alexander N, et al. Performance verification of the Abbott SARS-CoV-2 test for qualitative detection of IgG in Cali, Colombia. PLOS ONE. 2021;16(9):e0256566.
- 73. Anti-SARS-CoV-2 S1 Curve ELISA (IgG) Instructions for Use; Accessed: March 9, 2024. https://www.fda.gov/media/152747/download.