

LIPSCHITZ TRUNCATION METHOD FOR PARABOLIC DOUBLE-PHASE SYSTEMS AND APPLICATIONS

WONTAE KIM, JUHA KINNUNEN, AND LAURI SÄRKIÖ

ABSTRACT. We discuss a Lipschitz truncation technique for parabolic double-phase problems of p -Laplace type in order to prove energy estimates and uniqueness results for the Dirichlet problem. Moreover, we show existence for a non-homogeneous double-phase problem. The Lipschitz truncation method is based on a Whitney-type covering result and a related partition of unity in the intrinsic geometry for the double-phase problem.

1. INTRODUCTION

This paper discusses weak solutions $u = u(z) = u(x, t)$ to parabolic double-phase systems of type

$$u_t - \operatorname{div}(|\nabla u|^{p-2} \nabla u + a(z)|\nabla u|^{q-2} \nabla u) = - \operatorname{div}(|F|^{p-2} F + a(z)|F|^{q-2} F)$$

in $\Omega_T = \Omega \times (0, T)$, where Ω is a bounded domain in \mathbb{R}^n and $T > 0$. Here $2 \leq p < q < \infty$ and the coefficient $a(\cdot)$ is non-negative and Hölder continuous. It is natural to assume that the gradient of a weak solution and the source term satisfy

$$\iint_{\Omega_T} (|\nabla u|^p + a(z)|\nabla u|^q) dz + \iint_{\Omega_T} (|F|^p + a(z)|F|^q) dz < \infty.$$

Observe that the coefficient function $a(\cdot)$ may vanish in a subset of the domain and thus the condition above does not imply that $|\nabla u| \in L^q(\Omega_T)$. One of the challenges is that the structure of the double-phase systems is not dominated by the p -growth or the q -growth. Indeed, the intrinsic geometry depends on the location and it switches between two phases.

Energy estimates are of fundamental importance in existence and regularity results for partial differential equations. There is a delicate issue which does not occur in the elliptic case: A weak solution to the parabolic double-phase system under the natural function space assumption above is not an admissible test function after mollification in time, see Remark 2.3. A similar problem arises in the uniqueness of the Dirichlet problem since the difference of two solutions cannot be applied as a test function. Proposition 4.11 and Proposition 6.2 contain two versions of Lipschitz truncation for parabolic double-phase problems. These are applied to prove the energy estimate in Theorem 2.4 and the uniqueness of solutions in Theorem 2.6. As a corollary we obtain the higher integrability result for the gradient in [17] without the additional assumption $|\nabla u| \in L^q(\Omega_T)$, see Remark 2.5. The existence of solutions to a non-homogeneous Dirichlet problem (2.6) is discussed in Theorem 2.7.

The main idea of the Lipschitz truncation is to construct an admissible test function by keeping it unchanged in the set where the strong maximal function of the gradient is small and redefining it in the complement by applying a partition of unity related to a Whitney covering. The Whitney covering in Proposition 3.2 is constructed in an intrinsic geometry for the parabolic double-phase problem and it involves the p -intrinsic and (p, q) -intrinsic phases as in [17]. A careful analysis is needed to combine the phases. Since the intrinsic geometry depends on the location, it is challenging to obtain uniform bounds in a Vitali covering argument. The partition of unity in Proposition 3.2 is subordinate to the Whitney covering and reflects the double-phase structure. Maximal function estimates imply the convergence of the Lipschitz truncation to the original test

2020 *Mathematics Subject Classification.* 35K55, 35K65, 35D30, 35A01, 35A02.

Key words and phrases. Parabolic double-phase systems, existence theory, Lipschitz truncation method.

function. The existence proof is based on an approximation with regularized problems, Calderón-Zygmund estimates in [16], compactness arguments and the Minty-Browder approach.

The Lipschitz truncation was introduced for the elliptic setting in [1, 2]. The corresponding technique for parabolic p -Laplace systems was developed in [19] and discussed further in [3, 5, 6, 14]. It has also been used to prove existence result for parabolic problems in [7, 13]. The regularity theory of elliptic double-phase problems has been studied in [4, 9, 10, 11, 15]. Very little is known about the parabolic double-phase problems. Local higher integrability of the gradient has been studied in [17], the existence of weak solutions to parabolic double-phase systems was proved in the homogeneous case in [21] and in the scalar case with non-divergence data in [8]. Our existence result seems to be new in the vectorial case with non-homogeneous data. We are confident that our results can be applied to solve further regularity problems for the parabolic double-phase systems.

2. NOTATION AND MAIN RESULTS

2.1. Notation. Let Ω be a bounded open subset in \mathbb{R}^n and $\Omega_T = \Omega \times (0, T)$, $T > 0$. A point in \mathbb{R}^{n+1} is denoted as $z = (x, t)$ or $w = (y, s)$, where $x, y \in \mathbb{R}^n$ and $t, s \in \mathbb{R}$. A ball with center $x_0 \in \mathbb{R}^n$ and radius $\rho > 0$ is denoted as

$$B_\rho(x_0) = \{x \in \mathbb{R}^n : |x - x_0| < \rho\}$$

and a cube centered at the origin is denoted as

$$D_\rho = \{(x_1, \dots, x_n) \in \mathbb{R}^n : |x_i| < \rho, i \in \{1, \dots, n\}\}.$$

Let $t_0 \in \mathbb{R}$ and $\lambda > 0$. Parabolic cylinders with quadratic scaling in time are denoted as

$$Q_\rho(z_0) = B_\rho(x_0) \times I_\rho(t_0) \quad \text{and} \quad C_\rho = D_\rho \times I_\rho(0),$$

where

$$I_\rho(t_0) = (t_0 - \rho^2, t_0 + \rho^2).$$

We use the following notation for the double-phase functional. For the non-negative coefficient function $a(\cdot)$ extended to \mathbb{R}^{n+1} , see (3.2), we define a function $H(z, s) : \mathbb{R}^{n+1} \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$ as

$$H(z, s) = s^p + a(z)s^q.$$

For $\lambda \geq 1$, a p -intrinsic cylinder centered at $z_0 = (x_0, t_0)$ is

$$Q_\rho^\lambda(z_0) = B_\rho(x_0) \times I_\rho^\lambda(t_0), \quad I_\rho^\lambda(t_0) = (t_0 - \lambda^{2-p}\rho^2, t_0 + \lambda^{2-p}\rho^2) \quad (2.1)$$

and a (p, q) -intrinsic cylinder centered at $z_0 = (x_0, t_0)$ is

$$G_\rho^\lambda(z_0) = B_\rho(x_0) \times J_\rho^\lambda(t_0), \quad J_\rho^\lambda(t_0) = \left(t_0 - \frac{\lambda^2}{H(z_0, \lambda)}\rho^2, t_0 + \frac{\lambda^2}{H(z_0, \lambda)}\rho^2 \right).$$

The dependence from z_0 is omitted in the notation $J_\rho^\lambda(t_0)$ since λ will be chosen so that the denominator $H(z_0, \lambda)$ remains fixed regardless of z_0 whenever it appears. Here, λ^{2-p} and $\frac{\lambda^2}{H(z_0, \lambda)}$ are called the scaling factors in time. Note that if $\lambda = 1$, then $Q_\rho(z_0) = Q_\rho^\lambda(z_0)$. Moreover, we write

$$cQ_\rho^\lambda(z_0) = Q_{c\rho}^\lambda(z_0) \quad \text{and} \quad cG_\rho^\lambda(z_0) = G_{c\rho}^\lambda(z_0)$$

for $c > 0$. We also consider parabolic cylinders with arbitrary scaling in time and denote

$$Q_{r,s}(z_0) = B_r(x_0) \times \ell_s(t_0), \quad \ell_s(t_0) = (t_0 - s, t_0 + s), \quad r, s > 0. \quad (2.2)$$

The center point, height and radius may be omitted if irrelevant. In this case Q means an arbitrary cylinder of the above type.

The $(n+1)$ -dimensional Lebesgue measure of a set $E \subset \mathbb{R}^{n+1}$ is denoted as $|E|$. For $f \in L^1(\Omega_T, \mathbb{R}^N)$ and a measurable set $E \subset \Omega_T$ with $0 < |E| < \infty$, we denote the integral average of f over E as

$$(f)_E = \frac{1}{|E|} \iint_E f \, dz = \fint_E f \, dz.$$

2.2. Main results. Let $2 \leq p < q < \infty$. In this paper we consider energy estimates of weak solutions to parabolic problems of double-phase type,

$$u_t - \operatorname{div} \mathcal{A}(z, \nabla u) = -\operatorname{div}(|F|^{p-2}F + a(z)|F|^{q-2}F) \quad \text{in } \Omega_T, \quad (2.3)$$

where $\mathcal{A}(z, \xi) : \Omega_T \times \mathbb{R}^{Nn} \rightarrow \mathbb{R}^{Nn}$ with $N \geq 1$ is a Carathéodory vector field satisfying the structure assumptions

$$\mathcal{A}(z, \xi) \cdot \xi \geq \nu(|\xi|^p + a(z)|\xi|^q) \quad \text{and} \quad |\mathcal{A}(z, \xi)| \leq L(|\xi|^{p-1} + a(z)|\xi|^{q-1}) \quad (2.4)$$

for a.e. $z \in \Omega_T$ and every $\xi \in \mathbb{R}^{Nn}$ with constants $0 < \nu \leq L < \infty$. The source term $F : \Omega_T \rightarrow \mathbb{R}^{Nn}$ is a vector field satisfying

$$\iint_{\Omega_T} H(z, |F|) dz < \infty.$$

Throughout this paper we assume that the coefficient function $a : \Omega_T \rightarrow \mathbb{R}^+$ satisfies

$$q \leq p + \frac{2\alpha}{n+2}, \quad 0 \leq a \in C^{\alpha, \alpha/2}(\Omega_T) \quad \text{for some } \alpha \in (0, 1].$$

Here $a \in C^{\alpha, \alpha/2}(\Omega_T)$ means that $a \in L^\infty(\Omega_T)$ and there exists a constant $[a]_{\alpha, \alpha/2; \Omega_T} = [a]_\alpha > 0$, such that

$$|a(x, t) - a(y, t)| \leq [a]_{\alpha, \alpha/2; \Omega_T} |x - y|^\alpha \quad \text{and} \quad |a(x, t) - a(x, s)| \leq [a]_{\alpha, \alpha/2; \Omega_T} |t - s|^{\frac{\alpha}{2}} \quad (2.5)$$

for every $x, y \in \Omega$ and $t, s \in (0, T)$. It is not clear whether the range for q is optimal. However, considering the corresponding elliptic theory this condition with the parabolic scaling deficit $\frac{p}{2}$ seems natural.

Definition 2.1. A function $u \in C(0, T; L^2(\Omega, \mathbb{R}^N)) \cap L^1(0, T; W^{1,1}(\Omega, \mathbb{R}^N))$ satisfying

$$\iint_{\Omega_T} (H(z, |u|) + H(z, |\nabla u|)) dz < \infty$$

is a weak solution to (2.3) if

$$\iint_{\Omega_T} (-u \cdot \varphi_t + \mathcal{A}(z, \nabla u) \cdot \nabla \varphi) dz = \iint_{\Omega_T} (|F|^{p-2}F \cdot \nabla \varphi + a(z)|F|^{q-2}F \cdot \nabla \varphi) dz$$

for every $\varphi \in C_0^\infty(\Omega_T, \mathbb{R}^N)$.

In general, the time derivative of a weak solution exists only in the distributional sense. Steklov averages are applied to mollify the function in the time direction. Let $f \in L^1(\Omega_T)$ and $0 < h < T$. We define the Steklov average $[f]_h(x, t)$ for all $0 < t < T$ as

$$[f]_h(x, t) = \begin{cases} \int_t^{t+h} f(x, s) ds, & 0 < t < T - h, \\ 0, & T - h \leq t. \end{cases}$$

We refer to [12] for standard properties of the Steklov average. In particular we note that for any $f \in L^2(\Omega_T)$, we have $[f]_h \in W^{1,2}((0, T-h); L^2(\Omega))$ with

$$\partial_t [f]_h = \frac{f(x, t+h) - f(x, t)}{h}$$

for $(x, t) \in \Omega_{T-h}$. The proof of the following lemma can be found for example in [5, Lemma 8.1].

Lemma 2.2. Let $f \in L^1(\Omega_T)$ and $h > 0$. Then there exists a constant $c = c(n)$ such that

$$\iint_{Q_{r,s}(z_0)} [f]_h dz \leq c \iint_{[Q_{r,s}(z_0)]_h} f dz,$$

where $[Q_{r,s}(z_0)]_h = Q_{r,s+h}(z_0)$.

Remark 2.3. *The Steklov averages are applied in order to have*

$$\begin{aligned} & \int_{\Omega \times \{t\}} (\partial_t [u]_h \cdot \varphi + [|\nabla u|^{p-2} \nabla u + a |\nabla u|^{q-2} \nabla u]_h \cdot \nabla \varphi) dx \\ &= \int_{\Omega \times \{t\}} [|F|^{p-2} F + a |F|^{q-2} F]_h \cdot \nabla \varphi dx \end{aligned}$$

for every $\varphi \in C_0^\infty(\Omega, \mathbb{R}^N)$ and for a.e. $0 < t < T - h$. To obtain an energy estimate, we would like to apply the solution itself as a test function in the definition of weak solutions. This is possible if the right-hand side of

$$\begin{aligned} & \int_{\Omega \times \{t\}} |[|\nabla u|^{p-2} \nabla u + a |\nabla u|^{q-2} \nabla u]_h | |\nabla \varphi| dx \\ & \leq \int_{\Omega \times \{t\}} (|[|\nabla u|^{p-2} \nabla u]_h | |\nabla \varphi| + |[a |\nabla u|^{q-2} \nabla u]_h | |\nabla \varphi|) dx \end{aligned}$$

is finite for test functions in the function space of the weak solution. The first term in the integral is finite from Hölder's inequality and the property of the Steklov average as in the p -Laplace system case ($a(\cdot) \equiv 0$). However, if we write open the second term in the integral,

$$\int_{\Omega \times \{t\}} \left| \int_t^{t+h} a(x, \tau) |\nabla u(x, \tau)|^{q-2} \nabla u(x, \tau) d\tau \right| |\nabla \varphi(x)| dx,$$

and let $\varphi = [u]_h \zeta$ for any cutoff function ζ in Ω_T , it becomes apparent that this is difficult to bound since $[u]_h(\cdot, t)$ does not have the same time variable as $a(\cdot)$. The same problem occurs also for the source term. The Lipschitz truncation technique is applied to overcome this problem.

Our first main result is an energy estimate for weak solutions as in Definition 2.1.

Theorem 2.4. *Assume that u is a weak solution of (2.3). Then there exists a constant $c = c(p, q, \nu, L)$ such that*

$$\begin{aligned} & \sup_{t \in \ell_{S_1}(t_0)} \int_{B_{R_1}(x_0)} \frac{|u - (u)_{Q_{R_1, S_1}(z_0)}|^2}{S_1} dx + \iint_{Q_{R_1, S_1}(z_0)} H(z, |\nabla u|) dz \\ & \leq c \iint_{Q_{R_2, S_2}(z_0)} \left(H\left(z, \frac{|u - (u)_{Q_{R_2, S_2}(z_0)}|}{R_2 - R_1}\right) + \frac{|u - (u)_{Q_{R_2, S_2}(z_0)}|^2}{S_2 - S_1} + H(z, |F|) \right) dz \end{aligned}$$

for every $Q_{R_2, S_2}(z_0) \subset \Omega_T$ with $R_2, S_2 > 0$, $R_1 \in [R_2/2, R_2)$ and $S_1 \in [S_2/2^2, S_2)$.

Remark 2.5. *Theorem 2.4 and [17, Remark 2.2] imply that [17, Theorem 1.1] holds for weak solutions (2.3) as in Definition 2.1. In [17] the result was proved under a technical assumption $|\nabla u| \in L^q(\Omega_T)$.*

We discuss existence and uniqueness results for the Dirichlet problem

$$\begin{cases} u_t - \operatorname{div} \mathcal{A}(z, \nabla u) = -\operatorname{div}(|F|^{p-2} F + a(z)|F|^{q-2} F) & \text{in } C_R \\ u = g & \text{on } \partial_p C_R, \end{cases} \quad (2.6)$$

where $\mathcal{A}(z, \xi) = b(z)(|\xi|^{p-2} \xi + a(z)|\xi|^{q-2} \xi)$ and $b(z) : C_R \rightarrow \mathbb{R}^+$ is a measurable function satisfying the ellipticity condition

$$\nu \leq b(z) \leq L \quad \text{for a.e. } z \in C_R, \quad (2.7)$$

for some constants $0 < \nu \leq L < \infty$. Here $C_R = D_R \times I_R$ and $\partial_p C_R$ is the parabolic boundary defined as the union of $D_R \times \{-R^2\}$ and $\partial D_R \times (-R^2, R^2)$. Moreover, $g \in C^\infty(\mathbb{R}^{n+1}, \mathbb{R}^N)$ and $u = g$ on $\partial_p C_R$ means that

$$u - g \in L^p(I_R, W_0^{1,p}(D_R, \mathbb{R}^N)) \quad \text{and} \quad \lim_{\epsilon \rightarrow 0^+} \int_{-R^2}^{-R^2+\epsilon} \int_{D_R} |u - g|^2 dx dt = 0.$$

Assume that u and w are weak solutions of (2.6). Then

$$(u - w)_t - \operatorname{div}(\mathcal{A}(z, \nabla u) - \mathcal{A}(z, \nabla w)) = 0 \quad \text{in } C_R \quad (2.8)$$

and $u - w = 0$ on $\partial_p C_R$ in the sense that on the lateral boundary we have

$$u - w \in L^p(I_R; W_0^{1,p}(D_R, \mathbb{R}^N)), \quad (2.9)$$

and on the initial boundary we have

$$\lim_{\epsilon \rightarrow 0^+} \int_{-R^2}^{-R^2+\epsilon} \int_{D_R} |u - w|^2 dx dt = 0. \quad (2.10)$$

In the case of the parabolic p -Laplace system ($a(\cdot) \equiv 0$), $u - w$ is an admissible test function for (2.8) and the uniqueness follows in a straightforward manner. However, $u - w$ may not be an admissible test function when $a(\cdot) \not\equiv 0$. Again, we will apply the Lipschitz truncation method to settle this problem. The uniqueness result is stated in the following theorem.

Theorem 2.6. *Assume that u and w are weak solutions to the Dirichlet boundary problem (2.6). Then $u = w$ a.e. in C_R .*

Then we discuss the existence of a weak solution to the Dirichlet problem above. In order to be able to apply the estimates in [16], we assume that the coefficient $b(\cdot)$ satisfies a VMO condition

$$\lim_{r \rightarrow 0^+} \sup_{|I| \leq r^2} \sup_{x_0 \in D_R} \int_{I \cap I_R} \int_{B_r(x_0) \cap D_R} |b(x, t) - (b)_{(B_r(x_0) \times I) \cap C_R}| dx dt = 0. \quad (2.11)$$

Theorem 2.7. *Assume that $g \equiv 0$ and that (2.11) holds true. Then there exists a weak solution to the Dirichlet problem (2.6). Moreover, there exists a sequence $(u_l)_{l \in \mathbb{N}}$ of weak solutions $u_l \in L^q(I_R; W_0^{1,q}(D_R, \mathbb{R}^N))$, $l \in \mathbb{N}$, to the problem*

$$\begin{cases} \partial_t u_l - \operatorname{div} \mathcal{A}_l(z, \nabla u_l) = -\operatorname{div}(|F_l|^{p-2} F_l + a_l(z)|F_l|^{q-2} F_l) & \text{in } C_R \\ u_l = 0 & \text{on } \partial_p C_R, \end{cases}$$

such that

$$\lim_{l \rightarrow \infty} \iint_{C_R} H(z, |\nabla u - \nabla u_l|) dz = 0.$$

Here $(F_l)_{l \in \mathbb{N}}$ is a sequence of truncations $F_l \in L^\infty(C_R, \mathbb{R}^{N^n})$, $l \in \mathbb{N}$, of F satisfying

$$\lim_{l \rightarrow \infty} \iint_{C_R} H(z, |F - F_l|) dz = 0,$$

and \mathcal{A}_l is a perturbed q -Laplace operator from \mathcal{A} , defined with a positive and decreasing sequence $(\epsilon_l)_{l \in \mathbb{N}}$ as

$$\mathcal{A}_l(z, \xi) = b(z) (|\xi|^{p-2} \xi + a_l(z)|\xi|^{q-2} \xi), \quad a_l(z) = a(z) + \epsilon_l, \quad \lim_{l \rightarrow \infty} \epsilon_l = 0.$$

3. WHITNEY DECOMPOSITION

In this section we provide a Whitney covering result in Proposition 3.2 which is applied in the Lipschitz truncation method for parabolic double-phase problems in Section 4. The classical Whitney covering lemma decomposes an open set U into a family of open sets $\{U_i\}_{i \in \mathbb{N}}$ according to a given metric $d(\cdot, \cdot)$. It has been employed for parabolic p -Laplace systems with the parabolic p -Laplace metric to prove the Lipschitz truncation method, see [5, 19]. We extend this by decomposing U into $\{U_i\}_{i \in \mathbb{N}}$ according to pointwise metrics depending on the coefficient a .

The decomposition is based on the strong maximal function defined for $f \in L^1_{\text{loc}}(\mathbb{R}^{n+1})$ as

$$Mf(z) = \sup_{z \in Q} \iint_Q |f| dw, \quad (3.1)$$

where the supremum is taken over all parabolic cylinders of the type in (2.2) and satisfying $z \in Q$. It follows immediately that

$$Mf(z) \leq \sup_{t \in (t_1, t_2)} \int_{t_1}^{t_2} \sup_{x \in B} \int_B |f| dy ds = M_t(M_x(|f|))(z),$$

where M_x and M_t are the noncentered Hardy–Littlewood maximal operators with respect to space and time. By repeated application of the Hardy–Littlewood–Wiener maximal function theorem we obtain the following bound for the strong maximal function.

Lemma 3.1. *Let $1 < \sigma < \infty$ and $f \in L^\sigma(\mathbb{R}^{n+1})$. Then there exists a constant $c = c(n, \sigma)$ such that*

$$\iint_{\mathbb{R}^{n+1}} |Mf|^\sigma dz \leq c \iint_{\mathbb{R}^{n+1}} |f|^\sigma dz.$$

3.1. Initial definitions. We start by extending a to \mathbb{R}^{n+1} . Since $s \mapsto s^\alpha$ is a concave and subadditive function on \mathbb{R}^+ , the extended function

$$\tilde{a}(z) = \inf_{w=(y,s) \in \Omega_T} \left\{ a(w) + [a]_\alpha (|x-y|^\alpha + |t-s|^{\frac{\alpha}{2}}) \right\}, \quad z = (x, t) \in \mathbb{R}^{n+1}, \quad (3.2)$$

belongs to $C^{\alpha, \alpha/2}(\mathbb{R}^{n+1})$ (see [18, Theorem 2.7] for the details). With slight abuse of notation we denote $a(z) = \min\{\tilde{a}(z), \|a\|_{L^\infty(\Omega_T)}\}$. Note that $\|a\|_{L^\infty(\mathbb{R}^{n+1})} = \|a\|_{L^\infty(\Omega_T)}$.

Let $f \in L^p(\mathbb{R}^{n+1})$, $f \geq 0$, be such that

$$\iint_{\mathbb{R}^{n+1}} (f^p + af^q) dz < \infty, \quad (3.3)$$

and let $d = \frac{q-1+p}{2}$. By Lemma 3.1 there exists $\Lambda_0 > 1 + \|a\|_{L^\infty(\mathbb{R}^{n+1})}$ such that

$$\iint_{\mathbb{R}^{n+1}} \left(M(f^d + (af^q)^{\frac{d}{p}})(z) \right)^{\frac{p}{d}} dz \leq c(n, p, q) \iint_{\mathbb{R}^{n+1}} (f^p + af^q) dz \leq \Lambda_0.$$

Let $\Lambda > \Lambda_0$. We consider the set

$$E(\Lambda) = \left\{ z \in \mathbb{R}^{n+1} : M(f^d + (af^q)^{\frac{d}{p}})(z) \leq \Lambda^{\frac{d}{p}} \right\}, \quad (3.4)$$

where $M(\cdot)$ is the strong maximal operator defined in (3.1). Since the maximal function is lower semicontinuous, $E(\Lambda)^c = \mathbb{R}^{n+1} \setminus E(\Lambda)$ is an open subset of \mathbb{R}^{n+1} . Moreover, Chebyshev's inequality implies that

$$\lim_{\Lambda \rightarrow \infty} \Lambda |E(\Lambda)^c| \leq \lim_{\Lambda \rightarrow \infty} \iint_{E(\Lambda)^c} \left(M(f^d + (af^q)^{\frac{d}{p}})(z) \right)^{\frac{p}{d}} dz = 0. \quad (3.5)$$

Assuming $E(\Lambda)^c \neq \emptyset$, we construct a Whitney decomposition of $E(\Lambda)^c$. Since $s \mapsto s^p + a(z)s^q$ is an increasing function on \mathbb{R}^+ , for every $z \in E(\Lambda)^c$ there exists a unique $\lambda_z > 1$ such that $\lambda_z^p + a(z)\lambda_z^q = \Lambda$. Let

$$K = 2^{\frac{1}{2}} + 800[a]_\alpha \left(\frac{1}{|B_1|} \iint_{\mathbb{R}^{n+1}} \left(M(f^d + (af^q)^{\frac{d}{p}})(z) \right)^{\frac{p}{d}} dz \right)^{\frac{\alpha}{n+2}}. \quad (3.6)$$

We consider a family of metrics $\{d_z(\cdot, \cdot)\}_{z \in E(\Lambda)^c}$ given by

$$d_z(z_1, z_2) = \begin{cases} \max \left\{ |x_1 - x_2|, \sqrt{\lambda_z^{p-2} |t_1 - t_2|} \right\}, & \text{if } K^2 \lambda_z^p \geq a(z) \lambda_z^q, \\ \max \left\{ |x_1 - x_2|, \sqrt{\Lambda \lambda_z^{-2} |t_1 - t_2|} \right\}, & \text{if } K^2 \lambda_z^p < a(z) \lambda_z^q. \end{cases}$$

For every $z \in E(\Lambda)^c$ we define a distance from z to $E(\Lambda)$ as

$$4r_z = d_z(z, E(\Lambda)) = \inf_{w \in E(\Lambda)} d_z(z, w). \quad (3.7)$$

Using this distance, we construct a family of open subsets $\{U_z^\Lambda\}_{z \in E(\Lambda)^c}$ defined as

$$U_z^\Lambda = \begin{cases} Q_z^{\lambda_z}, & \text{if } K^2 \lambda_z^p \geq a(z) \lambda_z^q, \\ G_z^{\lambda_z}, & \text{if } K^2 \lambda_z^p < a(z) \lambda_z^q, \end{cases}$$

where $Q_z^{\lambda_z} = B_{r_z}(x) \times I_{r_z}^{\lambda_z}(t)$ is a p -intrinsic cylinder and $G_z^{\lambda_z} = B_{r_z}(x) \times J_{r_z}^{\lambda_z}(t)$ is a (p, q) -intrinsic cylinder. The case $K^2 \lambda_z^p \geq a(z) \lambda_z^q$ is called the p -intrinsic case and $K^2 \lambda_z^p < a(z) \lambda_z^q$ the (p, q) -intrinsic case.

In the rest of this section we use a Vitali type argument to find a countable collection of points $z_i \in E(\Lambda)^c$ such that the corresponding subfamily $\{U_{z_i}^\Lambda\}_{i \in \mathbb{N}}$ satisfies the properties in Proposition 3.2. The proof of the proposition is divided into Lemmas 3.3 - 3.13. Here, and in the remaining of the paper, we denote

$$\lambda_i = \lambda_{z_i}, \quad r_i = r_{z_i}, \quad d_i(\cdot, \cdot) = d_{z_i}(\cdot, \cdot), \quad Q_i = Q_{z_i}^{\lambda_i}, \quad G_i = G_{z_i}^{\lambda_i},$$

and

$$U_i = B_i \times I_i = \begin{cases} Q_i, & \text{if } K^2 \lambda_i^p \geq a(z_i) \lambda_i^q, \\ G_i, & \text{if } K^2 \lambda_i^p < a(z_i) \lambda_i^q. \end{cases} \quad (3.8)$$

Moreover, we write

$$d_i(U_i, E(\Lambda)) = \inf_{z \in U_i, w \in E(\Lambda)} d_i(z, w), \quad \mathcal{I}_i = \{j \in \mathbb{N} : 2U_i \cap 2U_j \neq \emptyset\} \quad (3.9)$$

and

$$K_i = \begin{cases} 50K^2 & \text{if } U_i = Q_i, \\ 50K & \text{if } U_i = G_i. \end{cases} \quad (3.10)$$

Proposition 3.2. *Let K be as in (3.6) and $E(\Lambda)$ as in (3.4). There exists a collection $\{U_i\}_{i \in \mathbb{N}}$ of cylinders defined as in (3.8) satisfying the following properties:*

- (i) $\cup_{i \in \mathbb{N}} U_i = E(\Lambda)^c$.
- (ii) $\frac{1}{6K} U_i \cap \frac{1}{6K} U_j = \emptyset$ for every $i, j \in \mathbb{N}$ with $i \neq j$.
- (iii) $3r_i \leq d_i(U_i, E(\Lambda)) \leq 4r_i$ for every $i \in \mathbb{N}$.
- (iv) $4U_i \subset E(\Lambda)^c$ and $5U_i \cap E(\Lambda) \neq \emptyset$ for every $i \in \mathbb{N}$.
- (v) $(12K)^{-1} r_j \leq r_i \leq 12K r_j$ for every $i \in \mathbb{N}, j \in \mathcal{I}_i$.
- (vi) $K^{-\frac{2}{p}} \lambda_j \leq \lambda_i \leq K^{\frac{2}{p}} \lambda_j$ for every $j \in \mathcal{I}_i$. Moreover, if $U_i = G_i$, then $2^{-\frac{1}{p}} \lambda_j \leq \lambda_i \leq 2^{\frac{1}{p}} \lambda_j$.
- (vii) There exists a constant $c = c(n, K)$ such that $|U_i| \leq c|U_j|$, for every $i \in \mathbb{N}$ and $j \in \mathcal{I}_i$.
- (viii) $2U_j \subset K_i U_i$ for every $i \in \mathbb{N}, j \in \mathcal{I}_i$.
- (ix) If $U_i = Q_i$, then there exists a constant $c = c([a]_\alpha, \alpha, K)$ such that $r_i^\alpha \lambda_i^q \leq c \lambda_i^p$.
- (x) If $U_i = G_i$, then $\frac{a(z_i)}{2} \leq a(z) \leq 2a(z_i)$ for every $z \in 200K Q_{r_i}(z_i)$.
- (xi) For any $i \in \mathbb{N}$, the cardinality of \mathcal{I}_i , denoted by $|\mathcal{I}_i|$, is finite. Moreover, there exists a constant $c = c(n, K)$, such that $|\mathcal{I}_i| \leq c$.

Additionally, there exists a partition of unity $\{\omega_i\}_{i \in \mathbb{N}}$ subordinate to the Whitney decomposition $\{2U_i\}_{i \in \mathbb{N}}$ with the following properties:

$$(xii) \quad 0 \leq \omega_i \leq 1, \quad \omega_i \in C_0^\infty(2U_i) \text{ for every } i \in \mathbb{N} \text{ and } \sum_{i \in \mathbb{N}} \omega_i = 1 \text{ on } E(\Lambda)^c.$$

$$(xiii) \quad \text{There exists a constant } c = c(n, K) \text{ such that } \|\nabla \omega_j\|_\infty \leq cr_i^{-1}, \text{ for every } i \in \mathbb{N} \text{ and } j \in \mathcal{I}_i.$$

$$(xiv) \quad \text{There exists a constant } c = c(n, K) \text{ such that}$$

$$\|\partial_t \omega_j\|_\infty \leq \begin{cases} cr_i^{-2} \lambda_i^{p-2}, & \text{if } U_i = Q_i, \\ cr_i^{-2} \lambda_i^{-2} \Lambda, & \text{if } U_i = G_i, \end{cases}$$

for every $i \in \mathbb{N}$ and $j \in \mathcal{I}_i$.

3.2. Estimates for scaling factors. In this subsection, we provide lemmas that allow us to compare the scaling factors of neighboring cylinders. The first lemma states that coefficient function $a(\cdot)$ is comparable in the standard cylinder for (p, q) -intrinsic case.

Lemma 3.3. *Assume $z \in E(\Lambda)^c$ and $K^2\lambda_z^p < a(z)\lambda_z^q$. Then $\frac{a(z)}{2} \leq a(w) \leq 2a(z)$ for every $w \in 200KQ_{r_z}(z)$.*

Proof. We claim that

$$2[a]_\alpha (200Kr_z)^\alpha < a(z). \quad (3.11)$$

Assume for contradiction that $2[a]_\alpha (200Kr_z)^\alpha \geq a(z)$. Since $G_z^{\lambda_z} \subset E(\Lambda)^c$ and $a(z)\lambda_z^q < \Lambda$, we have

$$a(z)\lambda_z^q < \iint_{G_z^{\lambda_z}} \left(M(f^d + (af^q)^{\frac{d}{p}})(w) \right)^{\frac{p}{d}} dw.$$

Using the facts that $\Lambda = \lambda_z^p + a(z)\lambda_z^q$ and $\lambda_z^p < a(z)\lambda_z^q$, we obtain

$$\begin{aligned} a(z)\lambda_z^q &< \iint_{G_z^{\lambda_z}} \left(M(f^d + (af^q)^{\frac{d}{p}})(w) \right)^{\frac{p}{d}} dw \\ &= \frac{\Lambda\lambda_z^{-2}}{2|B_1|r_z^{n+2}} \iint_{G_z^{\lambda_z}} \left(M(f^d + (af^q)^{\frac{d}{p}})(w) \right)^{\frac{p}{d}} dw \\ &< \frac{a(z)\lambda_z^{q-2}}{|B_1|r_z^{n+2}} \iint_{G_z^{\lambda_z}} \left(M(f^d + (af^q)^{\frac{d}{p}})(w) \right)^{\frac{p}{d}} dw. \end{aligned}$$

Dividing both sides by $a(z)\lambda_z^{q-2}r_z^{-n-2}$ and raising to power $\frac{\alpha}{n+2}$, it follows from (3.6) that

$$r_z^\alpha \lambda_z^{\frac{2\alpha}{n+2}} \leq \left(\frac{1}{|B_1|} \iint_{\mathbb{R}^{n+1}} \left(M(f^d + (af^q)^{\frac{d}{p}})(w) \right)^{\frac{p}{d}} dw \right)^{\frac{\alpha}{n+2}} \leq \frac{1}{800[a]_\alpha} K. \quad (3.12)$$

Since $K^2\lambda_z^p < a(z)\lambda_z^q$ and $q \leq p + \frac{2\alpha}{n+2}$, the counter assumption and (3.12) imply

$$K^2\lambda_z^p < a(z)\lambda_z^q \leq 2[a]_\alpha (200Kr_z)^\alpha \lambda_z^p \lambda_z^{\frac{2\alpha}{n+2}} < 400[a]_\alpha K \lambda_z^p r_z^\alpha \lambda_z^{\frac{2\alpha}{n+2}} \leq \frac{1}{2} K^2 \lambda_z^p.$$

This is a contradiction and thus (3.11) holds true.

By (3.11) we have

$$2[a]_\alpha (200Kr_z)^\alpha < a(z) \leq \inf_{w \in 200KQ_{r_z}(z)} a(w) + [a]_\alpha (200Kr_z)^\alpha$$

and thus

$$[a]_\alpha (200Kr_z)^\alpha < \inf_{w \in 200KQ_{r_z}(z)} a(w).$$

It follows that

$$\sup_{w \in 200KQ_{r_z}(z)} a(w) \leq \inf_{w \in 200KQ_{r_z}(z)} a(w) + [a]_\alpha (200Kr_z)^\alpha < 2 \inf_{w \in 200KQ_{r_z}(z)} a(w).$$

This completes the proof. \square

The second lemma states that if $a(\cdot)$ is comparable at two points, then $\lambda_{(\cdot)}$ is comparable at these points.

Lemma 3.4. *Assume $z, w \in E(\Lambda)^c$ and $\frac{a(z)}{2} \leq a(w) \leq 2a(z)$. Then $2^{-\frac{1}{p}}\lambda_w \leq \lambda_z \leq 2^{\frac{1}{p}}\lambda_w$. Moreover, the above inequality holds provided $K^2\lambda_z^p < a(z)\lambda_z^q$ and $w \in 200KQ_{r_z}(z)$.*

Proof. The second statement of the lemma follows from Lemma 3.3 and the first statement of this lemma. To prove the first statement, it suffices to show $\lambda_z \leq 2^{\frac{1}{p}}\lambda_w$. Assume for contradiction that $\lambda_z > 2^{\frac{1}{p}}\lambda_w$. Then

$$\Lambda = \lambda_z^p + a(z)\lambda_z^q > 2\lambda_w^p + 2^{\frac{q}{p}}a(z)\lambda_w^q,$$

and using the assumption on $a(\cdot)$, we get

$$\Lambda > 2\lambda_w^p + 2^{\frac{q}{p}-1}a(w)\lambda_w^q > \lambda_w^p + a(w)\lambda_w^q = \Lambda.$$

This is a contradiction and the proof is completed. \square

We need the estimate between λ_z and λ_w also in the p -intrinsic case. For this, we will use the continuity of $a(\cdot)$ and the following lemma.

Lemma 3.5. *Assume $z \in E(\Lambda)^c$ and $K^2\lambda_z^p \geq a(z)\lambda_z^q$. Then $[a]_\alpha(50Kr_z)^\alpha\lambda_z^q \leq (K^2 - 1)\lambda_z^p$.*

Proof. We argue as in the proof of Lemma 3.3. Observe that $Q_z^{\lambda_z} \subset E(\Lambda)^c$ and therefore

$$\begin{aligned} \lambda_z^p &< \iint_{Q_z^{\lambda_z}} \left(M(f^d + (af^q)^{\frac{d}{p}})(w) \right)^{\frac{p}{d}} dw \\ &= \frac{\lambda_z^{p-2}}{2|B_1|r_z^{n+2}} \iint_{Q_z^{\lambda_z}} \left(M(f^d + (af^q)^{\frac{d}{p}})(w) \right)^{\frac{p}{d}} dw. \end{aligned}$$

After dividing both sides and raising to power $\frac{\alpha}{n+2}$, it follows from (3.6) that

$$r_z^\alpha \lambda_z^{\frac{2\alpha}{n+2}} < \left(\frac{1}{|B_1|} \iint_{\mathbb{R}^{n+1}} \left(M(f^d + (af^q)^{\frac{d}{p}})(w) \right)^{\frac{p}{d}} dw \right)^{\frac{\alpha}{n+2}} \leq \frac{1}{50[a]_\alpha} (K-1).$$

Using this and the fact that $q \leq p + \frac{2\alpha}{n+2}$ we obtain

$$\begin{aligned} [a]_\alpha(50Kr_z)^\alpha\lambda_z^q &\leq 50[a]_\alpha \left(\frac{1}{|B_1|} \iint_{\mathbb{R}^{n+1}} \left(M(f^d + (af^q)^{\frac{d}{p}})(w) \right)^{\frac{p}{d}} dw \right)^{\frac{\alpha}{n+2}} K\lambda_z^p \\ &\leq (K-1)K\lambda_z^p. \end{aligned}$$

Since $(K-1)K \leq K^2 - 1$, the proof is completed. \square

Finally, we state the comparability of $\lambda_{(\cdot)}$ in the p -intrinsic case.

Lemma 3.6. *Assume $z \in E(\Lambda)^c$ and $K^2\lambda_z^p \geq a(z)\lambda_z^q$. If $w \in 50KQ_{r_z}(z)$, then $\lambda_z \leq K^{\frac{2}{p}}\lambda_w$.*

Proof. Assume for contradiction that $\lambda_w < K^{-\frac{2}{p}}\lambda_z$. By the Hölder regularity in (2.5) we have

$$\Lambda = \lambda_w^p + a(w)\lambda_w^q \leq \lambda_w^p + a(z)\lambda_w^q + [a]_\alpha(50Kr_z)^\alpha\lambda_w^q.$$

Therefore the counter assumption and Lemma 3.5 imply that

$$\begin{aligned} \Lambda &\leq \frac{1}{K^2}(\lambda_z^p + a(z)\lambda_z^q) + \frac{[a]_\alpha(50Kr_z)^\alpha}{K^2}\lambda_z^q \\ &\leq \frac{1}{K^2}(\lambda_z^p + a(z)\lambda_z^q) + \frac{K^2 - 1}{K^2}\lambda_z^p < \lambda_z^p + a(z)\lambda_z^q = \Lambda. \end{aligned}$$

This is a contradiction and the proof is completed. \square

3.3. Vitali-type covering argument. We select a countable family of intrinsic cylinders from $\{U_z^\Lambda\}_{z \in E(\Lambda)^c}$ such that they cover $E(\Lambda)^c$ and $\frac{1}{6K}$ -times the chosen cylinders are pairwise disjoint. We start by claiming that $r_z, z \in E(\Lambda)^c$, are uniformly bounded. Since $|E(\Lambda)^c| < \infty$ and $\Lambda > 1 + \|a\|_{L^\infty(\mathbb{R}^{n+1})}$, there exist $\lambda = \lambda(\Lambda, \|a\|_{L^\infty(\mathbb{R}^{n+1})}) > 1$ and $R = R(\Lambda, \|a\|_{L^\infty(\mathbb{R}^{n+1})}) > 0$ such that

$$\lambda^p + \|a\|_{L^\infty(\mathbb{R}^{n+1})}\lambda^q = \Lambda \tag{3.13}$$

and $|B_R \times (-\lambda^2\Lambda^{-1}R^2, \lambda^2\Lambda^{-1}R^2)| = |E(\Lambda)^c|$. It is easy to see that $\lambda \leq \lambda_z \leq \Lambda$ for every $z \in E(\Lambda)^c$. Therefore, if there exists $z \in E(\Lambda)^c$ such that $r_z > R$, then

$$|E(\Lambda)^c| = |B_R \times (-\lambda^2\Lambda^{-1}R^2, \lambda^2\Lambda^{-1}R^2)| < |U_z^\Lambda| \leq |E(\Lambda)^c|.$$

This is a contradiction and we conclude that $r_z \leq R$ for every $z \in E(\Lambda)^c$.

Let $\mathcal{F} = \{\frac{1}{6K}U_z^\Lambda\}_{z \in E(\Lambda)^c}$ and

$$\mathcal{F}_j = \left\{ \frac{1}{6K}U_z^\Lambda \in \mathcal{F} : \frac{R}{2^j} < r_z \leq \frac{R}{2^{j-1}} \right\}, \quad j \in \mathbb{N}.$$

Since $r_z, z \in E(\Lambda)^c$, are uniformly bounded, we have $\mathcal{F} = \cup_{j \in \mathbb{N}} \mathcal{F}_j$. We obtain a disjoint subcollection $\mathcal{G}_j \subset \mathcal{F}_j$ as follows. Let \mathcal{G}_1 be a maximal disjoint collection of cylinders in \mathcal{F}_1 . We proceed recursively. If $\mathcal{G}_1, \mathcal{G}_2, \dots, \mathcal{G}_{k-1}$ have been selected, let \mathcal{G}_k be any maximal disjoint collection of

$$\left\{ \frac{1}{6K} U_z^\Lambda \in \mathcal{F}_k : \frac{1}{6K} U_z^\Lambda \cap \frac{1}{6K} U_w^\Lambda = \emptyset \text{ for every } \frac{1}{6K} U_w^\Lambda \in \bigcup_{j=1}^{k-1} \mathcal{G}_j \right\}.$$

Note that for λ defined in (3.13) and $z \in \mathcal{G}_j$, we have

$$\frac{1}{2^j 6K} (B_R(x) \times (t - \lambda^2 \Lambda^{-1} R^2, t + \lambda^2 \Lambda^{-1} R^2)) \subset \frac{1}{6K} U_z^\Lambda \subset E(\Lambda)^c.$$

Since \mathcal{G}_j is a pairwise disjoint collection and $|E(\Lambda)^c| < \infty$, this implies that the cardinality of \mathcal{G}_j is finite. Let $\mathcal{G} = \cup_{j \in \mathbb{N}} \mathcal{G}_j$. By the construction, \mathcal{G} is a countable subfamily of pairwise disjoint cylinders.

In the rest of this subsection we prove that the cylinders in \mathcal{G} cover $E(\Lambda)^c$. Let $\frac{1}{6K} U_w^\Lambda \in \mathcal{F}$. There exists $i \in \mathbb{N}$ such that $\frac{1}{6K} U_w^\Lambda \in \mathcal{F}_i$ and by the maximal disjointness of \mathcal{G}_j , there exists a cylinder $\frac{1}{6K} U_z^\Lambda \in \cup_{j=1}^i \mathcal{G}_j$ such that

$$\frac{1}{6K} U_w \cap \frac{1}{6K} U_z \neq \emptyset. \quad (3.14)$$

Note that since $\frac{1}{6K} U_w \in \mathcal{F}_i$ and $\frac{1}{6K} U_z \in \mathcal{F}_j$ for some $j \in \{1, \dots, i\}$, we have

$$r_w \leq 2r_z. \quad (3.15)$$

We claim that $6K$ is the constant in the Vitali-type covering, that is,

$$\frac{1}{6K} U_w \subset U_z. \quad (3.16)$$

Let $z = (x, t)$ and $w = (y, s)$. For the inclusion in the spatial direction, observe that (3.14) implies

$$\frac{1}{6K} B_{r_w}(y) \cap \frac{1}{6K} B_{r_z} \neq \emptyset.$$

From (3.15) and the proof of the standard Vitali covering lemma, we obtain

$$\frac{1}{6K} B_{r_w}(y) \subset \frac{5}{6K} B_{r_z}(x) \subset B_{r_z}(x).$$

On the other hand, it follows from (3.14) that $Q_{r_w}(w) \cap Q_{r_z}(z) \neq \emptyset$. Again, it easily follows from the standard Vitali covering argument with (3.15) that $Q_{r_w}(w) \subset 5Q_{r_z}(z)$. In particular, we have

$$w \in 5Q_{r_z}(z) \subset 50KQ_{r_z}(z). \quad (3.17)$$

It remains to prove the inclusion in (3.16) in the time direction. We divide the proof into four cases.

Case 1: $U_w^\Lambda = Q_w^{\lambda_w}$ and $U_z^\Lambda = Q_z^{\lambda_z}$. By (3.15), we get

$$\begin{aligned} |\tau - t| &\leq |\tau - s| + |s - t| \leq \left| \frac{1}{6K} I_{r_w}^{\lambda_w}(s) \right| + \frac{1}{2} \left| \frac{1}{6K} I_{r_z}^{\lambda_z}(t) \right| \\ &= 2\lambda_w^{2-p} \left(\frac{r_w}{6K} \right)^2 + \lambda_z^{2-p} \left(\frac{r_z}{6K} \right)^2 \\ &\leq (8\lambda_w^{2-p} + \lambda_z^{2-p}) \left(\frac{r_z}{6K} \right)^2 \end{aligned}$$

for every $\tau \in \frac{1}{6K} I_{r_w}^{\lambda_w}(s)$. Lemma 3.6 and (3.17) imply $\lambda_z \leq K^{\frac{2}{p}} \lambda_w$ and thus we have

$$\begin{aligned} |\tau - t| &\leq (8K^{\frac{2(p-2)}{p}} + 1) \lambda_z^{2-p} \left(\frac{r_z}{6K} \right)^2 \\ &\leq (9K^2) \lambda_z^{2-p} \left(\frac{r_z}{6K} \right)^2 \leq \lambda_z^{2-p} r_z^2 = |I_{r_z}^{\lambda_z}(t)|, \end{aligned}$$

where we also applied the fact that $\frac{p-2}{p} \leq 1$. Therefore, the conclusion in (3.16) holds in this case.

Case 2: $U_w^\Lambda = G_w^{\lambda_w}$ and $U_z^\Lambda = Q_z^{\lambda_z}$. In this case we have $G_w^{\lambda_w} \subset Q_w^{\lambda_w}$. The conclusion follows from the argument in the previous case.

Case 3: $U_w^\Lambda = Q_w^{\lambda_w}$ and $U_z^\Lambda = G_z^{\lambda_z}$. By (3.15) we get

$$\begin{aligned} |\tau - t| &\leq |\tau - s| + |s - t| \leq \left| \frac{1}{6K} I_{r_w}^{\lambda_w}(s) \right| + \frac{1}{2} \left| \frac{1}{6K} J_{r_z}^{\lambda_z}(t) \right| \\ &= 2\lambda_w^{2-p} \left(\frac{r_w}{6K} \right)^2 + \lambda_z^2 \Lambda^{-1} \left(\frac{r_z}{6K} \right)^2 \\ &\leq (8\lambda_w^{2-p} + \lambda_z^2 \Lambda^{-1}) \left(\frac{r_z}{6K} \right)^2 \end{aligned}$$

for every $\tau \in \frac{1}{6K} I_{r_w}^{\lambda_w}(s)$. In this case we have $a(w)\lambda_w^q \leq K^2\lambda_w^p$, and thus we obtain

$$\lambda_w^{2-p} = 2 \frac{\lambda_w^2}{\lambda_w^p + \lambda_w^q} \leq 2K^2 \frac{\lambda_w^2}{\lambda_w^p + a(w)\lambda_w^q} = 2K^2 \lambda_w^2 \Lambda^{-1}.$$

Lemma 3.4 and (3.17) imply that

$$2K^2 \lambda_w^2 \Lambda^{-1} \leq 2K^2 (2^{\frac{1}{p}} \lambda_z)^2 \Lambda^{-1} \leq 4K^2 \lambda_z^2 \Lambda^{-1}.$$

It follows that

$$|\tau - t| \leq (32K^2 + 1) \lambda_z^2 \Lambda^{-1} \left(\frac{r_z}{6K} \right)^2 \leq 33K^2 \lambda_z^2 \Lambda^{-1} \left(\frac{r_z}{6K} \right)^2 \leq |J_{r_z}^{\lambda_z}(t)|,$$

and thus (3.16) holds in this case.

Case 4: $U_w^\Lambda = G_w^{\lambda_w}$ and $U_z^\Lambda = G_z^{\lambda_z}$. By (3.15) we get

$$\begin{aligned} |\tau - t| &\leq |\tau - s| + |s - t| \leq \left| \frac{1}{6K} J_{r_w}^{\lambda_w}(s) \right| + \frac{1}{2} \left| \frac{1}{6K} J_{r_z}^{\lambda_z}(t) \right| \\ &= 2\lambda_w^2 \Lambda^{-1} \left(\frac{r_w}{6K} \right)^2 + \lambda_z^2 \Lambda^{-1} \left(\frac{r_z}{6K} \right)^2 \\ &\leq (8\lambda_w^2 \Lambda^{-1} + \lambda_z^2 \Lambda^{-1}) \left(\frac{r_z}{6K} \right)^2 \end{aligned}$$

for every $\tau \in \frac{1}{6K} I_{r_w}^{\lambda_w}(s)$. Applying (3.17) and Lemma 3.4 gives

$$|\tau - t| \leq 17\lambda_z^2 \Lambda^{-1} \left(\frac{r_z}{6K} \right)^2 \leq |J_{r_z}^{\lambda_z}(t)|.$$

This completes the proof of (3.16) in the final case. Thus we have found a countable covering family $\{U_i\}_{i \in \mathbb{N}}$ of intrinsic cylinders defined as in (3.8) and with $\frac{1}{6K}U_i$ being pairwise disjoint.

3.4. Properties of the covering. In this subsection we show that the Whitney decomposition $\{U_i\}_{i \in \mathbb{N}}$ defined in the previous subsection satisfies the remaining properties in Proposition 3.2

Lemma 3.7. *We have $3r_i \leq d_i(U_i, E(\Lambda)) \leq 4r_i$ for every $i \in \mathbb{N}$.*

Proof. From the choice of r_i in (3.7), we have

$$d_i(U_i, E(\Lambda)) \leq d_i(z_i, E(\Lambda)) = 4r_i.$$

Therefore the second inequality holds. To show the first inequality, note that for any $z \in U_i$, triangle inequality gives

$$d_i(z, E(\Lambda)) \geq d_i(z_i, E(\Lambda)) - d_i(z, z_i) \geq 4r_i - r_i = 3r_i.$$

Taking infimum over $z \in U_i$, we get $d_i(U_i, E(\Lambda)) \geq 3r_i$. This completes the proof. \square

Next we will discuss properties (v)-(viii) in Proposition 3.2. First we show that r_j , λ_j and $d_j(\cdot, \cdot)$ are comparable in $j \in \mathcal{I}_i$, defined as in (3.9), by using the lemmas in Section 3.1.

Lemma 3.8. *We have $(12K)^{-1}r_j \leq r_i \leq 12Kr_j$ for every $i \in \mathbb{N}$ and $j \in \mathcal{I}_i$. Moreover, if $U_i = G_i$, then $r_j \leq 12r_i$.*

Proof. To prove the first statement it is enough to show $r_i \leq 12Kr_j$ as there is no difference in the roles of i and j . Let $w \in 2U_i \cap 2U_j$. Using Lemma 3.7 and triangle inequality, we have

$$3r_i \leq d_i(z_i, E(\Lambda)) \leq d_i(z_i, w) + d_i(w, E(\Lambda)) \leq 2r_i + d_i(w, E(\Lambda)).$$

On the other hand, Lemma 3.7 and triangle inequality give

$$d_j(w, E(\Lambda)) \leq d_j(w, z_j) + d_j(z_j, E(\Lambda)) \leq 2r_j + 4r_j = 6r_j.$$

Combining these estimates, we have

$$r_i \leq d_i(w, E(\Lambda)) \quad \text{and} \quad d_j(w, E(\Lambda)) \leq 6r_j. \quad (3.18)$$

To finish the proof we consider four cases to compare the $d_i(z, w)$ and $d_j(z, w)$.

Case 1: $U_i = Q_i$ and $U_j = Q_j$. Using $\lambda_i^p \leq \Lambda$ and $p \geq 2$, we have

$$d_i(z, w) = \max \left\{ |x - y|, \sqrt{\lambda_i^{p-2} |s - t|} \right\} \leq \max \left\{ |x - y|, \sqrt{\Lambda^{\frac{p-2}{p}} |s - t|} \right\}.$$

Since $\Lambda = \lambda_j^p + a(z_j)\lambda_j^q$ and $a(z_j)\lambda_j^q \leq K^2\lambda_j^p$, we have

$$\Lambda^{\frac{p-2}{p}} = (\lambda_j^p + a(z_j)\lambda_j^q)^{\frac{p-2}{p}} \leq (2K^2\lambda_j^p)^{\frac{p-2}{p}}$$

and therefore

$$d_i(z, w) \leq (2K)^{\frac{p-2}{p}} \max \left\{ |x - y|, \sqrt{\lambda_j^{p-2} |s - t|} \right\} \leq 2Kd_j(z, w).$$

The above inequality and (3.18) imply that $r_i \leq 12Kr_j$.

Case 2: $U_i = Q_i$ and $U_j = G_j$. Since $\lambda_i^p + a(z_i)\lambda_i^q = \lambda_j^p + a(z_j)\lambda_j^q = \Lambda$ and $p \geq 2$, we have

$$\lambda_i^{p-2} \leq \Lambda^{\frac{p-2}{p}} = \Lambda(\lambda_j^p + a(z_j)\lambda_j^q)^{-\frac{2}{p}} \leq \Lambda\lambda_j^{-2}.$$

Thus, it follows that

$$\begin{aligned} d_i(z, w) &= \max \left\{ |x - y|, \sqrt{\lambda_i^{p-2} |s - t|} \right\} \\ &\leq \max \left\{ |x - y|, \sqrt{\Lambda\lambda_j^{-2} |s - t|} \right\} = d_j(z, w), \end{aligned}$$

and the above inequality along with (3.18) gives $r_i \leq 6r_j$.

Case 3: $U_i = G_i$ and $U_j = Q_j$. Note that $2U_i \cap 2U_j \neq \emptyset$ implies $2Q_{r_i}(z_i) \cap 2Q_{r_j}(z_j) \neq \emptyset$. This fact together with the previous case implies that $z_j \in 14Q_{r_i}(z_i)$. It follows from Lemma 3.4 that $\lambda_j \leq 2^{\frac{1}{p}}\lambda_i$, which, together with the fact that $K^2\lambda_j^p \geq a(z_j)\lambda_j^q$, gives

$$\Lambda\lambda_i^{-2} = (\lambda_j^p + a(z_j)\lambda_j^q)\lambda_i^{-2} \leq 2K^2\lambda_j^p\lambda_i^{-2} \leq 4K^2\lambda_j^{p-2}.$$

Therefore, we obtain

$$\begin{aligned} d_i(z, w) &= \max \left\{ |x - y|, \sqrt{\Lambda\lambda_i^{-2} |s - t|} \right\} \\ &\leq 2K \max \left\{ |x - y|, \sqrt{\lambda_j^{p-2} |s - t|} \right\} = 2Kd_j(z, w) \end{aligned}$$

and conclude with (3.18) that $r_i \leq 12Kr_j$.

Case 4: $U_i = G_i$ and $U_j = G_j$. We may assume $r_j \leq r_i$. Then $z_j \in 4Q_{r_i}(z_i)$ and therefore Lemma 3.4 gives $2^{-\frac{1}{p}}\lambda_j \leq \lambda_i \leq 2^{\frac{1}{p}}\lambda_j$. Thus

$$\begin{aligned} d_i(z, w) &= \max \left\{ |x - y|, \sqrt{\Lambda\lambda_i^{-2} |t - s|} \right\} \\ &\leq 2 \max \left\{ |x - y|, \sqrt{\Lambda\lambda_j^{-2} |t - s|} \right\} = 2d_j(z, w). \end{aligned}$$

Combining this estimate with (3.18), we obtain $r_i \leq 12r_j$ in the final case. The second statement of the lemma follows the second and fourth cases and we are finished. \square

It follows from the previous lemma that $2Q_{r_j}(z_j) \subset 50KQ_{r_i}(z_i)$ for all $j \in \mathcal{I}_i$. We write again the conclusions of Lemma 3.4 and Lemma 3.6.

Lemma 3.9. *We have $K^{-\frac{2}{p}}\lambda_j \leq \lambda_i \leq K^{\frac{2}{p}}\lambda_j$ for every $i \in \mathbb{N}$ and $j \in \mathcal{I}_i$. Moreover, if $U_i = G_i$, then $2^{-\frac{1}{p}}\lambda_j \leq \lambda_i \leq 2^{\frac{1}{p}}\lambda_j$.*

Proof. It is enough to prove the first inequality of the second statement. If $U_j = G_j$, then the first statement gives $\lambda_j \leq 2^{\frac{1}{p}} \lambda_i \leq K^{\frac{2}{p}} \lambda_i$. If $U_j = Q_j$, then $\lambda_j \leq K^{\frac{2}{p}} \lambda_i$ by Lemma 3.6, since $z_i \in 50KQ_{r_j}(z_j)$. \square

It follows from the previous two lemmas that the measures of neighboring cylinders are comparable.

Lemma 3.10. *There exists $c = c(n, K)$ such that $|U_i| \leq c|U_j|$ for any $i \in \mathbb{N}$ and $j \in \mathcal{I}_i$.*

Proof. If $U_j = Q_j$, it follows from $I_i \subset I_{r_i}^{\lambda_i}(t_i)$, Lemma 3.8 and Lemma 3.9 that

$$|U_i| \leq |Q_i| = 2|B_1|r_i^{n+2}\lambda_i^{2-p} \leq 2|B_1|(12K)^{n+2}K^{\frac{2(p-2)}{p}}r_j^{n+2}\lambda_j^{2-p} \leq (12K)^{n+4}|Q_j|.$$

The case $U_i = G_i$ and $U_j = G_j$ follows similarly from Lemma 3.8 and Lemma 3.9.

Finally, if $U_i = Q_i$ and $U_j = G_j$, we have by $a(z_i)\lambda_i^q \leq K^2\lambda_i^p$, Lemma 3.8 and Lemma 3.9 that

$$\begin{aligned} |Q_i| &= 2|B_1|r_i^{n+2}\lambda_i^{2-p} = \frac{4|B_1|r_i^{n+2}\lambda_i^2}{\lambda_i^p + \lambda_i^q} \leq \frac{4K^2|B_1|r_i^{n+2}\lambda_i^2}{\lambda_i^p + a(z_i)\lambda_i^q} \\ &\leq \frac{8 \cdot (12)^{n+2}K^2|B_1|r_j^{n+2}\lambda_j^2}{\Lambda} = c(n, K)|G_j|. \end{aligned}$$

This completes the proof. \square

In the following two lemmas we consider the inclusion property of U_j for $j \in \mathcal{I}_i$. The first lemma is for the p -intrinsic case and the second lemma is for the (p, q) -intrinsic case. We remark that the covering constant for p -intrinsic case is $50K^2$ while for (p, q) -intrinsic case it is $50K$.

Lemma 3.11. *Let $i \in \mathbb{N}$ be such that $U_i = Q_i$. We have $2U_j \subset 50K^2Q_i$ for every $j \in \mathcal{I}_i$.*

Proof. Since $2B_i \cap 2B_j \neq \emptyset$ and Lemma 3.8 gives $r_j \leq 12Kr_i$, it is easy to see that $2B_j \subset 50KB_i$. It remains to show the inclusion in the time direction. As $I_j \subset I_{r_j}^{\lambda_j}(t_j)$, we have for $\tau \in 2I_j$ that

$$|\tau - t_i| \leq |\tau - t_j| + |t_j - t_i| \leq |2I_{r_j}^{\lambda_j}(t_j)| + \frac{1}{2}|2I_{r_i}^{\lambda_i}(t_i)| = 2\lambda_j^{2-p}(2r_j)^2 + \lambda_i^{2-p}(2r_i)^2.$$

Applying $r_j \leq 12Kr_i$ and $\lambda_i \leq K^{\frac{2}{p}}\lambda_j$, which follow from Lemma 3.8 and Lemma 3.9, we get

$$|\tau - t_i| \leq 2(K^{-\frac{2}{p}}\lambda_i)^{2-p}(24Kr_i)^2 + \lambda_i^{2-p}(2r_i)^2 = (2 \cdot 24^2 \cdot K^{\frac{2(p-2)}{p}}K^2 + 2^2)\lambda_i^{2-p}r_i^2.$$

Since $\frac{2(p-2)}{p} + 2 \leq 4$ and

$$2 \cdot 24^2 \cdot K^{\frac{2(p-2)}{p}}K^2 + 2^2 \leq (48K^2)^2 + 2^2 \leq (48K^2 + 2)^2 \leq (50K^2)^2,$$

we get $|\tau - t_i| \leq \lambda_i^{2-p}(50K^2r_i)^2$ and thus $2U_j \subset 50K^2Q_i$. \square

Lemma 3.12. *Let $i \in \mathbb{N}$ be such that $U_i = G_i$. We have $2U_j \subset 50KG_i$ for every $j \in \mathcal{I}_i$.*

Proof. From Lemma 3.8, we have $r_j \leq 12r_i$. It follows that $2B_j \subset 50B_i$. To prove the inclusion in the time direction, we suppose first that $U_j = Q_j$. For $\tau \in 2I_{r_j}^{\lambda_j}(t_j)$, we have

$$|\tau - t_i| \leq |\tau - t_j| + |t_j - t_i| \leq |2I_{r_j}^{\lambda_j}(t_j)| + \frac{1}{2}|2J_{r_i}^{\lambda_i}(t_i)| = 2\lambda_j^{2-p}(2r_j)^2 + \lambda_i^2\Lambda^{-1}(2r_i)^2.$$

Since $K^2\lambda_j^p \geq a(z_j)\lambda_j^q$, $\Lambda = \lambda_j^p + a(z_j)\lambda_j^q$ and Lemma 3.9 implies $\lambda_j \leq 2^{\frac{1}{p}}\lambda_i$, we have

$$\lambda_j^{2-p} = 2\lambda_j^2 \frac{1}{\lambda_j^p + \lambda_j^q} \leq 2K^2\lambda_j^2 \frac{1}{\lambda_j^p + a(z_j)\lambda_j^q} \leq 2K^2\lambda_j^2\Lambda^{-1} \leq 2 \cdot 2^{\frac{2}{p}}K^2\lambda_i^2\Lambda^{-1},$$

and using that $r_j \leq 6r_i$ by the second case in the proof of Lemma 3.8, we obtain

$$|\tau - t_i| \leq (48^2K^2 + 2^2)\lambda_i^2\Lambda^{-1}r_i^2 \leq \lambda_i^2\Lambda^{-1}(50Kr_i)^2.$$

Therefore, $2Q_j \subset 50KG_i$. On the other hand, if $U_j = G_j$, then for any $\tau \in 2J_{r_j}^{\lambda_j}(t_j)$ we have

$$|\tau - t_i| \leq |2J_{r_j}^{\lambda_j}(t_j)| + \frac{1}{2}|2J_{r_i}^{\lambda_i}(t_i)| = 2\lambda_j^2\Lambda^{-1}(2r_j)^2 + \lambda_i^2\Lambda^{-1}(2r_i)^2.$$

Using $r_j \leq 12r_i$ and $\lambda_j^2 \leq 2^{\frac{2}{p}} \lambda_i^2 \leq 2\lambda_i^2$, we get

$$|\tau - t_i| \leq (48^2 + 2^2) \lambda_i^2 \Lambda^{-1} r_i^2 \leq \lambda_i^2 \Lambda^{-1} (50r_i)^2.$$

Therefore, $2G_j \subset 50KG_i$ and the proof is completed. \square

Finally, we show that the cardinality of \mathcal{I}_i , denoted by $|\mathcal{I}_i|$, is uniformly bounded.

Lemma 3.13. *There exists a constant $c = c(n, K)$ such that $|\mathcal{I}_i| \leq c$ for every $i \in \mathbb{N}$.*

Proof. Observe that $2U_j \subset 50K^2U_i$ for all $j \in \mathcal{I}_i$ by Lemma 3.11 and Lemma 3.12. Since the cylinders $\{\frac{1}{6K}U_j\}_{j \in \mathcal{I}_i}$ are disjoint, we obtain with Lemma 3.10 that

$$|50K^2U_i| \geq \left| \bigcup_{j \in \mathcal{I}_i} \frac{1}{6K}U_j \right| = \sum_{j \in \mathcal{I}_i} \left| \frac{1}{6K}U_j \right| \geq \sum_{j \in \mathcal{I}_i} c(n, K) |U_i| = c(n, K) |\mathcal{I}_i| |U_i|,$$

and thus $|\mathcal{I}_i| \leq c(n, K)$. \square

3.5. Partition of unity. To finish the proof of Proposition 3.2, we construct a partition of unity subordinate to the Whitney decomposition $\{2U_i\}_{i \in \mathbb{N}}$. For each $i \in \mathbb{N}$, there exists $\psi_i \in C_0^\infty(2U_i)$ such that

$$0 \leq \psi_i \leq 1, \quad \psi_i \equiv 1 \text{ in } U_i, \quad \|\nabla \psi_i\|_\infty \leq 2r_i^{-1}$$

and

$$\|\partial_t \psi_i\|_\infty \leq \begin{cases} 2\lambda_i^{p-2} r_i^{-2}, & \text{if } U_i = Q_i, \\ 2\Lambda \lambda_i^{-2} r_i^{-2}, & \text{if } U_i = G_i. \end{cases}$$

Since $E(\Lambda)^c \subset \bigcup_{i \in \mathbb{N}} 2U_i$ and $|\mathcal{I}_i|$ is finite for each $i \in \mathbb{N}$, the function

$$\omega_i(z) = \frac{\psi_i(z)}{\sum_{j \in \mathbb{N}} \psi_j(z)} = \frac{\psi_i(z)}{\sum_{j \in \mathcal{I}_i} \psi_j(z)}$$

is well-defined and satisfies

$$\omega_i \in C_0^\infty(2U_i), \quad 0 \leq \omega_i \leq 1 \text{ in } 2U_i, \quad \sum_{i \in \mathbb{N}} \omega_i(z) \equiv 1.$$

The family $\{\omega_i\}_{i \in \mathbb{N}}$ is a partition of unity subordinate to $\{2U_i\}_{i \in \mathbb{N}}$. It follows from Lemma 3.8, Lemma 3.9 and Lemma 3.13 that there exists $c = c(n, K)$ such that

$$\|\nabla \omega_i\|_\infty \leq cr_i^{-1} \quad \text{and} \quad \|\partial_t \omega_i\|_\infty \leq \begin{cases} c\lambda_i^{p-2} r_i^{-2}, & \text{if } U_i = Q_i, \\ c\Lambda \lambda_i^{-2} r_i^{-2}, & \text{if } U_i = G_i. \end{cases}$$

Moreover, for any $j \in \mathcal{I}_i$ we have

$$\|\nabla \omega_j\|_\infty \leq cr_i^{-1} \quad \text{and} \quad \|\partial_t \omega_j\|_\infty \leq \begin{cases} c\lambda_i^{p-2} r_i^{-2}, & \text{if } U_i = Q_i, \\ c\Lambda \lambda_i^{-2} r_i^{-2}, & \text{if } U_i = G_i. \end{cases}$$

As the proof of the above display is based on the comparability of the scaling factors of i and j , which is already proved in the proof of Lemma 3.8, we omit the details.

4. LIPSCHITZ TRUNCATION

In this section we discuss the Lipschitz truncation method which allows us to apply weak solutions to (2.3) as test functions in Definition 2.1. The main idea of the Lipschitz truncation technique is to keep a function unchanged in the good set $E(\Lambda)$, defined as in (3.4), and redefine it in the bad set $\mathbb{R}^{n+1} \setminus E(\Lambda)$ by applying the partition of unity in Proposition 3.2. The properties of the Lipschitz truncation are summarized in Proposition 4.11. For simplicity we have written the argument in this section with respect to Theorem 2.4, while the modifications to show Theorem 2.6 are explained in Section 6.1. In this section u is a weak solution to (2.3) and we consider the domain $Q_{R_2, S_2}(z_0)$ and R_1, S_1 as in the statement of Theorem 2.4.

4.1. Construction of the Lipschitz truncation. Let $f \in L^p(\mathbb{R}^{n+1})$ be

$$f = (|\nabla u| + |u - u_0| + |F|)\chi_{Q_{R_2, S_2}(z_0)}, \quad u_0 = (u)_{Q_{R_2, S_2}(z_0)},$$

where u and F are extended to zero outside $Q_{R_2, S_2}(z_0)$. By Definition 2.1 the assumption (3.3) is satisfied. With respect to f , let Λ , $E(\Lambda)$ and K be defined as in Section 3.1. There exists a Whitney covering $\{U_i\}_{i \in \mathbb{N}}$ as in Proposition 3.2

Let $0 < h_0 < \frac{S_2 - S_1}{4}$ be small enough and $\eta \in C_0^\infty(B_{R_2}(x_0))$ and $\zeta \in C_0^\infty(\ell_{S_2 - h_0}(t_0))$ be standard cutoff functions satisfying $0 \leq \eta \leq 1$, $0 \leq \zeta \leq 1$, $\eta \equiv 1$ in $B_{R_1}(x_0)$, $\zeta \equiv 1$ in $\ell_{S_1}(t_0)$ and

$$\|\nabla \eta\|_\infty \leq \frac{3}{R_2 - R_1}, \quad \|\partial_t \zeta\|_\infty \leq \frac{3}{S_2 - S_1}. \quad (4.1)$$

For $0 < h < h_0$, we would like to apply

$$v_h = [u - u_0]_h \eta \zeta$$

as a test function in the proof of the energy estimate. We define the Lipschitz truncation of v_h as

$$v_h^\Lambda(z) = v_h(z) - \sum_{i \in \mathbb{N}} (v_h(z) - v_h^i) \omega_i(z), \quad (4.2)$$

where $z \in \mathbb{R}^{n+1}$ and

$$v_h^i = \begin{cases} (v_h)_{2U_i}, & \text{if } 2U_i \subset Q_{R_2, S_2}(z_0), \\ 0, & \text{otherwise.} \end{cases}$$

As v_h vanishes outside $Q_{R_2, S_2}(z_0)$, the definition of v_h^i implies that $v_h^\Lambda \equiv 0$ in $Q_{R_2, S_2}(z_0)^c$. Note as well that each term on the right-hand side of (4.2) belongs to $W^{1,2}(\ell_{S_2 - h_0}(t_0); L^2(B_{R_2}(x_0), \mathbb{R}^N))$ and by Proposition 3.2 (xi) only finitely many of these terms are nonzero on a neighborhood of any point. Thus

$$v_h^\Lambda \in W_0^{1,2}(\ell_{S_2 - h_0}(t_0); L^2(B_{R_2}(x_0), \mathbb{R}^N)).$$

Similarly, we denote

$$v = (u - u_0) \eta \zeta \quad \text{and} \quad v^\Lambda(z) = v(z) - \sum_{i \in \mathbb{N}} (v - v^i) \omega_i, \quad (4.3)$$

where

$$v^i = \begin{cases} (v)_{2U_i}, & \text{if } 2U_i \subset Q_{R_2, S_2}(z_0), \\ 0, & \text{otherwise.} \end{cases} \quad (4.4)$$

Remark 4.1. If $z \in E(\Lambda)^c$, then $z \in 2U_i$ for some $i \in \mathbb{N}$ by Proposition 3.2 (i). By Proposition 3.2 (xii) we therefore have

$$v_h^\Lambda(z) = v_h(z) - v_h(z) \sum_{j \in \mathbb{N}} \omega_j + \sum_{j \in \mathbb{N}} v_h^j \omega_j(z) = \sum_{j \in \mathbb{N}} v_h^j \omega_j(z) = \sum_{j \in \mathcal{I}_i} v_h^j \omega_j(z).$$

On the other hand, if $z \in E(\Lambda)$, then $\omega_j(z) = 0$ for every $j \in \mathbb{N}$ by Proposition 3.2 (i). Thus

$$v_h^\Lambda(z) = \begin{cases} \sum_{j \in \mathcal{I}_i} v_h^j \omega_j(z), & \text{if } z \in 2U_i \text{ for some } i \in \mathbb{N}, \\ v_h(z), & \text{if } z \in E(\Lambda). \end{cases} \quad (4.5)$$

As the cardinality of \mathcal{I}_i is finite, this implies that for any $z \in E(\Lambda)^c$ there is a neighborhood where v_h^Λ is a finite linear combination of smooth functions and therefore $v_h^\Lambda \in C^\infty(E(\Lambda)^c)$. By the same argument without the Steklov average, we have $v^\Lambda \in C^\infty(E(\Lambda)^c)$.

4.2. Fundamental estimates. In this subsection we provide lemmas used in Section 4.3 to estimate terms containing v_h^Λ by Λ . To achieve the desired convergence of the Lipschitz truncation, we need estimates based on (4.7). However, (4.7) does not hold for $4K_i[U_i]_h$ as a might not be comparable in $[U_i]_h$. On the other hand, to estimate v_h^Λ by f we need to increase the domain of integration by h because of Lemma 2.2. Therefore, we show the Lipschitz regularity of v_h^Λ with crude estimates based on (4.6). This allows us to use v_h^Λ as a test function, and after taking the limit $h \rightarrow 0$ we can use for v^Λ the finer estimates based on (4.7).

Apart from estimating Λ , the following lemma is a standard consequence of the definition of $E(\Lambda)$ and the Whitney decomposition. Unless specified otherwise, in this section the constants c depend only on

$$data = n, N, p, q, \alpha, \nu, L, [a]_\alpha, R_1, R_2, S_1, S_2, K, \|a\|_{L^\infty(\mathbb{R}^{n+1})}.$$

Lemma 4.2. *Let $1 \leq s \leq d$. For any cylinder $Q \subset \mathbb{R}^{n+1}$ such that $Q \cap E(\Lambda) \neq \emptyset$, we have*

$$\iint_Q f^s dz \leq \Lambda^{\frac{s}{p}}. \quad (4.6)$$

Moreover, there exists a constant $c = c(data)$ such that

$$\iint_{4K_i U_i} f^s dz \leq c \lambda_i^s, \quad (4.7)$$

where $K_i U_i$ is defined in (3.8) and (3.10).

Proof. Let $w \in Q \cap E(\Lambda)$. By the definitions in (3.4) and (3.1), we have

$$\begin{aligned} \iint_Q f^s dz &\leq \left(\iint_Q f^d dz \right)^{\frac{s}{d}} \leq (M(f^d)(w))^{\frac{s}{d}} \\ &\leq \left(M(f^d + (af^q)^{\frac{d}{p}})(w) \right)^{\frac{s}{d}} \leq \Lambda^{\frac{s}{p}}, \end{aligned}$$

which proves the first claim. Since $4K_i U_i \cap E(\Lambda) \neq \emptyset$ by Proposition 3.2 (iv), it follows that

$$\iint_{4K_i U_i} f^s dz \leq \Lambda^{\frac{s}{p}}.$$

If $U_i = Q_i$, we have $a(z_i)\lambda_i^q \leq K^2\lambda_i^p$ and therefore $\Lambda^{\frac{s}{p}} \leq ((K^2 + 1)\lambda_i^p)^{\frac{s}{p}} = c\lambda_i^s$, which finishes the proof in the p -intrinsic case. On the other hand, if $U_i = G_i$, we have by Proposition 3.2 (x) that

$$[a(z_i)]^{\frac{d}{p}} \iint_{4K_i U_i} (f^q)^{\frac{d}{p}} dz = \iint_{4K_i U_i} (a(z_i)f^q)^{\frac{d}{p}} dz \leq 2^{\frac{d}{p}} \iint_{4K_i U_i} (a(z)f^q)^{\frac{d}{p}} dz.$$

Again, by Proposition 3.2 (iv) there exists $w \in 4K_i U_i \cap E(\Lambda)$ and we get

$$\iint_{4K_i U_i} (a(z)f^q)^{\frac{d}{p}} dz \leq M((af^q)^{\frac{d}{p}})(w) \leq M(f^d + (af^q)^{\frac{d}{p}})(w) \leq \Lambda^{\frac{d}{p}}.$$

Because $U_i = G_i$, we have $\lambda_i^p < a(z_i)\lambda_i^q$ and it follows that

$$\begin{aligned} a(z_i)^{\frac{d}{p}} \iint_{4K_i U_i} (f^q)^{\frac{d}{p}} dz &\leq c\Lambda^{\frac{d}{p}} = c(\lambda_i^p + a(z_i)\lambda_i^q)^{\frac{d}{p}} \\ &\leq c(a(z_i)\lambda_i^q)^{\frac{d}{p}} = ca(z_i)^{\frac{d}{p}}\lambda_i^{\frac{qd}{p}}. \end{aligned}$$

As $a(z_i) > 0$, this implies

$$\iint_{4K_i U_i} f^{\frac{qd}{p}} dz \leq c\lambda_i^{\frac{qd}{p}}$$

and we have

$$\iint_{4K_i U_i} f^s dz \leq \left(\iint_{4K_i U_i} f^{\frac{qd}{p}} dz \right)^{\frac{sp}{qd}} \leq c\lambda_i^s,$$

which completes the proof. \square

A key tool in the estimates for the Lipschitz truncation is the Poincaré type inequality in Lemma 4.5. To prove it, we use Lemma 4.2 and the following lemma which is a standard consequence of elliptic Poincaré inequality and the fact that $u - u_0$ is a weak solution to (2.3).

Lemma 4.3. *Let $Q = B_r \times \ell_s$ be a cylinder defined as in (2.2) and satisfying $B_r \subset B_{R_2}(x_0)$.*

(i) *There exists a constant $c = c(\text{data})$ such that*

$$\begin{aligned} \iint_Q |v_h - (v_h)_Q| dz &\leq csr^{-1} \iint_{[Q]_h} (f^{p-1} + a(z)f^{q-1}) dz \\ &\quad + cs \iint_{[Q]_h} f dz + cr \iint_{[Q]_h} f dz. \end{aligned} \quad (4.8)$$

(ii) *If in addition $\ell_s \cap \ell_{S_2}(t_0)^c \neq \emptyset$, then there exists a constant $c = c(\text{data})$ such that*

$$\begin{aligned} \iint_Q |v_h| dz &\leq csr^{-1} \iint_{[Q]_h} (f^{p-1} + a(z)f^{q-1}) dz \\ &\quad + cs \iint_{[Q]_h} f dz + cr \iint_{[Q]_h} f dz. \end{aligned} \quad (4.9)$$

Moreover, the above estimates hold with v_h and $[Q]_h$ replaced by v and Q .

Proof. For $t_1, t_2 \in \ell_s$, $t_1 \leq t_2$, let $\zeta_\delta \in W_0^{1,\infty}(\ell_s)$ be a piecewise linear cutoff function

$$\zeta_\delta(t) = \begin{cases} 0, & t \in (-\infty, t_1 - \delta), \\ 1 + \frac{t - t_1}{\delta}, & t \in [t_1 - \delta, t_1] \\ 1, & t \in (t_1, t_2), \\ 1 - \frac{t - t_2}{\delta}, & t \in [t_2, t_2 + \delta], \\ 0, & t \in (t_2 + \delta, \infty). \end{cases}$$

Moreover, let $\varphi \in C_0^\infty(B_r)$ be a nonnegative function satisfying

$$\int_{B_r} \varphi dx = 1, \quad \|\nabla \varphi\|_\infty \leq \frac{c(n)}{r}, \quad \|\varphi\|_\infty \leq c(n),$$

and let $\psi = \varphi \eta \zeta_\delta \in W_0^{1,\infty}(Q \cap Q_{R_2, S_2-h}(z_0))$, where η and ζ are defined in (4.1). Taking ψ as a test function in the Steklov averaged weak formulation of (2.3), we obtain

$$\iint_Q -[u - u_0]_h \cdot \varphi \eta \partial_t(\zeta \zeta_\delta) dz = \iint_Q [-A(\cdot, \nabla u) + |F|^{p-2} F + a|F|^{q-2} F]_h \cdot \nabla \psi dz.$$

After using the product rule of time derivative and rearranging terms, we get

$$\begin{aligned} \text{I} &= \left| \iint_Q -[u - u_0]_h \cdot \eta \varphi \zeta \partial_t(\zeta_\delta) dz \right| \leq \left| \iint_Q [u - u_0]_h \cdot \eta \varphi \zeta_\delta \partial_t \zeta dz \right| \\ &\quad + (L+1) \left| \iint_Q [|\nabla u|^{p-1} + a|\nabla u|^{q-1} + |F|^{p-1} + a|F|^{q-1}]_h \cdot \nabla \psi dz \right| \\ &\leq \left| \iint_Q [f]_h \eta \varphi \zeta_\delta \partial_t \zeta dz \right| + c \left| \iint_Q [f^{p-1} + af^{q-1}]_h |\nabla \psi| dz \right| = \text{II} + \text{III}. \end{aligned}$$

Lemma 2.2 implies that

$$\text{II} \leq \iint_Q [f]_h |\partial_t \zeta| \|\varphi\|_\infty dz \leq csr^n \iint_{[Q]_h} f dz.$$

Again by Lemma 2.2 and properties of the test functions, we have

$$\begin{aligned}
\text{III} &\leq c \iint_Q [f^{p-1} + af^{q-1}]_h |\varphi \nabla \eta + \eta \nabla \varphi| dz \\
&\leq c \iint_Q [f^{p-1} + af^{q-1}]_h (\varphi + \eta) (|\nabla \varphi| + |\nabla \eta|) dz \\
&\leq csr^n \left(\frac{1}{R_2 - R_1} + \frac{1}{r} \right) \iint_{[Q]_h} (f^{p-1} + a(z)f^{q-1}) dz.
\end{aligned}$$

Since $B_r \subset B_{R_2}(x_0)$, we have $r \leq R_2$ and therefore

$$\frac{1}{R_2 - R_1} + \frac{1}{r} = \frac{R_2}{R_2 - R_1} \frac{1}{R_2} + \frac{1}{r} \leq 2 \max \left\{ \frac{R_2}{R_2 - R_1}, 1 \right\} \frac{1}{r}.$$

We conclude that

$$\text{III} \leq csr^{n-1} \iint_{[Q]_h} (f^{p-1} + a(z)f^{q-1}) dz.$$

By the one-dimensional Lebesgue differentiation theorem we have

$$\begin{aligned}
\lim_{\delta \rightarrow 0^+} \text{I} &= \left| \lim_{\delta \rightarrow 0^+} \iint_Q [u - u_0]_h \eta \varphi \zeta \partial_t(\zeta_\delta) dz \right| \\
&= \left| \int_{B_r \times \{t_1\}} v_h \varphi dz - \int_{B_r \times \{t_2\}} v_h \varphi dz \right| \\
&= r^n |(v_h \varphi)_{B_r}(t_1) - (v_h \varphi)_{B_r}(t_2)|.
\end{aligned}$$

As the estimates of II and III are independent of δ , we conclude that

$$\begin{aligned}
\text{ess sup}_{t_1, t_2 \in \ell_s} |(v_h \varphi)_{B_r}(t_1) - (v_h \varphi)_{B_r}(t_2)| &\leq cs \iint_{[Q]_h} f dz \\
&+ csr^{-1} \iint_{[Q]_h} (f^{p-1} + a(z)f^{q-1}) dz.
\end{aligned} \tag{4.10}$$

To complete the proof we estimate the left-hand sides of (4.8) and (4.9) with the supremum above. We start with (4.8). Using the standard Poincaré inequality, we get

$$\begin{aligned}
\iint_Q |v_h - (v_h)_Q| dz &\leq \iint_Q |v_h - (v_h)_{B_r}| dz + \iint_Q |(v_h)_{B_r} - (v_h)_Q| dz \\
&\leq cr \iint_Q |\nabla v_h| dz + \iint_Q |(v_h)_{B_r} - (v_h)_Q| dz.
\end{aligned} \tag{4.11}$$

We estimate the first term with Lemma 2.2 to obtain

$$\iint_Q |\nabla v_h| dz = \iint_Q |[\nabla u]_h \eta \zeta + [u - u_0]_h \nabla \eta \zeta| dz \leq c \iint_{[Q]_h} f dz. \tag{4.12}$$

For the second term on the right-hand side of (4.11), we have

$$\begin{aligned}
\iint_Q |(v_h)_{B_r}(t) - (v_h)_Q| dz &= \int_{\ell_s} \left| \int_{\ell_s} (v_h)_{B_r}(t) - (v_h)_{B_r}(\tau) dt \right| d\tau \\
&\leq \int_{\ell_s} 2|(v_h)_{B_r}(t) - (v_h \varphi)_{B_r}(t)| dt + \text{ess sup}_{t, \tau \in \ell_s} |(v_h \varphi)_{B_r}(t) - (v_h \varphi)_{B_r}(\tau)|.
\end{aligned}$$

The second term above is the same as in (4.10). It remains to estimate the first term. We use the fact that $(\varphi)_{B_r} = 1$, the standard Poincaré inequality and (4.12) to get

$$\begin{aligned}
\int_{\ell_s} |(v_h)_{B_r}(t) - (v_h\varphi)_{B_r}(t)| dt &= \int_{\ell_s} \left| \int_{B_r \times \{t\}} v_h \varphi dx - (v_h)_{B_r}(t) \int_{B_r} \varphi dx \right| dt \\
&= \int_{\ell_s} \left| \int_{B_r \times \{t\}} \varphi (v_h - (v_h)_{B_r}) dx \right| dt \\
&\leq \|\varphi\|_\infty \int_{\ell_s} \int_{B_r} |v_h - (v_h)_{B_r}| dx dt \\
&\leq cr \iint_Q |\nabla v_h| dz \leq c \iint_{[Q]_h} f dz.
\end{aligned} \tag{4.13}$$

This completes the proof of (4.8).

To prove (4.9) we argue in a similar way. As in (4.11), we have

$$\begin{aligned}
\iint_Q |v_h| dz &\leq \iint_Q |v_h - (v_h)_{B_r}| dz + \int_{\ell_s} |(v_h)_{B_r}| dz \\
&\leq cr \iint_Q |\nabla v_h| dz + \int_{\ell_s} |(v_h)_{B_r}| dz,
\end{aligned}$$

where the first term was already estimated in (4.12). For the second term we have

$$\int_{\ell_s} |(v_h)_{B_r}| dz \leq \int_{\ell_s} |(v_h)_{B_r} - (v_h\varphi)_{B_r}| dt + \int_{\ell_s} |(v_h\varphi)_{B_r}| dt.$$

The first term above was estimated in (4.13). For the second term we can choose $t_2 \in \ell_{S_2}(t_0)^c$ in (4.10) because of the assumption $\ell_s \cap \ell_{S_2}(t_0)^c \neq \emptyset$. Since ζ vanishes outside $\ell_{S_2}(t_0)$, we have

$$\int_{\ell_s} |(v_h\varphi)_{B_r}| dt \leq \operatorname{ess\,sup}_{t \in \ell_s} |(v_h\varphi)_{B_r}(t)| \leq \operatorname{ess\,sup}_{t_1, t_2 \in \ell_s} |(v_h\varphi)_{B_r}(t_1) - (v_h\varphi)_{B_r}(t_2)|,$$

which can be estimated by (4.10). This completes the proof of (4.9). Proofs for the corresponding statements without the Steklov average are analogous and we omit the details. \square

To prove Lemma 4.5 near the lateral boundary, we use the following boundary version of the Poincaré inequality. For a proof we refer to [18, Example 6.18 and Theorem 6.22].

Lemma 4.4. *Let $B_\rho(x_0) \subset \mathbb{R}^n$ and $B_r \subset \mathbb{R}^n$ be balls satisfying $B_r \cap B_\rho(x_0)^c \neq \emptyset$. Assume that $v \in W_0^{1,s}(B_\rho(x_0))$ with $1 < s < \infty$. Moreover, let $1 \leq \sigma \leq \frac{ns}{n-s}$ for $1 < s < n$ and $1 \leq \sigma < \infty$ for $n \leq s < \infty$. Then there exists a constant $c = c(n, s, \sigma)$ such that*

$$\left(\int_{B_{4r}} |v|^\sigma dx \right)^{\frac{1}{\sigma}} \leq cr \left(\int_{B_{4r}} |\nabla v|^s dx \right)^{\frac{1}{s}}.$$

Now we are ready to show the Poincaré type inequality.

Lemma 4.5. *The following holds for $K_i U_i$ defined in (3.8) and (3.10):*

(i) *If $K_i U_i \subset Q_{R_2, S_2}(z_0)$, then*

$$\iint_{K_i U_i} |v_h - (v_h)_{K_i U_i}| dz \leq c(\text{data}, \Lambda) r_i \tag{4.14}$$

and

$$\iint_{K_i U_i} |v - (v)_{K_i U_i}| dz \leq c(\text{data}) r_i \lambda_i. \tag{4.15}$$

(ii) *If $K_i U_i \not\subset Q_{R_2, S_2}(z_0)$, then*

$$\iint_{K_i U_i} |v_h| dz \leq c(\text{data}, \Lambda) r_i \tag{4.16}$$

and

$$\iint_{K_i U_i} |v| dz \leq c(\text{data}) r_i \lambda_i. \quad (4.17)$$

Proof. We start by proving (4.15). By the assumption $K_i U_i \subset Q_{R_2, S_2}(z_0)$, we may apply Lemma 4.3 with $Q = K_i U_i$. Assume first $U_i = Q_i$. Then Lemma 4.3 gives

$$\iint_{K_i U_i} |v - (v)_{K_i U_i}| dz \leq c \lambda_i^{2-p} r_i \iint_{K_i U_i} (f^{p-1} + a(z) f^{q-1}) dz + c r_i \iint_{K_i U_i} f dz, \quad (4.18)$$

where we already estimated the third term in (4.8) with the fourth one as $r_i \leq R_2$ and therefore

$$r_i^2 \lambda_i^{2-p} \iint_{K_i U_i} f dz \leq R_2 r_i \iint_{K_i U_i} f dz. \quad (4.19)$$

To estimate the term containing $a(z)$, observe that (2.5) implies

$$\iint_{K_i U_i} a(z) f^{q-1} dz \leq a(z_i) \iint_{K_i U_i} f^{q-1} dz + [a]_\alpha (K_i r_i)^\alpha \iint_{K_i U_i} f^{q-1} dz. \quad (4.20)$$

As $U_i = Q_i$, we have $a(z_i) \leq K^2 \lambda_i^{p-q}$ and Lemma 4.2 gives

$$a(z_i) \iint_{K_i U_i} f^{q-1} dz \leq c \lambda_i^{p-q} \iint_{K_i U_i} f^{q-1} dz \leq c \lambda_i^{p-1}.$$

To estimate the other term in (4.20), we apply Lemma 4.2 and Proposition 3.2 (ix) to get

$$r_i^\alpha \iint_{K_i U_i} f^{q-1} dz \leq r_i^\alpha \left(\iint_{K_i U_i} f^d dz \right)^{\frac{q-1}{d}} \leq r_i^\alpha \lambda_i^{q-1} \leq c \lambda_i^{p-1}.$$

Inserting these estimates and Lemma 4.2 to (4.18), we have the desired estimate

$$\iint_{K_i U_i} |v - (v)_{K_i U_i}| dz \leq c r_i \lambda_i.$$

In case $U_i = G_i$, Lemma 4.3 gives

$$\iint_{K_i U_i} |v - (v)_{K_i U_i}| dz \leq c \frac{\lambda_i^2 r_i}{\Lambda} \iint_{K_i U_i} (f^{p-1} + a(z) f^{q-1}) dz + c r_i \iint_{K_i U_i} f dz,$$

where we again used (4.19). The second term is again estimated by Lemma 4.2. To estimate the first term, we apply Proposition 3.2 (x) to cancel $a(z)$, which along with Lemma 4.2 gives

$$\begin{aligned} & \frac{\lambda_i^2 r_i}{\lambda_i^p + a(z_i) \lambda_i^q} \iint_{K_i G_i} (f^{p-1} + a(z) f^{q-1}) dz \\ & \leq \frac{\lambda_i^2 r_i}{\lambda_i^p} \iint_{K_i G_i} f^{p-1} dz + \frac{\lambda_i^2 r_i}{a(z_i) \lambda_i^q} \iint_{K_i G_i} a(z) f^{q-1} dz \\ & \leq c r_i \lambda_i + \frac{2 \lambda_i^2 r_i}{\lambda_i^q} \iint_{K_i G_i} f^{q-1} dz \leq c r_i \lambda_i. \end{aligned}$$

This finishes the proof of (4.15).

To prove (4.17), we assume $K_i U_i \not\subset Q_{R_2, S_2}(z_0)$. If in addition $K_i B_i \subset B_{R_2}(x_0)$, then $K_i I_i \cap \ell_{S_2}(t_0)^c \neq \emptyset$ and we may apply (4.9). Since the right-hand sides of (4.8) and (4.9) are the same, (4.17) follows from the proof of (4.15). On the other hand, if $K_i B_i \cap B_{R_2}^c \neq \emptyset$, then we may apply Lemma 4.4 with $\sigma = 1$ and $s = d$, since $v(\cdot, t) \in W_0^{1,d}(B_{R_2}(x_0), \mathbb{R}^N)$. It follows that

$$\iint_{K_i U_i} |v| dz \leq c r_i \left(\iint_{4K_i U_i} |\nabla v|^d dz \right)^{\frac{1}{d}},$$

and as $|\nabla v| \leq c f$, we have by Lemma 4.2 that

$$\left(\iint_{4K_i U_i} |\nabla v|^d dz \right)^{\frac{1}{d}} \leq \left(\iint_{4K_i U_i} f^d dz \right)^{\frac{1}{d}} \leq c \lambda_i.$$

This gives the desired estimate and we have proven (4.17).

Finally, we show the corresponding estimates with Steklov averages, that is, (4.14) and (4.16). If $K_i U_i \subset Q_{R_2, S_2}(z_0)$, then Lemma 4.3 gives

$$\begin{aligned} \iint_{K_i U_i} |v_h - (v_h)_{K_i U_i}| dz &\leq c |I_i| r_i^{-1} \iint_{[K_i U_i]_h} (f^{p-1} + a(z) f^{q-1}) dz \\ &\quad + c |I_i| \iint_{[K_i U_i]_h} f dz + c r_i \iint_{[K_i U_i]_h} f dz. \end{aligned}$$

Note that $|I_i| \leq r_i^2 \leq r_i R_2$ and $a(z) \leq \|a\|_\infty$. Using Lemma 4.2 with $s = d$ gives

$$\iint_{K_i U_i} |v_h - (v_h)_{K_i U_i}| dz \leq c r_i \max \left\{ 1, \iint_{[K_i U_i]_h} f^d dz \right\}^{\frac{q-1}{d}} \leq c r_i \Lambda^{\frac{q-1}{p}},$$

which finishes the proof of (4.14).

We get (4.16) in the case $K_i B_i \subset B_{R_2}(x_0)$ and $K_i I_i \not\subset \ell_{S_2}(t_0)$, from (4.9) and the proof of (4.14). On the other hand, if $K_i B_i \cap B_{R_2}(x_0)^c \neq \emptyset$, we apply Lemma 4.4 to get

$$\iint_{K_i U_i} |v_h| dz \leq c r_i \left(\iint_{4K_i U_i} |\nabla v_h|^d dz \right)^{\frac{1}{d}}.$$

By Lemma 2.2 and Lemma 4.2 we have

$$\begin{aligned} \left(\iint_{4K_i U_i} |\nabla v_h|^d dz \right)^{\frac{1}{d}} &\leq c \left(\iint_{4K_i U_i} [|\nabla v|^d]_h dz \right)^{\frac{1}{d}} \leq c \left(\iint_{[4K_i U_i]_h} |\nabla v|^d dz \right)^{\frac{1}{d}} \\ &\leq c \left(\iint_{[4K_i U_i]_h} f^d dz \right)^{\frac{1}{d}} \leq c \Lambda^{\frac{1}{p}}, \end{aligned}$$

and combining the estimates gives (4.17). This completes the proof. \square

Remark 4.6. *It follows from Lemma 4.5 that*

$$\iint_{2U_i} |v - v^i| dz \leq c(\text{data}) r_i \lambda_i. \quad (4.21)$$

Indeed, if $K_i U_i \subset Q_{R_2, S_2}(z_0)$, we have by (4.15) that

$$\begin{aligned} \iint_{2U_i} |v - v^i| dz &\leq \iint_{2U_i} |v - (v)_{K_i U_i}| dz + \iint_{2U_i} |(v)_{K_i U_i} - (v)_{2U_i}| dz \\ &\leq c \iint_{K_i U_i} |v - (v)_{K_i U_i}| dz \leq c r_i \lambda_i. \end{aligned}$$

On the other hand, if $K_i U_i \not\subset Q_{R_2, S_2}(z_0)$, we apply (4.17) to have

$$\iint_{2U_i} |v - v^i| dz \leq 2 \iint_{2U_i} |v| dz \leq c \iint_{K_i U_i} |v| dz \leq c r_i \lambda_i.$$

4.3. Properties of the Lipschitz truncation. In the following lemmas we provide pointwise estimates for v_h^Λ , v^Λ and their gradients. They allow us to show the Lipschitz regularity of v_h^Λ in Lemma 4.10 and the estimates in Proposition 4.11.

Lemma 4.7. *We have $|v_h^\Lambda(z)| \leq c(\text{data}, \Lambda)$ and $|v^\Lambda(z)| \leq c(\text{data}) \lambda_i$ for every $z \in U_i$.*

Proof. By the definition of v^j and Lemma 4.2, we have for every $j \in \mathcal{I}_i$ that

$$|v^j| \leq \iint_{2U_j} v dz \leq c \iint_{K_i U_i} f dz \leq c \lambda_i. \quad (4.22)$$

Since $v^\Lambda(z) = \sum_{j \in \mathcal{I}_i} v^j \omega_j(z)$ by (4.5), we get from the above estimate and Proposition 3.2 (xi) that

$$|v^\Lambda(z)| \leq \left| \sum_{j \in \mathcal{I}_i} v^j \omega_j(z) \right| \leq \sum_{j \in \mathcal{I}_i} |v^j| |\omega_j(z)| \leq c |v^j| \leq c \lambda_i.$$

The claim with the Steklov averages is proved in the same way, the only difference is that we apply Lemma 2.2 and (4.6) in (4.22). \square

The next lemma is an auxiliary tool to show Lemma 4.9.

Lemma 4.8. *We have $|v_h^i - v_h^j| \leq c(\text{data}, \Lambda)r_i$ and $|v^i - v^j| \leq c(\text{data})r_i\lambda_i$ for every $i \in \mathbb{N}$ and $j \in \mathcal{I}_i$.*

Proof. We only prove the claim without the Steklov averages since the corresponding claim with the Steklov averages is similar. First consider the case $K_i U_i, K_j U_j \subset Q_{R_2, S_2}(z_0)$. Proposition 3.2 (vii) and Proposition 3.2 (viii) imply that

$$\begin{aligned} |v^i - v^j| &\leq |v^i - (v)_{K_i U_i}| + |(v)_{K_i U_i} - v^j| \\ &= \left| \iint_{2U_i} v - (v)_{K_i U_i} dz \right| + \left| \iint_{2U_j} v - (v)_{K_i U_i} dz \right| \\ &\leq c \iint_{K_i U_i} |v - (v)_{K_i U_i}| dz \leq cr_i \lambda, \end{aligned}$$

where the last inequality follows from Lemma 4.5.

On the other hand, if $K_i U_i \not\subset Q_{R_2, S_2}(z_0)$, we apply Proposition 3.2 (vii) and Proposition 3.2 (viii) and (4.17) to have

$$|v^i - v^j| \leq \iint_{2U_i} |v| dz + \iint_{2U_j} |v| dz \leq c \iint_{K_i U_i} |v| dz \leq cr_i \lambda_i.$$

Since r_i, r_j and λ_i, λ_j are comparable by Proposition 3.2 (v) and Proposition 3.2 (vi), the case $K_j U_j \not\subset Q_{R_2, S_2}(z_0)$ is proven similarly and we are done. \square

Lemma 4.9. *For any $z \in U_i$, we have*

$$|\nabla v_h^\Lambda(z)| \leq c(\text{data}, \Lambda) \quad \text{and} \quad |\nabla v^\Lambda(z)| \leq c(\text{data})\lambda_i. \quad (4.23)$$

Moreover, we have

$$|\partial_t v_h^\Lambda(z)| \leq c(\text{data}, \Lambda)r_i^{-1}, \quad (4.24)$$

and

$$|\partial_t v^\Lambda(z)| \leq \begin{cases} c(\text{data})r_i^{-1}\lambda_i^{p-1}, & \text{if } U_i = Q_i, \\ c(\text{data})r_i^{-1}\lambda_i^{-1}\Lambda, & \text{if } U_i = G_i. \end{cases} \quad (4.25)$$

Proof. We start with the proof of the second inequality in (4.23). By (4.5) we have

$$\nabla v^\Lambda(z) = \nabla \left(\sum_{j \in \mathcal{I}_i} v^j \omega_j(z) \right) = \sum_{j \in \mathcal{I}_i} v^j \nabla \omega_j(z).$$

Since

$$0 = \nabla \left(\sum_{j \in \mathbb{N}} \omega_j(z) \right) = \nabla \left(\sum_{j \in \mathcal{I}_i} \omega_j(z) \right) = \sum_{j \in \mathcal{I}_i} \nabla \omega_j$$

by Proposition 3.2 (xii), we get

$$\nabla v^\Lambda(z) = \sum_{j \in \mathcal{I}_i} v^j \nabla \omega_j(z) = \sum_{j \in \mathcal{I}_i} v^j \nabla \omega_j(z) - v^i \sum_{j \in \mathcal{I}_i} \nabla \omega_j(z) = \sum_{j \in \mathcal{I}_i} (v^j - v^i) \nabla \omega_j(z).$$

Therefore Lemma 4.8 and Proposition 3.2 (xiii), (v), (xi) give

$$|\nabla v^\Lambda(z)| \leq \sum_{j \in \mathcal{I}_i} |v^j - v^i| |\nabla \omega_j| \leq c \sum_{j \in \mathcal{I}_i} r_j^{-1} r_i \lambda_i \leq c \lambda_i.$$

Exactly the same argument with v_h^Λ instead of v^Λ proves the first inequality in (4.23).

To show (4.24), we have by the same argument as above that

$$\partial_t v_h^\Lambda(z) = \sum_{j \in \mathcal{I}_i} v_h^j \partial_t \omega_j(z) = \sum_{j \in \mathcal{I}_i} (v_h^j - v_h^i) \partial_t \omega_j(z).$$

Again, Lemma 4.8 and Proposition 3.2 (xiv), (v), (xi) give

$$|\partial_t v_h^\Lambda(z)| \leq \sum_{j \in \mathcal{I}_i} |v_h^j - v_h^i| |\partial_t \omega_j| \leq c(\text{data}, \Lambda) \sum_{j \in \mathcal{I}_i} r_j |\partial_t \omega_j| \leq c(\text{data}, \Lambda) r_i^{-1}.$$

The same argument with v^Λ instead of v_h^Λ shows (4.25) and the proof is completed. \square

We are ready to show that v_h^Λ is Lipschitz continuous with respect to the metric

$$d_{\lambda^p}(z, w) = \max \left\{ |x - y|, \sqrt{\lambda^{p-2}|t - s|} \right\},$$

where $z, w \in Q_{R_2, S_2}(z_0)$ with $z = (x, t)$ and $w = (y, s)$, and λ is chosen to satisfy $\lambda^p + \|a\|_\infty \lambda^q = \Lambda$.

Lemma 4.10. *There exists a constant $c_\Lambda = c(\text{data}, \Lambda)$ such that*

$$|v_h^\Lambda(z) - v_h^\Lambda(w)| \leq c_\Lambda d_{\lambda^p}(z, w)$$

for every $z, w \in \mathbb{R}^{n+1}$.

Proof. Observe that a cylinder $Q_l^\lambda(w)$, as defined in (2.1), is a ball of radius l associated with the metric d_{λ^p} . Since $\lambda_i \geq \lambda$ and $p \geq 2$, it is clear that

$$d_{\lambda^p}(\cdot, \cdot) \leq d_i(\cdot, \cdot), \quad (4.26)$$

for all $i \in \mathbb{N}$. Therefore, $U_i \subset Q_{r_i}^\lambda(z_i)$ for any $i \in \mathbb{N}$. By the Campanato characterization of Lipschitz continuity, it is enough to show that there exists a constant $c_\Lambda = c(\text{data}, \Lambda)$, such that

$$\iint_{Q_l^\lambda(w)} \frac{|v_h^\Lambda - (v_h^\Lambda)_{Q_l^\lambda(w)}|}{l} dz \leq c_\Lambda \quad (4.27)$$

for all $Q_l^\lambda(w) \subset \mathbb{R}^{n+1}$. We fix $Q_l^\lambda(w)$ and prove the above estimate, first in the case when $2Q_l^\lambda(w)$ is inside the bad set, then when it is touching the good set.

Case 1: $2Q_l^\lambda(w) \subset E(\Lambda)^c$. Let $z \in Q_l^\lambda(w)$ and let $i \in \mathbb{N}$ be such that $z \in U_i$. Note that such i exists by Proposition 3.2(i). Triangle inequality and (4.26) give

$$l \leq d_{\lambda^p}(z, E(\Lambda)) \leq d_{\lambda^p}(z, z_i) + d_{\lambda^p}(z_i, E(\Lambda)) \leq d_i(z, z_i) + d_i(z_i, E(\Lambda)),$$

where z_i is the centre of U_i . Therefore, Proposition 3.2(iv) and the fact that U_i is a ball of radius r_i with respect to the metric $d_i(\cdot, \cdot)$ implies that

$$l \leq d_i(z, z_i) + d_i(z_i, E(\Lambda)) \leq r_i + 5r_i = 6r_i.$$

Combining the above inequality with Lemma 4.9, we obtain the uniform estimate

$$|\partial_t v_h^\Lambda(z)| \leq c(\text{data}, \Lambda) r_i^{-1} \leq c(\text{data}, \Lambda) l^{-1} \quad (4.28)$$

for all $z \in Q_l^\lambda(w)$.

To show (4.27), let $z_1, z_2 \in Q_l^\lambda(w)$ with $z_1 = (x_1, t_1)$ and $z_2 = (x_2, t_2)$. Since $v_h^\Lambda \in C^\infty(E(\Lambda)^c, \mathbb{R}^N)$ and $Q_l^\lambda(w) \subset E(\Lambda)^c$, by the mean value theorem we obtain

$$\begin{aligned} |v_h^\Lambda(z_1) - v_h^\Lambda(z_2)| &\leq |v_h^\Lambda(x_1, t_1) - v_h^\Lambda(x_2, t_1)| + |v_h^\Lambda(x_2, t_1) - v_h^\Lambda(x_2, t_2)| \\ &\leq cl \sup_{z \in Q_l^\lambda(w)} |\nabla v_h^\Lambda(z)| + c\lambda^{2-p}l^2 \sup_{z \in Q_l^\lambda(w)} |\partial_t v_h^\Lambda(z)| \leq c(\text{data}, \Lambda)l, \end{aligned}$$

where the last inequality follows from (4.23) and (4.28). The estimate above gives

$$\iint_{Q_l^\lambda(w)} \frac{|v_h^\Lambda - (v_h^\Lambda)_{Q_l^\lambda(w)}|}{l} dz \leq \iint_{Q_l^\lambda(w)} \iint_{Q_l^\lambda(w)} \frac{|v_h^\Lambda(z_1) - v_h^\Lambda(z_2)|}{l} dz_2 dz_1 \leq c(\text{data}, \Lambda).$$

This proves (4.27) in the case $2Q_l^\lambda(w) \subset E(\Lambda)^c$.

Case 2: $2Q_l^\lambda(w) \cap E(\Lambda) \neq \emptyset$ and $Q_l^\lambda(w) \cap E(\Lambda)^c = \emptyset$. In this case $v_h^\Lambda = v_h$ and therefore

$$\iint_{Q_l^\lambda(w)} \frac{|v_h^\Lambda - (v_h^\Lambda)_{Q_l^\lambda(w)}|}{l} dz = \iint_{Q_l^\lambda(w)} \frac{|v_h - (v_h)_{Q_l^\lambda(w)}|}{l} dz. \quad (4.29)$$

We denote $w = (y, s)$. If $B_l(y) \subset B_{R_2}(x_0)$, then $l \leq R_2$ holds and we apply Lemma 4.3 and Lemma 4.2 to get

$$\iint_{Q_l^\lambda(w)} \frac{|v_h - (v_h)_{Q_l^\lambda(w)}|}{l} dz \leq c(\text{data}, \Lambda)(\lambda^{2-p} + \lambda^{2-p}l + 1) \leq c(\text{data}, \Lambda).$$

On the other hand, if $B_l(y) \not\subset B_{R_2}(x_0)$, we use Lemma 4.4 with $\sigma = 1$ and $s = d$ to get

$$\iint_{Q_l^\lambda(w)} \frac{|v_h - (v_h)_{Q_l^\lambda(w)}|}{l} dz \leq 2 \iint_{Q_l^\lambda(w)} \frac{|v_h|}{l} dz \leq c \iint_{4Q_l^\lambda(w)} |\nabla v_h| dz.$$

Recalling that in this case $2Q_l^\lambda(w) \cap E(\Lambda) \neq \emptyset$, we conclude with Lemma 4.2 that

$$\iint_{4Q_l^\lambda(w)} |\nabla v_h| dz \leq c \iint_{[4Q_l^\lambda(w)]_h} f dz \leq c(\text{data}, \Lambda),$$

which finishes the proof of (4.27) in the case $2Q_l^\lambda(w) \cap E(\Lambda) \neq \emptyset$ and $Q_l^\lambda(w) \cap E(\Lambda)^c = \emptyset$.

Case 3: $2Q_l^\lambda(w) \cap E(\Lambda) \neq \emptyset$ and $Q_l^\lambda(w) \cap E(\Lambda)^c \neq \emptyset$. To estimate in the region where $Q_l^\lambda(w)$ touches $E(\Lambda)^c$, we define the index set

$$P = \{i \in \mathbb{N} : Q_l^\lambda(w) \cap 2U_i \neq \emptyset\}.$$

We claim that the radii r_i are bounded uniformly by l when $i \in P$. Let $i \in P$, $w_1 \in Q_l^\lambda(w) \cap 2U_i$ and $w_2 \in 2Q_l^\lambda(w) \cap E(\Lambda)$ with $w_1 = (y_1, s_1)$ and $w_2 = (y_2, s_2)$. By Proposition 3.2 (iii) and $w_1 \in 2U_i$, we have

$$3r_i \leq d_i(U_i, E(\Lambda)) \leq d_i(z_i, w_2) \leq d_i(z_i, w_1) + d_i(w_1, w_2) \leq 2r_i + d_i(w_1, w_2),$$

and therefore $r_i \leq d_i(w_1, w_2)$. Furthermore, since $\lambda \leq \lambda_i$ we have

$$\begin{aligned} d_i(w_1, w_2) &\leq \max \left\{ |y_1 - y_2|, \sqrt{\Lambda \lambda_i^{-2} |s_1 - s_2|} \right\} \\ &\leq \Lambda^{\frac{1}{2}} \max \left\{ |y_1 - y_2|, \sqrt{\lambda^{p-2} |s_1 - s_2|} \right\} \\ &= \Lambda^{\frac{1}{2}} d_{\lambda^p}(w_1, w_2) \leq 4\Lambda^{\frac{1}{2}} l, \end{aligned}$$

where the last inequality follows from the fact that $w_1, w_2 \in 2Q_l^\lambda(w)$. We conclude that $r_i \leq c(\Lambda)l$.

To prove (4.27) we add and subtract twice to obtain

$$\begin{aligned} \iint_{Q_l^\lambda(w)} \frac{|v_h^\Lambda - (v_h^\Lambda)_{Q_l^\lambda(w)}|}{l} dz &\leq 2 \iint_{Q_l^\lambda(w)} \frac{|v_h^\Lambda - (v_h)_{Q_l^\lambda(w)}|}{l} dz \\ &\leq 2 \iint_{Q_l^\lambda(w)} \frac{|v_h^\Lambda - v_h|}{l} dz + 2 \iint_{Q_l^\lambda(w)} \frac{|v_h - (v_h)_{Q_l^\lambda(w)}|}{l} dz. \end{aligned} \tag{4.30}$$

The second term on the right-hand side is the same as in (4.29) and can be estimated in the same way. To estimate the first term, observe from (4.2) and the fact that w_i is supported in $2U_i$, that

$$\begin{aligned} \iint_{Q_l^\lambda(w)} \frac{|v_h^\Lambda - v_h|}{l} dz &= \iint_{Q_l^\lambda(w)} \frac{|\sum_{i \in P} (v_h - v_h^i) \omega_i|}{l} dz \\ &\leq \iint_{Q_l^\lambda(w)} \sum_{i \in P} \frac{|v_h - v_h^i| \omega_i}{l} dz \\ &= \frac{1}{|Q_l^\lambda(w)|} \sum_{i \in P} \iint_{Q_l^\lambda(w) \cap 2U_i} \frac{|v_h - v_h^i| \omega_i}{l} dz. \end{aligned}$$

Since $\omega_i \leq 1$ and $r_i \leq c(\Lambda)l$, we have

$$\frac{1}{|Q_l^\lambda(w)|} \sum_{i \in P} \iint_{Q_l^\lambda(w) \cap 2U_i} \frac{|v_h - v_h^i| \omega_i}{l} dz \leq \frac{c(\Lambda)}{|Q_l^\lambda(w)|} \sum_{i \in P} \iint_{2U_i} \frac{|v_h - v_h^i|}{r_i} dz,$$

and combining the previous inequalities with (4.21) gives

$$\iint_{Q_l^\lambda(w)} \frac{|v_h^\Lambda - v_h|}{l} dz \leq \frac{c(\text{data}, \Lambda)}{|Q_l^\lambda(w)|} \sum_{i \in P} |U_i|.$$

Since $r_i \leq c(\Lambda)l$, we have $U_i \subset c(\text{data}, \Lambda)Q_l^\lambda(w)$ for every $i \in P$. From this and Proposition 3.2 (ii), we conclude that

$$\iint_{Q_l^\lambda(w)} \frac{|v_h^\Lambda - v_h|}{l} dz \leq \frac{c(\text{data}, \Lambda)}{|Q_l^\lambda(w)|} \sum_{i \in P} |\frac{1}{6K}U_i| \leq c(\text{data}, \Lambda).$$

Combining this estimate with (4.30), we have (4.27) in the final case and the proof is completed. \square

In the following proposition we collect the properties of Lipschitz truncation that will be used in the proof of Theorem 2.4.

Proposition 4.11. *Let $E(\Lambda)$ be as in (3.4). There exist functions $\{v_h^\Lambda\}_{h>0}$ and a function v^Λ satisfying the following properties:*

- (i) $v_h^\Lambda \in W_0^{1,2}(\ell_{S_2-h_0}(t_0); L^2(B_{R_2}(x_0), \mathbb{R}^N)) \cap L^\infty(\ell_{S_2-h_0}(t_0); W_0^{1,\infty}(B_{R_2}(x_0), \mathbb{R}^N))$.
- (ii) $v^\Lambda \in L^\infty(\ell_{S_2}(t_0); W_0^{1,\infty}(B_{R_2}(x_0), \mathbb{R}^N))$.
- (iii) $v_h^\Lambda = v_h$, $v^\Lambda = v$, $\nabla v_h^\Lambda = \nabla v_h$, $\nabla v^\Lambda = \nabla v$ a.e. in $E(\Lambda)$.
- (iv) $v_h^\Lambda \rightarrow v^\Lambda$ in $L^\infty(Q_{R_2, S_2}(z_0), \mathbb{R}^N)$ as $h \rightarrow 0^+$, taking a subsequence if necessary.
- (v) $\nabla v_h^\Lambda \rightarrow \nabla v^\Lambda$ and $\partial_t v_h^\Lambda \rightarrow \partial_t v^\Lambda$ a.e. in $E(\Lambda)^c$ as $h \rightarrow 0^+$.

Moreover, there exist constants $c = c(\text{data})$ and $c_\Lambda = c_\Lambda(\text{data}, \Lambda)$ such that

- (vi) $\iint_{E(\Lambda)^c} |v - v^\Lambda| |\partial_t v^\Lambda| dz \leq c_\Lambda |E(\Lambda)^c|$,
- (vii) $\iint_{E(\Lambda)^c} |v_h - v_h^\Lambda| |\partial_t v_h^\Lambda| dz \leq c_\Lambda |E(\Lambda)^c|$,
- (viii) $H(z, |v^\Lambda(z)|) + H(z, |\nabla v^\Lambda(z)|) \leq c_\Lambda$ for a.e. $z \in \mathbb{R}^{n+1}$,
- (ix) $H(z, |v_h^\Lambda(z)|) + H(z, |\nabla v_h^\Lambda(z)|) \leq c_\Lambda$ for a.e. $z \in \mathbb{R}^{n+1}$.

Proof. We stated (i) already at the beginning of this section, while (ii) is deduced from Lemma 4.10. From the construction $v_h^\Lambda = v_h$ and $v^\Lambda = v$ in $E(\Lambda)$. Therefore, as $v_h^\Lambda, v_h, v^\Lambda, v$ belong to $L^p(\ell_{S_2}(t_0); W_0^{1,p}(B_{R_2}(z_0), \mathbb{R}^N))$, we also have $\nabla v_h^\Lambda = \nabla v_h$ and $\nabla v^\Lambda = \nabla v$ a.e. in $E(\Lambda)$, which proves (iii).

Next we prove (iv). Note that $\{v_h^\Lambda\}_{h>0}$ is equicontinuous and uniformly bounded by Lemma 4.10. Therefore, it follows from the Arzela–Ascoli theorem that, by passing to a subsequence if necessary, there exists w such that v_h^Λ converges to w in $L^\infty(Q_{R_2, S_2}(z_0), \mathbb{R}^N)$. On the other hand, v_h^Λ converges to v^Λ a.e. as $h \rightarrow 0^+$, which means $w = v^\Lambda$ a.e. This shows that $v_h^\Lambda \rightarrow v^\Lambda$ in $L^\infty(Q_{R_2, S_2}(z_0), \mathbb{R}^N)$ as $h \rightarrow 0^+$.

If $z \in E(\Lambda)^c$, then there exists $i \in \mathbb{N}$ such that $z \in U_i$. It follows from (4.5) that

$$\nabla v_h^\Lambda(z) = \sum_{j \in \mathcal{I}_i} v_h^j \nabla \omega_j(z), \quad \partial_t v_h^\Lambda(z) = \sum_{j \in \mathcal{I}_i} v_h^j \partial_t \omega_j(z).$$

By the properties of the Steklov average we have $v_h^j \rightarrow v^j$ as $h \rightarrow 0^+$, which implies the a.e. convergence of ∇v_h^Λ and $\partial_t v_h^\Lambda$ to ∇v^Λ and $\partial_t v^\Lambda$. This shows that $\nabla v_h^\Lambda \rightarrow \nabla v^\Lambda$ and $\partial_t v_h^\Lambda \rightarrow \partial_t v^\Lambda$ a.e. in $E(\Lambda)^c$ as $h \rightarrow 0^+$. This completes the proof of (v).

Then we discuss (vi) and (vii). It follows from (4.3) that

$$\begin{aligned} \iint_{E(\Lambda)^c} |v - v^\Lambda| |\partial_t v^\Lambda| dz &\leq \iint_{E(\Lambda)^c} \sum_{i \in \mathbb{N}} |v - v^i| |\omega_i| |\partial_t v^\Lambda| dz \\ &\leq \sum_{i \in \mathbb{N}} \iint_{2U_i} |v - v^i| |\partial_t v^\Lambda| dz \\ &\leq \sum_{i \in \mathbb{N}} \|\partial_t v^\Lambda\|_{L^\infty(2U_i)} \iint_{2U_i} |v - v^i| dz. \end{aligned}$$

By (4.21) and (4.25) we have

$$\begin{aligned} \|\partial_t v^\Lambda\|_{L^\infty(2U_i)} \iint_{2U_i} |v - v^i| dz &\leq cr_i \lambda_i |U_i| \|\partial_t v^\Lambda\|_{L^\infty(2U_i)} \\ &\leq c\Lambda |U_i| = c\Lambda \left| \frac{1}{6K} U_i \right|, \end{aligned}$$

and using the disjointedness of $\{\frac{1}{6K}U_i\}_{i \in \mathbb{N}}$ in Proposition 3.2 (ii) and $U_i \subset E(\Lambda)^c$, (vi) follows. Moreover, replacing (4.15), (4.17) and (4.25) by (4.14), (4.16) and (4.24) respectively, we conclude that (vii) holds.

Then we consider (viii). If $z \in E(\Lambda)$, it is easy to see from the definition of $E(\Lambda)$, that

$$\begin{aligned} &|v^\Lambda(z)|^p + a(z)|v^\Lambda(z)|^q + |\nabla v^\Lambda(z)|^p + a(z)|\nabla v^\Lambda(z)|^q \\ &= |v(z)|^p + a(z)|v(z)|^q + |\nabla v(z)|^p + a(z)|\nabla v(z)|^q \\ &\leq c(f^p(z) + a(z)f^q(z)) \leq c\Lambda. \end{aligned}$$

On the other hand, if $z \in E(\Lambda)^c$, then $z \in U_i$ for some $i \in \mathbb{N}$ and Lemma 4.7 and Lemma 4.9 give

$$|v^\Lambda(z)|^p + a(z)|v^\Lambda(z)|^q + |\nabla v^\Lambda(z)|^p + a(z)|\nabla v^\Lambda(z)|^q \leq c(\lambda_i^p + a(z)\lambda_i^q).$$

We estimate the right-hand side in two cases. If $U_i = Q_i$, we use (2.5), Proposition 3.2 (ix) and the fact that $\Lambda = \lambda_i^p + a(z_i)\lambda_i^q$, to get

$$\lambda_i^p + a(z)\lambda_i^q \leq \lambda_i^p + a(z_i)\lambda_i^q + [a]_\alpha r_i^\alpha \lambda_i^q \leq c\Lambda.$$

If $U_i = G_i$, we have by Proposition 3.2 (viii), that

$$\lambda_i^p + a(z)\lambda_i^q \leq c(\lambda_i^p + 2a(z_i)\lambda_i^q) \leq c\Lambda.$$

This completes the proof of (viii).

Finally, we prove (ix). From Lemma 4.10 follows that $|\nabla v_h^\Lambda(z)| \leq c\Lambda$ for a.e. $z \in \mathbb{R}^{n+1}$, and therefore

$$|\nabla v_h^\Lambda(z)|^p + a(z)|\nabla v_h^\Lambda(z)|^q \leq (1 + \|a\|_\infty)(1 + |\nabla v_h^\Lambda(z)|)^q \leq c\Lambda.$$

Since $v_h^\Lambda = 0$ in $Q_{R_2, S_2}(z_0)^c$, Lemma 4.10 implies that $|v_h^\Lambda(z)| \leq c\Lambda$ for a.e. $z \in \mathbb{R}^{n+1}$ and

$$|v_h^\Lambda(z)|^p + a(z)|v_h^\Lambda(z)|^q \leq (1 + \|a\|_\infty)(1 + |v_h^\Lambda(z)|)^q \leq c\Lambda.$$

This finishes the proof. \square

5. ENERGY ESTIMATE

In this section we use the Lipschitz truncation to prove the energy estimate in Theorem 2.4. The main idea of the proof is to use the estimates in Proposition 4.11 and conclude the desired convergence from (3.5). A similar argument will be used to prove Theorem 2.6 in Section 6.1.

5.1. Proof of Theorem 2.4. For $\tau \in \ell_{S_2-h}(t_0)$ and sufficiently small $\delta > 0$, let

$$\zeta_\delta(t) = \begin{cases} 1, & t \in (-\infty, \tau - \delta), \\ 1 - \frac{t - \tau + \delta}{\delta}, & t \in [\tau - \delta, \tau], \\ 0, & t \in (\tau, t_0 + S_2 - h). \end{cases} \quad (5.1)$$

Proposition 4.11 (i) implies that $v_h^\Lambda \eta^q \zeta_\delta(\cdot, t) \in W_0^{1,\infty}(B_{R_2}(x_0), \mathbb{R}^N)$ for every $t \in \ell_{S_2-h}(t_0)$. Therefore, $v_h^\Lambda \eta^q \zeta_\delta(\cdot, t)$ can be used as a test function in the Steklov averaged weak formulation of (2.3), and we obtain

$$\begin{aligned} \text{I} + \text{II} &= \iint_{Q_{R_2, S_2}(z_0)} \partial_t [u - u_0]_h \cdot v_h^\Lambda \eta^q \zeta_\delta dz \\ &\quad + \iint_{Q_{R_2, S_2}(z_0)} [\mathcal{A}(\cdot, \nabla u)]_h \cdot \nabla (v_h^\Lambda \eta^q \zeta_\delta) dz \\ &= \iint_{Q_{R_2, S_2}(z_0)} [|F|^{p-2} F + a|F|^{q-2} F]_h \cdot \nabla (v_h^\Lambda \eta^q \zeta_\delta) dz = \text{III}. \end{aligned}$$

We show the desired convergence of each term by splitting the domain of integration into the good set $E(\Lambda)$ and the bad set $E(\Lambda)^c$. First letting $h \rightarrow 0^+$ and then $\Lambda \rightarrow \infty$, the integral on the good set converges to the corresponding integral without a truncation, while the integral over the bad set converges to zero.

We start with term I. Integration by parts and the product rule give

$$\begin{aligned} \text{I} &= \iint_{Q_{R_2, S_2}(z_0)} (-v_h \cdot v_h^\Lambda \eta^{q-1} \partial_t \zeta_\delta - v_h \cdot \partial_t v_h^\Lambda \eta^{q-1} \zeta_\delta) dz \\ &\quad - \iint_{Q_{R_2, S_2}(z_0)} [u - u_0]_h \cdot v_h^\Lambda \eta^q \zeta_\delta \partial_t \zeta dz = \text{I}_1 + \text{I}_2. \end{aligned}$$

First we consider I_1 and write it as

$$\begin{aligned} \text{I}_1 &= \iint_{Q_{R_2, S_2}(z_0)} -|v_h|^2 \eta^{q-1} \partial_t \zeta_\delta dz + \iint_{Q_{R_2, S_2}(z_0)} v_h \cdot (v_h - v_h^\Lambda) \eta^{q-1} \partial_t \zeta_\delta dz \\ &\quad - \iint_{Q_{R_2, S_2}(z_0)} (v_h - v_h^\Lambda) \cdot \partial_t v_h^\Lambda \eta^{q-1} \zeta_\delta dz - \iint_{Q_{R_2, S_2}(z_0)} v_h^\Lambda \cdot \partial_t v_h^\Lambda \eta^{q-1} \zeta_\delta dz. \end{aligned}$$

Integration by parts in the last term above gives

$$\begin{aligned} \text{I}_1 &= \iint_{Q_{R_2, S_2}(z_0)} -\left(|v_h|^2 - \frac{1}{2}|v_h^\Lambda|^2\right) \eta^{q-1} \partial_t \zeta_\delta dz \\ &\quad + \iint_{Q_{R_2, S_2}(z_0)} v_h \cdot (v_h - v_h^\Lambda) \eta^{q-1} \partial_t \zeta_\delta dz - \iint_{E(\Lambda)^c} (v_h - v_h^\Lambda) \cdot \partial_t v_h^\Lambda \eta^q \zeta_\delta dz, \end{aligned}$$

where we also observed that the domain in the third integral can be written as $E(\Lambda)^c$ since $v_h = v_h^\Lambda$ in $E(\Lambda)$ and $v_h = v_h^\Lambda = 0$ in $Q_{R_2, S_2}(z_0)^c$.

We claim that the above terms converge to the corresponding integrals without Steklov averages as $h \rightarrow 0^+$. Indeed, in the first integral the term v_h converges to v in $L^2(Q_{R_2, S_2}(z_0), \mathbb{R}^N)$ and v_h^Λ converges to v^Λ in $L^\infty(Q_{R_2, S_2}(z_0), \mathbb{R}^N)$ by the properties of Steklov averages and Proposition 4.11 (iv). The second integral converges similarly. The third integral is bounded uniformly in $L^1(E(\Lambda)^c)$ for all $h > 0$ by Proposition 4.11 (vii) and since v_h, v_h^Λ and $\partial_t v_h^\Lambda$ converge a.e. to v, v^Λ and $\partial_t v^\Lambda$ respectively, we get the desired convergence. Therefore, we obtain

$$\begin{aligned} \lim_{h \rightarrow 0} \text{I}_1 &= \iint_{Q_{R_2, S_2}(z_0)} -\left(|v|^2 - \frac{1}{2}|v^\Lambda|^2\right) \eta^{q-1} \partial_t \zeta_\delta dz + \iint_{Q_{R_2, S_2}(z_0)} v \cdot (v - v^\Lambda) \eta^{q-1} \partial_t \zeta_\delta dz \\ &\quad - \iint_{E(\Lambda)^c} (v - v^\Lambda) \cdot \partial_t v^\Lambda \eta^q \zeta_\delta dz = \text{I}_{11} + \text{I}_{12} + \text{I}_{13}. \end{aligned}$$

Again, we estimate each term separately. Since the Lipschitz truncation is done only in the bad set, we have

$$\text{I}_{11} = \iint_{E(\Lambda)} -\frac{1}{2}|v|^2 \eta^{q-1} \partial_t \zeta_\delta dz - \iint_{E(\Lambda)^c} \left(|v|^2 - \frac{1}{2}|v^\Lambda|^2\right) \partial_t \zeta_\delta dz.$$

In order to conclude that the second term on the right-hand side vanishes as $\Lambda \rightarrow \infty$, we write

$$\begin{aligned} & - \iint_{E(\Lambda)^c} \left(|v|^2 - \frac{1}{2}|v^\Lambda|^2\right) \partial_t \zeta_\delta dz \\ &= - \iint_{E(\Lambda)^c} |v|^2 \partial_t \zeta_\delta dz + \frac{1}{2} \iint_{E(\Lambda)^c} |v^\Lambda|^2 \partial_t \zeta_\delta dz. \end{aligned} \tag{5.2}$$

Since $v \in L^2(Q_{R_2, S_2}(z_0), \mathbb{R}^N)$, by the absolute continuity of integral the first term on the right-hand side of (5.2) vanishes as $\Lambda \rightarrow \infty$. For the second term on the right-hand side of (5.2), by Proposition 4.11 (viii) we obtain

$$\iint_{E(\Lambda)^c} |v^\Lambda|^2 |\partial_t \zeta_\delta| dz \leq \iint_{E(\Lambda)^c} \Lambda^{\frac{2}{p}} |\partial_t \zeta_\delta| dz \leq c(\delta) \Lambda |E(\Lambda)^c|.$$

By (3.5) we conclude that

$$\lim_{\Lambda \rightarrow \infty} I_{11} = \iint_{Q_{R_2, S_2}(z_0)} -\frac{1}{2}|v|^2 \eta^{q-1} \partial_t \zeta_\delta \, dz.$$

To estimate the second term, I_{12} , note that by Young's inequality $v \cdot (v - v^\Lambda) \leq 2|v|^2 + |v^\Lambda|^2$ and thus

$$|I_{12}| \leq 2 \iint_{E(\Lambda)^c} (|v|^2 + |v^\Lambda|^2) |\partial_t \zeta_\delta| \, dz.$$

As the integral above contains the same terms as (5.2), we get $\lim_{\Lambda \rightarrow \infty} I_{12} = 0$ as before. Finally, it follows from Proposition 4.11 (vi) and (3.5) that $\lim_{\Lambda \rightarrow \infty} I_{13} = 0$. Combining the convergences for I_{11} , I_{12} and I_{13} , we conclude that

$$\lim_{\Lambda \rightarrow \infty} \lim_{h \rightarrow 0^+} I_1 = \iint_{Q_{R_2, S_2}(z_0)} -\frac{1}{2}|v|^2 \eta^{q-1} \partial_t \zeta_\delta \, dz.$$

It remains to estimate I_2 . As in the argument for I_1 , we have

$$\lim_{h \rightarrow 0} I_2 = - \iint_{Q_{R_2, S_2}(z_0)} (u - u_0) \cdot v^\Lambda \eta^q \zeta_\delta \partial_t \zeta \, dz.$$

Estimating with the absolute value and dividing into the good and bad sets, we obtain

$$\begin{aligned} & - \iint_{Q_{R_2, S_2}(z_0)} (u - u_0) \cdot v^\Lambda \eta^q \zeta_\delta \partial_t \zeta \, dz \\ & \geq - \iint_{Q_{R_2, S_2}(z_0) \cap E(\Lambda)} |u - u_0|^2 |\partial_t \zeta| \, dz - \iint_{E(\Lambda)^c} |u - u_0| |v^\Lambda| |\partial_t \zeta| \, dz. \end{aligned}$$

We show that the integral on the bad set vanishes as $\Lambda \rightarrow \infty$. Hölder's inequality gives

$$\begin{aligned} & \iint_{E(\Lambda)^c} |u - u_0| |v^\Lambda| |\partial_t \zeta| \, dz \\ & \leq \left(\iint_{Q_{R_2, S_2}(z_0) \cap E(\Lambda)^c} |u - u_0|^2 |\partial_t \zeta|^2 \, dz \right)^{\frac{1}{2}} \left(\iint_{E(\Lambda)^c} |v^\Lambda|^2 \, dz \right)^{\frac{1}{2}}, \end{aligned}$$

and the first integral vanishes when $\Lambda \rightarrow \infty$ as $|u| \in L^2(\Omega_T)$. For the second integral on the right-hand side, by Proposition 4.11 (viii) and (3.5) we have

$$\lim_{\Lambda \rightarrow \infty} \iint_{E(\Lambda)^c} |v^\Lambda|^2 \, dz \leq \lim_{\Lambda \rightarrow \infty} \Lambda^{\frac{2}{p}} |E(\Lambda)^c| \leq \lim_{\Lambda \rightarrow \infty} \Lambda |E(\Lambda)^c| = 0.$$

Therefore the integral over the bad set vanishes at the limit and we obtain

$$\lim_{\Lambda \rightarrow \infty} \lim_{h \rightarrow 0^+} I_2 \geq - \iint_{Q_{R_2, S_2}(z_0)} |u - u_0|^2 |\partial_t \zeta| \, dz.$$

Combining the estimates for I_1 and I_2 , we conclude that

$$\lim_{\Lambda \rightarrow \infty} \lim_{h \rightarrow 0^+} I \geq \iint_{Q_{R_2, S_2}(z_0)} -\frac{1}{2}|v|^2 \eta^{q-1} \partial_t \zeta_\delta \, dz - \iint_{Q_{R_2, S_2}(z_0)} |u - u_0|^2 |\partial_t \zeta| \, dz.$$

Next we estimate

$$\Pi = \iint_{Q_{R_2, S_2}(z_0)} [\mathcal{A}(\cdot, \nabla u)]_h \cdot \nabla (v_h^\Lambda \eta^q \zeta_\delta) \, dz.$$

We justify the convergence of the above terms as $h \rightarrow 0^+$. Observe that the structure condition (2.4) gives

$$|[\mathcal{A}(\cdot, \nabla u)]_h(z)| \leq L[|\nabla u|^{p-1} + a|\nabla u|^{q-1}]_h(z).$$

Moreover, we note that $|\nabla (v_h^\Lambda \eta^q \zeta_\delta)|$ is uniformly bounded in $L^\infty(Q_{R_2, S_2}(z_0))$ for $h > 0$ by Lemma 2.2 and Proposition 4.11 (ix). By convergence properties of the Steklov average, $[\mathcal{A}(\cdot, \nabla u)]_h \rightarrow$

$\mathcal{A}(\cdot, \nabla u)$, $\nabla u_h \rightarrow \nabla u$ and $[u - u_0]_h \rightarrow u - u_0$ a.e. as $h \rightarrow 0^+$. This implies the convergence to the integral without Steklov averages and we obtain

$$\begin{aligned} \lim_{h \rightarrow 0^+} \Pi &= \iint_{Q_{R_2, S_2(z_0)} \cap E(\Lambda)} \mathcal{A}(z, \nabla u) \cdot \nabla ((u - u_0) \eta^{q+1} \zeta^2 \zeta_\delta) dz \\ &\quad + \iint_{Q_{R_2, S_2(z_0)} \cap E(\Lambda)^c} \mathcal{A}(z, \nabla u) \cdot \nabla (v^\Lambda \eta^q \zeta \zeta_\delta) dz = \Pi_1 + \Pi_2. \end{aligned}$$

There is no Λ inside the integral over the good set and we can immediately let $\Lambda \rightarrow 0$. With the product rule of the gradient and estimating with (2.4) and (4.1), we get

$$\begin{aligned} \lim_{\Lambda \rightarrow \infty} \lim_{h \rightarrow 0^+} \Pi_1 &= \iint_{Q_{R_2, S_2(z_0)}} (\mathcal{A}(z, \nabla u) \cdot \nabla u) \eta^{q+1} \zeta^2 \zeta_\delta dz \\ &\quad + \iint_{Q_{R_2, S_2(z_0)}} \mathcal{A}(z, \nabla u) \cdot (u - u_0) \nabla (\eta^{q+1}) \zeta^2 \zeta_\delta dz \\ &\geq \nu \iint_{Q_{R_2, S_2(z_0)}} (|\nabla u|^p + a(z) |\nabla u|^q) \eta^{q+1} \zeta^2 \zeta_\delta dz \\ &\quad - (q+1)L \iint_{Q_{R_2, S_2(z_0)}} (|\nabla u|^{p-1} + a(z) |\nabla u|^{q-1}) \frac{|u - u_0|}{R_2 - R_1} \eta^q \zeta^2 \zeta_\delta dz. \end{aligned}$$

Applying Young's inequality twice with conjugate pairs $(\frac{p}{p-1}, p)$ and $(\frac{q}{q-1}, q)$, we obtain

$$\begin{aligned} (q+1)L \iint_{Q_{R_2, S_2(z_0)}} (|\nabla u|^{p-1} + a(z) |\nabla u|^{q-1}) \frac{|u - u_0|}{R_2 - R_1} \eta^q \zeta^2 \zeta_\delta dz \\ \leq \frac{\nu}{2} \iint_{Q_{R_2, S_2(z_0)}} (|\nabla u|^p + a(z) |\nabla u|^q) \eta^{q+1} \zeta^2 \zeta_\delta dz \\ + c(p, q, \nu, L) \iint_{Q_{R_2, S_2(z_0)}} \left(\frac{|u - u_0|^p}{(R_2 - R_1)^p} + a(z) \frac{|u - u_0|^q}{(R_2 - R_1)^q} \right) dz, \end{aligned} \quad (5.3)$$

where we were also estimated η and ζ by one in the last term. After absorbing terms we arrive at the desired estimate

$$\begin{aligned} \lim_{\Lambda \rightarrow \infty} \lim_{h \rightarrow 0^+} \Pi_1 &\geq \frac{\nu}{2} \iint_{Q_{R_2, S_2(z_0)}} (|\nabla u|^p + a(z) |\nabla u|^q) \eta^{q+1} \zeta^2 \zeta_\delta dz \\ &\quad - c(p, q, \nu, L) \iint_{Q_{R_2, S_2(z_0)}} \left(\frac{|u - u_0|^p}{(R_2 - R_1)^p} + a(z) \frac{|u - u_0|^q}{(R_2 - R_1)^q} \right) dz. \end{aligned} \quad (5.4)$$

We claim that Π_2 vanishes at the limit. Indeed, by the product rule and (2.4) we have

$$\begin{aligned} |\Pi_2| &\leq \iint_{Q_{R_2, S_2(z_0)} \cap E(\Lambda)^c} |\mathcal{A}(z, \nabla u)| |(\nabla v^\Lambda) \eta^q \zeta \zeta_\delta + v^\Lambda q (\nabla \eta) \eta^{q-1} \zeta \zeta_\delta| dz \\ &\leq c(L, q, R_2, R_1) \iint_{Q_{R_2, S_2(z_0)} \cap E(\Lambda)^c} (|\nabla u|^{p-1} + a(z) |\nabla u|^{q-1}) (|\nabla v^\Lambda| + |v^\Lambda|) dz. \end{aligned}$$

Applying Young's inequality, Proposition 4.11 (viii) and (3.5), we conclude that

$$\lim_{\Lambda \rightarrow \infty} |\Pi_2| \leq c \lim_{\Lambda \rightarrow \infty} \left(\iint_{Q_{R_2, S_2(z_0)} \cap E(\Lambda)^c} (|\nabla u|^p + a(z) |\nabla u|^q) dz + \Lambda |E(\Lambda)|^c \right) = 0,$$

where the convergence of the integral is justified by the integrability assumption of a weak solution. Therefore, we got from (5.4) the desired estimate

$$\begin{aligned} \lim_{\Lambda \rightarrow \infty} \lim_{h \rightarrow 0^+} \Pi &\geq \frac{\nu}{2} \iint_{Q_{R_2, S_2(z_0)}} H(z, |\nabla u|) \eta^{q+1} \zeta^2 \zeta_\delta dz \\ &\quad - c \iint_{Q_{R_2, S_2(z_0)}} H \left(z, \frac{|u - u_0|}{R_2 - R_1} \right) dz. \end{aligned}$$

The argument to estimate III is very similar to the argument for II. Again, we split into good and bad parts to get

$$\begin{aligned} \lim_{h \rightarrow 0^+} \text{III} &= \iint_{Q_{R_2, S_2}(z_0) \cap E(\Lambda)} (|F|^{p-2} F + a(z)|F|^{q-2} F) \cdot \nabla ((u - u_0)\eta^{q+1}\zeta^2\zeta_\delta) dz \\ &\quad + \iint_{Q_{R_2, S_2}(z_0) \cap E(\Lambda)^c} (|F|^{p-2} F + a(z)|F|^{q-2} F) \cdot \nabla (v^\Lambda \eta^q \zeta \zeta_\delta) dz = \text{III}_1 + \text{III}_2. \end{aligned}$$

Applying Young's inequality as in (5.3), we get

$$\begin{aligned} \lim_{\Lambda \rightarrow \infty} \lim_{h \rightarrow 0^+} \text{III}_1 &\leq c \iint_{Q_{R_2, S_2}(z_0)} \left(H\left(z, \frac{|u - u_0|}{R_2 - R_1}\right) + H(z, |F|) \right) dz \\ &\quad + \frac{\nu}{4} \iint_{Q_{R_2, S_2}(z_0)} H(z, |\nabla u|) \eta^{q+1} \zeta^2 \zeta_\delta dz, \end{aligned}$$

where $c = c(p, q, \nu)$. Using the same arguments with $|\nabla u|$ replaced by $|F|$, we can estimate III_2 in the same way as II_2 . Therefore, we conclude that $\lim_{\Lambda \rightarrow \infty} \lim_{h \rightarrow 0^+} \text{III}_2 = 0$ and

$$\begin{aligned} \lim_{\Lambda \rightarrow \infty} \lim_{h \rightarrow 0^+} \text{III} &\leq c \iint_{Q_{R_2, S_2}(z_0)} \left(H\left(z, \frac{|u - u_0|}{R_2 - R_1}\right) + H(z, |F|) \right) dz \\ &\quad + \frac{\nu}{4} \iint_{Q_{R_2, S_2}(z_0)} H(z, |\nabla u|) \eta^{q+1} \zeta^2 \zeta_\delta dz. \end{aligned}$$

Combining the estimates for I, II and III at the limit of Λ and h , reordering and absorbing the last term of III with the first term of II, we get

$$\begin{aligned} &\iint_{Q_{R_2, S_2}(z_0)} -\frac{1}{2} |v|^2 \eta^{q-1} \partial_t \zeta_\delta dz + \frac{\nu}{4} \iint_{Q_{R_2, S_2}(z_0)} H(z, |\nabla u|) \eta^{q+1} \zeta^2 \zeta_\delta dz \\ &\leq c \iint_{Q_{R_2, S_2}(z_0)} \left(H\left(z, \frac{|u - u_0|}{R_2 - R_1}\right) + |u - u_0|^2 |\partial_t \zeta| + H(z, |F|) \right) dz. \end{aligned}$$

Taking $\delta \rightarrow 0^+$ and recalling that $\tau \in \ell_{S_1}(t_0)$ in (5.1) is arbitrary, we conclude that

$$\begin{aligned} &\sup_{t \in \ell_{S_1}(t_0)} \int_{B_{R_1}(x_0)} \frac{|u - u_0|^2}{S_1} dx + \iint_{Q_{R_1, S_1}(z_0)} H(z, |\nabla u|) dz \\ &\leq c \iint_{Q_{R_2, S_2}(z_0)} \left(H\left(z, \frac{|u - u_0|}{R_2 - R_1}\right) + \frac{|u - u_0|^2}{S_2 - S_1} + H(z, |F|) \right) dz, \end{aligned}$$

where we made the integration domain smaller on the left hand side and used the definitions for the cutoff functions in (4.1). Also note the use of the fact that $|Q_{R_2, S_2}| \approx |Q_{R_1, S_1}|$ by $R_1 \in [R_2/2, R_2]$ and $S_1 \in [S_2/2^2, S_2]$. The proof is completed by replacing u_0 with $(u)_{Q_{R_1, S_1}(z_0)}$ on the first term on the left-hand side.

6. EXISTENCE AND UNIQUENESS

In this section, we consider the Dirichlet problem (2.6). We apply the following standard estimates for the p -Laplace structure with $p \geq 2$. There exists a constant $c = c(n, N, p)$ such that

$$c|\xi_1 - \xi_2|^p \leq (|\xi_1|^{p-2}\xi_1 - |\xi_2|^{p-2}\xi_2) \cdot (\xi_1 - \xi_2) \quad (6.1)$$

and

$$\| |\xi_1|^{p-2}\xi_1 - |\xi_2|^{p-2}\xi_2 \| \leq c(|\xi_1|^{p-2} + |\xi_2|^{p-2})|\xi_1 - \xi_2| \quad (6.2)$$

for every $\xi_1, \xi_2 \in \mathbb{R}^{Nn}$.

6.1. Proof of Theorem 2.6. In this subsection, we provide the uniqueness of weak solutions to (2.6). Assume that u and w are weak solutions of (2.6). Then we have (2.8) with (2.9) and (2.10). Note that since $(u - w)(\cdot, R^2) \not\equiv 0$ in D_R may happen, the truncated function is not able to be defined as zero on the bad set as in (4.4). We overcome this difficulty by extending the system (2.8) to $D_R \times \mathbb{R}$. We first extend $u \equiv w \equiv g$ in $D_R \times (-\infty, -R^2]$. Then we have

$$(u - w)_t - \operatorname{div}(\mathcal{A}(z, \nabla u) - \mathcal{A}(z, \nabla w)) = 0 \quad \text{in } D_R \times (-\infty, R^2). \quad (6.3)$$

Indeed, for any $\varphi \in C_0^\infty(D_R \times (-\infty, R^2), \mathbb{R}^N)$, we have

$$- \iint_{D_R \times (-\infty, R^2)} (u - w) \cdot \varphi_t \, dz = - \lim_{\epsilon \rightarrow 0^+} \iint_{C_R} (u - w) \cdot (\varphi \zeta_\epsilon)_t \, dz,$$

where ζ_ϵ is a function defined as

$$\zeta_\epsilon(t) = \begin{cases} 0, & t \in (-\infty, -R^2), \\ \frac{1}{\epsilon} t, & t \in [-R^2, -R^2 + \epsilon], \\ 1, & t \in (-R^2 + \epsilon, \infty). \end{cases}$$

Here we used (2.10) to have

$$\lim_{\epsilon \rightarrow 0^+} \iint_{C_R} (u - w) \cdot \varphi(\zeta_\epsilon)_t \, dz = \lim_{\epsilon \rightarrow 0^+} \int_{-R^2}^{-R^2 + \epsilon} \int_{D_R} (u - w) \cdot \varphi \, dx \, dt = 0.$$

Taking $\varphi \zeta_\epsilon$ as a test function in (2.8), it follows that

$$\begin{aligned} - \iint_{D_R \times (-\infty, R^2)} (u - w) \cdot \varphi_t \, dz &= - \lim_{\epsilon \rightarrow 0^+} \iint_{C_R} (u - w) \cdot (\varphi \zeta_\epsilon)_t \, dz \\ &= - \lim_{\epsilon \rightarrow 0^+} \iint_{C_R} (\mathcal{A}(z, \nabla u) - \mathcal{A}(z, \nabla w)) \cdot \nabla \varphi \zeta_\epsilon \, dz \\ &= - \iint_{D_R \times (-\infty, R^2)} (\mathcal{A}(z, \nabla u) - \mathcal{A}(z, \nabla w)) \cdot \nabla \varphi \, dz. \end{aligned}$$

This means that (6.3) holds. For the range $D_R \times \mathbb{R}$, we remark that there is no possible extension to have $(u - w)_t - \operatorname{div}(\mathcal{A}(z, \nabla u) - \mathcal{A}(z, \nabla w)) = 0$ in $D_R \times \mathbb{R}$. In fact, by considering $G(z) = (\mathcal{A}(z, \nabla u) - \mathcal{A}(z, \nabla w))$ and extending u, w and G to be

$$(u - w)(x, t) = (u - w)(x, 2R^2 - t), \quad G(x, t) = -G(x, 2R^2 - t) \quad (6.4)$$

for $(x, t) \in D_R \times (R^2, \infty)$, we have $(u - w)_t - \operatorname{div} G = 0$ in $D_R \times \mathbb{R}$. For any $\varphi \in C_0^\infty(D_R \times \mathbb{R}, \mathbb{R}^N)$, consider

$$\psi(x, t) = \varphi(x, t) - \varphi(x, 2R^2 - t).$$

Note that $\psi \in W_0^{1, \infty}(D_R \times (-\infty, R^2), \mathbb{R}^N)$. Taking ψ as a test function in (6.3), we obtain

$$\begin{aligned} 0 &= \iint_{D_R \times (-\infty, R^2)} -(u - w) \cdot \psi_t \, dz + \iint_{B_R \times (-\infty, R^2)} G \cdot \nabla \psi \, dz \\ &= \iint_{D_R \times (-\infty, R^2)} (-(u - w) \cdot \varphi_t + G \cdot \nabla \varphi) \, dz \\ &\quad + \iint_{D_R \times (-\infty, R^2)} (-(u - w)(x, t) \cdot \varphi_t(x, 2R^2 - t) - G(x, t) \cdot \nabla \varphi(x, 2R^2 - t)) \, dz, \end{aligned}$$

and a change of variables gives

$$\iint_{D_R \times \mathbb{R}} (-(u - w) \cdot \varphi_t + G \cdot \nabla \varphi) \, dz = 0.$$

Therefore, $u - w \equiv 0$ in $(D_R \times (-R^2, 3R^2))^c$ and

$$(u - w)_t - \operatorname{div} G = 0 \quad \text{in } D_R \times \mathbb{R}. \quad (6.5)$$

We also observe that if we extend

$$u(x, t) = u(x, 2R^2 - t), \quad w(x, t) = w(x, 2R^2 - t), \quad a(x, t) = a(x, 2R^2 - t)$$

in $(x, t) \in D_R \times (-R^2, 3R^2)$ then the first extension of (6.4) is satisfied and

$$\nabla u(x, t) = \nabla u(x, 2R^2 - t), \quad \nabla w(x, t) = \nabla w(x, 2R^2 - t)$$

is well-defined in $(x, t) \in D_R \times (-R^2, 3R^2)$ with

$$H(z, |\nabla u|), H(z, |\nabla w|) \in L^1(D_R \times (-R^2, 3R^2)).$$

With these extensions we define a function $f : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ as

$$f(z) = (|\nabla u(z)| + |\nabla w(z)| + |u(z)| + |w(z)|)\chi_{D_R \times (-R^2, 3R^2)}(z).$$

It is easy to see that $H(z, f) \in L^1(\mathbb{R}^{n+1})$. The function f plays the same role in constructing an open subset in \mathbb{R}^{n+1} as in Section 4. To employ the Lipschitz truncation method, it needs to be shown that $|G|$ is bounded by $f^{p-1} + af^{q-1}$. Indeed, using (6.2) and (2.7), there exists $c = c(n, N, p, q, L)$ such that

$$|G| \leq c(|\nabla u|^{p-2} + |\nabla w|^{p-2})|\nabla u - \nabla w| + ca(|\nabla u|^{q-2} + |\nabla w|^{q-2})|\nabla u - \nabla w|,$$

and applying triangle inequality, to have $|\nabla u - \nabla w| \leq |\nabla u| + |\nabla w|$, and Young's inequality, we obtain

$$|G| \leq c((|\nabla u| + |\nabla w|)^{p-1} + a(|\nabla u| + |\nabla w|)^{q-1}) \leq c(f^{p-1} + af^{q-1}) \quad (6.6)$$

in $D_R \times \mathbb{R}$ as G is extended 0 below $\{t = -R^2\}$ and then extended oddly above $\{t = R^2\}$. Denoting $v = u - w$, we have that v_h is well defined in $D_R \times \mathbb{R}$ and

$$\partial_t v_h - \operatorname{div} G_h = 0 \quad \text{in } D_R \times \{t\} \quad (6.7)$$

for a.e. $t \in \mathbb{R}$. We follow the proof of Lemma 4.3 by taking the test function $\varphi\psi_\delta$ in (6.5) and (6.7) with (6.6) to obtain the following estimates. We omit the details. We keep track of the dependencies and denote

$$\text{data} = n, N, p, q, \alpha, \nu, L, [a]_\alpha, R, K, \|a\|_{L^\infty(\mathbb{R}^{n+1})}.$$

Lemma 6.1. *Let $Q = B_r \times \ell_s$ be a cylinder and satisfying $B_r \subset D_R$. There exists a constant $c = c(\text{data})$ such that*

$$\iint_Q |v_h - (v_h)_Q| dz \leq csr^{-1} \iint_{[Q]_h} (f^{p-1} + af^{q-1}) dz + cr \iint_{[Q]_h} f dz.$$

Moreover, the above estimate holds with v_h and $[Q]_h$ replaced by v and Q .

As (6.7) is defined on \mathbb{R} , the second estimate in Lemma 4.3 is not necessary. On the other hand, we have used Lemma 4.4 to deal with the case of B_r intersecting with the complement of the given domain in Section 4. Since a cube D_R satisfies the measure density condition, we may replace $B_\rho(x_0)$ by D_R in the statement of Lemma 4.4. We refer to [18, Example 6.18 and Theorem 6.22] for details. It is also noteworthy to mention that D_R has a Lipschitz boundary and this fact along with (2.9) guarantees that $u - w \in L^s(I_{3R}; W_0^{1,s}(D_R; \mathbb{R}^N))$ for $1 < s \leq p$. Therefore, we may apply Lemma 4.4 with $1 < s < p$ as in Section 4. With the Whitney decomposition $\{U_i = B_i \times I_i\}_{i \in \mathbb{N}}$ and partition of unity $\{\omega_i\}_{i \in \mathbb{N}}$, we define the Lipschitz truncation of v_h and v as

$$v_h^\Lambda(z) = v_h(z) - \sum_{i \in \mathbb{N}} (v_h(z) - v_h^i) \omega_i(z) \quad \text{and} \quad v^\Lambda(z) = v(z) - \sum_{i \in \mathbb{N}} (v(z) - v^i) \omega_i(z),$$

where

$$v_h^i = \begin{cases} (v_h)_{2U_i}, & \text{if } 2B_i \subset D_R, \\ 0, & \text{otherwise,} \end{cases} \quad v^i = \begin{cases} (v)_{2U_i}, & \text{if } 2B_i \subset D_R, \\ 0, & \text{otherwise.} \end{cases}$$

We now state the Lipschitz truncation result analogous to Proposition 4.11 in Section 4. Again, the proof is omitted.

Proposition 6.2. *Let $E(\Lambda)$ be as in (3.4). There exist functions $\{v_h^\Lambda\}_{h>0}$ and a function v^Λ satisfying the following properties:*

- (i) $v_h^\Lambda \in W^{1,2}(I_{3R}; L^2(D_R; \mathbb{R}^N)) \cap L^\infty(I_{3R}; W_0^{1,\infty}(D_R, \mathbb{R}^N))$.
- (ii) $v^\Lambda \in L^\infty(I_{3R}; W_0^{1,\infty}(D_R, \mathbb{R}^N))$.
- (iii) $v_h^\Lambda = v_h$, $v^\Lambda = v$, $\nabla v_h^\Lambda = \nabla v_h$, $\nabla v^\Lambda = \nabla v$ a.e. in $E(\Lambda)$.
- (iv) $v_h^\Lambda \rightarrow v^\Lambda$ in $L^\infty(D_R \times I_{3R}, \mathbb{R}^N)$ as $h \rightarrow 0^+$, taking a subsequence if necessary.
- (v) $\nabla v_h^\Lambda \rightarrow \nabla v^\Lambda$ and $\partial_t v_h^\Lambda \rightarrow \partial_t v^\Lambda$ a.e. in $E(\Lambda)^c$ as $h \rightarrow 0^+$.

Moreover, there exist constants $c = c(\text{data})$ and $c_\Lambda = c_\Lambda(\text{data}, \Lambda)$ such that

- (vi) $\iint_{E(\Lambda)^c} |v - v^\Lambda| |\partial_t v^\Lambda| dz \leq c\Lambda |E(\Lambda)^c|$,
- (vii) $\iint_{E(\Lambda)^c} |v_h - v_h^\Lambda| |\partial_t v_h^\Lambda| dz \leq c_\Lambda |E(\Lambda)^c|$,
- (viii) $H(z, |v^\Lambda(z)|) + H(z, |\nabla v^\Lambda(z)|) \leq c\Lambda$ for a.e. $z \in \mathbb{R}^{n+1}$,
- (ix) $H(z, |v_h^\Lambda(z)|) + H(z, |\nabla v_h^\Lambda(z)|) \leq c_\Lambda$ for a.e. $z \in \mathbb{R}^{n+1}$.

In the remaining of this subsection, we apply the energy estimate and conclude Theorem 2.6. For $\tau_1, \tau_2 \in I_R$ with $\tau_1 < \tau_2$, let $\delta > 0$ and $h > 0$ be sufficiently small so that $2\delta < \tau_2 - \tau_1$ and $\tau_1, \tau_2 \in I_{R-h}$. We let $\zeta_\delta \in W_0^{1,\infty}(I_{R-h})$ be a function defined as

$$\zeta_\delta(t) = \begin{cases} \frac{1}{\delta}(t - \tau_1), & \tau_1 \leq t \leq \tau_1 + \delta, \\ 1, & \tau_1 + \delta \leq t \leq \tau_2 - \delta, \\ 1 - \frac{1}{\delta}(t - \tau_2 + \delta), & \tau_2 - \delta \leq t \leq \tau_2, \\ 0, & \text{otherwise.} \end{cases}$$

Let $t \in I_R$. We take $v_h^\Lambda \zeta_\delta$ as a test function in (6.7), that is equivalently

$$\partial_t [u - w]_h - \operatorname{div} [\mathcal{A}(\cdot, \nabla u) - \mathcal{A}(\cdot, \nabla w)]_h = 0 \quad \text{in } D_R \times \{t\}.$$

Recalling that $u - w = v$, we obtain

$$0 = \iint_{C_R} \partial_t v_h \cdot v_h^\Lambda \zeta_\delta dz + \iint_{C_R} [\mathcal{A}(\cdot, \nabla u) - \mathcal{A}(\cdot, \nabla w)]_h \cdot \nabla v_h^\Lambda \zeta_\delta dz. \quad (6.8)$$

To estimate the first term on the right-hand side, we add and subtract $\partial_t v_h^\Lambda$, then apply integration by parts to have

$$\begin{aligned} \iint_{C_R} \partial_t v_h \cdot v_h^\Lambda \zeta_\delta dz &= \iint_{C_R} \partial_t (v_h - v_h^\Lambda) \cdot v_h^\Lambda \zeta_\delta dz + \iint_{C_R} \frac{1}{2} (\partial_t |v_h^\Lambda|^2) \zeta_\delta dz \\ &= - \iint_{C_R} (v_h - v_h^\Lambda) \cdot \partial_t v_h^\Lambda \zeta_\delta dz - \iint_{C_R} (v_h - v_h^\Lambda) \cdot v_h^\Lambda \partial_t \zeta_\delta dz - \iint_{C_R} \frac{1}{2} |v_h^\Lambda|^2 \partial_t \zeta_\delta dz. \end{aligned}$$

Following the argument in the proof of Theorem 2.4 by using Proposition 6.2, we deduce

$$\begin{aligned} &\lim_{\Lambda \rightarrow \infty} \lim_{h \rightarrow 0^+} \iint_{C_R} \partial_t v_h \cdot v_h^\Lambda \zeta_\delta dz \\ &= \int_{\tau_2 - \delta}^{\tau_2} \int_{D_R} \frac{1}{2} |v|^2 dx dt - \int_{\tau_1}^{\tau_1 + \delta} \int_{D_R} \frac{1}{2} |v|^2 dx dt. \end{aligned}$$

The last term in (6.8) is estimated as in the proof of Theorem 2.4 with Proposition 6.2. Then we obtain

$$\begin{aligned} &\lim_{\Lambda \rightarrow \infty} \lim_{h \rightarrow 0^+} \iint_{C_R} [\mathcal{A}(\cdot, \nabla u) - \mathcal{A}(\cdot, \nabla w)]_h \cdot \nabla v_h^\Lambda \zeta_\delta dz \\ &= \iint_{C_R} (\mathcal{A}(\cdot, \nabla u) - \mathcal{A}(\cdot, \nabla w)) \cdot (\nabla u - \nabla w) \zeta_\delta dz. \end{aligned}$$

By (2.7) and (6.1) there exists a constant $c = c(n, N, p, q, \nu)$ such that

$$\begin{aligned} & \iint_{C_R} (\mathcal{A}(\cdot, \nabla u) - \mathcal{A}(\cdot, \nabla w)) \cdot (\nabla u - \nabla w) \zeta_\delta \, dz \\ & \geq c \iint_{C_R} (|\nabla u - \nabla w|^p + a|\nabla u - \nabla w|^q \zeta_\delta) \, dz \\ & = c \iint_{C_R} H(z, |\nabla u - \nabla w|) \zeta_\delta \, dz \geq 0. \end{aligned}$$

Hence, it follows from (6.8) that

$$\int_{\tau_2 - \delta}^{\tau_2} \int_{D_R} |u - w|^2 \, dx \, dt \leq \int_{\tau_1}^{\tau_1 + \delta} \int_{D_R} |u - w|^2 \, dx \, dt,$$

where $\tau_1, \tau_2 \in I_R$ are arbitrary and δ is sufficiently small. We set $\tau_1 = \delta$ to have

$$\int_{\tau_2 - \delta}^{\tau_2} \int_{D_R} |u - w|^2 \, dx \, dt \leq \int_{\delta}^{2\delta} \int_{D_R} |u - w|^2 \, dx \, dt \leq 2 \int_0^{2\delta} \int_{D_R} |u - w|^2 \, dx \, dt.$$

Letting $\delta \rightarrow 0^+$, it follows from (2.10) that

$$\int_{D_R} |u - w|^2(x, \tau_2) \, dx \leq 0$$

for a.e. $\tau_2 \in I_R$. We conclude $u = w$ a.e. in C_R . This completes the proof of Theorem 2.6.

6.2. Proof of Theorem 2.7. In this subsection we assume $g \equiv 0$ and $b(\cdot)$ satisfies VMO condition (2.11). We construct a sequence of weak solutions of the Dirichlet boundary problem for perturbed q -Laplace type systems. This sequence converges to a weak solution of (2.6).

6.2.1. *Approximation.* For $k \in \mathbb{N}$, let F^k be the truncated function

$$F^k(z) = F(z) \chi_{\{|F(z)| \leq k\}}.$$

For $\epsilon > 0$, we consider the function

$$F_\epsilon^k(z) = F^k(z) \chi_{\{|F^k(z)| \leq \epsilon^{-\frac{1}{q-p}}\}}. \quad (6.9)$$

and the perturbed operator and map

$$H_\epsilon(z, |\xi|) = |\xi|^p + a_\epsilon(z)|\xi|^q, \quad \mathcal{A}_\epsilon(z, \xi) = b(z)(|\xi|^{p-2}\xi + a_\epsilon(z)|\xi|^{q-2}\xi),$$

where $a_\epsilon(z) = a(z) + \epsilon$. We observe $\mathcal{A}_\epsilon(z, \xi)$ is q -Laplace type operator as it satisfies

$$\epsilon|\xi|^q \leq \mathcal{A}_\epsilon(z, \xi) \cdot \xi, \quad |\mathcal{A}_\epsilon(z, \xi)| \leq 2^q L(\|a\|_\infty + 1)(|\xi|^{q-1} + 1)$$

and (6.1) implies the monotonicity condition

$$(\mathcal{A}_\epsilon(z, \xi_1) - \mathcal{A}_\epsilon(z, \xi_2)) \cdot (\xi_1 - \xi_2) > 0 \quad \text{for any } \xi_1, \xi_2 \in \mathbb{R}^{Nn} \text{ with } \xi_1 \neq \xi_2.$$

For each $k \in \mathbb{N}$ and $\epsilon > 0$, there exists by standard existence theory for q -parabolic equations the unique weak solution $u_\epsilon^k \in C(I_R; L^2(D_R, \mathbb{R}^N)) \cap L^q(I_R; W_0^{1,q}(D_R, \mathbb{R}^N))$ to the Dirichlet problem

$$\begin{cases} \partial_t u_\epsilon^k - \operatorname{div} \mathcal{A}_\epsilon(z, \nabla u_\epsilon^k) = -\operatorname{div}(|F_\epsilon^k|^{p-2} F_\epsilon^k + a_\epsilon |F_\epsilon^k|^{q-2} F_\epsilon^k) & \text{in } C_R, \\ u_\epsilon^k = 0 & \text{on } \partial_p C_R. \end{cases} \quad (6.10)$$

6.2.2. *First convergence.* By the standard energy estimate for the q -Laplace problems, there exists a constant $c = c(p, q, \nu)$ such that

$$\sup_{t \in I_R} \int_{D_R} |u_\epsilon^k(x, t)|^2 dx + \iint_{C_R} H_\epsilon(z, |\nabla u_\epsilon^k|) dz \leq c \iint_{C_R} H_\epsilon(z, |F_\epsilon^k|) dz. \quad (6.11)$$

We observe from (6.9) that

$$\epsilon |F_\epsilon^k(z)|^q = \epsilon |F_\epsilon^k(z)|^{q-p} |F_\epsilon^k(z)|^p \leq |F^k(z)|^p \leq |F(z)|^p. \quad (6.12)$$

Therefore, (6.11) becomes

$$\sup_{t \in I_R} \int_{D_R} |u_\epsilon^k(x, t)|^2 dx + \iint_{C_R} H_\epsilon(z, |\nabla u_\epsilon^k|) dz \leq c \iint_{C_R} H(z, |F|) dz. \quad (6.13)$$

Let $k \in \mathbb{N}$. First we investigate the weak limit of $\{u_\epsilon^k\}_{\epsilon > 0}$ when $\epsilon \rightarrow 0$. To apply a compactness result, we estimate $\{\partial_t u_\epsilon^k\}_{\epsilon > 0}$. Let $\varphi \in C_0^\infty(C_R, \mathbb{R}^N)$ be a test function in (6.10). By (2.7) there exists a constant $c = c(L)$ such that

$$\begin{aligned} \left| \iint_{C_R} \partial_t u_\epsilon^k \cdot \varphi dz \right| &\leq c \iint_{C_R} (|\nabla u_\epsilon^k|^{p-1} + |F_\epsilon^k|^{p-1}) |\nabla \varphi| dz \\ &+ c \iint_{C_R} a_\epsilon(z) (|\nabla u_\epsilon^k|^{q-1} + |F_\epsilon^k|^{q-1}) |\nabla \varphi| dz. \end{aligned}$$

By Hölder's inequality, there exists a constant $c = c(n, p, q, L, \|a\|_\infty, R)$ such that

$$\begin{aligned} \left| \iint_{C_R} \partial_t u_\epsilon^k \cdot \varphi dz \right| &\leq c \left(\iint_{C_R} |\nabla u_\epsilon^k|^p + |F_\epsilon^k|^p dz \right)^{\frac{1}{p'}} \left(\iint_{C_R} |\nabla \varphi|^q dz \right)^{\frac{1}{q}} \\ &+ c \left(\iint_{C_R} a_\epsilon(z) (|\nabla u_\epsilon^k|^q + |F_\epsilon^k|^q) dz \right)^{\frac{1}{q'}} \left(\iint_{C_R} |\nabla \varphi|^q dz \right)^{\frac{1}{q}}, \end{aligned}$$

and using (6.11) and (6.13), we obtain

$$\left| \iint_{C_R} \partial_t u_\epsilon^k \cdot \varphi dz \right| \leq c \left(\iint_{C_R} H(z, |F^k|) dz + 1 \right)^{\frac{1}{q'}} \left(\iint_{C_R} |\nabla \varphi|^q dz \right)^{\frac{1}{q}}.$$

Thus we have

$$\|\partial_t u_\epsilon^k\|_{L^{q'}(I_R; W^{-1, q'}(D_R, \mathbb{R}^N))} \leq c \left(\iint_{C_R} H(z, |F|) dz + 1 \right)^{\frac{1}{q'}}. \quad (6.14)$$

Since the function $b(\cdot)$ satisfies the VMO condition (2.11), we may apply a global Calderón-Zygmund estimate in [16, Theorem 2.5] for a weak solution to (6.10). Since (6.12) holds, $[a + \epsilon]_\alpha = [a]_\alpha$ and $\|a + \epsilon\|_{L^\infty(C_R)} \leq \|a\|_{L^\infty(C_R)} + 1$, there exists a constant

$$c_k = c_k(n, N, p, q, [a]_\alpha, R, \|H(z, |F^k|)\|_{L^{\frac{q}{p}}(C_R)}, \|a\|_{L^\infty(C_R)})$$

such that

$$\iint_{C_R} H_\epsilon(z, |\nabla u_\epsilon^k|)^{\frac{q}{p}} dz \leq c_k \quad \text{for every } \epsilon > 0.$$

In particular, we have

$$\iint_{C_R} |\nabla u_\epsilon^k|^q dz \leq c_k \quad \text{for every } \epsilon > 0. \quad (6.15)$$

Since estimates (6.14) and (6.15) hold for every $\epsilon > 0$ and we have

$$L^q(I_R; W_0^{1, q}(D_R, \mathbb{R}^N)) \Subset L^q(C_R, \mathbb{R}^N) \Subset L^2(C_R, \mathbb{R}^N) \subset L^{q'}(I_R; W^{-1, q'}(D_R, \mathbb{R}^N)),$$

a compactness result in [20, Corollary 4, Section 8] implies that there exists a limit function $u^k \in C(I_R; L^2(D_R, \mathbb{R}^N)) \cap L^q(I_R; W_0^{1,q}(D_R, \mathbb{R}^N))$ such that, by passing to a subsequence if necessary, we have

$$\begin{aligned} u_\epsilon^k &\rightarrow u^k \quad \text{strongly in } C(I_R; L^2(D_R, \mathbb{R}^N)) \cap L^q(C_R, \mathbb{R}^N), \\ u_\epsilon^k &\rightarrow u^k \quad \text{weakly in } L^q(I_R; W_0^{1,q}(D_R, \mathbb{R}^N)), \\ \partial_t u_\epsilon^k &\rightarrow \partial_t u^k \quad \text{weakly in } L^{q'}(I_R; W^{-1,q'}(D_R, \mathbb{R}^N)), \end{aligned}$$

as $\epsilon \rightarrow 0^+$. We claim that u^k is the weak solution of

$$\begin{cases} \partial_t u^k - \operatorname{div} \mathcal{A}(z, \nabla u^k) = -\operatorname{div}(|F^k|^{p-2} F^k + a|F^k|^{q-2} F^k) & \text{in } C_R, \\ u^k = 0 & \text{on } \partial_p C_R. \end{cases} \quad (6.16)$$

It suffices to show that

$$\iint_{C_R} H(z, |\nabla u_\epsilon^k - \nabla u^k|) dz \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0^+. \quad (6.17)$$

Indeed, the above strong convergence provides the weak convergence of operators. For any $\varphi \in C_0^\infty(C_R, \mathbb{R}^N)$ there holds

$$\begin{aligned} &\left| \iint_{C_R} (\mathcal{A}_\epsilon(z, \nabla u_\epsilon^k) - \mathcal{A}(z, \nabla u^k)) \cdot \nabla \varphi dz \right| \\ &\leq \iint_{C_R} |\mathcal{A}(z, \nabla u_\epsilon^k) - \mathcal{A}(z, \nabla u^k)| |\nabla \varphi| dz + L \epsilon \iint_{C_R} |\nabla u_\epsilon^k|^{q-1} |\nabla \varphi| dz. \end{aligned}$$

By Hölder's inequality and (6.13), we obtain

$$\lim_{\epsilon \rightarrow 0^+} \epsilon \iint_{C_R} |\nabla u_\epsilon^k|^{q-1} |\nabla \varphi| dz \leq \lim_{\epsilon \rightarrow 0^+} \epsilon |C_R|^{1 - \frac{q-1}{p}} \left(\iint_{C_R} |\nabla u_\epsilon^k|^p dz \right)^{\frac{q-1}{p}} \|\nabla \varphi\|_\infty = 0.$$

We apply (6.2) to estimate the remaining term. There exists a constant $c = c(n, N, p, q)$ such that

$$\begin{aligned} &\iint_{C_R} |\mathcal{A}(z, \nabla u_\epsilon^k) - \mathcal{A}(z, \nabla u^k)| |\nabla \varphi| dz \\ &\leq c \iint_{C_R} (|\nabla u_\epsilon^k|^{p-2} + |\nabla u^k|^{p-2}) |\nabla u_\epsilon^k - \nabla u^k| |\nabla \varphi| dz \\ &\quad + c \iint_{C_R} a(z) (|\nabla u_\epsilon^k|^{q-2} + |\nabla u^k|^{q-2}) |\nabla u_\epsilon^k - \nabla u^k| |\nabla \varphi| dz. \end{aligned}$$

Applying Hölder's inequality and (6.13), there exists $c = c(n, N, p, q, \|a\|_\infty, R)$ such that

$$\begin{aligned} &\iint_{C_R} |\mathcal{A}(z, \nabla u_\epsilon^k) - \mathcal{A}(z, \nabla u^k)| |\nabla \varphi| dz \\ &\leq c \left(\iint_{C_R} H(z, |F|) dz \right)^{\frac{p-2}{p}} \left(\iint_{C_R} |\nabla u_\epsilon^k - \nabla u^k|^p dz \right)^{\frac{1}{p}} \|\nabla \varphi\|_\infty \\ &\quad + c \left(\iint_{C_R} H(z, |F|) dz \right)^{\frac{q-2}{q}} \left(\iint_{C_R} a(z) |\nabla u_\epsilon^k - \nabla u^k|^q dz \right)^{\frac{1}{q}} \|\nabla \varphi\|_\infty. \end{aligned}$$

Therefore, (6.17) leads to

$$\lim_{\epsilon \rightarrow 0^+} \iint_{C_R} |\mathcal{A}(z, \nabla u_\epsilon^k) - \mathcal{A}(z, \nabla u^k)| |\nabla \varphi| dz = 0,$$

which means that $\mathcal{A}_\epsilon(z, \nabla u_\epsilon^k) \rightarrow \mathcal{A}(z, \nabla u^k)$ weakly in $\mathcal{D}'(Q_R, \mathbb{R}^N)$ as $\epsilon \rightarrow 0^+$.

To prove (6.17), we take $u_\epsilon^k - u_{\epsilon'}^k \in C(I_R; L^2(D_R, \mathbb{R}^N)) \cap L^q(I_R; W_0^{1,q}(D_R, \mathbb{R}^N))$ as a test function to the difference of corresponding systems

$$\begin{aligned} & \partial_t(u_\epsilon^k - u_{\epsilon'}^k) - \operatorname{div}(\mathcal{A}(z, \nabla u_\epsilon^k) - \mathcal{A}(z, \nabla u_{\epsilon'}^k)) \\ &= -\operatorname{div}(|F_\epsilon^k|^{p-2} F_\epsilon^k - |F_{\epsilon'}^k|^{p-2} F_{\epsilon'}^k + a(z)(|F_\epsilon^k|^{q-2} F_\epsilon^k - |F_{\epsilon'}^k|^{q-2} F_{\epsilon'}^k)) \\ & \quad - \operatorname{div}(\epsilon |F_\epsilon^k|^{q-2} F_\epsilon^k - \epsilon' |F_{\epsilon'}^k|^{q-2} F_{\epsilon'}^k) \\ & \quad + \operatorname{div}(b(z)(\epsilon |\nabla u_\epsilon^k|^{q-2} \nabla u_\epsilon^k - \epsilon' |\nabla u_{\epsilon'}^k|^{q-2} \nabla u_{\epsilon'}^k)). \end{aligned}$$

The standard comparison estimate gives a constant $c = c(n, N, p, q, \nu, L)$ such that

$$\begin{aligned} & \iint_{C_R} H(z, |\nabla u_\epsilon^k - \nabla u_{\epsilon'}^k|) dz \\ & \leq c \iint_{C_R} \left(||F_\epsilon^k|^{p-2} F_\epsilon^k - |F_{\epsilon'}^k|^{p-2} F_{\epsilon'}^k|^{\frac{p}{p-1}} + a(z) ||F_\epsilon^k|^{q-2} F_\epsilon^k - |F_{\epsilon'}^k|^{q-2} F_{\epsilon'}^k|^{\frac{q}{q-1}} \right) dz \\ & \quad + c(\epsilon + \epsilon') \iint_{C_R} (|F_\epsilon^k|^q + |F_{\epsilon'}^k|^q + |\nabla u_\epsilon^k|^q + |\nabla u_{\epsilon'}^k|^q) dz. \end{aligned}$$

The last term also converges to zero as $\epsilon, \epsilon' \rightarrow 0^+$, because of $|F_\epsilon^k|, |F_{\epsilon'}^k| \leq k$ and (6.15). Furthermore, the Lebesgue dominated convergence theorem implies that the first term on the right-hand side also converges to zero. Therefore, the claim in (6.17) is proved. Since $k \in \mathbb{N}$ was arbitrary, we obtain a sequence $(u^k)_{k \in \mathbb{N}} \subset C(I_R; L^2(D_R, \mathbb{R}^N)) \cap L^q(I_R; W_0^{1,q}(D_R, \mathbb{R}^N))$ of weak solutions to (6.16). Moreover, from (6.13) and (6.14) we have

$$\begin{aligned} & \sup_{t \in I_R} \int_{D_R} |u^k(x, t)|^2 dx + \iint_{C_R} H(z, |\nabla u^k|) dz + \|\partial_t u^k\|_{L^{q'}(I_R; W^{-1,q'}(D_R, \mathbb{R}^N))} \\ & \leq c \left(\iint_{C_R} H(z, |F|) dz + 1 \right). \end{aligned} \tag{6.18}$$

6.2.3. *Second convergence.* Note that (6.18) guarantees $(\nabla u^k)_{k \geq 1}$ is bounded only in $L^p(C_R, \mathbb{R}^{Nn})$. Since

$$L^p(I_R; W_0^{1,p}(D_R, \mathbb{R}^N)) \Subset L^p(C_R, \mathbb{R}^N) \Subset L^2(C_R, \mathbb{R}^N) \subset L^{q'}(I_R; W^{-1,q'}(D_R, \mathbb{R}^N)),$$

there exist limit functions $u \in C(I_R; L^2(D_R, \mathbb{R}^N)) \cap L^p(I_R; W_0^{1,p}(D_R, \mathbb{R}^N))$ and $v \in L^q(C_R, \mathbb{R}^{Nn})$ such that

$$\begin{aligned} & u^k \rightarrow u \quad \text{strongly in } C(I_R; L^2(D_R, \mathbb{R}^N)) \cap L^p(C_R, \mathbb{R}^N), \\ & u^k \rightarrow u \quad \text{weakly in } L^p(I_R; W_0^{1,p}(D_R, \mathbb{R}^N)), \\ & \partial_t u_\epsilon^k \rightarrow \partial_t u^k \quad \text{weakly in } L^{q'}(I_R; W^{-1,q'}(D_R, \mathbb{R}^N)), \\ & a^{\frac{1}{q}} \nabla u^k \rightarrow v \quad \text{weakly in } L^q(C_R, \mathbb{R}^{Nn}), \end{aligned}$$

as $k \rightarrow \infty$. As in the first convergence result, it is enough to show

$$\iint_{C_R} H(z, |\nabla u^k - \nabla u|) dz \rightarrow 0 \quad \text{as } k \rightarrow \infty. \tag{6.19}$$

We take $u^k - u^{k'} \in C(I_R; L^2(D_R, \mathbb{R}^N)) \cap L^q(I_R; W_0^{1,q}(D_R, \mathbb{R}^N))$ as a test function in

$$\begin{aligned} & \partial_t(u^k - u^{k'}) - \operatorname{div}(\mathcal{A}(z, \nabla u^k) - \mathcal{A}(z, \nabla u^{k'})) \\ &= -\operatorname{div}(|F^k|^{p-2} F^k - |F^{k'}|^{p-2} F^{k'} + a(z)(|F^k|^{q-2} F^k - |F^{k'}|^{q-2} F^{k'})) \end{aligned}$$

in C_R and obtain a constant $c = c(n, N, p, q, \nu)$ such that

$$\begin{aligned} & \iint_{C_R} H(z, |\nabla u^k - \nabla u^{k'}|) dz \\ & \leq c \iint_{C_R} \left(||F^k|^{p-2} F^k - |F^{k'}|^{p-2} F^{k'}|^{\frac{p}{p-1}} + a(z) ||F^k|^{q-2} F^k - |F^{k'}|^{q-2} F^{k'}|^{\frac{q}{q-1}} \right) dz. \end{aligned}$$

As we have $|F^k|, |F^{k'}| \leq |F|$, we conclude that

$$\lim_{k, k' \rightarrow \infty} \iint_{C_R} H(z, |\nabla u^k - \nabla u^{k'}|) dz = 0.$$

This shows that ∇u^k is a Cauchy sequence with respect to the double phase functional and by the weak convergence mentioned above the limit is ∇u . This proves (6.19) and therefore u is a weak solution to (2.6) with $g \equiv 0$.

To conclude the approximation statement of Theorem 2.7, we use (6.17). For every $k \in \mathbb{N}$, we have $\epsilon_k = \epsilon_k(k)$ small enough so that

$$\iint_{C_R} H(z, |\nabla u_{\epsilon_k}^k - \nabla u^k|) dz \leq \frac{1}{k}.$$

It follows from (6.19) that

$$\lim_{k \rightarrow \infty} \iint_{C_R} H(z, |\nabla u_{\epsilon_k}^k - \nabla u|) dz = 0.$$

Letting $u_l = u_{\epsilon_l}^l$ for every $l \in \mathbb{N}$, the proof is completed.

REFERENCES

- [1] E. Acerbi and N. Fusco. Semicontinuity problems in the calculus of variations. *Arch. Rational Mech. Anal.*, 86(2):125–145, 1984.
- [2] E. Acerbi and N. Fusco. An approximation lemma for $W^{1,p}$ functions. In *Material instabilities in continuum mechanics (Edinburgh, 1985–1986)*, Oxford Sci. Publ., pages 1–5. Oxford Univ. Press, New York, 1988.
- [3] K. Adimurthi, S. Byun, and J. Park. End point gradient estimates for quasilinear parabolic equations with variable exponent growth on nonsmooth domains. *Calc. Var. Partial Differential Equations*, 60(4):Paper No. 145, 67, 2021.
- [4] P. Baroni, M. Colombo, and G. Mingione. Harnack inequalities for double phase functionals. *Nonlinear Anal.*, 121:206–222, 2015.
- [5] V. Bögelein. *Regularity results for weak and very weak solutions of higher order parabolic systems*. ProQuest LLC, Ann Arbor, MI, 2007. Thesis (Ph.D.)—Friedrich-Alexander-Universität Erlangen-Nürnberg (Germany).
- [6] V. Bögelein, F. Duzaar, and G. Mingione. The regularity of general parabolic systems with degenerate diffusion. *Mem. Amer. Math. Soc.*, 221(1041):vi+143, 2013.
- [7] M. Bulíček, J. Burczak, and S. Schwarzacher. Well posedness of nonlinear parabolic systems beyond duality. *Ann. Inst. H. Poincaré C Anal. Non Linéaire*, 36(5):1467–1500, 2019.
- [8] I. Chlebicka, P. Gwiazda, and A. Zatorska-Goldstein. Parabolic equation in time and space dependent anisotropic Musielak-Orlicz spaces in absence of Lavrentiev’s phenomenon. *Ann. Inst. H. Poincaré C Anal. Non Linéaire*, 36(5):1431–1465, 2019.
- [9] M. Colombo and G. Mingione. Regularity for double phase variational problems. *Arch. Ration. Mech. Anal.*, 215(2):443–496, 2015.
- [10] M. Colombo and G. Mingione. Calderón-Zygmund estimates and non-uniformly elliptic operators. *J. Funct. Anal.*, 270(4):1416–1478, 2016.
- [11] C. De Filippis and G. Mingione. A borderline case of Calderón-Zygmund estimates for nonuniformly elliptic problems. *St. Petersburg Math. J.*, 31(3):455–477, 2020.
- [12] E. DiBenedetto. *Degenerate parabolic equations*. Universitext. Springer-Verlag, New York, 1993.
- [13] L. Diening, M. Růžička, and J. Wolf. Existence of weak solutions for unsteady motions of generalized Newtonian fluids. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 9(1):1–46, 2010.
- [14] L. Diening, S. Schwarzacher, B. Stroffolini, and A. Verde. Parabolic Lipschitz truncation and caloric approximation. *Calc. Var. Partial Differential Equations*, 56(4):Paper No. 120, 27, 2017.
- [15] L. Esposito, F. Leonetti, and G. Mingione. Sharp regularity for functionals with (p, q) growth. *J. Differential Equations*, 204(1):5–55, 2004.
- [16] W. Kim. Calderón-Zygmund type estimate for the parabolic double-phase system. *arXiv*, 2023.
- [17] W. Kim, J. Kinnunen, and K. Moring. Gradient higher integrability for degenerate parabolic double-phase systems. *arXiv*, 2022.
- [18] J. Kinnunen, J. Lehrbäck, and A. Vähäkangas. *Maximal function methods for Sobolev spaces*, volume 257 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, [2021] ©2021.
- [19] J. Kinnunen and J. L. Lewis. Very weak solutions of parabolic systems of p -Laplacian type. *Ark. Mat.*, 40(1):105–132, 2002.
- [20] J. Simon. Compact sets in the space $L^p(0, T; B)$. *Ann. Mat. Pura Appl. (4)*, 146:65–96, 1987.
- [21] T. Singer. Existence of weak solutions of parabolic systems with p, q -growth. *Manuscripta Math.*, 151(1-2):87–112, 2016.

(Wontae Kim) DEPARTMENT OF MATHEMATICS, AALTO UNIVERSITY, P.O. BOX 11100, 00076 AALTO, FINLAND
Email address: `wontae.kim@aalto.fi`

(Juha Kinnunen) DEPARTMENT OF MATHEMATICS, AALTO UNIVERSITY, P.O. BOX 11100, 00076 AALTO, FINLAND
Email address, Corresponding author: `juha.k.kinnunen@aalto.fi`

(Lauri Särkiö) DEPARTMENT OF MATHEMATICS, AALTO UNIVERSITY, P.O. BOX 11100, 00076 AALTO, FINLAND
Email address: `lauri.sarkio@aalto.fi`