

SUPERCONGRUENCES ARISING FROM RAMANUJAN-SATO SERIES

ANGELICA BABEL, MANAMI ROY, HOLLY SWISHER, BELLA TOBIN, AND FANG-TING TU

ABSTRACT. Recently, the authors with Lea Beneish established a recipe for constructing Ramanujan-Sato series for $1/\pi$, and used this to construct 11 explicit examples of Ramanujan-Sato series arising from modular forms for arithmetic triangle groups of non-compact type. Here, we use work of Chisholm, Deines, Long, Nebe and the third author to prove a general p -adic supercongruence theorem through an explicit connection to CM hypergeometric elliptic curves that provides p -adic analogues of these Ramanujan-Sato series. We further use this theorem to construct explicit examples related to each of our explicit Ramanujan-Sato series examples.

1. INTRODUCTION AND STATEMENT OF RESULTS

Using infinite series to approximate π dates back to Madhava of Sangamagrama circa 1400, and was revolutionized by Ramanujan in 1914 [51] when he stated several rapidly converging series formulas for $1/\pi$ of the form

$$\sum_{k=0}^{\infty} \frac{(\frac{1}{2})_k (\frac{1}{d})_k (\frac{d-1}{d})_k}{k!^3} (ak+1)(\lambda_d)^k = \frac{\delta}{\pi},$$

for $d \in \{2, 3, 4, 6\}$, where λ_d are singular values that correspond to elliptic curves with complex multiplication (CM), a, δ are explicit algebraic numbers, and $(a)_k$ denotes the rising factorial $(a)_k = a(a+1)\cdots(a+k-1)$.¹ Ramanujan's formulas are all examples of special evaluations of hypergeometric functions. They were proved in 1987 by brothers J. and P. Borwein [11] as well as D. and G. Chudnovsky [17], and both approaches rely on the arithmetic of elliptic integrals of the first and second kind, including the Legendre relation at singular values. Since the 1980's series of this type have been at the forefront of algorithms to compute decimal approximations of π . In particular, D. and G. Chudnovsky [17] established the record winning series

$$\frac{426880\sqrt{10005}}{13591409\pi} = \sum_{k=0}^{\infty} \left(\frac{a}{b}k+1\right) \frac{(\frac{1}{2})_k (\frac{1}{6})_k (\frac{5}{6})_k}{(1)_k^3} \left(\frac{-1}{53360^3}\right)^k = {}_4F_3 \left[\begin{matrix} \frac{1}{2} & \frac{1}{6} & \frac{5}{6} & \frac{a+b}{a} \\ & 1 & 1 & \frac{b}{a} \end{matrix} ; \frac{-1}{53360^3} \right],$$

where $a = 545140134$, $b = 13591409$. To date the current record of computing 10^{14} digits of π was set by Emma Haruka Iwao on June 8, 2022 [26] using (1). Generalized series of this form, in which the parameters are related to modular forms, are now called Ramanujan-Sato series after Sato [55], whose formulation in 2002 of a new example of this type renewed interest in the area.

In 1997, van Hamme [62] developed p -adic analogs of several Ramanujan type series for $1/\pi$. Analogs of this type are called Ramanujan type supercongruences, and relate truncated sums of hypergeometric series to values of the p -adic gamma function. Ramanujan type supercongruences have been a rich area of study, see for example [67, 24, 40, 60, 50, 9]. In 2013, Chisholm, Deines, Long, Nebe, and the third author [16] prove a general p -adic analog of Ramanujan type supercongruences modulo p^2 for suitable truncated hypergeometric series arising from CM elliptic curves.

2020 *Mathematics Subject Classification.* Primary 11F23, 11A07, 11G05, 11G15, 33C05, 33C20, 11F11, 44A20 .

Key words and phrases. hypergeometric functions, modular forms, elliptic curves, supercongruences, complex multiplication, Galois representations, L -values.

¹We note that in fact, a similar identity was given even earlier by Bauer [8].

Recently, the authors with Lea Beneish [6] established a recipe for constructing Ramanujan-Sato series for $1/\pi$, and used this to construct 11 explicit examples of Ramanujan-Sato series arising from modular forms for arithmetic triangle groups of non-compact type. In this article, we establish supercongruences related to Ramanujan-Sato series given in [6] motivated by the work of Chisholm et al. [16].

For series of $1/\pi$ that arise via modular forms for a group Γ such as in [13, 15, 6], there is a connection between the series and known p -adic analogs through families of elliptic curves whose monodromy group is Γ . Inspired by Chisholm et al. [16], we make this connection explicit through hypergeometric evaluations. In fact, it motivated us to revisit [16, Theorem 2] using arithmetic properties of elliptic families.

1.1. Notations for main results. We use the following notation for the hypergeometric functions appearing in this work. Given $\alpha \in \mathbb{C}$, $n \in \mathbb{N}$, $\mathbf{b} = (b_1, \dots, b_n) \in (\mathbb{C}^\times)^n$, define

$$(1.1) \quad f_{\alpha, \mathbf{b}}(z) = {}_{n+1}F_n \left[\begin{matrix} \frac{1+\alpha}{\alpha} & b_1 & b_2 & \cdots & b_n \\ \frac{1}{\alpha} & 1 & 1 & \cdots & 1 \end{matrix} ; z \right] = \sum_{k \geq 0} (\alpha k + 1) \frac{(b_1)_k \cdots (b_n)_k}{(1)_k^n} z^k,$$

where the ${}_{n+1}F_n$ is taken to be the limit as $\alpha \rightarrow 0$ when $\alpha = 0$ (yielding an ${}_nF_{n-1}$). We further write $[f_{\alpha, \mathbf{b}}]_N(z)$ to denote the truncation of the series (1.1) at z^N .

When we evaluate this truncation $[f_{\alpha, \mathbf{b}}]_N$ for some λ , i.e., the function f written as a series, truncated at the z^N term, and then evaluated at λ , we write

$$[f_{\alpha, \mathbf{b}}]_N(\lambda) = \sum_{k=0}^N (\alpha k + 1) \frac{(b_1)_k \cdots (b_n)_k}{(1)_k^n} \lambda^k.$$

For $d \in \{2, 3, 4, 6\}$ let $\tilde{E}_d(t)$ denote the following families of elliptic curves parameterized by t :

$$\begin{aligned} \tilde{E}_2(t) &: y^2 = x(x-1)(x-t), \\ \tilde{E}_3(t) &: y^2 + xy + \frac{t}{27}y = x^3, \\ \tilde{E}_4(t) &: y^2 = x(x^2 + x + \frac{t}{4}), \\ \tilde{E}_6(t) &: y^2 + xy = x^3 - \frac{t}{432}. \end{aligned}$$

Remark 1.1. *With this notation we have for $d \in \{2, 3, 4, 6\}$ that $\pi \cdot f_{0, (\frac{1}{d}, \frac{d-1}{d})}(t)$ is a fundamental period for $\tilde{E}_d(t)$.*

A curve E over a number field K has CM if its geometric endomorphism ring $\text{End}(E(\overline{\mathbb{Q}}))$ is an order of an imaginary quadratic field. For t such that $\tilde{E}_d(t)$ has CM, set $\lambda = -4t(t-1)$ and assume $|\lambda| < 1$ for any embedding. Then as discussed in Chisholm et al. [16], there exists unique algebraic numbers α, δ depending on d and λ such that

$$(1.2) \quad \sum_{k \geq 0} (\alpha k + 1) \frac{(\frac{1}{2})_k (\frac{1}{d})_k (\frac{d-1}{d})_k}{k!^3} \lambda^k = \frac{\delta}{\pi}.$$

Moreover, specific CM values of λ as well as the corresponding α, δ can be computed from data of the Borweins [11]. Moreover, (1.2) can also be rewritten as the following product by Lemma 2.8,

$$f_{0, (\frac{1}{d}, \frac{d-1}{d})}(t) \cdot f_{a_2, (\frac{1}{d}, \frac{d-1}{d})}(t) = \frac{\delta}{\pi},$$

where $t = \frac{1-\sqrt{1-\lambda}}{2}$, $a_2 = \frac{2\alpha(1-t)}{1-2t}$, and δ is a unique algebraic number depending on t .

Recently, the authors with Beneish [6] establish a general theorem which provides a recipe for constructing Ramanujan-Sato series, and provide 11 explicit examples. Each of these examples can be written in the form

$$(1.3) \quad \frac{\delta}{\pi} = \sum_{k \geq 0} (\alpha k + 1) A_k \lambda^k = f_{0, \mathbf{b}}(\lambda) \cdot f_{2\alpha, \mathbf{b}}(\lambda),$$

for parameters $\delta, \alpha, \mathbf{b}, \lambda$, where A_k is defined by $\sum_{k \geq 0} A_k z^k = f_{0, \mathbf{b}}(z)^2$ (see Lemma 2.7). The 11 explicit examples of Ramanujan-Sato series given in [6], rewritten from this perspective, are described by the data in Table 1, which uses some notation from Corollary 1.2 below.

TABLE 1. Ramanujan-Sato Example Data

Ex. in [6]	δ	N	α_N	\mathbf{b}	λ	μ	d
4.2	$2\sqrt{3}$	3	$3\sqrt{2} + 4$	$\frac{1}{4}, \frac{3}{4}$	$\frac{1}{6}(3 - 2\sqrt{2})$	$\frac{1}{6}(3 - 2\sqrt{2})$	4
4.3	$\frac{9}{4}\sqrt{2}$	5	$\frac{1}{4}(20 + 9\sqrt{5})$	$\frac{1}{4}, \frac{3}{4}$	$\frac{1}{18}(9 - 4\sqrt{5})$	$\frac{1}{18}(9 - 4\sqrt{5})$	4
4.4	$\sqrt{3}$	3	4	$\frac{1}{4}, \frac{3}{4}$	$-\frac{1}{3}$	$-\frac{1}{3}$	4
4.5	$\frac{8}{3}$	5	$\frac{1}{3}(10 + 4\sqrt{5})$	$\frac{1}{4}, \frac{3}{4}$	$\frac{1}{4}(2 - \sqrt{5})$	$\frac{1}{4}(2 - \sqrt{5})$	4
4.6	$3\sqrt{3}$	2	$3(\sqrt{2} + 1)$	$\frac{1}{3}, \frac{2}{3}$	$\frac{1}{4}(2 - \sqrt{2})$	$\frac{1}{4}(2 - \sqrt{2})$	3
4.7	$2\sqrt{2} + 3$	2	$4 + 2\sqrt{2}$	$\frac{1}{2}, \frac{1}{2}$	$3 - 2\sqrt{2}$	$3 - 2\sqrt{2}$	2
5.2	$\frac{5^2 \Gamma(\frac{1}{8})^2 \Gamma(\frac{3}{8})^2}{29^{9/2} \cdot 3\pi^3}$	2	$\frac{14}{3}$	$\frac{1}{2}, \frac{1}{6}, \frac{5}{6}$	$\frac{27}{125}$	$\frac{1}{2} - \frac{7}{5\sqrt{10}}$	6
5.3	$\frac{5^2 \sqrt{3} \Gamma(\frac{1}{3})^6}{2^{17/3} \pi^4}$	3	$\frac{11}{2}$	$\frac{1}{2}, \frac{1}{6}, \frac{5}{6}$	$\frac{4}{125}$	$\frac{1}{2} - \frac{11}{10\sqrt{5}}$	6
5.4	$\frac{\Gamma(\frac{1}{24})\Gamma(\frac{5}{24})\Gamma(\frac{7}{24})\Gamma(\frac{11}{24})}{2^{5/2} \pi^3}$	3	4	$\frac{1}{2}, \frac{1}{4}, \frac{3}{4}$	$\frac{1}{9}$	$\frac{1}{2} - \frac{\sqrt{2}}{3}$	4
5.5	$\frac{3^2 \Gamma(\frac{1}{24})\Gamma(\frac{5}{24})\Gamma(\frac{7}{24})\Gamma(\frac{11}{24})}{2^5 \pi^3}$	2	3	$\frac{1}{2}, \frac{1}{3}, \frac{2}{3}$	$\frac{1}{2}$	$\frac{1}{2} - \frac{1}{2\sqrt{2}}$	3
5.6	$\frac{3^2 \cdot 5 \Gamma(\frac{1}{15})\Gamma(\frac{2}{15})\Gamma(\frac{4}{15})\Gamma(\frac{8}{15})}{2^8 \pi^3}$	5	$\frac{33}{8}$	$\frac{1}{2}, \frac{1}{3}, \frac{2}{3}$	$\frac{4}{125}$	$\frac{1}{2} - \frac{11}{10\sqrt{5}}$	3

Corollary 1.2 (Corollary of [6, Theorem 1.1]). *Let Γ be a discrete subgroup of $\mathrm{SL}_2(\mathbb{R})$ commensurable with $\mathrm{SL}_2(\mathbb{Z})$ such that $(\begin{smallmatrix} 1 & \\ & 1 \end{smallmatrix}) \in \Gamma$, and let $X(\tau)$ be a Hauptmodul of Γ . Let $Z(\tau)$ be a weight- k modular form for Γ such that $\frac{1}{2\pi i} \frac{dX}{d\tau} = U(\tau)X(\tau)Z(\tau)$ and when $\tau \in D$, a domain of \mathbb{H} , $Z(\tau) = f_{0, \mathbf{b}}(X(\tau))^2$ for $\mathbf{b} \in \mathbb{Q}^n$. Further assume there exist $\varepsilon \in \mathbb{C}$ and $\gamma = \begin{pmatrix} a & b \\ c & -a \end{pmatrix} \in \mathrm{SL}_2(\mathbb{R})$ such that*

$$(Z|_k \gamma)(\tau) = \varepsilon Z(\tau).$$

Set $M_N(\tau) := Z(\tau)/Z(N\tau)$ for $N \in \mathbb{N}$ satisfying $\frac{a}{c}(1 - N) \in \mathbb{Z}$, and let $\tau_0 = \frac{a}{c} + \frac{i}{c\sqrt{N}}$. Set $v(\tau_0)$ as the special value $v(\tau_0) = U(\tau_0)X(\tau_0) \left(\frac{dM_N}{dX} \right) \Big|_{X=X(\tau_0)}$.

Then if $\gamma\tau_0 = \frac{a}{c} + \frac{i\sqrt{N}}{c} \in D$, we have²

$$\frac{\delta_N}{\pi} = f_{0, \mathbf{b}}(X(\gamma\tau_0)) \cdot f_{2\alpha_N, \mathbf{b}}(X(\gamma\tau_0)),$$

where $\delta_N = \frac{ck\sqrt{N}}{2v(\tau_0)}$, and $\alpha_N = \frac{2NU(\gamma\tau_0)}{v(\tau_0)}$.

²For the case where $\tau_0 \in D$ a similar result can be stated, but we omit this here.

Remark 1.3. We note that the form given in [6, Ex. 4.2 - 4.6] is different than (1.3), but using Corollary 1.2 we can rewrite them in the form of (1.3). For instance for [6, Ex. 4.2] we do this by setting $Z(\tau) = f_{0,\mathbf{b}}(t_2(\tau)/(t_2(\tau) - 1))^2$ with $\mathbf{b} = (1/4, 3/4)$ and $X(\tau) = \frac{t_2(\tau)}{t_2(\tau) - 1}$, using the notation in [6], which yields that $U(\tau) = \frac{1}{1 - t_2(\tau)} = 1 - X(\tau)$ and

$$\frac{dM_N}{dX} = N \left(\frac{d\left(\frac{dX}{dY}\right)}{dX} \cdot \frac{Y(1-Y)}{X(1-X)} + \frac{(1-2Y)}{X(1-X)} - \frac{dX}{dY} \cdot \frac{Y(1-Y)(1-2X)}{X^2(1-X)^2} \right).$$

This works similarly for [6, Ex. 4.6] with $\mathbf{b} = (1/3, 2/3)$ and $t_3(\tau)$ in the place of $t_2(\tau)$. The form given in [6, Ex. 4.7] matches (1.3), so we apply Corollary 1.2 with $X(\tau) = t_\infty(\tau)$ and $U(\tau) = 1 - X(\tau)$.

1.2. Statement of main results. We can now state our main theorem.

Theorem 1.4. Let $d \in \{2, 3, 4, 6\}$ and set $\mathbf{b} = (\frac{1}{d}, \frac{d-1}{d})$, or $\mathbf{b} = (\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})$. Suppose $X(\tau)$ and $Z(\tau) = f_{0,\mathbf{b}}(X(\tau))^2$ satisfy all of the conditions in Corollary 1.2 with $\gamma = \begin{pmatrix} a & b \\ c & -a \end{pmatrix}$ and $\tau_0 = \frac{a}{c} + \frac{i}{c\sqrt{N}}$, and furthermore if $\lambda = X(\gamma\tau_0)$, then $f_{0,\mathbf{b}}(\lambda) = f_{0,(\frac{1}{d}, \frac{d-1}{d})}(\mu)^r$ for some r , where $\mu = \lambda$ when $\mathbf{b} = (\frac{1}{d}, \frac{d-1}{d})$, and $\mu = \frac{1 \pm \sqrt{1-\lambda}}{2}$ when $\mathbf{b} = (\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})$.

Assume that $\mathbb{Q}(\lambda)$ is a totally real field, $|\lambda|, |\mu| < 1$, and $\tilde{E}_d(\mu)$ has CM (equivalently τ_0 is a CM point). Let p be a prime such that $X(\tau_0), \lambda, \alpha_N$ can be embedded into \mathbb{Z}_p^\times , and p is unramified in $\mathbb{Q}(\mu)$. Suppose that $\tilde{E}_d(\mu)$ has good reduction modulo p .

Then, it follows that if $\mathbf{b} = (\frac{1}{d}, \frac{d-1}{d})$,

$$[f_{0,(\frac{1}{d}, \frac{d-1}{d})} \cdot f_{2\alpha_N, (\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(X(\gamma\tau_0)) \equiv \begin{cases} p & \text{if } \left(\frac{-c^2N}{p}\right) = 1 \\ -p & \text{otherwise} \end{cases} \pmod{p^2},$$

if $\mathbf{b} = (\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})$,

$$[f_{0,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})} \cdot f_{2\alpha_N, (\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}]_{p-1}(X(\gamma\tau_0)) \equiv \begin{cases} pu_p^2 & \text{if } \left(\frac{-c^2N}{p}\right) = 1 \text{ and } \left(\frac{1-\lambda}{p}\right) = 1 \\ -\left(\frac{k_d}{p}\right) pu_p^2 & \text{if } \left(\frac{-c^2N}{p}\right) = 1 \text{ and } \left(\frac{1-\lambda}{p}\right) = -1 \\ 0 & \text{otherwise} \end{cases} \pmod{p^2},$$

where u_p is a fixed p -adic unit root of the geometric Frobenius at p acting on the first cohomology of the elliptic curve $\tilde{E}_d(\mu)$, and $k_2 = -1$, $k_3 = -3$, $k_4 = -2$.

Remark 1.5. We note that work of Beukers (see Theorem 2.5) as well as computational observations indicate that the series obtained from Theorem 1.4 holds modulo p^3 in some cases but not for others. See Examples 5.8 and 5.9 and Section 7 for further discussion.

In addition to Theorem 1.4, which is a general statement, we provide we provide specific examples using the Ramanujan-Sato series given in [6] (as in Table 1).

Hypergeometric functions, such as those appearing in Ramanujan-Sato series as well as their p -adic truncations and also those over finite fields, encode abundant information on the arithmetic invariants of hypergeometric motives such as their periods, Galois representations, and the p -adic unit roots of the local L -functions due to Dwork's unit roots theory. They are also closely connected to modular forms through the modularity theorem and the differential equations satisfied by certain modular forms (see [14, 22, 52, 64, 15], for example).

Using Theorem 1.4, we prove a few results in Section 7 connecting the hypergeometric functions appearing in the series in [6, Ex. 5.2 - 5.6] with Galois representations and special L -values of modular forms of weight 3. For example, we obtain the following corollary.

Corollary 1.6. *For primes $p \geq 7$*

$$\begin{aligned} \left[f_{0,(\frac{1}{2},\frac{1}{6},\frac{5}{6})} \right]_{p-1} \left(\frac{27}{125} \right) &\equiv a_p(f_{800.3.g.a}) \pmod{p^2} \\ \left[f_{0,(\frac{1}{2},\frac{1}{6},\frac{5}{6})} \right]_{p-1} \left(\frac{4}{125} \right) &\equiv a_p(f_{300.3.g.b}) \pmod{p^2} \\ \left[f_{0,(\frac{1}{2},\frac{1}{4},\frac{3}{4})} \right]_{p-1} \left(\frac{1}{9} \right) &\equiv a_p(f_{24.3.h.a}) \pmod{p^2} \\ \left[f_{0,(\frac{1}{2},\frac{1}{3},\frac{2}{3})} \right]_{p-1} \left(\frac{1}{2} \right) &\equiv a_p(f_{24.3.h.b}) \pmod{p^2} \\ \left[f_{0,(\frac{1}{2},\frac{1}{3},\frac{2}{3})} \right]_{p-1} \left(\frac{4}{125} \right) &\equiv a_p(f_{15.3.d.b}) \pmod{p^2}, \end{aligned}$$

where the subscripts on the modular forms f above are their LMFDB labels [38] and $a_p(f)$ is the p th Fourier coefficient of f .

This result follows from the more general Corollary 7.1, proved in Section 7, that uses modularity of CM elliptic curves and L -functions. While Corollary 1.6 gives values of the hypergeometric functions in terms of Fourier coefficients of modular forms, we further obtain in Section 7.2 special values in terms of periods Ω_K and special L -values of modular forms.

1.3. Outline of the paper. The rest of this paper is organized as follows. In Section 2, we provide preliminary results used throughout the rest of the paper. In Section 3, we describe the connection between the hypergeometric evaluations for $1/\pi$ given by Beneish and the authors in [6] and their p -adic counterparts provided in Theorem 1.4 through the study of hypergeometric elliptic curves and modular forms associated to certain triangle groups. We also provide several lemmas needed for the proof of Theorem 1.4. In Section 4, we prove Theorem 1.4. In Section 5, we give 11 examples of Theorem 1.4, which stem from applications of results in [6, Thm. 2]. In Section 6, we give 4 different examples of supercongruences by directly applying Theorem 2.1 to the series in [6, Ex. 4.2 - 4.5]. In Section 7, we describe connections between hypergeometric functions of the form $f_{0,(\frac{1}{2},\frac{1}{d},\frac{d-1}{d})}$ and weight 3 modular forms, modular Galois representations, and special L -values of weight 3 modular forms. We further discuss a modulo p^3 supercongruence conjecture for $f_{0,(\frac{1}{2},\frac{1}{d},\frac{d-1}{d})}$. In the Section 8 Appendix, we provide an explanation of Theorem 2.3 through the arithmetic properties of hypergeometric elliptic families, and further record some parallel results in terms of finite hypergeometric functions introduced by Beukers, Cohen, and Mellit [10], which allow us to see the supercongruence in Theorem 2.3 from a character sum perspective.

1.4. Acknowledgements. This material is based upon work supported by the National Science Foundation grant DMS-1928930, while the authors were in residence at the Simons Laufer Mathematical Sciences Research Institute in Berkeley, California, during the summer of 2023. The third author is supported by NSF grant DMS-2101906, the fifth author is supported by NSF grant DMS-230253, and the fourth author is partially supported by an AMS-Simons travel grant. The fifth author was hosted at Oregon State University during the fall of 2023 while working on this project. The authors would also like to thanks Frits Beukers and Ling Long for their helpful discussions.

2. PRELIMINARIES

The following is the main theorem of Chisholm, et al. [16], written using the notation introduced in Section 1.

Theorem 2.1 (Chisholm, et al. [16, Theorem 1]). *For $d \in \{2, 3, 4, 6\}$, let $\lambda \in \overline{\mathbb{Q}}$ such that $\mathbb{Q}(\lambda)$ is totally real, the elliptic curve $\tilde{E}_d(\frac{1 \pm \sqrt{1-\lambda}}{2})$ has CM, and $|\lambda| < 1$ for an embedding of λ to \mathbb{C} . For each prime p that is unramified in $\mathbb{Q}(\sqrt{1-\lambda})$ and coprime to the discriminant of $\tilde{E}_d(\frac{1-\sqrt{1-\lambda}}{2})$ such that α, λ give a series of the form (1.2) and can be embedded in \mathbb{Z}_p^\times (and we fix such embeddings), then*

$$[f_{\alpha, (\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\lambda) \equiv \text{sgn} \cdot \left(\frac{1-\lambda}{p} \right) \cdot p \pmod{p^2},$$

where $\left(\frac{1-\lambda}{p} \right)$ is interpreted as a Legendre symbol over \mathbb{Z}_p , which is possible since λ can be embedded in \mathbb{Z}_p , and $\text{sgn} = \pm 1$, equaling 1 (or -1) if and only if $\tilde{E}_d(\frac{1 \pm \sqrt{1-\lambda}}{2})$ is ordinary (or supersingular) at p .

Remark 2.2. *We note that as discussed in [16, Remarks 1 and 2] this congruence also holds when $|\lambda| \geq 1$ and has been conjectured by Zudilin to hold modulo p^3 [24, 25, 67]. See for example [9, 46, 67] for some confirmed cases.*

A companion theorem to Theorem 2.1 is the following

Theorem 2.3. (Revision of [16, Theorem 2]) *Assume the notation and assumptions in Theorem 2.1. Then,*

$$[f_{0, (\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\lambda) \equiv L \pmod{p^2},$$

where

$$L = \begin{cases} 0 & \text{if } \tilde{E}_d(\frac{1-\sqrt{1-\lambda}}{2}) \text{ is supersingular at } p \\ u_p^2 & \text{if } \tilde{E}_d(\frac{1-\sqrt{1-\lambda}}{2}) \text{ is ordinary at } p \text{ and } \left(\frac{1-\lambda}{p} \right) = 1, \\ \binom{k_d}{p} u_p^2 & \text{if } \tilde{E}_d(\frac{1-\sqrt{1-\lambda}}{2}) \text{ is ordinary at } p \text{ and } \left(\frac{1-\lambda}{p} \right) = -1, \end{cases}$$

and in the ordinary case, u_p is a fixed p -adic unit root of the geometric Frobenius at p acting on the first cohomology of the elliptic curve $\tilde{E}_d(\frac{1-\sqrt{1-\lambda}}{2})$ and $k_2 = k_6 = -1$, $k_3 = -3$, $k_4 = -2$.

We note that the statement of [16, Thm. 2] is missing the case when $\left(\frac{1-\lambda}{p} \right) = -1$. A more detailed proof of this missing case is supplied in the Appendix §8.1, however we briefly offer two approaches here. The key observation is the following congruence

$$(2.1) \quad [f_{0, (\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(v_+) [f_{0, (\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(v_-) \equiv u_{p,+} u_{p,-} \pmod{p\mathbb{Z}_p[\sqrt{1-\lambda}]},$$

where $v_- := \frac{1-\sqrt{1-\lambda}}{2}$, $v_+ := \frac{1+\sqrt{1-\lambda}}{2}$, and $u_{p,+}$ (resp. $u_{p,-}$) is a fixed p -adic unit root of the geometric Frobenius associated to $\tilde{E}_d(v_+)$ (resp. $\tilde{E}_d(v_-)$) while fixing an embedding of $\sqrt{1-\lambda}$. One way (2.1) can be obtained is from the perspective of hypergeometric character sums, as discussed in §8.2 (see Theorem 8.8 and Remark 8.9). Another approach is via the commutative formal group law, where (2.1) can be induced by [16, Lem. 12], [59, Thm. A.8(v)], and [30, Thm. 6.1].³ From these perspectives,

$$[f_{0, (\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(v_+) \equiv \varepsilon u_p \pmod{p\mathbb{Z}_p[\sqrt{1-\lambda}]},$$

³Here $P_{1,\rho}(T)$ is given by the zeta function over finite field \mathbb{F}_q with $q = p$ when $\left(\frac{1-\lambda}{p} \right) = 1$ and $q = p^2$ when $\left(\frac{1-\lambda}{p} \right) = -1$.

for a root of unity ε which can be determined by the CM theory of elliptic curves and elliptic curves over local fields (see [58, 57, 59]).

Combined with the truncated hypergeometric Clausen formula in Lemma 8.2, (2.1) will give the congruence in Theorem 2.3 modulo p . Then, similar to the discussion in [16], based on the work of Coster and Van-Hame [19] on CM elliptic curves, Katz [31], and Long-Ramakrishna [40], one obtains the supercongruence modulo p^2 .

Remark 2.4. *We note that computationally Theorem 2.3 appears to hold modulo p^3 for ordinary primes, however for supersingular primes the result only holds modulo p^2 generically. Some progress on this have been made, see Long-Ramakrishna [40] and Beukers [9] for example. In particular, [9, Theorem 1.27] gives the cases of $d = 6$ through the properties of the corresponding modular forms.*

We now state a modulo p^3 theorem in the case when $d = 6$ that follows from recent work of Beukers [9]. To set some notation, let D be a positive integer such that $D \equiv 0$ or $3 \pmod{4}$, and define

$$(2.2) \quad \omega_D = \begin{cases} \frac{\sqrt{-D}}{2} & \text{if } D \equiv 0 \pmod{4}, \\ \frac{1+\sqrt{-D}}{2} & \text{if } D \equiv 3 \pmod{4}. \end{cases}$$

Theorem 2.5. *(Theorems 2.1, 2.3, and [9, Theorem 1.27]) Assume the notation and assumptions in Theorem 2.1 and (2.2). Suppose $\lambda = 1728/j(\omega_D)$. Then, for an ordinary prime p ,*

$$[f_{\alpha, (\frac{1}{2}, \frac{1}{6}, \frac{5}{6})}]_{p-1}(\lambda) \equiv \left(\frac{1-\lambda}{p} \right) \cdot p \pmod{p^3},$$

and

$$[f_{0, (\frac{1}{2}, \frac{1}{6}, \frac{5}{6})}]_{p-1}(\lambda) \equiv \begin{cases} u_p^2 & \text{if } \left(\frac{1-\lambda}{p} \right) = 1, \\ \left(\frac{-1}{p} \right) u_p^2 & \text{if } \left(\frac{1-\lambda}{p} \right) = -1, \end{cases} \pmod{p^3},$$

where u_p is a fixed p -adic unit root of the geometric Frobenius at p acting on the first cohomology of the elliptic curve $\tilde{E}_d(\frac{1-\sqrt{1-\lambda}}{2})$.

We next state the following theorem of Deuring [34, Ch. 13 Thm. 12] which provides a test for whether E is ordinary/supersingular at p depending on the decomposition of p in k .

Theorem 2.6 (Deuring). *Let E be an elliptic curve over a number field F with CM, i.e., $\text{End}(E(\overline{\mathbb{Q}}))$ is an order in an imaginary quadratic field K . Suppose \mathfrak{P} is a place above a prime p where E has good reduction modulo \mathfrak{P} . Then E is supersingular at p if and only if p either ramifies or is inert in K , i.e., E is ordinary modulo p if and only if p splits in K .*

2.1. Power series and derivatives. In general for a power series $f(z) = \sum_{k \geq 0} c_k z^k \in \mathbb{C}[[z]]$, we have $z \frac{d}{dz} \sum_{k \geq 0} c_k z^k = \sum_{k \geq 0} k c_k z^k$, and thus

$$(2.3) \quad \sum_{k \geq 0} (ak + 1) c_k z^k = f(z) + az \frac{d}{dz} f(z).$$

In particular, we observe that

$$(2.4) \quad f_{a, \mathbf{b}}(z) = f_{0, \mathbf{b}}(z) + az \frac{d}{dz} f_{0, \mathbf{b}}(z).$$

We further obtain the following lemma.

Lemma 2.7. *Define $A_k \in \mathbb{C}$ by $f_{0, \mathbf{b}}(z)^2 = \sum_{k \geq 0} A_k z^k$. Then,*

$$\sum_{k \geq 0} (ak + 1) A_k z^k = f_{0, \mathbf{b}}(z) \cdot f_{2a, \mathbf{b}}(z).$$

Proof. Applying (2.3) with the chain rule, and (2.4), we obtain

$$\sum_{k \geq 0} (ak + 1) A_k z^k = f_{0,\mathbf{b}}(z)^2 + 2az f_{0,\mathbf{b}}(z) \frac{d}{dz} f_{0,\mathbf{b}}(z) = f_{0,\mathbf{b}}(z) \cdot f_{2a,\mathbf{b}}(z).$$

□

In [16, equation (3.5)], the authors show that for λ such that both sides converge,

$$(2.5) \quad f_{0,(\frac{1}{2},c,1-c)}(\lambda) = f_{0,(c,1-c)} \left(\frac{1 \pm \sqrt{1-\lambda}}{2} \right)^2.$$

Clausen's identity (2.5) enables us to show the following, which will be useful in Section 3.2.

Lemma 2.8. *Set $t = \frac{1-\sqrt{1-\lambda}}{2}$, so that $\lambda = -4t(t-1)$. Then, when both sides converge,*

$$f_{a_1,(\frac{1}{2},c,1-c)}(\lambda) = f_{0,(c,1-c)}(t) \cdot f_{a_2,(c,1-c)}(t),$$

where $a_2 = \frac{2a_1(1-t)}{1-2t}$.

Proof. By (2.4) and (2.5), we have that

$$f_{a_1,(\frac{1}{2},c,1-c)}(\lambda) = f_{0,(c,1-c)}(t)^2 + a_1 \lambda \frac{d}{d\lambda} f_{0,(c,1-c)}(t)^2.$$

Thus differentiating and applying (2.3) gives

$$\begin{aligned} f_{a_1,(\frac{1}{2},c,1-c)}(\lambda) &= f_{0,(c,1-c)}(t)^2 + 2a_1 f_{0,(c,1-c)}(t) \cdot \lambda \frac{dt}{d\lambda} \cdot \frac{d}{dt} f_{0,(c,1-c)}(t) \\ &= f_{0,(c,1-c)}(t)^2 + \frac{2a_1(1-t)}{1-2t} f_{0,(c,1-c)}(t) \cdot t \frac{d}{dt} f_{0,(c,1-c)}(t) \\ &= f_{0,(c,1-c)}(t) f_{a_2,(c,1-c)}(t), \end{aligned}$$

where $a_2 = \frac{2a_1(1-t)}{1-2t}$ as desired. □

2.2. Truncating hypergeometric products modulo p^2 and p^3 . We use the following notation. For $d \in \{2, 3, 4, 6\}$, and $p \geq 3$ prime, define positive integers r_d and s_d by

$$r_d := \begin{cases} \frac{p-1}{d} & \text{if } p \equiv 1 \pmod{d} \\ \frac{p-(d-1)}{d} & \text{if } p \equiv -1 \pmod{d} \end{cases}$$

$$s_d := \begin{cases} \frac{(d-1)p-(d-1)}{d} & \text{if } p \equiv 1 \pmod{d} \\ \frac{(d-1)p-1}{d} & \text{if } p \equiv -1 \pmod{d}. \end{cases}$$

We first observe that for p an odd prime and k a nonnegative integer, $\left(\frac{1}{2}\right)_k \equiv 0 \pmod{p}$ if and only if $k \geq \frac{p+1}{2}$. Thus when $\frac{p+1}{2} \leq k \leq p-1$,

$$(2.6) \quad \frac{\left(\frac{1}{2}\right)_k}{(1)_k} \equiv 0 \pmod{p}.$$

The following lemma gives further conditions for p -divisibility of our hypergeometric coefficients.

Lemma 2.9 (Chisholm, et al. [16]). *Let $d \in \{2, 3, 4, 6\}$ and p an odd prime coprime to d . Then⁴ when $r_d < k \leq s_d$,*

$$\frac{\left(\frac{1}{d}\right)_k \left(\frac{d-1}{d}\right)_k}{(1)_k^2} \equiv 0 \pmod{p} \quad \text{and} \quad \frac{\left(\frac{1}{d}\right)_k \left(\frac{d-1}{d}\right)_k}{(1)_k^2} \not\equiv 0 \pmod{p^2},$$

⁴We have reworded [16, Lemma 17] in terms of r_d and s_d .

and when $s_d < k \leq p-1$,

$$\frac{\left(\frac{1}{d}\right)_k \left(\frac{d-1}{d}\right)_k}{(1)_k^2} \equiv 0 \pmod{p^2}.$$

In general, we observe that $(\sum_{k=0}^{\infty} a_k z^k)(\sum_{k=0}^{\infty} b_k z^k) = \sum_{k=0}^{\infty} (\sum_{j=0}^k a_j b_{k-j}) z^k = \sum_{k=0}^{\infty} c_k z^k$. So truncating this series at $p-1$ gives

$$\sum_{k=0}^{p-1} c_k z^k = \sum_{k=0}^{p-1} \sum_{j=0}^k a_j b_{k-j} z^k.$$

Thus when we truncate individually and compare, we have

$$\begin{aligned} (2.7) \quad \left(\sum_{k=0}^{p-1} a_k z^k\right) \left(\sum_{k=0}^{p-1} b_k z^k\right) &= \sum_{k=0}^{2(p-1)} \sum_{\substack{0 \leq j \leq p-1 \\ 0 \leq k-j \leq p-1}} a_j b_{k-j} z^k \\ &= \sum_{k=0}^{p-1} \sum_{j=0}^k a_j b_{k-j} z^k + \sum_{k=p}^{2(p-1)} \sum_{j=k-(p-1)}^{p-1} a_j b_{k-j} z^k = \sum_{k=0}^{p-1} c_k z^k + \sum_{k=p}^{2(p-1)} \sum_{j=k-(p-1)}^{p-1} a_j b_{k-j} z^k \\ &= \sum_{k=0}^{p-1} c_k z^k + \sum_{k=1}^{p-1} \sum_{j=k}^{p-1} a_j b_{(p-1)-(j-k)} z^{(p-1)+k} = \sum_{k=0}^{p-1} c_k z^k + \sum_{k=1}^{p-1} \sum_{j=0}^{p-1-k} a_{k+j} b_{(p-1)-j} z^{(p-1)+k}. \end{aligned}$$

The following lemma is necessary for our proof of Theorem 1.4.

Lemma 2.10. *Using the notation in Theorem 1.4, we have that if $p \geq 5$ prime and $\alpha_N \in \overline{\mathbb{Q}}$ such that α_N can be embedded into \mathbb{Z}_p (and fixing such an embedding),*

$$[f_{0,\mathbf{b}} \cdot f_{2\alpha_N,\mathbf{b}}]_{p-1}(z) \equiv [f_{0,\mathbf{b}}]_{p-1}(z) \cdot [f_{2\alpha_N,\mathbf{b}}]_{p-1}(z) \begin{cases} \pmod{p^2} & \text{if } \mathbf{b} = \left(\frac{1}{d}, \frac{d-1}{d}\right), \\ \pmod{p^3} & \text{if } \mathbf{b} = \left(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d}\right). \end{cases}$$

Proof. Let $d \in \{2, 3, 4, 6\}$ and set $\mathbf{b} = \left(\frac{1}{d}, \frac{d-1}{d}\right)$, or $\mathbf{b} = \left(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d}\right)$. Write $f_{0,\mathbf{b}}(z) = \sum_{k \geq 0} a_k z^k$ and $f_{2\alpha_N,\mathbf{b}}(z) = \sum_{k \geq 0} b_k z^k$. Then by (2.7), we have

$$[f_{0,\mathbf{b}} \cdot f_{2\alpha_N,\mathbf{b}}]_{p-1}(z) = [f_{0,\mathbf{b}}]_{p-1}(z) \cdot [f_{2\alpha_N,\mathbf{b}}]_{p-1}(z) - \sum_{k=1}^{p-1} \sum_{j=0}^{p-1-k} a_{k+j} b_{(p-1)-j} z^{(p-1)+k}.$$

It follows from (1.1) that $b_k = (2\alpha_N k + 1)a_k$. Since α_N can be embedded into \mathbb{Z}_p it suffices to show for $1 \leq k \leq p-1$ and $0 \leq j \leq p-1-k$ that $a_{k+j} a_{(p-1)-j}$ is congruent to 0 modulo our desired power of p .

Using the notation in Lemma 2.9, we consider when $p-1-j$ is in the intervals $(0, r_d]$, $(r_d, s_d]$, and $(s_d, p-1]$ separately.

First, we consider the case when $\mathbf{b} = \left(\frac{1}{d}, \frac{d-1}{d}\right)$, so that $a_k = \frac{\left(\frac{1}{d}\right)_k \left(\frac{d-1}{d}\right)_k}{(1)_k^2}$. When $p-1-j \in (s_d, p-1]$, Lemma 2.9 directly gives that $a_{p-1-j} \equiv 0 \pmod{p^2}$. When $p-1-j \in (0, r_d]$, then $s_d = p-1-r_d \leq j < p-1$, which implies that $s_d < j+k \leq p-1$, so Lemma 2.9 gives that $a_{k+j} \equiv 0 \pmod{p^2}$. Now suppose $p-1-j \in (r_d, s_d]$. Then, we have

$$r_d = p-1-s_d \leq j < p-1-r_d = s_d.$$

Thus, we have that $p-1-j \in (r_d, s_d]$ and also that $j+k \in (r_d, p-1]$, so Lemma 2.9 gives that $a_{p-1-j} \equiv 0 \pmod{p}$ and $a_{k+j} \equiv 0 \pmod{p}$.

Next, we consider the case when $\mathbf{b} = (\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})$, so that $a_k = \frac{(\frac{1}{2})_k (\frac{1}{d})_k (\frac{d-1}{d})_k}{(1)_k^3}$. From the previous case, we have seen that $\frac{(\frac{1}{d})_k (\frac{d-1}{d})_k}{(1)_k^2} \equiv 0 \pmod{p^2}$ for $p-1-j \in (0, p-1]$. Thus it suffices to show that when $p-1-j \in (0, p-1]$, either $\frac{(\frac{1}{2})_{p-1-j}}{(1)_{p-1-j}} \equiv 0 \pmod{p}$ or $\frac{(\frac{1}{2})_{k+j}}{(1)_{k+j}} \equiv 0 \pmod{p}$. If $p-1-j \in (\frac{p-1}{2}, p-1]$, then from (2.6) we have $\frac{(\frac{1}{2})_{p-1-j}}{(1)_{p-1-j}} \equiv 0 \pmod{p}$. If $p-1-j \in (0, \frac{p-1}{2}]$, then $\frac{p-1}{2} \leq j < p-1$, so we have that $\frac{p-1}{2} < k+j \leq p-1$, and so from (2.6) we have $\frac{(\frac{1}{2})_{k+j}}{(1)_{k+j}} \equiv 0 \pmod{p}$. □

3. HYPERGEOMETRIC ELLIPTIC CURVES AND THE TRIANGLE MODULAR CURVES

In this section, we recall the modular curves arising from certain triangle groups. The connection between the hypergeometric evaluations for $1/\pi$ in [6] and their p -adic version in this paper arises from the moduli interpretation of such hypergeometric elliptic curves.

In particular, a group $\Gamma \in \{\Gamma(2), \Gamma_0(3), \Gamma_0(2)\}$ is a monodromy group for a family of elliptic curves as given in Table 2, where t is the Hauptmodul of Γ taking values at 0, 1 at two of the cusps and having a pole at the elliptic point of the other cusp, and $\tilde{E}_d(t)$ is a family of elliptic curves over points in X_Γ for $t \neq 0, 1, \infty$. In Table 2 we also provide the element W lying in the normalizer of the group Γ that switches the elements t and $1-t$.

TABLE 2. Families of hypergeometric elliptic curves

$\tilde{E}_d(t)$	\mathbf{b}	Γ	W
$\tilde{E}_2(t) : y^2 = x(x-1)(x-t)$	$(\frac{1}{2}, \frac{1}{2})$	$\Gamma(2)$	$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$
$\tilde{E}_3(t) : y^2 + xy + \frac{t}{27}y = x^3$	$(\frac{1}{3}, \frac{2}{3})$	$\Gamma_0(3)$	$\frac{1}{\sqrt{3}} \begin{pmatrix} 0 & -1 \\ 3 & 0 \end{pmatrix}$
$\tilde{E}_4(t) : y^2 = x(x^2 + x + \frac{t}{4})$	$(\frac{1}{4}, \frac{3}{4})$	$\Gamma_0(2)$	$\frac{1}{\sqrt{2}} \begin{pmatrix} 0 & -1 \\ 2 & 0 \end{pmatrix}$
$\tilde{E}_6(t) : y^2 + xy = x^3 - \frac{t}{432}$	$(\frac{1}{6}, \frac{5}{6})$	–	–

For the $d=6$ case, the function t is related to the modular group $\mathrm{PSL}_2(\mathbb{Z})$ by $4t(1-t) = 1728/j$, where j is the usual j -invariant, the Hauptmodul on $\mathrm{PSL}_2(\mathbb{Z})$ with $j(i) = 1728$, $j(\zeta_3) = 0$, and $j(i\infty) = \infty$.

3.1. Hypergeometric CM Elliptic Curves. Following [16], we summarize the connection between periods of elliptic curves and formulas for $1/\pi$ and their p -adic versions. Recall that the de Rham space $H_{DR}^1(\tilde{E}_d(t)/\mathbb{C}[[t]])$ is 2-dimensional, where one of its spanning vectors is given by the class ω_t of the standard invariant holomorphic differential 1-form (modulo exact forms). Another spanning vector can be obtained via the Gauss-Manin connection. For the families of elliptic curves $\tilde{E}_d(t)$, such a vector is given by $\partial_t \omega_t$, where ∂_t denotes the partial derivative with respect to t [59, 28, 29, 30, 42, 65].

On the other hand, if $\tilde{E}_d(t)$ has CM, then its endomorphism ring R corresponds to an order in an imaginary quadratic field $K = \mathbb{Q}(\sqrt{-D})$ with $-D$ a fundamental discriminant. Moreover, R induces an action on $H_{DR}^1(\tilde{E}_d(t); \mathbb{C}[[t]])$ for which ω_t is a common eigenvector. By Chowla and

Selberg [56], for any $C \in H_1(\tilde{E}_d(t), \mathbb{Z})$, the period $\int_C \omega_t$ satisfies

$$(3.1) \quad \int_C \omega_t \sim \pi \Omega_K,$$

where \sim denotes equivalence up to multiplication by an algebraic number. Here, $\Omega_K \in \mathbb{C}^\times$ is uniquely determined by K provided by the so-called Chowla-Selberg formula (for example, see [56] or [12, §6.3]),

$$(3.2) \quad \Omega_K := \frac{1}{\sqrt{\pi}} \left(\prod_{j=1}^{D-1} \Gamma\left(\frac{j}{D}\right)^{\chi(j)} \right)^{n/4h},$$

where n is the order of the unit group in K , χ is the quadratic Dirichlet character modulo D for K , and h is the class number of K .

There is another common eigenvector ν_t for the action of R on

$$H_{DR}^1(\tilde{E}_d(t); \mathbb{C}[[t]]) = \text{Span}_{\mathbb{C}}(\{\omega_t, [\partial_t \omega_t]\}),$$

which is coming from a differential of the second kind. From the Legendre relation as well as work of both Masser and Chowla-Selberg [56],

$$(3.3) \quad \int_C \omega_t \cdot \int_C \nu_t \sim \pi,$$

and there is a unique algebraic number c_t such that $\nu_t = \omega_t + c_t \cdot t \partial_t \omega_t$.

The p -adic analog of (3.3) follows from the formal group of the elliptic curve. We omit the details; for a background on the formal group, see [58, Ch. IV], and for some details on the role formal group laws play in the existence of congruences, see [36, 59]. Expand

$$(3.4) \quad \omega_t = \sum_{n \geq 0} a_t(n) \xi^n d\xi \quad \text{and} \quad \nu_t = \sum_{n \geq 0} b_t(n) \xi^n d\xi,$$

where $\xi = -\frac{x}{y}$ is a local uniformizer of $\tilde{E}_d(t)$ at infinity over \mathbb{Z}_p . For a fixed algebraic integer t of degree n over \mathbb{F}_p , we denote by

$$(3.5) \quad A := W(\mathbb{F}_p(t)),$$

the ring of Witt vectors of $\mathbb{F}_p(t)$, which is isomorphic to $\mathbb{Z}_p[\zeta_{p^n-1}]$, the ring of integers of the unique unramified extension of degree n over \mathbb{Q}_p . For the cases in this paper, we either have $A = \mathbb{Z}_p$, or A is the maximal order in the quadratic unramified extension of \mathbb{Q}_p .

When the parameter t is specified at values μ , the elliptic curve $\tilde{E}_d(\mu)$ has eigenvectors for the action of R given by $\omega := \omega_\mu$ and $\nu := \nu_\mu$. In this setting we drop the subscripts to simplify the notation. Moreover, via the isomorphism in [31, Lem. 5.1.2], we can identify ω and ν via their formal integrals by

$$(3.6) \quad \omega = \sum_{n \geq 1} \frac{a(n-1)}{n} \xi^n \quad \text{and} \quad \nu = \sum_{n \geq 1} \frac{b(n-1)}{n} \xi^n.$$

Following [16, Prop. 16] and [31, Thm. 6.1], we obtain the following p -adic version of (3.3).

Lemma 3.1. *Suppose $\mathbf{b} = (\frac{1}{d}, \frac{d-1}{d})$, and $\tilde{E}_d(\lambda)$ satisfies the conditions of Theorem 1.4. Then,*

$$a(p-1)b(p-1) \equiv \begin{cases} p & \text{if } \tilde{E}_d(\lambda) \text{ is ordinary at } p \\ -p & \text{if } \tilde{E}_d(\lambda) \text{ is supersingular at } p \end{cases} \pmod{p^2}.$$

Proof. We observe that using the notation of Chisholm et al. [16], $A = \mathbb{Z}_p$, ω, ν are eigenfunctions of the endomorphism ring R of the elliptic curve, and from the proof of [16, Thms.1,2], there exists a degree p (resp. p^2) Frobenius lifting Φ that commutes with the induced action of R on the deRham cohomology in the ordinary (resp. supersingular) case. Thus, we can apply Katz [31, Thm. 6.1] to obtain

$$a(p^2 - 1) - \text{tr}(\Phi)a(p - 1) + p \equiv 0 \pmod{p^2},$$

Here this corresponds with [16, (5.1)] with σ the identity map, $r = 2$, $n = 1$ and furthermore, ω is annihilated by $\Phi^2 - \text{tr}(\Phi)\Phi + p$. As this is the same setting as [16, Prop. 16], this time $\tilde{E}_d(\lambda)$ being defined over $\mathbb{Q}(\lambda)$ and $p > 3$ such that $\mathbb{Q}_p(\lambda) \cong \mathbb{Q}_p$, following the proof directly yields that

$$a(p - 1)b(p - 1) \equiv \begin{cases} p & \text{if } \tilde{E}_d(\lambda) \text{ is ordinary at } p \\ -p & \text{if } \tilde{E}_d(\lambda) \text{ is supersingular at } p \end{cases} \pmod{p^2}.$$

□

3.2. Connection with Modular Forms. Due to the modularity of the functions $Z(\tau)$ satisfying the conditions of Corollary 1.2, we are able to obtain information related to the elliptic curves in Table 2. In (3.3), we write π in terms of periods of elliptic curves. We can make this connection explicit via modular forms expressed as hypergeometric series. The Picard-Fuchs equation of $\tilde{E}_d(t)$ has holomorphic solution around $t = 0$ given by $f(t) := f_{0,(\frac{1}{d}, \frac{d-1}{d})}(t)$ (see [5]). Suppose we choose a Hauptmodul $t(\tau)$ of Γ such that $f(t(\tau))$ is a weight 1 modular form, and consider the elliptic curve $\tilde{E}_d(t(z_0))$ for a fixed CM point $z_0 \in \mathbb{H}$. It follows from (3.1) and the definition of Ω_K that we can choose a suitable 1-cycle (path) C such that given ω , i.e. the standard invariant holomorphic differential 1-form on $\tilde{E}_d(t(z_0))$, then

$$\int_C \omega \sim \pi \cdot f(t(z_0)),$$

where \sim denotes equivalence up to multiplication by an algebraic number. One can restate (3.3) by considering the quasi-period

$$\int_C v = \left(B \int_C \omega + t \partial_t \int_C \omega \right) |_{t=t(z_0)}$$

for a specific algebraic number $B \in \overline{\mathbb{Q}}$. Thus with z_0 as above such that $f(t(z_0)) \neq 0$, there exists a unique algebraic number $B(z_0) \in \overline{\mathbb{Q}}$ such that

$$(3.7) \quad f(t(z_0)) \left(t \frac{df}{dt} |_{t=t(z_0)} + B(z_0) f(t(z_0)) \right) \sim \frac{1}{\pi}.$$

From the proof of [16, Lemma 9], the specific value of $B(z_0)$ is given by

$$(3.8) \quad B(z_0) = -t(z_0) \frac{d\tau}{dt} \left(\frac{1}{f(t(z_0))} \frac{df}{dt} |_{t=t(z_0)} - \frac{1}{4\pi \text{Im}(z_0)} \right).$$

By the uniqueness of $B(z_0)$, (3.7) is the hypergeometric analog of (3.3) and we can explicitly write the quasi-period $\int_C \nu$ as

$$\frac{1}{\pi} \int_C \nu \sim t(z_0) \frac{df}{dt} |_{t=t(z_0)} + B(z_0) f(t(z_0)),$$

with $B(z_0)$ as above.

We further obtain the following lemma to certify that our hypergeometric evaluations are the periods and quasi-periods coming from the eigenvectors of the endomorphism ring of the associated CM elliptic curve $\tilde{E}_d(X(z_0))$ given in Corollary 1.2.

Lemma 3.2. *Suppose $X(\tau)$ and $Z(\tau) = f_{0,(\frac{1}{d}, \frac{d-1}{d})}(X(\tau))^2$ satisfy all of the conditions in Corollary 1.2 with $\gamma = \begin{pmatrix} a & b \\ c & -a \end{pmatrix}$, and $\tau_0 = \frac{a}{c} + \frac{i}{c\sqrt{N}}$ is a CM point with $\gamma\tau_0 \in D$. Then,*

$$2\alpha_N = \frac{1}{B(\gamma\tau_0)} \in \mathbb{Q}(X(\gamma\tau_0), X(\tau_0)).$$

Proof. Recall we define our derivative as $f' := \frac{1}{2\pi i} \frac{df}{d\tau}$. Set $f(X) := f_{0,(\frac{1}{d}, \frac{d-1}{d})}(X)$, and $Z(X) = f(X)^2$. Then $\frac{df}{f} = \frac{1}{2} \frac{dZ}{Z}$. Since $\gamma\tau_0 \in D$, we have from (3.8), that

$$-B(\gamma\tau_0) = \frac{X(\gamma\tau_0)}{X'(\gamma\tau_0)} \left(\frac{1}{2} \frac{Z'(\gamma\tau_0)}{Z(\gamma\tau_0)} - \frac{1}{4\pi \text{Im}(\gamma\tau_0)} \right).$$

Thus using [6, (17)], we obtain

$$\begin{aligned} -B(\gamma\tau_0) &= \frac{X(\gamma\tau_0)}{X'(\gamma\tau_0)} \left(\frac{1}{2} \cdot \frac{1}{2N} \left(\frac{c\sqrt{N}}{\pi} + \varepsilon \frac{1}{N} M'_N(\tau_0) \right) - \frac{c}{4\pi\sqrt{N}} \right) \\ &= \frac{X(\gamma\tau_0)}{X'(\gamma\tau_0)} \frac{1}{4N} \frac{\varepsilon}{N} \frac{dM_N}{dX} \Big|_{X=X(\tau_0)} \quad X'(\tau_0) = X(\gamma\tau_0) \frac{X'(\tau_0)}{X'(\gamma\tau_0)} \frac{1}{4N} \frac{\varepsilon}{N} \frac{dM_N}{dX} \Big|_{X=X(\tau_0)} \\ &= X(\gamma\tau_0) \frac{XUZ(\tau_0)}{XUZ(\gamma\tau_0)} \frac{1}{4N} \frac{\varepsilon}{N} \frac{dM_N}{dX} \Big|_{X=X(\tau_0)} = X(\gamma\tau_0) \frac{-NZ(\tau_0)XU(\tau_0)}{\varepsilon Z(\tau_0)XU(\gamma\tau_0)} \frac{1}{4N} \frac{\varepsilon}{N} \frac{dM_N}{dX} \Big|_{X=X(\tau_0)} \\ &= -\frac{1}{4N} \frac{U(\tau_0)}{U(\gamma\tau_0)} X(\tau_0) \frac{dM_N}{dX} \Big|_{X=X(\tau_0)} = -\frac{v(\tau_0)}{4NU(\gamma\tau_0)} \\ &= -\frac{1}{2\alpha_N}. \end{aligned}$$

As in Remark 1.3, we have in these cases that $U = 1 - X$, and thus the value α_N is indeed in the field $\mathbb{Q}(X(\gamma\tau_0), X(\tau_0))$. \square

We next observe the following useful lemma.

Lemma 3.3. *Suppose $X(\tau)$ and $Z(\tau) = f_{0,(\frac{1}{2}, \frac{1}{2}, \frac{d-1}{d})}(X(\tau))^2$ satisfy the conditions in Corollary 1.2 with $\gamma = \begin{pmatrix} a & b \\ c & -a \end{pmatrix}$, and $\tau_0 = \frac{a}{c} + \frac{i}{c\sqrt{N}}$ is a CM point with $\gamma\tau_0 \in D$. Then*

$$\frac{\delta_N}{f_{0,(\frac{1}{2}, \frac{1}{2}, \frac{d-1}{d})}(X(\gamma\tau_0))} \in \overline{\mathbb{Q}}.$$

Proof. Recall that from the classical result of modular forms with CM-values (see [16, Proposition 5], [18, §5.10], or [66, Proposition 26] for example), if f is a meromorphic modular form of weight k with algebraic Fourier coefficients at a CM point $z_0 \in K \cap \mathbb{H}$, then

$$f(z_0) \sim \Omega_K^k.$$

In the cases under consideration, the modular form $Z(\tau)$ is a weight-4 modular form whose Fourier expansion has integer coefficients. Therefore $Z(\gamma\tau_0) \sim \Omega_K^4$, where $K = \mathbb{Q}(c\sqrt{-N})$. Also, from the expression of δ_N in Corollary 1.2 we have

$$U(\tau_0)X(\tau_0) \left(\frac{dM_N}{dX} \right) \Big|_{X=X(\tau_0)} \cdot \delta_N \sim 1,$$

where $X(\tau_0)$, $\left(\frac{dM_N}{dX} \right) \Big|_{X=X(\tau_0)}$ are algebraic numbers, and $UZ(\tau) = \frac{1}{X} \cdot \frac{1}{2\pi i} \frac{dX}{d\tau}$. Since $UZ(\tau) = \frac{1}{X} \cdot \frac{1}{2\pi i} \frac{dX}{d\tau}$ is a meromorphic modular form of weight-2 and has integer Fourier coefficients, we have that

$$UZ(\tau_0) \sim \Omega_K^2.$$

Together with the fact $Z(\tau_0) \sim \Omega_K^4$, the identity tells us that $U(\tau_0) \sim \Omega_K^{-2}$ and hence

$$\delta_N \sim \Omega_K^2 \sim f_{0,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}(X(\gamma\tau_0)).$$

□

We now prove the following result which is an analogue to [16, Prop. 16] and used in the proof of Theorem 1.4.

Lemma 3.4. *Suppose $X(\tau)$ and $Z(\tau) = f_{0,\mathbf{b}}(X(\tau))^2$ satisfy all of the conditions in Corollary 1.2 with $\gamma = \begin{pmatrix} a & b \\ c & -a \end{pmatrix}$, and $\tau_0 = \frac{a}{c} + \frac{i}{c\sqrt{N}}$ is a CM point with $\gamma\tau_0 \in D$. Suppose further that τ_0, λ, μ, p satisfy all of the conditions of Theorem 1.4. Let $a(n)$ and $b(n)$ be defined by (3.6). Then, we have that*

$$\begin{aligned} a(p-1) &\equiv [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\mu) \pmod{p\mathbb{Z}_p[\mu]}, \\ [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\mu) \cdot [f_{c\mu,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\mu) &\equiv a(p-1)b(p-1) \pmod{p^2\mathbb{Z}_p[\mu]}, \end{aligned}$$

where

$$c_\mu = \begin{cases} 2\alpha_N & \text{if } \mu = \lambda \\ \frac{4\alpha_N(1-\mu)}{1-2\mu} = \frac{4\alpha_N(1-\mu)(1-2\mu)}{1-\lambda} & \text{if } \mu = \frac{1 \pm \sqrt{1-\lambda}}{2}. \end{cases}$$

Proof. As in §3.1 and (3.4), consider the differentials $\omega_t = \sum_{n \geq 0} a_t(n)\xi^n d\xi$ and $\nu_t = \omega_t + c_t \cdot t \partial_t \omega_t = \sum_{n \geq 0} b_t(n)\xi^n d\xi$. By [16, Lemma 12],

$$(3.9) \quad a_t(p-1) \equiv [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(t) \pmod{p}.$$

On the other hand,

$$\begin{aligned} (3.10) \quad b_t(p-1) &= a_t(p-1) + c_t \cdot t \frac{d}{dt} a_t(p-1) \\ &\equiv [f_{0,(\frac{1}{d}, \frac{d-1}{d})} + c_t \cdot t \frac{d}{dt} f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(t) \pmod{p} \\ &\stackrel{(2.4)}{\equiv} [f_{c_t,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(t) \pmod{p}. \end{aligned}$$

Next, we show the congruence relations at fixed $t = \mu$ where $\tilde{E}_d(\mu)$ has CM. The differentials

$$\omega := \omega_\mu = \sum_{n \geq 0} a(n-1)\xi^n d\xi, \quad \text{and} \quad \nu := \nu_\mu = \sum_{n \geq 0} b(n-1)\xi^n d\xi$$

are non-parallel common eigenvectors for the action of $\text{End } \tilde{E}_d(\mu)$ on the first de Rham cohomology group $H_{\text{dR}}^1(\tilde{E}_d(\mu)/A)$. Here A is the ring of Witt vectors $W(\mathbb{F}_p(\mu))$ as defined in (3.5), which is isomorphic to the ring of integers $\mathbb{Z}_p[\mu]$ of $\mathbb{Q}_p(\mu)$.

The first congruence relation then follows immediately from (3.9), and we obtain

$$(3.11) \quad a(p-1) \equiv [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\mu) \pmod{pA},$$

where $A = \mathbb{Z}_p[\mu]$ and we note that $A = \mathbb{Z}_p$ when $\mu = \lambda$.

To show the second congruence relation, we examine the unique value $c_\mu = \frac{1}{B(z_0)}$, where $\tilde{E}_d(\mu)$ has CM by $z_0 = \gamma\tau_0$ and $B(z_0)$ is as in (3.7). In order to write c_μ as a function of α_N , note that since the value α_N depends on $Z(\tau)$ (see Corollary 1.2), and therefore on \mathbf{b} , then the expression of c_μ as a function of α_N will depend on whether $\mu = \lambda$ or $\mu = \frac{1 \pm \sqrt{1-\lambda}}{2}$. When $\mu = \lambda$, the value $c_\mu = \frac{1}{B(z_0)} = 2\alpha_N$ is given by Lemma 3.2 and (3.8). When $\mu = \frac{1 \pm \sqrt{1-\lambda}}{2}$, from (3.7) we have that c_μ is the unique algebraic number such that

$$\frac{1}{\pi} \sim f_{0,(\frac{1}{d}, \frac{d-1}{d})}(\mu) \cdot f_{c_\mu,(\frac{1}{d}, \frac{d-1}{d})}(\mu) \\ \stackrel{\text{Lem. 2.8}}{=} f_{c'_\mu,(\frac{1}{2}, \frac{d-1}{d})}(\lambda),$$

where $c'_\mu = \frac{c_\mu(1-2\mu)}{2(1-\mu)}$. From Corollary 1.2, we obtain

$$(3.12) \quad \frac{1}{\pi} \cdot \frac{\delta_N}{f_{0,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}(\lambda)} = f_{2\alpha_N,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}(\lambda).$$

By Lemma 3.3, $\frac{\delta_N}{f_{0,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}(\lambda)}$ is algebraic, and the uniqueness of c_μ gives that $c'_\mu = 2\alpha_N$, and thus $c_\mu = \frac{4\alpha_N(1-\mu)}{1-2\mu}$. Therefore, 3.11 together with the specific expression of c_μ gives

$$[f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\mu) \cdot [f_{c_\mu,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\mu) \equiv a(p-1)b(p-1) \pmod{pA}.$$

In order to show the second congruence relation modulo p^2 , we consider separately the cases when $\tilde{E}_d(\mu)$ is supersingular or ordinary at p . When $\tilde{E}_d(\mu)$ is supersingular, the congruence follows as in [16, Proof of Proposition 16] by using the action of the multiplication by $-p$ on $\tilde{E}_d(\mu)$. When $\tilde{E}_d(\mu)$ is ordinary at p we use the action of the multiplication by u_p (the unit root given in Theorem 1.4) on $\tilde{E}_d(\mu)$ as in Remark 8.3. Then the congruence follows from Case 1 of the proof of Theorem 2.3 when μ can be embedded in \mathbb{Z}_p , and from Case 2 of the proof of Theorem 2.3 when μ can not be embedded in \mathbb{Z}_p . In each case, we obtain

$$[f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\mu) \cdot [f_{c_\mu,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\mu) \equiv a(p-1)b(p-1) \equiv \text{sgn} \cdot \varepsilon_\mu p \pmod{p^2A},$$

where $\text{sgn} = 1$ (or -1) if and only if $\tilde{E}_d(\mu)$ is ordinary (or supersingular) at p , and $\varepsilon_\mu = -1$ if μ can not be embedded in \mathbb{Z}_p and $\varepsilon_\mu = 1$ otherwise. \square

Remark 3.5. Let $\mu = \frac{1 \pm \sqrt{1-\lambda}}{2}$. By Lemmas 2.8, 2.10, and 3.4,

$$[f_{2\alpha_N,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\lambda) = [f_{0,(\frac{1}{d}, \frac{d-1}{d})} \cdot f_{c_\mu,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\mu) \\ \equiv [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\mu) \cdot [f_{c_\mu,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\mu) \equiv \text{sgn} \cdot \varepsilon_\mu p \pmod{p^2A}.$$

Thus, since $[f_{2\alpha_N,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\lambda)$ and $\text{sgn} \cdot \varepsilon_\mu p$ are both in \mathbb{Z}_p , from Lemma 3.4 it follows that

$$[f_{2\alpha_N,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\lambda) \equiv \text{sgn} \cdot \varepsilon_\mu p \pmod{p^2}.$$

4. PROOF OF THEOREM 1.4

We now prove our main theorem.

Proof of Theorem 1.4. Let $d \in \{2, 3, 4, 6\}$ and set $\mathbf{b} = (\frac{1}{d}, \frac{d-1}{d})$, or $\mathbf{b} = (\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})$.

Suppose $Z(\tau) = f_{0,\mathbf{b}}(X(\tau))^2$ satisfies all of the conditions in Corollary 1.2, and that if $\lambda = X(\gamma\tau_0)$, then $f_{0,\mathbf{b}}(\lambda) = f_{0,(\frac{1}{d}, \frac{d-1}{d})}(\mu)^r$ for some r , where $\mu \in \{\lambda, \frac{1 \pm \sqrt{1-\lambda}}{2}\}$.

Assume that $\mathbb{Q}(\mu) \subseteq \mathbb{Q}(\sqrt{1-\lambda})$ is a totally real field, $|\lambda|, |\mu| < 1$, and that $\tilde{E}_d(\mu)$ has CM. Let p be an odd prime such that $X(\tau_0)$, λ, μ can be embedded into \mathbb{Z}_p , and p is unramified in $\mathbb{Q}(\sqrt{1-\lambda})$. Suppose that $\tilde{E}_d(\mu)$ has good reduction modulo p .

First, we consider the case when $\mathbf{b} = (\frac{1}{d}, \frac{d-1}{d})$ and $\mu = \lambda = X(\gamma\tau_0)$, so that $f_{0,\mathbf{b}}(X(\gamma\tau_0)) = f_{0,(\frac{1}{d}, \frac{d-1}{d})}(\mu)$. Then by Lemma 2.10 and Lemma 3.4,

$$\begin{aligned} [f_{0,\mathbf{b}} \cdot f_{2\alpha_N,\mathbf{b}}]_{p-1}(X(\gamma\tau_0)) &= [f_{0,(\frac{1}{d}, \frac{d-1}{d})} \cdot f_{2\alpha_N,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\mu) \\ &\equiv [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\mu) \cdot [f_{2\alpha_N,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\mu) \pmod{p^2} \\ &\equiv a(p-1)b(p-1) \pmod{p^2}. \end{aligned}$$

Applying Lemma 3.1, we thus obtain

$$[f_{0,\mathbf{b}} \cdot f_{2\alpha_N,\mathbf{b}}]_{p-1}(X(\gamma\tau_0)) \equiv \begin{cases} p & \text{if } \tilde{E}_d(\lambda) \text{ is ordinary at } p \\ -p & \text{if } \tilde{E}_d(\lambda) \text{ is supersingular at } p \end{cases} \pmod{p^2}.$$

We next apply Theorem 2.6. Here $k = \mathbb{Q}(\tau_0) = \mathbb{Q}(c\sqrt{-N})$. Thus by the splitting of primes in imaginary quadratic fields (see Marcus [43, Thm. 25] for example), we obtain that $\tilde{E}_d(\lambda)$ is ordinary if and only if $\left(\frac{-c^2N}{p}\right) = 1$, as desired.

Next, we consider the case when $\mathbf{b} = (\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})$ and $\mu = \frac{1 \pm \sqrt{1-\lambda}}{2} = \frac{1 \pm \sqrt{1-X(\gamma\tau_0)}}{2}$, so that $f_{0,\mathbf{b}}(X(\gamma\tau_0)) = f_{0,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}\left(\frac{1 \pm \sqrt{1-X(\gamma\tau_0)}}{2}\right)^2$. Then by Lemma 2.10,

$$\begin{aligned} [f_{0,\mathbf{b}} \cdot f_{2\alpha_N,\mathbf{b}}]_{p-1}(X(\gamma\tau_0)) &= [f_{0,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})} \cdot f_{2\alpha_N,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\lambda) \\ &\equiv [f_{0,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\lambda) \cdot [f_{2\alpha_N,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\lambda) \pmod{p^3}. \end{aligned}$$

From (3.12) we know that α, λ admit a series of the form (1.2) which satisfies the conditions of Theorems 2.1 and 2.3. Thus we have that

$$[f_{2\alpha_N,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\lambda) \equiv \begin{cases} \left(\frac{1-\lambda}{p}\right) \cdot p & \text{if } \tilde{E}_d(\mu) \text{ is ordinary at } p \\ -\left(\frac{1-\lambda}{p}\right) \cdot p & \text{if } \tilde{E}_d(\mu) \text{ is supersingular at } p \end{cases} \pmod{p^2},$$

and that

$$[f_{0,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\lambda) \equiv \begin{cases} u_p^2 & \text{if } \tilde{E}_d\left(\frac{1-\sqrt{1-\lambda}}{2}\right) \text{ is ordinary at } p \text{ and } \left(\frac{1-\lambda}{p}\right) = 1 \\ \left(\frac{kd}{p}\right) u_p^2 & \text{if } \tilde{E}_d\left(\frac{1-\sqrt{1-\lambda}}{2}\right) \text{ is ordinary at } p \text{ and } \left(\frac{1-\lambda}{p}\right) = -1 \\ 0 & \text{if } \tilde{E}_d(\mu) \text{ is supersingular at } p \end{cases} \pmod{p^2}.$$

Putting these together and again applying Theorem 2.6 gives the desired result. \square

5. EXAMPLES OF THEOREM 1.4

Below we give 11 examples of Theorem 1.4. Each of these examples stem from applications of [6, Thm. 2]. As such, it is clear that Corollary 1.2 will hold for each of these examples, however in an effort to keep this paper self-contained we will demonstrate that the conditions of Corollary 1.2 are satisfied for each example, giving that

$$\frac{\delta}{\pi} = f_{0,\mathbf{b}}(\lambda) \cdot f_{2\alpha,\mathbf{b}}(\lambda)$$

where δ, α, λ are given in Table 1.

This table also lists the values of N, μ, d necessary for Theorem 1.4 and additionally from this table we can see that for each example $\mathbb{Q}(\lambda)$ is totally real and $|\lambda|, |\mu| < 1$. We then determine

primes p for which Theorem 1.4 holds. Data for the elliptic curves $\tilde{E}_d(\mu)$ is obtained using **Sagemath** [53]. We note that each elliptic curve $\tilde{E}_d(\mu)$ has CM.

We will use notation from [6] to refer to the Hauptmoduln for the examples. Below we give the Hauptmoduln in terms of the j -function and Dedekind η -function. For a discussion regarding our choice of Hauptmoduln, see [6].

$$\begin{aligned} t_2(\tau) &= -64\eta(2\tau)^{24}/\eta(\tau)^{24} \\ t_3(\tau) &= -27\eta(3\tau)^{12}/\eta(\tau)^{12} \\ t_\infty(\tau) &= 16\eta(\tau)^8\eta(4\tau)^{16}/\eta(2\tau)^{24} \\ t_{2,3}(\tau) &= 1728/j(\tau) \\ t_{2,4}(\tau) &= 256\eta(\tau)^{24}\eta(2\tau)^{24}/(\eta(\tau)^{24} + 64\eta(2\tau)^{24})^2 \\ t_{2,6}(\tau) &= 108\eta(\tau)^{12}\eta(3\tau)^{12}/(\eta(\tau)^{12} + 27\eta(3\tau)^{12})^2 \end{aligned}$$

5.1. Examples from [6, §4]. For each example stemming from [6, §4] we have that $Z(\tau) = f_{0,(\frac{1}{d}, \frac{d-1}{d})}(X(\tau))^2$ so $\mu = \lambda$. Furthermore, we either have that $\mathbb{Q}(\lambda) = \mathbb{Q}$ or $\mathbb{Q}(\lambda) = \mathbb{Q}(\sqrt{D})$ for some non-square $D \in \mathbb{Z}$. In the case where $\mathbb{Q}(\lambda)$ is a quadratic extension, requiring that λ can be embedded into \mathbb{Z}_p gives us that $\left(\frac{D}{p}\right) = 1$, which in turn implies that p splits in $\mathbb{Q}(\lambda)$. Thus, requiring that λ can be embedded into \mathbb{Z}_p is a stronger condition than requiring that p is unramified in $\mathbb{Q}(\lambda)$.

Examples 5.1-5.7 relate to examples 4.2-4.7. from [6], respectively. In Examples 5.1-5.4, we have $\Gamma = \Gamma_0(2)$ and consider $X(\tau) = \frac{t_2(\tau)}{t_2(\tau)-1}$. With $Z(\tau) = f_{0,(\frac{1}{4}, \frac{3}{4})}(X(\tau))^2$ we have that $\frac{1}{2\pi i} \frac{dX}{d\tau} = Z(\tau)X(\tau)U(\tau)$ for $U(\tau) = \frac{1}{1-t_2(\tau)}$ as in Remark 1.3.

For Examples 5.1,5.2 we will consider $\gamma = \begin{pmatrix} 0 & -\frac{1}{\sqrt{2}} \\ \sqrt{2} & 0 \end{pmatrix}$ and we have that $(Z|_\gamma)(\tau) = -Z(\tau)$ so $\varepsilon = -1$.

For Examples 5.3, 5.4 we will consider $\gamma = \begin{pmatrix} 1 & -1 \\ 2 & -1 \end{pmatrix}$ and we have that $(Z|_\gamma)(\tau) = Z(\tau)$.

Example 5.1. Let $N = 3$ so $\tau_0 = \frac{i}{\sqrt{6}}$ and the conditions of Corollary 1.2 are satisfied. From [16] we have that $X(\tau_0) = \frac{3+2\sqrt{2}}{6}$ and $\lambda = X(\gamma\tau_0) = \frac{3-2\sqrt{2}}{6}$.

We have that $X(\tau_0), \lambda$ can be embedded into \mathbb{Z}_p if $p \neq 2, 3$ and $\left(\frac{2}{p}\right) = 1$.

The conductor norm of $\tilde{E}_4(\mu)$ is $2^{10} \cdot 3^4$ so $\tilde{E}_4(\mu)$ has good reduction at $p \neq 2, 3$. Therefore, by Theorem 1.4, for primes $p \neq 2, 3$ with $p \equiv \pm 1 \pmod{8}$

$$\left[f_{0,(\frac{1}{4}, \frac{3}{4})} \cdot f_{6\sqrt{2}+8,(\frac{1}{4}, \frac{3}{4})} \right]_{p-1} \left(\frac{3-2\sqrt{2}}{6} \right) \equiv \begin{cases} p & \text{if } \left(\frac{-6}{p}\right) = 1 \\ -p & \text{otherwise} \end{cases} \pmod{p^2}.$$

Example 5.2. Let $N = 5$ so $\tau_0 = \frac{i}{\sqrt{10}}$ which satisfies the conditions of Corollary 1.2. From [6] we have that $X(\tau_0) = \frac{9+4\sqrt{5}}{18}$ and $\lambda = X(\gamma\tau_0) = \frac{9-4\sqrt{5}}{18}$.

We have that $X(\tau_0), \lambda$ can be embedded into \mathbb{Z}_p if $p \neq 2, 3$ and $\left(\frac{5}{p}\right) = 1$, which occurs if and only if $p \equiv \pm 1 \pmod{5}$. The conductor norm of $\tilde{E}_4(\mu)$ is $2^{16}3^4$ so $\tilde{E}_4(\mu)$ has good reduction for $p \neq 2, 3$.

Therefore, by Theorem 1.4 for primes $p \geq 5$ with $p \equiv \pm 1 \pmod{5}$

$$\left[f_{0,(\frac{1}{4},\frac{3}{4})} \cdot f_{\frac{20+9\sqrt{5}}{2},(\frac{1}{4},\frac{3}{4})} \right]_{p-1} \left(\frac{9-4\sqrt{5}}{18} \right) \equiv \begin{cases} p & \text{if } \left(\frac{-10}{p} \right) = 1 \\ -p & \text{otherwise} \end{cases} \pmod{p^2}.$$

Example 5.3. Let $N = 3$ so

$\tau_0 = \frac{1}{2} + \frac{i}{2\sqrt{3}}$ which satisfies the conditions of Corollary 1.2. From [6] we have that $X(\tau_0) = X(\gamma\tau_0) = \frac{-1}{3}$ so $\lambda = \frac{-1}{3}$.

We have that $X(\tau_0) = X(\lambda) = \frac{-1}{3}$ can be embedded into \mathbb{Z}_p for $p \neq 3$. The conductor norm of $\tilde{E}_4(\lambda)$ is $2^3 \cdot 3^2$ so $\tilde{E}_4(\lambda)$ has good reduction for $p \neq 2, 3$. Therefore, by Theorem 1.4 for primes $p \geq 5$

$$\left[f_{0,(\frac{1}{4},\frac{3}{4})} \cdot f_{8,(\frac{1}{4},\frac{3}{4})} \right]_{p-1} \left(\frac{-1}{3} \right) \equiv \begin{cases} p & \text{if } \left(\frac{-12}{p} \right) = 1 \\ -p & \text{otherwise} \end{cases} \pmod{p^2}.$$

Example 5.4. Let $N = 5$ so $\tau_0 = \frac{1}{2} + \frac{i}{2\sqrt{5}}$ which satisfies the conditions of Corollary 1.2. From [6] we have that $X(\tau_0) = X(\gamma\tau_0) = \frac{2-\sqrt{5}}{4}$ so $\lambda = \frac{2-\sqrt{5}}{4}$.

We have that $X(\tau_0), \lambda$ can be embedded into \mathbb{Z}_p if $p \neq 2$ and $\left(\frac{5}{p} \right) = 1$ which occurs if and only if $p \equiv \pm 1 \pmod{5}$. The conductor norm of $\tilde{E}_4(\lambda)$ is 2^{12} so $\tilde{E}_4(\lambda)$ has good reduction at $p \neq 2$. Therefore, by Theorem 1.4 for primes $p \neq 2$ with $p \equiv \pm 1 \pmod{5}$

$$\left[f_{0,(\frac{1}{4},\frac{3}{4})} \cdot f_{\frac{2}{3}(10+4\sqrt{5}),(\frac{1}{4},\frac{3}{4})} \right]_{p-1} \left(\frac{2-\sqrt{5}}{4} \right) \equiv \begin{cases} p & \text{if } \left(\frac{-20}{p} \right) = 1 \\ -p & \text{otherwise} \end{cases} \pmod{p^2}.$$

Remark 5.5. We can obtain supercongruences related to the series used for 5.1-5.4 above by applying 2.1 directly. These supercongruences are given and discussed in Section 6 below.

In [6, Ex. 4.6] we have that $\Gamma = \Gamma_0(3)$ and consider $X(\tau) = \frac{t_3(\tau)}{t_3(\tau)-1}$. With $Z(\tau) = f_{0,(\frac{1}{3},\frac{2}{3})}(X(\tau))^2$ we have that $\frac{1}{2\pi i} \frac{dX}{d\tau} = Z(\tau)X(\tau)U(\tau)$ for $U(\tau) = \frac{1}{1-t_3(\tau)}$ as in Remark 1.3.

Example 5.6. We consider $\gamma = \begin{pmatrix} 0 & -\frac{1}{\sqrt{3}} \\ \sqrt{3} & 0 \end{pmatrix}$ and we have that $(Z|_\gamma)(\tau) = Z(\tau)$ so $\varepsilon = 1$. Letting $N = 2$ so $\tau_0 = \frac{i}{\sqrt{6}}$ we see that the conditions of Corollary 1.2 are satisfied. From [6] we have that $X(\tau_0) = \frac{2+\sqrt{2}}{4}$ and $\lambda = X(\gamma\tau_0) = \frac{2-\sqrt{2}}{4}$.

We have that $X(\tau_0), \lambda$ can be embedded into \mathbb{Z}_p if $p \neq 2$ and $\left(\frac{2}{p} \right) = 1$ which occurs if and only if $p \equiv \pm 1 \pmod{8}$. The conductor norm of $\tilde{E}_3(\lambda)$ is 3^4 so $\tilde{E}_3(\lambda)$ has good reduction at $p \neq 3$. Therefore, by Theorem 1.4 for primes $p \neq 2, 3$ with $p \equiv \pm 1 \pmod{8}$

$$\left[f_{0,(\frac{1}{3},\frac{2}{3})} \cdot f_{6(\sqrt{2}+1),(\frac{1}{3},\frac{2}{3})} \right]_{p-1} \left(\frac{2-\sqrt{2}}{4} \right) \equiv \begin{cases} p & \text{if } \left(\frac{-6}{p} \right) = 1 \\ -p & \text{otherwise} \end{cases} \pmod{p^2}.$$

In [6, Ex. 4.7] we have that $\Gamma = \Gamma_0(4)$ and consider $X(\tau) = t_\infty(\tau)$. With $Z(\tau) = f_{0,(\frac{1}{2},\frac{1}{2})}(X(\tau))^2$ we have that $\frac{1}{2\pi i} \frac{dX}{d\tau} = Z(\tau)X(\tau)U(\tau)$ for $U(\tau) = 1 - t_\infty(\tau)$.

Example 5.7. We consider $\gamma = \begin{pmatrix} 0 & -\frac{1}{2} \\ 2 & 0 \end{pmatrix}$ and we have that $Z(\tau|_\gamma) = -Z(\tau)$ so $\varepsilon = -1$. Letting $N = 2$ so $\tau_0 = \frac{i}{2\sqrt{2}}$ satisfies the conditions of Corollary 1.2. From 1.2 we have that $X(\tau_0) = 2\sqrt{2}-2$ and $\lambda = X(\gamma\tau_0) = 3-2\sqrt{2}$. We have that $X(\tau_0), \lambda$ can be embedded into \mathbb{Z}_p if $\left(\frac{2}{p} \right) = 1$ which occurs

if and only if $p \equiv \pm 1 \pmod{8}$. The conductor norm of $\tilde{E}_2(\lambda)$ is 2^6 so $\tilde{E}_2(\lambda)$ has good reduction at $p \neq 2$. Therefore, by Theorem 1.4 for primes $p \neq 2$ with $p \equiv \pm 1 \pmod{8}$ we have

$$\left[f_{0,(\frac{1}{2},\frac{1}{2})} \cdot f_{2(4+2\sqrt{2}),(\frac{1}{2},\frac{1}{2})} \right]_{p-1} \left(3 - 2\sqrt{2} \right) \equiv \begin{cases} p & \text{if } \left(\frac{-8}{p} \right) = 1 \\ -p & \text{otherwise} \end{cases} \pmod{p^2}.$$

One may notice that the examples above arise from series that are of the form $\frac{\delta}{\pi} = f_{\alpha,\mathbf{b}}(\lambda)$, and thus one could apply Theorem 2.1 directly to obtain a supercongruence. See Section 6 for discussion and explicit examples arising from this method.

5.2. Examples from [6, §5]. Examples 5.8-5.12 below arise from [6, Examples 5.2-5.6]. For these examples, we have $X(\tau) = t_{2,m}$ for $m \in \{3, 4, 6\}$. Particularly, $m = 3$ and $\Gamma = \text{PSL}_2(\mathbb{Z})$ for Examples 5.8 and 5.9, $m = 4$ and $\Gamma = \Gamma_0^+(2)$ for Example 5.10, and $m = 6$ and $\Gamma = \Gamma_0^+(3)$ for Examples 5.11 and 5.12. With $Z(\tau) = f_{0,(\frac{1}{2},\frac{1}{2}-\frac{1}{m},\frac{1}{2}+\frac{1}{m})}(X(\tau))^2$ we have that $\frac{1}{2\pi i} \frac{dX}{d\tau} = Z(\tau)X(\tau)U(\tau)$ where $U(\tau)$ is given in [6, §5, (42)-(44)]. For the following examples, we consider $\gamma = \frac{1}{\sqrt{s}} \begin{pmatrix} 0 & -1 \\ s & 0 \end{pmatrix}$ where $s = 1$ for Examples 5.8 and 5.9, $s = 2$ for Example 5.10, and $s = 3$ for Examples 5.11 and 5.12 and we have $(Z|_{\gamma})(\tau) = Z(\tau)$. Note that $\varepsilon = 1$ for the remaining examples.

We now proceed with each examples individually to obtain supercongruence in each case as a consequence of Theorem 1.4. We note that

$$\begin{aligned} Z(\tau) &= f_{0,(\frac{1}{2},\frac{1}{2}-\frac{1}{m},\frac{1}{2}+\frac{1}{m})}(X(\tau))^2 \\ &= f_{0,(\frac{1}{4}-\frac{1}{2m},\frac{1}{4}+\frac{1}{2m})}(X(\tau))^4 \quad (\text{by [16, (34)]}) \\ &= f_{0,(\frac{1}{2}-\frac{1}{m},\frac{1}{2}+\frac{1}{m})} \left(\frac{1 \pm \sqrt{1-X(\tau)}}{2} \right)^4 \quad (\text{by [16, (31)]}). \end{aligned}$$

In each example we have $\mathbf{b} = (\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})$ for $d = \frac{1}{2} - \frac{1}{m} \in \{3, 4, 6\}$ and $\mu = \frac{1 - \sqrt{1-\lambda}}{2}$ as in Table 1. Below u_p is the unit root of the geometric Frobenius at p acting on the first cohomology of the elliptic curve $\tilde{E}_d(\mu)$ as defined in Theorem 1.4.

Example 5.8. In [6, Example 5.2], letting $N = 2$, we see that Corollary 1.2 is satisfied for $\tau_0 = \frac{i}{\sqrt{2}}$. From [6] we have that, $X(\tau_0) = \frac{27}{125}$ and $\lambda = X(\gamma\tau_0) = \frac{27}{125}$. We have that $X(\tau_0)$ and λ can be embedded into \mathbb{Z}_p if $p \neq 5$.

Also, p is unramified in $\mathbb{Q}(\mu) = \mathbb{Q}(\sqrt{10})$ if $p \neq 2, 5$. Furthermore, the conductor norm of $\tilde{E}_6(\mu)$ is $2^8 \cdot 5^2$. So, $\tilde{E}_6(\mu)$ has good reduction for $p \neq 2, 5$. Therefore, by Theorem 1.4, for $p \neq 2, 5$, we get that

$$\left[f_{0,(\frac{1}{2},\frac{1}{6},\frac{5}{6})} \cdot f_{\frac{28}{3},(\frac{1}{2},\frac{1}{6},\frac{5}{6})} \right]_{p-1} \left(\frac{27}{125} \right) \equiv \begin{cases} pu_p^2 & \text{if } \left(\frac{-2}{p} \right) = 1 \text{ and } \left(\frac{1-\lambda}{p} \right) = 1 \\ - \left(\frac{-1}{p} \right) pu_p^2 & \text{if } \left(\frac{-2}{p} \right) = 1 \text{ and } \left(\frac{1-\lambda}{p} \right) = -1 \end{cases} \pmod{p^3},$$

and $\left[f_{0,(\frac{1}{2},\frac{1}{6},\frac{5}{6})} \cdot f_{\frac{28}{3},(\frac{1}{2},\frac{1}{6},\frac{5}{6})} \right]_{p-1} \equiv 0 \pmod{p^2}$, otherwise.

Example 5.9. In [6, Example 5.3], letting $N = 3$, we see that Corollary 1.2 is satisfied for $\tau_0 = \frac{i}{\sqrt{3}}$. From [6] we have that, $X(\tau_0) = \frac{4}{125}$ and $\lambda = X(\gamma\tau_0) = \frac{4}{125}$.

We have that $X(\tau_0)$ and λ can be embedded into \mathbb{Z}_p if $p \neq 5$. Also, p is unramified in $\mathbb{Q}(\mu) = \mathbb{Q}(\sqrt{5})$ if $p \neq 5$. Furthermore, the conductor norm of $\tilde{E}_6(\mu)$ is $2^8 \cdot 3^4 \cdot 5^2$. So, $\tilde{E}_6(\mu)$ has good

reduction for $p \neq 2, 3, 5$. Therefore, since $\left(\frac{1-\lambda}{p}\right) = \left(\frac{5}{p}\right)$, then by Theorem 1.4, for $p \neq 2, 3, 5$, we get that

$$\left[f_{0,(\frac{1}{2}, \frac{1}{6}, \frac{5}{6})} \cdot f_{11,(\frac{1}{2}, \frac{1}{6}, \frac{5}{6})} \right]_{p-1} \left(\frac{4}{125} \right) \equiv \begin{cases} pu_p^2 & \text{if } \left(\frac{-3}{p}\right) = 1 \text{ and } \left(\frac{5}{p}\right) = 1 \\ -\left(\frac{-1}{p}\right) pu_p^2 & \text{if } \left(\frac{-3}{p}\right) = 1 \text{ and } \left(\frac{5}{p}\right) = -1 \end{cases} \pmod{p^3},$$

and $\left[f_{0,(\frac{1}{2}, \frac{1}{6}, \frac{5}{6})} \cdot f_{11,(\frac{1}{2}, \frac{1}{6}, \frac{5}{6})} \right]_{p-1} \left(\frac{4}{125} \right) \equiv 0 \pmod{p^2}$, otherwise.

Example 5.10. In [6, Example 5.4], letting $N = 3$, we see that Corollary 1.2 is satisfied for $\tau_0 = \frac{i}{\sqrt{6}}$. From [6] we have that, $X(\tau_0) = \frac{1}{9}$ and $\lambda = X(\gamma\tau_0) = \frac{1}{9}$.

We have that $X(\tau_0)$ and λ can be embedded into \mathbb{Z}_p if $p \neq 3$. Also, p is unramified in $\mathbb{Q}(\mu) = \mathbb{Q}(\sqrt{2})$ if $p \neq 2$. Furthermore, the conductor norm of $\tilde{E}_4(\mu)$ is $2^{10} \cdot 3^4$. So, $\tilde{E}_4(\mu)$ has good reduction for $p \neq 2, 3$. Therefore, since $\left(\frac{1-\lambda}{p}\right) = \left(\frac{2}{p}\right)$ then by Theorem 1.4, for $p \neq 2, 3$, we get that

$$\left[f_{0,(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})} \cdot f_{8,(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})} \right]_{p-1} \left(\frac{1}{9} \right) \equiv \begin{cases} pu_p^2 & \text{if } \left(\frac{-6}{p}\right) = 1 \text{ and } \left(\frac{2}{p}\right) = 1 \\ -\left(\frac{-2}{p}\right) pu_p^2 & \text{if } \left(\frac{-6}{p}\right) = 1 \text{ and } \left(\frac{2}{p}\right) = -1 \\ 0 & \text{otherwise} \end{cases} \pmod{p^2}.$$

Example 5.11. In [6, Example 5.5], letting $N = 2$, we see that Corollary 1.2 is satisfied for $\tau_0 = \frac{i}{\sqrt{6}}$. From [6] we have that, $X(\tau_0) = \frac{1}{2}$ and $\lambda = X(\gamma\tau_0) = \frac{1}{2}$.

We have that $X(\tau_0)$ and λ can be embedded into \mathbb{Z}_p if $p \neq 2$. Also, p is unramified in $\mathbb{Q}(\mu) = \mathbb{Q}(\sqrt{2})$ if $p \neq 2$. Furthermore, the conductor norm of $\tilde{E}_3(\mu)$ is 3^4 . So, $\tilde{E}_3(\mu)$ has good reduction for $p \neq 3$. Now, $\left(\frac{1-\lambda}{p}\right) = \left(\frac{2}{p}\right)$, and if both $\left(\frac{-6}{p}\right) = 1$ and $\left(\frac{2}{p}\right) = -1$, then $\left(\frac{-3}{p}\right) = -1$ so $-\left(\frac{-3}{p}\right) pu_p^2 = pu_p^2$. Therefore, by Theorem 1.4, for $p \neq 2, 3$, we get that

$$\left[f_{0,(\frac{1}{2}, \frac{1}{3}, \frac{2}{3})} \cdot f_{6,(\frac{1}{2}, \frac{1}{3}, \frac{2}{3})} \right]_{p-1} \left(\frac{1}{2} \right) \equiv \begin{cases} pu_p^2 & \text{if } \left(\frac{-6}{p}\right) = 1 \\ 0 & \text{otherwise} \end{cases} \pmod{p^2}.$$

Example 5.12. In [6, Example 5.6], letting $N = 5$, we see that Corollary 1.2 is satisfied for $\tau_0 = \frac{i}{\sqrt{15}}$. From [6] we have that, $X(\tau_0) = \frac{4}{125}$ and $\lambda = X(\gamma\tau_0) = \frac{4}{125}$.

We have that $X(\tau_0)$ and λ can be embedded into \mathbb{Z}_p if $p \neq 5$. Also, p is unramified in $\mathbb{Q}(\mu) = \mathbb{Q}(\sqrt{5})$ if $p \neq 5$. Furthermore, the conductor norm of $\tilde{E}_3(\mu)$ is 3^4 . So, $\tilde{E}_3(\mu)$ has good reduction for $p \neq 3$. Now, $\left(\frac{1-\lambda}{p}\right) = \left(\frac{5}{p}\right)$, and if both $\left(\frac{-15}{p}\right) = 1$ and $\left(\frac{5}{p}\right) = -1$, then $\left(\frac{-3}{p}\right) = -1$ so $-\left(\frac{-3}{p}\right) pu_p^2 = pu_p^2$. Therefore, by Theorem 1.4, for $p \neq 3, 5$, we get that

$$\left[f_{0,(\frac{1}{2}, \frac{1}{3}, \frac{2}{3})} \cdot f_{\frac{33}{4},(\frac{1}{2}, \frac{1}{3}, \frac{2}{3})} \right]_{p-1} \left(\frac{4}{125} \right) \equiv \begin{cases} pu_p^2 & \text{if } \left(\frac{-15}{p}\right) = 1 \\ 0 & \text{otherwise} \end{cases} \pmod{p^2}.$$

6. EXAMPLES OF THEOREM 2.1

As mentioned in Remark 5.5 we can directly apply Theorem 2.1 to the series in [6, Ex. 4.2-4.5] to obtain supercongruences. These supercongruences are genuinely different than the supercongruences obtained from Theorem 1.4.

We will formally list each example below, but note that applying Theorem 2.1 directly produces (finitely) more restrictions on the primes for which the supercongruences hold, however the result is expected to be stronger as Theorem 2.1 is conjectured to hold modulo p^3 .

As described in Remark 1.3, the examples in [6, Ex. 4.2-4.5], which are obtained from applying [6, Cor. 4.1], differ in form from their counterparts in Table 1 which are obtained by applying Corollary 1.2. However, each pair is related by an algebraic constant scaling factor.

To elaborate, when applying Corollary 1.2 we obtain a series of the form $f_{c,\hat{\mathbf{b}}}(\hat{\lambda})$ that is a product $f_{0,\mathbf{b}}(\lambda) \cdot f_{a,\mathbf{b}}(\lambda)$ such that there is a quadratic relationship between $\hat{\lambda}$ and λ (which can be obtained through hypergeometric quadratic formulas), as in Lemma 2.8. There is some algebraic number \mathbf{c} such that

$$f_{c,\hat{\mathbf{b}}}(\hat{\lambda}) = \mathbf{c} f_{0,\mathbf{b}} \cdot f_{a,\mathbf{b}}(\lambda),$$

however, as the series have different coefficients,

$$[f_{c,\hat{\mathbf{b}}}]_{p-1}(\hat{\lambda}) \neq [f_{0,\mathbf{b}} \cdot f_{a,\mathbf{b}}]_{p-1}(\lambda).$$

Further, due to the quadratic relationship between λ and $\hat{\lambda}$, the truncated series $[f_{c,\hat{\mathbf{b}}}]_{p-1}(\hat{\lambda})$ yields a higher exponent of λ for the n^{th} term. We also note that it is not always true that

$$[f_{c,\hat{\mathbf{b}}}]_{p-1}(\hat{\lambda}) \equiv [f_{0,\mathbf{b}} \cdot f_{a,\mathbf{b}}]_{p-1}(\lambda) \pmod{p^2}.$$

In addition, the truncations are related to periods for elliptic curves, and the truncations $[f_{c,\hat{\mathbf{b}}}]_{p-1}(\hat{\lambda})$ and $[f_{0,\mathbf{b}} \cdot f_{a,\mathbf{b}}]_{p-1}(\lambda)$ are related to isogenous, yet distinct, elliptic curves.

In the following examples, recall that $\text{sgn} = \pm 1$, equaling 1 (or -1) if and only if $E_d(\lambda) = \tilde{E}_d(\frac{1 \pm \sqrt{1-\lambda}}{2})$ is ordinary (or supersingular) at p . Since $d = 2$ for each example, let $\mathbf{b} = (\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$. As in Section 5, data for each elliptic curve $E_2(\lambda)$ is found using Sagemath [53].

Example 6.1. Consider the series

$$\frac{2}{\pi(4\sqrt{2}-5)} = f_{\frac{12}{3-2\sqrt{2}}, \mathbf{b}}(12\sqrt{2}-17)$$

given in [6, Example 4.2]. Here we have $d = 2$, $a = \frac{12}{3-2\sqrt{2}}$ and $\lambda = 12\sqrt{2} - 17$. Note that $\mathbb{Q}(\lambda) = \mathbb{Q}(\sqrt{2})$ is totally real, $|\lambda| = |12\sqrt{2} - 17| < 1$ and $E_2(\lambda) : y^2 = x(x-1)(x - \frac{1-\sqrt{18-12\sqrt{2}}}{2})$ has CM with CM discriminant -24 , which is coprime to all primes $p \neq 2, 3$. Further, all primes $p \neq 2, 3$ are unramified in $\mathbb{Q}(\sqrt{1-\lambda}) = \mathbb{Q}(\sqrt{18-12\sqrt{2}}) = \mathbb{Q}(\sqrt{2}, \sqrt{3})$. Further, $a = \frac{12(3+\sqrt{2})}{7}$ can be embedded into \mathbb{Z}_p for all $p \neq 7$ with $\left(\frac{2}{p}\right) = 1$ and $\lambda = 12\sqrt{2} - 17$ can be embedded into \mathbb{Z}_p for all primes $p \neq 2$.

Therefore, for all primes $p \neq 2, 7$ with $\left(\frac{2}{p}\right) = 1$

$$\left[f_{\frac{12}{3-2\sqrt{2}}, (\frac{1}{2}, \frac{1}{2}, \frac{1}{2})} \right]_{p-1} (12\sqrt{2}-17) \equiv \text{sgn} \cdot \left(\frac{18-12\sqrt{2}}{p} \right) \cdot p \pmod{p^2}.$$

Example 6.2. [6, Example 4.3] gives us the series

$$\frac{2}{\pi(23-10\sqrt{5})} = f_{\frac{12(15+4\sqrt{5})}{29}, \mathbf{b}}(72\sqrt{5}-161).$$

Applying Theorem 2.1 with $d = 2$, $a = \frac{12(15+4\sqrt{5})}{29}$, $\lambda = 72\sqrt{5} - 161$, we have that $\mathbb{Q}(\lambda) = \mathbb{Q}(\sqrt{5})$ is totally real, $|\lambda| = |72\sqrt{5} - 161| < 1$ and $E_2(\lambda) : y^2 = x(x-1)\left(x - \frac{1-3\sqrt{18-8\sqrt{5}}}{2}\right)$ has CM with CM discriminant -40 which is coprime to all primes $p \neq 2, 5$. All primes $p \neq 2, 5, 19$ are unramified in $\mathbb{Q}(\sqrt{1-\lambda}) = \mathbb{Q}(\sqrt{18-8\sqrt{5}})$.

We see that a, λ can be embedded in \mathbb{Z}_p^\times for all primes $p \neq 29$ with Legendre symbol $\left(\frac{5}{p}\right) = 1$, and we fix such an embedding. So for all primes p such that $p \equiv \pm 1 \pmod{5}$ and $p \neq 29$ we have that

$$\left[f_{\frac{12(15+4\sqrt{5})}{29}, \mathbf{b}} \right]_{p-1} (72\sqrt{5} - 161) \equiv \text{sgn} \cdot \left(\frac{-10(162 - 72\sqrt{5})}{p} \right) \cdot p \pmod{p^2}.$$

Example 6.3. [6, Example 4.4] gives us the series

$$\frac{4}{\pi} = f_{6, \mathbf{b}} \left(\frac{1}{4} \right).$$

Applying Theorem 2.1 with $d = 2$, $a = 6$, $\lambda = \frac{1}{4}$, we have that $\mathbb{Q}(\lambda) = \mathbb{Q}$ which is totally real, $|\lambda| = |\frac{1}{4}| < 1$, and $E_2(\lambda) : y^2 = x(x-1)(x - \frac{2-\sqrt{3}}{4})$ has CM with CM discriminant -12 , which is coprime to primes $p \neq 2, 3$. All primes $p \neq 3$ are unramified in $\mathbb{Q}(\sqrt{1-\lambda}) = \mathbb{Q}(\sqrt{3})$.

We see that a, λ can be embedded into \mathbb{Z}_p^\times for all primes $p \neq 2$ and we fix such an embedding. Then for all primes $p \neq 2, 3$ we have that

$$[f_{6, \mathbf{b}}]_{p-1} \left(\frac{1}{4} \right) \equiv \text{sgn} \cdot \left(\frac{3/4}{p} \right) \cdot p \pmod{p^2}$$

Example 6.4. [6, Example 4.5] gives us the series

$$\frac{10(\sqrt{5} + 1)}{\pi \sqrt{\sqrt{5} - 2}} = f_{5+\sqrt{5}, \mathbf{b}}(9 - 4\sqrt{5}).$$

Applying Theorem 2.1 with $d = 1$, $a = 5 + \sqrt{5}$, $\lambda = 9 - 4\sqrt{5}$, we have that $\mathbb{Q}(\lambda) = \mathbb{Q}(\sqrt{5})$ is totally real, $|\lambda| = |9 - 4\sqrt{5}| < 1$, and $E_2(\lambda) = x(x-1)(x - \frac{1-2\sqrt{\sqrt{5}-2}}{2})$ has CM with CM discriminant -20 , which is coprime to all primes $p \neq 2, 5$. Further, all primes $p \neq 2, 5$ are unramified in $\mathbb{Q}(\sqrt{1-\lambda})$.

We see that a, λ can be embedded into \mathbb{Z}_p^\times for primes p with $\left(\frac{5}{p}\right) = 1$. Therefore, for all primes $p \neq 2, 5$ with $\left(\frac{5}{p}\right) = 1$ we have that

$$\left[f_{5+\sqrt{5}, \mathbf{b}} \right]_{p-1} (9 - 4\sqrt{5}) \equiv \text{sgn} \cdot \left(\frac{4\sqrt{5} - 8}{p} \right) \cdot p \pmod{p^2}.$$

7. APPLICATIONS AND FUTURE DIRECTIONS.

In this section, we describe several relevant properties of hypergeometric functions of the form $f_{0, (\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}$. In §7.1, we describe connections between these hypergeometric functions and weight 3 modular forms, including a connection to modular Galois representations for certain cases. In §7.2, we interpret special values of these hypergeometric functions which are associated with rows [5.2 - 5.6] in Table 1, and relate them to special L -values of weight 3 modular forms. Finally, in §7.3, we discuss a modulo p^3 supercongruence conjecture for $f_{0, (\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}$. Interested readers may consult works such as [2, 48, 49, 52, 68] for relevant discussions.

7.1. Connections with modular Galois representations. By work of Freitas, Hung, and Siksek [21], elliptic curves over real quadratic fields are modular. The symmetric square $\rho_{\lambda, \ell} := \text{Sym}^2(\rho_{E, \ell})$ of the compatible family of Galois representations $\rho_{E, \ell}$ attached to $E := \tilde{E}_d \left(\frac{1-\sqrt{1-\lambda}}{2} \right)$ is also modular. When E has CM, $\rho_{\lambda, \ell}$ can be described in terms of a Grossencharacter attached to E and corresponds to a weight-3 CM cusp form by Weil's Converse Theorem [63, 47]. In this

subsection, we numerically identify the weight-3 forms whose Fourier coefficients satisfy the congruences derived from Theorem 2.3 attached to Examples 5.8 - 5.12. Further details are discussed in Appendix 8.2.

The following is a corollary to Theorem 2.3 using the theory of CM elliptic curves (see [57, Theorem 10.5] for example).

Corollary 7.1. *Assume the notation and assumptions in Theorem 2.3 and that $\lambda \in \mathbb{Q}$. Then there exists a weight-3 Hecke eigenform f , depending on d and λ , such that for almost all primes $p \nmid 2d$,*

$$\sum_{k=0}^{p-1} \frac{(\frac{1}{2})_k (\frac{1}{d})_k (\frac{d-1}{d})_k}{k!^3} \lambda^k \equiv a_p(f) \pmod{p^2},$$

where $a_p(f)$ is the p th Fourier coefficient of f .

We note that the supercongruence arising from $\lambda = 1$ in Corollary 7.1 is given by Mortenson [45].

Proof. By the modularity of CM elliptic curves over real quadratic fields, there exist Grossencharacters ψ_+ and ψ_- of such that the L -functions of ψ_+ and $E_+ := \tilde{E}_d\left(\frac{1+\sqrt{1-\lambda}}{2}\right)$, and respectively ψ_- and $E_- := \tilde{E}_d\left(\frac{1-\sqrt{1-\lambda}}{2}\right)$, coincide. Namely,

$$\begin{aligned} L(\psi_+, s) &= L(E_+/\mathbb{Q}(\sqrt{1-\lambda}), s), \\ L(\psi_-, s) &= L(E_-/\mathbb{Q}(\sqrt{1-\lambda}), s). \end{aligned}$$

Let ψ_{k_d} be the character induced by the quadratic character $\left(\frac{k_d}{\cdot}\right)$. There exists a weight-3 CM modular form f with an imprimitive Dirichlet character χ determined by the CM discriminant corresponding to $\psi_{k_d}\psi_-\psi_+$. Hence, by Theorem 2.3 one has for supersingular primes ($\chi(p) = -1$) that

$$\sum_{k=0}^{p-1} \frac{(\frac{1}{2})_k (\frac{1}{d})_k (\frac{d-1}{d})_k}{k!^3} \lambda^k \equiv 0 = a_p(f) \pmod{p^2},$$

and for ordinary primes ($\chi(p) = 1$),

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{(\frac{1}{2})_k (\frac{1}{d})_k (\frac{d-1}{d})_k}{k!^3} \lambda^k &\equiv \begin{cases} u_p^2(E_-) & \text{if } \left(\frac{1-\lambda}{p}\right) = 1, \\ \left(\frac{k_d}{p}\right) u_p^2(E_-) & \text{if } \left(\frac{1-\lambda}{p}\right) = -1, \end{cases} \\ &\equiv \begin{cases} \left(\frac{k_d}{p}\right) u_p(E_-) u_p(E_+) & \text{if } \left(\frac{1-\lambda}{p}\right) = 1, \\ \left(\frac{k_d}{p}\right) u_p(E_-) u_p(E_+) & \text{if } \left(\frac{1-\lambda}{p}\right) = -1, \end{cases} \\ &\equiv u_p(f) + p^2/u_p(f) \pmod{p^2}, \end{aligned}$$

where $u_p(f)$ and $p^2/u_p(f)$ are the roots of the Hecke polynomial $T^2 - a_p(f)T + p^2$ attached to f at p . \square

Remark 7.2. *By the above discussion, we have modularity for the Galois representations of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ attached to $f_{0,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}(\lambda)$ arising from the tensor product of the Galois representations of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\sqrt{1-\lambda}))$ attached to the CM curves E_- and E_+ . In particular, denote $\rho_\ell := \rho_{E_-} \otimes \rho_{E_+}$. Then ρ_ℓ decomposes as*

$$\rho_\ell = \sigma_\ell \oplus \epsilon_\ell \oplus \epsilon_\ell,$$

where ϵ_ℓ is the cyclotomic character tensored by the CM quadratic character and σ_ℓ is a 2-dimensional representation having trace of Frobenius $\text{Tr}(\sigma_\ell(\text{Frob}_p)) = a_p(g)$ for a weight-3 form g with CM arising from the Grossencharacter $\psi_-\psi_+$ defined in the proof of Corollary 7.1. The relationship between

g and the cusp form f in the proof of Corollary 7.1 is $f = \left(\frac{k_d}{d}\right) \otimes g$, which follows from Theorem 2.3. In particular, if p is supersingular, since g has CM it follows that $a_p(g) = 0$. If p is ordinary, since the eigenvalues of $\rho_{E,\ell}$ are εu_p and $\varepsilon p/u_p$, with $\varepsilon^2 = 1$, the eigenvalues of ρ_ℓ are u_p^2 , p , p , and p^2/u_p^2 . Then

$$\text{Tr}(\sigma_\ell(\text{Frob}_p)) = u_p^2 + \frac{p^2}{u_p^2},$$

and therefore $a_p(g) = u_p^2 + p^2/u_p^2$.

We obtain Corollary 1.6 from specific examples of Corollary 7.1. When $a_p(f) \neq 0$ we conjecture further congruences modulo p^3 , which were alluded to in Remark 1.5.

Conjecture 7.3. *Assume the hypotheses in Corollary 7.1. For each $\left[f_{0,(\frac{1}{2}, \frac{1}{d}, 1-\frac{1}{d})}\right]_{p-1}(\lambda)$ appearing in Corollary 7.1 such that $a_p(f) \neq 0$,*

$$\left[f_{0,(\frac{1}{2}, \frac{1}{d}, 1-\frac{1}{d})}\right]_{p-1}(\lambda) \equiv u_p(f) \pmod{p^3},$$

where $u_p(f)$ is the unit root of the Hecke polynomial $T^2 - a_p(f)T + p^2$ associated to f .

Remark 7.4. *For the case $d = 6$, some of the modulo p^3 supercongruences in Conjecture 7.3 are confirmed by Beukers' work as expressed in Theorem 2.5 and the modularity coming from Corollary 7.1.*

7.2. Special values. In [6, Examples 5.2-5.6], some special values $Z(\gamma\tau_0) = f_{0,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}(X(\gamma\tau_0))^2$ are computed in terms of Gamma functions. Here we interpret these values in terms of Ω_K defined in (3.2).

In each of these cases, $Z(\tau) = f_{0,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}(X(\tau))^2$ is a weight 4 modular form. Thus as discussed in the proof of Lemma 3.3,

$$f_{0,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}(X(\gamma\tau_0))^2 \sim \Omega_K^4,$$

where K is the CM-field of the elliptic curve $\tilde{E}_d \left(\frac{1-\sqrt{1-X(\gamma\tau_0)}}{2}\right)$. In particular, the evaluations of $f_{0,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}(X(\gamma\tau_0))^2$ given in [6, Examples 5.2-5.6] yield the following equalities,

$$(7.1) \quad f_{0,(\frac{1}{12}, \frac{5}{12})} \left(\frac{27}{125}\right)^2 = f_{0,(\frac{1}{2}, \frac{1}{6}, \frac{5}{6})} \left(\frac{27}{125}\right) = \frac{\sqrt{5}}{8} \Omega_{\mathbb{Q}(\sqrt{-8})}^2$$

$$(7.2) \quad f_{0,(\frac{1}{12}, \frac{5}{12})} \left(\frac{4}{125}\right)^2 = f_{0,(\frac{1}{2}, \frac{1}{6}, \frac{5}{6})} \left(\frac{4}{125}\right) = \frac{\sqrt{15}}{3 \cdot \sqrt[3]{32}} \Omega_{\mathbb{Q}(\sqrt{-3})}^2$$

$$(7.3) \quad f_{0,(\frac{1}{8}, \frac{3}{8})} \left(\frac{1}{9}\right)^2 = f_{0,(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})} \left(\frac{1}{9}\right) = \frac{1}{\sqrt{24}} \Omega_{\mathbb{Q}(\sqrt{-24})}^2$$

$$(7.4) \quad f_{0,(\frac{1}{3}, \frac{1}{6})} \left(\frac{1}{2}\right)^2 = f_{0,(\frac{1}{2}, \frac{1}{3}, \frac{2}{3})} \left(\frac{1}{2}\right) = \frac{\sqrt{3}}{8} \Omega_{\mathbb{Q}(\sqrt{-24})}^2$$

$$(7.5) \quad f_{0,(\frac{1}{3}, \frac{1}{6})} \left(\frac{4}{125}\right)^2 = f_{0,(\frac{1}{2}, \frac{1}{3}, \frac{2}{3})} \left(\frac{4}{125}\right) = \frac{\sqrt{3}}{8} \Omega_{\mathbb{Q}(\sqrt{-15})}^2.$$

Remark 7.5. *We expect that the special values in (7.1)-(7.5) are each related to the L -value of a weight-3 CM modular form through the modularity of CM elliptic curves. For example, by [37,*

Theorem 2] we have

$$\pi^2 \cdot f_{0,(\frac{1}{2}, \frac{1}{6}, \frac{5}{6})} \left(\frac{4}{125} \right) = 6\sqrt{5} \cdot L(\eta(2\tau)^3 \eta(6\tau)^3, 2) = 9\sqrt{5} \cdot L(\eta(6\tau)^4, 1)^2,$$

where for the elliptic curve $E : y^2 = x^3 + 1$,

$$L(\eta(6\tau)^4, 1) = L(E, 1) = \frac{\pi}{3^4 \sqrt{3} \sqrt[6]{32}} \Omega_{\mathbb{Q}(\sqrt{-3})}.$$

In particular, the following relation between weight-3 cusp forms, $f_{300.3.g.b} = \left(\frac{5}{\cdot} \right) \otimes \eta(2\tau)^3 \eta(6\tau)^3$, provides the algebraic relation

$$L(f_{300.3.g.b}, 2) \sim \sqrt{5} L(\eta(2\tau)^3 \eta(6\tau)^3, 2),$$

where $f_{300.3.g.b}$ is the weight-3 cusp form identified in Corollary 7.1. The exact identity can be determined explicitly, for example, with a modified \mathcal{G}_k Eisenstein series in [37], but we will not emphasize this here. This gives the identity

$$\pi^2 \cdot f_{0,(\frac{1}{2}, \frac{1}{6}, \frac{5}{6})} \left(\frac{4}{125} \right) = \frac{25}{4} L(f_{300.3.g.b}, 2).$$

For the case $\lambda = \pm 1$, some special L -values of the corresponding cusp forms at 1 and 2 are obtained in work of Zagier [66], Osburn and Straub [49], as well as Allen, Grove, Long, and the fifth author [3, Appendix] from a different approach. Here we list special values for the case $d = 2$,

$$\begin{aligned} f_{0,(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})}(1) &= \frac{8}{\pi} L(f_{16.3.c.a}, 1) = \frac{16}{\pi^2} L(f_{16.3.c.a}, 2), \\ f_{0,(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})}(-1) &= \frac{12}{\pi} L(f_{8.3.d.a}(\tau), 1) = \frac{12\sqrt{2}}{\pi^2} L(f_{8.3.d.a}(\tau), 2). \end{aligned}$$

In light of Remark 7.5 we make the following conjecture based on numeric evidence.

Conjecture 7.6. *The following identities hold,*

$$\begin{aligned} \pi^2 \cdot f_{0,(\frac{1}{2}, \frac{1}{6}, \frac{5}{6})} \left(\frac{27}{125} \right) &= \frac{25}{2} L(f_{800.3.g.a}, 2), \\ \pi^2 \cdot f_{0,(\frac{1}{2}, \frac{1}{6}, \frac{5}{6})} \left(\frac{4}{125} \right) &= \frac{25}{4} L(f_{300.3.g.b}, 2), \\ \pi^2 \cdot f_{0,(\frac{1}{2}, \frac{1}{4}, \frac{3}{4})} \left(\frac{1}{9} \right) &= 12 L(f_{24.3.h.a}, 2), \\ \pi^2 \cdot f_{0,(\frac{1}{2}, \frac{1}{3}, \frac{2}{3})} \left(\frac{1}{2} \right) &= 9 L(f_{24.3.h.b}, 2), \\ \pi^2 \cdot f_{0,(\frac{1}{2}, \frac{1}{3}, \frac{2}{3})} \left(\frac{4}{125} \right) &= \frac{45}{4} L(f_{15.3.d.b}, 2), \end{aligned}$$

where the subscripts on the modular forms f above are their LMFDB labels [38].

7.3. Conjectures modulo p^3 . We now examine which of the supercongruences presented in this paper appear to hold modulo p^3 and discuss some implications. As stated in Remark 2.2, Theorem 2.1 is conjectured to hold modulo p^3 . We further conjecture in the following that a special case of Theorem 1.4 holds modulo p^3 .

Conjecture 7.7. *Assume the hypotheses in Theorem 1.4 for $\mathbf{b} = (\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})$. Then for $p \geq 5$ prime,*

$$[f_{0,\mathbf{b}} \cdot f_{2\alpha_N, \mathbf{b}}]_{p-1}(\lambda) \equiv \begin{cases} 0 & \text{if } \tilde{E}_d\left(\frac{1-\sqrt{1-\lambda}}{2}\right) \text{ is supersingular at } p \\ u_p^2 & \text{if } \tilde{E}_d\left(\frac{1-\sqrt{1-\lambda}}{2}\right) \text{ is ordinary at } p \text{ and } \left(\frac{1-\lambda}{p}\right) = 1 \pmod{p^3}, \\ -\left(\frac{k_d}{p}\right) u_p^2 & \text{if } \tilde{E}_d\left(\frac{1-\sqrt{1-\lambda}}{2}\right) \text{ is ordinary at } p \text{ and } \left(\frac{1-\lambda}{p}\right) = -1 \end{cases}$$

where u_p is the unit root of the geometric Frobenius at p acting on the first cohomology of the elliptic curve $\tilde{E}_d\left(\frac{1-\sqrt{1-\lambda}}{2}\right)$ and $k_2 = k_6 = -1$, $k_3 = -3$, $k_4 = -2$.

We note that some of the cases of Conjecture 7.7 have been verified due to Theorem 2.5, see Examples 5.8 and 5.9. As stated in Remark 2.2, Zudilin has conjectured that Theorem 2.1 holds modulo p^3 . We next show that if that is true, it implies Conjecture 7.7.

Proposition 7.8. *If Theorem 2.1 holds modulo p^3 , then Conjecture 7.7 is true.*

Proof. If Theorem 2.1 holds modulo p^3 , then using Lemma 2.10 we have

$$[f_{0,\mathbf{b}} \cdot f_{2\alpha_N, \mathbf{b}}]_{p-1}(\lambda) \equiv \begin{cases} -\left(\frac{1-\lambda}{p}\right) [f_{0,\mathbf{b}}]_{p-1}(\lambda) \cdot p & \text{if } \tilde{E}_d\left(\frac{1-\sqrt{1-\lambda}}{2}\right) \text{ is supersingular at } p \\ [f_{0,\mathbf{b}}]_{p-1}(\lambda) \cdot p & \text{if } \tilde{E}_d\left(\frac{1-\sqrt{1-\lambda}}{2}\right) \text{ is ordinary at } p \text{ and } \left(\frac{1-\lambda}{p}\right) = 1 \pmod{p^3}. \\ -[f_{0,\mathbf{b}}]_{p-1}(\lambda) \cdot p & \text{if } \tilde{E}_d\left(\frac{1-\sqrt{1-\lambda}}{2}\right) \text{ is ordinary at } p \text{ and } \left(\frac{1-\lambda}{p}\right) = -1 \end{cases}$$

Thus applying Theorem 2.3 to $[f_{0,\mathbf{b}}]_{p-1}(\lambda) \cdot p$ to obtain congruences modulo p^3 , we obtain the result. \square

Motivated by Examples 5.8 - 5.12, we numerically verified that the following congruences hold modulo p^3 for applicable primes $p < 150$.

$$(7.6) \quad \left[f_{8, \left(\frac{1}{2}, \frac{1}{4}, \frac{3}{4}\right)} \right]_{p-1} \left(\frac{1}{9} \right) \equiv \text{sgn} \cdot \left(\frac{1-\lambda}{p} \right) \cdot p \pmod{p^3},$$

for $\lambda = \frac{1}{9}$ and primes $5 \leq p < 150$;

$$(7.7) \quad \left[f_{6, \left(\frac{1}{2}, \frac{1}{3}, \frac{2}{3}\right)} \right]_{p-1} \left(\frac{1}{2} \right) \equiv \text{sgn} \cdot \left(\frac{1-\lambda}{p} \right) \cdot p \pmod{p^3},$$

for $\lambda = \frac{1}{2}$ and primes $5 \leq p < 150$;

$$(7.8) \quad \left[f_{\frac{33}{4}, \left(\frac{1}{2}, \frac{1}{3}, \frac{2}{3}\right)} \right]_{p-1} \left(\frac{4}{125} \right) \equiv \text{sgn} \cdot \left(\frac{1-\lambda}{p} \right) \cdot p \pmod{p^3},$$

for $\lambda = \frac{4}{125}$ and primes $7 \leq p < 150$, $p \neq 11$.

Remark 7.9. *By Proposition 7.8, if (7.6), (7.7), or (7.8) hold modulo p^3 , then Examples 5.10, 5.11, or 5.12, respectively hold modulo p^3 . However, we note that Theorem 2.3 only seems to hold modulo p^3 for ordinary primes. For supersingular primes, the result only holds modulo p^2 . We have checked this for the $f_{0,\mathbf{b}}(z)$ corresponding to Examples 5.10, 5.11, and 5.12 for the appropriate primes $p < 150$.*

Finally, when $\mathbf{b} = \left(\frac{1}{d}, \frac{d-1}{d}\right)$ in Theorem 1.4, the congruences do not seem to generally extend to a higher power of p . For instance, Examples 5.1, 5.2, 5.3, 5.4, 5.6 do not hold modulo p^3 for

appropriate primes $p < 150$. However, for appropriate primes $p < 150$ Example 5.7 does hold modulo p^3 .

8. APPENDIX — PROOF OF THEOREM 2.3 AND HYPERGEOMETRIC CHARACTER SUMS

In this appendix, we provide an explanation of Theorem 2.3 through the arithmetic properties of the hypergeometric elliptic families, and further record some parallel results in terms of finite hypergeometric functions introduced in [10], which allow us to see the supercongruence in Theorem 2.3 from the character sum perspective.

8.1. Proof of Theorem 2.3. When E is an elliptic curve defined over a finite field \mathbb{F}_q containing q elements, define

$$a_q(E) := q + 1 - \#E(\mathbb{F}_q).$$

Proposition 8.1. *Let $d \in \{2, 3, 4, 6\}$, $p \geq 5$ prime, and \mathbb{F}_q a finite field with $q = p^r$ elements. For $t \in \mathbb{F}_q$ such that both $\tilde{E}_d(t)$ and $\tilde{E}_d(1-t)$ are elliptic curves defined over \mathbb{F}_q , one has*

$$a_q(\tilde{E}_d(t)) = \phi_q(k_d) a_q(\tilde{E}_d(1-t)),$$

where ϕ_q is the quadratic character of \mathbb{F}_q , and $k_2 = k_6 = -1$, $k_3 = -3$, $k_4 = -2$.

We believe this proposition has been documented in literature, however we find the following approach enlightening as we provide a detailed description of isogenies among the elliptic curves. One may see [27, Lemma 4] for a hypergeometric approach. The way we approach this proposition is through the moduli interpretation of elliptic curves as complex tori, and the action of W provided in Table 2. The action of the normalizer

$$W : t(W\tau) = 1 - t(\tau)$$

gives an isogeny from the complex torus $\mathbb{C}/\mathbb{Z} + \mathbb{Z}\tau$ to $\mathbb{C}/\mathbb{Z} + \mathbb{Z}(W\tau)$ and the composition of this isogeny and its dual gives $W^2 = -k_d$ -multiplication. These isogenies induce the isogenies on the reduction of the given elliptic curves. For each case, the isogeny is a composition of a isogeny defined over \mathbb{F}_q and a quadratic twist. In the proof of Proposition 8.1 below, we provide the explicit isogenies and quadratic twists.

Proof of Prop. 8.1. Fix $t \in \mathbb{F}_q$, $d \in \{2, 3, 4, 6\}$ such that $\tilde{E}_d(t)$ and $\tilde{E}_d(1-t)$ are defined over \mathbb{F}_q . When $d = 2$, $\tilde{E}_2(t)$ is isomorphic to $\tilde{E}'_2(t) : -y^2 = x(x-1)(x-(1-t))$ by the map

$$(x, y) \mapsto (-x + 1, y).$$

Therefore,

$$\begin{aligned} a_q(\tilde{E}_2(t)) &= a_q(\tilde{E}'_2(t)) = q + 1 - \#(\tilde{E}'_2(t)(\mathbb{F}_q)) \\ &= \sum_{x \in \mathbb{F}_q} \phi_q(-x(x-1)(x-(1-t))) = \phi_q(-1) \sum_{x \in \mathbb{F}_q} \phi_q(x(x-1)(x-(1-t))) \\ &= \phi_q(-1) a_q(\tilde{E}_2(1-t)). \end{aligned}$$

When $d = 6$, $\tilde{E}_d(t)$ is isomorphic to $\tilde{E}'_d(t) : y^2 = x^3 - 27x + 54(1-2t)$ by the map

$$(x, y) \mapsto (6^2x + 3, 6^3y + 3 \cdot 6^2x),$$

and the curve $-y^2 = x^3 - 27x + 54(1-2t)$ is isomorphic to $\tilde{E}'_d(1-t) : y^2 = x^3 - 27x - 54(1-2t)$ via $(x, y) \mapsto (-x, -y)$. This gives

$$a_q(\tilde{E}_6(t)) = \phi_q(-1) a_q(\tilde{E}_6(1-t)).$$

When $d = 4$, $\tilde{E}_4(t)$ is 2-isogenous to the curve $\tilde{E}'_4(t) : y^2 = x(x^2 - 2x + (1 - t))$ by the map

$$(x, y) \mapsto \left(x + 1 + \frac{t}{4x}, y \left(1 - \frac{t}{4x^2} \right) \right),$$

and the quadratic twist of the curve $\tilde{E}'_4(t)$ by -2 is isomorphic to $\tilde{E}_4(1 - t)$ by the isomorphism from $E : -2y^2 = x(x^2 - 2x + (1 - t)) \rightarrow \tilde{E}_d(1 - t)$ given by

$$(x, y) \mapsto (-x/2, y/2).$$

Putting this all together, we get

$$\begin{aligned} a_q(\tilde{E}_d(t)) &= a_q(\tilde{E}'_d(t)) = q + 1 - \#(\tilde{E}'_d(t)(\mathbb{F}_q)) \\ &= \sum_{x \in \mathbb{F}_q} \phi_q(x(x^2 - 2x + (1 - t))) = \phi_q(-2) \sum_{x \in \mathbb{F}_p} \phi_q(-2x(x^2 - 2x + (1 - t))) \\ &= \phi_q(-2) a_q(\tilde{E}_d(1 - t)). \end{aligned}$$

Similarly, in the case of $d = 3$, the curve $\tilde{E}_3(t)$ has a Weierstrass equation $y^2 = x^3 + \frac{1}{4} \left(x + \frac{t}{27} \right)^2$ through the map $(x, y) \mapsto \left(x, y + \frac{1}{2} \left(x + \frac{t}{27} \right) \right)$, which is 3-isogenous to $\tilde{E}'_3(t) : y^2 = x^3 - \frac{3}{4} \left(x - \frac{1-t}{9} \right)^2$. The curve $\tilde{E}_3(1 - t)$ is isomorphic to the quadratic twist of $\tilde{E}'_3(t)$ by -3 via the isomorphism from $E : -3y^2 = x^3 - \frac{3}{4} \left(x - \frac{1-t}{9} \right)^2 \rightarrow \tilde{E}_3(1 - t)$ given by

$$(x, y) \mapsto (-x/3, y/3).$$

This gives the desired identity

$$a_q(\tilde{E}_3(t)) = \phi_q(-3) a_q(\tilde{E}_3(1 - t)).$$

□

For our next lemma, we p -adically relate the evaluations of the truncation of $f_{0,(\frac{1}{d}, \frac{d-1}{d})}$ at λ and $1 - \lambda$, respectively.

Lemma 8.2. *Let $d \in \{2, 3, 4, 6\}$, $p \geq 5$ prime, and $k_2 = k_6 = -1$, $k_3 = -3$, $k_4 = -2$. Then for $\lambda \in \overline{\mathbb{Q}}$ such that λ can be embedded in \mathbb{Z}_p ,*

$$[f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\lambda) \equiv \left(\frac{k_d}{p} \right) [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(1 - \lambda) \pmod{p^2}.$$

Furthermore, as polynomials in t ,

$$[f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(t) - \left(\frac{k_d}{p} \right) [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(1 - t) \in p^2 \mathbb{Z}_p[t],$$

$$[f_{0,(\frac{1}{2}, \frac{1}{2}, \frac{d-1}{d})}]_{p-1}(t) - \left(\frac{k_d}{p} \right) [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1} \left(\frac{1 + \sqrt{1-t}}{2} \right) \cdot [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1} \left(\frac{1 - \sqrt{1-t}}{2} \right) \in p^2 \mathbb{Z}_p[t].$$

Proof. Trivially, $[f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(0) = 1$. Moreover, one can show $[f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(1) \equiv \left(\frac{k_d}{p} \right) \pmod{p}$ by first using Lemma 2.9 and the Chu-Vandermonde identity to obtain

$$[f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(1) \equiv [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{r_d}(1) = \frac{\left(\frac{1}{d} \right)_{r_d}}{(1)_{r_d}} \pmod{p},$$

where $r_d \equiv -1/d \pmod{p}$ is defined in Section 2.2, then interpreting the right hand side in terms of p -adic Gamma functions and [40, Thm. 14], then using quadratic reciprocity. From counting

points on classical hypergeometric elliptic families (see [27, 39, 59] for example) it follows from Proposition 8.1 that as polynomials in $\mathbb{Z}_p[t]$,

$$(8.1) \quad [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(t) \equiv \left(\frac{k_d}{p}\right) [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(1-t) \pmod{p}.$$

Thus there exists $n(t) \in \mathbb{Z}_p[t]$ such that

$$(8.2) \quad [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(t) = \left(\frac{k_d}{p}\right) [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(1-t) + pn(t),$$

and squaring both sides shows that

$$(8.3) \quad [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}^2(t) \equiv [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}^2(1-t) + 2p \left(\frac{k_d}{p}\right) n(t) [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(1-t) \pmod{p^2}.$$

From [16, Lemma 18] and the fact that $4t(1-t)$ is invariant under the change of variable $t \mapsto 1-t$, it follows that

$$(8.4) \quad [f_{0,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}]_{p-1}(4t(1-t)) \equiv [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}^2(t) \equiv [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}^2(1-t) \pmod{p^2}.$$

We conclude from (8.4) and (8.3) that as polynomials in t , $n(t)[f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(1-t) \equiv 0 \pmod{p}$. However, clearly $[f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(t) \not\equiv 0 \pmod{p}$ as it has constant term 1. Thus, $n(t) \equiv 0 \pmod{p}$, so (8.2) gives that

$$(8.5) \quad [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(t) \equiv \left(\frac{k_d}{p}\right) [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(1-t) \pmod{p^2},$$

which yields the first and second statements. Furthermore, (8.5) and (8.4) give that

$$(8.6) \quad [f_{0,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}]_{p-1}(4t(1-t)) \equiv \left(\frac{k_d}{p}\right) [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(t) \cdot [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(1-t) \pmod{p^2}.$$

From here it remains to substitute $\frac{1+\sqrt{1-t}}{2}$ for t , but we need to establish that we obtain a polynomial in only t . This follows from the fact that

$$F(u, v) := [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(u) \cdot [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(v)$$

is a symmetric polynomial in $\mathbb{Z}_p[u, v]$, and hence it can be written as a polynomial in the elementary symmetric polynomials $u+v, uv$. Thus setting $u = \frac{1+\sqrt{1-t}}{2}$ and $v = \frac{1-\sqrt{1-t}}{2}$, there exists $G(x, y) \in \mathbb{Z}_p[x, y]$ such that $F(u, v) = G(u+v, uv) = G(1, t/4) \in \mathbb{Z}_p[t]$. Thus we can substitute $\frac{1+\sqrt{1-t}}{2}$ for t in (8.6) to conclude

$$[f_{0,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}]_{p-1}(t) - \left(\frac{k_d}{p}\right) [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1} \left(\frac{1+\sqrt{1-t}}{2}\right) \cdot [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1} \left(\frac{1-\sqrt{1-t}}{2}\right) \in p^2\mathbb{Z}_p[t].$$

□

We now give a proof of Theorem 2.3 for the ordinary cases. Some useful information and terminology can be found in [57, Chapter 2] and [31, Chapter 6] for example.

Proof of Theorem 2.3 for ordinary primes p . Let λ be a totally real number satisfying the assumptions of Theorem 2.3. Set $\mu = v_- := \frac{1-\sqrt{1-\lambda}}{2}$, $v_+ := \frac{1+\sqrt{1-\lambda}}{2}$, as well as $E_- := \tilde{E}_d(v_-) = \tilde{E}_d(\mu)$ and $E_+ := \tilde{E}_d(v_+)$. For a given prime p such that $\tilde{E}_d(\mu)$ has good reduction and is ordinary at p , let \mathfrak{p} be a prime ideal in the ring of integers of $\mathbb{Q}(\mu)$ lying above p , and $\mathbb{Q}_{\mathfrak{p}}(\mu)$ be the completion of $\mathbb{Q}(\mu)$ at the place \mathfrak{p} . The ring of integers of $\mathbb{Q}_{\mathfrak{p}}(\mu)$ is isomorphic to the ring A of Witt vectors of $\mathbb{F}_p(\sqrt{1-\lambda})$. In this proof, we view E_- and E_+ as elliptic curves over A .

We examine cases based on whether the prime p splits in $\mathbb{Q}(\mu)$, i.e. when $\left(\frac{1-\lambda}{p}\right) = 1$, or p is inert in $\mathbb{Q}(\mu)$, i.e. when $\left(\frac{1-\lambda}{p}\right) = -1$.

Case 1: $\left(\frac{1-\lambda}{p}\right) = 1$. In this case $\mathbb{Q}_p(\mu) = \mathbb{Q}_p(\sqrt{1-\lambda}) \simeq \mathbb{Q}_p$ and we consider E_- as an elliptic curve over $A = W(\mathbb{F}_p) \simeq \mathbb{Z}_p$. Recall that u_p and p/u_p are roots of $T^2 - a_p(E_-)T + p$, where $a_p(E_-)$ is defined in Proposition 8.1. The endomorphism ring $\text{End}(E_-)$ embeds in \mathbb{Z}_p , and since p splits in this case, $\sqrt{1-\lambda}$ is also embedded in \mathbb{Z}_p . We set π^* to represent the image of p/u_p in \mathbb{Z}_p .

Multiplication on E_- by π^* is a rational function of the local uniformizer $\xi := -\frac{x}{y}$, and there is an algebraic number $\Delta := \Delta_{d,\pi^*}$ with $\Delta^2 \equiv 1 \pmod{pA}$ such that $[\pi^*](\xi) \equiv \Delta \xi^p \pmod{pA}$. This gives rise to a degree- p Frobenius map Φ on A , which induces a morphism

$$\begin{aligned} \tilde{\Phi}^* : H_{DR}^1(\widehat{E}_-/A, (p)) &\longrightarrow H_{DR}^1(\widehat{E}_-/A, (p)) \\ \sum_{n \geq 1} \frac{c(n-1)}{n} \xi^n &\mapsto \sum_{n \geq 1} \frac{c(n-1)^\sigma}{n} \Phi(\xi)^n = \sum_{n \geq 1} \frac{c(n-1)}{n} \Delta^n \xi^{pn}, \end{aligned}$$

where \widehat{E}_- is defined as in Katz [31].

The characteristic polynomial of $\tilde{\Phi}^*$ is $T^2 - a_p(E_-)T + p$ which has π^* as a root, and thus π^* is one of the eigenvalues of $\tilde{\Phi}^*$. Analogous to the construction in Section 3.1, the induced action of $\text{End}(E_-)$ on the de Rham space $H_{DR}^1(E_-/A, (p))$ has two common eigenvectors, one of which corresponds to the class of the standard invariant holomorphic differential one form ω , which as in (3.6) can be represented via $\omega = \sum_n \frac{a(n-1)}{n} \xi^n$. By the Cayley-Hamilton Theorem, $\tilde{\Phi}^*(\omega) = \pi^* \omega$. Comparing the coefficients of ξ^p from the left and the right hand sides of $\sum_n \frac{a(n-1)}{n} \Delta^n \xi^{pn} = \pi^* \sum_n \frac{a(n-1)}{n} \xi^n$, we get

$$\Delta \equiv \pi^* \frac{a(p-1)}{p} \pmod{pA},$$

which gives

$$a(p-1) \equiv \Delta u_p \pmod{pA} \equiv \Delta u_p \pmod{p\mathbb{Z}_p}.$$

According to [19, 16, 59], we have

$$(8.7) \quad [f_{0, \frac{1}{d}, \frac{d-1}{d}}]_{p-1}(v_-) \equiv \Delta u_p \pmod{p^2}.$$

Since we are assuming $\left(\frac{1-\lambda}{p}\right) = 1$, we get from Lemma 8.2 and (8.5) that

$$(8.8) \quad [f_{0, (\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\lambda) \equiv [f_{0, \frac{1}{d}, \frac{d-1}{d}}]_{p-1}^2(v_-) \equiv \Delta^2 u_p \equiv u_p^2 \pmod{p^2},$$

which proves the claim in this case.

Case 2: $\left(\frac{1-\lambda}{p}\right) = -1$. In this case $\mathbb{Q}_p(\mu) = \mathbb{Q}_p(\sqrt{1-\lambda}) \simeq \mathbb{Q}_p(\zeta_{p^2-1})$, which is the unique unramified quadratic extension of \mathbb{Q}_p , and we view E_- and E_+ as elliptic curves over $A = W(\mathbb{F}_p) \simeq \mathbb{Z}_p[\sqrt{1-\lambda}]$. As before, the endomorphism ring $\text{End}(E_-)$ is embedded in $\mathbb{Z}_p \subset \mathbb{Z}_p[\sqrt{1-\lambda}]$, however in this case, $\sqrt{1-\lambda}$ can not be embedded in \mathbb{Z}_p . Let $\sigma \in \text{Aut}(A)$ be the automorphism of A lifted from the Frobenius automorphism $z \mapsto z^p$ on \mathbb{F}_p . Then, $a^\sigma := \sigma(a) \equiv a^p \pmod{pA}$ for all $a \in A$.

Analogous to Case 1, u_p^2 and p^2/u_p^2 are roots of $T^2 - a_{p^2}(E_-)T + p^2$ and belong to the CM field K , whose completion at p is \mathbb{Q}_p . Denoting from now on u_p and p/u_p as the embedding of these elements in $\mathbb{Q}_p(\sqrt{1-\lambda})$, we have $u_p^2 \in \mathbb{Z}_p$ and moreover, $u_p^2 \equiv a_{p^2}(E_-) \pmod{p\mathbb{Z}_p}$. Furthermore, since p is inert in this case, u_p is a p -adic unit root of the geometric Frobenius associated to both E_- as well as E_+ .

Note that the multiplication map $[p^2/u_p^2]$ on E_- or E_+ gives rise to a degree- p^2 Frobenius map for which there exists a degree- p map $[\pi^*]$ satisfying $[\pi^*]^2 = [p^2/u_p^2]$. Then $[\pi^*]$, which is an isogeny

from $E_+ \rightarrow E_-$ and from $E_- \rightarrow E_+$, serves as a rational function of ξ as stated in [16] (see §4.2 and §5). Moreover, associated to each of the curves E_+ and E_- there is an algebraic number $\Delta_{\pm} := \Delta_{d, \pi^*}$ with $\Delta_{\pm}^4 \equiv 1 \pmod{pA}$ and $\Delta_- \Delta_+ \equiv 1 \pmod{pA}$ such that on E_- , $[\pi^*](\xi) \equiv \Delta_- \xi^p \pmod{pA}$ and on E_+ , $[\pi^*](\xi) \equiv \Delta_+ \xi^p \pmod{pA}$. This gives rise to a degree- p map Φ on A that induces morphisms

$$H_{DR}^1(\widehat{E}_-/A, (p)) \xrightarrow{\tilde{\Phi}^*} H_{DR}^1(\widehat{E}_+/A, (p)) \xrightarrow{\tilde{\Phi}^*} H_{DR}^1(\widehat{E}_-/A, (p)),$$

defined by

$$\sum_{n \geq 1} \frac{c(n-1)}{n} \xi^n \mapsto \sum_{n \geq 1} \frac{c(n-1)^\sigma}{n} \Phi(\xi)^n = \sum_{n \geq 1} \frac{c(n-1)^\sigma}{n} \Delta_-^n \xi^{pn} \mapsto \sum_{n \geq 1} \frac{c(n-1)}{n} \Delta_-^n \Delta_+^n \xi^{p^2 n}.$$

Let $\omega_- := \sum_n \frac{a(n-1)}{n} \xi^n$ represent the class of the standard invariant holomorphic differential one form on E_-/A . Then $\tilde{\Phi}^*(\omega_-)$ belongs to the subspace of the holomorphic 1-forms in $H_{DR}^1(\widehat{E}_+/A, (p))$ generated by $\omega_+ := \sum_n \frac{a(n-1)^\sigma}{n} \xi^n$. In particular, for some $\varepsilon \in \{\pm 1\}$,

$$\tilde{\Phi}^*(\omega_-) = \frac{\varepsilon p}{u_p} \omega_+ = \frac{\varepsilon p}{u_p} \sum_n \frac{a(n-1)^\sigma}{n} \xi^n.$$

Hence, by comparing the p -th power coefficients, we get

$$a(p-1)^\sigma \equiv \varepsilon \Delta_- u_p \pmod{pA}.$$

Further by [40, 16], one has

$$(8.9) \quad \left([f_{0, \frac{1}{d}, \frac{d-1}{d}}]_{p-1}(v_-) \right)^\sigma = [f_{0, \frac{1}{d}, \frac{d-1}{d}}]_{p-1}(v_+) \equiv \varepsilon \Delta_- u_p \pmod{p^2 A}.$$

Similarly, since $\tilde{\Phi}^*(\omega_+) = \frac{\varepsilon p}{u_p} \omega_-$, the maps

$$H_{DR}^1(\widehat{E}_+/A, (p)) \xrightarrow{\tilde{\Phi}^*} H_{DR}^1(\widehat{E}_-/A, (p)) \xrightarrow{\tilde{\Phi}^*} H_{DR}^1(\widehat{E}_+/A, (p))$$

give that

$$(8.10) \quad [f_{0, \frac{1}{d}, \frac{d-1}{d}}]_{p-1}(v_-) \equiv \varepsilon \Delta_+ u_p \pmod{p^2 A}.$$

Combining (8.9) and (8.10), we obtain

$$[f_{0, \frac{1}{d}, \frac{d-1}{d}}]_{p-1}(v_+) [f_{0, \frac{1}{d}, \frac{d-1}{d}}]_{p-1}(v_-) \equiv u_p^2 \pmod{p^2 A}.$$

Since $u_p^2 \in \mathbb{Z}_p$, and using Lemma 8.2, it follows that

$$[f_{0, \frac{1}{d}, \frac{d-1}{d}}]_{p-1}(v_+) [f_{0, \frac{1}{d}, \frac{d-1}{d}}]_{p-1}(v_-) \in \mathbb{Z}_p[\lambda] = \mathbb{Z}_p.$$

Thus,

$$(8.11) \quad [f_{0, \frac{1}{d}, \frac{d-1}{d}}]_{p-1}(v_+) [f_{0, \frac{1}{d}, \frac{d-1}{d}}]_{p-1}(v_-) \equiv u_p^2 \pmod{p^2}.$$

Finally, from Lemma 8.2 and (8.11),

$$[f_{0, (\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\lambda) \equiv \left(\frac{k_d}{p} \right) [f_{0, (\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(v_-) \cdot [f_{0, (\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(v_+) \equiv \left(\frac{k_d}{p} \right) u_p^2 \pmod{p^2},$$

which completes the proof. \square

Remark 8.3. We note that a second common eigenvector ν of the action of $\text{End}(E_-)$ on $H_{DR}^1(\widehat{E}_-/A, (p))$ corresponds to the class of a differential of the second kind, and similarly, $\tilde{\Phi}^*(\nu) = \varepsilon u_p \nu$ for $\varepsilon \in \{\pm 1\}$. One can then use similar steps as in the proof above to show the claims in Proposition 3.4.

We note that the following arises as a corollary of Lemma 8.2.

Corollary 8.4. *Fix a prime $p \geq 5$. As a polynomial in t ,*

$$[f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1} \left(\frac{1 + \sqrt{1-t}}{2} \right) - \left(\frac{k_d}{p} \right) [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1} \left(\frac{1 - \sqrt{1-t}}{2} \right) \in \begin{cases} \sqrt{1-t} \cdot p^2 \mathbb{Z}_p[t], & \text{if } \left(\frac{k_d}{p} \right) = 1, \\ p^2 \mathbb{Z}_p[t], & \text{if } \left(\frac{k_d}{p} \right) = -1. \end{cases}$$

In particular, for $1 \neq t \in \mathbb{Z}_p$, in the case when $\left(\frac{k_d}{p} \right) = 1$,

$$\sum_{k=1}^{p-1} \frac{\left(\frac{1}{d} \right)_k \left(\frac{d-1}{d} \right)_k}{k!^2} \left(\frac{1}{2} \right)^k \sum_{j=0}^{\lfloor \frac{k-1}{2} \rfloor s} \binom{k}{2j+1} (1-t)^j \equiv 0 \pmod{p^2},$$

and when $\left(\frac{k_d}{p} \right) = -1$,

$$1 + \sum_{k=1}^{p-1} \frac{\left(\frac{1}{d} \right)_k \left(\frac{d-1}{d} \right)_k}{k!^2} \left(\frac{1}{2} \right)^k \sum_{j=0}^{\lfloor \frac{k}{2} \rfloor} \binom{k}{2j} (1-t)^j \equiv 0 \pmod{p^2}.$$

Proof. Set $s = \sqrt{1-t}$ and $u = \frac{1+s}{2}$ so that $1-u = \frac{1-s}{2}$. From the second statement of Lemma 8.2, we have

$$g(s) := [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(u) - \left(\frac{k_d}{p} \right) [f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(1-u) \in p^2 \mathbb{Z}_p[u] = p^2 \mathbb{Z}_p[s].$$

On the other hand, expanding $g(s)$ as a series in s yields

$$\begin{aligned} g(s) &= \sum_{k=0}^{p-1} \frac{\left(\frac{1}{d} \right)_k \left(\frac{d-1}{d} \right)_k}{k!^2} \left(u^k - \left(\frac{k_d}{p} \right) (1-u)^k \right) \\ &= \sum_{k=0}^{p-1} \frac{\left(\frac{1}{d} \right)_k \left(\frac{d-1}{d} \right)_k}{k!^2} \left(\frac{1}{2} \right)^k \sum_{j=0}^k \left((1+s)^k - \left(\frac{k_d}{p} \right) (1-s)^k \right) \\ &= \sum_{k=0}^{p-1} \frac{\left(\frac{1}{d} \right)_k \left(\frac{d-1}{d} \right)_k}{k!^2} \left(\frac{1}{2} \right)^k \sum_{j=0}^k \binom{k}{j} s^j \left(1 - (-1)^j \left(\frac{k_d}{p} \right) \right). \end{aligned}$$

Observe that

$$\sum_{j=0}^k \binom{k}{j} s^j \left(1 - (-1)^j \left(\frac{k_d}{p} \right) \right) = \begin{cases} 2 \sum_{\substack{j=0 \\ j \text{ odd}}}^k \binom{k}{j} s^j = 2s \sum_{j=0}^{\lfloor \frac{k-1}{2} \rfloor} \binom{k}{2j+1} s^{2j} & \text{if } \left(\frac{k_d}{p} \right) = 1, \\ 2 \sum_{\substack{j=0 \\ j \text{ even}}}^k \binom{k}{j} s^j = 2 \sum_{j=0}^{\lfloor \frac{k}{2} \rfloor} \binom{k}{2j} s^{2j} & \text{if } \left(\frac{k_d}{p} \right) = -1. \end{cases}$$

Hence,

$$g(s) = \begin{cases} 2\sqrt{1-t} \sum_{k=1}^{p-1} \frac{\left(\frac{1}{d}\right)_k \left(\frac{d-1}{d}\right)_k}{k!^2} \left(\frac{1}{2}\right)^k \sum_{j=0}^{\lfloor \frac{k-1}{2} \rfloor} \binom{k}{2j+1} (1-t)^j \in \sqrt{1-t} \cdot p^2 \mathbb{Z}_p[t] & \text{if } \left(\frac{kd}{p}\right) = 1, \\ 2 \sum_{k=1}^{p-1} \frac{\left(\frac{1}{d}\right)_k \left(\frac{d-1}{d}\right)_k}{k!^2} \left(\frac{1}{2}\right)^k \sum_{j=0}^{\lfloor \frac{k}{2} \rfloor} \binom{k}{2j} (1-t)^j \in p^2 \mathbb{Z}_p[t] & \text{if } \left(\frac{kd}{p}\right) = -1, \end{cases}$$

and the result follows. \square

8.2. A quick survey on corresponding hypergeometric character sums and supercongruences. In this section, we summarize the relationship between hypergeometric character sums and truncated hypergeometric functions, and the modularity of hypergeometric Galois representations (see [35, 27, 52, et al.] for example) attached to the data provided in this paper. We will use the H_q -character sums described in the work of Beukers, Cohen, and Mellit [10].

Let \mathbb{F}_q be a finite field of odd characteristic and use $\widehat{\mathbb{F}_q^\times}$ to denote the group of multiplicative characters of \mathbb{F}_q^\times . For a given pair of multisets $\alpha = \{a_1, \dots, a_n\}$ and $\beta = \{1, b_2, \dots, b_n\}$ with entries in \mathbb{Q}^\times , we denote the positive least common denominator of all a_i, b_j by $M := \text{lcd}(\alpha \cup \beta)$. For a finite field \mathbb{F}_q , we fix ω a generator of $\widehat{\mathbb{F}_q^\times}$. Following [10], for $q \equiv 1 \pmod{M}$ and any $t \in \mathbb{F}_q$, define

$$H_q(\alpha, \beta; t; \omega) := \frac{1}{1-q} \sum_{k=0}^{q-2} \omega^k ((-1)^{n-t}) \prod_{j=1}^n \frac{g(\omega^{(q-1)a_j+k})}{g(\omega^{(q-1)a_j})} \frac{g(\bar{\omega}^{(q-1)b_j+k})}{g(\bar{\omega}^{(q-1)b_j})},$$

where $\bar{\omega} = \omega^{-1}$ and $g(\chi) := \sum_{x \in \mathbb{F}_q} \Psi(x) \chi(x)$ is the Gauss sum of χ with respect to a fixed choice of nontrivial additive character Ψ of \mathbb{F}_q .

We say that the datum $HD = \{\alpha, \beta\}$ is defined over \mathbb{Q} if both $\prod_{j=1}^n (X - e^{2\pi i a_j})$ and $\prod_{j=1}^n (X - e^{2\pi i b_j})$ are in $\mathbb{Z}[X]$. In such a special situation, one can rewrite $H_q(\alpha, \beta; t; \omega)$ by using the reflection and multiplication formulas of Gauss sums [10, Theorem 1.3] and drop the condition $q \equiv 1 \pmod{M}$. Based on results of Katz and Beukers-Cohen-Mellit, there exist explicit algebraic varieties attached to such data and their corresponding Galois representations can be determined by $H_p(HD; t)$. For simplicity, we state these results for $\beta = \{1, \dots, 1\}$ below.

Theorem 8.5 (See [30, 32, 10, 35]). *Let ℓ be a prime. Given a datum $HD := \{\alpha = \{a_1, \dots, a_n\}, \beta = \{1, \dots, 1\}\}$ with $a_i \neq 1$, $M = \text{lcd}(\alpha)$, for any $\lambda \in \mathbb{Z}[\zeta_M, 1/M] \setminus \{0\}$, there exists an ℓ -adic Galois representation*

$$\rho_{HD, \ell} : G(M) := \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\zeta_M)) \rightarrow \text{GL}(W_\lambda),$$

unramified almost everywhere, such that at each prime ideal \wp of $\mathbb{Z}[\zeta_M, 1/(M\ell\lambda)]$ with norm $N(\wp)$,

$$\text{Tr} \rho_{HD, \ell}(\text{Frob}_\wp) = H_{N(\wp)}(\alpha, \beta; 1/\lambda; \omega_\wp),$$

where Frob_\wp stands for the geometric Frobenius conjugacy class of $G(M)$ at \wp . Moreover,

- (1) *When $\lambda \neq 1$, $\dim_{\overline{\mathbb{Q}_\ell}} W_\lambda = n$ and all roots of the characteristic polynomial of $\rho_{HD, \ell}(\text{Frob}_\wp)$ are algebraic with the same absolute value $N(\wp)^{(n-1)/2}$ under all Archimedean embeddings.*
- (2) *When $\lambda = 1$, $\dim_{\overline{\mathbb{Q}_\ell}} W_\lambda = n - 1$.*

Furthermore, if HD is defined over \mathbb{Q} and $\lambda \in \mathbb{Q}$, then there exists an ℓ -adic representation $\rho_{HD, \ell}^{BCM}$ of $G_{\mathbb{Q}} := \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ such that

$$\rho_{HD, \ell}^{BCM} |_{G(M)} \simeq \rho_{HD, \ell}$$

and for each prime $p \nmid M\ell$,

$$\text{Tr} \rho_{HD, \ell}^{BCM}(\text{Frob}_p) = H_p(\alpha, \beta; 1/\lambda).$$

We now summarize the relations among the hypergeometric functions in different settings based on the works of [2, 10, 16, 27, 45, et.al.] and Proposition 8.1 for the specific data in this paper. Letting ϕ_q be the quadratic character on \mathbb{F}_q , we rewrite the relevant H_q as

$$H_q \left(\left\{ \frac{1}{2}, \frac{1}{2} \right\}, \{1, 1\}; t; \omega \right) := \frac{\phi_q(-1)}{q(1-q)} \sum_{k=0}^{q-2} g(\phi_q \omega^k)^2 g(\bar{\omega}^k)^2 \omega^k(t),$$

$$H_q \left(\left\{ \frac{1}{2}, \frac{1}{2}, \frac{1}{2} \right\}, \{1, 1, 1\}; t; \omega \right) := \frac{g(\phi_q)}{q^2(q-1)} \sum_{k=0}^{q-2} g(\phi_q \omega^k)^3 g(\bar{\omega}^k)^3 \omega^k(-t),$$

and for $d \in \{3, 4, 6\}$,

$$H_q \left(\left\{ \frac{1}{d}, \frac{d-1}{d} \right\}, \{1, 1\}; t; \omega \right) := \frac{1}{1-q} \sum_{k=0}^{q-2} g(\bar{\omega}^k)^2 S_d(\omega^k) \omega^k(t),$$

$$H_q \left(\left\{ \frac{1}{2}, \frac{1}{d}, \frac{d-1}{d} \right\}, \{1, 1, 1\}; t; \omega \right) := \frac{\phi_q(-1)g(\phi_q)}{q(q-1)} \sum_{k=0}^{q-2} g(\phi_q \omega^k) g(\bar{\omega}^k)^3 S_d(\omega^k) \omega^k(-t),$$

where $S_d(\chi) := \prod_{n|d} \left(\frac{g(\chi^n)}{g(\chi)} \chi(n^{-n}) \right)^{\mu(d/n)}$ and $\mu(\cdot)$ is the Möbius μ -function. Precisely, we obtain the values in Table 3.

TABLE 3. $S_d(\chi)$ for $d \in \{3, 4, 6\}$

d	3	4	6
$S_d(\chi)$	$\frac{g(\chi^3)}{g(\chi)} \chi(3^{-3})$	$\frac{g(\chi^4)}{g(\chi^2)} \chi(2^{-6})$	$\frac{g(\chi)g(\chi^6)}{g(\chi^2)g(\chi^3)} \chi(2^{-4}3^{-3})$

Theorem 8.6. ([10, 27, 68] and Proposition 8.1) For $q = p^k$ with $p \geq 5$ prime, let $t \in \mathbb{F}_q$ and $t \neq 0, 1$ such that $\tilde{E}_d(t)$ in Table 2 is an elliptic curve defined over \mathbb{F}_q . Then

$$\#\tilde{E}_d(t)(\mathbb{F}_q) = q + 1 - H_q \left(\left\{ d, \frac{d-1}{d} \right\}, \{1, 1\}; t \right).$$

Furthermore,

$$H_q \left(\left\{ d, \frac{d-1}{d} \right\}, \{1, 1\}; t \right) = \phi_q(k_d) H_q \left(\left\{ d, \frac{d-1}{d} \right\}, \{1, 1\}; 1-t \right),$$

$$H_p \left(\left\{ d, \frac{d-1}{d} \right\}, \{1, 1\}; 1 \right) = \left(\frac{k_d}{p} \right),$$

where ϕ_q is the quadratic character of \mathbb{F}_q , and $k_2 = k_6 = -1$, $k_3 = -3$, $k_4 = -2$.

Building on the work of Evans-Greene [20] and Theorem 8.6, we have the following H_q -version of Clausen's formula. The Galois interpretation can be found in [35], for example.

Theorem 8.7 (H_p -Clausen formula [20, 27]). Let $p > 3$ be a prime and $t \in \mathbb{F}_p \setminus \{0, 1\}$. Let $\sqrt{1-t}$ be a fixed element of \mathbb{F}_{p^2} such that $\sqrt{1-t}^2 = 1-t$. Then

$$\left(\frac{1-t}{p} \right) H_p \left(\left\{ \frac{1}{2}, \frac{1}{2}, \frac{1}{2} \right\}, \{1, 1, 1\}; t \right) = H_p \left(\left\{ \frac{1}{4}, \frac{3}{4} \right\}, \{1, 1\}; \frac{t}{t-1} \right)^2 - p.$$

For $d = 2, 3, 4, 6$, we have

$$\begin{aligned} & H_p \left(\left\{ \frac{1}{2}, \frac{1}{d}, 1 - \frac{1}{d} \right\}, \{1, 1, 1\}; t \right) \\ &= \begin{cases} H_p \left(\left\{ \frac{1}{d}, 1 - \frac{1}{d} \right\}, \{1, 1\}; \frac{1 - \sqrt{1-t}}{2} \right)^2 - p, & \text{if } 1-t \text{ is a square in } \mathbb{F}_p, \\ \left(\frac{k_d}{p} \right) H_{p^2} \left(\left\{ \frac{1}{d}, 1 - \frac{1}{d} \right\}, \{1, 1\}; \frac{1 - \sqrt{1-t}}{2} \right) - p, & \text{if } 1-t \text{ is not a square in } \mathbb{F}_p, \end{cases} \\ & H_p \left(\left\{ \frac{1}{2}, \frac{1}{d}, 1 - \frac{1}{d} \right\}, \{1, 1, 1\}; 1 \right) = H_p \left(\left\{ \frac{1}{d}, 1 - \frac{1}{d} \right\}, \{1, 1\}; \frac{1}{2} \right)^2 - \left(1 + \left(\frac{k_d}{p} \right) \right) p, \end{aligned}$$

where $k_2 = k_6 = -1$, $k_3 = -3$, $k_4 = -2$.

Combining the previous information gives the following.

Theorem 8.8. *Assume the notation and assumptions in Theorem 2.1 and that $\lambda \in \mathbb{Q}$ is a CM-value. Further let K_{CM} be the discriminant of the corresponding CM field. Then for primes $p \geq 7$ such that $\lambda \in \mathbb{Z}_p^\times$,*

$$\begin{aligned} & H_p \left(\left\{ \frac{1}{2}, \frac{1}{d}, 1 - \frac{1}{d} \right\}, \{1, 1, 1\}; \lambda \right) - \left(\frac{K_{CM}}{p} \right) \left(\frac{1 - \lambda}{p} \right) p \\ &= \begin{cases} 0 & \text{if } \tilde{E}_d(u) \text{ is supersingular at } p \\ u_p^2 + p^2/u_p^2 & \text{if } \tilde{E}_d(u) \text{ is ordinary at } p \text{ and } \left(\frac{1-\lambda}{p} \right) = 1, \\ \left(\frac{k_d}{p} \right) (u_p^2 + p^2/u_p^2) & \text{if } \tilde{E}_d(u) \text{ is ordinary at } p \text{ and } \left(\frac{1-\lambda}{p} \right) = -1, \end{cases} \end{aligned}$$

and

$$[f_{0,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}]_{p-1}(\lambda) \equiv H_p \left(\left\{ \frac{1}{2}, \frac{1}{d}, 1 - \frac{1}{d} \right\}, \{1, 1, 1\}; \lambda \right) - \left(\frac{K_{CM}}{p} \right) \left(\frac{1 - \lambda}{p} \right) p \pmod{p^2},$$

where $u = \frac{1 - \sqrt{1-\lambda}}{2}$, u_p is a fixed unit root of the geometric Frobenius at p acting on the first cohomology of the elliptic curve $\tilde{E}_d(u)$, and $k_2 = k_6 = -1$, $k_3 = -3$, $k_4 = -2$.

Remark 8.9. *In general, one has*

$$[f_{0,(\frac{1}{d}, \frac{d-1}{d})}]_{p-1}(t) \equiv H_p \left(\left\{ \frac{1}{d}, 1 - \frac{1}{d} \right\}, \{1, 1\}; t \right) \pmod{p},$$

$$[f_{0,(\frac{1}{2}, \frac{1}{d}, \frac{d-1}{d})}]_{p-1}(t) \equiv H_p \left(\left\{ \frac{1}{2}, \frac{1}{d}, 1 - \frac{1}{d} \right\}, \{1, 1, 1\}; t \right) \pmod{p},$$

for $t \in \mathbb{Q}^\times$ with $t \in \mathbb{Z}_p^\times$, see [39, 52] for example.

Corollary 8.10. *Assume the notation and assumptions as above. Then, for primes $p \nmid 2d$ such that $\lambda \in \mathbb{Z}_p^\times$, there exists a weight-3 Hecke eigenform f depending on d and λ such that*

$$H_p \left(\left\{ \frac{1}{2}, \frac{1}{d}, 1 - \frac{1}{d} \right\}, \{1, 1, 1\}; \lambda \right) - \left(\frac{K_{CM}}{p} \right) \left(\frac{1 - \lambda}{p} \right) p = a_p(f),$$

where $a_p(f)$ is the p -th Fourier coefficient of f .

The modularity of the corresponding Galois representations in this corollary is essentially coming from the modularity of CM hypergeometric elliptic curves (or K3 surfaces [33]) in the background. The modularity of the hypergeometric representations arising from datum defined over \mathbb{Q} has been developed by numerous researchers (see [1, 7, 23, 35, 41, 44, 61, 46, 52, 54]). In recent work, Allen, Grove, Long, and the fifth author [4] establish the modularity of a certain type of representations, those that can be lifted to $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, from the commutative formal group law and an explicit

construction of the corresponding cusp forms via the modular forms on arithmetic triangle groups as discussed in this paper and for example in [9, 6, 27].

REFERENCES

- [1] Scott Ahlgren and Ken Ono. Modularity of a certain Calabi-Yau threefold. *Monatsh. Math.*, 129(3):177–190, 2000.
- [2] Scott Ahlgren, Ken Ono, and David Penniston. Zeta functions of an infinite family of $K3$ surfaces. *Amer. J. Math.*, 124(2):353–368, 2002.
- [3] Michael Allen, Brian Grove, Ling Long, and Fang-Ting Tu. The Explicit Hypergeometric-Modularity Method II, 2024 (preprint).
- [4] Michael Allen, Brian Grove, Ling Long, and Fang-Ting Tu. The Explicit Hypergeometric-Modularity Method I, arXiv:2404.00711 (2024).
- [5] George E Andrews, Richard Askey, Ranjan Roy, Ranjan Roy, and Richard Askey. *Special functions*, volume 71. Cambridge university press Cambridge, 1999.
- [6] Angelica Babei, Lea Beneish, Manami Roy, Holly Swisher, Bella Tobin, and Fang-Ting Tu. Generalized Ramanujan-Sato series arising from modular forms. In *Research directions in number theory—Women in Numbers V*, volume 33 of *Assoc. Women Math. Ser.*, pages 87–131. Springer, Cham, 2024.
- [7] Rupam Barman and Neelam Saikia. Certain character sums and hypergeometric series. *Pacific J. Math.*, 295(2):271–289, 2018.
- [8] Gustav Bauer. Von den Coefficienten der Reihen von Kugelfunctionen einer Variablen. *J. Reine Angew. Math.*, 56:101–121, 1859.
- [9] Frits Beukers. Supercongruences using modular forms, arXiv: 2403.03301 (2024).
- [10] Frits Beukers, Henri Cohen, and Anton Mellit. Finite hypergeometric functions. *Pure Appl. Math. Q.*, 11(4):559–589, 2015.
- [11] Jonathan M. Borwein and Peter B. Borwein. *Pi and the AGM*. Canadian Mathematical Society Series of Monographs and Advanced Texts. John Wiley & Sons Inc., New York, 1987. A study in analytic number theory and computational complexity, A Wiley-Interscience Publication.
- [12] Jan Hendrik Bruinier, Gerard van der Geer, Günter Harder, and Don Zagier. *The 1-2-3 of modular forms*. Universitext. Springer-Verlag, Berlin, 2008. Lectures from the Summer School on Modular Forms and their Applications held in Nordfjordeid, June 2004, Edited by Kristian Ranestad.
- [13] Heng Huat Chan, Song Heng Chan, and Zhiguo Liu. Domb’s numbers and Ramanujan-Sato type series for $1/\pi$. *Adv. Math.*, 186(2):396–410, 2004.
- [14] Imin Chen and Gleb Glebov. On Chudnovsky–Ramanujan type formulae. *The Ramanujan Journal*, 46(3):677–712, 2018.
- [15] Imin Chen, Gleb Glebov, and Ritesh Goenka. Chudnovsky–Ramanujan type formulae for non-compact arithmetic triangle groups. *J. Number Theory*, 241:603–654, 2022.
- [16] Sarah Chisholm, Alyson Deines, Ling Long, Gabriele Nebe, and Holly Swisher. p -adic analogues of Ramanujan type formulas for $1/\pi$. *Mathematics*, 1(1):9–30, 2013.
- [17] David V. Chudnovsky and Gregory V. Chudnovsky. Approximations and complex multiplication according to Ramanujan. In *Ramanujan revisited (Urbana-Champaign, Ill., 1987)*, pages 375–472. Academic Press, Boston, MA, 1988.
- [18] Henri Cohen and Fredrik Strömberg. *Modular forms*, volume 179 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2017. A classical approach.
- [19] M. J. Coster and L. Van Hamme. Supercongruences of Atkin and Swinnerton-Dyer type for Legendre polynomials. *J. Number Theory*, 38(3):265–286, 1991.
- [20] Ron Evans and John Greene. Clausen’s theorem and hypergeometric functions over finite fields. *Finite Fields Appl.*, 15(1):97–109, 2009.
- [21] Nuno Freitas, Bao V Le Hung, and Samir Siksek. Elliptic curves over real quadratic fields are modular. *Inventiones mathematicae*, 201(1):159–206, 2015.
- [22] Jenny Fuselier, Ling Long, Ravi Ramakrishna, Holly Swisher, and Fang-Ting Tu. Hypergeometric functions over finite fields. *Mem. Amer. Math. Soc.*, 280(1382):vii+124, 2022.
- [23] Jenny G. Fuselier and Dermot McCarthy. Hypergeometric type identities in the p -adic setting and modular forms. *Proc. Amer. Math. Soc.*, 144(4):1493–1508, 2016.
- [24] Jesús Guillera and Wadim Zudilin. “Divergent” Ramanujan-type supercongruences. *Proc. Amer. Math. Soc.*, 140(3):765–777, 2012.
- [25] Victor JW Guo and Wadim Zudilin. Ramanujan-type formulae for $1/\pi$: q -analogues. *Integral Transforms and Special Functions*, 29(7):505–513, 2018.

- [26] Emma Haruka Iwao. Even more pi in the sky: Calculating 100 trillion digits of pi on google cloud, 2022. <https://cloud.google.com/blog/products/compute/calculating-100-trillion-digits-of-pi-on-google-cloud>.
- [27] Jerome W. Hoffman, Wen-Ching Winnie Li, Ling Long, and Fang-Ting Tu. Traces of Hecke Operators via Hypergeometric Character Sums, 2024z; arXiv: 2408.02918.
- [28] Nicholas M. Katz. The regularity theorem in algebraic geometry. In *Actes du Congrès International des Mathématiciens (Nice, 1970), Tome 1,*, pages 437–443. , 1971.
- [29] Nicholas M. Katz. Algebraic solutions of differential equations (p -curvature and the Hodge filtration). *Invent. Math.*, 18:1–118, 1972.
- [30] Nicholas M. Katz. p -adic properties of modular schemes and modular forms. In *Modular functions of one variable, III (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972)*, volume Vol. 350 of *Lecture Notes in Math.*, pages 69–190. Springer, Berlin-New York, 1973.
- [31] Nicholas M. Katz. Crystalline cohomology, Dieudonné modules, and Jacobi sums. In *Automorphic forms, representation theory and arithmetic (Bombay, 1979)*, volume 10 of *Tata Inst. Fundam. Res. Stud. Math.*, pages 165–246. Springer, Berlin-New York, 1981.
- [32] Nicholas M. Katz. Another look at the Dwork family. In *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II*, volume 270 of *Progr. Math.*, pages 89–126. Birkhäuser Boston, Boston, MA, 2009.
- [33] A. Klemm, W. Lerche, and P. Mayr. K_3 -fibrations and heterotic–type II string duality. *Phys. Lett. B*, 357(3):313–322, 1995.
- [34] Serge Lang. Elliptic functions. In *Elliptic Functions*, pages 5–21. Springer, 1987.
- [35] Wen-Ching W. Li, Ling Long, and Fang-Ting Tu. A Whipple ${}_7F_6$ formula revisited. *La Matematica*, 1-51, 2022.
- [36] Wen-Ching Winnie Li and Ling Long. Atkin and Swinnerton-Dyer congruences and noncongruence modular forms. In *Algebraic number theory and related topics 2012*, volume B51 of *RIMS Kôkyûroku Bessatsu*, pages 269–299. Res. Inst. Math. Sci. (RIMS), Kyoto, 2014.
- [37] Wen-Ching Winnie Li, Ling Long, and Fang-Ting Tu. Computing special L-values of certain modular forms with complex multiplication. *SIGMA 14 (2018), 090*, August 2018.
- [38] The LMFDB Collaboration. The L-functions and modular forms database. <http://www.lmfdb.org>, 2021. [Online; accessed 25 August 2021].
- [39] Ling Long. Some numeric hypergeometric supercongruences. In *Vertex operator algebras, number theory and related topics*, volume 753 of *Contemp. Math.*, pages 139–156. Amer. Math. Soc., Providence, RI, 2020.
- [40] Ling Long and Ravi Ramakrishna. Some supercongruences occurring in truncated hypergeometric series. *Advances in Mathematics*, 290:773–808, 2016.
- [41] Ling Long, Fang-Ting Tu, Noriko Yui, and Wadim Zudilin. Supercongruences for rigid hypergeometric Calabi-Yau threefolds. *Adv. Math.*, 393:Paper No. 108058, 49, 2021.
- [42] Yuri Ivanovich Manin. Algebraic curves over fields with differentiation. *Izvestiya Rossiiskoi Akademii Nauk. Seriya Matematicheskaya*, 22(6):737–756, 1958.
- [43] Daniel A. Marcus. *Number fields*. Universitext. Springer, Cham, 2018. Second edition of [MR0457396], With a foreword by Barry Mazur.
- [44] Dermot McCarthy and Matthew A. Papanikolas. A finite field hypergeometric function associated to eigenvalues of a Siegel eigenform. *Int. J. Number Theory*, 11(8):2431–2450, 2015.
- [45] Eric Mortenson. Supercongruences for truncated ${}_n+1F_n$ hypergeometric series with applications to certain weight three newforms. *Proc. Amer. Math. Soc.*, 133(2):321–330, 2005.
- [46] Eric Mortenson. A p -adic supercongruence conjecture of van Hamme. *Proc. Amer. Math. Soc.*, 136(12):4321–4328, 2008.
- [47] Andrew Ogg. *Modular forms and Dirichlet series*. W. A. Benjamin, Inc., New York-Amsterdam, 1969.
- [48] Ken Ono. *The web of modularity: arithmetic of the coefficients of modular forms and q -series*, volume 102 of *CBMS Regional Conference Series in Mathematics*. Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2004.
- [49] Robert Osburn and Armin Straub. Interpolated sequences and critical L -values of modular forms. In *Elliptic integrals, elliptic functions and modular forms in quantum field theory*, Texts Monogr. Symbol. Comput., pages 327–349. Springer, Cham, 2019.
- [50] Robert Osburn and Wadim Zudilin. On the (K.2) supercongruence of Van Hamme. *J. Math. Anal. Appl.*, 433(1):706–711, 2016.
- [51] Srinivasa Ramanujan. Modular equations and approximations to π [Quart. J. Math. **45** (1914), 350–372]. In *Collected papers of Srinivasa Ramanujan*, pages 23–39. AMS Chelsea Publ., Providence, RI, 2000.
- [52] David P. Roberts and Fernando Rodriguez Villegas. Hypergeometric motives. *Notices Amer. Math. Soc.*, 69(6):914–929, 2022.
- [53] Inc. SageMath. *CoCalc Collaborative Calculation Online*, 2016. <https://cocalc.com/>.

- [54] Adriana Salerno. Counting points over finite fields and hypergeometric functions. *Funct. Approx. Comment. Math.*, 49(1):137–157, 2013.
- [55] T Sato. Apéry numbers and ramanujan’s series for $1/\pi$. In *Abstract of a talk presented at the Annual meeting of the Mathematical Society of Japan*, pages 28–31, 2002.
- [56] Atle Selberg and S. Chowla. On Epstein’s zeta-function. *J. Reine Angew. Math.*, 227:86–110, 1967.
- [57] Joseph H. Silverman. *Advanced topics in the arithmetic of elliptic curves*, volume 151 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1994.
- [58] Joseph H Silverman. *The arithmetic of elliptic curves*, volume 106. Springer Science & Business Media, 2009.
- [59] Jan Stienstra and Frits Beukers. On the Picard-Fuchs equation and the formal Brauer group of certain elliptic $K3$ -surfaces. *Math. Ann.*, 271(2):269–304, 1985.
- [60] Holly Swisher. On the supercongruence conjectures of van Hamme. *Research in the Mathematical Sciences*, 2(1):1–21, 2015.
- [61] Mohit Tripathi and Jaban Meher. ${}_4F_3$ -Gaussian hypergeometric series and traces of Frobenius for elliptic curves. *Res. Math. Sci.*, 9(4):Paper No. 63, 24, 2022.
- [62] L. van Hamme. Some conjectures concerning partial sums of generalized hypergeometric series. In *p-adic functional analysis (Nijmegen, 1996)*, volume 192 of *Lecture Notes in Pure and Appl. Math.*, pages 223–236. Dekker, New York, 1997.
- [63] André Weil. Über die Bestimmung Dirichletscher Reihen durch Funktionalgleichungen. *Math. Ann.*, 168:149–156, 1967.
- [64] Yifan Yang. Schwarzian differential equations and Hecke eigenforms on Shimura curves. *Compositio Mathematica*, 149(1):1–31, 2013.
- [65] Masaaki Yoshida. *Fuchsian differential equations*, volume E11 of *Aspects of Mathematics*. Friedr. Vieweg & Sohn, Braunschweig, 1987. With special emphasis on the Gauss-Schwarz theory.
- [66] Don Zagier. Traces of singular moduli. In *Motives, polylogarithms and Hodge theory, Part I (Irvine, CA, 1998)*, volume 3 of *Int. Press Lect. Ser.*, pages 211–244. Int. Press, Somerville, MA, 2002.
- [67] Wadim Zudilin. Ramanujan-type supercongruences. *J. Number Theory*, 129(8):1848–1857, 2009.
- [68] Wadim Zudilin. A hypergeometric version of the modularity of rigid Calabi-Yau manifolds. *SIGMA Symmetry Integrability Geom. Methods Appl.*, 14:Paper No. 086, 16, 2018.

Email address: `babeiangelica@gmail.com`

DEPARTMENT OF MATHEMATICS, LAFAYETTE COLLEGE, EASTON, PA 18042, USA

Email address: `royma@lafayette.edu`

DEPARTMENT OF MATHEMATICS, OREGON STATE UNIVERSITY, CORVALLIS, OR 97301, USA

Email address: `swisherh@oregonstate.edu`

DEPARTMENT OF MATHEMATICS, AGNES SCOTT COLLEGE, DECATUR, GA 30030 USA

Email address: `btobin@agnesscott.edu`

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LA 70803, USA

Email address: `ftu@lsu.edu`