
Consistent estimation of generative model representations in the data kernel perspective space

Aranyak Acharyya
Johns Hopkins University
aachary6@jhu.edu

Michael W. Trosset
Indiana University
mtrosset@indiana.edu

Carey E. Priebe
Johns Hopkins University
cep@jhu.edu

Hayden S. Helm
Helivan Research
hayden@helivan.io

Abstract

Generative models, such as large language models and text-to-image diffusion models, produce relevant information when presented a query. Different models may produce different information when presented the same query. As the landscape of generative models evolves, it is important to develop techniques to study and analyze differences in model behaviour. In this paper we present novel theoretical results for embedding-based representations of generative models in the context of a set of queries. We establish sufficient conditions for the consistent estimation of the model embeddings in situations where the query set and the number of models grow.

1 Introduction

Generative models have gained popularity in natural language processing [4, 20], text-to-image generation [7], and code generation [24], as well as in other domains [1, 21]. The key feature of these models is their ability to generate relevant responses to wide-ranging queries. Based on differences such as pre-training data mixture, architecture, size, etc., the responses of different models (or of different generation configurations of the same model) may vary widely. Recently, empirical investigations have demonstrated the potential of embedding-based vector representations of models for capturing meaningful differences in model behavior in the context of a set of queries [10, 9, 12]. Following [12], the resulting representations are referred to as *perspectives* of the models and the space in which the representations reside is referred to as the *perspective space* of the models.

In this paper we provide theoretical justification for these successful empirical investigations by arguing that the induced perspective space is consistent for a population-level perspective space. In particular, we analyze three different – progressively more complicated – settings in which the perspective space can be used: i) fixed collection of models and fixed set of queries; ii) fixed collection of models and growing set of queries; and iii) growing collection of models and growing set of queries. For each setting we show that the multi-dimensional scaling with the raw stress criterion of a collection of matrix representations yields vectors that are consistent for an appropriately defined limiting configuration. Further, we provide sufficient conditions for consistency as a function of the generative properties of each model.

Our work is a part of two major bodies of literature: embeddings of inputs and outputs of generative models [15, 19, 16, 18] and embedding complex objects in suitable Euclidean spaces via multi-dimensional scaling [3, 23, 13, 5]. In particular, the perspective space is a minimizer of the raw stress criterion [14] for a dissimilarity matrix or function [22] defined on a set of matrix representations of generative models whose rows correspond to the average embedded response for a particular query.

Preprint.

The rest of the paper is organized as follows. In *Section 2* we introduce notation and review multi-dimensional scaling by the raw stress embedding criterion. We describe our setting in *Section 3* and then establish our main theoretical results in *Section 4*. We provide numerical evidence to support the theoretical results in *Section 5* and discuss our results, potential applications, and extensions in *Section 6*.

2 Preliminaries

2.1 Notations

Bold letters (such as \mathbf{B} or $\boldsymbol{\mu}$) are used to represent vectors and matrices. Any vector by default is a column vector. For a matrix \mathbf{B} , the j -th row is denoted by $(\mathbf{B})_{j\cdot}$, and the (i, i') -th entry is denoted by $\mathbf{B}_{ii'}$. Moreover, $\|\mathbf{B}\|_F$ denotes the Frobenius norm of the matrix \mathbf{B} . For any two vectors \mathbf{x} and \mathbf{y} , $\|\mathbf{x} - \mathbf{y}\|$ denotes the Euclidean distance between \mathbf{x} and \mathbf{y} . The set of $d \times d$ orthogonal matrices is denoted by $\mathcal{O}(d)$.

2.2 Multidimensional scaling by raw stress embedding

Multidimensional scaling ([3]) refers to the family of methods that produce vector representations of a set of objects from their pairwise dissimilarities. Raw stress embedding ([14, 22]) is a popular method for multidimensional scaling. For our purposes, we discuss raw stress embedding in the context of two scenarios: finite sample size and infinite sample size. The following is adapted from [21] for our specific setting of collections of generative models.

Finite sample size: Let us assume there are n objects and that the matrix $\boldsymbol{\Delta}^{(\infty)} \in \mathbb{R}^{n \times n}$ is such that $\boldsymbol{\Delta}_{ij}^{(\infty)}$ is the dissimilarity between the i -th object and the j -th object for $i, j \in [n]$. For a given embedding dimension $d \in \mathbb{N}$, our goal is to find vectors $\boldsymbol{\psi}_1, \dots, \boldsymbol{\psi}_n \in \mathbb{R}^d$ such that $\|\boldsymbol{\psi}_i - \boldsymbol{\psi}_j\| \approx \boldsymbol{\Delta}_{ij}^{(\infty)}$. We define solutions to this problem as:

$$(\boldsymbol{\psi}_1, \dots, \boldsymbol{\psi}_n) = \arg \min_{\mathbf{z}_i \in \mathbb{R}^d} \sum_{i, i'=1}^n \left(\|\mathbf{z}_i - \mathbf{z}_{i'}\| - \boldsymbol{\Delta}_{ii'}^{(\infty)} \right)^2. \quad (1)$$

We let $\boldsymbol{\psi} \in \mathbb{R}^{n \times d}$ denote the matrix whose i -th row is $\boldsymbol{\psi}_i$ and write $\boldsymbol{\psi} = \text{mds}(\boldsymbol{\Delta}^{(\infty)})$.

Remark 1. Given dissimilarity matrix $\boldsymbol{\Delta}^{(\infty)}$, $\boldsymbol{\psi} = \text{mds}(\boldsymbol{\Delta}^{(\infty)})$ is not necessarily unique. We denote the set of all solutions $\text{mds}(\boldsymbol{\Delta}^{(\infty)})$ by $\text{MDS}(\boldsymbol{\Delta}^{(\infty)})$.

Remark 2. Note that $\min_{\mathbf{z}_i \in \mathbb{R}^d} \sum_{i, i'=1}^n \left(\|\mathbf{z}_i - \mathbf{z}_{i'}\| - \boldsymbol{\Delta}_{ii'}^{(\infty)} \right)^2 = \min_{\tilde{\boldsymbol{\Delta}} \in \mathcal{Y}_n} \left\| \tilde{\boldsymbol{\Delta}} - \boldsymbol{\Delta}^{(\infty)} \right\|_F^2$, where \mathcal{Y}_n is the set of all $n \times n$ Euclidean distance matrices with Frobenius norm less than or equal to $\left\| \boldsymbol{\Delta}^{(\infty)} \right\|_F$, which is closed, bounded, and complete [22]. This guarantees the existence of a solution to Eq. (1).

Infinite sample size: In this case, the goal is to find Euclidean representations of all objects from a compact set. Since the number of objects is not necessarily finite, we may not be able to arrange the pairwise dissimilarities in a matrix and, instead, rely on the notion of a dissimilarity function. Let \mathcal{M} be a compact metric space, $\boldsymbol{\Delta}^{(\infty)} : \mathcal{M} \times \mathcal{M} \rightarrow \mathbb{R}_{\geq 0}$ be a dissimilarity function on \mathcal{M} , and $h : \mathcal{M} \rightarrow \mathbb{R}^d$ be a Borel-measurable embedding function. The *continuous raw stress criterion* [22] is defined by

$$\sigma((\boldsymbol{\Delta}^{(\infty)}, \mathcal{P}), h) = \int_{\mathcal{M}} \int_{\mathcal{M}} \left(\|h(\mathbf{m}') - h(\mathbf{m}'')\| - \boldsymbol{\Delta}^{(\infty)}(\mathbf{m}', \mathbf{m}'') \right)^2 \mathcal{P}(d\mathbf{m}') \mathcal{P}(d\mathbf{m}''). \quad (2)$$

We let $\text{mds} : \mathcal{M} \rightarrow \mathbb{R}^d$ denote the embedding function that minimizes $\sigma((\boldsymbol{\Delta}^{(\infty)}, \mathcal{P}), h)$ by varying h over the set of all possible Borel-measurable embedding functions from \mathcal{M} to \mathbb{R}^d .

Remark 3. Note that the function $\tilde{\boldsymbol{\Delta}} : \mathcal{M} \times \mathcal{M} \rightarrow \mathbb{R}$ defined as $\tilde{\boldsymbol{\Delta}}(\mathbf{m}', \mathbf{m}'') = \|h(\mathbf{m}') - h(\mathbf{m}'')\|$ is an Euclidean pseudometric on \mathcal{M} to \mathbb{R}^d . Since the set \mathcal{Y} of all Euclidean pseudometrics from \mathcal{M} to \mathbb{R}^d is closed and complete, $\sigma((\boldsymbol{\Delta}^{(\infty)}, \mathcal{P}), h)$ can be minimized by varying h over the set of all Borel-measurable embedding functions from $\mathcal{M} \rightarrow \mathbb{R}^d$.

3 Description of the setting

We consider a generative model f to be a (random) mapping from a query space \mathcal{Q} to an output space \mathcal{X} . Let f_1, \dots, f_n be n generative models with shared query and output spaces and $\{q_1, \dots, q_m\}$ be a collection of m queries. For our purposes, every model f_i responds to every query q_j exactly r times and we let $f_i(q_j)_k$ denote the k -th replicate of the response of f_i to q_j . We assume that there exists an embedding function g that maps a response to a vector in \mathbb{R}^s . For every (model, query) pair, we let F_{ij} be the distribution of embedded responses and assume $g(f_i(q_j)_k) \sim^{iid} F_{ij}$.

Let $\bar{\mathbf{X}}_i$ be the $m \times s$ matrix whose j^{th} row is the mean over replicates of the i^{th} model's response to the j^{th} query: $(\bar{\mathbf{X}}_i)_j = \frac{1}{r} \sum_{k=1}^r g(f_i(q_j)_k) \approx (\boldsymbol{\mu}_i)_j \equiv \mathbb{E}_{F_{ij}} [g(f_i(q_j))]$.

Further, let \mathbf{D} be the $n \times n$ pairwise distance matrix with entries $\mathbf{D}_{ii'} = \frac{1}{m} \|\bar{\mathbf{X}}_i - \bar{\mathbf{X}}_{i'}\|_F$. Then, as $r \rightarrow \infty$, $\mathbf{D}_{ii'}$ converges to the model mean discrepancy $\boldsymbol{\Delta}_{ii'} = \frac{1}{m} \|\boldsymbol{\mu}_i - \boldsymbol{\mu}_{i'}\|_F$.

Multidimensional scaling by raw stress minimization yields $\hat{\boldsymbol{\psi}} = \text{mds}(\mathbf{D}) \in \mathbb{R}^{n \times d}$, representing our collection of models as a configuration of n points in finite-dimensional Euclidean space \mathbb{R}^d . The geometry of $\hat{\boldsymbol{\psi}}$ approximates the model discrepancy geometry configuration matrix $\boldsymbol{\psi} = \text{mds}(\boldsymbol{\Delta}) \in \mathbb{R}^{n \times d}$. We refer to the i -th row $\boldsymbol{\psi}_i$ (respectively $\hat{\boldsymbol{\psi}}_i$) of the matrix $\boldsymbol{\psi}$ (respectively $\hat{\boldsymbol{\psi}}$) as (our estimate of) the *data kernel perspective space* (DKPS) representation of model f_i with respect to the set of queries $\{q_j\}$. The remainder of this paper details the consistency of the estimated DKPS as n, m and/or r grow.

Remark 4. In every setting we discuss, $r \rightarrow \infty$. For quantities whose definition depends on r , such as $\hat{\boldsymbol{\psi}}$, we sometimes emphasize the dependence by writing $\hat{\boldsymbol{\psi}}^{(r)}$.

4 Consistency of estimated model embeddings

The consistency results herein are presented in progressively more complicated settings where the number of queries and/or the number of models grow with a growing number of replicates from each F_{ij} .

4.1 Fixed set of models and fixed set of queries

We start with the simplest setting where the number of models and the number of queries both remain fixed and the number of replicates increases. In this setting, we can establish the consistency of $\hat{\boldsymbol{\psi}}$ (up to an affine transformation) by direct use of *Theorem 2* from [22], which establishes the convergence guarantees for raw stress embeddings of a sequence of dissimilarity matrices of a fixed size approaching another dissimilarity matrix.

Theorem 1. Let $\hat{\boldsymbol{\psi}} \in \text{MDS}(\mathbf{D}) \subset \mathbb{R}^{n \times d}$. Then there exists a subsequence $\{r_u\}_{u=1}^\infty$ of $\{r\}_{r=1}^\infty$ such that for all pairs $(i, i') \in [n]^2$,

$$\left(\left\| \hat{\boldsymbol{\psi}}_i^{(r_u)} - \hat{\boldsymbol{\psi}}_{i'}^{(r_u)} \right\| - \|\boldsymbol{\psi}_i - \boldsymbol{\psi}_{i'}\| \right) \xrightarrow{P} 0 \text{ as } u \rightarrow \infty,$$

where $\boldsymbol{\psi} = [\boldsymbol{\psi}_1 | \boldsymbol{\psi}_2 | \dots | \boldsymbol{\psi}_n]^T \in \text{MDS}(\boldsymbol{\Delta}) \subset \mathbb{R}^{n \times d}$.

4.2 Fixed set of models and growing set of queries

We next address the consistency of $\hat{\boldsymbol{\psi}}$ in settings where the number of queries grows but the collection of models is fixed. That is, n remains fixed but $m \rightarrow \infty$ as $r \rightarrow \infty$. Our results are adapted from the results from [22], where they show that if a sequence of dissimilarity matrices of a fixed size converges, then the raw stress embeddings of each term in the sequence approaches the raw stress embeddings of the limiting dissimilarity matrix. While we could directly use this result to argue that $\hat{\boldsymbol{\psi}}$ approaches $\boldsymbol{\psi}$ for fixed n and m where $\mathbf{D}_{ii'} \rightarrow \boldsymbol{\Delta}_{ii'}$ in *Theorem 1*, we cannot do so in the context of growing m . In particular, as m increases, so too does the dimensionality of $\boldsymbol{\Delta}$. Hence, in order to satisfy the condition that \mathbf{D} approaches a specific dissimilarity matrix as $m, r \rightarrow \infty$, we need an additional assumption:

Assumption 1. For some $q \in \mathbb{N}$, there exist vectors $\phi_1, \dots, \phi_n \in \mathbb{R}^q$, such that for every pair $(i, i') \in [n]^2$, $\boldsymbol{\Delta}_{ii'} = \frac{1}{m} \|\boldsymbol{\mu}_i - \boldsymbol{\mu}_{i'}\| \rightarrow \|\phi_i - \phi_{i'}\|$ as $r \rightarrow \infty$.

Assumption 1 presumes the existence of high-dimensional vector representations for each generative model. Our results will show that the sequence of low-dimensional raw stress configurations converges to a low-dimensional approximation of the $\{\phi_i\}$ under appropriate conditions. Let $\Delta^{(\infty)} = \{\|\phi_i - \phi_{i'}\|\}_{i,i'=1}^n \in \mathbb{R}^{n \times n}$ and $\psi = \text{mds}(\Delta^{(\infty)}) \in \text{MDS}(\Delta^{(\infty)}) \subset \mathbb{R}^{n \times d}$. Note that the vectors ϕ_i being independent of m and r ensures that the dissimilarity matrices \mathbf{D} approach a specific limit $\Delta^{(\infty)}$ as $m, r \rightarrow \infty$. Using results from [22], we show that if \mathbf{D} approaches the model mean discrepancy matrix $\Delta^{(\infty)}$ then $\hat{\psi}$ approaches ψ .

Lemma 1. ([22]) *Let $\hat{\psi} \in \text{MDS}(\mathbf{D}) \subset \mathbb{R}^{n \times d}$. Let n be fixed and let m grow with r . If $\|\mathbf{D} - \Delta^{(\infty)}\|_F \xrightarrow{P} 0$ as $r \rightarrow \infty$, then there exists a subsequence $\{r_u\}_{u=1}^\infty$ of $\{r\}_{r=1}^\infty$ such that for all pairs $(i, i') \in [n]^2$,*

$$\left(\left\| \hat{\psi}_i^{(r_u)} - \hat{\psi}_{i'}^{(r_u)} \right\| - \|\psi_i - \psi_{i'}\| \right) \xrightarrow{P} 0 \text{ as } u \rightarrow \infty,$$

where $\psi = [\psi_1 | \psi_2 | \dots | \psi_n]^T \in \text{MDS}(\Delta^{(\infty)}) \subset \mathbb{R}^{n \times d}$.

We note that by the weak law of large numbers we have $(\bar{\mathbf{X}}_i)_j \xrightarrow{P} (\mu_i)_j$ as $r \rightarrow \infty$. However, if m is large while r is relatively small, then there is non-zero probability that many rows in $\bar{\mathbf{X}}_i$ are far away from the corresponding rows in μ_i . Thus, if m grows too fast with respect to r , \mathbf{D} may not approach $\Delta^{(\infty)}$ and the condition of Lemma 1 will not be met. We next establish a sufficient condition for the rate at which m can grow with respect to r for the condition of Lemma 1.

Theorem 2. *Let $\Sigma_{ij} = \text{cov}[g(f_i(q_j)_k)]$ be the covariance matrix associated with the distribution F_{ij} and $\gamma_{ij} = \text{trace}(\Sigma_{ij})$. If, for all $i \in [n]$,*

$$\lim_{r \rightarrow \infty} \frac{\frac{1}{m} \sum_{j=1}^m \gamma_{ij}}{r} = 0$$

then $\|\mathbf{D} - \Delta^{(\infty)}\|_F \xrightarrow{P} 0$ as $r \rightarrow \infty$.

As a consequence of Lemma 1 and Theorem 2, a sufficient condition for consistency of $\hat{\psi}$ is $\frac{1}{m} \sum_{j=1}^m \gamma_{ij} = o(r)$ for all $i \in [n]$.

Theorem 3. *In the setting of Theorem 2, suppose for all $i \in [n]$, $\frac{1}{m} \sum_{j=1}^m \gamma_{ij} = o(r)$. Also, let $\hat{\psi} \in \text{MDS}(\mathbf{D}) \subset \mathbb{R}^{n \times d}$. There exists a subsequence $\{r_u\}_{u=1}^\infty$ of $\{r\}_{r=1}^\infty$ such that for all $(i, i') \in [n]^2$,*

$$\left(\left\| \hat{\psi}_i^{(r_u)} - \hat{\psi}_{i'}^{(r_u)} \right\| - \|\psi_i - \psi_{i'}\| \right) \xrightarrow{P} 0 \text{ as } u \rightarrow \infty,$$

where $\psi = [\psi_1 | \psi_2 | \dots | \psi_n]^T \in \text{MDS}(\Delta^{(\infty)}) \subset \mathbb{R}^{n \times d}$.

As a consequence of Theorem 3, each $\hat{\psi}_i$ will accumulate to some affine transformation of ψ_i , which is stated in the following corollary.

Corollary 1. *In the setting of Theorem 3, suppose for all $i \in [n]$, $\frac{1}{m} \sum_{j=1}^m \gamma_{ij} = o(r)$. Then there exist sequences $\{\mathbf{W}^{(u)}\}_{u=1}^\infty$ and $\{\mathbf{a}^{(u)}\}_{u=1}^\infty$, satisfying $\mathbf{W}^{(u)} \in \mathcal{O}(d)$ and $\mathbf{a}^{(u)} \in \mathbb{R}^d$ for all $u \in \mathbb{N}$, such that for all $i \in [n]$,*

$$\left\| \hat{\psi}_i^{(r_u)} - \left(\mathbf{W}^{(u)} \psi_i + \mathbf{a}^{(u)} \right) \right\| \xrightarrow{P} 0 \text{ as } u \rightarrow \infty.$$

Thus, Theorem 3 gives us an idea of how fast m can grow with respect to r as a function of F_{ij} , in order for $\hat{\psi}$ to be consistent. Observe that if $\max_{j \in [m]} \gamma_{ij} = O(1)$ for all $i \in [n]$, then m can grow arbitrarily fast with respect to r , and $\hat{\psi}$ will still be consistent.

4.3 Growing set of models and growing set of queries

We now let both the number of models and the number of queries grow as the number of replicates grow and establish that under appropriate conditions $\hat{\psi}$ is consistent. In this setting, we need an adaptation of *Assumption 1* that further assumes that the vectors that induce the limiting dissimilarity are elements of a compact Riemannian manifold:

Assumption 2. *Let \mathcal{M} be a compact Riemannian manifold. For every model f_i , there exists a vector $\phi_i \in \mathcal{M} \subset \mathbb{R}^q$, such that for all pairs $(i, i') \in \mathbb{N} \times \mathbb{N}$, $\frac{1}{m} \|\mu_i - \mu_{i'}\| \rightarrow \|\phi_i - \phi_{i'}\|$ as $r \rightarrow \infty$.*

Recall that in the growing sample size scenario, the definition of raw stress relies on a dissimilarity function. In our setting, we define the following dissimilarity function $\Delta^{(\infty)} : \mathcal{M} \times \mathcal{M} \rightarrow \mathbb{R}_{\geq 0}$ to be $\Delta^{(\infty)}(\phi_i, \phi_{i'}) = \|\phi_i - \phi_{i'}\|$. Results from [22] tell us that $\hat{\psi}$ is consistent if $\mathbf{D}_{ii'}$ approaches $\Delta^{(\infty)}(\phi_i, \phi_{i'})$ for all pairs $i, i' \in [n]$.

Lemma 2. ([22]) *Let $\phi_i \sim^{iid} \mathcal{P}$. Assume that for all pairs $(i, i') \in \mathbb{N} \times \mathbb{N}$, $\mathbf{D}_{ii'} \xrightarrow{P} \Delta^{(\infty)}(\phi_i, \phi_{i'})$ as $r \rightarrow \infty$. Then, for some subsequence $\{r_u\}_{u=1}^\infty$ of $\{r\}_{r=1}^\infty$, for all $p \geq 1$,*

$$\int_{\mathcal{M}} \int_{\mathcal{M}} \left(\left\| \hat{\psi}_1^{(r_u)} - \hat{\psi}_2^{(r_u)} \right\| - \|\text{mds}(\phi_1) - \text{mds}(\phi_2)\| \right)^p \mathcal{P}(d\phi_1) \mathcal{P}(d\phi_2) \xrightarrow{P} 0 \text{ as } u \rightarrow \infty.$$

The above theorem establishes that if the pairwise distances $\mathbf{D}_{ii'}$ approach the dissimilarities $\Delta^{(\infty)}(\phi_i, \phi_{i'})$, then $\hat{\psi}$ is consistent. In order for $\mathbf{D}_{ii'}$ to approach $\Delta^{(\infty)}(\phi_i, \phi_{i'})$, a result similar to *Theorem 2* is needed, which holds in the regime of growing n . Such a result can be shown to hold true by the same equipment used to establish *Theorem 2*.

Theorem 4. *In our setting where $m, n \rightarrow \infty$ as $r \rightarrow \infty$, $|\mathbf{D}_{ii'} - \Delta^{(\infty)}(\phi_i, \phi_{i'})| \xrightarrow{P} 0$ for every pair $(i, i') \in \mathbb{N} \times \mathbb{N}$, if*

$$\lim_{r \rightarrow \infty} \frac{\frac{1}{m} \sum_{j=1}^m \gamma_{ij}}{r} = 0 \text{ for all } i.$$

It may seem a bit surprising at first that the sufficient conditions in both scenarios are the same: The setting with growing m and n was expected to be more restrictive. Note that in both cases (of fixed n and growing n), it is pointwise convergence of $\mathbf{D}_{ii'}$ to $\Delta^{(\infty)}(\phi_i, \phi_{i'})$ that we need. That is why in each of the cases it suffices to ensure that $\frac{1}{m} \sum_{j=1}^m \gamma_{ij} = o(r)$ for all i . Further, the condition that for all i $\frac{1}{m} \sum_{j=1}^m \gamma_{ij} = o(r)$ is more difficult to ensure when n grows than when n is fixed.

As in *Section 4.2*, the lemmas in this section allow us to deduce a sufficient condition for the consistency of $\hat{\psi}$:

Theorem 5. *In the setting of Lemma 2, suppose for all $i \in \mathbb{N}$, $\frac{1}{m} \sum_{j=1}^m \gamma_{ij} = o(r)$. Then, for all $p \geq 1$, for some subsequence $\{r_u\}_{u=1}^\infty$ of $\{r\}_{r=1}^\infty$,*

$$\int_{\mathcal{M}} \int_{\mathcal{M}} \left(\left\| \hat{\psi}_1^{(r_u)} - \hat{\psi}_2^{(r_u)} \right\| - \|\text{mds}(\phi_1) - \text{mds}(\phi_2)\| \right)^p \mathcal{P}(d\phi_1) \mathcal{P}(d\phi_2) \xrightarrow{P} 0 \text{ as } u \rightarrow \infty.$$

Thus, even in the case where n grows, m is allowed to grow arbitrarily fast with respect to r if $\max_{j \in [m]} \gamma_{ij} = O(1)$ for all i . Moreover, since all we need is pointwise convergence of \mathbf{D} to $\Delta^{(\infty)}$, n can grow arbitrarily fast with respect to r .

5 Numerical experiments

We next provide empirical support of the consistency of the representations of models in the three settings we analyzed above. For each setting, we use LLaMA-3-8B-Chat [8] as a base model and consider up to 50 models. The models are parameterized by different fixed context augmentations a_i , i.e., $f_i = f(\cdot; a_i)$. Each a_i is a text string related to RA Fisher such as $a_i = \text{“RA Fisher pioneered the principles of the design of experiments.”}$ or $a_{i'} = \text{“RA Fisher’s view on eugenics were primarily based on anecdotes and prejudice.”}$ written by us. Given a query q_j , the base model is prompted with the appropriately formatted prompt $\text{“}a_i q_j\text{”}$ 50 times. In our experiments we consider up to 200 questions about RA Fisher such as $q_j = \text{“What is R.A. Fisher’s most well-known statistical theorem?”}$




Figure 1: Numerical evidence of the consistency of $\hat{\psi}$ to ψ for fixed $n = 10$ and fixed $m = 50$. The left figure shows the 2-d DKPS of 10 models induced with 20 queries, 50 replicates each, and the embedding function `NomicEmbedv1.5`. The red circles have radius equal to the average L_2 between $\hat{\psi}^*$ and model representations estimated with $r \in \{1, 5, 20\}$ replicates for each query. The right figure shows the distribution of the average L_2 norm in the DKPS across models for various r .

as queries. The questions were sampled from ChatGPT with the prompt “Provide 200 questions related to RA Fisher”. We use `NomicEmbedv1.5` [17] as the embedding model and `GrasPy`’s [6] implementation of multi-dimensional scaling throughout.

We do not know the “ground truth” DKPS ψ in any setting, and thus treat $\hat{\psi}$ estimated with $r = 50$ as a proxy for the ground truth. For settings with growing n and/or growing m , we similarly treat $\hat{\psi}^{(50)}$ estimated with $n = 50$ and $m = 200$ as the ground truth. We refer to these estimates as $\hat{\psi}^*$. We let $d = 2$ in all settings for visualization purposes. Finally, in settings where n , m , or r grow, we sample from the appropriate set of models, queries, or replicates with replacement to calculate $\hat{\psi}$ and report the average L_2 norm or average two-to-infinity norm of the difference between $\hat{\psi}^*$ and $\hat{\psi}\mathbf{W}$ of 10 bootstrap samples, where $\mathbf{W} \in \mathcal{O}(2)$ is the Procrustes solution [11].

5.1 Fixed collection of models and fixed set of queries

As with the theoretical analysis, our empirical analysis starts in the simplest setting where $n = 10$ and $m = 20$ are fixed while r grows. The collection of 10 models and set of 20 queries were selected at random without replacement. The target DKPS $\hat{\psi}^*$ is shown on the left of Figure 1. Each black dot represents a model. The red circles have radius equal to the average (across bootstrap samples) L_2 norm between $\hat{\psi}^*$ and $\hat{\psi}^{(r)}\mathbf{W}$ for selected r . The right figure of Figure 1 shows the distribution of the average (across bootstrap samples) L_2 norm for each model for more values of r . The decreasing average L_2 norm for all models as r increases supports Theorem 1.

5.2 Fixed collection of models and growing set of queries

We next consider the setting where $n = 10$ is fixed and both m and r grow. We use the same 10 models as above. The target DKPS $\hat{\psi}^*$ is shown on the left of Figure 2 and was estimated with $m = 200$ and $r = 50$. As with Figure 1, the red circles have radius equal to the average (across bootstrap samples) L_2 norm of the difference between $\hat{\psi}^*$ and $\hat{\psi}\mathbf{W}$ for selected (m, r) pairs. The right figure of Figure 2 shows the average two-to-infinity norm of the difference between $\hat{\psi}^*$ and $\hat{\psi}$ across bootstrap samples for more values of (m, r) . The two-to-infinity norm decreases as both (m, r) increase, which supports Theorem 3.

The right figure of Figure 2 shows the possibility of a computational trade-off between increasing m and increasing r . In particular, the computational cost of each iteration of the experiment is approximately $O(mr)$: getting more replicates of a fixed number of queries is approximately the same as getting a small set of replicates for more queries from a compute stand point. From an




Figure 2: Numerical evidence of the consistency of $\hat{\psi}$ to ψ for fixed $n = 20$ and growing m . The left figure shows the 2-d DKPS of 20 models induced with 200 queries and 50 replicates each. The red circles have radius equal to the average L_2 norm between the “ground truth” and model representations estimated for selected (m, r) pairs. The right figure shows the average maximum row L_2 norm for various (m, r) pairs.

estimation stand point, however, the right figure of Figure 2 shows that getting more queries may be more beneficial. Per Theorem 2, this observation depends on the distributional properties of each F_{ij} .

5.3 Growing collection of models and growing set of queries

Finally, we consider the setting with growing n and growing m . The target DKPS $\hat{\psi}^*$ was estimated using all 50 models, all 200 queries and all 50 replicates per query. For each (n, m, r) triple, we randomly sample n models, m queries, and r replicates without replacement and estimate the DKPS $\hat{\psi}$. Figure 3 shows the average two-to-infinity norm of the difference between $\hat{\psi}^*$ and $\hat{\psi}W$ across bootstrap samples for the n sampled models. As each element of the triple increases, the reported difference gets close to 0, which supports Theorem 5.

Similar to the difference in the effect on estimation between r and m , we observe a similar difference in the effect of estimation between r and n in Figure 3. In particular, increasing the number of models appears to have a larger effect on estimation than increasing r (or, even increasing m). As above, this observation depends on the F_{ij} , per Theorem 4. Unlike the observation in Figure 2, however, the computational cost of increasing n is larger than increasing m or r since $m \cdot r$ samples from F_{ij} are required for each additional n . Thus, there is a more delicate trade-off between computational cost and estimation quality in this setting than in the fixed n case.

6 Discussion

In [12], a novel method was proposed for embedding generative models into a finite-dimensional Euclidean space in the context of queries – the *perspective space*. In this paper, we analyzed the estimated perspective space in regimes where the number of models and/or the number of queries can remain fixed or grow and demonstrated that it is consistent for a population-level perspective space. Importantly, we also describe sufficient conditions for consistent estimation of the perspective space. In this regard, we establish that if the number of queries and/or models grow adequately slowly with the number of replicates relative to distributional properties of the F_{ij} then the perspective space can be estimated consistently.

Low-dimensional representations of collections of generative models enable the use of classical methods to understand differences in model behavior and, eventually, to make sense of model evolution. For example, [12] uses the perspective space to demonstrate that the communication structure underlying a system of interacting language models has an impact on system-level and model-level evolution. Going further, we could use the perspective space for entire classes of model-level inference problems such as predicting the the pre-training data mixture, predicting model safety,




Figure 3: Numerical evidence of the consistency of $\hat{\psi}$ to ψ for growing n and growing m . The target DKPS $\hat{\psi}$ was estimated using $n = 50$, $m = 200$, and $r = 50$.

predicting the model’s score on a benchmark, etc. Deeper investigations into these applications are warranted and our results support the perspective space as a principled foundation for these pursuits.

While the consistency of the estimated perspective space holds for all choices of the dimensionality of the raw stress embeddings, we note that the choice will impact practical properties of the estimate. In particular, choosing large d may result in slower convergence. Conversely, choosing small d may result in fast convergence but to a limiting set of vectors that poorly approximate the high-dimensional $\{\phi_i\}$.

Further, the consistency of the estimated perspectives is the least we could ask for [2]. We expect concentration inequalities and distributional results for the estimated perspectives to be important theoretical extensions to support applications in non-asymptotic regimes. For example, uniform convergence of the perspectives enable use of the estimated perspective space as a principled substitute for its population counterpart after a particular amount of models, queries, and replicates. Similarly, establishing distributional properties of the estimated perspectives, such as asymptotic normality, in particular settings as to practicable inference methods.

We required Assumptions 1 and 2 to establish consistency of $\hat{\psi}$ in regimes for growing m with either fixed or growing n . In particular, we assume the existence of a collection of vectors in a finite and fixed q -dimensional Euclidean space whose inner point distances are the limiting dissimilarities between the perspectives. These two assumptions put implicit constraints on how the queries grow and on the properties of each F_{ij} . An important and more general follow-on investigation may relax this assumption such that $\{\phi_i\}$ exist in a reproducing kernel Hilbert space.

We note that in most generative models in practice such as large language models with a maximum context size and maximum output sequence length and diffusion models that output RGB images, there are only a finite many possible input and output sequences or images. Hence, the maximum trace of the covariance matrices of $\{F_{ij}\}$ does not grow without bound and, importantly, we are in the regime where m or n can grow arbitrarily fast with respect to r .

Lastly, our results are for representations of models in the context of a set of queries. There are other potential vector representations of models that do not depend on a set of queries, such as model weights. The geometry of these quantities can similarly be obtained in settings where an appropriate dissimilarity is defined. In our setting, different sets of queries will approximate the non-query dependent geometry to different degrees. Investigations into how well different query sets approximate more inherent features of the models are necessary, as well as investigations into trade-offs between query-independent and query-dependent model representations.

Acknowledgements.

We would like to thank Avanti Athreya, Brandon Duderstadt, Youngser Park, and Zekun Wang for helpful discussions and comments throughout the development of this manuscript.

References

- [1] Sercan Ö Arık, Mike Chrzanowski, Adam Coates, Gregory Diamos, Andrew Gibiansky, Yongguo Kang, Xian Li, John Miller, Andrew Ng, Jonathan Raiman, et al. Deep voice: Real-time neural text-to-speech. In *International conference on machine learning*, pp. 195–204. PMLR, 2017.
- [2] Peter J Bickel and Kjell A Doksum. *Mathematical statistics: basic ideas and selected topics, volumes I-II package*. Chapman and Hall/CRC, 2015.
- [3] Ingwer Borg and Patrick JF Groenen. *Modern multidimensional scaling: Theory and applications*. Springer Science & Business Media, 2005.
- [4] Tom Brown, Benjamin Mann, Nick Ryder, Melanie Subbiah, Jared D Kaplan, Prafulla Dhariwal, Arvind Neelakantan, Pranav Shyam, Girish Sastry, Amanda Askell, et al. Language models are few-shot learners. *Advances in neural information processing systems*, 33:1877–1901, 2020.
- [5] Guodong Chen, Hayden S Helm, Kate Lytvynets, Weiwei Yang, and Carey E Priebe. Mental state classification using multi-graph features. *Frontiers in Human Neuroscience*, 16:930291, 2022.
- [6] Jaewon Chung, Benjamin D. Pedigo, Eric W. Bridgeford, Bijan K. Varjavand, Hayden S. Helm, and Joshua T. Vogelstein. Grasp: Graph statistics in python. *Journal of Machine Learning Research*, 20(158):1–7, 2019. URL <http://jmlr.org/papers/v20/19-490.html>.
- [7] Katherine Crowson, Stella Biderman, Daniel Kornis, Dashiell Stander, Eric Hallahan, Louis Castricato, and Edward Raff. Vqgan-clip: Open domain image generation and editing with natural language guidance. In *European Conference on Computer Vision*, pp. 88–105. Springer, 2022.
- [8] Abhimanyu Dubey, Abhinav Jauhri, Abhinav Pandey, Abhishek Kadian, Ahmad Al-Dahle, Aiesha Letman, Akhil Mathur, Alan Schelten, Amy Yang, Angela Fan, et al. The llama 3 herd of models. *arXiv preprint arXiv:2407.21783*, 2024.
- [9] Brandon Duderstadt, Hayden S. Helm, and Carey E. Priebe. Comparing foundation models using data kernels, 2024. URL <https://arxiv.org/abs/2305.05126>.
- [10] Guglielmo Faggioli, Laura Dietz, Charles LA Clarke, Gianluca Demartini, Matthias Hagen, Claudia Hauff, Noriko Kando, Evangelos Kanoulas, Martin Potthast, Benno Stein, et al. Perspectives on large language models for relevance judgment. In *Proceedings of the 2023 ACM SIGIR International Conference on Theory of Information Retrieval*, pp. 39–50, 2023.
- [11] Colin Goodall. Procrustes methods in the statistical analysis of shape. *Journal of the Royal Statistical Society: Series B (Methodological)*, 53(2):285–321, 1991.
- [12] Hayden Helm, Brandon Duderstadt, Youngser Park, and Carey E Priebe. Tracking the perspectives of interacting language models. *arXiv preprint arXiv:2406.11938*, 2024.
- [13] Hayden S Helm, Weiwei Yang, Sujeeth Bharadwaj, Kate Lytvynets, Oriana Riva, Christopher White, Ali Geisa, and Carey E Priebe. Inducing a hierarchy for multi-class classification problems. *arXiv preprint arXiv:2102.10263*, 2021.
- [14] Joseph B Kruskal. Multidimensional scaling by optimizing goodness of fit to a nonmetric hypothesis. *Psychometrika*, 29(1):1–27, 1964.
- [15] Tomas Mikolov. Efficient estimation of word representations in vector space. *arXiv preprint arXiv:1301.3781*, 2013.
- [16] Arvind Neelakantan, Tao Xu, Raul Puri, Alec Radford, Jesse Michael Han, Jerry Tworek, Qiming Yuan, Nikolas Tezak, Jong Wook Kim, Chris Hallacy, et al. Text and code embeddings by contrastive pre-training. *arXiv preprint arXiv:2201.10005*, 2022.
- [17] Zach Nussbaum, John X. Morris, Brandon Duderstadt, and Andriy Mulyar. Nomic embed: Training a reproducible long context text embedder, 2024. URL <https://arxiv.org/abs/2402.01613>.

- [18] Rajvardhan Patil, Sorio Boit, Venkat Gudivada, and Jagadeesh Nandigam. A survey of text representation and embedding techniques in nlp. *IEEE Access*, 11:36120–36146, 2023. doi: 10.1109/ACCESS.2023.3266377.
- [19] N Reimers. Sentence-bert: Sentence embeddings using siamese bert-networks. *arXiv preprint arXiv:1908.10084*, 2019.
- [20] Victor Sanh, Albert Webson, Colin Raffel, Stephen H Bach, Lintang Sutawika, Zaid Alyafeai, Antoine Chaffin, Arnaud Stiegler, Teven Le Scao, Arun Raja, et al. Multitask prompted training enables zero-shot task generalization. *arXiv preprint arXiv:2110.08207*, 2021.
- [21] Uriel Singer, Adam Polyak, Thomas Hayes, Xi Yin, Jie An, Songyang Zhang, Qiyuan Hu, Harry Yang, Oron Ashual, Oran Gafni, et al. Make-a-video: Text-to-video generation without text-video data. *arXiv preprint arXiv:2209.14792*, 2022.
- [22] Michael W Trosset and Carey E Priebe. Continuous multidimensional scaling. *arXiv preprint arXiv:2402.04436*, 2024.
- [23] Nian Wang, Robert J. Anderson, David G. Ashbrook, Vivek Gopalakrishnan, Youngser Park, Carey E. Priebe, Yi Qi, Rick Laoprasert, Joshua T. Vogelstein, Robert W. Williams, and G. Allan Johnson. Variability and heritability of mouse brain structure: Microscopic mri atlases and connectomes for diverse strains. *NeuroImage*, 222:117274, 2020. ISSN 1053-8119. doi: <https://doi.org/10.1016/j.neuroimage.2020.117274>. URL <https://www.sciencedirect.com/science/article/pii/S1053811920307606>.
- [24] Shun Zhang, Zhenfang Chen, Yikang Shen, Mingyu Ding, Joshua B Tenenbaum, and Chuang Gan. Planning with large language models for code generation. *arXiv preprint arXiv:2303.05510*, 2023.

A Proofs of lemmas and theorems

A.1 Fixed set of models and fixed set of queries

Theorem 1. Let $\widehat{\boldsymbol{\psi}} \in \text{MDS}(\mathbf{D}) \subset \mathbb{R}^{n \times d}$. Then there exists a subsequence $\{r_u\}_{u=1}^\infty$ of $\{r\}_{r=1}^\infty$ such that for all pairs $(i, i') \in [n]^2$,

$$\left(\left\| \widehat{\boldsymbol{\psi}}_i^{(r_u)} - \widehat{\boldsymbol{\psi}}_{i'}^{(r_u)} \right\| - \|\boldsymbol{\psi}_i - \boldsymbol{\psi}_{i'}\| \right) \xrightarrow{P} 0 \text{ as } u \rightarrow \infty,$$

where $\boldsymbol{\psi} = [\boldsymbol{\psi}_1 | \boldsymbol{\psi}_2 | \dots | \boldsymbol{\psi}_n]^T \in \text{MDS}(\boldsymbol{\Delta}) \subset \mathbb{R}^{n \times d}$.

Proof. See Theorem 2 of [22].

A.2 Fixed set of models and growing set of queries

Lemma 1. Let $\widehat{\boldsymbol{\psi}} \in \text{MDS}(\mathbf{D}) \subset \mathbb{R}^{n \times d}$. Let n be fixed and let m grow with r . If $\left\| \mathbf{D} - \boldsymbol{\Delta}^{(\infty)} \right\|_F \rightarrow 0$ as $r \rightarrow \infty$, then there exists a subsequence $\{r_u\}_{u=1}^\infty$ of $\{r\}_{r=1}^\infty$ such that for all pairs $(i, i') \in [n]^2$,

$$\left(\left\| \widehat{\boldsymbol{\psi}}_i^{(r_u)} - \widehat{\boldsymbol{\psi}}_{i'}^{(r_u)} \right\| - \|\boldsymbol{\psi}_i - \boldsymbol{\psi}_{i'}\| \right) \rightarrow 0 \text{ as } u \rightarrow \infty,$$

where $\boldsymbol{\psi} = [\boldsymbol{\psi}_1 | \boldsymbol{\psi}_2 | \dots | \boldsymbol{\psi}_n]^T \in \text{MDS}(\boldsymbol{\Delta}^{(\infty)}) \subset \mathbb{R}^{n \times d}$.

Proof. See Theorem 2 of [22].

Theorem 2. Let $\boldsymbol{\Sigma}_{ij} = \text{cov}[g(f_i(q_j)_k)]$ be the covariance matrix associated with the distribution F_{ij} and $\gamma_{ij} = \text{trace}(\boldsymbol{\Sigma}_{ij})$. If, for all $i \in [n]$,

$$\lim_{r \rightarrow \infty} \frac{\frac{1}{m} \sum_{j=1}^m \gamma_{ij}}{r} = 0$$

then $\left\| \mathbf{D} - \boldsymbol{\Delta}^{(\infty)} \right\|_F \xrightarrow{P} 0$ as $r \rightarrow \infty$.

Proof. Observe that by triangle inequality,

$$\begin{aligned} \|\bar{\mathbf{X}}_i - \bar{\mathbf{X}}_{i'}\|_F &\leq \|(\bar{\mathbf{X}}_i - \boldsymbol{\mu}_i) - (\bar{\mathbf{X}}_{i'} - \boldsymbol{\mu}_{i'})\|_F + \|\boldsymbol{\mu}_i - \boldsymbol{\mu}_{i'}\|_F \\ \implies \|\bar{\mathbf{X}}_i - \bar{\mathbf{X}}_{i'}\|_F - \|\boldsymbol{\mu}_i - \boldsymbol{\mu}_{i'}\|_F &\leq \|\bar{\mathbf{X}}_i - \boldsymbol{\mu}_i\|_F + \|\bar{\mathbf{X}}_{i'} - \boldsymbol{\mu}_{i'}\|_F. \end{aligned}$$

Similarly, we obtain,

$$\begin{aligned} \|\boldsymbol{\mu}_{i'} - \boldsymbol{\mu}_i\|_F &\leq \|(\bar{\mathbf{X}}_i - \boldsymbol{\mu}_i) - (\bar{\mathbf{X}}_{i'} - \boldsymbol{\mu}_{i'})\|_F + \|\bar{\mathbf{X}}_{i'} - \bar{\mathbf{X}}_i\|_F \\ \implies \|\boldsymbol{\mu}_{i'} - \boldsymbol{\mu}_i\|_F - \|\bar{\mathbf{X}}_{i'} - \bar{\mathbf{X}}_i\|_F &\leq \|\bar{\mathbf{X}}_i - \boldsymbol{\mu}_i\|_F + \|\bar{\mathbf{X}}_{i'} - \boldsymbol{\mu}_{i'}\|_F. \end{aligned}$$

Thus, from the above two equations combined, we get,

$$\left| \|\bar{\mathbf{X}}_i - \bar{\mathbf{X}}_{i'}\|_F - \|\boldsymbol{\mu}_i - \boldsymbol{\mu}_{i'}\|_F \right| \leq \|\bar{\mathbf{X}}_i - \boldsymbol{\mu}_i\|_F + \|\bar{\mathbf{X}}_{i'} - \boldsymbol{\mu}_{i'}\|_F. \quad (3)$$

From eqn (1),

$$|\mathbf{D}_{ii'} - \boldsymbol{\Delta}_{ii'}| \leq \frac{1}{m} \|\bar{\mathbf{X}}_i - \boldsymbol{\mu}_i\|_F + \frac{1}{m} \|\bar{\mathbf{X}}_{i'} - \boldsymbol{\mu}_{i'}\|_F. \quad (4)$$

Eqn (2) tells us that, in order to get $\|\mathbf{D} - \mathbf{\Delta}\|_F \rightarrow^P 0$, it suffices to have $\frac{1}{m} \|\bar{\mathbf{X}}_i - \boldsymbol{\mu}_i\|_F \rightarrow^P 0$ for all $i \in [n]$. Now, for an arbitrary ϵ ,

$$\begin{aligned} & \mathbb{P} \left[\frac{1}{m} \|\bar{\mathbf{X}}_i - \boldsymbol{\mu}_i\|_F > \epsilon \right] \\ & \leq \mathbb{P} \left[\sum_{j=1}^m \left\| (\bar{\mathbf{X}}_i)_{j\cdot} - (\boldsymbol{\mu}_i)_{j\cdot} \right\|^2 > m^2 \epsilon^2 \right] \\ & \leq \mathbb{P} \left[\bigcup_{j=1}^m \left\{ \left\| (\bar{\mathbf{X}}_i)_{j\cdot} - (\boldsymbol{\mu}_i)_{j\cdot} \right\|^2 > m \epsilon^2 \right\} \right] \\ & \leq \sum_{j=1}^m \mathbb{P} \left[\left\| (\bar{\mathbf{X}}_i)_{j\cdot} - (\boldsymbol{\mu}_i)_{j\cdot} \right\|^2 > m \epsilon^2 \right] \\ & \leq \sum_{j=1}^m \frac{\mathbb{E} \left\| (\bar{\mathbf{X}}_i)_{j\cdot} - (\boldsymbol{\mu}_i)_{j\cdot} \right\|^2}{m \epsilon^2}. \end{aligned}$$

Recall that $(\bar{\mathbf{X}}_i)_{j\cdot} = \frac{1}{r} \sum_{k=1}^r \mathbf{X}_{ijk}$ where $\mathbf{X}_{ijk} \sim^{iid} F_{ij}$ and the distribution F_{ij} has mean $(\boldsymbol{\mu}_i)_j$ and covariance matrix $\boldsymbol{\Sigma}_{ij}$. Thus, $\mathbb{E}[(\bar{\mathbf{X}}_i)_{j\cdot}] = (\boldsymbol{\mu}_i)_j$ and $\text{cov}[(\bar{\mathbf{X}}_i)_{j\cdot}] = \frac{1}{r} \boldsymbol{\Sigma}_{ij}$. Hence,

$$\mathbb{E} \left\| (\bar{\mathbf{X}}_i)_{j\cdot} - (\boldsymbol{\mu}_i)_{j\cdot} \right\|^2 = \text{trace}(\text{cov}[(\bar{\mathbf{X}}_i)_{j\cdot}]) = \text{trace} \left(\frac{1}{r} \boldsymbol{\Sigma}_{ij} \right) = \frac{\gamma_{ij}}{r}$$

where $\gamma_{ij} = \text{trace}(\boldsymbol{\Sigma}_{ij})$. Hence,

$$\mathbb{P} \left[\frac{1}{m} \|\bar{\mathbf{X}}_i - \boldsymbol{\mu}_i\|_F > \epsilon \right] \leq \sum_{j=1}^m \frac{\mathbb{E} \left\| (\bar{\mathbf{X}}_i)_{j\cdot} - (\boldsymbol{\mu}_i)_{j\cdot} \right\|^2}{m \epsilon^2} \leq \frac{\sum_{j=1}^m \gamma_{ij}}{r m \epsilon^2}.$$

Recall that as $r \rightarrow \infty$, $\|\mathbf{D} - \mathbf{\Delta}\|_F \rightarrow^P 0$ if $\frac{1}{m} \|\bar{\mathbf{X}}_i - \boldsymbol{\mu}_i\|_F \rightarrow^P 0$ for all $i \in [n]$.

Thus, $\lim_{r \rightarrow \infty} \frac{\frac{1}{m} \sum_{j=1}^m \gamma_{ij}}{r} = 0$ for all $i \in [n]$ ensures that $\|\mathbf{D} - \mathbf{\Delta}\|_F \rightarrow^P 0$, which again implies (using *Assumption 1* and Triangle Inequality) $\|\mathbf{D} - \mathbf{\Delta}^{(\infty)}\|_F \rightarrow^P 0$ as $r \rightarrow \infty$.

Theorem 3. Let $\boldsymbol{\Sigma}_{ij} = \text{cov}[g(f_i(q_j)_k)]$ be the covariance matrix associated with the distribution F_{ij} and $\gamma_{ij} = \text{trace}(\boldsymbol{\Sigma}_{ij})$. If, for all $i \in [n]$,

$$\lim_{r \rightarrow \infty} \frac{\frac{1}{m} \sum_{j=1}^m \gamma_{ij}}{r} = 0$$

then $\|\mathbf{D} - \mathbf{\Delta}^{(\infty)}\|_F \rightarrow^P 0$ as $r \rightarrow \infty$.

Proof. Obvious consequence of *Theorem 2* and *Theorem 3*.

A.3 Growing set of models and growing set of queries

Lemma 2. Let $\phi_i \sim^{iid} \mathcal{P}$. Assume that for all pairs $(i, i') \in \mathbb{N} \times \mathbb{N}$, $\mathbf{D}_{ii'} \rightarrow^P \mathbf{\Delta}^{(\infty)}(\phi_i, \phi_{i'})$ as $r \rightarrow \infty$. Then, for some subsequence $\{r_u\}_{u=1}^\infty$ of $\{r\}_{r=1}^\infty$, for all $p \geq 1$,

$$\int_{\mathcal{M}} \int_{\mathcal{M}} \left(\left\| \widehat{\boldsymbol{\psi}}_1^{(r_u)} - \widehat{\boldsymbol{\psi}}_2^{(r_u)} \right\| - \|\text{mds}(\phi_1) - \text{mds}(\phi_2)\| \right)^p \mathcal{P}(d\phi_1) \mathcal{P}(d\phi_2) \rightarrow^P 0 \text{ as } u \rightarrow \infty.$$

Proof. See *Theorem 3* of [22].

Theorem 4. In our setting where $m, n \rightarrow \infty$ as $r \rightarrow \infty$, $|\mathbf{D}_{ii'} - \mathbf{\Delta}^{(\infty)}(\phi_i, \phi_{i'})| \rightarrow^P 0$ for every pair $(i, i') \in \mathbb{N} \times \mathbb{N}$, if

$$\lim_{r \rightarrow \infty} \frac{\frac{1}{m} \sum_{j=1}^m \gamma_{ij}}{r} = 0 \text{ for all } i.$$

Proof. Same argument as *Theorem 3*.

Theorem 5. *In the setting of Lemma 2, suppose for all $i \in \mathbb{N}$, $\frac{1}{m} \sum_{j=1}^m \gamma_{ij} = o(r)$. Then, for all $p \geq 1$, for some subsequence $\{r_u\}_{u=1}^\infty$ of $\{r\}_{r=1}^\infty$,*

$$\int_{\mathcal{M}} \int_{\mathcal{M}} \left(\left\| \widehat{\psi}_1^{(r_u)} - \widehat{\psi}_2^{(r_u)} \right\| - \|\text{mds}(\phi_1) - \text{mds}(\phi_2)\| \right)^p \mathcal{P}(d\phi_1) \mathcal{P}(d\phi_2) \rightarrow^P 0 \text{ as } u \rightarrow \infty.$$

Proof. Obvious consequence of *Lemma 2* and *Theorem 4*.