

MUTATION-ACYCLIC QUIVERS ARE TOTALLY PROPER

SCOTT NEVILLE

ABSTRACT. Totally proper quivers, introduced by S. Fomin and the author [11], have many useful properties including powerful mutation invariants. We show that every mutation-acyclic quiver (i.e., a quiver that is mutation equivalent to an acyclic one) is totally proper. This yields new necessary conditions for a quiver to be mutation-acyclic.

1. INTRODUCTION

Quivers are finite directed graphs without oriented cycles of length 1 or 2. *Mutations* are operations that transform a quiver, based on a choice of a vertex. These notions are foundational in the study of cluster algebras [15]. A *mutation invariant* is a characteristic of a quiver that is preserved under mutations. Mutation invariants are helpful for deciding whether two quivers are mutation equivalent or not, i.e., whether there is a sequence of mutations that transforms one quiver into the other. See [5, 9, 14, 20, 22] for examples of known mutation invariants.

A *cyclically ordered quiver* (COQ) is a pair (Q, σ) where Q is a quiver and σ a cyclic ordering of its vertices. Cyclically ordered quivers were introduced in [11] to develop new powerful mutation invariants. A mutation in a COQ (Q, σ) transforms Q by the usual mutation rule, while simultaneously changing the cyclic ordering σ in a prescribed way. We note that mutations of COQs are only allowed at the vertices that satisfy a certain *properness* condition. It is this condition that ultimately enables the introduction of new mutation invariants.

In this paper, we prove that for one important class of quivers, the properness requirement can be lifted, so that mutations at all vertices are allowed and all invariants developed in [11] become true mutation invariants. Specifically, we show that in any *mutation-acyclic* quiver Q (i.e., a quiver that is mutation equivalent to an acyclic quiver), every vertex v is proper, for a particular canonical cyclic ordering σ on Q . After a mutation at v , we obtain a new COQ (Q', σ') , with canonical cyclic ordering σ' , that again has the same property: all the vertices are proper, so we can mutate at any one of them, and the process continues.

This theorem yields new mutation invariants of mutation-acyclic quivers, as well as new tools for proving that various quivers are not mutation-acyclic. One such quiver is shown in Figure 1. We give a short proof that this quiver is not mutation-acyclic in Example 6.11.

Date: September 27, 2024.

2020 Mathematics Subject Classification. Primary 13F60, Secondary 05E16, 15B36.

Key words and phrases. Quiver, mutation, cyclic ordering, Alexander polynomial.

Partially supported by NSF grants DMS-1840234, DMS-2054231, DMS-2348501.




Figure 1: An (unlabeled) triangular grid quiver with 4 vertices on each side. Readers may recognize it as the default quiver in B. Keller’s mutation applet [16].

As mentioned above, mutation at a vertex v in a COQ (Q, σ) is only allowed if v satisfies a combinatorial condition of “properness.” Informally, v is proper if every 2-arrow oriented path through v travels “clockwise,” i.e., in the direction of the cyclic ordering σ . A COQ is *proper* if every vertex v in it is proper (possibly after applying a sequence of transpositions called *wiggles*). Thus, we can mutate at any vertex in a proper COQ (Q, σ) and get a new COQ (Q', σ') —but (Q', σ') will not necessarily be proper. If it is the case that applying any sequence of mutations to (Q, σ) results in a proper COQ, then we say that (Q, σ) is *totally proper*.

We have shown in [11, Theorem 11.3, Remark 11.11] that a quiver Q can be upgraded to a totally proper COQ (Q, σ) in at most one way (up to wiggles). Moreover, the (essentially unique) candidate cyclic ordering $\sigma = \sigma_Q$ can be constructed efficiently. We say that a quiver Q is totally proper if the COQ (Q, σ_Q) is totally proper.

As shown in [11, Corollary 11.2], totally proper quivers have a powerful mutation invariant, which we recall next. This invariant is constructed as follows.

Input: a totally proper quiver Q .

Step 1. Construct the canonical cyclic ordering $\sigma = \sigma_Q$.

Step 2. “Tear” σ into a linear ordering $<$.

Step 3. Construct the skew-symmetric exchange matrix $B = B_Q$, with rows and columns ordered according to $<$.

Step 4. Construct the *unipotent companion* U , the unique unipotent upper-triangular matrix such that $B = U^T - U$.

Output: the integral congruence class of U , i.e., the set $\{GUG^T \mid G \in \mathrm{GL}_n(\mathbb{Z})\}$.

We have shown in [11] that wiggles, cyclic shifts of the linear ordering, and (proper) mutations of COQs preserve the integral congruence class of the unipotent companion. While the resulting invariant of proper mutations is very powerful, it is not easy to use in practice, since the problem of deciding whether two upper-triangular matrices are congruent over $\mathrm{GL}_n(\mathbb{Z})$ seems to be rather difficult. In [11], we bypassed this difficulty as follows. It is well known (and easy to see) that the integral congruence class of a matrix $U \in \mathrm{GL}_n(\mathbb{Z})$ uniquely determines the conjugacy class of its *cosquare* $U^{-T}U$. It follows that whenever two COQs are related by proper mutations, the cosquares of their respective unipotent companions must be conjugate in $\mathrm{GL}_n(\mathbb{Z})$. This conjugacy condition can be verified efficiently, though the algorithm is quite nontrivial [3, 7].

The cosquare $U^{-T}U$ can be used to construct other invariants of proper mutations, such as the monic characteristic polynomial of $U^{-T}U$, which we call the *Alexander polynomial* of the COQ Q . (While much simpler to compute, the Alexander polynomial is less powerful than the conjugacy class of $U^{-T}U$: integer matrices may have the same characteristic polynomial while not being conjugate in $\mathrm{GL}_n(\mathbb{Z})$.)

The main result of this paper is the following theorem that settles [11, Conjecture 12.8].

Theorem 1.1. *Any mutation-acyclic quiver is totally proper.*

Remark 1.2. It suffices to show that any acyclic quiver is totally proper. Furthermore, as noted in [11, Observation 10.7], any acyclic quiver has a proper cyclic ordering obtained by “closing up” any linear ordering compatible with the orientations of the quivers’ arrows. (All such cyclic orderings are related by wiggles.) Our proof of Theorem 1.1 shows that these cyclic orderings are totally proper.

In the course of proving Theorem 1.1, we establish a few results that may be of independent interest. In particular, Theorem 4.4 gives a simple combinatorial constraint on the orientations of arrows in mutation-acyclic quivers. Both Theorem 4.4 and the ensuing Lemma 5.5 assert that certain subquivers of a mutation-acyclic quiver must be acyclic. Should someone wish to compute with COQs and unipotent companions, Lemmas 4.1 and 4.2 generalize and remove conditions from [11, Lemma 7.2 and Remark 7.3]. This can speed up and simplify the computations.

An alternative proof of Theorem 1.1 was independently discovered by Hugh Thomas, using categorification [24].

Overview of the paper. Section 2 establishes notation and reviews some material that does not involve COQs. Much of this section is devoted to restating the results of A. Seven [21], who constructed a distinguished symmetric matrix A associated with an arbitrary mutation-acyclic quiver. Section 3 is a condensed summary of the necessary results and notation from [11]. It contains new examples, and Proposition 3.21 gives an alternative description of the Markov invariant. In Section 4 we establish a handful of technical lemmas. Lemmas 4.1 and 4.2 make it easier to compute the unipotent companion of a COQ after a proper mutation. Theorem 4.4 generalizes [11, Corollary 10.12] using the results of A. Seven [21]. Section 5 contains the proof of Theorem 1.1. We begin by defining $U = (A - B)/2$, where A comes from Seven’s construction and B is the usual exchange matrix. The fact that the integral congruence class of U is a mutation invariant follows quickly from Proposition 2.30. The bulk of the effort is then spent showing that U is indeed a unipotent companion. In Section 6, we give some applications and corollaries of Theorem 1.1. This includes a description of invariants for some small quivers, inequalities satisfied by the coefficients of the Alexander polynomial for mutation-acyclic quivers (Corollary 6.6), and several examples of quivers that are shown to be not mutation-acyclic.

Acknowledgements. I would like to thank my advisor, Sergey Fomin, for his guidance, comments, and suggestions; Grayson Moore, for reading and commenting on an early draft; and Danielle Ensign for her software assistance. I am also grateful to Roger Casals, Tucker Ervin, and Hugh Thomas for stimulating discussions.

I have used MAGMA and Sage for various computations. This research was supported in part through computational resources and services provided by Advanced Research Computing at the University of Michigan, Ann Arbor.

2. QUIVERS, MUTATION, AND QUASI-CARTAN COMPANIONS

We begin by reviewing notation, definitions and results from the literature which do not involve cyclically ordered quivers. This includes a summary of much from [21].

Definition 2.1. A *quiver* is a finite directed graph with parallel edges allowed, but no directed 1 or 2-cycles. Directed edges in a quiver are called *arrows*. Each vertex is marked as *mutable* or *frozen*. Unless otherwise indicated, all vertices are mutable.

We use the notation $u \rightarrow v$ to assert that there is at least one arrow from u to v . We use the notation $u \xrightarrow{x} v$ to denote that there are x arrows from vertex u to vertex v . Let $\text{In}(v) = \{u \mid u \rightarrow v\}$ and $\text{Out}(v) = \{u \mid u \leftarrow v\}$ respectively denote the inset and outset of v in a quiver.

Remark 2.2. By default, our quivers have labeled vertices. Thus, we distinguish between isomorphic quivers on the same set of vertices that differ from each other by a permutation of this set.

Definition 2.3. To *mutate* a quiver Q at a mutable vertex v , do the following:

- (1) for each oriented path $u \rightarrow v \rightarrow w$, draw a new arrow $u \rightarrow w$ (thus, if there are x arrows from $u \rightarrow v$ and y arrows $v \rightarrow w$, we add xy arrows $u \rightarrow w$);
- (2) reverse all arrows adjacent to v ;
- (3) remove oriented 2-cycles (one cycle at a time) until we again have a quiver.

We denote the resulting quiver by $\mu_v(Q)$. Mutation is an involution, $\mu_v(\mu_v(Q)) = Q$. Mutation does not change which vertices are mutable or frozen.

Definition 2.4. Two quivers are *mutation equivalent* if they are related by a sequence of mutations at mutable vertices. The *mutation class* $[Q]$ is the set of quivers mutation equivalent to Q .

Definition 2.5. A quiver is *acyclic* if it does not contain any directed cycles. A quiver is *mutation-acyclic* if it is mutation equivalent to an acyclic quiver.

Example 2.6. Only one of the two quivers in the Figure 2 is mutation-acyclic. Can you guess which is which?




Figure 2: Two quivers on 3-vertices. The quiver on the left *is not* mutation-acyclic, while the quiver on the right *is* mutation-acyclic (by mutating at v_2 and then v_1).

Definition 2.7. For a quiver Q , the *B-matrix* $B_Q = (b_{uv})$ (also known as the *exchange matrix*) is the skew-symmetric adjacency matrix of Q . By convention we have $b_{uv} > 0$ when there are b_{uv} arrows $u \rightarrow v$, and $b_{uv} < 0$ when there are $-b_{uv}$ arrows $u \leftarrow v$.

Definition 2.8. A (*full*) *subquiver* is an induced subgraph of a quiver. We call the remaining vertices of a subquiver its *support*. The *mutable part* of a quiver Q is the subquiver supported by its mutable vertices.

Definition 2.9. The *principal extension* (also known as *framing*) of a quiver Q is a quiver \widehat{Q} formed by adding a new frozen vertex v' and a single arrow $v' \rightarrow v$ for each vertex v . Thus Q is the mutable part of \widehat{Q} and \widehat{Q} has twice as many vertices as Q .




Figure 3: The black circle vertices v_i support an acyclic quiver Q . With the additional square (frozen) vertices v'_i , we get the principal framing \widehat{Q} .

We will use the following notation for mutation classes and associated matrices and data. (Cf. also Definitions 2.24 and 5.1.)

Definition 2.10. Let \mathbb{T}_n be an n -regular tree whose edges are labeled by the integers $1, \dots, n$, so that each edge label appears next to each vertex. We write $t \overset{i}{-} t'$ to indicate that an edge labeled i joins the vertices t and t' in \mathbb{T}_n . The tree \mathbb{T}_n always comes with a distinguished vertex t_0 .

Fix a quiver Q_0 with n mutable linearly ordered vertices $v_1 < \dots < v_n$ (and no frozen vertices). We index the mutation class $[\widehat{Q}_0]$ by simultaneously assigning a quiver $\tilde{Q}_t \in [\widehat{Q}_0]$ to each vertex t in \mathbb{T}_n , so that $\tilde{Q}_{t_0} = \widehat{Q}_0$ and $\tilde{Q}_t = \mu_{v_i}(\tilde{Q}_{t'})$ whenever $t \overset{i}{-} t'$. We call Q_0 the *initial quiver*. For each vertex $t \in \mathbb{T}_n$, let Q_t be the mutable part of \tilde{Q}_t . Let $B_t = (b_{v_i v_j; t})$ be the $n \times n$ exchange matrix of Q_t . Similarly, let $C_t = (c_{v'_i v_j; t})$ be $n \times n$ bipartite adjacency matrix between frozen vertices and mutable vertices. Thus

$$b_{v_i v_j; t} = \#\{\text{arrows } v_i \rightarrow v_j \text{ in } Q_t\} - \#\{\text{arrows } v_i \leftarrow v_j \text{ in } Q_t\},$$

$$c_{v'_i v_j; t} = \#\{\text{arrows } v'_i \rightarrow v_j \text{ in } \tilde{Q}_t\} - \#\{\text{arrows } v'_i \leftarrow v_j \text{ in } \tilde{Q}_t\}.$$

The matrix C_t is called the *C-matrix*, and its columns $\mathbf{c}_{v_i; t}$ are called *c-vectors* [18]. Note that B_t is skew-symmetric, while C_t need not have any symmetries.

Whenever we use a quiver or matrix indexed by t we will specify an initial quiver, or relevant conditions on that choice. We will often take an acyclic initial quiver Q_0 .

Example 2.11. Let Q_0 be the acyclic quiver with vertices v_1, v_2, v_3, v_4 , and arrows $v_1 \rightarrow v_2 \xrightarrow{2} v_4$, $v_1 \rightarrow v_3 \rightarrow v_4$, and $v_1 \xrightarrow{3} v_4$ (see Figure 3 for \widehat{Q}_0). With initial quiver Q_0 (and linear order $v_1 < v_2 < v_3 < v_4$), we have

$$B_{t_0} = \begin{pmatrix} 0 & 1 & 1 & 3 \\ -1 & 0 & 0 & 2 \\ -1 & 0 & 0 & 1 \\ -3 & -2 & -1 & 0 \end{pmatrix}, \quad C_{t_0} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Indeed, by construction C_{t_0} is the identity matrix regardless of the initial quiver.

If $t \xrightarrow{2} t_0$ then

$$B_t = \begin{pmatrix} 0 & -1 & 1 & 5 \\ 1 & 0 & 0 & -2 \\ -1 & 0 & 0 & 1 \\ -5 & 2 & -1 & 0 \end{pmatrix}, \quad C_t = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 2 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Remark 2.12. The map $t \mapsto (C_t, B_t)$ is called a (principal, tropical) Y -seed pattern in [21], as it is essentially equivalent to an assignment of tropical coefficient variables (usually denoted \mathbf{y}) and B -matrix to each vertex $t \in \mathbb{T}_n$, cf. [14, Proposition 3.6.5].

Definition 2.13. A (mutable) vertex v in a quiver Q_t (or \tilde{Q}_t) is green (resp., red) if every entry of $\mathbf{c}_{v;t}$ is nonnegative (resp., nonpositive).

Example 2.14. Every vertex in Q_{t_0} is green. Mutation at vertex v toggles the color of v . (Other vertices may change color as well!)

Theorem 2.15 ([6]). *Every vertex in Q_t is either green or red (never both), regardless of the choice of initial quiver.*

We next re-interpret Theorem 2.15 using the following definition.

Definition 2.16. For t in \mathbb{T}_n , let $\overset{\square}{Q}_t$ denote the quiver obtained from \tilde{Q}_t as follows:

- remove all frozen vertices and the arrows incident to them;
- add a single new frozen vertex \hat{v} ;
- for each green mutable vertex v_j , add $\sum_i c_{v'_i v_j; t}$ arrows $\hat{v} \rightarrow v_j$;
- for each red mutable vertex v_j , add $-\sum_i c_{v'_i v_j; t}$ arrows $\hat{v} \leftarrow v_j$.

Note that, by Theorem 2.15, all arrows connecting frozen vertices to a given mutable vertex are oriented in the same direction. It also follows from Theorem 2.15 that the “bundling” operation $\tilde{Q}_t \mapsto \overset{\square}{Q}_t$ commutes with mutation:

Corollary 2.17. *If $t \xrightarrow{i} t'$, or equivalently $\mu_{v_i}(\tilde{Q}_t) = \tilde{Q}_{t'}$, then $\mu_{v_i}(\overset{\square}{Q}_t) = \overset{\square}{Q}_{t'}$.*

Remark 2.18. More generally, C -matrices can be used to compute the number of arrows between frozen and mutable vertices for “triangular extensions” other than the principal framing, such as those that involve adding a frozen vertex \hat{v} and any number of arrows $\hat{v} \rightarrow v$, cf. [4, Theorem 3.2].

Definition 2.19. For a quiver Q we define the underlying unoriented *simple* graph K_Q . The graph K_Q has the same vertex set as Q , and an edge $u - v$ whenever there are any arrows $u \rightarrow v$ or $v \rightarrow u$ in Q . There are no parallel edges.

By a cycle in K_Q , we mean a sequence of vertices in K_Q where each consecutive pair is joined by an edge $\mathcal{O} = (w_0 - w_1 - \dots - w_\ell = w_0)$, considered up to cyclic shifts. In particular, cycles in K_Q have a direction of traversal. We will distinguish between two cycles with the same vertices traversed in the opposite order. Cycles thus correspond to an element of the first homology group $H_1(K_Q, \mathbb{Z})$.

Definition 2.20 (cf. [11, Proposition 2.14]). A cycle $\mathcal{O} = (w_0 - w_1 - \cdots - w_\ell = w_0)$ in K_Q is *chordless* if there are no edges between w_i and w_j for $i \neq j \pm 1$. If the arrows between w_i and w_{i+1} (in Q) are directed with the indexing of the cycle, $w_i \rightarrow w_{i+1}$, we say the cycle is *forward-oriented*. If instead the arrows are directed against the indexing, $w_i \leftarrow w_{i+1}$, then we say the cycle is *backward-oriented*. A cycle is *oriented* if it is either forward-oriented or backward-oriented.

Remark 2.21. What we call a “chordless cycle in K_Q ” is called a “cycle” in [21]. Thus, if a chordless cycle in K_Q is oriented, then it is called an “oriented cycle.” Similarly, cycles which are not oriented are called “nonoriented.”

In [21], cycles do not have an order of traversal, and so the forward/backward distinction does not arise. The order of traversal of cycles plays an important role in [11] (cf. Definition 3.8).



Figure 4: Two quivers Q and Q' . The cycle $(v_1 - v_2 - v_3 - v_4 - v_5 - v_6 - v_1)$ in K_Q is chordless and forward-oriented, while in $K_{Q'}$ it is not chordless. The chordless cycles $(v_1 - v_2 - v_3 - v_6 - v_1)$ and $(v_3 - v_4 - v_5 - v_6 - v_3)$ in $K_{Q'}$ are not oriented.

Definition 2.22 ([2]). Fix a quiver Q with B -matrix $B_Q = (b_{vw})$. A *quasi-Cartan companion* of Q is a symmetric matrix $A = (a_{vw})$ such that $a_{vv} = 2$ and $a_{vw} = \pm b_{vw}$ otherwise.

A quasi-Cartan companion of a given matrix B_Q is determined by a choice of $\binom{n}{2}$ signs. The way these signs are chosen along chordless cycles will play a special role:

Definition 2.23 ([21]). A quasi-Cartan companion $A = (a_{vw})$ of a quiver Q is *admissible* if for every chordless cycle $\mathcal{O} = (w_0 - \cdots - w_\ell = w_0)$ in K_Q the number $\#\{i \mid a_{w_i w_{i+1}} > 0 \text{ and } i < \ell\}$ is odd when \mathcal{O} is oriented, and is even otherwise.

Definition 2.24 (Cf. Definition 2.10). Fix an acyclic initial quiver Q_0 , with exchange matrix $B_{Q_0} = (b_{vw})$. We define the *initial quasi-Cartan companion* $A_{t_0} = (a_{vw})$ by $a_{vv} = 2$ and $a_{vw} = -|b_{vw}|$ for $v \neq w$. For each $t \in \mathbb{T}_n$ we define the matrix $A_t = C_t^T A_{t_0} C_t$.

The following theorems of A. Seven will play an important role in our proof of Theorem 1.1.

Theorem 2.25 ([21, Theorem 1.3]). *Suppose the initial quiver Q_0 is acyclic. Then for every $t \in \mathbb{T}_n$, the matrix $A_t = C_t^T A_{t_0} C_t$ (see Definition 2.24) is an admissible quasi-Cartan companion of Q_t .*

We will also need the following result.

Theorem 2.26 ([21, Theorems 1.3 and 1.4]). *Let Q_0 be acyclic, and fix $t \in \mathbb{T}_n$. Let $A_t = (a_{uv;t})$.*

- *For an arrow $u \rightarrow v$ in Q_t , we have $a_{uv;t} > 0$ if and only if u is red and v is green in Q_t .*
- *Every oriented path of mutable vertices $w_1 \rightarrow \cdots \rightarrow w_m$ in Q_t has at most one positive entry in A_t (thus $a_{w_i w_{i+1}; t} > 0$ for at most one $i < m$).*

An illustration of Theorem 2.26 can be found in Example 2.32.

Remark 2.27. To see how [21, Theorem 1.3] implies the first statement of Theorem 2.26, assume that $b_{ji} < 0$ and check the four combinations of $\text{sgn}(\mathbf{c}_i)$ and $\text{sgn}(\mathbf{c}_j)$.

Remark 2.28. For quivers mutation equivalent to acyclic quivers with sufficiently many arrows, a number of properties of the matrices B_t and C_t matrices have been recently established by T. Ervin, see [8, Proposition 5.2, Lemma 5.4]. These results may potentially be used to obtain enhancements of Theorem 2.26.

Definition 2.29 ([14, Theorem 2.8.3]). We define a pair of square matrices $M_Q(v, \pm 1)$, associated to mutating a quiver Q at a given vertex v . Let $B_Q = (b_{pq})$ be the exchange matrix of Q and choose $\varepsilon = \pm 1$. Let J be the diagonal matrix with $J_{vv} = -1$ and $J_{qq} = 1$ for $w \neq v$. Let $E = (e_{pq})$ be the matrix with v th column $e_{qv} = \max(0, -\varepsilon b_{qv})$ and 0 otherwise. Then $M_Q(v, \varepsilon) = J + E$.

The matrix M_Q can be used to mutate the B and C -matrices (and sometimes the quasi-Cartan companion) associated with Q . Recall that red and green vertices are determined by the signs of the C -matrix (Definition 2.13).

Proposition 2.30 ([14, Theorem 2.8.3], cf. [21, Lemma 3.1]). *Fix an initial quiver Q_0 . Choose a quiver $\tilde{Q}_t \in [\tilde{Q}_0]$, and let $Q = Q_t$ (the mutable part of \tilde{Q}_t). Select a vertex v_i of Q , and let $t \xrightarrow{i} t'$. Then:*

- *for either value of $\varepsilon = \pm 1$,*

$$(2.1) \quad B_{t'} = M_Q(v_i, \varepsilon) B_t M_Q(v_i, \varepsilon)^T;$$

- *the C -matrices satisfy*

$$C_{t'} = \begin{cases} C_t M_Q(v_i, 1)^T & \text{if } v_i \text{ is green,} \\ C_t M_Q(v_i, -1)^T & \text{if } v_i \text{ is red;} \end{cases}$$

- *if the initial quiver Q_0 is acyclic then*

$$A_{t'} = \begin{cases} M_Q(v_i, 1) A_t M_Q(v_i, 1)^T & \text{if } v_i \text{ is green,} \\ M_Q(v_i, -1) A_t M_Q(v_i, -1)^T & \text{if } v_i \text{ is red.} \end{cases}$$

The latter two formulas, involving C_t and A_t , follow from their definitions and (2.1).

Example 2.31. We demonstrate Proposition 2.30. Take our initial quiver Q_0 to be the acyclic quiver with vertices and arrows $v_1 \xrightarrow{2} v_3 \rightarrow v_2$ and $v_1 \rightarrow v_2$. Let $t \in \mathbb{T}_n$ satisfy $t_0 \xrightarrow{3} t_1 \xrightarrow{1} t_2 \xrightarrow{2} t$.

Consider the matrices B_t , and C_t , and their associated A_t shown below.

$$B_t = \begin{pmatrix} 0 & 3 & -13 \\ -3 & 0 & 5 \\ 13 & -5 & 0 \end{pmatrix}, C_t = \begin{pmatrix} 8 & -3 & 0 \\ 3 & -1 & 0 \\ 3 & -1 & -1 \end{pmatrix},$$

$$A_t = \begin{pmatrix} 2 & -3 & 13 \\ -3 & 2 & -5 \\ 13 & -5 & 2 \end{pmatrix} = C_t^T \begin{pmatrix} 2 & -1 & -2 \\ -1 & 2 & -1 \\ -2 & -1 & 2 \end{pmatrix} C_t.$$

By inspecting the signs of (the columns of) C_t , we see that v_1 is green while v_2 and v_3 are red in Q_t .

Let $t' \stackrel{1}{\leftarrow} t$ (so $Q_{t'} = \mu_{v_1}(Q_t)$). We compute the mutation at v_1 and find

$$B_{t'} = \begin{pmatrix} 0 & -3 & 13 \\ 3 & 0 & -34 \\ -13 & 34 & 0 \end{pmatrix}.$$

The B-matrix $B_{t'}$ for $\mu_{v_1}(Q_t)$ can be computed from B_t by performing either of two different congruences (corresponding to $M_{Q_t}(v_1, 1)$, and $M_{Q_t}(v_1, -1)$ respectively):

$$\begin{pmatrix} -1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} B_t \begin{pmatrix} -1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}^T = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 13 & 0 & 1 \end{pmatrix} B_t \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 13 & 0 & 1 \end{pmatrix}^T = B_{t'}.$$

However the same operations applied to A_t do not both result in quasi-Cartan companions. In this case, because v_1 is green, congruence with $M_{Q_t}(v_1, 1)$ gives the quasi-Cartan companion $A_{t'}$:

$$\begin{pmatrix} -1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & -3 & 13 \\ -3 & 2 & -5 \\ 13 & -5 & 2 \end{pmatrix} \begin{pmatrix} -1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}^T = \begin{pmatrix} 2 & -3 & -13 \\ -3 & 2 & 34 \\ -13 & 34 & 2 \end{pmatrix}.$$

If instead we multiply with the matrix $M_{Q_t}(v_1, -1)$:

$$\begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 13 & 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & -3 & 13 \\ -3 & 2 & -5 \\ 13 & -5 & 2 \end{pmatrix} \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 13 & 0 & 1 \end{pmatrix}^T = \begin{pmatrix} 2 & 3 & -39 \\ 3 & 2 & -44 \\ -39 & -44 & 678 \end{pmatrix}.$$

Note in particular that one of the diagonal entries of the last matrix is not 2.

If we instead consider $t'' \stackrel{3}{\leftarrow} t$, then the associated B and C matrices are:

$$B_{t''} = \begin{pmatrix} 0 & -62 & 13 \\ 62 & 0 & -5 \\ -13 & 5 & 0 \end{pmatrix}, C_{t''} = \begin{pmatrix} 8 & -3 & 0 \\ 3 & -1 & 0 \\ 3 & -6 & 1 \end{pmatrix}.$$

In this case, only congruence with $M_{Q_t}(v_3, -1)$ gives a quasi-Cartan companion:

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 5 \\ 0 & 0 & -1 \end{pmatrix} A_t \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 5 \\ 0 & 0 & -1 \end{pmatrix}^T = \begin{pmatrix} 2 & 62 & -13 \\ 62 & 2 & -5 \\ -13 & -5 & 2 \end{pmatrix}.$$

Example 2.32. Taking A_t, B_t and Q_t as in Example 2.31, we see the only positive off-diagonal entry of A_t is $a_{v_1 v_3; t} = 13$. As predicted by Theorem 2.26 we see $v_3 \rightarrow v_1$, v_3 is red, and v_1 is green in Q_t .

3. CYCLICALLY ORDERED QUIVERS

We recall many definitions, notations, and results from [11]. We include some new examples, but do not repeat proofs.

Definition 3.1. A *cyclic ordering* of a set V is a linear ordering considered up to cyclic shifts—that is, taking the minimal element v and forming a new linear ordering where $v > u$ for all $u \neq v$ (the order is otherwise unchanged). By repeated cyclic shifts, any element can be made maximal or minimal. The cyclic ordering associated to a linear ordering $v_1 < \dots < v_n$ will be denoted (v_1, \dots, v_n) .

Definition 3.2. A *cyclically ordered quiver* (Q, σ) (abbreviated COQ) is a quiver Q equipped with a cyclic ordering of its (mutable) vertices σ . Likewise, a *linearly ordered quiver* is a quiver equipped with a linear ordering of its (mutable) vertices. We may omit the ordering when it is clear from context or not needed, and simply denote a COQ by Q . There are n linearly ordered quivers associated to each COQ, given by “tearing” the cyclic order between different consecutive vertices in the cyclic ordering.

We say an oriented path $u \rightarrow v \rightarrow w$ makes a *right turn* at v if $u < v < w$ in some linear order associated to σ .




Figure 5: The $4!/4 = 6$ distinct COQs on the 4-vertex acyclic quiver with vertices and arrows $v_1 \rightarrow v_2 \xrightarrow{2} v_4$, $v_1 \rightarrow v_3 \rightarrow v_4$, and $v_1 \xrightarrow{3} v_4$.

Definition 3.3. The *unipotent companion* of a linearly ordered quiver Q is the unipotent upper triangular matrix U such that $U^T - U = B_Q$, where the rows and columns of B_Q are written in the linear ordering. The matrix U can be constructed by negating B_Q , then setting the lower triangular part to 0 and the diagonal entries to 1. For a COQ Q , the unipotent companion of any associated linearly ordered quiver is a unipotent companion of Q .

Example 3.4. Consider the leftmost COQ Q depicted in Figure 5. This COQ has 4 associated linearly ordered quivers. We list their respective unipotent companions and linear orderings:

$$\begin{pmatrix} 1 & -1 & -1 & -3 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & -2 & 1 \\ 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & -1 & 1 & 0 \\ 0 & 1 & 3 & 2 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 3 & 2 & 1 \\ 0 & 1 & -1 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$v_1 < v_2 < v_3 < v_4 \quad v_2 < v_3 < v_4 < v_1 \quad v_3 < v_4 < v_1 < v_2 \quad v_4 < v_1 < v_2 < v_3$$

Proposition 3.5 ([11, Proposition 4.4]). *All unipotent companions of a COQ are integrally congruent.*

Recall that the integral congruence class of a unipotent companion U is the set of matrices $\{GUG^T \mid G \in \mathrm{GL}_n(\mathbb{Z})\}$.

Definition 3.6. A *wiggle* is a transformation of a COQ which fixes the quiver Q , and transposes two adjacent vertices in the cyclic ordering that are not adjacent in Q . We say that two COQs are wiggle equivalent if they have the same quiver, and their cyclic orderings are related by a sequence of wiggles. The wiggle equivalence class of a COQ is the set of all wiggle equivalent COQs, see Figure 6. If U is a unipotent companion of some COQ in a wiggle equivalence class, we say that U is a unipotent companion of the class.




Figure 6: The 4 wiggle equivalence classes of the COQs in Figure 5.

Proposition 3.7 ([11, Proposition 4.5]). *All unipotent companions of wiggle equivalent COQs are integrally congruent.*

Definition 3.8 (Cf. [11, Definition 2.10]). Fix a COQ (Q, σ) . Let $\mathcal{O} = (w_0 - w_1 - \dots - w_k = w_0)$ be a cycle in K_Q (Definition 2.20). Fix a linear order $<$ associated to σ . The *winding number* $\mathrm{wind}_\sigma(\mathcal{O})$ is the (signed) number of times we “wrap around” the linear order:

$$(3.1) \quad \mathrm{wind}_\sigma(\mathcal{O}) = \#\{w_i \mid w_i \rightarrow w_{i+1}, w_i > w_{i+1}\} - \#\{w_i \mid w_i \leftarrow w_{i+1}, w_i < w_{i+1}\}.$$

We omit the (straightforward) proofs that $\mathrm{wind}_\sigma(\mathcal{O})$ does not depend on the choice of linear order associated to σ , and that this definition agrees with [11, Definition 2.10].

Example 3.9. Let Q be the underlying quiver of the COQs shown in Figure 6, and consider the (chordless) cycle $\mathcal{O} = (v_1 - v_2 - v_4 - v_1)$ in K_Q . The winding number of this undirected cycle is 0 in the three COQs in the top row and 1 in the three COQs in the bottom row.

Remark 3.10. The winding number corresponds to the winding number of certain maps from the cycle (as a cell complex) to the circle. Specifically, map the vertices onto the circle according to the cyclic ordering (*not* the order of traversal), and then map each edge to an arc between the endpoints, starting from the tail of an associated arrow and traveling clockwise to the head. See Figure 7.




Figure 7: A COQ Q and the image of the cycle $\mathcal{O} = (v_1 - v_2 - v_4 - v_3 - v_1)$ in K_Q , with the arrows drawn on the circle. In this case the image is contractible, and so the winding number is 0.

Theorem 3.11 ([11, Theorem 2.12]). *Two COQs (Q, σ) and (Q, σ') are wiggle equivalent if and only if $\text{wind}_\sigma(\mathcal{O}) = \text{wind}_{\sigma'}(\mathcal{O})$ for every undirected cycle \mathcal{O} in K_Q .*

We now turn to mutations of COQs.

Definition 3.12. A vertex v in a COQ (Q, σ) is *proper* if every oriented path $u \rightarrow v \rightarrow w$ makes a right turn at v (Definition 3.2). The wiggle equivalence class is *proper* if for every vertex v there is a COQ in the wiggle equivalence class in which v is proper.

Example 3.13. In the wiggle equivalence classes shown in Figure 6, vertices v_1, v_4 are proper vertices in each class, as they have no oriented paths through them. Vertex v_2 is proper only in the three COQs in the top row, while vertex v_3 is proper only in the three COQs in the leftmost column(s). Thus the wiggle equivalence class in the top left, using the cyclic orderings (v_1, v_2, v_3, v_4) and (v_1, v_3, v_2, v_4) , is the only class which is proper.

Definition 3.14. To mutate a COQ Q at a proper vertex v , mutate the quiver as usual and reposition v in the cyclic ordering so that v is still proper (that is, move v clockwise in the cyclic ordering past all the elements of $\text{In}(v)$ in the mutated quiver, without passing any vertices in $\text{Out}(v)$; all choices give wiggle equivalent COQs). We denote the resulting wiggle equivalence class by $\mu_v(Q)$. The *proper mutation class* of a COQ is the set of all COQs which can be obtained from the original by a sequence of proper mutations and wiggles.




Figure 8: A sequence of proper mutations.

Proposition 3.15 ([11, Proposition 6.4]). *Proper mutation is well defined up to wiggle equivalence—that is, if two COQs are wiggle equivalent and a vertex v is proper in each, then performing the proper mutation μ_v to each results in the same wiggle equivalence class of COQs.*

Theorem 3.16 ([11, Theorem 7.1]). *The unipotent companions of COQs which are related by a proper mutation are integrally congruent.*

Definition 3.17 ([11, Definition 11.1]). A (wobble equivalence class of) COQ(s) is *totally proper* if every COQ in its proper mutation class is proper. We say a cyclic ordering σ is *totally proper* for a quiver Q if (Q, σ) is totally proper. If Q has a totally proper cyclic ordering, we say Q is *totally proper*.

Certain unipotent companions are transformed by matrix conjugation with the *same* matrices (see Definition 2.29) that transform the B -matrix. One case was established in [11], and follows. We generalize this result in the next section.

Lemma 3.18 ([11, Lemma 7.2 and Remark 7.3]). *Suppose that v_j is a proper vertex in the COQ Q . Let U be a unipotent companion of Q with linear order $<$ such that $v_i < v_j$ for $v_i \in \text{In}(v_j)$ and $v_i < v_k$ for $v_k \in \text{Out}(v_j)$. Then the matrix $M_Q(v_j, -1)UM_Q(v_j, -1)^T$ is a unipotent companion of $\mu_{v_j}(Q)$.*

We conclude this section by recalling two easily expressed invariants of COQs, along with some examples. Definition 3.19 and Proposition 3.21 do not play any role in the proof of Theorem 1.1, but do appear prominently in applications and corollaries.

Definition 3.19 ([11][Definitions 8.1, 8.4]). Given a COQ Q with unipotent companion U , the *Alexander polynomial* of Q is given by $\Delta_Q(t) = \det(tU - U^T)$. The *Markov invariant* M_Q is defined by $M_Q = n + (\text{coefficient of } t^{n-1} \text{ in } \Delta_Q(t))$.

Remark 3.20. As the exchange matrix B and unipotent companion U of a linearly ordered quiver Q are related by $B = U^T - U$, it follows that

$$\det(B) = \det(U^T - U) = (-1)^n \Delta_Q(1).$$

Proposition 3.21. *Let Q be an n -vertex acyclic quiver. Fix a linear order $<$ of the vertices of Q so that $v_i < v_j$ whenever $v_i \rightarrow v_j$. Let U be the unipotent companion of the linearly ordered quiver $(Q, <)$. For a cycle $\mathcal{O} = (w_0 - \dots - w_\ell = w_0)$ in K_Q , let $wt(\mathcal{O}) = \prod_i |b_{w_i w_{i+1}}|$ if there is exactly one location i with $w_i \leftarrow w_{i+1}$ (thus $w_j \rightarrow w_{j+1}$ for all $j \neq i$), and $wt(\mathcal{O}) = 0$ otherwise. Then the Markov invariant*

$$M_Q = \sum_{v_i < v_j} b_{v_i v_j}^2 + \sum_{\mathcal{O}} wt(\mathcal{O}).$$

Proof. The t^{n-1} coefficient of $\det(tU - U^T)$ is negative of the trace of $U^T U^{-1}$. Let $N = I - U$, note N is positive and strictly upper triangular. Thus

$$U^{-1} = I + N + N^2 + \dots + N^{n-1},$$

and so $\text{tr}(U^T U^{-1}) = \sum_{k=0}^{n-1} \text{tr}(U^T N^k)$. We compute $\text{tr}(U^T I) = n$, $\text{tr}(U^T N) = -\sum_{v_i < v_j} b_{v_i v_j}^2$ and $\text{tr}(U^T N^k) = -\sum_{\mathcal{O}} wt(\mathcal{O})$, where the sum is over the cycles \mathcal{O} of length k . \square

Example 3.22. Let Q be the leftmost COQ shown in Figure 5. Then

$$\Delta_Q(t) = t^4 + 21t^3 - 43t^2 + 21t + 1,$$

and $M_Q = 1^2 + 1^2 + 1^2 + 2^2 + 3^2 + 1 \cdot 2 \cdot 3 + 1 \cdot 1 \cdot 3 = 4 + 21 = 25$.

Remark 3.23. Amanda Schwartz has recently developed a combinatorial description for *all* of the coefficients of the Alexander polynomial $\Delta_Q(t)$ in the case that Q is a tree quiver (that is, K_Q is a tree) [19].

Further examples appear in Section 6.

4. TECHNICAL LEMMAS

In this section we establish lemmas and propositions which we will need in the proof of Theorem 1.1. We begin with generalizations of Lemma 3.18.

Lemma 4.1. *Suppose that v_j is a proper vertex in the COQ Q . Let $U = (u_{v_i v_k})$ be a unipotent companion of Q with linear order $<$ such that $v_i < v_j$ for $v_i \in \text{In}(v_j)$. Then the matrix $M_Q(v_j, -1)UM_Q(v_j, -1)^T$ is a unipotent companion of $\mu_{v_j}(Q)$.*

Proof. We follow the proof of [11, Lemma 7.2] closely. Throughout $\text{In}(v_j)$ and $\text{Out}(v_j)$ respectively refer to the inset and outset of v_j in Q .

For the mutated quiver $Q' = \mu_{v_j}(Q)$ let $<'$ be a linear order such that

- $v_j <' v_i$ for all $v_i \in \text{In}(v_j)$,
- if both $v_k \in \text{Out}(v_j)$ and $v_k < v_j$ then $v_k <' v_j$, and
- otherwise $<'$ agrees with $<$ (so we have only changed the relative position of v_j).

Such an order exists because v_j is proper in Q . Let $U' = (u'_{v_i v_k})$ be the unipotent companion of Q' with linear ordering $<'$.

Let $B_Q = (b_{v_i v_k})$, set $n_{v_i v_j} = -\max(0, b_{v_i v_j})$ and also

$$\delta_{v_i} = \begin{cases} -1 & \text{if } v_i = v_j; \\ 1 & \text{else.} \end{cases}$$

We wish to show that:

$$(4.1) \quad u'_{v_i v_k} = \delta_{v_i} \delta_{v_k} u_{v_i v_k} - n_{v_i v_j} u_{v_j v_k} \delta_{v_k} - \delta_{v_i} u_{v_i v_j} n_{v_k v_j} + n_{v_i v_j} n_{v_k v_j}.$$

Let $u''_{v_i v_k}$ denote the right hand side of (4.1).

By construction, we have

$$u'_{v_i v_k} = \begin{cases} 1 & \text{if } v_i = v_k; \\ b_{v_j v_k} & \text{if } v_i = v_j <' v_k; \\ b_{v_i v_j} & \text{if } v_i <' v_j = v_k; \\ -b_{v_i v_k} - n_{v_i v_j} n_{v_k v_j} + \max(0, -b_{v_i v_j}) \max(0, -b_{v_j v_k}) & \text{if } v_j \neq v_i <' v_k \neq v_j; \\ 0 & \text{if } v_i >' v_k. \end{cases}$$

Every case where both $v_j <' v_i$, and $v_j \leq' v_k$ appears in [11, Lemma 7.2], and the arguments therein apply. We consider the cases where v_j appears after v_i or v_k in $<'$. Note by construction of $<'$ that if $w <' v_j$ then $u_{v_j w} = 0 = n_{w v_j}$ and $b_{w v_j} = -u_{w v_j}$.

- If $v_i = v_k <' v_j$ then $u'_{v_i v_i} = 1 - 0 - 0 + 0 = u''_{v_i v_i}$.
- If $v_i <' v_j = v_k$ then, $u'_{v_i v_j} = b_{v_i v_j} = -u_{v_i v_j} - 0 - 0 + 0 = u''_{v_i v_j}$.
- We split $v_j \neq v_i <' v_k \neq v_j$ into 2 cases, depending on the relative position of v_j .
 - If $v_i <' v_j <' v_k$ then $n_{v_k v_j} = -\max(0, b_{v_k v_j})$ and $u_{v_i v_j} = \max(0, -b_{v_i v_j})$. So

$$u'_{v_i v_k} = -b_{v_i v_k} + \max(0, -b_{v_i v_j}) \max(0, -b_{v_j v_k}) = u_{v_i v_k} - 0 - u_{v_i v_j} n_{v_k v_j} + 0 = u''_{v_i v_k}.$$

$$\text{– If } v_i <' v_k <' v_j, \text{ then } u'_{v_i v_k} = -b_{v_i v_k} = u_{v_i v_k} - 0 - 0 + 0 = u''_{v_i v_k}.$$

- If $v_k <' v_i$ and $v_k <' v_j$ then $u'_{v_i v_k} = 0 - 0 - 0 + 0 = u''_{v_i v_k}$.

Let π be the permutation matrix which transforms $<$ into $<'$. Equation (4.1) can be rewritten as $U' = \pi^T M_Q(v_j, -1) U M_Q(v_j, -1)^T \pi$ (recall the matrices E and J defined in Definition 2.29). The claim follows. \square

We establish a corresponding version of Lemma 4.1 for $\varepsilon = 1$.

Lemma 4.2. *Suppose that v_j is a proper vertex in the COQ Q . Let U be a unipotent companion of Q with linear order $<$ such that $v_j < v_k$ for $v_k \in \text{Out}(v_j)$. Then the matrix $M_Q(v_j, 1) U M_Q(v_j, 1)^T$ is a unipotent companion of $\mu_{v_j}(Q)$.*

Proof. We reduce to Lemma 4.1 by introducing the *opposite* linearly ordered quiver R^{op} of another linearly ordered R . The quiver of R^{op} has the same vertices as R , but each arrow is reversed ($v \rightarrow w$ becomes $v \leftarrow w$) and the linear ordering is reversed ($v < w$ becomes $w < v$). It is easy to see that if U_R is a unipotent companion of R , then U_R^T is a unipotent companion of R^{op} (note that U_R^T is upper triangular in the reversed linear order). We also have $M_R(v_j, 1) = M_{R^{op}}(v_j, -1)$.

Now we compute:

$$\begin{aligned} M_Q(v_j, 1) U M_Q(v_j, 1)^T &= (M_Q(v_j, 1) U^T M_Q(v_j, 1)^T)^T \\ &= (M_{Q^{op}}(v_j, -1) U^T M_{Q^{op}}(v_j, -1)^T)^T. \end{aligned}$$

Lemma 4.1 implies that $M_{Q^{op}}(v_j, -1) U^T M_{Q^{op}}(v_j, -1)^T$ is a unipotent companion of Q^{op} . Thus its transpose is a unipotent companion of Q , as desired. \square

Example 4.3. Let Q be the leftmost COQ depicted in Figure 5. The unipotent companions of Q appear in Example 3.4, let U be the rightmost unipotent companion corresponding to the linear order $v_4 < v_1 < v_2 < v_3$. Vertex v_2 is proper (and both Q and $\mu_{v_2}(Q)$ appear in the center of Figure 8). We compute the unipotent companion $M_Q(v_2, -1) U M_Q(v_2, -1)^T$ of $\mu_{v_2}(Q)$:

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 3 & 2 & 1 \\ 0 & 1 & -1 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}^T = \begin{pmatrix} 1 & 5 & -2 & 1 \\ 0 & 1 & 0 & -1 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

This matrix is upper triangular with the linear order $v_4 < v_2 < v_1 < v_3$ (thus we have swapped the 2nd and 3rd rows and columns), in agreement with Lemma 4.1.

If we instead take U' to be the leftmost unipotent companion depicted in Example 3.4 (linear order $v_1 < v_2 < v_3 < v_4$), then we may apply Lemma 4.2. We find another unipotent companion $M_Q(v, 1) U' M_Q(v, 1)^T$:

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 2 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 & -1 & -3 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 2 & 0 & 1 \end{pmatrix}^T = \begin{pmatrix} 1 & 1 & -1 & -5 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & -2 & 0 & 1 \end{pmatrix}.$$

This matrix is upper triangular with order $v_1 < v_3 < v_4 < v_2$.

We next establish that certain subquivers in a mutation-acyclic quiver are acyclic.

Theorem 4.4. *Suppose Q is mutation-acyclic, and v is a vertex in Q . Then the subquiver of Q supported by $\text{Out}(v)$ (resp. $\text{In}(v)$) is acyclic.*

This theorem generalizes an example provided in [21, Figure 1].

Proof. As Q is mutation-acyclic, by Theorem 2.25, Q has an admissible quasi-Cartan companion $A = (a_{pq})$ (Definition 2.23). Thus every subquiver of Q also has an admissible quasi-Cartan companion. (By restricting the rows and columns of A .)

Thus we may reduce to the case that v is a source (resp. sink), and v is adjacent to all other vertices in Q . If $\text{Out}(v)$ (resp. $\text{In}(v)$) contains an oriented cycle then it also contains an oriented chordless cycle (by a standard subdivision argument, any chord results in a smaller oriented cycle; cf. Lemma 4.6).

Assume for contradiction that the undirected simple graph K_Q consists only of a chordless forward-oriented cycle $\mathcal{O} = (w_0 - \dots - w_\ell = w_0)$ and the source vertex v with edges $w_i - v$ (the case with v a sink is identical). Then an odd number of the $a_{w_i w_{i+1}}$ are positive. In particular, at least one $a_{w_i w_{i+1}}$ is positive.

Without loss of generality, say $a_{w_0 w_1} > 0$. Note that A remains an admissible quasi-Cartan companion if we simultaneously toggle the signs of the row and column corresponding to w_i . By repeatedly toggling signs, we may assume that $a_{w_i w_{i+1}} < 0$ for all $i > 0$ [why?]. However, each of the chordless cycles $(w_i - w_{i+1} - v - w_i)$ in K_Q are not oriented, and so any admissible companion must satisfy an even number of the inequalities $a_{w_i w_{i+1}} > 0$, $a_{w_{i+1} v} > 0$, and $a_{w_i v} > 0$. In particular, one of $a_{w_0 v}$ or $a_{w_1 v}$ is positive, without loss of generality say $a_{w_1 v} > 0$. As $a_{w_1 w_2} < 0$, we must have $a_{w_2 v} > 0$ for the chordless cycle $(w_1 - w_2 - v - w_1)$ to have an even number of positive signs, and by induction we find that $a_{w_i v} > 0$ for all i . But then the chordless cycle $(w_0 - w_1 - v - w_0)$ has three positive entries in A , a contradiction. \square

Remark 4.5. Suppose Q is mutation-acyclic, and v is a vertex in Q . Partition the other vertices of Q into three sets $\text{Out}(v)$, $\text{In}(v)$ and the set of vertices S which are not adjacent to v . Consider the subquivers supported by $\text{Out}(v) \cup S$ (resp. $\text{In}(v) \cup S$). Essentially the same argument as in Theorem 4.4 implies that every oriented cycle in these subquivers contains at most one vertex (and thus no arrows) in $\text{Out}(v)$ (resp. $\text{In}(v)$). Thus, for example, the quiver in Figure 9 is not mutation-acyclic.




Figure 9: A quiver on 6 vertices which does not have an admissible quasi-Cartan companion, and so cannot be mutation-acyclic. The subquiver supported by v_1, \dots, v_5 is an oriented cycle, contains two vertices in $\text{Out}(v_6)$, but no vertices in $\text{In}(v_6)$.

The following elementary lemma lets us convert a cycle in the underlying unoriented graph (Definition 2.19) into a chordless cycle, while keeping a given vertex on an oriented path.

Lemma 4.6. *Fix a vertex v in a quiver Q . Let \mathcal{O} be a cycle in K_Q , which contains the path $u \rightarrow v \rightarrow w$ for some vertices u, w . Then there exists a chordless cycle that is supported by a subset of the vertices of \mathcal{O} and contains a path $u' \rightarrow v \rightarrow w'$ for some vertices u', w' , not necessarily distinct from u, w .*

Proof. We induct on the size of \mathcal{O} . If \mathcal{O} is chordless (i.e., if \mathcal{O} only contains u, v , and w), then there is nothing to prove. Otherwise, \mathcal{O} contains a chord. If \mathcal{O} contains a chord which does not involve v , then we get a shorter chordless cycle by including the chord edge, and removing vertices and edges so that u, v , and w remain. Suppose instead that \mathcal{O} contains a chord with the vertex v , without loss of generality say $v \rightarrow w'$ for a vertex w' in \mathcal{O} and distinct from u, w . We can replace the edge $v \rightarrow w$ by $v \rightarrow w'$ and remove vertices so that u, v and w' remain (and w does not). The claim follows by induction. \square

Finally, we give a new way to check that two COQs are wiggle equivalent.

Lemma 4.7. *Fix a quiver Q and two linear orderings of its vertices $<$, and $<'$. Let $U = (u_{vw})$ and $U' = (u'_{vw})$ be the unipotent companions of the linearly ordered quivers $(Q, <)$ and $(Q, <')$ respectively. If $u_{vw} = u'_{vw}$ for every pair of vertices v, w , then the COQs associated to $(Q, <)$ and $(Q, <')$ are wiggle equivalent.*

Remark 4.8. Note that the condition $u_{vw} = u'_{vw}$ for each pair of vertices is *not* the same as requiring that the matrices U and U' are equal when they are written in their own respective linear orderings. For example, the vertex v in the first subscript may refer to a different row than in the second.

Proof of Lemma 4.7. By Theorem 3.11, it suffices to check that both COQs have the same winding numbers. From (3.1), it suffices to identify the arrows which “wrap around” the respective linear orderings (and, if they differ, their orientations). For any two vertices v, w we have $v < w$ if and only if $u_{vw} \neq 0$. Thus the same arrows contribute regardless of if we use the linear ordering $<$ or $<'$, and so the winding numbers agree. \square

Example 4.9. Let Q be the acyclic quiver described in Figure 5. Consider the two unipotent companions from different linear orderings of Q below.

$$U = \begin{pmatrix} 1 & 3 & 2 & 1 \\ 0 & 1 & -1 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad U' = \begin{pmatrix} 1 & 3 & 1 & 2 \\ 0 & 1 & -1 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$v_4 < v_1 < v_2 < v_3 \quad v_4 < v_1 < v_3 < v_2$$

Each entry of U matches U' ; for example, the 2 in the first row and third column of U corresponds to u_{v_4, v_2} and agrees with the entry in the first row and fourth column of U' . Thus U and U' are wiggle equivalent, indeed they are related by the wiggle $(v_2 v_3)$.

5. PROOF OF THEOREM 1.1

We begin by establishing and recalling notation for this section. Using the notation from Definition 2.10, suppose our initial quiver Q_0 is acyclic with (mutable) vertices v_1, \dots, v_n . Fix a quiver $\tilde{Q}_t \in [\widehat{Q_0}]$, with mutable part Q_t . Thus C_t denotes the C -matrix of Q_t , and B_t denotes the B -matrix of Q_t . Finally, recall from Definition 2.24 the associated quasi-Cartan companions A_{t_0} and $A_t = (a_{ij;t}) = C_t^T A_{t_0} C_t$.

Definition 5.1. We associate to each $t \in \mathbb{T}_n$ an additional matrix

$$U_t = (u_{vw;t}) = (A_t - B_t)/2.$$

Observe that $u_{vv;t} = 1$ and $U_t^T - U_t = B_t$.

Example 5.2. Let Q_0 be the acyclic quiver of type E_6 shown (as a COQ) in Figure 10, and let $t \in \mathbb{T}_n$ be the vertex such that

$$t_0 \xrightarrow{3} t_1 \xrightarrow{2} t_2 \xrightarrow{5} t_3 \xrightarrow{4} t_4 \xrightarrow{2} t_5 \xrightarrow{1} t_6 = t.$$

In this case,

$$B_t = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & -1 \\ 0 & 0 & -1 & -1 & 1 & 1 \\ 0 & 1 & 0 & 0 & -1 & -1 \\ -1 & 1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 \\ 1 & -1 & 1 & 0 & 0 & 0 \end{pmatrix}, \quad C_t = \begin{pmatrix} -1 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 & 0 & 1 \\ 0 & -1 & 0 & 0 & 0 & 1 \\ 0 & -1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

and therefore

$$A_t = \begin{pmatrix} 2 & 0 & 0 & -1 & 0 & -1 \\ 0 & 2 & -1 & -1 & 1 & 1 \\ 0 & -1 & 2 & 0 & 1 & -1 \\ -1 & -1 & 0 & 2 & 0 & 0 \\ 0 & -1 & 1 & 0 & 2 & 0 \\ -1 & 1 & -1 & 0 & 0 & 2 \end{pmatrix}, \quad U_t = \begin{pmatrix} 1 & 0 & 0 & -1 & 0 & 0 \\ 0 & 1 & 0 & 0 & -1 & 0 \\ 0 & -1 & 1 & 0 & 1 & 0 \\ 0 & -1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ -1 & 1 & -1 & 0 & 0 & 1 \end{pmatrix}.$$

All of the above is written using the linear order $v_1 < v_2 < \dots < v_6$.




Figure 10: The quiver Q_0 of type E_6 , and a mutation equivalent quiver Q_t .

We will construct a family of linear orderings such that U_t is upper triangular when written using one of them (see Lemma 5.8). To facilitate the construction, we first create several new directed graphs from Q_t .

Definition 5.3. Fix a vertex v_j of Q_t . Let $L_{v_j}(Q_t)$ be the directed graph obtained by performing the following operations on Q_t :

- (1) if v_j is red (resp., green), then for each oriented 2 path $v_i \rightarrow v_j \rightarrow v_k$ with v_i, v_k both green (resp., red) in Q_t , add an arrow $v_k \xrightarrow{v_j} v_i$ labeled with the vertex v_j ;
- (2) reverse all arrows from a red vertex directed toward a green vertex. Equivalently, by Theorem 2.26, reverse all arrows $v \rightarrow w$ with $a_{vw;t} > 0$.

Example 5.4. Continuing with Example 5.2, we see from the columns of C_t that v_3, v_6 are green while v_1, v_2, v_4 , and v_5 are red in Q_t . Therefore each $L_{v_i}(Q_t)$ will reverse the arrows $v_2 \rightarrow v_6$ and $v_5 \rightarrow v_3$ (agreeing with the two positive off-diagonal entries of A_t), and we only add a new arrow to $L_{v_i}(Q_t)$ if $i \in \{2, 3, 6\}$.




Figure 11: The six directed graphs $L_{v_i}(Q_t)$ associated to Q_t in Example 5.2. Vertices v_3, v_6 are green, while vertices v_1, v_2, v_4, v_5 are red.

Lemma 5.5. *Each directed graph $L_v(Q_t)$ is acyclic. (In particular, there are no oriented 2-cycles; $L_v(Q_t)$ is an acyclic quiver with some marked arrows.) Furthermore, the subquiver of Q_t supported by all red (resp., green) vertices is acyclic.*

Proof. By construction, there are no directed paths from a red vertex to a green vertex in $L_v(Q_t)$. So any oriented cycle in $L_v(Q_t)$ must have all vertices of the same color (red or green).

We first consider oriented cycles which do not use any arrows labeled v , i.e. oriented cycles in Q_t which all have the same color. Let $\overset{\square}{Q}_t$ be constructed from \tilde{Q}_t as in Definition 2.16. By construction of $\overset{\square}{Q}_t$, the set $\text{In}(\overset{\square}{v})$ (resp., $\text{Out}(\overset{\square}{v})$) is precisely the red (resp., green) vertices in Q_t . As $\overset{\square}{Q}_0$ is acyclic, by Corollary 2.17, $\overset{\square}{Q}_t$ is mutation-acyclic. By Theorem 4.4, $\text{In}(\overset{\square}{v})$ (resp., $\text{Out}(\overset{\square}{v})$) is an acyclic subquiver of $\overset{\square}{Q}_t$. Thus there are no oriented cycles of all red (resp., green) vertices in Q_t .

Without loss of generality, suppose v is red. The case where v is green is similar. Then we do not add any arrows labeled v between red vertices, so we have shown there are no oriented cycles of red vertices. Further, any directed cycle involving green vertices must use at least one arrow labeled v (those created in Step 1 above). As all arrows labeled v in $L_v(Q_t)$ are oriented from $\text{Out}(v)$ towards $\text{In}(v)$, any oriented cycle must also involve at least one arrow which is not labeled v . We will now argue that any other cycles in $L_v(Q_t)$ imply a problematic cycle \mathcal{O} in the underlying unoriented simple graph K_{Q_t} of Q_t .

Suppose there is an oriented cycle $w_0 \xrightarrow{v} w_1 \rightarrow \cdots \rightarrow w_k = w_0$ in $L_v(Q_t)$ that involves just one arrow $w_0 \xrightarrow{v} w_1$ labeled v . By construction of $L_v(Q_t)$, there is a path $w_0 \leftarrow v \leftarrow w_1$ in Q_t . Thus there is a cycle $\mathcal{O} = (w_0 - v - w_1 - w_2 - \cdots - w_k)$ in K_{Q_t} . If \mathcal{O} is not chordless then replace it with a chordless cycle containing v by Lemma 4.6. Because the subquiver of green vertices in Q_t is acyclic, any chord

between the green vertices w_i, w_j has the same orientation as the directed path it replaced. Therefore \mathcal{O} is a chordless cycle which is not oriented. But the only edge on \mathcal{O} which has a positive entry in A_t is $w_0 - v$, contradicting Theorem 2.26. So there are no cycles in $L_v(Q_t)$ with a single arrow labeled v .

Suppose there is an oriented cycle in $L_v(Q_t)$ with at least 2 arrows labeled v , say

$$w_0 \xrightarrow{v} w_1 \rightarrow \cdots \rightarrow w_j \xrightarrow{v} w_{j+1} \rightarrow \cdots \rightarrow w_k = w_0$$

with $j > 1$ and none of the arrows on the path $w_1 \rightarrow w_2 \rightarrow \cdots \rightarrow w_j$ labeled v (the other arrows may or may not be labeled v). Thus $\mathcal{O} = (v - w_1 - \cdots - w_j - v)$ is a cycle in K_{Q_t} which is not oriented. After applying Lemma 4.6 if necessary, we may assume \mathcal{O} is a chordless cycle. But the only edge on \mathcal{O} which has a positive entry in A_t is $w_j - v$, contradicting Theorem 2.26. Thus $L_v(Q_t)$ is acyclic. \square

Definition 5.6. A *linear extension* of $L_v(Q_t)$ is a linear order $<$ on the vertices such that $v_i < v_j$ whenever $v_i \rightarrow v_j$ (it does not matter whether the arrow is labeled v). Let σ_v be a cyclic ordering associated to a linear extension of $L_v(Q_t)$.

Example 5.7. Continuing with Examples 5.2 and 5.4, there are several possible choices for the cyclic orderings σ_v for each fixed v , though all of them result in wiggle equivalent COQs (Q_t, σ_v) . We could take

$$\begin{aligned} \sigma_{v_1} = \sigma_{v_2} = \sigma_{v_3} &= (v_6, v_1, v_4, v_3, v_2, v_5), \text{ and} \\ \sigma_{v_4} = \sigma_{v_5} = \sigma_{v_6} &= (v_6, v_3, v_1, v_4, v_2, v_5). \end{aligned}$$

Lemma 5.8. *The matrix $U_t = (u_{v_i v_j; t})$ is upper triangular when written with a linear extension $<$ of $L_v(Q_t)$. Thus U_t is a unipotent companion of (Q_t, σ_v) .*

Proof. Suppose v_i, v_j are vertices such that $u_{v_i v_j; t} \neq 0$. We want to show that $v_i < v_j$.

Because $|a_{v_i v_j; t}| = |b_{v_i v_j; t}|$, and $u_{v_i v_j; t} = \frac{a_{v_i v_j; t} - b_{v_i v_j; t}}{2} \neq 0$, we have $a_{v_i v_j; t} = -b_{v_i v_j; t}$.

If $a_{v_i v_j; t} > 0$ then $v_i \leftarrow v_j$ (as $b_{v_i v_j; t} < 0$). By Definition 5.3, the graph $L_v(Q_t)$ has an arrow $v_i \rightarrow v_j$. Thus $v_i < v_j$ in this case.

Otherwise $a_{v_i v_j; t} < 0$ and $v_i \rightarrow v_j$ in $L_v(Q_t)$, so we have $v_i < v_j$. \square

Example 5.9. Continuing with Example 5.2, we find that U_t is a unipotent companion of (Q_t, σ_v) , for any vertex v . As suggested by Example 5.7, if we use the linear order $v_6 < v_3 < v_1 < v_4 < v_2 < v_5$, then

$$U_t = \begin{pmatrix} 1 & -1 & -1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & -1 & 1 \\ 0 & 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

Lemma 5.10. *The COQs $(Q_t, \sigma_v), (Q_t, \sigma_w)$ are wiggle equivalent for all vertices v, w .*

Proof. By Lemma 5.8, the matrix U_t is a unipotent companion of both (Q_t, σ_v) and (Q_t, σ_w) . The claim follows from Lemma 4.7. \square

Lemma 5.11. *The COQ (Q_t, σ_v) is proper.*

Proof. It suffices to check that each vertex q is proper in the COQ (Q_t, σ_q) by Lemma 5.10. Let $<$ be a linear extension of $L_q(Q_t)$. We consider all oriented paths $p \rightarrow q \rightarrow r$ in cases, based on the color (green or red) of p, q , and r in Q_t . As noted in Lemma 5.5, the subquiver of all red or all green vertices is acyclic. We have chosen our order $<$ to be compatible with these acyclic subquivers. Thus if all three of p, q, r are green (resp., red) then $p \rightarrow q \rightarrow r$ is a right turn at q . A similar argument applies if p or r is a different color than the other two vertices.

This leaves us with the alternating patterns: red-green-red or green-red-green. By construction of $<$, if p, r are both a different color than q , then there is an arrow $r \xrightarrow{q} p$ in $L_q(Q_t)$, and so $r < p$. Thus the path $p \rightarrow q \rightarrow r$ is again a right turn at q . \square

Example 5.12. Continuing with Examples 5.2 and 5.7, Figure 12 shows the COQ (Q_t, σ_{v_1}) . Note that every oriented path of length 2 is a right turn.




Figure 12: The proper COQ (Q_t, σ_{v_1}) .

Proof of Theorem 1.1. Let $\sigma_t = \sigma_{v_1}$ be a cyclic ordering constructed from Q_t as in Definition 5.6. By Lemma 5.8, U_t is a unipotent companion of the COQ (Q_t, σ_t) . By Lemma 5.11, (Q_t, σ_t) is proper. It remains to show that these COQs (as $t \in \mathbb{T}_n$ varies) are all proper mutation equivalent.

Fix an arbitrary $t' \stackrel{i}{\sim} t$. It suffices to show that the COQ $(\mu_{v_i}(Q_t), \sigma_{t'})$ is in the wobble equivalence class $\mu_{v_i}((Q_t, \sigma_t))$. Let $<$ be a linear extension of $L_{v_i}(Q_t)$. Say that v_i is green (resp., red) in Q_t . For every vertex $w \in \text{Out}(v_i)$ (resp., $\text{In}(v_i)$), we have $v_i < w$ (resp. $w < v_i$). By Proposition 2.30, we have

$$B_{t'} = M_{Q_t}(v_i, \varepsilon) B_t M_{Q_t}(v_i, \varepsilon)^T,$$

$$A_{t'} = M_{Q_t}(v_i, \varepsilon) A_t M_{Q_t}(v_i, \varepsilon)^T$$

for $\varepsilon = 1$ (resp., -1). Thus

$$\begin{aligned} U_{t'} &= \frac{1}{2} (M_{Q_t}(v_i, \varepsilon) A_t M_{Q_t}(v_i, \varepsilon)^T - M_{Q_t}(v_i, \varepsilon) B_t M_{Q_t}(v_i, \varepsilon)^T) \\ &= M_{Q_t}(v_i, \varepsilon) U_t M_{Q_t}(v_i, \varepsilon)^T. \end{aligned}$$

In particular, $U_{t'}$ is exactly the unipotent companion of $\mu_{v_i}((Q_t, \sigma_t))$ produced by Lemma 4.2 (resp. 4.1). The theorem then follows from Lemma 4.7. \square

6. APPLICATIONS

As noted in Theorem 3.16, the integral congruence class of a unipotent companion is a proper mutation invariant of COQs. If a COQ is totally proper, then every mutation is proper and so we have a mutation invariant of the underlying quiver.

Theorem 6.1 ([11, Theorem 11.3, Remark 11.11]). *A quiver Q has at most one totally proper cyclic ordering (up to wiggle equivalence). There is an efficient algorithm that computes a cyclic ordering σ_Q such that if Q has a totally proper cyclic ordering, then (Q, σ_Q) is totally proper.*

If a quiver Q is mutation equivalent to a totally proper quiver Q' , then the integral congruence class of a unipotent companion of the COQ (Q, σ_Q) will agree with that of $(Q', \sigma_{Q'})$. In particular, their Alexander polynomials and Markov invariants (Definition 3.19) will agree.

By Theorem 1.1, we now have many new totally proper quivers. Let us discuss a few examples.

Corollary 6.2. *Suppose Q is an acyclic quiver whose underlying unoriented simple graph is a chordless cycle \mathcal{O} . Then a cyclic ordering σ of the vertices of Q is totally proper if and only if $\text{wind}_\sigma(\mathcal{O}) = 0$.*

Note that Corollary 6.2 makes no assumption about the multiplicities of the arrows.

Example 6.3. As an illustration, we compute some mutation invariants for a few families. Let $\tilde{A}(r, \ell)$ denote a quiver with $n = r + \ell$ vertices v_1, \dots, v_n which has r locations with a single arrow $v_i \rightarrow v_{i+1}$ and ℓ locations with a single arrow $v_i \leftarrow v_{i+1}$ (see [13, Example 6.9]). While there are many ways to place the arrows, all are related by sink and/or source mutations. Thus we may assume that $\tilde{A}(r, \ell)$ has arrows

$$v_1 \rightarrow v_2 \rightarrow \cdots \rightarrow v_r \leftarrow v_{r+1} \leftarrow \cdots \leftarrow v_n \leftarrow v_1.$$

For $r, \ell > 0$, the quiver $\tilde{A}(r, \ell)$ is acyclic, and thus totally proper. One totally proper cyclic ordering is $\sigma_{r\ell} = (v_1, \dots, v_{r-1}, v_n, v_{n-1}, \dots, v_{r+1}, v_r)$.

The Alexander polynomial of $(\tilde{A}(r, \ell), \sigma_{r\ell})$ is

$$\Delta_{\tilde{A}(r, \ell)}(t) = (t^\ell - (-1)^\ell)(t^r - (-1)^r).$$

Every quiver mutation equivalent to $(\tilde{A}(r, \ell), \sigma_{r\ell})$ has integrally congruent unipotent companions, and in particular a matching Alexander polynomial.

Thus, if $r = 3$ and $\ell = 2$ then one unipotent companion is

$$U = \begin{pmatrix} 1 & -1 & 0 & -1 & 0 \\ 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

and we can compute the Alexander polynomial

$$\Delta_{\tilde{A}(3,2)}(t) = \det(tU - U^T) = t^5 - t^3 + t^2 - 1.$$

Example 6.4. We can expand Example 6.3 by considering quivers whose simple undirected graphs are cycles but have multiple arrows between vertices. Choose nonnegative integers r, ℓ and positive integers $a_1, \dots, a_{r+\ell}$. Let $C(r, \ell)$ denote the quiver with $n = r + \ell$ vertices v_i , and arrows

$$v_1 \xrightarrow{a_1} v_2 \xrightarrow{a_2} \dots \xrightarrow{a_{r-1}} v_r \xleftarrow{a_r} v_{r+1} \xleftarrow{a_{r+1}} \dots \xleftarrow{a_{\ell-1}} v_n \xleftarrow{a_\ell} v_1.$$

For $r, \ell > 0$, the cyclic ordering $\sigma_{r\ell}$ is totally proper.

We compute the Alexander polynomials for several small values of r, ℓ . Recall that for a 4-vertex quiver Q we can write the Alexander polynomial as

$$\Delta_Q(t) = (t-1)^4 + M_Q \cdot t(t-1)^2 + \det(B_Q) \cdot t^2.$$

In particular,

$$\begin{aligned} \Delta_{C(4,0)}(t) &= (t-1)^4 + (a_1^2 + a_2^2 + a_3^2 + a_4^2 - a_1 a_2 a_3 a_4) t(t-1)^2 \\ &\quad + (a_1^2 a_3^2 + a_2^2 a_4^2 - 2a_1 a_2 a_3 a_4) t^2; \\ \Delta_{C(3,1)}(t) &= (t-1)^4 + (a_1^2 + a_2^2 + a_3^2 + a_4^2 + a_1 a_2 a_3 a_4) t(t-1)^2 \\ &\quad + (a_1^2 a_3^2 + a_2^2 a_4^2 + 2a_1 a_2 a_3 a_4) t^2; \\ \Delta_{C(2,2)}(t) &= (t-1)^4 + (a_1^2 + a_2^2 + a_3^2 + a_4^2) t(t-1)^2 \\ &\quad + (a_1^2 a_3^2 + a_2^2 a_4^2 - 2a_1 a_2 a_3 a_4) t^2. \end{aligned}$$

For a 5-vertex quiver Q , we can write the Alexander polynomial as

$$\Delta_Q(t) = (t-1)^5 + M_Q \cdot t(t-1)^3 + d \cdot t^2(t-1)$$

for some integer coefficient d . In particular,

$$\begin{aligned} \Delta_{C(4,1)}(t) &= (t-1)^5 + (a_1^2 + a_2^2 + a_3^2 + a_4^2 + a_5^2 + a_1 a_2 a_3 a_4 a_5) t(t-1)^3 \\ &\quad + (a_1^2 a_3^2 + a_1^2 a_4^2 + a_2^2 a_4^2 + a_2^2 a_5^2 + a_3^2 a_5^2 - 3a_1 a_2 a_3 a_4 a_5) t^2(t-1); \\ \Delta_{C(3,2)}(t) &= (t-1)^5 + (a_1^2 + a_2^2 + a_3^2 + a_4^2 + a_5^2) t(t-1)^3 \\ &\quad + (a_1^2 a_3^2 + a_1^2 a_4^2 + a_2^2 a_4^2 + a_2^2 a_5^2 + a_3^2 a_5^2 - a_1 a_2 a_3 a_4 a_5) t^2(t-1). \end{aligned}$$

These examples suggest a pattern for the Markov invariant, agreeing with Proposition 3.21. Without loss of generality, assume $r > 0$. Then

$$M_{C(r,\ell)} = \begin{cases} \sum_i^n a_i^2 & \text{if } \ell \geq 2; \\ \sum_i^n a_i^2 + \prod_i^n a_i & \text{if } \ell = 1; \\ \sum_i^n a_i^2 - \prod_i^n a_i & \text{if } \ell = 0. \end{cases}$$

Remark 6.5. Recall Lagrange's four squares theorem, which states that every non-negative integer is the sum of at most four squares (of integers). Thus there is a totally proper 4-vertex COQ with Markov invariant x for every $x \geq 0$. However some integers are not the sum of exactly four *positive* squares. For example, 2^{2k+1} , or 29 (see [23]). Thus any quiver whose associated Markov invariant is one of these values must not be mutation equivalent to $C(2, 2)$ (for any positive values a_i).

We have the following corollaries of Proposition 3.21.

Corollary 6.6. *Let Q be a connected, totally proper and mutation-acyclic COQ on n vertices. Then the Markov invariant $M_Q \geq n - 1$. Further, if $M_Q = n - 1$ then Q is mutation equivalent to a quiver whose underlying undirected graph is a tree (with no parallel arrows).*

Corollary 6.7. *Let Q be a totally proper and mutation-acyclic COQ on n vertices. Then the Markov invariant and the determinant of the exchange matrix satisfy the following inequality:*

$$\det(B_Q) \leq (2M_Q)^{n/2}.$$

Proof. It suffices to consider acyclic COQs Q with vertices v_1, \dots, v_n . By Proposition 3.21, $\sum_{i < j} b_{v_i v_j}^2 \leq M_Q$. Recall the *Frobenius norm* $\|B_Q\|_F = (\sum_{i,j} b_{v_i v_j}^2)^{1/2}$ of B_Q (see also the Hilbert-Schmidt norm or Schur norm), thus $\|B_Q\|_F^2 \leq 2M_Q$.

As $\det(B_Q) = 0$ if n is odd, we assume n is even. As B_Q is skew-symmetric, its eigenvalues are all imaginary and come in signed pairs, say they have norms $|\lambda_1|, \dots, |\lambda_{n/2}|$. Let $\lambda = \max(|\lambda_j|)$. Then $\det(B_Q) \leq \lambda^n$. As B_Q is skew-symmetric the spectral norm $\rho(B_Q) = \lambda$. It is a classical fact that the spectral norm is a lower bound on the Frobenius norm: $\lambda \leq \|B_Q\|_F$. Therefore we have

$$\det(B_Q) \leq \lambda^n \leq \|B_Q\|_F^n \leq (2M_Q)^{n/2}. \quad \square$$

Corollaries 6.6 and 6.7 give short proofs that certain quivers are not mutation-acyclic.

Example 6.8. The *Somos sequences* are integer sequences, the first 5 of which are associated to cluster algebras with particularly symmetric quivers. The Somos-4 quiver S is shown in Figure 13. The distinguished cyclic ordering $\sigma_S = (v_1, v_4, v_2, v_3)$ (see [11][Remark 11.11]) is the only potentially totally proper ordering for S . This quiver is not mutation-acyclic though, which is quickly determined by computing the Markov invariant of the COQ (S, σ_S) and applying Corollary 6.6:

$$M_S = 1^2 + 1^2 + 2^2 + 3^2 + 2^2 + 1^2 - 6 - 6 + 2 + 2 - 12 = 0.$$





Figure 13: The Somos-4 quiver S .

Remark 6.9. Fix an integer $a \geq 1$. Example 6.8 generalizes to the 4-vertex quiver aS , which has the same vertices as S , and a arrows for every arrow in S (thus, for example, there are $3a$ arrows from v_2 to v_3). Each has a nonpositive Markov invariant.

Example 6.10. Continuing with Example 6.4, consider the COQ $C(4, 0)$ with cyclic ordering (v_1, v_2, v_3, v_4) , shown below.



Some such quivers are mutation-acyclic, for example when $a_i = 1$ for all i then $C(4,0)$ has type D_4 . Others are not. Warkentin showed in [26][Example 6.11] that if $a_1 = a_3 \geq 2$ and $a_2 = a_4 \geq 2$ then this quiver is not mutation-acyclic (by a precise description of the mutation class). The same can be observed by Corollary 6.6, noting that $M_{C(4,0)} \leq 0$ for such quivers.

Suppose that $a_4 = a_1 a_2 a_3$. Then $M_{C(4,0)} = a_1^2 + a_2^2 + a_3^2 \geq 3$. Thus we cannot determine whether the quiver is mutation-acyclic or not from Corollary 6.6. However,

$$\det(B_{C(4,0)}) = (a_1 a_3 + a_1 a_2^2 a_3)^2 = (a_2^2 + 1)^2 (a_1 a_3)^2.$$

So we have $\det(B_{C(4,0)}) > (2M_{C(4,0)})^2$ whenever (for example) $a_2 \geq \max(a_1, a_3)$ and $a_1 a_3 \geq 6$. Thus, for these values of a_i , the quiver $C(4,0)$ is again not mutation-acyclic by Corollary 6.7.

Example 6.11. Consider the quivers associated with the elliptic root system of double extended type E_7 or E_8 , denoted $E_7^{(1,1)}$ or $E_8^{(1,1)}$ and shown in Figure 14. (Note that the quiver in Figure 1 is mutation equivalent to $E_8^{(1,1)}$.) These quivers have finite mutation classes, and can be shown to be totally proper by exhaustive computation. (We emphasize the exhaustion; $E_8^{(1,1)}$ has 5739 quivers, up to isomorphism, in its class.) These quivers are not mutation-acyclic. This has been shown by brute force, or by representation theory [17][Section 2.3]. We sketch a new proof.

The canonical candidate cyclic orderings $\sigma_{E_7^{(1,1)}}$ and $\sigma_{E_8^{(1,1)}}$ for the quivers shown in Figure 1 are (v_1, \dots, v_9) and (v_1, \dots, v_{10}) respectively. (While $(E_7^{(1,1)}, \sigma_{E_7^{(1,1)}})$ and $(E_8^{(1,1)}, \sigma_{E_8^{(1,1)}})$ are totally proper, we do not rely on this fact.) So we treat $E_7^{(1,1)}$ and $E_8^{(1,1)}$ as COQs with their respective cyclic ordering. Their Markov invariants are:

$$M_{E_7^{(1,1)}} = 10 + 4 - 6 = 8, \quad M_{E_8^{(1,1)}} = 11 + 4 - 6 = 9.$$

By Corollary 6.6, the only acyclic quivers that $E_7^{(1,1)}$ (resp., $E_8^{(1,1)}$) could be mutation equivalent to are tree quivers (with no parallel arrows).

Up to isomorphism, there are 47 (unoriented) trees with 9 vertices and 106 trees with 10 vertices. All orientations of each of these trees are mutation equivalent. By Theorem 3.11, all cyclic orderings of a tree are wiggle equivalent. Thus any quiver mutation equivalent to a tree on 9 (resp., 10) vertices must have one of the 47 (resp., 106) Alexander polynomials. (Actually the list is slightly smaller, as some non-isomorphic trees have the same Alexander polynomial.)

So it suffices to check if $\Delta_{E_7^{(1,1)}}(t)$ and $\Delta_{E_8^{(1,1)}}(t)$ appear in these lists. A (still tedious, but straightforward and human verifiable) computation shows that they are not. The computation can be further simplified using the work of Amanda Schwartz, see Remark 3.23.





Figure 14: The extended affine quivers $E_7^{(1,1)}$ and $E_8^{(1,1)}$.

Example 6.12. Choose positive integers a, b, c . Let Q be the 4-vertex COQ below:



Then $M_Q = a^2 + 2b^2 + 2c^2 - 2abc$. Let Q' be the subquiver supported by v_1, v_2, v_3 , thus $M_{Q'} = a^2 + b^2 + c^2 - abc$. Note that $M_Q = 2M_{Q'} - a^2$. Thus $M_Q < 0$ whenever $M_{Q'} < \frac{a^2}{2}$. (This last inequality is generally satisfied for mutation-infinite quivers; as $M_{Q'}$ is constant on the mutation class, but the number of arrows grows arbitrarily large, we can find many quivers with some weight a much larger than $M_{Q'}$.) Thus by Corollary 6.6, such Q cannot be mutation-acyclic.

Example 6.13. We next illustrate a way of combining Corollary 6.6 with another mutation invariant [20], the multiset of GCDs of all the integers in the columns (or rows) of the B -matrix. We construct a COQ as in Example 6.12 with $a = b = 2$ and $c = 4$. Thus $M_Q = 12 \geq 3$, and we cannot use Corollary 6.6 to conclude that Q is not mutation-acyclic. However, note that the GCD of the entries of B_Q is 2. Thus if Q is mutation-acyclic, it must be mutation equivalent to a connected acyclic quiver with the same GCD and $M_Q = 12$. There are two such quivers, both trees with 2 arrows between each pair of neighboring vertices. The B -matrices of these acyclic quivers have determinant 0 and 16, but $\det(B_Q) = 144$. Thus Q is not mutation-acyclic.

Finally, we note that Lemmas 4.1 4.2 extend to the admissible quasi-Cartan companion $A_U = U + U^T$ (discussed in [11][Remark 13.2]).

Corollary 6.14. Fix a totally proper COQ Q and a proper vertex v_j . Let U_Q be a unipotent companion with linear order $<$ so that $v_i < v_j$ (resp., $v_j < v_i$) for all $v_i \in \text{In}(v_j)$ (resp., $\text{Out}(v_j)$). Then $M_Q(v_j, \epsilon)(U_Q + U_Q^T)M_Q(v_j, \epsilon)$ is an admissible quasi-Cartan companion of $\mu_{v_j}(Q)$ when $\epsilon = -1$ (resp., 1).

REFERENCES

- [1] A. Buan, B. Marsh, and I. Reiten, Cluster mutation via quiver representations. *Comment. Math. Helv.* **83** (2008), 143–177.
- [2] M. Barot, C. Geiss, and A. Zelevinsky, Cluster algebras of finite type and positive symmetrizable matrices, *J. London Math. Soc.* **73** (2006), 545–564.
- [3] W. Bley, T. Hofmann, and H. Johnston, Computation of lattice isomorphisms and the integral matrix similarity problem, *Forum Math. Sigma* **10** (2022), Paper No. e87, 36 pp.
- [4] P. Cao and F. Li, Uniform column sign-coherence and the existence of maximal green sequences, *J. Algebr. Comb.* **50** (2019), 403–417.
- [5] R. Casals, A binary invariant of matrix mutation, [arXiv:2311.03601](https://arxiv.org/abs/2311.03601), to appear in *J. Comb. Algebra*.
- [6] H. Derksen, J. Weyman, and A. Zelevinsky, Quivers with potentials and their representations II: applications to cluster algebras. *J. Amer. Math. Soc.* **23** (2010), 749–790.
- [7] B. Eick, T. Hofmann, and E. A. O’Brien, The conjugacy problem in $GL(n, \mathbb{Z})$, *J. Lond. Math. Soc.* **100** (2019), 731–756.
- [8] T. Ervin, All connected quivers have an unrestricted red size of $n - 1$ or n , [arXiv:2401.14958](https://arxiv.org/abs/2401.14958).
- [9] T. Ervin, New hereditary and mutation-invariant properties arising from forks, *Electron. J. Combin.* **31** (2024), no. 1.16, 1–30.
- [10] T. Ervin, private communication, June 2024.
- [11] S. Fomin and S. Neville, Cyclically ordered quivers, [arXiv:2304.11505](https://arxiv.org/abs/2304.11505).
- [12] S. Fomin and S. Neville, Long mutation cycles, [arXiv:2304.11505](https://arxiv.org/abs/2304.11505).
- [13] S. Fomin, M. Shapiro, and D. Thurston, Cluster algebras and triangulated surfaces. Part I: Cluster complexes, *Acta Math.* **201** (2008), 83–146.
- [14] S. Fomin, L. Williams, and A. Zelevinsky, Introduction to cluster algebras. Chapters 1–3, [arXiv:1608.05735](https://arxiv.org/abs/1608.05735).
- [15] S. Fomin and A. Zelevinsky, Cluster algebras I: Foundations, *J. Amer. Math. Soc.* **15** (2002), 497–529.
- [16] B. Keller, Quiver mutation in Java. Java applet available at <https://webusers.imj-prg.fr/~bernhard.keller/quivermutation/> (2006).
- [17] B. Keller and I. Reiten, Acyclic Calabi-Yau categories, *Compos. Math.* **144.5** (2008), 1332–1348.
- [18] T. Nakanishi and A. Zelevinsky, On tropical dualities in cluster algebras, *Algebraic groups and quantum groups*, 217–226, *Contemp. Math.* **565**, American Mathematical Society, Providence, RI, 2012.
- [19] A. Schwartz, private communication, June 2024.
- [20] A. I. Seven, Mutation classes of skew-symmetrizable 3×3 matrices, *Proc. Amer. Math. Soc.* **141** (2013), 1493–1504.
- [21] A. I. Seven, Cluster algebras and symmetric matrices, *Proc. Amer. Math. Soc.* **143**, 469–478 (2015).
- [22] A. Seven and İ. Ünal, Congruence invariants of matrix mutation, [arXiv:2403.10626](https://arxiv.org/abs/2403.10626).
- [23] N. J. A. Sloane and J. H. Conway, Numbers that are not the sum of 4 nonzero squares, Entry A000534 in *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org/A000534>.
- [24] H. Thomas, private communication, August 2024.
- [25] D. Vatne, The mutation class of D_n quivers, *Comm. Algebra.* **38** (2010), 1137–1146.
- [26] M. Warkentin, *Exchange graphs via quiver mutation*, Ph.D. thesis, Technische Universität Chemnitz, 2014, <https://nbn-resolving.org/urn:nbn:de:bsz:ch1-qucosa-153172>.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109, USA
Email address: nevilles@umich.edu