

BRILL-NOETHER LOCI ON AN ENRIQUES SURFACE COVERED BY A JACOBIAN KUMMER SURFACE

I. MACÍAS TARRÍO, C. SPIRIDON, A. STOENICĂ

ABSTRACT. The aim of this note is to exhibit proper first Brill-Noether loci inside the moduli spaces $M_{Y,H}(2; c_1, c_2)$ of H -stable rank 2 vector bundles with fixed Chern classes of a certain type on an Enriques surface Y which is covered by a Jacobian Kummer surface X .

1. INTRODUCTION

In the last decades, classical Brill-Noether theory, concerning line bundles on projective curves, has been extended to higher rank bundles on varieties of arbitrary dimension. In [4], Costa and Miró-Roig introduced the Brill-Noether loci W_H^k , whose set theoretic support is formed by those bundles in M_H having at least k independent sections, where M_H denotes the moduli space of H -stable vector bundles on the smooth polarized algebraic variety (X, H) of some rank and some fixed Chern classes.

Standard questions of generalized Brill-Noether theory naturally arose from the ones of the classical theory and concern non-emptiness, irreducibility, dimension of the irreducible components or singularities of W_H^k . These problems are very far from being solved for an arbitrary variety. Little is known even in the surface case, despite the progress which has been made for the projective plane \mathbb{P}^2 [7], K3 surfaces [13, 14], Hirzebruch surfaces [5], ruled surfaces [3] or fibered surfaces [16]. We indicate [2] for a survey regarding higher rank Brill-Noether theory for surfaces.

In this note, we analyze the case of an Enriques surface Y such that its K3 cover is a Jacobian Kummer surface and provide a way of choosing the Chern classes c_1, c_2 such that first Brill-Noether locus $W_H^1(2; c_1, c_2)$ is non-empty and strictly contained in the moduli space $M_{Y,H}(2; c_1, c_2)$ for any polarization H . The reasons for considering the above setup are a good understanding of the Picard lattice of a Jacobian Kummer surface and a theorem of F. Takemoto as reinterpreted by H. Kim [12, Section 2] which gives a recipe to construct rank 2 stable vector bundles on an Enriques surface by pushing forward line bundles on its K3 cover.

The outline of this note is as follows. We start by recalling in the first part of Section 2 the basics of higher rank Brill-Noether theory and then we discuss Jacobian Kummer surfaces. In Section 3 we present sufficient

Date: September 27, 2024.

2020 Mathematics Subject Classification. 14J60, 14J26.

conditions for line bundles on a Jacobian Kummer surface in order to obtain a proper first Brill-Noether locus on the corresponding Enriques surface. This is the content of Theorem 3.3. After this, in Propositions 3.6 and 3.9, we give general examples of line bundles satisfying the hypothesis of Theorem 3.3. Some specific examples are also discussed.

Acknowledgements. The authors would like to thank professor Marian Aprodu for suggesting the problem and helpful discussions.

I. Macías Tarrío was partly supported by PID2020-113674GB-I00. C. Spiridon and A. Stoenică were supported by the PNRR grant CF 44/14.11.2022 *Cohomological Hall algebras of smooth surfaces and applications*.

2. PRELIMINARIES

2.1. Generalized Brill-Noether theory. For the purpose of this note, we will place ourselves in the surface case and work over \mathbb{C} . Let us fix a smooth projective surface S and a polarization (i.e. ample line bundle) H on S . The pair (S, H) is usually called a *polarized surface*. Recall first the definition of slope stability according to which we construct the moduli space of stable bundles on S .

Definition 2.1. If \mathcal{E} is a vector bundle on S , we say that \mathcal{E} is stable with respect to H (or H -stable) if for any coherent subsheaf \mathcal{F} of \mathcal{E} with $0 < \text{rk}(\mathcal{F}) < \text{rk}(\mathcal{E})$ the following inequality holds true

$$\mu_H(\mathcal{F}) := \frac{c_1(\mathcal{F}) \cdot H}{\text{rk}(\mathcal{F})} < \frac{c_1(\mathcal{E}) \cdot H}{\text{rk}(\mathcal{E})} := \mu_H(\mathcal{E})$$

We denote by $M_H := M_{S,H}(r; c_1, c_2)$ the moduli space of H -stable vector bundles on S of rank r with fixed Chern classes c_i , for $i = 1, 2$. We recall the following (see e.g. [9, Section 4.5]):

Theorem 2.2. *If non-empty, M_H is a quasi-projective variety and each irreducible component of it has dimension at least $2rc_2 - (r-1)c_1^2 - (r^2 - 1)\chi(\mathcal{O}_S)$.*

Theorem 2.3. [4, Theorem 2.3] *Let (S, H) and $M_H = M_{S,H}(r; c_1, c_2)$ as above. We assume the following further condition:*

(*) $h^2(\mathcal{E}) = 0$ for any $\mathcal{E} \in M_H$;

Then, for any $k \geq 0$, there exists a determinantal variety $W_H^k(r; c_1, c_2)$, called the k -th Brill-Noether loci, such that

$$\text{Supp}(W_H^k(r; c_1, c_2)) = \{\mathcal{E} \in M_H \mid h^0(\mathcal{E}) \geq k\}.$$

Moreover, each non-empty irreducible component of $W_H^k(r; c_1, c_2)$ has dimension at least

$$\begin{aligned} \rho_H^k(r; c_1, c_2) &:= \dim(M_H) - k(k - \chi(r; c_1, c_2)) = \\ &= \dim(M_H) - k(k - (r\chi(\mathcal{O}_X) + \frac{1}{2}c_1 \cdot K_S - \frac{1}{2}c_1^2 + c_2)) \end{aligned}$$

and

$$W_H^{k+1}(r; c_1, c_2) \subseteq \text{Sing}(W_H^k(r; c_1, c_2))$$

whenever $W_H^k(r; c_1, c_2) \neq M_H$.

As one might expect, the number $\rho_H^k(r; c_1, c_2)$ is called the *generalized Brill-Noether number* and it is the *expected dimension* of $W_H^k(r; c_1, c_2)$.

The theorem above shows in particular that the Brill-Noether loci form a filtration of the moduli space $M_{S,H}(r; c_1, c_2)$:

$$M_H = W_H^0(r; c_1, c_2) \supseteq W_H^1(r; c_1, c_2) \supseteq \dots \supseteq W_H^k(r; c_1, c_2) \supseteq \dots$$

As noticed in [4, Corollary 2.3], the vanishing assumption (*) of Theorem 2.3 holds true when $(c_1.H) \geq (rK_S.H)$. The latter is true in particular if the first Chern class c_1 is effective and the canonical divisor on S is numerically trivial. Thus, for the purpose of our note, we record the following:

Remark 2.4. Let Y be an Enriques surface (i.e. a smooth projective surface with $q(Y) = p_a(Y) = 0$ and $2K_Y = 0$) and $c_i \in H^{2i}(Y, \mathbb{Z})$, for $i = 1, 2$, such that c_1 is effective. Then, for any polarization H , the vanishing condition (*) holds true. Consequently, for any $r \geq 2$ and for any $k \geq 0$ there exists the k -th Brill-Noether locus $W_H^k(r; c_1, c_2)$.

2.2. Jacobian Kummer surfaces. In the sequel, we recall the construction of a Jacobian Kummer surface (see e.g. [10], [11]). Let C be a smooth complex projective curve of genus 2 and $\mathcal{A} = J(C)$ its Jacobian variety. Note that \mathcal{A} is an abelian surface which comes with an involution $\iota : \mathcal{A} \rightarrow \mathcal{A}$ with 16 fixed points. The quotient $\mathcal{K} = \mathcal{A}/\iota$ can be identified with the so-called Kummer quartic, which is a surface of degree 4 in \mathbb{P}^3 with the maximal possible number of 16 singular points. Each singular point is a node. Moreover, there are precisely 16 planes in \mathbb{P}^3 touching \mathcal{K} along a conic. Each of these conics is called a trope. An important feature of \mathcal{K} is the so-called (16_6) -configuration, that is any node lies on exactly 6 tropes and each trope passes through exactly 6 nodes.

The Jacobian Kummer surface $X = \text{Kum}(\mathcal{A})$ is the minimal desingularization of \mathcal{K} . The exceptional curves on X lying over the nodes of \mathcal{K} and the proper transforms of the tropes of \mathcal{K} will be also called nodes and respectively tropes of X . The nodes and the tropes of the Jacobian Kummer surface X form two families of 16 mutually disjoint (-2) -curves on X . They can be described more explicitly in order to have a better understanding of the Picard lattice of X . For references we indicate [15, Section 4], [1, Section 2].

Take p_1, \dots, p_6 the Weierstrass points of C . The 2-torsion points of \mathcal{A} are $[0]$ and $[p_i - p_j]$, $1 \leq i < j \leq 6$. Accordingly, we denote the nodes of X by E_0 and E_{ij} , $1 \leq i < j \leq 6$. The curve C has also precisely 16 theta characteristics (i.e. divisor classes $D \in \text{Pic}(C)$ such that $2D \sim K_C$), namely $[p_i]$, $1 \leq i \leq 6$ and $[p_i + p_j - p_6]$ with $1 \leq i < j \leq 5$, giving rise to 16 theta divisors on \mathcal{A} , whose proper transforms are the tropes of X . Accordingly, we denote the tropes of X by T_i , $1 \leq i \leq 6$ and T_{ij6} , $1 \leq i < j \leq 5$.

Let (\cdot) be the intersection form on X . The following lemma describes the Picard lattice of X (cf. [15]):

Lemma 2.5. *The Picard group of X is generated over \mathbb{Z} by E_0 , E_{ij} , T_i , T_{ij6} and*

- $(E_0.T_i) = 1$ for any $1 \leq i \leq 6$;
- $(E_0.T_{ij6}) = 0$ for any $1 \leq i < j \leq 5$;
- $(E_{ij}.T_k) = \begin{cases} 1 & \text{if } k \in \{i, j\} \\ 0 & \text{otherwise} \end{cases}$
- $(E_{ij}.T_{kl6}) = \begin{cases} 1 & \text{if } \{i, j\} \subseteq \{k, l, 6\} \text{ or } \{i, j\} \cap \{k, l, 6\} = \emptyset \\ 0 & \text{otherwise} \end{cases}$

Due to a result of Keum [10, Theorem 2], there exists a fixed-point-free involution θ of X , giving rise to an Enriques surface $Y = X/\theta$. Let us denote by $\pi : X \rightarrow Y$ the étale double cover of Y . The set of line bundles on X which come from line bundles on Y via pullback is described by a result of Horikawa [8, Theorem 5.1] and recasted in the following form in [1, Lemma 2.2]:

Lemma 2.6. *The image of the map $\pi^* : \text{Pic}(Y) \rightarrow \text{Pic}(X)$ is the set of line bundles $M \in \text{Pic}(X)$ such that $\theta^*M \sim M$.*

The action of θ^* is presented in the table below [1]:

Nodes	Tropes	Nodes	Tropes
E_0	\longleftrightarrow	T_{456}	
E_{12}	\longleftrightarrow	T_3	
E_{13}	\longleftrightarrow	T_2	
E_{14}	\longleftrightarrow	T_{156}	
E_{15}	\longleftrightarrow	T_{146}	
E_{16}	\longleftrightarrow	T_{236}	
E_{23}	\longleftrightarrow	T_1	
E_{24}	\longleftrightarrow	T_{256}	
		E_{25}	\longleftrightarrow
		E_{26}	\longleftrightarrow
		E_{34}	\longleftrightarrow
		E_{35}	\longleftrightarrow
		E_{36}	\longleftrightarrow
		E_{45}	\longleftrightarrow
		E_{46}	\longleftrightarrow
		E_{56}	\longleftrightarrow
		T_{246}	
		T_{136}	
		T_{356}	
		T_{346}	
		T_{126}	
		T_6	
		T_5	
		T_4	

3. BRILL-NOETHER LOCI

3.1. Main result. Let X be a Jacobian Kummer surface, θ its fixed-point-free involution, $Y = X/\theta$ and $\pi : X \rightarrow Y$ the double cover of Y . In the sequel, we will state and prove the main result of this note. But first, let us recall the following important result of Takemoto, as recast by H. Kim [12, Section 2]:

Theorem 3.1. *Let H be a polarization of Y . If a π^*H -stable bundle \mathcal{E} is not isomorphic to $\pi^*\mathcal{F}$ for any bundle \mathcal{F} on Y , then $\pi_*\mathcal{E}$ is H -stable.*

Combining Lemma 2.6 and Theorem 3.1 we obtain a recipe to construct stable rank 2 bundles on the Enriques surface Y :

Corollary 3.2. *If $M \in \text{Pic}(X)$ is a line bundle on X such that $M \not\sim \theta^*M$, then $\pi_*\mathcal{O}_X(M)$ is a stable rank 2 bundle on Y with respect to any polarization of Y .*

Theorem 3.3. *Let $\pi : X \rightarrow Y$ be the double cover map of an Enriques surface Y by a Jacobian Kummer surface X . Let also D be a nonzero effective divisor on X such that:*

- (i) $h^0(\mathcal{O}_X(D)) = 1$;
- (ii) $\theta^*D' \not\sim D'$ for any nonzero effective subdivisor $D' \subseteq D$;
- (iii) $(D)^2 < -4$.

Take $\mathcal{V} = \pi_*\mathcal{O}_X(D)$ and denote by $c_i := c_i(\mathcal{V})$ for $i = 1, 2$. Then for any ample line bundle H on Y one has

$$\emptyset \neq W_H^1(2; c_1, c_2) \subsetneq M_H(2; c_1, c_2)$$

Proof. First of all, condition (ii) implies that $\theta^*D \not\sim D$ and therefore, using Corollary 3.2, we get that $\mathcal{V} = \pi_*\mathcal{O}_X(D)$ is H -stable. We also know that $h^0(\mathcal{V}) = h^0(\mathcal{O}_X(D)) = 1$ since π is a finite map. Then, by definition, $[\mathcal{V}] \in W_H^1(2; c_1, c_2)$, ensuring us about the non-emptiness of this Brill-Noether locus.

Secondly, let us prove that $W_H^1(2; c_1, c_2) \subsetneq M_H(2; c_1, c_2)$. Suppose by way of contradiction that $W_H^1(2; c_1, c_2) = M_H(2; c_1, c_2)$ and let M_0 be the irreducible component of the moduli space $M_H(2; c_1, c_2)$ containing $[\mathcal{V}]$. Since $h^0(\mathcal{V}) = 1$, there exists an open subset $U \subseteq M_0$ such that $h^0(\mathcal{E}) = 1$ for any $[\mathcal{E}] \in U$. Note that for any $[\mathcal{E}] \in U$, the vanishing locus $V(s)$ of the unique nonzero section $s \in H^0(\mathcal{E})$ has a divisorial part $C_{\mathcal{E}}$, which may be zero, and a zero-dimensional part $Z_{\mathcal{E}}$. Since the Picard group of Y is discrete, we may take an open subset $U' \subseteq U$ such that $C_{\mathcal{E}} = C$ for any $[\mathcal{E}] \in U'$.

Claim 1: $C = 0$.

Proof of Claim 1: Any vector bundle \mathcal{E} whose class belongs to U' sits in an exact sequence of the form:

$$0 \rightarrow \mathcal{O}_Y(C) \rightarrow \mathcal{E} \rightarrow \mathcal{I}_{Z_{\mathcal{E}}}(c_1 - C) \rightarrow 0$$

Note that $h^0(\mathcal{E} \otimes \mathcal{O}_Y(-C)) = 1$. As a consequence, since $h^0(\mathcal{V}) = 1$, we get that $h^0(\mathcal{V} \otimes \mathcal{O}_Y(-C)) = 1$. Now from the projection formula:

$$\begin{aligned} 1 = h^0(\mathcal{V} \otimes \mathcal{O}_Y(-C)) &= h^0(\pi_*(\mathcal{O}_X(D) \otimes \pi^*\mathcal{O}_Y(-C))) \\ &= h^0(\mathcal{O}_X(D) \otimes \pi^*\mathcal{O}_Y(-C)) \\ &= h^0(\mathcal{O}_X(D - \pi^*C)) \end{aligned}$$

Consequently, π^*C is an effective subdivisor of D . However, π^*C has to be invariant under the action of θ^* , by Lemma 2.6, but having in mind condition (ii), this is not possible unless $\pi^*C = 0$ and therefore $C = 0$. \square

Now let \mathcal{P} be the projective bundle over $\text{Hilb}_{c_2}(Y)$, whose fiber over the point $Z \in \text{Hilb}_{c_2}(Y)$ is the space of weak isomorphism classes of extensions of type:

$$(1) \quad 0 \rightarrow \mathcal{O}_Y \rightarrow \mathcal{E} \rightarrow \mathcal{I}_Z(c_1) \rightarrow 0$$

Since $h^0(\mathcal{E}) = 1$ for any $[\mathcal{E}] \in U'$, there is a well-defined injective map $U' \rightarrow \mathcal{P}$ sending $[\mathcal{E}]$ to the class of the extension (1). In particular, $\dim U' \leq \dim \mathcal{P}$.

Claim 2: $\dim \mathcal{P} \leq 3c_2 - c_1^2/2 - 1$.

Proof of Claim 2: Let us take a generic point $[\mathcal{E}]$ in U' and $Z \in \text{Hilb}_{c_2}(Y)$ the vanishing locus of the unique section of \mathcal{E} . Let us also denote $\mathcal{J} = \mathcal{I}_Z(c_1) \otimes K_Y$. After twisting the exact sequence

$$0 \rightarrow \mathcal{I}_Z \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_Z \rightarrow 0$$

by $K_Y(c_1)$ we get that

$$\chi(\mathcal{J}) = \chi(K_Y(c_1)) - \chi(\mathcal{O}_Z)$$

Consequently,

$$(2) \quad h^1(\mathcal{J}) = h^0(\mathcal{J}) + h^2(\mathcal{J}) - \frac{c_1^2}{2} - 1 + c_2$$

On the other hand, from the exact sequence

$$0 \rightarrow K_Y \rightarrow \mathcal{E} \otimes K_Y \rightarrow \mathcal{I}_Z(c_1) \otimes K_Y \rightarrow 0$$

we deduce that $h^0(\mathcal{J}) = h^0(\mathcal{E} \otimes K_Y)$. But since \mathcal{E} is generic

$$h^0(\mathcal{E} \otimes K_Y) \leq h^0(\pi_* \mathcal{O}_X(D) \otimes K_Y)$$

and then

$$\begin{aligned} h^0(\pi_* \mathcal{O}_X(D) \otimes K_Y) &= h^0(\pi_*(\mathcal{O}_X(D) \otimes \pi^* K_Y)) \\ &= h^0(\mathcal{O}_X(D) \otimes \pi^* K_Y) \\ &= h^0(\mathcal{O}_X(D) \otimes \mathcal{O}_X) = 1 \end{aligned}$$

We also note that $h^2(\mathcal{E} \otimes K_Y) = h^0(\mathcal{E}^\vee) = 0$ and thus, using the last exact sequence, we obtain the vanishing of $h^2(\mathcal{J})$. Therefore, formula (2) ensures us that:

$$h^1(\mathcal{J}) \leq c_2 - \frac{c_1^2}{2}$$

Now, the claim follows since $\dim \mathcal{P} = \dim \text{Hilb}_{c_2}(Y) + \text{ext}^1(\mathcal{I}_Z(c_1), \mathcal{O}_Y) - 1 = 2c_2 + h^1(\mathcal{I}_Z(c_1) \otimes K_Y) - 1$. \square

On the other hand, $\dim U' = \dim U \geq 4c_2 - c_1^2 - 3\chi(\mathcal{O}_Y) = 4c_2 - c_1^2 - 3$, by Theorem 2.2. Putting everything together we obtain:

$$4c_2 - c_1^2 - 3 \leq \dim U \leq \dim \mathcal{P} \leq 3c_2 - c_1^2/2 - 1$$

which implies that $c_2 - \frac{1}{2}c_1^2 \leq 2$.

Finally, by [6, Propositions 27, 28] one has that $c_1^2 = [\pi_* D]^2$ and $c_2 = \frac{1}{2}([\pi_* D]^2 - \pi_*(D^2)) = \frac{1}{2}[\pi_* D]^2 - \frac{1}{2}(D^2)$. Hence we obtain that $c_2 - \frac{1}{2}c_1^2 = -\frac{1}{2}D^2 > 2$, which is a contradiction. \square

Note that in order to show the strict inclusion of the first Brill-Noether locus $W_H^1(2; c_1, c_2)$ in the moduli space $M_H(2; c_1, c_2)$, we actually proved that the irreducible component of $M_H(2; c_1, c_2)$ to which $[\mathcal{V}]$ belongs is not contained in $W_H^1(2; c_1, c_2)$. This fact raises the following question:

Question 3.4. Is every irreducible component of $M_H(2; c_1, c_2)$ not contained in the first Brill-Noether locus?

Another natural question would concern the non-emptiness of the second Brill-Noether locus, $W_H^2(2; c_1, c_2)$. Our previous result does prove the non-triviality of the first locus if we have a bundle with certain properties, but going further to the second locus would require finding a bundle with exactly two global sections and whose pushforward has the same Chern classes as the previous one. It seems possible to find some specific bundles on the Jacobian Kummer surface with exactly two global sections, but controlling the Chern classes of the pushforwards and making sure that they are equal to the first ones is a much more difficult task.

The relevance of the last condition (iii) of Theorem 3.3 is underlined in the following example.

Example 3.5. Let D be a nonzero effective divisor on X and let $\mathcal{V} = \pi_* \mathcal{O}_X(D)$ and $c_i = c_i(\mathcal{V})$ for $i = 1, 2$. Then $h^0(\mathcal{E}) \geq (D^2)/2 + 2$, for any $[\mathcal{E}] \in M_H(2; c_1, c_2)$. Indeed, by Remark 2.4, $h^2(\mathcal{E}) = 0$, hence $h^0(\mathcal{E}) \geq \chi(\mathcal{E}) = \chi(\mathcal{V}) = (D^2)/2 + 2$. Therefore, if D is either a node or a trope, then $(D)^2 = -2$ and so $h^0(\mathcal{E}) \geq 1$ for any $[\mathcal{E}] \in M_H(2; c_1, c_2)$, which shows that $W_H^1(2; c_1, c_2) = M_H(2; c_1, c_2)$ in this case.

3.2. Examples. As stated before, the theorem does prove the non-emptiness and nontriviality of the first Brill-Noether locus for some specific values of the Chern classes. The next task should obviously be to show that the class of divisors with the properties mentioned in the hypothesis is non-empty. In this section, we construct general examples of divisors on a Jacobian Kummer surface X satisfying the conditions of Theorem 3.3.

Proposition 3.6. *Let $D = \sum_{i=1}^n E_i$, with the E_i 's not necessarily distinct nodes. Then:*

- (i) $h^0(\mathcal{O}_X(D)) = 1$;
- (ii) *for any nonzero effective subdivisor $D' \subseteq D$ one has $\theta^* D' \not\sim D'$;*
- (iii) $(D)^2 < 4$ *if and only if $n \geq 3$.*

Proof. (i) We write $D_n = \sum_{i=1}^n E_i$. We will first prove by induction on the number of terms in the sum the following claim: $h^1(\mathcal{O}_{D_n}) = 0$ for all $n \geq 1$.

For $n = 1$, consider the following exact sequence:

$$0 \rightarrow \mathcal{O}_X(-E_1) \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_{E_1} \rightarrow 0$$

Since E_1 is a (-2) -curve on a K3 surface, we know it is smooth and rational, therefore $\mathcal{O}_{E_1} \simeq \mathcal{O}_{\mathbb{P}^1}$ and $h^0(\mathcal{O}_{E_1}) = 1$, $h^1(\mathcal{O}_{E_1}) = h^2(\mathcal{O}_{E_1}) = 0$. Since $D_1 = E_1$, we obtain what we wanted.

Suppose now that $n \geq 2$ and that the previous claim is true for $n - 1$. In this case, $D_n = E_n + D_{n-1}$. Consider the exact sequence:

$$0 \rightarrow \mathcal{O}_{E_n}(-D_{n-1}) \rightarrow \mathcal{O}_{D_n} \rightarrow \mathcal{O}_{D_{n-1}} \rightarrow 0$$

Again, $E_n \simeq \mathbb{P}^1$ and $\mathcal{O}_{E_n}(-D_{n-1}) \simeq \mathcal{O}_{\mathbb{P}^1}(-E_n \cdot D_{n-1})$. We have that $-E_n \cdot D_{n-1} \geq 0$ because the product of E_n with any node is either 0 (if they are distinct) or -2 (if they are equal), so $h^1(\mathcal{O}_{E_n}(-D_{n-1})) = 0$ and from the long exact sequence in cohomology we obtain $h^1(\mathcal{O}_{D_n}) = h^1(\mathcal{O}_{D_{n-1}})$, which is zero by the induction hypothesis.

Now, consider the following exact sequence:

$$0 \rightarrow \mathcal{O}_X(-D_n) \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_{D_n} \rightarrow 0$$

We now know that $h^1(\mathcal{O}_{D_n}) = h^2(\mathcal{O}_{D_n}) = 0$, therefore, from the associated long exact sequence we obtain the equality $h^2(\mathcal{O}_X(-D_n)) = h^2(\mathcal{O}_X)$. But from Serre duality $h^2(\mathcal{O}_X(-D_n)) = h^0(\mathcal{O}_X(D_n))$, and since $h^2(\mathcal{O}_X) = 1$, we obtain the desired result.

(ii) For the second part, it is sufficient to prove it only for D , since any nonzero effective subdivisor would also be a sum of nodes and the same argument could be applied in the respective case. Write $D = \sum_{i \in I} a_i E_i$ with $a_i > 0$ and E_i distinct nodes. Take some $j \in I$ and simply note that $(D.E_j) = -2a_j < 0$, whereas $(\theta^* D.E_j) \geq 0$.

(iii) Since the nodes don't intersect, $(D)^2 = \sum_{i=1}^n (E_i)^2 = -2n$. \square

Remark 3.7. The last proposition remains obviously true if we replace nodes by tropes. Notice that for (i) the same argument works for sums of nodes and tropes, as long as there is no intersection between the nodes and the tropes, or if there exists either a node which intersects no trope, or a trope which intersects no node. For the former case, just choose in the proof of the claim E_n to be, for example, that node which intersects no trope.

Example 3.8. According to Proposition 3.6, some divisors which satisfy the conditions of Theorem 3.3 should look like $E_0 + E_{13} + E_{13}$, or $3E_{23} + E_{14} + 2E_{56}$. The only thing that we should care about is having at least 3 terms in our sum (counting multiplicities). The same idea applies for sums of tropes, as mentioned in the previous remark.

For the other types of divisors mentioned in the remark, we need to see how to choose some which fit the other conditions of Theorem 3.3. The last condition is the easiest to handle by far, since we know very well how the intersection product behaves. The second one is a bit trickier, and the following two results will tackle the problem of finding divisors which satisfy it.

Proposition 3.9. *Let $D = \sum_{i \in I} a_i E_i + T$ with the E_i 's distinct nodes, $a_i > 0$ and $(E_i.T) = 0$ for all $i \in I$. Then $D \not\sim \theta^* D$ if and only if D has one of the following properties:*

a) for all i , $E_i \neq \theta^* T$;

b) there exists $i_0 \in I$ such that $E_{i_0} = \theta^*T$ and either $a_{i_0} \geq 2$, or $a_{i_0} = 1$, but $I \setminus \{i_0\} \neq \emptyset$.

Proof. Let us first remark that $(D.T) = -2$.

" \Leftarrow " If a) holds, then $(\theta^*D.T) = (D.\theta^*T) = 0$, so they can not be equivalent. If b) holds, then $(\theta^*D.T) = -2a_i$. In the first situation, we will again obtain that $(\theta^*D.T) \neq (D.T)$. In the second one, assume by contradiction that $D \sim \theta^*D$. Then $D - (T + \theta^*T) \sim \theta^*D - (T + \theta^*T)$. But the left hand side contains only nodes and the right hand side only tropes, and both are nonzero due to our hypothesis, therefore we arrive at a contradiction, so $D \not\sim \theta^*D$.

" \Rightarrow " Suppose there exists $i_0 \in I$ such that $E_{i_0} = \theta^*T$. Also, suppose that $a_{i_0} = 1$. If $I = \{i_0\}$, then $D = T + \theta^*T = \theta^*D$ and we obtain a contradiction. \square

Example 3.10. Let us first consider the divisor $D = E_{12} + T_3 = \theta^*T_3 + T_3$. We know that the node and the trope don't intersect, but it is clear that $\theta^*D = D$. But if we had instead $D_1 = 2E_{12} + T_3$ or $D_2 = E_{12} + E_{14} + T_3$, the previous proposition implies that D_1 and D_2 are not θ^* -invariant.

Corollary 3.11. *Let $D = \sum_{i \in I} a_i E_i + T$ with the E_i 's distinct, $a_i > 0$ and $(E_i.T) = 0$ for all $i \in I$. Then D has no θ^* -invariant nonzero effective subdivisor if and only if for all i , $E_i \neq \theta^*T$.*

Proof. Let us first remark that in this case we have three types of nonzero effective subdivisors: T , some sum of nodes or T plus some nodes. We can ignore the first two cases since those subdivisors are clearly not θ^* -invariant due to how θ^* turns nodes into tropes and vice versa. Therefore, the only subdivisors which are relevant for us are those from the last category and we will only focus on them.

" \Rightarrow " In this case, D is not θ^* -invariant and we know its structure via the previous proposition. Assume by contradiction that we are in the second case, *i.e.* there exists $i_0 \in I$ such that $E_{i_0} = \theta^*T$. But then D would have a nonzero θ^* -invariant effective subdivisor, namely $\theta^*T + T$, and we would have a contradiction. Therefore we are in the first case, which is what we wanted.

" \Leftarrow " Due to our initial remark, let us consider a nonzero effective subdivisor of D of the form T plus some nodes which appear in D . Since for all i , $E_i \neq \theta^*T$, this subdivisor is not θ^* -invariant due to the previous proposition. In consequence, D has no θ^* -invariant nonzero effective subdivisor, which is what we wanted to prove. \square

Remark 3.12. Let D be as in the corollary with the property that $E_i \neq \theta^*T$ for all $i \in I$. Then, because of the corollary and the previous remark, we know that D satisfies the first two conditions of 3.3. For the last one, notice that $(D)^2 = \sum_{i \in I} (E_i)^2 + (T)^2 = (-2) \cdot (\#I + 1)$, so the third condition is satisfied if and only if $\#I \geq 2$.

REFERENCES

- [1] M. Aprodu and Y. Kim, *On the Borisov–Nuer conjecture and the image of the Enriques-to- $K3$ map*, Math. Ann. **293** (2020), 1044–1052.
- [2] I. Coskun, J. Huizenga, and H. Nuer, *Brill-Noether Theory of moduli spaces of sheaves on surfaces*, [arXiv:2306.11033](https://arxiv.org/abs/2306.11033).
- [3] L. Costa and I. Macías Tarrío, *Brill-Noether Theory of stable vector bundles on ruled surfaces*, Mediterr. J. Math. **21**, No. 3 (2024).
- [4] L. Costa and R. M. Miró-Roig, *Brill-Noether Theory for moduli spaces of sheaves on algebraic varieties*, Forum Math. **22**, No. 3 (2010), 411–432.
- [5] L. Costa and R. M. Miró-Roig, *Brill–Noether theory on Hirzebruch surfaces*, J. Pure Appl. Algebra **214**, No. 9 (2010), 1612–1622.
- [6] R. Friedman, *Algebraic surfaces and holomorphic vector bundles*, Universitext, Springer, New-York (1998).
- [7] B. Gould, W. Lee, and Y. Liu, *Higher rank Brill-Noether theory on \mathbb{P}^2* , Int. Math. Res. Not. **24** (2022), 22096–22137.
- [8] E. Horikawa, *On the Periods of Enriques Surfaces. I*, Mathematische Nachrichten **234** (1978), 73–88.
- [9] D. Huybrechts and M. Lehn, *The geometry of moduli spaces of sheaves, 2nd ed.*, Cambridge University Press (2010).
- [10] J. H. Keum, *Every algebraic Kummer surface is the $K3$ -cover of an Enriques surface*, Nagoya Math. J. **118** (1990), 99–110.
- [11] J. H. Keum, *Automorphisms of Jacobian Kummer surfaces*, Compositio Math. **107** (1997), 269–288.
- [12] H. Kim, *Moduli spaces of stable vector bundles on Enriques surfaces*, Nagoya Math. J. **150** (1998), 85–94.
- [13] M. Leyenson, *On the Brill-Noether Theory for $K3$ surfaces*, Cent. Eur. J. Math. **10**, No. 4 (2012), 1486–1540.
- [14] M. Leyenson, *On the Brill-Noether Theory for $K3$ surfaces II*, [arXiv:math/0602358](https://arxiv.org/abs/math/0602358).
- [15] H. Ohashi, *Enriques surfaces covered by Jacobian Kummer surfaces*, Nagoya Math. J. **195** (2009), 165–186.
- [16] G. Reyes-Ahumada, L. Roa-Leguizamón, and H. Torres-López, *A note on rank two stable bundles over surfaces*, Glasgow Mathematical Journal **64**, No. 2 (2022), 397–410.

I. MACÍAS TARRÍO: FACULTAT DE MATEMÀTIQUES I INFORMÀTICA, DEPARTAMENT DE MATEMÀTIQUES I INFORMÀTICA,
 GRAN VIA DE LES CORTS CATALANES 585, 08007 BARCELONA, SPAIN
Email address: irene.macias@ub.edu

C. SPIRIDON: SIMION STOILOW INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, RESEARCH GROUP OF THE PROJECT "COHOMOLOGICAL HALL ALGEBRAS OF SMOOTH SURFACES AND APPLICATIONS" - C.F. 44/14.11.2022
 P.O. BOX 1-764, RO-014700 BUCHAREST, ROMANIA, AND
 FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF BUCHAREST, ROMANIA
Email address: cspiridon@imar.ro

A. STOENICĂ: FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF BUCHAREST, ROMANIA, AND SIMION STOILOW INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, RESEARCH GROUP OF THE PROJECT "COHOMOLOGICAL HALL ALGEBRAS OF SMOOTH SURFACES AND APPLICATIONS" - C.F. 44/14.11.2022
Email address: andrei.stoenica@my.fmi.unibuc.ro