

A BOUND ON THE μ -INVARIANTS OF SUPERSINGULAR ELLIPTIC CURVES

RYLAN GAJEK-LEONARD

ABSTRACT. Let E/\mathbb{Q} be an elliptic curve and let p be a prime of good supersingular reduction. Attached to E are pairs of Iwasawa invariants μ_p^\pm and λ_p^\pm which encode arithmetic properties of E along the cyclotomic \mathbb{Z}_p -extension of \mathbb{Q} . A well-known conjecture of B. Perrin-Riou and R. Pollack asserts that $\mu_p^\pm = 0$. We provide support for this conjecture by proving that for all but finitely many primes, if $\lambda_p^\pm = 0$ then μ_p^\pm is at most 1 or 2, and if $\lambda_p^\pm = 1$ then μ_p^\pm is at most 3 or $p+1 + \lceil \frac{p}{\log p} \rceil$, depending on sign. As a corollary, we show that if E has rank 0 then $\mu_p^- \leq 1$ and $\mu_p^+ \leq 2$ hold on density 1 set of good supersingular primes.

1. INTRODUCTION

Let E/\mathbb{Q} be an elliptic curve and fix a prime p of good reduction. Attached to E is the p -primary Selmer group $\text{Sel}(E/\mathbb{Q}_{\text{cyc}})$, where \mathbb{Q}_{cyc} denotes the cyclotomic \mathbb{Z}_p -extension of \mathbb{Q} . This group fits into the exact sequence

$$(1) \quad 0 \rightarrow E(\mathbb{Q}_{\text{cyc}}) \otimes \mathbb{Q}_p/\mathbb{Z}_p \rightarrow \text{Sel}(E/\mathbb{Q}_{\text{cyc}}) \rightarrow \text{III}(E/\mathbb{Q}_{\text{cyc}}) \rightarrow 0,$$

where III denotes the p -part of the Shafarevich-Tate group, and therefore encodes many arithmetic properties of E along the cyclotomic line.

If p is a prime of ordinary reduction then $\text{Sel}(E/\mathbb{Q}_{\text{cyc}})$ is cotorsion as an Iwasawa module (see [7, Theorem 17.4]) and its characteristic ideal is therefore generated by a polynomial $L_p^{\text{alg}} \in \mathbb{Z}_p[T]$. The algebraic Iwasawa invariants λ_p^{alg} and μ_p^{alg} measure the degree and p -divisibility of L_p^{alg} , respectively. If $E[p]$ is irreducible as a $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ -module, a well-known conjecture of Greenberg [5, Conjecture 1.11] asserts that $\mu_p^{\text{alg}} = 0$.

If p is a prime of supersingular reduction then $\text{Sel}(E/\mathbb{Q}_{\text{cyc}})$ is no longer cotorsion, however Kobayashi [9] introduced signed Selmer groups $\text{Sel}^\pm(E/\mathbb{Q}_{\text{cyc}})$ which are cotorsion and encode analogous arithmetic data. In particular, the characteristic ideals of $\text{Sel}^\pm(E/\mathbb{Q}_{\text{cyc}})$ are generated by polynomials $L_{p,\text{alg}}^\pm \in \mathbb{Z}_p[T]$ which have associated pairs of Iwasawa invariants μ_{alg}^\pm and λ_{alg}^\pm . In the supersingular setting, $E[p]$ is automatically irreducible and it is similarly conjectured (see [12, Conjecture 6.3] and [11, Conjecture 7.1]) that $\mu_{\text{alg}}^\pm = 0$.

Recently, Chakravarthy [3, Theorem 1.4] made significant progress towards Greenberg's conjecture by proving that $\mu_p^{\text{alg}} = 0$ for all but finitely many primes of good ordinary reduction. In this article, we employ similar methods in the supersingular setting and offer a modest bound on the size of the signed *analytic* μ -invariants.

2020 *Mathematics Subject Classification*. Primary 11R23.

Key words and phrases. Iwasawa theory, elliptic curves, p -adic L -functions.

Theorem 1.1. *For all but finitely many primes p of good supersingular reduction for E , we have:*

- (1) *If $\lambda_{an}^- = 0$ then $\mu_{an}^- \leq 1$.*
- (2) *If $\lambda_{an}^+ = 0$ then $\mu_{an}^+ \leq 2$.*
- (3) *If $\lambda_{an}^- = 1$ then $\mu_{an}^- \leq p + 1 + \lceil \frac{p}{\log p} \rceil$.*
- (4) *If $\lambda_{an}^+ = 1$ then $\mu_{an}^+ \leq 3$.*

In the above theorem, the Iwasawa invariants λ_{an}^\pm and μ_{an}^\pm are those attached to Pollack's plus/minus p -adic L -functions $L_{p,an}^\pm \in \mathbb{Z}_p[[T]]$ defined in [12]. The construction of these p -adic L -functions requires $a_p = 0$, which is automatically true when $p > 3$. The main conjecture of Iwasawa theory in this setting asserts that $L_{p,an}^\pm$ and $L_{p,alg}^\pm$ generate the same ideal in $\mathbb{Z}_p[[T]]$, and in particular that

$$(2) \quad \lambda_{alg}^\pm = \lambda_{an}^\pm \quad \text{and} \quad \mu_{alg}^\pm = \mu_{an}^\pm.$$

Thus, Theorem 1.1 provides support for the vanishing of μ_{alg}^\pm .

Remark 1.2. The main conjecture is known to hold in many cases: the CM case was established by Pollack and Rubin [14], and Kobayashi [9, Theorem 1.3] proved the containment $(L_{p,an}^\pm) \subseteq (L_{p,alg}^\pm)$ for non CM curves. A proof of the full supersingular main conjecture was recently announced by Burungale, Skinner, Tian, and Wan [2, Theorem 1.2].

We henceforth assume (2) and write λ^\pm, μ^\pm to mean either analytic or algebraic invariants. We note that the assumption $\lambda^\pm \in \{0, 1\}$ is relatively mild in that standard conjectures predict it should hold for 'most' elliptic curves at 'most' supersingular primes. More specifically, a classical folklore conjecture (see [1]) asserts that 100% of rational elliptic curves have Mordell-Weil rank 0 or 1. Additionally, Kundu and Ray conjecture [8, pg. 2] that λ^\pm coincides with the Mordell-Weil rank on a density 1 set of good supersingular primes. Thus, one expects that for 100% of elliptic curves E/\mathbb{Q} , we have $\lambda^\pm \in \{0, 1\}$ for 100% of good supersingular primes. In fact, the conjecture of Kundu and Ray is known to hold in the rank 0 case by [8, Theorem 3.8]. Together with Theorem 1.1, we therefore obtain the following.

Corollary 1.3. *Suppose E/\mathbb{Q} has Mordell-Weil rank $r = 0$. Then the inequalities $\mu^+ \leq 2$ and $\mu^- \leq 1$ hold on a density 1 set of good supersingular primes.*

The crux of Chakravarty's proof in the ordinary setting is [3, Lemma 4.3], which bounds the size of the modular symbols defining the p -adic L -function just enough to conclude $\mu = 0$. In the supersingular setting, the signed p -adic L -functions are defined via a decomposition theorem of Pollack [12, Theorem 5.6], and in particular they are not as immediately understood in terms of modular symbols. Instead, the approach taken here is to apply Chakravarty's bound to the sequence of Mazur-Tate elements for E (which are defined using modular symbols), and then relate the Iwasawa invariants of the Mazur-Tate elements to those of the signed p -adic L -functions. Using our argument, Chakravarty's bound alone does not appear sharp enough to deduce the vanishing of μ^\pm .

The assumption that $\lambda^\pm = 0$ (resp., $\lambda^\pm = 1$) is used primarily to simplify our argument, as it allows one to bound μ^\pm in terms of the valuation of the constant (resp., linear or degree p) term of the Mazur-Tate elements for E . In particular, insisting on small λ -invariants enables one (via Proposition 3.1) to focus on Mazur-Tate elements 'near the bottom' of the cyclotomic \mathbb{Z}_p -extension. Roughly speaking,

the fact that base-layer Mazur-Tate elements live in smaller group algebras is crucial for obtaining sharper bounds on μ -invariants. For larger λ -invariants, one could certainly bound μ -invariants in terms of higher-degree coefficients of the Mazur-Tate elements, however one would also be forced to look higher up the \mathbb{Z}_p -tower, leading to even weaker bounds for μ^\pm .

1.1. Acknowledgements. We are grateful to A. Chakravarthy and J. Hatley for their careful reading and helpful suggestions in the preparation of this article.

2. IWASAWA INVARIANTS

Fix a prime p and let $F \in \Lambda = \mathbb{Z}_p[[T]]$. By the Weierstrass preparation theorem [17, Theorem 7.3] there are unique nonnegative integers λ and μ for which

$$F = p^\mu(T^\lambda + pF_0)U,$$

for some $F_0 \in \mathbb{Z}_p[T]$ of degree $< \lambda$ and some $U \in \Lambda^\times$. In terms of the coefficients of $F = \sum_{i \geq 0} a_i T^i$, we have

$$\begin{aligned} \mu &= \min\{\text{ord}_p(a_i) \mid i \geq 0\}, \\ \lambda &= \min\{i \mid \text{ord}_p(a_i) = \mu\}. \end{aligned}$$

Let $\omega_n = (1+T)^{p^n} - 1$ and $\Lambda_n = \Lambda/(\omega_n)$. Since $\Lambda = \varprojlim \Lambda_n$, there are projection maps

$$\pi_n : \Lambda \rightarrow \Lambda_n, \quad F \mapsto F \bmod \omega_n.$$

and we can define the *refined* Iwasawa invariants of $\pi_n(F)$ as follows. Since ω_n is a distinguished polynomial, the division algorithm for distinguished polynomials in Λ allows us to write

$$F = \omega_n Q_n + F_n,$$

for some unique $Q_n \in \Lambda$ and a polynomial $F_n \in \mathbb{Z}_p[T]$ of degree $< p^n$. Define

$$\begin{aligned} \lambda(\pi_n(F)) &= \lambda(F_n), \\ \mu(\pi_n(F)) &= \mu(F_n). \end{aligned}$$

The following lemma is known in the literature (see [13, Remark 4.3] or [4, Corollary A.7]), though we outline a proof for completeness.

Lemma 2.1. *Let $n \geq 0$. If $\lambda(F) < p^n$ then the Iwasawa invariants of F and $\pi_n(F)$ agree.*

Proof. We may assume $\mu(F) = 0$. Use the division algorithm to write

$$(3) \quad F = \omega_n Q_n + F_n.$$

If $\mu(F_n)$ were positive then (3) implies

$$F \equiv T^{p^n} Q_n \bmod p,$$

which contradicts the fact that $\lambda(F) < p^n$. Hence we must have $\mu(F_n) = 0$. Combining (3) with the Weierstrass decomposition $F = (T^{\lambda(F)} + pF_0)U$ now yields

$$F_n \equiv T^{\lambda(F)}(U - T^{p^n - \lambda(F)} Q_n) \bmod p,$$

but since U is a unit and $\lambda(F) < p^n$, $U - T^{p^n - \lambda(F)} Q_n$ must also be a unit. It follows that $\lambda(F_n) = \lambda(F)$. \square

3. MAIN RESULT

Let E/\mathbb{Q} be an elliptic curve and fix a prime p of good reduction such that $a_p = 0$. Let $L_p^\pm \in \Lambda$ and $\theta_n \in \mathbb{Q}[T]$ be the plus/minus p -adic L -functions and Mazur-Tate elements for E , as defined in §§2.9 and 6.15 of [12], respectively. The definition of both L_p^\pm and θ_n depend on a choice of complex periods $\Omega_E^\pm \in \mathbb{C}$. We henceforth assume that Ω_E^\pm are p -cohomological periods for E , in the sense of [15, §2.2]. The choice of cohomological periods ensures that the coefficients of θ_n are p -integral, i.e., we can view θ_n as elements of the localization $\mathbb{Z}_{(p)}[T] \subseteq \mathbb{Q}[T]$.

We now relate the Iwasawa invariants of θ_n to those of L_p^\pm . Let $q_1 = q_0 = 0$ and define for $n \geq 2$ the sequence

$$q_n = \begin{cases} p^{n-1} - p^{n-2} + \cdots + p - 1 & n \text{ even,} \\ p^{n-1} - p^{n-2} + \cdots + p^2 - p & n \text{ odd.} \end{cases}$$

Let λ^\pm and μ^\pm denote the Iwasawa invariant of L_p^\pm .

Proposition 3.1. *If $n \geq 0$ is even (resp., odd) and $\lambda^- < p^n - q_n$ (resp., $\lambda^+ < p^n - q_n$), then*

$$\begin{aligned} \mu(\theta_n) &= \mu^\pm, \\ \lambda(\theta_n) &= \lambda^\pm + q_n, \end{aligned}$$

where \pm is opposite the parity of n .

Proof. This follows from the argument of [4, page 3], which we reproduce in brief here. Let $\varepsilon_n = \text{sgn}(-1)^n$ denote the parity of n . By [12, Proposition 6.18], we have

$$(4) \quad \theta_n \equiv \omega_n^{-\varepsilon_n} L_p^{-\varepsilon_n} \pmod{\omega_n}.$$

Here

$$\omega_n^+ = \prod_{\substack{1 \leq i \leq n \\ i \text{ even}}} \Phi_{p^i}(1+T) \quad \text{and} \quad \omega_n^- = \prod_{\substack{1 \leq i \leq n \\ i \text{ odd}}} \Phi_{p^i}(1+T),$$

where $\Phi_{p^i}(T)$ is the p^i th cyclotomic polynomial. Since $\lambda(\Phi_{p^n}(1+T)) = p^n - p^{n-1}$, we have $\lambda(\omega_n^{-\varepsilon_n} L_p^{-\varepsilon_n}) = q_n + \lambda^{-\varepsilon_n}$. The sequence $p^n - q_n$ tends to infinity, thus we may take n large enough so that $\lambda(\omega_n^{-\varepsilon_n} L_p^{-\varepsilon_n}) < p^n$. The result now follows from Lemma 2.1. \square

Remark 3.2. The formula for λ -invariants in Proposition 3.1 can also be found in [16, Corollary 8.9], [11, §5], and [15, Theorem 4.1].

Let f be the cuspidal newform attached to E via modularity. For $r \in \mathbb{Q}$, recall the modular symbols of [10] defined by

$$[r]^\pm = \frac{\pi i}{\Omega_E^\pm} \left(\int_r^{i\infty} f(z) dz \pm \int_{-r}^{i\infty} f(z) dz \right).$$

For odd primes p we have by definition

$$(5) \quad \theta_n(T) = \sum_{a \in (\mathbb{Z}/p^{n+1}\mathbb{Z})^\times} [a/p^{n+1}]^+(1+T)^{\log_\gamma a}.$$

Here $0 \leq \log_\gamma(a) \leq p^n - 1$ is the unique integer for which $a \equiv \omega(a)(1+p)^{\log_\gamma(a)} \pmod{p^{n+1}}$, where ω is the mod p cyclotomic character.

Corollary 3.3.

- (1) If $\lambda^- = 0$ then $\mu^- = \text{ord}_p \left(\sum_{a \in (\mathbb{Z}/p\mathbb{Z})^\times} [a/p] \right) < \infty$.
- (2) If $\lambda^+ = 0$ then $\mu^+ = \text{ord}_p \left(\sum_{a \in (\mathbb{Z}/p^2\mathbb{Z})^\times} [a/p^2] \right) < \infty$.
- (3) If $\lambda^- = 1$ then $\mu^- = \text{ord}_p \left(\sum_{a \in (\mathbb{Z}/p^3\mathbb{Z})^\times} \binom{\log_\gamma(a)}{p} [a/p^3] \right) < \infty$.
- (4) If $\lambda^+ = 1$ then $\mu^+ = \text{ord}_p \left(\sum_{a \in (\mathbb{Z}/p^2\mathbb{Z})^\times} \log_\gamma(a) [a/p^2] \right) < \infty$.

Proof. If $\lambda^\pm = 0$ then by Proposition 3.1 we have $\mu^- = \mu(\theta_0)$, $\mu^+ = \mu(\theta_1)$, and $\lambda(\theta_0) = \lambda(\theta_1) = 0$. In particular, the μ -invariants of θ_0 and θ_1 are equal to the valuations their constant coefficients, which are both nonzero as $\lambda(\theta_0) = \lambda(\theta_1) = 0$. The constant coefficient of θ_n can be obtained by setting $T = 0$ in equation (5), yielding the desired result when $\lambda^\pm = 0$.

If $\lambda^+ = 1$ then Proposition 3.1 yields $\mu^+ = \mu(\theta_1)$ and $\lambda(\theta_1) = 1$. The μ -invariant of θ_1 is therefore equal to the valuation of its linear term, which is nonzero as $\lambda(\theta_1) = 1$. The linear term of θ_n can be found by setting $T = 0$ in the derivative

$$\frac{d\theta_n}{dT} = \sum_{\substack{a \in (\mathbb{Z}/p^{n+1}\mathbb{Z})^\times \\ a \neq 1}} \log_\gamma(a) [a/p^{n+1}]^+ (1+T)^{\log_\gamma(a)-1},$$

yielding the result when $\lambda^+ = 1$ (note $\log_\gamma(1) = 0$).

If $\lambda^- = 1$ then Proposition 3.1 yields $\mu^- = \mu(\theta_2)$ and $\lambda(\theta_2) = p$. The μ -invariant of θ_2 is therefore equal to the valuation of its degree p coefficient, which is nonzero as $\lambda(\theta_2) = p$. The p th derivative of θ_n is

$$\theta_n^{(p)} = \sum_{\substack{a \in (\mathbb{Z}/p^{n+1}\mathbb{Z})^\times \\ \log_\gamma(a) \notin \{0, \dots, p-1\}}} \frac{\log_\gamma(a)!}{(\log_\gamma(a) - p)!} [a/p^{n+1}]^+ (1+T)^{\log_\gamma(a)-p},$$

thus Taylor's formula yields

$$(p\text{th coefficient of } \theta_2) = \frac{\theta_2^{(p)}(0)}{p!} = \sum_{a \in (\mathbb{Z}/p^3\mathbb{Z})^\times} \binom{\log_\gamma(a)}{p} [a/p^3]^+.$$

The result when $\lambda^- = 1$ follows (note $\binom{\log_\gamma(a)}{p} = 0$ when $\log_\gamma(a) < p$). \square

Proof of Theorem 1.1. By [3, Lemma 4.3], there are constants c_1 and c_2 depending only on the conductor of E (and not p) such that for any $r \in \mathbb{Q}$,

$$|[r]^+| \leq c_1 + c_2 \log(\text{denominator}(r)).$$

In particular, for any fixed n we can take p large enough so that:

$$(6) \quad |[a/p^n]^+| < p.$$

Recalling that $0 \leq \log_\gamma(a) \leq p^n - 1$ for $a \in (\mathbb{Z}/p^{n+1}\mathbb{Z})^\times$, it follows that for sufficiently large p we have

$$\begin{aligned} \left| \sum_{a \in (\mathbb{Z}/p\mathbb{Z})^\times} [a/p]^+ \right| &\leq \sum_{a \in (\mathbb{Z}/p\mathbb{Z})^\times} |[a/p]^+| < (p-1)p < p^2, \\ \left| \sum_{a \in (\mathbb{Z}/p^2\mathbb{Z})^\times} [a/p^2]^+ \right| &\leq \sum_{a \in (\mathbb{Z}/p^2\mathbb{Z})^\times} |[a/p^2]^+| < (p^2-p)p < p^3, \\ \left| \sum_{a \in (\mathbb{Z}/p^2\mathbb{Z})^\times} \log_\gamma(a)[a/p^2]^+ \right| &\leq \sum_{a \in (\mathbb{Z}/p^2\mathbb{Z})^\times} (p-1) \cdot |[a/p^2]^+| < (p^2-p)(p-1)p < p^4. \end{aligned}$$

By Corollary 3.3, the above inequalities establish the result when $\lambda^\pm = 0$ and when $\lambda^+ = 1$.

To deal with the case $\lambda^- = 1$, note that

$$\begin{aligned} \left| \sum_{a \in (\mathbb{Z}/p^3\mathbb{Z})^\times} \binom{\log_\gamma(a)}{p} [a/p^3]^+ \right| &\leq \sum_{a \in (\mathbb{Z}/p^3\mathbb{Z})^\times} \binom{\log_\gamma(a)}{p} |[a/p^3]^+| \\ (7) \qquad \qquad \qquad &< p \sum_{i=p}^{p^2-1} \binom{i}{p} \\ (8) \qquad \qquad \qquad &= p \binom{p^2}{p+1} \\ (9) \qquad \qquad \qquad &\leq \frac{p^{2p+3}}{(p+1)!} \\ &< \frac{p^{2p+2}}{p!} \\ (10) \qquad \qquad \qquad &\leq p^{p+2+\lceil \frac{p}{\log p} \rceil} \end{aligned}$$

where (7) follows from (6), (8) follows the well-known ‘‘hockey-stick’’ identity [6, (2)], (9) follows from the fact that $\binom{n}{k} \leq \frac{n^k}{k!}$, and (10) follows from Stirling’s approximation (namely, $\log n! \geq n \log n - n$). The desired bound on μ^- now follows from Corollary 3.3. \square

REFERENCES

- [1] M. Bhargava and A. Shankar, *Binary quartic forms having bounded invariants, and the boundedness of the average rank of elliptic curves*, Ann. Math., 181 (2015), no. 1, 191–242.
- [2] A. Burungale, C. Skinner, Y. Tian, and X. Wan, *Zeta elements for elliptic curves and applications*, preprint (2024), [arXiv:2409.01350](https://arxiv.org/abs/2409.01350).
- [3] A. Chakravarthy, *The Iwasawa μ -invariants of elliptic curves over \mathbb{Q}* , preprint (2024), [arXiv:2408.07826](https://arxiv.org/abs/2408.07826).
- [4] R. Gajek-Leonard, *Iwasawa invariants of modular forms with $a_p = 0$* , preprint (2023), [arXiv:2302.05748](https://arxiv.org/abs/2302.05748).
- [5] R. Greenberg, *Iwasawa theory for elliptic curves*, Arithmetic theory of elliptic curves (Cetraro, 1997), Lecture Notes in Math., vol. 1716 (1999), Springer, Berlin, pp. 51–144.
- [6] C. H. Jones, *Generalized Hockey Stick Identities and N -Dimensional Block Walking*, Fibonacci Quarterly, 34 (1996), no. 3, 280–288.
- [7] K. Kato, *p -adic Hodge theory and values of zeta functions of modular forms*, Astérisque, no. 295 (2004), 117–290.
- [8] D. Kundu and A. Ray, *Statistics for Iwasawa Invariants of Elliptic Curves*, Trans. Amer. Math. Soc., 374 (2021), no. 11, 7945–7965.

- [9] S. Kobayashi, *Iwasawa theory for elliptic curves at supersingular primes*, Invent. Math., 152 (2003), no. 1, 1–36.
- [10] B. Mazur, J. Tate, *Refined conjectures of the “Birch and Swinnerton-Dyer type”*, Duke Math. J., 54 (1987), no. 2, 711–750.
- [11] B. Perrin-Riou, *Arithmétique des courbes elliptiques à réduction supersingulière en p* , Experiment. Math., 12 (2003), no. 2, 155–186.
- [12] R. Pollack, *On the p -adic L -function of a modular form at a supersingular prime*, Duke Math. J. 118 (2003), no. 3, 523–558.
- [13] R. Pollack, *An algebraic version of a theorem of Kurihara*, J. Number Theory, 110 (2005), pp. 164–177.
- [14] R. Pollack and K. Rubin, *The main conjecture for CM elliptic curves at supersingular primes*, Ann. of Math., 159 (2004), no. 1, 447–464.
- [15] R. Pollack and T. Weston, *Mazur-Tate elements of nonordinary modular forms*, Duke Math. J., 156 (2011), no. 3, 349–385.
- [16] F. Sprung, *On pairs of p -adic L -functions for weight-two modular forms*, Algebra & Number Theory, 11 (2017), no. 4, 885–928.
- [17] L. C. Washington, *Introduction to cyclotomic fields*, Graduate Texts in Mathematics 83, Springer, New York, 1982.

DEPARTMENT OF MATHEMATICS, UNION COLLEGE, BAILEY HALL 206B, SCHENECTADY, NY 12308, USA

Email address: gajekler@union.edu