The XMM-LSS survey: optical assessment and properties of different X-ray selected cluster classes *

- C. Adami¹, A. Mazure¹, M. Pierre², P.G. Sprimont³, C. Libbrecht², F. Pacaud^{2,8}, N. Clerc², T. Sadibekova³, J. Surdej³
 - , B. Altieri⁴, P.A. Duc², G. Galaz⁷, A. Gueguen², L. Guennou¹, G. Hertling⁷, O. Ilbert¹, J.P. LeFèvre¹⁴, H. Quintana⁷, I. Valtchanov⁴, J.P. Willis⁹
- , M. Akiyama¹², H. Aussel², L. Chiappetti¹⁰, A. Detal³, B. Garilli¹⁰, V. LeBrun¹, O. LeFèvre¹, D. Maccagni¹⁰, J.B. Melin¹³, T.J. Ponman¹¹, D. Ricci³, and L. Tresse¹
- LAM, OAMP, Pôle de l'Etoile Site Château-Gombert 38, Rue Frédréric Juliot-Curie, 13388 Marseille, Cedex 13, France
- ² Laboratoire AIM, CEA/DSM/IRFU/Sap, CEA-Saclay, F-91191 Gif-sur-Yvette Cedex, France
- ³ Institut d'Astrophysique et de Géophysique, Université de Liège, Allée du 6 Août, 17, B5C, 4000 Sart Tilman, Belgium
- ⁴ ESA, Villafranca del Castillo, Spain
- ⁵ UPMC Université Paris 06, UMR 7095, Institut d'Astrophysique de Paris, F-75014, Paris, France
- ⁶ CNRS, UMR 7095, Institut d'Astrophysique de Paris, F-75014, Paris, France
- Departamento de Astronomía y Astrofísica, Pontificia Universidad Católica de Chile, Casilla 306, Santiago 22, Chile
- 8 Argelander-Institut f
 ür Astronomie, University of Bonn, Auf dem H
 ügel 71, 53121 Bonn, Germany
- Department of Physics and Astronomy, University of Victoria, Elliot Building, 3800 Finnerty Road, Victoria, V8V 1A1, BC, Canada
- ¹⁰ INAF-IASF Milano, via Bassini 15, I-20133 Milano, Italy
- School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK
- Astronomical Institute, Tohoku University 6-3 Aramaki, Aoba-ku, Sendai, 980-8578, Japan
- 13 CEA/DSM/IRFU/SPP, CEA Saclay, F-91191 Gif-sur-Yvette, France.
- ¹⁴ CEA/DSM/IRFU/SEDI, CEA Saclay, F-91191 Gif-sur-Yvette, France

Accepted . Received; Draft printed: October 9, 2018

ABSTRACT

Context. XMM and Chandra opened a new area for the study of clusters of galaxies. Not only for cluster physics but also, for the detection of faint and distant clusters that were inaccessible with previous missions.

Aims. This article presents 66 spectroscopically confirmed clusters $(0.05 \le z \le 1.5)$ within an area of 6 deg2 enclosed in the XMM-LSS survey. Almost two thirds have been confirmed with dedicated spectroscopy only and 10% have been confirmed with dedicated spectroscopy supplemented by literature redshifts.

Methods. Sub-samples, or classes, of extended-sources are defined in a two-dimensional X-ray parameter space allowing for various degrees of completeness and contamination. We describe the procedure developed to assess the reality of these cluster candidates using the CFHTLS photometric data and spectroscopic information from our own follow-up campaigns.

Results. Most of these objects are low mass clusters, hence constituting a still poorly studied population. In a second step, we quantify correlations between the optical properties such as richness or velocity dispersion and the cluster X-ray luminosities. We examine the relation of the clusters to the cosmic web. Finally, we review peculiar structures in the surveyed area like very distant clusters and fossil groups.

Key words. Surveys; Galaxies: clusters: general; Cosmology: large-scale structure of Universe.

Send offprint requests to: C. Adami e-mail: christophe.adami@oamp.fr

of France, and the University of Hawaii. This work is based in part on data products produced at TERAPIX and the Canadian Astronomy Data Centre as part of the Canada-France-Hawaii Telescope Legacy Survey, a collaborative project of NRC and CNRS. This work is also based on observations collected at TNG (La Palma, Spain), Magellan (Chile), and at ESO Telescopes at the La Silla and

^{*} Based on observations obtained with MegaPrime/MegaCam, a joint project of CFHT and CEA/DAPNIA, at the Canada-France-Hawaii Telescope (CFHT) which is operated by the National Research Council (NRC) of Canada, the Institut National des Science de l'Univers of the Centre National de la Recherche Scientifique (CNRS)

1. Introduction

With the quest for the characterisation of the Dark Energy properties and the upcoming increasingly large instruments (JWST, ALMA, LSST, EUCLID, etc. ...) the beginning of the 21st century is to be an exciting time for cosmology. In this respect, a new era was already open for X-ray astronomy by the XMM-Newton and Chandra observatories in 1999. The increasing amount of high quality multi-wavelength observations along with the concept of "multi-probe" approach is expected to provide strong constraints on the cosmological models. In this context, X-ray surveys have an important role to play, as it was already the case in the 80s and 90s (e.g. Romer et al. 1994, Castander et al. 1995, Collins et al. 1997, Henry et al. 1997, Bohringer et al. 1998, Ebeling et al. 1998, Jones et al. 1998, Rosati et al. 1998, Vikhlinin et al. 1998, De Grandi et al. 1999, Romer et al. 2000, and ref. therein). New cluster surveys are presently coming to birth (e.g. Romer et al. 2001, Pierre et al. 2004, Finoguenov et al 2007).

One of them, the XMM-LSS survey, covers $11 \ deg^2$ at a sensitivity of $\sim 10^{-14} \ erg/s/cm^2$ at 0.5-2keV for spatially-extended X-ray sources and is currently the largest contiguous deep XMM cluster survey. This sky region is covered by parallel surveys in multiple complementary wavebands ranging from radio to the γ -ray wavelengths (Pierre et al., 2004) and therefore constitutes a unique area for pioneering studies. It can for instance detect a Coma-like cluster at $z\sim2$. A number of articles describing the properties of the XMM-LSS source population have been published by e.g. Pierre et al (2006) and Pacaud et al (2007) for clusters of galaxies and Gandhi et al (2006) for AGNs; the complete X-ray source catalog along with optical identifications for the first 5 deg² of the survey was published by Pierre et al (2007).

One of the major goals of the XMM-LSS survey is to provide samples of galaxy clusters with well defined selection criteria, in order to enable cosmological studies out to redshift $z \sim 1.5$. Indeed, monitoring selection effects is mandatory not only to study the evolution of the cluster X-ray luminosity (i.e. mass) function or of the 3-D cluster distribution but also, as shown by Pacaud et al. (2007), to characterise the evolution of the cluster scaling laws such as the luminosity-temperature relation. We have put special emphasis on the X-ray selection criteria in the XMM-LSS survey. The procedure enables the construction of samples having various degrees of completeness and allows for given rates of contamination by non cluster sources. The subsequent optical spectroscopic observations constitute the ultimate assessment of the clusters, thus operates the purification of the samples.

In a first paper, Pacaud et al (2007) presented the Class One (C1) clusters pertaining to the first 5 deg² of the survey (the ones with the highest apriori probability to be real clusters). The C1 selection yields a purely X-ray selected cluster sample with an extremely low contamination level and corresponds to rather high surface brightness objects. The present article summarises these former findings including now the

Paranal Observatories under programmes ID 072.A-0312, 074.A-0476, 076.A-0509, 070.A-0283, 072.A-0104, and 074.A-0360.

clusters selected from less stringent X-ray criteria (C2 and C3) and including the contiguous Subaru Deep Survey (SXDS, e.g. Ueda et al. 2008). The C2 and C3 objects presented here come from an initial sample with a higher degree of contamination, but have all passed the final spectroscopic tests. Compared to the C1 clusters, they are fainter and correspond a-priori to less massive clusters or to groups at a redshift of ~0.5: this is a population that is for the first time systematically unveiled by the XMM-LSS survey. A few massive very distant clusters are falling into this category too.

The present study is the first attempt to give a comprehensive census (X-ray and optical properties) of the low-mass cluster population within the 0 < z < 1 range. Search for correlations between optical and X-ray properties has already been a long story from, e.g, Smith et al. (1979) or Quintana & Melnick (1982). However, with more than 60 spectroscopically confirmed clusters, the current sample constitutes, by far, the spectroscopically confirmed cluster sample with the highest surface density ever published. The article is organised as follows. Next section describes the X-ray cluster selection. Section 3 presents the available optical photometric and spectroscopic data. Section 4 explains the adopted cluster validation procedure, the new X-ray luminosity computations, and presents the resulting catalog. Then, the global properties of each cluster class and category are examined in Section 5 and, subsequently, the properties of the cluster galaxy population in Section 6. Section 7 details the z=1.53 cluster and investigates possible peculiar structures in the survey. Finally Section 8 gathers the conclusions. The two appendixes discuss the accuracy of photometric redshifts in the context of dense environments and lists additional redshift structures found in the course of the study.

Throughout the paper we assume $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.27$, and $\Omega_{\Lambda} = 0.73$ (Dunkley et al 2009). All magnitudes are in the AB system.

2. The initial cluster candidate selection

The clusters presented in this paper are for the great majority X-ray selected. The XMM-LSS pipeline (Pacaud et al 2006) provides for each detected source some 20 parameters (coordinates, count rate, etc..). Out of these, two are especially relevant for the characterisation of extended sources: the extent measurement (EXT) and the likelihood of extent (EXT_LH). We recall (as defined by Pacaud et al 2006) that the "extent" parameter is the core radius of the beta-profile fit by the survey pipeline to each source, assuming a fixed beta of 2/3. The cluster selection basically operates in this two-dimensional space and has been extensively adjusted and tested using simulations of hundreds of XMM images. This allows the definition of three cluster samples.

- The C1 class is defined such that ~ no point sources are misclassified as extended (i.e. less than 1% of the cluster candidates are point sources) and is described by EXT > 5", EXT_LH > 33 plus an additional boundary on the detection likelihood, set to be greater than 32.
- The C2 class is limited by EXT > 5", EXT_LH > 15 and displays an a-priori contamination rate of about 50%.

- The C3 clusters are faint objects and thus, have less-well characterized X-ray properties. They may be located at the very edge of the XMM field of view or suffer contamination by point sources. They therefore result from a subjective selection mostly based on a visual inspection of the X-ray and optical data; their selection function is up to now undefined.

More details about the classification can be found in Pacaud et al (2006) and Pierre et al (2006).

In this paper, we have presented a large sample of X-ray clusters, including the 29 C1 confirmed clusters published by Pacaud et al. (2007). These C1 clusters were already unambiguously confirmed, but we take the occasion of this publication to reprocess the associated optical spectroscopic data following the standard method developed in the present paper. This will provide a unique homogeneous cluster sample. The clusters pertaining to this paper are, for most of them, located in the first 5 deg² of the XMM-LSS region, supplemented by the Subaru Deep Survey. The validated C1, C2, and C3 samples are presented in Tables 2, 3, and 4. In these Tables, XLSS catalog names refer to sources published in Pierre et al. (2007). XLSSU catalog names refer to sources whose fields were not yet considered in XLSS (for example flagged bad or in SXDS fields) and reobserved (or reprocessed later), or which were below the detection likelihood threshold in the input data set used as source for XLSS.

In the course of the data inspection, we have also identified a few clusters using optically based criteria such as the red sequence or the gapper method. Our spectroscopic data set allowed us to confirm them as bona fide clusters, although these objects are not detected in the X-rays by the current version of the XMM-LSS pipeline or the association between X-ray detected sources and optical clusters is not straightforward. We thought of interest to publish these objects and they are listed in Tables 5 and B.1.

We now describe the involved optical data and the general identification processes.

3. The optical data

3.1. The optical spectroscopic data

We have been performing a dedicated spectroscopic follow-up of all C1 clusters and of a number of C2 and C3 clusters. These PI observations are listed in Table 1 and provide about 2000 redshifts to date. We supplemented this data set with the VVDS deep (e.g. LeFèvre et al. 2005: ~11000 redshifts in 0.49 deg²) and ultradeep (LeFèvre et al. in preparation) data, and with a redshift compilation pertaining to the Subaru Deep Survey (Ueda et al 2008) included in the XMM-LSS area. Some 200 other redshifts were also available from NED for part of the area. We show in Fig. 1 the location of these different surveys, as well as the exposure time of the different XMM fields.

Individual redshift measurements of spectra resulting from the PI data were made following a procedure similar to that adopted by the VVDS survey. Each spectrum was independently measured by several people and the redshift subsequently validated by a moderator. Quality flags were assigned




Fig. 1. Map showing the different involved surveys. The grey level disks are the 11' central areas of the XMM pointings (exposure time depends on the greyness). Large squares show the spectroscopic VVDS-deep and Subaru Deep surveys, and the CFHTLS D1 field. C1, C2, and C3 clusters are also shown. Above a declination of -3.6deg, only g',r',z' coverage is available, hence no photometric redshifts are derived for this zone.

Table 1. PI spectroscopic runs involved in the present paper.

Telescope	Instrument	Year	Nights	Run ID
Magellan	LDSS2	2002	2	-
Magellan	LDSS2	2003	4	-
NTT	EMMI	2003	3	72.A-0312
NTT	EMMI	2004	4	74.A-0476
NTT	EMMI	2005	3	76.A-0509
TNG	DOLORES	2007	4	AOT16/CAT_75
VLT	FORS2	2002	3	70.A-0283
VLT	FORS2	2003	4	72.A-0104
VLT	FORS2	2004	4.5	74.A-0360

to each measurement following the VVDS rules: flag 0 indicates an inconclusive result, flag 1 means a probability of 50% that the assigned redshift is wrong, flag 2 means a probability of 25%, flag 3 means a probability of 5%, flag 4 means a probability of 1%, and flag 9 means we have assigned a redshift with a single line using absent lines in order to limit the possibilities. These percentage levels proved to be reliable in the VVDS survey (LeFèvre et al. 2005).

Our spectroscopic redshifts having quite heterogeneous origins (different telescopes, instrumentations, and resolutions), it is therefore useful to compute the ability to measure a redshift and the achieved velocity resolution. In order to achieve such a goal, we chose to compare the PI data to the VVDS survey, which provides a well qualified set of data. Only 26 galaxies

both measured by the VVDS and our dedicated follow-up have a quality flag greater or equal than 2. For these objects, given the VVDS quality flags (6 flags 2, 6 flags 3, and 14 flags 4), we expect to have 3.2 wrong redshifts. We indeed find 3 redshifts differing by more than 0.05 between the PI and VVDS data. VVDS spectroscopic redshifts are expected to have a typical uncertainty of 280 km/s (from repeated VVDS redshift measurements, Le Fèvre et al. 2005). Excluding all redshifts with differences greater than 0.02, we find a typical uncertainty between PI and VVDS redshifts of 340 km/s. Even with a comparison done on a rather limited size sample, the PI redshifts appear thus reliable in the [0.,1.] redshift range and in the [18,23] I VVDS magnitude range.

Finally, it has to be mentioned that, for the spectroscopic sample, no completeness, neither spatial nor in luminosity, can be globally defined because of the various data origins.

3.2. The optical photometric data

Most of the XMM-LSS area is covered by the Canada-France-Hawaii Telescope Legacy Wide Survey (CFHTLS-Wide¹). This survey, performed by means of the MegaCam camera, covers some 171 deg² in 4 independent patches with five filters (u*, g', r', i' and z'). Resulting catalogs are 80% complete down to i'_{AB} =24. The Wide survey encloses a sample of about 20×10^6 galaxies inside a volume size of ~ 1 Gpc³, with a median redshift of $z\sim0.92$ (Coupon et al 2009). Northern of Dec= -3.6 deg, the CFHTLS data were complemented by PI MegaCam observations (3 deg²) performed in g', r', z' at the same depth as the CFHTLS; they were reduced following the same procedure.

The optical images and catalogs were primarily used to check for the presence of galaxy concentrations coinciding with the extended X-ray emission. The CFHTLS data (only the T0004 release was available at the beginning of the present study) enabled the determination of photometric redshifts in the best fit template (Coupon et al. 2009). These photometric redshifts cover 35 deg^2 in the T0004 partially overlapping with the XMM-LSS area. They were computed using a template-fitting method, calibrated with public spectroscopic catalogs. The method includes correction of magnitude systematic offsets. The achieved photometric redshift precision $\sigma_z/(1+z)$ is of the order of 0.04 with a catastrophic error percentage of less than 5% at $i^* \le 23$ (the magnitude limit we adopted for the photometric redshifts).

4. "Cluster candidate" validation process

4.1. General method

Extragalactic extended X-ray emission is the signature of a deep gravitational potential well. Apart from the hypothetical "dark clusters", this potential well coincides with a galaxy over-density. The system (cluster or group) is therefore detectable using optical information only. In this article, we aim at assessing the presence of optical structures corresponding

to the X-ray cluster candidates. Such systems are expected to manifest themselves as compact structures in redshift space (both spectroscopic and photometric ones) and as localized excess in projected galaxy density maps.

To perform such an analysis, we make use of the two optical data sets mentioned above. The investigated lines of sight (centered on the X-ray emissions) were initially selected from the condition that at least two spectroscopic redshift measurements (whatever their values) are available within the X-ray isophotes. The subsequent conditions were more stringent depending on the cluster nature (see below).

The CFHTLS Wide survey and subsequent analyses (e.g. Coupon et al. 2009) provide us with galaxy positions as well as their apparent and absolute magnitudes, photometric redshifts and the corresponding "galaxy types" T (from the spectral fitting performed during the photometric redshift computation). With the exception of the usual "masking problems" due to bright stars or CCD defaults, photometric data are homogeneous and allow us to define complete sub-samples in terms of spatial extension or in magnitudes. Limitations to these data are the redshift range within which photometric redshifts are reliable and the adopted magnitude limit. Here we restrict ourselves to 0.2 < z < 1.2 and i' = 23 (see Coupon et al. 2009). This limiting magnitude will affect partly the use of photometric data to detect structures. Indeed, the characteristic magnitude m* of the Schechter luminosity function is about i' = 20at z = 0.5 and i' = 22.5 at z = 1 leading to sampled luminosity function ranges of about m^*+3 to $m^*+0.5$ at these respective redshifts. One drawback is therefore that for z > 0.5 the number of galaxies actually belonging to a structure will be rapidly overcome by the background contamination (see e.g Table 1 of Adami et al 2005). One way to fight this contamination will be to use redshift slices defined on a photometric redshift basis (see Mazure et al. 2007) but the range covered in magnitude by structure members will remain limited.

In order not to bias the optical characterisation of the X-ray sources, the information concerning the C1, C2, C3 classification was used only at the very final stage.

4.2. Different analysis steps

The first step concerns the expected compactness in spectroscopic redshift space. To reveal such compact associations, we used the already well tested and used "gap method" (e.g. Biviano et al 1997, Rizzo et al 2004). It looks for significant gaps between successive galaxy velocities within the ordered redshift distribution obtained along a given line of sight. As in Adami et al (2005), we use a gap defined by $g = 600 \, (1+z) \, \text{km/s}$ which was optimum for the considered redshift range. When the velocity difference between 2 successive galaxies is smaller than g, they are assigned to belong to a common structure, otherwise they are put in different groups.

Since the lines of sight most of the time sample redshifts up to at least z=1, this first step of the analysis ends in general with several groups. Thus, with the mean redshift of every group, a cosmological distance was assigned, a physical region of 500 kpc (radius) defined and the galaxies within this radius

¹ http://terapix.iap.fr/cplt/T0006/T0006-doc.pdf

are selected as potential real cluster members. We choose this size as being representative of clusters in terms of membership of galaxies w.r.t the field. Taking larger regions would decrease any real contrast, while taking smaller regions would decrease the number of true members. As a second step we then apply the usual ROSTAT tools (Beers et al. 1990) on individual redshift groups to test for final membership and definition of the group properties (robust redshift locations and scales with their corresponding bootstrap errors).

As already mentioned, several groups are in general identified along the lines of sight. Before comparing the galaxy distribution and the X-ray isophotes, we used then, when available, the CFHTLS photometric redshift information. As a third step, we selected galaxies in photometric redshift slices (of width: \pm 0.04 (1+z), see Coupon et al. 2009) around the mean redshift of the considered group and produced iso-contours of numerical galaxy density (see Mazure et al 2007 for details and previous application). It is expected that the optical group physically associated with the X-ray emission will show up with a clear density contrast located next to the position of the X-ray center. This is because the use of photometric redshift slices removes a large part of the fore and background contaminations. We also look, as another check, at the photometric redshift distribution within various central regions compared to the one in the largest available region, conveniently renormalized and defined as the "field". Again, one expects a clear contrast at the redshift values given by the spectroscopy.

An illustration is given with the source XLSSC 013 in the XMM-LSS database. Three main groups were identified along the line of sight (z ~ 0.2 with 9 redshifts, z ~ 0.3 with 26 redshifts, z ~ 0.6 with 5 redshifts). Consecutive examination of both the photometric redshift distribution and the numerical density histograms gives strong evidence for the z ~ 0.3 group to be chosen (see Figs. 2 and 3).

However, as mentioned above, photometric redshift data were not always available and spectroscopic data could be very sparse (our velocity dispersion measurements are then subject to very complex selection functions in the target selection when measuring and collecting galaxy redshifts). The final selection is then done by a visual inspection of X-ray and optical maps taking into account all the informations available. Fig. 3 shows the group at z=0.3 chosen for XLSSC 013. As an extreme contrary case, we show in Fig. 4 XLSSC 035 for which only a few redshifts were available. The fact that a giant galaxy at z=0.069 lies at the center finally pleads in favor of that redshift (Fig. 4) in the present paper. We note however that a $z\sim0.17$ galaxy layer is also detected along this line of sight and we could deal with a superposition effect.

4.3. Results

We examined 34 C1 candidate X-ray sources. Identification fails for only two lines of sight mainly because very few redshifts were available in the X-ray region and/or no photometric redshifts. All identified sources were classied as galaxy clusters; this means that at least 95% of the C1 objects are real clusters (when obvious nearby galaxies - which show also a diffuse




Fig. 2. Isodensity maps of the numerical density of galaxies within photometric slices of width \pm 0.04 (1+z) around the group redshifts. From top to bottom: $z \sim 0.2$, $z \sim 0.3$, $z \sim 0.6$. The best agreement with the X-ray emission of XLSSC 013 is obtained at z = 0.3. Large red circles are the same as in Fig. 3.

Fig. 3. XMM-LSS X-ray contours for system XLSSC 013 with cluster member galaxies with a measured redshift (between z=0.3049 and 0.3112) superimposed. The red circle corresponds to a radius of 500 kpc at z = 0.3.

Fig. 4. XMM-LSS X-ray contours for system XLSSC 035 with galaxies with measured redshifts superimposed. The red circle corresponds to a radius of 500 kpc at z = 0.069.

X-ray emission - are excluded). Among the C2 and C3 candidates, only those having 2 redshifts within the X-ray isophotes were selected for the present analysis. As our current spectroscopic data set is heterogeneous and does not provide a systematic targeting of all C2 and C3 cluster candidates, it is not possible to draw firm conclusions about the effective contamination rate (in terms of non-cluster sources) for these populations. We may only state that for all C2 (resp. C3) sources having yet at least 2 spectroscopic redshifts within the X-ray isophotes, more than 80% (resp. 50%) of the examined sources turned out to be real clusters.

An additional potential X-ray source was also discovered (C555 in Table 4). Not listed in Pierre et al. (2007), this source is merged with XLSSU J022533.8-042540. We detected a very clear associated galaxy structure in optical. A manual extraction of the X-ray source gives a count rate of 0.003±0.001 counts per second ([0.5-2keV]).

For seven of the analysed lines of sight, the association between X-ray source and optical galaxy concentration was not obvious or the X-ray source was not significantly different from the background. However, these clusters are identified on the basis of the color magnitude relation (for 2 of them) or are detected as significant galaxy overdensities in Adami et al. (2010) using photometric redshifts during the analysis. All these objects have been classified as C0 clusters.

C1, C2, C3, and C0 clusters are presented in Tables 2, 3, 4, and 5. Almost two thirds have been confirmed with dedicated spectroscopy only and 10% have been confirmed with dedicated spectroscopy supplemented by literature redshifts.

We compared the cluster redshifts listed in the present paper (see also next section) with the estimates already published within the XMM-LSS framework (from Pacaud et al. 2007 and Bremer et al. 2006: 29 C1 clusters and 1 C2 cluster), and we found the expected good agreement. This is not surprising as Pacaud et al. (2007) and Bremer et al. (2006) are included in the presently used spectroscopic redshift sample. However, redshift measurements have been redone on a more homogeneous basis and sometimes with new data. The difference is only 0.00075±0.00329 when excluding XLSSC 035. For this cluster, we detected a possible error in the individual redshifts measurement process. The central galaxy seems to be at z=0.069 and not 0.17 as stated in Pacaud et al. (2007: cluster redshift changed to z=0.069). We are in the process of acquiring more data in order to definitively solve this case. We also note that the central galaxies of XLSSC 028 are also at z~0.3 and not at z~0.08 as stated in Pacaud et al. (2007: cluster redshift unchanged at $z\sim0.3$).




Fig. 5. Previous and present cluster flux (in a 0.5Mpc radius) comparisons.

The agreement is very good for the [0.5-2 keV] fluxes measured in a 500kpc radius (Fig. 5).

For the Subaru Deep Survey region, we compared our detections with the extended X-ray source catalog of Ueda et al. (2008) and with the structure catalog of Finoguenov et al. (2010). Nine of our X-ray clusters are inside the area covered by these catalogs and six are also detected by these authors. Redshifts are always in very good agreement. Finoguenov et al. (2010) list in their paper 57 structures inside this area. However, their selection function (completeness/contamination) for the X-ray extended sources as well as the characteristics of these sources (extent-measurement along with error or likelihood) are not published, thus preventing any meaningful comparison between the two samples. Moreover, as shown by Pacaud et al (2006) a flux limit, unless it is set very high, cannot define a complete uncontaminated sample of extended sources.

We finally performed a comparison with independently optically detected clusters in the literature. Limiting ourselves to studies giving a galaxy velocity dispersion estimate, we have five detections in common with Hamana et al. (2009: see Table 2). All these clusters are C1 structures. Redshifts are always in good agreement. Galaxy velocity dispersions are also consistent within error bars with an exception for XLSSC 050 where we find 408±96 km/s and where Hamana et al. (2009) find 739⁺¹⁵⁰₋₈₆km/s. This structure being very complex, the galaxy velocity dispersion is however very dependent on the selected galaxies and on the exact center choice.

4.4. Updated X-ray luminosities

We apply the principle of "aperture photometry" to the flux measurement of the X-ray clusters, which avoids any other assumption than spherical symmetry as to the cluster shape. We note that Pacaud et al. (2007) used a beta-model fitting, which is not possible for the larger sample presented here, that comprises faint objects. For these C2 and C3 objects, having sometimes at most some hundred counts, it is not possible to perform a semi-interactive spatial fit as in Pacaud et al (2007), i.e. let-




Fig. 6. Redshift distribution for the 3 classes C1 (black histogram), C2 (red histogram), C3 (blue histogram).

ting the core radius and the beta value as free parameters. The resulting uncertainty would be very large.

We integrate the count rate in concentric annuli and derive the uncertainties by using Poisson statistic. Then, considering the count rate in each annulus, we stop the integration at the radius of the annulus for which the corresponding count-rate increase is comparable to the background 1-sigma fluctuation. This program operates in semi-interactive mode leaving the possibility to optimize the determination of the X-ray centroid and of the background level. The measurement yields for each cluster the total MOS1 + MOS2 + PN count-rate within a radius 500 kpc. The fluxes have been obtained assuming a fixed conversion factor into the [0.5-2] keV band using a constant conversion factor of 9×10^{-13} [(ergs/cm2/s)/(cnts/s)]. This value was calculated using Xspec from an APEC emission model with the following parameters: z=0.5, T=2 keV, $Nh=2.6\times10^{20}$ cm⁻², Ab=0.3. Bolometric luminosities (also within a 500kpc radius) listed in the tables were also calculated with Xspec from the measured fluxes using the Pacaud et al. (2007) and the Bremer et al. (2006) temperatures when available. We used redshifts described in the next section. For clusters not listed in these papers (probably low mass structures), we used T=1.5keV.

5. Global properties of the various classes

We will consider from this section to the end of the paper only clusters successfully identified.

5.1. Rich and poor structures

For X-ray sources unambiguously identified with optical velocity structures, one has then to address the question: has the C1, C2, C3 classification a physical basis, or is it only reflecting the X-ray selection process?

As a first step, we look at the redshift distribution of the cluster C1, C2, and C3 classes (Fig. 6). For the 32 C1, the 9 C2, and the 17 C3 the mean redshift is 0.41, 0.66, and 0.38.

Comparing the C1 and C3 distributions and their almost similar mean redshifts and letting aside for a while the $z \ge 0.5$




Fig. 7. Redshift distribution for the *most luminous* (Most 1.), *luminous* (L.), *moderately luminous* (Mod. 1.), and C0 clusters.

C3 structures, it is tempting to consider C1 to be in the most cases "X-ray bright and optical nearby ($z \le 0.4$) rich systems" and most of the C3 as "faint and poor" at $z \sim 0.4$ redshift. The more distant C3 clusters would be rather distant C1-like and therefore "rich". C2 clusters would be a mix of nearby poor and distant rich clusters.

We can define alternative categories to the C1, C2, C3 classification. For instance, we chose to group the clusters as a function of their X-ray luminosity. Clusters more luminous than 10^{44} erg/s were called the X-ray *most luminous* sample. Clusters between 10^{43} and 10^{44} erg/s were called the X-ray *luminous* sample. Clusters below 10^{43} erg/s were called the X-ray *moderately luminous* sample. Finally, clusters without any X-ray detection (C0 clusters) were considered separately. We give in Fig. 7 the redshift distribution of these 4 categories. As expected because of the relatively small angular coverage of the XMM-LSS survey, the *most luminous* clusters are mainly distant objects. Similarly, *moderately luminous* clusters are quite nearby objects because our X-ray selection function does not allow us to detect them when they are distant, according to the well known Malmquist bias.

We show in Fig. 8 a synthethic view of the clusters listed in Tables 2, 3, and 4 allowing the reader to visualize the different classes (C1, C2, C3, *most luminous*, *luminous*, and *moderately luminous*) in a redshift versus X-ray luminosity diagram.

5.2. Optical richness

We know (e.g. Edge & Stewart 1991) that optical and X-ray cluster properties should be relatively well correlated. It is then necessary to characterize the optical richness (N_{Rich}) of our clusters. This is done by taking first the number of galaxies in the region of 500 kpc (radius), within the photometric redshift slice zmean \pm 0.04 (1+z) and with magnitude less than $m^* + 3$. That number is then corrected by the "field contribution" estimated in the same manner within 1 Mpc to give the final estimate. This richness value is probably not accurate enough in terms of absolute value, but can be used in a relative way when comparing a structure to another one. We also note that, given




Fig. 8. Present paper cluster distribution in a $log 10(L_X)$ versus redshift diagram. The two vertical green lines separate the *most luminous*, *luminous*, and *moderately luminous* clusters. Black disks are C1 clusters, red disks are C2 clusters, blue disks are C3 clusters. We also show as black, red, and blue curves the detection limit of the lowest X-ray flux cluster in C1, C2, and C3 classes.

the CFHTLS wide magnitude limit we adopted (i'=23), only $z\le0.5$ clusters are sampled deeply enough to reach $m^* + 3$. We therefore only considered these clusters in order to avoid to have biased optical richnesses.

Fitting a richness-velocity dispersion for all $z \le 0.5$ structures for which both data were available, we get: $\log(\sigma) = (0.45 \pm 0.24) \log(N_{Rich}) + (1.96 \pm 0.38)$.

This is compatible, within the uncertainties, with the value of Yee and Ellingson (2003), for similar kind of data:

$$\log(\sigma) = (0.55 \pm 0.09) \log(B_{cg}) + (1.26 \pm 0.30).$$

We now test richness and velocity dispersions versus X-ray properties. We first consider $z\le0.5$ structures with known X-ray luminosity and optical richness. We selected only C1, C2 and C3 clusters with X-ray luminosity at least two times larger than the associated uncertainty. We show in Fig. 9 the possible relation between the logarithm of N_{Rich} and of Lx. The linear regression between the two parameters has a slope of 0.84 ± 0.51 . We note that this value only appears poorly significantly different from a null slope.

There is a single clear interloper: XLSSC 006 at $z\sim0.43$ (outside of the box shown in Fig. 9). This is one of the most massive clusters in our sample. The observed spectra in the cluster center do not show any sign of AGN activity, so we have no reason to believe that the X-ray flux is polluted by a point source. This cluster shows signs of major substructures in the velocity distribution and this may explain its relatively high Lx value compared to its optical richness. Resulting compression in the intracluster medium could increase the gas density resulting in an enhanced X-ray luminosity.

Considering now clusters at z \leq 0.5 with a known X-ray temperature (from Pacaud et al. 2007) and a measured galaxy velocity dispersion, we searched for a relation between N_{Rich} , velocity dispersion, and X-ray temperature. Fig. 10 shows the relation between $\log(N_{Rich}\sigma^2)$ and $\log(T_X)$. We expect a linear relation as $(N_{Rich}\sigma^2)$ is at least a qualitative measurement of




Fig. 9. $\log(N_{Rich})$ versus $\log(L_X)$. Crosses are clusters at signal to noise lower than 2 regarding the X-ray luminosity. Disks are clusters at signal to noise greater than 2 and at z \leq 0.5 (black: C1, red: C2, blue: C3). They give the following fit: $\log(N_{Rich}) = (0.84 \pm 0.51) \log(L_X) + (7.2 \pm 0.81)$.

the kinetic energy of the clusters, therefore close to the X-ray temperature. Error bars on $(N_{Rich}\sigma^2)$ are 68% uncertainties and are computed assuming a perfect knowledge of the richness and the error bars on σ^2 given in Tables 2, 3, and 4. As quoted in Table 2, these uncertainties are computed with a bootstrap technique.

We have two outliers: XLSSC 027 and XLSSC 018. XLSSC 027 is known to have strong discrepancies between galaxy and weak lensing equivalent velocity dispersions $(898^{+523}_{-527} \text{km/s})$ from Gavazzi & Soucail (2007) against 323 ± 78 km/s for our own galaxy velocity dispersion and $447^{+82}_{-52} \text{km/s}$ for the Hamana et al. (2009) galaxy velocity dispersion). We note that using the weak lensing equivalent velocity dispersion puts XLSSC 027 close to the best fit relation. We also note that this cluster has close contaminants at z=0.31 and 0.38 detected along the line of sight. This could also affect the measurement of the optical richness via the background estimate.

XLSSC 018 (without any sign of major substructures: see below) would need a larger optical richess and/or a larger galaxy velocity dispersion, or a lower X-ray temperature to fall on the best fit relation. The last solution in unlikely as only an X-ray temperature of the order of 0.4 keV would place XLSSC 018 on the best fit relation. A possible explanation would be that we are dealing with a structure close to a fossil group (even if it does not satisfy the characteristics of this class of structures). A significant part of the cluster member galaxies could have merged with the central galaxy, then depopulating the $\leq m^* + 3$ magnitude range and diminishing the measured optical richness.

In conclusion, and despite a few detected interlopers, the global agreements show the statistical reliability of our optical richness and galaxy velocity dispersion estimates.

Fig. 11 shows the histograms of the richness for the 3 classes. C1 has a mean N_{Rich} of 45, C2 a mean N_{Rich} of 37 and C3 a median N_{Rich} of 35.




Fig. 10. $\log(N_{Rich}\sigma^2)$ versus $\log(T_X)$ for the z≥0.5 clusters (all C1 but XLSSC 046 which is C2). The + sign indicates XLSSC 027 and translates to the red disk when replacing the galaxy velocity dispersion by the weak lensing estimate from Gavazzi & Soucail (2007). The continuous line is the fit (computed without XLSSC 027 and XLSSC 018): $\log(N_{Rich}\sigma^2) = (1.05 \pm 0.13) \log(T_X) + (7.05 \pm 0.07)$.

5.3. Substructure level in velocity space

Our spectroscopic catalogs are generally too sparse to allow precise substructure analyses. However, limiting ourselves to the confirmed clusters with available CFHTLS data and with more than 9 redshifts in the structure (10 clusters: XLSSC 013, XLSSC 025, XLSSC 022, XLSSC 006, XLSSC 008, XLSSC 001, XLSSC 000, XLSSC 018, XLSSC 044, and XLSSC 058), we applied the Serna-Gerbal method (Serna & Gerbal, 1996: SG hereafter) to these spectroscopic catalogs. Two of them (XLSSC 006 and XLSSC 001) are from the most luminous cluster category. All the others (but XLSSC 000 which is a C0 cluster) are members of the luminous cluster category. The SG method is widely used in order to characterize the substructure level in clusters of galaxies (e.g. Adami et al. 2009). Basically, the method allows galaxy subgroups to be extracted from a catalog containing positions, magnitudes, and redshift, based on the calculation of their relative binding energies. The output of the SG method is a list of galaxies belonging to each group, as well as information on the binding energy, and mass estimate of the galaxy structures.

The spectroscopic catalogs being still relatively sparse, we will only be able to detect very prominent substructures, but this is a good way for example to check if the analysed clusters are in the process of a major merging event.

Over the 10 analysed clusters, only two (which both belong to the *most luminous* cluster category) present sign of substructures (XLSSC 006 with two dominant galaxies in its center and XLSSC 001) with 2 detected sub-groups. We checked if these two clusters were atypically sampled in terms of number of available redshifts. XLSSC 001 has 17 redshifts and XLSSC 006 16 redshifts. Three other clusters without detected signs of substructures are equally sampled: XLSSC 013 has 19 redshifts, XLSSC 022 has 15 redshifts, and XLSSC 044 has 17 redshifts. The substructure detection therefore does not seem to be entirely due to selection effects depending on the

available number of redshifts. As a conclusion, all the tested *most luminous* clusters show signs of substructures while none of the other tested clusters show similar signs. This would be in good agreement with the most massive clusters being regularly fed by their surrounding large scale structure in terms of infalling groups. Less luminous clusters would already be close to their equilibrium, with a less intense infalling activity. This has, however, to be confirmed with larger spectroscopic samples.

5.4. Relation between XMM-LSS clusters and their parent cosmic web portion

The previous subsection naturally raises the question of the characteristics of cosmological surrounding filaments. Numerical simulations place clusters of galaxies at the nodes of the cosmic web. Clusters are then growing via accretion of matter flowing along the cosmic filaments. This unquestionable scenario for massive clusters is less evident for low mass structures as groups. Such groups could also form along the cosmic filaments as for example suggested for fossil groups by Adami et al. (2007a). Moreover, even for the most massive structures, the precise process of filament matter accretion is assessed most of the time only by individual cluster studies (e.g. Boué et al. 2008). The XMM-LSS cluster sample presented in this paper offers a unique opportunity to investigate the cluster-filament connection with a well controled sample.

5.4.1. General counting method

We first have to detect the filaments connected to a given cluster. These filaments are very low mass and young dynamical structures. It is therefore very difficult to detect them through X-ray observations. This is possible only in a few peculiar cases (e.g. Boué et al. 2008, Werner et al. 2008) and with very long integration times. The XMM-LSS exposure times are anyway not well suited to such detections. We, therefore, used optical CFHTLS photometric redshift catalogs.

Table 2. List of the XMM-LSS C1 systems having successfully passed the spectroscopic identification. Name refers to the official IAU name. XLSSC refers to the official XMM-LSS name. PH gives the availability of CFHTLS T0004 photometric redshifts: 0/1= not available/available. RA and DEC are the decimal J2000 coordinates of the X-ray emission center. N is the number of galaxies with spectroscopic redshifts belonging to the identified structure and within a radius of 500 kpc. ZBWT is the biweight estimate of the mean redshift of the structure (at a 0.001 precision). ERRZ is the upper value of the bootstrap uncertainty on ZBWT, at a precision of 0.001. It was computed only when more than 5 redshifts were available. SIG-GAP is the "Gapper" estimate of the velocity dispersion given in km/s. ERR is the bootstrap uncertainty on SIG (computed when >5 redshifts were available). Flux is the value of the [0.5;2] keV flux in 0.5Mpc (radius) in 10^{-14} ergs/cm²/s. Lx is the bolometric (0.1 to 50 keV) X-ray luminosity (in 10^{43} erg/s) derived from the observed flux, N_{NP} is the net number of photons in 0.5 Mpc. The two last columns give name and redshift (when available) when the considered cluster was also detected by Hamana et al. (2009), Ueda et al. (2008), or Finoguenov et al. (2010). When coming from Hamana et al. (2009), the cluster name has the SL Jhhmm.mddmm format. When coming from Ueda et al. (2008), the cluster name refers to this paper (4 digits number). When coming from Finoguenov et al. (2010), the cluster name refers to this paper (with the SXDF root). The * symbols indicate that the cluster validation was made with one or two spectroscopic redshifts. The (I) attached to the cluster id means that we have a lack of precision in the measured galaxy redshifts, preventing us to compute uncertainty of the mean cluster redshift, and velocity dispersions.

Name	XLSSC	PH	RA	DEC	N	ZBWT	ERRZ	SIG	ERR	Flux [0.5;2]keV	L_{Bol}	N _{NP}	Lit Id	Lit z
			deg	deg				km/s	km/s	10 ⁻¹⁴ ergs/cm ² /s	10 ⁴³ erg/s			
										in 0.5Mpc	in 0.5Mpc	in 0.5Mpc		
XLSSU J021735.2-051325	059	1	34.391	-5.223	8	0.645	0.001	513	151	1.6±0.3	6.0±1.1	104±19	SXDF69XGG/0514	0.645
XLSSU J021945.1-045329	058	1	34.938	-4.891	9	0.333	0.001	587	236	2.0 ± 0.2	1.5 ± 0.1	601 ± 54	SXDF36XGG/1176	0.333
XLSS J022023.5-025027	039	0	35.098	-2.841	4	0.231				2.2 ± 0.4	0.7 ± 0.1	183 ± 34		
XLSS J022045.4-032558	023	0	35.189	-3.433	3	0.328				4.1 ± 0.4	3.1 ± 0.3	465±43		
XLSS J022145.2-034617	006	1	35.438	-3.772	16	0.429	0.001	977	157	24.1 ± 0.8	46.0 ± 1.6	1409 ± 52		
XLSS J022205.5-043247	040*	1	35.523	-4.546	2	0.317				2.0 ± 0.3	1.4 ± 0.2	265 ± 35		
XLSS J022206.7-030314	036*	0	35.527	-3.054	2	0.494				10.2 ± 0.5	23.8 ± 1.3	551±32		
XLSS J022210.7-024048	047	0	35.544	-2.680	14	0.790	0.001	765	163	1.4 ± 0.3	9.7 ± 1.9	92±18		
XLSS J022253.6-032828	048*	0	35.722	-3.474	2	1.005				1.1 ± 0.2	13.1 ± 2.9	81±18		
XLSS J022348.1-025131	035*	0	35.950	-2.858	1	0.069				6.1 ± 0.6	0.2 ± 0.1	478 ± 48		
XLSS J022356.5-030558	028	0	35.985	-3.100	8	0.296	0.001	281	46	2.0 ± 0.4	1.1 ± 0.2	177±31		
XLSS J022357.4-043517	049*	1	35.989	-4.588	1	0.494				1.9 ± 0.2	3.9 ± 0.5	287 ± 34		
XLSS J022402.0-050525	018	1	36.008	-5.090	9	0.324	0.001	364	69	1.7 ± 0.2	1.3 ± 0.1	382 ± 44		
XLSS J022404.1-041330	029(1)	1	36.017	-4.225	5	1.050				3.1 ± 0.2	43.7 ± 3.1	323 ± 24		
XLSS J022433.8-041405	044	1	36.141	-4.234	17	0.262	0.001	483	100	2.6 ± 0.3	1.1 ± 0.1	510 ± 54	SL J0224.5-0414	0.2627
XLSS J022456.2-050802	021	1	36.234	-5.134	7	0.085	0.001	231	64	4.2 ± 0.7	0.1 ± 0.1	664±115		
XLSS J022457.1-034856	001	1	36.238	-3.816	17	0.614	0.001	940	141	7.8 ± 0.4	29.3±1.6	671±36		
XLSS J022520.8-034805	800	1	36.337	-3.801	11	0.299	0.001	544	124	1.8 ± 0.4	1.0 ± 0.2	196±39		
XLSS J022524.8-044043	025	1	36.353	-4.679	10	0.266	0.001	702	178	8.7 ± 0.5	4.2 ± 0.2	1098±62	SL J0225.3-0441	0.2642
XLSS J022530.6-041420	041	1	36.377	-4.239	6	0.140	0.002	899	218	21.8±1.1	2.2 ± 0.1	1143±62	SL J0225.4-0414	0.1415
XLSS J022532.2-035511	002	1	36.384	-3.920	8	0.772	0.001	296	56	2.6 ± 0.3	16.4 ± 1.7	238 ± 25		
XLSSU J022540.7-031123	050	0	36.419	-3.189	13	0.140	0.001	408	96	54.0 ± 1.1	7.8 ± 0.2	4929±103	SL J0225.7-0312	0.1395
XLSS J022559.5-024935	051	0	36.498	-2.826	6	0.279	0.001	369	99	1.1 ± 0.3	0.5 ± 0.1	160 ± 41		
XLSS J022609.9-045805	011	1	36.541	-4.968	8	0.053	0.001	83	16	13.1 ± 1.5	0.2 ± 0.1	1706±192		
XLSS J022616.3-023957	052	0	36.568	-2.665	5	0.056	0.001	194	60	16.4±1.8	0.2 ± 0.1	1297±146		
XLSS J022709.2-041800	005*	1	36.788	-4.300	2	1.053				1.0 ± 0.1	13.4 ± 1.8	165 ± 23		
XLSS J022722.4-032144	010	0	36.843	-3.362	5	0.331	0.001	315	56	5.7 ± 0.4	4.7 ± 0.4	459±38		
XLSS J022726.0-043216	013	1	36.858	-4.538	19	0.307	0.001	397	58	2.6 ± 0.3	1.5 ± 0.2	374 ± 48		
XLSS J022738.3-031758	003	0	36.909	-3.299	9	0.836	0.001	784	189	3.7 ± 0.4	29.0 ± 3.1	196±22		
XLSS J022739.9-045127	022	1	36.916	-4.857	15	0.293	0.001	535	106	9.6±0.3	5.5 ± 0.2	1741±58		
XLSS J022803.4-045103	027	1	37.014	-4.851	6	0.295	0.001	323	78	6.3±0.4	4.2 ± 0.3	653±41	SL J0228.1-0450	0.2948
XLSS J022827.0-042547 /	012	1	37.114	-4.432	5	0.434	0.002	726	95	3.4 ± 0.3	4.7 ± 0.4	444±37		
XLSS J022827.8-042601														

Adami et al.: Optical assessment and comparative study of the C1, C2, and C3 cluster classes

Table 3. Same as Table 2 but for C2 XMM-LSS systems. XLSS J022756.3-043119 would require more spectroscopy for confirmation. XLSSC 009, 064, and 065 were originally classified as C2, but would be classified as C1 using more recent pipeline version. The * symbols indicate that the cluster validation was made with one or two spectroscopic redshifts. The (l) attached to the cluster id means that we have a lack of precision in the measured galaxy redshifts, preventing us to compute uncertainty of the mean cluster redshift, and velocity dispersions.

Name	XLSSC	PH	RA	DEC	N	ZBWT	ERRZ	SIG	ERR	Flux [0.5;2]keV	L_{Bol}	N_{NP}	Lit Id	Lit z
			deg	deg				km/s	km/s	$10^{-14} \text{ ergs/cm}^2/\text{s}$	10 ⁴³ erg/s			
										in 0.5Mpc	in 0.5Mpc	in 0.5Mpc		
XLSSU J021658.9-044904	065	1	34.245	-4.821	3	0.435				1.1±0.4	1.6±0.6	50±18	285/287	
XLSSU J021832.0-050105	064	1	34.633	-5.016	3	0.875				1.6 ± 0.1	13.5 ± 0.8	937±56	SXDF46XGG/829	0.875
XLSSU J021837.0-054028	063*	1	34.654	-5.675	2	0.275				4.1 ± 0.5	2.0 ± 0.2	255 ± 32		
XLSS J022303.3-043621	046	1	35.764	-4.606	8	1.213	0.001	595	121	0.7 ± 0.1	14.0 ± 3.2	82±19		
XLSS J022307.2-041259	030	0	35.780	-4.216	5	0.631	0.001	520	158	0.6 ± 0.1	2.1 ± 0.4	129 ± 27		
XLSS J022405.9-035512	007	1	36.024	-3.920	5	0.559	0.001	369	179	1.2 ± 0.3	3.2 ± 0.7	113±25		
XLSSU J022644.2-034107	009	1	36.686	-3.684	8	0.328	0.001	261	53	1.7 ± 0.4	1.2 ± 0.3	87±21		
XLSS J022725.1-041127	038	1	36.854	-4.191	4	0.584				0.1 ± 0.1	0.4 ± 0.3	31±28		
XLSS J022756.3-043119*	-	1	36.985	-4.522	2	1.050?				0.2 ± 0.1	2.9 ± 1.5	49 ± 26		

Table 4. Same as Table 3 but for C3 XMM-LSS systems. The ** symbols indicate that we merged by hand 2 groups separated by the gapper technique in the final analysis. The last line (C555 class) is the new cluster (see text). The * symbols indicate that the cluster validation was made with one or two spectroscopic redshifts. The (1) attached to the cluster id means that we have a lack of precision in the measured galaxy redshifts, preventing us to compute uncertainty of the mean cluster redshift, and velocity dispersions.

Name	XLSSC	PH	RA	DEC	N	ZBWT	ERRZ	SIG	ERR	Flux [0.5;2]keV	L_{Bol}	N_{NP}	Lit Id	Lit z
			deg	deg				km/s	km/s	$10^{-14} \text{ ergs/cm}^2/\text{s}$	10 ⁴³ erg/s			
										in 0.5Mpc	in 0.5Mpc	in 0.5Mpc		
XLSSU J021651.3-042328*	-	1	34.214	-4.392	1	0.273				0.6 ± 0.3	0.3 ± 0.2	58±32		
XLSSU J021754.6-052655(1)	066	1	34.478	-5.447	6	0.250				0.3 ± 0.3	0.1 ± 0.1	68±59	SXDF85XGG/621	0.25
XLSSU J021842.8-053254	067	1	34.678	-5.548	5	0.380	0.001	847	279	1.0 ± 0.4	1.0 ± 0.4	84±34	SXDF01XGG/876	0.378
XLSSU J021940.3-045103*	-	1	34.919	-4.852	1	0.454				0.8 ± 0.2	1.2 ± 0.3	193±44		
XLSS J022258.4-04070	024	1	35.744	-4.121	5	0.293	0.001	452	98	1.3 ± 0.2	0.7 ± 0.1	254 ± 41		
XLSS J022341.8-043051	026	1	35.925	-4.514	3	0.436				1.6 ± 0.2	2.2 ± 0.3	255 ± 34		
XLSSU J022509.2-043239	037	1	36.286	-4.542	3	0.767				0.1 ± 0.1	0.3 ± 0.7	11±28		
XLSSU J022510.5-040147	043**	1	36.294	-4.029	3	0.170				2.6 ± 0.9	0.4 ± 0.1	142±49		
XLSS J022522.8-042649	042	1	36.345	-4.447	6	0.462	0.003	1009	257	0.6 ± 0.2	0.9 ± 0.3	111±33		
XLSS J022542.2-042434	068	1	36.424	-4.410	4	0.585				0.2 ± 0.2	0.5 ± 0.6	23 ± 28		
XLSS J022610.0-043120	069	1	36.542	-4.523	8	0.824	0.001	398	114	-0.4 ± 0.2	-	-		
XLSSU J022627.3-050001	017	1	36.615	-5.003	5	0.383	0.001	352	132	1.5 ± 0.4	1.6 ± 0.4	134±35		
XLSSU J022632.5-040314	014	1	36.635	-4.054	6	0.345	0.001	304	84	1.4 ± 0.5	1.1 ± 0.4	74 ± 29		
XLSSU J022632.4-050003	020	1	36.638	-5.007	3	0.494				2.0 ± 0.4	3.8 ± 0.8	159 ± 32		
XLSSU J022726.8-045412	070	1	36.860	-4.904	5	0.301	0.001	180	47	0.2 ± 0.2	0.1 ± 0.1	32±39		
XLSS J022754.1-035100*	-	1	36.974	-3.851	2	0.140				0.7 ± 0.5	0.1 ± 0.1	107 ± 72		
XLSSU J022828.9-045939	016*	1	37.121	-4.994	2	0.335				0.4 ± 0.5	0.3 ± 0.4	33±44		
C555	-	1	36.375	-4.429	7	0.921	0.001	759	248	0.3±0.1	2.9±1.1	47±18		




Fig. 11. Distribution of the optical richness for the 3 classes C1 (thin black), C2 (thick red), and C3 (thick blue).

- We first selected only clusters at $z \le 0.5$ in order to be able to sample deeply enough the galaxy population to potentially detect the filaments, given the i'=23 magnitude limit for the photometric redshift catalog as demonstrated in Adami et al. (2010).
- For a given cluster, we selected galaxies with photometric redshifts in a $0.04 \times (1+z)$ slice around the cluster redshift.
- We then computed the number of galaxies in the slice in 72 angular sectors 10 degrees wide each, with position angles between 0 and 360 degrees. Each sector was overlapping the previous one by 5 degrees. We did this exercise for galaxies in a circle of 2.5Mpc radius, and in an annulus between 2.5 and 5 Mpc.

5.4.2. Filament detection and signal enhancement

Intuitively, if a given sector is significantly more populated than other sectors, it means that this sector is including a galaxy overdensity which could be explained by a filament or by a group along a filament. The question is then to define a significance level. For a given cluster (and then a given redshift slice) and a given radius, we chose to compute the mean and dispersion of the galaxy numbers in the 72 considered sectors. If a given sector had a number of galaxies larger than the mean + 3 times the dispersion, we considered this sector as hosting a potential cosmic filament portion.

However, individual clusters exhibit at best a single $3\text{-}\sigma$ significant candidate filament. This is due to the intrinsic very low galaxy density in filaments. Moreover, the goal of the present section is not to make individual cluster studies, but to draw statistical tendencies. In order to enhance the significance of the filament detections, we therefore stacked different clusters, considering two categories: the *luminous* and the *moderately luminous* clusters. Other categories did not have enough cluster members in the selected redshift range. This technique is based on the assumption that the angular separation between different filaments feeding a given cluster is more or less constant. In order to make such a stack we now need to homogeneize the cluster position angles. We chose the position angle of the highest galaxy overdensity (the PA), limiting our-

Fig. 12. Stacked X-ray images with position angles defined by the highest galaxy overdensities aligned along the 180deg arbitary angle. Images were rescaled to physical units according to cluster redshift. Image size is 1 Mpc×650 kpc. Contours were drawn with a 20×20 pixel smoothing. Upper figure: mean stacking. Lower figure: median stacking.

Fig. 13. Same as Fig. 12 with a median stacking and without any position angle correction.

selves to clusters exhibiting a detection more significant than the 3- σ level. Selected clusters were rotated to have their most significant filament at an arbitrary position angle of 180 degrees (east-west direction). In order to check if this alignment technique has a physical meaning, we superposed in the same way the X-ray images using the position angles defined by the highest galaxy overdensities (more significant than the 3- σ level). After rotating these X-ray images, we spatially rescaled them to physical units (kpc) according to redshift, and we simply added them together, taking into account the corresponding weight maps. The resulting point spread function is a mean of the individual values and remains small compared to cluster typical sizes. Fig. 12 shows that we generate in this way a clearly elongated synthethic X-ray cluster along the 180deg direction. The measured ellipticity of the external isophote is equal to 0.41. If instead of a simple sum we compute the median of the images (Fig. 12), the resulting ellipticity of the external isophote is still 0.36. Finally, if we combine the X-ray images without correcting by the optically determined orientation, we produce Fig. 13, showing a basically null ellipticity.

If the galaxy-defined prefered orientations are valid, the detected elongation in X-rays is an expected behaviour as X-ray emiting groups are also expected to fall onto clusters coming from surrounding filaments (see e.g. Boué et al. 2008).

We have to take into account the cluster redshift before merging their galaxy populations. A single catalog magnitude limit would evidently increase the weight of nearby clusters as compared to more distant ones. We therefore limited the galaxy catalogs to i'=23 at z=0.5. The limits were brighter by D magnitudes for nearer clusters with D being the distance moduli difference between the cluster redshifts and z=0.5.

Renormalizing finally the galaxy counts by the number of selected clusters in a given class, we are able to produce figures giving the mean galaxy counts as a function of the angular position.

5.4.3. Results

We first draw stacked (using the previously defined PA of each cluster) angular galaxy counts for *luminous* and *moderately luminous* clusters in the annulus [2.5,5]Mpc (Fig. 14). The minimal and maximal radii have been choosen to be close to the mean virial radius of clusters (e.g. Carlberg et al. 1996) and not too large in order to limit the contamination by other clusters. These annuli will therefore mainly sample the infalling galaxy layers, just before the cluster dominated

Table 5. Same as Table 3 but for C0 clusters. An approximate upper limit for the X-ray luminosity of these clusters would be the faintest detected value for C3 clusters: $\sim 0.08 \ 10^{43} \ \text{erg/s}$.

Name	XLSSC	PH	RA	DEC	N	ZBWT	ERRZ	SIG	ERR
			deg	deg				km/s	km/s
022207.9-042808*	-	1	35.533	-4.469	2	0.316			
022402.4-051753	000	1	36.010	-5.298	11	0.496	0.001	435	88
022405.0-041612	-	1	36.021	-4.270	8	0.862	0.001	457	70
022528.3-041536	045	1	36.369	-4.261	4	0.556			
022550.4-044500*	-	1	36.460	-4.750	2	1.529			
022647.5-041428*	-	1	36.698	-4.241	1	0.742			
022829.7-031257*	-	0	37.124	-3.216	2	0.313			




Fig. 14. Stacked angular galaxy counts (in arbitray units) for *luminous* (black line) and *moderately luminous* (red line) clusters in annuli of [2.5,5]Mpc. The horizontal lines show the $3-\sigma$ detection levels.

areas. As expected, the signal from the most significant filament candidate is drastically increased, but no other features are detected at the $3-\sigma$ level besides the main filament.

We redo now the same exercise inside a 2.5 Mpc radius central area (Fig. 15). This area is mainly dominated by the clusters themselves (the few hundreds of kpc central areas) and by the galaxy layers just beginning to experience the cluster influence (close to the virial radius). We therefore investigate the cluster region as fed by the connected filaments. The signal from the main filaments is still increased. Other significant filament candidates are detected at the $3-\sigma$ level mainly for the *moderately luminous* cluster sample.

This difference between the 2.5 Mpc radius central area and the [2.5,5]Mpc annulus could be explained if the immediate vicinity of the considered clusters would be depopulated by the potential well of the clusters, diminishing the contrast between cosmic filaments and voids. Larger spectroscopic redshift samples will soon become available in the area and will allow us to refine our results in future works.

6. Cluster galaxy populations characteristics

We now investigate the optical properties of the galaxy populations in association with the X-ray clusters. We refer the reader




Fig. 15. Stacked angular galaxy counts (in arbitray units) for *luminous* (black line) and *moderately luminous* (red line) clusters inside a circle of 2.5 Mpc radius. The horizontal lines show the $3-\sigma$ detection levels.

to Urquhart et al. (2010) for individual studies of the clusters providing a temperature measurement.

6.1. Rest frame Red Sequences

The so-called red sequence (RS hereafter) commonly shows up to at least $z\sim1.2$ (e.g. Stanford et al. 2002) in the massive structures. It is also detected in a less compact state in field galaxy populations up to $z\sim2$ (Franzetti et al. 2007). We therefore searched RSs in our sample of clusters. This sample does not provide enough statistics per cluster in order to perform individual studies. The optimal strategy is therefore to build synthetic clusters by gathering galaxies for clusters of the same category. We therefore considered 4 classes of clusters: the most luminous, the luminous, the moderately luminous, and the C0 clusters. The RS being a powerful tool to characterize the evolutionary stage of the cluster galaxy populations (e.g. Adami et al. 2007b), such a study will allow us to assess the properties of these 4 cluster classes.

In order to be able to stack different clusters at different redshifts, rest frame absolute magnitudes were computed in the process of getting photometric redshifts with LePhare (e.g. Ilbert et al. 2006) and we used these magnitudes to compute colors. Basically the method consists in selecting the observed band which is the closest of the requested rest frame band to

compute the magnitude in this band applying correction factors. They are described in the annex of Ilbert et al. (2005), including for example k-correction. This method is the closest of the observations and minimizes our dependence on the assumed spectral energy distributions, which could not be exactly the same in clusters and in the field (see also annex I of the present paper).

6.1.1. Red sequence using spectrocopic redshifts

In a first step we look only at galaxy members using spectroscopic redshifts rather than photometric ones in order to remove potential interloper galaxies which are non cluster members but close to the cluster redshift. Such galaxies could be interpreted as cluster members considering only photometric redshifts because of their limited precision. We here consider u*-r' rest frame colours and look at their behaviour versus rest frame r' absolute magnitude.

Fig. 16 shows that a RS is present with $u^*-r^*\sim 2$ for all clusters. The slopes of the RS appear in good agreement with literature estimates (e.g. Adami et al. 2007b: between -0.1 and -0.02 for the Coma cluster) and are given in Table 6. As expected, RSs are populated by early type galaxies, while later type objects are grouped in a much less compact bluer sequence.

There are potential differences between C0 clusters (without detectable X-ray emission) and other classes. The slope of the RS appears nearly flat for C0 clusters while being more negative for more luminous clusters. This effect is only poorly significant when considering the uncertainty of the slope of these RS's. We performed however a bi-dimensional Kolmogorov-Smirnov statistical test on the early type galaxies of Fig. 16. The probability that the C0 and the *most luminous* cluster early type galaxies come from the same population is only 0.6%. The probability that the C0 and the *luminous* cluster early type galaxies come from the same population is only 0.1%. Finally, the probability that the C0 and the moderately luminous cluster early type galaxies come from the same population is 3.1%. At least for the most luminous and luminous cluster populations, early type galaxies therefore seem to be differently distributed in a color magnitude relation compared to C0 cluster early type galaxies. If these differences come from the slope of the RS, this effect could be interpreted as a metallicity effect (Kodama & Arimoto, 1997). The more massive a galaxy, the more easily it will retain metals against dissipative processes. The more metals present in a galaxy, the redder the galaxy will be. Massive galaxies are therefore expected to be redder than lower mass objects. A possible explanation would be that the faint early type C0 cluster galaxies would originate from depleted cores of larger galaxies, so being metal rich before becoming faint (see e.g. Adami et al. 2006). This is possible for example in small groups where velocity dispersion is low enough to favor galaxy-galaxy encounters.

Galaxie members of the *most luminous* clusters also appear to exhibit a more pronounced dichotomy between early and late type objects. Blue members of the *most luminous* clusters are clearly bluer than blue members of the less luminous clusters.




Fig. 16. Rest frame u*-r' versus absolute r' magnitude relation for clusters using spectroscopic redshifts to compute absolute magnitudes. We only plot cluster members in these figures. From top to bottom, figures are for the *most luminous*, the *luminous*, the *moderately luminous*, and the C0 clusters. Red symbols correspond to early type galaxies ($T \le 21$), blue symbols correspond to late type non starburst galaxies ($58 \ge T \ge 21$), and pink symbols correspond to starburst galaxies ($T \le 59$). Black continuous lines are computed using only $T \le 21$ galaxies (see Table 6).

Table 6. Slopes of the red sequences for the four classes of clusters: *most luminous, luminous, moderately luminous* and CO.

Category	Slope
most luminous	-0.04 ± 0.04
luminous	-0.04 ± 0.02
moderately luminous	-0.10 ± 0.04
C0	-0.01 ± 0.05

6.1.2. Age of formation of the cluster galaxy stellar populations

We expect distant clusters to naturally exhibit younger galaxy star populations compared to nearby structures. In order to investigate this question, we computed with LePhare ages of stellar population in galaxies with a spectroscopic redshift lying inside the considered clusters. The templates used to generate public photometric redshifts in the CFHTLS does not allow to provide this information, so we used in LePhare the Bruzual & Charlot (2003) templates, fixing the redshifts to the spectroscopic values. The metallicity was let free to vary between 0.004, 0.008, and 0.02 Z_{\odot} . In C0 clusters, z=[0.3;0.6] galaxies have a stellar population aged of 6.2 ± 1.9 Gyr, and z=[0.7;0.9] galaxies have a stellar population aged of only 2.7±1.3 Gyr. Considering members of luminous clusters, z=[0.25;0.35] galaxies have a stellar population aged of 7.4 ± 1.0 Gyr, and z=[0.35;0.65] galaxies have a stellar population aged of only 5.3±2.1 Gyr. Finally, members of the most luminous clusters, z=[0.4;0.65] galaxies have a stellar population aged of 5.2 ± 2.1 Gyr, and z=[0.75;1.25] galaxies have a stellar population aged of 3.3±1.1 Gyr.

Taking the mean redshift of the highest redshift bin for each of these 3 categories and diminishing the corresponding elapsed time since the beginning of the Universe by the mean age of the early type galaxy stellar populations leads us to estimate the mean age of formation of the star populations in these galaxies. Galaxy stellar populations probably formed at $z\sim1.6$ in C0 clusters, at $z\sim2$ in *luminous* clusters, and at $z\sim2.5$ in the most *most luminous*. These values are in good agreement with general expectations for the massive clusters to form early than low mass structures, up to redshifts close to $z\sim2$.

6.1.3. Red sequence using photometric redshifts and color-color diagrams

In order to study larger samples and detect possible weak effects, we used photometric redshifts to define a cluster membership, and compute absolute magnitudes and colors as provided by the CFHTLS data. Given its photometric redshift, a galaxy was assigned to a cluster when closer than 500 kpc from the cluster center and at less than 0.08 from the cluster redshift. This corresponds to the values quoted in Table A.1 for cluster galaxies. We then were able to search for RSs in the *most luminous*, the *luminous*, and the *moderately luminous* clusters. Selecting all available clusters in these three categories Fig. 17 clearly shows red sequences in each case. They all are

consistent with a u*-r' color of 1.9, the most massive clusters exhibiting the more negative RS slope (computed with $T \leq 21$ galaxies). On the contrary, the C0 clusters (no X-ray detection) only exhibit a very low number of early type galaxies (but still consistent with u*-r' ~1.9). These structures therefore appear as quite young structures, with modest early type galaxy populations.

However, we are merging in Fig. 17 clusters with quite different redshifts and evolutionary effects could play an important role. We therefore selected only the *luminous* clusters (the only category providing enough clusters) and we divided this population in 3 different redshifts bins (\leq 0.3,]0.3,0.5], and]0.5,0.8]) in Fig. 18. This figure only shows $T \leq 21$ galaxies (early types). RSs appear very similar, with the most negative slope occuring for z=]0.5,0.8] clusters. If evolutionary effects are present, they are therefore rather weak, besides the most distant clusters appearing to have the most negative RS slope (-0.069 ± 0.017). This is consistent with the slope computed for the *most luminous* clusters (which are also nearly all at redshift greater than 0.5): -0.052 ± 0.015 .

It could be argued that the use of photometric redshifts could introduce a bias due for example to SEDs not adapted to high density regions. In order to check the previous results, we therefore simply draw u*-r' versus r'-z' color-color diagrams for the same sets of clusters. Fig. 19 shows that in both cases, early types still occupy well defined loci in the color-color space, confirming the existence of an old galaxy population in these cluster classes.

We therefore confirm that both massive and less massive X-ray structures in our sample exhibit quite similar red sequences, making them overall quite old structures. Non X-ray clusters are probably minor structures with a poor spectral early type population.

Fig. 17 also shows a slightly larger percentage of starburst galaxies (as determined during the photometric redshift computation process: see Coupon et al., 2009) in low luminosity clusters. C0, *moderately luminous*, and *luminous* clusters exhibit 20% more starburst galaxies compared to the *most luminous* clusters. This is an expected behaviour, for example in good qualitative agreement with Urquhart et al. (2010).

6.2. Luminosity Functions

In the same spirit, we checked whether our structures behave as genuine clusters or groups concerning their galaxy luminosity functions. For a detailed study of the individual XMM-LSS C1 cluster luminosity functions, we refer the reader to Alshino et al. (2010). We computed luminosity functions using galaxies within the cluster bins (according to photometric redshifts). The Schechter function fitting was performed allowing a constant background to take into account galaxies included in the photometric redshift slice but not part of the clusters.

Selecting all clusters (C1+C2+C3+C0), stacking their luminosity functions, and only limiting absolute magnitude to $i' \le -17.5$ in order to not be too affected by incompleteness, we got a best fit of a Schechter function with alpha = -1.15 ± 0.09




Fig. 17. u*-r' versus r' with red dots being $T \le 21$ galaxies (early types), blue dots being $58 \ge T \ge 21$ galaxies (late types), and pink dots being starburst galaxies. From top to bottom, figures are for the *most massive*, the *massive*, the *moderately massive*, and the $z \le 1$ C0 clusters. Absolute magnitude computations are based on photometric redshifts. Black continuous lines are the RSs computed with $T \le 21$ galaxies, except for C0 clusters where we had not enough available early type galaxies.




Fig. 18. u*-r' versus r' for $T \le 21$ galaxies (early types). Red dots: z=]0.5,0.8], green dots: z=[0.3,0.5], and blue dots: z=[0.,0.3]. Black continuous lines are the RSs computed with $T \le 21$ galaxies.

and M*i'=-23.8 \pm 0.8. This is consistent within error bars with the estimates of Alshino et al. (2010) at z \sim 0.3. Playing the same game with the C3 clusters, we get a slightly shallower Schechter fit: alpha = -0.96 \pm 0.14 and M*i'=-22.1 \pm 0.6.

If we use the luminosity categories, we can compute similarly Schechter fits for the *luminous* and the *moderately luminous* clusters (*most luminous* clusters are too distant and therefore undersampled toward the faint magnitudes, and C0 and C2 clusters are not numerous enough). We get alpha = -1.1 ± 0.03 and M*i'= -23.4 ± 0.3 for the *moderately luminous* and we get alpha = -1.1 ± 0.03 and M*i'= -23.2 ± 0.2 for the *luminous* clusters.

The fitted Schechter functions are in agreement with those of bona fide clusters at similar depth (e.g. Lumsden et al. 1997). Slopes are also similar within error bars between all cluster classes. C1 clusters seem to exhibit however, brighter M_* than C3 clusters, in good agreement with the fact that C1 clusters would be older and more massive systems than C3 clusters.

7. Peculiar structures in the XMM-LSS

7.1. Distant cluster candidates

Several structures with redshifts ~ 1 or greater have already been found in Class 1 (Pacaud et al 2007) or 2 (Bremer et al. 2006). Some other candidates appear among the C2's (e.g. XLSS J022756.3-043119 at z ~ 1) and the C0's (022550.4-044500 at z ~ 1.53). This last structure (Fig. 20) is just below the X-ray detection limit. It has an extension of ~ 13 arcsec and its extension maximum likelihood is ~ 10 . We note that the measured flux is $0.2\pm 0.1\ 10^{-14}\ erg/s/cm^2$. At z=1.53 and for a temperature of 1.5keV, this would lead to an X-ray luminosity of $8.6e+43\ ergs/s$.

The weakness of the evidences for an X-ray detection leads us, however, to classify this source as C0 and then to investigate it from the optical side. The regular CFHTLS photometric redshifts (based on u*g'r'i'z' magnitudes) are not well suited to study this potential structure because of a lack of near infrared photometric bands. This candidate is however




Fig. 19. u*-r' versus r'-z diagrams with red dots being $T \le 21$ galaxies (early types), blue dots being $58 \ge T \ge 21$ galaxies (late types), and pink dots being starburst galaxies. The upper figure is for the *most luminous*, the middle figure is for the *luminous*, and the lower figure is for the *moderately luminous* clusters.

included in the WIRDS survey (near infrared imaging from CFHT-WIRCAM). Photometric redshifts have been computed combining these near infrared data (McCracken et al., private communication) and the CFHTLS deep magnitudes. Fig. 20 shows a clear concentration of z=[1.43;1.63] galaxies inside the XMMLSS contours. We therefore may have detected one of the most distant known clusters of galaxies. A near infrared spectroscopic follow up of this candidate is however mandatory in order to confirm the nature of this very weak X-ray source.

Fig. 20. A distant cluster candidate at z=1.53. Large red circle is a 500 kpc radius circle. Blue circles are galaxies with spectroscopic redshifts outside the z=[1.52;1.54] interval. The two magenta squares are the two known spectroscopic redshifts inside the z=[1.52;1.54] interval. Small red circles are the near-infrared-based photometric redshifts inside the z=[1.43;1.63] interval. White contours are the XMM-LSS contours.

7.2. Structures with discrepant optically and X-ray contents

XLSSC 000 is a C0 structure not detected in the X-rays. Its velocity dispersion is however relatively large (435±88 km/s). The Serna-Gerbal analysis does not detect any sign of substructures with the 11 known spectroscopic redshifts, so this velocity dispersion does not appear as obviously biaised. The photometric redshift distribution also presents excesses at the structure redshift. This structure is finally populated with a significant number of early types galaxies: among the 21 objects within the z=0.49 photometric redshift slice, 9 have type T≤10. The optical content is therefore similar to what we could expect if considering a massive cluster. This case with clear discrepancies between X-ray and optical content remains quite puzzling and both deeper X-ray observations and additional spectroscopic followup are required to explain the observed behaviour.

We also have detected a prominent X-ray structure which is much less evident in optical and which could be a fossil group (XLSSU J021754.6-052655). Described for example in Jones et al. (2003) or Mendes de Oliveira et al. (2006 and references therein), these structures are considered as the ultimate stage of group evolution: the nearly complete fusion of all the bright and intermediate magnitude galaxies of the group into a single bright galaxy. The resulting galaxy is brighter than the second remaining group galaxy (within half the projected virial radius) by at least 2 magnitudes (in the R band). However, the extended X-ray gas envelope is still present and more luminous than $10^{42} h_{50}^{-2}$ erg s⁻¹ (Jones et al. 2003). The origin of these structures is however being still widely debated. They could find their origin in the small impact parameter of $L \sim L_*$ galaxies travelling along filaments (e.g. D'Onghia et al. 2005), or simply in their highly isolated status (e.g. Adami et al. 2007a) so that no galaxies will then have fallen into them lately.

In our survey, XLSSU J021754.6-052655 (classified as C2) is quite similar to such fossil groups. Fig. 21 shows the field covered by this galaxy structure. The X-ray source is clearly extended. Available spectroscopic redshifts only show 2 galaxies at the structure redshift which are only slightly too bright to satisfy the 2 magnitudes criteria (one is satisfying the criterion in i' and z' band). Photometric redshifts from Coupon et al. (2009) also exhibit only two other similar galaxies at less than 0.15 from the structure redshift. Considering error bars on magnitude, the magnitude difference between the brightest galaxy and the second brightest object could be consistent with the requested 2 magnitudes gap at the 3- σ level. We therefore conclude that this object is similar to the structure described in Ulmer et al. (2005) and is very close to the fossil group status.

Fig. 21. CFHTLS i' band image of the XLSSU J021754.6-052655 XMM-LSS source. White contours are XMM X-ray emission. Pink squares are galaxies with a spectroscopic redshift inside the structure. Red circles are galaxies with a photometric redshift at less than 0.15 from the structure redshift. Large symbols (circles or squares) are galaxies not satisfying the 2 magnitudes criteria in g', r', i', or z' bands. For these objects we also give the magnitude difference with the brightest galaxy in g'/r'/i'/z'.

We investigate if this group is the dominant structure of its cosmological bubble (similarly to Adami et al. 2007a). For this, we selected all known spectroscopic redshifts in the range [0.241,0.261] and at less than 1.5deg from the group (about 20 Mpc at the structure redshift, close to the average size of known voids: e.g. Hoyle & Vogeley 2004). Contrary to the results of Adami et al. (2007a), our group does not appear as an isolated structure (Fig. 22). The ratio of galaxies with a spectroscopic redshift inside and outside the range [0.241,0.261] is not significantly different when considering the 1.5deg region or the complete spectroscopic sample.




Fig. 22. α , δ map of the immediate vicinity of XLSS J021754.6-052655. Black dots are all galaxies with a known spectroscopic redshift in a 1.5deg radius region (large blue filled circle). Red dots are galaxies in the redshift range [0.241,0.261].

8. Conclusions

Starting from known XMM-LSS sources, we considered 75 of them for which at least two spectroscopic redshifts were available within the X-ray isophotes. We then generated a catalog of 59 groups or clusters of galaxies in the z=[0.05;1.53] redshift range associated with an X-ray source as well as 7 other real structures for which X-ray association is not clear. Finally, 11 redshift structures (named C999) detected along the various lines of sight were detected in addition of the main systems and are listed in Appendix. In 3 cases the X-ray sources are in fact associated with QSO's identified from their optical spectra.

The assessment of the clusters and groups as actual massive structures has been based on various spectroscopic data (including PI observations) associated to photometric data from the CFHTLS T0004 release (when available) and some PI data. The analysis (without a-priori knowledge of their X-ray class) of the optical lines of sight centered on the X-ray emission was based on criteria such as compactness in redshift space (spectroscopic and photometric), and significant excess in galaxy density obtained within photometric redshift slices and final visual inspection.

All the detected systems exhibit "bona fide" clusters or groups optical properties in terms of red sequence, color-color clumping, luminosity function, and morphological segregation. Considering X-ray luminosity classes does not change the results. From the X-ray and optical properties of the structures now associated with the XMM extended sources, the C1 clusters can be considered in most cases as relatively nearby, X-ray bright and optically rich and regular (no sub-clustering) clusters, while C3's appear faint and poor at the same redshift and quite rich at high redshift. C2's are a mix with the exception of some distant possible candidates. Finally, looking at larger scales using the CFHTLS-W, these clusters statistically appear as clear nodes of the galactic cosmic web, reinforcing therefore their true existence. The full sample of X-ray clusters with associated optical spectroscopic data is available via the L3SDB database (http://l3sdb.in2p3.fr:8080/l3sdb/). The optical images as well as the details of the redshift determination for all clusters presented in this article will also be publicly available at this place.

Finally, we investigated the photometric redshift precision in our sample as a function of the environment and of the galaxy spectral types (see appendix). We show for example that the galaxy photometric redshift accuracy is degraded in the most massive clusters for early and late type galaxies.

Acknowledgements. The authors thank R. Bielby, J. Coupon, Y. Mellier, and H.J. McCracken for help. AD, TS and JS acknowledge support from the ESA PRODEX Programme "XMM-LSS", from the Belgian Federal Science Policy Office and from the Communauté francaise de Belgique - Actions de recherche concertées - Académie universitaire Wallonie-Europe. HQ acknowledges the support of FONDAP Center for Astrophysics 15010003. GG is supported by FONDECYT 1085267. The authors thank the referee for useful and constructive comments.

References

Adami, C., Mazure, A., Ilbert, O. et al., 2005, A&A 443, 805 Adami, C., Scheidegger R., Ulmer M.P., et al., 2006, A&A 459, 679

Adami C., Durret F., Mazure, 2007, A&A 462, 411 (b) Adami C., Russeil D., Durret F., 2007, A&A 467, 459 (a)

Adami C., Ilbert O. Pelló R., et al., 2008, A&A 491, 681

Adami C., LeBrun V., Biviano A., et al., 2009, A&A 507, 1225

Adami C., Durret F., Benoist C., et al., 2010, A&A 509, 81 Alshino A., Khosroshahi H., Ponman T., et al., 2010, MNRAS 401, 941

Beers T., Flyn K., Gebhardt K., 1990, AJ, 119, 2645

Biviano A., Katgert P., Mazure A. et al., 1997, A&A 321, 84 Bohringer H., et al. 1998, in Proc. 14th IAP Conf., Wide Field Surveys in Cosmology, ed. S. Colombi, Y. Mellier, & B. Raban (Paris: Editions Frontires), 261

Bolzonella M., Miralles J-M., Pelló R., 2000, A&A 363, 476 Boué G., Durret F., Adami C., et al., 2008, A&A 489, 11 Bremer M.N., Valtchanov I., Willis J., et al., 2006, MNRAS 371, 1427

Bruzual G., Charlot S., 2003, MNRAS 344, 1000

Carlberg R.G., Yee H.K.C., Ellingson E., et al. 1996, ApJ 462, 32.

Castander F. J., Bower R.G., Ellis R.S., et al. 1995, Nature 377, 39

Collins C. A., Burke D. J., Romer A. K., Sharples R. M., Nichol R. C. 1997, ApJ 479, L117

Coupon J., Ilbert O., Kilbinger M. et al. 2009, A&A 500, 981 De Grandi S., Bohringer H., Guzzo L., et al. 1999, ApJ 514, 148

D'Onghia E., Sommer-Larsen J., Romeo A.D. et al. 2005, ApJ 630, L109

Dunkley J., Komatsu E., Nolta M. R., et al., 2009, ApJS 180, 306

Ebeling H., Edge A. C., Bohringer H., et al. 1998, MNRAS 301, 881

Edge A.C., Stewart G.C., 1991, MNRAS 252, 428

Finoguenov A., Watson M.G., Tanaka M., et al., 2010, MNRAS 403, 2063

Finoguenov A., Guzzo L., Hasinger G., et al., 2007, ApJS 172, 182

Franzetti P., Scodeggio M., Garilli B., et al., 2007, A&A 465, 711

Gavazzi R., Soucail G., 2007, A&A 462, 459

Gandhi P., Garcet O., Disseau L., et al., 2006, A&A 457, 393 Guennou L., Adami C., Ulmer M.P., et al., A&A in press, astroph: 10085318

Hamana T., Miyazaki S., Kashikawa N., et al., 2009, PASJ 61, 833

Henry J. P. 1997, ApJ 489, L1

Hoyle F., Vogeley M. S., 2004, ApJ 607, 751

Ilbert O., Arnouts S., McCracken H.J., et al., 2006, A&A 457, 841

Ilbert O., Tresse L., Zucca E., et al., 2005, A&A 439, 863 Jones L. R., Scharf C., Ebeling H., et al. 1998, ApJ 495, 100 Jones L.R., Ponman T.J., Horton A., et al., 2003, MNRAS 343, 627

Kodama T., Arimoto N., 1997, A&A 320, 41

Le Fèvre, O.; Vettolani, G.; Garilli, B., et al., 2005, A&A 439, 845

Lumsden S.L., Collins C.A., Nichol R.C., et al., 1997, MNRAS 290, 119

Mazure A., Adami C., Pierre M., et al., 2007, A&A 467,49 Mendes de Oliveira C., Cypriano E.S., Sodré Jr L., 2006, AJ 131, 158

Pacaud F., Pierre M., Refregier A. et al. 2006, MNRAS 372, 578

Pacaud F., Pierre M., Adami C. et al. 2007, MNRAS 382, 1289

Pierre M., Valtchanov I, Altieri B., et al., 2004, JCAP 9, 11

Pierre M., Pacaud F., Duc P.-A. . et al. 2006, MNRAS 372, 591

Pierre M., Chiappetti L., Pacaud F. et al. 2007, MNRAS 382, 279

Quintana H., Melnick J., 1982, AJ 87, 972

Rizzo, D., Adami, C., Bardelli, S. et al., 2004, A&A 413, 453 Romer A.K., Collins C. A., Bohringer H., et al., 1994, Nature 372, 75

Romer A. K., Nichol R. C., Holden B. P., et al. 2000, ApJS 126, 209

Romer A.K., Viana P.T.P., Liddle A.R., Mann R.G., 2001, ApJ 547, 594

Rosati P., Della Ceca R., Norman C., Giacconi R., 1998, ApJ 492, L21

Serna A., Gerbal D., 1996, A&A 309, 65

Smith B.W., Mushotzky R.F., Serlemitsos P.J., 1979, ApJ 227, 37

Stanford S.A., Holden B., Rosati P., et al., 2002, AJ 123, 619 Ueda Y., Watson M., Stewart G., et al 2008 ApJS 179, 124 Ulmer M.P., Adami C., Covone G., et al., 2005, ApJ 624, 124 Urquhart S. A., Willis J. P., Hoekstra H., Pierre M., 2010, MNRAS in press, astroph: 10040020

Vikhlinin A., McNamara B. R., Forman W., et al. 1998, ApJ 502, 558

Werner N., Finoguenov A., Kaastra J.S., et al. 2008, A&A 482, L29

Yee H. K. C., Ellington, E., 2003, ApJ 585, 226

Appendix A: Photometric redshifts in dense environments

A by-product of the present paper is the test of photometric redshift precision in dense environments. Photometric redshift technique is widely used for several cosmological purposes, and is mainly based on synthetic energy distributions (SEDs hereafter) fits to observed magnitudes. The available SEDs in the literature are however mainly selected in low density environments, outside clusters. Applying these SEDs to cluster galaxies is then potentially problematic. Several papers (e.g Adami et al. 2008) seem to show various photometric uncertainties as a function of the galaxy spectral type in these dense environements. If confirmed, this could be due in massive structures to environmental effects driving peculiar color galaxy evolutions. Degeneracies could then be induced between photometric redshift value and galaxy spectral type when applying classical photometric redshift codes as LePhare or HyperZ (Bolzonella et al. 2000). However, these tendencies are still based on very sparse samples for clusters of galaxies and before embarking in the very demanding task of building cluster-dedicated SEDs, we have to put on a firmer ground the photometric redshift uncertainty variation as a function of the environment and of the galaxy spectral type.

The XMM-LSS survey offers such a unique opportunity, both providing X-ray and optical characterizations of the clusters, and photometric redshift informations from the CFHTLS. We selected all spectroscopic redshifts included in the present clusters and located in the 500 kpc (radius) central area. This

insures us to have galaxies really located in the densest areas of the clusters. Then, we extracted informations (photometric redshift itself and spectral type) from the CFHTLS T0004 photometric redshift release. Finally, we considered separately clusters brighter than 10^{44} erg/s, between 10^{43} and 10^{44} erg/s, and fainter than 10^{43} erg/s. We also considered the C0 class, acting as the low mass cluster category (we remind that these structures are real but without clear X-ray emission).

A.1. General agreement between spectroscopic and photometric redshifts

We first checked that the general agreement between spectroscopic and photometric redshifts was acceptable inside clusters of galaxies. Fig. A.1 shows a good agreement. Selecting a priori galaxies with a $|zphot-zspec| \leq (0.15 \times (1+zspec))$, the whole cluster galaxy sample exhibits a σ of 0.06. This is only slightly larger than the estimates of Coupon et al. (2009) for the whole CFHTLS W1 field. This shows that from a general point of view, CFHTLS T0004 photometric redshifts are not clearly worst in clusters than in the field. We have now to investigate in more details the behaviour of the photometric redshift uncertainty as a function of the galaxy spectral type and as a function of the cluster characteristics.




Fig. A.1. Photometric versus spectroscopic redshifts for the cluster galaxies in our sample. Black lines give the perfect relation of slope 1 and the $\pm 0.15 \times (1+z)$ classical uncertainty (see e.g. Ilbert et al. 2006). Red filled circles are early spectral type galaxies and blue filled circles are late spectral type galaxies (see text).

A.2. Photometric redshifts in dense environments and galaxy spectral types

We redid the previous analysis splitting our samples into early and late type galaxies and considering the C0, the *most luminous*, the *luminous*, and the *moderately luminous* clusters. Table A.1 gives the values of σ (computed in the same way as in the previous subsection) in these different cases.

We first detect a clear tendency to have higher uncertainties in photometric redshift calculations in the *most luminous*

Table A.1. σ between photometric and spectroscopic redshifts as a function of the environment and of the galaxy spectral type. The last column indicates the percentage of spectral types $T \leq 21$ galaxies.

Cluster class	Early	Late	% of early types
most luminous	0.081	0.092	69%
luminous	0.036	0.082	81%
moderately luminous	0.047	0.069	72%
C0	0.043	0.064	75%
Global cluster sample	0.048	0.096	76%

clusters. Second, late type galaxies in *luminous* and *moderately luminous* clusters (as well as in C0 clusters) also exhibit higher uncertainties than early type galaxies, by a factor of 2 in *luminous* clusters and by 50% in *moderately luminous* and C0 clusters. This behaviour was already detected in Guennou et al. (2010).

This can be explained if galaxies were undergoing peculiar evolutions in clusters of galaxies, depending of the mass of the considered clusters, making them different from field galaxies. This would occur for all galaxy types in the most massive clusters, while less massive clusters would only affect late type galaxies. These various environments do not seem to strongly affect the percentage of early type galaxies which stays high anyway (see Table A.1). A finer analysis shows however, as expected, a regular increase of the mean type of $T \le 21$ galaxies, from 1.4 for the *most luminous* clusters, to 2.2 for the *luminous* clusters, and finally to 3.0 for the *moderately luminous* clusters.

As a conclusion, we can then say that photometric redshift values are globally correct in clusters of galaxies of the present sample (as compared to field environments). However, all galaxies in the most massive clusters and late type galaxies in all other clusters have their photometric redshift uncertainty increased by a factor of 50 to 100%. Depending on the science goals, this can significantly affect the cluster population definition by photometric redshift criteria, for example for galaxy luminosity function purposes. It would therefore be useful to create cluster-dedicated spectroscopic SEDs.

Appendix B: Additional redshift structures

As a bonus of the general cluster detection process, for a given line of sight, other real galaxy groups are detected besides the ones associated with the X-ray emission (C999: see Table B.1). So, if the identification with an optical group would appear wrong in the future, or if more data become available, trace is kept to re-examine other possibilities.

Table B.1. Same as Table 3 but for other real groups (C999) detected along the lines of sight.

XLSSC	PH	RA	DEC	N	ZBWT	ERRZ	SIG	ERR
			deg	deg			km/s	km/s
065	1	34.245	-4.821	3	0.138			
039	0	35.098	-2.841	3	0.183			
044	1	36.141	-4.234	11	0.317	0.001	410	87
-	1	36.424	-4.410	4	0.142			
-	1	36.424	-4.410	4	0.632			
-	1	36.424	-4.410	3	0.915			
-	1	36.460	-4.750	3				
-	1	36.698	-4.241	3	0.210			
-	1	36.698	-4.241	3	0.432			
-	1	36.698	-4.241	3	0.705			
013	1	36.858	-4.538	9	0.254	0.001	346	75

