

Generation of finite simple groups by an involution and an element of prime order

Carlisle S. H. King
Imperial College
London
United Kingdom
SW7 2AZ
 carlisle.king10@ic.ac.uk

Abstract

We prove that every non-abelian finite simple group is generated by an involution and an element of prime order.

1. Introduction

Given a finite simple group G , it is natural to ask which elements generate G . Results of Miller [29], Steinberg [39], Aschbacher and Guralnick [2] prove that every finite simple group is generated by a pair of elements. A natural refinement is then to ask whether the orders of the generating elements may be restricted: given a finite simple group G and a pair of positive integers (a, b) , does there exist a pair of elements $x, y \in G$ with x of order a and y of order b such that $G = \langle x, y \rangle$? If such a pair exists, we say G is (a, b) -generated.

As two involutions generate a dihedral group, the smallest pair of interest is $(2, 3)$. The question of which finite simple groups are $(2, 3)$ -generated has been studied extensively. All alternating groups A_n except for $n = 3, 6, 7, 8$ are $(2, 3)$ -generated by [29]. All but finitely many simple classical groups not equal to $PSp_4(2^a), PSp_4(3^a)$ are $(2, 3)$ -generated by [24]. In fact, recent work by Pellegrini [32] completes the classification of the $(2, 3)$ -generated finite simple projective special linear groups, which shows that $PSL_n(q)$ is $(2, 3)$ -generated for $(n, q) \neq (2, 9), (3, 4), (4, 2)$. There is also literature on the $(2, 3)$ -generation of many other simple classical groups $Cl_n(q)$, showing a positive result for large n explicitly listed (for example, see [36]). All simple exceptional groups except for ${}^2B_2(2^{2m+1})$ (which contain no elements of order 3) are $(2, 3)$ -generated by [25]. And all sporadic simple groups except for M_{11}, M_{22}, M_{23} and McL are $(2, 3)$ -generated by [45].

Nevertheless, the problem of determining exactly which finite simple groups are $(2, 3)$ -generated, or more generally $(2, p)$ -generated for some prime p , remains open. In this paper, we prove:

Theorem 1. *Every non-abelian finite simple group G is generated by an involution and an element of prime order.*

By [29], for $n \geq 5$ and $n \neq 6, 7, 8$, the alternating groups A_n are $(2, 3)$ -generated, and by [30] these exceptions are $(2, 5)$ -generated. By [25] the exceptional groups not equal to ${}^2B_2(2^{2m+1})$ are $(2, 3)$ -generated, and by [11] the Suzuki groups are $(2, 5)$ -generated. By [45] the sporadic groups not equal to $M_{11}, M_{22}, M_{23}, McL$, are $(2, 3)$ -generated, and by [44] these exceptions are $(2, p)$ -generated for $p = 11, 5, 23, 5$ respectively (in fact, all of these exceptions are $(2, 5)$ -generated, which can be seen using GAP). By Lemma 2.4 below, the 4-dimensional symplectic groups $PSp_4(2^a)$ ($a > 1$), $PSp_4(3^a)$ are $(2, 5)$ -generated, and when combined with Lemmas 2.1 and 2.2, this shows that all finite simple classical groups with natural module of dimension $n \leq 7$ (and $P\Omega_8^+(2)$) are $(2, p)$ -generated for some $p \in \{3, 5, 7\}$.

By Zsigmondy's theorem [46], for $q, e > 1$ with $(q, e) \neq (2^a - 1, 2), (2, 6)$, there exists a prime divisor $r = r_{q,e}$ of $q^e - 1$ such that r does not divide $q^i - 1$ for $i < e$. We call r a primitive prime divisor of $q^e - 1$. Notice that, in general, $r_{q,e}$ is not uniquely determined by (q, e) . In the group $(\mathbb{F}_r)^\times$, q has order e , and so $r \equiv 1 \pmod{e}$. In view of the above discussion, Theorem 1 follows from the following result.

Theorem 2. *Let G be a finite simple classical group with natural module of dimension n over \mathbb{F}_{q^δ} , where $\delta = 2$ if G is unitary and $\delta = 1$ otherwise. Assume $n \geq 8$ and $G \neq P\Omega_8^+(2)$. Let r be a primitive prime divisor of $q^e - 1$, where e is listed in Table 1. Then G is $(2, r)$ -generated.*

Table 1: Values of e in Theorem 2

G	e
$PSL_n(q), PSp_n(q), P\Omega_n^-(q)$	n
$P\Omega_n^+(q)$	$n - 2$
$P\Omega_n(q)$ (nq odd)	$n - 1$
$PSU_n(q)$ (n odd)	$2n$
$PSU_n(q)$ (n even)	$2n - 2$

We note that r is well-defined for the groups described in Theorem 2.

There is a large literature on other aspects of the generation of finite simple groups, and we note just a few results. In [27], it is shown that every non-abelian finite simple group other than $PSU_3(3)$ is generated by three involutions. In [13], and proved independently in [38], it is shown that, given a finite simple group G , there exists a conjugacy class C of G such that, given an arbitrary non-identity element x in G , there exists an element y in C such that $G = \langle x, y \rangle$.

We now sketch our approach to proving Theorem 2. Let G be any finite group. Let $M <_{\max} G$ denote a maximal subgroup. For a group H let $i_m(H)$ denote the number of

elements of order m in H . Let $P_{2,p}(G)$ denote the probability that G is generated by a random involution and a random element of order p , and let $Q_{2,p}(G) = 1 - P_{2,p}(G)$. Then we have

$$Q_{2,p}(G) \leq \sum_{M <_{\max} G} \frac{i_2(M) i_p(M)}{i_2(G) i_p(G)}, \quad (1.1)$$

since the right-hand side is an upper bound for the probability that a random involution and a random element of order p lie in some maximal subgroup of G . To prove G is $(2, p)$ -generated, it suffices to prove $Q_{2,p}(G) < 1$.

In fact, for the proof of Theorem 2 we also need a refinement of (1.1). Let G, r be as in Theorem 2, and let $x \in G$ be an element of order r . Let $P_2(G, x)$ denote the probability that G is generated by x and a random involution, and let $Q_2(G, x) = 1 - P_2(G, x)$. Then by similar reasoning we have

$$Q_2(G, x) \leq \sum_{x \in M <_{\max} G} \frac{i_2(M)}{i_2(G)}. \quad (1.2)$$

To prove G is $(2, r)$ -generated, it suffices to prove $Q_2(G, x) < 1$. Our method in most cases is to determine the maximal subgroups M of G containing x , and then bound $i_2(M)$ and $i_2(G)$ in terms of n and q such that for n and q sufficiently large we have $Q_2(G, x) < 1$. For the remaining cases with small n and q we improve the bounds case by case using literature such as [4].

Acknowledgements. This paper is part of work towards a PhD degree under the supervision of Martin Liebeck, and the author would like to thank him for his guidance throughout. The author is also grateful for the financial support from EPSRC.

2. Preliminary results

The $(2, 3)$ -generation of classical groups has been studied extensively, and there are many results for groups of small dimension that we will make use of.

Lemma 2.1. *If G is a finite simple classical group listed in Table 2, then G is $(2, 3)$ -generated.*

We note that though there are many other $(2, 3)$ -generation results regarding classical groups, our method will not require them.

Lemma 2.2. *If G is listed in Table 3 then G is $(2, p)$ -generated, where $p \in \{5, 7\}$ is specified.*

Proof. Let $G = PSL_3(4)$ and $p = 7$. By [7] the only maximal subgroups M of G with order divisible by 7 are isomorphic to $PSL_2(7)$. The index of $M = PSL_2(7)$ in G is 120

Table 2: Some $(2, 3)$ -generated finite simple classical groups

G	Exceptions	References
$PSL_n(q), 2 \leq n \leq 7$	$(n, q) = (2, 9), (3, 4), (4, 2)$	[26, 34, 35, 42, 41]
$PSp_n(q), 4 \leq n \leq 6$	$n = 4, q = 2^a, 3^a$	[35, 31]
$P\Omega_n(q), n = 7$		[31]
$PSU_n(q), 3 \leq n \leq 7$	$(n, q) = (3, 3), (3, 5), (4, 2), (4, 3), (5, 2)$	[34, 35, 33, 31]

Table 3: Cases when G is small and $(2, p)$ -generated

G	(n, q)	p
$PSL_n(q)$	$(2, 9), (4, 2)$	5
	$(3, 4)$	7
$PSU_n(q)$	$(4, 2), (5, 2)$	5
	$(3, 3), (3, 5), (4, 3)$	7
$P\Omega_8^+(2)$		5

and there are 3 G -conjugacy classes of such subgroups. We find $i_2(PSL_2(7)) = 21, i_2(G) = 315, i_7(PSL_2(7)) = 48, i_7(G) = 5760$. Therefore by (1.1),

$$\begin{aligned}
 Q_{2,7}(G) &\leq \sum_{M <_{\max} G} \frac{i_2(M) i_7(M)}{i_2(G) i_7(G)} \\
 &= 360 \frac{i_2(PSL_2(7)) i_7(PSL_2(7))}{i_2(G) i_7(G)} \\
 &= \frac{1}{5},
 \end{aligned}$$

and so G is $(2, 7)$ -generated. The results for the remaining groups are proved similarly. \square

Remark We note that the groups in Table 3 are not $(2, 3)$ -generated: it is elementary to prove that $PSL_2(9)$ is not $(2, 3)$ -generated; for $PSL_3(4), PSU_3(3), PSU_3(5)$, this is proved in [34]; for $PSL_4(2), PSU_4(3)$, this is proved in [35]; for $PSU_4(2) \cong PSp_4(3)$, this is proved in [24]; for $PSU_5(2)$, this is proved in [43]; and for $P\Omega_8^+(2)$, this is proved by Vsemirnov. We also note that $PSL_3(4), PSU_3(5)$ and $PSU_4(3)$ are actually $(2, 5)$ -generated; this can be seen using GAP. However, the method discussed using (1.1) fails in these cases, and so we do not prove this statement.

As discussed in the preamble to Theorem 2, in order to reduce our study of $(2, p)$ -generation of finite simple groups to the groups covered by Theorem 2 we need to prove

that the 4-dimensional symplectic groups $PSp_4(2^a)$ ($a > 1$), $PSp_4(3^a)$ are $(2, p)$ -generated for some prime p . We first state [21, Proposition 1.3], which will be useful for later results.

Proposition 2.3. *Let Y be a simple algebraic group over K , an algebraically closed field of characteristic $p > 0$, and let N be the number of positive roots in the root system of Y . Suppose that F is a Frobenius morphism of Y such that $G = (Y^F)'$ is a finite simple group of Lie type over \mathbb{F}_q . Assume G is not of type 2F_4 , 2G_2 or 2B_2 , and define $N_2 = \dim Y - N$. Then*

$$i_2(\text{Aut } G) < 2(q^{N_2} + q^{N_2-1}).$$

Lemma 2.4. *Let $G = PSp_4(q)$ where $q = 2^a$ or 3^a , $q \neq 2$. Then G is $(2, 5)$ -generated.*

Remark We note that by [35, Theorem 1.2], $PSp_4(3^{2a})$ is $(2, 5)$ -generated. However, we do not use this result; we prove Lemma 2.4 by our usual method using (1.1).

Proof. Suppose $q = 2^a$. Details on the conjugacy classes of G can be found in [10]. We find $i_2(G) = (q^2 + 1)(q^4 - 1)$, $i_5(G) \geq q^3(q - 1)(q^2 + 1)(q^2 - q + 4)$. By [4] the possible maximal subgroups with order divisible by 5 are isomorphic to $[q^3] : GL_2(q)$, $Sp_2(q) \wr S_2$, $Sp_2(q^2).2$, $SO_4^\pm(q)$, $Sz(q)$ or $Sp_4(q^{\frac{1}{t}})$ for some prime t .

If $M \cong [q^3] : GL_2(q)$, then by [10] we compute $i_2(M) = (q - 1)(q^3 + 2q^2 + q + 1)$, $i_5(M) \leq 2q^3(q + 1)(2q + 5)$. By [4] there are two G -conjugacy classes of subgroups of this type, and $|G : M| = (q + 1)(q^2 + 1)$.

If $M \cong Sp_2(q) \wr S_2$ then $i_2(M) = (i_2(Sp_2(q)) + 1)^2 - 1 + |Sp_2(q)| = q^4 + q^3 - q - 1$, $i_5(M) = (i_5(Sp_2(q)) + 1)^2 - 1 \leq 4q(q^3 + 2q^2 + 2q + 1)$. There is a single G -conjugacy class of subgroups of this type, and $|G : M| = \frac{q^2(q^2+1)}{2}$.

If $M \cong Sp_2(q^2).2$, then by Proposition 2.3 we have $i_2(M) < 2q^2(q^2 + 1)$, and we also have $i_5(M) = i_5(Sp_2(q^2)) \leq 2q^2(q^2 + 1)$. There is a single G -conjugacy class of subgroups of this type, and $|G : M| = \frac{q^2(q^2-1)}{2}$.

Suppose $M \cong Sp_4(q^{\frac{1}{t}})$ for some prime t . Then by [10] we have $i_2(M) = (q^{\frac{2}{t}} + 1)(q^{\frac{4}{t}} - 1)$, $i_5(M) \leq q^{\frac{3}{t}}(q^{\frac{1}{t}} + 1)(q^{\frac{2}{t}} + 1)(q^{\frac{1}{t}} + 4)$. There are fewer than $\log_2(q)$ G -conjugacy classes of such subgroups, each with $|G : M| = \frac{q^4(q^2-1)(q^4-1)}{q^{\frac{4}{t}}(q^{\frac{2}{t}}-1)(q^{\frac{4}{t}}-1)}$.

If $M \cong SO_4^+(q) \cong SL_2(q) \wr S_2$ then $i_2(M) = (i_2(SL_2(q)) + 1)^2 - 1 + |SL_2(q)| = q^4 + q^3 - q - 1$, $i_5(M) = (i_5(SL_2(q)) + 1)^2 - 1 \leq 4q(q^3 + 2q^2 + 2q + 1)$ as above. There is a single G -conjugacy class of subgroups of this type, and $|G : M| = \frac{q^2(q^2+1)}{2}$.

If $M \cong SO_4^-(q) \cong SL_2(q^2).2$, then by Proposition 2.3 we have $i_2(M) < 2q^2(q^2 + 1)$ and $i_5(M) = i_5(SL_2(q^2)) \leq 2q^2(q^2 + 1)$ as above. There is a single G -conjugacy class of subgroups of this type, and $|G : M| = \frac{q^2(q^2-1)}{2}$.

Finally suppose $M = Sz(q)$. Then by [40] we have $i_2(M) = (q - 1)(q^2 + 1)$, $i_5(M) \leq q^2(q + \sqrt{2q} + 1)(q - 1)$. There is a single G -conjugacy class of subgroups of this type, and $|G : M| = q^2(q + 1)(q^2 - 1)$.

Therefore by (1.1),

$$\begin{aligned}
Q_{2,5}(G) &\leq \sum_{M <_{\max} G} \frac{i_2(M) i_5(M)}{i_2(G) i_5(G)} \\
&= \left(\sum_{M \cong [q^3]:GL_2(q)} + \sum_{M \cong Sp_2(q)\wr S_2} + \sum_{M \cong Sp_2(q^2).2} + \sum_{M \cong Sp_4(q^{\frac{1}{2}})} \right. \\
&\quad \left. + \sum_{M \cong SO_4^+(q)} + \sum_{M \cong SO_4^-(q)} + \sum_{M \cong Sz(q)} \right) \frac{i_2(M) i_5(M)}{i_2(G) i_5(G)} \\
&\leq \left(4q^3(q-1)(q+1)^2(2q+5)(q^2+1)(q^3+2q^2+q+1) \right. \\
&\quad + 2q^3(q^2+1)(q^3+2q^2+2q+1)(q^4+q^3-q-1) \\
&\quad + 2q^6(q^2-1)(q^2+1)^2 \\
&\quad + 2\log_2(q)q^{\frac{7}{2}}(q^{\frac{1}{2}}+1)(q+1)(q+q^{\frac{1}{2}}+4)(q^2-1)(q^4-1) \\
&\quad + 2q^3(q^2+1)(q^3+2q^2+2q+1)(q^4+q^3-q-1) \\
&\quad + 2q^6(q^2-1)(q^2+1)^2 \\
&\quad \left. + q^4(q-1)^2(q+1)(q+\sqrt{2q}+1)(q^2-1)(q^2+1) \right) \frac{1}{q^3(q-1)(q^2+1)^2(q^2-q+4)(q^4-1)},
\end{aligned}$$

and it is straightforward to verify $Q_{2,5} < 1$ for $q \geq 8$. If $q = 4$ then a similar argument using [7] yields the result. This completes the proof for $q = 2^a$.

The argument for $q = 3^a$ is similar (using [37] instead of [10] for details on the conjugacy classes of G). \square

3. Involutions in classical groups

Let G be a finite simple classical group with natural module of dimension n defined over the field \mathbb{F}_{q^δ} , where $\delta = 2$ if G is unitary and $\delta = 1$ otherwise. In this section we find a lower bound for $i_2(G)$, the number of involutions in G .

Proposition 3.1. *The number $i_2(G)$ of involutions satisfies $i_2(G) \geq I_2(G)$, for $I_2(G)$ is given in Table 4.*

Proof. For each G we specify an involution y and calculate $|y^G|$ to give a lower bound for $i_2(G)$. The choices for y and values $|y^G|$ are listed in Table 5. It is then elementary to obtain the bounds stated in Table 4. We let $[d]$ denote an arbitrary group of order d . The notation of the elements in the table is as follows:

If q is odd define the following (projective) involutions in $GL_n(q)$:

$$s = \begin{pmatrix} iI_{\frac{n}{2}} & \\ & -iI_{\frac{n}{2}} \end{pmatrix} \text{ if } n \text{ is even, } q \equiv 1 \pmod{4} \text{ and } i^2 = -1,$$

$$t = \begin{pmatrix} & I_{\frac{n}{2}} \\ -I_{\frac{n}{2}} & \end{pmatrix} \quad \text{if } n \text{ is even,}$$

$$u_k = \begin{pmatrix} I_k & \\ & -I_{n-k} \end{pmatrix}, \quad 0 < k < n;$$

If q is even, define the following involutions in $GL_n(q)$:

$$j_k = \begin{pmatrix} I_k & & \\ & I_{n-2k} & \\ & & I_k \end{pmatrix}, \quad 0 < k \leq \frac{n}{2}.$$

In the case where q is odd, conjugacy class representatives of involutions are found in Table 5.5.1 of [12], listed alongside information on their respective centralizers.

In the case where q is even and $G = PSL_n^\epsilon(q)$, involutions are conjugate to j_k for some k . The structures of the centralizers of j_k are found in Sections 4 and 6 of [3], and exact conjugacy class sizes can be calculated using [23, Theorem 7.1]. If q is even and G is symplectic or orthogonal, involutions are of the form a_k, b_k or c_k with Jordan normal form j_k by [3]. In Sections 7 and 8 of the same paper the structure of the centralizer of each involution is given, and exact conjugacy class sizes can be calculated using [23, Theorem 7.2]. \square

Table 4: Lower bounds for $i_2(G)$

G	$I_2(G)$
$PSL_n^\epsilon(q)$	$\frac{1}{8}q^{\lfloor \frac{n^2}{2} \rfloor}$
$PSp_n(q)$	$\frac{1}{2}q^{\frac{n^2}{4} + \frac{n}{2}}$
$P\Omega_n^\epsilon(q)$	$\frac{1}{8}q^{\frac{n^2}{4} - 1}$
$P\Omega_n(q), nq \text{ odd}$	$\frac{1}{2}q^{\frac{n^2-1}{4}}$

Table 5: Values for $|y^G|$

G	Conditions	y	$ y^G $
$PSL_n^\epsilon(q), n \geq 2$	n even, q even	$j_{\frac{n}{2}}$	$ GL_n^\epsilon(q) : [q^{\frac{n^2}{4}}].GL_{\frac{n}{2}}^\epsilon(q) $
	n even, $q \equiv \epsilon \pmod{4}$	s	$ GL_n^\epsilon(q) : GL_{\frac{n}{2}}^\epsilon(q)^2.2 $
	n even, $q \equiv -\epsilon \pmod{4}$	t	$ GL_n^\epsilon(q) : GL_{\frac{n}{2}}(q^2).2 $
	n odd, q even	$j_{\frac{n-1}{2}}$	$ GL_n^\epsilon(q) : [q^{\frac{n^2+2n-7}{4}}].(GL_{\frac{n-1}{2}}^\epsilon(q) \times GL_1^\epsilon(q)) $
	n odd, q odd	$(-1)^{\frac{n+1}{2}}u_{\frac{n-1}{2}}$	$ GL_n^\epsilon(q) : GL_{\frac{n-1}{2}}^\epsilon(q) \times GL_{\frac{n+1}{2}}^\epsilon(q) $

$PSp_n(q), n \geq 4$	$\frac{n}{2}$ even, q even	$c_{\frac{n}{2}}$	$ Sp_n(q) : [q^{\frac{1}{2}(\frac{n^2}{4} + \frac{3n}{2} - 2)}]. Sp_{\frac{n}{2}-2}(q) $
	$\frac{n}{2}$ odd, q even	$b_{\frac{n}{2}}$	$ Sp_n(q) : [q^{\frac{n}{2}(\frac{n}{2} + 1)}]. Sp_{\frac{n}{2}-1}(q) $
	$q \equiv 1 \pmod{4}$	s	$ Sp_n(q) : GL_{\frac{n}{2}}(q).2 $
	$q \equiv 3 \pmod{4}$	t	$ Sp_n(q) : GU_{\frac{n}{2}}(q).2 $
$P\Omega_n^+(q), n \geq 8$	$\frac{n}{2}$ even, q even	$c_{\frac{n}{2}}$	$ \Omega_n^+(q) : [q^{\frac{1}{2}(\frac{n^2}{4} + \frac{n}{2} - 2)}]. Sp_{\frac{n}{2}-2}(q) \cap \Omega_n^+(q) $
	$\frac{n}{2}$ even, $q^{\frac{n}{4}} \equiv \epsilon \pmod{4}$	$u_{\frac{n}{2}}$	$ \Omega_n^+(q) : O_{\frac{n}{2}}^\epsilon(q)^2.2 \cap \Omega_n^+(q) $
	$\frac{n}{2}$ odd, q even	$c_{\frac{n}{2}-1}$	$ \Omega_n^+(q) : [q^{\frac{1}{2}(\frac{n^2}{4} + \frac{3n}{2} - 10)}]. (Sp_{\frac{n}{2}-3}(q) \times Sp_2(q)) \cap \Omega_n^+(q) $
	$\frac{n}{2}$ odd, q odd	$u_{\frac{n}{2}-1}$	$ \Omega_n^+(q) : O_{\frac{n}{2}-1}^+(q) \times O_{\frac{n}{2}+1}^+(q) \cap \Omega_n^+(q) $
$P\Omega_n^-(q), n \geq 8$	$\frac{n}{2}$ even, q even	$c_{\frac{n}{2}}$	$ \Omega_n^-(q) : [q^{\frac{1}{2}(\frac{n^2}{4} + \frac{n}{2} - 2)}]. Sp_{\frac{n}{2}-2}(q) \cap \Omega_n^-(q) $
	$\frac{n}{2}$ even, q odd	$u_{\frac{n}{2}}$	$ \Omega_n^-(q) : (O_{\frac{n}{2}}^+(q) \times O_{\frac{n}{2}}^-(q)).2 \cap \Omega_n^-(q) $
	$\frac{n}{2}$ odd, q even	$c_{\frac{n}{2}-1}$	$ \Omega_n^-(q) : [q^{\frac{1}{2}(\frac{n^2}{4} + \frac{3n}{2} - 10)}]. (Sp_{\frac{n}{2}-3}(q) \times Sp_2(q)) \cap \Omega_n^-(q) $
	$\frac{n}{2}$ odd, $q^{\frac{n-2}{4}} \equiv \epsilon \pmod{4}$	$u_{\frac{n}{2}-\epsilon}$	$ \Omega_n^-(q) : O_{\frac{n}{2}-1}^\epsilon(q) \times O_{\frac{n}{2}+1}^{-\epsilon}(q) \cap \Omega_n^-(q) $
$P\Omega_n(q)$ (nq odd), $n \geq 7$	$q^{\lfloor \frac{n+1}{4} \rfloor} \equiv \epsilon \pmod{4}$	$(-1)^{\frac{n+1}{2}} u_{\frac{n-1}{2}}$	$ \Omega_n(q) : O_{2\lfloor \frac{n+1}{4} \rfloor}^\epsilon(q) \times O_{n-2\lfloor \frac{n+1}{4} \rfloor}(q) \cap \Omega_n(q) $

4. Maximal subgroups with order divisible by r

Let G be a finite simple classical group described in Theorem 2 – that is, with natural module $n \geq 8$ and $G \neq P\Omega_8^+(2)$. If $G = P\Omega_n^\epsilon(q)$, let $|PSO_n^\epsilon(q) : P\Omega_n^\epsilon(q)| = a_\epsilon \in \{1, 2\}$ and let $|Z(\Omega_n^\epsilon(q))| = z_\epsilon \in \{1, 2\}$. The values a_ϵ, z_ϵ can be found in [19, § 2]. Recall the definition of r , a primitive prime divisor of $q^e - 1$ for e listed in Table 1.

The subgroup structure of G is well-understood due to a theorem of Aschbacher [1]. The theorem states that if M is a maximal subgroup of G , then M lies in a natural collection $\mathcal{C}_1, \dots, \mathcal{C}_8$, or $M \in \mathcal{S}$. Subgroups in \mathcal{C}_i are described in detail in [19], where the structure and number of conjugacy classes are given. Subgroups in class \mathcal{S} are almost simple groups which act absolutely irreducibly on the natural module V of G . The orders of subgroups in classes $\mathcal{C}_1, \dots, \mathcal{C}_8$ can easily be computed using [19, § 4], and this yields the result below.

Proposition 4.1. *Let M be a maximal subgroup of G with order divisible by r , and assume M lies in one of the Aschbacher classes $\mathcal{C}_1, \dots, \mathcal{C}_8$. Then M is conjugate to a subgroup listed in Table 6. The number of G -conjugacy classes of each type, c_M , is also listed.*

Table 6 also lists bounds $I_2(M)$ and N_M , where $i_2(M) \leq I_2(M)$ and $N_M \leq |N_M(\langle x \rangle)|$ where $x \in G$ is an element of order r contained in M . These bounds are justified in Propositions 5.1 and 6.4 below.

The subgroups $M \in \mathcal{S}$ with order divisible by r are given by [15]. We first list these subgroups with $\text{soc}(M) \in \text{Lie}(p')$, where $\text{Lie}(p')$ is the set of finite simple groups of Lie type with characteristic not equal to p .

Table 6: Maximal subgroups $M \notin \mathcal{S}$ with $r \mid |M|$

G	Class	Type of M	Conditions	c_M	$I_2(M)$	N_M
$PSL_n(q), n \geq 2$	\mathcal{C}_3	$GL_k(q^t).t$	$n = kt, t$ prime	1	$2\frac{q^{2t}-1}{q-1}q^{\frac{n^2}{2t}+\frac{n}{2}-2t}$	$ N_G(\langle x \rangle) $
	\mathcal{C}_6	$2^{2k}Sp_{2k}(2)$	$n = 2^k, r = 2^k + 1, q$ odd	$(q-1, n)$	$2^{\log_2(n)(2\log_2(n)+3)}$	r
	\mathcal{C}_8	$PSp_n(q)$	n even	$(q-1, \frac{n}{2})$	$2(q+1)q^{\frac{n^2+2n-4}{4}}$	$\frac{n(q^{\frac{n}{2}}+1)}{(2, q-1)}$
		$PSO_n^-(q)$	n even, q odd	$\frac{(q-1, n)}{2}$	$2(q+1)q^{\frac{n^2-4}{4}}$	$\frac{n(q^{\frac{n}{2}}+1)}{2a_-}$
		$PSU_n(q^{\frac{1}{2}})$	n odd, q square	$\frac{q-1}{[q^{\frac{1}{2}}+1, \frac{q-1}{(q-1, n)}]}$	$2(q^{\frac{1}{2}}+1)q^{\frac{n^2+n-4}{4}}$	$\frac{n(q^{\frac{n}{2}}+1)}{(q^{\frac{1}{2}}+1)(n, q^{\frac{1}{2}}+1)}$
$PSp_n(q), n \geq 4$	\mathcal{C}_3	$Sp_k(q^t).t$	$n = kt, k$ even, t prime	1	$2(q^t+1)q^{\frac{n^2}{4t}+\frac{n}{2}-t}$	$ N_G(\langle x \rangle) $
		$GU_{\frac{n}{2}}(q).2$	$\frac{n}{2}$ odd, q odd	1	$(q+1)^2q^{\frac{n^2+2n-16}{8}}$	$ N_G(\langle x \rangle) $
	\mathcal{C}_6	$2^{2k}O_{2k}^-(2)$	$n = 2^k, q = p$ odd, $r = n + 1$	1 or 2	$2^{2\log_2^2(n)+\log_2(n)+1}$	r
	\mathcal{C}_8	$PSO_n^-(q)$	q even	1	$2(q+1)q^{\frac{n^2-4}{4}}$	$\frac{n(q^{\frac{n}{2}}+1)}{2}$
$P\Omega_n^+(q), n \geq 8$	\mathcal{C}_1	$O_{n-2}^-(q) \times O_2^-(q)$		1	$2(q+1)^2q^{\frac{n^2-4n}{4}}$	$ N_G(\langle x \rangle) $
		$O_{n-1}(q) \times O_1(q)$	q odd	2	$\frac{4}{z_+}(q+1)q^{\frac{n^2-2n-4}{4}}$	$\frac{(n-2)(q^{\frac{n}{2}-1}+1)}{a_+}$
		$O_{n-1}(q)$	q even	1	$2(q+1)q^{\frac{n^2-2n-4}{4}}$	$\frac{(n-2)(q^{\frac{n}{2}-1}+1)}{2}$
	\mathcal{C}_2	$O_1(q)$ wr S_n	$q = p$ odd, $r = n - 1$	2 or 4	$2^{n-1}n!$	r
	\mathcal{C}_3	$O_{\frac{n}{2}}(q^2).2$	$\frac{n}{2}$ odd, q odd	1 or 2	$\frac{4}{z_+}(q+1)q^{\frac{n^2-20}{8}}$	$\frac{(n-2)(q^{\frac{n}{2}-1}+1)}{4a_+}$
		$GU_{\frac{n}{2}}(q).2$	$\frac{n}{2}$ even	2	$\frac{2}{z_+}(q+1)^2q^{\frac{n^2+2n-16}{8}}$	$ N_G(\langle x \rangle) $
	\mathcal{C}_6	$2^{2k}O_{2k}^+(2)$	$n = 2^k, q = p$ odd, $r = n - 1$	4 or 8	$2^{\log_2(n)(2\log_2(n)+1)}$	r
	$P\Omega_n^-(q), n \geq 8$	\mathcal{C}_3	$O_k^-(q^t).t$	$n = kt, t$ prime, $k \geq 4$	1	$2(q^t+1)q^{\frac{n^2}{4t}-t}$
$GU_{\frac{n}{2}}(q).2$			$\frac{n}{2}$ odd	1	$\frac{2}{z_-}(q+1)^2q^{\frac{n^2+2n-16}{8}}$	$ N_G(\langle x \rangle) $
$P\Omega_n(q), n \geq 7, nq$ odd	\mathcal{C}_1	$O_{n-1}^-(q) \times O_1(q)$		1	$4(q+1)q^{\frac{n^2-2n-3}{4}}$	$ N_G(\langle x \rangle) $
	\mathcal{C}_2	$O_1(q)$ wr S_n	$q = p, r = n$	1 or 2	$2^{n-1}n!$	r
$PSU_n(q), n \geq 3, n$ odd	\mathcal{C}_3	$GU_k(q^t).t$	$n = kt, t$ prime, $t \geq 3$	1	$\frac{2(q^t+1)^2}{q+1}q^{\frac{n^2}{2t}+\frac{n}{2}-2t}$	$ N_G(\langle x \rangle) $
$PSU_n(q), n \geq 4, n$ even	\mathcal{C}_1	$GU_{n-1}(q) \times GU_1(q)$		1	$2(q+1)q^{\frac{n^2-n-4}{2}}$	$ N_G(\langle x \rangle) $

To state the next result, let

$$e_G = \begin{cases} (n, q - \epsilon) & \text{if } G = PSL_n^\epsilon(q) \\ (2, q - 1) & \text{if } G = PSp_n(q) \\ a_\epsilon(2, q - 1)^2 & \text{if } G = P\Omega_n^\epsilon(q) \text{ (} n \text{ even)} \\ 2 & \text{if } G = P\Omega_n(q) \text{ (} nq \text{ odd)}. \end{cases}$$

Then e_G is the index of G in the projective similarity group of the same type (see [19, § 2]).

Proposition 4.2. *Let M be a maximal subgroup of G with order divisible by r , and assume $M \in \mathcal{S}$ with $\text{soc}(M) \in \text{Lie}(p')$. The possibilities for $\text{soc}(M)$ are listed in Table 7. Upper bounds C_M are given for the number of G -conjugacy classes of maximal subgroups with each listed socle.*

Proof. Let $M \in \mathcal{S}$ with $\text{soc}(M) \in \text{Lie}(p')$, and let V be the natural module of G . The possibilities for $\text{soc}(M)$ are determined in [15], and we list them in Table 7 alongside necessary conditions.

It remains to justify the bound C_M . Consider $\text{soc}(M) = S = PSL_d(s)$, $d \geq 3$. Here $\dim V = n = \frac{s^d - 1}{s - 1} - \delta$ where $\delta \in \{0, 1\}$. By [16, Theorem 1.1], in most cases the representation of S on V is one of at most $s - 1$ Weil representations. The exceptions are when $d = 3$, $s = 2$ or 4, and in each case it can be verified that the number of possible representations is again at most $s - 1$ using [7] and [18]. Hence the number of $PGL(V)$ -classes of such subgroups M is at most $s - 1$. If $G = PSL_n(q)$, it follows that that number of G -classes of subgroups M is at most $(s - 1)|PGL_n(q) : PSL_n(q)| = (s - 1)e_G$. If $G = P\Omega_n^\epsilon(q)$, Corollary 2.10.4 of [19] shows that the number of $PGO_n^\epsilon(q)$ -classes of subgroups M is also at most $s - 1$, and so the number of G -classes is at most $(s - 1)|PGO_n^\epsilon(q) : P\Omega_n^\epsilon(q)| = (s - 1)e_G$.

The remaining possibilities for $\text{soc}(M)$ and G are dealt with similarly, using [14], [17], [5] for $\text{soc}(M) = PSp_{2d}(s)$, $PSU_d(s)$, $PSL_2(s)$ respectively. \square

Table 7: Maximal subgroups $M \in \mathcal{S}$ with $\text{soc}(M) \in \text{Lie}(p')$, $r \mid |M|$

G	$\text{soc}(M)$	Conditions	C_M/e_G
$PSL_n(q), PSp_n(q),$	$PSL_d(s)$	$d \geq 3$ prime, $n = \frac{s^d - 1}{s - 1} - 1, r = n + 1$	$s - 1$
$P\Omega_n^-(q)$	$PSp_{2d}(s)$	$s \neq 3$ odd, $d = 2^b \geq 2, n = \frac{1}{2}(s^d - 1), r = n + 1$	4
	$PSU_d(s)$	d prime, $n = \frac{s^d + 1}{s + 1} - 1, r = n + 1$	$s + 1$
	$PSL_2(n)$	$n = 2^b, b = 2^{b'}, r = n + 1$	1
	$PSL_2(n + 1)$	$r = n + 1$	$\frac{n}{4}$
	$PSL_2(2n + 1)$	$r = n + 1$ or $2n + 1$	2
$P\Omega_n^+(q)$	$PSp_{2d}(3)$	$n = \frac{3^d + 1}{2}$ odd, $d \geq 3$ prime, $r = n - 1$	4
	$PSL_2(n - 1)$	$r = n - 1$	$\frac{n - 4}{4}$
	$PSL_2(n)$	$n = 2^b, b$ prime, $r = n - 1$	1
	$PSL_2(2n - 1)$	$r = n - 1$	2

$P\Omega_n(q), nq$ odd	$PSL_d(s)$	$d \geq 3$ prime, $n = \frac{s^d-1}{s-1}, r = n$	$s - 1$
	$PSp_{2d}(s)$	$s \neq 3$ odd, $d = 2^b \geq 2, n = \frac{1}{2}(s^d + 1), r = n$	4
	$PSp_{2d}(3)$	d an odd prime, $n = \frac{1}{2}(3^d - 1), r = n$	4
	$PSU_d(s)$	d prime, $n = \frac{s^d+1}{s+1}, r = n$	$s + 1$
	$PSL_2(n-1)$	$r = n$	$\frac{n-3}{2}$
	$PSL_2(n)$	$n = 2^b, b = 2^{b'}, r = n$	1
	$PSL_2(n+1)$	$r = n$	$\frac{n+1}{2}$
	$PSL_2(2n-1)$	$r = n$ or $2n-1$	2
	$PSL_2(2n+1)$	$r = n$	2
$PSU_n(q), n$ odd	$PSL_2(2n+1)$	$r = 2n+1$	2
$PSU_n(q), n$ even	$PSL_2(2n-1)$	$r = 2n-1$	2

We denote by $c_{\mathcal{S}}$ the number of G -classes of subgroups $M \in \mathcal{S}$ with order divisible by r and $\text{soc}(M)$ not an alternating group.

Corollary 4.3. *We have $c_{\mathcal{S}} \leq C_{\mathcal{S}}$, where $C_{\mathcal{S}}$ is given in Table 8 for each G .*

Table 8: Upper bounds for $c_{\mathcal{S}}$

G	$C_{\mathcal{S}}$
$PSL_n(q), PSp_n(q), P\Omega_n^-(q), n \geq 7$	$(n^2 + \frac{21}{4}n - 1)e_G$
$P\Omega_n^+(q), n \geq 10$	$(\frac{1}{4}n + 9)e_G$
$P\Omega_n(q), n \geq 9, nq$ odd	$(n^2 + 6n + 4)e_G$
$PSU_n(q), n \geq 7$	$3e_G$

Proof. Let $M \in \mathcal{S}$ with $\text{soc}(M)$ not an alternating group. If $\text{soc}(M) \notin \text{Lie}(p')$, the number c of $GL_n(q)$ -conjugacy classes of such M is given in [15, Example 2.7]. By [19, Corollary 2.10.4], the number of G -classes is therefore bounded above by ce_G . If $\text{soc}(M) \in \text{Lie}(p')$, the number of G -classes of M is bounded above by C_M in Proposition 4.2.

Suppose $G = PSL_n(q)$. In this case $c \leq 6$ by [15], and so the contribution to $c_{\mathcal{S}}$ from $M \in \mathcal{S}, \text{soc}(M) \notin \text{Lie}(p')$ is at most $6e_G = 6(n, q - 1)$. The contribution to $c_{\mathcal{S}}$ from $M \in \mathcal{S}$ with $\text{soc}(M) \in \text{Lie}(p')$ is as follows using Proposition 4.2:

- For $\text{soc}(M) = PSL_d(s), d \geq 3$ we have $s \leq n$. For each s there exists at most one d such that $n = \frac{s^d-1}{s-1} - 1$ as in Table 7. Therefore, by Proposition 4.2, the contribution is less than $\sum_{s=2}^n (s-1)e_G = \frac{1}{2}n(n-1)e_G$;

- For $\text{soc}(M) = PSp_{2d}(s)$, $d \geq 2$ we have $s \leq n$ and $s \neq 3$, and since for each s there exists at most one d as above, the contribution is less than $4(n-2)e_G$;
- For $\text{soc}(M) = PSU_d(s)$ we also have $s \leq n$, and, similar to above, the contribution is less than $\sum_{s=2}^n (s+1)e_G = \frac{1}{2}(n-1)(n+4)e_G$;
- For $\text{soc}(M) = PSL_2(s)$ we have $s = n, n+1$ or $2n+1$, contributing $e_G, \frac{1}{4}ne_G, 2e_G$ respectively.

Summing these contributions yields $C_{\mathcal{S}} = (n^2 + \frac{21}{4}n - 1)e_G$. The remaining bounds are found in a similar manner. \square

5. Involutions in maximal subgroups

Let G be a finite simple classical group described in Theorem 2. Recall our definition of r as a primitive prime divisor of $q^e - 1$ for e listed in Table 1.

In this section we deduce upper bounds for the number of involutions in maximal subgroups of G with order divisible by r .

Proposition 5.1. *Let $M <_{\max} G$ with $r \mid |M|$.*

- i) If $M \notin \mathcal{S}$, then $i_2(M) \leq I_2(M)$, where $I_2(M)$ is listed in Table 6;*
- ii) If $M \in \mathcal{S}$ and $\text{soc}(M)$ is not an alternating group then $i_2(M) < q^{2n+4}$;*
- iii) If $M \in \mathcal{S}$ and $\text{soc}(M) = A_{n'}$ for some $n' \geq 9$, then $n' = n+1$ or $n+2$ and $i_2(M) < (n+2)!$.*

Proof. We prove this case by case for each Aschbacher class. Subgroups $M <_{\max} G$ with $M \notin \mathcal{S}$ and $r \mid |M|$ are listed in Table 6.

First suppose $M \in \mathcal{C}_1$. Then M is of type $Cl_{n-k}(q) \times Cl_k(q)$ for some classical group $Cl_k(q)$ of the same type as G and $k = 1$ or 2 . We can bound the number of involutions in each factor using Proposition 2.3. If $G = P\Omega_n^+(q)$ and M is of type $O_{n-2}^-(q) \times O_2^-(q)$ we have $i_2(M) \leq i_2(PO_{n-2}^-(q))i_2(PO_2^-(q))$. By Proposition 2.3, $i_2(PO_{n-2}^-(q)) \leq i_2(\text{Aut}(P\Omega_{n-2}^-(q))) < 2(q+1)q^{\frac{n^2-4n}{4}}$. Using the fact that $O_2^-(q) \cong D_{2(q+1)}$ gives $i_2(PO_2^-(q)) \leq q+1$, yielding the result. The other values for $I_2(M)$ are calculated in a similar fashion.

Next suppose $M \in \mathcal{C}_2, \mathcal{C}_6$ or \mathcal{S} . In most cases we bound $i_2(M)$ by $|M|$. For $M \in \mathcal{S}$ with $\text{soc}(M) \neq A_{n'}$ we have $|M| < q^{3n\delta}$ by [22, Theorem 4.1]. In fact, using the fact that $r \mid |M|$, by Theorem 4.2 of the same paper we have $|M| < q^{(2n+4)\delta}$ unless $(G, \text{soc}(M)) = (PSL_{11}(2), M_{24}), (P\Omega_8^+(q), P\Omega_7(q))$. In the first case, $i_2(M_{24}) = 43263 < 2^{24}$ by [7], and in the second case, $i_2(M) < 2(q+1)q^{11} < q^{20}$ by Proposition 2.3. If G is unitary then M has socle $PSL_2(2n \pm 1)$ or J_3 by [15], and each has $i_2(M) < q^{2n+4}$, as can be seen from their respective character tables. If $\text{soc}(M) = A_{n'}$, $n' \geq 9$ then by [15] we have $n' = n+1$ or $n+2$. Therefore $i_2(M) \leq i_2(S_{n+2}) < (n+2)!$.

If $M \in \mathcal{C}_3$ then M is of type $Cl_k(q^t).t$ for some classical group $Cl_k(q^t)$. Suppose $G = PSL_n(q)$ and let M be of type $GL_k(q^t).t$. If $k \geq 2$ we have $i_2(M) \leq \frac{q^t-1}{q-1}i_2(\text{Aut}(PSL_k(q^t)))$, and applying Proposition 2.3 yields the result. If $k = 1$, M is of type $GL_1(q^n).n$ where n is prime, and since we are assuming $n > 2$ we have $i_2(M) = 0$. Other bounds $I_2(M)$ for $M \in \mathcal{C}_3$ are calculated in a similar fashion.

If $M \in \mathcal{C}_8$ then $\text{soc}(M)$ is a finite simple classical group, and so we can apply Proposition 2.3 to obtain the result. \square

6. Proof of Theorem 2 for n sufficiently large

Let G be a finite simple classical group described in Theorem 2. If $G = P\Omega_n^\epsilon(q)$ recall the definition of $a_\epsilon = |PSO_n^\epsilon(q) : P\Omega_n^\epsilon(q)|$. Also, recall that r is a primitive prime divisor of $q^e - 1$, where e is listed in Table 1. Let $x \in G$ be an element of order r .

In this section, except for a small number of possible exceptions (given in Proposition 6.5) we prove Theorem 2 holds in the following cases:

$$\begin{aligned} &PSL_n(q), n \geq 9 \\ &PSp_n(q), n \geq 12 \\ &P\Omega_n^\epsilon(q) \text{ (} n \text{ even), } n \geq 14 \\ &P\Omega_n(q) \text{ (} nq \text{ odd), } n \geq 13 \\ &PSU_n(q), n \geq 8. \end{aligned}$$

We first require a result on the number of conjugates of a maximal subgroup containing x .

Lemma 6.1. *If x lies in two conjugate maximal subgroups of G , say M and M^g for some $g \in G$, then $mg \in N_G(\langle x \rangle)$ for some $m \in M$.*

Proof. We first note that Sylow r -subgroups of G are cyclic (this follows from [12, Theorem 4.10.2]).

Suppose $x \in M \cap M^g$ where M is a maximal subgroup of G and $g \in G$, so $x, x^{g^{-1}} \in M$. We have $\langle x \rangle, \langle x^{g^{-1}} \rangle$ contained in Sylow r -subgroups of M , conjugate by some $m \in M$. As the Sylow r -subgroups are cyclic, $\langle x \rangle$ and $\langle x^{g^{-1}} \rangle$ are also conjugate by m . This implies that $mg \in N_G(\langle x \rangle)$. \square

Corollary 6.2. *If x lies in a maximal subgroup M of G , the number of G -conjugates of M containing x is $\frac{|N_G(\langle x \rangle)|}{|N_M(\langle x \rangle)|}$.*

Proof. Let $C = \{M^g : g \in G, x \in M^g\}$. Then $N_G(\langle x \rangle)$ acts on C by conjugation, and Lemma 6.1 implies this action is transitive. The stabilizer of M is $N_G(\langle x \rangle) \cap N_G(M) = N_M(\langle x \rangle)$ since M is maximal. Therefore $|C| = \frac{|N_G(\langle x \rangle)|}{|N_M(\langle x \rangle)|}$, completing the proof. \square

Lemma 6.3. *Let $T \leq G$ be a maximal torus containing x . Then $C_G(x) = T$.*

Proof. Let H be a simple algebraic group over $\overline{\mathbb{F}}_q$ such that $(H^F)' = G$ for a Steinberg morphism F . Also let $\mathbb{F}_{q^e}^\times = \langle \lambda \rangle$. If $\mu \in \mathbb{F}_{q^e}$ is an eigenvalue of x , then its Galois-conjugates $\mu^{q^\delta}, \mu^{q^{2\delta}}, \dots$ are also eigenvalues of x , where $\delta = 2$ if G is unitary and $\delta = 1$ otherwise. Therefore, as x is semisimple, x is conjugate in H to $\text{diag}(\lambda^a, \lambda^{aq^\delta}, \lambda^{aq^{2\delta}}, \dots, \lambda^{aq^{e-\delta}}, I_k)$, where a is such that x has order r , and $k \leq 2$ with equality if and only if $G = P\Omega_n^+(q)$. We find that $C_H(x)$ is contained in a unique maximal torus containing x : if $k \leq 1$ then x has distinct eigenvalues and the statement is clear, otherwise $k = 2, G = P\Omega_n^+(q)$, and the 1-space is a 2-dimensional torus of type $O_2^+(q)$. Therefore, taking fixed points yields the result. \square

Proposition 6.4. *Let $M <_{\max} G$ with $x \in M$, and assume $M \notin \mathcal{S}$. Then $|N_M(\langle x \rangle)| \geq N_M$, where N_M is listed in Table 6. The values of $|N_G(\langle x \rangle)|$ are listed in Table 9.*

Table 9: Orders of normalizers of x

G	$ N_G(\langle x \rangle) $
$PSL_n(q)$	$\frac{n(q^n-1)}{(q-1)(n,q-1)}$
$PSp_n(q)$	$\frac{n(q^{\frac{n}{2}}+1)}{(2,q-1)}$
$P\Omega_n^+(q)$	$\frac{(n-2)(q^{\frac{n}{2}-1}+1)(q+1)}{a_+(2,q-1)^2}$
$P\Omega_n^-(q)$	$\frac{n(q^{\frac{n}{2}}+1)}{a_-(2,q-1)}$
$P\Omega_n(q)$	$\frac{(n-1)(q^{\frac{n-1}{2}}+1)}{2}$
$PSU_n(q), n$ odd	$\frac{n(q^n+1)}{(q+1)(n,q+1)}$
$PSU_n(q), n$ even	$\frac{(n-1)(q^{n-1}+1)}{(n,q+1)}$

Proof. We first consider $N_G(\langle x \rangle)$. By Lemma 6.3 there is a unique maximal torus T of G containing x . We have $N_G(\langle x \rangle) \leq N_G(T)$ by Lemma 6.3, and conversely the normalizer of T will normalize the unique Sylow r -subgroup of T containing x , yielding $N_G(\langle x \rangle) = N_G(T)$. By [28, § 25], T corresponds to an element w in the Weyl group W of G , and Proposition 25.3 in the same section gives $|C_W(w)| = |N_G(T) : T|$. The values of $|T|$ and $|N_G(T) : T|$ can be computed from [6]. Hence $|N_G(\langle x \rangle)|$ as in Table 9.

We now consider $N_M(\langle x \rangle)$, and proceed case by case for each Aschbacher class of subgroups M containing x with $r \mid |M|$. Such classes are listed in Table 6.

Consider $M \in \mathcal{C}_1$ of type $Cl_{n-k}(q) \times Cl_k(q)$ for some classical group $Cl_m(q)$ of the same type as G with $k \leq 2$. Then $N_M(\langle x \rangle)$ is of the form $N_{Cl_{n-k}(q)}(\langle x \rangle) \times Cl_k(q)$. If the maximal torus in $Cl_{n-k}(q)$ containing x is T' , then, by the same reasoning as for $N_G(\langle x \rangle)$ in first paragraph, the normalizer of x in M is of the form $N_{Cl_{n-k}(q)}(T') \times Cl_k(q)$, and using [6] yields the result.

For $M \in \mathcal{C}_2$ or \mathcal{C}_6 we use the obvious bound $|N_M(\langle x \rangle)| \geq r$.

Consider $M \in \mathcal{C}_3$ of the form $M_0.t$ where $M_0 = Cl_k(q^t)$, a classical group of the same type as G . Then $T \leq M_0$, and $N_{M_0}(\langle x \rangle) = N_{M_0}(T)$ as above. It remains to consider $\frac{M}{M_0}$. By Lemma 6.3, $C_G(x) = T$, and so there is a unique Sylow r -subgroup P of M containing x . Therefore $N_M(\langle x \rangle) = N_M(P)$. By the Frattini argument, $M = M_0 N_M(P)$, and so

$$\frac{N_M(\langle x \rangle)}{N_{M_0}(\langle x \rangle)} = \frac{M_0 N_M(\langle x \rangle)}{M_0} = \frac{M_0 N_M(P)}{M_0} = \frac{M}{M_0} = C_t.$$

Hence $N_M(\langle x \rangle) = N_{M_0}(\langle x \rangle).t$, showing that for $M \in \mathcal{C}_3$ we have $N_G(\langle x \rangle) = N_M(\langle x \rangle)$. For the remaining cases $M \in \mathcal{C}_3$ a similar method yields the result.

For $M \in \mathcal{C}_8$, we use the same method as we did for $N_G(\langle x \rangle)$ in the first paragraph to obtain the lower bound N_M . \square

Proposition 6.5. *Apart from 4 possible exceptions, Theorem 2 holds for G in the following cases:*

$$\begin{aligned} &PSL_n(q), n \geq 9; \\ &PSp_n(q), n \geq 12; \\ &P\Omega_n^\epsilon(q) \text{ (} n \text{ even), } n \geq 14; \\ &P\Omega_n(q) \text{ (} nq \text{ odd), } n \geq 13; \\ &PSU_n(q), n \geq 8. \end{aligned}$$

The possible exceptions are $G = PSp_{12}(2), P\Omega_{14}^+(2), P\Omega_{16}^+(2), P\Omega_{18}^+(2)$.

Proof. To prove G is $(2, r)$ -generated, by (1.2) it suffices to prove $\sum_{x \in M <_{\max G}} \frac{i_2(M)}{i_2(G)} < 1$.

Let μ_i be a set of G -conjugacy class representatives of maximal subgroups $M \in \mathcal{C}_i$ with order divisible by r , and μ_0 be a set of G -conjugacy class representatives for $M \in \mathcal{S}$ with order divisible by r . By Corollary 6.2 we have

$$\sum_{x \in M <_{\max G}} \frac{i_2(M)}{i_2(G)} = \Sigma_1 + \Sigma_2 + \cdots + \Sigma_8 + \Sigma_0$$

where

$$\Sigma_i = \sum_{M \in \mu_i} \frac{|N_G(\langle x \rangle)|}{|N_M(\langle x \rangle)|} \frac{i_2(M)}{i_2(G)}.$$

The values $|\mu_i|$ and maximal subgroups M contributing to Σ_i , $1 \leq i \leq 8$ are found in Table 6 by Proposition 4.1. Upper bounds for the number of conjugacy class representatives $M \in \mu_0$ with $\text{soc}(M) \notin \{A_{n+1}, A_{n+2}\}$ are found in Table 8 by Corollary 4.3. For $M \in \mu_0$ with $\text{soc}(M) \in \{A_{n+1}, A_{n+2}\}$, $\text{soc}(M)$ has a unique irreducible n -dimensional representation over any field (where the characteristic $p \mid n+2$ in the $n+2$ case), preserving an orthogonal (or symplectic in characteristic 2) form (see [19, § 5]). Hence the number of such representations is 0 if $G = PSL_n^\epsilon(q)$, and is at most e_G otherwise.

By Proposition 6.4, if $M \notin \mathcal{S}$ then $\frac{|N_G(\langle x \rangle)|}{|N_M(\langle x \rangle)|} \leq \frac{|N_G(\langle x \rangle)|}{N_M}$ where N_M is found in Table 6 and $|N_G(\langle x \rangle)|$ is found in Table 9. If $M \in \mathcal{S}$ we use $|N_M(\langle x \rangle)| \geq r$ if $\text{soc}(M) \notin \{A_{n+1}, A_{n+2}\}$,

and $|N_M(\langle x \rangle)| \geq \frac{1}{2}r(r-1)$ otherwise. The ratio $\frac{i_2(M)}{i_2(G)}$ is bounded by $\frac{I_2(M)}{I_2(G)}$ by Proposition 3.1 and Proposition 5.1. This leads to an upper bound for each Σ_i which can be manipulated into a decreasing function in n and q .

As an illustration, consider $G = P\Omega_n^-(q)$. By Table 6,

$$\sum_{x \in M <_{\max} G} \frac{i_2(M)}{i_2(G)} = \Sigma_3 + \Sigma_0.$$

Consider first Σ_3 . In \mathcal{C}_3 there are fewer than $\frac{n}{2}$ classes of type $O_k^-(q^t).t$ and there is 1 class of type $GU_{\frac{n}{2}}(q).2$. By Corollary 6.2 and Table 6, if $M \in \mu_3$ then M is the unique conjugate containing x since $|N_G(\langle x \rangle)| = |N_M(\langle x \rangle)|$. By Table 6, for M of type $O_k^-(q^t).t$ we have $i_2(M) < 2(q^t + 1)q^{\frac{n^2}{4t} - t} \leq 2(q^2 + 1)q^{\frac{n^2}{8} - 2}$. For M of type $GU_{\frac{n}{2}}(q).2$ we have $i_2(M) < \frac{2}{z_-}(q+1)^2q^{\frac{n^2}{8} + \frac{n}{4} - 2}$. By Table 4, we have $i_2(G) \geq \frac{1}{8}q^{\frac{n^2}{4} - 1}$. Therefore

$$\begin{aligned} \Sigma_3 &= \sum_{x \in M \in \mathcal{C}_3} \frac{i_2(M)}{i_2(G)} \\ &\leq \sum_{M \in \mu_3} \frac{|N_G(\langle x \rangle)|}{|N_M(\langle x \rangle)|} \frac{i_2(M)}{i_2(G)} \\ &\leq \frac{\frac{n}{2} \cdot 2(q^2 + 1)q^{\frac{n^2}{8} - 2} + \frac{2}{z_-}(q+1)^2q^{\frac{n^2}{8} + \frac{n}{4} - 2}}{\frac{1}{8}q^{\frac{n^2}{4} - 1}} \\ &= \frac{2^3n(q^2 + 1)}{q^{\frac{n^2}{8} + 1}} + \frac{2^4(q+1)^2}{q^{\frac{n^2}{8} - \frac{n}{4} + 1}}. \end{aligned} \tag{1}$$

We now consider Σ_0 . By Corollary 4.3 there are at most $(n^2 + \frac{21}{4}n - 1)e_G$ classes of subgroups $M \in \mu_0$ such that $\text{soc}(M) \notin \{A_{n+1}, A_{n+2}\}$. As shown above, there are at most e_G classes of subgroups $M \in \mu_0$ such that $\text{soc}(M) \in \{A_{n+1}, A_{n+2}\}$. By Corollary 6.2, for M of the first type there are at most $\frac{n(q^{\frac{n}{2}} + 1)}{ra_-(2, q-1)}$ G -conjugates of M containing x , and for M of the second type there are at most $\frac{n(q^{\frac{n}{2}} + 1)}{\frac{1}{2}r(r-1)a_-(2, q-1)}$. By Proposition 5.1 we have $i_2(M) < q^{2n+4}$ for M of the first type, and $i_2(M) < (n+2)!$ for the second type. Therefore

$$\begin{aligned} \Sigma_0 &= \sum_{x \in M \in \mathcal{S}} \frac{i_2(M)}{i_2(G)} \\ &\leq \sum_{M \in \mu_0} \frac{|N_G(\langle x \rangle)|}{|N_M(\langle x \rangle)|} \frac{i_2(M)}{i_2(G)} \\ &\leq \frac{e_G n(n^2 + \frac{21}{4}n - 1)(q^{\frac{n}{2}} + 1)q^{2n+4}}{ra_-(2, q-1)\frac{1}{8}q^{\frac{n^2}{4} - 1}} + \frac{e_G n(q^{\frac{n}{2}} + 1)(n+2)!}{\frac{1}{2}r(r-1)a_-(2, q-1)\frac{1}{8}q^{\frac{n^2}{4} - 1}} \\ &\leq \frac{2^3(2, q-1)(n^2 + \frac{21}{4}n - 1)(q^{\frac{n}{2}} + 1)}{q^{\frac{n^2}{4} - 2n - 5}} + \frac{2^4(2, q-1)(n+2)!(q^{\frac{n}{2}} + 1)}{nq^{\frac{n^2}{4} - 1}}. \end{aligned} \tag{2}$$

Using (1), (2) it can then be verified that for $n \geq 16$ and all q , or for $n \geq 14$ and $q \geq 3$, we have $\Sigma_3 + \Sigma_0 < 1$. For $n = 14, q = 2$, we find $r = 43$, and so by Table 7 and [15] there are no subgroups $M \in \mathcal{S}$ with order divisible by r . Therefore using (1) we find $\Sigma_3 < 1$, and this proves the result for $G = P\Omega_n^-(q), n \geq 14$. Similar arguments deal with all other possibilities for G . \square

7. Proof of Theorem 2 for small n

We now consider G with n smaller than in Proposition 6.5. To prove Theorem 2 it suffices to prove the following groups are $(2, r)$ -generated:

- $PSL_8(q)$;
- (†) $PSp_n(q), n = 8, 10$ and $(n, q) = (12, 2)$;
- $P\Omega_n^\epsilon(q), n = 8$ ($q \neq 2$ for $\epsilon = +$), $10, 12$ and $(n, q, \epsilon) = (14, 2, +), (16, 2, +), (18, 2, +)$;
- $P\Omega_n(q)$ (q odd), $n = 9, 11$.

Proposition 7.1. *If G is a group listed in (†) and M is a maximal subgroup of G with order divisible by r , then M is conjugate to a group listed in Table 6 ($M \in \mathcal{C}_i$) or a group with socle listed in Table 10 ($M \in \mathcal{S}$). In Table 10, upper bounds C_M are given for the number of G -classes of subgroups M with the given socle.*

Table 10: Maximal subgroups $M \in \mathcal{S}$ with $r \mid |M|$ for G in (†)

G	$\text{soc}(M)$	Conditions	C_M
$PSp_8(q)$	$PSL_2(17)$	$q = p$ or $p^2, q \geq 9$ or $q = 2, r = 17$	2
$PSp_{10}(q)$	$PSL_2(11)$	$q = p$ or p^2, q odd, $r = 11$	6
	$PSU_5(2)$	$q = p$ odd, $r = 11$	2
$PSp_{12}(2)$	$PSL_2(25)$		1
	A_{14}		1
$P\Omega_8^+(q)$	$P\Omega_7(q)$	q odd	4
	$PSp_6(q)$	q even	2
	$PSU_3(q)$	$q \equiv 2 \pmod{3}$	$(2, q - 1)^2$
	$P\Omega_8^+(2)$	$q = p$ odd, $r = 7$	4
	$Sz(8)$	$q = 5$	8
	A_{10}	$q = 5$	12
$P\Omega_{12}^+(q)$	$PSL_2(11)$	$q = p \geq 19, r = 11$	8
	M_{12}	$q = p \geq 5, r = 11$	8
	A_{13}	$q = p$ odd, $r = 11$	4
$P\Omega_{14}^+(2)$	$PSL_2(13)$		$2e_G$
	$G_2(3)$		e_G
	A_{16}		e_G

$P\Omega_{10}^-(q)$	$PSL_2(11)$	$q = p \geq 11, r = 11$	$(q + 1, 4)$
	A_{11}	$q \neq 2, r = 11$	$(q + 1, 4)$
	A_{12}	$q = 2$	1
$P\Omega_{12}^-(q)$	$PSL_2(13)$	$q = p$ or $p^3, q \geq 8, r = 13$	6
	$PSL_3(3)$	$q = p, r = 13$	$2(q + 1, 2)$
	A_{13}	$q \neq 7, r = 13$	$(q + 1, 2)$
$P\Omega_9(q)$	$PSL_2(17)$	$q = p$ or $p^2, r = 17$	2
$P\Omega_{11}(q)$	A_{12}	$q = p, r = 11$	2

Proof. For $n \leq 12$, this is proved using [4]. For the groups $G = P\Omega_{14}^+(2), P\Omega_{16}^+(2), P\Omega_{18}^+(2)$ with $M \in \mathcal{S}$, a list of possibilities for $\text{soc}(M)$ and the number of $GL_n(q)$ -classes c for each $\text{soc}(M)$ is obtained from [15]. By [19, Corollary 2.10.4] we can bound the number of G -classes of M by ce_G . \square

Proposition 7.2. *Theorem 2 holds for G listed in (\dagger) .*

Proof. We prove the remaining cases of G are all $(2, r)$ -generated in the usual way, proceeding case by case for n . We use similar bounds for $\Sigma_i, 1 \leq i \leq 8$, to those found in the proof of Proposition 6.5, though since n is fixed in each case we are able to improve the bounds on Σ_2 and Σ_3 in the following way:

- Consider $M \in \mathcal{C}_2$ of the form $O_1(q) \text{ wr } S_n$. Then we have $i_2(M) \leq 2^{n-1}(i_2(S_n) + 1) = 2^{n-1} \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n!}{2^k k! (n-2k)!}$.
- Let $M \in \mathcal{C}_3$ be of the form $Cl_k(q^t).t$ for a classical group $Cl_k(q^t)$ of the same type as G . Instead of bounding the number of prime divisors of n (and hence the number of classes) by dn for some $d \leq 1$, we instead calculate the exact number of prime divisors and bound the number of involutions contained in each class separately using Table 6.

We can improve the bound on Σ_0 more significantly: Table 10 lists possible $\text{soc}(M)$ for subgroups M contributing to Σ_0 . If $\text{soc}(M) \notin \{A_{n+1}, A_{n+2}\}$ then we bound $i_2(M)$ using either $|M|$ or Proposition 2.3 if applicable, rather than q^{2n+4} from [22]. If $\text{soc}(M) = A_{n'}$ for $n' = n + 1$ or $n + 2$, then $i_2(M) \leq i_2(S_{n'}) \leq \sum_{k=0}^{\lfloor \frac{n'}{2} \rfloor} \frac{n'!}{2^k k! (n'-2k)!}$.

We note that for each n it may not be possible to prove $Q_2(G, x) < 1$ for all q using the lower bound $i_2(G) \geq I_2(G)$. However, for specific q we can prove $Q_2(G, x) < 1$ by instead using the lower bound for $i_2(G)$ listed in Table 5.

As an illustration, consider $G = P\Omega_{12}^+(q)$. By Table 6 and Table 10,

$$\sum_{x \in M <_{\max} G} \frac{i_2(M)}{i_2(G)} = \Sigma_1 + \Sigma_2 + \Sigma_3 + \Sigma_0.$$

We first consider Σ_1 . Let μ_1 be a set of conjugacy class representatives of subgroups $M \in \mathcal{C}_1$ such that $r \mid |M|$. There exists a unique $M \in \mu_1$ of type $O_{10}^-(q) \times O_2^-(q)$ and

either two representatives of type $O_{11}(q) \times O_1(q)$ (q odd) or one of type $O_{11}(q)$ (q even). Using Tables 4, 6 and 9 we bound $c_M \frac{|N_G(\langle x \rangle)| i_2(M)}{|N_M(\langle x \rangle)| i_2(G)}$ by $c_M \frac{|N_G(\langle x \rangle)| i_2(M)}{N_M} \frac{i_2(M)}{i_2(G)}$, and we find that the subgroups giving the largest contribution to Σ_1 occur when q is odd. This leads to

$$\begin{aligned}
\Sigma_1 &= \sum_{M \in \mu_1} \frac{|N_G(\langle x \rangle)| i_2(M)}{|N_M(\langle x \rangle)| i_2(G)} \\
&\leq \sum_{\substack{M \in \mu_1 \text{ of type} \\ O_{10}^-(q) \times O_2^-(q)}} \frac{|N_G(\langle x \rangle)| i_2(M)}{|N_M(\langle x \rangle)| i_2(G)} + \sum_{\substack{M \in \mu_1 \text{ of type} \\ O_{11}(q) \times O_1(q)}} \frac{|N_G(\langle x \rangle)| i_2(M)}{|N_M(\langle x \rangle)| i_2(G)} \\
&\leq \frac{2(q+1)^2 q^{24}}{\frac{1}{8} q^{35}} + \frac{2(q+1)}{(2, q-1)^2} \times \frac{\frac{4}{z_+} (q+1) q^{29}}{\frac{1}{8} q^{35}} \\
&= \frac{2^4 (q+1)^2}{q^{11}} + \frac{2^6 (q+1)^2}{(2, q-1)^3 q^6}.
\end{aligned}$$

We now consider Σ_2 . Let μ_2 be a set of conjugacy class representatives for $M \in \mathcal{C}_2$ such that $r \mid |M|$. By Table 6, $|\mu_2| \leq 4$, and for each $M \in \mu_2$ we have $i_2(M) \leq 2^{11} \sum_{k=0}^6 \frac{12!}{2^k k! (12-2k)!} = 2^{14} .17519$ from the above discussion. By Table 6 we require $q = p \neq 2$ for such subgroups to exist. This yields

$$\begin{aligned}
\Sigma_2 &= \sum_{M \in \mu_2} \frac{|N_G(\langle x \rangle)| i_2(M)}{|N_M(\langle x \rangle)| i_2(G)} \\
&\leq 4 \times \frac{10(q^5+1)(q+1)}{4a_+ r} \times \frac{2^{11} \sum_{k=0}^6 \frac{12!}{2^k k! (12-2k)!}}{\frac{1}{8} q^{35}} \\
&\leq \frac{2^{16} .17519 (q+1) (q^5+1)}{q^{35}}.
\end{aligned}$$

We now consider Σ_3 . Let μ_3 be a set of conjugacy class representatives for $M \in \mathcal{C}_3$. By Table 6 there are 2 such classes of the form $GU_6(q).2$, and for each we have $i_2(M) < \frac{2}{z_+} (q+1)^2 q^{19}$. Each class has a unique conjugate containing x by Corollary 6.2. Therefore

$$\begin{aligned}
\Sigma_3 &= \sum_{M \in \mu_3} \frac{|N_G(\langle x \rangle)| i_2(M)}{|N_M(\langle x \rangle)| i_2(G)} \\
&\leq 2 \times \frac{2(q+1)^2 q^{19}}{\frac{1}{8} q^{35}} \\
&= \frac{2^5 (q+1)^2}{q^{16}}.
\end{aligned}$$

We now consider Σ_0 . By Table 10, we can assume $r = 11$. Let μ_0 be a set of conjugacy class representatives for $M \in \mathcal{S}$ with $r \mid |M|$. By Table 10 there are at most 8 classes with socle $PSL_2(11)$, 8 classes with socle M_{12} , and 4 classes with socle A_{13} . By Corollary 6.2 and Table 9, for M of the first or second type there are at most $\frac{10(q^5+1)(q+1)}{11a_+(2, q-1)^2}$ G -conjugates of M also containing x , and for M of alternating type there are at most $\frac{10(q^5+1)(q+1)}{\frac{1}{2} 11(11-1)a_+(2, q-1)^2}$

conjugates containing x . Using [4] and [7] we calculate $i_2(M) \leq 55, 190080, 272415$ for $\text{soc}(M) = PSL_2(11), M_{12}, A_{13}$ respectively. Therefore, as $q = p$ is odd in all cases,

$$\begin{aligned} \Sigma_0 &= \sum_{M \in \mu_0} \frac{|N_G(\langle x \rangle)| i_2(M)}{|N_M(\langle x \rangle)| i_2(G)} \\ &= 8 \times \frac{10(q^5 + 1)(q + 1)}{ra_+(2, q - 1)^2} \times \frac{55}{\frac{1}{8}q^{35}} + 8 \times \frac{10(q^5 + 1)(q + 1)}{ra_+(2, q - 1)^2} \times \frac{190080}{\frac{1}{8}q^{35}} \\ &\quad + 4 \times \frac{10(q^5 + 1)(q + 1)}{\frac{1}{2}r(r - 1)a_+(2, q - 1)^2} \times \frac{272415}{\frac{1}{8}q^{35}} \\ &= \frac{2^3 \cdot 5 \cdot 11 \cdot 3593 (q^5 + 1)(q + 1)}{q^{35}}. \end{aligned}$$

We see that for $q \geq 3$ we have $Q_2(G, x) \leq \Sigma_1 + \Sigma_2 + \Sigma_3 + \Sigma_0 < 1$. It therefore suffices to prove $\Sigma < 1$ for $q = 2$. Computing similar bounds for Σ_i using the lower bound for $i_2(G)$ found in Table 5 instead of $I_2(G)$ from Table 4 yields the result.

Calculations for the remaining G are similar. □

References

- [1] M. Aschbacher, On the maximal subgroups of the finite classical groups. *Invent. Math.* **76** (1984), 469-514.
- [2] M. Aschbacher, R. Guralnick, Some applications of the first cohomology group. *J. Alg.* **90** (1984), 446-460.
- [3] M. Aschbacher, G. M. Seitz, Involutions in Chevalley groups over fields of even order. *Nagoya Math. J.* **63** (1976), 1-91.
- [4] J. N. Bray, D. F. Holt, C. M. Roney-Dougal, The Maximal Subgroups of the Low-Dimensional Finite Classical Groups. *London Math. Soc. Lecture Note Series* **407**, Cambridge Univ. Press (2013).
- [5] R. Burkhardt, Die Zerlegungsmatrizen der Gruppen $PSL(2, p^f)$. *J. Alg.* **40** (1976), 75-96.
- [6] A. A. Buturlakin, M. A. Grechkoseeva, The cyclic structure of maximal tori of the finite classical groups. *Alg. and Log.* **46** (2007), 73-89.
- [7] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, R. A. Wilson, An ATLAS of Finite Groups. *Clarendon Press, Oxford* (1985).
- [8] L. Di Martino, N. A. Vavilov, $(2, 3)$ -generation of $SL_n(q)$. I. Cases $n = 5, 6, 7$. *Comm. Alg.* **22** (1994), 1321-1347.
- [9] L. Di Martino, N. A. Vavilov, $(2, 3)$ -generation of $SL_n(q)$. II. Cases $n \geq 8$. *Comm. Alg.* **24** (1996), 487-515.

- [10] H. Enomoto, The characters of the finite symplectic group $Sp(4, q), q = 2^f$. *Osaka J. Math.* **9** (1972), 75-94.
- [11] M. J. Evans, A note on two-generator groups. *Rocky Mount. J. Math.* **17** (1987), 887-889.
- [12] D. Gorenstein, R. Lyons, R. Solomon, The Classification of the Finite Simple Groups, Number 3. *Math. Surveys and Monographs* **40**, Am. Math. Soc. (1998).
- [13] R. M. Guralnick, W. M. Kantor, Probabilistic generation of finite simple groups. Special issue in honor of Helmut Wielandt. *J. Alg.* **234** (2000), 743-792.
- [14] R. Guralnick, K. Magaard, J. Saxl, P. H. Tiep, Cross characteristic representations of symplectic and unitary groups. *J. Alg.* **257** (2002), 291-347.
- [15] R. Guralnick, T. Penttila, C. E. Praeger, J. Saxl, Linear groups with orders having certain large prime divisors. *Proc. London Math. Soc.* **78** (1999), 167-214.
- [16] R. Guralnick, P. H. Tiep, Low-dimensional representations of special linear groups in cross characteristics. *Proc. London Math. Soc.* **78** (1999), 116-138.
- [17] G. Hiss, G. Malle, Low dimensional representations of special unitary groups. *J. Alg.* **236** (2001), 745-767.
- [18] C. Jansen, K. Lux, R. A. Parker, R. A. Wilson, An ATLAS of Brauer characters. *Clarendon Press, Oxford* (1995).
- [19] P. B. Kleidman, M. W. Liebeck, The Subgroup Structure of the Finite Classical Groups. *London Math. Soc. Lecture Note Series* **129**, Cambridge Univ. Press (1990).
- [20] S. Lang, Algebraic groups over finite fields. *Am. Math. J.* **78** (1956), 555-563.
- [21] R. Lawther, M. W. Liebeck, G. M. Seitz, Fixed point ratios in actions of finite exceptional groups of lie type. *Pac. J. Math.* **205** (2002), 393-464.
- [22] M. W. Liebeck, On the orders of maximal subgroups of the finite classical groups. *Proc. London Math. Soc.* **50** (1985), 426-446.
- [23] M. W. Liebeck, G. M. Seitz, Unipotent and Nilpotent Classes in Simple Algebraic Groups and Lie Algebras. *Math. Surveys and Monographs* **180**, Am. Math. Soc. (2012).
- [24] M. W. Liebeck, A. Shalev, Classical groups, probabilistic methods and the (2, 3)-generation problem. *Ann. Math.* **144** (1996), 77-125.
- [25] F. Lübeck, G. Malle, (2, 3)-generation of exceptional groups. *J. London. Math. Soc.* **59** (1999), 101-122.
- [26] A. M. Macbeath, Generators of the linear fractional group. *Proc. Symp. Pure. Math.* **12** (1969), 14-32.

- [27] G. Malle, J. Saxl, T. Weigel, Generation of classical groups. *Geom. Dedicata* **49** (1994), 85-116.
- [28] G. Malle, D. Testerman, Linear Algebraic Groups and Finite Groups of Lie Type. *Cambridge studies in advanced math.* **133**, Cambridge Univ. Press (2011).
- [29] G. A. Miller, On the groups generated by two operators. *Bull. Am. Math. Soc.* **7** (1901), 424-426.
- [30] G. A. Miller, Possible orders of two generators of the alternating and of the symmetric group. *Trans. Am. Math. Soc.* **30** (1928), 24-32.
- [31] M. A. Pellegrini, The $(2, 3)$ -generation of the classical simple groups of dimension 6 and 7. *Bull. Aus. Math. Soc.* **93** (2016), 61-72.
- [32] M. A. Pellegrini, The $(2, 3)$ -generation of the special linear groups over finite fields. To appear in *Bull. Aus. Math. Soc.*, <https://doi.org/10.1017/S0004972716000617>.
- [33] M. A. Pellegrini, M. Prandelli, M. C. Tamburini Bellani, The $(2, 3)$ -generation of the special unitary groups of dimension 6. *J. Alg. and its Appl.* **15** (2016), Article 1650171.
- [34] M. A. Pellegrini, M. C. Tamburini Bellani, The simple classical groups of dimension less than 6 which are $(2, 3)$ -generated. *J. Alg. and its Appl.* **14** (2015), Article 1550148.
- [35] M. A. Pellegrini, M. C. Tamburini Bellani, M. A. Vsemirnov, Uniform $(2, k)$ -generation of the 4-dimensional classical groups. *J. Alg.* **369** (2012), 322-350.
- [36] P. Sanchini, M. C. Tamburini, Constructive $(2, 3)$ -generation: a permutational approach. *Rend. Sem. Mat. Fis. Milano* **64** (1994), 141-158.
- [37] M. A. Shahabi, H. Mohtadifar, The characters of finite projective symplectic group $PSp(4, q)$. Groups St. Andrews 2001 in Oxford. Vol. II. *London Math. Soc. Lecture Note Series* **305**, Cambridge Univ. Press (2003), 496-527.
- [38] A. Stein, $1\frac{1}{2}$ -generation of finite simple groups. *Beit. Alg. Geom.* **39** (1998), 349-358.
- [39] R. Steinberg, Generators for simple groups. *Canad. J. Math.* **14** (1962), 277-283.
- [40] M. Suzuki, On a class of doubly transitive groups. *Ann. of Math.* **75** (1962), 105-145.
- [41] K. Tabakov, $(2, 3)$ -generation of the groups $PSL_7(q)$. *Proc. of the Forty Second Spring Conf. of the Union of Bulg. Math., Borovetz, April 2-6* (2013), 260-264.
- [42] K. Tabakov, K. Tchakerian, $(2, 3)$ -generation of the groups $PSL_6(q)$. *Serdica Math. J.* **37** (2011), 365-370.
- [43] M. A. Vsemirnov, More classical groups which are not $(2, 3)$ -generated. *Arch. Math.* **96** (2011), 123-129.
- [44] R. A. Wilson, The Monster is a Hurwitz group. *J. Group Theory* **4** (2001), 367-374.

- [45] A. J. Woldar, On Hurwitz generation and genus actions of sporadic groups. *Ill. J. Math.* **33** (1989), 416-437.
- [46] K. Zsigmondy, Zur Theorie der Potenzreste. *Monatsh. Für Math. u. Phys.* **3** (1892), 265-284.