

A Tight Rate Bound and a Matching Construction for Locally Recoverable Codes with Sequential Recovery From Any Number of Multiple Erasures

S. B. Balaji, Ganesh R. Kini and P. Vijay Kumar, *Fellow, IEEE*

Department of Electrical Communication Engineering, Indian Institute of Science, Bangalore.

Email: balaji.profess@gmail.com, kiniganesh94@gmail.com, pvk1729@gmail.com

Abstract

An $[n, k]$ code \mathcal{C} is said to be locally recoverable in the presence of a single erasure, and with locality parameter r , if each of the n code symbols of \mathcal{C} can be recovered by accessing at most r other code symbols. An $[n, k]$ code is said to be a locally recoverable code with sequential recovery from t erasures, if for any set of $s \leq t$ erasures, there is an s -step sequential recovery process, in which at each step, a single erased symbol is recovered by accessing at most r other code symbols. This is equivalent to the requirement that for any set of $s \leq t$ erasures, the dual code contain a codeword whose support contains the co-ordinate of precisely one of the s erased symbols.

In this paper, a tight upper bound on the rate of such a code, for any value of number of erasures t and any value $r \geq 3$, of the locality parameter is derived. This bound proves an earlier conjecture due to Song, Cai and Yuen. While the bound is valid irrespective of the field over which the code is defined, a matching construction of *binary* codes that are rate-optimal is also provided, again for any value of t and any value $r \geq 3$.

Index Terms

Distributed storage, locally recoverable codes, codes with locality, locally repairable codes, sequential repair, multiple erasures, rate bound, proof of conjecture.

I. INTRODUCTION

An $[n, k]$ code \mathcal{C} is said to have locality r if each of the n code symbols of \mathcal{C} can be recovered by accessing at most r other code symbols. Equivalently, there exist n codewords $\underline{h}_1 \cdots \underline{h}_n$, not necessarily distinct, in the dual code \mathcal{C}^\perp , such that $i \in \text{supp}(\underline{h}_i)$ and $|\text{supp}(\underline{h}_i)| \leq r + 1$ for $1 \leq i \leq n$ where $\text{supp}(\underline{h}_i)$ denotes the support of the codeword \underline{h}_i .

a) *Codes with Sequential Recovery:* An $[n, k]$ code \mathcal{C} over a field \mathbb{F}_q is defined as a code with sequential recovery [1] from t erasures and with locality-parameter r , if for any set of $s \leq t$ erased symbols $\{c_{\sigma_1}, \dots, c_{\sigma_s}\}$, there exists a codeword \underline{h} in the dual code \mathcal{C}^\perp of Hamming weight $\leq r + 1$, such that $\text{supp}(\underline{h}) \cap \{\sigma_1, \dots, \sigma_s\} = 1$. We will formally refer to this class of codes as $(n, k, r, t)_{\text{seq}}$ codes. When the parameters (n, k, r, t) are clear from the context, we will simply refer to a code in this class as a code with sequential recovery.

A. Background

In [2] Gopalan et al. introduced the concept of codes with locality (see also [3], [4]), where an erased code symbol is recovered by accessing a small subset of other code symbols. The size of this subset denoted by r , is typically much smaller than the dimension of the code, making the repair process more efficient when compared with MDS codes. The focus of [2] was local recovery from single erasure (see also [2], [5], [6], [7]).

The sequential approach to recovery from erasures, introduced by Prakash et al. [8] is one of several approaches to locally recover from multiple erasures. Codes employing this approach have been shown to be better in terms of rate and minimum distance (see [8], [9], [10], [11], [1], [12]). Local recovery in the presence of two erasures is considered in [8] (see also [10]) where a tight rate bound for two erasure case and an optimal construction is provided. Codes with sequential recovery from three erasures can be found discussed in [10], [1], [13]. A bound on rate of an $(n, k, r, 3)_{\text{seq}}$ code was derived in [13]. A rate bound for $t = 4$ appears in [12].

There are several approaches to local recovery from multiple erasures:

- 1) *Stronger Local Codes:* Here, every code symbol is contained in a local code of length at most $r + t$ and minimum distance at least $t + 1$, see ([14]). While the recovery process is sequential, the erased symbols can be recovered in arbitrary order, see [6], [15].

P. Vijay Kumar is also an Adjunct Research Professor at the University of Southern California. This research is supported in part by the National Science Foundation under Grant 1421848 and in part by an India-Israel UGC-ISF joint research program grant.

S. B. Balaji would like to acknowledge the support of TCS research-scholarship program.

- 2) *Codes with Availability*: For every code symbol c_i in \mathcal{C} , there exist t codewords h_1^i, \dots, h_t^i in the dual of the code, each of Hamming weight $\leq r + 1$, such that $\text{supp}(h_g^i) \cap \text{supp}(h_j^i) = \{i\}$, $\forall 1 \leq g \neq j \leq t$; recovery can be carried out in parallel. For more details see [16], [17], [7], [18].
- 3) *Codes with Selectable Recovery*: Here, given any set of t erasures, every erased symbol has a parity check that involves that symbol and no other erased symbol; with these codes, one is free to choose the order in which to recover the erased symbols; recovery in parallel may or may not be possible, depending upon the specific construction. For details, see [19].
- 4) *Codes with Co-operative local recovery*: A code with co-operative local recovery has the property that for a given t , any set of t erased symbols can be co-operatively recovered by accessing at most r other symbols, see [9].
- 5) *Codes with Sequential Recovery*: This class of codes is the object of study in the present paper and a definition has been provided above. The class of sequential-recovery codes is a larger class of codes that contains all the four above-mentioned classes of codes as depicted in Fig. 1. For this reason, codes with sequential recovery can potentially achieve higher rate and have larger minimum distance.




Fig. 1: Figure showing the relationship of codes with sequential recovery to some other classes of codes designed for recovery from multiple erasures.

B. Contributions of the Paper

In this paper, we derive an upper bound on the rate of a code having locality-parameter r with sequential recovery from t erasures, for any $r \geq 3$ and any t . While the bound is valid irrespective of the field over which the code is defined, we provide here a matching construction of *binary* codes that are rate-optimal, i.e., binary codes achieving the rate bound for any t and any $r \geq 3$. These results are also shown to prove a conjecture on code rate due to Song, Cai and Yeun [10].

II. UPPER BOUND ON RATE OF AN $(n, k, r, t)_{seq}$ CODE

In this section we provide an upper bound on the rate of an $(n, k, r, t)_{seq}$ code for $r \geq 3$. The cases of t even and t odd are considered in two separate subsections.

A. Upper Bound on Rate of an $(n, k, r, t)_{seq}$ Code for t Even:

In this subsection we provide an upper bound on the rate of an $(n, k, r, t)_{seq}$ code for t even and $r \geq 3$.

Theorem 1. *Let \mathcal{C} be an $(n, k, r, t)_{seq}$ code over a field \mathbb{F}_q . Let t be a positive even integer and $r \geq 3$. Then*

$$\frac{k}{n} \leq \frac{r^{\frac{t}{2}}}{r^{\frac{t}{2}} + 2 \sum_{i=0}^{\frac{t}{2}-1} r^i}. \quad (1)$$

Proof. We begin by setting $\mathcal{B}_0 = \text{span}(\underline{c} \in \mathcal{C}^\perp : w_H(\underline{c}) \leq r + 1)$ ($w_H(\underline{c})$ denotes the Hamming weight of the vector \underline{c}). Let m be the dimension of \mathcal{B}_0 . Let $\underline{c}_1, \dots, \underline{c}_m$ (row vectors) be a basis of \mathcal{B}_0 such that $w_H(\underline{c}_i) \leq r + 1$.

$$\text{Let } H_1 = \begin{bmatrix} \underline{c}_1 \\ \vdots \\ \underline{c}_m \end{bmatrix}.$$

It follows that H_1 is a parity check matrix of an $(n, n - m, r, t)_{seq}$ code as its row space contains all the codewords of Hamming weight at most $r + 1$ which are contained in \mathcal{C}^\perp . Also,

$$\frac{k}{n} \leq 1 - \frac{m}{n}.$$

The idea behind the next few arguments in the proof is the following. The codes with largest rate will tend to have a larger value of n for fixed m . On the other hand, the Hamming weight of the matrix H_1 (i.e., the number of non-zero entries in the matrix) is bounded above by $m(r + 1)$. It follows that to make n large, one would like the columns of H_1 to have as small

a weight as possible. It is therefore quite natural to start building H_1 by picking as many columns of weight 1 as possible, then columns of weight 2 and so on. As one proceeds by following this approach, it turns out that the matrix H_1 is forced to have a certain sparse, block-diagonal, staircase form and an understanding of this structure is used to derive the upper bound on code rate. We now proceed to derive an upper bound on $1 - \frac{m}{n}$. It can be seen that the matrix H_1 , after permutation of rows and columns can be written in the following form

$$H_1 = \left[\begin{array}{c|c|c|c|c|c|c|c} D_0 & A_1 & 0 & 0 & \dots & 0 & 0 & 0 \\ \hline 0 & D_1 & A_2 & 0 & \dots & 0 & 0 & 0 \\ \hline 0 & 0 & D_2 & A_3 & \dots & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & D_3 & \dots & 0 & 0 & 0 \\ \hline \vdots & \vdots & \vdots & \vdots & \dots & \vdots & \vdots & \vdots \\ \hline 0 & 0 & 0 & 0 & \dots & A_{\frac{t}{2}-2} & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & \dots & D_{\frac{t}{2}-2} & A_{\frac{t}{2}-1} & 0 \\ \hline 0 & 0 & 0 & 0 & \dots & 0 & D_{\frac{t}{2}-1} & 0 \\ \hline 0 & 0 & 0 & 0 & \dots & 0 & 0 & C \end{array} \right] D, \quad (2)$$

where

- 1) the rows of H_1 are labeled $1, 2, \dots, m$ and columns are labeled $1, 2, \dots, n$,
- 2) A_i is a $\rho_{i-1} \times a_i$ matrix for $1 \leq i \leq \frac{t}{2} - 1$, D_i is a $\rho_i \times a_i$ matrix for $0 \leq i \leq \frac{t}{2} - 1$ for some $\{\rho_i\}, \{a_i\}$,
- 3) D_0 is a matrix with each column having weight (Hamming weight) 1 and each row having weight at least 1,
- 4) $\{A_i\}, \{D_i\}, \{B_i = \begin{bmatrix} A_i \\ D_i \end{bmatrix}\}$ are such that for $1 \leq i \leq \frac{t}{2} - 1$ each column of B_i has weight 2, each column of A_i has weight at least 1 and each row of D_i has weight at least 1 and each column of D_i has weight at most 1,
- 5) C is a matrix with each column having weight 2, D is a matrix which exactly contains all the columns of H_1 which have weight ≥ 3 ,
- 6) If J is the first index such that A_J, D_J are empty matrices ($0 \times L$, $L \times 0$, 0×0 matrix) ($J = 0$ if D_0 is an empty matrix) then we set A_i, D_i to be empty matrices for all $J \leq i \leq \frac{t}{2} - 1$ and set $a_i = 0, \rho_i = 0, \forall J \leq i \leq \frac{t}{2} - 1$. If none of $D_0, A_i, D_i, 1 \leq i \leq \frac{t}{2} - 1$ are empty matrices, then we simply set $J = \frac{t}{2}$. If $J < \frac{t}{2}$, let $\alpha_1, \dots, \alpha_l \in \{\sum_{j=0}^{J-1} a_j + 1, \dots, n\}$ be the indices of 2-weight columns in H_1 apart from the 2-weight columns corresponding to columns of B_1, \dots, B_{J-1} . We take the submatrix of H_1 with rows indexed by $\sum_{j=0}^{J-1} \rho_j + 1, \dots, m$ and columns indexed by $\alpha_1, \dots, \alpha_l$ and name the submatrix as matrix C . Let the number of columns in C be $a_{\frac{t}{2}}$. If C is an empty matrix then we set $a_{\frac{t}{2}} = 0$. If $J < \frac{t}{2}$, the matrix H_1 can be written in the form (3)

$$H_1 = \left[\begin{array}{c|c|c|c|c|c|c|c} D_0 & A_1 & 0 & 0 & \dots & 0 & 0 & 0 \\ \hline 0 & D_1 & A_2 & 0 & \dots & 0 & 0 & 0 \\ \hline 0 & 0 & D_2 & A_3 & \dots & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & D_3 & \dots & 0 & 0 & 0 \\ \hline \vdots & \vdots & \vdots & \vdots & \dots & \vdots & \vdots & \vdots \\ \hline 0 & 0 & 0 & 0 & \dots & A_{J-2} & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & \dots & D_{J-2} & A_{J-1} & 0 \\ \hline 0 & 0 & 0 & 0 & \dots & 0 & D_{J-1} & 0 \\ \hline 0 & 0 & 0 & 0 & \dots & 0 & 0 & C \end{array} \right] D, \quad (3)$$

The entire rate bound derivation is correct and all the bounds will hold with $a_i = 0, \rho_i = 0, J \leq i \leq \frac{t}{2} - 1$. If $l (< J)$ is the first index such that D_l is an empty matrix with A_l a non empty matrix then the proof of the following claim 1 (since in the proof of claim 1, we prove the claim for A_l first and then proceed to D_l and since D_0, A_j, D_j must be non empty $\forall j < l$) will imply that A_l has column weight 1 which will imply D_l cannot be empty. Hence the case A_l , a non empty matrix and D_l , an empty matrix cannot occur. Although we have to prove the following claim 1 for $A_i, D_i, 1 \leq i \leq J - 1, D_0$, we assume all $D_0, A_i, D_i, 1 \leq i \leq \frac{t}{2} - 1$ to be non-empty matrices and prove the claim. Since the proof is by induction, the induction can be made to stop at $J - 1$ (induction starts at 0) and the proof is unaffected by it.

Claim 1. For $1 \leq i \leq \frac{t}{2} - 1$, A_i is a matrix with each column having weight 1 and for $0 \leq j \leq \frac{t}{2} - 1$, D_j is a matrix with each row and each column having weight 1.

Proof. We use the fact that $d_{\min}(C) \geq t + 1$ to prove the above claim. Proof is given in Appendix A. \square

By Claim 1, after permutation of columns of H_1 (in (2) or (3) depending on J) within the columns labeled by the set $\{\sum_{l=0}^{j-1} a_l + 1, \dots, \sum_{l=0}^{j-1} a_l + a_j\}$ for $0 \leq j \leq J - 1$, the matrix $D_j, 0 \leq j \leq J - 1$ can be assumed to be a diagonal matrix with non-zero entries along the diagonal and hence $\rho_i = a_i, \forall 0 \leq i \leq \frac{t}{2} - 1$.

Since the sum of the weights of the columns of $A_i, 1 \leq i \leq \frac{t}{2} - 1$ must equal the sum of the row weights and since each row of A_i for $i \leq J - 1$ can have weight at most r and not $r + 1$ due to weight one rows in D_{i-1} , and since for $\frac{t}{2} - 1 \geq i \geq J$, A_i is an empty matrix and we have set $a_i = 0$, we obtain:

$$\begin{aligned} \text{For } 1 \leq i \leq \frac{t}{2} - 1 : \\ \rho_{i-1} r &\geq a_i, \\ a_{i-1} r &\geq a_i. \end{aligned} \quad (4)$$

For some $p \geq 0$,

$$\sum_{i=0}^{\frac{t}{2}-1} a_i + p = m. \quad (5)$$

By equating sum of row weights of C , with sum of column weights of C , we obtain:

$$2a_{\frac{t}{2}} \leq (a_{\frac{t}{2}-1} + p)(r + 1) - a_{\frac{t}{2}-1}. \quad (6)$$

Note that if C is an empty matrix then also the inequality (6) is true as we would have set $a_{\frac{t}{2}} = 0$. If $J < \frac{t}{2}$ and C a non-empty matrix then the number of rows in C is p with each column of C having weight 2, hence the inequality (6) is still true.

Substituting (5) in (6) we get:

$$\begin{aligned} 2a_{\frac{t}{2}} &\leq (m - \sum_{i=0}^{\frac{t}{2}-2} a_i)(r + 1) - (m - \sum_{i=0}^{\frac{t}{2}-2} a_i - p), \\ 2a_{\frac{t}{2}} &\leq (m - \sum_{i=0}^{\frac{t}{2}-2} a_i)r + p. \end{aligned} \quad (7)$$

By equating sum of row weights of H_1 , with sum of column weights of H_1 , we obtain:

$$m(r + 1) \geq a_0 + 2\left(\sum_{i=1}^{\frac{t}{2}} a_i\right) + 3\left(n - \sum_{i=0}^{\frac{t}{2}} a_i\right), \quad (8)$$

If $J < \frac{t}{2}$ then $a_i = 0, \forall J \leq i \leq \frac{t}{2} - 1$. If C is an empty matrix then $a_{\frac{t}{2}} = 0$. Hence the inequality (8) is true irrespective of whether $J = \frac{t}{2}$ or $J < \frac{t}{2}$ (even if C is an empty matrix).

From (8) :

$$m(r + 1) \geq 3n - 2a_0 - \left(\sum_{i=1}^{\frac{t}{2}} a_i\right). \quad (9)$$

Our basic inequalities are (4),(5),(6),(8). We manipulate these 4 inequalities to derive the bound on rate.

Substituting (5) in (9) we get:

$$\begin{aligned} m(r + 1) &\geq 3n - a_0 - a_{\frac{t}{2}} - (m - p), \\ m(r + 2) &\geq 3n + p - a_0 - a_{\frac{t}{2}}. \end{aligned} \quad (10)$$

Substituting (7) in (10), we get:

$$\begin{aligned} m(r + 2) &\geq 3n + p - a_0 - \left(\frac{(m - \sum_{i=0}^{\frac{t}{2}-2} a_i)r + p}{2}\right), \\ m\left(r + 2 + \frac{r}{2}\right) &\geq 3n + \frac{p}{2} - a_0 + \left(\sum_{i=0}^{\frac{t}{2}-2} a_i\right) \frac{r}{2}. \end{aligned} \quad (11)$$

From (5), for any $0 \leq j \leq \frac{t}{2} - 2$:

$$a_{\frac{t}{2}-2-j} = m - \sum_{i=0}^{\frac{t}{2}-2-j-1} a_i - \sum_{i=\frac{t}{2}-2-j+1}^{\frac{t}{2}-1} a_i - p. \quad (12)$$

Substituting (4) for $\frac{t}{2} - 2 - j + 1 \leq i \leq \frac{t}{2} - 1$ in (12), we get:

$$\begin{aligned} a_{\frac{t}{2}-2-j} &\geq m - \sum_{i=0}^{\frac{t}{2}-2-j-1} a_i - \sum_{i=\frac{t}{2}-2-j+1}^{\frac{t}{2}-1} a_{\frac{t}{2}-2-j} r^{i-(\frac{t}{2}-2-j)} - p, \\ a_{\frac{t}{2}-2-j} &\geq m - \sum_{i=0}^{\frac{t}{2}-2-j-1} a_i - \sum_{i=1}^{j+1} a_{\frac{t}{2}-2-j} r^i - p, \\ a_{\frac{t}{2}-2-j} &\geq \frac{m - \sum_{i=0}^{\frac{t}{2}-2-j-1} a_i - p}{1 + \sum_{i=1}^{j+1} r^i}. \end{aligned} \quad (13)$$

Let

$$\delta_0 = \frac{r}{2}, \quad (14)$$

$$\delta_{j+1} = \delta_j - \frac{\delta_j}{1 + \sum_{i=1}^{j+1} r^i}. \quad (15)$$

Let us prove the following inequality by induction for $0 \leq J_1 \leq \frac{t}{2} - 2$,

$$m(r + 2 + \delta_{J_1}) \geq 3n + p \left(\frac{1}{2} + \delta_{J_1} - \frac{r}{2} \right) - a_0 + \left(\sum_{i=0}^{\frac{t}{2}-2-J_1} a_i \right) \delta_{J_1}. \quad (16)$$

(16) is true for $J_1 = 0$ by (11). Hence (16) is proved for $t = 4$ and the range of J_1 is vacuous for $t = 2$. Hence assume $t > 4$. Hence let us assume (16) is true for J_1 such that $\frac{t}{2} - 3 \geq J_1 \geq 0$ and prove it for $J_1 + 1$. Substituting (13) for $j = J_1$ in (16), we get:

$$\begin{aligned} m(r + 2 + \delta_{J_1}) &\geq 3n + p \left(\frac{1}{2} + \delta_{J_1} - \frac{r}{2} \right) - a_0 + \left(\sum_{i=0}^{\frac{t}{2}-2-J_1-1} a_i \right) \delta_{J_1} + \left(\frac{m - \sum_{i=0}^{\frac{t}{2}-2-J_1-1} a_i - p}{1 + \sum_{i=1}^{J_1+1} r^i} \right) \delta_{J_1}, \\ m(r + 2 + \delta_{J_1} - \frac{\delta_{J_1}}{1 + \sum_{i=1}^{J_1+1} r^i}) &\geq 3n + p \left(\frac{1}{2} + \delta_{J_1} - \frac{\delta_{J_1}}{1 + \sum_{i=1}^{J_1+1} r^i} - \frac{r}{2} \right) - a_0 + \\ &\quad \left(\sum_{i=0}^{\frac{t}{2}-2-J_1-1} a_i \right) \left(\delta_{J_1} - \frac{\delta_{J_1}}{1 + \sum_{i=1}^{J_1+1} r^i} \right). \end{aligned} \quad (17)$$

Substituting (15) in (17), we obtain

$$m(r + 2 + \delta_{J_1+1}) \geq 3n + p \left(\frac{1}{2} + \delta_{J_1+1} - \frac{r}{2} \right) - a_0 + \left(\sum_{i=0}^{\frac{t}{2}-2-J_1-1} a_i \right) \delta_{J_1+1}. \quad (18)$$

Hence (16) is proved for any $0 \leq J_1 \leq \frac{t}{2} - 2$ for $t \geq 4$. Hence writing (16) for $J_1 = \frac{t}{2} - 2$ for $t \geq 4$, we obtain:

$$\begin{aligned} m(r + 2 + \delta_{\frac{t}{2}-2}) &\geq 3n + p \left(\frac{1}{2} + \delta_{\frac{t}{2}-2} - \frac{r}{2} \right) - a_0 + (a_0) \delta_{\frac{t}{2}-2}, \\ m(r + 2 + \delta_{\frac{t}{2}-2}) &\geq 3n + p \left(\frac{1}{2} + \delta_{\frac{t}{2}-2} - \frac{r}{2} \right) + a_0 (\delta_{\frac{t}{2}-2} - 1). \end{aligned} \quad (19)$$

It can be seen that δ_j for $r \geq 2$ has a product form as:

$$\delta_j = \frac{r}{2} \left(\frac{r^{j+1} - r^j}{r^{j+1} - 1} \right). \quad (20)$$

Hence for $r \geq 3, t \geq 4$:

$$\delta_{\frac{t}{2}-2} = \frac{r}{2} \left(\frac{r^{\frac{t}{2}-1} - r^{\frac{t}{2}-2}}{r^{\frac{t}{2}-1} - 1} \right) \geq 1.$$

Hence we can substitute (13) for $j = \frac{t}{2} - 2$ in (19) :

$$\begin{aligned} m(r+2 + \delta_{\frac{t}{2}-2}) &\geq 3n + p \left(\frac{1}{2} + \delta_{\frac{t}{2}-2} - \frac{r}{2} \right) + \left(\frac{m-p}{1 + \sum_{i=1}^{\frac{t}{2}-1} r^i} \right) (\delta_{\frac{t}{2}-2} - 1), \\ m(r+2 + \delta_{\frac{t}{2}-2} - \frac{\delta_{\frac{t}{2}-2}}{1 + \sum_{i=1}^{\frac{t}{2}-1} r^i} + \frac{1}{1 + \sum_{i=1}^{\frac{t}{2}-1} r^i}) &\geq 3n + p \left(\frac{1}{2} + \delta_{\frac{t}{2}-2} - \frac{\delta_{\frac{t}{2}-2}}{1 + \sum_{i=1}^{\frac{t}{2}-1} r^i} + \frac{1}{1 + \sum_{i=1}^{\frac{t}{2}-1} r^i} - \frac{r}{2} \right). \end{aligned} \quad (21)$$

Substituting (15) in (21), we obtain:

$$m \left(r+2 + \delta_{\frac{t}{2}-1} + \frac{1}{1 + \sum_{i=1}^{\frac{t}{2}-1} r^i} \right) \geq 3n + p \left(\frac{1}{2} + \delta_{\frac{t}{2}-1} + \frac{1}{1 + \sum_{i=1}^{\frac{t}{2}-1} r^i} - \frac{r}{2} \right). \quad (22)$$

Using (20), we obtain:

$$\left(\frac{1}{2} + \delta_{\frac{t}{2}-1} + \frac{1}{1 + \sum_{i=1}^{\frac{t}{2}-1} r^i} - \frac{r}{2} \right) \geq 0.$$

Hence (22) implies:

$$m \left(r+2 + \delta_{\frac{t}{2}-1} + \frac{1}{1 + \sum_{i=1}^{\frac{t}{2}-1} r^i} \right) \geq 3n. \quad (23)$$

(23) after some algebraic manipulations gives the required bound on $1 - \frac{m}{n}$ and hence on $\frac{k}{n}$ as stated in the theorem. Note that although the derivation is valid for $r \geq 3, t \geq 4$, the final bound given in the theorem is correct and tight for $t = 2$. The bound for $t = 2$ can be derived specifically by substituting $a_0 \leq m$ in (11) and noting that $p \geq 0$. \square

An alternative proof for Theorem 1 is given below by using linear programming:

Proof. The inequalities (4),(6) and (9) are linear inequalities and are written in matrix form as:

$$A\mathbf{x} \geq \mathbf{b}$$

where

$$A = \begin{bmatrix} r & -1 & 0 & \dots & 0 & 0 & 0 & 0 \\ 0 & r & -1 & \dots & 0 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & r & -1 & 0 & 0 \\ 0 & 0 & 0 & \dots & 0 & r & -2 & (r+1) \\ (r+3) & (r+2) & (r+2) & \dots & (r+2) & (r+2) & 1 & (r+1) \end{bmatrix} \quad (24)$$

which is a $(\frac{t}{2} + 1) \times (\frac{t}{2} + 2)$ matrix and

$$\mathbf{x} = [a_0 \ a_1 \ \dots \ a_{\frac{t}{2}} \ p]^T, \mathbf{b} = [0 \ 0 \ \dots \ 0 \ 3n]^T \quad (25)$$

where \mathbf{x} is a $(\frac{t}{2} + 2) \times 1$ matrix and \mathbf{b} is a $(\frac{t}{2} + 1) \times 1$ matrix. The problem of finding an upper bound on rate of the code now becomes one of minimizing $m = \mathbf{c}^T \mathbf{x}$, which is a linear objective function where $\mathbf{c} = [1 \ 1 \ \dots \ 1 \ 0 \ 1]^T$ is a $(\frac{t}{2} + 2) \times 1$ matrix. Also by definition of \mathbf{x} , $\mathbf{x} \geq 0$. This is now in a standard form of a linear program formulation as:

$$\begin{aligned} &\text{minimize } \mathbf{c}^T \mathbf{x} \\ &\text{s.t. } A\mathbf{x} \geq \mathbf{b}, \mathbf{x} \geq 0 \end{aligned}$$

The dual problem of the above is

$$\begin{aligned} &\text{maximize } \mathbf{b}^T \boldsymbol{\lambda} \\ &\text{s.t. } A^T \boldsymbol{\lambda} \leq \mathbf{c}, \boldsymbol{\lambda} \geq 0 \end{aligned}$$

We will solve the dual problem by writing it in standard minimize $-\underline{\mathbf{b}}^T \underline{\lambda}$ form and using the simplex method. Let us introduce slack variables $s_1, \dots, s_{\frac{t}{2}+2}$ and re-write the constraints as

$$B\underline{\mathbf{v}} = \underline{\mathbf{c}}, \quad \underline{\mathbf{v}} \geq 0,$$

where

$$B = \begin{bmatrix} r & 0 & 0 & \dots & 0 & (r+3) & 0 & 1 & 0 & 0 & \dots & 0 & 0 \\ -1 & r & 0 & \dots & 0 & (r+2) & 0 & 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & -1 & r & \dots & 0 & (r+2) & 0 & 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & -1 & r & (r+2) & 0 & 0 & 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & 0 & \dots & -2 & 1 & 0 & 0 & 0 & 0 & \dots & 0 & 1 \\ 0 & 0 & 0 & \dots & (r+1) & (r+1) & 1 & 0 & 0 & 0 & \dots & 0 & 0 \end{bmatrix}, \quad (26)$$

and

$$\underline{\mathbf{v}} = [\lambda_1 \quad \lambda_2 \quad \dots \quad \lambda_{\frac{t}{2}+1} \quad s_1 \quad \dots \quad s_{\frac{t}{2}+2}]^T. \quad (27)$$

With this, the objective function now is $\underline{\mathbf{d}}^T \underline{\mathbf{v}}$, where $\underline{\mathbf{d}} = [-\underline{\mathbf{b}}^T \quad 0 \quad 0 \quad \dots \quad 0]^T$ which is a $(t+3) \times 1$ matrix. Define $\underline{\mathbf{v}}_{BV} = [\lambda_1, \dots, \lambda_{\frac{t}{2}+1}, s_1]^T$.

We pick the variables $\beta_1 = \lambda_1, \dots, \beta_{\frac{t}{2}+1} = \lambda_{\frac{t}{2}+1}$, $\beta_{\frac{t}{2}+2} = s_1$ as “basic variables” and the rest, called “non-basic variables” $\alpha_1 = s_2, \dots, \alpha_{\frac{t}{2}+1} = s_{\frac{t}{2}+2}$ will be set to 0. The set of basic variables is chosen such that the matrix formed by the columns of B corresponding to these basic variables is a full-rank square matrix B_{BV} . The remaining columns of B will give a matrix B_{NBV} . The system of equations is now in the following form:

$$[B_{BV} \quad B_{NBV}] \begin{bmatrix} \underline{\mathbf{v}}_{BV} \\ \underline{\mathbf{0}} \end{bmatrix} = \underline{\mathbf{c}}$$

Therefore we will equivalently solve

$$B_{BV} \underline{\mathbf{v}}_{BV} = \underline{\mathbf{c}}$$

The above system of equations can be solved in closed form to get the following:

$$\lambda_{\frac{t}{2}+1} = \frac{2 \sum_{i=0}^{\frac{t}{2}-1} r^i}{3(r^{\frac{t}{2}} + 2 \sum_{i=0}^{\frac{t}{2}-1} r^i)}, \quad (28)$$

$$s_1 = \frac{1}{r^{\frac{t}{2}} + 2 \sum_{i=0}^{\frac{t}{2}-1} r^i}, \quad (29)$$

$$\lambda_{j+1} = \frac{r^{\frac{t}{2}} - 3r^{\frac{t}{2}-(j+1)} + 2}{3(r-1)(r^{\frac{t}{2}} + 2 \sum_{i=0}^{\frac{t}{2}-1} r^i)}, \quad \text{for } 0 \leq j \leq \frac{t}{2} - 1 \quad (30)$$

which are non-negative if $r \geq 3$. Hence the solution given by (28),(29),(30) is a basic feasible solution. Let the elements of the vector $\underline{\mathbf{d}}$ be indexed by the elements of the vector $\underline{\mathbf{v}}$ i.e., i^{th} component of $\underline{\mathbf{d}}$ is indexed by i^{th} component of $\underline{\mathbf{v}}$. To check for optimality we check if the “reduced cost coefficients” $r_{\alpha_i} = d_{\alpha_i} - z_{\alpha_i}$ are non-negative, for every non-basic variable α_i , $1 \leq i \leq \frac{t}{2} + 1$. We note that for the above made choice of basic and non-basic variables, in the vector $\underline{\mathbf{d}}$ only $d_{\beta_{\frac{t}{2}+1}} = -3n$ is non-zero. The quantity z_{α_i} is defined as follows:

$$z_{\alpha_i} = \sum_{j=1}^{\frac{t}{2}+2} d_{\beta_j} y_{(j, \alpha_i)} = d_{\beta_{\frac{t}{2}+1}} y_{(\frac{t}{2}+1, \alpha_i)} = -3n y_{(\frac{t}{2}+1, \alpha_i)}$$

where $y_{(\frac{t}{2}+1, \alpha_i)}$ are as shown in the row reduced echelon form of matrix B below:

$$B_{rref} = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 & 0 & 0 & y(1, \alpha_1) & y(1, \alpha_2) & y(1, \alpha_3) & \dots & y(1, \alpha_{\frac{t}{2}}) & y(1, \alpha_{\frac{t}{2}+1}) \\ 0 & 1 & 0 & \dots & 0 & 0 & 0 & y(2, \alpha_1) & y(2, \alpha_2) & y(2, \alpha_3) & \dots & y(2, \alpha_{\frac{t}{2}}) & y(2, \alpha_{\frac{t}{2}+1}) \\ 0 & 0 & 1 & \dots & 0 & 0 & 0 & y(3, \alpha_1) & y(3, \alpha_2) & y(3, \alpha_3) & \dots & y(3, \alpha_{\frac{t}{2}}) & y(3, \alpha_{\frac{t}{2}+1}) \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 0 & 0 & y(\frac{t}{2}, \alpha_1) & y(\frac{t}{2}, \alpha_2) & y(\frac{t}{2}, \alpha_3) & \dots & y(\frac{t}{2}, \alpha_{\frac{t}{2}}) & y(\frac{t}{2}, \alpha_{\frac{t}{2}+1}) \\ 0 & 0 & 0 & \dots & 0 & 1 & 0 & y(\frac{t}{2}+1, \alpha_1) & y(\frac{t}{2}+1, \alpha_2) & y(\frac{t}{2}+1, \alpha_3) & \dots & y(\frac{t}{2}+1, \alpha_{\frac{t}{2}}) & y(\frac{t}{2}+1, \alpha_{\frac{t}{2}+1}) \\ 0 & 0 & 0 & \dots & 0 & 0 & 1 & y(\frac{t}{2}+2, \alpha_1) & y(\frac{t}{2}+2, \alpha_2) & y(\frac{t}{2}+2, \alpha_3) & \dots & y(\frac{t}{2}+2, \alpha_{\frac{t}{2}}) & y(\frac{t}{2}+2, \alpha_{\frac{t}{2}+1}) \end{bmatrix}$$

We observe that in B to B_{rref} , to row- $(\frac{t}{2}+1)$ only non-negative linear combinations of the rows above it are added, entries of which are either 0 or 1. Therefore $r_{\alpha_i} \geq 0$ for α_i all non-basic variables. Hence $y_{(\frac{t}{2}+1, \alpha_1)}, \dots, y_{(\frac{t}{2}+1, \alpha_{\frac{t}{2}+1})} \geq 0$. Hence the basic solution given by (28), (29) and (30) is an ‘‘optimal basic feasible’’ solution.

By the theorem of strong duality the optimal solutions of the primal problem and the dual problem are equal. Therefore the minimum value of m is $3n\lambda_{\frac{t}{2}+1} = \frac{n2 \sum_{i=0}^{\frac{t}{2}-1} r^i}{(r^{\frac{t}{2}+2} + 2 \sum_{i=0}^{\frac{t}{2}-1} r^i)}$.

Hence we get the upper bound on the rate:

$$\frac{k}{n} \leq 1 - \frac{m}{n} \leq \frac{r^{\frac{t}{2}}}{r^{\frac{t}{2}+2} + 2 \sum_{i=0}^{\frac{t}{2}-1} r^i}$$

We now pick a solution for the primal problem and show that it is feasible and gives the optimal objective function value.

$$a_i = \frac{2nr^i}{r^{\frac{t}{2}+2} + 2 \sum_{i=0}^{\frac{t}{2}-1} r^i}, \text{ for } 0 \leq i \leq \frac{t}{2} - 1$$

$$a_{\frac{t}{2}} = \frac{nr^{\frac{t}{2}}}{r^{\frac{t}{2}+2} + 2 \sum_{i=0}^{\frac{t}{2}-1} r^i}, \quad p = 0.$$

It is easy to check that this solution satisfies the first $\frac{t}{2}$ constraints of the primal problem with equality. It remains to check the following:

$$(r+3)a_0 + (r+2) \sum_{i=1}^{\frac{t}{2}-1} a_i + a_{\frac{t}{2}} \geq 3n.$$

Upon simplification, it is seen that the above inequality is met with equality. Therefore the chosen solution is a feasible solution. It is also easy to check that the solution gives the optimal value of the objective function. Hence it is an optimal feasible solution. We thus conclude that a code having the above chosen values will have the optimal rate. \square

B. Upper Bound on Rate of an $(n, k, r, t)_{seq}$ Code for t Odd:

In this subsection we provide an upper bound on the rate of an $(n, k, r, t)_{seq}$ code whenever t is a positive odd integer and $r \geq 3$.

Theorem 2. *Let C be an $(n, k, r, t)_{seq}$ code over a field \mathbb{F}_q . Let $t = 2s - 1, s \geq 1$ and $r \geq 3$. Then*

$$\frac{k}{n} \leq \frac{r^s}{r^s + 2 \sum_{i=1}^{s-1} r^i + 1}. \quad (31)$$

Proof. As earlier, we again begin by setting $\mathcal{B}_0 = \text{span}(\underline{c} \in \mathcal{C}^\perp : w_H(\underline{c}) \leq r+1)$. Let m be the dimension of \mathcal{B}_0 . Let $\underline{c}_1, \dots, \underline{c}_m$ (row vectors) be a basis of \mathcal{B}_0 such that $w_H(\underline{c}_i) \leq r+1$.

$$\text{Let } H_1 = \begin{bmatrix} \underline{c}_1 \\ \vdots \\ \underline{c}_m \end{bmatrix}.$$

It follows that H_1 is a parity check matrix of an $(n, n - m, r, t)_{seq}$ code as its row space contains all the codewords of

Hamming weight at most $r + 1$ which are contained in \mathcal{C}^\perp . Also,

$$\frac{k}{n} \leq 1 - \frac{m}{n}.$$

We now proceed to derive an upper bound on $1 - \frac{m}{n}$. Again it can be seen that the matrix H_1 , after permutation of rows and columns can be written in the following form

$$H_1 = \left[\begin{array}{c|c|c|c|c|c|c|c|c} D_0 & A_1 & 0 & 0 & \dots & 0 & 0 & 0 & \\ \hline 0 & D_1 & A_2 & 0 & \dots & 0 & 0 & 0 & \\ \hline 0 & 0 & D_2 & A_3 & \dots & 0 & 0 & 0 & \\ \hline 0 & 0 & 0 & D_3 & \dots & 0 & 0 & 0 & \\ \hline \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \\ \hline 0 & 0 & 0 & 0 & \dots & A_{s-2} & 0 & 0 & \\ \hline 0 & 0 & 0 & 0 & \dots & D_{s-2} & A_{s-1} & 0 & \\ \hline 0 & 0 & 0 & 0 & \dots & 0 & D_{s-1} & 0 & \\ \hline 0 & 0 & 0 & 0 & \dots & 0 & 0 & C & \end{array} \right] D, \quad (32)$$

where

- 1) the rows of H_1 are labeled $1, 2, \dots, m$ and columns are labeled $1, 2, \dots, n$,
- 2) A_i is a $\rho_{i-1} \times a_i$ matrix for $1 \leq i \leq s-1$, D_i is a $\rho_i \times a_i$ matrix for $0 \leq i \leq s-1$ for some $\{\rho_i\}, \{a_i\}$,
- 3) D_0 is a matrix with each column having weight 1 and each row having weight at least 1,
- 4) $\{A_i\}, \{D_i\}, \{B_i = \begin{bmatrix} A_i \\ D_i \end{bmatrix}\}$ are such that for $1 \leq i \leq s-1$ each column of B_i has weight 2, each column of A_i has weight at least 1, each row of D_i has weight at least 1 and each column of D_i has weight at most 1,
- 5) C is a matrix with each column having weight 2, D is a matrix which exactly contains all the columns of H_1 which have weight at least 3,
- 6) If J is the first index such that A_J, D_J are empty matrices ($0 \times L$, $L \times 0$, 0×0 matrix) ($J = 0$ if D_0 is an empty matrix) then we set A_i, D_i to be empty matrices for all $s-1 \geq i \geq J$ and set $a_i = 0, \rho_i = 0, \forall s-1 \geq i \geq J$. If none of $D_0, A_i, D_i, 1 \leq i \leq s-1$ are empty matrices, then we simply set $J = s$. If $J < s$, let $\alpha_1, \dots, \alpha_l \in \{\sum_{j=0}^{J-1} a_j + 1, \dots, n\}$ be the indices of 2-weight columns in H_1 apart from the 2-weight columns corresponding to columns of B_1, \dots, B_{J-1} . We take the submatrix of H_1 with rows indexed by $\sum_{j=0}^{J-1} \rho_j + 1, \dots, m$ and columns indexed by $\alpha_1, \dots, \alpha_l$ and name the submatrix as matrix C . Let the number of columns in C be a_s . If C is an empty matrix then we set $a_s = 0$. If $J < s$, the matrix H_1 can be written in the form (33).

$$H_1 = \left[\begin{array}{c|c|c|c|c|c|c|c|c} D_0 & A_1 & 0 & 0 & \dots & 0 & 0 & 0 & \\ \hline 0 & D_1 & A_2 & 0 & \dots & 0 & 0 & 0 & \\ \hline 0 & 0 & D_2 & A_3 & \dots & 0 & 0 & 0 & \\ \hline 0 & 0 & 0 & D_3 & \dots & 0 & 0 & 0 & \\ \hline \vdots & \vdots & \vdots & \vdots & \dots & \vdots & \vdots & \vdots & \\ \hline 0 & 0 & 0 & 0 & \dots & A_{J-2} & 0 & 0 & \\ \hline 0 & 0 & 0 & 0 & \dots & D_{J-2} & A_{J-1} & 0 & \\ \hline 0 & 0 & 0 & 0 & \dots & 0 & D_{J-1} & 0 & \\ \hline 0 & 0 & 0 & 0 & \dots & 0 & 0 & C & \end{array} \right] D, \quad (33)$$

The entire rate bound derivation is correct and all the bounds will hold with $a_i = 0, \rho_i = 0, J \leq i \leq s-1$. If $l (< J)$ is the first index such that D_l is an empty matrix with A_l a non empty matrix then the proof of the following claim 2 (since in the proof of claim 2, we prove the claim for A_l first and then proceed to D_l and since D_0, A_j, D_j must be non empty $\forall j < l$) will imply that A_l has column weight 1 which will imply D_l cannot be empty. Hence the case A_l , a non empty matrix and D_l , an empty matrix cannot occur. Although we have to prove the following claim 2 for $A_i, D_i, 1 \leq i \leq J-1, D_0$, we assume all $D_0, A_i, D_i, 1 \leq i \leq s-1$ to be non-empty matrices and prove the claim. Since the proof is by induction, the induction can be made to stop at $J-1$ (induction starts at 0) and the proof is unaffected by it.

Claim 2. For $1 \leq i \leq s-1$, A_i is a matrix with each column having weight 1 and for $0 \leq i \leq s-2$, D_i is a matrix with each row and each column having weight 1. D_{s-1} is a matrix with each column having weight 1.

Proof. Proof is exactly similar to the proof of claim 1. So we skip the proof. \square

By Claim 2, after permutation of columns of H_1 (in (32) or (33) depending on J) within the columns labeled by the set $\{\sum_{l=0}^{j-1} a_l + 1, \dots, \sum_{l=0}^{j-1} a_l + a_j\}$ for $0 \leq j \leq \min(J-1, s-2)$, the matrix $D_j, 0 \leq j \leq \min(J-1, s-2)$ can be assumed to be a diagonal matrix with non-zero entries along the diagonal and hence $\rho_i = a_i$, for $0 \leq i \leq s-2$.

Since the sum of the weights of the columns of $A_i, 1 \leq i \leq s-1$ must equal the sum of the row weights and since each row of A_i for $i \leq J-1$ can have weight at most r and not $r+1$ due to weight one rows in D_{i-1} , and since for $s-1 \geq i \geq J$, A_i is an empty matrix and we have set $a_i = 0$, we obtain:

$$\begin{aligned} \text{For } 1 \leq i \leq s-1 : \\ \rho_{i-1}r &\geq a_i, \\ a_{i-1}r &\geq a_i. \end{aligned} \quad (34)$$

For some $p \geq 0$,

$$\rho_{s-1} + \sum_{i=0}^{s-2} a_i + p = m. \quad (35)$$

By equating sum of row weights of $[\frac{D_{s-1}}{0}|C]$, with sum of column weights of $[\frac{D_{s-1}}{0}|C]$, we obtain:

$$2a_s + a_{s-1} \leq (\rho_{s-1} + p)(r+1). \quad (36)$$

If $J < s$ then the number of rows in C is p with each column of C having weight 2 and $a_{s-1} = \rho_{s-1} = 0$ (and $a_s = 0$ if C is also an empty matrix), hence the inequality (36) is true. If $J = s$ and C is an empty matrix then also the inequality (36) is true as we would have set $a_s = 0$.

Substituting (35) in (36):

$$2a_s \leq (m - \sum_{i=0}^{s-2} a_i)(r+1) - a_{s-1}. \quad (37)$$

By equating sum of row weights of D_{s-1} , with sum of column weights of D_{s-1} , we obtain (Note that if D_{s-1} is an empty matrix then also the following inequality is true as we would have set $a_{s-1} = 0$):

$$a_{s-1} \leq \rho_{s-1}(r+1). \quad (38)$$

By equating sum of row weights of H_1 , with sum of column weights of H_1 , we obtain

$$m(r+1) \geq a_0 + 2\left(\sum_{i=1}^s a_i\right) + 3\left(n - \sum_{i=0}^s a_i\right), \quad (39)$$

If $J < s$ then $a_i = 0 \forall J \leq i \leq s-1$. If C is an empty matrix then $a_s = 0$. Hence the inequality (39) is true irrespective of whether $J = s$ or $J < s$ (even if C is an empty matrix).

$$m(r+1) \geq 3n - 2a_0 - \left(\sum_{i=1}^s a_i\right). \quad (40)$$

Our basic inequalities are (34),(35),(36),(38),(39). We manipulate these 5 inequalities to derive the bound on rate. Substituting (37) in (40):

$$m(r+1) \geq 3n - 2a_0 - \left(\sum_{i=1}^{s-1} a_i\right) - \left(\frac{(m - \sum_{i=0}^{s-2} a_i)(r+1) - a_{s-1}}{2}\right). \quad (41)$$

For $s = 1$, (41) becomes:

$$\begin{aligned} m(r+1) &\geq 3n - 2a_0 - \left(\frac{m(r+1) - a_0}{2}\right), \\ m\frac{3(r+1)}{2} &\geq 3n - \frac{3}{2}a_0. \end{aligned} \quad (42)$$

Substituting (38) in (42):

$$m\frac{3(r+1)}{2} \geq 3n - \frac{3}{2}\rho_0(r+1), \quad (43)$$

Substituting (35) in (43):

$$m \frac{3(r+1)}{2} \geq 3n - \frac{3}{2}(m-p)(r+1),$$

Since $p \geq 0$, $3m(r+1) \geq 3n$. (44)

(44) implies,

$$\frac{k}{n} \leq \frac{r}{r+1}. \quad (45)$$

(45) proves the bound (31) for $s = 1$. Hence from now on we assume $s \geq 2$.

For $s \geq 2$, (41) implies:

$$m \frac{3(r+1)}{2} \geq 3n + a_0 \left(\frac{r+1}{2} - 2 \right) + \left(\sum_{i=1}^{s-2} a_i \right) \left(\frac{r+1}{2} - 1 \right) - \frac{a_{s-1}}{2}. \quad (46)$$

Substituting (34) in (46) and since $r \geq 3$:

$$\begin{aligned} m \frac{3(r+1)}{2} &\geq 3n + \frac{a_{s-1}}{r^{s-1}} \left(\frac{r+1}{2} - 2 \right) + \left(\sum_{i=1}^{s-2} \frac{a_{s-1}}{r^{s-1-i}} \right) \left(\frac{r+1}{2} - 1 \right) - \frac{a_{s-1}}{2}, \\ m \frac{3(r+1)}{2} &\geq 3n + a_{s-1} \left(\left(\sum_{i=1}^{s-1} \frac{1}{r^i} \right) \left(\frac{r+1}{2} - 1 \right) - \frac{1}{r^{s-1}} - \frac{1}{2} \right), \\ m \frac{3(r+1)}{2} &\geq 3n - a_{s-1} \left(\frac{3}{2r^{s-1}} \right). \end{aligned} \quad (47)$$

Rewriting (35):

$$\rho_{s-1} + \sum_{i=0}^{s-2} a_i + p = m. \quad (48)$$

Substituting (38),(34) in (48):

$$\begin{aligned} \rho_{s-1} + \sum_{i=0}^{s-2} a_i + p &= m, \\ \frac{a_{s-1}}{r+1} + \sum_{i=0}^{s-2} \frac{a_{s-1}}{r^{s-1-i}} &\leq m - p, \\ a_{s-1} &\leq \frac{m-p}{\frac{1}{r+1} + \sum_{i=1}^{s-1} \frac{1}{r^i}}, \\ a_{s-1} &\leq \frac{(m-p)(r+1)}{1 + \frac{(r^{s-1}-1)(r+1)}{(r-1)}}. \end{aligned} \quad (49)$$

Substituting (49) in (47):

$$m \frac{3(r+1)}{2} \geq 3n - \frac{(m-p)(r+1)}{1 + \frac{(r^{s-1}-1)(r+1)}{(r-1)}} \left(\frac{3}{2r^{s-1}} \right),$$

Since $p \geq 0$, $m \frac{3(r+1)}{2} \left(1 + \frac{1}{r^{s-1} + \frac{(r^{s-1}-1)(r+1)}{(r-1)}} \right) \geq 3n$. (50)

(50) after some algebraic manipulations gives the required bound on $1 - \frac{m}{n}$ and hence on $\frac{k}{n}$ as stated in the theorem. \square

Again an alternative proof for Theorem 2 is given below by using linear programming:

Proof. The inequalities (34),(36),(38) and (40) are linear inequalities and are written in matrix form as:

$$A\mathbf{x} \geq \mathbf{b}$$

where

$$A = \begin{bmatrix} r & -1 & 0 & \dots & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & r & -1 & \dots & 0 & 0 & 0 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & r & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \dots & 0 & -1 & (r+1) & -2 & (r+1) & 0 \\ 0 & 0 & 0 & \dots & 0 & -1 & (r+1) & 0 & 0 & 0 \\ (r+3) & (r+2) & (r+2) & \dots & (r+2) & 1 & (r+1) & 0 & (r+1) & 0 \end{bmatrix} \quad (51)$$

which is a $(s+2) \times (s+3)$ matrix and

$$\underline{\mathbf{x}} = [a_0 \ a_1 \ \dots \ a_{s-1} \ \rho_{s-1} \ a_s \ p]^T, \underline{\mathbf{b}} = [0 \ 0 \ \dots \ 0 \ 3n]^T \quad (52)$$

where $\underline{\mathbf{x}}$ is an $(s+3) \times 1$ matrix and $\underline{\mathbf{b}}$ is an $(s+2) \times 1$ matrix. The problem of finding an upper bound on rate of the code now becomes one of minimizing $m = \underline{\mathbf{c}}^T \underline{\mathbf{x}}$, which is a linear objective function where $\underline{\mathbf{c}} = [1 \ 1 \ \dots \ 1 \ 0 \ 1 \ 0 \ 1]^T$ is an $(s+3) \times 1$ matrix. Also by definition of $\underline{\mathbf{x}}$, $\underline{\mathbf{x}} \geq 0$. This is now in a standard form of a linear program formulation as:

$$\begin{aligned} & \text{minimize } \underline{\mathbf{c}}^T \underline{\mathbf{x}} \\ & \text{s.t. } A \underline{\mathbf{x}} \geq \underline{\mathbf{b}} \\ & \quad \underline{\mathbf{x}} \geq 0 \end{aligned}$$

The dual problem of the above is

$$\begin{aligned} & \text{maximize } \underline{\mathbf{b}}^T \underline{\lambda} \\ & \text{s.t. } A^T \underline{\lambda} \leq \underline{\mathbf{c}} \\ & \quad \underline{\lambda} \geq 0 \end{aligned}$$

We will solve the dual problem by writing it in standard minimize $-\underline{\mathbf{b}}^T \underline{\lambda}$ form and using the simplex method. Let us introduce slack variables h_1, \dots, h_{s+3} and re-write the constraints as

$$B \underline{\mathbf{v}} = \underline{\mathbf{c}}, \underline{\mathbf{v}} \geq 0,$$

where

$$B = \begin{bmatrix} r & 0 & 0 & \dots & 0 & 0 & 0 & (r+3) & 0 & 1 & 0 & 0 & \dots & 0 & 0 & 0 & 0 & 0 \\ -1 & r & 0 & \dots & 0 & 0 & 0 & (r+2) & 0 & 0 & 1 & 0 & \dots & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & r & \dots & 0 & 0 & 0 & (r+2) & 0 & 0 & 0 & 1 & \dots & 0 & 0 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & r & 0 & 0 & (r+2) & 0 & 0 & 0 & 0 & \dots & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \dots & -1 & -1 & -1 & 1 & 0 & 0 & 0 & 0 & \dots & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & \dots & 0 & (r+1) & (r+1) & (r+1) & 0 & 0 & 0 & 0 & \dots & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & \dots & 0 & -2 & 0 & 0 & 0 & 0 & 0 & 0 & \dots & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & \dots & 0 & (r+1) & 0 & (r+1) & 1 & 0 & 0 & 0 & \dots & 0 & 0 & 0 & 0 & 0 \end{bmatrix}, \quad (53)$$

and

$$\underline{\mathbf{v}} = [\lambda_1 \ \lambda_2 \ \dots \ \lambda_{s+2} \ h_1 \ \dots \ h_{s+3}]^T. \quad (54)$$

With this, the objective function now is $\underline{\mathbf{d}}^T \underline{\mathbf{v}}$, where $\underline{\mathbf{d}} = [-\underline{\mathbf{b}}^T \ 0 \ 0 \ \dots \ 0]^T$ which is an $(2s+5) \times 1$ matrix. Define $\underline{\mathbf{v}}_{BV} = [\lambda_1, \dots, \lambda_{s+2}, h_1]^T$.

We pick the variables $\beta_1 = \lambda_1, \dots, \beta_{s+2} = \lambda_{s+2}, \beta_{s+3} = h_1$ as “basic variables” and the rest, called “non-basic variables” $\alpha_1 = h_2, \dots, \alpha_{s+2} = h_{s+3}$ will be set to 0. The set of basic variables is chosen such that the columns of B corresponding to these basic variables is a full-rank square matrix B_{BV} . The remaining columns of B will give a matrix B_{NBV} . The system of equations is now in the following form:

$$[B_{BV} \ B_{NBV}] \begin{bmatrix} \underline{\mathbf{v}}_{BV} \\ \underline{\mathbf{0}} \end{bmatrix} = \underline{\mathbf{c}}$$

Therefore we will equivalently solve

$$B_{BV} \underline{\mathbf{v}}_{BV} = \underline{\mathbf{c}}$$

The above system of equations can be solved in closed form to get the following:

$$\lambda_{s+2} = \frac{2 \sum_{i=1}^{s-1} r^i + 1}{3(r^s + 2 \sum_{i=1}^{s-1} r^i + 1)}, \quad (55)$$

$$h_1 = \frac{\sum_{i=1}^{s-1} r^i + 2}{3(\sum_{i=0}^{s-1} r^i)}, \quad (56)$$

$$\lambda_{j+1} = \frac{r^s - 3r^{s-n-1} + r + 1}{3(r+1)(r^s - 1)}, \quad \text{for } 0 \leq j \leq s-2 \quad (57)$$

$$\lambda_s = 0, \quad (58)$$

$$\lambda_{s+1} = \frac{\sum_{i=1}^{s-1} r^i + 2}{3(r+1)(\sum_{i=0}^{s-1} r^i)}. \quad (59)$$

which are non-negative if $r \geq 3$.

Hence the solution given by (55),(56),(57),(58) and (59) is a basic feasible solution. Let the elements of the vector \underline{d} be indexed by the elements of the vector \underline{v} i.e., i^{th} component of \underline{d} is indexed by i^{th} component of \underline{v} . To check for optimality we check if the ‘‘reduced cost coefficients’’ $r_{\alpha_i} = d_{\alpha_i} - z_{\alpha_i}$ are non-negative, for every non-basic variable $\alpha_i, 1 \leq i \leq s+2$. We note that for the above made choice of basic and non-basic variables, in the vector \underline{d} only $d_{\beta_{s+2}} = -3n$ is non-zero. The quantity z_{α_i} is defined as follows:

$$z_{\alpha_i} = \sum_{j=1}^{s+3} d_{\beta_j} y_{(j,\alpha_i)} = d_{\beta_{s+2}} y_{(s+2,\alpha_i)} = -3n y_{(s+2,\alpha_i)}$$

where $y_{(s+2,\alpha_i)}$ are as shown in the row reduced echelon form of matrix B below:

$$B_{rref} = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 & 0 & 0 & y_{(1,\alpha_1)} & y_{(1,\alpha_2)} & y_{(1,\alpha_3)} & \dots & y_{(1,\alpha_{s+1})} & y_{(1,\alpha_{s+2})} \\ 0 & 1 & 0 & \dots & 0 & 0 & 0 & y_{(2,\alpha_1)} & y_{(2,\alpha_2)} & y_{(2,\alpha_3)} & \dots & y_{(2,\alpha_{s+1})} & y_{(2,\alpha_{s+2})} \\ 0 & 0 & 1 & \dots & 0 & 0 & 0 & y_{(3,\alpha_1)} & y_{(3,\alpha_2)} & y_{(3,\alpha_3)} & \dots & y_{(3,\alpha_{s+1})} & y_{(3,\alpha_{s+2})} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 0 & 0 & y_{(s+1,\alpha_1)} & y_{(s+1,\alpha_2)} & y_{(s+1,\alpha_3)} & \dots & y_{(s+1,\alpha_{s+1})} & y_{(s+1,\alpha_{s+2})} \\ 0 & 0 & 0 & \dots & 0 & 1 & 0 & y_{(s+2,\alpha_1)} & y_{(s+2,\alpha_2)} & y_{(s+2,\alpha_3)} & \dots & y_{(s+2,\alpha_{s+1})} & y_{(s+2,\alpha_{s+2})} \\ 0 & 0 & 0 & \dots & 0 & 0 & 1 & y_{(s+3,\alpha_1)} & y_{(s+3,\alpha_2)} & y_{(s+3,\alpha_3)} & \dots & y_{(s+3,\alpha_{s+1})} & y_{(s+3,\alpha_{s+2})} \end{bmatrix}$$

We observe that in reducing B to B_{rref} , to row- $(s+2)$ only non-negative linear combinations of the rows above it are added, entries of which are either 0 or 1. Therefore $r_{\alpha_i} \geq 0$ for α_i all non-basic variables. Hence $y_{(s+2,\alpha_1)}, \dots, y_{(s+2,\alpha_M)} \geq 0$. Hence the basic solution is an ‘‘optimal basic feasible’’ solution.

By the theorem of strong duality the optimal solutions of the primal problem and the dual problem are equal. Therefore the minimum value of m is $3n\lambda_{s+2} = \frac{n(2 \sum_{i=1}^{s-1} r^i + 1)}{(r^s + 2 \sum_{i=1}^{s-1} r^i + 1)}$.

Hence we get the upper bound on the rate:

$$\frac{k}{n} \leq 1 - \frac{m}{n} \leq \frac{r^s}{(r^s + 2 \sum_{i=1}^{s-1} r^i + 1)}$$

We now pick a solution for the primal problem and show that it is feasible and gives the optimal objective function value.

$$a_i = \frac{nr^i(r+1)}{r^s + 2 \sum_{i=1}^{s-1} r^i + 1}, \quad \text{for } 0 \leq i \leq s-1$$

$$\rho_{s-1} = \frac{nr^{s-1}}{r^s + 2 \sum_{i=1}^{s-1} r^i + 1}, \quad a_s = 0, \quad p = 0.$$

It is easy to check that this solution satisfies the constraints of the primal problem with equality. Therefore the chosen solution is a feasible solution. It is also easy to check that the solution gives the optimal value of the objective function. Hence it is an optimal feasible solution. We thus conclude that a code having the above chosen values will have the optimal rate. \square

From the two derivations, it becomes clear that a code achieving the rate bound given in (1) will have parity check matrix of the form (2) with D being an empty matrix and all the inequalities (4),(5),(6),(9) met with equality. A similar observation holds for t odd also. It may be noted here that our bound, for the special cases of $t = 2, 3, 4$, matches with the rate bound given in [8],[13],[12] respectively. In the rest of the paper, the codes achieving the bounds (1) or (31) depending on t will be referred to as ‘‘rate-optimal codes’’.

Remark 1. We now make a remark on the blocklength of the rate-optimal codes. From the proofs it can be seen that, for t even, for the optimal values of $a_0, \dots, a_{\frac{t}{2}}$ to be integral, $2n$ needs to be an integer multiple of $r^{\frac{t}{2}} + 2 \sum_{i=0}^{\frac{t}{2}-1} r^i$. Similarly for t odd, for the optimal values of $a_0, \dots, a_{s-1}, \rho_{s-1}$ to be integral, $(r+1)n$ needs to be an integer multiple of $r^s + 2 \sum_{i=1}^{s-1} r^i + 1$ and nr^{s-1} needs to be an integer multiple of $r^s + 2 \sum_{i=1}^{s-1} r^i + 1$.

Remark 2. The linear program was numerically solved for integer values of the variables and objective function. The solution for the case of $t = 4$ and $r = 5$ is shown in figure 2. Notice that integer solutions achieving our rate bound are possible for a set of blocklengths in a periodic fashion as predicted by theoretical solution.




Fig. 2: Plot showing integer solution for the linear program formulation. Circled points are the values of blocklength that allow for integer solution achieving our rate bound.




Fig. 3: Comparison of rate bounds on codes with availability and codes with sequential recovery for an example case $t = 10$.

Remark 3. In general, one would expect an $(n, k, r, t)_{seq}$ code to achieve a higher rate than a counterpart code having t -availability. The achievable upper bound on rate of a code with sequential recovery is significantly larger than the upper bound on the rate of codes having availability given in [16] as shown in Fig. 3.

III. CONSTRUCTION OF CODES ACHIEVING THE UPPER BOUND ON RATE

In this section we give a construction of codes with sequential recovery achieving the rate bound (1) for any $r \geq 3, t = 2s, s \geq 1$ and also give a construction achieving (31) for any $r \geq 3, t = 2s - 1, s \geq 1$.

A. Construction of Binary Rate-Optimal Codes for t Even

In this subsection we give a construction of rate-optimal codes for any $r \geq 3$ and t -even. While the rate bound is independent of the size of the field over which the code is defined, our construction provides optimal binary codes. A construction achieving the bound (1) for the special case of $t = 2$ was provided in [8], [10], [1] and for the case of $t = 4$, in [12].

We provide an iterative, graph-based construction of rate-optimal codes for a given $r \geq 3$ and t even. We build a graph $G_{\frac{t}{2}-1}$ iteratively, starting from G_0 , by adding nodes in vertical, layer-by-layer fashion. At every stage of the iteration, the graph G_i thus constructed, will always have girth $\geq t + 1$. We then define our rate-optimal code based on the graph $G_{\frac{t}{2}-1}$.

Construction of graph $G_{\frac{t}{2}-1}$: Denote the vertex set of a graph G by $V(G)$. Let $N(v)$ denote the neighbors of a node v in a graph. We make use of graphs G_0 and B_i , for $1 \leq i \leq \frac{t}{2} - 1$ described below as the ingredients:

- 1) We pick G_0 as follows: Let G_0 be an r -regular graph with girth $\geq t + 1^\dagger$. Define $U_0 = V(G_0)$, $|U_0| = u_0$.
- 2) Next, for i in the range $1 \leq i \leq \frac{t}{2} - 1$, we iteratively pick an (r, u_{i-1}) -biregular bipartite graph B_i with girth $g_i \geq \left\lceil \frac{t+1}{i+\frac{1}{2}} \right\rceil$.
 \dagger . Let U_i, L_i be the two (upper and lower) sets of nodes in the bipartite graph with $\text{degree}(x) = r, \forall x \in U_i$ and $\text{degree}(y) = u_{i-1}, \forall y \in L_i$. Hence $V(B_i) = U_i \cup L_i$. and Let $|U_i| = u_i$ and $|L_i| = l_i$. By edge counting, we have $ru_i = u_{i-1}l_i$.

The flowchart shown in Fig. 4 describes the iterative construction of the graph $G_{\frac{t}{2}-1}$. An example construction of G_1 for $t = 4, r = 3$ depicting the steps of the flowchart for the case of a single iteration appears in Fig. 5. We make the following observations:

- 1) For $0 \leq i \leq \frac{t}{2} - 1$, there are $f_i = \prod_{j=i+1}^{\frac{t}{2}-1} l_j$ disjoint copies of G_i that reside inside $G_{\frac{t}{2}-1}$ as subgraphs due to the replication steps in the construction of $G_{\frac{t}{2}-1}$ (given in the flowchart Fig 4). For all $\frac{t}{2} - 1 > j > i$, since copies of G_i reside inside G_j as subgraphs, when we replicate G_j in the $(j + 1)^{\text{th}}$ replication step, all the copies of G_i contained within G_j also get replicated; we refer to all such copies of G_i here. Let us denote the disjoint union of these subgraphs corresponding to all copies of G_i formed due to replication steps by $G_i^{\text{rep}} = \bigcup_{j=1}^{f_i} (G_i)_j$, where $(G_i)_j$ is the j^{th} copy of G_i in $G_{\frac{t}{2}-1}$ formed due to some replication step. We view G_i^{rep} as a subgraph of $G_{\frac{t}{2}-1}$.
- 2) Note that the construction of $G_{\frac{t}{2}-1}$ proceeds by adding nodes layer by layer with each layer connecting to the layer below it in a tree-like fashion while maintaining the girth of the overall graph to be atleast $t + 1$. Let $N(v)$ represent neighbors of a node v in $G_{\frac{t}{2}-1}$. Let $(U_i)_j$ be the copy of U_i in $V((G_i)_j), \forall 1 \leq j \leq f_i$. For $1 \leq i \leq \frac{t}{2} - 1$, the nodes $T_i = \bigcup_{j=1}^{f_i} (U_i)_j$ represent the nodes in the i^{th} layer and they connect in a tree-like fashion to the $(i - 1)^{\text{th}}$ layer nodes $T_{i-1} = \bigcup_{j=1}^{f_{i-1}} (U_{i-1})_j$. The connection is tree-like because for $v \in T_i$, let $S_v = N(v) \cap T_{i-1}$ then $S_v \cap S_w = \emptyset$ and $|S_v| = |S_w| = r, \forall v, w \in T_i, v \neq w$ and $N(v) \cap T_p = \emptyset, \forall p \notin \{i - 1, i + 1\}, \forall v \in T_i$.
- 3) **Girth of G_i :** Let the girth of G_{i-1} be at least $t + 1$. Consider the construction of graph G_i from l_i replicated copies of G_{i-1} and the biregular bipartite graph B_i . $(U_{i-1})_j$ is the copy of U_{i-1} in the j^{th} copy of G_{i-1} i.e., $(G_{i-1})_j, 1 \leq j \leq l_i$ appearing at the replication step in the construction of G_i from G_{i-1} . Consider a cycle in G_i . If the cycle does not involve any nodes in U_i then it must be completely contained in $(G_{i-1})_j$ for some $1 \leq j \leq l_i$ and hence of length at least $t + 1$ because the girth of G_{i-1} is assumed to be at least $t + 1$. If the cycle involves a node $v_1 \in U_i$ then the cycle must be of the form $v_1 P_{a_1 b_1} v_2 \dots v_q P_{a_q b_q} v_1$ (as shown in fig 8) where $v_m \in U_i, a_m, b_m \in \bigcup_{j=1}^{l_i} (U_{i-1})_j$ and $P_{a_m b_m}$ (shown in green in fig 8) is a path in $(G_{i-1})_p$ for some $1 \leq p \leq l_i$ starting at a_m and ending at b_m for $1 \leq m \leq q$. This is because when we start from v_1 and move to a node say a_1 in $\bigcup_{j=1}^{l_i} (U_{i-1})_j$ the only way to come back to v_1 for completing the cycle is by traversing a path $P_{a_1 b_1}$ in $(G_{i-1})_p$ for some $1 \leq p \leq l_i$ ending at some node $b_1 \in (U_{i-1})_p$. By repeating the above argument it can be seen that the cycle must be of the form $v_1 P_{a_1 b_1} v_2 \dots v_q P_{a_q b_q} v_1$. Now by above argument, for $1 \leq m \leq q$, let $a_m, b_m \in (U_{i-1})_{\sigma_m}, 1 \leq \sigma_m \leq l_i$. Hence a_m, b_m must have been merged with some two nodes in $V_{\sigma_m}^i$ (nodes coming from splitting $v_{\sigma_m}^i \in L_i$ into degree 1 nodes) in the construction of G_i . Hence $v_1 v_{\sigma_1}^i v_2 v_{\sigma_2}^i \dots v_q v_{\sigma_q}^i v_1$ must contain a non trivial cycle in B_i . Also the length of the path $P_{a_m b_m}$ is at least $2(i - 1) + 1$ due to the tree-like structure of $(G_{i-1})_{\sigma_m}$. Hence if the girth of B_i is g_i then the length of the cycle $v_1 P_{a_1 b_1} v_2 \dots v_q P_{a_q b_q} v_1$ is at least $2q + q(2(i - 1) + 1) \geq g_i + \frac{q}{2}(2(i - 1) + 1) = g_i(i + \frac{1}{2}) \geq \frac{t+1}{i+\frac{1}{2}}(i + \frac{1}{2}) = t + 1$. Hence girth of G_i is at least $t + 1$.

Description of a rate-optimal code \mathcal{C} based on $G_{\frac{t}{2}-1}$:

Code \mathcal{C} is defined on the graph $G_{\frac{t}{2}-1}$ as follows:

- 1) The edges of G_0^{rep} represent information symbols. It can be seen that the number of information symbols is $k = \frac{u_0 r}{2} \prod_{j=1}^{\frac{t}{2}-1} l_j$.
- 2) Every node of $G_{\frac{t}{2}-1}$ represents a distinct parity symbol described as follows.
- 3) A node $v \in V(G_0^{\text{rep}})$ represents a parity symbol which is the binary sum of information symbols that are represented by edges in G_0^{rep} incident on v .

\dagger Such a graph can be constructed, due to [20]




Fig. 4: Flowchart showing the iterative construction of $G_{\frac{t}{2}-1}$ for $t \geq 4$. For $t = 2$, $G_{\frac{t}{2}-1} = G_0$. Hence the graph G_0 mentioned in the first step of the flow chart will be our required graph construction for $t = 2$.

- 4) For $1 \leq i \leq \frac{t}{2} - 1$, a node $v \in T_i$ in $G_{\frac{t}{2}-1}$ represents a parity symbol which is the binary sum of code symbols that are represented by the nodes in $N(v) \cap T_{i-1}$.
- 5) It can be seen that the total number of nodes is $n - k = \sum_{i=0}^{\frac{t}{2}-1} u_i \prod_{j=i+1}^{\frac{t}{2}-1} l_j = \sum_{i=0}^{\frac{t}{2}-1} \frac{u_0}{r^i} \prod_{j=1}^{\frac{t}{2}-1} l_j$.
- 6) \mathcal{C} is defined by the information symbols represented by edges in G_0^{rep} and parity symbols represented by nodes in $G_{\frac{t}{2}-1}$.
- 7) From the above counts, the rate of the code \mathcal{C} can be seen to be equal to the bound given by (1).

We now prove that \mathcal{C} can correct t erasures sequentially.

- 1) The Tanner graph of our code can be viewed as follows: We now view $G_{\frac{t}{2}-1}$ differently to give an alternative description of our code. In the graph $G_{\frac{t}{2}-1}$, let every edge represent a code symbol and every node represent a parity check of the code symbols corresponding to the edges incident on it. Apart from this, for every node in $U_{\frac{t}{2}-1} (\subseteq V(G_{\frac{t}{2}-1}))$ there is one distinct degree-1 variable node attached to it. Each of these degree-1 variable nodes corresponds to a distinct code symbol. The code symbol corresponding to a degree-1 variable node is part of the parity check represented by the node




Fig. 5: Depiction of steps mentioned in the flowchart Fig 4 for an example construction of G_1 for $t = 4$, $r = 3$. Here G_0 is the Petersen graph. For the purpose of representation the merger of nodes is shown only for 9 nodes by the dashed lines.




Fig. 6: Example of splitting a node into degree-1 nodes

in $U_{\frac{t}{2}-1}$ to which it is attached. Our code is now defined by the code symbols represented by the edges of $G_{\frac{t}{2}-1}$ and degree-1 variable nodes attached to $U_{\frac{t}{2}-1}$. This completes the alternative description of our code.

The alternative description of the code can be illustrated using the figures 9, 10 and 11.

- Consider a node $v \in T_i$ for $1 \leq i \leq \frac{t}{2} - 2$ in $G_{\frac{t}{2}-1}$. In the original description of the code, the node v represents a code symbol say c_p which is the binary sum of r symbols say c_1, c_2, \dots, c_r where c_1, c_2, \dots, c_r are represented by nodes in $N(v) \cap T_{i-1}$; in the alternative description of the code the same node v now represents the parity check $c_p \oplus c_1 \oplus c_2 \oplus \dots \oplus c_r = 0$ and as shown in figure 9 the edge going out upward from the node v represents the code symbol c_p and the edges going out downward from v represent the code symbols c_1, c_2, \dots, c_r .
- Consider a node $v \in U_{\frac{t}{2}-1}$ in $G_{\frac{t}{2}-1}$. In the original description of the code, the node v represents a code symbol say c_p which is the binary sum of r symbols say c_1, c_2, \dots, c_r where c_1, c_2, \dots, c_r are represented by nodes in $N(v) \cap T_{\frac{t}{2}-2}$; in the alternative description of the code the same node v now represents the parity check $c_p \oplus c_1 \oplus c_2 \oplus \dots \oplus c_r = 0$ and as shown in figure 10 the degree-1 node attached to v represents the code symbol c_p and the edges going out downward from v represent the code symbols c_1, c_2, \dots, c_r . Hence there is a unique distinct degree-1 node attached to every node in $U_{\frac{t}{2}-1}$. These are exactly the degree-1 variable nodes we described above in the alternative description of the code.
- Consider a node $v \in T_0$ in $G_{\frac{t}{2}-1}$. In the original description of the code, the node v represents a code symbol say c_p which is the binary sum of r information symbols say i_1, i_2, \dots, i_r where i_1, i_2, \dots, i_r are represented by edges incident on v in G_0^{rep} ; in the alternative description of the code the same node v now represents the parity check





Fig. 7: Example of merging two nodes

Fig. 8: Figure showing a cycle in the graph G_i . Only the edges constituting a cycle are shown.

$c_p \oplus i_1 \oplus i_2 \oplus \dots \oplus i_r = 0$ and as shown in figure 11 the edge going out upward from v represents the code symbol c_p and the edges going out downward from v still represents the information symbols i_1, i_2, \dots, i_r .

- We transform the meaning of nodes and edges in $G_{\frac{t}{2}-1}$ from original to alternative description of the code as described above across nodes and edges of the graph $G_{\frac{t}{2}-1}$ and it can be done maintaining consistency across the transformations of meaning of nodes and edges from original to alternative description.
- By the above explanation, all the parity checks necessary to describe our code are represented by nodes in the graph $G_{\frac{t}{2}-1}$ in the alternative description.

Note that degree-1 variable nodes are not part of $G_{\frac{t}{2}-1}$ and these degree-1 variable nodes are the reason for rate optimality.

- 2) Let E be a set of erased symbols. Let the code symbols corresponding to degree-1 variable nodes be D_1 . Let $E \cap D_1 = \emptyset$. Let an edge (v_1, v_2) represent an erased symbol in E . Construct a maximal walk $P = v_1 v_2 \dots v_w, v_i \in V(G_{\frac{t}{2}-1}), \forall i \in [w]$ such that $v_i \neq v_j, \forall w \geq i \neq j \geq 1$, and the edge (v_i, v_{i+1}) represents an erased code symbol in $E \forall i \in [w-1]$ (walk P is maximal in terms of number of vertices). Since the walk is maximal any edge of the form (v_w, v) for $v \in N(v_w) - \{v_{w-1}\}, v \in V(G_{\frac{t}{2}-1})$:
 - a) either represents an unerased symbol for all $v \in N(v_w) - \{v_{w-1}\}, v \in V(G_{\frac{t}{2}-1})$ in which case the symbol represented by the edge (v_{w-1}, v_w) can be recovered by using the parity check represented by v_w .
 - b) or represents an erased symbol and $v \in \{v_1, \dots, v_w\}$ in which case there is a cycle. Since girth of $G_{\frac{t}{2}-1}$ is $\geq t+1$, there must at least $t+1$ distinct erased symbols.

Hence either an erased symbol can be recovered or there are more than t erasures. The argument can be repeated for recovering subsequent erased symbols sequentially one by one if $|E| < t$.

- 3) If $E \cap D_1 \neq \emptyset$, then start with a vertex $v_1 \in U_{\frac{t}{2}-1}$ to which a degree-1 variable node corresponding to an erased code symbol is attached. Now construct a maximal walk, $P = v_1 \dots v_w, v_i \in V(G_{\frac{t}{2}-1}), \forall i \in [w]$, on the erased symbols as before. Now either as before, one of the symbols in E can be recovered or there are more than t erased symbols in E due to a cycle or the walk ends in another node in $U_{\frac{t}{2}-1}$ i.e., $v_w \in U_{\frac{t}{2}-1}$ with the code symbol corresponding to degree-1 variable node attached to v_w also being erased. Since $G_{\frac{t}{2}-1}$ is tree-like, any two nodes in $U_{\frac{t}{2}-1}$ are separated by a path of length at least $t-1$ in $G_{\frac{t}{2}-1}$. Hence there are at least $t-1+2 = t+1$ erasures (+2 because $|E \cap D_1| \geq 2$).
- 4) Thus the entire point of constructing $G_{\frac{t}{2}-1}$ in a tree-like fashion is for protection from more than one erasure in the code symbols corresponding to degree-1 variable nodes attached to $U_{\frac{t}{2}-1}$.




Fig. 9: Figure showing two descriptions of our code at a node T_i for $1 \leq i \leq \frac{t}{2} - 2$




Fig. 10: Figure showing two descriptions of our code at a node in $U_{\frac{t}{2}-1}$

B. Construction of codes achieving the upper bound on rate for t odd:

In this subsection we give a construction of codes achieving the rate bound (31) for any $r \geq 3$ and $t = 2s - 1$, $s \geq 1$. A construction achieving the bound (31) for $t = 5$ can be found in [12]. For $t = 3$, the bound (31) can be achieved by taking the product of two $[r + 1, r]$ single parity check code (2 dimensional product code). For $t = 1$, the bound (31) can be achieved by taking the $[r + 1, r]$ single parity check code. Hence we give a rate optimal construction for any $r \geq 3$ and $t = 2s - 1$, $s \geq 2$ in the following.

We provide an iterative, graph-based construction of rate-optimal codes for a given $r \geq 3$ and $t = 2s - 1$, $s \geq 2$. We build a new graph G_{s-1} iteratively, starting from G_0 (different from the one considered for t even case.), by adding nodes in vertical, layer-by-layer fashion. At every stage of the iteration, the graph G_i (again different from the one constructed for t even case.) thus constructed, will always have girth $\geq t + 1$. We then define our rate-optimal code based on the graph G_{s-1} .

Construction of graph G_{s-1} : Denote the vertex set of a graph G by $V(G)$. We make use of graphs G_0 and B_i , for $1 \leq i \leq s - 1$ described below as the ingredients:

- 1) We pick G_0 as follows: Let G_0 be an r -regular bipartite graph with girth $\geq t + 1$. Let $U_{0,1}, U_{0,2}$ be the two (upper and lower) sets of nodes in the bipartite graph with $\text{degree}(x) = r, \forall x \in U_{0,1}$ and $\text{degree}(y) = r, \forall y \in U_{0,2}$. Hence $V(G_0) = U_{0,1} \cup U_{0,2}$. Define $|U_{0,1}| = |U_{0,2}| = u_0$.
- 2) Next, for i in the range $1 \leq i \leq s - 1$, we iteratively construct an $(r, 2u_{i-1})$ -biregular bipartite graph B_i with girth $g_i \geq \left\lceil \frac{t+1}{i+\frac{1}{2}} \right\rceil$ with the following properties. Let U_i, L_i be the two (upper and lower) sets of nodes in the bipartite graph with $\text{degree}(x) = r, \forall x \in U_i$ and $\text{degree}(y) = 2u_{i-1}, \forall y \in L_i$. Hence $V(B_i) = U_i \cup L_i$. We have that $U_i = U_{i,1} \cup U_{i,2}$ with $|U_{i,1}| = |U_{i,2}|$ and $U_{i,1} \cap U_{i,2} = \emptyset$. Let $|U_{i,1}| = |U_{i,2}| = u_i$ and $|L_i| = l_i$. Therefore $|U_i| = 2u_i$. By edge counting, we have $2ru_i = 2u_{i-1}l_i$ i.e. $ru_i = u_{i-1}l_i$. Additionally we have that, $\forall x \in L_i, |N(x) \cap U_{i,1}| = |N(x) \cap U_{i,2}| = u_{i-1}$ where $N(x)$ denotes the neighbors of node x in B_i . Construction of such graphs will be described separately.

Construction of B_i :

- Consider a (r, u_{i-1}) -biregular bipartite graph $G^1 = (X_1 \cup X_2, E)$ with girth at least $\left\lceil \frac{t+1}{i+\frac{1}{2}} \right\rceil$. Let $X_1 = \{b_1, \dots, b_{\hat{m}}\}$, $X_2 = \{c_1, \dots, c_{\lambda}\}$ for some \hat{m}, λ with $\text{degree}(b_j) = r$, for $1 \leq j \leq \hat{m}$ and $\text{degree}(c_j) = u_{i-1}$, for $1 \leq j \leq \lambda$. Such a graph G^1 can be constructed due to [20].
- Consider a λ -regular bipartite graph $G^2 = (X \cup Y, E)$ with girth at least $\frac{1}{2} \left\lceil \frac{t+1}{i+\frac{1}{2}} \right\rceil$. Let $X = \{x_1, \dots, x_{l'}\}$ and $Y = \{y_1, \dots, y_{l'}\}$ for some l' with each node of degree λ . Such a graph G^2 can be constructed due to [20].
- Take $2l'$ disjoint copies of G^1 . Let G^3 be the disjoint union of the $2l'$ copies named $G^1_1, \dots, G^1_{2l'}$ of G^1 . Denote the nodes b_j, c_j in the l^{th} copy G^1_l by b^l_j, c^l_j respectively.
- Replace the node x_f in G^2 by the nodes $x_{1,f}, \dots, x_{\lambda,f}$ for $1 \leq f \leq l'$. If the neighbors of x_f in G^2 are the nodes in the set $\{y_{q_1}, \dots, y_{q_\lambda}\}$ then connect $x_{j,f}$ to y_{q_j} for $1 \leq j \leq \lambda$. Now replace the node y_f by the nodes $y_{1,f}, \dots, y_{\lambda,f}$ for $1 \leq f \leq l'$. If the neighbors of y_f after replacing the nodes in X were the nodes in the set $\{x_{\sigma_1, \beta_1}, \dots, x_{\sigma_\lambda, \beta_\lambda}\}$ then connect $y_{j,f}$ to x_{σ_j, β_j} for $1 \leq j \leq \lambda$. Denote this graph by G^4 . For every edge $(x_{j,f}, y_{\hat{j}, \hat{f}})$ in G^4 merge the nodes c^f_j and $c^{\hat{f}+f}_{\hat{j}}$ in G^3 and label the resulting merged node as $v^i_{(f-1)\lambda+j}$ in G^3 . Relabel b^l_j as $c^{(i,1)}_{(l-1)\hat{m}+j}$, $\forall 1 \leq l \leq l', \forall 1 \leq j \leq \hat{m}$




Fig. 11: Figure showing two descriptions of our code at a node in T_0

in G^3 . Relabel b_j^l as $c_{(l-l'-1)\hat{m}+j}^{(i,2)}$, $\forall l'+1 \leq l \leq 2l', \forall 1 \leq j \leq \hat{m}$ in G^3 . After the merging and relabeling of nodes in G^3 as described, resulting graph is named as B_i .

- The constructed graph B_i is a $(r, 2u_{i-1})$ -biregular bipartite graph $B_i = (U_i \cup L_i, E_i)$ with $U_i = U_{i,1} \cup U_{i,2}$ where $U_{i,1} = \{c_1^{(i,1)}, \dots, c_{u_i}^{(i,1)}\}$, $U_{i,2} = \{c_1^{(i,2)}, \dots, c_{u_i}^{(i,2)}\}$ and $u_i = l'\hat{m}$ and $L_i = \{v_1^i, \dots, v_{l_i}^i\}$ where $l_i = l'\lambda$ and $\text{degree}(c_j^{(i,p)}) = r$ for $1 \leq j \leq u_i, 1 \leq p \leq 2$ and $\text{degree}(v_j^i) = 2u_{i-1}$ for $1 \leq j \leq l_i$. It can be seen that B_i has girth at least $\left\lceil \frac{t+1}{i+\frac{1}{2}} \right\rceil$. It can also be seen from the construction of B_i that $|N(v_j^i) \cap U_{i,1}| = |N(v_j^i) \cap U_{i,2}| = u_{i-1}$, for $1 \leq j \leq l_i$.

Construction of G_{s-1} : The flowchart shown in Fig. 12 describes the iterative construction of the graph G_{s-1} .

We make the following observations (similar observations compared to t even case):

- 1) For $0 \leq i \leq s-1$, there are $f_i = \prod_{j=i+1}^{s-1} l_j$ disjoint copies of G_i that reside inside G_{s-1} as subgraphs due to the replication steps in the construction of G_{s-1} (given in the flowchart Fig 12). For all $s-1 > j > i$, since copies of G_i reside inside G_j as subgraphs, when we replicate G_j in the $(j+1)^{th}$ replication step, all the copies of G_i contained within G_j also get replicated; we refer to all such copies of G_i here. Let us denote the disjoint union of these subgraphs corresponding to all copies of G_i formed due to replication steps by $G_i^{rep} = \cup_{j=1}^{f_i} (G_i)_j$, where $(G_i)_j$ is the j^{th} copy of G_i in G_{s-1} formed due to some replication step. We view G_i^{rep} as a subgraph of G_{s-1} .
- 2) Note that the construction of G_{s-1} proceeds by adding nodes layer by layer with each layer connecting to the layer below it in a tree-like fashion while maintaining the girth of overall graph to be atleast $t+1$. Let $N(v)$ represent neighbors of a node v in G_{s-1} . Let $(U_i)_j$ be the copy of U_i in $V((G_i)_j), \forall 1 \leq j \leq f_i$. For $1 \leq i \leq s-1$, the nodes $T_i = \cup_{j=1}^{f_i} (U_i)_j$ represent the nodes in the i^{th} layer and they connect in a tree-like fashion to the $(i-1)^{th}$ layer nodes $T_{i-1} = \cup_{j=1}^{f_{i-1}} (U_{i-1})_j$. The connection is tree-like because for $v \in T_i$, let $S_v = N(v) \cap T_{i-1}$ then $S_v \cap S_w = \emptyset$ and $|S_v| = |S_w| = r, \forall v, w \in T_i, v \neq w$ and $N(v) \cap T_p = \emptyset, \forall p \notin \{i-1, i+1\}, \forall v \in T_i$.
- 3) For $1 \leq p \leq 2$, let $(U_{i,p})_j$ be the copy of $U_{i,p}$ in $V((G_i)_j), \forall 1 \leq j \leq f_i$. In fact it can be seen that for $1 \leq i \leq s-1$, the nodes in $T_i^p = \cup_{j=1}^{f_i} (U_{i,p})_j$ connect in a tree-like fashion to nodes in $T_{i-1}^p = \cup_{j=1}^{f_{i-1}} (U_{i-1,p})_j$ for $p \in \{1, 2\}$. The connection is tree-like because for $v \in T_i^p$, let $S_v^p = N(v) \cap T_{i-1}^p$ then $S_v^p \cap S_w^p = \emptyset$ and $|S_v^p| = |S_w^p| = r, \forall v, w \in T_i^p, v \neq w$ and $N(v) \cap T_q^p = \emptyset, \forall q \notin \{i-1, i+1\}, \forall v \in T_i^p$. This implies a tree-like graph is built starting from bottom layer T_0^p (leaf-like nodes) towards T_{s-1}^p . It can be seen that the tree-like graph built from T_0^1 does not intersect or disjoint with tree-like graph built from T_0^2 . The only interaction between these 2 tree-like graphs is by edges representing information symbols connecting nodes in T_0^1 with T_0^2 .
- 4) The fact that G_i has girth at least $t+1$ can be proved using arguments similar to those used in t even case. Hence we skip the proof.

Description of a rate-optimal code \mathcal{C} based on G_{s-1} :


Code \mathcal{C} is defined on the graph G_{s-1} . We will first describe a code \mathcal{C}_1 based on G_{s-1} . On puncturing the code \mathcal{C}_1 by dropping few parity symbols, we get our desired rate-optimal code \mathcal{C} .

Code \mathcal{C}_1 is defined on the graph G_{s-1} as follows:

- 1) The edges of G_0^{rep} represent information symbols. It can be seen that the number of information symbols is $k = u_0 r \prod_{j=1}^{s-1} l_j$.
- 2) Every node of G_{s-1} represents a distinct parity symbol described as follows.
- 3) A node $v \in V(G_0^{rep})$ represents a parity symbol which is the binary sum of information symbols that are represented by edges in G_0^{rep} incident on v .
- 4) For $1 \leq i \leq s-1$, a node $v \in T_i$ in G_{s-1} represents a parity symbol which is the binary sum of code symbols that are represented by the nodes in $N(v) \cap T_{i-1}$.
- 5) It can be seen that the total number of nodes is $n' - k = \sum_{i=0}^{s-1} 2u_i \prod_{j=i+1}^{s-1} l_j = 2 \sum_{i=0}^{s-1} \frac{u_0}{r^i} \prod_{j=1}^{s-1} l_j$.
- 6) \mathcal{C}_1 is an $[n', k]$ code defined by the information symbols represented by edges in G_0^{rep} and parity symbols represented by nodes in G_{s-1} .

Code \mathcal{C} is defined as follows:

- 1) Puncture the code \mathcal{C}_1 by dropping the parity symbols represented by nodes in $U_{s-1,2}$. Hence \mathcal{C} is an $[n, k]$ code defined by the information symbols represented by edges in G_0^{rep} and parity symbols represented by nodes in $V(G_{s-1}) - U_{s-1,2}$.

Fig. 12: Flowchart showing the iterative construction of G_{s-1} .

- 2) Hence $n - k = n' - k - u_{s-1} = 2 \sum_{i=0}^{s-2} u_i \prod_{j=i+1}^{s-1} l_j + u_{s-1} = 2 \sum_{i=0}^{s-2} \frac{u_0}{r^i} \prod_{j=1}^{s-1} l_j + \frac{u_0}{r^{s-1}} \prod_{j=1}^{s-1} l_j$.
- 3) From the above counts, the rate of the code \mathcal{C} can be seen to be equal to the bound given by (31).

The proof that \mathcal{C} can correct t erasures sequentially is similar to the t even case. Hence we skip the proof but we give the main idea as follows. The only point to be noted for sequential recovery of t erasures considering the cases we have dealt with in t even case is that nodes in the top most layer which are part of the code \mathcal{C} i.e., nodes in $T_{s-1}^1 \cup T_{s-2}^2$ are separated from each other with just enough distance that a path of erased symbols connecting any two nodes (representing erased symbols) at this top-most layer (similar to t even case) must have at least $t + 1$ erased symbols. Since the path between 2 nodes in T_{s-2}^2 is potentially smaller than the the path between two nodes in T_{s-1}^1 , we need a little extra distance separating two nodes in T_{s-2}^2 which is given by the bipartite nature of the graph at bottom most layer since the path from a node in T_{s-2}^2 can reach another node in T_{s-2}^2 only through two nodes T_0^2 we get 1 extra distance as any two nodes in T_0^2 is separated by a distance of atleast 2 due to the bipartite nature of G_0^{rep} . This is precisely the reason why we built tree-like graph from T_0^1 disjointly from T_0^2 , so that when we puncture which drops the distance between nodes in top most layer, we get a little extra distance at the bottom layer.

Remark 4. The following conjecture on the rate of an $(n, k, r, t)_{seq}$ code appeared in [13].

$$\frac{k}{n} \leq \frac{1}{1 + \sum_{i=1}^m \frac{a_i}{r^i}},$$

$$a_i \geq 0, a_i \in \mathbb{Z}, \sum_{i=1}^m a_i = t,$$

$$m = \lceil \log_r(k) \rceil.$$

Our rate bound verifies the conjecture as our rate bound can be written in the form:

For t even:

$$\frac{k}{n} \leq \frac{1}{1 + \sum_{i=1}^{\frac{t}{2}} \frac{2}{r^i}}.$$

For t Odd:

$$\frac{k}{n} \leq \frac{1}{1 + \sum_{i=1}^{s-1} \frac{2}{r^i} + \frac{1}{r^s}}.$$

More specific conjecture was given for $t = 5, 6$. While our bound proves the conjecture for $t = 5$, the conjectured upper bound for $t = 6$ does not hold as our upper bound on code rate is both larger and achievable).

For $t = 6$, Our bound takes the form:

$$\frac{k}{n} \leq \frac{r^3}{r^3 + 2r^2 + 2r + 2}$$

$$\frac{k}{n} \leq \frac{1}{1 + \frac{2}{r} + \frac{2}{r^2} + \frac{2}{r^3}}$$

The conjecture for $t = 6$ given in [13]:

$$\frac{k}{n} \leq \frac{1}{1 + \frac{2}{r} + \frac{3}{r^2} + \frac{1}{r^3}}$$

Hence the conjecture given in [13] has a smaller rate than our bound for $t = 6$. And further our upper bound on rate for $t = 6$ can be achieved.

Remark 5. In [9] the authors provide a construction of a code with sequential recovery for any r and t having rate $\frac{r-1}{r+1} + \frac{1}{n}$. $\frac{r-1}{r+1} + \frac{1}{n}$ will meet our rate bound exactly when a Moore graph of degree $r + 1$ and girth $t + 1$ exists. Moore graphs (degree = $r + 1$, girth = $t + 1$) are shown to not exist for any $t \notin \{2, 3, 4, 5, 7, 11\}$ for any $r \geq 2$ (see [21]). Hence the construction in [9] is not rate-optimal for most of the cases.

APPENDIX A PROOF OF CLAIM 1

It is enough to show that:

- $\{A_i\}$ are matrices with each column having Hamming weight 1.

- $\{D_i\}$ are matrices with each row having Hamming weight 1.

As the point 1 written above combined with the fact that each column of $\left[\frac{A_i}{D_i}\right]$ has Hamming weight 2 implies $D_i, i \geq 1$ are matrices with each column having Hamming weight 1 and D_0 by definition is a matrix with each column having Hamming weight 1.

Let us denote the j^{th} column of matrix H_1 by $\underline{h}_j, 1 \leq j \leq n$. Let us show the claim by induction as follows:

(Note that in the arguments that follow $\underline{h}_{p_1}, \dots, \underline{h}_{p_\psi}, \underline{h}_{y_1}, \dots, \underline{h}_{y_\phi}, \underline{h}_{z_1}, \dots, \underline{h}_{z_\theta}$ and $\underline{h}_{w_1}, \dots, \underline{h}_{w_\gamma}$ will be repeatedly used each time denoting some set of column vectors not necessarily related to a different instance of the same notation.)

Induction Hypothesis:

- Property P_i : any $m \times 1$ vector having Hamming weight at most 2 with support contained in $\{\sum_{l=0}^{i-1} \rho_l + 1, \dots, \sum_{l=0}^{i-1} \rho_l + \rho_i\}$ can be written as some linear combination of at most $2(i+1)$ column vectors of H_1 say $\underline{h}_{p_1}, \dots, \underline{h}_{p_\psi}$ for some $\{p_1, \dots, p_\psi\} \subseteq \{1, \dots, \sum_{l=0}^i a_l\}$ and $0 < \psi \leq 2(i+1)$.
- Let us assume as induction hypothesis that the property P_i is true and the Claim 1 is true for $A_1, \dots, A_i, D_0, \dots, D_i$.

Initial step:

- We show that each row of D_0 has Hamming weight exactly 1.
Suppose there exists a row of D_0 with Hamming weight more than 1; let the support set of the row be $\{i_1, i_2, \dots\}$. Then the columns $\underline{h}_{i_1}, \underline{h}_{i_2}$ of H_1 can be linearly combined to give a zero column. This contradicts the fact that $d_{\min}(C) \geq t+1, t > 0$ and t is even. Hence, all rows of D_0 have Hamming weight exactly 1.
- If $t = 2$, then the claim is already proved. So let $t \geq 4$.
- We show that each column of A_1 has Hamming weight exactly 1.
Suppose j^{th} column of A_1 for some $1 \leq j \leq a_1$ has Hamming weight 2; let the support of the column be $\{j_1, j_2\}$ in A_1 . Then the column \underline{h}_{a_0+j} in H_1 along with the 2 column vectors of H_1 say $\underline{h}_{p_1}, \underline{h}_{p_2}$ where $\{p_1, p_2\} \subseteq \{1, \dots, a_0\}$ where \underline{h}_{p_1} has exactly one non-zero entry in j_1^{th} co-ordinate and \underline{h}_{p_2} has exactly one non-zero entry j_2^{th} co-ordinate, can be linearly combined to give a zero column again leading to a contradiction on minimum distance. Such columns with one column having only one non-zero entry exactly in j_1^{th} co-ordinate and another column having only one non-zero entry exactly in j_2^{th} co-ordinate with column labels in $\{1, \dots, a_0\}$ exist due to the single weight columns in the matrix D_0 .
- The above argument also shows that any $m \times 1$ vector having Hamming weight at most 2 with support contained in $\{1, \dots, \rho_0\}$ can be written as some linear combination of at most 2 column vectors of H_1 say $\underline{h}_{p_1}, \dots, \underline{h}_{p_\psi}$ for some $\{p_1, \dots, p_\psi\} \subseteq \{1, \dots, a_0\}$ ($\psi = 1$ or 2). Hence Property P_0 is true.
- We now show that each row of D_1 has Hamming weight exactly 1. Suppose j^{th} row of D_1 has Hamming weight more than 1; let the support set of the row be $\{l_1, l_2, \dots, l_z\}$ in D_1 . Now there is some linear combination of columns $\underline{h}_{a_0+l_1}$ and $\underline{h}_{a_0+l_2}$ in H_1 that gives a zero in $(\rho_0 + j)^{\text{th}}$ co-ordinate and thus this linear combination has support contained in $\{1, \dots, \rho_0\}$ with Hamming weight at most 2. Now applying Property P_0 on this linear combination implies that there is a non-empty set of at most 4 linearly dependent columns in H_1 leading to a contradiction on minimum distance.
- Now we show that Property P_1 is true: We have to prove that any $m \times 1$ vector with Hamming weight at most 2 with support contained in $\{\rho_0 + 1, \dots, \rho_0 + \rho_1\}$ can be written as linear combination of at most $2(1+1) = 4$ column vectors of H_1 say $\underline{h}_{p_1}, \dots, \underline{h}_{p_\psi}$ for some $\{p_1, \dots, p_\psi\} \subseteq \{1, \dots, \sum_{l=0}^1 a_l\}$ and $0 < \psi \leq 4$. This can be easily seen using arguments similar to ones presented before. Let an $m \times 1$ vector have non-zero entries exactly in co-ordinates $\rho_0 + j_1, \rho_0 + j_2$ or $\rho_0 + j_1$. Then this vector can be linearly combined with at most 2 column vectors in H_1 say $\underline{h}_{y_1}, \dots, \underline{h}_{y_\phi}$ where $\{y_1, \dots, y_\phi\} \subseteq \{a_0 + 1, \dots, a_0 + a_1\}$ ($\phi = 1$ or 2) (2 columns $\underline{h}_{y_1}, \underline{h}_{y_2}$ with first and second column having a non-zero entry in co-ordinates $\rho_0 + j_1, \rho_0 + j_2$ respectively or a column \underline{h}_{y_1} with a non-zero entry in $(\rho_0 + j_1)^{\text{th}}$ co-ordinate. These columns $\underline{h}_{y_1}, \underline{h}_{y_2}$ or \underline{h}_{y_1} exist due to D_1 .) to form a $m \times 1$ vector with Hamming weight at most 2 with support contained in $\{1, \dots, \rho_0\}$ which in turn can be written as linear combination of at most 2 column vectors in H_1 say $\underline{h}_{z_1}, \dots, \underline{h}_{z_\theta}$ for some $\{z_1, \dots, z_\theta\} \subseteq \{1, \dots, a_0\}$ ($\theta = 1$ or 2) by property P_0 . Hence the given $m \times 1$ vector is written as linear combination of at most $2(1+1) = 4$ column vectors in H_1 say $\underline{h}_{w_1}, \dots, \underline{h}_{w_\gamma}$ for some $\{w_1, \dots, w_\gamma\} \subseteq \{1, \dots, \sum_{l=0}^1 a_l\}$ and $0 < \gamma \leq 4$.

Induction step :

- Let us assume by induction hypothesis that Property P_i is true and the Claim 1 is true for $A_1, \dots, A_i, D_0, \dots, D_i$ for some $i \leq \frac{t}{2} - 2$ and prove the induction hypothesis for $i+1$. For $t = 4$, the initial step of induction completes the proof of Claim 1. Hence assume $t > 4$.
- Now we show that each column of A_{i+1} has Hamming weight exactly 1: suppose j^{th} column of A_{i+1} for some $1 \leq j \leq a_{i+1}$ has Hamming weight 2; let the support of the column be j_1, j_2 in A_{i+1} . It is clear that the corresponding column vector $\underline{h}_{\sum_{l=0}^i a_l + j}$ in H_1 is a vector with support contained in $\{\sum_{l=0}^{i-1} \rho_l + 1, \dots, \sum_{l=0}^{i-1} \rho_l + \rho_i\}$ and Hamming weight 2.

Now applying Property P_i on this column vector $\underline{h}_{\sum_{l=0}^i a_l + j}$ implies that there is a non-empty set of at most $2(i+1) + 1$

columns in H_1 which are linearly dependent; hence contradicts the minimum distance as $2(i+1) + 1 \leq t - 1$. Hence each column of A_{i+1} has Hamming weight exactly 1.

- Now we show that each row of D_{i+1} has Hamming weight exactly 1: suppose j^{th} row of D_{i+1} has Hamming weight more than 1; let the support set of the row be $\{l_1, \dots, l_z\}$ in D_{i+1} . Now some linear combination of columns $\underline{h}_{\sum_{j=0}^i a_j + l_1}$ and $\underline{h}_{\sum_{j=0}^i a_j + l_2}$ in H_1 will make the resulting vector have a 0 in $(\sum_{l=0}^i \rho_l + j)^{th}$ co-ordinate and the resulting vector also has Hamming weight at most 2 with support contained in $\{\sum_{l=0}^{i-1} \rho_l + 1, \dots, \sum_{l=0}^{i-1} \rho_l + \rho_i\}$ and hence applying Property P_i on this resulting vector implies that there is a non-empty set of at most $2(i+1) + 2$ columns in H_1 which are linearly dependent; hence contradicts the minimum distance as $2(i+1) + 2 \leq t$; thus proving that each row of D_{i+1} has Hamming weight exactly 1.
- Now we show that Property P_{i+1} is true: We have to prove that any $m \times 1$ vector with Hamming weight at most 2 with support contained in $\{\sum_{l=0}^i \rho_l + 1, \dots, \sum_{l=0}^i \rho_l + \rho_{i+1}\}$ can be written as linear combination of at most $2(i+2)$ column vectors of H_1 say $\underline{h}_{p_1}, \dots, \underline{h}_{p_\psi}$ for some $\{p_1, \dots, p_\psi\} \subseteq \{1, \dots, \sum_{l=0}^{i+1} a_l\}$ and $0 < \psi \leq 2(i+2)$. This can be easily seen using arguments similar to ones presented before. Let an $m \times 1$ vector have non-zero entries in co-ordinates $\sum_{l=0}^i \rho_l + j_1, \sum_{l=0}^i \rho_l + j_2$ or $\sum_{l=0}^i \rho_l + j_1$. Then this vector can be linearly combined with at most 2 column vectors in H_1 say $\underline{h}_{y_1}, \dots, \underline{h}_{y_\phi}$ where $\{y_1, \dots, y_\phi\} \subseteq \{\sum_{l=0}^i a_l + 1, \dots, \sum_{l=0}^i a_l + a_{i+1}\}$ with $\phi = 1$ or 2, (2 columns $\underline{h}_{y_1}, \underline{h}_{y_2}$ with first column and second column having a non-zero entry in co-ordinates $\sum_{l=0}^i \rho_l + j_1, \sum_{l=0}^i \rho_l + j_2$ respectively or a column \underline{h}_{y_1} with a non-zero entry in $(\sum_{l=0}^i \rho_l + j_1)^{th}$ co-ordinate. These columns $\underline{h}_{y_1}, \underline{h}_{y_2}$ or \underline{h}_{y_1} exist due to D_{i+1} .) to form a $m \times 1$ vector with Hamming weight at most 2 with support contained in $\{\sum_{l=0}^{i-1} \rho_l + 1, \sum_{l=0}^{i-1} \rho_l + \rho_i\}$ which in turn can be written as linear combination of at most $2(i+1)$ column vectors in H_1 say $\underline{h}_{z_1}, \dots, \underline{h}_{z_\theta}$ for some $\{z_1, \dots, z_\theta\} \subseteq \{1, \dots, \sum_{l=0}^i a_l\}$ and $0 < \theta \leq 2(i+1)$ by property P_i . Hence the given $m \times 1$ vector is written as linear combination of at most $2(i+2)$ column vectors in H_1 say $\underline{h}_{w_1}, \dots, \underline{h}_{w_\gamma}$ for some $\{w_1, \dots, w_\gamma\} \subseteq \{1, \dots, \sum_{l=0}^{i+1} a_l\}$ and $0 < \gamma \leq 2(i+2)$.

REFERENCES

- [1] S. B. Balaji, K. P. Prasanth, and P. V. Kumar, "Binary codes with locality for multiple erasures having short block length," *CoRR*, 2016. [Online]. Available: <http://arxiv.org/abs/1601.07122/>
- [2] P. Gopalan, C. Huang, H. Simitci, and S. Yekhanin, "On the Locality of Codeword Symbols," *IEEE Trans. Inf. Theory*, vol. 58, no. 11, pp. 6925–6934, Nov. 2012.
- [3] D. Papailiopoulos and A. Dimakis, "Locally repairable codes," in *Information Theory Proceedings (ISIT), 2012 IEEE International Symposium on*, July 2012, pp. 2771–2775.
- [4] F. Oggier and A. Datta, "Self-repairing homomorphic codes for distributed storage systems," in *INFOCOM, 2011 Proceedings IEEE*, April 2011, pp. 1215–1223.
- [5] C. Huang, M. Chen, and J. Li, "Pyramid codes: Flexible schemes to trade space for access efficiency in reliable data storage systems," in *Network Computing and Applications, 2007. NCA 2007. Sixth IEEE International Symposium on*, July 2007, pp. 79–86.
- [6] G. Kamath, N. Prakash, V. Lalitha, and P. Kumar, "Codes with local regeneration," in *Information Theory and Applications Workshop (ITA), 2013, Feb 2013*, pp. 1–5.
- [7] I. Tamo and A. Barg, "A family of optimal locally recoverable codes," *IEEE Trans. Inf. Theory*, vol. 60, no. 8, pp. 4661–4676, 2014.
- [8] N. Prakash, V. Lalitha, and P. V. Kumar, "Codes with locality for two erasures," *CoRR*, vol. abs/1401.2422, 2014. [Online]. Available: <http://arxiv.org/abs/1401.2422>
- [9] A. Rawat, A. Mazumdar, and S. Vishwanath, "On cooperative local repair in distributed storage," in *Information Sciences and Systems (CISS), 2014 48th Annual Conference on*, 2014, pp. 1–5.
- [10] W. Song and C. Yuen, "Locally repairable codes with functional repair and multiple erasure tolerance," *CoRR*, vol. abs/1507.02796, 2015. [Online]. Available: <http://arxiv.org/abs/1507.02796>
- [11] —, "Binary locally repairable codes - sequential repair for multiple erasures," *CoRR*, vol. abs/1511.06034, 2015. [Online]. Available: <http://arxiv.org/abs/1511.06034>
- [12] S. Balaji, K. Prasanth, and P. V. Kumar, "Binary codes with locality for four erasures," *arXiv preprint arXiv:1607.02817*, 2016.
- [13] W. Song, K. Cai, and C. Yuen, "On sequential locally repairable codes," *arXiv preprint arXiv:1610.09767*, 2016.
- [14] N. Prakash, G. Kamath, V. Lalitha, and P. Kumar, "Optimal linear codes with a local-error-correction property," in *Proc. IEEE Int. Symp. Inf. Theory (ISIT)*, 2012, pp. 2776–2780.
- [15] W. Song, S. H. Dau, C. Yuen, and T. Li, "Optimal locally repairable linear codes," *Selected Areas in Communications, IEEE Journal on*, vol. 32, no. 5, pp. 1019–1036, May 2014.
- [16] I. Tamo, A. Barg, and A. Frolov, "Bounds on the parameters of locally recoverable codes," *CoRR*, vol. abs/1506.07196, 2015. [Online]. Available: <http://arxiv.org/abs/1506.07196>
- [17] A. Wang and Z. Zhang, "Repair locality from a combinatorial perspective," in *Proc. IEEE Int. Symp. Inf. Theory (ISIT)*, 2014, pp. 1972–1976.
- [18] —, "Repair locality with multiple erasure tolerance," *IEEE Trans. Inf. Theory*, vol. 60, no. 11, pp. 6979–6987, 2014.
- [19] L. Pamies-Juarez, H. Hollmann, and F. Oggier, "Locally repairable codes with multiple repair alternatives," in *Information Theory Proceedings (ISIT), 2013 IEEE International Symposium on*, July 2013, pp. 892–896.
- [20] Z. Furedi, F. Lazebnik, A. Seress, V. A. Ustimenko, and A. J. Woldar, "Graphs of prescribed girth and bi-degree," *Journal of Combinatorial Theory Ser. B* 64 (2), pp. 228–239, 1995.
- [21] G. Exoo and R. Jajcay, "Dynamic cage survey," *Electron. J. Combin, Dynamic Survey*, 2013.