ON A BILINEAR STRICHARTZ ESTIMATE ON IRRATIONAL TORI

CHENJIE FAN[‡]*, GIGLIOLA STAFFILANI[†]*, HONG WANG[§], AND BOBBY WILSON[#]

ABSTRACT. We prove a bilinear Strichartz type estimate for irrational tori via a decoupling type argument, [4], recovering and generalizing the result of [7]. As a corollary, we derive a global well-posedness result for the cubic defocusing NLS on two dimensional irrational tori with data of infinite energy.

1. INTRODUCTION

In [4] Bourgain and Demeter proved the full range of Strichartz estimates for the Schrödingier equation on tori as a consequence of the L^2 decoupling theorem. In this paper we prove in full generality the analog of the improved Strichartz estimate that first appeared in [7] for rational tori.

1.1. Statement of the problem and main results. Let $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ be the one dimensional torus, and let $\alpha_1, ..., \alpha_{d-1} \in [1/2, 1]$, we define *d*-dimensional torus \mathbb{T}^d as $\mathbb{T}^d = \mathbb{T} \times \alpha_1 \mathbb{T} \times \cdots \times \alpha_{d-1} \mathbb{T}$. We say that the torus is irrational if at least on α_i is irrational. The torus is rational otherwise. For any $\lambda \geq 1$, we define \mathbb{T}^d_{λ} as a rescaling of \mathbb{T}^d by λ , i.e. $\mathbb{T}^d_{\lambda} = \lambda \mathbb{T}^d = (\lambda \mathbb{T}) \times (\alpha_1 \lambda \mathbb{T}) \dots \times (\alpha_{d-1} \lambda \mathbb{T})$. When $\lambda \to \infty$, one should think \mathbb{T}_{λ} as a large torus approximating \mathbb{R}^d . We consider the following linear

Schrödinger equation on \mathbb{T}_{λ} , We consider the following Cauchy problem for the linear Schrödinger equation on \mathbb{T}_{λ}^{d} ,

$$\begin{cases} iu_t - \Delta u = 0, \ (t, x) \in \mathbb{R} \times \mathbb{T}^d_{\lambda}; \\ u(0, x) = u_0, \ u_0 \in L^2(\mathbb{T}^d_{\lambda}). \end{cases}$$
(1.1)

Let $U_{\lambda}(t)u_0$ be the solution to (1.1), and let $\Lambda_{\lambda} := \frac{1}{\lambda} (\mathbb{Z} \times \frac{1}{\alpha_1} \mathbb{Z} \times \cdots \times \frac{1}{\alpha_{d-1}} \mathbb{Z})$. One has

$$U_{\lambda}(t)u_{0}(x) = \frac{1}{\lambda^{d/2}} \sum_{k \in \Lambda_{\lambda}} e^{2\pi k i x - |2\pi k|^{2} i t} \widehat{u}_{0}(k).$$
(1.2)

Our main theorem is the following bi-linear refined Strichartz estimate.

Theorem 1.1. Let $\phi_1, \phi_2 \in L^2(\mathbb{T}_{\lambda})$ be two initial data such that $supp \hat{\phi}_i \subset \{k : |k| \sim N_i\}, i = 1, 2$, for some large $N_1 \geq N_2$, and let $\eta(t)$ be a time cut-off function, $supp \eta \in [0,1]$. Then when d = 2,

$$\|\eta(t)U_{\lambda}\phi_{1}\cdot\eta(t)U_{\lambda}\phi_{2}\|_{L^{2}_{x,t}} \lesssim N_{2}^{\epsilon} \left(\frac{1}{\lambda} + \frac{N_{2}}{N_{1}}\right)^{1/2} \|\phi_{1}\|_{L^{2}} \|\phi_{2}\|_{L^{2}},$$
(1.3)

when $d \geq 3$

$$\|\eta(t)U_{\lambda}\phi_{1}\cdot\eta(t)U_{\lambda}\phi_{2}\|_{L^{2}_{x,t}} \lesssim N_{2}^{\epsilon} \left(\frac{N_{2}^{d-3}}{\lambda} + \frac{N_{2}^{d-1}}{N_{1}}\right)^{1/2} \|\phi_{1}\|_{L^{2}} \|\phi_{2}\|_{L^{2}}.$$
(1.4)

We note that when $d = 2, N_1 = N_2, \lambda = 1$, estimate (1.3) recovers the Strichartz inequality for the (irrational) torus after an application of Hölder's inequality, up to an N_2^{ϵ} -loss. When $\lambda \to \infty$, estimate (1.3), (1.4) consistent with the billinear Strichartz inequality in \mathbb{R}^{d+1} , [2]. Up to the N_2^{ϵ} -loss, inequality (1.3) is sharp.

Furthermore, when $\lambda \geq N_1$, the estimates fall into the so-called semiclassical regime in which the geometry of \mathbb{T}_{λ} is irrelevant. We refer to the work of Hani, [11], for same estimate (without N_2^{ϵ} loss) on general compact manifolds.

^{*} Chenjie Fan and Gigliola Staffilani are partially supported by NSF DMS 1362509 and DMS 1462401.

[‡]Department of Mathematics, University of Chicago, 5734 S University Ave, Chicago, IL 60637, USA. email: cjfan@math.uchicago.edu.

[†] Department of Mathematics, Massachusetts Institute of Technology, 77 Massachusetts Ave, Cambridge, MA 02139-4307 USA. email: gigliola@math.mit.edu.

[§]Department of Mathematics, Massachusetts Institute of Technology, 77 Massachusetts Ave, Cambridge, MA 02139-4307 USA. email: hongwang@mit.edu.

[#]Department of Mathematics, Massachusetts Institute of Technology, 77 Massachusetts Ave, Cambridge, MA 02139-4307 USA. email: blwilson@mit.edu.

On the torus, our results improves the estimate in [11] for $\lambda \leq N_1$. Estimate (1.3), (1.4) rely on the geometry of torus and cannot hold on general compact manifolds.

Remark 1.2. It may also be interesting to consider trilinear estimates. In fact when one considers the quintic nonlinear Schrödinger equation as in [12] and [13], trilinear estimates are fundamental. See also [15].

We will derive Theorem 1.1 from some bilinear decoupling type estimates. We first introduce some basic notations.

Let P be the truncated paraboloid in \mathbb{R}^{d+1} ,

$$P = \{ (\xi, |\xi|^2) : \xi \in \mathbb{R}^d, |\xi| \lesssim 1 \}.$$
(1.5)

For any function f supported on P, we define

$$Ef = \widehat{f}d\widehat{\sigma},\tag{1.6}$$

1 /0

where σ is the measure on P.

Note a function supported on P can be naturally understood as a function supported on the ball $B = \{\xi \in \mathbb{R}^d, |\xi| \leq 1\}$.

By a slight abuse of notation, for a function f supported in the ball B in \mathbb{R}^d , we also define

$$Ef(x,t) = \int_{B} e^{-2\pi i(\xi \cdot x + |\xi|^{2}t)} f(\xi) d\xi.$$
 (1.7)

One can see that the two definitions of Ef are essentially the same since P projects onto B.

We decompose P as a finitely overlapping union of caps θ of radius δ . Here a cap θ of radius δ is the set $\theta = \{\xi \in P, |\xi - \xi_0| \leq \delta\}$ for some fixed $\xi_0 \in P$. We define $Ef_{\theta} = \widehat{f_{\theta}d\sigma}$, where f_{θ} is f restricted to θ . We use a similar definition also when f is a function supported on the unit ball in \mathbb{R}^d . We have $Ef = \sum_{\theta} Ef_{\theta}$.

Now, we are ready to state our main decoupling type estimate.

Theorem 1.3. Given $\lambda \geq 1$, $N_1 \geq N_2 \geq 1$. Let f_1 be supported on P where $|\xi| \sim 1$, and let f_2 be supported on P where $|\xi| \sim \frac{N_2}{N_1}$. Let $\Omega = \{(t, x) \in [0, N_1^2] \times [0, (\lambda N_1)^2]^d\}$. For a finitely overlapping covering of the ball $B = \{|\xi| \leq 1\}$ of caps $\{\theta\}$, $|\theta| = \frac{1}{\lambda N_1}$, we have the following estimate. For any small $\epsilon > 0$, when d = 2,

$$\|Ef_1Ef_2\|_{L^2_{avg}(w_{\Omega})} \lesssim_{\epsilon} (N_2)^{\epsilon} \lambda^{d/2} \left(\frac{1}{\lambda} + \frac{N_2^{d-1}}{N_1}\right)^{1/2} \prod_{j=1}^2 \left(\sum_{|\theta| = \frac{1}{\lambda N_1}} \|Ef_{j,\theta}\|_{L^4_{avg}(w_{\Omega})}^2\right)^{1/2},$$
(1.8)

when $d \geq 3$,

$$\|Ef_1Ef_2\|_{L^2_{avg}(w_{\Omega})} \lesssim_{\epsilon} (N_2)^{\epsilon} \lambda^{d/2} \left(\frac{N_2^{d-3}}{\lambda} + \frac{N_2^{d-1}}{N_1}\right)^{1/2} \prod_{j=1}^2 \left(\sum_{|\theta| = \frac{1}{\lambda N_1}} \|Ef_{j,\theta}\|_{L^4_{avg}(w_{\Omega})}^2\right)^{1/2}, \tag{1.9}$$

where w_{Ω} is a weight adapted to Ω .

The presence of weight w in these estimates is standard. We list the basic property of w in Section 1.5, and one can refer to [5] for more details. The notation $L_{avg}(w_{\Omega})^2$ is explained in notation subsection below, subsection 1.4.

The proof of Theorem 1.3 gives another proof of the linear decoupling theorem in [4] in dimension d = 2, and does not rely on multilinear-Kakeya or multilinear restriction theorems in \mathbb{R}^3 . The proof of Theorem 1.3 in dimension $d \geq 3$ relies instead on linear decoupling in \mathbb{R}^{d+1} , [4].

Remark 1.4. The estimate in Theorem 1.1, Theorem 1.3 is sharp up to an N_2^{ϵ} . See Section A for examples.

Remark 1.5. The N_2^{ϵ} loss in Theorem 1.1 is typical if one wants to directly use a decoupling type argument. It may be possible to remove N_2^{ϵ} in the mass supercritical setting, (in our case, this means $d \ge 3$), using the approach in [14], where the scale invariant Strichartz estimates are studied.

Remark 1.6. Similar bilinear estimates for dimension $d \ge 3$ were also considered in [14] for non-rescaled tori, see Lemma 3.3. On the other hand in this work we also consider the d = 2 case which is mass critical.

1.2. Acknowledgment. We thank Larry Guth for very helpful discussions during the course of this work.

$$\begin{cases} iu_t + \Delta u = |u|^2 u, \\ u(0) = u_0 \in H^s(\mathbb{T}^d). \end{cases}$$
(1.10)

Let us focus for a moment on the d = 2 case. The Cauchy problem is said to be locally well-posed in $H^s(\mathbb{T}^d)$ if for any initial data $u_0 \in H^s(\mathbb{T}^d)$ there exists a time $T = T(||u_0||_s)$ such that a unique solution to the initial value problem exists on the time interval [0, T]. We also require that the data to solution map is continuous from $H^s(\mathbb{T}^d)$ to $C_t^0 H_x^s([0, T] \times \mathbb{T}^d)$. If $T = \infty$, we say that a Cauchy problem is globally well-posed.

The initial value problem (1.10) is locally well-posed for initial data $u_0 \in H^s$, s > 0 via Strichartz estimates. Note that using iteration, by the energy conservation law, i.e.

$$E(u(t)) = E(u_0) = \frac{1}{2} \int |\nabla u|^2 + \frac{1}{4} \int |u|^4,$$

all initial data in $H^1(\mathbb{T}^2)$ give rise to a global solution. Next, by the nowadays standard I-method, [6], by considering a modified version of the energy, in the rational torus case, it was proved in [7] that (1.10) is indeed globally wellposedness for initial data in H^s , s > 2/3. The key estimate there was in fact (1.3) for linear solutions on **rescaled** tori, which we prove here to be available also for irrational tori.

The proof for (1.3) presented in [7] is only for rational tori since it relies on certain types of counting lemmata that cannot directly work on irrational tori. One of the main purpose of this work in fact is to extend results on rational tori to irrational ones.

Based on the discussion we just made, as a corollary of Theorem 1.1, we have

Corollary 1.7. The initial value problem (1.10) defined on any torus \mathbb{T}^2 is globally well-posed for initial data in $H^s(\mathbb{T}^2)$ with s > 2/3.

Remark 1.8. Results such as Corollary 1.7 usually also give a control on the growth of Sobolev norms of the global solutions. We do not address this particular question here. We instead refer the reader to the recent work [8].

The original Strichartz estimates needed to prove the local well-posedness of Cauchy problems such as (1.10) were first obtained in [1] via number theoretical related counting arguments for rational tori. Recently, the striking proof of the L^2 decoupling Theorem, [4], provided a completely different approach from which all the desired Strichartz estimates on tori, both rational and irrational, follow. This approach in particular does not depend on counting lattice points. See also the work [10] and [9]. The method of proof we implement in this present work is mostly inspired by [4] and the techniques used to prove the L^2 decoupling Theorem.

We quickly recall the main result in [4]. Let P be a unit parabola in \mathbb{R}^{d+1} , covered by finitely overlapping caps θ of radius $\frac{1}{R}$. Let f be a function defined on P, then one has for any $\epsilon > 0$ small,

$$\|Ef\|_{L^p(w_{B_{R^2}})} \lesssim_{\epsilon} R^{\epsilon} (R^2)^{d/4 - \frac{d+2}{2p}} \left(\sum_{\theta} \|Ef_{\theta}\|_{L^p(B_{R^2})}^2 \right)^{1/2}, \quad p \ge \frac{2(d+2)}{d}.$$
(1.11)

Note that (1.11) corresponds to Theorem 1.1 in [4], and the dimension n in the estimate (2) there corresponds to our d+1. Also note that the linear decoupling (1.11) not only works for those f exactly supported on P, but those f supported in a R^{-2} neighborhood of P, and in this case, cap θ would be replaced by the R^{-2} neighborhood of the original θ , see Theorem 1.1 in [4].

We remark that one key feature of this decoupling type estimate is that one needs to work on a larger scale in physical space, i.e. the scale R^2 rather than R, in order to observe the decoupling phenomena. The proper observational scale dictated by Heisenberg's uncertainty principle is R.

Indeed, one principle, which is usually called *parallel decoupling*, indicates that if decoupling happens in a small region, then decoupling happens in a large region as well. We state a bilinear version the parallel decoupling below.

Lemma 1.9 ([4], [5]). Let D be a domain, and $D = D_1 \cup D_2 \dots \cup D_J$, $D_i \cap D_j = \emptyset$. If for some constant A > 0 and for function h_1, h_2 , defined on the unit parabola, one has

$$\|Eh_1Eh_2\|_{L^2_{avg}(w_{D_i})} \le A \prod_{j=1}^2 \left(\sum_{|\theta| = \frac{1}{\lambda N_1}} \|Eh_{j,\theta}\|_{L^4_{avg}(w_{D_i})}^2 \right)^{1/2} \quad i = 1, \dots, J,$$
(1.12)

then one also has

$$\|Eh_1Eh_2\|_{L^2_{avg}(w_D)} \le A \prod_{j=1}^2 \left(\sum_{|\theta| = \frac{1}{\lambda N_1}} \|Eh_{j,\theta}\|_{L^4_{avg}(w_D)}^2 \right)^{1/2}.$$
(1.13)

The proof of this particular formulation of parallel decoupling follows by Minkowski's inequality.

As it exists, parallel decoupling is a principle rather than a concrete lemma. We state the version here solely for concreteness. It should be easy to generalize the lemma under different conditions.

1.4. Notation. We write $A \leq B$ if $A \leq CB$, for a constant C > 0, $A \sim B$ if both $A \leq B$ and $B \leq A$. We say $A \leq_{\epsilon} B$ if the constant C depends on ϵ . Similarly for $A \sim_{\epsilon} B$. For a Borel set, $E \subset \mathbb{R}^d$, we denote that diameter of E by |E| and the Lebesgue measure of E by m(E).

We will use the usual function space L^p . We also use a (weighted) average version of L^p space, i.e

$$\|g\|_{L^{p}_{avg}(A)} = \left(\int_{A} |g|^{p}\right)^{1/p} := \left(\frac{1}{m(A)} \int_{A} |g|^{p}\right)^{1/p}$$

and

$$||g||_{L^p_{avg}(w_A)} = \left(\frac{1}{m(A)}\int |g|^p w_A\right)^{1/p},$$

where w_A is a weight function described below.

For any function f, we use \tilde{f} to denote its Fourier transform. When we say *unit ball*, we refer to a ball of radius $r \sim 1$. We will often identify a torus as a bounded domain in Euclidean space, for example, we will view $(\mathbb{R}/\mathbb{Z})^d$ as $[0,1]^d \subset \mathbb{R}^d$. In this work, Ω is used to denote the domain $[0, N_1^2] \times [0, (\lambda N_1)^2]^d \subset \mathbb{R}^{d+1}$.

1.5. The weight w_A . If h is a Schwartz function whose Fourier transform, \hat{h} , is supported in a ball of radius 1/R, we expect h be essentially constant on balls of radius R, and morally

$$\|h\|_{L^p_{avg}(B_R)} \sim \|h\|_{L^2_{avg}(B_R)} \sim \|h\|_{L^\infty(B_R)}.$$
(1.14)

Expression (1.14) is not rigorous, and the introduction of the weight w_{B_R} is a standard way to overcome this technical difficulty. We refer to Lemma 4.1 in [5] for more detailed discussion of the weight function.

For any bounded open convex set A, the weight function w_A , might change from line to line, from the left hand side of the inequality to the right hand side, satisfies the same properties:

- $\int w_A \sim m(A)$.
- $w_A \gtrsim 1$ on A, and rapidly (polynomial type) decay outside A.

We will usually define A to be a ball, or the product of balls in this paper.

Furthermore, let B_R be a ball centered at 0, and let μ_{B_R} be a function such that $\widehat{\mu_{B_R}}$ is about $\frac{1}{m(B_{1/R})}$ on $B_{1/R}$, and supported in $B_{2/R}$, then μ_{B_R} is about 1 on B_R , decays faster than any polynomial outside of B_R . $\mu_{B_R}^2$ is positive, decays faster than any polynomial outside of B_R and fourier supported in $B_{4/R}$, We take translations B'of B_R to cover the whole space, we note $\mu_{B'}$ as the corresponding translation of μ_{B_R} and $w_{B_R}(B') = \max_{x \in B'} w_{B_R}$, we have the following useful property,

$$w_{B_R}(x) \le \sum_{B'} w_{B_R}(B') 1_{B'}(x) \lesssim \sum_{B'} w_{B_R}(B') \mu_{B'}^2(x) \lesssim w_{B_R}(x).$$
(1.15)

The last inequality follows from the fact that $\mu_{B'}^2$ decays faster than any polynomial outside of B'.

Lemma 1.10. For a function f supported in $B_{1/R}$, for any $p < \infty$,

$$|Ef||_{L^{\infty}(B_R)} \lesssim ||Ef||_{L^p_{avg}(\mu_{B_R})}.$$

We refer the proof to Corollary 4.3 in [5] with the weight on the left hand side being 1_{B_R} so that on the right hand side we have a fast decay weight.

Remark 1.11. In general, Lemma 1.10 should hold for any convex set A and the dual convex body A^* .

2. PROOF OF THEOREM 1.1 ASSUMING THEOREM 1.3

Assume Theorem 1.3, let us prove Theorem 1.1. The argument below comes from the proof of discrete restriction and Strichartz estimate on irrational tori assuming the L^2 decoupling estimate, see Theorem 2.2, Theorem 2.3 in [4]. The argument originally comes as observation due to Bourgain [3]. We record it here for completeness.

Let ϕ_1, ϕ_2 be as in Theorem 1.1. We rescale ϕ_1 to be supported in the unit ball and rescale ϕ_2 to be supported in a ball of radius $\sim \frac{N_2}{N_1}$. Recall,

$$U_{\lambda}(t)\phi_j(x,t) = \frac{1}{\lambda^{d/2}} \sum_{k \in \Lambda_{\lambda}, k \sim N_1} e^{2\pi i k \cdot x - |2\pi k|^2 t} \widehat{\phi}_j(k).$$
(2.1)

We perform a change of variables $\xi = \frac{k}{N_1}$ and we let

$$h_j(\tau) = \frac{1}{\lambda^{d/2}} \sum_{\xi \in \Lambda_{\lambda N_1}, |\xi| \sim 1} \widehat{\phi}_j(\xi N_1) \delta_{\xi}(\tau), \qquad j = 1, 2.$$

$$(2.2)$$

Note one can directly check that

$$U_{\lambda}(t)\phi_j(x,t) = Eh_j(-2\pi N_1 x, (2\pi)^2 N_1^2 t).$$
(2.3)

Without loss of generality, we suppress the constants -2π and $(2\pi)^2$.

Let $Q_0 = [0, N_1^2] \times \mathbb{T}_{\lambda N_1}^d$ and let us view $\mathbb{T}_{\lambda N_1}^d$ as a compact set in \mathbb{R}^d . In particular, one can construct the associated weight function w_{Q_0} . Direct computation (via change of variables) gives

$$\|U_{\lambda}(t)\phi_{1})U_{\lambda}(t)\phi_{2}\|_{L^{2}([0,1]\times\mathbb{T}^{d}_{\lambda})} \sim N_{1}^{-\frac{d+2}{2}}m(Q_{0})^{1/2}\|Eh_{1}Eh_{2}\|_{L^{2}_{avg}(Q_{0})}$$
(2.4)

and due to the the periodicity of Eh_i , i = 1, 2, one has

$$||Eh_1Eh_2||_{L^2_{avg}(\Omega)} = ||Eh_1Eh_2||_{L^2_{avg}(Q_0)}.$$
(2.5)

For a covering $\{\theta\}$ of caps of radius $\frac{1}{\lambda N_1}$, each cap θ contains at most one $\xi_{\theta} \in \Lambda_{\lambda N_1}$, corresponding to $k_{\theta} = N_1 \xi_{\theta} \in \Lambda_{\lambda}$, then

$$\|Eh_{j,\theta}\|_{L^4_{avg}(w_{Q_0})} \sim h_j(\xi_{\theta}) \sim \frac{1}{\lambda^d} \widehat{\phi}_j(k_{\theta})$$

and

$$\prod_{j=1}^{2} \left(\sum_{|\theta| = \frac{1}{\lambda N_{1}}} \|Eh_{j}\|_{L^{4}_{avg}(w_{Q_{0}})}^{2} \right)^{1/2} \sim \lambda^{-d} \prod_{j=1}^{2} \left(\frac{1}{\lambda^{d}} \sum_{k \in \Lambda_{\lambda}} |\widehat{\phi}_{j}(k)|^{2} \right)^{1/2} \sim \lambda^{-d} \|\phi_{1}\|_{L^{2}} \|\phi_{2}\|_{L^{2}}.$$

For convenience of notation let

$$D_{\lambda,N_1,N_2} := \begin{cases} \frac{1}{\lambda} + \frac{N_2}{N_1}, \text{ when } d = 2, \\ \frac{N_2^{d-3}}{\lambda} + \frac{N_2^{d-1}}{N_1}, \text{ when } d \ge 3. \end{cases}$$
(2.6)

Recall that $\Omega = [0, N_1]^2 \times [0, (\lambda N_1)^2]^d$, we apply Theorem 1.3 with $f_j = h_j$, and we have

$$\|Eh_{1}Eh_{2}\|_{L^{2}_{avg}(w_{\Omega})} \lesssim_{\epsilon} (N_{2})^{\epsilon} \lambda^{d/2} D^{1/2}_{\lambda,N_{1},N_{2}} \prod_{j=1}^{2} \left(\sum_{|\theta|=\frac{1}{\lambda N_{1}}} \|Eh_{j,\theta}\|^{2}_{L^{4}_{avg}(w_{\Omega})} \right)^{1/2}.$$
(2.7)

Note that Ω can be covered by Q such that $\{Q\}$ are finitely overlapping and each Q is a translation of Q_0 . Since Eh_j are periodic on x, estimate (2.7) is equivalent to

$$\|Eh_{1}Eh_{2}\|_{L^{2}_{avg}(w_{Q_{0}})} \lesssim_{\epsilon} (N_{2})^{\epsilon} \lambda^{d/2} D^{1/2}_{\lambda,N_{1},N_{2}} \prod_{j=1}^{2} \left(\sum_{|\theta|=\frac{1}{\lambda N_{1}}} \|Eh_{j,\theta}\|_{L^{4}(w_{Q_{0}})}^{2} \right)^{1/2}.$$
(2.8)

Plugging (2.8) into (2.4) gives

$$\begin{split} \|U_{\lambda}(t)\phi_{1}U_{\lambda}(t)\phi_{2}\|_{L^{2}([0,1]\times\mathbb{T}^{d}_{\lambda})} \\ \lesssim N_{1}^{-\frac{d+2}{2}} \cdot N_{1}m(\mathbb{T}^{d}_{\lambda N_{1}})^{1/2}\lambda^{-d} \cdot (N_{2})^{\epsilon}\lambda^{d/2}D_{\lambda,N_{1},N_{2}}^{1/2}\|\phi_{1}\|_{L^{2}}\|\phi_{2}\|_{L^{2}} \\ \sim (N_{2})^{\epsilon}D_{\lambda,N_{1},N_{2}}^{1/2}\|\phi_{1}\|_{L^{2}}\|\phi_{2}\|_{L^{2}} \end{split}$$

and Theorem 1.1 follows.

The rest of the paper details the proof of Theorem 1.3.

3. An overview of the proof of Theorem 1.3

First, we reduce the proof of Theorem 1.3 to the following proposition.

Proposition 3.1. Let τ_1 be a cap of radius $\frac{N_2}{N_1}$ supported at ξ and $|\xi| \sim 1$. Let τ_2 be a cap of radius $\frac{N_2}{N_1}$ supported at ξ with $|\xi| \sim \frac{N_2}{N_1}$. Let f_j be a function supported in τ_j , then for any small $\epsilon > 0$, when d = 2

$$\|Ef_1 Ef_2\|_{L^2_{avg}(w_{\Omega})} \lesssim_{\epsilon} (N_2)^{\epsilon} \lambda^{d/2} \left(\frac{1}{\lambda} + \frac{N_2}{N_1}\right)^{1/2} \prod_{j=1}^2 \left(\sum_{|\theta| = \frac{1}{\lambda N_1}, \theta \subset \tau_j} \|Ef_{j,\theta}\|_{L^4_{avg}(w_{\Omega})}^2\right)^{1/2}$$
(3.1)

when $d \geq 3$,

$$\|Ef_1Ef_2\|_{L^2_{avg}(w_{\Omega})} \lesssim_{\epsilon} (N_2)^{\epsilon} \lambda^{d/2} \left(\frac{N_2^{d-3}}{\lambda} + \frac{N_2^{d-1}}{N_1}\right)^{1/2} \prod_{j=1}^2 \left(\sum_{|\theta| = \frac{1}{\lambda N_1}, \theta \subset \tau_j} \|Ef_{j,\theta}\|_{L^4_{avg}(w_{\Omega})}^2\right)^{1/2}.$$
 (3.2)

Now, let f_1, f_2 be as in Proposition 3.1. We define $K_0(\lambda, N_1, N_2)$ to be the best constant such that

$$\|Ef_1Ef_2\|_{L^2_{avg}(w_{\Omega})} \le \lambda^{d/2} K_0(\lambda, N_1, N_2) \prod_{j=1}^2 \left(\sum_{|\theta| = \frac{1}{\lambda N_1}} \|Ef_{j,\theta}\|_{L^4_{avg}(w_{\Omega})}^2 \right)^{1/2}.$$
(3.3)

We also let $\tilde{K}(\lambda, N_1, N_2)$ and $K(\lambda, N_1, N_2)$ be defined as the best constants such that

$$\|Ef_1Ef_2\|_{L^2_{avg}(w_{[0,N_1^2]\times[0,\lambda N_1]^d})} \le \lambda^{d/2}\tilde{K}(\lambda,N_1,N_2)\prod_{j=1}^2 \left(\sum_{|\theta|=\frac{1}{\lambda N_1}} \|Ef_{j,\theta}\|_{L^4_{avg}(w_{[0,N_1^2]\times[0,\lambda N_1]^d})}^2\right)^{1/2}, \quad (3.4)$$

$$\|Ef_1 Ef_2\|_{L^2_{avg}(w_{B_{N_1^2}})} \le \lambda^{d/2} K(\lambda, N_1, N_2) \prod_{j=1}^2 \left(\sum_{|\theta| = \frac{1}{\lambda N_1}} \|Ef_{j,\theta}\|_{L^4_{avg}(w_{B_{N_1^2}})}^2 \right)^{1/2}.$$
(3.5)

Below we will prove that

$$K_0(\lambda, N_1, N_2) \lesssim N_2^{\epsilon} (\frac{1}{\lambda} + \frac{N_2}{N_1})^{1/2}, \qquad d = 2,$$

$$K_0(\lambda, N_1, N_2) \lesssim N_2^{\epsilon} (\frac{N_2^{d-3}}{\lambda} + \frac{N_2^{d-1}}{N_2})^{1/2}, \qquad d \ge 3.$$
(3.6)

We point out here that by parallel decoupling and Lemma 1.9 one always has

$$K_0(\lambda, N_1, N_2) \lesssim K(\lambda, N_1, N_2), \qquad K_0(\lambda, N_1, N_2) \lesssim \tilde{K}(\lambda, N_1, N_2).$$
(3.7)

The proof of Proposition 3.1 or equivalently (3.6) proceeds as follows. We first show **Lemma 3.2.** When $\lambda \geq N_1$,

$$\tilde{K}(\lambda, N_1, N_2) \lesssim N_2^{\epsilon} \frac{N_2^{(d-1)/2}}{N_1^{1/2}}.$$
(3.8)

Note that when $\lambda \geq N_1$, Proposition 3.1 follows from (3.7) and Lemma 3.2. Then, we show Lemma 3.3. When $\lambda \leq N_1$,

$$K(\lambda, N_1, N_2) \lesssim N_2^{\epsilon} (\frac{1}{\lambda} + \frac{N_2}{N_1})^{1/2}, \quad d = 2,$$

$$K(\lambda, N_1, N_2) \lesssim N_2^{\epsilon} (\frac{N_2^{d-3}}{\lambda} + \frac{N_2^{d-1}}{N_1})^{1/2}, \quad d = 3.$$
(3.9)

From (3.7), clearly Proposition 3.1 follows from Lemma 3.2 and Lemma 3.3.

The proof of Lemma 3.3 in dimension d = 2 relies on induction (of scale N_2). The proof of Lemma 3.3 in dimension in $d \ge 3$ is easier and more straightforward, (in some sense, it also relies on induction, but it is enough to induct only once.)

We first show the base case:

Lemma 3.4. When $\lambda \leq N_1$ and $N_2 \lesssim 1$, , $K(\lambda, N_1, N_2) \lesssim \frac{1}{\lambda^{1/2}}$.

Lemma 3.4 is not as useful in dimension $d \ge 3$, we indeed have a better estimate:

Lemma 3.5. When $d \ge 3$, $\lambda \le N_1$ and $\lambda \le \frac{N_1}{N_2^2}$, $K(\lambda, N_1, N_2) \lesssim \left(\frac{N_2^{d-3}}{\lambda}\right)^{1/2}$.

We then show the following lemma, which ensures that we only need to induct until $\lambda \leq \frac{N_1}{N_2}$, when d = 2, and until $\frac{N_1}{N_2}$ when $d \geq 3$.

Lemma 3.6. Let $\lambda \leq N_1$.

Let d = 2. Assume we have that $K(\lambda, N_1, N_2) \leq \lambda^{-1/2}$ when $\lambda < \frac{N_1}{N_2}$. Then

$$K(\lambda, N_1, N_2) \le N_2^{\epsilon} \frac{N_2^{\frac{d-1}{2}}}{N_1^{1/2}} \qquad when \quad \lambda \ge \frac{N_1}{N_2}.$$

Let $d \ge 3$. Assume we have that $K(\lambda, N_1, N_2) \le \left(\frac{N_2^{d-3}}{\lambda}\right)^{1/2}$ when $\lambda < \frac{N_1}{N_2^2}$. Then

$$K(\lambda, N_1, N_2) \le N_2^{\epsilon} \frac{N_2^{\frac{d-1}{2}}}{N_1^{1/2}} \quad when \quad \lambda \ge \frac{N_1}{N_2^2}.$$

Note that when $d \ge 3$, Lemma 3.5 and Lemma 3.6 imply Lemma 3.3. In dimension d = 2, we use induction (we rely on the so-called parabolic rescaling) to finish the proof of Lemma 3.3.

We end this section with an outline of the structure of the rest of the paper. We show that Proposition 3.1 implies Theorem 1.3 in Section 4. Lemma 3.2, Lemma 3.4, Lemma 3.6 all rely on the exploration of the so-called *transversality* which essentially allow us to reduce the dimensionality of the problem. We first explore *transversality* in Section 5 and then we prove Lemma 3.2, Lemma 3.4, Lemma 3.6 in Section 6.

The detail of the induction procedure, (which is non trivial), that is used to prove Lemma 3.3 in dimension d = 2 is given in Section 5. We remark here the proof of Lemma 3.3 relies on Lemma 3.2.

Finally, we prove Lemma 3.5 at the end of Section 7, which, together with Lemma 3.6 will conclude the proof of Lemma 3.3 in dimension $d \ge 3$.

4. Proposition 3.1 implies Theorem 1.3

We first introduce one standard but important tool in the following lemma.

Lemma 4.1. [[4], [5]] Let $\{g_{\alpha}\}$ be a family of functions such that $supp \widehat{g}_{\alpha}$ are finitely overlapped cubes of length ρ . Let A be bounded convex open set tiled by finitely overlapped cubes Q of side length $\geq \rho^{-1}$, then for the w_A adapted to A, the following holds,

$$\int_{A} |\sum g_{\alpha}|^{2} w_{A} \lesssim \sum \frac{1}{m(A)} \int |g_{\alpha}|^{2} w_{A}.$$

Proof. Since we can sum up the weight function over a finitely overlapping cover $\{Q\}$ of A: $w_A = \sum_{Q \subset A} w_Q$, it suffices to prove for A = Q. Recall by the inequality 1.15, we cover the whole space \mathbb{R}^n by translations Q' of Q,

$$\begin{split} \oint_{Q} |\sum g_{\alpha}|^{2} w_{Q} dx &\leq \frac{1}{m(Q)} \sum_{Q'} \int_{Q'} w_{Q}(Q') |\sum g_{\alpha}|^{2} \\ &\leq \frac{1}{m(Q)} \sum_{Q'} w_{Q}(Q') \int |\sum g_{\alpha}|^{2} \mu_{Q'}^{2} \\ &= \frac{1}{m(Q)} \sum_{Q'} w_{Q}(Q') \int |\widehat{g}_{\alpha} * \widehat{\mu_{Q'}}|^{2} \\ &\lesssim \frac{1}{m(Q)} \sum_{Q'} w_{Q}(Q') \sum_{\alpha} \int |g_{\alpha}|^{2} \mu_{Q'}^{2} \\ &\lesssim \frac{1}{m(Q)} \sum_{\alpha} \int |g_{\alpha}|^{2} w_{Q} \end{split}$$

Now we can reduce Theorem 3.1 to a bilinear decoupling on two $\frac{N_2}{N_1}$ -diameter caps.

Lemma 4.2. Theorem 1.3 is equivalent to Proposition 3.1.

Proof. Let f_1, f_2 be as in Theorem 1.3. Then $f_1 = \sum_{|\tau| = \frac{N_2}{N_1}} f_{1,\tau}$ and $f_{1,\tau}$'s are supported on finitely overlapping caps of diameter $\frac{N_2}{N_1}$.

Since $|f_2|$ is supported in a cap of diameter $\frac{N_2}{N_1}$, the supports of $\{\widehat{Ef}_{1,\tau} * \widehat{Ef}_2\}_{\tau}$ are in finitely overlapping cubes of length $\frac{N_2}{N_1}$. Since the scale of Ω is larger than N_1/N_2 , i.e. it contains a ball of radius $> N_1/N_2$, By Lemma 4.1,

$$\int_{\Omega} |Ef_1 Ef_2|^2 w_{\Omega} dx \leq \sum_{|\tau|=\frac{N_2}{N_1}} \left| \int_{\Omega} Ef_{1,\tau} Ef_2 \right|^2 w_{\Omega} dx$$

Now apply Proposition 3.1 for $f_{1,\tau}$ and f_2 for each τ , Theorem 1.3 follows.

5. Transversality

Let f_1, f_2 be as in Proposition 3.1, then f_1 is supported around $(0, 0, \ldots, 0, 1, 1)$ and f_2 is supported around $(0, 0, \ldots, 0)$. The main goal of this section is to explore the transversality between $(0, 0, \ldots, 0, 1)$ and $(0, 0, 0, \ldots, 0)$, or more precisely, the transversality between the unit normal vectors of the truncated parabola at these two points. The main lemma in this section is Lemma 5.1 below, and Corollary 5.7 which essentially follows from Lemma 5.1.

We first introduce some basic notation. Let (e_1, \ldots, e_d) be the standard basis of \mathbb{R}^d . We will encounter caps of radius v around $(0, 0, \ldots, 0)$ and $(0, \ldots, 0, 1, 1)$ on the parabola. Note around those two points, when v is small (which is always the case in our work), one may view those caps as their natural projection to \mathbb{R}^{d-1} . And their image is essentially a square/cap of radius v. We say that a (v, v^2) -plate is a d-dimensional rectangle with the short side on e_{d-1} direction such that its image under under the orthogonal projection to \mathbb{R}^{d-1} is a $v \times v \times \cdots \times v \times v^2$ -rectangle.

Lemma 5.1. Given |v| < 1, let f_1 be a function supported on a cap of radius v, centered at $(0, \ldots, 0, 1, 1)$ on the truncated parabola P, and let f_2 be a function supported on a cap of radius v centered at $(0, \ldots, 0, 0, 0)$ on the paraboloid. For a covering $\{\tau_i\}$ of supp f_i with (v, v^2) -plates, with the shorter side on e_{d-1} direction. We have the following decoupling inequality, for any $R > v^{-2}$,

$$\int |Ef_1 Ef_2|^2 w_{B_R} \lesssim \sum_{\tau_1, \tau_2} \int |Ef_{1, \tau_1} Ef_{2, \tau_2}|^2 w_{B_R}.$$
(5.1)

Remark 5.2. We thank J. Ramos for pointing out that Lemma 5.1 is a particular case of Proposition 2 in his work [15]. We still write a proof in this paper for clarity.

Proof. The proof is similar to the proof of the L^4 Strichartz estimate on the one dimensional torus. From the inequality 1.15, we only need to prove that

$$\int_{B'} |Ef_1 Ef_2|^2 \lesssim \sum_{\tau_1, \tau_2} \int |Ef_{1, \tau_1} Ef_{2, \tau_2}|^2 \mu_{B'}^2$$

for all translation B' of B_R .

$$\int_{B'} |Ef_1 Ef_2|^2 \le \sum_{\tau_1, \tau_2, \tau_3, \tau_4} \int_{B'} Ef_{1, \tau_1} Ef_{2, \tau_2} \overline{Ef_{1, \tau_3} Ef_{2, \tau_4}} \mu_{B'}^2$$
(5.2)

Let $\xi_i \in \tau_i, \, \xi_i = (\xi_{i,1}, \dots, \xi_{i,d-1}, \sum_{j=1}^{d-1} (\xi_i^j)^2) \equiv (\bar{\xi}_i, \xi_{i,d-1}, |\bar{\xi}_i|^2 + (\xi_i^{d-1})^2), \, i = 1, 2, 3, 4.$ We have

$$\begin{aligned} |\bar{\xi}_i| &\lesssim \upsilon, & i = 1, 2, 3, 4. \\ |\xi_{i,d-1} - 1| &\lesssim \upsilon, & i = 1, 3. \\ |\xi_{i,d-1}| &\lesssim \upsilon, & i = 2, 4. \end{aligned}$$
(5.3)

Essentially, for any $\tau_1, \tau_2, \tau_3, \tau_4$ such that

$$\int Ef_{1,\tau_1} Ef_{2,\tau_2} \overline{Ef_{1,\tau_3} Ef_{2,\tau_4}} \mu_{B'}^2 \neq 0,$$

one must have for some $\xi_i \in \tau_i$,

$$\xi_1 - \xi_3 = \xi_2 - \xi_4 + O(R^{-1}),$$

$$|\xi_1|^2 - |\xi_3|^2 = |\xi|_2^2 - |\xi|_4^2 + O(R^{-1}),$$
(5.4)

and the second formula in (5.4) implies

$$(\xi_{1,d-1} - \xi_{3,d-1})(\xi_{1,d-1} + \xi_{3,d-1}) = O(|\xi_2|^2 + |\xi_4|^2) + O(|\bar{\xi}_1|^2 + |\bar{\xi}_3|^2) + O(R^{-1}).$$
(5.5)

Plugging into (5.3), one has $|\xi_{1,d-1} - \xi_{3,d-1}| \leq v^2$, which again implies $|\xi_{2,d-1} - \xi_{4,d-1}| \leq v^2$.

To summarize, $\int Ef_{1,\tau_1} Ef_{2,\tau_2} \overline{Ef_{1,\tau_3}} Ef_{2,\tau_4} \mu_{B'}^2 \neq 0$ implies the distance between τ_1 and τ_3 and the distance between τ_2 and τ_4 are both bounded by v^2 , which essentially means $\tau_i = \tau_{i+2}$, i = 1, 2. Applying this fact to (5.2), Lemma 5.1 follows.

Remark 5.3. A quantitative version of estimate (5.1) can be stated as follows: assume that the support of f_1 is centered at $(0, 1/K, (1/K)^2)$ rather than (0, 0, 1), from the proof we can attain the same estimate as in (5.1) by introducing an additional constant K,

$$\int |Ef_1 Ef_2|^2 w_{B_R} \lesssim K \sum_{\tau_1, \tau_2} \int |Ef_{1, \tau_1} Ef_{2, \tau_2}|^2 w_{B_R}$$
(5.6)

Indeed, the proof essentially only relies on the fact that for $\xi_i \in supp f_i, i = 1, 2$, the difference between the d-1 components is at least $\frac{1}{K}$. Similar arguments also hold for estimate in Lemma 5.5, Cor 5.7 below.

Remark 5.4. We remark that for any $\alpha < v$, a function which is supported on a cap of radius α can be naturally understood as a function supported on a cap of radius v.

Lemma 5.1 facilitates the decomposition of caps of radius v into plates of size (v, v^2) , we can further decompose those into caps of radius v^2 .

Lemma 5.5. With same notation as in Lemma 5.1, $R \ge v^{-2}$, let $suppf_i$ be the covered by finitely overlapping caps θ_i of radius v^2 , i = 1, 2. Then

$$\int |Ef_1 Ef_2|^2 w_{B_R} \lesssim v^{-(d-1)} \sum_{|\theta_i| = v^2} \int |Ef_{1,\theta_1} Ef_{2,\theta_2}|^2 w_{B_R}.$$
(5.7)

Proof. Clearly, we need only to prove (5.7) for every ball of radius v^{-2} contained in B_R , and then sum them together. (This is in the same principle of parallel decoupling, Lemma 1.9.)

Fix a pair of (v, v^2) -plates τ_1, τ_2 .

$$\int |Ef_{1,\tau_1} Ef_{2,\tau_2}|^2 w_{B_R} = \int |\sum_{\theta_2 \subset \tau_2, |\theta_2| = v^2} Ef_{1,\tau_1} Ef_{2,\theta_2}|^2 w_{B_R}$$

$$\leq v^{-(d-1)} \sum_{\theta_2 \subset \tau_2, |\theta_2| = v^2} |Ef_{1,\tau_1} Ef_{2,\theta_2}|^2 w_{B_R}$$

$$\lesssim \sum_{\theta_j \subset \tau_j, |\theta_j| = v^2} |Ef_{1,\theta_1} Ef_{2,\theta_2}|^2 w_{B_R}$$
(5.8)

The last inequality follows from Lemma 4.1 and Lemma 4.2.

Remark 5.6. Similar to Remark 5.4, for $v^2 < \alpha < v$, a cap of scale v naturally lies in a cap of scale $\sqrt{\alpha}$. Thus if we let f_1 be a function supported on a cap of radius α , centered at $(0, \ldots, 0, 1, 1)$ on the paraboloid and we let f_2 be a function supported on a cap of radius α centered at $(0, \ldots, 0, 0, 0)$ on the paraboloid, then by arguing similar to the proof of Lemma 5.5, we have for $R \ge \alpha^{-1}$,

$$\int |Ef_1 Ef_2|^2 w_{B_R} \lesssim (\nu/\alpha)^{(d-1)} \sum_{|\theta_i|=\alpha} \int |Ef_{1,\theta_1} Ef_{2,\theta_2}|^2 w_{B_R}.$$
(5.9)

If we directly use Holder inequality for all caps in the support of f_i to estimate as in (5.8), then the interpolation in the proof of Lemma 5.5 will give us a constant v^{-d} rather than $v^{-(d-1)}$ in (5.7), since one has v^{-d} caps for each f_i . The bilinear transversality, i.e. the transversality between $(0, 0, \ldots, 0)$ and $(0, \ldots, 0, 1, 1)$ helps in reducing the dimension by one since in one direction we can use L^4 orthogonality, as shown in Lemma 5.1. Thus here we are able to improve the constant in (5.7) to $v^{-(d-1)}$.

Corollary 5.7. Same notation as in Lemma 5.1, there exists a constant C, such that for any $v, \delta, R^{-1} \leq \delta \leq v$,

$$\int |Ef_1 Ef_2|^2 w_{B_R} \lesssim \left(\frac{\upsilon}{\delta}\right)^{d-1} \left|\frac{\log\delta}{\log\upsilon}\right|^C \sum_{|\theta_i|=\delta} \int |Ef_{1,\theta_1} Ef_{2,\theta_2}|^2 w_{B_R}.$$

Proof. The proof is most clear when $\delta = v^{2^n}$ for some *n*, let us first handle this case and then go to the general case. One may use induction. (This induction, however, does not rely on parabolic rescaling.) If n = 0, there is nothing to prove.

Assume the result holds for the case n = k, let us turn to the case n = k + 1, where $\delta = v^{2^{k+1}}$, $\delta^{1/2} = v^{2^k}$, thus by induction assumption, we have

$$\int |Ef_1 Ef_2|^2 w_{B_R} \lesssim \left(\frac{\upsilon}{\delta^{1/2}}\right)^{d-1} 2^{Ck} \sum_{|\eta_i| = \delta^{1/2}} \int |Ef_{1,\eta_1} Ef_{2,\eta_2}|^2 w_{B_R}.$$
(5.10)

Now note $R \ge (\delta^{-1/2})^2$, by Lemma 5.5, we have for each pair (η_1, η_2) in (5.10) that

$$\int |Ef_{1,\eta_1} Ef_{2,\eta_2}|^2 w_{B_R} \lesssim (\delta^{1/2})^{-(d-1)} \sum_{\theta_i \subset \eta_i, |\theta_i| = \delta} \int |Ef_{1,\theta_1} Ef_{2,\theta_2}|^2 w_{B_R}.$$
(5.11)

The case n = k + 1 clearly follows if one plugs (5.11) into (5.10), taking the constant C large enough.

Now we turn to the general case, we only need to work on the case $v^{2^{n+1}} < \delta < v^{2^n}$. Recall that previously, when $\delta = v^{2^n}$, we used induction as $v \to v^2 \to v^{2^2} \dots \to v^{2^n} = \delta$, and in each step we used Lemma 5.5 to finish the induction $v^{2^k} \to v^{2^{k+1}}$.

the induction $v^{2^k} \to v^{2^{k+1}}$. In the case $v^{2^{n+1}} < \delta < v^{2^n}$ we have $v^{2^n} < \delta^{1/2}$, and we use induction as before for $v \to v^2 \to v^{2^2} \dots \to v^{2^n}$, and we use (5.9) to use induction again from v^{2^n} to δ . This ends the proof.

6. Proof of Lemma 3.2, Lemma 3.4 and Lemma 3.6

We are now prepared to use transversality to prove Lemma 3.2, Lemma 3.4, and Lemma 3.6. Recall Lemma 3.2 concerns $\tilde{K}(\lambda, N_1, N_2)$ defined in (3.4). Furthermore, Lemma 3.4 and Lemma 3.6 refer to $K(\lambda, N_1, N_2)$ defined in (3.5).

6.1. **Proof of Lemma 3.2.** For convenience of notation, we let $\Omega_1 := [0, N_1^2] \times [0, \lambda N_1]^d$. Note that one can use finite overlapped balls of radius N_1^2 to cover Ω_1 since $\lambda \ge N_1$. We want to prove

$$\|Ef_1Ef_2\|_{L^2_{avg}(\omega_{\Omega_1})} \lesssim_{\epsilon} \lambda^{d/2} N_2^{\epsilon} \frac{N_2^{d-1}}{N_1} \prod_{j=1}^2 \left(\sum_{|\theta| = \frac{1}{\lambda N_1}} \|Ef_{j,\theta}\|_{L^4_{avg}(w_{\Omega_1})}^2 \right)^{1/2}.$$
(6.1)

We first apply Corollary 5.7 with $\delta = N_1^{-2}$, $v = \frac{N_2}{N_1}$, $R = N_1^2$. Note that $\delta \leq v$. Then we have

$$\int |Ef_1 Ef_2|^2 w_{B_{N_1^2}} \lesssim (N_1 N_2)^{d-1} \left| \frac{\log N_1}{\log N_1 - \log N_2} \right|^C \sum_{|\theta_j| = \frac{1}{N_1^2}} \int |Ef_{1,\theta_1} Ef_{2,\theta_2}|^2 w_{B_{N_1^2}}$$

$$\lesssim (N_1 N_2)^{d-1} N_2^{\epsilon} \sum_{|\theta_j| = \frac{1}{N_1^2}} \prod_{j=1}^2 ||Ef_{j,\theta_j}||_{L^4(w_{B_{N_1^2}})}^2.$$
(6.2)

Remark 6.1. We avoid the case when $N_1 = N_2$, and thus $\ln N_1 - \ln N_2 = 0$, by first decomposing caps of diameter N_2/N_1 into caps of diameter $N_2/2N_1$ with loss of a fixed constant, then continuing with the proof as above. In all of the text that follows, one may assume, without loss of generality, that $N_1 \ge 2N_2$.

Via the principle of parallel decoupling, Lemma 1.9, or by summing different $B_{N_1^2}$ together, we have

$$\int |Ef_1 Ef_2|^2 w_{\Omega_1} \lesssim (N_1 N_2)^{d-1} N_2^{\epsilon} \sum_{|\theta_j| = \frac{1}{N_1^2}} \prod_{j=1}^2 ||Ef_{j,\theta_j}||_{L^4(w_{\Omega_1})}^2.$$
(6.3)

Next we would like to show that

$$\|Ef_{j,\theta_j}\|_{L^4(w_{\Omega_1})}^2 \le \left(\frac{\lambda}{N_1}\right)^{d/2} \sum_{\theta_j' \subset \theta_j, |\theta_j'| = \frac{1}{\lambda N_1}} \|Ef_{j,\theta_j'}\|_{L^4(w_{\Omega_1})}^2.$$
(6.4)

It suffices to show

$$\|Ef_{j,\theta_j}\|_{L^4_{avg}(\Omega_1)}^2 \le \left(\frac{\lambda}{N_1}\right)^{d/2} \sum_{\theta_j' \subset \theta_j, |\theta_j'| = \frac{1}{\lambda N_1}} \|Ef_{j,\theta_j'}\|_{L^4_{avg}(w_{\Omega_1})}^2$$

and sum up as in Lemma 4.1

Each function Ef_{j,θ'_j} is fourier supported in θ'_j , in particular, fourier supported in a cylinder of radius $\frac{1}{\lambda N_1}$, height $\frac{1}{N_1^2}$. Ω_1 is tiled by cylinders of radius λN_1 , height N_1^2 in *t*-direction. The proof of Lemma 4.1 works the same,

$$\begin{split} \|Ef_{j,\theta_{j}}\|_{L^{2}_{avg}(\Omega_{1})}^{2} \lesssim \sum_{\theta_{j}^{\prime} \subset \theta_{j}, |\theta_{j}^{\prime}| = \frac{1}{\lambda N_{1}}} \|Ef_{j,\theta_{j}^{\prime}}\|_{L^{2}_{avg}(w_{B_{R}})}^{2} \\ \lesssim \sum_{\theta_{j}^{\prime} \subset \theta_{j}, |\theta_{j}^{\prime}| = \frac{1}{\lambda N_{1}}} \|Ef_{j,\theta_{j}^{\prime}}\|_{L^{4}_{avg}(w_{B_{R}})}^{2} \end{split}$$

For the L^{∞} -estimate, we apply Cauchy Schwartz inequality:

$$\begin{split} \|Ef_{j,\theta_j}\|_{L^{\infty}(\Omega_1)}^2 &\leq \left(\frac{\lambda}{N_1}\right)^d \sum_{\substack{\theta_j' \subset \theta_j, |\theta_j'| = \frac{1}{\lambda N_1}}} \|Ef_{j,\theta_j'}\|_{L^{\infty}(\Omega_1)}^2 \\ &\lesssim \left(\frac{\lambda}{N_1}\right)^d \sum_{\substack{\theta_j' \subset \theta_j, |\theta_j'| = \frac{1}{\lambda N_1}}} \|Ef_{j,\theta_j'}\|_{L^4_{avg}(w_{\Omega_1})}^2 \end{split}$$

The last inequality is an application of Lemma 1.10. Note $f_{\theta'_j}$ is supported in a ball of scale $\frac{1}{\lambda N_1}$, and inside a box C of size $\frac{1}{N_1^2} \times \frac{1}{\lambda N_1} \times \cdots \times \frac{1}{\lambda N_1}$. We can make a affine transform of C into a cube Q^* of scale λ_{N_1} , which on the physical side would transform Ω_1 into a cube of scale λN_1 . We apply Lemma 1.10 after the affine transformation and then transform back. (Note in those setting, cube is no different than a ball.)

We apply Hölder's inequality to conclude the argument.

6.2. **Proof of Lemma 3.4.** Let $\lambda \leq N_1$. We first note that we can use finitely overlapping balls $B_{\lambda N_1}$ to cover Ω and that $N_2 \lesssim 1$. Applying Corollary 5.7 with $\delta = \frac{1}{\lambda N_1}$ and $v = \frac{N_2}{N_1}$ we have

$$\begin{split} \int |Ef_1 Ef_2|^2 w_{B_{\lambda N_1}} &\lesssim (\lambda N_2)^{d-1} \Big| \frac{\log \lambda + \log N_1}{\log N_1 - \log N_2} \Big|^C \sum_{|\theta_j| = \frac{1}{\lambda N_1}} \int |Ef_{1,\theta_1} Ef_{2,\theta_2}|^2 w_{B_{\lambda N_1}} \\ &\lesssim (\lambda N_2)^{d-1} N_2^{\epsilon} \sum_{|\theta_j| = \frac{1}{\lambda N_1}} \prod_{j=1}^2 \|Ef_{j,\theta_j}\|_{L^4(w_{B_{\lambda N_1}})}^2. \end{split}$$

With parallel decoupling, Lemma 1.9, then the desired estimate follows. (As remarked in Remark 6.1, one can assume $N_1 \geq 2N_2$.)

6.3. Proof of Lemma 3.6. Let $\lambda < N_1$.

We have the following two cases:

- Case 1: d = 2, N₁ ≥ λ ≥ N₁/N₂, and N'₂ = (N₁/λ),
 Case 2: d ≥ 3, N₁ ≥ λ ≥ N₁/N₂², and N'₂ = (N₁/λ)^{1/2}.

It is easy to check that we only need to show that

$$K(\lambda, N_1, N_2) \lesssim K(\lambda, N_1, N_2') \left(\frac{N_1}{N_2'} \frac{N_2}{N_1}\right)^{\frac{d-1}{2}}.$$
 (6.5)

We claim that

$$\|Ef_1Ef_2\|_{L^4_{avg}(w_{B_{N_1^2}})} \lesssim \left(\frac{N_2/N_1}{N_2'/N_1}\right)^{d-1/2} \prod_{j=1}^2 \left(\sum_{|\theta|=\frac{N_2'}{N_1}} \|Ef_{j,\theta}\|_{L^4_{avg}(w_{B_{N_1^2}})}^2\right)^{1/2}.$$
(6.6)

Since $\lambda \leq N_1$, we cover $B_{N_1^2}$ with balls of radius λN_1 . Thus by parallel decoupling, to prove (6.6), we only need to show \ 1/2

$$\|Ef_1Ef_2\|_{L^4_{avg}(w_{B_{\lambda N_1}})} \lesssim \left(\frac{N_2/N_1}{N_2'/N_1}\right)^{d-1/2} \prod_{j=1}^2 \left(\sum_{|\theta|=\frac{1}{\lambda N_1}} \|Ef_{j,\theta}\|_{L^4_{avg}(w_{B_{\lambda N_1}})}^2\right)^{1/2}.$$
(6.7)

Note that since $\lambda N_1 \geq \frac{1}{N'_2/N_1}$, estimate (6.7) follows from Corollary (5.7) by setting $\delta = N_2/N_1$, $\upsilon = N'_2/N_1$ via interpolation and local constant arguments as in Section 6.1.

By the definition of $K(\lambda, N_1, N_2)$, we have that for any θ_1, θ_2 in (6.7),

$$\|Ef_{1,\theta_1}Ef_{2,\theta_2}\|_{L^4_{avg}(w_{B_{\lambda N_1}})} \lesssim \lambda^{d/2} K(\lambda, N_1, N_2') \prod_{j=1}^2 \left(\sum_{|\theta_j'| = \frac{1}{\lambda N_1}, \theta_j \subset \theta_j} \|Ef_{j,\theta_j'}\|_{L^4_{avg}(w_{\Omega})}^2 \right)^{1/2}.$$
(6.8)

Plugging (6.8) into (6.7), clearly (6.5) follows.

7. Induction procedure and proof of Lemma 3.3

To conclude the proof of Proposition 3.1, we are left with the proof of Lemma 3.3. For this lemma the proof relies on induction on N_2 . The base case $N_2 \leq 1$ is resolved by Lemma 3.4, and by Lemma 3.6, so we need only to induct until $\lambda = \frac{(N_2)^{d-1}}{N_1}$.

Let f_1, f_2 be as in Lemma 3.3. Applying Lemma 5.1, taking $v = N_1/N_2$ and $R = N_1^2$, we could decouple the $\frac{N_2}{N_1}$ caps into $\left(\frac{N_2}{N_1}, \frac{N_2^2}{N_1^2}\right)$ plates without any loss, i.e.

$$\int |Ef_1 Ef_2|^2 w_{B_{N_1^2}} \lesssim \sum_{\tau_1, \tau_2} \int |Ef_{1, \tau_1} Ef_{2, \tau_2}|^2 w_{B_{N_1}^2}.$$
(7.1)

Here τ_i are plates as described in Lemma 5.1. We focus on the case when d = 2 in \mathbb{R}^3 , the high dimensional case would be explained in the end. When d = 2, the underlying plates become strips. We start with some preparation before the induction.

7.1. Preliminary preparation for the induction. We fix a pair of $\left(\frac{N_2}{N_1}, \frac{N_2^2}{N_1^2}\right)$ strips τ_1, τ_2 from estimate (7.1). We decompose τ_j into a union of $\frac{N_2}{KN_1} \times \frac{N_2^2}{N_1^2}$ strips $\{s_j\}$.

Using the notation *nonadj* short for *nonadjacent*, and *adj* short for *adjacent*, we have

$$\begin{split} |Ef_{\tau_j}|^2 &= \sum_{s_j} |Ef_{s_j}|^2 + \sum_{s_j, s'_j adj} |Ef_{s_j} Ef_{s'_j}| + \sum_{s_j, s'_j nonadj} |Ef_{s_j} Ef_{s'_j}| \\ &\leq 10 \sum_{s_j} |Ef_{s_j}|^2 + \sum_{s_j, s'_j nonadj} |Ef_{s_j} Ef_{s'_j}| \\ &= I_{j,1} + I_{j,2} \end{split}$$

$$\int |Ef_{\tau_1} Ef_{\tau_2}|^2 w_{B_{N_1^2}} \le \int |(Ef_{\tau_1}^2 - I_{1,1})(|Ef_{\tau_2}^2 - I_{2,1})| + Ef_{\tau_1}^2 I_{2,1} + Ef_{\tau_2}^2 I_{1,1} + I_{1,1} I_{2,1} w_{B_{N_1^2}}$$
(7.2)

$$\lesssim \sum_{s_j, s'_j nonadj} \int |Ef_{s_1} Ef_{s_1} Ef_{s_2} Ef_{s'_2} | w_{B_{N_1^2}} + \sum_{s_1, s_2} \int |Ef_{s_1} Ef_{s_2}|^2 w_{B_{N_1^2}}$$
(7.3)

The last inequality follows from Lemma 4.1 and Lemma 4.2.

The reason why we want to have non-adjacent parts is that we would like transversality (after rescaling) on the other direction. Formula (7.3) will the starting point of our induction.

For the second term in (7.3), we will later directly use induction (not relying on parallel rescaling) on N_2 and reduce everything to the known base case $N_2 = 1$.

For the first term, using Cauchy-Schwartz

$$\int |Ef_{s_1}Ef_{s_1'}Ef_{s_2}Ef_{s_2'}|w_{B_{N_1^2}} \le \left(\int |Ef_{s_1}Ef_{s_1'}|^2 w_{B_{N_1^2}}\right)^{1/2} \left(\int |Ef_{s_2}Ef_{s_2'}|^2 w_{B_{N_1^2}}\right)^{1/2}.$$
(7.4)

We point out here that in what follows we do not rely on the bilinear transversality between s_1 and s_2 (or s_1 and s'_2), which is already handled in Lemma 5.1. Instead we will rely on the bilinear transversality between s_1 and s'_1 , (or s_2, s'_2), since they are not adjacent. This transversality is most clear when one applies parabolic rescaling.

Let us now turn to the term $\int |Ef_{s_2}Ef_{s'_2}|^2 w_{\Omega}$, when s_2, s'_2 are non adjacent. The term with s_1, s'_1 is handled similarly, though one may need to rotate the coordinates.

Finally we point out here that K would be chosen large later and any (fixed) power of K will not impact the final estimate. In particular, in the following estimates we would not worry about losing powers of K.

Without loss of generality, we assume

- s_2 is the strip that $\{(a_1, a_2, a_1^2 + a_2^2) | |a_1| \le N_2^2/N_1^2, |a_2| \le N_2/KN_1\}$
- s'_2 is the strip that $\{(b_1, b_2, b_1^2 + b_2^2) | |b_1| \le N_2^2/N_1^2, |b_2 CN_2/KN_1| \le N_2/KN_1\}, C \ge 10$. (Here 10 is of course just some universal constant.)

7.2. **Parabolic rescaling.** The next step, parabolic scaling, is standard in decoupling types results; we give the details here for the convenience of the reader.

Note s_2, s'_2 lie on the same $\frac{N_2}{N_1}$ cap. We rescale the $\frac{N_2}{N_1}$ cap to radius 1. By a slight abuse of notation, we regard f_{s_i} as a function depending only on two variables $(\xi_{i,1}, \xi_{i,2})$. For convenience notation, we let $h_1 = f_{s_2}, h_2 = f_{s'_2}$. Let also $g_i(\eta_{i,1}, \eta_{i,2}) := h_i((N_2/N_1)\eta_{i,1}, (N_2/N_1)\eta_{i,2})$.

Note now

- g_1 is supported in the strip of $\{(a_1, a_2, a_1^2 + a_2^2) | |a_1| \le N_2/N_1, |a_2| \le 1/K\}$
- g_2 is supported in the strip of $\{(b_1, b_2, b_1^2 + b_2^2) \mid |b_1| \le N_2/N_1, |b_2 C/K| \le 1/K\}, C \ge 10$

Note g_1, g_2 are supported on a pair of transverse $\frac{N_2}{N_1} \times 1$ strips¹ due to the non adjacency of s_2, s'_2 . We point out here the transversality between g_1, g_2 is not as in the assumption of Lemma 5.1, but it is in the sense of Remark 5.3, which usually cause a loss of K in the estimate, but this does not matter.

The *parabolic scaling* says the following:

¹ Strictly speaking, we need them to support on a pair of $\frac{N_2}{N_1} \times \frac{1}{100}$ strips, we neglect this technical point here.

Claim 7.1. Let $Eg_i(y_1, y_2, y_3) = Eh_i(N_1/N_2y_1, N_1/N_2y_2, N_1^2/N_2^2y_3)$, and let D be domain in \mathbb{R}^3 and let

$$\hat{D} := \{ (y_1, y_2, y_3) : N_1 / N_2 y_1, N_1 / N_2 y_2, N_1^2 / N_2^2 y_3 \in D \}$$

then it follows from standard change of variables technique that the following two estimates, with the same constant A, are equivalent:

$$\|Eh_1Eh_2\|_{L^2_{avg}(w_D)} \lesssim A \prod_{j=1}^2 \left(\sum_{|\theta| = \frac{1}{\lambda N_1}} \|Ef_{s_j,\theta}\|_{L^4_{avg}(w_D)}^2 \right)^{1/2},$$
(7.5)

$$\|Eg_1Eg_2\|_{L^2_{avg}(w_{\bar{D}})} \lesssim A \prod_{j=1}^2 \left(\sum_{|\tilde{\theta}| = \frac{1}{\lambda N_2}} \|Eg_{j,\tilde{\theta}}\|_{L^4_{avg}(w_{\bar{D}})}^2 \right)^{1/2}.$$
(7.6)

we then concentrate on (7.6).

Take $D = B_{N_1^2}$, then $\tilde{D} = [0, N_2^2] \times [0, N_1 N_2]^2$. (Here, without loss of generality, we regard $B_{N_1^2}$ as $[0, N_1^2]^3$.) For convenience of notation, we set $\tilde{\Omega} = [0, N_2^2] \times [0, N_1 N_2]^2$. The parabolic rescaling gives

Lemma 7.2. Assume g_1, g_2 are two general functions defined on the parabola. Let g_1 be supported in a strip of size $N_2/N_1 \times 1$ around (0,0,0), and g_2 be supported in a strip of size $N_2/N_1 \times 1$ around (0,1,1). If for some constant A, one has (for all such g_1, g_2),

$$\|Eg_1Eg_2\|_{L^2_{avg}(w_{\bar{\Omega}})} \lesssim A\left(\sum_{|\tilde{\theta}|=\frac{1}{\lambda N_2}} \|Eg_{j,\tilde{\theta}}\|_{L^4_{avg}(w_{\bar{\Omega}})}^2\right)^{1/2},\tag{7.7}$$

then for the same constant A, one has

$$\|Ef_{s_2}Ef_{s'_2}\|_{L^2_{avg}(w_{B_{N_1^2}})} \lesssim K^C A\left(\sum_{|\theta|=\frac{1}{\lambda N_1}} \|Ef_{s_2,\theta}\|_{L^4_{avg}(w_{B_{N_1^2}})}^2\right)^{1/2} \left(\sum_{|\theta|=\frac{1}{\lambda N_1}} \|Ef_{s'_2,\theta}\|_{L^4_{avg}(w_{B_{N_1^2}})}^2\right)^{1/2}.$$
 (7.8)

Remark 7.3. After rescaling, the relevant g_1, g_2 should be supported around (0,0,0) and $(0,1/K,1/K^2)$ rather than (0,0,0) and (0,0,1). We state our lemma for g_1, g_2 supported around (0,0,0) and (0,1,1) to be consistent with the statement in Lemma 5.1. This causes a loss of K^C , but we emphasize again that any loss due to a power of K would be irrelevant in the proof.

We end this section by introducing some notation.

Let g_1, g_2 be as in Lemma 7.2, we define $A(\lambda, N_1, N_2)$ to be the best constant such that

$$\|Eg_1Eg_2\|_{L^2_{avg}(w_{\bar{\Omega}})} \lesssim A(\lambda, N_1, N_2) \left(\sum_{|\tilde{\theta}| = \frac{1}{\lambda N_2}} \|Eg_{j,\tilde{\theta}}\|_{L^4_{avg}(w_{\bar{\Omega}})}^2 \right)^{1/2}.$$
(7.9)

Then we can restate Lemma (7.2).

Lemma 7.4. For j = 1, 2, we have

$$\|Ef_{s_{j}}Ef_{s_{j}'}\|_{L^{2}_{avg}(w_{B_{N^{2}_{1}}})}$$

$$\lesssim K^{C}A(\lambda, N_{1}, N_{2}) \left(\sum_{|\theta| = \frac{1}{\lambda N_{1}}} \|Ef_{s_{j},\theta}\|_{L^{4}_{avg}(w_{B_{N^{2}_{1}}})}^{2} \right)^{1/2} \left(\sum_{|\theta| = \frac{1}{\lambda N_{1}}} \|Ef_{s_{j}',\theta}\|_{L^{4}_{avg}(w_{B_{N^{2}_{1}}})}^{2} \right)^{1/2} .$$

$$(7.10)$$

7.3. The induction procedure.

7.3.1. Before induction. Now we are ready to start the induction for the proof of Lemma 3.3. We emphasize here the induction is on N_2 , (though mixed with induction on K). Note we are now in dimension d = 2.

We need to show that for all $1 \leq N_2 \leq N_1$ and $\lambda \leq N_1$, one has

$$K(\lambda, N_1, N_2) \lesssim N_2^{\epsilon} (\frac{1}{\lambda} + \frac{N_2}{N_1})^{1/2}.$$

Note the base case $N_2 = 1$ is already established in Corollary 3.4. And with Lemma 3.6, we need only to perform induction until $\lambda = N_2/N_1$.

We will work on $A(\lambda, N_1, N_2)$ defined in (7.9) to explore the transversality between nonadjacent strips. The induction process is two fold in some sense. We will induct on N_2 to better understand $K(\lambda, N_1, N_2)$, and in turn we find more information about $A(\lambda, N_1, N_2)$, which in turn gives a better understanding of $K(\lambda, N_1, N_2)$.

This is a final summary before we start the induction. Recall, we have (7.1) and (7.3), thus we have

$$\int |Ef_{1}Ef_{2}|^{2} w_{B_{N_{1}^{2}}}
\lesssim \int \sum_{s_{j},s_{j}'nonadj} \int |Ef_{s_{1}}Ef_{s_{1}}Ef_{s_{2}}Ef_{s_{2}'}| w_{B_{N_{1}^{2}}}
+ \int_{s_{1},s_{2}} \int |Ef_{s_{1}}Ef_{s_{2}}| w_{B_{N_{1}^{2}}}.$$
(7.11)

Also recall that s_1, s'_1, s_2, s'_2 are all $(N_2/N_1)^2 \times N_2/KN_1$ strips. The second term can be easily handled by direct induction, (which is not the main point of the induction procedure explained later). Indeed, if there were only the second term in (7.11), since s_1, s_2 are both contained in caps of radius (N_2/KN_1) , then (7.11) already reduces the decoupling problem for f_i supported in caps of size N_2/N_1 into the decoupling problem for f_i supported in caps of size N_2/K_1 .

We will focus on the first term of (7.11). Hölder inequality gives

$$\int |Ef_{s_1} Ef_{s_1'} Ef_{s_2} Ef_{s_2'}| w_{B_{N_1^2}} \le \prod_{j=1}^2 \left(\int |Ef_{s_j} Ef_{s_j'}|^2 w_{B_{N_1^2}} \right)^{1/2}.$$
(7.12)

Estimate (7.12) is the start point of the analysis in the following Subsections.

We summarize in the lemma below how (7.12) and (7.11) come together to highlight the relevance of $A(N_1, N_2, \lambda)$ in the induction procedure.

Lemma 7.5. When $\lambda \leq N_1/N_2$ and $\lambda \leq N_1$, we have

$$K(N_1, N_2, \lambda) \lesssim K^C \frac{1}{\lambda} A(N_1, N_2, \lambda) + K(N_1, N_2/K, \lambda)$$
(7.13)

Note that the assumption of Lemma 7.5 always holds during the induction procedure to prove Lemma 3.3.

Proof of Lemma 7.5. Applying Lemma 7.4, we have

$$\|Ef_{s_{j}}Ef_{s_{j}'}\|_{L^{2}_{avg}(w_{B_{N_{1}^{2}}})}$$

$$\lesssim K^{C}A(N_{1}, N_{2}, \lambda) \left(\sum_{|\theta| = \frac{1}{\lambda N_{1}}} \|Ef_{s_{j}, \theta}\|_{L^{4}_{avg}(w_{B_{N_{1}^{2}}})}^{2} \right)^{1/2} \left(\sum_{|\theta| = \frac{1}{\lambda N_{1}}} \|Ef_{s_{j}', \theta}\|_{L^{4}_{avg}(w_{B_{N_{1}^{2}}})}^{2} \right)^{1/2} .$$

$$(7.14)$$

Plugging (7.14) into (7.12), and then plugging into (7.11), we derive

$$\begin{split} &\|Ef_{1}Ef_{2}\|\|_{l^{2}(w_{B_{N_{1}^{2}}})} \\ &\lesssim K^{C}\lambda\left(\frac{1}{\lambda}\right)^{1/2}\prod_{i=1}^{2}\left(\sum_{|\theta|=\frac{1}{\lambda N_{1}}}\|Ef_{i,\theta}\|_{L^{4}_{avg}(w_{B_{N_{1}^{2}}})}^{2}\right)^{1/2} \\ &+\left(\sum_{|\theta|=\frac{N_{2}}{\lambda K N_{1}}}\|Ef_{i,\theta}\|_{L^{4}_{avg}(w_{B_{N_{1}^{2}}})}^{2}\right)^{1/2}. \end{split}$$

$$(7.15)$$

Thus we derive

$$K(N_1, N_2, \lambda) \lesssim K^C A(N_1, N_2, \lambda) + \lambda K(N_1, N_2/K, \lambda)$$
(7.16)

Thus, Lemma 7.5 follows.

Now we are ready to start with the induction procedure on N_2 . We emphasize again that by Lemma 3.6 we only need to consider the case $\lambda \leq N_1/N_2$.

7.3.2. First induction: Case $N_2^2 \leq N_1$. It will become clear in the following proof why we choose the first splitting point at $N_1 = N_2^2$. We start with an estimate for $A(\lambda, N_1, N_2)$. We have

Lemma 7.6. When $N_2 \leq N_1^2$, $\lambda \leq N_1$, $\lambda \leq N_1/N_2$,

 λ

$$A(\lambda, N_1, N_2) \lesssim \lambda^{1/2} \equiv \lambda \lambda^{-1/2}.$$
(7.17)

Assuming Lemma 7.6 for the moment, let us finish the proof of Lemma 3.3 when $N_1 \ge N_2^2$. Applying Lemma 7.6 with Lemma 7.5, we derive

$$K(N_1, N_2, \lambda) \lesssim K^C \lambda \left(\frac{1}{\lambda}\right)^{1/2} + K(N_1, N_2/K, \lambda)$$
(7.18)

when $N_1 \ge N_2^2$ and $\lambda \le N_1/N_2$. Choosing $1 \ll K \sim N_2^{\epsilon^{10}}$, performing induction on N_2 again, and recalling that the case $N_2 \lesssim 1$ is covered by Lemma 3.4, then Lemma 3.3 follows when $N_1 \ge N_2^2$.

Now, we turn to the proof of Lemma 7.6.

Proof of Lemma 7.6. Since $N_1 \leq N_2^2$, thus $\frac{N_2}{N_1} \leq \frac{1}{N_2}$. (It is exactly because of this that we decided our first splitting point $N_1 \leq N_2^2$). Thus, the support of g_1, g_2 appearing in (7.9) are (contained in) strips of size $\frac{1}{N_2} \times 1$. Thus, in a ball of radius N_2^2 , we have

$$\int |Eg_1 Eg_2| w_{B_{N_2^2}} \lesssim \sum_{|\theta_i| = \frac{1}{N_2}, \theta_i \subset supp \ g_i} \int |E_{g_1, \theta_1} Eg_{2, \theta_2}| w_{B_{N_2^2}}.$$
(7.19)

The proof of (7.19) is essentially the same as the proof of Lemma 5.1 and we leave it to reader.

Note one can use balls $B_{N_2^2}$ to cover $\tilde{\Omega} := [0, N_2^2] \times [0, N_1 N_2]^2$, (since $N_1 \ge N_2$) thus we extend (7.19) to

$$\int |Eg_1 Eg_2| w_{\tilde{\Omega}} \lesssim \sum_{|\theta_i| = \frac{1}{N_2}, \theta_i \subset supp \, g_i} \int |E_{g_1, \theta_1} Eg_{2, \theta_2}| w_{\tilde{\Omega}}.$$
(7.20)

We claim for any fixed θ_1, θ_2 , one has

$$\|Eg_{1,\theta_1}Eg_{2,\theta_2}\|_{L^2(w_{\bar{\Omega}})} \lesssim \lambda \lambda^{-1/2} \prod_{i=1}^2 \left(\sum_{\tilde{\theta}_i \subset \theta_i, \|\tilde{\theta}_i = \frac{1}{\lambda N_2}\|} \|Eg_{i,\tilde{\theta}_i}\|_{L^4(w_{\bar{\Omega}})} \right)^{1/2}$$
(7.21)

Plugging (7.21) into (7.20), we have

$$A(N_1, N_2, \lambda) \lesssim \lambda(\frac{1}{\lambda})^{1/2}.$$
(7.22)

and the Lemma 7.6 follows.

Now we are left with the proof of (7.21). Let $N'_1 = N_2, N'_2 = N_2^2/N_1 \leq 1$. When $N'_1 = N_2 \leq \lambda$, recall the definition of $\tilde{K}(\lambda, N_1, N_2)$ in (3.4) and apply Lemma 3.2, we have

$$K(N'_1, N'_2, \lambda) \lesssim (N'_2)^{\epsilon} \left(\frac{N'_2}{N'_1}\right)^{1/2} \lesssim \left(\frac{1}{\lambda} + \frac{N_2}{N_1}\right)^{1/2} \lesssim \lambda^{-1/2}.$$
 (7.23)

The last inequality in (7.23) follows because we always have $\lambda \leq N_1/N_2$ in the whole induction process. Note (7.23) implies

$$\|Eg_{1,\theta_{1}}Eg_{2,\theta_{2}}\|_{L^{2}(w_{[0,N_{2}^{2}]\times[0,\lambda N_{2}]^{2}})}$$

$$\lesssim \quad \lambda \tilde{K}(N_{1}',N_{2}',\lambda) \prod_{i=1}^{2} \left(\sum_{\tilde{\theta}_{i}\subset\theta_{i},\|\tilde{\theta}_{i}=\frac{1}{\lambda N_{2}}\|} \|Eg_{i,\tilde{\theta}_{i}}\|_{L^{4}(w_{[0,N_{2}^{2}]\times[0,\lambda N_{2}]^{2}})} \right)^{1/2}.$$

$$(7.24)$$

Since $\lambda \leq N_1$, (which is also always the case during the induction process), Ω can be covered by the translations of $[0, N_2^2] \times [0, \lambda N_2]$, thus (7.24) implies (7.21) by the parallel decoupling Lemma 1.9.

When $\lambda \leq N'_1$, since $N'_2 \lesssim 1$, by Lemma 3.4, we have

$$K(\lambda, N_1', N_2') \lesssim \lambda^{-1/2}.$$
(7.25)

Thus,

$$\|Eg_{1,\theta_1}Eg_{2,\theta_2}\|_{L^2(w_{B_{N_2^2}})} \lesssim \lambda \lambda^{-1/2} \prod_{i=1}^2 \left(\sum_{\tilde{\theta}_i \subset \theta_i, \|\tilde{\theta}_i = \frac{1}{\lambda N_2} \|} \|Eg_{i,\tilde{\theta}_i}\|_{L^4(w_{B_{N_2^2}})} \right)^{1/2}.$$
(7.26)

Since one can use $B_{N_2^2}$ and its translations to cover $\tilde{\Omega}$, (7.26) implies (7.21) by the parallel decoupling Lemma 1.9.

7.3.3. Second induction: Case $N_2^{3/2} \le N_1 \le N_2^2$.

Lemma 7.7. When $N_2^{3/2} \le N_1 \le N_2^2$, $\lambda \le N_1$ and $\lambda \le N_1/N_2$, we have

$$A(\lambda, N_1, N_2) \lesssim \lambda^{1/2} \equiv \lambda \lambda^{-1/2}.$$
(7.27)

Clearly, using Lemma 7.5 and arguing as in Section 7.3.2, Lemma 3.3 follows from Lemma 7.7 when $N_2^{3/2} \leq N_1 \leq N_2^2$.

Now we are left with proof of Lemma 7.7, i.e. the estimate (7.27). We will prove that estimate (7.27), in case $N_2^{3/2} \leq N_1 \leq N_2^2$, follows from the fact that Lemma 3.3 holds when $N_2^2 \geq N_1$, (given Lemma 3.2).

Proof of Lemma 7.7. The proof starts similarly as the proof of Lemma 7.6, note now we have $N_2/N_1 \ge 1/N_2$. As we derived (7.19), we have in a ball of radius N_1^2/N_2^2 ,

$$\int |Eg_1 Eg_2| w_{B_{(\frac{N_1}{N_2})^2}} \lesssim \sum_{|\theta_i| = \frac{N_2}{N_1}, \theta_i \subset supp \, g_i} \int |E_{g_1, \theta_1} Eg_{2, \theta_2}| w_{B_{(N_1/N_2)^2}}.$$
(7.28)

Note one can use $B_{(\frac{N_1}{N_2})^2}$ and its translations to cover $\hat{\Omega}$, thus we have

$$\int |Eg_1 Eg_2| w_{\tilde{\Omega}} \lesssim \sum_{|\theta_i| = \frac{N_2}{N_1}, \theta_i \subset supp \ g_i} \int |E_{g_1, \theta_1} Eg_{2, \theta_2}| w_{\tilde{\Omega}}.$$
(7.29)

1 /0

The following procedure is essentially the same as in the first induction. Note that to prove (7.27) we only need to further show that for fix θ_1, θ_2 ,

$$\|Eg_{1,\theta_{1}}Eg_{2,\theta_{2}}\|_{L^{2}(w_{\tilde{\Omega}})} \lesssim \lambda \lambda^{-1/2} \prod_{i=1}^{2} \left(\sum_{\tilde{\theta}_{i} \subset \theta_{i}, |\tilde{\theta}_{i}| = \frac{1}{\lambda N_{2}}} \|Eg_{i,\tilde{\theta}_{i}}\|_{L^{4}(w_{\tilde{\Omega}})} \right)^{1/2},$$
(7.30)

where now $|\theta_i| = \frac{N_2}{N_1}$.

Let $N'_1 = N_2$, $N'_2 = \frac{N_2^2}{N_1}$, note we have $N'_1 \ge (N'_2)^2$ since $N_1 \ge N_2^{3/2}$. When $\lambda \ge N'_1$, we have by Lemma 3.2

$$\|Eg_{1,\theta_1}Eg_{2,\theta_2}\|_{L^2(w_{[0,N_2^2]\times[0,\lambda N_2]^2})} \lesssim \lambda \left(\frac{N_2'}{N_1'}\right)^{-1/2} \prod_{i=1}^2 \left(\sum_{\tilde{\theta_i}\subset\theta_i, |\tilde{\theta_i}|=\frac{1}{\lambda N_2}} \|Eg_{i,\tilde{\theta_i}}\|_{L^4(w_{[0,N_2]\times[0,\lambda N_2]^2})}\right)^{1/2}.$$
 (7.31)

Since one can use $[0, N_2^2] \times [0, \lambda N_2]^2$ to cover $\tilde{\Omega}$, (7.30) follows from (7.31), (note $\frac{N'_2}{N'_1} = \frac{N_2}{N_1} \le \lambda^{-1}$).

When $\lambda \leq N'_1$, since one can use $B_{N_2^2}$ to cover $\tilde{\Omega}$, to prove (7.30), we need only to show

$$\|Eg_{1,\theta_1}Eg_{2,\theta_2}\|_{L^2(w_{B_{N_2^2}})} \lesssim \lambda \lambda^{-1/2} \prod_{i=1}^2 \left(\sum_{\tilde{\theta_i} \subset \theta_i, \|\tilde{\theta_i} = \frac{1}{\lambda N_2}\|} \|Eg_{i,\tilde{\theta_i}}\|_{L^4(w_{B_{N_2^2}})} \right)^{1/2},$$
(7.32)

which is equivalent to $K(N'_1, N'_2, \lambda) \leq \frac{1}{\lambda}$. But recall that $N'_1 \geq (N'_2)^2$, thus this is exactly what we proved in first induction, i.e. Lemma 3.3 holds when $N_1 \geq N_2^2$.

7.3.4. Later inductions and the conclusion of the induction process. Recall that the first induction covers the case $N_1 \ge N_2^2$ and the second inductions covers the case $N_2^{\alpha} \le N_1 \le N_2^2$, $\alpha = 3/2$. The goal now is to use induction to cover the case $N_2^{\alpha} \le N_1$, all the way to $\alpha = 1$. The arguments here are similar to those for the second induction presented in Section 7.3.3. Let $N_1' = N_2, N_2' = N_2^2/N_1$, then $N_1' \ge (N_2')^{\alpha}$ is equivalent to $N_1 \ge N_2^{\frac{2\alpha-1}{\alpha}}$. Once we show that Lemma 3.3 holds when $N_2^{\alpha} \le N_1 \le N_2^2$, we would be able to extend Lemma 7.7 to the case when $N_2^{\frac{2\alpha-1}{\alpha}} \le N_1$, which in turn proves that Lemma 3.3 holds when $N_2^{\frac{2\alpha-1}{\alpha}} \le N_1 \le N_2^2$. The induction would not end until $\alpha = 1$. We finally point out, that only an induction with finite steps is involved.

To show Lemma 3.3 for a fixed ϵ_0 , we may pick an $\tilde{\epsilon} \ll \epsilon_0$, then we perform the induction for $\tilde{\epsilon}$ as above.

After we prove Lemma 3.3 for $N_1 \ge N_2^{1+\tilde{\epsilon}}$, we are left with the case $N_1 \le N_2^{1+\tilde{\epsilon}}$. We first use Hölder inequality to shrink the size of the cap from N_2/N_1 to $N_2^{1-2\tilde{\epsilon}}/N_1$, which only gives a loss of $N_2^{C\tilde{\epsilon}} \ll N_2^{\epsilon_0}$. Then we use Lemma (3.3) in the case $N_1 \ge N_2^{1+\tilde{\epsilon}}$ again.

Thus, Lemma 3.3 holds for all the case for our fixed ϵ_0 .

7.4. The high dimension case. To handle the case $d \ge 3$, we are left with the proof of Lemma 3.5. The proof is indeed similar to previous arguments in this section and easier. The proof relies on the linear decoupling estimate in [4].

As aforementioned, applying Lemma 5.1, taking $v = N_2/N_1$ and $R = N_1^2$, we can decouple the $\frac{N_2}{N_1}$ caps into $\left(\frac{N_2}{N_1}, \frac{N_2^2}{N_1^2}\right)$ plates without any loss, i.e. (7.1). However, since we are in the case $\lambda \leq N_1/N_2^2$, indeed $N_2^2/N_1^2 \leq \frac{1}{\lambda N_1}$, we only need a weaker version of (7.1), i.e. we only want to decouple the $\frac{N_2}{N_1}$ caps into $\left(\frac{N_2}{N_1}, \frac{1}{\lambda N_1}\right)$ plates:

$$\int |Ef_1 Ef_2|^2 w_{B_{N_1^2}} \lesssim \sum_{\tau_1, \tau_2} \int |Ef_{1, \tau_1} Ef_{2, \tau_2}|^2 w_{B_{N_1}^2}$$
(7.33)

Here τ_i are $(\frac{N_2}{N_1}, \frac{1}{\lambda N_1})$ plates as described in Lemma 5.1. Note (7.33) follows from (7.1).

Now, for each τ_i fixed, we further decouple τ_i into $(\frac{1}{N_1}, \frac{1}{\lambda N_1})$ plates via linear decoupling in [4], here recalled in (1.11). Note direct application of linear decoupling in dimension d gives us

$$\|Ef_{\tau_i}\|_{L^4(w_{B_{N_1^2}})} \lesssim N_2^{\epsilon}(N_2^2)^{\frac{d}{4} - \frac{d+2}{8}} \left(\sum_{v_i \subset \tau_i} \|Ef_{v_i}\|_{L^4(w_{B_{N_1^2}})}^2 \right)^{1/2},$$
(7.34)

However, we are able to use (1.11) when the dimension is d-1 rather than d, because our plates are so thin (of scale $\frac{1}{\lambda N_1} \leq \frac{1}{N_1}$), which reduce the dimension by 1. Indeed, Linear decoupling (1.11) not only work for those functions which are exactly supported in parabola P but also those which are supported in a N_1^{-2} neighborhood of P. This is consistent in uncertainty principle, since in physical space we of scale N_1^2 , in frequency space any scale of N_1^{-2} cannot be differentiated. Since our plate are so thin, of scale $\frac{1}{\lambda N_1} \leq N_1^{-2}$, one could indeed view it as a N_1^{-2} neighborhood of some d-1 dimensional parabola. To be more specific and use τ_2 as example, since τ_2 is supported at the origin. Let $\pi_t^{-1}(\tau_2)$ be the pull back image of τ_2 to the paraboloid. The fourier inverse transform of Ef_{τ_2} is supported on $\pi_t^{-1}(\tau_2)$. One can see that if we project along x_1 -axis, the projection image of $\pi_t^{-1}(\tau_2)$ is the $(\frac{1}{\lambda N_1})^2$ -neighborhood of a (d-1)-dimensional paraboloid (a piece of length $\frac{N_2}{N_1}$).

Now, apply d-1 dimensional linear decoupling, we improve (7.34) into

$$\|Ef_{\tau_i}\|_{L^4(w_{B_{N_1^2}})} \lesssim N_2^{\epsilon}(N_2^2)^{\frac{d-1}{4} - \frac{d+1}{8}} \left(\sum_{v_i \subset \tau_i} \|Ef_{v_i}\|_{L^4(w_{B_{N_1^2}})}^2 \right)^{1/2},$$
(7.35)

where v_i are $(\frac{1}{N_1}, \frac{1}{\lambda N_1})$ plates.

Finally, similarly to the derivation of (6.4), we decouple v_i into caps of radius $\frac{1}{\lambda N_1}$,

$$\|Ef_{v_i}\|_{L^4(w_{B_{N_1^2}})}^4 \lesssim \lambda^{(d-1)} \left(\sum_{\theta_i \subset v_i} \|Ef_{\theta_i}\|_{L^4(w_{B_{N_1^2}})}^2\right)^2.$$
(7.36)

We remark that each v_i can be coved by λ^{d-1} rather than λ^d caps of radius $\frac{1}{\lambda N_1}$. Plugging (7.36) into (7.35), then plugging it into (7.33), we derive

$$\|Ef_1Ef_2\|_{L^2_{avg}(w_{B_{N_1^2}})} \le \lambda^{d-1/2} N_2^{\frac{d-3}{2}} \prod_{j=1}^2 \left(\sum_{|\theta| = \frac{1}{\lambda N_1}} \|Ef_{j,\theta}\|_{L^4_{avg}(w_{B_{N_1^2}})}^2 \right)^{1/2}.$$
(7.37)

Thus, the desired estimate for $K(\lambda, N_1, N_2)$ follows.

Appendix A. Sharpness of Theorem 1.3 and Theorem 1.1

The sharpness (up to N_2^{ϵ}) of Theorem 1.3 is provided by the following examples. One can also re-scale those example to show the sharpness of Theorem 1.1.

We take
$$Ef_1 = \sum_{\xi \in \Lambda_{\lambda N_1}, |\xi| \le \frac{N_2}{N_1}} e^{2\pi i (\xi \cdot x + |\xi|^2 t)}$$
 and $f_2 = f_1(\cdot - (1, 0, \dots, 0))$. Then $|Ef_1|$ is about $(\lambda N_2)^d$ at $B(0, \frac{N_1}{N_2})$

in \mathbb{R}^{d+1} . Note that it follows from uncertainty principle, it is locally constant in any ball of size $\frac{N_1}{N_2}$ and one can easily compute $|Ef_1(0)| \sim (\lambda N_2)^d$. Also note $|Ef_1|$ has periodicity around λN_1 in all components of x, (not necessarily in t). The same is true for $|Ef_2|$. Thus,

$$\begin{aligned} \|Ef_1 Ef_2\|_{L^2(w_{\Omega})}^2 \gtrsim (\lambda N_2)^{4d} |B(0, \frac{N_1}{N_2})| (\lambda N_1)^d \\ \gtrsim \lambda^{5d} N_1^{2d+1} N_2^{3d-1} \end{aligned}$$

Each cap θ_j of radius $\frac{1}{\lambda N_1}$ contains at most one point $\xi \in \Lambda_{\lambda N_1}$. Hence $\|Ef_{j,\theta_j}\|_{L^4(w_\Omega)}^4 \lesssim |\Omega| = N_1^2 (\lambda N_1)^{2d}$.

$$\Pi_{j=1}^{2} \left(\sum_{|\theta_{j}|=\frac{1}{\lambda N_{1}}} \|Ef_{j,\theta_{j}}\|_{L^{4}(w_{\Omega})}^{2}\right) \lesssim (\lambda N_{2})^{2d} N_{1}^{2} (\lambda N_{1})^{2d}$$
$$\lesssim \lambda^{4d} N_{1}^{2d+2} N_{2}^{2d}$$

This example shows that the term with $\frac{N_2^{d-1}}{N_1}$ is sharp for both d = 2 and $d \ge 3$. When d = 2, we consider the example when

$$Ef_1 = \sum_{\xi \in \Lambda_{\lambda N_1}, \xi_1 = 1, |\xi_2| \le \frac{1}{N_1}} e^{2\pi i (\xi \cdot x + |\xi|^2 t)}$$
$$Ef_2 = \sum_{\xi \in \Lambda_{\lambda N_1}, \xi_1 = 0, |\xi_2| \le \frac{1}{N_1}} e^{2\pi i (\xi \cdot x + |\xi|^2 t)}.$$

 $|Ef_1|$ is about λ in the box of height N_1^2 (i.e. the *t* direction), width N_1 , (i.e the x_2 direction) and length $(\lambda N_1)^2$, (i.e. the x_1 direction) centered at origin. $|Ef_2|$ is the same size in the same box. Moreover, Ef_1 and Ef_2 both have periodicity around λN_1 in x_2 .

$$\begin{split} \|Ef_1Ef_2\|_{L^2(w_{\Omega})}^2 \gtrsim \lambda^4 N_1^2 \cdot N_1 \cdot (\lambda N_1)^2 \cdot \lambda N_1 \\ \gtrsim \lambda^7 N_1^6 \end{split}$$

As calculated previously, $||Ef_{j,\theta_j}||_{L^4(w_\Omega)}^4 = |\Omega|.$

$$\Pi_{j=1}^{2} \left(\sum_{|\theta_{j}| = \frac{1}{\lambda N_{1}}} \|Ef_{j,\theta_{j}}\|_{L^{4}(w_{\Omega})}^{2} \right) \lesssim \lambda^{2} \cdot |\Omega|$$
$$\lesssim \lambda^{6} N_{1}^{6}.$$

This example shows that when d = 2, the term with $\frac{1}{\lambda}$ is sharp.

When $d \geq 3$, we consider the example when

$$Ef_1 = \sum_{\xi \in \Lambda_{\lambda N_1}, \xi_1 = 1, |(\xi_2, \dots, \xi_d)| \le \frac{N_2}{N_1}} e^{2\pi i (\xi \cdot x + |\xi|^2 t)}$$
$$Ef_2 = \sum_{\xi \in \Lambda_{\lambda N_1}, \xi_1 = 0, |(\xi_2, \dots, \xi_d)| \le \frac{N_2}{N_1}} e^{2\pi i (\xi \cdot x + |\xi|^2 t)}.$$

Notice that we construct the example in $d \ge 3$ differently, the support of f_j is in a thin plate of radius $\frac{N_2}{N_1}$ instead of the $\frac{1}{N_1}$ as in 2-dimensional example.

 $|Ef_1|$ is about $(\lambda N_2)^{d-1}$ in a box of size $(\frac{N_1}{N_2}) \times \cdots \times \frac{N_1}{N_2} \times (\frac{N_1}{N_2})^2 \times (\lambda N_1)^2$. $|Ef_2|$ is about $(\lambda N_2)^{d-1}$ in the same box. Both Ef_1 and Ef_2 has periodicity around λN_1 in x_2, \ldots, x_d -directions.

$$\begin{split} \|Ef_1Ef_2\|_{L^2(w_{\Omega})}^2 \gtrsim (\lambda N_1)^{4(d-1)} (\frac{N_1}{N_2})^{d+1} (\lambda N_1)^2 (\lambda N_1)^{d-1} \\ \gtrsim \lambda^{5d-3} N_1^{2d+2} N_2^{3d-5} \\ \Pi_{j=1}^2 (\sum_{|\theta_j|=\frac{1}{\lambda N_1}} \|Ef_{j,\theta_j}\|_{L^4(w_{\Omega})}^2) \lesssim (\lambda N_2)^{2(d-1)} \cdot |\Omega| \\ \lesssim \lambda^{4d-2} N_1^{2d+2} N_2^{2d-2} \end{split}$$

This example shows that when $d \ge 3$, the term with $\frac{N_2^{d-3}}{\lambda}$ is sharp.

References

- J. Bourgain. Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. Geometric and Functional Analysis, 3(3):209–262, 1993.
- J. Bourgain. Refinements of Strichartz inequality and applications to 2D-NLS with critical nonlinearity. International Mathematics Research Notices, 1998(5):253-283, 1998.
- [3] J. Bourgain. Moment inequalities for trigonometric polynomials with spectrum in curved hypersurfaces. Israel Journal of Mathematics, 193(1):441-458, 2013.
- [4] J. Bourgain and C. Demeter. The proof of the L^2 decoupling conjecture. Annals of mathematics, 182(1):351–389, 2015.
- [5] J. Bourgain and C. Demeter. A study guide for the L^2 decoupling theorem. arXiv preprint arXiv:1604.06032, 2016.
- [6] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. Almost conservation laws and global rough solutions to a nonlinear Schrödinger equation. *Mathematical Research Letters*, 9(5), 2002.
- [7] D. De Silva, N. Pavlovic, G. Staffilani, and N. Tzirakis. Global well-posedness for a periodic nonlinear schrodinger equation in 1d and 2d. DYNAMICAL SYSTEMS, 19(1):37–65, 2007.
- [8] Y. Deng and P. Germain. Growth of solution of NLS on irrational tori. *Private communication*.
- [9] Y. Deng, P. Germain, and L. Guth. Strichartz estimates on irrational tori. Private communication.
- [10] Z. Guo, T. Oh, and Y. Wang. Strichartz estimates for Schrödinger equations on irrational tori. Proceedings of the London Mathematical Society, page pdu025, 2014.
- [11] Z. Hani. A bilinear oscillatory integral estimate and bilinear refinements to Strichartz estimates on closed manifolds. Analysis & PDE, 5(2):339–363, 2012.
- [12] S. Herr, D. Tataru, N. Tzvetkov, et al. Global well-posedness of the energy-critical nonlinear Schrödinger equation with small initial data in $H^1(\mathbb{T}^3)$. Duke Mathematical Journal, 159(2):329–349, 2011.
- [13] A. D. Ionescu, B. Pausader, et al. The energy-critical defocusing NLS on T³. Duke Mathematical Journal, 161(8):1581–1612, 2012.
- [14] R. Killip and M. Visan. Scale-invariant strichartz estimates on tori and applications. Mathematical Research Letters, 23(2):445–472, 2016.
- [15] J. Ramos. The trilinear restriction estimate with sharp dependence on the transversality. arXiv preprint arXiv:1601.05750, 2016.