Avoiding Conjugacy Classes on the 5-letter Alphabet

Golnaz Badkobeh^{a,1}, Pascal Ochem^{b,2,*}

^a Goldsmiths, University of London ^bLIRMM, Université de Montpellier, CNRS, Montpellier, France

Abstract

We construct an infinite word w over the 5-letter alphabet such that for every factor f of w of length at least two, there exists a cyclic permutation of f that is not a factor of w. In other words, w does not contain a non-trivial conjugacy class. This proves the conjecture in Gamard et al. [TCS 2018]

Keywords: Combinatorics on words, Conjugacy classes

1. Introduction

We consider infinite words over a finite alphabet that do not contain all the conjugates of the same word w, with the necessary condition that $|w| \ge 2$. A recent interest in such words appeared in the context of pattern avoidance [1] and of iterative algebras [2]. Bell and Madill [2] obtained a pure morphic word with this property (and some additional properties) over the 12-letter alphabet. Gamard et al. [1] independently obtained a morphic word over the 6-letter alphabet. They also conjectured that the alphabet size can be lowered to 5, which is best possible. In this paper, we prove this conjecture using a morphic word.

Together with the construction of a morphic binary word avoiding every conjugacy class of length at least 5 and a morphic ternary word avoiding every conjugacy class of length at least 3 [1], this settles the topic of the smallest alphabet needed to avoid every conjugacy class of length at least k.

2. Main result

Let ε denote the empty word. We consider the morphic word $w_5 = G(F^{\omega}(0))$ defined by the following morphisms.

^{*}Corresponding author

 $[\]label{eq:constraint} Email \ addresses: \ {\tt g.badkobeh@gold.ac.uk} \ (Golnaz \ Badkobeh), \ {\tt ochem@lirmm.fr} \ (Pascal \ Ochem)$

 $^{^1\}mathrm{Golnaz}$ Badkobeh is supported by the Leverhulme Trust on the Leverhulme Early Career Scheme.

²This work is supported by the ANR project CoCoGro (ANR-16-CE40-0005).

F(0) = 01 ,	$G(0) = \mathtt{abcd},$
F(1) = 2,	$G(1) = \varepsilon,$
F(2) = 03,	$G(2) = \mathtt{eacd},$
F(3) = 24 ,	$G(3) = \mathtt{becd},$
F(4) = 23.	G(4) = be.

Theorem 2.1. The morphic word $w_5 \in \Sigma_5^*$ avoids every conjugacy class of length at least 2.

In order to prove this theorem, it is convenient to express w_5 with the larger morphisms $f = F^3$ and $g = G \circ F^2$ given below. Clearly, $w_5 = g(f^{\omega}(0))$.

f(0) = 01203,	$g(\mathtt{0})=\mathtt{abcdeacd},$
f(1) = 0124,	$g(\mathtt{1}) = \mathtt{abcdbecd},$
f(2) = 0120323,	$g({\tt 2})={\tt abcdeacdbe},$
f(3) = 01240324,	$g(3) = \mathtt{abcdbecdeacdbecd},$
f(4) = 01240323.	$g(\mathtt{4})=\mathtt{abcdbecdeacdbe}.$

2.1. Avoiding conjugacy classes in $F^{\omega}(0)$

Here we study the pure morphic word and the conjugacy classes it contains.

Lemma 2.2. The infinite word $F^{\omega}(0)$ contains only the conjugacy classes listed in $C = \{F(2), F^2(2), F^d(4), f^d(0)\}, \text{ for all } d \ge 1.$

Proof. Notice that the factor 01 only occurs as the prefix of the f-image of every letter in $F^{\omega}(0)$. Moreover, every letter 1 only occurs in $F^{\omega}(0)$ as the suffix of the factor 01. Let us say that the *index* of a conjugacy class is the number of occurrences of 1 in any of its elements. An easy computation shows that the set of complete conjugacy classes in $F^{\omega}(0)$ with index at most one is $C_1 = \{F(2), F^2(2), F(4), F^2(4), f(4), f(0)\}$. Let us assume that $F^{\omega}(0)$ contains a conjugacy class c with index at least two. Let $w \in c$ be such that 01 is a prefix of w. We write w = ps such that the leftmost occurrence of 01 in w is the prefix of s. Then the conjugate sp of w also belongs to c and thus is a factor of $F^{\omega}(0)$. This implies that the pre-image $v = f^{-1}(w)$ is a factor of $F^{\omega}(0)$, and so does every conjugate of v. Thus, $F^{\omega}(0)$ contains a conjugacy class c' such that the elements of c with prefix 01 are the f-images of the elements of c'. Moreover, the index of c' is strictly smaller than the index of c.

Using this argument recursively, we conclude that every complete conjugacy class in $F^{\omega}(0)$ has a member of the form $f^{i}(x)$ such that x is an element of a conjugacy class in C_{1} .

Now we show that F(2) does not generate larger conjugacy classes in $F^{\omega}(0)$. We thus have to exhibit a conjugate of $f(F(2)) = F^4(2) = 0120301240324$ that is not a factor of $F^{\omega}(0)$. A computer check shows that the conjugate 4012030124032 is not a factor of $F^{\omega}(0)$. Similarly, $F^2(2)$ does not generate larger conjugacy classes in $F^{\omega}(0)$ since the conjugate 301203012401203230124032of $f(F^2(2)) = F^5(2) = 012030124012032301240323$ is not a factor of $F^{\omega}(0)$.

2.2. Avoiding conjugacy classes in w_5

We are ready to prove Theorem 2.1. Notice that **ab** only appears in w_5 as the prefix of the *g*-image of every letter. A computer check shows that w_5 avoids every conjugacy class of length at most 100. Consider a word *w* with length at least 101 whose conjugacy class is complete in w_5 . So *w* contains at least two occurrences of **ab**.

Let us assume that w_5 contains a conjugacy class c. Let $w \in c$ be such that ab is a prefix of w. Since $|w| \ge 101$, w contains at least 2 occurrences of ab and we write w = ps such that the leftmost occurrence of ab in w is the prefix of s. Then the conjugate sp of w also belongs to c and thus is a factor of w_5 . This implies that the pre-image $v = g^{-1}(w)$ is a factor of $F^{\omega}(0)$, and so does every conjugate of v. Thus, $F^{\omega}(0)$ contains a conjugacy class c' such that the elements of c with prefix ab are the f-images of the elements of c'.

To finish the proof, it is thus sufficient to show that for every $c' \in C$, there exists a conjugate of g(c') that is not a factor of w_5 . Recall that $C = \{F(2), F^2(2), F^d(4), f^d(0)\}$ for all $d \ge 1$. The computer check mentioned above settles the case of F(2) and $F^2(2)$ since $|g(F(2))| < |g(F^2(2))| = 40 \le 100$.

The next four lemmas handle the remaining cases:

- $g(f^d(F(4))) = g(f^d(23))$
- $g(f^d(F^2(4))) = g(f^d(0324))$
- $g(f^{d+1}(4)) = g(f^d(01240323))$
- $g(f^{d+1}(0)) = g(f^d(01203))$

Lemma 2.3. Let $p_{23} = e.g(3f(3)...f^{d-1}(3).f^d(3))$ and $s_{23} = g(f^{d-1}(01203).f^{d-2}(01203)...f(01203)01203).abcdeacdb.$ For every $d \ge 0$, the word $T_{23} = p_{23}s_{23}$ is a conjugate of $g(f^d(23))$ that is not a factor of w_5 .

Proof. Let us assume that T_{23} appears in w_5 .

The letter 3 in $f^{\omega}(0)$ appears after either 0 or 2. However **e** is a suffix of g(2) and not of g(0). Therefore, **e**.g(3) is a suffix of g(23) only. Since 23 is a suffix of f(2) and not of f(0), then g(23f(3)) is a suffix of g(f(23)) only. Using this argument recursively, p_{23} is a suffix of $g(f^d(23))$ only.

Now, the letter 3 in $f^{\omega}(0)$ appears before either 0 or 2, however abcdeacdb is a prefix of g(2) and not of g(0). Thus g(01203).abcdeacdb is a prefix of g(012032) only. Since 012032 is a prefix of f(2) and not of f(0), then g(f(01203)012032) is a prefix of g(f(012032)) only. Using this argument recursively, s_{23} is a prefix of $g(f^{d-1}(012032))$ only. Thus T_{23} is a factor of $g(f^d(232))$ but 232 is not a factor of $f^{\omega}(0)$, contradiction.

Lemma 2.4. Let $p_{0324} = acdbecd.g(24f(24)...f^{d-1}(24)).f^d(24))$ and $s_{0324} = g(f^d(0).f^{d-1}(01240)...f(01240).01240).abcdbecde.$ For every $d \ge 0$, the word $T_{0324} = p_{0324}g(f^d(0))s_{0324}$ is a conjugate of $g(f^d(0324))$ that is not a factor of w_5 .

Proof. Let us assume that T_{0324} appears in w_5 .

The letter 2 in $f^{\omega}(0)$ appears after either 1 or 3. However acdbecd is a suffix of g(3) and not of g(1). Therefore acdbecd.g(24) is a suffix of g(324) only. Since 324 is a suffix of f(3) and not of f(1), then g(324f(24)) is a suffix of g(f(324)) only. Using this argument recursively, p_{3240} is a suffix of $g(f^d(324))$ only.

Now, the letter 0 in $f^{\omega}(0)$ appears before either 1 or 3. However abcdbecde is a prefix of g(3) and not of g(1). Thus g(01240).abcdbecde is a prefix of g(012403) only. Since 012403 is a prefix of f(3) and not of f(1), then g(f(01240)012403) is a prefix of g(f(012403)) only. Using this argument recursively, s_{3240} is a prefix of $g(f^{d-1}(012403))$ only. Thus T_{3240} is a factor of $g(f^d(32403))$ but 32403 is not a factor of $f^{\omega}(0)$, contradiction.

Lemma 2.5. Let $p_{01240323} = ecdeacdbe.g(0323f(0323) \cdots f^{d-1}(0323).f^d(0323))$ and $s_{01240323} = g(f^d(012)f^{d-1}(012).\cdots f(012)012).abcdb$. For every $d \ge 0$, the word $T_{01240323} = p_{01240323}s_{01240323}$ is a conjugate of $g(f^d(0323))$ that is not a factor of w_5 .

Proof. Let us assume that $T_{01240323}$ appears in w_5 .

The factor 03 in $f^{\omega}(0)$ appears after either 2 or 4. However ecdeacdbe is a suffix of g(4) and not of g(2). Therefore ecdeacdbe.g(0323) is a suffix of g(40323) only. Since 40323 is a suffix of f(4) and not of f(2), then g(40323f(0323)) is a suffix of g(f(40323)), using this argument recursively, $p_{01240323}$ is a suffix of $g(f^d(40323))$ only.

Now, the factor 12 in $f^{\omega}(0)$ appears before either 0 or 4. However abcdb is a prefix of g(4) and not of g(0). Thus g(012).abcdb must only be a prefix of g(0124) and since 0323 is a prefix of f(4) and not of f(0) then g(f(012)0124)is a prefix of g(f(0124)) only. Using this argument recursively, $s_{01240323}$ is a prefix of $g(f^d(0124))$ only. Thus $T_{01240323}$ is a factor of $g(f^d(403230124))$ but 403230124 is not a factor of $f^{\omega}(0)$, contradiction.

Lemma 2.6. Let $p_{01203} = d.g(3f(3) \dots f^{d-1}(3).f^d(3))$ and $s_{01203} = g(f^d(012)f^{d-1}(012).f^{d-2}(012) \dots f(012)012).abcdeac.$ For every $d \ge 0$, the word $T_{01203} = p_{01203}s_{01203}$ is a conjugate of $g(f^d(01203))$ that is not a factor of w_5 .

Proof. Let us assume that T_{01203} appears in w_5 .

The letter 3 in $f^{\omega}(0)$ appears after either 0 or 2. however d is a suffix of g(0) and not of g(2). Therefore d.g(2) is a suffix of g(12) only. Since 12 is a suffix of f(1) and not of f(3), then g(12f(2)) is a suffix of g(f(12)) only. Using this argument recursively, p_{01203} is a suffix of $g(f^d(12))$ only.

Now, 012 in $f^{\omega}(0)$ appears before either 1 or 4, however abcdeac is only a prefix of g(1) and not of g(4). Thus g(012).abcdeac is a prefix of g(0120)only. Since 0120 is a prefix of f(1) and not of f(4), then g(f(012)0120) is a prefix of g(f(0120)) only. Using this argument recursively, s_{01203} is a prefix of $g(f^d(0120))$. Thus T_{01203} is a factor of $g(f^d(030120))$ but 030120 is not a factor of $f^{\omega}(0)$, contradiction.

References

- G. Gamard, P. Ochem, G. Richomme, and P. Séébold. Avoidability of circular formulas. *Theoret. Comput. Sci* 726 (2018), 1–4.
- [2] J.P. Bell and B.W. Madill. Iterative Algebras. Algebr. Represent. Theor. 18(6) (2015), 1533–1546. https://doi.org/10.1007/s10468-015-9550-y