

Approximate Multi-Matroid Intersection via Iterative Refinement

André Linhares^{‡,1}, Neil Olver^{§,2}, Chaitanya Swamy^{‡,1}, and Rico Zenklusen^{¶,3}

¹ Dept. of Combinatorics and Optimization, Univ. Waterloo, Waterloo, Canada.
{alinhare, cswamy}@uwaterloo.ca

² Dept. of Econometrics & Operations Research, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands; CWI, Amsterdam, The Netherlands. n.olver@vu.nl

³ Department of Mathematics, ETH Zurich, Zurich, Switzerland.
ricoz@math.ethz.ch

Abstract. We introduce a new iterative rounding technique to round a point in a matroid polytope subject to further matroid constraints. This technique returns an independent set in one matroid with limited violations of the other ones. On top of the classical steps of iterative relaxation approaches, we iteratively refine/split involved matroid constraints to obtain a more restrictive constraint system, that is amenable to iterative relaxation techniques. Hence, throughout the iterations, we both tighten constraints and later relax them by dropping constraints under certain conditions. Due to the refinement step, we can deal with considerably more general constraint classes than existing iterative relaxation/rounding methods, which typically round on one matroid polytope with additional simple cardinality constraints that do not overlap too much.

We show how our rounding method, combined with an application of a matroid intersection algorithm, yields the first 2-approximation for finding a maximum-weight common independent set in 3 matroids. Moreover, our 2-approximation is LP-based, and settles the integrality gap for the natural relaxation of the problem. Prior to our work, no better upper bound than 3 was known for the integrality gap, which followed from the greedy algorithm. We also discuss various other applications of our techniques, including an extension that allows us to handle a mixture of matroid and knapsack constraints.

1 Introduction

Matroids are among the most fundamental and well-studied structures in combinatorial optimization. Recall that a *matroid* M is a pair $M = (N, \mathcal{I})$, where N is a finite ground set and $\mathcal{I} \subseteq 2^N$ is a family of sets, called *independent sets*,

[‡] Research supported by NSERC grant 327620-09 and an NSERC DAS Award.

[§] Supported by NWO VIDI grant 016.Vidi.189.087.

[¶] Supported by Swiss National Science Foundation grant 200021_165866.

such that (i) $\emptyset \in \mathcal{I}$, (ii) if $A \in \mathcal{I}$ and $B \subseteq A$, then $B \in \mathcal{I}$, and (iii) if $A, B \in \mathcal{I}$ with $|A| > |B|$, then there is an element $e \in A \setminus B$ such that $B \cup \{e\} \in \mathcal{I}$. As is common when dealing with matroids, we assume that a matroid is specified via an *independence oracle*, that, given $S \subseteq N$ as input, returns if $S \in \mathcal{I}$. Matroids capture a variety of interesting problems, and matroid-optimization algorithms provide a powerful tool in the design and analysis of efficient algorithms. A key matroid-optimization problem is the *matroid intersection* problem, wherein we seek a maximum-weight set that is independent in *two* matroids, for which various efficient algorithms are known, and we also have a celebrated min-max theorem and a polyhedral understanding of the problem. The versatility of matroid intersection comes from the fact that the intersection of matroids allows for describing a very broad family of constraints.

Unfortunately, as soon as the intersection of 3 or more matroids is considered, already the unweighted version of determining a maximum cardinality common independent set becomes *APX-hard*. Due to its fundamental nature, and many natural special cases, the problem of optimizing over 3 or more matroids has received considerable attention. In particular, there is extensive prior work ranging from the study of maximum cardinality problems [15], the maximization of submodular functions over the intersection of multiple matroids (see [8,16,17,11,4] and the references therein), to various interesting special cases like k -dimensional matching (see [13,12,3,7,6] and the references therein; many of these results apply also to the k -set packing problem which generalizes k -dimensional matching).

Nevertheless, there are still basic open questions regarding the approximability of the optimization over 3 or more matroids. Perhaps the most basic problem of this type is the *weighted 3-matroid intersection* problem, defined as follows.

Weighted 3-matroid intersection. Given matroids $M_i = (N, \mathcal{I}_i)$, for $i = 1, 2, 3$, on a common ground set N , and a weight vector $w \in \mathbb{R}^N$, solve

$$\max \{w(I) : I \in \mathcal{I}_1 \cap \mathcal{I}_2 \cap \mathcal{I}_3\},$$

where we use the shorthand $w(S) := \sum_{e \in S} w(e)$ for any set $S \subseteq N$.

The *unweighted 3-matroid intersection* problem, which is also sometimes called the *cardinality* version of 3-matroid intersection, is the special case where $w(e) = 1$ for all $e \in N$, so $w(S) = |S|$ for $S \subseteq N$.

The 3-matroid intersection problem has a natural and canonical LP-relaxation:

$$\max \{w^T x : x \in P_{\mathcal{I}_1} \cap P_{\mathcal{I}_2} \cap P_{\mathcal{I}_3}\}, \quad (\text{LP}_{3\text{-mat}})$$

where, for a matroid $M = (N, \mathcal{I})$, we denote by $P_{\mathcal{I}} \subseteq [0, 1]^N$ the matroid polytope of M , which is the convex hull of all characteristic vectors of sets in \mathcal{I} . It has a well known inequality description given by

$$P_{\mathcal{I}} = \{x \in \mathbb{R}_{\geq 0}^N : x(S) \leq r(S) \quad \forall S \subseteq N\},$$

where $r : 2^N \rightarrow \mathbb{Z}_{\geq 0}$ is the *rank function* of M , which, for $S \subseteq N$, is defined by $r(S) := \max\{|I| : I \in \mathcal{I}, I \subseteq S\}$. The rank function is submodular, and $r(S)$ can

be computed for any $S \subseteq N$ using an independence oracle. It will therefore often be convenient to assume that a matroid M is specified via its *rank oracle* that given $S \subseteq N$ as input, returns $r(S)$. In particular, one can efficiently optimize any linear function over $P_{\mathcal{I}}$ given a rank oracle (or equivalently an independence oracle). The above LP-relaxation extends naturally to the *k-matroid intersection* problem, which is the extension of 3-matroid intersection to k matroids.

Whereas $(\text{LP}_{3\text{-mat}})$, and its extension $(\text{LP}_{k\text{-mat}})$ to k -matroid intersection, are well-known LP-relaxations, there remain various gaps in our understanding of these relaxations. It is widely known that the greedy algorithm is a k -approximation for k -matroid intersection. Moreover, this approximation is relative to the optimal value of $(\text{LP}_{k\text{-mat}})$, which leads to the current-best upper bound of k on the integrality gap of $(\text{LP}_{k\text{-mat}})$, for all $k \geq 3$. However, the best lower bound on the integrality gap of $(\text{LP}_{k\text{-mat}})$ is $k - 1$ (also for all $k \geq 3$); this is known to be achievable in instances where the involved matroids are partition matroids, and for unweighted instances [9,15].

Significant progress on approximating k -matroid intersection was achieved by Lee, Sviridenko, and Vondrák [17], who presented, for any fixed $\epsilon > 0$, a local search procedure with running time exponential in ϵ that leads to a $k - 1 + \epsilon$ -approximation (i.e., the weight of the set returned is at least $(\text{optimum}) / (k - 1 + \epsilon)$). Unfortunately, apart from its high running time dependence on ϵ , this approach does not shed any insights on $(\text{LP}_{k\text{-mat}})$, as the above guarantee is not relative to $\text{OPT}_{\text{LP}_{k\text{-mat}}}$. Further progress on understanding the quality of the LP-relaxations has only been achieved in special cases. In particular, for *unweighted k-matroid intersection*, Lau, Ravi and Singh [15] give an LP-based $(k - 1)$ -approximation through iterative rounding. Their proof is based on identifying an element with “large” fractional value, picking it, and altering the fractional solution so that it remains feasible; the last step crucially uses the fact that the instance is unweighted to control the loss in the LP objective value. For the intersection of k partition matroids, a problem also known as *k-dimensional matching*, Chan and Lau [3] were able to obtain a $(k - 1)$ -approximation based on $(\text{LP}_{k\text{-mat}})$.

Although it is generally believed that a $(k - 1)$ -approximation for k -matroid intersection should exist, and that the integrality gap $(\text{LP}_{k\text{-mat}})$ is equal to the known lower bound of $k - 1$, this has remained open even for 3-matroid intersection (prior to our work). Recall that in this case, the best-known upper and lower bounds on the integrality gap of $(\text{LP}_{3\text{-mat}})$ are 3 (via the classical greedy algorithm) and 2 respectively. Moreover, the only method to beat the trivial 3-approximation of the greedy algorithm is the non-LP based and computationally quite expensive $(2 + \epsilon)$ -approximation in [17]. One main reason for the limited progress is the lack of techniques for rounding points in the intersection of multiple matroid polytopes with sufficiently strong properties. In particular, one technical difficulty that is encountered is that the tight constraints (even at an extreme point) may have large overlap, and we do not know of ways for dealing with this.

Our results. We introduce a new iterative-rounding approach to handle the above difficulties, that allows for dealing with a very general class of optimization problems involving matroids. Before delving into the details of this technique, we highlight its main implication in the context of 3-matroid intersection.

Theorem 1. *There is an LP-relative 2-approximation for weighted 3-matroid intersection. That is, for any instance, we can efficiently find a common independent set R with $w(R) \geq \text{OPT}_{\text{LP}_{3\text{-mat}}}/2$; thus, the integrality gap of $(\text{LP}_{3\text{-mat}})$ is at most 2.*

This is the *first* 2-approximation for 3-matroid intersection (with general weights). Moreover, our result *settles* the integrality gap of $(\text{LP}_{3\text{-mat}})$ due to the known matching integrality-gap lower bound of 2.

The chief new technical ingredient that leads to Theorem 1, and results for other applications discussed in Section 3, is an approximation result based on a novel *iterative refinement* technique (see Section 2) for problems of the following type. Let $N = N_0$ be a finite ground set, and $M_i = (N_i, \mathcal{I}_i)$ for $i = 0, \dots, k$ be $k+1$ matroids with rank functions $\{r_i\}$, where $N_i \subseteq N$ and $w \in \mathbb{R}^N$ be a weight vector (note that *negative* weights are allowed). We consider the problem

$$\max \{w(I) : I \in \mathcal{B}_0, \quad I \cap N_i \in \mathcal{I}_i \quad \forall i \in [k]\}, \quad (1)$$

where \mathcal{B}_0 is the set of all bases of M_0 and $[k] := \{1, \dots, k\}$. The reason we consider matroids M_i for $i \in [k]$ defined on ground sets N_i that are subsets of N , is because, as we show below, we obtain guarantees depending on how strongly the sets N_i overlap; intuitively, problem (1) becomes easier as the overlap between N_1, \dots, N_k decreases, and our guarantee improves correspondingly.

We cannot hope to solve (1) optimally, as this would enable one to solve the NP -hard k -matroid intersection problem. Our goal will be to find a basis of M_0 of large weight that is “approximately independent” in the matroids M_1, \dots, M_k .

How to quantify “approximate independence”? Perhaps the two notions that first come to mind are additive and multiplicative violation of the rank constraints. Whereas additive violations are common in the study of degree-bounded MST problems, which can be cast as special cases of (1), it turns out that such a guarantee is impossible to obtain (in polytime) for (1). More precisely, we show in Appendix A (via a replication idea) that, even for $k = 2$, if we could find in polytime a basis B of M_0 satisfying $|B| \leq r_i(B) + \alpha$ for $i = 1, 2$ for $\alpha = O(|N|^{1-\epsilon})$ for any $\epsilon > 0$, then we could efficiently find a basis of M_0 that is independent in M_1, M_2 ; the latter problem is easily seen to be NP -hard via a reduction from Hamiltonian path. We therefore consider multiplicative violation of the rank constraints. We say that $S \subseteq N$ is α -*approximately independent*, or simply α -*independent*, for a matroid $M = (N, \mathcal{I})$, if $|T| \leq \alpha \cdot r(T) \quad \forall T \subseteq S$ (equivalently, $\chi^S \in \alpha P_{\mathcal{I}}$, where χ^S is the characteristic vector of S). This is much stronger than simply requiring that $|S| \leq \alpha \cdot r(S)$, and it is easy to give examples where this weaker notion admits sets that one would consider to be quite far from being independent. An appealing feature of the stronger definition is that, using the min-max result for matroid-intersection (or via matroid partition; see, e.g., [5]),

it follows easily that if $\alpha \in \mathbb{Z}_{\geq 0}$, then S is α -independent iff S can be partitioned into at most α independent sets of M . We now state the guarantee we obtain for (1) precisely. We consider the following canonical LP-relaxation of (1):

$$\max \{w^T x : x \in \mathbb{R}_{\geq 0}^N, \quad x \in P_{\mathcal{B}_0}, \quad x|_{N_i} \in P_{\mathcal{I}_i} \quad \forall i \in [k]\}, \quad (\text{LP}_{\text{mat}})$$

where for a set $S \subseteq N$, we use $x|_S \in \mathbb{R}^S$ to denote the restriction of x to S . For ease of notation, we will sometimes write $x \in P_{\mathcal{I}_i}$ and $R \in \mathcal{I}_i$ instead of $x|_{N_i} \in P_{\mathcal{I}_i}$ and $R \cap N_i \in \mathcal{I}_i$, respectively. Our main result for (1), based on a new iterative rounding algorithm for (LP_{mat}) described in Section 2, is the following.

Theorem 2. *Let $q_1, \dots, q_k \in \mathbb{Z}_{\geq 1}$ such that*

$$\sum_{i \in [k]: e \in N_i} q_i^{-1} \leq 1 \quad \forall e \in N. \quad (2)$$

If (LP_{mat}) is feasible, then one can efficiently compute $R \subseteq N$ such that

- (i) $R \in \mathcal{B}_0$;
- (ii) $w(R) \geq \text{OPT}_{\text{LP}_{\text{mat}}}$; and
- (iii) R is q_i -independent in $M_i \quad \forall i \in [k]$.

Note that, in particular, taking $q_i = \max_{e \in N} |\{j \in [k] : e \in N_j\}|$ for all $i \in [k]$ satisfies (2). Thus, we violate the constraints imposed by the other matroids M_1, \dots, M_k by a multiplicative factor depending on how strongly the N_i s overlap.

While we have stated Theorem 2 in terms of *bases* of M_0 , the following natural variant is easily deduced from it (we defer the proof to the Appendix B).

Corollary 3. *Theorem 2 also holds when R is required only to be an independent set in M_0 (as opposed to a basis), and (LP_{mat}) is replaced by*

$$\max \{w^T x : x \in \mathbb{R}_{\geq 0}^N, \quad x|_{N_i} \in P_{\mathcal{I}_i} \quad \forall i = 0, 1, \dots, k\}. \quad (3)$$

A variety of problem settings can be handled via Theorem 2 and Corollary 3 in a unified way. We first show how to obtain a crisp, simple proof of Theorem 1.

Proof of Theorem 1. Given matroids $M_i = (N, \mathcal{I}_i)$ for $i = 0, 1, 2$, and a weight vector $w \in \mathbb{R}^N$, we first solve $(\text{LP}_{3\text{-mat}})$ to obtain an optimal solution x^* . Now we utilize Corollary 3 with the same three matroids, and $q_1 = q_2 = 2$. Clearly, these q -values satisfy (2), and x^* is a feasible solution to (3). Thus we obtain a set $A \in \mathcal{I}_0$ with $w(A) \geq w^T x^*$ and $\chi^A \in 2P_{\mathcal{I}_1} \cap 2P_{\mathcal{I}_2}$.

It is well known that $P_{\mathcal{I}_1} \cap P_{\mathcal{I}_2}$ is a polytope with integral extreme points (see, e.g., [5]). So since $\chi^A/2 \in P_{\mathcal{I}_1} \cap P_{\mathcal{I}_2}$, by using an algorithm for (weighted) matroid intersection applied to matroids M_1 and M_2 restricted to A , we can find a set $R \subseteq A$ such that $R \in \mathcal{I}_1 \cap \mathcal{I}_2$ and $w(R) \geq w^T \chi^A/2 \geq w^T x^*/2$. Finally, since $R \subseteq A$ and $A \in \mathcal{I}_0$, we also have that $R \in \mathcal{I}_0$. \square

Beyond 3-matroid intersection, Theorem 2 is applicable to various constrained (e.g., degree-bounded) spanning tree problems; we expand on this below. In Section 3, we discuss an application in this direction, wherein we seek a min-cost spanning tree satisfying matroid-independence constraints on the edge-sets of a given disjoint collection of node sets. Using Theorem 2, we obtain a spanning tree with a multiplicative factor-2 violation of the matroid constraints.

In Section 3, we also present a noteworthy extension of Theorem 2 with t *knapsack constraints* in addition to k matroid constraints, and show that we can obtain multiplicative violations of both the matroid and knapsack constraints. The only other such result we are aware of that applies to a *mixture* of matroid and knapsack constraints is by Gupta et al. [10]; their result in our setting yields an $O(kt)$ -approximation with no constraint violation, which is incomparable to our result.

Related work and connections. We note that by choosing M_0 to be the graphic matroid, problem (1) generalizes a variety of known constrained spanning tree problems. This includes degree-bounded spanning trees, and generalizations thereof considered by Bansal et al. [2], Király et al. [14], and Zenklusen [20]. Theorem 2 thus yields a unified way to deal with various spanning tree problems considered in the literature, where the soft/degree constraints are violated by at most a constant factor. However, as noted earlier, whereas the above works obtain stronger, additive-violation results, for the various constrained spanning tree problems they consider, such guarantees are not possible for our general problem (1) (see Appendix A). This hardness (of obtaining small additive violations) carries over to the spanning tree application that we consider in Section 3 (which generalizes the matroidal degree-bounded spanning tree problem considered by [20]).

To showcase how Theorem 2 can be used for such problems, consider the minimum degree-bounded spanning tree problem, where given is a graph $G = (V, E)$ with edge weights $w : E \rightarrow \mathbb{R}$ and degree bounds $B_v \in \mathbb{Z}_{\geq 1}$ for $v \in V$. The nominal problem asks to find a spanning tree $T \subseteq E$ with $|T \cap \delta(v)| \leq B_v$ for $v \in V$ minimizing $w(T)$, where $\delta(v)$ denotes the set of edges incident with v . Here one can apply Theorem 2 with M_0 being the graphic matroid of G , and for each $v \in V$ we define a uniform matroid M_v with ground set $\delta(v)$ and rank B_v . Theorem 2 with $q_v = 2 \forall v \in V$ and negated edge weights leads to a spanning tree T with $|T \cap \delta(v)| \leq 2B_v \forall v \in V$ and weight no more than the optimal LP-weight. Whereas this is a simple showcase example, Theorem 2 can be used in a similar way for considerably more general constraints than just degree constraints.

Finally, we highlight a main difference of our approach compared to prior techniques. Prior techniques for related problems, as used for example by Singh and Lau [19], Király et al. [14], and Bansal et al. [2], successively drop constraints of a relaxation. Also, interesting variations have been suggested that do not just drop constraints but may relax constraints by replacing a constraint by a weaker family (see work by Bansal et al. [1]). In contrast, our method does not just relax constraints, but also strengthens the constraint family in some iterations, so as to simplify it and enable one to drop constraints later on.

2 Our rounding technique

Our rounding technique heavily relies on a simple yet very useful “splitting” procedure for matroids, which we call *matroid refinement*.

Matroid refinement. Let $M = (N, \mathcal{I})$ be a matroid with rank function $r : 2^N \rightarrow \mathbb{Z}_{\geq 0}$, and let $S \subsetneq N$, $S \neq \emptyset$. The *refinement* of M with respect to S are the two matroids $M_1 = M|_S$ obtained by restricting M to S , and $M_2 = M/S$ obtained by contracting S in M . Formally, the independent sets of the two matroids $M_1 = (S, \mathcal{I}_1), M_2 = (N \setminus S, \mathcal{I}_2)$ are given by $\mathcal{I}_1 = \{I \subseteq S : I \in \mathcal{I}\}$, and $\mathcal{I}_2 = \{I \subseteq N \setminus S : I \cup I_S \in \mathcal{I}\}$, where $I_S \in \mathcal{I}$ is a maximum cardinality independent subset of S . It is well-known that the definition of \mathcal{I}_2 does not depend on which set I_S is chosen. The rank functions $r_1 : 2^S \rightarrow \mathbb{Z}_{\geq 0}$ and $r_2 : 2^{N \setminus S} \rightarrow \mathbb{Z}_{\geq 0}$ of M_1 and M_2 , respectively, are given by

$$r_1(A) = r(A) \quad \forall A \subseteq S, \quad \text{and} \quad r_2(B) = r(B \cup S) - r(S) \quad \forall B \subseteq N \setminus S. \quad (4)$$

We refer the reader to [18, Volume B] for more information on matroid restrictions and contractions. The following lemma describes two basic yet important relations between a matroid $M = (N, \mathcal{I})$ and its refinements $M_1 = M|_S$ and $M_2 = M/S$. These relations easily follow from well-known properties of matroids; we include the proofs in Appendix B for completeness.

Lemma 4. *Let $x \in \mathbb{R}^N$ such that $x|_S \in P_{\mathcal{I}_1}$ and $x|_{N \setminus S} \in P_{\mathcal{I}_2}$. Then $x \in P_{\mathcal{I}}$.*

Lemma 5. *Let $x \in P_{\mathcal{I}}$ such that $x(S) = r(S)$. Then $x|_S \in P_{\mathcal{I}_1}$ and $x|_{N \setminus S} \in P_{\mathcal{I}_2}$.*

Intuitively, the benefit of matroid refinement is that it serves to partly decouple the matroid independence constraints for M , thereby allowing one to work with somewhat “simpler” matroids subsequently, and we leverage this carefully in our algorithm.

An algorithm based on iterative refinement and relaxation. Algorithm 1 describes our method to prove Theorem 2. Recall that the input is an instance of problem (1), which consists of $k + 1$ matroids $M_i = (N_i, \mathcal{I}_i)$ for $i = 0, \dots, k$, where each N_i is a subset of a finite ground set $N = N_0$, and a weight vector $w \in \mathbb{R}^N$. We are also given integers $q_i \geq 1$ for $i \in [k]$ satisfying (2).

Algorithm 1 starts by solving the natural LP-relaxation in step 2 to obtain a point x^* . As is common in iterative rounding algorithms, we delete all elements of value 0 and fix all elements of value 1 through contractions in step 3. Apart from these standard operations, we *refine* the matroids in step 5, as long as there is a nontrivial x^* -tight set in some matroid in our collection. Notice that after step 5, the q -values for the matroids in the new collection \mathcal{M} continue to satisfy (2). Step 6 is our relaxation step, where we drop a matroid $M' = (N', \mathcal{I}')$ if $|N'| - x^*(N') < q_{M'}$. This is the step that results in a violation of the matroid constraints, but, as we show, the above condition ensures that even if we select

Algorithm 1: Iterative refinement/relaxation algorithm for Theorem 2

-
1. Initialize $\mathcal{M} \leftarrow \{M_1, \dots, M_k\}$, $q_{M_i} \leftarrow q_i$ for all $i \in [k]$.
 2. Compute an optimal basic solution x^* to (LP_{mat}) for the matroids $\{M_0\} \cup \mathcal{M}$.
 3. Delete all $e \in N$ with $x^*(e) = 0$ and contract all $e \in N$ with $x^*(e) = 1$ from all relevant matroids, updating also the ground set N .
 4. If $N = \emptyset$: **return** the set of all elements contracted so far.
 5. **While** there is a matroid $M' = (N', \mathcal{I}') \in \mathcal{M}$ with associated rank function r' , s.t. $\exists \emptyset \neq S \subsetneq N'$ with $x^*(S) = r'(S)$:
 (Refinement.) Set $M'_1 = M'|_S$, $M'_2 = M'/S$, and $q_{M'_1} = q_{M'_2} = q_{M'}$.
 Update $\mathcal{M} \leftarrow (\mathcal{M} \setminus \{M'\}) \cup \{M'_1, M'_2\}$.
 6. Find a matroid $M' = (N', \mathcal{I}') \in \mathcal{M}$ with associated rank function r' , such that $x^*(N') = r'(N')$ and $|N'| - x^*(N') < q_{M'}$; remove M' from \mathcal{M} . Go to step 2.
-

all elements of N' in the solution, the violation is still within the prescribed bounds.

In order to find an x^* -tight set $\emptyset \neq S \subsetneq N'$ (if one exists) in step 5, one can, for example, minimize the submodular function $r'(A) - x^*(A)$ over the sets $\emptyset \neq A \subsetneq N'$. Depending on the matroids involved, faster specialized approaches can be employed.

It is perhaps illuminating to consider the combined effect of all the refinement steps and step 6 corresponding to a given basic optimal solution x^* . Using standard uncrossing techniques, one can show that for each matroid $M' = (N', \mathcal{I}') \in \mathcal{M}$, there is a nested family of sets $\emptyset \subsetneq S_1 \subsetneq \dots \subsetneq S_p \subseteq N'$ whose rank constraints span the x^* -tight constraints of M' , and so any S_i can be used to refine M' . The combined effect of steps 5 for M' can be seen as replacing M' by the matroids $(M'|_{S_\ell})/S_{\ell-1}$ for $\ell = 1, \dots, p+1$, where $S_0 := \emptyset$, $S_{p+1} := N'$. Step 6 chooses some $M' \in \mathcal{M}$ and a “ring” $S_\ell \setminus S_{\ell-1}$ of its nested family satisfying $|S_\ell \setminus S_{\ell-1}| - x^*(S_\ell \setminus S_{\ell-1}) < q_{M'}$, and drops the matroid created for this ring.

Analysis. Lemma 6 shows that if Algorithm 1 terminates, then it returns a set with the desired properties. In Lemma 7, we show that the algorithm terminates in a polynomial number of iterations. In particular, we show that in step 6, there will always be a matroid in our collection that we can drop.

Lemma 6. *Suppose that Algorithm 1 returns a set $R \subseteq N$. Then, R satisfies the properties stated in Theorem 2.*

Proof. Note that $R \in \mathcal{B}_0$, as M_0 is only modified via deletions or contractions. Moreover, $w(R) \geq \text{OPT}$, where OPT is the optimal value of (LP_{mat}) for the input instance. Indeed, if x^* is the current optimal solution, and we update our instance (via deletions, contractions, refinements, or dropping matroids), then x^* restricted to the new ground set remains feasible for (LP_{mat}) for the new instance. This is immediate for deletions and contractions, and if we drop a matroid; it holds for refinements due to Lemma 5. So if the optimal value of (LP_{mat}) decreases, this is only because we contract elements with $x^*(e) = 1$, which we include in R . It follows that $w(R) \geq \text{OPT}$.

We now show that R is q_i -independent in M_i for all $i \in [k]$. Consider the state of the algorithm at a point during its execution right before performing step 2. Hence, the instance may already have been modified through prior refinements, contractions, deletions, and relaxations. We claim that the following invariant holds throughout the algorithm:

If R' satisfies the properties of Theorem 2 with respect to the current instance, then the set R consisting of R' and all elements contracted so far fulfills the properties of Theorem 2 with respect to the original instance.

To show the claim, it suffices to show that the invariant is preserved whenever we change the instance in the algorithm. First, one can observe that if the instance changes by deleting an element of value 0 or contracting an element of value 1, then the invariant is preserved. Next, consider step 5, where we refine $M' = (N', \mathcal{I}') \in \mathcal{M}$ to obtain $M'|_S = (S, \mathcal{I}'_1)$ and $M'/S = (N' \setminus S, \mathcal{I}'_2)$ whose q -values are set to $q_{M'}$. We are given that $\chi^{R'}|_S \in q_{M'} P_{\mathcal{I}'_1}$ and $\chi^{R'}|_{N' \setminus S} \in q_{M'} P_{\mathcal{I}'_2}$. So by Lemma 4, we have $\chi^{R'}/q_{M'} \in P_{\mathcal{I}'}$, or equivalently $\chi^{R'} \in q_{M'} P_{\mathcal{I}'}$.

Finally, consider the case where a matroid $M' = (N', \mathcal{I}') \in \mathcal{M}$ gets dropped in step 6. We need to show that $\chi^{R'}|_{N'} \in q_{M'} \cdot P_{\mathcal{I}'}$. Let x^* be the optimal solution used in the algorithm when M' was dropped. We have $|N'| - x^*(N') < q_{M'}$, and since $x^*(N') = r'(N')$, $|N'|$, and $q_{M'}$ are integral, this implies $|N'| - x^*(N') \leq q_{M'} - 1$. So N' can be partitioned into a basis of M' , which has size $r'(N') = x^*(N') \geq |N'| - (q_{M'} - 1)$, and at most $q_{M'} - 1$ other singleton sets. Each singleton $\{e\}$ is independent in M' , since $0 < x^*(e) \leq r'(\{e\})$ as $x^*|_{N'} \in P_{\mathcal{I}'}$. Therefore, N' can be partitioned into at most $q_{M'}$ independent sets of M' . Intersecting these sets with R' shows that $R' \cap N'$ can be partitioned into at most $q_{M'}$ independent sets of M' . \square

We now prove that the algorithm terminates. Note that refinements guarantee that whenever the algorithm is at step 6, then for any $M' = (N', \mathcal{I}') \in \mathcal{M}$, only the constraint of $P_{\mathcal{I}'}$ corresponding to N' may be x^* -tight. This allows us to leverage ideas similar to those in [2,14] to show that step 6 is well defined.

Lemma 7. *Algorithm 1 terminates in at most $(2k + 1)|N|$ iterations.*

Proof. We show that whenever the algorithm is at step 6, then at least one matroid in our collection can be dropped. This implies the above bound on the number of iterations as follows. There can be at most $|N|$ deletions or contractions. Each matroid $M_i = (N_i, \mathcal{I}_i)$ in our input spawns at most $|N_i|$ refinements, as each refinement of a matroid creates two matroids with disjoint (nonempty) ground sets. This also means that step 6 can be executed at most $k|N|$ times.

We focus on showing that step 6 is well defined. Consider the current collection of matroids \mathcal{M} . (Recall that \mathcal{M} does not contain the current version of M_0 .) Let x^* be the current basic solution, which is not integral; otherwise every element would have been deleted or contracted in step 3 and we would have terminated in step 4. Since we deleted all elements e with $x^*(e) = 0$, the current ground set N satisfies $N = \text{supp}(x^*)$.

Consider a full-rank subsystem of (LP_{mat}) , $Ax = b$, consisting of linearly independent, x^* -tight constraints. By standard uncrossing arguments, we may assume that the constraints of $Ax = b$ coming from a single matroid correspond to a nested family of sets. The system $Ax = b$ must contain some constraint corresponding to a matroid $M' \in \mathcal{M}$. Otherwise, we would have a full-rank system consisting of constraints coming from only one matroid, namely M_0 , which would yield a unique integral solution; but x^* is not integral. Furthermore, for a matroid $M' = (N', \mathcal{I}')$ $\in \mathcal{M}$, the only constraint of $P_{\mathcal{I}'}$ that can be x^* -tight corresponds to N' , as otherwise, M' would have been refined in step 5. So a matroid $M' \in \mathcal{M}$ gives rise to at most one row of A , which we denote by $A_{M'}$ if it exists. Let $\emptyset \subsetneq S_1 \subsetneq \dots \subsetneq S_p \subseteq N_0 = N$ denote the nested family of sets that give rise to the constraints of M_0 in our full-rank system.

Consider the following token-counting argument. Each $e \in N$ gives $x^*(e)$ tokens to the row of A corresponding to the smallest set S_ℓ containing e (if one exists). It also supplies $(1 - x^*(e))/q_{M'}$ tokens to every row $A_{M'}$ corresponding to a matroid $M' \in \mathcal{M}$ whose ground set contains e . Since the q -values satisfy (2), every $e \in N$ supplies at most one unit of token in total to the rows of A . Every row of A corresponding to a set S_ℓ receives $x^*(S_\ell) - x^*(S_{\ell-1})$ tokens, where $S_0 := \emptyset$. This is positive and integer, and thus at least 1. We claim that there is some $e \in N$ that supplies strictly less than one token unit. Given this, it must be that there is a row $A_{M'}$ corresponding to a matroid $M' = (N', \mathcal{I}') \in \mathcal{M}$ that receives less than 1 token-unit; thus $|N'| - x^*(N') < q_{M'}$ as desired.

Finally, we prove the claim. If every element supplies *exactly* one token-unit, then it must be that: (i) $S_p = N$, (ii) inequality (2) is tight for all $e \in N$, and (iii) for every $e \in N$, every matroid $M' = (N', \mathcal{I}') \in \mathcal{M}$ with $e \in N'$ gives rise to a row $A_{M'}$. But then $\sum_{M' \in \mathcal{M}} \frac{1}{q_{M'}} \cdot A_{M'} = \chi^N$, which is the row of A corresponding to the constraint of M_0 for the set S_p . This contradicts that A has full rank. \square

3 Further applications and extensions

3.1 Generalized Matroidal Degree-Bounded Spanning Trees

We introduce the generalized matroidal degree-bounded spanning tree problem (GMDST): given an undirected graph $G = (V, E)$ with edge costs $c \in \mathbb{R}^E$, disjoint node-sets S_1, \dots, S_k , and matroids $M_i = (\delta(S_i), \mathcal{I}_i) \forall i \in [k]$, we want to find a spanning tree T of minimum cost such that $T \cap \delta(S_i) \in \mathcal{I}_i \forall i \in [k]$. Here, $\delta(S_i)$ is the set of edges of G that cross S_i . This generalizes matroidal degree-bounded MSTs considered by [20], wherein each node $\{v\}$ is an S_i set.

Using Theorem 2, we obtain an algorithm for GMDST that violates the matroid constraints by a factor of at most 2. Let OPT denote the optimal value for this problem.

Theorem 8. *There exists an efficient algorithm for GMDST that computes a spanning tree $T \subseteq E$ such that $c(T) \leq 2 \cdot OPT$ and $T \cap \delta(S_i)$ is 2-independent in M_i for every $i \in [k]$.*

Proof. Let M_0 be the graphic matroid of G . We apply Theorem 2 to the matroids $\{M_i\}_{i \in \{0, \dots, k\}}$, with weight function $w := -c$, setting $q_i = 2$ for all $i \in [k]$. Note that these q -values satisfy (2), since each edge belongs to the boundary of at most two sets S_i .

Note that (LP_{mat}) is a relaxation of our instance of GMDST (with a flipped-sign objective). If (LP_{mat}) is infeasible, then so is our instance. Otherwise, we obtain a set T of edges such that $c(T) = -w(T) \leq -OPT_{\text{LP}_{\text{mat}}} \leq OPT$, and such that $T \cap \delta(S_i)$ is 2-independent in M_i for every $i \in [k]$. Further, T is a spanning tree of G since it is a basis of M_0 . \square

Whereas [20] obtains an $O(1)$ additive violation of the matroid constraints for the matroidal degree-bounded MST problem, we show in Appendix A (see Theorem 13) that, unless $P = NP$, it is impossible to obtain such an additive guarantee for GMDST. This follows from the same replication idea used to rule out small additive violations for (1).

3.2 Extension to knapsack constraints

We can consider a generalization of (1), where, in addition to the matroids M_0, \dots, M_k (over subsets of N) and the weight vector $w \in \mathbb{R}^N$, we have t knapsack constraints, indexed by $i = k+1, \dots, k+t$. The i -th knapsack constraint is specified by a ground set $N_i \subseteq N$, a cost vector $c^i \in \mathbb{R}_{\geq 0}^{N_i}$, and a budget $U_i \geq 0$. The goal is to find a maximum-weight set R such that $R \in \mathcal{B}_0 \cap \mathcal{I}_1 \cap \dots \cap \mathcal{I}_k$, and satisfying $c^i(R \cap N_i) \leq U_i$ for all $i = k+1, \dots, k+t$.

We assume without loss of generality that $\max_{e \in N_i} c_e^i \leq U_i$ for $i = k+1, \dots, k+t$. (If this inequality did not hold, then we could identify and drop some elements that do not belong to any feasible solution.)

We obtain the problem

$$\max \left\{ w(I) : I \in \mathcal{B}_0, \quad I \in \mathcal{I}_i \quad \forall i \in [k], \right. \\ \left. c^i(I \cap N_i) \leq U_i \quad \forall i = k+1, \dots, k+t \right\},$$

and its LP-relaxation

$$\max \left\{ w^T x : x \in \mathbb{R}_{\geq 0}^N, \quad x \in P_{\mathcal{B}_0}, \quad x|_{N_i} \in P_{\mathcal{I}_i} \quad \forall i \in [k], \right. \\ \left. (c^i)^T x|_{N_i} \leq U_i \quad \forall i = k+1, \dots, k+t \right\}. \quad (\text{LP}_{\text{matkn}})$$

We obtain the following generalization of Theorem 2.

Theorem 9. *Let $q_1, \dots, q_{k+t} \in \mathbb{Z}_{\geq 1}$ such that*

$$\sum_{i \in [k+t]: e \in N_i} q_i^{-1} \leq 1 \quad \forall e \in N. \quad (5)$$

If $(\text{LP}_{\text{matkn}})$ is feasible, then one can efficiently compute $R \subseteq N$ such that

- (i) $R \in \mathcal{B}_0$;
- (ii) $w(R) \geq OPT_{\text{LP}_{\text{matkn}}}$;
- (iii) R is q_i -independent in M_i for all $i \in [k]$; and
- (iv) $c^i(R \cap N_i) \leq U_i + q_i \cdot (\max_{e \in N_i} c_e^i) \leq (q_i + 1)U_i$ for all $i \in \{k+1, \dots, k+t\}$.

Again, taking $q_i = \Delta := \max_{e \in N} |\{j \in [k+t] : e \in N_j\}|$ for all $i \in [k+t]$ satisfies (5). So we can obtain Δ -independence for all the matroid constraints, and violate each knapsack constraint by at most a factor of $\Delta + 1$.

We prove Theorem 9 utilizing Algorithm 2, obtained via a small modification of Algorithm 1. The result follows from Lemmas 11 and 10.

Algorithm 2: Iterative refinement/relaxation algorithm for Theorem 9

1. Initialize $\mathcal{M} \leftarrow \{M_1, \dots, M_k\}$, $q_{M_i} \leftarrow q_i$ for all $i \in [k]$, $\mathcal{K} \leftarrow \{k+1, \dots, k+t\}$.
 2. Compute an optimal basic solution x^* to $(\text{LP}_{\text{matkn}})$ for the matroids $\{M_0\} \cup \mathcal{M}$ and the knapsack constraints indexed by \mathcal{K} .
 3. Delete all $e \in N$ with $x^*(e) = 0$ and contract all $e \in N$ with $x^*(e) = 1$ from all relevant matroids, updating also the ground set N . For every $i \in \mathcal{K}$, update $U_i \leftarrow U_i - c^i(\{e \in N_i : x^*(e) = 1\})$ and $N_i \leftarrow N_i \setminus \{e \in N_i : x^*(e) = 1\}$.
 4. If $N = \emptyset$: **return** the set of all elements contracted so far.
 5. **While** there is a matroid $M' = (N', \mathcal{I}')$ in \mathcal{M} with associated rank function r' , s.t. $\exists \emptyset \neq S \subsetneq N'$ with $x^*(S) = r'(S)$:
 (Refinement.) Set $M'_1 = M'|_S$, $M'_2 = M'/S$, and $q_{M'_1} = q_{M'_2} = q_{M'}$.
 Update $\mathcal{M} \leftarrow (\mathcal{M} \setminus \{M'\}) \cup \{M'_1, M'_2\}$.
 6. If there exists a matroid $M' = (N', \mathcal{I}')$ in \mathcal{M} with associated rank function r' , such that $x^*(N') = r'(N')$ and $|N'| - x^*(N') < q_{M'}$, then remove M' from \mathcal{M} . Otherwise, find $i \in \mathcal{K}$ such that $|N_i| - x^*(N_i) \leq q_i$; remove i from \mathcal{K} . Go to step 2.
-

Lemma 10. *Suppose that Algorithm 2 returns a set $R \subseteq N$. Then, R satisfies the properties stated in Theorem 9.*

Proof. Parts (i)–(iii) follow from the arguments used in the proof of Lemma 6. To prove part (iv), consider the i -th knapsack constraint. Note that the only place where we possibly introduce a violation in the knapsack constraint is when we drop the constraint. If x^* is the optimal solution just before we drop the constraint, then we know that $(c^i)^T x^*|_{N_i} \leq U_i$. (Note that N_i and U_i refer to the updated ground set and budget.) It follows that if S denotes the set of elements included from this residual ground set N_i , then the additive violation

in the knapsack constraint is at most

$$\begin{aligned}
 c^i(S) - U_i &\leq c^i(N_i) - U_i = (c^i)^T x^*|_{N_i} + \sum_{e \in N_i} c_e^i(1 - x_e^*) - U_i \\
 &\leq \left(\max_{e \in N_i} c_e^i\right) \sum_{e \in N_i} (1 - x_e^*) = \left(\max_{e \in N_i} c_e^i\right) (|N_i| - x^*(N_i)) \\
 &\leq q_i \cdot \left(\max_{e \in N_i} c_e^i\right) \leq q_i \cdot U_i \quad . \quad \square
 \end{aligned}$$

Lemma 11. *Algorithm 2 terminates in at most $(2k + 1)|N| + t$ iterations.*

Proof. We claim that whenever the algorithm is at step 6, there is at least one matroid constraint, or knapsack constraint that can be dropped. Assuming this, it is clear that we drop a knapsack constraint at most t times. The number of the remaining types of operations can be bounded by $(2k + 1)|N|$, as explained in the proof of Lemma 7; the bound in the lemma statement follows.

It remains to prove the claim, which follows from the token-counting argument used in the proof of Lemma 7. Recall that if $Ax = b$ is a full-rank subsystem of $(\text{LP}_{\text{matkn}})$ consisting of linearly independent x^* -tight constraints, then we may assume that the rows of A corresponding to the M_0 -constraints form a nested family \mathcal{C} . We define a token-assignment scheme, where each $e \in N$ supplies $x^*(e)$ tokens to the row of A corresponding to the smallest set in \mathcal{C} containing e (if one exists), and $(1 - x^*(e))/q_{M'}$ to each row $A_{M'}$ coming from a matroid $M' \in \mathcal{M}$ in our collection whose ground set contains e . *Additionally*, every $e \in N$ now also supplies $(1 - x^*(e))/q_i$ tokens to each row of A originating from a knapsack constraint whose ground set contains e . Under this scheme, as before, given the constraint on our q -values, it follows that every $e \in N$ supplies at most 1 token unit. Also, as before, each row of A corresponding to an M_0 constraint receives at least 1 token unit. So either there is some row $A_{M'}$ coming from a matroid in \mathcal{M} that receives strictly less than 1 token-unit, or there must be some row of A corresponding to a knapsack constraint that receives at most 1 token-unit; the latter case corresponds to a knapsack constraint i with $|N_i| - x^*(N_i) \leq q_i$. \square

Acknowledgments

We are thankful to Lap Chi Lau for pointing us to relevant literature.

References

1. N. Bansal, R. Khandekar, J. Könemann, V. Nagarajan, and B. Peis. On generalizations of network design problems with degree bounds. In *Proceedings of Integer Programming and Combinatorial Optimization (IPCO)*, pages 110–123, 2010.
2. N. Bansal, R. Khandekar, and V. Nagarajan. Additive guarantees for degree-bounded directed network design. *SIAM Journal on Computing*, 39(4):1413–1431, 2009.

3. Y. H. Chan and L. C. Lau. On linear and semidefinite programming relaxations for hypergraphic matching. *Mathematical Programming, Series A*, 135:123–148, 2012.
4. C. Chekuri, J. Vondrák, and R. Zenklusen. Submodular function maximization via the multilinear relaxation and contention resolution schemes. *SIAM Journal on Computing*, 43(6):1831–1879, 2014.
5. W. J. Cook, W. H. Cunningham, W. R. Pulleyblank, and A. Schrijver. Combinatorial optimization, 1997.
6. M. Cygan. Improved approximation for 3-dimensional matching via bounded path-width local search. In *Proceedings of 54th IEEE Symposium on Foundations of Computer Science*, pages 509–518, 2013.
7. M. Cygan, F. Grandoni, and M. Mastrolilli. How to sell hyperedges: The hypermatching assignment problem. In *Proceedings of the 24th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 342–351, 2013.
8. M. L. Fisher, G.L. Nemhauser, and L. A. Wolsey. An analysis of approximations for maximizing submodular set functions - II. *Mathematical Programming Study*, 8:73–87, 1978.
9. Z. Füredi. Maximum degree and fractional matchings in uniform hypergraphs. *Combinatorica*, 1(2):155–162, 1981.
10. A. Gupta, V. Nagarajan, and R. Ravi. Robust and maxmin optimization under matroid and knapsack uncertainty sets. *ACM Transactions on Algorithms*, 12(1):121, Nov 2015.
11. A. Gupta, A. Roth, G. Schoenebeck, and K. Talwar. Constrained non-monotone submodular maximization: offline and secretary algorithms. In *Proceedings of the 6th International Conference on Internet and Network Economics (WINE)*, pages 246–257, 2010.
12. M. M. Halldórsson. Approximating discrete collections via local improvements. In *Proceedings of the 6th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 160–169, 1995.
13. C. A. J. Hurkens and A. Schrijver. On the size of systems of sets every t of which have an SDR, with an application to the worst-case ratio of heuristics for packing problems. *SIAM Journal on Discrete Mathematics*, 2(1):68–72, 1989.
14. T. Király, L. C. Lau, and M. Singh. Degree bounded matroids and submodular flows. In *Proceedings of Integer Programming and Combinatorial Optimization (IPCO)*, pages 259–272, 2008.
15. L. C. Lau, R. Ravi, and M. Singh. *Iterative Methods in Combinatorial Optimization*. Cambridge University Press, New York, NY, USA, 1st edition, 2011.
16. J. Lee, V. Mirrokni, V. Nagarajan, and M. Sviridenko. Maximizing nonmonotone submodular functions under matroid or knapsack constraints. *SIAM Journal on Discrete Mathematics*, 23(4):2053–2078, 2010.
17. J. Lee, M. Sviridenko, and J. Vondrák. Submodular maximization over multiple matroids via generalized exchange properties. *Mathematics of Operations Research*, 35(4):795–806, 2010.
18. A. Schrijver. *Combinatorial Optimization, Polyhedra and Efficiency*. Springer, 2003.
19. M. Singh and L. C. Lau. Approximating minimum bounded degree spanning trees to within one of optimal. In *Proceedings of the 39th Annual ACM Symposium on Theory of Computing (STOC)*, pages 661–670, 2007.
20. R. Zenklusen. Matroidal degree-bounded minimum spanning trees. In *Proceedings of the 23rd Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1512–1521, 2012.

A Impossibility of achieving small additive violations

We show that Theorem 2 for problem (1) cannot be strengthened to yield a basis of M_0 that has small additive violation for the matroid constraints of M_1, \dots, M_k , even when $k = 2$.

We first define additive violation precisely. Given a matroid $M = (N, \mathcal{I})$ with rank function r , we say that a set $R \subseteq N$ is μ -additively independent in M if $|R| - r(R) \leq \mu$; equivalently, we can remove at most μ elements from R to obtain an independent set in M . Unlike results for degree-bounded spanning trees, or matroidal degree-bounded MST [20], we show that small additive violation is not possible in polytime (assuming $P \neq NP$) even for the special case of (1) where $k = 2$, so we seek a basis of M_0 that is independent in M_1, M_2 .

Theorem 12. *Let $f(n) = O(n^{1-\varepsilon})$, where $\varepsilon > 0$ is a constant. Suppose we have a polytime algorithm \mathcal{A} for (1) that returns a basis of M_0 that is $f(|N|)$ -additively independent in M_i for $i = 1, 2$. Then we can find in polytime a basis of M_0 that is independent in M_1, M_2 .*

The problem of finding a basis of M_0 that is independent in M_1, M_2 is NP -hard, as shown by an easy reduction from the directed Hamiltonian path problem. Thus, Theorem 12 shows that it is NP -hard to obtain an additive violation for problem (1) that is substantially better than linear violation.

Proof of Theorem 12. Choose t large enough so that $t > 2f(t|N|)$. Since $f(n) = O(n^{1-\varepsilon})$, this is achieved by some $t = \text{poly}(|N|)$. For each $i \in \{0, 1, 2\}$, let M'_i be the direct sum of t copies of M_i . Let N' be the ground set of these matroids, which consists of t disjoint copies of N , which we label N_1, \dots, N_t .

Clearly, the instance (M'_0, M'_1, M'_2) is feasible iff the original instance is feasible. Suppose that running \mathcal{A} on the replicated instance yields a basis R' of M'_0 that has the stated additive violation for the matroids M'_1, M'_2 . Hence, there are two sets $Q_1, Q_2 \subseteq R'$ with $|Q_1|, |Q_2| \leq f(t|N|)$, such that $R' \setminus Q_i$ is independent in M'_i for $i = 1, 2$. Hence, $R' \setminus (Q_1 \cup Q_2)$ is independent in both M'_1 and M'_2 . Because $|Q_1 \cup Q_2| \leq 2f(t|N|) < t$, we have by the pigeonhole principle that there is one $j \in [t]$ such that $(Q_1 \cup Q_2) \cap N_j = \emptyset$. This implies that $R = R' \cap N_j = (R' \setminus (Q_1 \cup Q_2)) \cap N_j$, when interpreted on the ground set N , is independent in both M_1 and M_2 . Moreover, the elements of R , when interpreted on the ground set N , are a basis in M_0 because R' is a basis in M'_0 . Hence, R is the desired basis without any violations. \square

We can mimic the above proof to show that one cannot achieve small additive violations to the matroid constraints for another special case of (1), namely, the problem GMDST introduced in Section 3: we are given $(G = (V, E), c \in \mathbb{R}^E, \{M_i = (\delta(S_i), \mathcal{I}_i)\}_{i \in [k]})$, where the S_i s are pairwise-disjoint node sets, and we seek a min-cost spanning tree T such that $T \cap \delta(S_i) \in \mathcal{I}_i$ form all $i \in [k]$. As above, we show that if we can find a spanning tree that has small additive violation for the matroids M_1, \dots, M_k , then we can find a feasible solution to GMDST. The latter is NP -hard, even when the M_i s are uniform matroids, as this

captures the degree-bounded spanning tree, and hence the Hamiltonian path problem.

This provides an alternative proof of why one cannot achieve additive guarantees for (1). (Note, however, that the hardness results from Theorems 12 and 13 are orthogonal.)

Theorem 13. *Let $f(n) = O(n^{1-\varepsilon})$, where $\varepsilon > 0$ is a constant. Suppose we have a polytime algorithm for GMDST that returns a spanning tree $T \subseteq E$ such that $T \cap \delta(S_i)$ is $f(|E|)$ -additively independent in M_i for all $i \in [k]$. Then we can find a feasible solution to GMDST in polytime.*

Proof. Let $t = \text{poly}(|E|)$ be such that $t > kf(t(|E| + 1))$. We construct a graph by taking the union of t copies of G . To connect the copies with each other, we utilize an additional vertex z , connected by an edge to all the copies of an arbitrarily chosen vertex of G . Let $G' = (V', E')$ denote the graph thus obtained, and note that $|E'| = t(|E| + 1)$. For each $i \in [k]$, let S'_i be the union of all copies of S_i . The matroid M'_i on $\delta_{G'}(S'_i)$ is the union of the direct sum of t copies of M_i , and the free matroid on the edges of $\delta_{G'}(S'_i)$ incident to z .

Note that the resulting GMDST instance is feasible iff the original GMDST instance is feasible. Furthermore, if T' is a spanning tree of G' , then T' restricted to each copy of G yields a spanning tree of G . The choice of t ensures that in some copy, the resulting tree satisfies the matroid constraints for M_1, \dots, M_k , and is therefore a feasible solution to the original GMDST instance. \square

B Omitted proofs

Proof of Corollary 3. Extend N by adding a set F of $r(N_0)$ additional elements with 0 weight, where r is the rank function of M_0 . We modify M_0 to a matroid \widehat{M}_0 on the ground set $N_0 \cup F$, given by the rank function $\widehat{r}(S) := \min\{r(S \cap N_0) + |S \cap F|, r(N_0)\}$. That is, \widehat{M}_0 is the union of M_0 with a free matroid on F , but then truncated to have rank $r(N_0)$. It is now easy to see that if $x \in \mathbb{R}^{N \cup F}$ lies in $P_{\widehat{B}_0}$, then $x|_{N_0} \in P_{\mathcal{I}_0}$. Moreover, we can extend $x \in \mathbb{R}^N$ with $x \in P_{\mathcal{I}_0}$ to $x' \in \mathbb{R}^{N \cup F}$ so that $x'|_{N_0 \cup F} \in P_{\widehat{B}_0}$ and $x'|_N = x$. The corollary thus follows by applying Theorem 2 to $\widehat{M}_0, M_1, \dots, M_k$. \square

Proof of Lemma 4. For any set $A \subseteq N$, we have

$$x(A) = x(A \cap S) + x(A \setminus S) \leq r_1(A \cap S) + r_2(A \setminus S) = r(A \cap S) + r(A \cup S) - r(S).$$

Using submodularity of r , we have $r(A \cup S) - r(S) \leq r(A) - r(A \cap S)$, so $x(A) \leq r(A)$. This holds for every $A \subseteq N$, so $x \in P_{\mathcal{I}}$. \square

Proof of Lemma 5. Let $N_1 = S$ and $N_2 = N \setminus S$. For $i \in \{1, 2\}$, to show $x|_{N_i} \in P_{\mathcal{I}_i}$ we have to verify that $x|_{N_i}$ fulfills all constraints of the matroid polytope $P_{\mathcal{I}_i} = \{y \in \mathbb{R}_{\geq 0}^{N_i} : y(Q) \leq r_i(Q) \forall Q \subseteq N_i\}$. For $i = 1$ this immediately

follows from the fact that r_1 is the restriction of r to subsets of S , and $x \in P_{\mathcal{I}}$; for any $A \subseteq S$, we have

$$x|_{N_1}(A) = x(A) \leq r(A) = r_1(A),$$

where the inequality follows from $x \in P_{\mathcal{I}}$, and the second equation from (4).

Moreover, $x|_{N_2} \in P_{\mathcal{I}_2}$ holds because for any $B \subseteq N \setminus S$, we have

$$x(B) = x(B) + x(S) - r(S) = x(B \cup S) - r(S) \leq r(B \cup S) - r(S) = r_2(B),$$

where the first equation is a consequence of $x(S) = r(S)$, the inequality is implied by $x \in P_{\mathcal{I}}$, and the last equation holds due to (4). \square