

New MDS Euclidean Self-orthogonal Codes

Xiaolei Fang Meiqing Liu Jinquan Luo*

Abstract: In this paper, a criterion of MDS Euclidean self-orthogonal codes is presented. New MDS Euclidean self-dual codes and self-orthogonal codes are constructed via this criterion. In particular, among our constructions, for large square q , about $\frac{1}{8} \cdot q$ new MDS Euclidean (almost) self-dual codes over \mathbb{F}_q can be produced. Moreover, we can construct about $\frac{1}{4} \cdot q$ new MDS Euclidean self-orthogonal codes with different even lengths n with dimension $\frac{n}{2} - 1$.

Key words: MDS code, Euclidean (almost) self-dual code, Euclidean self-orthogonal code, Generalized Reed-Solomon(GRS) code.

1 Introduction

Let q be a prime power and \mathbb{F}_q be a finite field of cardinality q . A linear code C over \mathbb{F}_q , denote by $[n, k, d]_q$, is a linear \mathbb{F}_q -subspace of \mathbb{F}_q^n with dimension k and minimal (Hamming) distance d . If the parameters can reach the Singleton bound, that is $n + 1 = k + d$, then we call C a maximum distance separable(MDS) code. Denote the Euclidean dual code of C by C^\perp . If $C \subseteq C^\perp$, we call C a Euclidean self-orthogonal code. If $C = C^\perp$, then C is called a Euclidean self-dual code.

Both MDS codes and Euclidean self-dual codes have theoretical and practical significance. The study on MDS Euclidean self-dual codes have attracted a lot of interest, especially for the constructions of MDS Euclidean self-dual codes. Such codes can be constructed in various ways, which mainly are: (1). orthogonal designs, see [3, 7, 8]; (2). building up technique, see [11, 12]; (3). constacyclic codes, see [10, 16, 18]; (4). (extended) GRS codes, see [1, 6, 9, 13, 16, 17, 19].

It is noted that the parameters of MDS Euclidean self-dual codes can be completely determined by the code length n . For the finite field of even characteristic, Grassl and Gulliver completely determined the MDS Euclidean self-dual codes of length n for all $1 \leq n \leq q$ in [4]. In [5, 10, 16], the authors obtained some new MDS Euclidean self-dual codes through cyclic, constantcyclic and negacyclic codes. In [9], Jin

The authors are with School of Mathematics and Statistics & Hubei Key Laboratory of Mathematical Sciences, Central China Normal University, Wuhan China 430079.
E-mail: fangxiaolei@mails.cnu.edu.cn(X.Fang), 15732155720@163.com(M.Liu), luojinquan@mail.cnu.edu.cn(J.Luo)

and Xing firstly presented a general and efficient method to construct MDS Euclidean self-dual codes by utilizing GRS codes. Afterwards, many researchers started to construct MDS Euclidean self-dual codes via GRS codes. In [19], Zhang and Feng stated that when $q \equiv 3 \pmod{4}$ and $n \equiv 2 \pmod{4}$, MDS Euclidean self-dual code of length n does not exist. We list all the known results on the systematic constructions of MDS Euclidean self-dual codes, which are depicted in Table 1.

Table 1: Known systematic construction on MDS Euclidean self-dual codes of length n (η is the quadratic character of \mathbb{F}_q)

q	n even	Reference
q even	$n \leq q$	[4]
q odd	$n = q + 1$	[4]
q odd	$(n-1) (q-1), \eta(1-n) = 1$	[17]
q odd	$(n-2) (q-1), \eta(2-n) = 1$	[17]
$q = r^s \equiv 3 \pmod{4}$	$n-1 = p^m (q-1)$, prime $p \equiv 3 \pmod{4}$ and m odd	[5]
$q = r^s, r \equiv 1 \pmod{4}, s$ odd	$n-1 = p^m (q-1)$, m odd and prime $p \equiv 1 \pmod{4}$	[5]
$q = r^s, r$ odd, $s \geq 2$	$n = lr, l$ even and $2l (r-1)$	[17]
$q = r^s, r$ odd, $s \geq 2$	$n = lr, l$ even, $(l-1) (r-1)$ and $\eta(1-l) = 1$	[17]
$q = r^s, r$ odd, $s \geq 2$	$n = lr + 1, l$ odd, $l (r-1)$ and $\eta(l) = 1$	[17]
$q = r^s, r$ odd, $s \geq 2$	$n = lr + 1, l$ odd, $(l-1) (r-1)$ and $\eta(l-1) = \eta(-1) = 1$	[17]
$q = r^2$	$n \leq r$	[9]
$q = r^2, r \equiv 3 \pmod{4}$	$n = 2tr$ for any $t \leq \frac{r-1}{2}$	[9]
$q = r^2, r$ odd	$n = tr, t$ even and $1 \leq t \leq r$	[17]
$q = r^2, r$ odd	$n = tr + 1, t$ odd and $1 \leq t \leq r$	[17]
$q \equiv 1 \pmod{4}$	$n (q-1), n < q-1$	[17]
$q \equiv 1 \pmod{4}$	$4^n \cdot n^2 \leq q$	[9]
$q = p^k, k$ odd prime	$n = p^r + 1, r k$	[17]
$q = p^k, k$ odd prime	$n = 2p^e, 1 \leq e < k, \eta(-1) = 1$	[17]
$q = r^2, r$ odd	$n = tm, 1 \leq t \leq \frac{r-1}{\gcd(r-1, m)}, \frac{q-1}{m}$ even	[13]
$q = r^2, r$ odd	$n = tm + 1, tm$ odd, $1 \leq t \leq \frac{r-1}{\gcd(r-1, m)}$ and $m (q-1)$	[13]
$q = r^2, r$ odd	$n = tm + 2, tm$ even, $1 \leq t \leq \frac{r-1}{\gcd(r-1, m)}$ and $m (q-1)$	[13]
$q = r^2, r$ odd	$n = tm, 1 \leq t \leq \frac{r+1}{\gcd(r+1, m)}, \frac{q-1}{m}$ even	[2]
$q = r^2, r$ odd	$n = tm + 2, tm$ even (except when t is even, m is even and $r \equiv 1 \pmod{4}$), $1 \leq t \leq \frac{r+1}{\gcd(r+1, m)}$ and $m (q-1)$	[2]
$q = r^2, r$ odd	$n = tm + 1, tm$ odd, $2 \leq t \leq \frac{r+1}{2\gcd(r+1, m)}$ and $m (q-1)$	[2]
$q = r^2, r$ odd	$n = tm, 1 \leq t \leq \frac{s(r-1)}{\gcd(s(r-1), m)}, s$ even, $s m$, $\frac{r+1}{s}$ even and $\frac{q-1}{m}$ even	[2]
$q = r^2, r$ odd	$n = tm + 2, 1 \leq t \leq \frac{s(r-1)}{\gcd(s(r-1), m)}, s$ even, $s m$, $s r+1$ and $m (q-1)$	[2]
$q = p^m, m$ even, odd prime p	$n = 2tr^l$ with $r = p^s, s \frac{m}{2}, 0 \leq l \leq \frac{m}{s}$ and $1 \leq t \leq \frac{r-1}{2}$	[1]

$q = p^m, m \text{ even, odd prime } p$	$n = (2t + 1)r^l + 1$ with $r = p^s, s \mid \frac{m}{2}, 0 \leq l < \frac{m}{s}$ and $0 \leq t \leq \frac{r-1}{2}$ or $l = \frac{m}{s}, t = 0$	[1]
$q = p^m \equiv 1 \pmod{4}$	$n = p^l + 1$ with $0 \leq l \leq m$	[1]

For large q , the number of different lengths for MDS self-dual codes in Table 1 is much smaller than $\frac{q+1}{2}$. However, in this paper, we produce approximately $\frac{q}{8}$ many q -ary MDS Euclidean self-dual codes with different lengths. In particular, for odd length, there does not exist self-dual codes. The extreme case is almost self-dual. Self-orthogonality is a nature generalization of self-duality. A criterion of MDS Euclidean self-orthogonal via (extended) GRS codes is presented and applied to construct new MDS Euclidean self-orthogonal codes in this paper.

The rest of this paper is organized as follows. In Section 2, we introduce some basic notations and useful results on self-dual codes, self-orthogonal codes and GRS codes. A criterion of MDS self-orthogonal codes is also presented. In Section 3, we give constructions of new MDS Euclidean self-dual codes utilizing GRS codes. In Section 4, some new MDS self-orthogonal codes are constructed by applying the criterion of MDS self-orthogonal codes. We conclude the results in Section 5.

2 Preliminaries

In this section, we introduce some basic notations and useful results on self-dual codes, self-orthogonal codes and GRS codes.

For two vectors $\vec{b} = (b_1, b_2, \dots, b_n)$ and $\vec{c} = (c_1, c_2, \dots, c_n)$ of \mathbb{F}_q^n , the Euclidean inner product is

$$\langle \vec{b}, \vec{c} \rangle = b_1c_1 + b_2c_2 + \dots + b_nc_n.$$

Let C be a linear code over \mathbb{F}_q . The Euclidean dual of C is defined as follows:

$$C^\perp = \{ \vec{u} \in \mathbb{F}_q^n \mid \text{for all } \vec{c} \in C, \langle \vec{u}, \vec{c} \rangle = 0 \}.$$

If $C \subseteq C^\perp$, we call C a Euclidean self-orthogonal code. If $C = C^\perp$, then C is called a Euclidean self-dual code. It is noted that if C is Euclidean self-dual, the length of C is even. For odd length, when $C \subseteq C^\perp$ and $\dim(C^\perp) = \dim(C) + 1$, we call C almost self-dual.

For $1 < n < q$, we choose two n -tuples $\vec{v} = (v_1, v_2, \dots, v_n)$ and $\vec{a} = (a_1, a_2, \dots, a_n)$, where $v_i \in \mathbb{F}_q^*$, $1 \leq i \leq n$ (v_i may not be distinct) and $a_i, 1 \leq i \leq n$ are distinct elements in \mathbb{F}_q . Then the GRS code of

length n associated with \vec{v} and \vec{a} is defined below:

$$\mathbf{GRS}_k(\vec{a}, \vec{v}) = \{(v_1 f(a_1), \dots, v_n f(a_n)) : f(x) \in \mathbb{F}_q[x], \deg(f(x)) \leq k-1\}, \quad (1)$$

where $1 \leq k \leq n$.

It is well-known that the code $\mathbf{GRS}_k(\vec{a}, \vec{v})$ is a q -ary $[n, k]$ MDS code and its dual code is also MDS [15, Chapter 11].

Moreover, the extended GRS code associated with \vec{v} and \vec{a} is defined by:

$$\mathbf{GRS}_k(\vec{a}, \vec{v}, \infty) = \{(v_1 f(a_1), \dots, v_n f(a_n), f_{k-1}) : f(x) \in \mathbb{F}_q[x], \deg(f(x)) \leq k-1\}, \quad (2)$$

where $1 \leq k \leq n$ and f_{k-1} is the coefficient of x^{k-1} in $f(x)$.

It is also well known that the code $\mathbf{GRS}_k(\vec{a}, \vec{v}, \infty)$ is a q -ary $[n+1, k]$ MDS code and its dual is also MDS [15, Chapter 11].

We define

$$L_{\vec{a}}(a_i) = \prod_{1 \leq j \leq n, j \neq i} (a_i - a_j)$$

and let QR_q be the set of nonzero squares of \mathbb{F}_q . We give the following lemmas, which are useful in the proof of the main results.

Lemma 2.1. ([14], Lemma 5) *Let a_1, a_2, \dots, a_n be distinct elements of \mathbb{F}_q . Then we have*

$$\sum_{i=1}^n a_i^m \cdot L_{\vec{a}}(a_i)^{-1} = \begin{cases} 0, & 0 \leq m \leq n-2; \\ 1, & m = n-1. \end{cases}$$

A criterion of MDS Euclidean self-orthogonal codes is presented in the following lemmas, which can be regarded as generalizations of Corollary 2.4 in [9] and Lemma 2 in [17], and be applied to construct new MDS Euclidean self-orthogonal codes.

Lemma 2.2. *Assume $1 \leq k \leq \lfloor \frac{n}{2} \rfloor$. The code $\mathbf{GRS}_k(\vec{a}, \vec{v})$ is Euclidean self-orthogonal if and only if there exists a nonzero polynomial $\lambda(x) = \lambda_{n-2k}x^{n-2k} + \dots + \lambda_1x + \lambda_0 \in \mathbb{F}_q[x]$ such that $v_i^2 = \lambda(a_i)L_{\vec{a}}(a_i)^{-1} \neq 0$ for any $1 \leq i \leq n$.*

Proof. Let $(v_1 a_1^i, v_2 a_2^i, \dots, v_n a_n^i)$ be a basis of $\mathbf{GRS}_k(\vec{a}, \vec{v})$ with $0 \leq i \leq k-1$.

$\mathbf{GRS}_k(\vec{a}, \vec{v})$ is self-orthogonal

$$\begin{aligned} &\Leftrightarrow v_1^2 a_1^m + \dots + v_n^2 a_n^m = 0, \text{ for } 0 \leq m \leq 2k-2 \\ &\Leftrightarrow \begin{pmatrix} 1 & 1 & \dots & 1 \\ a_1 & a_2 & \dots & a_n \\ \vdots & \vdots & \ddots & \vdots \\ a_1^{2k-2} & a_2^{2k-2} & \dots & a_n^{2k-2} \end{pmatrix} \begin{pmatrix} v_1^2 \\ v_2^2 \\ \vdots \\ v_n^2 \end{pmatrix} = \vec{0}. \end{aligned}$$

Denote by $v_i^2 = x_i$ with $1 \leq i \leq n$. By Lemma 2.1, the solution of

$$\begin{pmatrix} 1 & 1 & \cdots & 1 \\ a_1 & a_2 & \cdots & a_n \\ \vdots & \vdots & \ddots & \vdots \\ a_1^{n-2} & a_2^{n-2} & \cdots & a_n^{n-2} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = \vec{0}$$

is

$$\lambda \cdot \left(\frac{1}{L_{\vec{a}}(a_1)}, \frac{1}{L_{\vec{a}}(a_2)}, \dots, \frac{1}{L_{\vec{a}}(a_n)} \right)^T,$$

where $\lambda \in \mathbb{F}_q$. Let

$$A = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ a_1 & a_2 & \cdots & a_n \\ \vdots & \vdots & \ddots & \vdots \\ a_1^{2k-2} & a_2^{2k-2} & \cdots & a_n^{2k-2} \end{pmatrix} \text{ and } X = \begin{pmatrix} v_1^2 \\ v_2^2 \\ \vdots \\ v_n^2 \end{pmatrix} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}.$$

Since $\text{rank}(A) = n - 2k + 1$, the vectors

$$\left(\frac{1}{L_{\vec{a}}(a_1)}, \dots, \frac{1}{L_{\vec{a}}(a_n)} \right)^T, \left(\frac{a_1}{L_{\vec{a}}(a_1)}, \dots, \frac{a_n}{L_{\vec{a}}(a_n)} \right)^T, \dots, \left(\frac{a_1^{n-2k}}{L_{\vec{a}}(a_1)}, \dots, \frac{a_n^{n-2k}}{L_{\vec{a}}(a_n)} \right)^T \quad (3)$$

form a basis of the solution space of $AX = \vec{0}$. Therefore,

$$v_i^2 = \sum_{h=0}^{n-2k} \lambda_h a_i^h \cdot L_{\vec{a}}(a_i)^{-1} \neq 0 \text{ for any } 1 \leq i \leq n,$$

where $\lambda_h \in \mathbb{F}_q$ for $0 \leq h \leq n - 2k$. As a result, $v_i^2 = \lambda(a_i)L_{\vec{a}}(a_i)^{-1} \neq 0$ for any $1 \leq i \leq n$. \square

As a corollary of this result, Corollary 2.4 of [9] can be deduced directly with even n and $k = \frac{n}{2}$ by choosing $\lambda(x) = \lambda$ to be a constant.

Corollary 2.1. ([9], Corollary 2.4) *Let n be an even integer and $k = \frac{n}{2}$. If there exists $\lambda \in \mathbb{F}_q^*$ such that $\lambda L_{\vec{a}}(a_i)^{-1} \in QR_q$ for all $1 \leq i \leq n$, then there exists $\vec{v} = (v_1, \dots, v_n)$ with $v_i^2 = \lambda L_{\vec{a}}(a_i)^{-1}$ such that the code $\mathbf{GRS}_k(\vec{a}, \vec{v})$ defined in (1) is an MDS self-dual code of length n .*

Lemma 2.3. *Assume $1 \leq k \leq \lfloor \frac{n+1}{2} \rfloor$. The code $\mathbf{GRS}_k(\vec{a}, \vec{v}, \infty)$ is Euclidean self-orthogonal if and only if there exists a polynomial $\lambda(x) = -x^{n-2k+1} + \lambda_{n-2k}x^{n-2k} + \dots + \lambda_0 \in \mathbb{F}_q[x]$ such that $v_i^2 = \lambda(a_i)L_{\vec{a}}(a_i)^{-1} \neq 0$ for any $1 \leq i \leq n$.*

Proof. Since $\mathbf{GRS}_k(\vec{a}, \vec{v}, \infty)$ is self-orthogonal, one has

$$\begin{cases} \sum_{i=1}^n v_i^2 a_i^m = 0, & 0 \leq m \leq 2k-3; \\ \sum_{i=1}^n v_i^2 a_i^{2k-2} + 1 = 0. \end{cases} \quad (4)$$

Similarly as in Lemma 2.2, from the first equation of (4),

$$v_i^2 = \sum_{h=0}^{n-2k+1} \lambda_h a_i^h \cdot L_{\vec{a}}(a_i)^{-1} \text{ for any } 1 \leq i \leq n.$$

Substituting to the last equation of (4),

$$\lambda_{n-2k+1} \sum_{i=1}^n a_i^{n-1} \cdot L_{\vec{a}}(a_i)^{-1} + 1 = 0,$$

which implies $\lambda_{n-2k+1} = -1$ from Lemma 2.1. Hence there exists a polynomial

$$\lambda(x) = -x^{n-2k+1} + \lambda_{n-2k}x^{n-2k} + \cdots + \lambda_0 \in \mathbb{F}_q[x] \quad (5)$$

such that $v_i^2 = \lambda(a_i)L_{\vec{a}}(a_i)^{-1} \neq 0$ for any $1 \leq i \leq n$. \square

As a corollary of this result, Lemma 2 in [17] can be deduced directly with odd n and $k = \frac{n+1}{2}$ by setting $\lambda(x) = -1$.

Corollary 2.2. ([17], Lemma 2) *Let n be odd and $k = \frac{n+1}{2}$. If $-L_{\vec{a}}(a_i)^{-1} \in QR_q$ for all $1 \leq i \leq n$, then there exists $\vec{v} = (v_1, \dots, v_n)$ with $v_i^2 = -L_{\vec{a}}(a_i)^{-1}$ such that $\mathbf{GRS}_k(\vec{a}, \vec{v}, \infty)$ defined in (2) is an MDS self-dual code of length $n+1$.*

Lemma 2.4. ([17], Lemma 3) *Let $m \mid q-1$ be a positive integer and $\alpha \in \mathbb{F}_q$ be a primitive m -th root of unity. Then for any $1 \leq i \leq m$, we have*

$$\prod_{1 \leq j \leq m, j \neq i} (\alpha^i - \alpha^j) = m\alpha^{-i}.$$

3 MDS Euclidean Self-dual Codes

In this section, we present new MDS Euclidean self-dual codes from GRS codes. Throughout the remaining of this paper, we fix g as a primitive element of \mathbb{F}_q^* .

Theorem 1. *Let $q = r^2$, where r is an odd prime power. For any $1 \leq s \leq \frac{r+1}{2}$ and $1 \leq t \leq \frac{r-1}{2}$, assume that $n = s(r-1) + t(r+1)$. There exists a q -ary $[n, \frac{n}{2}]$ MDS Euclidean self-dual code, if r and s satisfy one of the following conditions:*

- (i). $r \equiv 1 \pmod{4}$ and s is even.
- (ii). $r \equiv 3 \pmod{4}$ and s is odd.

Proof. (i). Let $\alpha = g^{r+1}$, $\beta = g^{r-1}$ and $\gamma = g^{\frac{r+1}{2}}$. Denote by $\langle \alpha \rangle = \{1, \alpha, \dots, \alpha^{r-2}\}$ and $\langle \beta \rangle = \{1, \beta, \dots, \beta^r\}$ and choose

$$\vec{a} = \left(\langle \alpha \rangle, \beta \langle \alpha \rangle, \dots, \beta^{s-1} \langle \alpha \rangle, \gamma \langle \beta \rangle, \gamma^3 \langle \beta \rangle, \dots, \gamma^{2(t-1)+1} \langle \beta \rangle \right).$$

Since $r \equiv 1 \pmod{4}$, it follows that $\alpha, \beta \in QR_q$ and $\gamma \notin QR_q$. Therefore, $\beta^i \langle \alpha \rangle \cap \gamma^{2j+1} \langle \beta \rangle = \emptyset$ for any $0 \leq i \leq s-1$ and $0 \leq j \leq t-1$. It is necessary to consider two cases.

- For $0 \leq i \leq s-1$ and $0 \leq j \leq r-2$,

$$\begin{aligned} L_{\vec{a}}(\beta^i \alpha^j) &= \prod_{h=0, h \neq j}^{r-2} (\beta^i \alpha^j - \beta^i \alpha^h) \cdot \prod_{l=0, l \neq i}^{s-1} \prod_{h=0}^{r-2} (\beta^i \alpha^j - \beta^l \alpha^h) \cdot \prod_{l=0}^{t-1} \prod_{h=0}^r (\beta^i \alpha^j - \beta^h \gamma^{2l+1}) \\ &= \beta^{i(r-2)} \cdot (r-1) \cdot \alpha^{-j} \cdot \prod_{l=0, l \neq i}^{s-1} (\beta^{-2i} - \beta^{-2l}) \cdot \prod_{l=0}^{t-1} \left(\alpha^{j(r+1)} - \gamma^{(2l+1)(r+1)} \right). \end{aligned} \quad (6)$$

Here, the second equality follows from Lemma 2.4.

Note that $\beta, \alpha, r-1 \in QR_q$ and $\prod_{l=0}^{t-1} \left(\alpha^{j(r+1)} - \gamma^{(2l+1)(r+1)} \right) \in \mathbb{F}_r^* \subseteq QR_q$. We only need to consider

$u := \prod_{l=0, l \neq i}^{s-1} (\beta^{-2i} - \beta^{-2l})$. Then

$$\begin{aligned} u^r &= \prod_{l=0, l \neq i}^{s-1} (\beta^{2i} - \beta^{2l}) = \prod_{l=0, l \neq i}^{s-1} (\beta^{2(i+l)} \cdot (\beta^{-2l} - \beta^{-2i})) \\ &= \beta^{2 \left((s-1)i + \sum_{l=0, l \neq i}^{s-1} l \right)} \cdot (-1)^{s-1} \cdot \prod_{l=0, l \neq i}^{s-1} (\beta^{-2i} - \beta^{-2l}) \\ &= (-1)^{s-1} \cdot \beta^{2 \left((s-2)i + \frac{s(s-1)}{2} \right)} \cdot u. \end{aligned}$$

So

$$u^{r-1} = (-1)^{s-1} \cdot \beta^{2 \left((s-2)i + \frac{s(s-1)}{2} \right)} = g^{\frac{(r+1)(s-1)}{2} \cdot (r-1) + 2 \left((s-2)i + \frac{s(s-1)}{2} \right) \cdot (r-1)},$$

that is

$$u = g^{\frac{(r+1)(s-1)}{2} + 2 \left((s-2)i + \frac{s(s-1)}{2} \right) + m \cdot (r+1)} \quad (7)$$

with some integer m .

- For $0 \leq i \leq t-1$ and $0 \leq j \leq r$,

$$\begin{aligned}
L_{\vec{\alpha}}(\gamma^{2i+1}\beta^j) &= \prod_{h=0, h \neq j}^r (\gamma^{2i+1}\beta^j - \gamma^{2i+1}\beta^h) \cdot \prod_{l=0, l \neq i}^{t-1} \prod_{h=0}^r (\gamma^{2i+1}\beta^j - \gamma^{2l+1}\beta^h) \\
&\quad \cdot \prod_{h=0}^{s-1} \prod_{l=0}^{r-2} (\gamma^{2i+1}\beta^j - \alpha^l\beta^h) \\
&= \left(\gamma^{r(2i+1)} \cdot (r+1) \cdot \beta^{-j} \right) \cdot \prod_{l=0, l \neq i}^{t-1} \left(\gamma^{(2i+1)(r+1)} - \gamma^{(2l+1)(r+1)} \right) \\
&\quad \cdot \left((-1)^s \cdot \prod_{h=0}^{s-1} (\beta^{-2j} + \beta^{-2h}) \right). \tag{8}
\end{aligned}$$

Consider $u := \prod_{h=0}^{s-1} (\beta^{-2j} + \beta^{-2h})$. Then

$$\begin{aligned}
u^r &= \prod_{h=0}^{s-1} (\beta^{2j} + \beta^{2h}) = \prod_{h=0}^{s-1} \beta^{2(j+h)} (\beta^{-2j} + \beta^{-2h}) \\
&= \beta^{2\left(sj + \sum_{h=0}^{s-1} h\right)} \cdot \prod_{h=0}^{s-1} (\beta^{-2j} + \beta^{-2h}) \\
&= \beta^{2\left(sj + \frac{s(s-1)}{2}\right)} \cdot u.
\end{aligned}$$

Hence

$$u^{r-1} = g^{(r-1) \cdot 2\left(sj + \frac{s(s-1)}{2}\right)},$$

that is, $u = g^{2\left(sj + \frac{s(s-1)}{2}\right) + m(r+1)}$ for some integer m .

Choose $\lambda = \gamma = g^{\frac{r+1}{2}}$.

- In (6), all of $\beta, \alpha, r-1$ and $\prod_{l=0}^{t-1} (\alpha^{j(r+1)} - \gamma^{(2l+1)(r+1)})$ are squares in \mathbb{F}_q^* . From (7) and s is even,

$$\lambda u^{-1} = g^{-\left(\frac{(r+1)(s-2)}{2} + 2\left((s-2)i + \frac{s(s-1)}{2}\right) + m \cdot (r+1)\right)}$$

is a nonzero square. Thus $\lambda L_{\vec{\alpha}}(\beta^i \alpha^j)^{-1} \in QR_q$.

- In (8), it is clear that all elements except $\gamma^{r(2i+1)}$ are nonzero squares. So $\lambda L_{\vec{\alpha}}(\gamma^{2i+1}\beta^j)^{-1} \in QR_q$.

In summary, by Corollary 2.1, there exists a q -ary $\left[n, \frac{n}{2}\right]$ MDS Euclidean self-dual code over \mathbb{F}_q .

(ii). Since the proof of case (ii) is similar to the proof of case (i), we omit some details. Let $\alpha = g^{r+1}$, $\beta = g^{r-1}$ and $\gamma = g^{\frac{r-1}{2}}$. Choose

$$\vec{\alpha} = \left(\langle \beta \rangle, \alpha \langle \beta \rangle, \dots, \alpha^{t-1} \langle \beta \rangle, \gamma \langle \alpha \rangle, \gamma^3 \langle \alpha \rangle, \dots, \gamma^{2(s-1)+1} \langle \alpha \rangle \right).$$

Since $r \equiv 3 \pmod{4}$, it follows that $\alpha, \beta \in QR_q$ and $\gamma \notin QR_q$. Therefore, $\alpha^i \langle \beta \rangle \cap \gamma^{2j+1} \langle \alpha \rangle = \emptyset$ for any $0 \leq i \leq t-1$ and $0 \leq j \leq s-1$. We also consider two cases.

- For $0 \leq i \leq t-1$ and $0 \leq j \leq r$,

$$\begin{aligned} L_{\vec{\alpha}}(\alpha^i \beta^j) &= \prod_{h=0, h \neq j}^r (\alpha^i \beta^j - \alpha^i \beta^h) \cdot \prod_{l=0, l \neq i}^{t-1} \prod_{h=0}^r (\alpha^i \beta^j - \alpha^l \beta^h) \cdot \prod_{h=0}^{s-1} \prod_{l=0}^{r-2} (\alpha^i \beta^j - \alpha^l \gamma^{2h+1}) \\ &= \alpha^{ir} \cdot (r+1) \cdot \beta^{-j} \cdot \prod_{l=0, l \neq i}^{t-1} (\alpha^{i(r+1)} - \alpha^{l(r+1)}) \cdot \prod_{h=0}^{s-1} (\beta^{j(r-1)} - \gamma^{(2h+1)(r-1)}). \end{aligned} \quad (9)$$

Since $\beta, \alpha, r+1 \in QR_q$ and $\prod_{l=0, l \neq i}^{t-1} (\alpha^{i(r+1)} - \alpha^{l(r+1)}) \in \mathbb{F}_r^* \subseteq QR_q$, we only need to consider

$u := \prod_{h=0}^{s-1} (\beta^{j(r-1)} - \gamma^{(2h+1)(r-1)})$. Let $\xi = \gamma^{r-1}$. Then $u = \prod_{h=0}^{s-1} (\xi^{2j} - \xi^{2h+1})$. Therefore,

$$u^r = (-1)^s \cdot \xi^{-2sj-s^2} \cdot u.$$

It implies

$$u^{r-1} = (-1)^s \cdot \xi^{-2sj-s^2} = g^{\frac{(r+1)s}{2} \cdot (r-1) + (-2sj-s^2) \cdot \frac{(r-1)^2}{2}},$$

that is,

$$u = g^{\frac{(r+1)s}{2} - (s^2+2sj) \cdot \frac{r-1}{2} + m \cdot (r+1)} \quad (10)$$

for some integer m .

- For $0 \leq i \leq s-1$ and $0 \leq j \leq r-2$,

$$\begin{aligned} L_{\vec{\alpha}}(\gamma^{2i+1} \alpha^j) &= \prod_{h=0, h \neq j}^{r-2} (\gamma^{2i+1} \alpha^j - \gamma^{2i+1} \alpha^h) \cdot \prod_{l=0, l \neq i}^{s-1} \prod_{h=0}^{r-2} (\gamma^{2i+1} \alpha^j - \gamma^{2l+1} \alpha^h) \\ &\quad \cdot \prod_{h=0}^{t-1} \prod_{l=0}^r (\gamma^{2i+1} \alpha^j - \beta^l \alpha^h) \\ &= (\gamma^{(r-2)(2i+1)} \cdot (r-1) \cdot \alpha^{-j}) \cdot \prod_{l=0, l \neq i}^{s-1} (\gamma^{(2i+1)(r-1)} - \gamma^{(2l+1)(r-1)}) \\ &\quad \cdot \prod_{h=0}^{t-1} (-\alpha^{j(r+1)} - \alpha^{h(r+1)}). \end{aligned} \quad (11)$$

Let $u = \prod_{l=0, l \neq i}^{s-1} (\gamma^{(2i+1)(r-1)} - \gamma^{(2l+1)(r-1)})$ and $\xi = \gamma^{r-1} = g^{\frac{(r-1)^2}{2}}$. Then $u = \prod_{l=0, l \neq i}^{s-1} (\xi^{2i+1} - \xi^{2l+1})$.

Therefore,

$$u^r = (-1)^{s-1} \cdot \xi^{-(s-2)(2i+1)-s^2} \cdot u.$$

It follows that

$$u^{r-1} = g^{(r-1) \cdot \left(\frac{(r+1)(s-1)}{2} - \frac{r-1}{2} \cdot ((s-2)(2i+1) + s^2) \right)},$$

that is,

$$u = g^{\frac{(r+1)(s-1)}{2} - \frac{r-1}{2} \cdot ((s-2)(2i+1) + s^2) + m(r+1)}$$

with some integer m . Since $r \equiv 3 \pmod{4}$ and s is odd, it is easy to verify that

$$\frac{(r+1)(s-1)}{2} - \frac{r-1}{2} \cdot ((s-2)(2i+1) + s^2) + m(r+1)$$

is even, which yields $u \in QR_q$.

Choose $\lambda = \gamma = g^{\frac{r-1}{2}}$.

- In (9), all of $\beta, \alpha, r+1$ and $\prod_{l=0, l \neq i}^{t-1} (\alpha^{i(r+1)} - \alpha^{l(r+1)})$ are nonzero squares in \mathbb{F}_q . From (10), $r \equiv 3 \pmod{4}$ and s is odd,

$$\lambda u^{-1} = g^{-\left(\frac{(r+1)s}{2} - (s^2+1+2sj) \cdot \frac{r-1}{2} + m \cdot (r+1) \right)}$$

is a nonzero square, which yields $\lambda L_{\vec{\alpha}}(\alpha^i \beta^j)^{-1} \in QR_q$.

- In (11), from the above analysis, it is clear that all factors except $\gamma^{(r-2)(2i+1)}$ are nonzero squares. As a consequence, $\lambda L_{\vec{\alpha}}(\gamma^{2i+1} \alpha^j)^{-1} \in QR_q$.

In short, by Corollary 2.1, there exists a q -ary $\left[n, \frac{n}{2} \right]$ MDS Euclidean self-dual code over \mathbb{F}_q . \square

Remark 3.1. In (i) (resp. (ii)) of Theorem 1, the number of s is $\frac{r-1}{4}$ (resp. $\frac{r+1}{4}$) and the number of t is $\frac{r-1}{2}$. Thus the number of MDS Euclidean self-dual codes is $\frac{r-1}{4} \times \frac{r-1}{2} = \frac{1}{8} \cdot (r-1)^2 \approx \frac{1}{8}q$ (resp. $\frac{r+1}{4} \times \frac{r-1}{2} = \frac{1}{8} \cdot (q-1) \approx \frac{1}{8}q$).

Example 3.1. By utilizing all the previous results in Table 1, 1320 many MDS Euclidean self-dual codes over \mathbb{F}_{149^2} of different lengths can be constructed. However, by Theorem 1, there are $\frac{1}{8} \cdot (\sqrt{q}-1)^2 = 2738$ MDS Euclidean self-dual codes of different lengths.

Example 3.2. For $q = 151^2$, by Theorem 1, much more MDS Euclidean self-dual codes can be constructed than all previous works. The comparison can be presented in the following table.

Table 2: Comparison of MDS Euclidean self-dual codes of length n over \mathbb{F}_{151^2}

Source	the number of different lengths n
All previous constructions	1500
Theorem 1	2850

Remark 3.2. For all the previous results, the number of possible lengths in most cases is bounded by r or some divisor of $r - 1$, $r + 1$ or $q - 1$. Therefore, it is expected that when q is large, the number of possible lengths is much less than $c \cdot q$ for some constant c . However, Theorem 1 will produce about $q/8$ MDS Euclidean self-dual codes of different lengths which contribute $\frac{q/8}{q/2} = 25\%$ to all possible lengths for MDS Euclidean self-dual codes.

4 MDS Euclidean Self-orthogonal Codes

In this section, we apply the criterion (Lemma 2.2) to construct new MDS Euclidean self-orthogonal codes.

Theorem 2. Let $q = r^2$, where r is an odd prime power. For any $1 \leq s \leq \frac{r+1}{2}$ and $1 \leq t \leq \frac{r-1}{2}$, assume $n = s(r-1) + t(r+1)$ and $1 \leq k \leq \frac{n}{2} - 1$. Then there exists a q -ary $[n, k]$ MDS Euclidean self-orthogonal code.

Proof. **Case 1:** $r \equiv 1 \pmod{4}$. Recall α , β and γ in the proof of Theorem 1(i). We also choose

$$\vec{a} = (\langle \alpha \rangle, \beta \langle \alpha \rangle, \dots, \beta^{s-1} \langle \alpha \rangle, \gamma \langle \beta \rangle, \gamma^3 \langle \beta \rangle, \dots, \gamma^{2(t-1)+1} \langle \beta \rangle).$$

By the proof of Theorem 1(i),

$$L_{\vec{a}}(\beta^i \alpha^j) = \beta^{i(r-2)} \cdot (r-1) \cdot \alpha^{-j} \cdot g^{\frac{(r+1)(s-1)}{2} + 2((s-2)i + \frac{s(s-1)}{2}) + k \cdot (r+1)} \cdot \prod_{l=0}^{t-1} (\alpha^{j(r+1)} - \gamma^{(2l+1)(r+1)})$$

and

$$\begin{aligned} L_{\vec{a}}(\gamma^{2i+1} \beta^j) &= (\gamma^{r(2i+1)} \cdot (r+1) \cdot \beta^{-j}) \cdot \prod_{l=0, l \neq i}^{t-1} (\gamma^{(2i+1)(r+1)} - \gamma^{(2l+1)(r+1)}) \\ &\quad \cdot \left((-1)^s \cdot g^{2(sj + \frac{s(s-1)}{2}) + m(r+1)} \right). \end{aligned}$$

- When s is even, choose $\lambda(x) = \gamma$. Then

$$\lambda(\beta^i \alpha^j) L_{\vec{a}}(\beta^i \alpha^j)^{-1}, \lambda(\gamma^{2i+1} \beta^j) L_{\vec{a}}(\gamma^{2i+1} \beta^j)^{-1} \in QR_q.$$

- When s is odd, choose $\lambda(x) = x$. Then

$$\lambda(\beta^i \alpha^j) L_{\vec{a}}(\beta^i \alpha^j)^{-1}, \lambda(\gamma^{2i+1} \beta^j) L_{\vec{a}}(\gamma^{2i+1} \beta^j)^{-1} \in QR_q.$$

By Lemma 2.2, it follows that there exists a q -ary $[n, k]$ MDS Euclidean self-orthogonal code.

Case 2: $r \equiv 3 \pmod{4}$. Recall α, β and γ in Theorem 1(ii). We choose

$$\vec{a} = \left(\langle \beta \rangle, \alpha \langle \beta \rangle, \dots, \alpha^{t-1} \langle \beta \rangle, \gamma \langle \alpha \rangle, \gamma^3 \langle \alpha \rangle, \dots, \gamma^{2(s-1)+1} \langle \alpha \rangle \right)$$

Utilizing the proof of Theorem 1(ii),

$$L_{\vec{a}}(\alpha^i \beta^j) = \alpha^{ir} \cdot (r+1) \cdot \beta^{-j} \cdot \prod_{l=0, l \neq i}^{t-1} (\alpha^{i(r+1)} - \alpha^{l(r+1)}) \cdot g^{\frac{(r+1)s}{2} - (s^2 + 2sj) \cdot \frac{r-1}{2} + m \cdot (r+1)}$$

and

$$L_{\vec{a}}(\gamma^{2i+1} \alpha^j) = \left(\gamma^{(r-2)(2i+1)} \cdot (r-1) \cdot \alpha^{-j} \right) \cdot \prod_{h=0}^{t-1} \left(-\alpha^{j(r+1)} - \alpha^{h(r+1)} \right) \cdot g^{\frac{(r+1)(s-1)}{2} - \frac{r-1}{2} \cdot ((s-2)(2i+1) + s^2) + m(r+1)}.$$

- When s is odd, choose $\lambda(x) = \gamma$. Then

$$\lambda(\beta^i \alpha^j) L_{\vec{a}}(\beta^i \alpha^j)^{-1}, \lambda(\gamma^{2i+1} \beta^j) L_{\vec{a}}(\gamma^{2i+1} \beta^j)^{-1} \in QR_q.$$

- When s is even, choose $\lambda(x) = x$. Then

$$\lambda(\beta^i \alpha^j) L_{\vec{a}}(\beta^i \alpha^j)^{-1}, \lambda(\gamma^{2i+1} \beta^j) L_{\vec{a}}(\gamma^{2i+1} \beta^j)^{-1} \in QR_q.$$

By Lemma 2.2, it follows that there exists a q -ary $[n, k]$ MDS Euclidean self-orthogonal code. \square

Remark 4.1. In Theorem 2, the conditions that s is even in Theorem 1(i) and s is odd in Theorem 1(ii) can be removed. So we can construct approximately $\frac{1}{4} \cdot q$ MDS Euclidean self-orthogonal codes C with length n and dimension $\frac{n}{2} - 1$. which means $C \subseteq C^\perp$ and $\dim(C^\perp) = \dim(C) + 2$.

For odd length, Euclidean almost self-dual codes can be constructed via Lemma 2.2.

Theorem 3. Let $q = r^2$, where r is an odd prime power. For any $1 \leq s \leq \frac{r+1}{2}$ and $1 \leq t \leq \frac{r-1}{2}$, assume that $n = s(r-1) + t(r+1) + 1$. Then there exists a q -ary $[n, \frac{n-1}{2}]$ MDS Euclidean almost self-dual code, provided that r and s satisfy one of the following conditions:

- (i). $r \equiv 1 \pmod{4}$ and s is odd.
- (ii). $r \equiv 3 \pmod{4}$ and s is even.

Proof. (i). Recall α, β, γ and g in Theorem 1(i). Choose

$$\vec{a} = \left(\langle \alpha \rangle, \beta \langle \alpha \rangle, \dots, \beta^{s-1} \langle \alpha \rangle, \gamma \langle \beta \rangle, \gamma^3 \langle \beta \rangle, \dots, \gamma^{2(t-1)+1} \langle \beta \rangle, 0 \right).$$

Similarly as the proof of Theorem 1(i),

$$L_{\vec{\alpha}}(\beta^i \alpha^j) = \beta^{i(r-1)} \cdot (r-1) \cdot g^{\frac{(r+1)(s-1)}{2} + 2((s-2)i + \frac{s(s-1)}{2}) + m \cdot (r+1)} \cdot \prod_{l=0}^{t-1} \left(\alpha^{j(r+1)} - \gamma^{(2l+1)(r+1)} \right),$$

and

$$\begin{aligned} L_{\vec{\alpha}}(\gamma^{2i+1} \beta^j) &= \left(\gamma^{(r+1)(2i+1)} \cdot (r+1) \right) \cdot \prod_{l=0, l \neq i}^{t-1} \left(\gamma^{(2i+1)(r+1)} - \gamma^{(2l+1)(r+1)} \right) \\ &\quad \cdot \left((-1)^s \cdot g^{2(sj + \frac{s(s-1)}{2}) + m(r+1)} \right) \end{aligned}$$

A routine calculation shows that

$$\begin{aligned} L_{\vec{\alpha}}(0) &= \prod_{l=0}^{s-1} \prod_{h=0}^{r-2} (0 - \beta^l \alpha^h) \cdot \prod_{l=0}^{t-1} \prod_{h=0}^r (0 - \beta^h \gamma^{2l+1}) \\ &= \alpha^{\frac{(r-2)(r-1)s}{2}} \cdot \beta^{\frac{(r-1)(s-1)s + r(r+1)t}{2}} \cdot \gamma^{(r+1)t^2}. \end{aligned}$$

When s is odd, it is easy to see

$$L_{\vec{\alpha}}(\beta^i \alpha^j)^{-1}, L_{\vec{\alpha}}(\gamma^{2i+1} \beta^j)^{-1}, L_{\vec{\alpha}}(0)^{-1} \in QR_q.$$

By Lemma 2.2, it follows that there exists a q -ary $[n, \frac{n-1}{2}]$ MDS Euclidean almost self-dual code.

(ii). Recall α, β, γ and g in Theorem 1(ii). Choose

$$\vec{\alpha} = \left(\langle \beta \rangle, \alpha \langle \beta \rangle, \dots, \alpha^{t-1} \langle \beta \rangle, \gamma \langle \alpha \rangle, \gamma^3 \langle \alpha \rangle, \dots, \gamma^{2(s-1)+1} \langle \alpha \rangle, 0 \right)$$

Similarly as the proof of Theorem 1(ii),

$$L_{\vec{\alpha}}(\alpha^i \beta^j) = \alpha^{i(r+1)} \cdot (r+1) \cdot \prod_{l=0, l \neq i}^{t-1} \left(\alpha^{i(r+1)} - \alpha^{l(r+1)} \right) \cdot g^{\frac{(r+1)s}{2} - (s^2 + 2sj) \cdot \frac{r-1}{2} + m \cdot (r+1)}$$

and

$$\begin{aligned} L_{\vec{\alpha}}(\gamma^{2i+1} \alpha^j) &= \left(\gamma^{(r-1)(2i+1)} \cdot (r-1) \right) \cdot \prod_{l=0, l \neq i}^{s-1} \left(\gamma^{(2i+1)(r-1)} - \gamma^{(2l+1)(r-1)} \right) \\ &\quad \cdot g^{\frac{(r+1)(s-1)}{2} - \frac{r-1}{2} \cdot ((s-2)(2i+1) + s^2) + m(r+1)}. \end{aligned}$$

Meanwhile,

$$\begin{aligned} L_{\vec{\alpha}}(0) &= \prod_{l=0}^{t-1} \prod_{h=0}^r (0 - \alpha^l \beta^h) \cdot \prod_{l=0}^{s-1} \prod_{h=0}^{r-2} (0 - \gamma^{2l+1} \alpha^h) \\ &= \alpha^{\frac{(r+1)(t-1)t + (r-2)(r-1)s}{2}} \cdot \beta^{\frac{r(r+1)t}{2}} \cdot \gamma^{(r-1)s^2}. \end{aligned}$$

When s is even, it is obvious that

$$L_{\vec{\alpha}}(\alpha^i \beta^j)^{-1}, L_{\vec{\alpha}}(\gamma^{2i+1} \alpha^j)^{-1}, L_{\vec{\alpha}}(0)^{-1} \in QR_q.$$

By Lemma 2.2, it follows that there exists a q -ary $[n, \frac{n-1}{2}]$ MDS Euclidean almost self-dual code. \square

Remark 4.2. In Theorem 3, the number of s is about $\frac{r}{4}$ and the number of t is $\frac{r-1}{2}$. Therefore, there exist about $\frac{1}{8} \cdot q$ MDS Euclidean almost self-dual codes with different lengths.

5 Conclusion

We propose a criterion of Euclidean self-orthogonal GRS codes. Based on the criterion, we construct new MDS Euclidean (almost) self-dual codes and new MDS Euclidean self-orthogonal codes by utilizing GRS codes. For any large square q , about $\frac{1}{8} \cdot q$ new MDS Euclidean (almost) self-dual codes with different lengths can be produced. Our results contribute about 25% possible lengths. But for nonsquare odd prime power q , the possible lengths for MDS Euclidean self-dual codes are very restricted and there are still a lot of cases to be explored.

References

- [1] W. Fang and F. Fu, “New constructions of MDS Euclidean self-dual codes from GRS codes and extended GRS codes,” *IEEE Trans. Inf. Theory*, vol. 65, no. 9, pp. 5574–5579, 2019.
- [2] X. Fang, K. Lebed, H. Liu, and J. Luo, “New MDS Euclidean self-dual codes over finite fields of odd characteristic,” arXiv:1811.02802v9 [cs.IT], Sep. 2019.
- [3] S. Georgion and C. Koukouvinos, “MDS Euclidean self-dual codes over large prime fields,” *Finite Fields and Their Appl.*, vol. 8, no. 4, pp. 455-470, 2002.
- [4] M. Grassl and T. A. Gulliver, “On self-dual MDS codes,” in *Proc. of ISIT*, pp. 1954-1957, 2008.
- [5] K. Guenda, “New MDS Euclidean self-dual codes over finite fields,” *Des. Codes Cryptogr.*, vol. 62, no. 1, pp. 31-42, 2012.
- [6] T. A. Gulliver, J. L. Kim, and Y. Lee, “New MDS or near-MDS Euclidean self-dual codes,” *IEEE Trans. Inf. Theory*, vol. 54, no. 9, pp. 4354-4360, 2008.
- [7] M. Harada and H. Kharaghani, “Orthogonal designs, self-dual codes and the Leech lattice,” *J. Combin. Designs*, vol. 13, no. 3, pp. 184-194, 2005.
- [8] M. Harada and H. Kharaghani, “Orthogonal designs and MDS Euclidean self-dual codes,” *Australas. J. Combin.*, vol. 35, pp. 57-67, 2006.

- [9] L. Jin and C. Xing, "New MDS self-dual codes from generalized Reed-Solomon codes," *IEEE Trans. Inf. Theory*, vol. 63, no. 3, pp. 1434-1438, 2017.
- [10] X. Kai, S. Zhu S, and Y. Tang, "Some constacyclic self-dual codes over the integers modulo 2^m ," *Finite Fields and Their Appl.* vol. 18, no. 2, pp. 258-270, 2012.
- [11] J. L. Kim and Y. Lee, "MDS Euclidean self-dual codes," in *Proc. of ISIT*, pp. 1872-1877, 2004.
- [12] J. L. Kim and Y. Lee, "Euclidean and Hermitian self-dual MDS codes over large finite fields," *J. Combin. Theory, Series A*, vol. 105, no. 1, pp. 79-95, 2004.
- [13] K. Lebed, H. Liu, and J. Luo, "Construction of MDS Euclidean self-dual codes over finite field," *Finite Fields and Their Appl.*, vol. 59, pp. 199-207, 2019.
- [14] Z. Li, L. Xing, and X. Wang, "Quantum generalized Reed-Solomon codes: unified framework for quantum MDS codes," *Phys. Rev. A* , vol. 77, no. 1, pp. 012308-1-012308-4, 2008.
- [15] F. J. MacWilliams and N. J. A. Sloane, *The Theory of Error-correcting Codes*. The Netherlands: North Holland, Amsterdam, 1977.
- [16] H. Tong and X. Wang, "New MDS Euclidean and Hermitian self-dual codes over finite fields," *Advances in Pure Mathematics.*, vol. 7, no. 5, 2016.
- [17] H. Yan, "A note on the construction of MDS Euclidean self-dual codes," *Cryptogr. Commun.*, vol. 11, no. 2, pp. 259-268, 2019.
- [18] Y. Yang and W. Cai, "On self-dual constacyclic codes over finite fields," *Des. Codes Cryptogr.*, vol. 74, no. 2, pp. 355-364, 2015.
- [19] A. Zhang and K. Feng, "An unified approach on constructing of MDS self-dual codes via Reed-Solomon codes," arXiv: 1905.06513v1 [cs.IT], May 2019.