Reducing Path TSP to TSP

Vera Traub¹

Jens Vygen¹

Rico Zenklusen²

July 24, 2019

Abstract

We present a black-box reduction from the path version of the Traveling Salesman Problem (Path TSP) to the classical tour version (TSP). More precisely, we show that given an α -approximation algorithm for TSP, then, for any $\varepsilon > 0$, there is an $(\alpha + \varepsilon)$ -approximation algorithm for the more general Path TSP. This reduction implies that the approximability of Path TSP is the same as for TSP, up to an arbitrarily small error. This avoids future discrepancies between the best known approximation factors achievable for these two problems, as they have existed until very recently.

A well-studied special case of TSP, Graph TSP, asks for tours in unit-weight graphs. Our reduction shows that any α -approximation algorithm for Graph TSP implies an $(\alpha + \varepsilon)$ -approximation algorithm for its path version. By applying our reduction to the 1.4-approximation algorithm for Graph TSP by Sebő and Vygen, we obtain a polynomial-time $(1.4 + \varepsilon)$ -approximation algorithm for Graph Path TSP, improving on a recent 1.497-approximation algorithm of Traub and Vygen.

We obtain our results through a variety of new techniques, including a novel way to set up a recursive dynamic program to guess significant parts of an optimal solution. At the core of our dynamic program we deal with instances of a new generalization of (Path) TSP which combines parity constraints with certain connectivity requirements. This problem, which we call Φ -TSP, has a constant-factor approximation algorithm and can be reduced to TSP in certain cases when the dynamic program would not make sufficient progress.

¹Research Institute for Discrete Mathematics and Hausdorff Center for Mathematics, University of Bonn, Bonn, Germany. Email: traub@or.uni-bonn.de, vygen@or.uni-bonn.de. This research was initiated while the first two authors visited FIM at ETH Zurich.

²Department of Mathematics, ETH Zurich, Zurich, Switzerland. Email: ricoz@math.ethz.ch. Research supported in part by the Swiss National Science Foundation grants 200021_184622 and 200021_165866.

1 Introduction

The Traveling Salesman Problem (TSP) is one of the most fundamental and well-studied problems in Combinatorial Optimization with a multitude of applications. The common denominator of the numerous variants of the problem is that a set of cities have to be visited on a shortest possible tour. Its best-known variant, often just dubbed TSP, assumes that the distances between the cities are non-negative and symmetric, and the task is to find a tour beginning and ending in the same city. For our purposes it will be useful to work with an undirected graph G = (V, E) with edge lengths $\ell : E \to \mathbb{R}_{\geq 0}$. While it is often assumed that G is complete and ℓ fulfills the triangle-inequality, we do not assume this, but allow the tour to visit cities more than once; this is easily seen (and well-known) to be equivalent. So a tour is a closed walk in G visiting all vertices.

One of the best-studied extensions of TSP is Path TSP, where in addition to G and ℓ , a fixed start $s \in V$ and end $t \in V$ are given and the task is to find a shortest walk from s to t visiting all vertices.

TSP and its variants are well-known to be APX-hard (see [20, 14] and references therein) and they have been studied very extensively under the viewpoint of approximation algorithms. While TSP and Path TSP look quite similar, there are fundamental differences. First, there is a classical 3/2-approximation algorithm for TSP by Christofides [4] and Serdjukov [25]. This algorithm can easily be adapted to Path TSP, but then has only approximation ratio 5/3 as Hoogeveen [12] showed in the early 90's. Second, the integrality gaps of the classical LP relaxations seem to be different. For TSP it is widely believed to be 4/3, while for Path TSP it is at least 3/2. In both cases there are well-known instances attaining these lower bounds on the integrality gaps, and these instances have unit lengths ($\ell(e) = 1$ for all $e \in E$). Therefore the unit-length special cases, Graph TSP and Graph Path TSP, have received considerable attention [1, 19, 16, 17, 24, 8, 26]. In these special cases, the integrality gaps are known to be different: it is at most 7/5 for Graph TSP and exactly 3/2for Graph Path TSP as shown by Sebő and Vygen [24].

While Christofides' algorithm for TSP is still unbeaten after more than four decades, the approximation ratio for Path TSP has been improved. The first improvement, about 20 years after Hoogeveen [12], was obtained by An, Kleinberg, and Shmoys [1], who devised an elegant $(1 + \sqrt{5})/2 \approx 1.618$ -approximation algorithm. A sequence of successive improvements [22, 28, 10, 23, 27] culminated in Zenklusen's recent $^{3}/_{2}$ -approximation algorithm [30]. Hence, at the moment, the best known approximation ratios for TSP and Path TSP are the same.

For Graph TSP and Graph Path TSP the situation is different. The best known approximation ratios are $^{7/5}$ for Graph TSP [24] and 1.497 for Graph Path TSP [26]. Since the latter result achieves an approximation ratio better than the integrality gap $^{3/2}$, one might hope that Graph Path TSP is actually no harder than Graph TSP although the integrality gaps differ.

These recent developments naturally lead to the following general question regarding the relation between the approximability of Path TSP and TSP, which we address in this paper:

Is (Graph) Path TSP substantially harder to approximate than its well-known special case (Graph) TSP?

The answer is no. The main contribution of this paper is to show in a constructive way that Path TSP can be approximated equally well as TSP (up to an arbitrarily small error), by presenting a black-box reduction that transforms approximation algorithms for TSP into ones for Path TSP.

1.1 Our results

The main consequence of our reduction can be summarized as follows.

Theorem 1. Let \mathcal{A} be an α -approximation algorithm for TSP. Then, for any $\varepsilon > 0$, there is an $(\alpha + \varepsilon)$ approximation algorithm for Path TSP that, for any instance (G, ℓ, s, t) , calls \mathcal{A} a strongly polynomial

number of times on TSP instances defined on subgraphs of (G, ℓ) , and performs further operations taking strongly polynomial time.

The following two statements are immediate consequences of the above theorem.

Corollary 2. Let $\varepsilon > 0$ and $\alpha > 1$. If there is a (strongly) polynomial-time α -approximation algorithm for *TSP*, then there is a (strongly) polynomial-time ($\alpha + \varepsilon$)-approximation algorithm for Path TSP.

Corollary 3. Let $\varepsilon > 0$ and $\alpha > 1$. If there is a polynomial-time α -approximation algorithm for Graph *TSP*, then there is a polynomial-time $(\alpha + \varepsilon)$ -approximation algorithm for Graph Path TSP.

Notice that since Graph (Path) TSP does not involve any large numbers in its input, the notions of polynomial-time and strongly polynomial-time algorithm are identical in this context.

The above statements create a strong link between the approximability of Path TSP and TSP, as well as its graph versions. More precisely, Theorem 1 implies that such a link exists for any class of TSP instances that is closed under taking instances on subgraphs of the original instance (without changing the edge lengths). In particular, any potential future progress on the approximability of (Graph) TSP will immediately carry over to (Graph) Path TSP.

Moreover, Corollary 3 allows us to make significant progress on the currently best approximation factor of 1.497 for Graph Path TSP [26], through a black-box reduction to the 1.4-approximation algorithm for Graph TSP by Sebő and Vygen [24].

Corollary 4. For any $\varepsilon > 0$, there is a polynomial-time $(1.4 + \varepsilon)$ -approximation algorithm for Graph Path *TSP*.

Our reduction technique is quite versatile. In particular, it applies to a pretty general problem class (the Φ -tour problem with interfaces of bounded size; see Definition 6 and Theorem 25). This includes the *T*-tour problem for bounded |T| (see [24, 3, 22] for a definition) and certain uncapacitated vehicle routing problems such as the one with a fixed number of depots studied in [29].

1.2 Organization of the paper

After some brief preliminaries in Section 2 to fix basic terminology and notation, we provide an overview of our approach in Section 3. Here, we first focus on some key aspects of our approach, which is based on a new way to employ dynamic programming by using a well-chosen auxiliary problem, which we call Φ -TSP. Moreover, we break down the problem of finding a short solution to Φ -TSP into two cases. Combining the two cases, applying the same algorithm recursively, and using a constant-factor approximation algorithm for Φ -TSP on the final recursion level will imply our main reduction result, Theorem 1.

For one case, we show in Section 4 how to reduce the problem to TSP. For the other case, we show in Section 5 how to guess a constant fraction of an optimum solution via dynamic programming. The detailed proof of Theorem 1 is in Section 6. Finally, Section 7 contains a 4-approximation algorithm for Φ -TSP, which is followed by concluding remarks in Section 8.

2 Preliminaries

A weighted graph is a tuple (V, E, ℓ) , where V is the vertex set, E is the edge set, which we assume w.l.o.g. not to have loops or parallel edges, and $\ell : E \to \mathbb{R}_{\geq 0}$ denotes the edge lengths. We only consider undirected graphs with non-negative edge lengths and do not always state this explicitly.

We often deal with multi-sets of edges. Although E does not contain parallel edges, when we write $F \subseteq E$, we mean a multi-set F that can contain several copies of the same edge. We use the operator \cup to

designate the multi-union. For a vertex set $W \subseteq V$, we denote by $\delta(W) \subseteq E$ all edges with exactly one endpoint in W, and, for $v \in V$, we use $\delta(v)$ as a shorthand for $\delta(\{v\})$. For a multi-set $F \subseteq E$, we define

 $\operatorname{odd}(F) \coloneqq \{v : v \text{ is a vertex with } | \delta(v) \cap F | \text{ is odd} \}$.

For a vertex set $T \subseteq V$, a T-join is a multi-set of edges $F \subseteq E$ with odd(F) = T.

For a set $I \subseteq V$ and a graph G, we denote by G/I the graph obtained from G by contracting the vertex set I. If I is empty, we define $G/\emptyset := G$. For a vertex set W and an edge set F, we define $F[W] := \{e \in F : both endpoints of e are in W\}$. Moreover, G[W] denotes the induced subgraph with vertex set W and edge set E[W].

Instead of describing tours as walks in G, it is convenient to consider them as multi-edge sets. Then a solution to (Graph) TSP is a multi-edge set F such that (V, F) is connected and $odd(F) = \emptyset$. A solution to (Graph) Path TSP (with $s \neq t$) is a multi-edge set F such that (V, F) is connected and $odd(F) = \{s, t\}$. From such multi-edge sets—which we also call *tours* or *s*-*t tours*, respectively—we can easily recover walks by Euler's theorem.

We often use OPT as an arbitrary (but fixed) optimal solution for the problem in question. Finally, when using the notion of *approximation algorithm* we will not assume that the algorithm is polynomial time, but state it explicitly if this is the case.

In the interest of clarity and simplicity of the presentation, we did not try to optimize the running times of our procedures. Consequently, we often opt for weaker constants that are easier to obtain.

3 Overview of approach

A key novelty of our approach is a new way to set up a dynamic program to successively strengthen a basic algorithm by combining it with a stronger algorithm for TSP. Every time we apply our dynamic program to obtain a stronger algorithm, we end up with a more difficult problem, slowly approaching problem settings for which it is very challenging to find strong approximation algorithms. However, as we show, by guessing a well-chosen set of edges through the dynamic program, we can limit the recursion depth by a constant, which allows us to stay in a regime where our approach runs efficiently.

To introduce our approach, we start with a brief discussion of a much more basic dynamic programming idea that has previously been used in related settings. We explain the challenges this procedure faces when trying to extend it for our purposes, and outline how we overcome the barriers encountered by existing methods.

3.1 Key challenges and high-level approach

Assume we are given an α -approximation algorithm \mathcal{A} for TSP. Then finding a short Path TSP solution using \mathcal{A} as an oracle would be easy if the distance d(s,t) between the start s and the end t was short compared to $\ell(\text{OPT})$, i.e., the length of a shortest Path TSP solution OPT. Indeed, in this case the length of a shortest TSP tour OPT_{TSP} and a shortest s-t tour OPT do not differ by much because any solution of one problem can be converted to a solution of the other one by adding a shortest s-t path P. More precisely, $\text{OPT}_{\text{TSP}} \cup P$ is an s-t tour and $\text{OPT} \cup P$ is a TSP tour. Hence, one can simply compute an α -approximate TSP tour F and a shortest s-t path P and return $F \cup P$.

Consequently, a canonical plan would be to try to transform the Path TSP instance to another one with small s-t distance. It turns out that if the distance between s and t is very large, then such a reduction is indeed possible by using a technique based on dynamic programming that goes back to Blum, Chawla, Karger, Lane, Meyerson, and Minkoff [2], who studied variants of the Orienteering Problem. Their approach was later extended by Traub and Vygen [26] in the context of Graph Path TSP. This approach allows for

reducing to Path TSP instances where the distance between s and t is at most $(1/3 + \varepsilon) \cdot \ell(\text{OPT})$, for some arbitrarily small constant $\varepsilon > 0$ (see [26]).

However, this technique faces significant barriers when aiming at a reduction to smaller s-t distances. Thus our approach follows a different path. Nevertheless, it is on a high level inspired by the dynamic program in [2] and later variations and extensions thereof [27, 30, 26, 18]. We therefore start with a brief discussion of this prior technique in the context of Path TSP as used in [26], which will be helpful for the understanding of our approach.

For simplicity of exposition, consider a Graph Path TSP instance and assume that $d(s,t) \ge (1/3 + \varepsilon) \cdot |\text{OPT}|$ for some constant $\varepsilon > 0$. The idea is to study the structure of edges of OPT in the d(s,t) many *s*-*t* cuts $\delta(L_0), \ldots, \delta(L_{d(s,t)-1})$ defined by

$$L_i := \{ v \in V : d(s, v) \le i \} \qquad \forall i \in 0, \dots, d(s, t) - 1 .$$

The key observation is that a constant fraction of these *s*-*t* cuts will only contain a single edge of OPT, and, hence, one can try to "guess" these edges through a dynamic program. Indeed, every edge can be in at most one of the cuts $\delta(L_0), \delta(L_1), \ldots, \delta(L_{d(s,t)-1})$. Hence, the average number of OPT-edges in a cut $\delta(L_i)$ can be no higher than |OPT|/d(s,t). Using that every *s*-*t* cut must have an odd intersection with OPT, because OPT is an *s*-*t* tour, this implies that a constant fraction of the cuts contains only one edge of OPT. For brevity, we call a cut $\delta(L_i)$ with $|\delta(L_i) \cap OPT| = 1$ a 1-*cut*. Assume we knew all edges of OPT contained in 1-cuts. Then the problem decomposes into smaller Path TSP instances. See Figure 1 for an illustration.



Figure 1: Illustration of an optimal solution OPT of a Graph Path TSP instance with $9 = d(s,t) > 1/3 \cdot |OPT| = 2^3/3$. Thick blue edges are OPT-edges that are the only one in some cut $\delta(L_i)$. Knowing these edges, the problem breaks into six smaller instances: a trivial $s \cdot u_0$ Path TSP problem in $G[L_0]$, one $v_0 \cdot u_2$ Path TSP problem in $G[L_2 \setminus L_0]$ and so on, with the last instance being a trivial v_8 -t Path TSP problem in $G[V \setminus L_8]$.

Of course, the OPT-edges in 1-cuts are not known upfront, and hence, the problem cannot be decomposed so easily. However, one can use a dynamic program to guess the 1-cuts from left to right, i.e., from $\delta(L_0)$ to $\delta(L_{d(s,t)-1})$, together with the OPT-edge in each of them. Notice that the sub-instances may not have a short start-to-end distance (e.g. $d(v_6, u_8)$ in Figure 1 may be substantially larger than 1/3 times an optimum v_6 - v_8 -tour in $G[L_8 \setminus L_6]$). As shown in [26], this issue can be addressed by applying the dynamic program recursively to the sub-instances. A key observation in [26] is that a constant recursion depth is enough to ensure that the total cost of the remaining sub-instances becomes negligible compared to the edges guessed through the recursive dynamic program.

Notice that to apply this dynamic programming idea, one crucially needs $d(s,t) \ge (1/3 + \varepsilon) \cdot |OPT|$ for some constant $\varepsilon > 0$. Indeed, otherwise none of the cuts $\delta(L_i)$ may be a 1-cut, and no decomposition into smaller Path TSP instances as above is possible. This is the reason why this techniques has only been applied to reduce the start-to-end distance to about a third of OPT.

If we could guess not only 1-cuts, but also cuts with a larger constant number of OPT-edges, say up to 5, then we could handle instances with an *s*-*t* distance below $1/3 \cdot \ell$ (OPT). (This idea is inspired by a recent dynamic programming approach in [18] in the context of chain-constrained spanning trees.) Our approach aims at realizing this high-level plan. However, this ostensibly simple algorithmic idea comes with several significant technical hurdles. Most importantly, if we guess more than one edge, the resulting sub-problems are not Path TSP problems anymore. More precisely, if we guess 5 edges in each of two consecutive 5-cuts, then we have up to 10 *interface vertices*, i.e., endpoints of guessed edges. See Figure 2 for an example.



Figure 2: Illustration for guessing edges in cuts $\delta(L_i)$ that contain at most five OPT-edges. In this example, $\delta(L_0), \delta(L_1)$, and $\delta(L_3)$ are these cuts and the edges crossing them are highlighted as thick blue edges. We now define a sub-problem in $G[L_3 \setminus L_1]$. We call the endpoints of the thick blue edges *interface vertices* (red). The gray sets show the connectivity requirements on the interfaces vertices in this sub-problem: in a solution of the sub-problem, vertices in the same gray set must be in the same connected component. Interface vertices shown as squares must have odd degree; all other vertices must have even degree.

An optimum s-t tour is not necessarily connected inside the vertex set of a sub-problem but every connected component must contain at least one interface vertex. Moreover, OPT needs to connect some of the interface vertices to each other. This induces connectivity constraints for the sub-problem, shown as gray sets in Figure 2. They can also be guessed since the number of interface vertices is constant. Note, however, that we cannot guess the entire connected components, as there are exponentially many options.

Clearly, these sub-problems become significantly more difficult than the original Path TSP problem. Moreover, if we try to apply such a procedure recursively, then the sub-problems can become more complex with each recursion step, because of an increase in the number of interface vertices per sub-problem. Another important issue in a recursive application to our more complex sub-problem is to identify good cuts in which we should guess edges of OPT. Our cuts will result from the dual of a T-join problem. They will no longer form a chain, but their laminar structure still allows for a dynamic programming approach.

Moreover, it is not obvious how to reduce the problem to TSP in the case when we cannot guess edges by dynamic programming, and this will involve a careful guessing of further edges of OPT.

We will now describe our approach in detail. We start by defining a new problem class around which our method is centered, and which we call Φ -TSP.

3.2 Φ-TSP

As described above, when guessing edges, the endpoints of those edges play a special role in terms of how we have to connect things. We capture this through the notion of an interface. We define this notion for a general graph G below and will typically use it for subgraphs of the instance we are interested in.

Definition 5 (interface). An interface Φ of a graph G = (V, E) is a triple $\Phi = (I, T, C)$ with

(i) $T \subseteq I \subseteq V$, where |T| is even, and

(ii) $C \subseteq 2^I$ is a partition of I.

For an interface Φ of G, we denote by $(I_{\Phi}, T_{\Phi}, C_{\Phi})$ its corresponding triple and call $|I_{\Phi}|$ its size.

For a given interface, we are interested in finding what we call Φ -tours, which are defined as follows.

Definition 6 (Φ -tour). Let G = (V, E) be a graph. Let $\Phi = (I, T, C)$ be an interface of G. A Φ -tour in G is a multi-set $F \subseteq E$ with

- (i) T = odd(F), i.e., F is a T-join,
- (ii) (V, F)/I is connected, and
- (iii) for any $C \in C$, the vertices in C lie in the same connected component of (V, F).

Figure 3 exemplifies the notation of an interface Φ and a Φ -tour.



Figure 3: Example of an interface $\Phi = (I, T, C)$ and a Φ -tour. The partition $C = \{C_1, C_2, C_3, C_4\}$ of I is highlighted as gray sets. Hence, $I = \bigcup_{i=1}^{4} C_i$. Moreover, the vertices in T are drawn as red rectangles. This defines a Φ -TSP instance that results as a sub-problem in Figure 2. The shown edges F are a Φ -tour.

The problem we focus on in the following, which we call Φ -TSP, seeks to find a shortest Φ -tour.

Definition 7 (Φ -TSP). *Given a weighted graph* $G = (V, E, \ell)$ *and an interface* Φ *of* G*, compute a shortest* Φ *-tour in* G *or decide that none exists. In short,*

$$\min \left\{ \ell(F) : F \text{ is a } \Phi \text{-tour in } G \right\} \quad . \tag{\Phi-TSP}$$

Note that for any distinct $s, t \in V$, by choosing the interface $\Phi = (I, T, C)$ with $I = T = \{s, t\}$ and $\{C\} = \{\{s, t\}\}$, we have that Φ -tours correspond to solutions to s-t Path TSP. Analogously, for larger sets T, one captures the T-tour problem (see [24, 3, 22]). Another special case is the uncapacitated vehicle routing problem with a fixed number of depots, for which Xu and Rodrigues [29] gave a 3/2-approximation. Here, I is the set of depots, $T = \emptyset$, and C is the partition into singletons.

Depending on the structure of the graph G and the interface Φ , it may be that no Φ -tour exists. We call an interface Φ of G feasible if G admits a Φ -tour. The existence of a Φ -tour admits the following easy characterization, which can be checked in linear time.

Lemma 8. Let $G = (V, E, \ell)$ be a weighted graph. Let $\Phi = (I, T, C)$ be an interface of G. Then G admits a Φ -tour if and only if all of the following conditions hold.

- (i) Each connected component of G contains an even number of vertices in T,
- (ii) G/I is connected, and
- (iii) for every $C \in C$, the vertices in C lie in the same connected component of G.

Proof. The three mentioned conditions are clearly necessary for G to admit a Φ -tour. Moreover, if they are satisfied then, due to (i), there exists a T-join $J \subseteq E$, and points (ii) and (iii) guarantee that $E \cup (E \setminus J)$ is a Φ -tour in G.

It is crucial for our approach to start with a polynomial-time constant-factor approximation algorithm, which we will successively strengthen as discussed in the following.

A 7-approximation algorithm for Φ -TSP can be obtained easily as follows. Compute a minimum cost edge set F_1 satisfying (i) (*T*-join), a minimum cost edge set F_2 satisfying (ii) (spanning tree in G/I), and a 2-approximation F_3 of a minimum cost edge set satisfying (iii) (Steiner forest). Then the disjoint union $F_1 \cup F_2 \cup F_3 \cup F_3 \cup F_3$ is a 7-approximation.

With a little more care we can obtain a 4-approximation algorithm, using Jain's iterative rounding framework [13]:

Theorem 9. Φ -*TSP admits a strongly polynomial* 4-*approximation algorithm.*

We defer the proof to Section 7. In the rest of this paper, we will derive a strongly polynomial $(\alpha + \varepsilon)$ -approximation algorithm for Φ -TSP instances with bounded interface size, where α is the approximation guarantee for TSP; see Theorem 25.

3.3 Iterative improvement of basic algorithm

For a TSP algorithm \mathcal{A} , we denote for every weighted graph G by $f_{\mathcal{A}}(G)$ the maximum runtime of algorithm \mathcal{A} on any subgraph of G. Similarly, for a Φ -TSP algorithm \mathcal{B} , we denote for every weighted graph G and any $k \in \mathbb{R}_{\geq 0}$ by $f_{\mathcal{B}}(G, k)$ the maximum runtime of algorithm \mathcal{B} on any instance (G', Φ) , where G' is a subgraph of G and $|I_{\Phi}| \leq k$.

Our plan is to start with the 4-approximation algorithm for Φ -TSP guaranteed by Theorem 9, and successively improve it through a TSP algorithm with an approximation guarantee α . The following Boosting Theorem is the main technical result towards this goal and quantifies the improvement in terms of approximation factor that we are able to obtain in one improvement step.

Theorem 10 (Boosting Theorem). Let $\alpha, \beta > 1$. Suppose we are given:

(a) an α -approximation algorithm \mathcal{A} for TSP, and

(b) a β -approximation algorithm \mathcal{B} for Φ -TSP.

Then there is an algorithm that, for any $\varepsilon \in (0, 1]$, any weighted graph $G = (V, E, \ell)$, and any feasible interface $\Phi = (I, T, \mathcal{C})$ of G, returns a Φ -tour F in G of length

$$\ell(F) \le \max\left\{(1+\varepsilon)\alpha, \ \beta - \frac{\varepsilon}{8}(\beta-1)\right\} \cdot \ell(\text{OPT})$$
(1)

in time $|V|^{O\left(\frac{|I|}{\varepsilon}\right)} \cdot \left(f_{\mathcal{A}}(G) + f_{\mathcal{B}}\left(G, \frac{9|I|}{\varepsilon}\right)\right)$, where OPT is a shortest Φ -tour in G. In particular, the algorithm makes calls to \mathcal{B} only on instances with interfaces of size bounded by $\frac{9|I|}{\varepsilon}$.

To prove Theorem 1, we start with $\beta = 4$ (Theorem 9) and apply Theorem 10 repeatedly, but only a constant number of times. The approximation guarantee β decreases until it reaches $(1 + \varepsilon)\alpha$. All interfaces will have constant size. We defer the details to Section 6.

3.4 **Proof outline of Boosting Theorem (Theorem 10)**

Theorem 10 is obtained by designing two algorithms to obtain a Φ -tour and then returning the better of the Φ -tours computed by these algorithms. Each of the following two theorems summarizes the guarantee we obtain with one of the two algorithms. After that, Algorithm 1, described below, combines these two sub-procedures to obtain an algorithm that implies Theorem 10.

The following theorem yields a short Φ -tour if the length of a minimum T_{Φ} -join is small.

Theorem 11. Let $\alpha > 1$. Assume we are given an α -approximation algorithm \mathcal{A} for TSP. Then, for any $\delta > 0$, any weighted graph $G = (V, E, \ell)$, and any feasible interface $\Phi = (I, T, \mathcal{C})$ of G, one can determine a Φ -tour F in G with

$$\ell(F) \le (1+\delta) \cdot \alpha \cdot \ell(\text{OPT}) + (\alpha+1) \cdot \ell(J)$$

in time $|V|^{O\left(\frac{|I|}{\delta}\right)} \cdot f_{\mathcal{A}}(G)$, where J is a shortest T-join in G and OPT is a shortest Φ -tour in G.

We will give the proof in Section 4. The next theorem, proven in Section 5, states that we also obtain a short Φ -tour if the length of a minimum T-join is large.

Theorem 12. Let $\beta > 1$. Assume we are given a β -approximation algorithm \mathcal{B} for Φ -TSP. Then, for any $\delta > 0$, any weighted graph $G = (V, E, \ell)$, and any feasible interface $\Phi = (I, T, \mathcal{C})$ of G, one can determine a Φ -tour F in G with

 $\ell(F) \leq \left(\beta + \delta \cdot (\beta - 1)\right) \cdot \ell(\mathrm{OPT}) - (\beta - 1) \cdot \ell(J)$

in time $|V|^{O\left(|I|+\frac{|T|}{\delta}\right)} \cdot f_{\mathcal{B}}\left(G, |I|+\frac{|T|}{\delta}\right)$, where J is a shortest T-join in G and OPT is a shortest Φ -tour in G.

Algorithm 1: Approximation algorithm for Φ -TSP to prove Theorem 10

- 1. Run algorithm guaranteed by Theorem 11 with $\delta = \varepsilon/2$ to obtain Φ -tour F_1 .
- 2. Run algorithm guaranteed by Theorem 12 with $\delta = \varepsilon/8$ to obtain Φ -tour F_2 .
- 3. Return the shorter Φ -tour among F_1 and F_2 .

Lemma 13. Given a weighted graph G and a feasible interface $\Phi = (I, T, C)$ of G, Algorithm 1 returns a Φ -tour F in G with the guarantees stated in Theorem 10.

Proof. The running time guarantee stated in Theorem 10 immediately follows from Theorem 11 and Theorem 12, using $|I| + \frac{|T|}{\varepsilon/8} \le \frac{9|I|}{\varepsilon}$.

Let $F \in \{F_1, F_2\}$ be the Φ -tour returned by Algorithm 1. To show that F fulfills the approximation guarantee stated in (1), we distinguish two cases.

If $\ell(J) \leq \frac{\varepsilon}{4} \cdot \ell(\text{OPT})$, then the solution F_1 will be short enough:

$$\ell(F) \le \ell(F_1) \le \left(1 + \frac{\varepsilon}{2}\right) \alpha \cdot \ell(\text{OPT}) + (\alpha + 1)\frac{\varepsilon}{4} \cdot \ell(\text{OPT}) \le (1 + \varepsilon) \cdot \alpha \cdot \ell(\text{OPT})$$

where we used $(\alpha + 1)/2 \leq \alpha$ for the last inequality, which holds because $\alpha \geq 1$.

If $\ell(J) \geq \frac{\varepsilon}{4} \cdot \ell(\text{OPT})$, then the Φ -tour F_2 will be short enough:

$$\ell(F) \le \ell(F_2) \le \left(\beta + \frac{\varepsilon}{8}(\beta - 1)\right) \cdot \ell(\text{OPT}) - (\beta - 1) \cdot \frac{\varepsilon}{4} \cdot \ell(\text{OPT}) = \left(\beta - \frac{\varepsilon}{8}(\beta - 1)\right) \cdot \ell(\text{OPT}) ,$$

thus completing the proof of Lemma 13.

For the proof of Theorem 10, it remains to show Theorem 11 and Theorem 12.

4 Finding a short Φ -tour if there is a short T-join

In this section we prove Theorem 11, i.e., how to get a short Φ -tour if the shortest T-join has small length compared to $\ell(\text{OPT})$.

We start by analyzing a simple algorithm for computing a Φ -tour. However, this simple algorithm will not be sufficient to prove Theorem 11. Thus in a second step, we will refine the algorithm to obtain the desired bound.

Algorithm	2: A	A sim	ple Φ-	TSP	algorithm
-----------	------	-------	--------	-----	-----------

Input: a weighted graph G, an interface $\Phi = (I, T, C)$ of G, and a T-join J in G. **Output:** an edge set F.

- 1. In each connected component of G apply A to get an α -approximate TSP-tour. Let Q be the union of these tours.
- 2. Return $F = Q \cup J$.

Notice that Algorithm 2 always returns an edge set F, even if the input is infeasible. We therefore show first that Algorithm 2 does return a Φ -tour whenever it is run with a feasible input.

Lemma 14. The set F returned by Algorithm 2 is a Φ -tour if and only if the input is feasible, i.e., G admits a Φ -tour.

Proof. Assume that G admits a Φ -tour, which implies by Lemma 8 that the three properties (i), (ii), and (iii) listed in Lemma 8 are fulfilled. Because the set Q computed in Algorithm 2 consists of TSP tours in each connected component of G, the vertex sets of the connected components of (V, Q) and G are the same. Because G fulfills (ii) and (iii), this implies that also (V, Q) and $(V, Q \cup J)$ fulfill these two conditions. Finally, $odd(Q \cup J) = odd(J) = T$, because $odd(Q) = \emptyset$ and J is a T-join, which shows that $Q \cup J$ is indeed a Φ -tour.

We now analyze the length of the Φ -tour returned by Algorithm 2.

Lemma 15. Assume we are given an α -approximation algorithm \mathcal{A} for TSP. Let $G = (V, E, \ell)$ be a weighted graph, $\Phi = (I, T, \mathcal{C})$ a feasible interface of G, and J a T-join in G. Then, Algorithm 2 computes a Φ -tour F in G with

 $\ell(F) \le \alpha \cdot \ell(\text{OPT}) + (\alpha + 1) \cdot \ell(J) + 2\alpha \cdot |I| \cdot \max\left\{\ell(e) : e \in E \setminus (\text{OPT} \cup J)\right\} ,$

in time $O(|V| \cdot f_{\mathcal{A}}(G))$, where OPT is a shortest Φ -tour. Here $\max \emptyset := 0$.

Proof. First, observe that the running time is indeed as claimed, because the bottleneck of the algorithm is calling \mathcal{A} for each connected component of G; moreover, the connected components can be found in linear time and there are at most |V| many of them.

To bound $\ell(Q)$, we transform a shortest Φ -tour OPT into a union of TSP solutions, one for each connected component of G. Let $L \subseteq E$ be a minimal edge set such that the vertex sets of the connected components of $(V, \text{OPT} \cup J \cup L)$ and G are the same. Observe that the multi-set $\text{OPT} \cup J \cup L \cup L$ is a union of TSP solutions, one for each connected component of G. Because the set Q determined in Algorithm 2 was obtained through \mathcal{A} , which is an α -approximation algorithm, we have

$$\ell(Q) \le \alpha \cdot \left(\ell(\text{OPT}) + \ell(J) + 2\ell(L)\right)$$
,

and, hence, the solution $F = Q \cup J$ returned by the algorithm satisfies

$$\ell(F) = \ell(Q) + \ell(J) \le \alpha \cdot \ell(\text{OPT}) + (\alpha + 1) \cdot \ell(J) + 2\alpha \cdot \ell(L) \quad (2)$$

Moreover, because OPT is a Φ -tour, we have that (V, OPT)/I must be connected, which implies that (V, OPT) has at most |I| connected components, and thus

$$|L| \leq |I| - 1 \; .$$

Together with (2), this leads to the desired guarantee:

$$\ell(F) \le \alpha \cdot \ell(\text{OPT}) + (\alpha + 1) \cdot \ell(J) + 2\alpha \cdot |I| \cdot \max\left\{\ell(e) : e \in E \setminus (\text{OPT} \cup J)\right\}$$

where the inequality follows from $L \subseteq E \setminus (OPT \cup J)$, which holds because the edges in L connect different connected components of $(V, OPT \cup J)$.

We now explain how to refine Algorithm 2 by a guessing step to obtain the guarantees claimed in Theorem 11. If all edges that are not contained in $OPT \cup J$ have length at most $\frac{\delta \cdot \ell(OPT)}{2 \cdot |I|}$, Lemma 15 already implies the desired bound. To obtain this property, we delete all edges from G that are *heavy*, i.e. have length at least $\frac{\delta \cdot \ell(OPT)}{2 \cdot |I|}$, and are not contained in $OPT \cup J$. We guess this set of edges to delete as follows. First we guess the set H of heavy edges, which can be done in polynomial time by guessing a minimum length edge in H. Then we guess the set $H^* = OPT \cap H$ of heavy edges contained in OPT. Algorithm 3 formalizes this procedure and, as we show next, indeed implies Theorem 11.

Algorithm 3: Φ -TSP algorithm to prove Theorem 11

Compute a shortest T-join J in G. For $f \in E$ define $H_f := \{e \in E : \ell(e) \ge \ell(f)\}$. for every edge set $H \in \{H_f : f \in E\} \cup \{\emptyset\}$ do for every set $H^* \subseteq H$ with $|H^*| \le 2|I|/\delta$ do Set $D := H \setminus (H^* \cup J)$. Apply Algorithm 2 to the graph $(V, E \setminus D)$ to obtain a multi-set F_D of edges, which is a Φ -tour in G if the input is feasible. Among all computed Φ tours F_D return a shortest one

Among all computed Φ -tours F_D , return a shortest one.

Proof of Theorem 11. We start by observing that the running time of Algorithm 3 is indeed bounded by $|V|^{O(|I|/\delta)} \cdot f_{\mathcal{A}}(G)$. There are at most $|V|^2$ possible edges f that are being considered in the outer for-loop. For each of them, there are $|V|^{O(|I|/\delta)}$ possible sets H^* considered in the inner for-loop. Thus, there are at most $|V|^{O(|I|/\delta)}$ calls to Algorithm 2. Finally, all other operations can be done in time $|V|^{O(1)}$.

We now show that Algorithm 3 returns a Φ -tour with the guarantees claimed by Theorem 11. Let OPT be a shortest Φ -tour and let $H := \{e \in E : \ell(e) \ge \delta \cdot \ell(\text{OPT})/2|I|\}$ be the set of heavy edges. Then in some iteration of the outer for-loop we consider the set H. Because

$$\ell(\text{OPT}) \ge \ell(H \cap \text{OPT}) \ge |H \cap \text{OPT}| \cdot \frac{\delta \cdot \ell(\text{OPT})}{2 \cdot |I|}$$

we have $|H \cap OPT| \leq 2|I|/\delta$, and thus, we consider the set $H^* := H \cap OPT$ in some iteration of the inner for-loop. As $D = H \setminus (H^* \cup J)$ does not contain any edge of OPT, the Φ -tour OPT is a feasible solution of the instance to which we apply Algorithm 2. Moreover, the set D contains all heavy edges not contained in $OPT \cup J$ and hence by Lemma 15, we obtain

$$\ell(F_D) \le \alpha \cdot \ell(\text{OPT}) + (\alpha + 1) \cdot \ell(J) + 2\alpha \cdot |I| \cdot \max\left\{\ell(e) : e \in (E \setminus D) \setminus (\text{OPT} \cup J)\right\}$$
$$\le \alpha \cdot \ell(\text{OPT}) + (\alpha + 1) \cdot \ell(J) + 2\alpha \cdot |I| \cdot \frac{\delta \cdot \ell(\text{OPT})}{2 \cdot |I|}$$
$$= (1 + \delta) \cdot \alpha \cdot \ell(\text{OPT}) + (\alpha + 1) \cdot \ell(J) .$$

5 Iterative improvement via dynamic programming

In this section, we show how to prove Theorem 12, i.e., how to obtain a short Φ -tour if the length of a shortest *T*-join is large. Here, our goal is to use dynamic programming to "guess" a significant portion, in terms of total length, of edges used in OPT. Very recently, dynamic programming has become a strong tool in the context of Path TSP, Chain-Constrained Spanning Trees, and related problems [27, 26, 30, 18], leading to the currently best known approximation factors for these settings. The dynamic programming idea we employ combines and extends elements used in these prior dynamic programming techniques.

What we aim to achieve with dynamic programming in the context of Φ -TSP, for some interface $\Phi = (I, T, C)$ of G, is the following. We can fix an arbitrary laminar family \mathcal{L} of subsets of V. Our goal is to guess what edges of OPT are crossing the cuts in \mathcal{L} . Clearly, if $OPT \cap \delta(L)$ contains many edges for some $L \in \mathcal{L}$, it seems computationally elusive to guess them. This is the reason why we fix some constant k and only guess OPT-edges in cuts $\delta(L)$ for $L \in \mathcal{L}$ if $|OPT \cap \delta(L)| \leq k$. We denote the sets inducing these cuts by $\mathcal{L}(OPT, k) \subseteq \mathcal{L}$ and the OPT-edges in these cuts by $OPT(\mathcal{L}, k) \subseteq OPT$. Formally, for any edge set $R \subseteq E$, we define

$$\begin{split} \mathcal{L}(R,k) &\coloneqq \{L \in \mathcal{L} : |R \cap \delta(L)| \leq k\} \ , \text{ and} \\ R(\mathcal{L},k) &\coloneqq \bigcup_{L \in \mathcal{L}(R,k)} \left(\delta(L) \cap R\right) \ . \end{split}$$

As we discuss in more detail later, guessing the edges $OPT(\mathcal{L}, k)$ can be achieved through a dynamic program that guesses the OPT-edges in the different cuts defined by \mathcal{L} step by step, from smaller to larger sets in \mathcal{L} . However, the running time of the propagation step of the dynamic program depends on the number of disjoint sets in \mathcal{L} that can be contained in some larger set $L \in \mathcal{L}$. We capture this dependency through the width width(\mathcal{L}) of the laminar family \mathcal{L} (see [18] for a similar use of this notion).

Definition 16 (width of a laminar family). The width width(\mathcal{L}) of a laminar family \mathcal{L} is the number of minimal sets contained in the family.

Observe that the number of minimal sets of a laminar family bounds the size of any subfamily of disjoint sets.

The following theorem formalizes what we can achieve through our dynamic program, which we present later in detail. Notice that for the algorithm to be efficient, we need \mathcal{L} to have width bounded by a constant.

Theorem 17. Let $\beta > 1$. Assume there is a β -approximation algorithm \mathcal{B} for Φ -TSP. Then there is an algorithm that computes for any feasible interface $\Phi = (I, T, C)$ of a weighted graph $G = (V, E, \ell)$, any $k \in \mathbb{Z}_{>0}$, and any laminar family \mathcal{L} over V, a Φ -tour F with

$$\ell(F) \le \min\left\{\beta \cdot \ell(R) - (\beta - 1) \cdot \ell(R(\mathcal{L}, k)) : R \text{ is a } \Phi \text{-tour}\right\}$$
(3)

in time $|V|^{O(|I|+k \cdot \text{width}(\mathcal{L}))} \cdot f_{\mathcal{B}}(G, |I|+k \cdot (\text{width}(\mathcal{L})+1))$. In particular, the algorithm calls \mathcal{B} only on instances with interfaces of size bounded by $|I|+k \cdot (\text{width}(\mathcal{L})+1)$.

Note that the guarantee stated in (3) for R = OPT indeed reflects the guessing of the edges in $OPT(\mathcal{L}, k)$. More precisely, by replacing R by OPT in (3), we obtain a Φ -tour F with an upper bound on its length $\ell(F)$ that decomposes into two terms:

- (i) a term $\ell(OPT(\mathcal{L}, k))$, i.e., each edge $e \in OPT(\mathcal{L}, k)$ contributes its length $\ell(e)$, and
- (ii) a term $\beta \cdot \ell(\text{OPT} \setminus \text{OPT}(\mathcal{L}, k))$, where the length of each other edge in OPT gets inflated by the approximation factor β of the algorithm \mathcal{B} .

5.1 Finding a suitable laminar family

To make significant progress through Theorem 17, we need to find a laminar family \mathcal{L} over V such that $\ell(\operatorname{OPT}(\mathcal{L}, k))$ is large. Let J be a shortest T-join. If $\ell(J)$ is large, then we will construct a family \mathcal{L} with the property that even for any T-join R, the length $\ell(R(\mathcal{L}, k))$ is large. Notice that this implies what we want because OPT is a T-join.

This statement is formalized in Lemma 19, which is derived from the dual of the natural linear program to find a shortest T-join. We exploit that there is an optimal dual solution whose support corresponds to a laminar family of subsets of V, which follows from combinatorial uncrossing arguments.

Lemma 18. Let $G = (V, E, \ell)$ be a weighted graph. Moreover, let $T \subseteq V$ such that G contains a T-join, and let $t \in T$. Then there is strongly polynomial algorithm that computes a laminar family \mathcal{L} over $V \setminus \{t\}$ and values $y \in \mathbb{R}_{\geq 0}^{\mathcal{L}}$ such that

$$\sum_{\substack{L \in \mathcal{L}:\\ e \in \delta(L)}} y_L \le \ell(e) \qquad \forall e \in E \quad , \tag{4}$$

$$\sum_{L \in \mathcal{L}} y_L = \ell(J), \text{ and}$$
(5)

$$|L \cap T| \text{ is odd} \qquad \forall L \in \mathcal{L} .$$
(6)

Proof. We start with a classical linear description to find a minimum length T-join, based on the dominant of the T-join polytope. To this end, let $\mathcal{F} = \{Q \subseteq V \setminus \{t\} : |Q \cap T| \text{ is odd}\}$; these vertex sets induce all T-cuts. Then, the following linear program computes the value $\ell(J)$ of a shortest T-join (see, e.g., [21]).

$$\min \sum_{\substack{e \in E \\ x(\delta(Q)) \ge 1 \\ x \in \mathbb{R}_{\ge 0}^{E}}} \forall Q \in \mathcal{F}$$
(7)

Its dual problem, which is a fractional T-cut packing problem, is given below.

$$\max \sum_{\substack{Q \in \mathcal{F}:\\ e \in \delta(Q)}} y_Q \leq \ell(e) \quad \forall \ e \in E$$

$$y \in \mathbb{R}_{>0}^{\mathcal{F}}$$
(8)

If $y \in \mathbb{R}_{\geq 0}^{\mathcal{F}}$ is an optimum dual solution with laminar support \mathcal{L} , then y and \mathcal{L} have the desired properties. Here (5) follows from strong duality and (6) follows from $L \in \mathcal{F}$.

A strongly polynomial algorithm to compute such an optimal dual solution with laminar support can be obtained by standard techniques: Using the framework of Frank and Tardos [7], one can first find in strongly polynomial time a vector $\hat{\ell} \in \mathbb{R}_{\geq 0}^{E}$ with encoding length polynomial in |E|, and such that the set of optimal solutions of (7) remains the same when replacing ℓ by $\hat{\ell}$. Moreover, also the set of optimal dual bases remains the same. This allows for solving (7) in strongly polynomial time through the ellipsoid method. To find an optimal dual basis, one can delete all variables from (8) that do not correspond to constraints encountered by the ellipsoid algorithm when solving (8). Now solving the reduced dual problem (8) with $\hat{\ell}$ instead of ℓ allows for finding an optimal dual basis, which, by the result of Frank and Tardos, remains an optimal dual basis for (8) without replacing ℓ by $\hat{\ell}$. Knowing an optimal dual basis, one can obtain an optimal dual basis for (8) in strongly polynomial time by solving a linear equation system.

Finally, this solution can be transformed into a laminar one by uncrossing: if $y_A > 0$ and $y_B > 0$ for $A, B \in \mathcal{F}$ with $A \setminus B \neq \emptyset$ and $B \setminus A \neq \emptyset$ and $A \cap B \neq \emptyset$, then either $A \cap B$ and $A \cup B$ belong to \mathcal{F} or $A \setminus B$ and $B \setminus A$ belong to \mathcal{F} ; we can increase the dual variables on these two sets by min $\{y_A, y_B\}$ and decrease the dual variables y_A and y_B by the same amount, maintaining a feasible dual solution. Karzanov [15] showed how to obtain a laminar family by a sequence of such uncrossing steps in strongly polynomial time.

We now show that the family \mathcal{L} from Lemma 18 has the desired properties.

Lemma 19. Let $G = (V, E, \ell)$ be a weighted graph. Moreover, let $T \subseteq V$ such that G admits a T-join. Then, there is a strongly polynomial algorithm that computes a laminar family \mathcal{L} over V with width $(\mathcal{L}) \leq \max\{0, |T| - 1\}$ such that for any T-join $R \subseteq E$, and any $k \in \mathbb{Z}_{>0}$, we have

$$\ell(R(\mathcal{L},k)) \ge \ell(J) - \frac{1}{k+1} \cdot \ell(R)$$
,

where J is a shortest T-join in G.

Proof. If $T = \emptyset$, we can simply set $\mathcal{L} = \emptyset$ because $\ell(J) = 0$. Otherwise, we compute \mathcal{L} and y as in Lemma 18 and show that \mathcal{L} has the desired properties. Since every set in \mathcal{L} must contain an element of $T \setminus \{t\}$, we have width $(\mathcal{L}) \leq |T| - 1$.

Let now $R \subseteq E$ be a *T*-join, and let $k \in \mathbb{Z}_{\geq 0}$. Since *R* is a *T*-join, it has a non-empty intersection with every cut $\delta(L)$ with $L \in \mathcal{L}$ because of (6). Hence, by (4),

$$\ell(R(\mathcal{L},k)) = \sum_{e \in R(\mathcal{L},k)} \ell(e) \ge \sum_{e \in R(\mathcal{L},k)} \sum_{\substack{L \in \mathcal{L}:\\ e \in \delta(L)}} y_L = \sum_{L \in \mathcal{L}} |R(\mathcal{L},k) \cap \delta(L)| \cdot y_L \ge \sum_{L \in \mathcal{L}(R,k)} y_L \quad .$$
(9)

Again using (4), we moreover obtain

$$\ell(R) \ge \sum_{e \in R} \sum_{\substack{L \in \mathcal{L}:\\ e \in \delta(L)}} y_L = \sum_{L \in \mathcal{L}} |R \cap \delta(L)| \cdot y_L \ge \sum_{L \in \mathcal{L} \setminus \mathcal{L}(R,k)} (k+1) \cdot y_L \quad .$$
(10)

Combining (5), (9), and (10), we obtain

$$\ell(J) = \sum_{L \in \mathcal{L}} y_L = \sum_{L \in \mathcal{L}(R,k)} y_L + \sum_{L \in \mathcal{L} \setminus \mathcal{L}(R,k)} y_L \le \ell(R(\mathcal{L},k)) + \frac{1}{k+1} \cdot \ell(R) ,$$

as desired.

Finally, Theorem 12 is a direct consequence of Theorem 17 and Lemma 19.

Proof of Theorem 12. If $T = \emptyset$, we simply call the given β -approximation algorithm \mathcal{B} . Otherwise, let $k = \lfloor 1/\delta \rfloor$. We apply Lemma 19 to obtain in strongly polynomial time a laminar family \mathcal{L} over V such that (i) $\ell(R(\mathcal{L}, k)) \ge \ell(J) - \frac{1}{k+1} \cdot \ell(R) \ge \ell(J) - \delta \cdot \ell(R) \quad \forall T$ -join $R \subseteq E$, and (ii) width $(\mathcal{L}) \le \max\{0, |T| - 1\} = |T| - 1$, where the equality follows from the assumption $T \neq \emptyset$.

Because a shortest Φ -tour OPT is a *T*-join, we have

$$\ell(\operatorname{OPT}(\mathcal{L}, k)) \geq \ell(J) - \delta \cdot \ell(\operatorname{OPT})$$
,

which, together with Theorem 17 implies the desired results, i.e., that one can find a Φ -tour F in G with

$$\ell(F) \leq \beta \cdot \ell(\text{OPT}) - (\beta - 1) \cdot \ell(\text{OPT}(\mathcal{L}, k))$$

$$\leq \beta \cdot \ell(\text{OPT}) - (\beta - 1) (\ell(J) - \delta \cdot \ell(\text{OPT}))$$

$$= (\beta + \delta \cdot (\beta - 1)) \cdot \ell(\text{OPT}) - (\beta - 1) \cdot \ell(J)$$

in time

$$|V|^{O\left(|I|+\frac{\operatorname{width}(\mathcal{L})}{\delta}\right)} \cdot f_{\mathcal{B}}\left(G, |I| + \frac{\operatorname{width}(\mathcal{L}) + 1}{\delta}\right) \le |V|^{O\left(|I|+\frac{|T|}{\delta}\right)} \cdot f_{\mathcal{B}}\left(G, |I| + \frac{|T|}{\delta}\right) \quad .$$

It remains to derive Theorem 17, which, as mentioned, we show through a dynamic programming approach.

5.2 Combining partial solutions

In the analysis of our dynamic programming algorithm we use the following notion of an *induced interface*, which allows us to analyze the algorithm with respect to interfaces coming from a shortest Φ -tour OPT.

Definition 20 (induced interface). Let $G = (V, E, \ell)$ be a weighted graph. Let $\Phi = (I, T, C)$ be an interface of G, and let F be a Φ -tour in G. For $W \subseteq V$, the interface $\Phi_W = (I_W, T_W, C_W)$ induced by (F, Φ) on W is defined by

- (i) $I_W = (I \cap W) \cup U$, where U is the set of vertices in W that are connected by an edge of F to a vertex in $V \setminus W$,
- (ii) $T_W = \text{odd}(F[W])$, and
- (iii) $C_W \subseteq 2^{I_W}$ contains, for each connected component of (W, F[W]), a set including all vertices of I_W contained in that connected component.

See Figure 4 for an example of an induced interface. Moreover, also Figure 2, which we used as an illustrative example in the introduction to showcase the guessing of multiple edges per cut, highlights an induced interface with $W = L_3 \setminus L_1$, which is induced by an *s*-*t* tour. We remark that the interface induced by (F, Φ) depends only on F and I, not on T or C.

The following lemma shows some basic properties of induced interfaces.

Lemma 21. Let $G = (V, E, \ell)$ be a weighted graph and $\Phi = (I, T, C)$ an interface of G. Let F be a Φ -tour in G and $W \subseteq V$. Let Φ_W be the interface induced by (F, Φ) on W. Then

- (i) Φ_W is an interface of G[W],
- (ii) F[W] is a Φ_W -tour in G[W], and
- (iii) for every $W' \subseteq W$, the interface induced by $(F[W], \Phi_W)$ on W' equals the interface induced by (F, Φ) on W'.

Proof. Let $\Phi_W = (I_W, T_W, C_W)$. As in Definition 20 (i), let U be the set of vertices in W that are connected by an edge of F to a vertex in $V \setminus W$.

To prove (i), we have to observe that $T_W \subseteq I_W$. (Notice that we clearly have that $|T_W|$ is even because $T_W = \text{odd}(F[W])$.) Let $u \in T_W$. If F contains an edge connecting u with $V \setminus W$, then $u \in U$ and hence $u \in I_W$. Otherwise we have $(\delta(u) \cap F) \subseteq E[W]$ and hence $u \in T_W = \text{odd}(F[W])$ implies $u \in \text{odd}(F)$. Since F is a Φ -tour, we conclude $u \in T \subseteq I$. Moreover, $u \in T_W = \text{odd}(F[W]) \subseteq W$, so $u \in I \cap W \subseteq I_W$.

To prove (ii), we have to show that $(W, F[W])/I_W$ is connected (the other two conditions of Definition 6 trivially hold). Suppose not. Then there is a set $W' \subseteq W \setminus I_W$ with $W' \neq W$ and $F[W] \cap \delta(W') = \emptyset$.



Figure 4: On the left-hand side, an interface $\Phi = (I, T, C)$ on a graph G = (V, E) is shown together with a Φ -tour $F \subseteq E$ (the black edges) and a set $W \subseteq V$. The right-hand side figure depicts the interface $\Phi_W = (I_W, T_W, C_W)$ induced by (F, Φ) on W.

This implies, together with $I_W = (I \cap W) \cup U$ —which holds by definition of I_W —that $W' \subseteq V \setminus I$ with $W' \neq V$ and $F \cap \delta(W') = F[W] \cap \delta(W') = \emptyset$. This contradicts the fact that (V, F)/I is connected, which has to hold because F is a Φ -tour.

To show (iii), let (I_1, T_1, C_1) be the interface induced by (F, Φ) on W' and let (I_2, T_2, C_2) be the interface induced by $(F[W], \Phi_W)$ on W'. Let U_1 be the set of vertices in W' that are connected by an edge of F to a vertex in $V \setminus W'$. Let U_2 be the set of vertices in W' that are connected by an edge of F[W] to a vertex in $W \setminus W'$. Then $U_1 = (U \cap W') \cup U_2$. Therefore,

$$I_1 = (I \cap W') \cup U_1$$

= $(I \cap W') \cup (U \cap W') \cup U_2$
= $(((I \cap W) \cup U) \cap W') \cup U_2$
= $(I_W \cap W') \cup U_2$
= I_2 .

Finally, because (F[W])[W'] = F[W'], which follows from $W' \subseteq W$, we have

$$T_2 = \text{odd}((F[W])[W']) = \text{odd}(F[W']) = T_1$$
,

and also $C_1 = C_2$, because these partitions of $I_1 = I_2$ are both defined with respect to the connected components of (W', F[W']), because (W', (F[W])[W']) = (W', F[W']).

Notice that given an interface $\Phi = (I, T, C)$ on a graph G = (V, E) and a Φ -tour $F \subseteq E$, then the interface Φ_V induced by (F, Φ) on V is not necessarily identical to Φ . More precisely, $\Phi_V = (I_V, T_V, C_V)$ always fulfills $I_V = I$ and $T_V = T$. However, F may connect different parts of the partition C, which, in the interface Φ_V , will then only appear as one set in C_V . See the left-hand side illustration in Figure 4 for such an example where the highlighted Φ -tour would induce an interface $\Phi_V \neq \Phi$ on V because $C_2 \cup C_3$ is a single set in C_V .

In our dynamic program we will combine solutions for different subgraphs with induced interfaces. The following lemma shows sufficient conditions under which this works out.



Figure 5: The dashed ellipsoids show the partition of V into W_0, \ldots, W_p . The thick blue edges are the edges in X. Only the vertices in \overline{I} are shown here, where the vertices with a red boundary are those contained in I.

Lemma 22. Let $G = (V, E, \ell)$ be a weighted graph. Let $\Phi = (I, T, C)$ be an interface of G and let F be a Φ -tour in G. Let W_0, \ldots, W_p be a partition of V. For $i \in \{0, \ldots, p\}$, let $\Phi_i = (I_i, T_i, C_i)$ be the interface induced by (F, Φ) on W_i , and let F_i be a Φ_i -tour in $G[W_i]$. Then

$$F' \coloneqq X \stackrel{\cdot}{\cup} \left(\bigcup_{i=0}^p F_i \right)$$

is a Φ -tour in G, where $X := F \cap \bigcup_{i=0}^{p} \delta(W_i)$.

Proof. We first show point (i) of Definition 6, i.e. odd(F') = T. For $i \in \{0, ..., p\}$, we have $odd(F_i) = T_i = odd(F[W_i])$ since F_i is a Φ_i -tour. Thus

$$\operatorname{odd}(F') = \operatorname{odd}(X) \bigtriangleup \operatorname{odd}(F_0) \bigtriangleup \cdots \bigtriangleup \operatorname{odd}(F_p)$$

= $\operatorname{odd}(X) \bigtriangleup \operatorname{odd}(F[W_0]) \bigtriangleup \cdots \bigtriangleup \operatorname{odd}(F[W_p])$
= $\operatorname{odd}(F)$
= T ,

where \triangle denotes the symmetric difference; we used $F = X \cup F[W_0] \cup \ldots \cup F[W_p]$. Before proving that F' also fulfills the remaining two properties of a Φ -tour, we show the following claim. See Figure 5 for an illustration.

Claim 23. Let $\overline{I} := I_0 \cup \ldots \cup I_p$ and $a, b \in \overline{I}$. Suppose (V, F) contains an a-b path. Then (V, F') contains an a-b path.

Proof of Claim 23. Suppose the claim is wrong. Then there exist vertices $a, b \in \overline{I}$ such that (V, F) contains an a-b path P, but (V, F') does not. We choose a, b, and P such that the number of edges of P is minimum. Consequently, P contains no vertex of $\overline{I} \setminus \{a, b\}$. We now distinguish two cases. **Case 1:** $X \cap E(P) = \emptyset$.

Then P is completely contained in a single set W_i for some $i \in \{0, ..., p\}$, by definition of X. Hence, $a, b \in W_i \cap \overline{I} = I_i$ and a and b are connected by the path P in $(W_i, F[W_i])$. Since Φ_i is the interface induced by (F, Φ) on W_i , the vertices a and b are contained in the same set of the partition C_i of I_i . This implies that every Φ_i -tour, and in particular F_i , must contain an a-b path, contradicting the assumption that (V, F') contains no a-b path. Case 2: $X \cap E(P) \neq \emptyset$.

Recall $X = F \cap \bigcup_{i=0}^{p} \delta(W_i)$. For $i \in \{0, \dots, p\}$, the set I_i contains all vertices of W_i that are an endpoint of an edge in X, by definition of the induced interface Φ_i . Thus all endpoints of edges in X are contained in \overline{I} . Since P contains no vertex of $\overline{I} \setminus \{a, b\}$, we have $X \cap E(P) = \{\{a, b\}\}$, i.e. the path P consists only of a single edge that is contained in X and thus also in F'. This contradicts our assumption that (V, F') contains no a-b path.

(proof of Claim 23)

To show point (iii) of Definition 6, we need to show that any two vertices a and b that are contained in the same set of the partition C of I are also contained in the same connected component of (V, F'). If a and b are contained in the same set of the partition C, they are contained in the same connected component of (V, F) because F is a Φ -tour. Hence by Claim 23 and $I \subseteq \overline{I}$, also (V, F') contains an a-b path.

It remains to show point (ii) of Definition 6, i.e., we prove that (V, F')/I is connected. First observe that if p = 0, then the result holds because then $F' = F_0$ is a Φ_0 -tour and $I_0 = I$. Hence, assume from now on p > 0. In this case, we first observe that

$$I_i \neq \emptyset \quad \forall i \in \{0, \dots, p\} \ . \tag{11}$$

Indeed, because (V, F)/I is connected, which follows from F being a Φ -tour, we have for each $i \in \{0, \ldots, p\}$ that either $I \cap W_i \neq \emptyset$ or $\delta(W_i) \cap F \neq \emptyset$, both of which imply $I_i \neq \emptyset$.

To conclude that (V, F')/I is connected, we will observe the following two properties, which immediately imply the result:

(a) For each $i \in \{0, ..., p\}$, each vertex $v \in W_i$ is connected to a vertex in I_i in the graph (W_i, F_i) .

(b) All vertices in $\bigcup_{i=0}^{p} I_i$ are connected in (V, F')/I.

Notice that (a) is a consequence of (11) and the fact that $(W_i, F_i)/I_i$ is connected, which holds because F_i is a Φ_i -tour in $G[W_i]$. Finally, (b) follows from Claim 23 due to the following. Either $I = \emptyset$, in which case (V, F)/I = (V, F) is connected—because F is a Φ -tour—which implies (b) by Claim 23. Or $I \neq \emptyset$, in which case the connectivity of (V, F)/I implies that in (V, F) each vertex $v \in \bigcup_{i=0}^p I_i$ is connected to a vertex of I, again implying (b) by Claim 23.

5.3 The dynamic program

We now expand on the dynamic program used to show Theorem 17. The dynamic program is formally described by Algorithm 4 below. See also Figure 6 for an illustration. Before formally proving that Algorithm 4 indeed returns a Φ -tour implying Theorem 17, we provide a brief explanatory discussion outlining the core ideas of the algorithm and the line of reasoning we employ to show its correctness.

To this end, let R be a Φ -tour (unknown to the algorithm), and we will show that the dynamic program returns a Φ -tour $F \subseteq E$ such that $\ell(F) \leq \beta \cdot \ell(R) - (\beta - 1) \cdot \ell(R(\mathcal{L}, k))$. Conceptually, we want to consider the elements of the laminar family $\mathcal{L}(R, k) \subseteq \mathcal{L}$ from smaller to larger ones. Since we do not know the laminar family $\mathcal{L}(R, k)$, we consider all sets in \mathcal{L} in an arbitrary fixed order of non-decreasing cardinality. We then guess, for every vertex set $L \in \mathcal{L}(R, k)$, the interface Φ_L induced by (R, Φ) on L. Now we compute a Φ_L -tour F_{L,Φ_L} in G[L] as follows.

First, we guess the children L_1, \ldots, L_p of L in the laminar family $\mathcal{L}(R, k)$. Then we guess the set $X \subseteq R[L]$ of edges that cross the cuts $\delta(L_1), \ldots, \delta(L_p)$. In other words, we guess all edges in $R(\mathcal{L}, k)$ that are contained in L, but not in any child of L. Moreover, for each child L_i with $i \in \{1, \ldots, p\}$, we guess the interface Φ_i induced by (R, Φ) on L_i . Because we consider the elements of the laminar family \mathcal{L} in an order of non-decreasing cardinality, we have already considered L_i before considering the current set L. Hence we have already computed some Φ_i -tour F_{L_i,Φ_i} for all $i \in \{1, \ldots, p\}$.

We now want to extend the union of these Φ_i -tours for all $i \in \{1, \ldots, p\}$ and the set X of edges crossing the boundaries of the children L_1, \ldots, L_p to a Φ_L -tour in G[L]. To this end we define $L_0 := L \setminus \bigcup_{i=1}^p L_i$. Then L_0, \ldots, L_p is a partition of L. We also guess the interface Φ_0 that (R, Φ) induces on L_0 . Then, by Lemma 22 applied to the graph G[L], the union of X and arbitrary Φ_i -tours in $G[L_i]$ for $i = \{0, \ldots, p\}$ is a Φ_L -tour in G[L]. Here we use that Φ_i is the interface induced by $(R[L], \Phi_L)$ on L_i for $i = \{0, \ldots, p\}$ (cf. Lemma 21 (iii)). Finally, we use the given algorithm \mathcal{B} to compute a β -approximation F_0 of a minimum length Φ_0 -tour in the subgraph $G[L_0]$ and combine X, F_0 , and the Φ_i -tours F_{L_i,Φ_i} for $i \in \{1, \ldots, p\}$ to a Φ_L -tour F_{L,Φ_L} .



Figure 6: Illustration of Algorithm 4. The dashed ellipses show the laminar family \mathcal{L} ; these sets are considered by the algorithm in an order of non-decreasing cardinality. Suppose we are considering $L \in \mathcal{L}(R, k)$; only subsets of L are shown in the figure. In the dynamic program we guess the children L_1, \ldots, L_p of L in the laminar family $\mathcal{L}(R, k)$. The sets L_1, \ldots, L_p are shown as blue ellipses with white interior. The light blue area shows the set $L_0 = L \setminus (L_1 \cup \cdots \cup L_p)$.

We also guess the set X of edges in $R[L] \cap (\delta(L_1) \cup \cdots \cup \delta(L_p))$; these are the thick blue edges. The thin gray edges are the edges in $R \cap \delta(L)$; these will be guessed only in a later step. However, we do guess the interface $\Phi_L = (I_L, T_L, C_L)$ that (R, Φ) induces on L, where I_L consists of all vertices in $I \cap L$ (shown with a thick red boundary) and all vertices in L that are endpoints of gray edges. Moreover, we guess the interfaces $\Phi_i = (I_i, T_i, C_i)$ that (R, Φ) induces on the sets L_i for $i \in \{0, \ldots, p\}$. The picture shows only the vertices in $I_0 \cup \cdots \cup I_p$; these are the vertices in L that are contained in the set I or are an endpoint of a thick blue or thin gray edge.

We compute a Φ_0 -tour in $G[L_0]$ and combine it with X and the Φ_i -tours in $G[L_i]$ for $i \in \{1, ..., p\}$ that we have computed in previous steps of the dynamic program. This yields a Φ_L -tour in G[L].

Algorithm 4: Algorithm computing a Φ -tour with the properties mentioned in Theorem 17

Let $\overline{\mathcal{L}} = \mathcal{L} \cup \{V\};$ for $L \in \overline{\mathcal{L}}$, in non-decreasing order of cardinality **do** for each interface $\Phi_L = (I_L, T_L, \mathcal{C}_L)$ of G[L] with $|I_L| \leq |I| + k$ do // No Φ_L -tour found yet. Let $F_{L,\Phi_L} \coloneqq \mathsf{Nil};$ // We set by convention: $\ell(Nil) = \infty$. for each subfamily $\{L_1, \ldots, L_p\} \subseteq \overline{\mathcal{L}}$ of disjoint proper subsets of L do Let $L_0 \coloneqq L \setminus \bigcup_{i=1}^p L_i$; for all $X \subseteq (\bigcup_{i=1}^p \delta(L_i)) \cap E[L]$ with $|X \cap \delta(L_i)| \leq k \ \forall i \in \{1, \dots, p\}$ do For $i \in \{0, \dots, p\}$, let $U_i \subseteq L_i$ be the set of all vertices in L_i that are an endpoint of some edge in X; Let $I_i \coloneqq (I_L \cap L_i) \cup U_i \quad \forall i \in \{0, \dots, p\};$ for all T_i, C_i such that $\Phi_i = (I_i, T_i, C_i)$ is an interface of $G[L_i]$ $(i \in \{0, \dots, p\})$ do Use Algorithm \mathcal{B} to find a Φ_0 -tour F_0 ; Let $F := X \stackrel{.}{\cup} F_0 \stackrel{.}{\cup} \bigcup_{i=1}^p F_{L_i, \Phi_i};$ if F is a Φ_L -tour and $\ell(F) \leq \ell(F_{L,\Phi_L})$ then $\left| \text{Set } F_{L,\Phi_L} = F; \right|$ return $F_{V,\Phi}$;

In what follows, we now provide a rigorous proof that Algorithm 4 implies Theorem 17 by leveraging the tools from Section 5.2.

5.4 Proof of Theorem 17

We start by showing that Algorithm 4 has indeed the claimed running time, before proving its correctness.

Running time

The running time of Algorithm 4 is dominated by the 5-fold nested for-loops. We first determine upper bounds on the number of iterations of each for-loop separately, whenever the algorithm reaches it.

- 1st for-loop: It goes over all sets in $\overline{\mathcal{L}}$. Because $\overline{\mathcal{L}}$ is a laminar family over V, it contains O(|V|) sets.
- 2nd for-loop: It goes over all interfaces $\Phi_L = (I_L, T_L, \mathcal{C}_L)$ of G[L] with $|I_L| \leq |I| + k$. There are no more than $(|L|+1)^{|I|+k} \leq (|V|+1)^{|I|+k}$ choices for choosing I_L . Moreover, there are at most $2^{|I_L|} \leq 2^{|I|+k}$ choices for $T_L \subseteq I_L$. Finally, the number of partitions \mathcal{C}_L of I_L can be upper bounded by $|I_L|^{|I_L|} \leq |V|^{|I|+k}$. Overall, the number of iterations of any run of the second for-loop is bounded by $|V|^{O(|I|+k)}$.
- 3rd for-loop: It iterates over subfamilies of $\overline{\mathcal{L}}$ of disjoint proper subsets of L. Because the sets are disjoint, such a family can have at most width $(\overline{\mathcal{L}}) \leq \text{width}(\mathcal{L}) + 1$ sets, and we can therefore bound the number of these subfamilies by $|\overline{\mathcal{L}}|^{\text{width}(\overline{\mathcal{L}})} = |V|^{O(\text{width}(\mathcal{L}))}$.
- 4th for-loop: It iterates over edge sets $X \subseteq (\bigcup_{i=1}^{p} \delta(L_i)) \cap E[L]$ with $|X \cap \delta(L_i)| \leq k$ for all $i \in \{1, \ldots, p\}$, and can be bounded as follows. Notice that $|X| \leq \sum_{i=1}^{p} |X \cap \delta(L_i)| \leq p \cdot k \leq \text{width}(\mathcal{L}) \cdot k$. Hence, there are at most $(|E| + 1)^{k \cdot \text{width}(\mathcal{L})} = |V|^{O(k \cdot \text{width}(\mathcal{L}))}$ options for X.
- 5th for-loop: This loop runs for all $i \in \{0, 1, ..., p\}$ over all interfaces $\Phi_i = (I_i, T_i, C_i)$ of $G[L_i]$, where L_i and I_i are fixed. The number of interfaces Φ_i for a fixed $i \in \{0, ..., p\}$ is thus bounded

by $(2|I_i|)^{|I_i|} \leq (2|V|)^{|I_i|}$ and, hence, the total number of combinations of such interfaces, and thus also on the number of iterations each time this for-loop is run, is bounded by

$$\prod_{i=0}^{p} (2|V|)^{|I_i|} = (2|V|)^{\sum_{i=0}^{p} |I_i|} .$$
(12)

Moreover, for $i \in \{1, ..., p\}$, we have $|I_i| \leq k + |I_L \cap L_i|$, which follows from the fact that each set I_i contains the elements of $I_L \cap L_i$ together with at most k endpoints of edges from X because $|X \cap \delta(L_i)| \leq k$. This implies

$$\sum_{i=1}^{p} |I_i| \le p \cdot k + |I_L| \le \operatorname{width}(\mathcal{L}) \cdot k + (|I| + k) = O(|I| + k \cdot \operatorname{width}(\mathcal{L})) \quad . \tag{13}$$

Similarly,

$$|I_0| \le |I_L| + k \cdot \operatorname{width}(\mathcal{L}) \le |I| + k + k \cdot \operatorname{width}(\mathcal{L}) = O(|I| + k \cdot \operatorname{width}(\mathcal{L})) \quad . \quad (14)$$

Combining (13) and (14) with (12), we can bound the number of iterations of the fifth for-loop by $|V|^{O(|I|+k \cdot \text{width}(\mathcal{L}))}$.

The most expensive single operation performed by Algorithm 4 is the call to Algorithm \mathcal{B} to find a Φ_0 -tour, which, by assumption, takes no more than $f_{\mathcal{B}}(G, |I_0|)$ time. Due to the bound on $|I_0| \leq |I| + k \cdot (\text{width}(\mathcal{L}) + 1)$ provided by (14), we have that the total running time is thus indeed bounded by $|V|^{O(|I|+k \cdot \text{width}(\mathcal{L}))} \cdot f_{\mathcal{B}}(G, |I| + k \cdot (\text{width}(\mathcal{L}) + 1))$.

Correctness

We now show that, whenever G admits a Φ -tour, then Algorithm 4 will find a Φ -tour $F_{V,\Phi}$ with the length guarantee claimed by Theorem 17. So let R be a Φ -tour. We have to show that $F_{V,\Phi}$ computed by the algorithm is a Φ -tour (instead of Nil) and that it satisfies

$$\ell(F_{V,\Phi}) \le \beta \cdot \ell(R \setminus R(\mathcal{L}, k)) + \ell(R(\mathcal{L}, k)) \quad . \tag{15}$$

We prove (15) by showing the following claim from smaller to larger sets $L \in \mathcal{L}(R, k) \cup \{V\}$.

Claim 24. Let $L \in \mathcal{L}(R,k) \cup \{V\}$. If L = V, let $\Phi_L = \Phi$. Otherwise, let $\Phi_L = (I_L, T_L, C_L)$ be the interface induced by (R, Φ) on L. Then Algorithm 4 computes a Φ_L -tour F_{L,Φ_L} such that

$$\ell(F_{L,\Phi_L}) \leq \beta \cdot \ell\left(R[L] \setminus R(\mathcal{L},k)\right) + \ell\left(R[L] \cap R(\mathcal{L},k)\right)$$

Observe that the claim immediately implies Theorem 17 by choosing L = V. Hence, it remains to prove the claim.

Proof of Claim 24. We prove the claim by induction from smaller to larger sets in $\mathcal{L}(R,k) \cup \{V\}$. Hence, let $L \in \mathcal{L}(R,k) \cup \{V\}$ and assume that the claim holds for sets in $\mathcal{L}(R,k) \cup \{V\}$ of strictly smaller cardinality than L. In particular, it holds for the children L_1, \ldots, L_p of L in the laminar family $\mathcal{L}(R,k) \cup \{V\}$. (Note that L may also not have any children.) Let $L_0 \coloneqq L \setminus \bigcup_{i=1}^p L_i$, and for $i \in \{0, \ldots, p\}$, let $\Phi_i = (I_i, T_i, \mathcal{C}_i)$ be the interface induced by (R, Φ) on L_i . By using Lemma 21 (iii) in the case $L \neq V$, we observe that Φ_i is also the interface induced by $(R[L], \Phi_L)$ on L_i . Let F_0 be a Φ_0 -tour obtained through Algorithm \mathcal{B} . Because L_0, L_1, \ldots, L_p partitions L, we have by Lemma 22 that

$$F \coloneqq X \stackrel{.}{\cup} F_0 \stackrel{.}{\cup} \bigcup_{i=1}^p F_{L_i,\Phi_i}$$

is a Φ_L -tour, where

$$X := R[L] \cap \bigcup_{i=1}^{p} \delta(L_i) \quad .$$
(16)

Before discussing that this Φ_L -tour F will indeed be considered by Algorithm 4, we bound its length. First, $\ell(F_0) \leq \beta \cdot \ell(R[L_0])$ because \mathcal{B} is a β -approximation algorithm and $R[L_0]$ is a Φ_0 -tour by Lemma 21 (ii). Moreover, for $i \in \{1, \ldots, p\}$ we apply the induction hypothesis to L_i and Φ_i , which is possible because $L_i \in \mathcal{L}(R, k)$ has strictly smaller cardinality than L. Hence, F_{L_i, Φ_i} is a Φ_i -tour and fulfills the length bound stated in the claim. We therefore get

$$\ell(F) = \ell(X) + \ell(F_0) + \sum_{i=1}^{p} \ell(F_{L_i,\Phi_i})$$

$$\leq \ell(X) + \beta \cdot \ell(R[L_0]) + \sum_{i=1}^{p} \left(\beta \cdot \ell(R[L_i] \setminus R(\mathcal{L}, k)) + \ell(R[L_i] \cap R(\mathcal{L}, k))\right)$$

$$= \beta \cdot \ell(R[L] \setminus R(\mathcal{L}, k)) + \ell(R[L] \cap R(\mathcal{L}, k)) \quad , \qquad (17)$$

where the last equality follows by observing that

$$R[L_0], R[L_1] \setminus R(\mathcal{L}, k), \dots, R[L_p] \setminus R(\mathcal{L}, k)$$
 partitions $R[L] \setminus R(\mathcal{L}, k)$, and
 $X, R[L_1] \cap R(\mathcal{L}, k), \dots, R[L_p] \cap R(\mathcal{L}, k)$ partitions $R[L] \cap R(\mathcal{L}, k)$.

Due to (17), the Φ_L -tour F fulfills the length bound of the claim. It remains to show that the Φ_L -tour F will indeed be considered by Algorithm 4. For this, we show that the following quantities are considered in the five nested for-loops:

 1^{st} for-loop: considers L,

2nd for-loop: considers the interface $\Phi_L = (I_L, T_L, C_L)$,

3rd for-loop: considers the children L_1, \ldots, L_p of L in the laminar family $\mathcal{L}(R, k) \cup \{V\}$,

 4^{th} for-loop: considers the set X,

5th for-loop: considers, for $i \in \{0, ..., p\}$, the interfaces Φ_i induced by $(R[L], \Phi_L)$ on L_i .

This run would indeed produce F. All that remains to be shown is that the above five quantities, to be considered within the five nested for-loops, fulfill the conditions set by the respective for-loops:

1st for-loop: Algorithm 4 considers all sets in \mathcal{L} and hence, also L.

- 2nd for-loop: If L = V, the interface Φ is obviously considered. Otherwise $\Phi_L = (I_L, T_L, C_L)$ is the interface induced by (R, Φ) on L, and we have $I_L = (I \cap L) \cup U$, where U is the set of vertices in L connected by an edge of R to a vertex in $V \setminus L$. As $L \in \mathcal{L}(R, k) \cup \{V\}$, we have $|\delta(L) \cap R| \leq k$, and hence $|U| \leq k$, which implies $|I_L| \leq |I| + k$ and shows that the interface Φ_L is considered in the second for-loop.
- 3rd for-loop: We have $\{L_1, \ldots, L_p\} \subseteq \overline{\mathcal{L}}$. Hence, the subfamily $\{L_1, \ldots, L_p\}$ will be considered in the third nested for-loop.
- 4th for-loop: The set X we want to consider is given by (16). This set clearly satisfies $X \subseteq (\bigcup_{i=1}^{p} \delta(L_i)) \cap E[L]$ because $R[L] \subseteq E[L]$. Moreover, for each $i \in \{1, \ldots, p\}$ we have

$$|X \cap \delta(L_i)| = |R[L] \cap \delta(L_i)| \le |R \cap \delta(L_i)| \le k ,$$

where the last inequality follows from $L_i \in \mathcal{L}(R, k)$. Hence, the set X will be considered during the fourth nested for-loop of the algorithm.

5th for-loop: For $i \in \{0, \ldots, p\}$ we have that $\Phi_i = (I_i, T_i, C_i)$ is the interface of $G[L_i]$ induced by $(R[L], \Phi_L)$ on L_i . Hence, $I_i = (I_L \cap L_i) \cup U_i$, where U_i are all vertices in L_i connected by an edge of R[L] to a vertex in $L \setminus L_i$. We have $R[L] \cap \delta(L_i) = X \cap \delta(L_i)$ by our choice of X as described in (16) and because $\{L_0, \ldots, L_p\}$ is a partition of L. Therefore, $I_i \coloneqq (I_L \cap L_i) \cup U_i$, as desired. Hence, the interfaces Φ_i for $i \in \{0, \ldots, p\}$ indeed get considered in the fifth nested for-loop of the algorithm.

As said, Claim 24 implies (15), completing the proof of Theorem 17.

We remark that Claim 24 can be slightly strengthened as follows. The statement also holds when replacing the induced interface $\Phi_L = (I_L, T_L, C_L)$ by any interface $\Phi'_L = (I_L, T_L, C'_L)$ where C'_L is a refinement of C_L . However, we do not need this for our purposes.

6 **Proof of the main theorem**

We finally prove that the Boosting Theorem (Theorem 10) implies Theorem 1. In fact, we prove a generalization, stated below as Theorem 25, which, for k = 2 and $\Phi = (I, T, C)$ with $I = T = \{s, t\}$ and $C = \{\{s, t\}\}$, yields Theorem 1.

Theorem 25. Let A be an α -approximation algorithm for TSP. Then, for any $\varepsilon > 0$ and any integer k, there is an $(\alpha + \varepsilon)$ -approximation algorithm for Φ -TSP restricted to instances with $|I_{\Phi}| \leq k$ that, for any instance (G, Φ) , calls A a strongly polynomial number of times on TSP instances defined on subgraphs of G, and performs further operations taking strongly polynomial time.

Proof. We obtain the result by repeatedly applying the Boosting Theorem, i.e., Theorem 10, to strengthen the 4-approximation algorithm for Φ -TSP guaranteed by Theorem 9 through the α -approximation algorithm for TSP which we assume to exist. Without loss of generality $\varepsilon \leq 1$. The Boosting Theorem will be repeated i_{max} many times with error parameter given by $\varepsilon' = \varepsilon/\alpha$, where

$$i_{\max} := \left\lceil \frac{4 - (\alpha + \varepsilon)}{\alpha - 1} \cdot \frac{8\alpha}{\varepsilon} \right\rceil$$

Notice that i_{\max} is constant, because both ε and α are fixed.

Let $\beta_0 := 4$ be the approximation factor for Φ -TSP before applying the Boosting Theorem. We assume $\alpha \leq 1.5 < \beta_0$ because Christofides' algorithm is a strongly polynomial 1.5-approximation algorithm for TSP [4, 25]. Let $i \in \{1, \ldots, i_{\text{max}}\}$. After *i* applications of the Boosting Theorem we obtain an algorithm \mathcal{B}_i for Φ -TSP with approximation ratio at most

$$\beta_i \coloneqq \max\left\{ (1+\varepsilon')\alpha, \ \beta_{i-1} - \frac{\varepsilon'}{8} \cdot (\beta_{i-1} - 1) \right\} = \max\left\{ \alpha + \varepsilon, \ \beta_{i-1} - \frac{\varepsilon}{8\alpha} \cdot (\beta_{i-1} - 1) \right\}$$

where we used $\varepsilon' = \varepsilon / \alpha$. We therefore have

$$\beta_i \le \max\left\{\alpha + \varepsilon, \ \beta_{i-1} - \frac{\varepsilon}{8\alpha}(\alpha - 1)\right\} \le \max\left\{\alpha + \varepsilon, \ \beta_0 - i \cdot \frac{\varepsilon}{8\alpha}(\alpha - 1)\right\},$$

where the last inequality follows by induction on i. Hence,

$$\beta_{i_{\max}} \leq \max\left\{\alpha + \varepsilon, \ 4 - i_{\max} \cdot \frac{\varepsilon}{8\alpha}(\alpha - 1)\right\}$$
$$= \max\left\{\alpha + \varepsilon, \ 4 - \left\lceil \frac{4 - (\alpha + \varepsilon)}{\alpha - 1} \cdot \frac{8\alpha}{\varepsilon} \right\rceil \cdot \frac{\varepsilon}{8\alpha}(\alpha - 1)\right\}$$
$$= \alpha + \varepsilon \ .$$

Moreover, we define real numbers $k_i > 0$ for $i \in \{0, \ldots, i_{\max}\}$ to upper bound the size of the interfaces we have to be able to handle after *i* boosting steps. We want the $\beta_{i_{\max}}$ -approximation algorithm $\mathcal{B}_{i_{\max}}$, obtained after i_{\max} many applications of the Boosting Theorem, to handle interfaces of size $k_{i_{\max}} := k$. Because $\mathcal{B}_{i_{\max}}$ was obtained by applying the Boosting Theorem to $\mathcal{B}_{i_{\max}-1}$, we obtain that $\mathcal{B}_{i_{\max}-1}$ needs to handle interfaces of size bounded by $k_{i_{\max}-1} := \frac{9}{\varepsilon'} \cdot k_{i_{\max}}$. Repeating this reasoning, we obtain upper bounds k_i on the size of the interfaces that we have to handle with \mathcal{B}_i that satisfy

$$k_i := \frac{9}{\varepsilon'} \cdot k_{i+1} \quad \forall i \in \{i_{\max} - 1, \dots, 1, 0\} ,$$

which implies

$$k_i = k \cdot \left(\frac{9}{\varepsilon'}\right)^{i_{\max}-i} \quad \forall i \in \{0, \dots, i_{\max}\}$$

Notice that because i_{\max} , k, and ε' are constant, also k_0 is constant.

For $i = i_{\text{max}}$, the following claim implies Theorem 25 because $\beta_{i_{\text{max}}} = \alpha + \varepsilon$ and i_{max} , k_0 , and ε' are constant and \mathcal{B}_0 is a strongly polynomial algorithm.

Claim 26. Let c > 0 be the hidden constant in the big-O notation in the runtime bound in Theorem 10. Let $i \in \{0, ..., i_{max}\}$ and let A be the given α -approximation algorithm for TSP. Then there is a β_i -approximation algorithm \mathcal{B}_i for Φ -TSP that, for every weighted graph G, runs in time at most

$$f_i(G) \coloneqq |V|^{i \cdot c \cdot \frac{\kappa_0}{\varepsilon'}} \cdot \left(i \cdot f_{\mathcal{A}}(G) + f_{\mathcal{B}_0}(G)\right)$$
(18)

on any instance (G', Φ) , where G' is a subgraph of G and $|I_{\Phi}| \leq k_i$.

We prove the claim by induction on *i*. By Theorem 9 we have a strongly polynomial β_0 -approximation algorithm \mathcal{B}_0 for Φ -TSP, implying the claim for i = 0.

Now let $i \in \{1, \ldots, i_{\max}\}$. By our induction hypothesis, there exists a β_{i-1} -approximation algorithm \mathcal{B}_{i-1} that runs in time $f_{i-1}(G)$ on every weighted graph G and every interface Φ of G with $|I_{\Phi}| \leq k_{i-1}$. Applying Theorem 10 to the algorithms \mathcal{A} and \mathcal{B}_{i-1} then yields a β_i -approximation algorithm \mathcal{B}_i for Φ -TSP that runs on every graph G and every interface Φ with $|I_{\Phi}| \leq k_i$ in time at most

$$|V|^{c \cdot \frac{k_i}{\varepsilon'}} \cdot \left(f_{\mathcal{A}}(G) + f_{\mathcal{B}_{i-1}} \left(G, \frac{9 \cdot k_i}{\varepsilon'} \right) \right)$$

$$\leq |V|^{c \cdot \frac{k_0}{\varepsilon'}} \cdot \left(f_{\mathcal{A}}(G) + f_{\mathcal{B}_{i-1}} \left(G, k_{i-1} \right) \right)$$

$$\leq |V|^{c \cdot \frac{k_0}{\varepsilon'}} \cdot \left(f_{\mathcal{A}}(G) + f_{i-1}(G) \right)$$

$$= |V|^{c \cdot \frac{k_0}{\varepsilon'}} \cdot f_{\mathcal{A}}(G) + |V|^{i \cdot c \cdot \frac{k_0}{\varepsilon'}} \cdot \left((i-1) \cdot f_{\mathcal{A}}(G) + f_{\mathcal{B}_0}(G) \right)$$

$$\leq f_i(G) .$$

7 Strongly polynomial 4-approximation algorithm for Φ -TSP

Theorem 9. Φ -*TSP admits a strongly polynomial* 4-*approximation algorithm.*

Proof. Let $\Phi = (I, T, C)$ be an interface of G = (V, E). By Lemma 8, we can assume that Φ is feasible. The main component of our algorithm is to obtain a strongly polynomial 2-approximation algorithm for the problem of finding a set (not a multi-set) $F \subseteq E$ of minimum length $\ell(F)$ that satisfies the following three conditions:

- (i) (V, F)/I is connected;
- (ii) (V, F) connects all vertices within any $C \in C$;
- (iii) each connected component of (V, F) contains an even number of vertices in T.

We will achieve this through an application of Jain's iterative rounding method for the Generalized Steiner Network Problem [13] together with the elegant framework of Frank and Tardos [7] to transform certain polynomial-time algorithms into strongly polynomial ones.

Before we discuss the details of Jain's method in our setting together with the framework of Frank and Tardos, we first assume that we can indeed find in strongly polynomial time a set $F \subseteq E$ fulfilling (i), (ii), and (iii) of length no larger than twice the length of a shortest edge set fulfilling these three conditions. Because a shortest Φ -tour OPT must fulfill these conditions, and removing parallel edges does not destroy them, there is a subset of OPT that contains no parallel edges and satisifies (i), (ii), and (iii). Therefore, $\ell(F) \leq 2\ell(\text{OPT})$.

Due to property (iii), the set F contains a T-join $J \subseteq F$, which we can find in linear time through standard techniques. We then return $F \cup (F \setminus J)$, which is indeed a Φ -tour and satisfies

$$\ell(F \cup (F \setminus J)) \le 2\ell(F) \le 4\ell(\text{OPT})$$
,

as desired. It remains to show how to obtain a strongly polynomial 2-approximation algorithm for finding a shortest edge set fulfilling (i), (ii), and (iii). We start by showing how an application of Jain's iterative rounding method leads to a polynomial-time, but not necessarily strongly polynomial, 2-approximation algorithm.

To this end, observe that a set $F \subseteq E$ satisfies (i), (ii), and (iii) if and only if

$$|F \cap \delta(S)| \ge f(S) \qquad \forall S \subseteq V \quad , \tag{19}$$

where the function $f: 2^V \to \{0, 1\}$ is defined as follows. For $S \subsetneq V$ with $S \neq \emptyset$, we set f(S) = 1 if at least one of the following three properties holds:

- (a) $S \cap I = \emptyset$;
- (b) $\exists C \in \mathcal{C} \text{ s.t. } S \cap C \neq \emptyset \text{ and } C \setminus S \neq \emptyset;$
- (c) $|S \cap T|$ is odd.

Otherwise we set f(S) = 0. (In particular, $f(\emptyset) = f(V) = 0$.) Indeed, the properties (a), (b), and (c) are just reformulations of (i), (ii), and (iii), respectively.

Jain's technique [13] leads to a polynomial-time 2-approximation algorithm for finding a shortest edge set F satisfying (19) if, first, the function f is *weakly supermodular*, which means

$$f(X) + f(Y) \le \max\left\{f(X \cup Y) + f(X \cap Y), f(X \setminus Y) + f(Y \setminus X)\right\} \qquad \forall X, Y \subseteq V ,$$
(20)

and, second, one can separate over the polytope

$$P = \left\{ x \in [0,1]^E : x(\delta(S)) \ge f(S) \ \forall S \subseteq V \right\}$$
(21)

in polynomial time.

We start by showing (20). Notice that (20) clearly holds if $X \subseteq Y$, because in this case we have $\{X, Y\} = \{X \cup Y, X \cap Y\}$. Hence, in what follows, we always assume that $X \setminus Y \neq \emptyset$ and $Y \setminus X \neq \emptyset$.

Let f_a , f_b , and f_c be the functions from 2^V to $\{0, 1\}$ that take a value of 1 precisely for sets $S \subsetneq V, S \neq \emptyset$ that satisfy (a), (b), or (c), respectively. Hence, $f(S) = \max\{f_a(S), f_b(S), f_c(S)\}$. First, one can observe that each of the functions f_a , f_b , and f_c is weakly supermodular. Consider first f_a and let $X, Y \subseteq V$ with $X \setminus Y \neq \emptyset$ and $Y \setminus X \neq \emptyset$. If $f_a(X) = 1$ then $f_a(X \setminus Y) = 1$. Similarly, if $f_a(Y) = 1$, then $f_a(Y \setminus X) = 1$. Hence, f_a satisfies (20). The function f_b corresponds to pairwise connectivity requirements and, as shown in [13], is therefore weakly supermodular. The function f_c is easily seen to be a so-called *proper* function, which means that $f_c(V) = 0$, f_c is symmetric, and $f_c(S_1 \cup S_2) \leq \max\{f_c(S_1), f_c(S_2)\}$ for any pair of disjoint sets $S_1, S_2 \subseteq V$. Finally, it is well-known that any proper function is weakly supermodular (see [9]).

We say that a set $S \subsetneq V$ with $S \neq \emptyset$ is of type (a), (b), or (c), if it satisfies (a), (b), or (c), respectively. Because each of the functions f_a , f_b , and f_c is weakly supermodular, the inequality (20) holds whenever the sets X and Y are of the same type, or if X or Y is none of the three types. Hence, it remains to consider sets X and Y of two different types among the types (a), (b), and (c). Let $S_a, S_b, S_c \subseteq V$ be sets of type (a), (b), and (c), respectively. Thus, we need to show that (20) holds for the three cases where (X, Y) is either $(S_a, S_b), (S_a, S_c)$, or (S_b, S_c) . Moreover, let $C \in C$ be a set such that $S_b \cap C \neq \emptyset$ and $C \setminus S_b \neq \emptyset$, which exists because S_b is of type (b).

We start by considering the case $(X, Y) = (S_a, S_b)$. As discussed, we assume that $S_a \not\subseteq S_b$ and $S_b \not\subseteq S_a$; for otherwise, (20) holds trivially. Notice that in this case we have

$$2 = f(S_a) + f(S_b) \le f(S_a \setminus S_b) + f(S_b \setminus S_a) = 2 ,$$

because $(S_a \setminus S_b) \cap I \subseteq S_a \cap I = \emptyset$, as well as $(S_b \setminus S_a) \cap I = S_b \cap I$ and $C \subseteq I$. Hence, $S_a \setminus S_b$ is of type (a) and $S_b \setminus S_a$ is of type (b).

Consider now the case $(X, Y) = (S_a, S_c)$. Here, we have

$$2 = f(S_a) + f(S_c) \le f(S_a \setminus S_c) + f(S_c \setminus S_a) = 2$$

because $(S_a \setminus S_c) \cap I \subseteq S_a \cap I = \emptyset$, implying that $S_a \setminus S_c$ is of type (a), and $|(S_c \setminus S_a) \cap T| = |S_c \cap T|$ due to $S_a \cap T \subseteq S_a \cap I = \emptyset$, which implies that $S_c \setminus S_a$ is of type (c).

It remains to consider the case $(X, Y) = (S_b, S_c)$. We first observe that

$$\max\left\{f(S_b \setminus S_c), f(S_b \cup S_c)\right\} \ge 1 \quad \text{, and} \tag{22}$$

$$\max\left\{f(S_b \cap S_c), f(S_c \setminus S_b)\right\} \ge 1 \quad , \tag{23}$$

due to the following. Inequality (22) holds because $S_b \cup S_c$ can be partitioned into S_c and $S_b \setminus S_c$. Because $|S_c \cap T|$ is odd, either $S_b \cup S_c$ or $S_b \setminus S_c$ must also have an odd intersection with T and is thus of type (c). Inequality (23) follows from an analogous reasoning using the partition of S_c into $S_b \cap S_c$ and $S_c \setminus S_b$. Moreover, we have

$$\max\left\{f(S_b \setminus S_c), f(S_b \cap S_c)\right\} \ge 1 \quad \text{, and} \tag{24}$$

$$\max\left\{f(S_b \cup S_c), f(S_c \setminus S_b)\right\} \ge 1 \quad , \tag{25}$$

because S_b is of type (b), i.e., $S_b \cap C \neq \emptyset$ and $C \setminus S_b \neq \emptyset$. Indeed, even without any assumptions on $S_c \subseteq V$, we have that either $S_b \setminus S_c$ or $S_b \cap S_c$ is also of type (b). The same holds for either $S_b \cup S_c$ or $S_c \setminus S_b$. Among the four expressions $f(S_b \cup S_c)$, $f(S_b \cap S_c)$, $f(S_b \setminus S_c)$, and $f(S_c \setminus S_b)$, consider any one of minimum value and sum up the two inequalities among (22), (23), (24), and (25) containing that expression. This gives the desired result. For example, if $f(S_b \setminus S_c)$ achieves minimum value among the four, then (22) implies $f(S_b \cup S_c) = 1$ and (24) implies $f(S_b \cap S_c) = 1$. Hence,

$$2 = f(S_b) + f(S_c) \le f(S_b \cup S_c) + f(S_b \cap S_c) = 2 ,$$

as desired. This completes the proof that f is weakly supermodular.

To apply Jain's method, it remains to show that we can separate over P, and we will in fact give a strongly polynomial algorithm. Given $y \in [0,1]^E$, we will either show that all constraints $y(\delta(S)) \ge f(S)$ for $S \subseteq V$ are fulfilled or return one of these constraints that is violated. Notice that, because $y \ge 0$, a

constraint $y(\delta(S)) \ge f(S)$ can only be violated if f(S) = 1, i.e., S is either of type (a), (b), (c). Hence, we can check these constraints for each type separately.

Whether there is a violated constraints of type (a) reduces to finding a minimizer of

$$\min \left\{ y(\delta(S)) : S \subseteq V \text{ with } S \cap I = \emptyset \right\}$$

This can be solved through a global minimum cut algorithm applied to the graph G/I with edge weights y. Indeed, this either leads to a cut S with $S \cap I = \emptyset$ as desired or one where $I \subseteq S$, in which we can replace S by $V \setminus S$.

To check whether there is a violated constraint of type (b) reduces to

$$\min \left\{ y(\delta(S)) : \exists C \in \mathcal{C} \text{ with } S \cap C \neq \emptyset \text{ and } C \setminus S \neq \emptyset \right\}$$

This can be solved by performing the following for all $C \in C$ with $|C| \geq 2$. Number the vertices in C arbitrarily $C = \{c_1, \ldots, c_k\}$, and solve a minimum $c_i c_{i+1}$ cut problem in G with edge weights y for each $i \in \{1, \ldots, k-1\}$. If any of these *s*-*t* cut problems leads to a cut of value strictly smaller than 1, then the minimizing cut corresponds to a violated constraint. Otherwise, there is no violated constraints of type $y(\delta(S)) \geq f(S)$ for any set S of type (b).

Finally, checking whether there is a violated constraint of type (c) reduces to

$$\min \left\{ y(\delta(S)) : S \subseteq V, |S \cap T| \text{ is odd} \right\}$$

This is a minimum weight T-cut problem, for which strongly polynomial algorithms are well known (see, e.g., [21]).

In summary, the separation problem over P can be solved in strongly polynomial time, and we can therefore apply Jain's technique as claimed.

It remains to show that the overall algorithm can be transformed into a strongly polynomial one. This is a consequence of the framework of Frank and Tardos [7] (see also [11, 5, 13] for similar applications). More precisely, the only step that is not strongly polynomial in Jain's iterative rounding method is solving linear programs on faces of P with objective function given by ℓ . Notice that the coefficients in the constraints describing P are all 0 or 1. Hence, they have small encoding length. For such cases, Frank and Tardos [7] show how ℓ can be replaced (in strongly polynomial time) by another objective $\hat{\ell}$ of encoding length polynomial in the dimension |E| of the problem such that the set of optimal solutions over any polytope in |E|dimensions with constraints of small encoding length is the same for the two objectives ℓ and $\hat{\ell}$. Hence, one can find an optimal linear programming solution with respect to $\hat{\ell}$ instead of ℓ , whenever a linear program has to be solved in Jain's procedure.

8 Conclusions and open problems

We showed that given a polynomial-time α -approximation algorithm for TSP we can obtain a polynomialtime ($\alpha + \varepsilon$)-approximation algorithm for Path TSP. Feige and Singh [6] proved a similar kind of result for the asymmetric traveling salesman problem (ATSP): given a polynomial-time α -approximation algorithm for ATSP, there is a polynomial-time ($2\alpha + \varepsilon$)-approximation algorithm for its path version. A natural question is whether our techniques can be used to improve on their result and avoid losing a factor of two in the approximation ratio.

For the Asymmetric Path TSP, the relatively simple dynamic program (sketched in Section 3.1) still works and could be used to reduce to the case where the distance d(s,t) from s to t is not much more than $1/2 \cdot \ell(\text{OPT})$. (We get 1/2 instead of 1/3 because a cut can contain two forward edges and one backward edge, and backward edges can belong to many cuts.) To make further progress, we might again try to guess edges

also in cuts in which OPT contains a larger, but constant number of edges. However, even if the distance d(s,t) is very small, the distance d(t,s) from t to s could be large. In this case we do not know how to reduce to ATSP or guess edges of significant length via dynamic programming. Another obstacle is the following: Our approach for reducing Path TSP to TSP required a constant-factor approximation algorithm for Φ -TSP. Thus, for the asymmetric case one would probably need a suitable constant-factor approximation algorithm for a directed version of Φ -TSP, and we do not know how to obtain this.

A special case of Φ -TSP that is more general than Path TSP is the *T*-tour problem. Here $I_{\Phi} = T_{\Phi}$ is the given set *T* and $C_{\Phi} = \{T\}$. None of the recent improvements for Path TSP seems to extend to general *T*-tours beyond constant |T|, so Sebő's ⁸/5-approximation [22] remains the best that we know. Another question is how well Φ -TSP can be approximated in general. We showed an approximation ratio of four, but a better ratio might be possible.

References

- [1] H.-C. An, R. Kleinberg, and D. B. Shmoys. Improving Christofides' algorithm for the *s*-*t* path TSP. *Journal of the ACM*, 62(5):34:1–34:28, 2015. Short version appeared in STOC 2012.
- [2] A. Blum, S. Chawla, D. R. Karger, T. Lane, A. Meyerson, and M. Minkoff. Approximation algorithms for orienteering and discounted-reward TSP. *SIAM Journal on Computing*, 37(2):653–670, 2007. Short version appeared in FOCS 2003.
- [3] J. Cheriyan, Z. Friggstad, and Z. Gao. Approximating minimum-cost connected *T*-joins. *Algorithmica*, 72:126–147, 2015. Short version appeared in APPROX/RANDOM 2012.
- [4] N. Christofides. Worst-case analysis of a new heuristic for the Travelling Salesman Problem. Technical Report 388, Graduate School of Industrial Administration, Carnegie Mellon University, 1976.
- [5] F. Eisenbrand. Integer programming and algorithmic geometry of numbers. In M. Jünger, T. M. Liebling, D. Naddef, G. L. Nemhauser, and W. R. Pulleyblank, editors, 50 Years of Integer Programming 1958–2008, pages 505–559. Springer, 2010.
- [6] U. Feige and M. Singh. Improved approximation algorithms for traveling salesperson tours and paths in directed graphs. In *Proceedings of the 10th International Workshop on Approximation Algorithms* for Combinatorial Optimization Problems (APPROX), pages 104–118, 2007.
- [7] A. Frank and É. Tardos. An application of simultaneous diophantine approximation in combinatorial optimization. *Combinatorica*, 7(1):49–65, 1987.
- [8] Z. Gao. An LP-based $\frac{3}{2}$ -approximation algorithm for the *s*-*t* path graph traveling salesman problem. *Operations Research Letters*, 41:615–617, 2013.
- [9] M. X. Goemans, A. V. Goldberg, S. Plotkin, D. B. Shmoys, É. Tardos, and D. P. Williamson. Improved approximation algorithms for network design problems. In *Proceedings of the 5th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 223–232, 1994.
- [10] C. Gottschalk and J. Vygen. Better s-t-tours by Gao trees. Mathematical Programming B, 172:191–207, 2018. Short version appeared at IPCO 2016.
- [11] M. Grötschel, L. Lovász, and A. Schrijver. Geometric Algorithms and Combinatorial Optimization, volume 2 of Algorithms and Combinatorics. Springer, second corrected edition, 1993.

- [12] J.A. Hoogeveen. Analysis of Christofides' heuristic: Some paths are more difficult than cycles. Operations Research Letters, 10(5):291–295, 1991.
- [13] K. Jain. A factor 2 approximation algorithm for the generalized Steiner Network Problem. *Combina-torica*, 21:39–60, 2001. Short version appeared in FOCS 1998.
- [14] M. Karpinski, M. Lampis, and R. Schmied. New inapproximability bounds for TSP. Journal of Computer and System Sciences, 81:1665–1677, 2015. Short version appeared in ISAAC 2013.
- [15] A. V. Karzanov. How to tidy up a symmetric set-system by use of uncrossing operations. *Theoretical Computer Science*, 157:215–225, 1996.
- [16] T. Mömke and O. Svensson. Removing and adding edges for the Traveling Salesman Problem. *Journal of the ACM*, 63(1):2:1–2:28, 2016. Short version appeared in FOCS 2011.
- [17] M. Mucha. $\frac{13}{9}$ -approximation for graphic TSP. *Theory of Computing Systems*, 55(4):640–657, 2014. Short version appeared in STACS 2012.
- [18] M. Nägele and R. Zenklusen. A new dynamic programming approach for spanning trees with chain constraints and beyond. In *Proceedings of the 30th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1550–1569, 2019.
- [19] S. Oveis Gharan, A. Saberi, and M. Singh. A randomized rounding approach to the Traveling Salesman Problem. In *Proceedings of the 52nd Annual IEEE Symposium on Foundations of Computer Science* (FOCS), pages 550–559, 2011.
- [20] C. H. Papadimitriou and M. Yannakakis. The Traveling Salesman Problem with distances one and two. *Mathematics of Operations Research*, 18(1):1–12, 1993.
- [21] A. Schrijver. Combinatorial Optimization, Polyhedra and Efficiency. Springer, 2003.
- [22] A. Sebő. Eight-fifth approximation for the Path TSP. In *Proceedings of 16th International Conference* on Integer Programming and Combinatorial Optimization (IPCO), pages 263–373, 2013.
- [23] A. Sebő and A. van Zuylen. The salesman's improved paths through forests. *Journal of the ACM*, 66(4):28:1–28:16, 2019. Short version appeared in FOCS 2016.
- [24] A. Sebő and J. Vygen. Shorter tours by nicer ears: 7/5-approximation for the graph-TSP, 3/2 for the path version, and 4/3 for two-edge-connected subgraphs. *Combinatorica*, 34(5):597–629, 2014.
- [25] A. I. Serdjukov. Some extremal bypasses in graphs [in Russian]. Upravlyaemye Sistemy, 17:76–79, 1978.
- [26] V. Traub and J. Vygen. Beating the integrality ratio for s-t-tours in graphs. In Proceedings of 59th Annual IEEE Symposium on Foundations of Computer Science (FOCS), pages 766–777, 2018.
- [27] V. Traub and J. Vygen. Approaching $\frac{3}{2}$ for the *s*-*t* path TSP. *Journal of the ACM*, 66(2):14:1–14:17, 2019. Short version appeared in SODA 2018.
- [28] J. Vygen. Reassembling trees for the traveling salesman. SIAM Journal on Discrete Mathematics, 30(2):875–894, 2016.
- [29] Z. Xu and B Rodrigues. A 3/2-approximation algorithm for the multiple TSP with a fixed number of depots. *INFORMS Journal on Computing*, 27(4):636–645, 2015.

[30] R. Zenklusen. A 1.5-approximation for Path TSP. In *Proceedings of 30th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1539–1549, 2019.