PROJECTIVE-UMBILIC POINTS OF CIRCULAR REAL HYPERSURFACES IN \mathbb{C}^2

DAVID E. BARRETT AND DUSTY E. GRUNDMEIER

ABSTRACT. We show that the boundary of any bounded strongly pseudoconvex complete circular domain in \mathbb{C}^2 must contain points that are exceptionally tangent to a projective image of the unit sphere.

1. BACKGROUND

A *vertex* of a smooth plane curve may be viewed as a point at which there is a circle exceptionally tangent to to the curve; that is, there is a circle with fourth-order (or better) contact with the curve at the vertex in contrast with with the situation of a non-vertex at which third-order contact is the best possible. (What we are calling the order of contact here is also known as the "point-contact order" [Rut,§5.1.1]. An alternate convention – used for example in the study of jets of functions – reduces the orders by one.)

The famous four-vertex theorem ([Muk], [Kne], [Oss]) says the every smooth Jordan curve in the plane has at least four vertices. There are corresponding results giving lower bounds for the size of the set of *affine vertices* where a curve is exceptionally tangent to a conic [Muk], for the set of *umbilic points* on a smooth non-toric compact surface in \mathbb{R}^3 where the surface is exceptionally tangent to a ball (see for example [Ber, pp. 389-390], noting the unresolved Carathéodory conjecture), for the set of points where an orientation-preserving diffeomorphism of the unit circle is exceptionally tangent to a holomorphic automorphism of the unit disk ([Ghy], [OvTa]), and for the set of *CR-umbilic points* on the boundary of a real ellipsoid or bounded complete circular domain in \mathbb{C}^2 that are exceptionally tangent to a local biholomorphic image of the unit sphere ([HuJi], [EbSo]; note also the counter-example in [ESZ]). (The order of contact at stake in the last batch of results is seven, and the points in question are those where a certain sixth-order tensor (due to Cartan [Car1], [Car2]) vanishes.)

In the current work we again consider real hypersurfaces in \mathbb{C}^2 or the projective space \mathbb{CP}^2 , looking at orders of contact with projective images of the unit sphere (equivalently, with \mathbb{C} -affine images of the unit sphere or the Heisenberg hypersurface {Im $z_2 = |z_1|^2$ }). The special points are the *projective-umbilic points* at which third-order contact is possible.

Due to the discrepancy in the collection of allowable maps on the one hand and the order of contact on the other hand, CR-umbilic points need not be projective-umbilic, nor vice versa (see Example 6 below.)

Date: March 6, 2020.

²⁰¹⁰ Mathematics Subject Classification: 32V10.

The first author was supported in part by NSF grant number DMS-1500142.

We show below that boundary of any bounded strongly pseudoconvex circular domain in \mathbb{C}^2 must contain projective-umbilic points.

2. A Beltrami-style tensor

The following tensorial object will prove useful: for S a smooth real strongly pseudoconvex hypersurface in \mathbb{C}^2 with defining function r we set

(2.1)
$$\mathscr{B}_{S} \stackrel{\text{def}}{=} -\frac{\det \begin{pmatrix} 0 & r_{1} & r_{2} \\ r_{1} & r_{11} & r_{21} \\ r_{2} & r_{12} & r_{22} \end{pmatrix}}{\det \begin{pmatrix} 0 & r_{1} & r_{2} \\ r_{\overline{1}} & r_{1\overline{1}} & r_{2\overline{1}} \\ r_{\overline{2}} & r_{1\overline{2}} & r_{2\overline{2}} \end{pmatrix}} \cdot \frac{dz_{1} \wedge dz_{2}}{d\overline{z_{1}} \wedge d\overline{z_{2}}}$$

on *S*, where the subscripts denote differentiation. (The non-vanishing of the denominator here is well-known to be equivalent to the strong pseudoconvexity of *S*.) The last factor above is to indicate that this object is to be viewed as a section of the product of the canonical bundle of (2,0)-forms with the conjugate-inverse of that bundle. We will refer to such objects as Beltrami differentials. (In a one-variable setting this reduces to the reciprocal of the Beltrami differentials $\mu(z) \frac{d\overline{z}}{dz}$ used in particular in the study of quasi-conformal mappings, as for example found in [Leh]. See also §5 below.)

Proposition 1. (2.2a) \mathscr{B}_S does not depend on the choice of defining function *r*. (2.2b) If ψ is an automorphism of \mathbb{CP}^2 then $\mathscr{B}_S = \psi^* \left(\mathscr{B}_{\psi(S)} \right)$ (where defined).

Proof. For (2.2a), check that if *r* is replaced by $\eta \cdot r$ with η non-vanishing then both the numerator and denominator above pick up a factor of $\eta(p)^3$ at $p \in S$.

For (2.2b), first recall that automorphisms of \mathbb{CP}^2 have the form

$$(z_1, z_2) \mapsto \left(\frac{D + Ez_1 + Fz_2}{A + Bz_1 + Cz_2}, \frac{G + Hz_1 + Iz_2}{A + Bz_1 + Cz_2}\right).$$

with $\begin{pmatrix} A & B & C \\ D & E & F \\ G & H & I \end{pmatrix}$ invertible. The group of these automorphisms is generated by the invertible affine

transformations together with the particular transformation

$$(z_1, z_2) \mapsto \left(\frac{1}{z_1}, \frac{z_2}{z_1}\right);$$

the transformation law can be verified by straightforward computation in either case.

In view of (2.2b), the construction of \mathscr{B}_S also makes sense on a strongly pseudoconvex real hypersurface in \mathbb{CP}^2 – see [Bar, §5.3] for a more directly projective approach to the construction and the transformation law.

Proposition 2. If
$$S_{sph}$$
 is the unit sphere $\{(z_1, z_2) \in \mathbb{C}^2 : |z_1|^2 + |z_2|^2 = 1\}$ then $\mathscr{B}_{S_{sph}} = 0 \cdot \frac{dz_1 \wedge dz_2}{d\overline{z_1} \wedge d\overline{z_2}}$.

Proof. This follows by direct computation with $r(z_1, z_2) = |z_1|^2 + |z_2|^2 - 1$.

Corollary 3. If S_{Heis} is the Heisenberg hypersurface $\{(z_1, z_2) \in \mathbb{C}^2 : \text{Im } z_2 = |z_1|^2\}$ then $\mathscr{B}_{S_{\text{Heis}}} = 0 \cdot \frac{dz_1 \wedge dz_2}{d\overline{z_1} \wedge d\overline{z_2}}$.

Proof. This can be handled either by direct computation as above or by applying (2.2b) to the projective automorphism

$$\begin{split} \psi \ : \ S_{\text{Heis}} & \rightarrow S_{\text{sph}} \\ & \left(z_1, z_2 \right) \mapsto \left(\frac{2z_1}{i + z_2}, \frac{i - z_2}{i + z_2} \right) \end{split}$$

In the other direction we have the following result.

Theorem 4. ([Jen], [DeTr], [Bol]).

 \mathscr{B}_{S} vanishes identically if and only if S is locally a projective image of the unit sphere S_{sph} (or equivalently, of the Heisenberg hypersurface S_{Heis}).

Proposition 5. Let S be a smooth strongly pseudoconvex real hypersurface in \mathbb{CP}^2 and let p be a point in S. Then there is an automorphism of \mathbb{CP}^2 moving p to $0 \in \mathbb{C}^2$ so that the transformed S takes the form

(2.3)
$$\operatorname{Im} z_2 = |z_1|^2 + \beta \operatorname{Re} z_1^2 + O\left(\|(z_1, \operatorname{Re} z_2)\|^3\right)$$

near 0 *with uniquely-determined* $\beta \in [0, \infty)$ *.*

Proof. See [Bar, Prop. 5] and the following discussion.

(For projective normalization of higher-order terms see [Ham].) For a hypersurface *S* of the form (2.3) we have (by direct calculation) that

$$\mathscr{B}_{S}(0) = \beta \, \frac{dz_{1} \wedge dz_{2}}{d\overline{z_{1}} \wedge d\overline{z_{2}}};$$

moreover the order of contact between S and S_{Heis} is $\begin{cases} \geq 3 & \text{if } \beta = 0 \\ 2 & \text{if } \beta \neq 0. \end{cases}$ Thus the *projective-umbilic*

points from the end of §1 are precisely the points where \mathscr{B}_S vanishes.

Example 6. The smooth portion $\{(z_1, z_2) : |z_1|^p + |z_2|^p = 1, z_1z_2 \neq 0\}$ of the boundary of the unit L^p ball in \mathbb{C}^2 is locally CR-equivalent to the sphere, using a branch of $(z_1^{2/p}, z_2^{2/p})$, but contains no projective-umbilic points when $p \neq 2$ since

$$\mathscr{B}_{S} = \frac{2-p}{p} \frac{\overline{z_{1}z_{2}}}{z_{1}z_{2}} \frac{dz_{1} \wedge dz_{2}}{d\overline{z_{1}} \wedge d\overline{z_{2}}}$$

in this case.

3

3. MAIN RESULT

Theorem 7. If S is the boundary of a bounded strongly pseudoconvex complete circular domain in \mathbb{C}^2 then S contains at least one circle of projective-umbilic points.

Proof. The complete circularity condition implies in particular that S intersects each complex line L through the origin in one circle C_L ; since S is strongly pseudoconvex, L cannot be completely tangent to S along C_L , so in fact L intersects S transversely along C_L .

It follows that we may write

(3.1)
$$S \setminus \{z_2 = 0\} = \left\{ (z_1, z_2) : z_2 \neq 0, z_2 \overline{z_2} = e^{\rho(z_1/z_2)} \right\}$$

where ρ is a smooth \mathbb{R} -valued function on \mathbb{C} . Similarly we may write

(3.2)
$$S \setminus \{z_1 = 0\} = \left\{ (z_1, z_2) : z_1 \neq 0, z_1 \overline{z_1} = e^{\widetilde{\rho}(z_2/z_1)} \right\}.$$

Setting $\zeta = z_1/z_2$ we have from (3.1) that

$$\mathscr{B}_{S} = b_{S}(\zeta) \frac{\overline{z_{2}}^{2}}{z_{2}^{2}} \frac{dz_{1} \wedge dz_{2}}{d\overline{z_{1}} \wedge d\overline{z_{2}}}$$

where $b_S(\zeta) = -\frac{\rho_{\zeta\zeta} - \rho_{\zeta}^2}{\rho_{\zeta\overline{\zeta}}}$ is C-valued (but not holomorphic). The strong pseudoconvexity of S guarantees that the denominator $\rho_{\zeta\overline{\zeta}}$ is non-vanishing for $\zeta \in \mathbb{C}$.

Using (3.2) instead we have the alternate formula

$$\mathscr{B}_{S} = \widetilde{b}_{S}(1/\zeta) \ \frac{\overline{z_{1}}^{2}}{z_{1}^{2}} \frac{dz_{1} \wedge dz_{2}}{d\overline{z_{1}} \wedge d\overline{z_{2}}}.$$

Comparing the formulae we find that

$$b_{S}(\zeta) = \widetilde{b}_{S}(1/\zeta) \cdot \frac{\overline{\zeta}^{2}}{\zeta^{2}} \overset{\zeta \text{ large}}{\approx} \widetilde{b}_{S}(0) \cdot \frac{\overline{\zeta}^{2}}{\zeta^{2}}$$

Assuming that S is not projective-umbilic at points lying on the z_1 -axis we have $\tilde{b}_S(0) \neq 0$. It follows that the logarithmic integral $\int_{|\zeta|=M} \frac{db_S}{b_S} \in 2\pi i\mathbb{Z}$ must equal $-8\pi i$ for M large. From Stokes' theorem we rescale that the Stokes' theorem we now see that b_S must have zeros in the disk $|\zeta| < M$.

4. COMMENTS AND EXAMPLES.

Example 8. Consider the hypersurface $S = \{(z_1, z_2) : (|z_1|^2 + |z_2|^2)^2 + |z_1|^4 + |z_2|^4 = 2\}.$ Computation reveals that

$$\mathscr{B}_{S} = -\frac{3\overline{z_{1}^{2}z_{2}^{2}}}{2\left(|z_{1}|^{4} + 4|z_{1}z_{2}|^{2} + |z_{2}|^{4}\right)} \frac{dz_{1} \wedge dz_{2}}{d\overline{z_{1}} \wedge d\overline{z_{2}}}$$

so \mathscr{B}_S has double (conjugate) zeros along each axis. In fact, S has fourth order contact with $2|z_1|^2 + |z_2|^2 = 2$ along the z_1 -axis and with $|z_1|^2 + 2|z_2|^2 = 2$ along the z_2 -axis.

Question 9. Suppose that $S \subset \mathbb{CP}^2$ is a not-necessarily-circular compact strongly pseudoconvex real hypersurface satisfying the strong \mathbb{C} -convexity condition $|\mathscr{B}_S| < 1$ ([APS, Def. 2.5.10, [Bar], §§5.2-3]). Must S have a projective-umbilic point?

Example 10. The answer to the above question is negative if the strong \mathbb{C} -convexity condition is dropped. In fact, the example

$$\left(\log |z_1|\right)^2 + \left(\log |z_2|\right)^2 = \varepsilon^2$$

from [ESZ] of a compact strongly pseudoconvex hypersurface in \mathbb{C}^2 without CR-umbilic points also has no projective-umbilic points when ε is small. (The latter claim follows from $\beta_S = -\frac{\overline{z_1 z_2}}{z_1 z_2}(1 + \varepsilon)$)

 $\mathcal{O}(\varepsilon)) \frac{dz_1 \wedge dz_2}{d\overline{z_1} \wedge d\overline{z_2}}.)$

Note: The strong \mathbb{C} -convexity condition appearing in Question 9 implies in particular that the domain bounded by *S* is homeomorphic to the unit ball [APS, Thm. 2.4.2].

Remark 11. The proof of Theorem 7 given above is essentially topological. In effect, it shows that any Beltrami differential on S^3 that is invariant under rotations R_{θ} : $(z_1, z_2) \mapsto (e^{i\theta_1}z_1, e^{i\theta_2}z_2)$ must vanish along at least one circle.

It will not be possible to resolve Question 9 by a purely topological argument; in fact, on any smooth real hypersurface in \mathbb{C}^2 we have the nowhere-vanishing Beltrami differential $\frac{dz_1 \wedge dz_2}{d\overline{z_1} \wedge d\overline{z_2}}$.

It is worth noting here that most of the results mentioned in §1 require a proof with genuine geometry, not just topology (though topological arguments often suffice to prove weaker versions).

5. COMPETING CR STRUCTURES WITH THE SAME MAXIMAL COMPLEX SUBSPACE

Let $S \subset \mathbb{C}^2$ be a smooth connected real hypersurface with defining function *r*. The maximal complex subbundle $HS \subset TS$ may be described as ker $\left(\frac{d^c r}{TS} \right)$ where $d^c = \frac{\partial -\overline{\partial}}{2i}$.

Suppose that we have an alternate CR structure on S with the same maximal complex subbundle HS.

Let ω be a nowhere-vanishing 1-form on S that is type (1, 0) on HS with respect to the alternate CR structure. Then ω may be uniquely decomposed as $\omega' + \omega''$ where ω' is type (1,0) on each H_pS with respect to the original CR structure and ω'' is type (0,1) on each H_pS with respect to the original CR structure. We have $|\omega''| < |\omega'|$ if the orientations on HS match and $|\omega''| > |\omega'|$ if they do not match.

The 2-forms $d^c r \wedge \omega'$ and $d^c r \wedge \omega''$ may be extended to forms on a neighborhood of S of types (2,0) and (0,2), respectively, with respect to the original CR structure.

Replacing ω by $\tilde{\omega} = \lambda \omega$ (with λ nowhere vanishing) has the effect of multiplying the (2,0)- and (0,2)-forms above by λ along S so that the ratio is unchanged along S.

If the orientations on HS match then to avoid a vanishing denominator we should take the (2,0)form to be the denominator. By the same reasoning, in the other case we should take the (0,2)-form to be the denominator; this is the situation arising behind the scenes earlier in this paper, where the alternate CR structure is the one induced by projective duality considerations as in §3 of [BaGr], leading to the ratio (2.1) above.

The alternate CR structure can also be defined by the orientation choice together with a dilationinvariant family of ellipses in each H_pS which correspond to circles for the alternate structure. In the case of non-matching orientations the magnitude $\left| b(p) \frac{dz_1 \wedge dz_2}{d\overline{z_1} \wedge d\overline{z_2}} \right| \stackrel{\text{def}}{=} |b(p)| < 1$ determines the major-to-minor axis ratio $\frac{1 + |b(p)|}{1 - |b(p)|}$ for the ellipses in H_pS , while a vector $X \in H_p(S) \setminus \{0\}$ points in the direction of the minor axes precisely when $b(p) \frac{dz_1 \wedge dz_2}{d\overline{z_1} \wedge d\overline{z_2}} (X, Y) > 0$ for some (equivalently, for all) $Y \in T_pS \setminus H_pS$ – see [Bar, §5.3]. (Here T_pS is the space of *real* tangent vectors to S.) In the case of matching orientations we reach corresponding conclusions starting with $b(p) \frac{d\overline{z_1} \wedge d\overline{z_2}}{dz_1 \wedge dz_2}$ (again with |b(p)| < 1).

To provide a concrete illustration we consider a famous example of Rossi [Ros] (see also [Bur]) of a family of competing CR structures on the unit sphere S^3 in C^2 .

The standard CR structure on S^3 can be described by the condition that the CR functions on S^3 are those annihilated by the (complex) tangential vector field $\overline{L} \stackrel{\text{def}}{=} z_2 \frac{\partial}{\partial \overline{z_1}} - z_1 \frac{\partial}{\partial \overline{z_2}}$. The CR functions for the alternate structure (which depends on a real parameter $t \in (-1, 1)$) are those annihilated by $\overline{L} + tL$. The two structures share the same $H_{(z_1, z_2)}S^3 = \{(\gamma \overline{z_2}, -\gamma \overline{z_2})\}$: $\gamma \in \mathbb{C}\}$, with matching orientation. (Here we are viewing $H_{(z_1, z_2)}S^3$ as a vector subspace of the real tangent space $T_{(z_1, z_2)}\mathbb{C}^2$, which we identify with \mathbb{C}^2 .)

The function $f = z_1^2 + z_2^2 + t\left(\overline{z}_1^2 + \overline{z}_2^2\right)$ is CR for the alternate structure, so the map

$$d_{(z_1,z_2)}f : H_{(z_1,z_2)}S^3 \to \mathbb{C}$$

$$\left(\gamma \overline{z}_2, -\gamma \overline{z}_1\right) \mapsto \gamma \left(z_1 \overline{z}_2 - \overline{z}_1 z_2\right) + t \overline{\gamma} \left(\overline{z}_1 z_2 - z_1 \overline{z}_2\right)$$

is C-linear for the alternate structure. So the ellipses we want are given by

$$\left|\gamma\left(z_1\overline{z}_2 - \overline{z}_1z_2\right) + \overline{\gamma}t\left(\overline{z}_1z_2 - z_1\overline{z}_2\right)\right| = C^{\text{constant}}$$

with the minor axis corresponding to $\gamma \in i\mathbb{R}$ and the major-to-minor axis ratio equal to $\frac{1+t}{1-t}$.

(The argument above runs into trouble at points where $d_{(z_1,z_2)}f = 0$ on $H_{(z_1,z_2)}S^3$; this happens in particular when one of the z_j is a real multiple of the other. However, the conclusions above can still be shown to hold at such points by replacing f by $g = z_1^2 - z_2^2 - t(\overline{z_1}^2 - \overline{z_2}^2)$ or by $h = z_1 z_2 - t \overline{z_1} \overline{z_2}$.)

be shown to hold at such points by replacing f by $g = z_1^2 - z_2^2 - t\left(\overline{z_1}^2 - \overline{z_2}^2\right)$ or by $h = z_1 z_2 - t \overline{z_1} \overline{z_2}$.) To compare this conclusions to the geometric discussion of Beltrami differentials above we see that we must have $\frac{1 + |b(p)|}{1 - |b(p)|} = \frac{1 + t}{1 - t}$ hence $|b| \equiv t$ on S^3 . To check the direction of the minor axes we note that the vector $\left\{ (\gamma \overline{z_2}, -\gamma \overline{z_1}) \right\}$: $\gamma \in \mathbb{C} \left\{ = H_{(z_1, z_2)} S^3 \right\}$ points in the direction of the minor axes when $\gamma (\overline{z_1} z_2 - z_1 \overline{z_2}) \in \mathbb{R}$, in particular when $\gamma = i$; thus the vector field $X \stackrel{\text{def}}{=} (i \overline{z_2}, -i \overline{z_1})$ describes the minor axes.

Taking Y to be the rotational vector field

$$S^3 \to TS^3$$
$$(z_1, z_2) \mapsto (iz_1, iz_2)$$

we may rewrite X and Y in operator form

$$X = i \left(\overline{z_2} \frac{\partial}{\partial z_1} - \overline{z_1} \frac{\partial}{\partial z_2} - z_2 \frac{\partial}{\partial \overline{z_1}} + z_1 \frac{\partial}{\partial \overline{z_2}} \right)$$
$$Y = i \left(z_1 \frac{\partial}{\partial z_1} + z_2 \frac{\partial}{\partial z_2} - \overline{z_1} \frac{\partial}{\partial \overline{z_1}} - \overline{z_2} \frac{\partial}{\partial \overline{z_2}} \right)$$

Thus $dz_1 \wedge dz_2(X, Y) = -1 = d\overline{z}_1 \wedge d\overline{z}_2$ and $\frac{d\overline{z_1} \wedge d\overline{z_2}}{dz_1 \wedge dz_2}(X, Y) > 0$.

Combining our conclusions, we see that the Beltrami differential for the Rossi example is $t \frac{d\overline{z_1} \wedge d\overline{z_2}}{dz_1 \wedge dz_2}$. (Compare Remark 11 above.)

REFERENCES

- [Bar] D. Barrett, *Holomorphic projection and duality for domains in complex projective space*, Trans. Amer. Math. Soc. **368** (2016), 827–850.
- [BaGr] D. Barrett and D. Grundmeier, Sums of CR functions from competing CR structures, Pacific J. Math. 293-2 (2018), 257–275.
- [Ber] M. Berger, Geometry revealed: a Jacob's ladder to modern higher geometry, Springer, 2010.
- [Bol] M. Bolt, The Möbius geometry of hypersurfaces, Michigan Math. J. 56 (2008), 603-622.
- [Bur] D. Burns, *Global behavior of some tangential Cauchy-Riemann equations. Partial differential equations and geometry*, Lecture Notes in Pure and Appl. Math., **48** (1979), Dekker, New York, pp.51–56.
- [Car1] É. Cartan, Sur la géométrie pseudo-conforme des hypersurfaces de l'espace de deux variables complexes, Ann. Mat. Pura Appl. **11** (1933), 17–90.
- [Car2] É. Cartan, Sur la géométrie pseudo-conforme des hypersurfaces de l'espace de deux variables complexes II, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (2) 1 (1932), 333Đ354.
- [DeTr] J. Detraz and J.M. Trépreau, *Une caractérisation des quadriques hermitiennes dans* \mathbb{C}^n , J. Analyse Math. 55, (1990), 51–58.
- [EbSo] P. Ebenfelt and D. Son, *Umbilical points on three dimensional strictly pseudoconvex CR manifolds I: manifolds with U*(1)-action, Math. Ann. **368** (2017), 537–560.
- [ESZ] P. Ebenfelt, D. Son, and D. Zaitsev, A family of compact strictly pseudoconvex hypersurfaces in C² without umbilical points, Math. Res. Lett. 25 (2018), 75–84.
- [Ghy] E. Ghys, *Cercles osculateurs et géométrie lorentzienne*, talk at the journée inaugurale du CMI, Marseille, February 1995.
- [Ham] C. Hammond, *Invariants of Transformation Groups Acting on Real Hypersurfaces of Complex Spaces*, University of Michigan PhD. dissertation, 2009.
- [HuJi] X. Huang and S. Ji, Every real ellipsoid in C² admits CR umbilical points, Trans. Amer. Math. Soc. 359 (2007), 1191–1204.
- [Jen] G. Jensen, *Projective deformation and biholomorphic equivalence of real hypersurfaces*. Ann. Global Anal. Geom. **1** (1983), 1–34.
- [Kne] A. Kneser, Bemerkungen über die Anzahl der Extrema des Krümmung auf geschlossenen Kurven und uber verwandte Fragen in einer nicht euklidischen Geometrie, Festschrift Heinrich Weber, Teubner (1912), 170– 180.
- [Leh] O. Lehto, Univalent functions and Teichmüller spaces, Graduate Texts in Mathematics 109, Springer, 1987.
- [Muk] S. Mukhopadhyaya, New Methods in the Geometry of a Plane Arc, Bull. Calcutta Math. Soc. 1 (1908), 31–38.
- [Oss] R. Osserman, The four-or-more vertex theorem, Amer. Math. Monthly 92 (1985), 332–337.

- [OvTa] V. Ovsienko and S. Tabachnikov, *Sturm theory, Ghys theorem on zeroes of the Schwarzian derivative and flattening of Legendrian curves,* Selecta Math. (N.S.) **2** (1996), 297–307.
- [Ros] H. Rossi, Attaching analytic spaces to an analytic space along a pseudoconcave boundary, 1965 Proc. Conf. Complex Analysis (Minneapolis, 1964) Springer, pp. 242–256.
- [Rut] J. Rutter, *Geometry of curves*, Chapman & Hall/CRC Mathematics, 2000.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109-1043 USA *E-mail address*: barrett@umich.edu

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE, MA 02138-2901 USA *E-mail address*: deg@math.harvard.edu