

The α -divergences associated with a pair of strictly comparable quasi-arithmetic means

Frank Nielsen*
 Sony Computer Science Laboratories Inc.
 Tokyo, Japan

Abstract

We generalize the family of α -divergences using a pair of strictly comparable weighted means. In particular, we obtain the 1-divergence in the limit case $\alpha \rightarrow 1$ (a generalization of the Kullback-Leibler divergence) and the 0-divergence in the limit case $\alpha \rightarrow 0$ (a generalization of the reverse Kullback-Leibler divergence). We state the condition for a pair of quasi-arithmetic means to be strictly comparable, and report the formula for the quasi-arithmetic α -divergences and its subfamily of bipower homogeneous α -divergences which belong to the Csisár's f -divergences. Finally, we show that these generalized quasi-arithmetic 1-divergences and 0-divergences can be decomposed as the sum of generalized cross-entropies minus entropies, and rewritten as conformal Bregman divergences using monotone embeddings.

Keywords: Kullback-Leibler divergence, α -divergences, comparable means, weighted quasi-arithmetic means, α -geometry, homogeneous divergences, conformal divergences, geometric divergence, monotone embeddings, conformal flattening.

1 Introduction

1.1 Statistical divergences

Consider a measurable space $(\mathcal{X}, \mathcal{F})$ (where \mathcal{F} denotes the σ -algebra and \mathcal{X} the sample space) equipped with a positive measure μ (e.g., usually the Lebesgue measure or the counting measure). The notion of *statistical dissimilarity* [4] $D[P : Q] = D_\mu[p_\mu : q_\mu]$ between two arbitrary probability measures with Radon-Nikodym (RN) densities $p_\mu = \frac{dP}{d\mu}$ and $q_\mu = \frac{dQ}{d\mu}$ with respect to μ is at the core of many algorithms in signal processing, information theory, information fusion, and machine learning among others. When those statistical dissimilarities are smooth, they are called *divergences* [2] in the literature. The most renowned statistical divergence rooted in information theory [9] is the *Kullback-Leibler divergence* (KLD):

$$\text{KL}_\mu[p_\mu : q_\mu] := \int_{\mathcal{X}} p_\mu(x) \log \frac{p_\mu(x)}{q_\mu(x)} d\mu(x). \quad (1)$$

Since the KLD is independent of the reference measure μ , i.e., $\text{KL}_\mu[p_\mu : q_\mu] = \text{KL}_\nu[p_\nu : q_\nu]$ for $p_\mu = \frac{dP}{d\mu}$ and $q_\mu = \frac{dQ}{d\mu}$, and $p_\nu = \frac{dP}{d\nu}$ and $q_\nu = \frac{dQ}{d\nu}$ the RN derivatives with respect to another

*E-mail: Frank.Nielsen@acm.org

positive measure ν , we write concisely in the remainder:

$$\text{KL}[p : q] = \int p \log \frac{p}{q} d\mu, \quad (2)$$

instead of $\text{KL}_\mu[p_\mu : q_\mu]$.

The KLD belongs to a parametric family of α -divergences [7] $I_\alpha[p : q]$ for $\alpha \in \mathbb{R}$:

$$I_\alpha[p : q] := \begin{cases} \frac{1}{\alpha(1-\alpha)} (1 - \int p^\alpha q^{1-\alpha} d\mu), & \alpha \in \mathbb{R} \setminus \{0, 1\} \\ I_1[p : q] = \text{KL}[p : q], & \alpha = 1 \\ I_0[p : q] = \text{KL}[q : p], & \alpha = 0 \end{cases}. \quad (3)$$

The α -divergences extended to positive densities (not necessarily normalized) play a central role in information geometry [2]:

$$I_\alpha[p : q] := \begin{cases} \frac{1}{\alpha(1-\alpha)} \int (\alpha p + (1-\alpha)q - p^\alpha q^{1-\alpha}) d\mu, & \alpha \in \mathbb{R} \setminus \{0, 1\} \\ I_1[p : q] = \text{KL}_e[p : q], & \alpha = 1 \\ I_0[p : q] = \text{KL}_e[q : p], & \alpha = 0 \end{cases}, \quad (4)$$

where KL_e denotes the *extended Kullback-Leibler divergence*:

$$\text{KL}_e[p : q] := \int \left(p \log \frac{p}{q} + q - p \right) d\mu. \quad (5)$$

The α -divergences are asymmetric for $\alpha \neq 0$ (i.e., $I_\alpha[p : q] \neq I_\alpha[q : p]$ for $\alpha \neq 0$) but exhibit the following *reference duality* [39]:

$$I_\alpha[q : p] = I_{1-\alpha}[p : q] = (I_\alpha)^*[p : q], \quad (6)$$

where we denoted by $D^*[p : q] := D[q : p]$, the *reverse divergence* for an arbitrary divergence D (e.g., $I_\alpha^*[p : q] = I_\alpha[q : p] = I_{1-\alpha}[p : q]$).

The α -divergences belong to the family of Csizár's f -divergences [10] which are defined for a convex function f satisfying by $f(1) = 0$ by:

$$I_f[p : q] := \int p f \left(\frac{q}{p} \right) d\mu. \quad (7)$$

We have

$$I_\alpha[p : q] = I_{f_\alpha}[p : q], \quad (8)$$

with

$$f_\alpha(u) = \begin{cases} \frac{1}{\alpha(1-\alpha)} (\alpha + (1-\alpha)u - u^{1-\alpha}), & \alpha \in \mathbb{R} \setminus \{0, 1\} \\ u - 1 - \log u, & \alpha = 1 \\ 1 - u + u \log u, & \alpha = -1 \end{cases} \quad (9)$$

In information geometry, α -divergences (and more generally f -divergences) are *invariant divergences* [2], and it is customary to rewrite the α -divergences using $\alpha_A = 1 - 2\alpha$ (i.e., $\alpha = \frac{1-\alpha_A}{2}$). Thus the *extended α_A -divergence* is defined by

$$\hat{I}_{\alpha_A}[p : q] = \begin{cases} \frac{4}{1-\alpha_A^2} \int \left(\frac{1-\alpha_A}{2} p + \frac{1+\alpha_A}{2} q - p^{\frac{1-\alpha_A}{2}} q^{\frac{1+\alpha_A}{2}} \right) d\mu, & \alpha \in \mathbb{R} \setminus \{-1, 1\} \\ \hat{I}_1[p : q] = \text{KL}_e[p : q], & \alpha = 1 \\ \hat{I}_{-1}[p : q] = \text{KL}_e[q : p], & \alpha = -1 \end{cases}, \quad (10)$$

and the reference duality is expressed by $\hat{I}_{\alpha_A}[q : p] = \hat{I}_{-\alpha_A}[p : q]$.

A statistical divergence $D[\cdot : \cdot]$ when evaluated on densities belonging to a given parametric family $\mathcal{P} = \{p_\theta : \theta \in \Theta\}$ of densities are equivalent to a corresponding *contrast function* [12]:

$$D_{\mathcal{P}}(\theta_1 : \theta_2) := D[p_{\theta_1} : p_{\theta_2}]. \quad (11)$$

Although quite confusing, those contrast functions have also been called recently divergences in the literature [2]. Thus to disambiguate whether the divergence is a statistical divergence or a parameter divergence (i.e., contrast function), we choose to use the brackets for encapsulating arguments in statistical divergences and the parenthesis to encapsulate parameter arguments in divergences which are contrast functions.

A smooth divergence $D(\theta_1 : \theta_2)$ induces a dualistic structure in information geometry [2]. For example, the KLD on the family Δ of probability mass functions on a finite alphabet \mathcal{X} with equivalent contrast function a Bregman divergence induces a *dually flat space* [2]. More generally, the α_A -divergences on the probability simplex Δ induce the α_A -*geometry*.

The α -divergences are widely used in information sciences, see [3, 8, 38, 22, 17, 1] just to cite a few applications. The singly-parametric α -divergences have also been generalized to bi-parametric families of divergences like the (α, β) -divergences [2] or the $\alpha\beta$ -divergences [37].

In this work, based on the observation that the term $\alpha p + (1 - \alpha)q - p^\alpha q^{1-\alpha}$ (in the extended $I_\alpha[p : q]$ divergence for $\alpha \in (0, 1)$ of Eq. 4) is a difference between a weighted arithmetic mean $A_{1-\alpha}(p, q) := \alpha p + (1 - \alpha)q$ and a weighted geometric mean $G_{1-\alpha}(p, q) := p^\alpha q^{1-\alpha}$, we investigate a generalization of α -divergences with respect to a pair of abstract strictly comparable weighted means [25].

1.2 Divergences and decomposable divergences

A statistical divergence D shall satisfy the following two axioms:

D1. non-negativity. $D[p : q] \geq 0$ for all densities p and q ,

D2. identity of indiscernibles. $D[p : q] = 0$ if and only if $p = q$ μ -almost everywhere.

These axioms are a subset of the metric axioms since we do not consider the symmetry axiom nor the triangular inequality axiom of metric distances. See [14] for some common examples of *probability metrics* (e.g., total variation or Wasserstein metrics).

A divergence $D[p : q]$ is said *decomposable* [2] when it can be written as an integral of a scalar divergence $d(\cdot, \cdot)$:

$$D[p : q] = \int d(p(x) : q(x)) d\mu(x), \quad (12)$$

or $D[p : q] = \int d(p : q) d\mu$ for short.

The α -divergences are decomposable divergences since we have

$$I_\alpha[p : q] = \int i_\alpha(p(x) : q(x)) d\mu \quad (13)$$

with the following *scalar α -divergence*:

$$i_\alpha(a : b) := \begin{cases} \frac{1}{\alpha(1-\alpha)} (\alpha a + (1-\alpha)b - a^\alpha b^{1-\alpha}), & \alpha \in \mathbb{R} \setminus \{0, 1\} \\ i_1(a : b) = a \log \frac{a}{b} + b - a & \alpha = 1 \\ i_0(a : b) = i_1(b : a), & \alpha = 0 \end{cases} \quad (14)$$

1.3 Contributions and paper outline

The outline of the paper and the contributions are summarized as follows:

We define the generic α -divergences in §2 for two families of strictly comparable means (Definition 2). Then section 2.2 reports a closed-form formula (Theorem 3) for the quasi-arithmetic α -divergences induced by two strictly comparable quasi-arithmetic means with monotonically increasing generators f and g such that $f \circ g^{-1}$ is *strictly convex and differentiable*. In §2.3, we study the divergences I_0 and I_1 obtained in the limit cases when $\alpha \rightarrow 0$ and $\alpha \rightarrow 1$, respectively. We obtain generalized Kullback-Leibler divergences when $\alpha \rightarrow 1$ and generalized reverse Kullback-Leibler divergences when $\alpha \rightarrow 0$, which can be factorized as generalized cross-entropies minus entropies. In §2.4, we show how to express these generalized I_1 -divergences and I_0 -divergences as conformal Bregman representational divergences and briefly explain their induced conformally flat statistical manifolds. Section 3 explicits the subfamily of bipower homogeneous α -divergences which belong to the family of Csiszár f -divergences [10]. Finally, Section 4 summarizes the work and present several opportunities for future research directions.

2 The α -divergences based on a pair of means

2.1 The abstract (M, N) α -divergences

The point of departure for generalizing the α -divergences is to rewrite Eq. 4 for $\alpha \in \mathbb{R} \setminus \{0, 1\}$ as

$$I_\alpha[p : q] = \frac{1}{\alpha(1-\alpha)} \int (A_{1-\alpha}(p : q) - G_{1-\alpha}(p : q)) d\mu, \quad (15)$$

where A_λ and G_λ for $\lambda \in (0, 1)$ stands for the weighted arithmetic and geometric means, respectively:

$$\begin{aligned} A_\lambda(x, y) &= (1-\lambda)x + \lambda y, \\ G_\lambda(x, y) &= x^{1-\lambda} y^\lambda. \end{aligned}$$

We choose the convention $A_0(x, y) = x$ and $A_1(x, y) = 1$ so that $\{A_\lambda(x, y)\}_{\lambda \in [0, 1]}$ smoothly interpolates between x ($\lambda = 0$) and y ($\lambda = 1$).

In general, a *mean* $M(x, y)$ aggregates two values x and y of an interval I to produce an intermediate quantity which satisfies the *innerness property* [5]:

$$\min\{x, y\} \leq M(x, y) \leq \max\{x, y\}, \quad \forall x, y \in I. \quad (16)$$

A mean is said *strict* if the inequalities of Eq. 16 are strict whenever $x \neq y$. A mean M is said *reflexive* iff $M(x, x) = x$ for all $x \in I$. In the remainder, we consider $I = (0, \infty)$. By using the unique dyadic representation of any real $\lambda \in (0, 1)$ (i.e., $\lambda = \sum_{i=1}^{\infty} \frac{d_i}{2^i}$ with $d_i \in \{0, 1\}$, a binary digit), one can build a *weighted mean* M_λ from any given mean M , see [25] for such a construction.

By analogy to the α -divergences, let us define the (decomposable) (M, N) α -divergences for a pair of weighted means $M_{1-\alpha}$ and $N_{1-\alpha}$ for $\alpha \in (0, 1)$ as

$$I_{\alpha}^{M,N}[p : q] := \frac{1}{\alpha(1-\alpha)} \int (M_{1-\alpha}(p : q) - N_{1-\alpha}(p : q)) d\mu. \quad (17)$$

The ordinary α -divergences for $\alpha \in (0, 1)$ are recovered as the (A, G) α -divergences:

$$I_{\alpha}^{A,G}[p : q] = \frac{1}{\alpha(1-\alpha)} \int (A_{1-\alpha}(p : q) - G_{1-\alpha}(p : q)) d\mu = I_{1-\alpha}[p : q] = I_{\alpha}[q : p] = I_{\alpha}^*[p : q]. \quad (18)$$

In order to define generalized α -divergences satisfying the axioms D1 and D2 of proper divergences, we need to characterize the class of acceptable means. We give a definition strengthening the notion of comparable means in [25]:

Definition 1 (Strictly comparable weighted means). *A pair (M, N) of means are said strictly comparable whenever $M_{\lambda}(x, y) \leq N_{\lambda}(x, y)$ for all $x, y \in (0, \infty)$ with equality if and only if $x = y$, and for all $\lambda \in (0, 1)$.*

For example, the inequality of the arithmetic and geometric means states that $A(x, y) \geq G(x, y)$: Means A and G are comparable, denoted by $A \geq G$. Furthermore, the arithmetic and geometric weighted means are distinct whenever $x \neq y$: Indeed, consider the equation $(1-\alpha)x + \alpha y = x^{1-\alpha}y^{\alpha}$ for $x, y > 0$ and $x \neq y$. By taking the logarithm on both sides, we get

$$\log((1-\alpha)x + \alpha y) = (1-\alpha)\log x + \alpha \log y. \quad (19)$$

Since the logarithm is a strictly convex function, the only solution is $x = y$. Thus (A, G) is a pair of strictly comparable weighted means.

For a weighted mean M , define $\bar{M}_{\alpha}(x, y) := M_{1-\alpha}(x, y)$. We are ready to state the definition of generalized α -divergences:

Definition 2 ((M, N) α -divergences). *The (M, N) α -divergences $I_{\alpha}^{M,N}[p : q]$ between two positive densities p and q for $\alpha \in (0, 1)$ is defined for a pair of strictly comparable weighted means M_{α} and N_{α} with $M_{\alpha} \geq N_{\alpha}$ by:*

$$I_{\alpha}^{M,N}[p : q] := \frac{1}{\alpha(1-\alpha)} \int (M_{1-\alpha}(p : q) - N_{1-\alpha}(p : q)) d\mu, \quad (20)$$

$$= \frac{1}{\alpha(1-\alpha)} \int (\bar{M}_{\alpha}(p : q) - \bar{N}_{\alpha}(p : q)) d\mu. \quad (21)$$

Using $\alpha = \frac{1-\alpha_A}{2}$, we can rewrite this divergence as

$$\hat{I}_{\alpha_A}^{M,N}[p : q] := \frac{4}{1-\alpha_A^2} \int \left(M_{\frac{1+\alpha_A}{2}}(p : q) - N_{\frac{1+\alpha_A}{2}}(p : q) \right) d\mu, \quad (22)$$

$$= \frac{4}{1-\alpha_A^2} \int \left(\bar{M}_{\frac{1-\alpha_A}{2}}(p : q) - \bar{N}_{\frac{1-\alpha_A}{2}}(p : q) \right) d\mu. \quad (23)$$

A weighted mean M_{α} is said *symmetric* iff $M_{\alpha}(x, y) = M_{1-\alpha}(y, x)$. When both the weighted means M and N are symmetric, we have the following *reference duality* [39]:

$$I_{\alpha}^{M,N}[p : q] = I_{1-\alpha}^{M,N}[q : p]. \quad (24)$$

We consider symmetric means in the remainder.

In the limit cases of $\alpha \rightarrow 0$ or $\alpha \rightarrow 1$, we define the 0-*divergence* $I_0^{M,N}[p : q]$ and the 1-*divergence* $I_1^{M,N}[p : q]$, respectively, by

$$I_0^{M,N}[p : q] = \lim_{\alpha \rightarrow 0} I_\alpha^{M,N}[p : q], \quad (25)$$

$$I_1^{M,N}[p : q] = \lim_{\alpha \rightarrow 1} I_\alpha^{M,N}[p : q] = I_0^{M,N}[q : p], \quad (26)$$

provided that those limits exist.

2.2 The quasi-arithmetic α -divergences

A *quasi-arithmetic mean* (QAM) is defined for a continuous and strictly monotonic function $f : I \subset \mathbb{R} \rightarrow J \subset \mathbb{R}$ as:

$$M^f(x, y) := f^{-1} \left(\frac{f(x) + f(y)}{2} \right). \quad (27)$$

Function f is called the generator of the quasi-arithmetic mean. These strict and reflexive quasi-arithmetic means are also called Kolmogorov means [19], Nagumo means [23] or de Finetti means [11], or quasi-linear means [15] in the literature. These means are called quasi-arithmetic means because they can be interpreted as arithmetic means on the arguments $f(x)$ and $f(y)$:

$$f(M^f(x, y)) = \frac{f(x) + f(y)}{2} = A(f(x), f(y)). \quad (28)$$

QAMs are strict, reflexive and symmetric means.

Without loss of generality, we assume strictly increasing functions f instead of monotonic functions since $M^{-f} = M^f$. Indeed, $M^{-f}(x, y) = (-f)^{-1}(-f(M^f(x, y)))$ and $((-f)^{-1} \circ (-f))(u) = u$, the identity function. Notice that the composition $f_1 \circ f_2$ of two strictly monotonic increasing functions f_1 and f_2 is a strictly monotonic increasing function. Furthermore, we consider $I = J = (0, \infty)$ in the remainder since we apply these means on positive densities. Two quasi-arithmetic means M^f and M^g coincide if and only if $f(u) = ag(u) + b$ for some $a > 0$ and $b \in \mathbb{R}$ see [15]. The quasi-arithmetic means were considered in the axiomatization of the entropies by Rényi to define the α -entropies (see Eq. 2.11 of [36]).

By choosing $f_A(u) = u$, $f_G(u) = \log u$ or $f_H(u) = \frac{1}{u}$, we obtain the Pythagorean's arithmetic A , geometric G , and harmonic H means, respectively:

- the *arithmetic mean* (A): $A(x, y) = \frac{x+y}{2} = M^{f_A}(x, y)$,
- the *geometric mean* (G): $G(x, y) = \sqrt{xy} = M^{f_G}(x, y)$, and
- the *harmonic mean* (H): $H(x, y) = \frac{2}{\frac{1}{x} + \frac{1}{y}} = \frac{2xy}{x+y} = M^{f_H}(x, y)$.

More generally, choosing $f_{P_r}(u) = u^r$, we obtain the parametric family of *power means* (also called *Hölder means* [16]):

$$P_r(x, y) = \left(\frac{x^r + y^r}{2} \right)^{\frac{1}{r}} = M^{f_{P_r}}(x, y), \quad r \in \mathbb{R} \setminus \{0\}. \quad (29)$$

In order to get a *smooth family* of power means, we define the geometric mean in the limit case of $r \rightarrow 0$:

$$P_0(x, y) = \lim_{r \rightarrow 0} P_r(x, y) = G(x, y) = \sqrt{xy}. \quad (30)$$

It is known that the positively homogeneous quasi-arithmetic means, i.e. $M^f(\lambda a, \lambda b) = \lambda M^f(a, b)$ for $\lambda > 0$, coincide exactly with the family of power means. The weighted QAMs are given by

$$M_\alpha^f(p, q) = f^{-1}((1 - \alpha)f(p) + \alpha f(q)) = f^{-1}(f(p) + \alpha(f(q) - f(p))) = M_{1-\alpha}^f(q, p). \quad (31)$$

The logarithmic mean $L(x, y)$ for $x > 0$ and $y > 0$:

$$L(x, y) = \frac{y - x}{\log y - \log x}$$

is an example of a homogeneous mean (i.e., $L(\lambda x, \lambda y) = \lambda L(x, y)$ for any $\lambda > 0$) that is *not* a QAM. Besides the family of QAMs, there exist many other families of means [5]: For example, let us mention the *Lagrangean means* [18] which intersect with the QAMs only for the arithmetic mean, or a generalization of the QAMs called the *Bajraktarević means* [35].

Let us strengthen a recent theorem of [21] (Theorem 1, 2010):

Theorem 1 (Strictly comparable weighted QAMs). *The pair (M^f, M^g) of quasi-arithmetic means obtained for two strictly increasing generators is strictly comparable provided that $f \circ g^{-1}$ is strictly convex.*

Proof. Since $f \circ g^{-1}$ is strictly convex, it is convex, and therefore it follows from Theorem 1 of [21] that $M_\alpha^f \geq M_\alpha^g$ for all $\alpha \in [0, 1]$. Thus the very nice property of QAMs is that $M^f \geq M^g$ implies that $M_\alpha^f \geq M_\alpha^g$ for *any* $\alpha \in [0, 1]$.

Now, let us consider the equation $M_\alpha^f(p, q) = M_\alpha^g(p, q)$ for $p \neq q$:

$$f^{-1}((1 - \alpha)f(p) + \alpha f(q)) = g^{-1}((1 - \alpha)g(p) + \alpha g(q)). \quad (32)$$

Since $f \circ g^{-1}$ is assumed strictly convex, and g is strictly increasing, we have $g(p) \neq g(q)$ for $p \neq q$, and we reach the following contradiction:

$$(1 - \alpha)f(p) + \alpha f(q) = (f \circ g^{-1})((1 - \alpha)g(p) + \alpha g(q)), \quad (33)$$

$$< (1 - \alpha)(f \circ g^{-1})(g(p)) + \alpha(f \circ g^{-1})(g(q)), \quad (34)$$

$$< (1 - \alpha)f(p) + \alpha f(q). \quad (35)$$

Thus $M_\alpha^f(p, q) \neq M_\alpha^g(p, q)$ for $p \neq q$, and $M_\alpha^f(p, q) = M_\alpha^g(p, q)$ for $p = q$. \square

Note that the (A, G) α -divergences (i.e., the ordinary α -divergences) is a proper divergence satisfying both the properties D1 and D2 because $f_A(u) = u$ and $f_G(u) = \log u$, and hence $(f_A \circ f_G^{-1})(u) = \exp(u)$ is strictly convex on $(0, \infty)$.

Let us denote by $I_\alpha^{f,g}[p : q] := I_\alpha^{M^f, M^g}[p : q]$ the *quasi-arithmetic α -divergences*. Since the QAMs are symmetric means, we have $I_\alpha^{f,g}[p : q] = I_{1-\alpha}^{f,g}[q : p]$.

2.3 Limit cases of 1-divergences and 0-divergences

We seek a closed-form formula of the limit divergence $\lim_{\alpha \rightarrow 0} I_{\alpha}^{f,g}[p : q]$ when $\alpha \rightarrow 0$.

Lemma 1. *A first-order Taylor approximation of the quasi-arithmetic mean [30] M_{α}^f for a C_1 strictly increasing generator f when $\alpha \simeq 0$ yields*

$$M_{\alpha}^f(p : q) = p + \frac{\alpha(f(q) - f(p))}{f'(p)} + o(\alpha(f(q) - f(p))). \quad (36)$$

Proof. By taking the first-order Taylor expansion of $f^{-1}(x)$ at x_0 (i.e., Taylor polynomial of order 1), we get:

$$f^{-1}(x) = f^{-1}(x_0) + (x - x_0)(f^{-1})'(x_0) + o(x - x_0). \quad (37)$$

Using the property of the derivative of an inverse function:

$$(f^{-1})'(x) = \frac{1}{(f'(f^{-1}(x)))}, \quad (38)$$

it follows that the first-order Taylor expansion of $f^{-1}(x)$ is:

$$f^{-1}(x) = f^{-1}(x_0) + (x - x_0) \frac{1}{(f'(f^{-1}(x_0)))} + o(x - x_0). \quad (39)$$

Plugging $x_0 = f(p)$ and $x = f(p) + \alpha(f(q) - f(p))$, we get a *first-order approximation* of the weighted quasi-arithmetic mean M_{α}^f when $\alpha \rightarrow 0$:

$$M_{\alpha}^f(p, q) = p + \frac{\alpha(f(q) - f(p))}{f'(p)} + o(\alpha(f(q) - f(p))). \quad (40)$$

□

Let us introduce the following bivariate function:

$$E_f(p, q) := \frac{f(q) - f(p)}{f'(p)}. \quad (41)$$

Thus we obtain closed-form formula for the I_1 -divergence and I_0 -divergence:

Theorem 2 (Quasi-arithmetic I_1 -divergence and I_0 -divergence). *The quasi-arithmetic I_1 -divergence induced by two strictly increasing and differentiable functions f and g such that $f \circ g^{-1}$ is strictly convex is*

$$I_1^{f,g}[p : q] = \lim_{\alpha \rightarrow 1} I_{\alpha}^{f,g}[p : q] = \int (E_f(p, q) - E_g(p, q)) \, d\mu \geq 0, \quad (42)$$

$$= \int \left(\frac{f(q) - f(p)}{f'(p)} - \frac{g(q) - g(p)}{g'(p)} \right) \, d\mu. \quad (43)$$

We have $I_0^{f,g}[p : q] = I_1^{f,g}[q : p]$.

Proof. Let us prove that $I_1^{f,g}$ is a proper divergence satisfying axioms D1 and D2. Note that a sufficient condition for $I_1^{f,g}[p : q] \geq 0$ is to check that

$$E_f(p, q) \geq E_g(p, q), \quad (44)$$

$$\frac{f(q) - f(p)}{f'(p)} \geq \frac{g(q) - g(p)}{g'(p)}. \quad (45)$$

If $p = q$ μ -a.e. then clearly $I_1^{f,g}[p : q] = 0$. Consider $p \neq q$ (i.e., at some observation x : $p(x) \neq q(x)$).

We shall use the following property of a strictly convex and differentiable function h for $x < y$ (sometimes called the chordal slope lemma, see [25]):

$$h'(x) \leq \frac{h(y) - h(x)}{y - x} \leq h'(y). \quad (46)$$

We consider $h(x) = (f \circ g^{-1})(x)$ so that $h'(x) = \frac{f'(g^{-1}(x))}{g'(g^{-1}(x))}$. There are two cases to consider:

- $p < q$ and therefore $g(p) < g(q)$. Let $y = g(q)$ and $x = g(p)$ in Eq. 46. We have $h'(x) = \frac{f'(p)}{g'(p)}$ and $h'(y) = \frac{f'(q)}{g'(q)}$, and the double inequality of Eq. 46 becomes

$$\frac{f'(p)}{g'(p)} \leq \frac{f(q) - f(p)}{g(q) - g(p)} \leq \frac{f'(q)}{g'(q)}.$$

Since $g(q) - g(p) > 0$ and $g'(p) > 0$ and $f'(p) > 0$, we get

$$\frac{g(q) - g(p)}{g'(p)} \leq \frac{f(q) - f(p)}{f'(p)}.$$

- $q < p$ and therefore $g(p) > g(q)$. Then the double inequality of Eq. 46 becomes

$$\frac{f'(q)}{g'(q)} \leq \frac{f(q) - f(p)}{g(q) - g(p)} \leq \frac{f'(p)}{g'(p)}$$

That is,

$$\frac{f(q) - f(p)}{f'(p)} \geq \frac{g(q) - g(p)}{g'(p)},$$

since $g(q) - g(p) < 0$.

Thus in both cases, we checked that $E_f(p(x), q(x)) \geq E_g(p(x), q(x))$. Therefore $I_1^{f,g}[p : q] \geq 0$ and since the QAMs are distinct $I_1^{f,g}[p : q] = 0$ iff $p(x) = q(x)$ μ -a.e. \square

We can interpret the I_1 divergences as generalized KL divergences, and define generalized notions of cross-entropies and entropies. Since the KL divergence can be written as the cross-entropy minus the entropy, we can also decompose the I_1 divergences as follows:

$$I_1^{f,g}[p : q] = \int \left(\frac{f(q)}{f'(p)} - \frac{g(q)}{g'(p)} \right) d\mu - \int \left(\frac{f(p)}{f'(p)} - \frac{g(p)}{g'(p)} \right) d\mu, \quad (47)$$

$$= h_{\times}^{f,g}(p : q) - h^{f,g}(p), \quad (48)$$

where $h_{\times}^{f,g}(p : q)$ denotes the (f, g) -cross-entropy (for a constant $c \in \mathbb{R}$):

$$h_{\times}^{f,g}(p : q) = \int \left(\frac{f(q)}{f'(p)} - \frac{g(q)}{g'(p)} \right) d\mu + c, \quad (49)$$

and $h^{f,g}(p)$ stands for the (f, g) -entropy (self cross-entropy):

$$h^{f,g}(p) = h_{\times}^{f,g}(p : p) = \int \left(\frac{f(p)}{f'(p)} - \frac{g(p)}{g'(p)} \right) d\mu + c. \quad (50)$$

We define the generalized (f, g) -Kullback-Leibler divergence:

$$\text{KL}_{f,g}[p : q] := h_{\times}^{f,g}(p : q) - h^{f,g}(p). \quad (51)$$

When $f = f_A$ and $g = f_G$, we resolve the constant to $c = 0$, and recover the ordinary Shannon cross-entropy and entropy:

$$h_{\times}^{f_A, f_G}(p : q) = \int (q - p \log q) d\mu = h_{\times}(p : q), \quad (52)$$

$$h^{f_A, f_G}(p : q) = h_{\times}^{f_A, f_G}(p : p) = \int (p - p \log p) d\mu = h(p), \quad (53)$$

and we have the (f_A, f_G) -Kullback-Leibler divergence that is the extended Kullback-Leibler divergence:

$$\text{KL}_{f_A, f_G}[p : q] = \text{KL}_e[p : q] = h_{\times}(p : q) - h(p) = \int (p \log \frac{p}{q} + q - p) d\mu. \quad (54)$$

Thus we have the (f, g) -cross-entropy and (f, g) -entropy expressed as

$$h_{\times}^{f,g}(p : q) = \int \left(\frac{f(q)}{f'(p)} - \frac{g(q)}{g'(p)} \right) d\mu, \quad (55)$$

$$h^{f,g}(p) = \int \left(\frac{f(p)}{f'(p)} - \frac{g(p)}{g'(p)} \right) d\mu. \quad (56)$$

In general, we can define the (f, g) -Jeffreys' divergence as:

$$J^{f,g}[p : q] = \text{KL}^{f,g}[p : q] + \text{KL}^{f,g}[q : p]. \quad (57)$$

Thus we define the quasi-arithmetic mean α -divergences as follows:

Theorem 3 (Quasi-arithmetic α -divergences). *Let f and g be two strictly continuously increasing and differentiable functions on $(0, \infty)$ such that $f \circ g^{-1}$ is strictly convex. Then the quasi-arithmetic α -divergences induced by (f, g) for $\alpha \in [0, 1]$ is*

$$I_{\alpha}^{f,g}[p : q] = \begin{cases} \frac{1}{\alpha(1-\alpha)} \int \left(M_{1-\alpha}^f(p : q) - M_{1-\alpha}^g(p : q) \right) d\mu, & \alpha \in \mathbb{R} \setminus \{0, 1\}. \\ I_1^{f,g}[p : q] = \int \left(\frac{f(q)-f(p)}{f'(p)} - \frac{g(q)-g(p)}{g'(p)} \right) d\mu & \alpha = 1, \\ I_0^{f,g}[p : q] = I_1^{f,g}(q : p) = \int \left(\frac{f(p)-f(q)}{f'(q)} - \frac{g(p)-g(q)}{g'(q)} \right) d\mu, & \alpha = 0. \end{cases} \quad (58)$$

When $f(u) = f_A(u) = u$ ($M^f = A$) and $g(u) = f_G(u) = \log u$ ($M^g = G$), we get:

$$I_1^{A,G}[p : q] = \int \left(q - p - p \log \frac{q}{p} \right) d\mu = \text{KL}_e[p : q] = I_1[p : q], \quad (59)$$

the Kullback-Leibler divergence (KLD) extended to positive densities, and $I_0 = \text{KL}_e^*$ the reverse extended KLD.

We can rewrite the α -divergence $I_\alpha^{f,g}[p : q]$ for $\alpha \in (0, 1)$ as

$$I_\alpha^{f,g}[p : q] = \frac{1}{\alpha(1-\alpha)} \left(S_{1-\alpha}^f(p : q) - S_{1-\alpha}^g(p : q) \right), \quad (60)$$

where

$$S_\lambda^h(p : q) := \int M_\lambda^h(p : q) d\mu. \quad (61)$$

Zhang [39] (p. 188-189) considered the (A, M^ρ) α_A -divergences:

$$D_\alpha^\rho[p : q] = \frac{4}{1-\alpha^2} \int \left(\frac{1-\alpha}{2}p + \frac{1+\alpha}{2}q - \rho^{-1} \left(\frac{1-\alpha}{2}\rho(p) + \frac{1+\alpha}{2}\rho(q) \right) \right) d\mu. \quad (62)$$

The formula he obtained for $D_{\pm 1}^\rho(p : q)$:

$$D_1^\rho[p : q] = \int \left(p - q - (\rho^{-1})'(\rho(q))(\rho(p) - \rho(q)) \right) d\mu = D_{-1}^\rho[q : p], \quad (63)$$

is in accordance with our generic formula of Eq. 42 since $(\rho^{-1}(x))' = \frac{1}{\rho'(\rho^{-1}(x))}$. Notice that $A_\alpha \geq P_\alpha^r$ for $r \leq 1$: The arithmetic weighted mean dominates the weighted power means P^r when $r \leq 1$.

Furthermore, by imposing the homogeneity condition $I_\alpha^{A, M^\rho}[\lambda p : \lambda q] = \lambda I_\alpha^{A, M^\rho}[p : q]$ for $\lambda > 0$, Zhang [39] obtained the class of the (α_A, β_A) -divergences for $(\alpha_A, \beta_A) \in [-1, 1]^2$:

$$D_{\alpha_A, \beta_A}[p : q] = \frac{4}{1-\alpha_A^2} \frac{2}{1+\beta_A} \int \left(\frac{1-\alpha_A}{2}p + \frac{1+\alpha_A}{2}q - \left(\frac{1-\alpha_A}{2}p^{\frac{1-\beta_A}{2}} + \frac{1+\alpha_A}{2}q^{\frac{1-\beta_A}{2}} \right)^{\frac{2}{1-\beta_A}} \right) d\mu. \quad (64)$$

2.4 Generalized KL divergences as conformal Bregman divergences on monotone embeddings

We can rewrite the generalized KLDs $I_1^{f,g}$ as a *conformal Bregman representational divergence* [32, 33, 34] as follows:

Theorem 4. *The generalized KLDs $I_1^{f,g}$ divergences are conformal Bregman representational divergences:*

$$I_1^{f,g}[p : q] = \int \frac{1}{f'(p)} B_F(g(q) : g(p)) d\mu, \quad (65)$$

with $F = f \circ g^{-1}$ a strictly convex and differentiable Bregman convex generator.

Proof. For the Bregman strictly convex and differentiable generator $F = f \circ g^{-1}$, we expand the following conformal divergence:

$$\frac{1}{f'(p)} B_F(g(q) : g(p)) = \frac{1}{f'(p)} (F(g(q)) - F(g(p)) - (g(q) - g(p))F'(g(p))), \quad (66)$$

$$= \frac{1}{f'(p)} \left((f(q) - f(p)) - (g(q) - g(p)) \frac{f'(p)}{g'(p)} \right), \quad (67)$$

since $(g^{-1} \circ g)(x) = x$ and $F'(g(x)) = \frac{f'(x)}{g'(x)}$. It follows that

$$\frac{1}{f'(p)} B_F(g(q) : g(p)) = \frac{f(q) - f(p)}{f'(p)} - \frac{g(q) - g(p)}{g'(p)}, \quad (68)$$

$$= E_f(p, q) - E_g(p, q) = I_1^{f,g}[p : q]. \quad (69)$$

Hence, we easily check that $I_1^{f,g}[p : q] = \int \frac{1}{f'(p)} B_F(g(q) : g(p)) d\mu \geq 0$ since $f'(p) > 0$ and $B_F \geq 0$. \square

In general, for a functional generator f and a strictly monotonic representational function r , we can define the representational Bregman divergence [29] $B_{f \circ r^{-1}}(r(p) : r(q))$ provided that $F = f \circ r^{-1}$ is a Bregman generator (i.e., strictly convex and differentiable).

In [30], a generalization of the Bregman divergences was obtained using the *comparative convexity* induced by two abstract means M and N to define (M, N) -Bregman divergences as limit of scaled (M, N) -Jensen divergences. The skew (M, N) -Jensen divergences are defined for $\alpha \in (0, 1)$ by:

$$J_{F,\alpha}^{M,N}(p : q) = \frac{1}{\alpha(1-\alpha)} (N_\alpha(F(p), F(q)) - F(M_\alpha(p, q))), \quad (70)$$

where M_α and N_α are weighted means that should be *regular* [30] (i.e., homogeneous, symmetric, continuous and increasing in each variable). Then we can define the (M, N) -Bregman divergence as

$$\begin{aligned} B_F^{M,N}[p : q] &= \lim_{\alpha \rightarrow 1^-} J_{F,\alpha}^{M,N}(p : q), \\ &= \lim_{\alpha \rightarrow 1^-} \frac{1}{\alpha(1-\alpha)} (N_\alpha(F(p), F(q)) - F(M_\alpha(p, q))). \end{aligned}$$

The formula obtained in [30] for the quasi-arithmetic means M^f and M^g and a functional generator F that is (M^f, M^g) -convex is:

$$B_F^{f,g}(p : q) = \frac{g(F(p)) - g(F(q))}{g'(F(q))} - \frac{f(p) - f(q)}{f'(q)} F'(q), \quad (71)$$

$$= \frac{1}{f'(F(q))} B_{g \circ F \circ f^{-1}}(f(p) : f(q)) \geq 0. \quad (72)$$

This is a conformal divergence [32] that can be written using the E_f terms as:

$$B_F^{f,g}(p : q) = E_g(F(q), F(p)) - E_f(q, p)F'(q). \quad (73)$$

A function F is (M^f, M^g) -convex iff $g \circ F \circ f^{-1}$ is (ordinary) convex [30].

The information geometry induced by a Bregman divergence (or equivalently by its convex generator) is a dually flat space [2, 27]. The dualistic structure induced by a conformal Bregman representational divergence is related to conformal flattening [33, 34].

Following the work of Ohara [33, 34], the *geometric divergence* $\rho(p, r)$ (a contrast function in affine differential geometry) induced by a pair (L, M) of strictly monotone smooth functions between two distributions p and r of the d -dimensional probability simplex Δ_d is defined by (Eq. 28 in [33]):

$$\rho(p : r) = \frac{1}{\Lambda(r)} \sum_{i=1}^{d+1} \frac{L(p_i) - L(r_i)}{L'(r_i)} = \frac{1}{\Lambda(r)} \sum_{i=1}^{d+1} E_L(r_i, p_i), \quad (74)$$

where $\Lambda(r) = \sum_{i=1}^{d+1} \frac{1}{L'(p_i)} p_i$. Affine immersions [20] can be interpreted as special embeddings.

Let ρ be a divergence (contrast function) and $(\rho g, \rho \nabla, \rho \nabla^*)$ be the induced statistical manifold structure with

$${}^\rho g_{ij}(p) := -(\partial_i)_p (\partial_j)_p \rho(p, q)|_{q=p}, \quad (75)$$

$$\Gamma_{ij,k}(p) := -(\partial_i)_p (\partial_j)_p (\partial_k)_q \rho(p, q)|_{q=p}, \quad (76)$$

$$\Gamma_{ij,k}^*(p) := -(\partial_i)_p (\partial_j)_q (\partial_k)_q \rho(p, q)|_{q=p}, \quad (77)$$

where $(\partial_i)_s$ denotes the tangent vector at s of a vector field ∂_i .

Consider a conformal divergence $\rho_\kappa(p : q) = \kappa(q) \rho(p : q)$ for a positive function $\kappa(q) > 0$, called the conformal factor. Then the induced statistical manifold [12, 2] $(\rho_\kappa g, \rho_\kappa \nabla, \rho_\kappa \nabla^*)$ is 1-conformally equivalent to $(\rho g, \rho \nabla, \rho \nabla^*)$ and we have

$${}^{\rho_\kappa} g = \kappa {}^\rho g, \quad (78)$$

$$\rho_\kappa g(\rho_\kappa \nabla_X Y, Z) = \rho g(\rho \nabla_X Y, Z) - d(\log \kappa)(Z) \rho g(X, Y). \quad (79)$$

The dual affine connections $\rho_\kappa \nabla^*$ and $\rho \nabla^*$ are projectively equivalent [20] (and $\rho \nabla^*$ is said -1 -conformally flat).

Conformal flattening [33, 34] consists in choosing the conformal factor κ such that $(\rho_\kappa g, \rho_\kappa \nabla, \rho_\kappa \nabla^*)$ becomes a dually flat space [2] equipped with a canonical Bregman divergence.

Therefore it follows that the statistical manifolds induced by the 1-divergence $I_1^{f,g}$ is a representational 1-conformally flat statistical manifold.

3 The subfamily of homogeneous (r, s) -power α -divergences

In particular, we can define the (r, s) -power α -divergences from two power means $P_r = M^{\text{pow}_r}$ and $P_s = M^{\text{pow}_s}$ with $r > s$ (and $P_r \geq P_s$) with the family of generators $\text{pow}_l(u) = u^l$. Indeed, we check that $f_{rs}(u) := \text{pow}_r \circ \text{pow}_s^{-1}(u) = u^{\frac{r}{s}}$ is strictly convex on $(0, \infty)$ since $f_{rs}''(u) = \frac{r}{s} \left(\frac{r}{s} - 1\right) u^{\frac{r}{s}-2} > 0$ for $r > s$. Thus P_r and P_s are two QAMs which are both comparable and distinct. Table 1 lists the expressions of $E_r(p, q) := E_{\text{pow}_r}(p, q)$ obtained from the power mean generators $\text{pow}_r(u) = u^r$.

We conclude with the definition of the (r, s) -power α -divergences:

Corollary 1 (power α -divergences). *Given $r > s$, the α -power divergences are defined for $r > s$ and $r, s \neq 0$ by*

$$I_\alpha^{r,s}[p : q] = \begin{cases} \frac{1}{\alpha(1-\alpha)} \int \left((\alpha p^r + (1-\alpha)q^r)^{\frac{1}{r}} - (\alpha p^s + (1-\alpha)q^s)^{\frac{1}{s}} \right) d\mu, & \alpha \in \mathbb{R} \setminus \{0, 1\}. \\ I_1^{r,s}[p : q] = \int \left(\frac{q^r - p^r}{rp^{r-1}} - \frac{q^s - p^s}{sp^{s-1}} \right) d\mu & \alpha = 1, \\ I_0^{r,s}[p : q] = I_1^{r,s}(q : p) & \alpha = 0. \end{cases} \quad (80)$$

Power mean	$E_r(p, q)$
$P_r(r \in \mathbb{R} \setminus \{0\})$	$\frac{q^r - p^r}{rp^{r-1}}$
$Q(r = 2)$	$\frac{q^2 - p^2}{2p}$
$A(r = 1)$	$q - p$
$G(r = 0)$	$p \log \frac{q}{p}$
$H(r = -1)$	$-p^2 \left(\frac{1}{q} - \frac{1}{p} \right) = p - \frac{p^2}{q}$

Table 1: Expressions of the terms E_r for the family of power means P_r , $r \in \mathbb{R}$.

When $r = 0$, we get the following power α -divergences for $s < 0$:

$$I_\alpha^{r,s}[p : q] = \begin{cases} \frac{1}{\alpha(1-\alpha)} \int \left(p^\alpha q^{1-\alpha} - (\alpha p^s + (1-\alpha)q^s)^{\frac{1}{s}} \right) d\mu, & \alpha \in \mathbb{R} \setminus \{0, 1\}. \\ I_1^{r,s}[p : q] = \int \left(p \log \frac{q}{p} - \frac{q^s - p^s}{sp^{s-1}} \right) d\mu & \alpha = 1, \\ I_0^{r,s}[p : q] = I_1^{r,s}[q : p] & \alpha = 0. \end{cases} \quad (81)$$

When $s = 0$, we get the following power α -divergences for $r > 0$:

$$I_\alpha^{r,s}[p : q] = \begin{cases} \frac{1}{\alpha(1-\alpha)} \int \left((\alpha p^r + (1-\alpha)q^r)^{\frac{1}{r}} - p^\alpha q^{1-\alpha} \right) d\mu, & \alpha \in \mathbb{R} \setminus \{0, 1\}. \\ I_1^{r,s}[p : q] = \int \left(\frac{q^r - p^r}{rp^{r-1}} - p \log \frac{q}{p} \right) d\mu & \alpha = 1, \\ I_0^{r,s}[p : q] = I_1^{r,s}[q : p] & \alpha = 0. \end{cases} \quad (82)$$

In particular, we get the following family of (A, H) α -divergences:

$$I_\alpha^{A,H}[p : q] = I_\alpha^{1,-1}[p : q] = \begin{cases} \frac{1}{\alpha(1-\alpha)} \int \left(\alpha p + (1-\alpha)q - \frac{pq}{\alpha q + (1-\alpha)p} \right) d\mu, & \alpha \in \mathbb{R} \setminus \{0, 1\}. \\ I_1^{1,-1}[p : q] = \int \left(q - 2p + \frac{p^2}{q} \right) d\mu & \alpha = 1, \\ I_0^{1,-1}[p : q] = I_1^{1,-1}(q : p) & \alpha = 0. \end{cases}, \quad (83)$$

and the family of (G, H) α -divergences:

$$I_\alpha^{G,H}[p : q] = I_\alpha^{0,-1}(p : q) = \begin{cases} \frac{1}{\alpha(1-\alpha)} \int \left(p^\alpha q^{1-\alpha} - \frac{pq}{\alpha q + (1-\alpha)p} \right) d\mu, & \alpha \in \mathbb{R} \setminus \{0, 1\}. \\ I_1^{0,-1}[p : q] = \int \left(p \log \frac{q}{p} - p + \frac{p^2}{q} \right) d\mu & \alpha = 1, \\ I_0^{0,-1}[p : q] = I_1^{0,-1}[q : p] & \alpha = 0. \end{cases}. \quad (84)$$

The (r, s) -power α -divergences for $r, s \neq 0$ yield homogeneous divergences: $I_\alpha^{r,s}[\lambda p : \lambda q] = \lambda I_\alpha^{r,s}[p : q]$ for any $\lambda > 0$ because the power means are homogeneous: $P_\alpha^r(\lambda x, \lambda y) = \lambda P_\alpha^r(x, y) = \lambda x P_\alpha^r(1, \frac{y}{x})$. Thus the $I_\alpha^{r,s}$ -divergences are Csiszár f -divergences [10]

$$I_\alpha^{r,s}[p : q] = \int p(x) f_{r,s} \left(\frac{q(x)}{p(x)} \right) d\mu \quad (85)$$

for the generator

$$f_{r,s}(u) = \frac{1}{\alpha(1-\alpha)} (P_\alpha^r(1, u) - P^s(1, u)). \quad (86)$$

Thus the family of (r, s) -power α -divergences are *homogeneous divergences*:

$$I_\alpha^{r,s}[\lambda p : \lambda q] = \lambda I_\alpha^{r,s}[p : q], \quad \forall \lambda > 0 \quad (87)$$

4 Conclusion, discussion and perspectives

For two comparable strict means $M \geq N$, one can define the (M, N) -divergence as

$$I^{M,N}[p : q] := 4 \int (M(p, q) - N(p, q)) d\mu. \quad (88)$$

When the property of strict comparable means extend to their induced weighted means M_α and N_α (i.e., $M_\alpha \geq N_\alpha$), one can further define the (M, N) α -divergences for $\alpha \in (0, 1)$:

$$I_\alpha^{M,N}[p : q] := \frac{1}{\alpha(1-\alpha)} \int (M_{1-\alpha}(p, q) - N_{1-\alpha}(p, q)) d\mu, \quad (89)$$

so that $I^{M,N}[p : q] = I_{\frac{1}{2}}^{M,N}[p : q]$. When the weighted means are symmetric, the reference duality holds (i.e., $I_\alpha^{M,N}[q : p] = I_{1-\alpha}^{M,N}[p : q]$), and we can define the (M, N) -equivalent of the Kullback-Leibler divergence, i.e., the (M, N) 1-divergence, as the limit case (when it exists): $I_1^{M,N}[p : q] = \lim_{\alpha \rightarrow 1} I_\alpha^{M,N}[p : q]$.

We proved that the quasi-arithmetic weighted means M_α^f and M_α^g are strictly comparable whenever $f \circ g^{-1}$ is *strictly* convex. In the limit cases of $\alpha \rightarrow 0$ and $\alpha \rightarrow 1$, we reported a closed-form formula for the equivalent of the Kullback-Leibler divergence and the reverse Kullback-Leibler divergence. We reported closed-form formula for the quasi-arithmetic α -divergences $I_\alpha^{f,g}(p : q)$ for $\alpha \in [0, 1]$ (Theorem 3), and for the subfamily of homogeneous (r, s) -power α -divergences $I_\alpha^{r,s}(p : q)$ induced by power means (Corollary 1). The ordinary (A, G) α -divergences, the (A, H) α -divergences and the (G, H) α -divergences are examples of (r, s) -power α -divergences for $(r, s) = (1, 0)$, $(r, s) = (1, -1)$ and $(r, s) = (0, -1)$, respectively.

Generalized α -divergences may prove useful in reporting closed-form formula between parametric densities: For example, consider the ordinary α -divergences between two scale Cauchy densities $p_1(x) = \frac{1}{\pi} \frac{s_1}{x^2 + s_1^2}$ and $p_2(x) = \frac{1}{\pi} \frac{s_2}{x^2 + s_2^2}$: There is no obvious closed-form for the ordinary α -divergences but we can report easily a closed-form for the (A, H) α -divergences following the calculus reported in [26]:

$$I_\alpha^{A,H}[p_1 : p_2] = \frac{1}{\alpha(1-\alpha)} \left(1 - \int H_{1-\alpha}(p_1(x), p_2(x)) d\mu(x) \right), \quad (90)$$

$$= \frac{1}{\alpha(1-\alpha)} \left(1 - \frac{s_1 s_2}{(\alpha s_1 + (1-\alpha) s_2) s_{1-\alpha}} \right), \quad (91)$$

with $s_\alpha = \sqrt{\frac{\alpha s_1 s_2^2 + (1-\alpha) s_2 s_1^2}{\alpha s_1 + (1-\alpha) s_2}}$. In general, instead of considering ordinary α -divergences in applications, one may consider the (r, s) -power α -divergences, and tune the three parameters (r, s, α) according to the various tasks (say, by cross-validation in supervised machine learning tasks). We note that the quasi-arithmetic means have been recently considered by Eguchi et al. [13] to define a novel non-parametric dualistic structure of information geometry via generalizations of the e -geodesics and the m -geodesics.

The elucidation the (f, g) α -geometry of these generalized α -divergences is left for future work. For the limit cases of $\alpha \rightarrow 0$ or of $\alpha \rightarrow 1$, we proved that the limit KL-type divergences amount to a conformal Bregman divergence on a monotone embedding, and briefly showed the connection with conformal divergences and conformal flattening. The geometry of conformal flattening [33, 34] and the relationships with the (ρ, τ) -monotone embeddings [24] shall be further studied.

Applications of (f, g) α -divergences to center-based clustering [31] and to generalized α -divergences in positive-definite matrix spaces [2] shall also be considered in future work. The quasi-arithmetic weighted means are convex if and only if the generators are differentiable with positive first derivatives with corresponding functions $-E_f$ of Eq. 41 convex (Theorem 4 of [6], i.e., convexity of the quasi-arithmetic weighted means does *not* depend on the weights). For example, when both quasiarithmetic means are convex means, the quasi-arithmetic α -divergence is the difference of two convex mean functions, and the k -means centroid computation amounts to solve a *Difference of Convex* (DC) program which can be solved by the smooth DC Algorithm, DCA, called the Convex-ConCave Procedure [28]. Similarly, when $\alpha \in \{0, 1\}$, we get a DC program since $I_\alpha^{f,g}$ writes as a difference of convex functions.

References

- [1] Vahideh Ahrari, Arezou Habibirad, and S Baratpour. Exponentiality test based on alpha-divergence and gamma-divergence. *Communications in Statistics-Simulation and Computation*, 48(4):1138–1152, 2019.
- [2] S. Amari. *Information Geometry and Its Applications*. Applied Mathematical Sciences. Springer Japan, 2016.
- [3] Shun-ichi Amari. Integration of stochastic models by minimizing α -divergence. *Neural computation*, 19(10):2780–2796, 2007.
- [4] Michèle Basseville. Divergence measures for statistical data processing — an annotated bibliography. *Signal Processing*, 93(4):621–633, 2013.
- [5] Peter S Bullen, Dragoslav S Mitrinovic, and M Vasic. *Means and their Inequalities*, volume 31. Springer Science & Business Media, 2013.
- [6] Jacek Chudziak, Dorota Glazowska, Justyna Jarczyk, and Witold Jarczyk. On weighted quasi-arithmetic means which are convex. *Mathematical Inequalities & Applications*, 22(4):1123–1136, 2019.
- [7] Andrzej Cichocki and Shun-ichi Amari. Families of alpha-beta-and gamma-divergences: Flexible and robust measures of similarities. *Entropy*, 12(6):1532–1568, 2010.
- [8] Andrzej Cichocki, Hyekyoung Lee, Yong-Deok Kim, and Seungjin Choi. Non-negative matrix factorization with α -divergence. *Pattern Recognition Letters*, 29(9):1433–1440, 2008.
- [9] Thomas M. Cover and Joy A. Thomas. *Elements of information theory*. John Wiley & Sons, 2012.
- [10] Imre Csiszár. Information-type measures of difference of probability distributions and indirect observation. *Studia Scientiarum Mathematicarum Hungarica*, 2:229–318, 1967.
- [11] Bruno De Finetti. Sul concetto di media. *Istituto italiano degli attuari*, 3:369–396, 1931.
- [12] Shinto Eguchi. Geometry of minimum contrast. *Hiroshima Mathematical Journal*, 22(3):631–647, 1992.

- [13] Shinto Eguchi, Osamu Komori, and Atsumi Ohara. Information geometry associated with generalized means. In *Information Geometry and its Applications IV*, pages 279–295. Springer, 2016.
- [14] Alison L. Gibbs and Francis Edward Su. On choosing and bounding probability metrics. *International statistical review*, 70(3):419–435, 2002.
- [15] G.H. Hardy, J.E. Littlewood, and G. Pólya. *Inequalities*. Cambridge Mathematical Library. Cambridge University Press, 1988.
- [16] Otto Ludwig Holder. Über einen Mittelwertssatz. *Nachr. Akad. Wiss. Gottingen Math.-Phys. Kl.*, pages 38–47, 1889.
- [17] Asif Iqbal and Abd-Krim Seghouane. An α -divergence-based approach for robust dictionary learning. *IEEE Transactions on Image Processing*, 28(11):5729–5739, 2019.
- [18] Justyna Jarczyk. When Lagrangean and quasi-arithmetic means coincide. *J. Inequal. Pure Appl. Math.*, 8:71, 2007.
- [19] Andrey Nikolaevich Kolmogorov. Sur la notion de moyenne. *Acad. Naz. Lincei Mem. Cl. Sci. His. Mat. Natur. Sez.*, 12:388–391, 1930.
- [20] Takashi Kurose. On the divergences of 1-conformally flat statistical manifolds. *Tohoku Mathematical Journal, Second Series*, 46(3):427–433, 1994.
- [21] Gyula Maksa and Zsolt Páles. Remarks on the comparison of weighted quasi-arithmetic means. *Colloquium Mathematicae*, 120(1):77–84, 2010.
- [22] Yuzo Maruyama, Takeru Matsuda, and Toshio Ohnishi. Harmonic Bayesian prediction under α -divergence. *IEEE Transactions on Information Theory*, 65(9):5352–5366, 2019.
- [23] Mitio Nagumo. Über eine klasse der mittelwerte. *Japanese journal of mathematics: transactions and abstracts*, 7(0):71–79, 1930.
- [24] Jan Naudts and Jun Zhang. Rho–tau embedding and gauge freedom in information geometry. *Information Geometry*, 1(1):79–115, 2018.
- [25] Constantin P. Niculescu and Lars-Erik Persson. *Convex functions and their applications: A contemporary approach*. Springer Science & Business Media, 2006. First edition.
- [26] Frank Nielsen. Generalized Bhattacharyya and Chernoff upper bounds on Bayes error using quasi-arithmetic means. *Pattern Recognition Letters*, 42:25–34, 2014.
- [27] Frank Nielsen. An elementary introduction to information geometry. *arXiv preprint arXiv:1808.08271*, 2018.
- [28] Frank Nielsen and Sylvain Boltz. The Burbea-Rao and Bhattacharyya centroids. *IEEE Transactions on Information Theory*, 57(8):5455–5466, 2011.
- [29] Frank Nielsen and Richard Nock. The dual Voronoi diagrams with respect to representational Bregman divergences. In *Sixth International Symposium on Voronoi Diagrams (ISVD)*, pages 71–78. IEEE, 2009.

- [30] Frank Nielsen and Richard Nock. Generalizing skew Jensen divergences and Bregman divergences with comparative convexity. *IEEE Signal Processing Letters*, 24(8):1123–1127, 2017.
- [31] Frank Nielsen, Richard Nock, and Shun-ichi Amari. On clustering histograms with k -means by using mixed α -divergences. *Entropy*, 16(6):3273–3301, 2014.
- [32] Richard Nock, Frank Nielsen, and Shun-ichi Amari. On conformal divergences and their population minimizers. *IEEE Transactions on Information Theory*, 62(1):527–538, 2015.
- [33] Atsumi Ohara. Conformal flattening for deformed information geometries on the probability simplex. *Entropy*, 20(3):186, 2018.
- [34] Atsumi Ohara. Conformal flattening on the probability simplex and its applications to voronoi partitions and centroids. In *Geometric Structures of Information*, pages 51–68. Springer, 2019.
- [35] Zsolt Páles and Amr Zakaria. On the equality of bajraktarević means to quasi-arithmetic means. *Results in Mathematics*, 75(1):19, 2020.
- [36] Alfréd Rényi. On measures of entropy and information. In *Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability*. The Regents of the University of California, 1961. Volume 1: Contributions to the Theory of Statistics.
- [37] Auxiliadora Sarmiento, Irene Fondón, Iván Durán-Díaz, and Sergio Cruces. Centroid-based clustering with $\alpha\beta$ -divergences. *Entropy*, 21(2):196, 2019.
- [38] Junichiro Wada and Yuta Kamahara. Studying malapportionment using α -divergence. *Mathematical Social Sciences*, 93:77–89, 2018.
- [39] Jun Zhang. Divergence function, duality, and convex analysis. *Neural Computation*, 16(1):159–195, 2004.