New binary self-dual codes of lengths 56, 62, 78, 92 and 94 from a bordered construction

J. Gildea, A. Korban and A. M. Roberts Department of Physical, Mathematical and Engineering Sciences University of Chester Exton Park Chester CH1 4AR United Kingdom

> A. Tylyshchak Department of Algebra Uzhgorod National University Uzhgorod Ukraine

Keywords: Binary self-dual codes, Bordered constructions, Gray maps, Extremal codes, Best known codes

2020 MSC: 94B05, 15B10, 15B33

Abstract

In this paper, we present a new bordered construction for self-dual codes which employs λ -circulant matrices. We give the necessary conditions for our construction to produce self-dual codes over a finite commutative Frobenius ring of characteristic 2. Moreover, using our bordered construction together with the well-known building-up and neighbour methods, we construct many binary self-dual codes of lengths 56, 62, 78, 92 and 94 with parameters in their weight enumerators that were not known in the literature before.

1 Introduction

The study of self-dual codes over finite fields and rings is an active area of research in coding theory. This is mainly due to their connections to other areas in mathematics such as combinatorics, design theory and number theory. In [1], it is shown that one can produce interesting designs using self-dual codes over fields. In [3], it is shown

E-mail addresses: j.gildea@chester.ac.uk (J. Gildea), adrian3@windowslive.com (A. Korban), adammichaelroberts@outlook.com (A. M. Roberts), alxtlk@bigmir.net (A. Tylyshchak)

that one of the most powerful techniques for producing optimal unimodular lattices uses self-dual codes over rings. Moreover, as seen in [29], the well-known proof of the non-existence of the projective plane of order 10 used the theory of binary self-dual codes.

Finding methods for constructing self-dual codes, classifying self-dual codes and determining the largest minimum distance among all self-dual codes are open problems in coding theory. Many researchers have employed different techniques to construct self-dual codes. For example, in [18], the authors establish a strong connection between group rings and self-dual codes. They prove that a group ring element corresponds to a self-dual code if and only if it is a unitary unit and they use this fact to construct many new extremal binary self-dual codes of length 68. In [35, 36, 37, 38, 39], the authors employ a particular technique which guarantees that the constructed self-dual codes have a fixed automorphism of odd prime order.

A powerful method for finding new binary self-dual codes is to consider bordered matrix constructions. There exist several papers in which such constructions are used to obtain binary self-dual codes of different lengths with new parameters in their weight enumerators. Examples of bordered constructions can be found in [15, 14, 20]. In this work, we continue in this direction and we present a new bordered matrix construction that we employ to obtain many binary self-dual codes of lengths 56, 62, 78, 92 and 94 with weight enumerators that were not known in the literature before. Moreover, we present the necessary conditions that the construction needs to satisfy in order to produce a self-dual code over a finite commutative Frobenius ring of characteristic 2.

The paper is organised as follows. In Section 2, we give the standard definitions and results on self-dual codes, the alphabets we use, Gray maps and some special matrices that we use in this work. In Section 3, we present the main construction and show under what conditions it produces self-dual codes over a finite commutative Frobenius ring of characteristic 2. In Section 4, we present our computational results. Namely, we tabulate all new binary self-dual codes that we obtain by a direct application of our main construction together with the building-up and neighbour constructions. We finish with concluding remarks and directions for possible future research.

2 Preliminaries

2.1 Self-Dual Codes

Let R be a commutative Frobenius ring (see [8] for a full description of Frobenius rings and codes over Frobenius rings). Throughout this work, we always assume Rhas unity. A code C of length n over R is a subset of R^n whose elements are called codewords. If C is a submodule of R^n , then we say that C is linear. Let $\mathbf{x}, \mathbf{y} \in R^n$ where $\mathbf{x} = (x_1, x_2, \ldots, x_n)$ and $\mathbf{y} = (y_1, y_2, \ldots, y_n)$. The (Euclidean) dual C^{\perp} of C is given by

$$\mathcal{C}^{\perp} = \{ \mathbf{x} \in \mathbb{R}^n : \langle \mathbf{x}, \mathbf{y} \rangle = 0, \forall \mathbf{y} \in \mathcal{C} \}$$

where $\langle \cdot, \cdot \rangle$ denotes the Euclidean inner product defined by

$$\langle \mathbf{x}, \mathbf{y} \rangle = \sum_{i=1}^{n} x_i y_i$$

and we say that \mathcal{C} is self-orthogonal if $\mathcal{C} \subseteq \mathcal{C}^{\perp}$ and self-dual if $\mathcal{C} = \mathcal{C}^{\perp}$.

An upper bound on the minimum (Hamming) distance of a doubly-even (Type II) binary self-dual code was given in [31] and likewise for a singly-even (Type I) binary self-dual code in [32]. Let $d_{\rm I}(n)$ and $d_{\rm II}(n)$ be the minimum distance of a Type I and Type II binary self-dual code of length n, respectively. Then

and

$$d_{\mathrm{II}}(n) \le 4\lfloor n/24 \rfloor + 4$$

$$d_{\mathrm{I}}(n) \leq \begin{cases} 4\lfloor n/24 \rfloor + 2, & \text{if } n \equiv 0 \pmod{24}, \\ 4\lfloor n/24 \rfloor + 4, & \text{if } n \not\equiv 22 \pmod{24}, \\ 4\lfloor n/24 \rfloor + 6, & \text{if } n \equiv 22 \pmod{24}. \end{cases}$$

A self-dual code whose minimum distance meets its corresponding bound is called *extremal*. A self-dual code with the highest minimum distance for its length is said to be *optimal*. Extremal codes are necessarily optimal but optimal codes are not necessarily extremal. A *best known* self-dual code is a self-dual code with the highest known minimum distance for its length.

2.2 Alphabets

In this paper, we consider the alphabets \mathbb{F}_2 , $\mathbb{F}_2 + u\mathbb{F}_2$, $\mathbb{F}_2 + u\mathbb{F}_2 + v\mathbb{F}_2 + uv\mathbb{F}_2$ and $\mathbb{F}_4 + u\mathbb{F}_4$.

Define

$$\mathbb{F}_2 + u\mathbb{F}_2 = \{a + bu : a, b \in \mathbb{F}_2, u^2 = 0\},\$$

then $\mathbb{F}_2 + u\mathbb{F}_2$ is a commutative ring of order 4 and characteristic 2 such that $\mathbb{F}_2 + u\mathbb{F}_2 \cong \mathbb{F}_2[u]/\langle u^2 \rangle$.

Define

$$\mathbb{F}_{2} + u\mathbb{F}_{2} + v\mathbb{F}_{2} + uv\mathbb{F}_{2} = \{a + bu + cv + duv : a, b, c, d \in \mathbb{F}_{2}, u^{2} = v^{2} = uv + vu = 0\}$$

then $\mathbb{F}_2 + u\mathbb{F}_2 + v\mathbb{F}_2 + uv\mathbb{F}_2$ is a commutative ring of order 16 and characteristic 2 such that $\mathbb{F}_2 + u\mathbb{F}_2 + v\mathbb{F}_2 + uv\mathbb{F}_2 \cong \mathbb{F}_2[u, v]/\langle u^2, v^2, uv + vu \rangle$. Note that $\mathbb{F}_2 + u\mathbb{F}_2 + v\mathbb{F}_2 + uv\mathbb{F}_2$ can be viewed as an extension of $\mathbb{F}_2 + u\mathbb{F}_2$ and so we can also define

$$\mathbb{F}_{2} + u\mathbb{F}_{2} + v\mathbb{F}_{2} + uv\mathbb{F}_{2} = \{a + bv : a, b \in \mathbb{F}_{2} + u\mathbb{F}_{2}, v^{2} = 0\}$$

We define $\mathbb{F}_4 \cong \mathbb{F}_2[\omega]/\langle \omega^2 + \omega + 1 \rangle$ so that

$$\mathbb{F}_4 = \{a\omega + b(1+\omega) : a, b \in \mathbb{F}_2, \omega^2 + \omega + 1 = 0\}.$$

Define

$$\mathbb{F}_4 + u\mathbb{F}_4 = \{a + bu : a, b \in \mathbb{F}_4, u^2 = 0\}$$

then $\mathbb{F}_4 + u\mathbb{F}_4$ is a commutative ring of order 16 and characteristic 2 such that $\mathbb{F}_4 + u\mathbb{F}_4 \cong \mathbb{F}_4[u]/\langle u^2 \rangle \cong \mathbb{F}_2[\omega, u]/\langle \omega^2 + \omega + 1, u^2, \omega u + u\omega \rangle$. Note that $\mathbb{F}_4 + u\mathbb{F}_4$ can be viewed as an extension of $\mathbb{F}_2 + u\mathbb{F}_2$ and so we can also define

$$\mathbb{F}_4 + u\mathbb{F}_4 = \{a\omega + b(1+\omega) : a, b \in \mathbb{F}_2 + u\mathbb{F}_2, \omega^2 + \omega + 1 = 0\}.$$

We recall the following Gray maps from [10, 17, 30, 12]:

$$\begin{split} \varphi_{\mathbb{F}_{2}+u\mathbb{F}_{2}} &: (\mathbb{F}_{2}+u\mathbb{F}_{2})^{n} \to \mathbb{F}_{2}^{2n} \\ & a+bu \mapsto (b,a+b), \, a, b \in \mathbb{F}_{2}^{n}, \\ \varphi_{\mathbb{F}_{4}+u\mathbb{F}_{4}} &: (\mathbb{F}_{4}+u\mathbb{F}_{4})^{n} \to \mathbb{F}_{4}^{2n} \\ & a+bu \mapsto (b,a+b), \, a, b \in \mathbb{F}_{4}^{n}, \\ \phi_{\mathbb{F}_{2}+u\mathbb{F}_{2}+v\mathbb{F}_{2}+uv\mathbb{F}_{2}} &: (\mathbb{F}_{2}+u\mathbb{F}_{2}+v\mathbb{F}_{2}+uv\mathbb{F}_{2})^{n} \to (\mathbb{F}_{2}+u\mathbb{F}_{2})^{2n} \\ & a+bv \mapsto (b,a+b), \, a, b \in (\mathbb{F}_{2}+u\mathbb{F}_{2})^{n}, \\ \psi_{\mathbb{F}_{4}} &: \mathbb{F}_{4}^{n} \to \mathbb{F}_{2}^{2n} \\ & a\omega + b(1+\omega) \mapsto (a,b), \, a, b \in \mathbb{F}_{2}^{n}, \\ \psi_{\mathbb{F}_{4}+u\mathbb{F}_{4}} &: (\mathbb{F}_{4}+u\mathbb{F}_{4})^{n} \to (\mathbb{F}_{2}+u\mathbb{F}_{2})^{2n} \\ & a\omega + b(1+\omega) \mapsto (a,b), \, a, b \in (\mathbb{F}_{2}+u\mathbb{F}_{2})^{n} \end{split}$$

and we note that these Gray maps preserve orthogonality in their respective alphabets (see [30] for details). If $C \subseteq (\mathbb{F}_4 + u\mathbb{F}_4)^n$, then the binary codes $\varphi_{\mathbb{F}_2+u\mathbb{F}_2} \circ \psi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$ and $\psi_{\mathbb{F}_4} \circ \varphi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$ are equivalent to each other (see [17, 30] for details). The Lee weight of a codeword is defined to be the Hamming weight of its binary image under any of the previously mentioned compositions of maps. A self-dual code in \mathbb{R}^n where \mathbb{R} is equipped with a Gray map to the binary Hamming space is said to be of Type II if the Lee weights of all codewords are multiples of 4, otherwise it is said to be of Type I.

Proposition 2.1. ([30]) Let C be a code over $\mathbb{F}_4 + u\mathbb{F}_4$. If C is self-orthogonal, then $\psi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$ and $\varphi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$ are self-orthogonal. The code C is a Type I (resp. Type II) code over $\mathbb{F}_4 + u\mathbb{F}_4$ if and only if $\varphi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$ is a Type I (resp. Type II) \mathbb{F}_4 -code if and only if $\psi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$ is a Type I (resp. Type II) ($\mathbb{F}_2 + u\mathbb{F}_2$)-code. Furthermore, the minimum Lee weight of C is the same as the minimum Lee weight of $\psi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$ and $\varphi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$.

The next corollary follows immediately from Proposition 2.1.

Corollary 2.2. Let C be a self-dual code over $\mathbb{F}_4 + u\mathbb{F}_4$ of length n and minimum Lee distance d. Then $\varphi_{\mathbb{F}_2+u\mathbb{F}_2} \circ \psi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$ is a binary self-dual [4n, 2n, d] code. Moreover, the Lee weight enumerator of C is equal to the Hamming weight enumerator of $\varphi_{\mathbb{F}_2+u\mathbb{F}_2} \circ$ $\psi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$. If C is a Type I (resp. Type II) code, then $\varphi_{\mathbb{F}_2+u\mathbb{F}_2} \circ \psi_{\mathbb{F}_4+u\mathbb{F}_4}(C)$ is a Type I (resp. Type II) code.

We also have the following proposition from [13]:

Proposition 2.3. ([13]) Let C be a self-dual code over $\mathbb{F}_2 + u\mathbb{F}_2 + v\mathbb{F}_2 + uv\mathbb{F}_2$ of length n and minimum Lee distance d. Then $\varphi_{\mathbb{F}_2+u\mathbb{F}_2} \circ \phi_{\mathbb{F}_2+u\mathbb{F}_2+v\mathbb{F}_2+uv\mathbb{F}_2}(C)$ is a binary [4n, 2n, d] self-dual code. Moreover, the Lee weight enumerator of C is equal to the Hamming weight enumerator of $\varphi_{\mathbb{F}_2+u\mathbb{F}_2} \circ \phi_{\mathbb{F}_2+u\mathbb{F}_2+v\mathbb{F}_2+uv\mathbb{F}_2}(C)$. If C is a Type I (resp. Type II) code, then $\varphi_{\mathbb{F}_2+u\mathbb{F}_2} \circ \phi_{\mathbb{F}_2+u\mathbb{F}_2+vv\mathbb{F}_2+uv\mathbb{F}_2}(C)$ is a Type II) code.

2.3 Special Matrices

We now define and discuss the properties of some special matrices which we use in our work. Let $\mathbf{a} = (a_0, a_1, \dots, a_{n-1}) \in \mathbb{R}^n$ where \mathbb{R} is a commutative ring and let

$$A = \begin{pmatrix} a_0 & a_1 & a_2 & \cdots & a_{n-1} \\ \lambda a_{n-1} & a_0 & a_1 & \cdots & a_{n-2} \\ \lambda a_{n-2} & \lambda a_{n-1} & a_0 & \cdots & a_{n-3} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \lambda a_1 & \lambda a_2 & \lambda a_3 & \cdots & a_0 \end{pmatrix}$$

where $\lambda \in R$. Then A is called the λ -circulant matrix generated by **a**, denoted by $A = \operatorname{circ}_{\lambda}(\mathbf{a})$. If $\lambda = 1$, then A is called the circulant matrix generated by **a** and is more simply denoted by $A = \operatorname{circ}(\mathbf{a})$. If we define the matrix

$$P_{\lambda} = \begin{pmatrix} \mathbf{0} & I_{n-1} \\ \lambda & \mathbf{0} \end{pmatrix},$$

then it follows that $A = \sum_{i=0}^{n-1} a_i P_{\lambda}^i$. Clearly, the sum of any two λ -circulant matrices is also a λ -circulant matrix. If $B = \operatorname{circ}_{\lambda}(\mathbf{b})$ where $\mathbf{b} = (b_0, b_1, \dots, b_{n-1}) \in \mathbb{R}^n$, then $AB = \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} a_i b_j P_{\lambda}^{i+j}$. Since $P_{\lambda}^n = \lambda I_n$ there exist $c_k \in \mathbb{R}$ such that $AB = \sum_{k=0}^{n-1} c_k P_{\lambda}^k$ so that AB is also λ -circulant. In fact, it is true that

$$c_k = \sum_{\substack{[i+j]_n = k \\ i+j < n}} a_i b_j + \sum_{\substack{[i+j]_n = k \\ i+j \ge n}} \lambda a_i b_j = \mathbf{x}_1 \mathbf{y}_{k+1}$$

for $k \in \{0, \ldots, n-1\}$, where \mathbf{x}_i and \mathbf{y}_i respectively denote the i^{th} row and column of A and B and $[i+j]_n$ denotes the smallest non-negative integer such that $[i+j]_n \equiv i+j \pmod{n}$. From this, we can see that λ -circulant matrices commute multiplicatively and in fact the set of λ -circulant matrices over a commutative ring of fixed size is itself a commutative ring. Moreover, if λ is a unit in R, then A^T is λ^{-1} -circulant such that $A^T = a_0 I_n + \lambda \sum_{i=1}^{n-1} a_{n-i} P_{\lambda^{-1}}^i$. It follows then that AA^T is λ -circulant if and only if λ is involutory in R, i.e. $\lambda^2 = 1$.

Let J_n be an $n \times n$ matrix over R whose $(i, j)^{\text{th}}$ entry is 1 if i + j = n + 1 and 0 if otherwise. Then J_n is called the $n \times n$ exchange matrix and corresponds to the row-reversed (or column-reversed) version of I_n . We see that J_n is both symmetric and involutory, i.e. $J_n = J_n^T$ and $J_n^2 = I_n$. For any matrix $A \in R^{m \times n}$, premultiplying A by J_m and postmultiplying A by J_n inverts the order in which the rows and columns of A appear, respectively. Namely, the $(i, j)^{\text{th}}$ entries of $J_m A$ and AJ_n are the $([1-i]_m, j)^{\text{th}}$ and $(i, [1-j]_n)^{\text{th}}$ entries of A, respectively. Note that $[i + j]_n$ corresponds to the $(i + 1, j + 1)^{\text{th}}$ entry of the matrix $J_n V$ where $V = \text{circ}(n-1, 0, 1, \ldots, n-2)$ for $i, j \in \{0, \ldots, n-1\}$.

For a λ -circulant matrix A of size $n \times n$, it is easy to see that the sum of entries in the i^{th} row of A is equal to the sum of entries in the $(n - i + 1)^{\text{th}}$ column of A. Thus, for $\boldsymbol{\xi} = (\xi, \xi, \dots, \xi) \in \mathbb{R}^n$, we have $\boldsymbol{\xi}A = \boldsymbol{\xi}A^T J_n$.

3 The Construction

In this section, we present our technique for constructing self-dual codes. We will hereafter always assume R is a finite commutative Frobenius ring of characteristic 2.

Lemma 3.1. Let

$$X = \begin{pmatrix} AC & B \\ B^T C & A^T \end{pmatrix},$$

where $A = \operatorname{circ}_{\lambda}(\mathbf{a})$, $B = \operatorname{circ}_{\lambda}(\mathbf{b})$ and $C = \operatorname{circ}_{\mu}(\mathbf{c})$ with $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{R}^{n}$ and $\lambda, \mu \in \mathbb{R}$: $\lambda^{2} = \mu^{2} = 1$. If $CC^{T} = I_{n}$, then $XX^{T} = I_{2n}$ if and only if $AA^{T} + BB^{T} = I_{n}$.

Proof. Since $\lambda^2 = 1$ by assumption, we have that A and B as well as their transpositions all commute with one another multiplicatively. We have

$$XX^{T} = \begin{pmatrix} AC & B \\ B^{T}C & A^{T} \end{pmatrix} \begin{pmatrix} C^{T}A^{T} & C^{T}B \\ B^{T} & A \end{pmatrix} = \begin{pmatrix} x_{1,1} & x_{1,2} \\ x_{1,2}^{T} & x_{2,2} \end{pmatrix}$$

where

$$x_{1,1} = ACC^{T}A^{T} + BB^{T},$$

$$x_{1,2} = ACC^{T}B + BA,$$

$$x_{2,2} = B^{T}CC^{T}B + A^{T}A$$

and since $CC^T = I_n$ by assumption, we get

$$x_{1,1} = AA^T + BB^T,$$

$$x_{1,2} = AB + BA = 2AB = \mathbf{0},$$

$$x_{2,2} = B^T B + A^T A = AA^T + BB^T$$

Therefore, $XX^T = I_{2n}$ if and only if $AA^T + BB^T = I_n$.

Theorem 3.2. Let $n \in \mathbb{Z}^+$ such that n is odd and let

$$G = \begin{pmatrix} \mathbf{v} & \mathbf{0} & \xi_3 & \xi_4 \\ \hline I_{2n} & X & \mathbf{v}^T & \mathbf{v}^T \end{pmatrix}, \quad where \ X = \begin{pmatrix} AC & B \\ B^TC & A^T \end{pmatrix}$$

where $\mathbf{v} = (\boldsymbol{\xi}_1, \boldsymbol{\xi}_2)$ with $\boldsymbol{\xi}_i = (\xi_i, \xi_i, \dots, \xi_i) \in \mathbb{R}^n$ for $i \in \{1, 2\}$ and $\xi_3, \xi_4 \in \mathbb{R}$ also with $A = \operatorname{circ}_{\lambda}(\mathbf{a})$, $B = \operatorname{circ}_{\lambda}(\mathbf{b})$ and $C = \operatorname{circ}_{\mu}(\mathbf{c})$ for $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{R}^n$ and $\lambda, \mu \in \mathbb{R}$: $\lambda^2 = \mu^2 = 1$. If $CC^T = I_n$, then G is a generator matrix of a self-dual code of length 2(2n+1) if and only if

$$\begin{cases} AA^{T} + BB^{T} = I_{n}, \\ \sum_{i=1}^{4} \xi_{i}^{2} = 0, \\ \xi_{j}(\xi_{3} + \xi_{4} + 1) = 0 \quad j \in \{1, 2\}, \end{cases}$$

and the free rank of $(\boldsymbol{\xi}_1A + \boldsymbol{\xi}_2B, \boldsymbol{\xi}_1B + \boldsymbol{\xi}_2A, \xi_3, \xi_4)$ is 1.

Proof. First, let us determine the conditions required for G to be a generator matrix of a self-orthogonal code. We have

$$GG^{T} = \begin{pmatrix} \mathbf{v} & | \mathbf{0} | \xi_{3} & \xi_{4} \\ \hline I_{2n} & | X | \mathbf{v}^{T} & \mathbf{v}^{T} \end{pmatrix} \begin{pmatrix} \mathbf{v}^{T} & | I_{2n} \\ \hline \mathbf{0} & | X^{T} \\ \hline \xi_{3} & | \mathbf{v} \\ \xi_{4} & | \mathbf{v} \end{pmatrix} = \begin{pmatrix} g_{1,1} & g_{1,2} \\ g_{1,2}^{T} & g_{2,2} \end{pmatrix},$$

where

$$g_{1,1} = \mathbf{v}\mathbf{v}^{T} + \xi_{3}^{2} + \xi_{4}^{2},$$

$$g_{1,2} = (\xi_{3} + \xi_{4} + 1)\mathbf{v},$$

$$g_{2,2} = XX^{T} + I_{2n} + 2\mathbf{v}^{T}\mathbf{v}$$

so that $GG^T = \mathbf{0}$ if and only if $g_{1,1} = 0$, $g_{1,2} = \mathbf{0}$ and $g_{2,2} = \mathbf{0}$. Since R is of characteristic 2 and n is odd, we have

$$g_{1,1} = \mathbf{v}\mathbf{v}^{T} + \xi_{3}^{2} + \xi_{4}^{2}$$

= $n(\xi_{1}^{2} + \xi_{2}^{2}) + \xi_{3}^{2} + \xi_{4}^{2}$
= $\xi_{1}^{2} + \xi_{2}^{2} + \xi_{3}^{2} + \xi_{4}^{2}$,

so $g_{1,1} = 0$ if and only if $\sum_{i=1}^{4} \xi_i^2 = 0$. We also have $2\mathbf{v}^T \mathbf{v} = \mathbf{0}$, so $g_{2,2} = \mathbf{0}$ if and only if $XX^T = I_{2n}$. Since $CC^T = I_n$ by assumption, it follows from Lemma 3.1 that $XX^T = I_{2n}$ if and only if $AA^T + BB^T = I_n$. Finally, we see that $g_{1,2} = \mathbf{0}$ if and only if $\xi_j(\xi_3 + \xi_4 + 1) = 0$ for $j \in \{1, 2\}$.

Assume now that G is a generator matrix of a self-orthogonal code. We need to prove that the free rank of G is 2n+1 if and only if the free rank of $(\boldsymbol{\xi}_1A + \boldsymbol{\xi}_2B, \boldsymbol{\xi}_1B + \boldsymbol{\xi}_2A, \boldsymbol{\xi}_3, \boldsymbol{\xi}_4)$ is 1. The free rank of G is unchanged by elementary row (or column) operations and premultiplication (or postmultiplication) by an invertible matrix of appropriate size. Let $\tilde{G} = GM$ where

$$M = \frac{\begin{pmatrix} I_{2n} \mid X \mid \mathbf{v}^T \quad \mathbf{v}^T \\ \hline \mathbf{0} \mid I_{2n} \mid \mathbf{0} \\ \hline \mathbf{0} \mid \mathbf{0} \mid I_2 \end{pmatrix}.$$

Let rank() denote the free rank of a matrix over R. It is clear that M is invertible and hence rank $(\tilde{G}) = \operatorname{rank}(G)$. We have that

$$GM = \begin{pmatrix} \mathbf{v} & | \mathbf{0} & | \xi_3 & \xi_4 \\ I_{2n} & | X & | \mathbf{v}^T & \mathbf{v}^T \end{pmatrix} \begin{pmatrix} I_{2n} & | X & | \mathbf{v}^T & \mathbf{v}^T \\ \hline \mathbf{0} & | I_{2n} & | \mathbf{0} \\ \hline \mathbf{0} & | \mathbf{0} & | & I_2 \end{pmatrix}$$
$$= \begin{pmatrix} \mathbf{v} & \mathbf{v}X & \mathbf{v}(\mathbf{v}^T, \mathbf{v}^T) + (\xi_3, \xi_4) \\ I_{2n} & \mathbf{0} & \mathbf{0} \end{pmatrix}$$
$$= \begin{pmatrix} \mathbf{v} & \mathbf{v}X & (\mathbf{v}\mathbf{v}^T + \xi_3, \mathbf{v}\mathbf{v}^T + \xi_4) \\ I_{2n} & \mathbf{0} & \mathbf{0} \end{pmatrix}.$$

Let $r = \operatorname{rank}((\mathbf{v}X, \mathbf{v}\mathbf{v}^T + \xi_3, \mathbf{v}\mathbf{v}^T + \xi_4))$. Then $\operatorname{rank}(\tilde{G}) = 2n + 1$ if and only if r = 1. We see that

$$\mathbf{v}X = (\boldsymbol{\xi}_1, \boldsymbol{\xi}_2) \begin{pmatrix} AC & B \\ B^T C & A^T \end{pmatrix}$$
$$= ((\boldsymbol{\xi}_1 A + \boldsymbol{\xi}_2 B^T) C, \boldsymbol{\xi}_1 B + \boldsymbol{\xi}_2 A^T)$$

and

$$(\mathbf{v}\mathbf{v}^T + \xi_3, \mathbf{v}\mathbf{v}^T + \xi_4) = (\xi_3 + \xi_1^2 + \xi_2^2, \xi_4 + \xi_1^2 + \xi_2^2).$$

Since G is a generator matrix of a self-orthogonal code, we have $\sum_{i=1}^{4} \xi_i^2 = 0$ so that $\xi_1^2 + \xi_2^2 = \xi_3^2 + \xi_4^2$. By elementary column operations we obtain

$$r = \operatorname{rank}((\mathbf{v}X, \mathbf{v}\mathbf{v}^{T} + \xi_{3}, \mathbf{v}\mathbf{v}^{T} + \xi_{4}))$$

= $\operatorname{rank}((\mathbf{v}X, \xi_{3} + \xi_{1}^{2} + \xi_{2}^{2}, \xi_{4} + \xi_{1}^{2} + \xi_{2}^{2}))$
= $\operatorname{rank}((\mathbf{v}X, \xi_{3} + \xi_{3}^{2} + \xi_{4}^{2}, \xi_{4} + \xi_{3}^{2} + \xi_{4}^{2}))$
= $\operatorname{rank}((\mathbf{v}X, \xi_{3} + \xi_{3}^{2} + \xi_{4}^{2}, \xi_{3} + \xi_{4}))$
= $\operatorname{rank}((\mathbf{v}X, \xi_{3} + \xi_{3}^{2} + \xi_{4}^{2} + (\xi_{3} + \xi_{4})^{2}, \xi_{3} + \xi_{4}))$
= $\operatorname{rank}((\mathbf{v}X, \xi_{3}, \xi_{3} + \xi_{4}))$
= $\operatorname{rank}((\mathbf{v}X, \xi_{3}, \xi_{3} + \xi_{4}))$

We also have that $CC^T = I_n$ so that C is invertible. We also recall that $\boldsymbol{\xi}_2 A = \boldsymbol{\xi}_2 A^T J_n$ so that $\boldsymbol{\xi}_2 A J_n = \boldsymbol{\xi}_2 A^T$ and likewise $\boldsymbol{\xi}_2 B J_n = \boldsymbol{\xi}_2 B^T$. Thus, we get

$$r = \operatorname{rank}((\mathbf{v}X, \xi_3, \xi_4))$$

= $\operatorname{rank}(((\boldsymbol{\xi}_1 A + \boldsymbol{\xi}_2 B^T) C, \boldsymbol{\xi}_1 B + \boldsymbol{\xi}_2 A^T, \xi_3, \xi_4))$
= $\operatorname{rank}((\boldsymbol{\xi}_1 A + \boldsymbol{\xi}_2 B J_n, \boldsymbol{\xi}_1 B + \boldsymbol{\xi}_2 A J_n, \xi_3, \xi_4))$
= $\operatorname{rank}((\boldsymbol{\xi}_1 A + \boldsymbol{\xi}_2 B, \boldsymbol{\xi}_1 B + \boldsymbol{\xi}_2 A, \xi_3, \xi_4))$

and so rank(G) = 2n + 1 if and only if rank $((\boldsymbol{\xi}_1 A + \boldsymbol{\xi}_2 B, \boldsymbol{\xi}_1 B + \boldsymbol{\xi}_2 A, \boldsymbol{\xi}_3, \boldsymbol{\xi}_4)) = 1$. \Box

4 Results

In this section, we apply Theorem 3.2 to obtain many new best known, optimal and extremal binary self-dual codes. In particular, we obtain 8 singly-even [56, 28, 10] codes, one [78, 39, 14] code, 3 [92, 46, 16] codes and 43 [94, 47, 16] codes.

We also apply the following well-known technique for constructing self-dual codes referred to as the building-up construction.

Theorem 4.1. ([11]) Let R be a commutative Frobenius ring. Let G' be a generator matrix of a self-dual code C' of length 2n over R and let \mathbf{r}_i denote the i^{th} row of G'. Let $\varepsilon \in R : \varepsilon^2 = -1$, $\boldsymbol{\delta} \in R^{2n} : \langle \boldsymbol{\delta}, \boldsymbol{\delta} \rangle = -1$ and $\gamma_i = \langle \mathbf{r}_i, \boldsymbol{\delta} \rangle$ for $i \in \{1, \ldots, n\}$. Then the matrix

$$G = \begin{pmatrix} 1 & 0 & \boldsymbol{\delta} \\ -\gamma_1 & \varepsilon\gamma_1 & \mathbf{r}_1 \\ -\gamma_2 & \varepsilon\gamma_2 & \mathbf{r}_2 \\ \vdots & \vdots & \vdots \\ -\gamma_n & \varepsilon\gamma_n & \mathbf{r}_n \end{pmatrix}$$

is a generator matrix of a self-dual code of length 2(n+1) over R.

By applying Theorem 4.1, we obtain 3 singly-even [56, 28, 10] codes.

Remark 4.2. Two binary self-dual codes of length 2n are said to be neighbours if their intersection has dimension n-1. Let \mathcal{C}^* be a binary self-dual code of length 2n and let $\mathbf{x} \in \mathbb{F}_2^{2n} \setminus \mathcal{C}^*$. Then $\mathcal{C} = \langle \langle \mathbf{x} \rangle^{\perp} \cap \mathcal{C}^*, \mathbf{x} \rangle$ is a neighbour of \mathcal{C}^* , where $\langle \mathbf{x} \rangle$ denotes the code generated by \mathbf{x} .

$\mathbb{F}_2 + u\mathbb{F}_2$	$\mathbb{F}_2 + u\mathbb{F}_2 + v\mathbb{F}_2 + uv\mathbb{F}_2$	$\mathbb{F}_4 + u\mathbb{F}_4$	Symbol
0	0	0	0
1	1	1	1
u	u	w	2
1+u	1+u	1+w	3
_	v	u	4
_	1 + v	1+u	5
_	u + v	w+u	6
_	1 + u + v	1 + w + u	7
_	uv	wu	8
_	1 + uv	1 + wu	9
_	u + uv	w + wu	Α
_	1 + u + uv	1 + w + wu	В
_	v + uv	u + wu	С
_	1 + v + uv	1 + u + wu	D
_	u + v + uv	w + u + wu	Е
	1+u+v+uv	1+w+u+wu	F

Table 1: Hexadecimal notation system for elements of $\mathbb{F}_2 + u\mathbb{F}_2$, $\mathbb{F}_2 + u\mathbb{F}_2 + v\mathbb{F}_2 + uv\mathbb{F}_2$ and $\mathbb{F}_4 + u\mathbb{F}_4$.

Using Remark 4.2, we obtain one new [62, 31, 12] code as a neighbour of a [62, 31, 12] code constructed by applying Theorem 3.2.

We conduct the search for these codes using MATLAB and Magma [4] and determine their properties using Q-extension [5] and Magma. In MATLAB, we employ an algorithm which randomly searches for the construction parameters that satisfy the necessary and sufficient conditions stated in Theorem 3.2. For such parameters, we then build the corresponding binary generator matrices and print them to text files. We then use Q-extension to read these text files and determine the minimum distance and partial weight enumerator of each corresponding code. Furthermore, we determine the automorphism group order of each code using Magma. We implement a similar procedure for Theorem 4.1. Magma is used to search for neighbours as described in Remark 4.2. A database of generator matrices of the new codes is given online at [23]. The database is partitioned into text files (interpretable by Q-extension) corresponding to each code type. In these files, specific properties of the codes including the construction parameters, weight enumerator parameter values and automorphism group order are formatted as comments above the generator matrices. Partial weight enumerators of the codes are also formatted as comments below the generator matrices. Table 1 gives the hexadecimal notation system we use to represent elements of $\mathbb{F}_2 + u\mathbb{F}_2$, $\mathbb{F}_2 + u\mathbb{F}_2 + v\mathbb{F}_2 + uv\mathbb{F}_2$ and $\mathbb{F}_4 + u\mathbb{F}_4$.

4.1 New Self-Dual Codes of Length 56

The possible weight enumerators of a singly-even binary self-dual [56, 28, 10] code are given in [27] as

$$W_{56,1} = 1 + (308 + 4\alpha)x^{10} + (4246 - 8\alpha)x^{12} + \cdots,$$

$$W_{56,2} = 1 + (308 + 4\alpha)x^{10} + (3990 - 8\alpha)x^{12} + \cdots,$$

where $\alpha \in \mathbb{Z}$. Previously known α values for weight enumerators $W_{56,1}$ and $W_{56,2}$ can be found online at [34] (see [27, 19, 20, 33, 22]).

Table 2: Code of length 54 over \mathbb{F}_2 from Theorem 3.2 to which we apply Theorem 4.1 to obtain the codes in Table 3, where $\boldsymbol{\xi} = (\xi_1, \xi_2, \xi_3, \xi_4)$.

$\mathcal{C}_{54,i}'$	а	b	с	ξ
1	(0100111100101)	(1111101111010)	(1011001111110)	(1010)

Table 3: New singly-even binary self-dual [56, 28, 10] codes from applying Theorem 4.1 to $C'_{54,i}$ as given in Table 2 with $\delta = (0, \delta_0)$.

$\mathcal{C}_{56,i}$	$\mathcal{C}_{54,j}'$	δ_0	$W_{56,k}$	α	$ \operatorname{Aut}(\mathcal{C}_{56,i}) $
1	1	(000101100101100011111000101)	1	-55	1
2	1	(001000100111010111011101110)	1	-47	1
3	1	(110101100010101111001101100)	2	-50	1

We obtain 11 new best known singly-even binary self-dual codes of length 56 of which 6 have weight enumerator $W_{56,1}$ for

 $\alpha \in \{-z : z = 45, 47, 49, 50, 54, 55\}$

and 5 have weight enumerator $W_{56,2}$ for

 $\alpha \in \{-z : z = 50, ..., 53, 55\}.$

Of the 11 new codes, 3 are constructed by first applying Theorem 3.2 to obtain codes of length 54 over \mathbb{F}_2 (Table 2) to which we then apply Theorem 4.1 (Table 3); 6 are constructed by applying Theorem 3.2 over $\mathbb{F}_2 + u\mathbb{F}_2 + v\mathbb{F}_2 + uv\mathbb{F}_2$ (Table 4) and 2 are constructed by applying Theorem 3.2 over $\mathbb{F}_4 + u\mathbb{F}_4$ (Table 5).

4.2 New Self-Dual Code of Length 62

The possible weight enumerators of a binary self-dual [62, 31, 12] code are given in [6] as

$$W_{62,1} = 1 + 2308x^{12} + 23767x^{14} + \cdots,$$

$$W_{62,2} = 1 + (1860 + 32\alpha)x^{12} + (28055 - 160\alpha)x^{14} + \cdots,$$

where $\alpha \in \mathbb{Z}$. Previously known α values for weight enumerator $W_{62,2}$ can be found online at [34] (see [25, 6, 35, 36, 38]).

Table 4: New singly-even binary self-dual [56, 28, 10] codes from Theorem 3.2 over $\mathbb{F}_2 + u\mathbb{F}_2 + v\mathbb{F}_2 + uv\mathbb{F}_2$, where $\boldsymbol{\xi} = (\xi_1, \xi_2, \xi_3, \xi_4)$.

$\mathcal{C}_{56,i}$	λ	μ	а	b	с	ξ	$W_{56,j}$	α	$ \operatorname{Aut}(\mathcal{C}_{56,i}) $
4	1	9	(B03)	(39D)	(344)	(7EBA)	1	-54	$2^2 \cdot 3$
5	F	1	(331)	(8F9)	(EE3)	(DEBA)	1	-50	2^{2}
6	1	3	(D52)	(F95)	(700)	(9EAB)	2	-55	$2^2 \cdot 3$
7	1	5	(DB9)	(D45)	(88D)	(D654)	2	-53	$2^{3} \cdot 3$
8	1	1	(31F)	(54D)	(00F)	(D6DC)	2	-52	$2^3 \cdot 3$
9	1	F	(FA7)	(FB5)	(C47)	(76AB)	2	-51	2^{2}

Table 5: New singly-even binary self-dual [56, 28, 10] codes from Theorem 3.2 over $\mathbb{F}_4 + u\mathbb{F}_4$, where $\boldsymbol{\xi} = (\xi_1, \xi_2, \xi_3, \xi_4)$.

$\mathcal{C}_{56,i}$	λ	μ	а	b	с	ξ	$W_{56,j}$	α	$ \operatorname{Aut}(\mathcal{C}_{56,i}) $
10	1	1	(48D)	(5F2)	(CC9)	(6F67)	1	-49	$2 \cdot 3$
11	D	9	(B5D)	(D61)	(900)	(6F67)	1	-45	2

Table 6: Code of length 62 over \mathbb{F}_2 from Theorem 3.2 to which we apply Remark 4.2 to obtain the code in Table 7, where $\boldsymbol{\xi} = (\xi_1, \xi_2, \xi_3, \xi_4)$.

$\mathcal{C}^*_{62,i}$	а	b	с	ξ
1	(000000100100101)	(000011101110111)	(10000000000000)	(0110)

We obtain one new extremal binary self-dual codes of length 62 which has weight enumerator $W_{62,2}$ for

 $\alpha = 2.$

The new code is constructed by first applying Theorem 3.2 to obtain a code of length 62 over \mathbb{F}_2 (Table 6) and then searching for neighbours of this code using Remark 4.2 (Table 7).

4.3 New Self-Dual Code of Length 78

The possible weight enumerators of a binary self-dual [78, 39, 14] code are given in [9, 24] as

$$W_{78,1} = 1 + (3705 + 8\alpha)x^{14} + (62244 - 24\alpha + 512\beta)x^{16} + (774592 - 64\alpha - 4608\beta)x^{18} + \cdots,$$

$$W_{78,2} = 1 + (3705 + 8\alpha)x^{14} + (71460 - 24\alpha)x^{16} + (658880 - 64\alpha)x^{18} + \cdots,$$

where $\alpha, \beta \in \mathbb{Z}$. Previously known (α, β) values for weight enumerator $W_{78,1}$ can be found online at [34] (see [9, 2, 24, 16, 40, 37, 20]).

We obtain one new optimal binary self-dual codes of length 78 which has weight enumerator $W_{78,1}$ for

 $\beta = 0$ and $\alpha = -76$.

The new code is constructed by applying Theorem 3.2 over \mathbb{F}_2 (Table 8).

Table 7: New binary self-dual [62, 31, 12] code from searching for neighbours of $C^*_{62,j}$ as given in Table 6 using Remark 4.2 with $\mathbf{x} = (\mathbf{0}, \mathbf{x}_0)$.

$\mathcal{C}_{62,i}$	$\mathcal{C}^*_{62,j}$	\mathbf{x}_0	$W_{62,k}$	α	$ \operatorname{Aut}(\mathcal{C}_{62,i}) $
1	1	(1111001101001110100110000100110)	2	2	2^{2}

Table 8: New binary self-dual [78, 39, 14] code from Theorem 3.2 over \mathbb{F}_2 , where $\boldsymbol{\xi} = (\xi_1, \xi_2, \xi_3, \xi_4)$.

$\mathcal{C}_{78,i}$	a	b	с	ξ
1	(0100101010100001000)	(1111101101011010000)	(001010111111101101)	(0101)

Table 8:	(continued)
----------	-------------

$\mathcal{C}_{78,i}$	$W_{78,j}$	α	β	$ \operatorname{Aut}(\mathcal{C}_{78,i}) $
1	1	-76	0	19

4.4 New Self-Dual Codes of Length 92

The possible weight enumerators of a binary self-dual $\left[92,46,16\right]$ code are given in $\left[9\right]$ as

$$W_{92,1} = 1 + (4692 + 4\alpha)x^{16} + (174800 - 8\alpha + 256\beta)x^{18} + (2425488 - 52\alpha - 2048\beta)x^{20} + \cdots,$$

$$W_{92,2} = 1 + (4692 + 4\alpha)x^{16} + (174800 - 8\alpha + 256\beta)x^{18} + (2441872 - 52\alpha - 2048\beta)x^{20} + \cdots,$$

$$W_{92,3} = 1 + (4692 + 4\alpha)x^{16} + (121296 - 8\alpha)x^{18} + (3213968 - 52\alpha)x^{20} + \cdots,$$

where $\alpha, \beta \in \mathbb{Z}$. Previously known (α, β) values for weight enumerator $W_{92,1}$ can be found online at [34] (see [7, 17, 26, 39, 21, 22]).

We obtain 3 new extremal binary self-dual codes of length 92 which have weight enumerator $W_{92,1}$ for

 $\beta = 0$ and $\alpha \in \{807, 862, 1038\}.$

The new codes are constructed by applying Theorem 3.2 over $\mathbb{F}_2 + u\mathbb{F}_2$ (Table 9).

Table 9: New binary self-dual [92, 46, 16] codes from Theorem 3.2 over $\mathbb{F}_2 + u\mathbb{F}_2$, where $\boldsymbol{\xi} = (\xi_1, \xi_2, \xi_3, \xi_4)$.

$\mathcal{C}_{92,i}$	λ	μ	а	b	с	ξ	$W_{92,j}$	α	β	$ \operatorname{Aut}(\mathcal{C}_{92,i}) $
1	1	1	(02223003031)	(02321323010)	(22232222222)	(2301)	1	807	0	$2 \cdot 11$
2	1	1	(22101022322)	(33123310002)	(02101130201)	(1201)	1	862	0	$2 \cdot 11$
3	1	1	(21200233200)	(31233320122)	(22232222222)	(2310)	1	1038	0	$2^{2} \cdot 11$

4.5 New Self-Dual Codes of Length 94

The possible weight enumerators of a binary self-dual $\left[94,47,16\right]$ code are given in $\left[28\right]$ as

$$W_{94,1} = 1 + 2\alpha x^{16} + (134044 - 2\alpha + 128\beta)x^{18} + (2010660 - 30\alpha - 896\beta)x^{20} + \cdots,$$

$$W_{94,2} = 1 + 2\alpha x^{16} + (134044 - 2\alpha + 128\beta)x^{18} + (2018852 - 30\alpha - 896\beta)x^{20} + \cdots,$$

$$W_{94,3} = 1 + 2\alpha x^{16} + (134044 - 2\alpha + 128\beta)x^{18} + (2190884 - 30\alpha - 896\beta)x^{20} + \cdots,$$

where $\alpha, \beta \in \mathbb{Z}$. Previously known (α, β) values for weight enumerator $W_{94,1}$ can be found online at [34] (see [28]).

We obtain 43 new best known binary self-dual codes of length 94 which have weight enumerator $W_{\rm 94,1}$ for

$$\begin{split} \beta &= -69 \text{ and } \alpha \in \{46z : z = 78\}; \\ \beta &= -46 \text{ and } \alpha \in \{46z : z = 69, 70, 76, 78, 79\}; \\ \beta &= -23 \text{ and } \alpha \in \{46z : z = 62, 63, 65, ..., 79\}; \\ \beta &= 0 \text{ and } \alpha \in \{46z : z = 57, ..., 65, 67, ..., 75, 77, 79\}. \end{split}$$

The new codes are constructed by applying Theorem 3.2 over \mathbb{F}_2 (Table 10).

5 Conclusion

In this work, we presented a new bordered construction for self-dual codes. We gave the necessary conditions for our method to produce self-dual codes over a finite commutative Frobenius ring of characteristic 2. Using our new bordered construction together with the well-known building-up and neighbour constructions, we were able to construct the following new singly-even binary self-dual codes:

Codes of length 56: We were able to construct new singly-even binary selfdual [56, 28, 10] codes which have weight enumerator $W_{56,1}$ for:

$$\alpha \in \{-z : z = 45, 47, 49, 50, 54, 55\}$$

and weight enumerator $W_{56,2}$ for:

$$\alpha \in \{-z : z = 50, ..., 53, 55\}.$$

Code of length 62: We were able to construct a new binary self-dual [62, 31, 12] code which has weight enumerator $W_{62,2}$ for:

 $\alpha = 2.$

Code of length 78: We were able to construct a new binary self-dual [78, 39, 14] code which has weight enumerator $W_{78,1}$ for:

$$\beta = 0$$
 and $\alpha = -76$.

$\mathcal{C}_{94,i}$	а	b	с	ξ
1	(10001010001010111110101)	(01000001011100010110100)	(00111001110111000101110)	(1001)
2	(10011110100001000100100)	(00011011100101010111010)	(00111010111010011111000)	(0110)
3	(00101010100010100100010)	(11100001010001000001101)	(11111111000100100011010)	(1010)
4	(00000100011100110110010)	(00011111001000001101111)	(00010101100111100100000)	(0110)
5	(10011000101101011101110)	(10110011100000101011110)	(11100010011010111010101)	(0110)
6	(01000001110110010100101)	(11100010110101000111000)	(10010110110010111001011)	(1001)
7	(11100100101101110010101)	(11001101011001100101010)	(00100111000000010110011)	(0101)
8	(01001110101010101000110)	(01110011110110110001110)	(11010001000011101111110)	(1001)
9	(01110111111000111010001)	(11000000111010011100011)	(10001100101100101001000)	(1001)
10	(00011101100001100010000)	(10011101010000001110000)	(10011011010001111100011)	(0101)
11	(111111111101010111110101)	(11010011100101011111101)	(10001011101100011111001)	(0110)
12	(01000100110100111011001)	(10110111110000110101101)	(00001011111100101101110)	(1010)
13	(10010111000100001000011)	(01111100011110001100100)	(10101011110110000101101)	(0101)
14	(11100001111011111011011)	(00101000010110010100011)	(10101110101100100011110)	(1010)
15	(00111000011101010101010)	(01000111100000001111001)	(01011011110011100111000)	(1010)
16	(10110010010001101000110)	(01100011101110101000100)	(10111110010111010100100)	(1010)
17	(01111110110101010010100)	(00100010010010101110000)	(10000011010000001011011)	(0110)
18	(10011001010010000100000)	(10110110000111011111001)	(11001011101011100000111)	(0110)
19	(10111110111001000111001)	(00011100011101100010101)	(10001110111001110001110)	(0101)
20	(11100000011100111000111)	(01011101000011000010100)	(10001001101011101010111)	(0101)
21	(11000100000010010011001)	(10011111110011000001011)	(00010110110011100011111)	(1001)
22	(0100000000001111111111)	(1110110111011111011110)	(00111110100100101011101)	(1010)
23	(11101110101111101010001)	(00111000100001100101110)	(11011010101111011000010)	(1010)
24	(10111110010111110101001)	(00001100100001100110111)	(00010001001010011010011)	(0110)
25	(011010111111001001011100)	(10011111010101001011000)	(10100111101000111111000)	(0110)
26	(00110001110000010000100)	(01000110110110010010001)	(01001110000110001011000)	(1010)
27	(11111100110100101000011)	(01100110011000101111010)	(10000011111011010010111)	(1001)
28	(10111000111101100010111)	(10111011110011100110110)	(10101010001001000001101)	(1001)
29	(01110010100111101110101)	(01101110001001100110001)	(00110100000010110111000)	(0101)
30	(01101001100010111101111)	(11101000101110111111001)	(01101001110110011101001)	(1010)
31	(01100100110000110010010)	(01110111110111011110010)	(01011011010000110111101)	(0110)
32	(00111110101010110101110)	(11010100110110100100100)	(01110101000101100111101)	(1010)
33	(11010001010100000110111)	(00010111100111001101111)	(10010011000011011101111)	(0110)
34	(00011101000001110100111)	(00110000101010111100001)	(00110000111001000011010)	(1010)
35	(11010110011101000101011)	(01001110111001101001100)	(11100011000011101110110)	(0101)
36	(10010001001001011011010)	(10100110101000001110110)	(00000011101111110001111)	(1001)
37	(00010011001011000010101)	(01100100010011101111010)	(00111111011100000011110)	(0110)
38	(01100001111100111001010)	(10101111000011101100101)	(11111010010101100010110)	(0110)
39	(11001101110110000000111)	(11101011010011000101011)	(11100011011101000110011)	(0110)
40	(10111110110110010101000)	(01110001000111110000111)	(00101101110000010111111)	(1010)
41	(0000001001010101111101)	(01011000110110010010011)	(011110111011010001001)	(1010)
42	(01110110010010100011011)	(01000011111001101101011)	(01000000101101110000011)	(0101)
43	(11101100100110110001001)	(0110101101100000000011)	(01000000000000000000000)	(1010)

Table 10: New binary self-dual [94, 47, 16] codes from Theorem 3.2 over \mathbb{F}_2 , where $\boldsymbol{\xi} = (\xi_1, \xi_2, \xi_3, \xi_4)$.

$\mathcal{C}_{94,i}$	$W_{94,j}$	α	β	$ \operatorname{Aut}(\mathcal{C}_{94,i}) $
1	1	3588	-69	23
2	1	3174	-46	23
3	1	3220	-46	23
4	1	3496	-46	23
5	1	3588	-46	23
6	1	3634	-46	23
7	1	2852	-23	23
8	1	2898	-23	23
9	1	2990	-23	23
10	1	3036	-23	23
11	1	3082	-23	23
12	1	3128	-23	23
13	1	3174	-23	23
14	1	3220	-23	23
15	1	3266	-23	23
16	1	3312	-23	23
17	1	3358	-23	23
18	1	3404	-23	23
19	1	3450	-23	23
20	1	3496	-23	23
21	1	3542	-23	23
22	1	3588	-23	23
23	1	3634	-23	23
24	1	2622	0	23
25	1	2668	0	23
26	1	2714	0	23
27	1	2760	0	23
28	1	2806	0	23
29	1	2852	0	23
30	1	2898	0	23
31	1	2944	0	23
32	1	2990	0	23
33	1	3082	0	23
34	1	3128	0	23
35	1	3174	0	23
36	1	3220	0	23
37	1	3266	0	23
38	1	3312	0	23
39	1	3358	0	23
40	1	3404	0	23
41	1	3450	0	23
42	1	3542	0	23
43	1	3634	0	$2 \cdot 23$

Table 10: (continued)

Codes of length 92: We were able to construct new binary self-dual [92, 46, 16] codes which have weight enumerator $W_{92,1}$ for:

$$\beta = 0$$
 and $\alpha \in \{807, 862, 1038\}.$

Codes of length 94: We were able to construct new binary self-dual [94, 47, 16] codes which have weight enumerator $W_{94,1}$ for:

$$\begin{aligned} \beta &= -69 \text{ and } \alpha \in \{46z : z = 78\}, \\ \beta &= -46 \text{ and } \alpha \in \{46z : z = 69, 70, 76, 78, 79\}, \\ \beta &= -23 \text{ and } \alpha \in \{46z : z = 62, 63, 65, \dots, 79\}, \\ \beta &= 0 \text{ and } \alpha \in \{46z : z = 57, \dots, 65, 67, \dots, 75, 77, 79\}. \end{aligned}$$

A suggestion for future work would be to consider our bordered construction with the λ -circulant matrices A, B and C being replaced with some other matrices, for example, matrices that come from group rings. This would give one many more possible choices for the matrices A, B and C which could lead to finding more new binary self-dual codes of different lengths.

References

- E. F. Assmus, Jr. and H. F. Mattson, Jr. New 5-designs. J. Combin. Theory, 6 (2):122–151, 1969. doi: 10.1016/S0021-9800(69)80115-8.
- [2] A. Baartmans and V. Yorgov. Some New Extremal Codes of Lengths 76 and 78. *IEEE Trans. Inform. Theory*, 49(5):1353–1354, 2003. doi: 10.1109/TIT.2003. 810653.
- [3] E. Bannai, S. T. Dougherty, M. Harada, and M. Oura. Type II Codes, Even Unimodular Lattices, and Invariant Rings. *IEEE Trans. Inform. Theory*, 45(4): 1194–1205, 1999. doi: 10.1109/18.761269.
- [4] W. Bosma, J. Cannon, and C. Playoust. The Magma Algebra System I: The User Language. J. Symbolic Comput., 24(3–4):235–265, 1997. doi: 10.1006/jsco.1996. 0125.
- [5] I. G. Bouyukliev. What is Q-extension? Serdica J. Comput., 1(2):115–130, 2007.
- [6] R. Dontcheva and M. Harada. New Extremal Self-Dual Codes of Length 62 and Related Extremal Self-Dual Codes. *IEEE Trans. Inform. Theory*, 48(7):2060– 2064, 2002. doi: 10.1109/TIT.2002.1013144.
- [7] R. Dontcheva, R. Russeva, and N. Ziapkov. On the binary extremal codes of length 92. "Proceedings of the International Workshop Optimal Codes and Related Topics", Sunny Beach, Bulgaria, pages 53–58, 2001.
- [8] S. T. Dougherty. Algebraic Coding Theory Over Finite Commutative Rings. Springer International Publishing, Cham, Switzerland, 1st edition, 2017.
- [9] S. T. Dougherty, T. A. Gulliver, and M. Harada. Extremal Binary Self-Dual Codes. *IEEE Trans. Inform. Theory*, 43(6):2036–2047, 1997. doi: 10.1109/18. 641574.

- [10] S. T. Dougherty, P. Gaborit, M. Harada, and P. Solé. Type II Codes over $\mathbb{F}_2 + u\mathbb{F}_2$. *IEEE Trans. Inform. Theory*, 45(1):32–45, 1999. doi: 10.1109/18.746770.
- [11] S. T. Dougherty, J.-L. Kim, H. Kulosman, and H. Liu. Self-dual codes over commutative Frobenius rings. *Finite Fields Appl.*, 16(1):14–26, 2010. doi: 10. 1016/j.ffa.2009.11.004.
- [12] S. T. Dougherty, B. Yildiz, and S. Karadeniz. Codes over R_k , Gray maps and their binary images. *Finite Fields Appl.*, 17(3):205–219, 2011. doi: 10.1016/j.ffa. 2010.11.002.
- [13] S. T. Dougherty, B. Yildiz, and S. Karadeniz. Self-dual codes over R_k and binary self-dual codes. *Eur. J. Pure Appl. Math.*, 6(1):89–106, 2013.
- [14] S. T. Dougherty, J. Gildea, A. Korban, A. Kaya, A. Tylyshchak, and B. Yildiz. Bordered constructions of self-dual codes from group rings and new extremal binary self-dual codes. *Finite Fields Appl.*, 57:108–127, 2019. doi: 10.1016/j.ffa. 2019.02.004.
- [15] S. T. Dougherty, J. Gildea, and A. Kaya. Quadruple bordered constructions of self-dual codes from group rings. *Cryptogr. Commun.*, 12(1):127–146, 2020. doi: 10.1007/s12095-019-00380-8.
- [16] P. Gaborit and A. Otmani. Experimental constructions of self-dual codes. *Finite Fields Appl.*, 9(3):372–394, 2003. doi: 10.1016/S1071-5797(03)00011-X.
- [17] P. Gaborit, V. Pless, P. Solé, and O. Atkin. Type II Codes over F₄. *Finite Fields Appl.*, 8(2):171–183, 2002. doi: 10.1006/ffta.2001.0333.
- [18] J. Gildea, A. Kaya, R. Taylor, and B. Yildiz. Constructions for self-dual codes induced from group rings. *Finite Fields Appl.*, 51:71–92, 2018. doi: 10.1016/j.ffa. 2018.01.002.
- [19] J. Gildea, A. Kaya, A. Tylyshchak, and B. Yildiz. A Group Induced Four-circulant Construction for Self-dual Codes and New Extremal Binary Self-dual Codes, 2019. https://arxiv.org/abs/1912.11758.
- [20] J. Gildea, A. Kaya, A. Tylyshchak, and B. Yildiz. A modified bordered construction for self-dual codes from group rings. J. Algebra Comb. Discrete Struct. Appl., 7(2):103–119, 2020. doi: 10.13069/jacodesmath.729402.
- [21] J. Gildea, A. Korban, A. Kaya, and B. Yildiz. Constructing self-dual codes from group rings and reverse circulant matrices. *Adv. Math. Commun.*, 2020. doi: 10.3934/amc.2020077.
- [22] J. Gildea, A. Korban, and A. M. Roberts. New binary self-dual codes of lengths 56, 58, 64, 80 and 92 from a modification of the four circulant construction. *Finite Fields Appl.*, 75, 2021. doi: 10.1016/j.ffa.2021.101876.
- [23] J. Gildea, A. Korban, A. M. Roberts, and A. Tylyshchak. Generator matrix database, 2021. https://amr4-f6sxz49ktd.netlify.app.
- [24] T. A. Gulliver, M. Harada, and J.-L. Kim. Construction of new extremal selfdual codes. *Discrete Math.*, 263(1–3):81–91, 2003. doi: 10.1016/S0012-365X(02) 00570-8.

- [25] M. Harada. Construction of an Extremal Self-Dual Code of Length 62. IEEE Trans. Inform. Theory, 45(4):1232–1233, 1999. doi: 10.1109/18.761274.
- [26] M. Harada and T. Nishimura. An extremal singly even self-dual code of length 88. Adv. Math. Commun., 1(2):261–267, 2007. doi: 10.3934/amc.2007.1.261.
- [27] M. Harada and K. Saito. Singly even self-dual codes constructed from Hadamard matrices of order 28. Australas. J. Combin., 70(2):288–296, 2018.
- [28] M. Harada, M. Kiermaier, A. Wassermann, and R. Yorgova. New Binary Singly Even Self-Dual Codes. *IEEE Trans. Inform. Theory*, 56(4):1612–1617, 2010. doi: 10.1109/TIT.2010.2040967.
- [29] C. W. H. Lam. The Search for a Finite Projective Plane of Order 10. Amer. Math. Monthly, 98(4):305–318, 1991. doi: 10.1080/00029890.1991.12000759.
- [30] S. Ling and P. Solé. Type II Codes over $\mathbb{F}_4 + u\mathbb{F}_4$. European J. Combin., 22(7): 983–997, 2001. doi: 10.1006/eujc.2001.0509.
- [31] C. L. Mallows and N. J. A. Sloane. An Upper Bound for Self-Dual Codes. Inf. Control, 22(2):188–200, 1973. doi: 10.1016/S0019-9958(73)90273-8.
- [32] E. M. Rains. Shadow Bounds for Self-Dual Codes. *IEEE Trans. Inform. Theory*, 44(1):134–139, 1998. doi: 10.1109/18.651000.
- [33] A. M. Roberts. Constructions of extremal and optimal self-dual and Hermitian self-dual codes over finite fields using circulant matrices. Master's thesis, University of Chester, Chester, UK, 2020. https://drive.google.com/file/d/ 1CMjnuBvQtrXOY8foy6_gfXOcFFuHAaFs/view.
- [34] A. M. Roberts. Weight enumerator parameter database for binary self-dual codes, 2021. https://amr-wepd-bsdc.netlify.app.
- [35] R. Russeva and N. Yankov. On binary self-dual codes of lengths 60, 62, 64 and 66 having an automorphism of order 9. Des. Codes Cryptogr., 45:335–346, 2007. doi: 10.1007/s10623-007-9127-9.
- [36] N. Yankov. Self-dual [62, 31, 12] and [64, 32, 12] codes with an automorphism of order 7. Adv. Math. Commun., 8(1):73–81, 2014. doi: 10.3934/amc.2014.8.73.
- [37] N. Yankov and D. Anev. New self-dual [78, 39, 14] codes with an automorphism of order 13. "Proceedings of the Eighth International Workshop on Optimal Codes and Related Topics", Sofia, Bulgaria, pages 128–133, 2017.
- [38] N. Yankov and D. Anev. On the self-dual codes with an automorphism of order 5. Appl. Algebra Engrg. Comm. Comput., 2019. doi: 10.1007/s00200-019-00403-0.
- [39] R. Yorgova and A. Wassermann. Binary self-dual codes with automorphisms of order 23. Des. Codes Cryptogr., 48(2):155–164, 2008. doi: 10.1007/ s10623-007-9152-8.
- [40] T. Zhang, J. Michel, T. Feng, and G. Ge. On the Existence of Certain Optimal Self-Dual Codes with Lengths Between 74 and 116. *Electron. J. Combin.*, 22(4), 2015. doi: 10.37236/5213.