IMPROVING CASAZZA-KALTON-CHRISTENSEN-VAN EIJNDHOVEN PERTURBATION WITH APPLICATIONS

K. MAHESH KRISHNA Department of Humanities and Basic Sciences Aditya College of Engineering and Technology Surampalem, East-Godavari Andhra Pradesh 533 437 India Email: kmaheshak@gmail.com September 7, 2021

Abstract: Let \mathcal{X}, \mathcal{Y} be Banach spaces and $S : \mathcal{X} \to \mathcal{Y}$ be an invertible Lipschitz map. Let $T : \mathcal{X} \to \mathcal{Y}$ be a map and there exist $\lambda_1, \lambda_2 \in [0, 1)$ such that

$$||Tx - Ty - (Sx - Sy)|| \le \lambda_1 ||Sx - Sy|| + \lambda_2 ||Tx - Ty||, \quad \forall x, y \in \mathcal{X}.$$

Then we prove that T is an invertible Lipschitz map. This improves 25 years old Casazza-Kalton-Christensen-van Eijndhoven perturbation. It also improves 28 years old Soderlind-Campanato perturbation and 2 years old Barbagallo-Ernst-Thera perturbation. We give applications to the theory of metric frames. The notion of Lipschitz atomic decomposition for Banach spaces is also introduced.

Keywords: Paley-Wiener perturbation, Lipschitz map, metric frame, atomic decomposition. **Mathematics Subject Classification (2020)**: 26A16, 47A55, 42C15.

1. INTRODUCTION

Let \mathcal{X} be a Banach space and $I_{\mathcal{X}}$ be the identity operator on \mathcal{X} . Carl Neumann's classical result says that if $T: \mathcal{X} \to \mathcal{X}$ is a bounded linear operator such that $||T - I_{\mathcal{X}}|| < 1$, then T is invertible [39]. Following two results are consequences of this result. They are known as **Paley-Wiener theorems**.

- (1) Sequences close to orthonormal bases in Hilbert spaces are Riesz bases [40, 52].
- (2) Sequences close to Schauder bases in Banach spaces are Schauder bases [4,44].

History of Paley-Wiener theorems are nicely presented in [1,43]. It was in the setting of Hilbert spaces, Paley-Wiener theorem was first generalized by Pollard [42], second generalized by Sz. Nagy [21] and third generalized by Hilding [33]. Hilding proved the following theorem.

Theorem 1.1. [33] (Hilding perturbation) Let \mathcal{H} be a Hilbert space. If a linear operator $T : \mathcal{H} \to \mathcal{H}$ is such that there exists $\lambda \in [0, 1)$ with

$$||Th - h|| \le \lambda ||Th|| + \lambda ||h||, \quad \forall h \in \mathcal{H},$$

then T is bounded, invertible and

$$\frac{1-\lambda}{1+\lambda} \|h\| \le \|Th\| \le \frac{1+\lambda}{1-\lambda} \|h\|, \quad \forall h \in \mathcal{H};$$
$$\frac{1-\lambda}{1+\lambda} \|h\| \le \|T^{-1}h\| \le \frac{1+\lambda}{1-\lambda} \|h\|, \quad \forall h \in \mathcal{H}.$$

It took around 50 years to improve Theorem 1.1 to the most generality for Banach spaces.

Theorem 1.2. [9,11,49] (Casazza-Kalton-Christensen-van Eijndhoven perturbation) Let \mathcal{X}, \mathcal{Y} be Banach spaces and $S : \mathcal{X} \to \mathcal{Y}$ be a bounded invertible operator. If a linear operator $T : \mathcal{X} \to \mathcal{Y}$ is

such that there exist $\lambda_1, \lambda_2 \in [0, 1)$ with

$$||Tx - Sx|| \le \lambda_1 ||Sx|| + \lambda_2 ||Tx||, \quad \forall x \in \mathcal{X},$$

then T is bounded, invertible and

$$\begin{aligned} \frac{1-\lambda_1}{1+\lambda_2} \|Sx\| &\leq \|Tx\| \leq \frac{1+\lambda_1}{1-\lambda_2} \|Sx\|, \quad \forall x \in \mathcal{X}; \\ \frac{1-\lambda_2}{1+\lambda_1} \frac{1}{\|S\|} \|y\| \leq \|T^{-1}y\| \leq \frac{1+\lambda_2}{1-\lambda_1} \|S^{-1}\| \|y\|, \quad \forall y \in \mathcal{Y} \end{aligned}$$

There is an improvement of Theorem 1.2 which is due to Guo with an extra assumption that T is bounded.

Theorem 1.3. [26] Let \mathcal{X}, \mathcal{Y} be Banach spaces and $S : \mathcal{X} \to \mathcal{Y}$ be a bounded invertible operator. If a bounded linear operator $T : \mathcal{X} \to \mathcal{Y}$ is such that there exist $\lambda_1 \in [0, 1)$ and $\lambda_2 \in [0, 1]$ with

$$|Tx - Sx|| \le \lambda_1 ||Sx|| + \lambda_2 ||Tx||, \quad \forall x \in \mathcal{X},$$

then T is invertible. Further, for every $\varepsilon > 0$ satisfying $1 > \lambda_2 - \varepsilon > 0$ and $\lambda_1 + \varepsilon ||TS^{-1}|| < 1$, we have

$$\begin{aligned} \frac{1-\lambda_1-\varepsilon\|TS^{-1}\|}{1+\lambda_2-\varepsilon}\|Sx\| &\leq \|Tx\| \leq \frac{1+\lambda_1+\varepsilon\|TS^{-1}\|}{1-\lambda_2+\varepsilon}\|Sx\|, \quad \forall x \in \mathcal{X}; \\ \frac{1-\lambda_2+\varepsilon}{1+\lambda_1+\varepsilon\|TS^{-1}\|}\frac{1}{\|S\|}\|y\| &\leq \|T^{-1}y\| \leq \frac{1+\lambda_2-\varepsilon}{1-\lambda_1-\varepsilon\|TS^{-1}\|}\|S^{-1}\|\|y\|, \quad \forall y \in \mathcal{Y}. \end{aligned}$$

Theorem 1.2 and its variants are useful in various studies such as stability of frames for Hilbert spaces [11], stability of frames and atomic decompositions for Banach spaces [47], stability of frames for Hilbert C*-modules [27], stability of G-frames [48], multipliers for Hilbert spaces [46], quantum detection problem [5], continuous frames [24], fusion frames [10], operator representations of frames (dynamics of frames) [15], pseudo-inverses of operators [22], outer inverses of operators [51], shift-invariant spaces [35], frame sequences [17], sampling [53] etc.

The main objective of this paper is to generalize Theorem 1.2 for Lipschitz functions between Banach spaces. We do this in Theorem 2.9. We show that our result generalizes Soderlind-Campanato Perturbation (Theorem 3.1) and Barbagallo-Ernst-Thera perturbation (Theorem 3.2). We then give an application to the theory of frames for metric spaces. Further, the notion of Lipschitz atomic decomposition for Banach spaces is introduced and a perturbation result is derived using Theorem 2.9.

2. Improving Casazza-Kalton-Christensen-van Eijndhoven perturbation

Let \mathcal{M} be a metric space and \mathcal{X} be a Banach space. Recall that a function $f : \mathcal{M} \to \mathcal{X}$ is said to be Lipschitz if there exists b > 0 such that

$$\|f(x) - f(y)\| \le b \, d(x, y), \quad \forall x, y \in \mathcal{M}.$$

A Lipschitz function $f: \mathcal{M} \to \mathcal{X}$ is said to be bi-Lipschitz if there exists a > 0 such that

$$a d(x, y) \le ||f(x) - f(y)||, \quad \forall x, y \in \mathcal{M}.$$

Definition 2.1. [50] Let \mathcal{X} be a Banach space.

(i) Let \mathcal{M} be a metric space. The collection $\operatorname{Lip}(\mathcal{M}, \mathcal{X})$ is defined as $\operatorname{Lip}(\mathcal{M}, \mathcal{X}) \coloneqq \{f : \mathcal{M} \to \mathcal{X} \text{ is Lipschitz}\}$. For $f \in \operatorname{Lip}(\mathcal{M}, \mathcal{X})$, the Lipschitz number is defined as

$$\operatorname{Lip}(f) \coloneqq \sup_{x,y \in \mathcal{M}, x \neq y} \frac{\|f(x) - f(y)\|}{d(x,y)}$$

IMPROVING CASAZZA-KALTON-CHRISTENSEN-VAN EIJNDHOVEN PERTURBATION ...

(ii) Let $(\mathcal{M}, 0)$ be a pointed metric space. The collection $\operatorname{Lip}_0(\mathcal{M}, \mathcal{X})$ is defined as $\operatorname{Lip}_0(\mathcal{M}, \mathcal{X}) \coloneqq \{f : \mathcal{M} \to \mathcal{X} \text{ is Lipschitz and } f(0) = 0\}$. For $f \in \operatorname{Lip}_0(\mathcal{M}, \mathcal{X})$, the Lipschitz norm is defined as

$$\|f\|_{\operatorname{Lip}_0} \coloneqq \sup_{x,y \in \mathcal{M}, x \neq y} \frac{\|f(x) - f(y)\|}{d(x,y)}.$$

Theorem 2.2. [50] Let \mathcal{X} be a Banach space.

- (i) If \mathcal{M} is a metric space, then $\operatorname{Lip}(\mathcal{M}, \mathcal{X})$ is a semi-normed vector space w.r.t. the semi-norm $\operatorname{Lip}(\cdot)$.
- (ii) If $(\mathcal{M}, 0)$ is a pointed metric space, then $\operatorname{Lip}_0(\mathcal{M}, \mathcal{X})$ is a Banach space w.r.t. the norm $\|\cdot\|_{\operatorname{Lip}_0}$. Further, $\operatorname{Lip}_0(\mathcal{X}) \coloneqq \operatorname{Lip}_0(\mathcal{X}, \mathcal{X})$ is a unital Banach algebra. In particular, if $T \in \operatorname{Lip}_0(\mathcal{X})$ satisfies $\|T - I_{\mathcal{X}}\|_{\operatorname{Lip}_0} < 1$, then T is invertible and $T^{-1} \in \operatorname{Lip}_0(\mathcal{X})$.

We now develop perturbation result for Lipschitz functions using various results. Our developments are motivated from the linear version of improvement of Paley-Wiener theorem by van Eijndhoven [49]. Since $\operatorname{Lip}_0(\mathcal{X})$ is a unital Banach algebra, we can talk about the notion of spectrum and resolvents. In the remaining part of the paper, the spectrum of an element $T \in \operatorname{Lip}_0(\mathcal{X})$ is denoted by $\sigma(T)$ and the resolvent by $\rho(T)$.

Lemma 2.3. Let \mathcal{X} be a Banach space and $T \in Lip_0(\mathcal{X})$. Suppose there are $\alpha_0 \in \mathbb{R}$ and $\beta_0 > 0$ such that

(1)
$$||Tx - Ty - \alpha(x - y)|| \ge \beta_0 ||x - y||, \quad \forall x, y \in \mathcal{X}, \forall \alpha \le \alpha_0.$$

Then $(-\infty, \alpha_0] \subseteq \rho(T)$.

Proof. If $\sigma(T) \cap \mathbb{R} = \emptyset$, then there is nothing to argue. So let $\sigma(T) \cap \mathbb{R} \neq \emptyset$. Since $\sigma(T)$ is non empty, define

$$\lambda_0 \coloneqq \min\{\lambda \in \mathbb{R} : \lambda \in \sigma(T) \cap \mathbb{R}\}.$$

Claim: $\lambda_0 \geq \alpha_0$. If this is false, we have $\lambda_0 < \alpha_0$. From the assumption we then have

$$||Tx - Ty - \lambda_0(x - y)|| \ge \beta_0 ||x - y||, \quad \forall x, y \in \mathcal{X}$$

which implies

$$\|(T - \lambda_0 I_{\mathcal{X}})x - (T - \lambda_0 I_{\mathcal{X}})y\| \ge \beta_0 \|x - y\|, \quad \forall x, y \in \mathcal{X}$$

which says that $T - \lambda_0 I_{\mathcal{X}}$ is injective. We now try to show that $T - \lambda_0 I_{\mathcal{X}}$ is surjective. Let $y \in \mathcal{X}$. Define

$$\alpha_n \coloneqq \lambda_0 - \frac{1}{n}, \quad \forall n \in \mathbb{N}$$

From the definition of λ_0 we then have $T - \alpha_n I_{\mathcal{X}}$ is invertible for all $n \in \mathbb{N}$. Using Inequality (1) we have

$$\|(T - \alpha_n I_{\mathcal{X}})x - (T - \alpha_n I_{\mathcal{X}})y\| \ge \beta_0 \|x - y\|, \quad \forall x, y \in \mathcal{X}, \forall n \in \mathbb{N}$$

which gives

$$||u - v|| \ge \beta_0 ||(T - \alpha_n I_{\mathcal{X}})^{-1} u - (T - \alpha_n I_{\mathcal{X}})^{-1} v||, \quad \forall u, v \in \mathcal{X}, \forall n \in \mathbb{N}$$

which implies

$$\|(T - \alpha_n I_{\mathcal{X}})^{-1}\|_{\operatorname{Lip}_0} \le \frac{1}{\beta_0}, \quad \forall n \in \mathbb{N}.$$

Now define

$$x_n \coloneqq (T - \alpha_n I_{\mathcal{X}})^{-1} y, \quad \forall n \in \mathbb{N}$$

Now using resolvent identity in Banach algebra [20] we get

$$||x_n - x_m|| = \left|\frac{1}{n} - \frac{1}{m}\right| ||(T - \alpha_n I_{\mathcal{X}})^{-1} (T - \alpha_m I_{\mathcal{X}})^{-1} y||$$

$$\leq \left|\frac{1}{n} - \frac{1}{m}\right| ||(T - \alpha_n I_{\mathcal{X}})^{-1}||_{\operatorname{Lip}_0} ||(T - \alpha_m I_{\mathcal{X}})^{-1}||_{\operatorname{Lip}_0} ||y||$$

$$\leq \left|\frac{1}{n} - \frac{1}{m}\right| \frac{1}{\beta_0^2} ||y||, \quad \forall n, m \in \mathbb{N}.$$

Thus $\{x_n\}_n$ is Cauchy and it converges to an element, say $x \in \mathcal{X}$. Finally

$$\begin{aligned} \|(T - \lambda_0 I_{\mathcal{X}})x - y\| &= \|(T - \lambda_0 I_{\mathcal{X}})x - (T - \alpha_n I_{\mathcal{X}})x_n\| \\ &\leq \|(T - \lambda_0 I_{\mathcal{X}})x - (T - \lambda_0 I_{\mathcal{X}})x_n\| + \|(T - \lambda_0 I_{\mathcal{X}})x_n - (T - \alpha_n I_{\mathcal{X}})x_n\| \\ &\leq \|(T - \lambda_0 I_{\mathcal{X}})^{-1}\|_{\text{Lip}_0} \|x_n - x\| + |\alpha_n - \lambda_0| \|x_n\| \\ &\leq \|(T - \lambda_0 I_{\mathcal{X}})^{-1}\|_{\text{Lip}_0} \|x_n - x\| + \frac{1}{n} \|x_n\|, \quad \forall n \in \mathbb{N}. \end{aligned}$$

Letting $n \to \infty$, we get $(T - \lambda_0 I_{\mathcal{X}})x = y$. Thus $T - \lambda_0 I_{\mathcal{X}}$ is invertible. Hence $\lambda_0 \in \sigma(T) \cap \rho(T) = \emptyset$ which is a contradiction which proves the claim as well as the lemma.

Remark 2.4. Note that Lemma 2.3 also holds for real Banach spaces. In fact, if spectrum $\sigma(T)$ is empty then the lemma holds trivially and if it is non empty, then it becomes a compact set and proof of Lemma 2.3 can be carried over. Due to this, all coming results in this paper are valid for real Banach spaces also.

Theorem 2.5. Let \mathcal{X} be a Banach space and $T \in Lip_0(\mathcal{X})$. Suppose there are $\alpha_1 \in \mathbb{R}$ and $\beta_1 > 0$ such that

(2)
$$||Tx - Ty - \alpha(x - y)|| \ge \beta_1 ||x - y||, \quad \forall x, y \in \mathcal{X}, \forall \alpha \le \alpha_1$$

Then $(-\infty, \alpha_1 + \beta_1) \subseteq \rho(T)$.

Proof. Let $0 < \eta < \beta_1$. Define $\alpha_0 \coloneqq \alpha_1 + \eta$ and $\beta_0 \coloneqq \beta_1 - \eta$. Let $\alpha \le \alpha_0$. Case (i): $\alpha \le \alpha_1$. Then from Inequality (2), $||Tx - Ty - \alpha(x - y)|| \ge \beta_1 ||x - y||, \forall x, y \in \mathcal{X}$. Case (ii): Let $\alpha_1 < \alpha \le \alpha_0$. Then

$$||Tx - Ty - \alpha(x - y)|| \ge ||Tx - Ty - \alpha_1(x - y)|| - (\alpha - \alpha_1)||x - y||$$
$$\ge (\beta_1 - \eta)||x - y|| = \beta_0 ||x - y||, \quad \forall x, y \in \mathcal{X}.$$

Lemma 2.3 now gives $(-\infty, \alpha_0] \subseteq \rho(T)$. Result follows by observing that

$$(-\infty, \alpha_1 + \beta_1) = \bigcup_{0 < \eta < \beta_1} (-\infty, \alpha_1 + \eta].$$

Theorem 2.6. Let \mathcal{X} be a Banach space, $T : \mathcal{X} \to \mathcal{X}$ be a map, T0 = 0 and there exist $\lambda_1, \lambda_2 \in [0, 1)$ such that

(3)
$$||Tx - Ty - (x - y)|| \le \lambda_1 ||x - y|| + \lambda_2 ||Tx - Ty||, \quad \forall x, y \in \mathcal{X}.$$

Then

(i) T is Lipschitz and

(4)
$$\frac{1-\lambda_1}{1+\lambda_2} \|x-y\| \le \|Tx-Ty\| \le \frac{1+\lambda_1}{1-\lambda_2} \|x-y\|, \quad \forall x, y \in \mathcal{X}.$$

(ii) We have

$$\left(-\infty, \frac{1-\lambda_1}{1+\lambda_2}\right) \subseteq \rho(T).$$

(iii) T is invertible and

$$\frac{1-\lambda_2}{1+\lambda_1} \|x-y\| \le \|T^{-1}x - T^{-1}y\| \le \frac{1+\lambda_2}{1-\lambda_1} \|x-y\|, \quad \forall x, y \in \mathcal{X}.$$

(iv) We have

$$\frac{1-\lambda_1}{1+\lambda_2} \le \|T\|_{Lip_0} \le \frac{1+\lambda_1}{1-\lambda_2} \quad and \quad \frac{1-\lambda_2}{1+\lambda_1} \le \|T^{-1}\|_{Lip_0} \le \frac{1+\lambda_2}{1-\lambda_1}.$$

Proof. Let $x, y \in \mathcal{X}$. Then using Inequality (3)

$$\begin{aligned} \|Tx - Ty\| &\leq \|Tx - Ty - (x - y)\| + \|x - y\| \leq \lambda_1 \|x - y\| + \lambda_2 \|Tx - Ty\| + \|x - y\| \\ &= (1 + \lambda_1) \|x - y\| + \lambda_2 \|Tx - Ty\| \\ &\implies \|Tx - Ty\| \leq \frac{1 + \lambda_1}{1 - \lambda_2} \|x - y\| \end{aligned}$$

and

$$\begin{aligned} \|x - y\| &\leq \|Tx - Ty - (x - y)\| + \|Tx - Ty\| \leq \lambda_1 \|x - y\| + \lambda_2 \|Tx - Ty\| + \|Tx - Ty\| \\ &= \lambda_1 \|x - y\| + (1 + \lambda_2) \|Tx - Ty\| \\ &\implies \frac{1 - \lambda_1}{1 + \lambda_2} \|x - y\| \leq \|Tx - Ty\|. \end{aligned}$$

Hence T is Lipschitz and (i) holds. Let $\alpha \leq 0.$ Then

$$\begin{split} \|Tx - Ty - \alpha(x - y)\| &= \|(1 - \alpha)(x - y) - (x - y - (Tx - Ty))\| \\ &\geq (1 - \alpha)\|x - y\| - \|Tx - Ty - (x - y)\| \\ &\geq (1 - \alpha)\|x - y\| - \lambda_1\|x - y\| - \lambda_2\|Tx - Ty\| \\ &= (1 - \alpha - \lambda_1)\|x - y\| - \lambda_2\|Tx - Ty\| \\ &\geq (1 - \alpha - \lambda_1)\|x - y\| - \lambda_2\|Tx - Ty - \alpha(x - y)\| + \lambda_2\alpha\|x - y\| \\ &= (1 - \alpha - \lambda_1 + \lambda_2\alpha)\|x - y\| - \lambda_2\|Tx - Ty - \alpha(x - y)\| \end{split}$$

which implies

$$\|Tx - Ty - \alpha(x - y)\| \ge \frac{1 - \alpha - \lambda_1 + \lambda_2 \alpha}{1 + \lambda_2} \|x - y\|$$
$$= \frac{1 - \lambda_1 - (1 - \lambda_2) \alpha}{1 + \lambda_2} \|x - y\| \ge \frac{1 - \lambda_1}{1 + \lambda_2} \|x - y\|.$$

By applying Theorem 2.5 for $\alpha_1 = 0$ and $\beta_1 = \frac{1-\lambda_1}{1+\lambda_2}$ we get $(-\infty, \frac{1-\lambda_1}{1+\lambda_2}) \subseteq \rho(T)$. Since $0 \in (-\infty, \frac{1-\lambda_1}{1+\lambda_2})$, T is invertible. Using Inequality (4) we then get

$$\frac{1-\lambda_1}{1+\lambda_2} \|T^{-1}u - T^{-1}v\| \le \|u - v\| \le \frac{1+\lambda_1}{1-\lambda_2} \|T^{-1}u - T^{-1}v\|, \quad \forall u, v \in \mathcal{X}$$

which gives (iii). Finally (iv) follows from (i) and (iii).

Theorem 2.7. Let \mathcal{X} , \mathcal{Y} be Banach spaces and $S \in Lip_0(\mathcal{X}, \mathcal{Y})$ be invertible. Let $T : \mathcal{X} \to \mathcal{Y}$ be a map, T0 = 0 and there exist $\lambda_1, \lambda_2 \in [0, 1)$ such that

(5)
$$||Tx - Ty - (Sx - Sy)|| \le \lambda_1 ||Sx - Sy|| + \lambda_2 ||Tx - Ty||, \quad \forall x, y \in \mathcal{X}$$

Then

(i) T is Lipschitz and

$$\frac{1-\lambda_1}{1+\lambda_2}\|Sx - Sy\| \le \|Tx - Ty\| \le \frac{1+\lambda_1}{1-\lambda_2}\|Sx - Sy\|, \quad \forall x, y \in \mathcal{X}.$$

- (ii) $\alpha S T$ is invertible for all $\alpha \in \left(-\infty, \frac{1-\lambda_1}{1+\lambda_2}\right)$.
- (iii) T is invertible and

$$\frac{1-\lambda_2}{1+\lambda_1} \frac{1}{\|S\|_{Lip_0}} \|u-v\| \le \|T^{-1}u - T^{-1}v\| \le \frac{1+\lambda_2}{1-\lambda_1} \|S^{-1}\|_{Lip_0} \|u-v\|, \quad \forall u, v \in \mathcal{Y}.$$

(iv) We have

$$\frac{1-\lambda_1}{1+\lambda_2} \|S\|_{Lip_0} \le \|T\|_{Lip_0} \le \frac{1+\lambda_1}{1-\lambda_2} \|S\|_{Lip_0} \quad and \quad \frac{1-\lambda_2}{1+\lambda_1} \frac{1}{\|S\|_{Lip_0}} \le \|T^{-1}\|_{Lip_0} \le \frac{1+\lambda_2}{1-\lambda_1} \|S^{-1}\|_{Lip_0}.$$

Proof. Define $R \coloneqq TS^{-1}$. Then Inequality (5) gives

$$||TS^{-1}u - TS^{-1}v - (SS^{-1}u - SS^{-1}v)|| \le \lambda_1 ||SS^{-1}u - SS^{-1}v|| + \lambda_2 ||TS^{-1}u - TS^{-1}v||, \quad \forall u, v \in \mathcal{Y}.$$

i.e.,

$$||Ru - Rv - (u - v)|| \le \lambda_1 ||u - v|| + \lambda_2 ||Ru - Rv||, \quad \forall u, v \in \mathcal{Y}.$$

By applying Theorem 2.6 to R we get the following.

(i) R is Lipschitz hence T is Lipschitz. Further,

$$\frac{1-\lambda_1}{1+\lambda_2}\|Sx-Sy\| \le \|R(Sx)-R(Sy)\| \le \frac{1+\lambda_1}{1-\lambda_2}\|Sx-Sy\|, \quad \forall x, y \in \mathcal{X}.$$

But $||R(Sx) - R(Sy)|| = ||Tx - Ty||, \forall x, y \in \mathcal{X}.$

- (ii) $\alpha I_{\mathcal{X}} R$ is invertible for all $\alpha \in \left(-\infty, \frac{1-\lambda_1}{1+\lambda_2}\right)$. Since S is invertible we then have $\alpha S T$ is invertible for all $\alpha \in \left(-\infty, \frac{1-\lambda_1}{1+\lambda_2}\right)$.
- (iii) R is invertible hence T is invertible. Further,

$$\begin{split} \frac{1-\lambda_2}{1+\lambda_1} \frac{1}{\|S\|_{\operatorname{Lip}_0}} \|u-v\| &\leq \frac{1}{\|S\|_{\operatorname{Lip}_0}} \|R^{-1}u - R^{-1}v\| \leq \|S^{-1}(R^{-1}u) - S^{-1}(R^{-1}v)\| \\ &= \|T^{-1}u - T^{-1}v\| \leq \|S^{-1}\|_{\operatorname{Lip}_0} \|R^{-1}u - R^{-1}v\| \\ &\leq \frac{1+\lambda_2}{1-\lambda_1} \|S^{-1}\|_{\operatorname{Lip}_0} \|u-v\|, \quad \forall u, v \in \mathcal{Y}. \end{split}$$

(iv) This follows easily from (i) and (iii).

Our next job is to derive the results by removing the condition T0 = 0.

Theorem 2.8. Let \mathcal{X} be a Banach space, $T : \mathcal{X} \to \mathcal{X}$ be a map and there exist $\lambda_1, \lambda_2 \in [0, 1)$ such that

$$||Tx - Ty - (x - y)|| \le \lambda_1 ||x - y|| + \lambda_2 ||Tx - Ty||, \quad \forall x, y \in \mathcal{X}.$$

Then

(i) T is Lipschitz and

$$\frac{1-\lambda_1}{1+\lambda_2}\|x-y\| \le \|Tx-Ty\| \le \frac{1+\lambda_1}{1-\lambda_2}\|x-y\|, \quad \forall x, y \in \mathcal{X}.$$

(ii) $\alpha I_{\mathcal{X}} - T$ is invertible for all $\alpha \in \left(-\infty, \frac{1-\lambda_1}{1+\lambda_2}\right)$. (iii) T is invertible and

$$\frac{1-\lambda_2}{1+\lambda_1} \|x-y\| \le \|T^{-1}x - T^{-1}y\| \le \frac{1+\lambda_2}{1-\lambda_1} \|x-y\|, \quad \forall x, y \in \mathcal{X}.$$

(iv) We have

$$\frac{1-\lambda_1}{1+\lambda_2} \le Lip(T) \le \frac{1+\lambda_1}{1-\lambda_2} \quad and \quad \frac{1-\lambda_2}{1+\lambda_1} \le Lip(T^{-1}) \le \frac{1+\lambda_2}{1-\lambda_1}$$

Proof. Define

$$\tilde{T}x \coloneqq Tx - T0, \quad \forall x \in \mathcal{X}.$$

Then $\tilde{T}0 = 0$ and

$$\|\tilde{T}x - \tilde{T}y - (x - y)\| = \|Tx - Ty - (x - y)\| \le \lambda_1 \|x - y\| + \lambda_2 \|Tx - Ty\|$$
$$= \lambda_1 \|x - y\| + \lambda_2 \|\tilde{T}x - \tilde{T}y\|, \quad \forall x, y \in \mathcal{X}.$$

Applying Theorem 2.6 and using the fact that 'a map is bijective if and only if its translate is bijective', proof is complete. \Box

Theorem 2.9. Let \mathcal{X}, \mathcal{Y} be Banach spaces and $S \in Lip(\mathcal{X}, \mathcal{Y})$ be invertible. Let $T : \mathcal{X} \to \mathcal{Y}$ be a map and there exist $\lambda_1, \lambda_2 \in [0, 1)$ such that

$$||Tx - Ty - (Sx - Sy)|| \le \lambda_1 ||Sx - Sy|| + \lambda_2 ||Tx - Ty||, \quad \forall x, y \in \mathcal{X}.$$

Then

(i) T is Lipschitz and

$$\frac{1-\lambda_1}{1+\lambda_2}\|Sx - Sy\| \le \|Tx - Ty\| \le \frac{1+\lambda_1}{1-\lambda_2}\|Sx - Sy\|, \quad \forall x, y \in \mathcal{X}.$$

(ii) $\alpha S - T$ is invertible for all $\alpha \in \left(-\infty, \frac{1-\lambda_1}{1+\lambda_2}\right)$.

(iii) T is invertible and

$$\frac{1-\lambda_2}{1+\lambda_1}\frac{1}{Lip(S)}\|u-v\| \le \|T^{-1}u - T^{-1}v\| \le \frac{1+\lambda_2}{1-\lambda_1}Lip(S^{-1})\|u-v\|, \quad \forall u, v \in \mathcal{Y}.$$

(iv) We have

$$\frac{1-\lambda_1}{1+\lambda_2}Lip(S) \le Lip(T) \le \frac{1+\lambda_1}{1-\lambda_2}Lip(S) \quad and \quad \frac{1-\lambda_2}{1+\lambda_1}\frac{1}{Lip(S)} \le Lip(T^{-1}) \le \frac{1+\lambda_2}{1-\lambda_1}Lip(S^{-1}).$$

Proof. Define $R \coloneqq TS^{-1}$ and the proof is similar to proof of Theorem 2.7.

Following two corollaries are motivated from [33].

Corollary 2.10. Let $p \ge 1$. Let \mathcal{X} , \mathcal{Y} be Banach spaces and $S \in Lip(\mathcal{X}, \mathcal{Y})$ be invertible. Let $T : \mathcal{X} \to \mathcal{Y}$ be a map and there exist $\lambda_1, \lambda_2 \in [0, 1)$ such that

(6)
$$||Tx - Ty - (Sx - Sy)|| \le ((\lambda_1 ||Sx - Sy||)^p + (\lambda_2 ||Tx - Ty||)^p)^{\frac{1}{p}}, \quad \forall x, y \in \mathcal{X}.$$

Then

(i) T is Lipschitz and

$$\frac{1-\lambda_1}{1+\lambda_2}\|Sx - Sy\| \le \|Tx - Ty\| \le \frac{1+\lambda_1}{1-\lambda_2}\|Sx - Sy\|, \quad \forall x, y \in \mathcal{X}.$$

(ii) $\alpha S - T$ is invertible for all $\alpha \in \left(-\infty, \frac{1-\lambda_1}{1+\lambda_2}\right)$. (iii) T is invertible and

$$\frac{1-\lambda_2}{1+\lambda_1}\frac{1}{Lip(S)}\|u-v\| \le \|T^{-1}u - T^{-1}v\| \le \frac{1+\lambda_2}{1-\lambda_1}Lip(S^{-1})\|u-v\|, \quad \forall u, v \in \mathcal{Y}.$$

(iv) We have

$$\frac{1-\lambda_1}{1+\lambda_2}Lip(S) \le Lip(T) \le \frac{1+\lambda_1}{1-\lambda_2}Lip(S) \quad and \quad \frac{1-\lambda_2}{1+\lambda_1}\frac{1}{Lip(S)} \le Lip(T^{-1}) \le \frac{1+\lambda_2}{1-\lambda_1}Lip(S^{-1}).$$

Proof. Note that if $r, s \ge 0$, then

$$(r^{p} + s^{p})^{\frac{1}{p}} \le r + s \text{ if } p \ge 1.$$

Hence Inequality (6) gives

$$\begin{aligned} \|Tx - Ty - (Sx - Sy)\| &\leq \left((\lambda_1 \|Sx - Sy\|)^p + (\lambda_2 \|Tx - Ty\|)^p \right)^{\frac{1}{p}} \\ &\leq \lambda_1 \|Sx - Sy\| + \lambda_2 \|Tx - Ty\|, \quad \forall x, y \in \mathcal{X}. \end{aligned}$$

Result follows by applying Theorem 2.9.

Corollary 2.11. Let $0 . Let <math>\mathcal{X}$, \mathcal{Y} be Banach spaces and $S \in Lip(\mathcal{X}, \mathcal{Y})$ be invertible. Let $T : \mathcal{X} \to \mathcal{Y}$ be a map and there exist $\lambda_1, \lambda_2 \in \left[0, 2^{1-\frac{1}{p}}\right)$ such that

$$||Tx - Ty - (Sx - Sy)|| \le ((\lambda_1 ||Sx - Sy||)^p + (\lambda_2 ||Tx - Ty||)^p)^{\frac{1}{p}}, \quad \forall x, y \in \mathcal{X}.$$

Then

(i) T is Lipschitz and

$$\frac{1-2^{\frac{1}{p}-1}\lambda_1}{1+2^{\frac{1}{p}-1}\lambda_2}\|Sx-Sy\| \le \|Tx-Ty\| \le \frac{1+2^{\frac{1}{p}-1}\lambda_1}{1-2^{\frac{1}{p}-1}\lambda_2}\|Sx-Sy\|, \quad \forall x, y \in \mathcal{X}.$$
(ii) $\alpha S - T$ is invertible for all $\alpha \in \left(-\infty, \frac{1-2^{\frac{1}{p}-1}\lambda_1}{1+2^{\frac{1}{p}-1}\lambda_2}\right).$

(iii) T is invertible and

$$\frac{1-2^{\frac{1}{p}-1}\lambda_2}{1+2^{\frac{1}{p}-1}\lambda_1}\frac{1}{Lip(S)}\|u-v\| \le \|T^{-1}u-T^{-1}v\| \le \frac{1+2^{\frac{1}{p}-1}\lambda_2}{1-2^{\frac{1}{p}-1}\lambda_1}Lip(S^{-1})\|u-v\|, \quad \forall u, v \in \mathcal{Y}.$$

(iv) We have

$$\frac{1-2^{\frac{1}{p}-1}\lambda_1}{1+2^{\frac{1}{p}-1}\lambda_2}Lip(S) \le Lip(T) \le \frac{1+2^{\frac{1}{p}-1}\lambda_1}{1-2^{\frac{1}{p}-1}\lambda_2}Lip(S) \quad and$$
$$\frac{1-2^{\frac{1}{p}-1}\lambda_2}{1+2^{\frac{1}{p}-1}\lambda_1}\frac{1}{Lip(S)} \le Lip(T^{-1}) \le \frac{1+2^{\frac{1}{p}-1}\lambda_2}{1-2^{\frac{1}{p}-1}\lambda_1}Lip(S^{-1}).$$

Proof. Note that if $r, s \ge 0$, then

$$(r^p + s^p)^{\frac{1}{p}} \le 2^{\frac{1}{p}-1}(r+s)$$
 if $p < 1$.

Hence Inequality (6) gives

$$||Tx - Ty - (Sx - Sy)|| \le ((\lambda_1 ||Sx - Sy||)^p + (\lambda_2 ||Tx - Ty||)^p)^{\frac{1}{p}}$$

$$\le 2^{\frac{1}{p} - 1} \lambda_1 ||Sx - Sy|| + 2^{\frac{1}{p} - 1} \lambda_2 ||Tx - Ty||, \quad \forall x, y \in \mathcal{X}.$$

Result follows by applying Theorem 2.9.

Next we generalize Corollary 1 in [11].

Corollary 2.12. Let \mathcal{X} , \mathcal{Y} be Banach spaces and $S \in Lip(\mathcal{X}, \mathcal{Y})$ be invertible. Let $T : \mathcal{X} \to \mathcal{Y}$ be a map and there exists $\lambda \in [0, 1)$ such that

$$\|Tx - Ty - (Sx - Sy)\| \le \lambda \|Sx - Sy\| + \|Tx - Ty\|, \quad \forall x, y \in \mathcal{X}$$

Then T is Lipschitz, invertible and

$$Lip(T^{-1}) \le \frac{2}{1-\lambda} Lip(S^{-1}).$$

Proof. Define $R \coloneqq TS^{-1}$. Then

$$||Ru - Rv - (u - v)|| \le \lambda ||u - v|| + ||Ru - Rv||, \quad \forall u, v \in \mathcal{Y}.$$

Note that $\operatorname{Lip}(R) \neq 0$. Let

$$0 < \varepsilon < \min\left\{1, \frac{1-\lambda}{\operatorname{Lip}(R)}\right\}.$$

Define $\lambda_1 \coloneqq \lambda - \varepsilon \operatorname{Lip}(R)$ and $\lambda_2 \coloneqq 1 - \varepsilon$. Then $\lambda_1, \lambda_2 \in [0, 1)$ and

$$\begin{aligned} \|Ru - Rv - (u - v)\| &\leq \lambda \|u - v\| + \|Ru - Rv\| \\ &\leq \lambda \|u - v\| + \|Ru - Rv\| + \varepsilon(\operatorname{Lip}(R)\|u - v\| - \|Ru - Rv\|) \\ &= (\lambda + \varepsilon \operatorname{Lip}(R))\|u - v\| + (1 - \varepsilon)\|Ru - Rv\|, \quad \forall u, v \in \mathcal{Y}. \end{aligned}$$

By applying Theorem 2.8 we get that R is Lipschitz, invertible and

$$\operatorname{Lip}(R^{-1}) \le \frac{2-\varepsilon}{1-(\lambda+\varepsilon\operatorname{Lip}(R))}$$

Since ε can be made arbitrarily small, we must have

$$\operatorname{Lip}(R^{-1}) \le \frac{2}{1-\lambda}.$$

Substituting the expression of R gives

$$\frac{1}{\text{Lip}(S^{-1})}\text{Lip}(T^{-1}) \le \text{Lip}(ST^{-1}) = \text{Lip}(R^{-1}) \le \frac{2}{1-\lambda}.$$

We finally derive the following non-linear version of Theorem 1.3.

Theorem 2.13. Let \mathcal{X} , \mathcal{Y} be Banach spaces and $S \in Lip(\mathcal{X}, \mathcal{Y})$ be invertible. Let $T : \mathcal{X} \to \mathcal{Y}$ be a Lipschitz map and there exist $\lambda_1 \in [0, 1)$ and $\lambda_2 \in [0, 1]$ such that

$$\|Tx - Ty - (Sx - Sy)\| \le \lambda_1 \|Sx - Sy\| + \lambda_2 \|Tx - Ty\|, \quad \forall x, y \in \mathcal{X}.$$

Then T is Lipschitz invertible. Further, for every $\varepsilon > 0$ satisfying $1 > \lambda_2 - \varepsilon > 0$ and $\lambda_1 + \varepsilon Lip(TS^{-1}) < 1$, we have

(i)

$$\frac{1-\lambda_1-\varepsilon Lip(TS^{-1})}{1+\lambda_2-\varepsilon}\|Sx-Sy\| \le \|Tx-Ty\| \le \frac{1+\lambda_1+\varepsilon Lip(TS^{-1})}{1-\lambda_2+\varepsilon}\|Sx-Sy\|, \quad \forall x,y \in \mathcal{X}.$$

- (ii) $\alpha S T$ is invertible for all $\alpha \in \left(-\infty, \frac{1-\lambda_1 \varepsilon Lip(TS^{-1})}{1+\lambda_2 \varepsilon}\right)$.
- (iii) T is invertible and

 $\frac{1-\lambda_2+\varepsilon}{1+\lambda_1+\varepsilon Lip(TS^{-1})}\frac{1}{Lip(S)}\|u-v\| \le \|T^{-1}u-T^{-1}v\| \le \frac{1+\lambda_2-\varepsilon}{1-\lambda_1-\varepsilon Lip(TS^{-1})}Lip(S^{-1})\|u-v\|, \quad \forall u,v \in \mathcal{Y}.$

$$\frac{1-\lambda_1-\varepsilon Lip(TS^{-1})}{1+\lambda_2-\varepsilon}Lip(S) \le Lip(T) \le \frac{1+\lambda_1+\varepsilon Lip(TS^{-1})}{1-\lambda_2+\varepsilon}Lip(S) \quad and \\ \frac{1-\lambda_2+\varepsilon}{1+\lambda_1+\varepsilon Lip(TS^{-1})}\frac{1}{Lip(S)} \le Lip(T^{-1}) \le \frac{1+\lambda_2-\varepsilon}{1-\lambda_1-\varepsilon Lip(TS^{-1})}Lip(S^{-1}).$$

Proof. Define $R := TS^{-1}$. Then for every $\varepsilon > 0$ satisfying $1 > \lambda_2 - \varepsilon > 0$ and $\lambda_1 + \varepsilon \operatorname{Lip}(TS^{-1}) < 1$,

$$\begin{aligned} \|Ru - Rv - (u - v)\| &\leq \lambda_1 \|u - v\| + \lambda_2 \|Ru - Rv\| \\ &= \lambda_1 \|u - v\| + (\lambda_2 - \varepsilon) \|Ru - Rv\| + \varepsilon \|Ru - Rv\| \\ &\leq \lambda_1 \|u - v\| + (\lambda_2 - \varepsilon) \|Ru - Rv\| + \varepsilon \operatorname{Lip}(R) \|u - v\| \\ &= (\lambda_1 + \varepsilon \operatorname{Lip}(R)) \|u - v\| + (\lambda_2 - \varepsilon) \|Ru - Rv\|, \quad \forall u, v \in \mathcal{Y}. \end{aligned}$$

Remaining parts of the proof is similar to the proof of Theorem 2.7.

It is an easy observation that the constant λ_1 in Theorem 2.13 can not be improved. We are therefore left with the following open problem.

Question 2.14. Can the constant λ_2 be improved further in Theorem 2.13?

3. Applications

Our first two applications of Theorem 2.9 are easy proofs of Soderlind-Campanato perturbation and Barbagallo-Ernst-Thera perturbation.

Theorem 3.1. [6,45] (Soderlind-Campanato perturbation) Let \mathcal{X} be a real Banach space, $A : \mathcal{X} \to \mathcal{X}$ be a map and there exist $\alpha > 0, 0 \le \beta < 1$ such that

$$\|\alpha Ax - \alpha Ay - (x - y)\| \le \beta \|x - y\|, \quad \forall x, y \in \mathcal{X}.$$

Then A is Lipschitz, invertible and $Lip(A^{-1}) \leq \frac{\alpha}{1-\beta}$.

Proof. Set $T = \alpha A$ and $\lambda_1 = \beta$ in Theorem 2.9. Then $\frac{1}{\alpha} \operatorname{Lip}(A^{-1}) = \operatorname{Lip}(T^{-1}) \leq \frac{1}{1-\lambda_1} = \frac{1}{1-\beta}$.

Theorem 3.2. [3] (Barbagallo-Ernst-Thera perturbation) Let \mathcal{X} be a real Banach space, $A : \mathcal{X} \to \mathcal{X}$ be a map and there exist $\alpha > 0, 0 \leq \beta < 1$ such that

(7)
$$\|Ax - Ay - (\alpha x - \alpha y)\| \le \beta \|Ax - Ay\|, \quad \forall x, y \in \mathcal{X}.$$

Then

(i) If $\beta < 1/2$, then A is Lipschitz, invertible and $Lip(A^{-1}) \leq \frac{1-\beta}{\alpha(1-2\beta)}$.

(ii) If \mathcal{X} is a Hilbert space, then A is Lipschitz, invertible and $Lip(A^{-1}) \leq \frac{1+\beta}{\alpha}$.

Proof. Set $T = \frac{1}{\alpha}A$ and $\lambda_2 = \beta$ in Theorem 2.9. Then

- (i) $\alpha \operatorname{Lip}(A^{-1}) = 1 + \lambda_2 = 1 + \beta \leq \frac{1-\beta}{1-2\beta}$.
- (ii) $\alpha \text{Lip}(A^{-1}) = 1 + \lambda_2 = 1 + \beta.$

Remark 3.3. In [3], it is stated that "the number 1/2 in Theorem 3.2 can not be improved" (see line -7, page 20, line -14, page 20, and line 14, page 18 in [3]). In an attempt of giving an example, an operator A is constructed which satisfies $||A - I_X|| = 1/2||A||$. Unfortunately this example does not satisfy Inequality (7). Reason is the following. The operator A is constructed as follows. Let \mathcal{X} be a Banach space and assume that $\mathcal{X} = \mathcal{Y} \oplus \mathcal{Z}$ for some closed subspaces \mathcal{Y} and \mathcal{Z} of \mathcal{X} . Then the projections $P: \mathcal{Y} \oplus \mathcal{Z} \ni y \oplus z \to y \in \mathcal{X}, Q: \mathcal{Y} \oplus \mathcal{Z} \ni y \oplus z \to z \in \mathcal{X}$ are bounded linear operators. Define $A \coloneqq 2P$ which is linear. Then

$$A - I_{\mathcal{X}} = 2P - (P + Q) = P - Q.$$

Now for $x = y \oplus z \in \mathcal{Y} \oplus \mathcal{Z}$, we have

$$||Ax - x|| = ||Px - Qx|| = ||y \oplus (-z)||$$
 and $||Ax|| = ||2Px|| = 2||y||$

Thus there is no $\beta \geq 0$ such that

$$\|Ax - x\| = \|y \oplus z\| \le 2\beta \|y\| = \beta \|Ax\|, \quad \forall x = y \oplus z \in \mathcal{Y} \oplus \mathcal{Z}.$$

In view of Theorem 2.9, we see that the statement given in [3] "the number 1/2 in Theorem 3.2 can not be improved" is false. However, the statement "1/2 is optimal for symmetry modulus" is true.

We now give applications to the theory of frames. Paley-Wiener theorem for orthonormal basis in Hilbert spaces inspired the study of perturbation of frames for Hilbert spaces. This was first derived by Christensen in his two papers [12,13]. This motivated the perturbation of frames and atomic decompositions for Banach spaces [16]. Crucial result used in all these perturbation results is the Neumann series. Later, using Theorem 1.2, Casazza and Christensen [11] improved the results obtained in paper [13]. Using Theorem 1.2 Stoeva made a systematic study of perturbations of frames for Banach spaces [47]. For the sake of completeness, we note that Theorem 1.2 was used in the study of perturbations of frames for Hilbert C*-modules [27].

Large body of work on frames for Hilbert spaces (see [14,23,28,29]) lead to the well developed theory of frames (known as Banach frames and \mathcal{X}_d -frames) for Banach spaces (see [7,8,18,25]) lead to the beginning of frames for frames for metric spaces (known as metric frames) [36].

For stating these definitions we need the definition of BK-space (Banach scalar valued sequence space or Banach co-ordinate space).

Definition 3.4. [2] A sequence space \mathcal{M}_d is said to be a BK-space if all the coordinate functionals are continuous, i.e., whenever $\{x_n\}_n$ is a sequence in \mathcal{M}_d converging to $x \in \mathcal{M}_d$, then each coordinate of x_n converges to each coordinate of x.

Definition 3.5. [36] Let \mathcal{M} be a metric space and \mathcal{M}_d be an associated BK-space. Let $\{f_n\}_n$ be a collection in $\operatorname{Lip}(\mathcal{M}, \mathbb{K})$ (where $\mathbb{K} = \mathbb{R}$ or \mathbb{C}) and $S : \mathcal{M}_d \to \mathcal{M}$. If:

(i) $\{f_n(x)\}_n \in \mathcal{M}_d$, for each $x \in \mathcal{M}$,

- (ii) There exist positive a, b such that $a d(x, y) \leq ||\{f_n(x) f_n(y)\}_n||_{\mathcal{M}_d} \leq b d(x, y), \forall x, y \in \mathcal{M},$
- (iii) S is Lipschitz and $S({f_n(x)}_n) = x$, for each $x \in \mathcal{M}$,

then we say that $(\{f_n\}_n, S)$ is a **metric frame** for \mathcal{M} with respect to \mathcal{M}_d . We say constant a as lower frame bound and constant b as upper frame bound.

As noted in [36], we observe that if $(\{f_n\}_n, S)$ is a metric frame for \mathcal{M} w.r.t. \mathcal{M}_d , then Definition 3.5 tells that the analysis map

$$\theta_f: \mathcal{M} \ni x \mapsto \theta_f x \coloneqq \{f_n(x)\}_n \in \mathcal{M}_d$$

is well-defined and bi-Lipschitz. Further, $S\theta_f = I_M$. Now we improve Theorem 4 in [11] to metric frames for Banach spaces.

Theorem 3.6. Let $(\{f_n\}_n, S)$ be a metric frame with lower frame bound a and upper frame bound b for a Banach space \mathcal{X} w.r.t. \mathcal{M}_d . Let $T : \mathcal{M}_d \to \mathcal{X}$ be a Lipschitz map and suppose that there exist $\lambda_1, \lambda_2, \mu \ge 0$ such that $\max\{\lambda_2, \lambda_1 + \mu b\} < 1$ and

$$||S\{c_n\}_n - S\{d_n\}_n - (T\{c_n\}_n - T\{d_n\}_n)|| \le \lambda_1 ||S\{c_n\}_n - S\{d_n\}_n|| + \lambda_2 ||T\{c_n\}_n - T\{d_n\}_n|| + \mu ||\{c_n - d_n\}_n||, \quad \forall \{c_n\}_n, \{d_n\}_n \in \mathcal{M}_d.$$
(8)

Then there exists a collection $\{g_n\}_n$ in $\operatorname{Lip}(\mathcal{X}, \mathbb{K})$ such that $(\{g_n\}_n, T)$ is a metric frame for \mathcal{X} with lower and upper frame bounds

$$\frac{a(1-\lambda_2)}{1+\lambda_1+\mu b}, \quad \frac{b(1+\lambda_2)}{1-(\lambda_1+\mu b)}$$

respectively.

Proof. Given $x, y \in \mathcal{X}$, by taking $\{c_n\}_n$ as $\theta_f x$ and $\{d_n\}_n$ as $\theta_f y$ in Inequality (8) we get $\|S\theta_f x - S\theta_f y - (T\theta_f x - T\theta_f y)\| \le \lambda_1 \|S\theta_f x - S\theta_f y\| + \lambda_2 \|T\theta_f x - T\theta_f y\| + \mu \|\theta_f x - \theta_f y\|, \quad \forall x, y \in \mathcal{X}.$

But $S\theta_f x = x, \forall x \in \mathcal{X}$ and hence

$$\begin{aligned} \|x - y - (T\theta_f x - T\theta_f y)\| &\leq \lambda_1 \|x - y\| + \lambda_2 \|T\theta_f x - T\theta_f y\| + \mu \|\theta_f x - \theta_f y\| \\ &\leq (\lambda_1 + \mu \operatorname{Lip}(\theta_f)) \|x - y\| + \lambda_2 \|T\theta_f x - T\theta_f y\| \\ &\leq (\lambda_1 + \mu b) \|x - y\| + \lambda_2 \|T\theta_f x - T\theta_f y\|, \quad \forall x, y \in \mathcal{X}. \end{aligned}$$

Theorem 2.9 now says that the map $T\theta_f$ is Lipschitz invertible and

$$\frac{1-\lambda_2}{1+\lambda_1+\mu b} \le \operatorname{Lip}(T\theta_f)^{-1} \le \frac{1+\lambda_2}{1-(\lambda_1+\mu b)}$$

Define $g_n \coloneqq f_n(T\theta_f)^{-1}$ for all $n \in \mathbb{N}$. Then g_n is Lipschitz for all $n, \{g_n(x)\}_n \in \mathcal{M}_d$, for each $x \in \mathcal{M}$ and

$$\begin{aligned} \|\{g_n(x) - g_n(y)\}_n\| &= \|\{f_n((T\theta_f)^{-1}(x)) - f_n((T\theta_f)^{-1}(y))\}_n\| \le b\|(T\theta_f)^{-1}(x) - (T\theta_f)^{-1}(y)\| \\ &\le b\frac{1+\lambda_2}{1-(\lambda_1+\mu b)}\|x-y\|, \quad \forall x, y \in \mathcal{X}, \end{aligned}$$
$$a\frac{1-\lambda_2}{1+\lambda_1+\mu b}\|x-y\| \le a\|(T\theta_f)^{-1}(x)) - (T\theta_f)^{-1}(y)\| \le \|\{f_n((T\theta_f)^{-1}(x)) - f_n((T\theta_f)^{-1}(y))\}_n\|$$

$$= \|\{g_n(x) - g_n(y)\}_n\|, \quad \forall x, y \in \mathcal{X}$$

which establish lower and upper frame bounds. Further,

$$T\{g_n(x)\}_n = T\{f_n(T\theta_f)^{-1}(x)\}_n = T\theta_f((T\theta_f)^{-1}(x)) = x, \quad \forall x \in \mathcal{X}$$

Therefore $(\{g_n\}_n, T)$ is a metric frame for \mathcal{X} .

We now give another application of Theorem 2.9. For this purpose, we introduce the notion of non-linear atomic decompositions.

Definition 3.7. Let \mathcal{X} be a Banach space \mathcal{X} and \mathcal{M}_d be a BK-space. Let $\{f_n\}_n$ be a sequence in $\operatorname{Lip}(\mathcal{X}, \mathbb{K})$ and $\{\tau_n\}_n$ to be a sequence in \mathcal{X} If:

- (i) $\{f_n(x)\}_n \in \mathcal{M}_d$, for each $x \in \mathcal{X}$,
- (ii) There exist positive a, b such that

$$a \|x - y\| \le \|\{f_n(x) - f_n(y)\}_n\|_{\mathcal{M}_d} \le b \|x - y\|, \quad \forall x, y \in \mathcal{X},$$

(iii) $x = \sum_{n=1}^{\infty} f_n(x)\tau_n$, for each $x \in \mathcal{X}$,

then we say that $(\{f_n\}_n, \{\tau_n\}_n)$ is a Lipschitz atomic decomposition for \mathcal{X} with respect to \mathcal{M}_d . We say constant a as lower Lipschitz atomic bound and constant b as upper Lipschitz atomic bound.

In [8] it is proved that not every Banach space admits an atomic decomposition. Motivated from this, we ask the following open problem.

Question 3.8. Classify Banach spaces which admit Lipschitz atomic decompositions.

Following proposition gives various examples of Lipschitz atomic decompositions.

Proposition 3.9. Let $(\{g_n\}_n, \{\omega_n\}_n)$ be an atomic decomposition for a Banach space \mathcal{Y} w.r.t. BK-space \mathcal{M}_d . Let \mathcal{X} be a Banach space and let $A : \mathcal{X} \to \mathcal{Y}$ be a bi-Lipschitz map such that there exists a linear map $A : \mathcal{Y} \to \mathcal{X}$ satisfying $BA = I_{\mathcal{X}}$. Then $(\{f_n \coloneqq g_n A\}_n, \{\tau_n \coloneqq B\omega_n\}_n)$ is a Lipschitz atomic decomposition \mathcal{X} w.r.t. \mathcal{M}_d . In particular, if a Banach space admits a Schauder basis, then it admits a Lipschitz atomic decomposition.

Particular case of Proposition 3.9 gives the following example.

Example 3.10. Let $(\{g_n\}_n, \{\omega_n\}_n)$ be an atomic decomposition for a Banach space \mathcal{X} w.r.t. a BKspace \mathcal{M}_d . Let $T : \mathcal{X} \to \mathcal{X}$ be any bi-Lipschitz map. Define $A : \mathcal{X} \ni x \mapsto (x, Tx) \in \mathcal{X} \oplus \mathcal{X}$ and $B : \mathcal{X} \oplus \mathcal{X} \ni (x, y) \mapsto x \in \mathcal{X}$. Then A is bi-Lipschitz, B is linear and satisfies $BA = I_{\mathcal{X}}$. Hence $(\{f_n \coloneqq g_n A\}_n, \{\tau_n \coloneqq B\omega_n\}_n)$ is a Lipschitz atomic decomposition for \mathcal{X} w.r.t. \mathcal{M}_d .

At this point it seems that it is best to develop some theory of Lipschitz atomic decompositions before giving an application of Theorem 2.9.

Proposition 2.3 in [8] shows that under certain conditions, there is a close relationship between Banach frames and atomic decompositions. Following is the non-linear version of that result.

Proposition 3.11. Let \mathcal{X} be a Banach space and \mathcal{M}_d be a BK-space. Let $\{f_n\}_n$ be a sequence in $\operatorname{Lip}(\mathcal{X}, \mathbb{K})$ and $S : \mathcal{M}_d \to \mathcal{X}$ be a bounded linear operator. If the standard unit vectors $\{e_n\}_n$ form a Schauder basis for \mathcal{M}_d , then the following are equivalent.

- (i) $(\{f_n\}_n, S)$ is a metric frame for \mathcal{X} .
- (ii) $(\{f_n\}_n, \{Se_n\}_n)$ is a Lipschitz atomic decomposition for \mathcal{X} w.r.t. \mathcal{M}_d .

Proof. We set $\tau_n = Se_n, \forall n \in \mathbb{N}$ and see that

$$\sum_{n=1}^{\infty} f_n(x)\tau_n = \sum_{n=1}^{\infty} f_n(x)Se_n = S\left(\sum_{n=1}^{\infty} f_n(x)e_n\right) = S\left(\{f_n(x)\}_n\right), \quad \forall x \in \mathcal{X}.$$

Well established dilation theory of frames for Hilbert spaces says that frames are images of Riesz bases under projections [19,29,34]. This result has been extended to frames and atomic decompositions for Banach spaces [8,30–32,37]. In the next theorem we derive a dilation result for Lipschitz atomic decompositions. We need a proposition to use in the theorem.

Proposition 3.12. [38] A sequence $\{\tau_n\}_n$ in a Banach space \mathcal{X} is a Schauder basis for \mathcal{X} if and only if the following three conditions hold.

- (i) $\tau_n \neq 0$ for all n.
- (ii) There exists b > 0 such that for every sequence $\{a_k\}_k$ of scalars and every pair of natural numbers n < m, we have

$$\left\|\sum_{k=1}^{n} a_k \tau_k\right\| \le b \left\|\sum_{k=1}^{m} a_k \tau_k\right\|$$

(iii) $\overline{\operatorname{span}}\{\tau_n\}_n = \mathcal{X}.$

Theorem 3.13. Let $(\{f_n\}_n, \{\tau_n\}_n)$ be a Lipschitz atomic decomposition for a Banach space \mathcal{X} w.r.t. \mathcal{M}_d . Then there is a Banach space \mathcal{Z} with a Schauder basis $\{\omega_n\}_n$, an injective map $\theta : \mathcal{X} \to \mathcal{Z}$ and a map $P : \mathcal{Z} \to \mathcal{Z}$ satisfying $P(\mathcal{Z}) = \mathcal{X}$, $P^2 = P$ and $P\omega_n = \theta\tau_n, \forall n \in \mathbb{N}$.

Proof. We generalize the idea of proof of Theorem 2.6 in [8] (which is motivated from the arguments in [41]) to non-linear setting. Let c_{00} be the vector space of scalar sequences with only finitely many non-zero terms. Let $\{e_n\}_n$ be the standard unit vectors in c_{00} .

Case (i): $\tau_n \neq 0$, for all n. We define a norm on c_{00} as follows. Let $\{a_n\}_n \in c_{00}$. Define

(9)
$$\left\|\sum_{n=1}^{\infty} a_n e_n\right\| \coloneqq \max_n \left\|\sum_{k=1}^n a_k \tau_k\right\|.$$

Proposition 3.12 then tells that $\{e_n\}_n$ is a Schauder basis for the completion of c_{00} , call as \mathcal{Z} w.r.t. just defined norm. Define

$$\theta: \mathcal{X} \ni x \mapsto \theta x \coloneqq \sum_{n=1}^{\infty} f_n(x) e_n \in \mathcal{Z}$$

From the first condition of the definition of Lipschitz atomic decomposition, from Definition 9 and from the construction of \mathcal{Z} , it follows that θ is well-defined. From the third condition of definition of atomic decomposition, θ is injective. We next define

$$\Gamma: \mathcal{Z} \ni \sum_{n=1}^{\infty} a_n e_n \mapsto \Gamma\left(\sum_{n=1}^{\infty} a_n e_n\right) \coloneqq \sum_{n=1}^{\infty} a_n \tau_n \in \mathcal{X}.$$

By verifying Γ is bounded linear on dense subset c_{00} of \mathcal{Z} , we see that Γ is bounded on \mathcal{Z} . Then

(10)
$$\Gamma\theta x = \Gamma\left(\sum_{n=1}^{\infty} f_n(x)e_n\right) = \sum_{n=1}^{\infty} f_n(x)\tau_n = x, \quad \forall x \in \mathcal{X}.$$

IMPROVING CASAZZA-KALTON-CHRISTENSEN-VAN EIJNDHOVEN PERTURBATION ...

So if we define $P := \theta \Gamma$, then $P^2 = \theta \Gamma \theta \Gamma = \theta \Gamma = P$. Equation (10) tells that $P(\mathcal{Z}) = \mathcal{X}$. We next see that $Pe_n = \theta \Gamma e_n = \theta \tau_n$, $\forall n$. Thus we can take $\omega_n = e_n$, for all n to get the result.

Case (ii): $\tau_n = 0$, for some n. Let $J = \{n : \tau_n \neq 0\}$. We now apply case (i) to the collection $\{a_n\}_{n \in J}$. Let θ , \mathcal{Z} , Γ and P be as in the case (i). Without affecting the definition of atomic decomposition, we can take $f_n = 0$ for all $n \in J^c$. Now consider the space $\mathcal{Z} \oplus \ell^2(J^c)$ and let $\{\rho_n\}_{n \in J^c}$ be an orthonormal basis for $\ell^2(J^c)$. Define $Q: \mathcal{Z} \oplus \ell^2(J^c) \ni z \oplus y \mapsto Q(z \oplus y) \coloneqq Pz \oplus 0 \in \mathcal{Z} \oplus \ell^2(J^c)$. Now the space $\mathcal{Z} \oplus \ell^2(J^c)$ has Schauder basis $\{\tau_n \oplus 0, 0 \oplus \rho_m\}_{n \in J, m \in J^c}$ and Q satisfy the conclusions. Thus we can take $\omega_n = e_n$, for all $n \in J$ and $\omega_n = \rho_n$, for all $n \in J^c$ to get the result.

We leave the further study of Lipschitz atomic decomposition to future work and end the paper with an application of Theorem 2.9.

Theorem 3.14. Let $(\{f_n\}_n, \{\tau_n\}_n)$ be a Lipschitz atomic decomposition with lower Lipschitz atomic bound a and upper Lipschitz atomic bound b for \mathcal{X} w.r.t. \mathcal{M}_d . Let $\{\omega_n\}_n$ be a collection in \mathcal{X} and suppose that there exist $\lambda_1, \lambda_2, \mu \geq 0$ such that the following conditions hold.

- (i) For each $x \in \mathcal{X}$, the series $\sum_{n=1}^{\infty} f_n(x)\omega_n$ converges in \mathcal{X} .
- (ii) $\max\{\lambda_2, \lambda_1 + \mu b\} < 1$ and

(iii)

(11)
$$\left\|\sum_{n=1}^{\infty} (c_n - d_n)(\tau_n - \omega_n)\right\| \leq \lambda_1 \left\|\sum_{n=1}^{\infty} (c_n - d_n)\tau_n\right\| + \lambda_2 \left\|\sum_{n=1}^{\infty} (c_n - d_n)\omega_n\right\| + \mu \|\{c_n - d_n\}_n\|, \quad \forall \{c_n\}_n, \{d_n\}_n \in \mathcal{M}_d.$$

Then there exists a collection $\{g_n\}_n$ in $\operatorname{Lip}(\mathcal{X}, \mathbb{K})$ such that $(\{g_n\}_n, \{\omega_n\}_n)$ is a Lipschitz atomic decomposition for \mathcal{X} with lower and upper Lipschitz atomic bounds

$$\frac{a(1-\lambda_2)}{1+\lambda_1+\mu b}, \quad \frac{b(1+\lambda_2)}{1-(\lambda_1+\mu b)}$$

respectively.

Proof. From the first condition we get that the map $T : \mathcal{X} \ni x \mapsto \sum_{n=1}^{\infty} f_n(x)\omega_n \in \mathcal{X}$ is well-defined. Now using Inequality (11),

$$\begin{aligned} \|x - y - (Tx - Ty)\| &= \left\| \sum_{n=1}^{\infty} (f_n(x) - f_n(y))\tau_n - \sum_{n=1}^{\infty} (f_n(x) - f_n(y))\omega_n \right\| \\ &= \left\| \sum_{n=1}^{\infty} (f_n(x) - f_n(y))(\tau_n - \omega_n) \right\| \\ &\leq \lambda_1 \left\| \sum_{n=1}^{\infty} (f_n(x) - f_n(y))\tau_n \right\| + \lambda_2 \left\| \sum_{n=1}^{\infty} (f_n(x) - f_n(y))\omega_n \right\| + \mu \|\{f_n(x) - f_n(y)\}_n\| \\ &= \lambda_1 \|x - y\| + \lambda_2 \|Tx - Ty\| + \mu \|\{f_n(x) - f_n(y)\}_n\| \\ &\leq (\lambda_1 + \mu b) \|x - y\| + \lambda_2 \|Tx - Ty\|, \quad \forall x, y \in \mathcal{X}. \end{aligned}$$

Theorem 2.9 then says that T is Lipschitz, invertible and

$$\frac{1-\lambda_2}{1+\lambda_1+\mu b} \le \operatorname{Lip}(T)^{-1} \le \frac{1+\lambda_2}{1-(\lambda_1+\mu b)}$$

Define $g_n \coloneqq f_n T^{-1}, \forall n \in \mathbb{N}$. Then $\{g_n(x)\}_n \in \mathcal{M}_d$, for each $x \in \mathcal{X}$ and

$$\begin{aligned} \|\{g_n(x) - g_n(y)\}_n\| &= \|\{f_n(T^{-1}x) - f_n(T^{-1}y)\}_n\|_n \le b\|T^{-1}x - T^{-1}y\|\\ &\le b\frac{1+\lambda_2}{1-(\lambda_1+\mu b)}\|x-y\|, \quad \forall x, y \in \mathcal{X},\\ a\frac{1-\lambda_2}{1+\lambda_1+\mu b}\|x-y\| \le a\|T^{-1}x - T^{-1}y\| \le \|\{f_n(T^{-1}x) - f_n(T^{-1}y)\}_n\| \end{aligned}$$

Finally

$$\sum_{n=1}^{\infty} g_n(x)\omega_n = \sum_{n=1}^{\infty} f_n(T^{-1}x)\omega_n = T(T^{-1}x) = x, \quad \forall x \in \mathcal{X}.$$

 $= \|\{g_n(x) - g_n(y)\}_n\|, \quad \forall x, y \in \mathcal{X}.$

We conclude the paper with the following remarks.

Remark 3.15. We can improve Theorems 3.1, 3.2, 3.6 and 3.14 using Theorem 2.13.

Remark 3.16. So far in the literature, there are three ways to prove Theorem 1.2 one given in [9], another in [49] and yet another in [11]. As we mentioned earlier, we have done the non-linear version of arguments used in [49]. We hope that arguments used in [9] and [11] can be generalized to give different proofs of Theorem 2.9.

4. Acknowledgements

I thank Dr. P. Sam Johnson, Department of Mathematical and Computational Sciences, National Institute of Technology Karnataka (NITK), Surathkal for several discussions.

References

- [1] Maynard G. Arsove. The Paley-Wiener theorem in metric linear spaces. Pacific J. Math., 10:365–379, 1960.
- [2] Jozef Banas and Mohammad Mursaleen. Sequence spaces and measures of noncompactness with applications to differential and integral equations. Springer, New Delhi, 2014.
- [3] Annamaria Barbagallo, Octavian-Emil Ernst, and Michel Théra. Orthogonality in locally convex spaces: two nonlinear generalizations of Neumann's lemma. J. Math. Anal. Appl., 484(1):123663, 18, 2020.
- [4] R. P. Boas, Jr. Expansions of analytic functions. Trans. Amer. Math. Soc., 48:467–487, 1940.
- [5] Sara Botelho-Andrade, Peter G. Casazza, Desai Cheng, and Tin T. Tran. The solution to the frame quantum detection problem. J. Fourier Anal. Appl., 25(5):2268–2323, 2019.
- [6] Sergio Campanato. On the condition of nearness between operators. Ann. Mat. Pura Appl. (4), 167:243–256, 1994.
- [7] Peter Casazza, Ole Christensen, and Diana T. Stoeva. Frame expansions in separable Banach spaces. J. Math. Anal. Appl., 307(2):710–723, 2005.
- [8] Peter G. Casazza, Deguang Han, and David R. Larson. Frames for Banach spaces. In The functional and harmonic analysis of wavelets and frames (San Antonio, TX, 1999), volume 247 of Contemp. Math., pages 149–182. Amer. Math. Soc., Providence, RI, 1999.
- [9] Peter G. Casazza and Nigel J. Kalton. Generalizing the Paley-Wiener perturbation theory for Banach spaces. Proc. Amer. Math. Soc., 127(2):519–527, 1999.
- [10] Peter G. Casazza, Gitta Kutyniok, and Shidong Li. Fusion frames and distributed processing. Appl. Comput. Harmon. Anal., 25(1):114–132, 2008.
- [11] Peter G. Cazassa and Ole Christensen. Perturbation of operators and applications to frame theory. J. Fourier Anal. Appl., 3(5):543–557, 1997.
- [12] Ole Christensen. Frame perturbations. Proc. Amer. Math. Soc., 123(4):1217–1220, 1995.

IMPROVING CASAZZA-KALTON-CHRISTENSEN-VAN EIJNDHOVEN PERTURBATION ...

- [13] Ole Christensen. A Paley-Wiener theorem for frames. Proc. Amer. Math. Soc., 123(7):2199–2201, 1995.
- [14] Ole Christensen. An introduction to frames and Riesz bases. Applied and Numerical Harmonic Analysis. Birkhäuser/Springer, [Cham], second edition, 2016.
- [15] Ole Christensen and Marzieh Hasannasab. Operator representations of frames: boundedness, duality, and stability. Integral Equations Operator Theory, 88(4):483–499, 2017.
- [16] Ole Christensen and Christopher Heil. Perturbations of Banach frames and atomic decompositions. Math. Nachr., 185:33–47, 1997.
- [17] Ole Christensen, Chris Lennard, and Christine Lewis. Perturbation of frames for a subspace of a Hilbert space. Rocky Mountain J. Math., 30(4):1237–1249, 2000.
- [18] Ole Christensen and Diana T. Stoeva. p-frames in separable Banach spaces. Adv. Comput. Math., 18(2-4):117–126, 2003.
- [19] Wojciech Czaja. Remarks on Naimark's duality. Proc. Amer. Math. Soc., 136(3):867-871, 2008.
- [20] H. G. Dales. Banach algebras and automatic continuity, volume 24 of London Mathematical Society Monographs. New Series. The Clarendon Press, Oxford University Press, New York, 2000.
- [21] Béla de Sz. Nagy. Expansion theorems of Paley-Wiener type. Duke Math. J., 14:975–978, 1947.
- [22] Jiu Ding. New perturbation results on pseudo-inverses of linear operators in Banach spaces. Linear Algebra Appl., 362:229–235, 2003.
- [23] R. J. Duffin and A. C. Schaeffer. A class of nonharmonic Fourier series. Trans. Amer. Math. Soc., 72:341–366, 1952.
- [24] Jean-Pierre Gabardo and Deguang Han. Frames associated with measurable spaces. Adv. Comput. Math., 18(2-4):127– 147, 2003.
- [25] Karlheinz Gröchenig. Describing functions: atomic decompositions versus frames. Monatsh. Math., 112(1):1–42, 1991.
- [26] Xunxiang Guo. Perturbations of invertible operators and stability of g-frames in Hilbert spaces. Results Math., 64(3-4):405-421, 2013.
- [27] Deguang Han, Wu Jing, and Ram N. Mohapatra. Perturbation of frames and Riesz bases in Hilbert C*-modules. Linear Algebra Appl., 431(5-7):746–759, 2009.
- [28] Deguang Han, Keri Kornelson, David Larson, and Eric Weber. Frames for undergraduates, volume 40 of Student Mathematical Library. American Mathematical Society, Providence, RI, 2007.
- [29] Deguang Han and David R. Larson. Frames, bases and group representations. Mem. Amer. Math. Soc., 147(697):x+94, 2000.
- [30] Deguang Han, David R. Larson, Bei Liu, and Rui Liu. Dilations of frames, operator-valued measures and bounded linear maps. In Operator methods in wavelets, tilings, and frames, volume 626 of Contemp. Math., pages 33–53. Amer. Math. Soc., Providence, RI, 2014.
- [31] Deguang Han, David R. Larson, Bei Liu, and Rui Liu. Operator-valued measures, dilations, and the theory of frames. Mem. Amer. Math. Soc., 229(1075):viii+84, 2014.
- [32] Deguang Han, David R. Larson, and Rui Liu. Dilations of operator-valued measures with bounded p-variations and framings on Banach spaces. J. Funct. Anal., 274(5):1466–1490, 2018.
- [33] Sven H. Hilding. Note on completeness theorems of Paley-Wiener type. Ann. of Math. (2), 49:953–955, 1948.
- [34] B. S. Kashin and T. Yu. Kulikova. A remark on the description of frames of general form. Mat. Zametki, 72(6):941–945, 2002.
- [35] Yoo Young Koo and Jae Kun Lim. Perturbation of frame sequences and its applications to shift-invariant spaces. *Linear Algebra Appl.*, 420(2-3):295–309, 2007.
- [36] K. Mahesh Krishna and P. Sam Johnson. Frames for metric spaces. http://arXiv.org/2011.01870v1 [math.FA] 3 November 2020.
- [37] David R. Larson and Franciszek Hugon Szafraniec. Framings and dilations. Acta Sci. Math. (Szeged), 79(3-4):529–543, 2013.
- [38] Joram Lindenstrauss and Lior Tzafriri. Classical Banach spaces. I. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 92. Springer-Verlag, Berlin-New York, 1977.
- [39] Carl Neumann. Untersuchungen uber das logarithmische und Newtonsche Potential. Teubner, Leipzig, 1877.
- [40] Raymond E. A. C. Paley and Norbert Wiener. Fourier transforms in the complex domain, volume 19 of American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, RI, 1987.
- [41] A. Peł czyński. Any separable Banach space with the bounded approximation property is a complemented subspace of a Banach space with a basis. *Studia Math.*, 40:239–243, 1971.

- [42] Harry Pollard. Completeness theorems of Paley-Wiener type. Ann. of Math. (2), 45:738–739, 1944.
- [43] James R. Retherford. Basic sequences and the Paley-Wiener criterion. Pacific J. Math., 14:1019–1027, 1964.
- [44] Friedrich Wilhelm Shäfke. Das Kriterium von Paley und Wiener im Banachschen Raum. Math. Nachr., 3:59–61, 1949.
- [45] Gustaf Söderlind. Bounds on nonlinear operators in finite-dimensional Banach spaces. Numer. Math., 50(1):27–44, 1986.
- [46] D. T. Stoeva and P. Balazs. Invertibility of multipliers. Appl. Comput. Harmon. Anal., 33(2):292–299, 2012.
- [47] Diana T. Stoeva. Perturbation of frames in Banach spaces. Asian-Eur. J. Math., 5(1):1250011, 15, 2012.
- [48] Wenchang Sun. Stability of g-frames. J. Math. Anal. Appl., 326(2):858–868, 2007.
- [49] S. J. L. van Eijndhoven. Hilding's theorem for banach spaces. RANA: reports on applied and numerical analysis, volume 9612: Technische Universiteit Eindhoven, pages 1–6, 1996.
- [50] Nik Weaver. Lipschitz algebras. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2018.
- [51] Xiaodan Yang and Yuwen Wang. Some new perturbation theorems for generalized inverses of linear operators in Banach spaces. *Linear Algebra Appl.*, 433(11-12):1939–1949, 2010.
- [52] Robert M. Young. An introduction to nonharmonic Fourier series, volume 93 of Pure and Applied Mathematics. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1980.
- [53] Ping Zhao, Chun Zhao, and Peter G. Casazza. Perturbation of regular sampling in shift-invariant spaces for frames. IEEE Trans. Inform. Theory, 52(10):4643–4648, 2006.