

Slow manifolds for stochastic Koper models with stable Lévy noises

Hina Zulfiqar^a, Shenglan Yuan^{b,*}, Muhammad Shoaib Saleem^a

^a*Department of Mathematics, University of Okara, Renala Khurd, Punjab, Pakistan*

^b*Institut für Mathematik, Universität Augsburg, 86135, Augsburg, Germany*

Abstract

The Koper model is a vector field in which the differential equations describe the electrochemical oscillations appearing in diffusion processes. This work focuses on the understanding of the slow dynamics of stochastic Koper model perturbed by stable Lévy noise. We establish the slow manifold for stochastic Koper model with stable Lévy noise and verify exponential tracking property. We also present a practical example to demonstrate the analytical results with numerical simulations.

Keywords: random slow manifold, Koper model, fast-slow stochastic system, Lévy motion.

2020 MSC: 37D10; 37H30; 60H10

1. Introduction

The Koper model [17] is an idealized model of the chemical reaction described in [3]. Invariant manifolds are useful in investigating the dynamical behavior of the multiscale systems [4, 14]. Invariant manifolds for investigating the dynamical behavior of deterministic systems without being influenced of stochastic forces are discussed in [8, 10, 23], while invariant manifolds for deterministic systems influenced by stochastic forces are constructed in [5, 9, 33]. An invariant manifold for a fast-slow stochastic system in which fast mode is indicated by the slow mode tends to slow manifold when the scale parameter goes to zero. Moreover, slow manifold converges to critical manifold as the scale parameter approaches zero.

Slow manifold for stochastic system driven by Brownian motion is demonstrated in [13, 24, 27], and its numerical simulations are presented in [21, 22]. Lévy motions arise from the models for fluctuations, they have independent, stationary increments and discontinuous paths. For example, Lévy processes affect the evolution of the state variables in the turbulent flow of fluids [32]. Some models about stochastic systems processed by Lévy noise are explained in [20, 31, 34]. The slow manifolds under Lévy noise are constructed in [35]. The existence of slow manifold for nonlocal fast-slow stochastic evolutionary equations is proved in [36, 37]. Invariant manifold of variable stability in the Koper model is established in [3]. It continues to be an active topic on the characterization of stochastic Koper model driven by Lévy process for both theoretical reasons and applications.

The goal of this article is to construct the three-dimensional stochastic Koper model in Euclidean space \mathbb{R}^3 and establish the existence of slow manifold for stochastic Koper model processed by α -stable Lévy

*Corresponding author

Email addresses: zhinazulfiqar@hotmail.com (Hina Zulfiqar), shenglan.yuan@math.uni-augsburg.de (Shenglan Yuan), shoaib83455@gmail.com (Muhammad Shoaib Saleem)

noise with $\alpha \in (1, 2)$. Namely, we consider the stochastic Koper system in the following version

$$\begin{cases} \dot{x} = \frac{1}{\epsilon}[ky - x^3 + 3x - \lambda(z)] + \sigma\epsilon^{-\frac{1}{\alpha}}\dot{L}_t^\alpha, \\ \dot{y} = x - 2y + z, \\ \dot{z} = \hat{\epsilon}(y - z), \end{cases} \quad (1)$$

where k and $\lambda(z)$ are the main bifurcation parameters. Here, $\hat{\epsilon} = 1$, but ϵ represents a small parameter with the property $0 < \epsilon \ll 1$, and it indicates the ratio of two times scales such that stochastic fast-slow system (1) has one fast variable x , and two slow variables y and z . The dot stands for the differentiation with respect to time t . The noise L_t^α is two-sided symmetric α -stable Lévy process taking real values, with the stability index $\alpha \in (1, 2)$ and the intensity $\sigma > 0$; see the references [1, 10].

We start from a random transformation such that a solution of stochastic Koper model (1) can be expressed as a transformed solution of some random system. The establishment of slow manifold for defined random system is proved by the utilization of Lyapunov-Perron method [9, 12].

The article is organized as follows. In Section 2, some concepts about random dynamical system, and stochastic differential equation processed by Lévy motion are discussed. In Section 3, the stability of the stochastic Koper system (1) is proved and a random transformation is defined, which converts stochastic Koper system (1) into random system. In Section 4, a short review about random invariant manifold and the existence of exponential tracking slow manifold for random system is provided. In Section 5, we present numerical results using an example from electrochemical oscillations to corroborate our analytical results. Finally, Section 6 summarizes our findings as well as directions for future study.

2. Preliminaries

In this section, some concepts about the random dynamical system are given.

Definition 1. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $\theta = \{\theta_t\}_{t \in \mathbb{R}}$ be a flow on Ω satisfying the conditions

- $\theta_0 = \text{Id}_\Omega$;
- $\theta_{t_1}\theta_{t_2} = \theta_{t_1+t_2}$, where $t_1, t_2 \in \mathbb{R}$;

and the mapping $\theta : \mathbb{R} \times \Omega \rightarrow \Omega$ can be defined by $(t, \omega) \mapsto \theta_t\omega$, which is $\mathcal{B}(\mathbb{R}) \otimes \mathcal{F} - \mathcal{F}$ measurable. Here, we consider that the probability measure \mathbb{P} is invariant with respect to the flow $\{\theta_t\}_{t \in \mathbb{R}}$, i.e., $\theta_t\mathbb{P} = \mathbb{P}$ for all $t \in \mathbb{R}$. Then $\Theta = (\Omega, \mathcal{F}, \mathbb{P}, \theta)$ is said to be a metric dynamical system [35].

Throughout this article, we use scalar Lévy process. Take L_t^α , $\alpha \in (1, 2)$ as a symmetric two-sided α -stable Lévy process with values in \mathbb{R} . Consider a canonical sample space for it. Let $\Omega = D(\mathbb{R}, \mathbb{R})$ be the space of càdlàg functions having zero value at $t = 0$, i.e.,

$$D(\mathbb{R}, \mathbb{R}) = \{\omega : \text{for } \forall t \in \mathbb{R}, \lim_{s \uparrow t} \omega(s) = \omega(t-), \lim_{s \downarrow t} \omega(s) = \omega(t) \text{ exist and } \omega(0) = 0\}.$$

If we use standard usual open-compact metric, then the space $D(\mathbb{R}, \mathbb{R})$ may not be separable and complete. But the space $D(\mathbb{R}, \mathbb{R})$ of real-valued càdlàg functions can be extended by introducing another metric d^0 since it can be made into the complete and separable space on unit interval or on \mathbb{R} [7]. For functions $\omega_1, \omega_2 \in D(\mathbb{R}, \mathbb{R})$, $d^0(\omega_1, \omega_2)$ is defined by

$$d^0(\omega_1, \omega_2) = \inf \left\{ \varepsilon > 0 : |\omega_1(t) - \omega_2(\lambda t)| \leq \varepsilon, \left| \ln \frac{\arctan(\lambda t) - \arctan(\lambda s)}{\arctan(t) - \arctan(s)} \right| \leq \varepsilon, \right. \\ \left. \text{for every } t, s \in \mathbb{R} \text{ and some } \lambda \in \Lambda^{\mathbb{R}} \right\},$$

where

$$\Lambda^{\mathbb{R}} = \{\lambda : \mathbb{R} \rightarrow \mathbb{R}; \lambda \text{ is injective increasing, } \lim_{t \rightarrow -\infty} \lambda(t) = -\infty, \lim_{t \rightarrow \infty} \lambda(t) = \infty\}.$$

By Theorem 3.2 in [28], the space $D(\mathbb{R}, \mathbb{R})$ equipped with Skorokhod's \mathcal{J}_1 -topology generated by the metric d^0 is a complete and separable space, i.e., Polish space. On this Polish space, we consider a measurable flow $\theta = \{\theta_t\}_{t \in \mathbb{R}}$ defined by mapping

$$\theta : \mathbb{R} \times D(\mathbb{R}, \mathbb{R}) \rightarrow D(\mathbb{R}, \mathbb{R}), \text{ such that, } \theta_t \omega(\cdot) = \omega(\cdot + t) - \omega(t),$$

where $\omega \in D(\mathbb{R}, \mathbb{R})$.

The sample paths of Lévy process are in $D(\mathbb{R}, \mathbb{R})$. Assume that \mathbb{P} is the probability measure on $\mathcal{F} = \mathcal{B}(D(\mathbb{R}, \mathbb{R}))$ introduced by the distribution of symmetric two-sided α -stable Lévy process. We consider the restriction of \mathbb{P} on \mathcal{F} , but still it is indicated by \mathbb{P} . Observe that \mathbb{P} is ergodic with respect to the shift $\{\theta_t\}_{t \in \mathbb{R}}$. Thus $(D(\mathbb{R}, \mathbb{R}), \mathcal{F}, \mathbb{P}, \{\theta_t\}_{t \in \mathbb{R}})$ is a metric dynamical system. It is worth pointing out that we can take a subset $\Omega_1 = D^0(\mathbb{R}, \mathbb{R}) \subset \Omega = D(\mathbb{R}, \mathbb{R})$ with \mathbb{P} -measure one instead of $D(\mathbb{R}, \mathbb{R})$. Here, $D^0(\mathbb{R}, \mathbb{R})$ is $\{\theta_t\}_{t \in \mathbb{R}}$ -invariant, which means that $\theta_t \Omega_1 = \Omega_1$ for $t \in \mathbb{R}$.

Definition 2. A cocycle ϕ satisfies that

$$\begin{aligned} \phi(0, \omega, u) &= u; \\ \phi(t_1 + t_2, \omega, u) &= \phi(t_2, \theta_{t_1} \omega, \phi(t_1, \omega, u)). \end{aligned}$$

It is $\mathcal{B}(\mathbb{R}) \otimes \mathcal{F} \otimes \mathcal{B}(\mathbb{R}^3) - \mathcal{F}$ measurable and defined by map:

$$\phi : \mathbb{R} \times \Omega \times \mathbb{R}^3 \rightarrow \mathbb{R}^3,$$

for $u \in \mathbb{R}^3$, $\omega \in \Omega$ and $t_1, t_2 \in \mathbb{R}$. Metric dynamical system $(\Omega, \mathcal{F}, \mathbb{P}, \theta)$ with the cocycle ϕ generates a random dynamical system [2].

The above cocycle property indicates that random dynamical system ϕ arrives at the same destination whether we consider the position $\phi(t_1 + t_2, \omega, u)$ of the path starting in u at time $t_1 + t_2$ or the position $\phi(t_2, \theta_{t_1} \omega, \phi(t_1, \omega, u))$ of the path with random initial state $\phi(t_1, \omega, u)$ at time t_2 . It is important to note that the path moves from u to $\phi(t_1, \omega, u)$, the underlying ω potentially may change as well over time t_1 . Instead of ω we need to use $\theta_{t_1} \omega$ for the new movement with the starting point $\phi(t_1, \omega, u)$, where θ_{t_1} indicates the new development of the underlying probability space over time t_1 .

If $u \mapsto \phi(t, \omega, u)$ is continuous or differentiable for $t \in \mathbb{R}$ and $\omega \in \Omega$, then random dynamical system $(\Omega, \mathcal{F}, \mathbb{P}, \theta, \phi)$ is also continuous or differentiable. The family of nonempty closed sets $\mathcal{M} = \{\mathcal{M}(\omega) \subset \mathbb{R}^3 : \omega \in \Omega\}$ is said to be a random set if for all $u' \in \mathbb{R}^3$ the map:

$$\omega \mapsto \inf_{u \in \mathcal{M}(\omega)} |u - u'|,$$

is a random variable.

Definition 3. If random variable $u(\omega)$ taking values in \mathbb{R}^3 , satisfies

$$\phi(t, \omega, u(\omega)) = u(\theta_t \omega), \quad a.s.$$

for all $t \in \mathbb{R}$. Then, random variable $u(\omega)$ is known as stationary orbit or random fixed point [11].

Definition 4. A random set $\mathcal{M} = \{\mathcal{M}(\omega) \subset \mathbb{R}^3 : \omega \in \Omega\}$ is called random positively invariant set [13] for random dynamical system ϕ , if

$$\phi(t, \omega, \mathcal{M}(\omega)) \subset \mathcal{M}(\theta_t \omega),$$

for all $\omega \in \Omega$ and $t \geq 0$.

Definition 5. Introduce a map

$$l : \mathbb{R}^2 \times \Omega \rightarrow \mathbb{R},$$

such that $v \mapsto l(v, \omega)$ is Lipschitz continuous for all $\omega \in \Omega$. Consider

$$\mathcal{M}(\omega) = \{(l(v, \omega), v) : v \in \mathbb{R}^2\},$$

such that random positively invariant set $\mathcal{M} = \{\mathcal{M}(\omega) \subset \mathbb{R}^3 : \omega \in \Omega\}$ can be expressed as a graph of Lipschitz continuous map l , then \mathcal{M} is known as Lipschitz continuous invariant manifold [35].

Moreover, \mathcal{M} posses the exponential tracking property, if there exists an $u' \in \mathcal{M}(\omega)$ for all $u \in \mathbb{R}^3$ satisfying

$$|\phi(t, \omega, u) - \phi(t, \omega, u')| \leq c_1(u, u', \omega)e^{c_2 t}|u - u'|,$$

for all $\omega \in \Omega$. Here, c_1 is positive random variable, and c_2 is negative constant.

3. Stability Analysis

Stochastic Koper system (1) is consist of one fast mode and two slow modes. The state space for the fast mode is \mathbb{R} , and the state space for the slow modes is \mathbb{R}^2 . For the construction of slow manifold, we assume the following hypotheses for the Koper system (1).

(H1) (Lipschitz continuity) With regard to nonlinear parts of (1), there are positive constants $L_1, L_2, L_3 > 0$ such that for all $(x_i, y_i, z_i)^T$ in \mathbb{R}^3 and for all $(x_j, y_j, z_j)^T$ in \mathbb{R}^3 ,

$$\begin{aligned} & |g_1(x_i, y_i, z_i) - g_1(x_j, y_j, z_j)| + |g_2(x_i, y_i, z_i) - g_2(x_j, y_j, z_j)| + |g_3(x_i, y_i, z_i) - g_3(x_j, y_j, z_j)| \\ & \leq L_1(|x_i - x_j| + |y_i - y_j| + |z_i - z_j|), \end{aligned}$$

where T is the transpose of the vector, and $g_m : \mathbb{R}^3 \rightarrow \mathbb{R}, m = 1, 2, 3$ are defined by $g_1(x, y, z) = ky - x^3 + 3x - \lambda(z), g_2(x, y, z) = x - 2y + z$, and $g_3(x, y, z) = y - z$.

(H2) (Growth) For all $(x, y, z) \in \mathbb{R}^3$, there exists a positive constant L such that

$$|g_1(x, y, z)|^2 + |g_2(x, y, z)|^2 + |g_3(x, y, z)|^2 \leq L(1 + |x|^2 + |y|^2 + |z|^2).$$

(H3) (Monotonocity) For all $x_1, x_2 \in \mathbb{R}$, there exists a positive constant L such that

$$(x_2 - x_1)(g_1(x_2) - g_1(x_1)) \leq -L|x_2 - x_1|^2.$$

Now let $\Theta_1 = (\Omega_1, \mathcal{F}_1, \mathbb{P}_1, \theta_t^1)$, $\Theta_2 = (\Omega_2, \mathcal{F}_2, \mathbb{P}_2, \theta_t^2)$ and $\Theta_3 = (\Omega_3, \mathcal{F}_3, \mathbb{P}_3, \theta_t^3)$ be three independent driving (metric) dynamical systems as mentioned in Section 2. Define

$$\Theta = \Theta_1 \times \Theta_2 \times \Theta_3 = (\Omega_1 \times \Omega_2 \times \Omega_3, \mathcal{F}_1 \otimes \mathcal{F}_2 \otimes \mathcal{F}_3, \mathbb{P}_1 \times \mathbb{P}_2 \times \mathbb{P}_3, (\theta_t^1, \theta_t^2, \theta_t^3)^T),$$

and

$$\theta_t \omega := (\theta_t^1 \omega_1, \theta_t^2 \omega_2, \theta_t^3 \omega_3)^T, \text{ for } \omega := (\omega_1, \omega_2, \omega_3)^T \in \Omega := \Omega_1 \times \Omega_2 \times \Omega_3.$$

Let $L_t^\alpha, \alpha \in (1, 2)$ be a two-sided symmetric α -stable Lévy process in \mathbb{R} with a generating triplet (a, \mathcal{Q}, ν) . We will prove the existence and uniqueness of solution for the stochastic Koper system (1).

Lemma 1. [19] Under Lipschitz condition **(H1)**, the equation

$$d\delta(t) = g_1(\delta(t))dt + \sigma dL_t^\alpha, \quad \delta(0) = \delta_0, \quad (2)$$

has the unique càdlàg solution

$$\delta(t) = \delta_0 + \int_0^t g_1(\delta(s))ds + \sigma L_t^\alpha.$$

According to the Lévy-Itô decomposition, L_t^α can be expressed as

$$L_t^\alpha = \int_{|w|<1} w\tilde{N}(t, dw) + \int_{|w|\geq 1} wN(t, dw).$$

It follows that

$$\delta(t) = \delta_0 + \int_0^t g_1(\delta(s))ds + \sigma \int_{|w|<1} w\tilde{N}(t, dw) + \sigma \int_{|w|\geq 1} wN(t, dw).$$

Remark 1. ([11, p.191]) L_{ct}^α and $c^{\frac{1}{\alpha}}L_t^\alpha$ have the same distribution for every $c > 0$, i.e., $L_{ct}^\alpha \stackrel{d}{=} c^{\frac{1}{\alpha}}L_t^\alpha$.

Lemma 2. Under the assumptions **(H1)**-**(H3)**, stochastic Koper system (1) has a unique solution.

Proof. Rewrite the stochastic Koper system (1) into the form

$$\begin{pmatrix} \dot{x} \\ \dot{y} \\ \dot{z} \end{pmatrix} = \begin{pmatrix} \frac{1}{\epsilon}g_1(x, y, z) \\ g_2(x, y, z) \\ g_3(x, y, z) \end{pmatrix} + \begin{pmatrix} \sigma\epsilon^{-\frac{1}{\alpha}}\dot{L}_t^\alpha \\ 0 \\ 0 \end{pmatrix}$$

From [29], under the assumptions **(H1)**-**(H3)**, it implies from [16, Theorem III.2.3.2] that there exists a unique solution of the equation (2). Moreover, there also exists one exponentially mixing invariant measure with respect to transition semigroup of $x(t)$. Then by [1, chapter 6], the assumptions **(H1)** – **(H3)** indicate that there exists a unique mild solution $(x(t), y(t), z(t))^T$ in \mathbb{R}^3 for the stochastic Koper system (1). \square

Define a random transformation

$$\begin{pmatrix} X \\ Y \\ Z \end{pmatrix} := \mu(\theta_t\omega, x, y, z) := \begin{pmatrix} x - \sigma\eta^\epsilon(\theta_t\omega) \\ y \\ z \end{pmatrix}.$$

where $\eta^\epsilon(\theta_t\omega) := \epsilon^{-\frac{1}{\alpha}}L_t^\alpha(\omega)$. Then $(X(t), Y(t), Z(t)) = \mu(\theta_t\omega, x, y, z)$ satisfies the random system

$$\begin{cases} dX = \frac{1}{\epsilon}g_1(X + \sigma\eta^\epsilon(\theta_t\omega), Y, Z)dt, \\ dY = g_2(X + \sigma\eta^\epsilon(\theta_t\omega), Y, Z)dt, \\ dZ = g_3(X + \sigma\eta^\epsilon(\theta_t\omega), Y, Z)dt. \end{cases} \quad (3)$$

The term $\sigma\eta^\epsilon(\theta_t\omega)$ do not change the Lipschitz constants of g_1 , g_2 and g_3 . So g_1 , g_2 and g_3 in random dynamical system (3) and in stochastic dynamical system (1) have the same Lipschitz constants. The random system (3) can be solved for any $\omega \in \Omega$. So, for any initial value $(X(0), Y(0), Z(0))^T = (X_0, Y_0, Z_0)^T$, the solution operator

$$\begin{aligned} (t, \omega, (X_0, Y_0, Z_0)^T) &\mapsto \Phi(t, \omega, (X_0, Y_0, Z_0)^T) \\ &= (X(t, \omega, (X_0, Y_0, Z_0)^T), Y(t, \omega, (X_0, Y_0, Z_0)^T), Z(t, \omega, (X_0, Y_0, Z_0)^T))^T, \end{aligned}$$

represents the random dynamical system for (3). Furthermore,

$$\phi(t, \omega, (X_0, Y_0, Z_0)^T) = \Phi(t, \omega, (X_0, Y_0, Z_0)^T) + (\sigma\eta^\epsilon(\theta_t\omega), 0, 0)^T,$$

characterizes the random dynamical system generated by stochastic Koper system (1).

4. Random slow manifolds

Introduce the Banach spaces of functions for investigating the random system (3). For any $\beta \in \mathbb{R}$,

$$C_{\beta}^{\mathbb{R},-} := \left\{ \Phi : (-\infty, 0] \rightarrow \mathbb{R} \text{ is continuous and } \sup_{t \in (-\infty, 0]} |e^{-\beta t} \Phi(t)| < \infty \right\},$$

$$C_{\beta}^{\mathbb{R},+} := \left\{ \Phi : [0, \infty) \rightarrow \mathbb{R} \text{ is continuous and } \sup_{t \in [0, \infty)} |e^{-\beta t} \Phi(t)| < \infty \right\},$$

with norms

$$\|\Phi\|_{C_{\beta}^{\mathbb{R},-}} := \sup_{t \in (-\infty, 0]} |e^{-\beta t} \Phi(t)|, \quad \text{and} \quad \|\Phi\|_{C_{\beta}^{\mathbb{R},+}} := \sup_{t \in [0, \infty)} |e^{-\beta t} \Phi(t)|.$$

Let $C_{\beta}^{\mathbb{R}^3, \pm}$ be the product of spaces $C_{\beta}^{\mathbb{R}^3, \pm} := C_{\beta}^{\mathbb{R}, \pm} \times C_{\beta}^{\mathbb{R}, \pm} \times C_{\beta}^{\mathbb{R}, \pm}$, with norm

$$\|U\|_{C_{\beta}^{\mathbb{R}^3, \pm}} = \|X\|_{C_{\beta}^{\mathbb{R}, \pm}} + \|Y\|_{C_{\beta}^{\mathbb{R}, \pm}} + \|Z\|_{C_{\beta}^{\mathbb{R}, \pm}}, \quad U = (X, Y, Z)^T \in C_{\beta}^{\mathbb{R}^3, \pm}.$$

For $U(\cdot, \omega) = (X(\cdot, \omega), Y(\cdot, \omega), Z(\cdot, \omega))^T \in C_{\beta}^{\mathbb{R}^3, -}$, it is the solution of (3) with initial value $U_0 = (X_0, Y_0, Z_0)^T$ iff $U(t, \omega)$ satisfies

$$\begin{pmatrix} X(t) \\ Y(t) \\ Z(t) \end{pmatrix} = \begin{pmatrix} \frac{1}{\epsilon} \int_{-\infty}^t g_1(X(s) + \sigma \eta^{\epsilon}(\theta_s \omega), Y(s), Z(s)) ds \\ Y_0 + \int_0^t g_2(X(s) + \sigma \eta^{\epsilon}(\theta_s \omega), Y(s), Z(s)) ds \\ Z_0 + \int_0^t g_3(X(s) + \sigma \eta^{\epsilon}(\theta_s \omega), Y(s), Z(s)) ds \end{pmatrix}. \quad (4)$$

Lemma 3. *Suppose that*

$$U(t, \omega, U_0) = \left(X(t, \omega, (X_0, Y_0, Z_0)^T), Y(t, \omega, (X_0, Y_0, Z_0)^T), Z(t, \omega, (X_0, Y_0, Z_0)^T) \right)^T$$

is the solution of (4) with $t \leq 0$. Then $U(t, \omega, U_0)$ is the unique solution in $C_{\beta}^{\mathbb{R}^3, -}$, where $U_0 = (X_0, Y_0, Z_0)^T$ is the initial value.

Proof. By using Banach fixed point theorem, we prove that

$$U(t, \omega, U_0) = \left(X(t, \omega, (X_0, Y_0, Z_0)^T), Y(t, \omega, (X_0, Y_0, Z_0)^T), Z(t, \omega, (X_0, Y_0, Z_0)^T) \right)^T$$

is the unique solution of (4). For the prove of it, define three operators for $t \leq 0$:

$$\begin{aligned} \mathfrak{I}_1(U)[t] &= \frac{1}{\epsilon} \int_{-\infty}^t g_1(X(s) + \sigma \eta^{\epsilon}(\theta_s \omega), Y(s), Z(s)) ds, \\ \mathfrak{I}_2(U)[t] &= Y_0 + \int_0^t g_2(X(s) + \sigma \eta^{\epsilon}(\theta_s \omega), Y(s), Z(s)) ds, \\ \mathfrak{I}_3(U)[t] &= Z_0 + \int_0^t g_3(X(s) + \sigma \eta^{\epsilon}(\theta_s \omega), Y(s), Z(s)) ds. \end{aligned}$$

Then Lyapunov-Perron transform is defined to be

$$\mathfrak{I}(U) = \begin{pmatrix} \mathfrak{I}_1(U) \\ \mathfrak{I}_2(U) \\ \mathfrak{I}_3(U) \end{pmatrix} = (\mathfrak{I}_1(U), \mathfrak{I}_2(U), \mathfrak{I}_3(U))^T.$$

Now, it is needed to prove that \mathfrak{S} maps $C_{\beta}^{\mathbb{R}^3, -}$ into itself. Take $U = (X, Y, Z)^T \in C_{\beta}^{\mathbb{R}^3, -}$ satisfying:

$$\begin{aligned}
\|\mathfrak{S}_1(U)[t]\|_{C_{\beta}^{\mathbb{R}, -}} &= \left\| \frac{1}{\epsilon} \int_{-\infty}^t g_1(X(s) + \sigma\eta^{\epsilon}(\theta_s\omega), Y(s), Z(s)) ds \right\|_{C_{\beta}^{\mathbb{R}, -}} \\
&\leq \frac{1}{\epsilon} \sup_{t \in (-\infty, 0]} \left\{ e^{-\beta t} \int_{-\infty}^t |g_1(X(s) + \sigma\eta^{\epsilon}(\theta_s\omega), Y(s), Z(s))| ds \right\} \\
&\leq \frac{K}{\epsilon} \sup_{t \in (-\infty, 0]} \left\{ e^{-\beta t} \int_{-\infty}^t (|X(s)| + |Y(s)| + |Z(s)|) ds \right\} + C_1 \\
&\leq \frac{K}{\epsilon} \sup_{t \in (-\infty, 0]} \left\{ \int_{-\infty}^t e^{-\beta(t-s)} ds \right\} \|U\|_{C_{\beta}^{\mathbb{R}^3, -}} + C_1 \\
&= \frac{K}{-\epsilon\beta} \|U\|_{C_{\beta}^{\mathbb{R}^3, -}} + C_1.
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
\|\mathfrak{S}_2(U)[t]\|_{C_{\beta}^{\mathbb{R}, -}} &= \|Y_0 + \int_0^t g_2(X(s) + \sigma\eta^{\epsilon}(\theta_s\omega), Y(s), Z(s)) ds\|_{C_{\beta}^{\mathbb{R}, -}} \\
&\leq \sup_{t \in (-\infty, 0]} \left\{ e^{-\beta t} \int_0^t |g_2(X(s) + \sigma\eta^{\epsilon}(\theta_s\omega), Y(s), Z(s))| ds \right\} + \|Y_0\|_{C_{\beta}^{\mathbb{R}, -}} \\
&\leq \sup_{t \in (-\infty, 0]} \left\{ e^{-\beta t} \int_0^t (|X(s)| + |Y(s)| + |Z(s)|) ds \right\} + C_2 \\
&\leq K \sup_{t \in (-\infty, 0]} \left\{ \int_0^t e^{-\beta(t-s)} ds \right\} \|U\|_{C_{\beta}^{\mathbb{R}^3, -}} + C_2 \\
&= \frac{K}{\beta} \|U\|_{C_{\beta}^{\mathbb{R}^3, -}} + C_2.
\end{aligned}$$

Furthermore,

$$\begin{aligned}
\|\mathfrak{S}_3(U)[t]\|_{C_{\beta}^{\mathbb{R}, -}} &= \|Z_0 + \int_0^t g_3(X(s) + \sigma\eta^{\epsilon}(\theta_s\omega), Y(s), Z(s)) ds\|_{C_{\beta}^{\mathbb{R}, -}} \\
&\leq \sup_{t \in (-\infty, 0]} \left\{ e^{-\beta t} \int_0^t |g_3(X(s) + \sigma\eta^{\epsilon}(\theta_s\omega), Y(s), Z(s))| ds \right\} + \|Z_0\|_{C_{\beta}^{\mathbb{R}, -}} \\
&\leq \sup_{t \in (-\infty, 0]} \left\{ e^{-\beta t} \int_0^t (|X(s)| + |Y(s)| + |Z(s)|) ds \right\} + C_3 \\
&\leq K \sup_{t \in (-\infty, 0]} \left\{ \int_0^t e^{-\beta(t-s)} ds \right\} \|U\|_{C_{\beta}^{\mathbb{R}^3, -}} + C_3 \\
&= \frac{K}{\beta} \|U\|_{C_{\beta}^{\mathbb{R}^3, -}} + C_3.
\end{aligned}$$

By using Lyapunov-Perron transform, the estimate of \mathfrak{S} in combined form is

$$\|\mathfrak{S}(U)\|_{C_{\beta}^{\mathbb{R}^3, -}} \leq \varrho(\beta, K, \epsilon) \|U\|_{C_{\beta}^{\mathbb{R}^3, -}} + C,$$

where

$$\varrho(\beta, K, \epsilon) = \frac{-K}{\epsilon\beta} + \frac{K}{\beta} + \frac{K}{\beta}.$$

Hence $\mathfrak{J}(U)$ is in $C_{\beta}^{\mathbb{R}^3, -}$ for all $U \in C_{\beta}^{\mathbb{R}^3, -}$, which means that \mathfrak{J} maps $C_{\beta}^{\mathbb{R}^3, -}$ into itself.

Now we should show that the map \mathfrak{J} is contractive. For this, take $U = (X, Y, Z)^T$, $\tilde{U} = (\tilde{X}, \tilde{Y}, \tilde{Z})^T \in C_{\beta}^{\mathbb{R}^3, -}$,

$$\begin{aligned} & \|\mathfrak{J}_1(U) - \mathfrak{J}_1(\tilde{U})\|_{C_{\beta}^{\mathbb{R}, -}} \\ & \leq \frac{1}{\epsilon} \sup_{t \in (-\infty, 0]} \left\{ e^{-\beta t} \int_{-\infty}^t \left| g_1(X(s) + \sigma \eta^{\epsilon}(\theta_s, \omega), Y(s), Z(s)) - g_1(\tilde{X}(s) + \sigma \eta^{\epsilon}(\theta_s, \omega), \tilde{Y}(s), \tilde{Z}(s)) \right| ds \right\} \\ & \leq \frac{K}{\epsilon} \sup_{t \in (-\infty, 0]} \left\{ e^{-\beta t} \int_{-\infty}^t (|X(s) - \tilde{X}(s)| + |Y(s) - \tilde{Y}(s)| + |Z(s) - \tilde{Z}(s)|) ds \right\} \\ & \leq \frac{K}{\epsilon} \sup_{t \in (-\infty, 0]} \left\{ \int_{-\infty}^t e^{-\beta(t-s)} ds \right\} \|U - \tilde{U}\|_{C_{\beta}^{\mathbb{R}^3, -}} \\ & = \frac{K}{-\epsilon \beta} \|U - \tilde{U}\|_{C_{\beta}^{\mathbb{R}^3, -}}. \end{aligned}$$

Using the same way,

$$\begin{aligned} \|\mathfrak{J}_2(U) - \mathfrak{J}_2(\tilde{U})\|_{C_{\beta}^{\mathbb{R}, -}} & \leq K \sup_{t \in (-\infty, 0]} \left\{ \int_t^0 e^{-\beta(t-s)} ds \right\} \|U - \tilde{U}\|_{C_{\beta}^{\mathbb{R}^3, -}} \\ & \leq \frac{K}{\beta} \|U - \tilde{U}\|_{C_{\beta}^{\mathbb{R}^3, -}}. \end{aligned}$$

Moreover,

$$\begin{aligned} \|\mathfrak{J}_3(U) - \mathfrak{J}_3(\tilde{U})\|_{C_{\beta}^{\mathbb{R}, -}} & \leq K \sup_{t \in (-\infty, 0]} \left\{ \int_t^0 e^{-\beta(t-s)} ds \right\} \|U - \tilde{U}\|_{C_{\beta}^{\mathbb{R}^3, -}} \\ & \leq \frac{K}{\beta} \|U - \tilde{U}\|_{C_{\beta}^{\mathbb{R}^3, -}}. \end{aligned}$$

Combing the three together,

$$\|\mathfrak{J}(U) - \mathfrak{J}(\tilde{U})\|_{C_{\beta}^{\mathbb{R}^3, -}} \leq \varrho(\beta, K, \epsilon) \|U - \tilde{U}\|_{C_{\beta}^{\mathbb{R}^3, -}},$$

where

$$\varrho(\beta, K, \epsilon) = \frac{K}{-\epsilon \beta} + \frac{K}{\beta} + \frac{K}{\beta}.$$

By setting $\beta = -\frac{\gamma}{\epsilon}$,

$$\varrho(\beta, K, \epsilon) \rightarrow \frac{K}{\gamma} \text{ for } \epsilon \rightarrow 0.$$

So, there exists a sufficiently small $\epsilon_0 \rightarrow 0$ with property

$$0 < \varrho(\beta, K, \epsilon) < 1, \text{ for } \epsilon \in (0, \epsilon_0).$$

Hence, the map \mathfrak{J} in $C_{-\frac{\gamma}{\epsilon}}^{\mathbb{R}^3, -}$ is contractive. By Banach fixed point theorem, every contractive mapping in Banach space has a unique fixed point. Thus, (4) has the unique solution

$$U(t, \omega, U_0) = (X(t, \omega, (X_0, Y_0, Z_0)^T), Y(t, \omega, (X_0, Y_0, Z_0)^T), Z(t, \omega, (X_0, Y_0, Z_0)^T))^T \text{ in } C_{-\frac{\gamma}{\epsilon}}^{\mathbb{R}^3, -}.$$

□

From Lemma 3 we obtain the following remark.

Remark 2. For any $(X_0, Y_0, Z_0)^T, (X'_0, Y'_0, Z'_0)^T \in \mathbb{R}^3$, and for all $\omega \in \Omega, Y_0, Y'_0, Z_0, Z'_0 \in \mathbb{R}$, there is an $\epsilon_0 > 0$ such that

$$\|U(t, \omega, (X_0, Y_0, Z_0)^T) - U(t, \omega, (X'_0, Y'_0, Z'_0)^T)\|_{C_{-\frac{\gamma}{\epsilon}}^{\mathbb{R}^3, -}} \leq \frac{1}{1 - \varrho(\beta, K, \epsilon)} (|Y_0 - Y'_0| + |Z_0 - Z'_0|). \quad (5)$$

Proof. Instead of writing $U(t, \omega, (X_0, Y_0, Z_0)^T)$ and $U(t, \omega, (X'_0, Y'_0, Z'_0)^T)$, we write $U(t, \omega, Y_0, Z_0)$ and $U(t, \omega, Y'_0, Z'_0)$. For all $\omega \in \Omega$ and $Y_0, Y'_0, Z_0, Z'_0 \in \mathbb{R}$, we determine an upper bound

$$\begin{aligned} & \|U(t, \omega, Y_0, Z_0) - U(t, \omega, Y'_0, Z'_0)\|_{C_{-\frac{\gamma}{\epsilon}}^{\mathbb{R}^3, -}} \\ &= \|X(t, \omega, Y_0, Z_0) - X(t, \omega, Y'_0, Z'_0)\|_{C_{-\frac{\gamma}{\epsilon}}^{\mathbb{R}, -}} + \|Y(t, \omega, Y_0, Z_0) - Y(t, \omega, Y'_0, Z'_0)\|_{C_{-\frac{\gamma}{\epsilon}}^{\mathbb{R}, -}} + \|Z(t, \omega, Y_0, Z_0) - Z(t, \omega, Y'_0, Z'_0)\|_{C_{-\frac{\gamma}{\epsilon}}^{\mathbb{R}, -}} \\ &\leq \left(\frac{K}{-\epsilon\beta} + \frac{2K}{\beta}\right) \times \|U(t, \omega, Y_0, Z_0) - U(t, \omega, Y'_0, Z'_0)\|_{C_{-\frac{\gamma}{\epsilon}}^{\mathbb{R}^3, -}} + |Y_0 - Y'_0| + |Z_0 - Z'_0| \\ &= \varrho(\beta, K, \epsilon) \|U(t, \omega, Y_0, Z_0) - U(t, \omega, Y'_0, Z'_0)\|_{C_{-\frac{\gamma}{\epsilon}}^{\mathbb{R}^3, -}} + |Y_0 - Y'_0| + |Z_0 - Z'_0|. \end{aligned}$$

Therefore (5) is valid. \square

Next, with the help of Lyapunov-Perron method we will construct the slow manifold as a random graph.

Theorem 1. Assume that the hypotheses **(H1)**-**(H3)** hold. Then for sufficiently small $\epsilon > 0$, random system (3) posses a Lipschitz random slow manifold:

$$\mathcal{M}^\epsilon(\omega) = \{(l^\epsilon(\omega, Y_0, Z_0), Y_0, Z_0)^T : Y_0, Z_0 \in \mathbb{R}\},$$

where

$$l^\epsilon(\cdot, \cdot) : \Omega \times \mathbb{R}^2 \rightarrow \mathbb{R},$$

is a Lipschitz graph map with Lipschitz constant

$$\text{Lip} l^\epsilon(\omega, \cdot) \leq \frac{K}{\gamma - K(1 - 2\epsilon)}.$$

Proof. For any $Y_0, Z_0 \in \mathbb{R}$ we define Lyapunov-Perron map l^ϵ by

$$l^\epsilon(\omega, Y_0, Z_0) = \frac{1}{\epsilon} \int_{-\infty}^0 g_1(X(s, \omega, Y_0, Z_0) + \sigma \eta^\epsilon(\theta_s \omega), Y(s, \omega, Y_0, Z_0), Z(s, \omega, Y_0, Z_0)) ds.$$

Then by (5), we obtain

$$|l^\epsilon(\omega, Y_0, Z_0) - l^\epsilon(\omega, Y'_0, Z'_0)| \leq \frac{K}{-\epsilon\beta [1 - \varrho(\beta, K, \epsilon)]} (|Y_0 - Y'_0| + |Z_0 - Z'_0|),$$

for all $Y_0, Y'_0, Z_0, Z'_0 \in \mathbb{R}$ and $\omega \in \Omega$. So

$$|l^\epsilon(\omega, Y_0, Z_0) - l^\epsilon(\omega, Y'_0, Z'_0)| \leq \frac{K}{\gamma [1 - \varrho(\beta, K, \epsilon)]} (|Y_0 - Y'_0| + |Z_0 - Z'_0|),$$

for every $Y_0, Y'_0, Z_0, Z'_0 \in \mathbb{R}$ and $\omega \in \Omega$. Utilizing Theorem III.9 in Casting and Valadier [6, p.67], $\mathcal{M}^\epsilon(\omega)$ is a random set, i.e., for any $U = (X, Y, Z)^T \in \mathbb{R}^3$,

$$\omega \mapsto \inf_{U' \in \mathbb{R}^3} |U - (I^\epsilon(\omega, \mathfrak{I}U'), \mathfrak{I}U')^T|, \quad (6)$$

is measurable. The space \mathbb{R}^3 has a countable dense subset \mathbb{Q}^3 . Then the infimum in (6) is equivalent to

$$\inf_{U' \in \mathbb{Q}^3} |U - (I^\epsilon(\omega, \mathfrak{I}U'), \mathfrak{I}U')^T|.$$

Under the infimum in (6) the measurability of any expression can be determined, since for all U' in \mathbb{R}^3 the map $\omega \mapsto I^\epsilon(\omega, \mathfrak{I}U')$ is measurable. The slow flow is found as a random graph of I^ϵ .

Finally we need to show that $\mathcal{M}^\epsilon(\omega)$ is positively invariant, i.e., for all $U_0 = (X_0, Y_0, Z_0)^T$ in $\mathcal{M}^\epsilon(\omega)$, $U(s, \omega, U_0)$ is in $\mathcal{M}^\epsilon(\theta_s \omega)$ for every $s \geq 0$. Note that $U(t + s, \omega, U_0)$ is a solution of

$$\begin{aligned} dX &= \frac{1}{\epsilon} g_1(X + \sigma \eta^\epsilon(\theta_t \omega), Y, Z) dt, \\ dY &= g_2(X + \sigma \eta^\epsilon(\theta_t \omega), Y, Z) dt, \\ dZ &= g_3(X + \sigma \eta^\epsilon(\theta_t \omega), Y, Z) dt, \end{aligned}$$

with initial value $U(s) = (X(s), Y(s), Z(s))^T = U(s, \omega, U_0)$. So, $U(t + s, \omega, U_0) = U(t, \theta_s \omega, U(s, \omega, U_0))$. Since $U(\cdot, \omega, U_0)$ is in $C_{-\frac{\gamma}{\epsilon}}^{\mathbb{R}^3, -}$, we gain $U(\cdot, \theta_s \omega, U(s, \omega, U_0)) \in C_{-\frac{\gamma}{\epsilon}}^{\mathbb{R}^3, -}$. Hence, $U(s, \omega, U_0) \in \mathcal{M}^\epsilon(\theta_s \omega)$. \square

5. Example

Namely, we consider the stochastic system

$$\begin{cases} \dot{x} = \frac{1}{\epsilon}(-x^3 + 3x - 10y + 5z + 3) + \frac{\sigma}{\sqrt{\epsilon}} \dot{L}_t^\alpha, & \text{in } \mathbb{R}, \\ \dot{y} = x - 2y + z, & \text{in } \mathbb{R}, \\ \dot{z} = y - z, & \text{in } \mathbb{R}, \end{cases} \quad (7)$$

where x is the ‘‘fast’’ component, (y, z) is the ‘‘slow’’ component, $k = -10$ and $\lambda(z) = -5z - 3$.




Figure 1: The evolution of system (7) for the choice of initial conditions $x(0) = y(0) = z(0) = 0$ at the fixed noise intensity $\sigma = 0.5$ with a scaling parameter $\epsilon = 0.05$ and a gradual increase in the stability index: (a) $\alpha = 0.8$; (b) $\alpha = 1.6$; (c) $\alpha = 1.9$.

If we fix $\sigma = 0.5$ and increase α in system (7) for the choice of initial conditions $x(0) = y(0) = z(0) = 0$ with a scaling parameter $\epsilon = 0.05$, we observe the following typical sequence of events in Figure 1.

The stochastic nonlinear dynamics with the stability index $\alpha = 0.8$ display the complex spatio-temporal oscillations accompanied by few big jumps as depicted in Fig. 1(a). With respect to $\alpha = 1.6$, the external Lévy noise leads to dramatically different dynamical behavior in the red trajectory, which is confirmed numerically in Fig. 1(b). As the stability index α is increased toward 1.9, the path of stochastic system (7) shows low frequency oscillations excited by Lévy noise as illustrated in Fig. 1(c). For three different values of α , stochastic noise of the fast variable x can significantly influence the dynamics of the whole Koper model. However, the shapes of different trajectories from the viewpoint of the slow surface (y, z) are consistent.




Figure 2: The evolution of system (7) starting from $x(0) = y(0) = z(0) = 0$ with a fixed stability index $\alpha = 1$ and a scaling parameter $\epsilon = 0.05$ as the noise intensity increases: (a) $\sigma = 0.1$; (b) $\sigma = 0.5$; (c) $\sigma = 0.8$.

Now we fix $\alpha = 1$ and consider a variation of σ in system (7) starting from $x(0) = y(0) = z(0) = 0$ with a scaling parameter $\epsilon = 0.05$; see Fig. 2. For small enough $\sigma = 0.1$, the path has spiral pattern with small-amplitude fluctuations, which is clearly demonstrated in Fig. 2(a). When σ -value just arrives at 0.5, the status of the path changes happens abruptly for stochastic system (7) as plotted in Fig. 2(b). As the noise intensity σ increases further to 0.8, large variations in the dynamics of the trajectory are indicated in Fig. 2(c) because of strong external noise. Based on the effects of increasing noise intensities, the curves become more and more sophisticated.

If we scale the time $t \rightarrow \epsilon t$ and use the self-similarity $\epsilon^{-1/\alpha} L_{\epsilon t}^\alpha \stackrel{d}{=} L_t^\alpha$, then stochastic system (7) in the sense of distribution is equal to

$$\begin{cases} dx = (-x^3 + 3x - 10y + 5z + 3)dt + \sigma dL_t^\alpha, & \text{in } \mathbb{R}, \\ dy = \epsilon(x - 2y + z)dt, & \text{in } \mathbb{R}, \\ dz = \epsilon(y - z)dt, & \text{in } \mathbb{R}. \end{cases}$$

When $\sigma = 0$, the deterministic system

$$\begin{cases} \dot{x} = (-x^3 + 3x - 10y + 5z + 3), & \text{in } \mathbb{R}, \\ \dot{y} = \epsilon(x - 2y + z), & \text{in } \mathbb{R}, \\ \dot{z} = \epsilon(y - z), & \text{in } \mathbb{R}. \end{cases} \quad (8)$$

has a unique fixed point $P = (1, 1, 1)$. Linearize by finding the Jacobian matrix. Hence

$$J = \begin{pmatrix} 3(1 - x^2) & -10 & 5 \\ \epsilon & -2\epsilon & \epsilon \\ 0 & \epsilon & -\epsilon \end{pmatrix}.$$

Therefore,

$$J_P = \begin{pmatrix} 0 & -10 & 5 \\ \epsilon & -2\epsilon & \epsilon \\ 0 & \epsilon & -\epsilon \end{pmatrix}.$$

The stability of the equilibrium point P is determined by the associated Jacobian matrix J_P . The trace $\text{tr}(J_P) = -3\epsilon$ is negative, and the determinant $\det(J_P) = -5\epsilon^2$ is also negative. It follows that three eigenvalues are negative. Thus, P is a stable fixed point.




Figure 3: When $x(0) = y(0) = z(0) = 0$ and $\epsilon = 0.05$: (a) one trajectory of system (8); (b) dynamical behaviors for the three variables $x(t)$, $y(t)$ and $z(t)$ in system (8).

Observe that in the beginning the blue curve of system (8) with $\epsilon = 0.05$ departs from $x(0) = y(0) = z(0) = 0$ along the direction of x -axis, shown in Figure 3(a). But some time later, there is a gentle growing tendency for the trajectory at $x(t) = 2$ to move upward. With the increase of time, the path encounters a crucial turning point and has a precipitous climb to the stable fixed point P . More significantly, it is parallel to the (y, z) -plane of the slow variables in this situation. By comparison with Fig. 1-2, the Koper system (7) exhibits sensitivity to stochastic disturbance.

We sketch three time series data for the system (8) when $x(0) = y(0) = z(0) = 0$ and $\epsilon = 0.05$, as clearly detailed in Figure 3(b). It is manifest that the fast variable $x(t)$ jumps to 2 instantaneously at first, and then decreases to 1 rapidly and suddenly. The green curve about $x(t)$ stays at the same level after $t = 15$. From the time series drawings of the slow variables $y(t)$ and $z(t)$, they both grow to 1 eventually. The red path about $y(t)$ increases much quicker than the purple trajectory about $z(t)$.

6. Conclusions and future challenges

We investigated three-dimensional stochastic slow-fast Koper system (1) driven by α -stable Lévy noise. We constructed the slow manifold in which the fast variable x can be expressed as the random function of the two slow variables y and z . Stochastic nonlinear dimensionality reduction helped us make more accurate

predictions by using stochastic differential equations for the slow variables. We carried out the computation of the practical Koper models (7) and (8).

When the scaling parameter $\hat{\varepsilon}$ is sufficiently small, i.e., $0 < \hat{\varepsilon} \ll 1$, stochastic Koper model (1) has three times scales: variable x is fast, y is slow, and z is slower. It indicates the ratio of three times scales such that $|\frac{dx}{dt}| \gg |\frac{dy}{dt}| \gg |\frac{dz}{dt}|$. So it is meaningful to project high-dimensional dynamics onto lower-dimensional effective manifolds in this case.

What happens if the main bifurcation parameters k and $\lambda(z) =: \lambda$ change? The folded node/focus and supercritical Hopf bifurcation are expected to occur in parameter space for the Koper model without noise. By classifying the type and stability of the equilibrium points, we obtain a bifurcation diagram under parameter variation.

We can generalize the consideration to a number of cases where the variables x , y and z both are perturbed by α -stable Lévy noises, even to the extent that the influences are the more general Lévy processes including multiplicative and additive effects. We may use Itô, Stratonovich or Marcus type stochastic differential equations.

The orbits can escape from the region of the metastable equilibrium if we face random perturbation [18]. It is interesting to characterize the most probable transition from one metastable state to another [15, 25]. The random slow manifold still depends on k and λ . It is a challenge to plot the bifurcation diagram in the (k, λ) -plane of the random slow flow. Luckily, we can explore stochastic bifurcations in dynamical systems driven by Lévy noises with support for statistical modeling and computation [26, 31, 34].

We often have difficulty comprehending data in phase space with the attracting or repelling random slow manifolds. Thus it is useful for visualization purposes by reducing data to a small number of dimensions.

DATA AVAILABILITY

Numerical algorithms source code that support the findings of this study are openly available in GitHub, Ref. [30].

References

References

- [1] Applebaum, D. *Lévy processes and stochastic calculus*; Cambridge university press, 2009.
- [2] Arnold, L. *Random dynamical systems*; Springer Science & Business Media, 2013.
- [3] Balabaev, M. Invariant manifold of variable stability in the Koper model. *J. Phys. Conf. Ser.* **2019**, 1368, 042003.
- [4] Bates, P.W.; Lu, K.; Zeng, C. *Existence and persistence of invariant manifolds for semiflows in Banach space*; American Mathematical Society, 1998.
- [5] Caraballo, T.; Chueshov, I.; Langa, J.A. Existence of invariant manifolds for coupled parabolic and hyperbolic stochastic partial differential equations. *Nonlinearity* **2005**, 18, 747-767.
- [6] Castaing, C.; Valadier, M. *Convex analysis and measurable multifunctions*; Springer, 2006.
- [7] Chao, Y.; Wei, P.; Yuan, S. Invariant foliations for stochastic dynamical systems with multiplicative stable Lévy noise. *Electron. J. Differ. Equ.* **2019**, 2019, 1-21.
- [8] Chicone, C.; Latushkin, Y. Center manifolds for infinite dimensional nonautonomous differential equations. *J. Differ. Equ.* **1997**, 141, 356-399.
- [9] Chow, S.N.; Lu, K. Invariant manifolds for flows in Banach spaces. *J. Differ. Equ.* **1988**, 74, 285-317.
- [10] Chow, S.N.; Lu, K.; Lin, X.B. Smooth foliations for flows in Banach space. *J. Differ. Equ.* **1991**, 94, 266-291.
- [11] Duan, J. *An introduction to stochastic dynamics*; Cambridge University Press, 2015.
- [12] Duan, J.; Lu, K.; Schmalfuss, B. Smooth stable and unstable manifolds for stochastic evolutionary equations. *J. Dyn. Differ. Equ.* **2004**, 16, 949-972.
- [13] Fu, H.; Liu, X.; Duan, J. Slow manifolds for multi-time-scale stochastic evolutionary systems. *Comm. Math. Sci.* **2013**, 11, 141-162.
- [14] Henry, D. *Geometric theory of semilinear parabolic equations*; Springer, 2006.

- [15] Huang, Y.; Chao, Y.; Yuan, S.; Duan, J. Characterization of the most probable transition paths of stochastic dynamical systems with stable Lévy noise. *J. Math. Phys.* **2019**, 2019, 063204.
- [16] Jacod, J.; Shiryaev, A. *Limit theorems for stochastic processes*; Springer Science & Business Media, 2013.
- [17] Koper, M.T.M. Bifurcations of mixed-mode oscillations in a three-variable autonomous Van der Pol-Duffing model with a cross-shaped phase diagram. *Physica D* **1995**, 80, 72-94.
- [18] Li, Y.; Yuan, S.; Xu, S. Controlling mean exit time of stochastic dynamical systems based on quasipotential and machine learning. *arXiv preprint* **2022**, arXiv:2209.13098.
- [19] Protter, P. *Stochastic Integration and Differential Equations*; Springer-Verlag, 2004.
- [20] Qiao, M.; Yuan, S. Analysis of a stochastic predator-prey model with prey subject to disease and Lévy noise. *Stoch. Dynam.* **2019**, 19, 1950038.
- [21] Ren, J.; Duan, J.; Jones, C.K.R.T. Approximation of random slow manifolds and settling of inertial particles under uncertainty. *J. Dyn. Differ. Equ.* **2015**, 27, 961-979.
- [22] Ren, J.; Duan, J.; Wang, X. A parameter estimation method based on random slow manifolds. *Appl. Math. Model.* **2015**, 39, 3721-3732.
- [23] Ruelle, D. Characteristic exponents and invariant manifolds in Hilbert space. *Ann. Math.* **1982**, 115, 243-290.
- [24] Schmalfuss, B.; Schneider, K.R. Invariant manifolds for random dynamical systems with slow and fast variables. *J. Dyn. Differ. Equ.* **2008**, 20, 133-164.
- [25] Tesfay, A.; Yuan, S.; Tesfay, D.; Brannan, J. Most Probable Dynamics of the Single-Species with Allee Effect under Jump-diffusion Noise. *arXiv preprint* **2021**, arXiv:2112.07234.
- [26] Tesfay, A.; Tesfay, D.; Yuan, S.; Brannan, J.; Duan, J. Stochastic bifurcation in single-species model induced by α -stable Lévy noise. *J. Stat. Mech.* **2021**, 2021, 103403.
- [27] Wang, W.; Roberts, A. Slow manifold and averaging for slow-fast stochastic differential system. *J. Math. Anal. Appl.* **2013**, 398, 822-839.
- [28] Wei, H.H. Weak convergence of probability measures on metric spaces of nonlinear operators. *Bull. Inst. Math. Acad. Sin.* **2016**, 11, 485-519.
- [29] Yang, X.; Xu, Y.; Wang, R.; Jiao, Z. The central limit theorem for slow-fast systems with Lévy noise. *Appl. Math. Lett.* **2022**, 128, 107897.
- [30] Yuan, S. Code. *Github* **2022**. <https://github.com/ShenglanYuan/Slow-manifolds-for-stochastic-Koper-models-with-stable-Lvy-noises>
- [31] Yuan, S.; Blömker, D. Modulation and amplitude equations on bounded domains for nonlinear SPDEs driven by cylindrical α -stable Lévy processes. *SIAM J. Appl. Dyn. Syst.* **2022**, 21, 1748-1777.
- [32] Yuan, S.; Blömker, D.; Duan, J. Stochastic turbulence for Burgers equation driven by cylindrical Lévy process. *Stoch. Dynam.* **2022**, 22, 2240004.
- [33] Yuan, S.; Li, Y.; Zeng, Z. Stochastic bifurcations and tipping phenomena of insect outbreak systems driven by α -stable Lévy processes. *Math. Model. Nat. Phenom.* **2022**, 17, 34.
- [34] Yuan, S.; Zeng, Z.; Duan, J. Stochastic bifurcation for two-time-scale dynamical system with α -stable Lévy noise. *J. Stat. Mech.* **2021**, 2021, 033204.
- [35] Yuan, S.; Hu, J.; Liu, X.; Duan, J. Slow manifolds for dynamical systems with non-Gaussian stable Lévy noise. *Anal. Appl.* **2019**, 17, 477-511.
- [36] Zulfıqar, H.; He, Z.; Yang, M.; Duan, J. Slow manifold and parameter estimation for a nonlocal fast-slow dynamical system with brownian motion. *Acta Math. Sci.* **2021**, 41, 1057-1080.
- [37] Zulfıqar, H.; Yuan, S.; He, Z.; Duan, J. Slow manifolds for a nonlocal fast-slow stochastic system with stable Lévy noise. *J. Math. Phys.* **2019**, 60, 091501.