

---

**FUNCTIONAL DONOHO-STARK-ELAD-BRUCKSTEIN-RICAUD-TORRÉSANI  
UNCERTAINTY PRINCIPLE**

---

**K. MAHESH KRISHNA**

Post Doctoral Fellow

Statistics and Mathematics Unit

Indian Statistical Institute, Bangalore Centre

Karnataka 560 059, India

Email: kmaheshak@gmail.com

Date: April 10, 2023

---

**Abstract:** Let  $(\{f_j\}_{j=1}^n, \{\tau_j\}_{j=1}^n)$  and  $(\{g_k\}_{k=1}^m, \{\omega_k\}_{k=1}^m)$  be p-Schauder frames for a finite dimensional Banach space  $\mathcal{X}$ . Then for every  $x \in \mathcal{X} \setminus \{0\}$ , we show that

$$(1) \quad \|\theta_f x\|_0^{\frac{1}{p}} \|\theta_g x\|_0^{\frac{1}{q}} \geq \frac{1}{\max_{1 \leq j \leq n, 1 \leq k \leq m} |f_j(\omega_k)|} \quad \text{and} \quad \|\theta_g x\|_0^{\frac{1}{p}} \|\theta_f x\|_0^{\frac{1}{q}} \geq \frac{1}{\max_{1 \leq j \leq n, 1 \leq k \leq m} |g_k(\tau_j)|}.$$

where

$$\theta_f : \mathcal{X} \ni x \mapsto (f_j(x))_{j=1}^n \in \ell^p([n]); \quad \theta_g : \mathcal{X} \ni x \mapsto (g_k(x))_{k=1}^m \in \ell^p([m])$$

and  $q$  is the conjugate index of  $p$ . We call Inequality (1) as **Functional Donoho-Stark-Elad-Bruckstein-Ricaud-Torresani Uncertainty Principle**. Inequality (1) improves Ricaud-Torresani uncertainty principle [*IEEE Trans. Inform. Theory, 2013*]. In particular, it improves Elad-Bruckstein uncertainty principle [*IEEE Trans. Inform. Theory, 2002*] and Donoho-Stark uncertainty principle [*SIAM J. Appl. Math., 1989*].

**Keywords:** Uncertainty Principle, Orthonormal Basis, Parseval Frame, Hilbert space, Banach space.

**Mathematics Subject Classification (2020):** 42C15.

---

## 1. INTRODUCTION

While Heisenberg's uncertainty principle [3] (English translation of original 1927 paper) is one of the greatest inequalities in the first half of the 20 century, Donoho-stark uncertainty principle [1] is one of the greatest inequalities in the second half of the 20 century. For  $h \in \mathbb{C}^d$ , let  $\|h\|_0$  be the number of nonzero entries in  $h$ . Let  $\hat{\cdot} : \mathbb{C}^d \rightarrow \mathbb{C}^d$  be the Fourier transform.

**Theorem 1.1.** (*Donoho-Stark Uncertainty Principle*) [1] *For every  $d \in \mathbb{N}$ ,*

$$(2) \quad \left( \frac{\|h\|_0 + \|\hat{h}\|_0}{2} \right)^2 \geq \|h\|_0 \|\hat{h}\|_0 \geq d, \quad \forall h \in \mathbb{C}^d \setminus \{0\}.$$

In 2002, Elad and Bruckstein generalized Inequality (2) to pairs of orthonormal bases [2]. To state the result we need some notations. Given a collection  $\{\tau_j\}_{j=1}^n$  in a finite dimensional Hilbert space  $\mathcal{H}$  over  $\mathbb{K}$  ( $\mathbb{R}$  or  $\mathbb{C}$ ), we define

$$\theta_\tau : \mathcal{H} \ni h \mapsto \theta_\tau h := (\langle h, \tau_j \rangle)_{j=1}^n \in \mathbb{K}^n.$$

**Theorem 1.2. (Elad-Bruckstein Uncertainty Principle)** [2] Let  $\{\tau_j\}_{j=1}^n, \{\omega_j\}_{j=1}^n$  be two orthonormal bases for a finite dimensional Hilbert space  $\mathcal{H}$ . Then

$$\left( \frac{\|\theta_\tau h\|_0 + \|\theta_\omega h\|_0}{2} \right)^2 \geq \|\theta_\tau h\|_0 \|\theta_\omega h\|_0 \geq \frac{1}{\max_{1 \leq j, k \leq n} |\langle \tau_j, \omega_k \rangle|^2}, \quad \forall h \in \mathcal{H} \setminus \{0\}.$$

Note that Theorem 1.1 follows from Theorem 1.2. In 2013, Ricaud and Torrésani showed that orthonormal bases in Theorem 1.2 can be improved to Parseval frames [6].

**Theorem 1.3. (Ricaud-Torrésani Uncertainty Principle)** [6] Let  $\{\tau_j\}_{j=1}^n, \{\omega_j\}_{j=1}^n$  be two Parseval frames for a finite dimensional Hilbert space  $\mathcal{H}$ . Then

$$\left( \frac{\|\theta_\tau h\|_0 + \|\theta_\omega h\|_0}{2} \right)^2 \geq \|\theta_\tau h\|_0 \|\theta_\omega h\|_0 \geq \frac{1}{\max_{1 \leq j, k \leq n} |\langle \tau_j, \omega_k \rangle|^2}, \quad \forall h \in \mathcal{H} \setminus \{0\}.$$

In this paper, we derive uncertainty principle for finite dimensional Banach spaces which contains Theorem 1.3 as a particular case.

## 2. FUNCTIONAL DONOHO-STARK-ELAD-BRUCKSTEIN-RICAUD-TORRÉSANI UNCERTAINTY PRINCIPLE

In the paper,  $\mathbb{K}$  denotes  $\mathbb{C}$  or  $\mathbb{R}$  and  $\mathcal{X}$  denotes a finite dimensional Banach space over  $\mathbb{K}$ . Identity operator on  $\mathcal{X}$  is denoted by  $I_{\mathcal{X}}$ . Dual of  $\mathcal{X}$  is denoted by  $\mathcal{X}^*$ . Whenever  $1 < p < \infty$ ,  $q$  denotes conjugate index of  $p$ . For  $d \in \mathbb{N}$ , standard finite dimensional Banach space  $\mathbb{K}^d$  over  $\mathbb{K}$  equipped with standard  $\|\cdot\|_p$  norm is denoted by  $\ell^p([d])$ . Canonical basis for  $\mathbb{K}^d$  is denoted by  $\{\delta_j\}_{j=1}^d$  and  $\{\zeta_j\}_{j=1}^d$  be the coordinate functionals associated with  $\{\delta_j\}_{j=1}^d$ . We need the following variant of p-approximate Schauder frames defined by Krishna and Johnson in [5].

**Definition 2.1.** Let  $\mathcal{X}$  be a finite dimensional Banach space over  $\mathbb{K}$ . Let  $\{\tau_j\}_{j=1}^n$  be a collection in  $\mathcal{X}$  and  $\{f_j\}_{j=1}^n$  be a sequence in  $\mathcal{X}^*$ . The pair  $(\{f_j\}_{j=1}^n, \{\tau_j\}_{j=1}^n)$  is said to be a **p-Schauder frame** ( $1 < p < \infty$ ) for  $\mathcal{X}$  if the following conditions hold.

(i) For every  $x \in \mathcal{X}$ ,

$$\|x\|^p = \sum_{j=1}^n |f_j(x)|^p.$$

(ii) For every  $x \in \mathcal{X}$ ,

$$x = \sum_{j=1}^n f_j(x) \tau_j.$$

We easily see that condition (i) in Definition 2.1 says that the map

$$\theta_f : \mathcal{X} \ni x \mapsto (f_j(x))_{j=1}^n \in \ell^p([n])$$

is a linear isometry. Like Holub's characterization of frames for Hilbert spaces [4], following theorem characterizes p-Schauder frames.

**Theorem 2.2.** A pair  $(\{f_j\}_{j=1}^n, \{\tau_j\}_{j=1}^n)$  is a p-Schauder frame for  $\mathcal{X}$  if and only if

$$f_j = \zeta_j U, \quad \tau_j = V \delta_j, \quad \forall j = 1, \dots, n,$$

where  $U : \mathcal{X} \rightarrow \ell^p([n])$ ,  $V : \ell^p([n]) \rightarrow \mathcal{X}$  are linear operators such that  $VU = I_{\mathcal{X}}$  and  $U$  is an isometry.

Following is the crucial result of this paper.

**FUNCTIONAL DONOHO-STARK-ELAD-BRUCKSTEIN-RICAUD-TORRÉSANI  
UNCERTAINTY PRINCIPLE**

**Theorem 2.3.** (*Functional Donoho-Stark-Elad-Bruckstein-Ricaud-Torresani Uncertainty Principle*) Let  $(\{f_j\}_{j=1}^n, \{\tau_j\}_{j=1}^n)$  and  $(\{g_k\}_{k=1}^m, \{\omega_k\}_{k=1}^m)$  be  $p$ -Schauder frames for a Banach space  $\mathcal{X}$ . Then

for every  $x \in \mathcal{X} \setminus \{0\}$ , we have

$$(3) \quad \|\theta_f x\|_0^{\frac{1}{p}} \|\theta_g x\|_0^{\frac{1}{q}} \geq \frac{1}{\max_{1 \leq j \leq n, 1 \leq k \leq m} |f_j(\omega_k)|} \quad \text{and} \quad \|\theta_g x\|_0^{\frac{1}{p}} \|\theta_f x\|_0^{\frac{1}{q}} \geq \frac{1}{\max_{1 \leq j \leq n, 1 \leq k \leq m} |g_k(\tau_j)|}.$$

*Proof.* Let  $x \in \mathcal{X} \setminus \{0\}$  and  $q$  be the conjugate index of  $p$ . First using  $\theta_f$  is an isometry and later using  $\theta_g$  is an isometry, we get

$$\begin{aligned} \|x\|^p &= \|\theta_f x\|^p = \sum_{j=1}^n |f_j(x)|^p = \sum_{j \in \text{supp}(\theta_f x)} |f_j(x)|^p \\ &= \sum_{j \in \text{supp}(\theta_f x)} \left| f_j \left( \sum_{k=1}^m g_k(x) \omega_k \right) \right|^p = \sum_{j \in \text{supp}(\theta_f x)} \left| \sum_{k=1}^m g_k(x) f_j(\omega_k) \right|^p \\ &= \sum_{j \in \text{supp}(\theta_f x)} \left| \sum_{k \in \text{supp}(\theta_g x)} g_k(x) f_j(\omega_k) \right|^p \leq \sum_{j \in \text{supp}(\theta_f x)} \left( \sum_{k \in \text{supp}(\theta_g x)} |g_k(x) f_j(\omega_k)| \right)^p \\ &\leq \left( \max_{1 \leq j \leq n, 1 \leq k \leq m} |f_j(\omega_k)| \right)^p \sum_{j \in \text{supp}(\theta_f x)} \left( \sum_{k \in \text{supp}(\theta_g x)} |g_k(x)| \right)^p \\ &= \left( \max_{1 \leq j \leq n, 1 \leq k \leq m} |f_j(\omega_k)| \right)^p \|\theta_f x\|_0 \left( \sum_{k \in \text{supp}(\theta_g x)} |g_k(x)| \right)^p \\ &\leq \left( \max_{1 \leq j \leq n, 1 \leq k \leq m} |f_j(\omega_k)| \right)^p \|\theta_f x\|_0 \left( \sum_{k \in \text{supp}(\theta_g x)} |g_k(x)|^p \right)^{\frac{p}{p}} \left( \sum_{k \in \text{supp}(\theta_g x)} 1^q \right)^{\frac{p}{q}} \\ &= \left( \max_{1 \leq j \leq n, 1 \leq k \leq m} |f_j(\omega_k)| \right)^p \|\theta_f x\|_0 \|\theta_g x\|^p \|\theta_g x\|_0^{\frac{p}{q}} \\ &= \left( \max_{1 \leq j \leq n, 1 \leq k \leq m} |f_j(\omega_k)| \right)^p \|\theta_f x\|_0 \|x\|^p \|\theta_g x\|_0^{\frac{p}{q}}. \end{aligned}$$

Therefore

$$\frac{1}{\max_{1 \leq j \leq n, 1 \leq k \leq m} |f_j(\omega_k)|} \leq \|\theta_f x\|_0^{\frac{1}{p}} \|\theta_g x\|_0^{\frac{1}{q}}.$$

On the other way, first using  $\theta_g$  is an isometry and  $\theta_f$  is an isometry, we get

$$\begin{aligned}
 \|x\|^p &= \|\theta_g x\|^p = \sum_{k=1}^m |g_k(x)|^p = \sum_{k \in \text{supp}(\theta_g x)} |g_k(x)|^p \\
 &= \sum_{k \in \text{supp}(\theta_g x)} \left| g_k \left( \sum_{j=1}^n f_j(x) \tau_j \right) \right|^p = \sum_{k \in \text{supp}(\theta_g x)} \left| \sum_{j=1}^n f_j(x) g_k(\tau_j) \right|^p \\
 &= \sum_{k \in \text{supp}(\theta_g x)} \left| \sum_{j \in \text{supp}(\theta_f x)} f_j(x) g_k(\tau_j) \right|^p \leq \sum_{k \in \text{supp}(\theta_g x)} \left( \sum_{j \in \text{supp}(\theta_f x)} |f_j(x) g_k(\tau_j)| \right)^p \\
 &\leq \left( \max_{1 \leq j \leq n, 1 \leq k \leq m} |g_k(\tau_j)| \right)^p \sum_{k \in \text{supp}(\theta_g x)} \left( \sum_{j \in \text{supp}(\theta_f x)} |f_j(x)| \right)^p \\
 &= \left( \max_{1 \leq j \leq n, 1 \leq k \leq m} |g_k(\tau_j)| \right)^p \|\theta_g x\|_0 \left( \sum_{j \in \text{supp}(\theta_f x)} |f_j(x)| \right)^p \\
 &\leq \left( \max_{1 \leq j \leq n, 1 \leq k \leq m} |g_k(\tau_j)| \right)^p \|\theta_g x\|_0 \left( \sum_{j \in \text{supp}(\theta_f x)} |f_j(x)|^p \right)^{\frac{p}{p}} \left( \sum_{j \in \text{supp}(\theta_f x)} 1^q \right)^{\frac{p}{q}} \\
 &= \left( \max_{1 \leq j \leq n, 1 \leq k \leq m} |g_k(\tau_j)| \right)^p \|\theta_g x\|_0 \|\theta_f x\|^p \|\theta_f x\|_0^{\frac{p}{q}} \\
 &= \left( \max_{1 \leq j \leq n, 1 \leq k \leq m} |g_k(\tau_j)| \right)^p \|\theta_g x\|_0 \|x\|^p \|\theta_f x\|_0^{\frac{p}{q}}
 \end{aligned}$$

Therefore

$$\frac{1}{\max_{1 \leq j \leq n, 1 \leq k \leq m} |g_k(\tau_j)|} \leq \|\theta_g x\|_0^{\frac{1}{p}} \|\theta_f x\|_0^{\frac{1}{q}}.$$

□

**Corollary 2.4.** *Theorem 1.3 follows from Theorem 2.3.*

*Proof.* Let  $\{\tau_j\}_{j=1}^n, \{\omega_j\}_{j=1}^n$  be two Parseval frames for a finite dimensional Hilbert space  $\mathcal{H}$ . Define

$$f_j : \mathcal{H} \ni h \mapsto \langle h, \tau_j \rangle \in \mathbb{K}; \quad g_j : \mathcal{H} \ni h \mapsto \langle h, \omega_j \rangle \in \mathbb{K}, \quad \forall 1 \leq j \leq n.$$

Then  $p = q = 2$ ,  $\theta_\tau = \theta_f$ ,  $\theta_\omega = \theta_g$  and

$$|f_j(\omega_k)| = |\langle \omega_k, \tau_j \rangle|, \quad \forall 1 \leq j, k \leq n.$$

□

Theorem 2.3 brings the following question.

**Question 2.5.** *Given  $p, m, n$  and a Banach space  $\mathcal{X}$ , for which pairs of  $p$ -Schauder frames  $(\{f_j\}_{j=1}^n, \{\tau_j\}_{j=1}^n)$  and  $(\{g_k\}_{k=1}^m, \{\omega_k\}_{k=1}^m)$ , we have equality in Inequality (3)?*

#### REFERENCES

- [1] David L. Donoho and Philip B. Stark. Uncertainty principles and signal recovery. *SIAM J. Appl. Math.*, 49(3):906–931, 1989.

## FUNCTIONAL DONOHO-STARK-ELAD-BRUCKSTEIN-RICAUD-TORRÉSANI UNCERTAINTY PRINCIPLE

- 
- [2] Michael Elad and Alfred M. Bruckstein. A generalized uncertainty principle and sparse representation in pairs of bases. *IEEE Trans. Inform. Theory*, 48(9):2558–2567, 2002.
  - [3] W. Heisenberg. The physical content of quantum kinematics and mechanics. In John Archibald Wheeler and Wojciech Hubert Zurek, editors, *Quantum Theory and Measurement*, Princeton Series in Physics, pages 62–84. Princeton University Press, Princeton, NJ, 1983.
  - [4] James R. Holub. Pre-frame operators, Besselian frames, and near-Riesz bases in Hilbert spaces. *Proc. Amer. Math. Soc.*, 122(3):779–785, 1994.
  - [5] K. Mahesh Krishna and P. Sam Johnson. Towards characterizations of approximate Schauder frame and its duals for Banach spaces. *J. Pseudo-Differ. Oper. Appl.*, 12(1):Paper No. 9, 13, 2021.
  - [6] Benjamin Ricaud and Bruno Torrésani. Refined support and entropic uncertainty inequalities. *IEEE Trans. Inform. Theory*, 59(7):4272–4279, 2013.