
The geometry of financial institutions - Wasserstein clustering of financial data

Lorenz Riess
University of Vienna and
Oesterreichische Nationalbank

Mathias Beiglboeck
University of Vienna

Johannes Temme
Oesterreichische Nationalbank

Andreas Wolf
Oesterreichische Nationalbank

Julio Backhoff-Veraguas
University of Vienna

Abstract

The increasing availability of granular and big data on various objects of interest has made it necessary to develop methods for condensing this information into a representative and intelligible map. Financial regulation is a field that exemplifies this need, as regulators require diverse and often highly granular data from financial institutions to monitor and assess their activities. However, processing and analyzing such data can be a daunting task, especially given the challenges of dealing with missing values and identifying clusters based on specific features.

To address these challenges, we propose a variant of Lloyd’s algorithm that applies to probability distributions and uses generalized Wasserstein barycenters to construct a metric space which represents given data on various objects in condensed form. By applying our method to the financial regulation context, we demonstrate its usefulness in dealing with the specific challenges faced by regulators in this domain. We believe that our approach can also be applied more generally to other fields where large and complex data sets need to be represented in concise form.

1 Introduction

The contribution of this article is to provide an algorithm which takes as input a number of (discrete) probability distributions in a high dimensional space which are then *clustered and represented* as points in a metric space, where the distance gauges differences between the individual distributions. A crucial feature of the algorithm is that we allow for missing data, which may be missing systematically. Specifically, individual distributions may be missing all information in one or more coordinates.

Our original motivation to devise this type of algorithm stems from a challenge faced by regulators of the financial industry. Data that are delivered from financial institutions to the regulator consist of various different formats, from highly aggregate data such as the total volume of the balance sheet, down to very granular data about individual credits described by characteristics such as volume, interest rate, etc. Two individual credits might then be considered as similar if those characteristics take similar numerical values, i.e. have small Euclidean distance when viewed as vectors in \mathbb{R}^d . To build a distance between financial institutions, one can view these institutions as probability distributions over a high dimensional space, e.g. by considering the empirical measure of reported credits, and determine the respective distance as a Wasserstein distance. One could then interpret the landscape of all financial institutions as an ensemble of points in the Wasserstein space, susceptible to familiar methods of clustering, outlier detection, etc. A particular challenge which renders the problem more complicated is the “missingness of data”. Data delivered by institutions may have missing values. More crucially, data may be missing systematically, as different institutions are required to deliver different data at varying levels of detail and granularity.

The algorithm we propose as a ramification simultaneously clusters probability measures with missing data and represents them as elements of a metric space. The principal structure follows the idea of Lloyd’s algorithm for k -means clustering and combines it with the concept of generalized Wasserstein barycenters, recently introduced by Delon, Gozlan and Saint-Dizier [7]. A classical approach to dealing with missing data would be to impute from (weighted) nearest neighbors or a probability distribution in the same cluster. However these types of imputation systematically skew results in the sense that probability distributions with less reported data can appear as being closer to other points which the imputation procedure is attempting to mimic. This type of bias may be highly undesirable, e.g. when one is trying to identify distributions that are “atypical”, where “atypical” could specifically refer to the manner in which data are missing. We devise a particular way of soft imputation which accounts for a random element in filling up missing values, and in particular avoids the above mentioned bias.

For technical reasons we will present our algorithm in a more general form in Section 3. That is, we cluster and perform soft imputation for points in an abstract metric space rather than a Wasserstein space of probability measures. This allows us to simplify the presentation and has the additional benefit that we obtain a version of classical Euclidean k -means clustering with missing data. This general perspective allows us in Section 7.1 to compare our method with existing solutions for clustering in \mathbb{R}^d with missing data. As for the important case of probability distributions with missing data our contribution is detailed in Section 5 and complemented experimentally in Section 7.2 with results using actual (anonymized) data stemming from the original motivation of regulation of financial institutions.

Related Literature Clustering distributions using Wasserstein distances has been explored in various works including [17, 25, 23, 12, 10]. A different financial context was also considered in this area, including [11] by Horvath et al., for clustering of market regimes. In [21], Staib et al. proposed the Wasserstein k -means++ and its initialization strategy, generalizing the classical k -means++ algorithm (introduced in [2]). However, to the best of our knowledge, clustering distributions with missing coordinates has not been investigated before. To address this problem, we rely heavily on the concept of the generalized Wasserstein barycenter, introduced by Delon, Gozlan, and Saint-Dizier [7], which extends the classical Wasserstein barycenter of Agueh and Carlier [1]. For computational aspects related to optimal transport and regularized Wasserstein distance, we refer to the work of Cuturi [5] and the book [18] by Cuturi and Peyré. Additionally, we utilize the POT (0.8.2) package (a Python package for optimal transport), as detailed in [8], extensively for implementation purposes.

2 k -means clustering in metric spaces: a summary

Let x_1, \dots, x_N be given points in a fixed metric space (\mathcal{X}, d) . The metric k -means problem is concerned with assigning the data points into k clusters. The clusters hereby are governed by k barycenters, which will be some type of average or barycenter of the points in the corresponding cluster. k -means was first introduced by MacQueen in [16]. The problem can be formalized as

$$\min \left\{ \sum_{i=1}^N d(x_i, c_{a_i})^2 : c_1, \dots, c_k \in \mathcal{X}, a \in \{1, \dots, k\}^N \right\}. \quad (\text{KM})$$

In this formulation a denotes a vector of assignments, i.e. a_i indicates the cluster membership of data point x_i . The points c_1, \dots, c_k serve as cluster barycenters. Furthermore a notion of barycenter, being typically a function of some of the data points, is needed. Then, problem (KM) can be tackled by the well-established Lloyd algorithm, see also [14].

Following an initialization phase, two steps are iterated:

- i) assignment step: given barycenters $c_1, \dots, c_k \in \mathcal{X}$, pick for each $i \in \{1, \dots, N\}$,

$$a_i \in \arg \min \{d(x_i, c_j) : j \in \{1, \dots, k\}\}, \quad (1)$$

- ii) barycenter step: given assignment a , update the barycenters, i.e. for each $j \in \{1, \dots, k\}$ set

$$c_j \in \text{barycenter}(\{x_i : a_i = j\}). \quad (2)$$

Let us give two examples with particular choices of data space \mathcal{X} , metric d , and barycenter operation, which will be of interest in the sequel:

Example 2.1. Let $\mathcal{X} = \mathbb{R}^d$ and $d(x, y) = \|x - y\|_2$, the Euclidean space and distance. Then (KM) simplifies to the classical k -means problem in Euclidean space,

$$\min \left\{ \sum_{i=1}^N \|x_i - c_{a_i}\|_2^2 : c_1, \dots, c_k \in \mathbb{R}^d, a \in \{1, \dots, k\}^N \right\}. \quad (k\text{-means})$$

The barycenter function is the Euclidean mean/average in this case.

Example 2.2. Take $\mathcal{X} = \mathcal{P}_2(\mathbb{R}^d)$, the space of all Borel probability measures on \mathbb{R}^d with finite second moments, and take for d the Wasserstein distance $d(\mu, \nu) := W_2(\mu, \nu)$, $\mu, \nu \in \mathcal{P}_2(\mathbb{R}^d)$ (recalled in (10) below). We will henceforth call $\mu_i = x_i$, $i = 1, \dots, N$. Then (KM) turns into

$$\min \left\{ \sum_{i=1}^N W_2^2(\mu_i, \nu_{a_i}) : \nu_1, \dots, \nu_k \in \mathcal{P}_2(\mathbb{R}^d), a \in \{1, \dots, k\}^N \right\}. \quad (\text{W-}k\text{-means})$$

As barycenter one takes the Wasserstein barycenter, i.e. the barycenter updating step (2) reads as

$$\nu_j \in \arg \min \left\{ \sum_{i:a_i=j} W_2^2(\mu_i, \nu) : \nu \in \mathcal{P}_2(\mathbb{R}^d) \right\}. \quad (3)$$

In what follows we want to generalize the previous examples, especially Example 2.2, to the case in which certain coordinates/marginals are missing for certain observations, for an illustration see Figure 1 in Section 5.

For Lloyd’s algorithm, both a distance function and a notion of barycenter are needed. In the case of a metric space one could always consider a Fréchet mean for the barycenter operator, see [9]. In the case of missing coordinates, however, the data does not come from one metric space but from several different ones, therefore several distances will be considered. For the same reason, in order to define a proper notion of barycenter, some generalization will be needed. We will introduce a general approach to this in the next section and discuss our specific examples in more detail in Example 3.1 and Section 5.

3 Clustering projected elements of metric spaces

Suppose we are working in a metric space (\mathcal{X}, d) and that there are points $x_1, \dots, x_N \in \mathcal{X}$ which we want to assign to k clusters. However, what we really observe are the points $\tilde{x}_i := \varphi_i(x_i)$, where $\varphi_i : \mathcal{X} \rightarrow \mathcal{X}_i$ is a known map into another metric space \mathcal{X}_i . One can think for example of a projection onto some coordinates in the case of $\mathcal{X} = \mathbb{R}^d$. Having observed $\tilde{x}_i \in \mathcal{X}_i$ define for $y \in \mathcal{X}$

$$d_i(\tilde{x}_i, y) := d(\varphi_i^{-1}(\tilde{x}_i), y) := \inf \{ d(x, y) : x \in \mathcal{X}, \varphi_i(x) = \tilde{x}_i \}, \quad (4)$$

which serves as a type of dissimilarity measure of a point $y \in \mathcal{X}$ in the “full” metric space.

3.1 Problem and algorithm

Having this at hand, we introduce the problem

$$\min \left\{ \sum_{i=1}^N d_i(\tilde{x}_i, c_{a_i})^2 : c_1, \dots, c_k \in \mathcal{X}, a \in \{1, \dots, k\}^N \right\}. \quad (5)$$

This says we want to find optimal cluster barycenters in \mathcal{X} , as well as a vector optimally assigning each observed point to a cluster. It is important to note that we are always looking for cluster barycenters in the “full” space \mathcal{X} .

We suggest to tackle (5) by the following two steps which are iterated in a Lloyd algorithm fashion:

- i) given c_1, \dots, c_k , set $a_i \in \arg \min \{ d_i(\tilde{x}_i, c_j) : j \in \{1, \dots, k\} \}$,
- ii) given a , put $c_j \in \arg \min \{ \sum_{i:a_i=j} d_i(\tilde{x}_i, y)^2 : y \in \mathcal{X} \}$.

Let us note that step ii) does not necessarily admit minimizers. However, in our two applications, i.e. NA k -means in Example 3.1 and NA Wasserstein k -means, which is discussed in Section 5, minimizers always exist. For a more precise discussion of this technical point see Appendix B.

3.2 Using clustering for imputation

After clustering we obtain a vector of assignments a and barycenters c_1, \dots, c_k . Define

$$I_f := \{i : \varphi_i = Id_{\mathcal{X}}\}, \quad I_m := \{1, \dots, N\} \setminus I_f, \quad (6)$$

i.e. respectively the set of indices being observed in \mathcal{X} (i.e. fully observed data) and the set of indices which are only observed after some non-trivial map (i.e. with missing data). Let us also define the final clusters, i.e. for each $j \in \{1, \dots, k\}$ define

$$C_j := \{i : a_i = j\}.$$

We want to use the clusters to find, for a point \tilde{x}_i with $i \in I_m$, a probability measure on \mathcal{X} , i.e. an element of $\mathcal{P}(\mathcal{X})$, which hopefully concentrates around the true unobserved point x_i . We define for $i \in I_m$ the set of indices we use for filling up the missing data of \tilde{x}_i , as

$$J_i := I_f \cap C_{a_i}.$$

This says that we want to use the “full” points in the same cluster as \tilde{x}_i in order to compensate the incomplete information that we have about \tilde{x}_i . It can of course happen that $J_i = \emptyset$. In this case we use the corresponding cluster barycenter c_{a_i} to complement the information about \tilde{x}_i . In the following we assume $J_i \neq \emptyset$. Taking now some x_ℓ with $\ell \in J_i$ we set

$$y_\ell^i \in \arg \min \{d(x, x_\ell) : x \in \mathcal{X}, \varphi_i(x) = \tilde{x}_i\}, \quad (7)$$

which will be one choice of “filling up”. We introduce the shorthand $D_\ell^i := d(y_\ell^i, x_\ell) = d(\varphi_i^{-1}(\tilde{x}_i), x_\ell)$. In order to determine the weights of a probability measure, we set $p_\ell^i := f(D_\ell^i)$ with $f : [0, \infty) \rightarrow [0, \infty)$ being some positive decreasing function fixed in advance, e.g. $f(x) := \exp(-x^2)$. This way $p_\ell^i \geq 0$ and also $\sum_{\ell \in J_i} p_\ell^i = 1$ after possibly renormalizing the weights by a positive constant. Having obtained the weights, we define the probability measure which should represent a *randomly reconstructed* x_i as

$$\theta_i := \sum_{\ell \in J_i} p_\ell^i \delta_{y_\ell^i}.$$

To embed also the fully observed points, for $i \in I_f$ we set $\theta_i := \delta_{x_i}$, i.e. the Dirac delta concentrated on x_i . For $i \in I_m$ with $J_i = \emptyset$, we set θ_i to be the Dirac delta concentrated on some minimizer of $d(\varphi_i^{-1}(\tilde{x}_i), c_{a_i})$. Thus we embed all observed points $\tilde{x}_1, \dots, \tilde{x}_N$ in the same space, i.e. $\mathcal{P}(\mathcal{X})$. In other words, we identify the possibly unobserved point x_i with a probability measure θ_i .

In order to compare the hitherto constructed probability measures, as we will need to do in Section 7, we define a (generalized) metric on $\mathcal{P}(\mathcal{X})$, see Lemma A.1, through the cost induced by the *product* or *independent coupling*. That is, for $\mu, \nu \in \mathcal{P}(\mathcal{X})$ we set $\rho(\mu, \nu) := 0$ if $\mu = \nu$ and otherwise

$$\rho(\mu, \nu) := \int_{\mathcal{X} \times \mathcal{X}} d(x, x') d\mu \otimes \nu(x, x') := \int_{\mathcal{X}} \int_{\mathcal{X}} d(x, x') d\mu(x) d\nu(x'). \quad (8)$$

With a Wasserstein distance, or a similar notion of distance, imputed points are biased to be closer than “fully” observed points. The metric ρ is designed to avoid this type of bias. Indeed, the choice in (8) formalises the idea that missing values are not imputed in a deterministic sense but in a *random* or *soft* fashion, as indicated in the introduction. The distance between two randomly imputed values is then estimated as an *independent* average of distances, following the rationale that the imputation for one point does not inform the imputation for a different point.

Example 3.1 (NA k -means). *Corresponding to classical k -means, i.e. Example 2.1, consider $\mathcal{X} = \mathbb{R}^d$. Let $\varphi_i = P_i : \mathbb{R}^d \rightarrow \mathbb{R}^{d_i}$ be projections to some of the coordinates. Then (5) becomes*

$$\min \left\{ \sum_{i=1}^N \|P_i(x_i) - P_i(c_{a_i})\|_2^2 : c_1, \dots, c_k \in \mathbb{R}^d, a \in \{1, \dots, k\}^N \right\}. \quad (9)$$

The cluster assignment iteration steps assigns each point to the cluster to whose barycenter it is closest to, by only considering the known coordinates. The barycenter updating step is solved by $(\sum_{i:a_i=j} P_i^T P_i)^{-1} \sum_{i:a_i=j} P_i^T P_i(x_i)$, which in each coordinate is the average of all the points in the cluster having that coordinate. See Section 5 for details if the inverse does not exist. By imputing according to the above discussion, we obtain measures in $\mathcal{P}_2(\mathbb{R}^d)$, on which we can then calculate pairwise distances using the metric ρ in (8). Experiments using this method may be found in Section 7.1. It is worth mentioning that Chi et. al. in [4] consider the same loss function, i.e. (9), but they propose a different algorithm for clustering Euclidean points with missing values.

Before discussing our second example in Section 5, corresponding to the generalization of Example 2.2, we recall the needed notions from optimal transport theory.

4 Wasserstein distance and generalized Wasserstein barycenter: a summary

Optimal transport and Wasserstein distance Let μ, ν be probabilities on Polish spaces X, Y resp. Denote by

$$\Pi(\mu, \nu) := \{\pi \in \mathcal{P}(X \times Y) : pr_{X\#}\pi = \mu, pr_{Y\#}\pi = \nu\}$$

the set of all measures on the product space having μ and ν as marginals. The Kantorovich problem for a cost function $c : X \times Y \rightarrow \mathbb{R}_+$, introduced in [13], is

$$\inf_{\pi \in \Pi(\mu, \nu)} \int_{X \times Y} c(x, y) d\pi(x, y). \quad (\text{KP})$$

We specialise to the case $X = Y = \mathbb{R}^d$ and $c(x, y) = \|x - y\|_p^p$ for $p \in [1, \infty)$. The Wasserstein distance on the space $\mathcal{P}_p(\mathbb{R}^d)$ of probability measures with finite p -moment is then defined as

$$W_p(\mu, \nu) := \left(\inf_{\pi \in \Pi(\mu, \nu)} \int_{\mathbb{R}^d \times \mathbb{R}^d} \|x - y\|_p^p d\pi(x, y) \right)^{1/p}. \quad (10)$$

Wasserstein barycenter A notion of averaging probabilities which has recently received significant attention is the notion of Wasserstein barycenters, introduced by Agueh and Carlier [1]. A Wasserstein barycenter of $\mu_1, \dots, \mu_n \in \mathcal{P}_2(\mathbb{R}^d)$ with weights $\lambda_1, \dots, \lambda_n \geq 0$, is a solution of

$$\inf \left\{ \sum_{i=1}^n \lambda_i W_2^2(\mu_i, \nu) : \nu \in \mathcal{P}_2(\mathbb{R}^d) \right\}.$$

Since we want to generalize the algorithm introduced in Sections 2-3 to the setting of distributions we need to have a variant of the Wasserstein barycenter, which is applicable if only some coordinates of the measures are known. An appropriate generalized Wasserstein barycenter was introduced and studied by Julien, Gozlan and Saint-Dizier in [7]. To formulate it, let probability measures $\mu_1, \dots, \mu_n \in \mathcal{P}_2(\mathbb{R}^d)$ be given and consider linear maps $P_i : \mathbb{R}^d \rightarrow \mathbb{R}^{d_i}$. In our intended applications, these will be projections onto some of the coordinates. A generalized Wasserstein barycenter of the push-forwarded measures $P_1\#\mu_1, \dots, P_n\#\mu_n$ with weights $\lambda_1, \dots, \lambda_n \geq 0$, is a solution of

$$\inf \left\{ \sum_{i=1}^n \lambda_i W_2^2(P_i\#\mu_i, P_i\#\nu) : \nu \in \mathcal{P}_2(\mathbb{R}^d) \right\}. \quad (11)$$

To solve the problem it is useful to translate it back to a classical Wasserstein barycenter problem. By associating to P_i a matrix $P_i \in \mathbb{R}^{d_i \times d}$, define $A := \sum_{i=1}^n \lambda_i P_i^T P_i$ and assume in the following that A is invertible. Then set $\tilde{\mu}_i := (A^{-1/2} P_i^T)\#\mu_i \in \mathcal{P}_2(\mathbb{R}^d)$ for $i = 1, \dots, n$, and consider the conventional Wasserstein barycenter problem

$$\inf \left\{ \sum_{i=1}^n \lambda_i W_2^2(\tilde{\mu}_i, \tilde{\nu}) : \tilde{\nu} \in \mathcal{P}_2(\mathbb{R}^d) \right\}. \quad (12)$$

Then, ν is a solution to (11) if and only if $A^{1/2}\#\tilde{\nu}$ is a solution to (12), thanks to [7, Proposition 3.1]. Concerning computational aspects, especially in the discrete case, efficient algorithms for calculating Wasserstein barycenters have already been developed; see the work [6] by Cuturi and Doucet, as well as [15] by Álvarez, del Barrio, Cuesta-Albertos and Matrán, to name but two.

5 NA Wasserstein k -means

Algorithm description Having the two operations needed in Section 3 at hand, we can discuss the case of $\mathcal{X} = \mathcal{P}_2(\mathbb{R}^d)$ in detail. Suppose we want to cluster probability measures $\mu_1, \dots, \mu_N \in \mathcal{P}_2(\mathbb{R}^d)$ into k clusters but we only observe the push forwards $P_i\#\mu_i \in \mathcal{P}_2(\mathbb{R}^{d_i})$ under projections P_i , $i = 1, \dots, N$. Hence we have $\varphi_i(\cdot) := P_i\#(\cdot)$. Then problem (5) reads as

$$\inf \left\{ \sum_{i=1}^N W_2^2(P_i\#\mu_i, P_i\#\nu_{a_i}) : \nu_1, \dots, \nu_k \in \mathcal{P}_2(\mathbb{R}^d), a \in \{1, \dots, k\}^N \right\}. \quad (\text{NA-W-}k\text{-means})$$

We can tackle problem (NA-W- k -means) by the suggested iterations in Section 3.1, which read as

- i) update cluster assignments given barycenters ν_1, \dots, ν_k , i.e. for each $i \in \{1, \dots, N\}$ set

$$a_i \in \arg \min \{ W_2(P_i\#\mu_i, P_i\#\nu_j) : j \in \{1, \dots, k\} \}, \quad (13)$$

- ii) update cluster barycenters given an assignment a , i.e. for each $j \in \{1, \dots, k\}$ set

$$\nu_j \in \arg \min \left\{ \sum_{i: a_i=j} W_2^2(P_i\#\mu_i, P_i\#\nu) : \nu \in \mathcal{P}_2(\mathbb{R}^d) \right\}.$$

Again, we want to find a vector a of optimally assigning each measure to one of the k clusters, as well as optimal barycenters, or in this case generalized Wasserstein barycenters, ν_1, \dots, ν_k . Note that once more we are looking for barycenters on the full space, i.e. having all coordinates.

It can happen that a cluster ends up consisting of measures which all miss the same coordinate, i.e. for some $j \in \{1, \dots, k\}$ we have $\bigcap_{i:a_i=j} \ker P_i \neq \{0\}$. In order not to run into numerical problems in that case, one can adapt step ii): Assume at iteration t the barycenters are $\nu_1^{(t)}, \dots, \nu_k^{(t)}$ and a new assignment vector $a^{(t+1)}$ has been calculated. Introduce a weight $\lambda^{(t)} \in [0, 1]$ and set

$$\nu_j^{(t+1)} \in \arg \min \left\{ (1 - \lambda^{(t)}) \sum_{i:a_i^{(t+1)}=j} W_2^2(P_i \# \mu_i, P_i \# \nu) + \lambda^{(t)} W_2^2(\nu_j^{(t)}, \nu) : \nu \in \mathcal{P}_2(\mathbb{R}^d) \right\}.$$

Experimentally we have observed that $\lambda^{(t)} := \frac{1}{(t+1)^{1/2}}$ works well. Having defined the two steps to be iterated, we can present the main computational contribution of this work, Algorithm 1:

Algorithm 1: NA Wasserstein k -means algorithm

Data: N observed measures $P_1 \# \mu_1, \dots, P_N \# \mu_N$, number of cluster k , maximum number of iterations T , weighting schedule $(\lambda^{(t)})_{t=0}^T$

Result: barycenters ν_1, \dots, ν_k , assignment vector $a \in \{1, \dots, K\}^N$

$t := 0$, initialize $\nu_1^{(0)}, \dots, \nu_k^{(0)}$, and $a^{(0)}$

while $t < T$ **do**

for $i = 1, \dots, N$ **do**

$$\quad \quad \quad a_i^{(t+1)} := \arg \min \{ W_2^2(P_i \# \mu_i, P_i \# \nu_j^{(t)}) : j \in \{1, \dots, k\} \} \quad (14)$$

for $j = 1, \dots, k$ **do**

 set $\nu_j^{(t+1)}$ as solution of

$$\quad \quad \quad \min \left\{ (1 - \lambda^{(t)}) \sum_{i:a_i^{(t+1)}=j} W_2^2(P_i \# \mu_i, P_i \# \nu) + \lambda^{(t)} W_2^2(\nu_j^{(t)}, \nu) : \nu \in \mathcal{P}_2(\mathbb{R}^d) \right\}$$

if $a^{(t)} == a^{(t+1)}$ **then**

break

$t := t + 1$

In (14) we use the entry of the old assignment vector $a_i^{(t)}$ if it is a minimizer. We can formulate a simple convergence result whose proof is reported in Appendix A. Let

$$L(\nu_1, \dots, \nu_k, a) := \sum_{i=1}^N W_2^2(P_i \# \mu_i, P_i \# \nu_{a_i}).$$

Theorem 5.1. *Given initialization $\nu_1^{(0)}, \dots, \nu_k^{(0)} \in \mathcal{P}_2(\mathbb{R}^d)$, $a \in \{1, \dots, k\}^N$, Algorithm 1 strictly decreases L until it terminates after finitely many steps, for any choice of $(\lambda^{(t)})_{t \in \mathbb{N}}$ (even for $T = \infty$). If in every iteration each cluster has a measure with all coordinates, then Algorithm 1 with $\lambda^{(t)} = 0$ yields a local minimum of L .*

A toy example To illustrate the algorithm, consider six measures on the plane, of which one misses the vertical coordinate (denoted by vertical lines). Each measure consists of three support points with equal weight. The measures are visualized by different colors and depicted in the left of Figure 1. If we think of clustering these measures into three clusters, it is quite clear which clusters we want. By applying the introduced algorithm this is indeed the case and leads to the results depicted in Figure 1. To obtain it we calculated free support barycenters in each step of the algorithm with a fixed support size of three. For the blue and brown barycenters each support point is precisely the average of support points of the two measures in that clusters. For the pink barycenter the situation is slightly different. In the y coordinate it inherits the values of the red measure, while as in the x coordinate the support points are averages of the x coordinates of a red and rose support point, respectively.



Figure 1: (left): 6 measures on the plane of which one misses the vertical coordinate of its points. (right) Clustering of the 6 measures into 3 clusters.

Imputation via NA Wasserstein k -means Let us look how the imputation procedure according to Section 3.2 works in the case where $\mathcal{X} = \mathcal{P}_2(\mathbb{R}^d)$. Having clustered measures $P_i \# \mu_i \in \mathcal{P}_2(\mathbb{R}^{d_i})$, $i = 1, \dots, N$, we obtain barycenters $\nu_1, \dots, \nu_k \in \mathcal{P}_2(\mathbb{R}^d)$ and a vector of assignments $a \in \{1, \dots, k\}^N$. For $i \in I_m$, see (6) for the definition, we then set

$$\mu_i := \sum_{\ell \in J_i} p_\ell^i \delta_{\tau_i \# (P_i \# \mu_i \otimes P_i^C \# \mu_\ell)}$$

to obtain a measure $\mu_i \in \mathcal{P}_2(\mathcal{P}_2(\mathbb{R}^d))$, i.e. a random measure. Here, by P_i^C we denote the projection on the missing coordinates of $P_i \# \mu_i$, and by τ_i we denote the unique permutation such that $\tau_i \circ (P_i, P_i^C) = Id_{\mathbb{R}^d}$. For $i \in I_f$ we set $\mu_i := \delta_{\mu_i}$. For $i \in I_m$ and $|J_i| > 1$ we suggest the weights

$$p_\ell^i \propto \exp\left(-\frac{\lambda}{2\sigma_i^2} W_2^2(P_i \# \mu_i, P_i \# \mu_\ell)\right), \quad \ell \in J_i.$$

Here, $\lambda > 0$ and the variance σ_i^2 is used to standardize the distances, i.e.

$$\sigma_i^2 := \frac{1}{|J_i|-1} \sum_{\ell \in J_i} W_2^2(P_i \# \mu_i, P_i \# \mu_\ell).$$

If $|J_i| = 1$ it is clear what to do, as we only have one measure to fill up with.

After imputation we obtain random measures $\mu_1, \dots, \mu_N \in \mathcal{P}_2(\mathcal{P}_2(\mathbb{R}^d))$. To calculate their distances, we use metric ρ from (8), i.e., for $\mu, \nu \in \mathcal{P}(\mathcal{P}_2(\mathbb{R}^d))$, $\rho(\mu, \nu) = 0$, if $\mu = \nu$, and

$$\rho(\mu, \nu) = \int_{\mathcal{P}_2(\mathbb{R}^d)} \int_{\mathcal{P}_2(\mathbb{R}^d)} W_2(\mu, \nu) d\mu(\mu) d\nu(\nu), \quad \text{if } \mu \neq \nu. \quad (15)$$

Limitations For the dataset of our application in Section 5 the method developed above is sufficient. However, for other applications it might be desirable to investigate the use of the *regularized Wasserstein distance* or *entropic optimal transport* in our method for reduced computational complexity. For this, a notion of a *generalized regularized Wasserstein barycenter*, as the NA-Wasserstein k -means algorithm relies heavily on the generalized Wasserstein barycenter has to be developed which we intend to do in future work.

6 Evaluation method

Our clustering/imputation method will be evaluated on Euclidean and distributional data. Hence we present here in abstract language the setting in which the method will be tested. Suppose we have elements x_1, \dots, x_N in a metric space $(\mathcal{X}, d_{\mathcal{X}})$, which we collect in the set $X := \{x_1, \dots, x_N\}$. We only observe data via the maps $\varphi_i : \mathcal{X} \rightarrow \mathcal{X}_i$ from Section 3, i.e. $\tilde{x}_i = \varphi_i(x_i)$. We collect the observed points in $(\tilde{x}_1, \dots, \tilde{x}_N) \in \mathcal{X}_{NA}$, where we define $\mathcal{X}_{NA} := \mathcal{X}_1 \times \dots \times \mathcal{X}_N$. Let us summarize this by saying that we only observe the values of (x_1, \dots, x_N) under the map $h : \mathcal{X}^N \rightarrow \mathcal{X}_{NA}$, i.e. $h_i(x_1, \dots, x_N) = \varphi_i(x_i) = \tilde{x}_i$.

The goal is to reconstruct the metric structure of X as accurately as possible. This means that we want to find a *good* metric space $(\mathcal{Y}, d_{\mathcal{Y}})$ and a reconstruction map $R : \mathcal{X}_{NA} \rightarrow \mathcal{Y}^N$.

Before finding a way of comparing the reconstructed points $R((\tilde{x}_1, \dots, \tilde{x}_N))$ to the original points X , we also want to consider another piece of information, namely that some observations of X might be more important for us than others. This is motivated by our main practical application of reconstructing a banking landscape, where we think of the importance of a bank being linked to attributes such as the size of that bank as measured, e.g. by its total assets. It is then more important to

reconstruct the characteristics of a very large bank than those of a much smaller bank. We model this by assigning weights proportional to the size of the bank. Formally, we define a probability measure μ_X on X by setting $\mu_X(x_i) := p_i \geq 0$, where we assume $\sum_{i=1}^N p_i = 1$. Naturally we define μ_Y , a probability measure on \mathcal{Y} , via $\mu_Y(R_i(\tilde{x}_1, \dots, \tilde{x}_N)) := p_i$.

To compare the metric measure spaces (X, d_X, μ_X) and $(R((\tilde{x}_1, \dots, \tilde{x}_N)), d_Y, \mu_Y)$, we use a *Gromov-Wasserstein distance* (see [22, 8] for resp. theory and implementation):

$$GW_2((\mathcal{X}, d_X, \mu), (\mathcal{Y}, d_Y, \nu))^2 := \inf_{\pi \in \Pi(\mu, \nu)} \int_{\mathcal{X}^2 \times \mathcal{Y}^2} |d_X(x, x') - d_Y(y, y')|^2 d\pi(x, y) d\pi(x', y').$$

7 Experimental results

7.1 Reconstructing points from a Gaussian Mixture Model

We start by comparing our clustering/reconstruction method to existing imputation methods in the Euclidean case, i.e. $\mathcal{X} = \mathbb{R}^d$, and evaluate the results using the Gromov-Wasserstein distance.

We simulate data from a Gaussian Mixture Model $\gamma := \sum_{j=1}^k \alpha_j \mathcal{N}(\mu_j, \Sigma_j)$, with $\sum_{j=1}^k \alpha_j = 1$. To simulate the importance of points, i.e. $\mu_X(x_i)$ from Section 6, we sample from Lognormal(μ, σ^2) to weigh each observation proportional to its sample value. In the simulation study we set $k = 5, d = 10$, and we always sample $N = 500$ points. To simulate missing values we use several missingness structures, i.e. several choices for the map h from Section 6. Before creating missing values, we adjust each dimension by its mean and standard deviation. In order to define h from Section 6, we calculate for each dimension the 25% quantile, draw a uniform random variable $h_j \sim U(0, 2/k)$ for each cluster j and choose a correlation parameter $\beta \in [0, 1]$. We then define h depending on β by the following properties: for each cluster $j \in \{1, \dots, k\}$ set $100 \min\{(\beta h_j + (1 - \beta)\alpha_j), 0.95\}\%$ to NA, and for each dimension set the values below the previously calculated 25% quantile to NA.

This means we are left with choosing the parameters $\alpha_1, \dots, \alpha_5, \mu_1, \dots, \mu_5, \Sigma_1, \dots, \Sigma_5$ for the Gaussian Mixture model, β for the map h , and μ, σ^2 for the lognormal which will govern the importance of sampled points. We set $\mu = 20, \sigma = 1.5$ as with these parameters the lognormal distribution models total assets of a financial institution reasonably well (empirically). We let β vary between 0 and 1 with steps 0.1. For the Gaussian mixture model we want to simulate the weights of the normal distributions, the means and the covariance, as non-informative as possible. Therefore, the α_j 's are uniformly sampled from the probability simplex. The μ_j 's are sampled uniformly on the cube $[-10, 10]^d$ and the Σ_j 's are sampled from a Wishart distribution with parameters (d, Id_d) .

For each choice of β we sample the explained parameters 100 times, create data, standardize it and calculate pairwise distances between data. Then, we create missing values according to h and compare different imputation procedures presented below to reconstruct the points and then calculate pairwise distances between the imputed points in the case if missing values are imputed by points. In the case of imputing the points by measures, we use the distance ρ from Section 3.2 to calculate pairwise distances. Afterwards we calculate the corresponding Gromov Wasserstein distance between original pairwise distances and imputed pairwise distances with the corresponding weights which come from the *Lognormal*(μ, σ^2). We look at the following imputation techniques:

- NA k -means: this is our method coming from Example 3.1,
- NA k -means-m: this corresponds to using NA k -means for clustering and imputation, but after obtaining the measures θ_i we calculate their expectation and use this as imputed point,
- mean imputation: impute each missing value by the average value of this attribute,
- median imputation: impute each missing value by the median value of this attribute,
- multiple imputation: a method [20, 3] trying out several values for each missing value,
- KNN: K-nearest-neighbour imputation which imputes each missing value by a weighted average of points close in the coordinates which are not missing, where we choose $K = 4$,
- LR: impute by regressing on points, with missing values and using predictions as imputation.

In Table 1 we report the estimated Gromov Wasserstein Distances \pm standard errors based on sampling 100 times for each scenario of missing values.

Table 1: Gromov Wasserstein Distance for Euclidean simulation.

settings	NA k -means.	NA k -means-m	mean imp.	median imp.	multiple imp.	KNN	LR
β 0.0	2.411 \pm 0.078	2.451 \pm 0.079	4.788 \pm 0.109	4.682 \pm 0.111	2.855 \pm 0.098	3.299 \pm 0.09	2.887 \pm 0.099
β 0.1	2.444 \pm 0.079	2.486 \pm 0.079	4.828 \pm 0.108	4.747 \pm 0.11	2.862 \pm 0.101	3.33 \pm 0.088	2.935 \pm 0.102
β 0.2	2.461 \pm 0.081	2.495 \pm 0.08	4.871 \pm 0.103	4.777 \pm 0.106	2.903 \pm 0.092	3.371 \pm 0.086	2.982 \pm 0.095
β 0.3	2.5 \pm 0.082	2.551 \pm 0.081	4.93 \pm 0.102	4.851 \pm 0.105	2.935 \pm 0.095	3.388 \pm 0.084	3.04 \pm 0.097
β 0.4	2.493 \pm 0.079	2.524 \pm 0.079	4.986 \pm 0.102	4.92 \pm 0.104	3.011 \pm 0.094	3.418 \pm 0.085	3.079 \pm 0.096
β 0.5	2.52 \pm 0.075	2.555 \pm 0.075	5.056 \pm 0.102	4.986 \pm 0.104	3.094 \pm 0.099	3.446 \pm 0.084	3.172 \pm 0.094
β 0.6	2.548 \pm 0.079	2.582 \pm 0.08	5.141 \pm 0.106	5.066 \pm 0.107	3.122 \pm 0.095	3.48 \pm 0.085	3.225 \pm 0.096
β 0.7	2.61 \pm 0.08	2.631 \pm 0.079	5.222 \pm 0.109	5.153 \pm 0.11	3.213 \pm 0.1	3.502 \pm 0.085	3.326 \pm 0.1
β 0.8	2.619 \pm 0.073	2.594 \pm 0.076	5.317 \pm 0.107	5.228 \pm 0.111	3.297 \pm 0.095	3.533 \pm 0.085	3.392 \pm 0.098
β 0.9	2.601 \pm 0.077	2.625 \pm 0.076	5.4 \pm 0.112	5.323 \pm 0.114	3.322 \pm 0.097	3.549 \pm 0.087	3.467 \pm 0.105
β 1.0	2.614 \pm 0.074	2.639 \pm 0.073	5.465 \pm 0.113	5.385 \pm 0.114	3.405 \pm 0.101	3.559 \pm 0.086	3.557 \pm 0.105

We see that, in the case of Euclidean points with missing values, the introduced method outperforms classical imputation methods when considering Gromov-Wasserstein as evaluation measure. Indeed, whether we apply *NA k -means* directly, or perform another averaging step, i.e. *NA k -means-m*, these two columns seem to outperform the remaining five for all settings of missing value creation. In Appendix C we use another way to evaluate the method, namely the Rand index, when viewing *NA k -means* solely as clustering algorithm and not as a method to reconstruct a metric structure. Also, further settings of missing value generations can be found there to accompany our results.

7.2 Reconstructing financial institutions

As mentioned in the introduction, our main motivation for developing the proposed method comes from clustering financial institutions based on granular loan data they are obliged to report to the central bank. Applying the proposed algorithm to such real data in a practical empirical example, we consider 500 financial institutions. For each of them we include information on 100 loans in 7 dimensions, i.e. each loan is described by up to 7 attributes. Among the attributes considered, we include e.g. credit volume, time to maturity, and interest rate. Out of the 500 institutions, 3 just reported 2 attributes, 9 institutions reported 3, 56 reported 4, 110 reported 5, 134 reported 6 and 188 reported all attributes. We cluster the institutions into 15 clusters using the *NA Wasserstein k -means* algorithm proposed in Section 5. Afterwards we use the imputation method described in the same section. Using the distance ρ on $\mathcal{P}_2(\mathcal{P}_2(\mathbb{R}^7))$, defined in (15), we calculate pairwise distances between the financial institutions. In order to get a visual representation of the landscape of financial institutions, and the clustering results, we use *multidimensional scaling* to represent the matrix of pairwise distances $(\rho(\theta_i, \theta_j))_{i,j=1}^{500}$ as points in \mathbb{R}^3 with a similar matrix of pairwise distances.




Figure 2: Visualization of pairwise distances.

In Figure 2 we see the representation of a banking landscape and clusters. Each color describes a cluster, and each point a bank. We observe that clusters are not strictly separated, due to multidimensional scaling, since we want to embed points via pairwise distances from a very high-dimensional into a very low-dimensional space. The visualisation may help analysts to further their understanding which financial institutions are similar to each other and also identify financial institutions which behave significantly different to others with respect to their loan structure. Detecting potential outliers is of course of particular interest.

8 Discussion

We have proposed a novel method for clustering elements in a metric space that are only observable under maps to potentially smaller metric spaces. The method can also reconstruct the metric structure of the original elements. In a simulation study in Euclidean space, our proposed method, boiling down to a new NA k -means algorithm, outperforms classical imputation methods when using the Gromov-Wasserstein distance as evaluation criterion. A particular variant of our method, the NA Wasserstein k -means algorithm, clusters probability measures that are only observable as push-forwards under projections. This algorithm is novel, and to our knowledge, there are no references in the literature dealing with partially observed distributions. In upcoming work with a central bank we will study, much more extensively than in Section 7 and from an enhanced economic point of view, the application to financial data for regulatory purposes.

References

- [1] M. Agueh and G. Carlier. Barycenters in the Wasserstein space. *SIAM Journal on Mathematical Analysis*, 43(2):904–924, 2011.
- [2] D. Arthur and S. Vassilvitskii. K-means++: The advantages of careful seeding. In *Proceedings of the Eighteenth Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA '07, page 1027–1035, USA, 2007. Society for Industrial and Applied Mathematics.
- [3] M. J. Azur, E. A. Stuart, C. Frangakis, and P. J. Leaf. Multiple imputation by chained equations: what is it and how does it work? *International Journal of Methods in Psychiatric Research*, 20(1):40–49, 2011.
- [4] J. T. Chi, E. C. Chi, and R. G. Baraniuk. k -pod: A method for k -means clustering of missing data. *The American Statistician*, 70(1):91–99, jan 2016.
- [5] M. Cuturi. Sinkhorn distances: Lightspeed computation of optimal transport. In C. J. C. Burges, L. Bottou, M. Welling, Z. Ghahramani, and K. Q. Weinberger, editors, *Advances in Neural Information Processing Systems*, volume 26. Curran Associates, Inc., 2013.
- [6] M. Cuturi and A. Doucet. Fast computation of wasserstein barycenters. In E. P. Xing and T. Jebara, editors, *Proceedings of the 31st International Conference on Machine Learning*, volume 32 of *Proceedings of Machine Learning Research*, pages 685–693, Beijing, China, 22–24 Jun 2014. PMLR.
- [7] J. Delon, N. Gozlan, and A. Saint-Dizier. Generalized Wasserstein barycenters between probability measures living on different subspaces. *To appear in Annals of Applied Probability*, 2021.
- [8] R. Flamary, N. Courty, A. Gramfort, M. Z. Alaya, A. Boisbunon, S. Chambon, L. Chapel, A. Corenflos, K. Fatras, N. Fournier, L. Gautheron, N. T. Gayraud, H. Janati, A. Rakotomamonjy, I. Redko, A. Rolet, A. Schutz, V. Seguy, D. J. Sutherland, R. Tavenard, A. Tong, and T. Vayer. Pot: Python optimal transport. *Journal of Machine Learning Research*, 22(78):1–8, 2021.
- [9] M. Fréchet. Les éléments aléatoires de nature quelconque dans un espace distancié. In *Annales de l'institut Henri Poincaré*, volume 10, pages 215–310, 1948.
- [10] N. Ho, X. L. Nguyen, M. Yurochkin, H. H. Bui, V. Huynh, and D. Phung. Multilevel clustering via wasserstein means. In *Proceedings of the 34th International Conference on Machine Learning - Volume 70*, ICML'17, page 1501–1509. JMLR.org, 2017.
- [11] B. Horvath, Z. Issa, and A. Muguruza. Clustering market regimes using the wasserstein distance. *arXiv:2110.11848*, 2021.
- [12] V. Huynh, N. Ho, N. Dam, X. Nguyen, M. Yurochkin, H. Bui, and D. Phung. On efficient multilevel clustering via wasserstein distances. *Journal of Machine Learning Research*, 2021.
- [13] L. Kantorovich. On the translocation of masses. *C. R. (Doklady) Acad. Sci. URSS (N.S.)*, 37:199–201, 1942.
- [14] S. Lloyd. Least squares quantization in pcm. *IEEE Transactions on Information Theory*, 28(2):129–137, 1982.
- [15] P. C. Álvarez Esteban, E. del Barrio, J. Cuesta-Albertos, and C. Matrán. A fixed-point approach to barycenters in wasserstein space. *Journal of Mathematical Analysis and Applications*, 441(2):744–762, 2016.

- [16] J. MacQueen. Some methods for classification and analysis of multivariate observations. *Proc. 5th Berkeley Symp. Math. Stat. Probab., Univ. Calif. 1965/66, 1, 281-297.*, 1967.
- [17] G. I. Papayiannis, G. N. Domazakis, D. Drivaliaris, S. Koukoulas, A. E. Tsekrekos, and A. N. Yannacopoulos. On clustering uncertain and structured data with wasserstein barycenters and a geodesic criterion for the number of clusters. *Journal of Statistical Computation and Simulation*, 91(13):2569–2594, mar 2021.
- [18] G. Peyré and M. Cuturi. Computational optimal transport. With applications to data sciences. *Found. Trends Mach. Learn.*, 11(5-6):1–262, 2018.
- [19] W. M. Rand. Objective criteria for the evaluation of clustering methods. *Journal of the American Statistical Association*, 66:846–850, 1971.
- [20] D. B. Rubin. *Multiple imputation for nonresponse in surveys*. Hoboken, NJ: John Wiley & Sons, reprint of the 1987 original edition, 2004.
- [21] M. Staib and S. Jegelka. Wasserstein k-means++ for cloud regime histogram clustering. In *Proceedings of the Seventh International Workshop on Climate Informatics: CI 2017*, 2017.
- [22] K.-T. Sturm. The space of spaces: curvature bounds and gradient flows on the space of metric measure spaces. *to appear in Memoirs AMS*, 2020.
- [23] I. Verdinelli and L. Wasserman. Hybrid Wasserstein distance and fast distribution clustering. *Electronic Journal of Statistics*, 13(2):5088 – 5119, 2019.
- [24] C. Villani. *Topics in optimal transportation*, volume 58 of *Grad. Stud. Math.* Providence, RI: American Mathematical Society (AMS), 2003.
- [25] Y. Zhuang, X. Chen, and Y. Yang. Wasserstein k -means for clustering probability distributions. In A. H. Oh, A. Agarwal, D. Belgrave, and K. Cho, editors, *Advances in Neural Information Processing Systems*, 2022.

A Proofs

Proof of Theorem 5.1. Let us first show that the loss function decreases monotonically. Calculate,

$$\begin{aligned}
& (1 - \lambda^{(t+1)})L(\nu_1^{(t)}, \dots, \nu_k^{(t)}, a^{(t)}) \\
&= (1 - \lambda^{(t+1)}) \sum_{i=1}^N W_2^2(P_i \# \mu_i, P_i \# \nu_{a_i^{(t)}}^{(t)}) \\
&\stackrel{(\star)}{\geq} (1 - \lambda^{(t+1)}) \sum_{i=1}^N W_2^2(P_i \# \mu_i, P_i \# \nu_{a_i^{(t+1)}}^{(t)}) \\
&= \sum_{j=1}^k \left[(1 - \lambda^{(t+1)}) \sum_{i: a_i^{(t+1)}=j} W_2^2(P_i \# \mu_i, P_i \# \nu_j^{(t)}) + \lambda^{(t+1)} \underbrace{W_2^2(\nu_j^{(t)}, \nu_j^{(t)})}_{=0} \right] \\
&\geq \sum_{j=1}^k \left[(1 - \lambda^{(t+1)}) \sum_{i: a_i^{(t+1)}=j} W_2^2(P_i \# \mu_i, P_i \# \nu_j^{(t+1)}) + \lambda^{(t+1)} \underbrace{W_2^2(\nu_j^{(t)}, \nu_j^{(t+1)})}_{\geq 0} \right] \\
&\geq (1 - \lambda^{(t+1)}) \sum_{i=1}^N W_2^2(P_i \# \mu_i, P_i \# \nu_{a_i^{(t+1)}}^{(t+1)}) \\
&= (1 - \lambda^{(t+1)})L(\nu_1^{(t+1)}, \dots, \nu_k^{(t+1)}, a^{(t+1)}).
\end{aligned}$$

Since $1 - \lambda^{(t+1)} > 0$ this shows the monotonicity. In inequality (\star) we have equality only if nothing changes in the assignment step, i.e. there is equality if and only if

$$a_i^{(t)} = a_i^{(t+1)}, \forall i \in \{1, \dots, N\}.$$

In that case the algorithm has converged and if there is some $i \in \{1, \dots, N\}$ such that $a_i^{(t)} \neq a_i^{(t+1)}$ we have a strict inequality, since we only change the assignment in Algorithm 1 if L decreases due to the new assignment.

Since only a finite number of possible assignments is possible because we are aiming to cluster finitely many measures into finitely many clusters, the algorithm strictly decreases L and converges after a finite number of steps, finishing the proof. \square

Let us also prove that ρ , defined in (8), is a distance, or to be more concrete just a generalized distance, i.e. we allow for ρ to take the value infinity.

Lemma A.1. *Let (\mathcal{X}, d) be a metric space and define on $\mathcal{P}(\mathcal{X})$ the map $\rho : \mathcal{P}(\mathcal{X}) \times \mathcal{P}(\mathcal{X}) \rightarrow [0, \infty]$ by*

$$\rho(\mu, \nu) := \begin{cases} \int_{\mathcal{X} \times \mathcal{X}} d(x, x') d(\mu \otimes \nu)(x, x'), & \text{if } \mu \neq \nu \\ 0, & \text{if } \mu = \nu. \end{cases}$$

Then, ρ is a generalized metric on $\mathcal{P}(\mathcal{X})$.

Proof. Symmetry of the generalized metric is clear due to the symmetry of the underlying distance d . Concerning the triangle inequality take $\mu, \nu, \theta \in \mathcal{P}(\mathcal{X})$ which we assume to be different from each other (otherwise there is nothing to prove). Furthermore, take three independent random variables $X \sim \mu, Y \sim \nu, Z \sim \theta$. Then,

$$\begin{aligned}
\rho(\mu, \theta) &= \int_{\mathcal{X} \times \mathcal{X}} d(x, x') d(\mu \otimes \theta)(x, x') = \mathbb{E}[d(X, Z)] \leq \\
&\leq \mathbb{E}[d(X, Y) + d(Y, Z)] = \rho(\mu, \nu) + \rho(\nu, \theta),
\end{aligned}$$

which proves the triangle inequality for ρ . Concerning definiteness, if $\mu = \nu$, we have $\rho(\mu, \nu) = 0$ by definition.

Suppose now $\mu \neq \nu$, but $\rho(\mu, \nu) = 0$. Furthermore, take two independent random variables $X \sim \mu, Y \sim \nu$. We then have,

$$0 = \int_{\mathcal{X} \times \mathcal{X}} d(x, x') d\mu \otimes \nu(x, x') = \mathbb{E}[d(X, Y)].$$

From this we conclude $d(X, Y) = 0$ almost surely, which translates to $X = Y$ almost surely. This implies $\mu = \nu$ which is a contradiction. Therefore, whenever $\mu \neq \nu$ we have $\rho(\mu, \nu) > 0$. By contraposition, $\rho(\mu, \nu) = 0$ implies $\mu = \nu$, which concludes the prove of definiteness. \square

B Existence of “barycenters”

In this section we want to discuss the existence of minimizers in the general barycenter updating step of Section 3.1. Thus a metric space (\mathcal{X}, d) and finitely many continuous maps into other metric spaces $\varphi_i : \mathcal{X} \rightarrow \mathcal{X}_i$ are given. We observe the image points $\tilde{x}_i = \varphi_i(x_i)$ for $i = 1, \dots, n$. The question is: under which conditions is

$$\inf\{\sum_{i=1}^n \lambda_i d(\varphi_i^{-1}(\tilde{x}_i), y)^2 : y \in \mathcal{X}\} \quad (16)$$

attained? Here, $\lambda_i > 0$, $\sum_{i=1}^n \lambda_i = 1$ denote some weights. Since in this general formulation the maps φ_i can be very arbitrary, minimizers do not have to exist. However, if one of the maps φ_i is the identity on \mathcal{X} , we can guarantee existence under mild assumptions.

Lemma B.1. *Assume there exists a metrizable topology τ on \mathcal{X} which is weaker than the topology induced by the metric d , and such that*

- i) $\varphi_i^{-1}(\tilde{x}_i)$ is closed in τ for all $i \in \{1, \dots, n\}$,
- ii) balls, i.e. $B_d(y, c) := \{x \in \mathcal{X} : d(x, y) \leq c\}$, are τ compact for all $y \in \mathcal{X}, c > 0$,
- iii) $d(\cdot, \cdot)$ is lower semi continuous w.r.t. the product topology $\tau \times \tau$,
- iv) there exists $j \in \{1, \dots, n\}$ such that $\varphi_j = Id_{\mathcal{X}}$.

Then (16) admits a minimizer.

Proof. Let us first prove that for $d(\varphi_i^{-1}(\tilde{x}_i), y)$ there exists $z \in \varphi_i^{-1}(\tilde{x}_i)$ such that

$$d(\varphi_i^{-1}(\tilde{x}_i), y) = \inf\{d(x, y) : x \in \varphi_i^{-1}(\tilde{x}_i)\} = d(z, y). \quad (17)$$

To this end, take a sequence $\{x_n\}_{n \geq 1} \subset \varphi_i^{-1}(\tilde{x}_i)$ such that

$$d(x_n, y) \searrow \inf\{d(x, y) : x \in \varphi_i^{-1}(\tilde{x}_i)\}.$$

We then set $c := d(x_1, y)$, so $\{x_n\}_{n \geq 1} \subset B_d(y, c)$. Since $B_d(y, c)$ is τ -compact we obtain a subsequence $(x_{n_k})_{k \geq 1}$ such that $x_{n_k} \xrightarrow{\tau} z \in B_d(y, c)$. Since $\{x_{n_k}\}_{k \geq 1} \subset \varphi_i^{-1}(\tilde{x}_i)$ and $\varphi_i^{-1}(\tilde{x}_i)$ is closed in τ , we also have $z \in \varphi_i^{-1}(\tilde{x}_i)$. By lower semicontinuity of d w.r.t. $\tau \times \tau$ we obtain

$$d(z, y) \leq \liminf_{k \rightarrow \infty} d(x_{n_k}, y) = \inf\{d(x, y) : x \in \varphi_i^{-1}(\tilde{x}_i)\},$$

which proves the existence of $z \in \varphi_i^{-1}(\tilde{x}_i)$ such that $d(z, y) = d(\varphi_i^{-1}(\tilde{x}_i), y)$.

Let us next prove that for a set $A = \varphi_i^{-1}(\tilde{x}_i)$ the map $x \mapsto d(A, x)$ is almost lower semi continuous w.r.t. τ . Precisely, let us show that for a sequence $\{x_n\}_{n \geq 1} \subset \mathcal{X}$ such that $x_n \xrightarrow{\tau} x \in \mathcal{X}$ and $d(x_n, x) \leq c$ we have

$$\liminf_{n \rightarrow \infty} d(A, x_n) \geq d(A, x). \quad (18)$$

Without loss of generality, assume the left hand side of (18) is finite, as otherwise there is nothing to prove. Therefore, we may assume the existence of a constant $c' > 0$ such that $d(A, x_n) \leq c'$. By what we already proved, for $n \geq 1$ there exists $z_n \in A$ such that $d(A, x_n) = d(z_n, x_n)$. We then have

$$d(z_n, x) \leq d(z_n, x_n) + d(x_n, x) = d(A, x_n) + d(x_n, x) \leq c' + c.$$

By τ compactness of $B_d(x, c + c')$ and since A is closed in τ , there exists $z \in A \cap B_d(x, c + c')$ such that $z_n \xrightarrow{\tau} z$ up to switching to a subsequence. By lower semi continuity of d w.r.t. $\tau \times \tau$, i.e. assumption iii), we have

$$\liminf_{n \rightarrow \infty} d(A, x_n) = \liminf_{n \rightarrow \infty} d(z_n, x_n) \geq d(z, x) \geq d(A, x),$$

which proves (18).

To prove the existence of minimizers for (16) we set

$$V := \inf\{\sum_{i=1}^n \lambda_i d(\varphi_i^{-1}(\tilde{x}_i), y)^2 : y \in \mathcal{X}\}$$

and take a sequence $\{y_n\}_{n \geq 1} \subset \mathcal{X}$ such that $\sum_{i=1}^n \lambda_i d(\varphi_i^{-1}(\tilde{x}_i), y_n)^2 \searrow V$. We assume without loss of generality that $d(\tilde{x}_j, y_n) \leq 2V$, since $\varphi_j = Id_{\mathcal{X}}$, so we have $\{y_n\}_{n \geq 1} \subset B_d(\tilde{x}_j, 2V)$. By compactness of $B_d(\tilde{x}_j, 2V)$ w.r.t. τ we have the existence of a subsequence $\{y_{n_k}\}_{k \geq 1} \subset B_d(\tilde{x}_j, 2V)$ such that $y_{n_k} \xrightarrow{\tau} y^*$ for some $y^* \in B_d(\tilde{x}_j, 2V)$. We can now use the proved property (18) to obtain

$$V = \liminf_{k \rightarrow \infty} \sum_{i=1}^n \lambda_i d(\varphi_i^{-1}(\tilde{x}_i), y_{n_k})^2 \geq \sum_{i=1}^n \lambda_i d(\varphi_i^{-1}(\tilde{x}_i), y^*)^2.$$

Therefore, we have found $y^* \in \mathcal{X}$ such that

$$\sum_{i=1}^n \lambda_i d(\varphi_i^{-1}(\tilde{x}_i), y^*)^2 = \inf\{\sum_{i=1}^n \lambda_i d(\varphi_i^{-1}(\tilde{x}_i), y)^2 : y \in \mathcal{X}\},$$

finishing the proof. □

In the proof we also showed the existence of minimizers in (7), i.e. points used for “filling up”, under the same assumptions. This is precisely (17).

Let us discuss the assumptions of Lemma B.1. Note that assumptions i), ii) and iii) are fulfilled in our two important examples of $\mathcal{X} = \mathbb{R}^d$ and $\mathcal{X} = \mathcal{P}_2(\mathbb{R}^d)$. Indeed, for $\mathcal{X} = \mathbb{R}^d$ this is clear, whereas for $\mathcal{X} = \mathcal{P}_2(\mathbb{R}^d)$ the role of the metric τ is played by the metric induced by weak convergence of probability measures. For details check Chapter 7, *The Metric Side of Optimal Transportation*, in [24]. As for Assumption iv), this is not a big restriction in our two examples, since this basically (in practical terms) means that for each cluster we need at least one fully observed data point. Even if this is not the case, we can still include the old barycenter with a small weight to enforce the existence of minimizers, as done in Section 5 and Algorithm 1. This also helps numerically, as otherwise infinite solutions for the generalized mean in the Euclidean case, and the generalized Wasserstein barycenter in the case of probability measures, may exist.

C Further simulation experiments

Let us discuss some additional simulation experiments to evaluate the proposed method. In Section 7.1 we evaluated our method as an imputation method, but without the imputation step it can also be viewed as a clustering algorithm. In the case of Euclidean data it can be viewed as a clustering algorithm for points with missing values. Thus, we can also compare it to classical k -means after imputing the missing data with the more standard methods for imputing missing values we used in 7.1. To recall, the standard methods for imputing missing values we consider are mean imputation, median imputation, multiple imputation, K -nearest neighbour imputation, and imputation through linear regression. However, in the case of clustering Euclidean points with missing data we also consider the k -pod method which was introduced in [4]. These authors consider the same loss function as we do in Example 3.1, i.e. (9), however they propose a different algorithm. To perform the comparison, we use the same simulation from Section 7.1, i.e. sampling from a Gaussian Mixture Model with $k = 5$ clusters in 10 dimensions and different settings of missing values. We use precisely the same simulated data, and for each of those we apply classical k -means to cluster the points after imputing them with a standard imputation method. For our proposed method, i.e. NA k -means, and for the k -pod method, no imputation step is needed as these directly cluster points with missing values. To evaluate the clustering results we use the so-called Rand index, also known as the Rand score, introduced in [19]. It compares a baseline clustering (in our case, the labels from the normal distribution of the Gaussian Mixture Model to which an observation belongs) to a clustering of the data using an imputation and/or clustering algorithm. The Rand score takes values in $[0, 1]$, with 0 meaning nothing is clustered the same, and 1 meaning the clusterings coincide except for renaming of clusters. In particular, a higher Rand score corresponds to a better clustering method. In Table 2 we report the corresponding Rand scores \pm standard errors for these simulations.

Table 2: Rand scores for Euclidean simulation.

settings	NA k -means	mean imp.	median imp.	multiple imp.	KNN	LR	k -pod
β 0.0	0.82 ± 0.008	0.792 ± 0.008	0.786 ± 0.009	0.823 ± 0.007	0.747 ± 0.006	0.822 ± 0.008	0.713 ± 0.008
β 0.1	0.817 ± 0.008	0.799 ± 0.008	0.783 ± 0.009	0.824 ± 0.008	0.744 ± 0.006	0.823 ± 0.008	0.711 ± 0.008
β 0.2	0.823 ± 0.008	0.799 ± 0.009	0.778 ± 0.009	0.82 ± 0.008	0.743 ± 0.006	0.82 ± 0.007	0.7 ± 0.009
β 0.3	0.828 ± 0.008	0.797 ± 0.008	0.777 ± 0.009	0.83 ± 0.008	0.743 ± 0.006	0.822 ± 0.008	0.701 ± 0.009
β 0.4	0.829 ± 0.008	0.799 ± 0.008	0.78 ± 0.009	0.824 ± 0.007	0.744 ± 0.006	0.824 ± 0.007	0.698 ± 0.009
β 0.5	0.826 ± 0.008	0.8 ± 0.008	0.774 ± 0.009	0.828 ± 0.007	0.743 ± 0.006	0.826 ± 0.007	0.698 ± 0.008
β 0.6	0.822 ± 0.008	0.8 ± 0.008	0.772 ± 0.009	0.825 ± 0.007	0.742 ± 0.006	0.819 ± 0.007	0.691 ± 0.008
β 0.7	0.819 ± 0.008	0.796 ± 0.008	0.776 ± 0.009	0.818 ± 0.007	0.742 ± 0.006	0.818 ± 0.007	0.688 ± 0.008
β 0.8	0.813 ± 0.008	0.795 ± 0.008	0.778 ± 0.009	0.826 ± 0.007	0.737 ± 0.005	0.814 ± 0.007	0.685 ± 0.008
β 0.9	0.804 ± 0.008	0.793 ± 0.009	0.777 ± 0.009	0.819 ± 0.007	0.734 ± 0.006	0.811 ± 0.007	0.683 ± 0.008
β 1.0	0.8 ± 0.008	0.788 ± 0.008	0.769 ± 0.009	0.814 ± 0.007	0.729 ± 0.006	0.808 ± 0.007	0.682 ± 0.008

We can see that the introduced method NA k -means either performs best or lies within the standard error of the best performing method for each setting of missing values when considering it solely as clustering algorithm for points with missing data. The naive approaches of mean and median imputation are outperformed.

Generalizing our results In our simulation scenarios we have two parameters concerning the creation of missing values in the data. One is the *quantile* below which we remove values for each dimension. The second one is h_j , which is a uniform random variable per cluster which is then correlated with the cluster size to decide what percentage of missing values per cluster is generated additionally. In Section 7.1 and in the previous paragraph, the *quantile* is chosen as 25% quantile and we set $h_j \sim U(0, 2/k)$ where k is the number of clusters, i.e. 5, in the considered simulations. Essentially, this fixes the share of missing values per simulated dataset. However, we want to vary these two parameters in order to show the robustness of our method. In Table 3 and Table 4 we report the corresponding results analogous to Table 1 and Table 2, respectively, when choosing 5 different choices of the *quantile* and h_j .

Table 3: Gromov Wasserstein distance for Euclidean simulation with varying share of missing values.

settings	NA k -means.	NA k -means-m	mean imp.	median imp.	multiple imp.	KNN	LR
10% quantile and $h_j \sim U(0, 0.1)$							
β 0.0	0.742 ± 0.04	0.804 ± 0.042	1.48 ± 0.05	1.466 ± 0.046	0.916 ± 0.046	0.874 ± 0.043	0.913 ± 0.048
β 0.25	0.747 ± 0.039	0.864 ± 0.046	1.763 ± 0.05	1.724 ± 0.051	0.975 ± 0.047	1.036 ± 0.045	0.939 ± 0.046
β 0.5	0.828 ± 0.045	0.928 ± 0.043	2.058 ± 0.054	2.032 ± 0.057	1.009 ± 0.047	1.187 ± 0.048	0.976 ± 0.048
β 0.75	0.967 ± 0.046	0.982 ± 0.044	2.458 ± 0.066	2.414 ± 0.065	1.123 ± 0.046	1.406 ± 0.047	1.1 ± 0.047
β 1.0	1.099 ± 0.052	1.163 ± 0.053	2.94 ± 0.082	2.861 ± 0.082	1.273 ± 0.056	1.645 ± 0.053	1.324 ± 0.06
20% quantile and $h_j \sim U(0, 0.2)$							
β 0.0	1.622 ± 0.063	1.742 ± 0.069	3.247 ± 0.078	3.182 ± 0.077	1.777 ± 0.068	2.233 ± 0.074	1.795 ± 0.068
β 0.25	1.711 ± 0.064	1.798 ± 0.064	3.542 ± 0.082	3.465 ± 0.081	1.904 ± 0.076	2.444 ± 0.075	1.947 ± 0.077
β 0.5	1.816 ± 0.067	1.875 ± 0.069	3.863 ± 0.083	3.781 ± 0.086	2.053 ± 0.074	2.63 ± 0.077	2.126 ± 0.076
β 0.75	1.931 ± 0.066	1.985 ± 0.065	4.237 ± 0.093	4.154 ± 0.095	2.329 ± 0.082	2.765 ± 0.071	2.403 ± 0.085
β 1.0	2.062 ± 0.064	2.073 ± 0.064	4.629 ± 0.103	4.549 ± 0.105	2.612 ± 0.087	2.929 ± 0.072	2.695 ± 0.092
30% quantile and $h_j \sim U(0, 0.3)$							
β 0.0	2.866 ± 0.084	2.939 ± 0.087	5.274 ± 0.105	5.172 ± 0.107	3.33 ± 0.1	3.79 ± 0.091	3.413 ± 0.097
β 0.25	3.035 ± 0.082	3.062 ± 0.082	5.518 ± 0.106	5.409 ± 0.107	3.551 ± 0.1	3.913 ± 0.092	3.658 ± 0.099
β 0.5	3.093 ± 0.079	3.105 ± 0.08	5.764 ± 0.111	5.665 ± 0.112	3.773 ± 0.102	4.034 ± 0.096	3.871 ± 0.102
β 0.75	3.183 ± 0.082	3.205 ± 0.08	6.067 ± 0.117	5.971 ± 0.118	4.043 ± 0.109	4.138 ± 0.095	4.179 ± 0.107
β 1.0	3.251 ± 0.081	3.264 ± 0.08	6.349 ± 0.121	6.26 ± 0.124	4.342 ± 0.113	4.203 ± 0.097	4.462 ± 0.114
40% quantile and $h_j \sim U(0, 0.4)$							
β 0.0	4.341 ± 0.104	4.354 ± 0.105	7.536 ± 0.133	7.431 ± 0.138	5.633 ± 0.128	5.333 ± 0.117	5.725 ± 0.129
β 0.25	4.398 ± 0.102	4.407 ± 0.102	7.671 ± 0.131	7.551 ± 0.134	5.791 ± 0.127	5.395 ± 0.115	5.86 ± 0.129
β 0.5	4.487 ± 0.106	4.489 ± 0.106	7.82 ± 0.131	7.713 ± 0.135	5.901 ± 0.128	5.44 ± 0.117	6.055 ± 0.13
β 0.75	4.531 ± 0.104	4.536 ± 0.105	8.021 ± 0.136	7.904 ± 0.14	6.058 ± 0.126	5.486 ± 0.119	6.275 ± 0.134
β 1.0	4.528 ± 0.1	4.531 ± 0.1	8.206 ± 0.139	8.102 ± 0.144	6.284 ± 0.135	5.513 ± 0.12	6.46 ± 0.141
50% quantile and $h_j \sim U(0, 0.5)$							
β 0.0	5.59 ± 0.129	5.6 ± 0.129	9.895 ± 0.155	9.802 ± 0.16	8.012 ± 0.15	6.883 ± 0.142	8.12 ± 0.152
β 0.25	5.693 ± 0.123	5.701 ± 0.124	9.924 ± 0.151	9.823 ± 0.156	8.038 ± 0.15	6.861 ± 0.142	8.198 ± 0.146
β 0.5	5.677 ± 0.127	5.677 ± 0.127	9.982 ± 0.148	9.872 ± 0.151	8.106 ± 0.144	6.884 ± 0.138	8.26 ± 0.145
β 0.75	5.72 ± 0.128	5.72 ± 0.128	10.035 ± 0.148	9.938 ± 0.152	8.159 ± 0.151	6.88 ± 0.14	8.307 ± 0.147
β 1.0	5.705 ± 0.125	5.705 ± 0.125	10.09 ± 0.148	10.0 ± 0.151	8.252 ± 0.152	6.898 ± 0.14	8.399 ± 0.153

From Table 3 we conclude that the proposed method NA k -means outperforms all the other considered methods independent of the parameters needed for the missing value generation, when considering the Gromov Wasserstein distance as evaluation measure.

Table 4: Rand scores for Euclidean simulation with various shares of missing values.

settings	NA k -means	mean imp.	median imp.	multiple imp.	KNN	LR	k -pod
10% quantile and $h_j \sim U(0, 0.1)$							
β 0.0	0.916 ± 0.009	0.911 ± 0.008	0.903 ± 0.007	0.923 ± 0.008	0.919 ± 0.008	0.922 ± 0.008	0.863 ± 0.008
β 0.25	0.907 ± 0.009	0.908 ± 0.007	0.902 ± 0.007	0.919 ± 0.007	0.92 ± 0.007	0.926 ± 0.007	0.833 ± 0.008
β 0.5	0.913 ± 0.007	0.901 ± 0.007	0.891 ± 0.008	0.923 ± 0.007	0.909 ± 0.007	0.916 ± 0.007	0.817 ± 0.009
β 0.75	0.906 ± 0.007	0.891 ± 0.008	0.882 ± 0.007	0.903 ± 0.007	0.891 ± 0.007	0.904 ± 0.007	0.784 ± 0.008
β 1.0	0.883 ± 0.008	0.875 ± 0.008	0.854 ± 0.008	0.899 ± 0.007	0.857 ± 0.007	0.902 ± 0.006	0.75 ± 0.008
20% quantile and $h_j \sim U(0, 0.2)$							
β 0.0	0.85 ± 0.008	0.856 ± 0.009	0.856 ± 0.008	0.877 ± 0.008	0.834 ± 0.008	0.877 ± 0.008	0.783 ± 0.009
β 0.25	0.851 ± 0.008	0.849 ± 0.008	0.855 ± 0.007	0.878 ± 0.007	0.818 ± 0.007	0.871 ± 0.008	0.759 ± 0.009
β 0.5	0.861 ± 0.007	0.846 ± 0.009	0.842 ± 0.007	0.869 ± 0.007	0.805 ± 0.007	0.867 ± 0.007	0.74 ± 0.008
β 0.75	0.85 ± 0.008	0.84 ± 0.008	0.831 ± 0.008	0.857 ± 0.007	0.792 ± 0.006	0.858 ± 0.007	0.718 ± 0.008
β 1.0	0.832 ± 0.008	0.823 ± 0.008	0.806 ± 0.008	0.845 ± 0.007	0.767 ± 0.007	0.845 ± 0.007	0.696 ± 0.007
30% quantile and $h_j \sim U(0, 0.3)$							
β 0.0	0.804 ± 0.008	0.787 ± 0.009	0.78 ± 0.009	0.813 ± 0.008	0.727 ± 0.006	0.811 ± 0.008	0.703 ± 0.008
β 0.25	0.797 ± 0.007	0.783 ± 0.008	0.777 ± 0.009	0.799 ± 0.009	0.716 ± 0.006	0.801 ± 0.008	0.692 ± 0.008
β 0.5	0.79 ± 0.007	0.774 ± 0.008	0.761 ± 0.009	0.799 ± 0.007	0.717 ± 0.005	0.794 ± 0.006	0.678 ± 0.008
β 0.75	0.774 ± 0.007	0.775 ± 0.008	0.758 ± 0.009	0.789 ± 0.007	0.711 ± 0.005	0.792 ± 0.006	0.666 ± 0.008
β 1.0	0.762 ± 0.008	0.769 ± 0.007	0.743 ± 0.008	0.783 ± 0.007	0.697 ± 0.005	0.773 ± 0.008	0.662 ± 0.007
40% quantile and $h_j \sim U(0, 0.4)$							
β 0.0	0.729 ± 0.006	0.708 ± 0.009	0.698 ± 0.01	0.731 ± 0.006	0.665 ± 0.005	0.728 ± 0.007	0.636 ± 0.007
β 0.25	0.723 ± 0.006	0.709 ± 0.008	0.7 ± 0.01	0.728 ± 0.006	0.666 ± 0.005	0.729 ± 0.006	0.629 ± 0.008
β 0.5	0.713 ± 0.006	0.712 ± 0.008	0.695 ± 0.01	0.732 ± 0.007	0.665 ± 0.005	0.723 ± 0.006	0.628 ± 0.007
β 0.75	0.702 ± 0.007	0.704 ± 0.009	0.7 ± 0.009	0.73 ± 0.007	0.662 ± 0.004	0.718 ± 0.007	0.625 ± 0.007
β 1.0	0.695 ± 0.007	0.705 ± 0.008	0.696 ± 0.009	0.728 ± 0.006	0.659 ± 0.004	0.715 ± 0.006	0.626 ± 0.007
50% quantile and $h_j \sim U(0, 0.5)$							
β 0.0	0.661 ± 0.005	0.632 ± 0.009	0.627 ± 0.009	0.663 ± 0.006	0.64 ± 0.004	0.655 ± 0.006	0.599 ± 0.008
β 0.25	0.658 ± 0.005	0.636 ± 0.01	0.633 ± 0.009	0.672 ± 0.006	0.637 ± 0.004	0.653 ± 0.006	0.598 ± 0.007
β 0.5	0.656 ± 0.006	0.641 ± 0.01	0.634 ± 0.009	0.674 ± 0.007	0.635 ± 0.004	0.657 ± 0.006	0.6 ± 0.007
β 0.75	0.653 ± 0.005	0.646 ± 0.009	0.635 ± 0.009	0.677 ± 0.006	0.638 ± 0.004	0.659 ± 0.006	0.601 ± 0.007
β 1.0	0.649 ± 0.006	0.643 ± 0.009	0.644 ± 0.008	0.683 ± 0.006	0.636 ± 0.004	0.659 ± 0.006	0.601 ± 0.007

In Table 4 we can see that also w.r.t. the Rand score, when NA k -means is solely viewed as clustering procedure for Euclidean points with missing values, it competes with the best performing imputation + clustering techniques.

D An example highlighting NA k -means strength

We give a toy example highlighting a scenario in which the proposed method works well, and which hopefully helps building some intuition. Therefore, we work in dimension 2 and consider Euclidean points with missing values. In particular, we consider three clusters, the first and third one with 200 points each and the second one with 100 points, coming from the three Gaussians

$$\mathcal{N}\left(\begin{pmatrix} 3 \\ 3 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}\right), \quad \mathcal{N}\left(\begin{pmatrix} 0 \\ -1.5 \end{pmatrix}, \begin{pmatrix} 1 & 0.6 \\ 0.6 & 1 \end{pmatrix}\right), \quad \mathcal{N}\left(\begin{pmatrix} -3 \\ 3 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}\right).$$

In order to model *structural missingness of data*, we assume that we only observe the y coordinate of the points if it is positive, so missing values are created for non-positive y coordinates. This can be seen in Figure 3.




Figure 3: (left): The true points with true labels. (right) The observed points with the true labels.

In order to cluster in this situation, most methods for imputation lead to a bias by shifting up the points with the unobserved y coordinates. Especially, as in this case there is essentially only one cluster with points having negative y coordinates and they are the majority of that cluster. NA k -means however, does not introduce this bias and therefore recovers the true cluster membership much better. The corresponding rand scores in this example are in Table 5.

Table 5: Rand scores.

NA k -means	mean imp.	median imp.	multiple imp.	KNN	LR	k -pod
0.963	0.927	0.925	0.927	0.928	0.927	0.904

In Figure 4 below we present the corresponding observed clusterings, i.e. we plot the true points and color them with the labels from the corresponding clustering method.




Figure 4: (first row, left to right): NA k -means, mean imputation, median imputation, multiple imputation. (second row, left to right): KNN, LR, k -pod. The reader is encouraged to zoom in.

We can see that most methods solely base their clustering on the values of the x -coordinate. The reason is that, by imputing the points coming from the turquoise cluster upwards, these points end up having y -values similar to the y -values of the points with no missing values. The clustering coming from NA k -means recovers the original cluster membership of the points the best.