Depth Dependence of μP Learning Rates in ReLU MLPs

Samy Jelassi¹^{*}, Boris Hanin¹, Ziwei Ji², Sashank J. Reddi², Srinadh Bhojanapalli², Sanjiv Kumar²

> ¹Princeton University ²Google Research, NYC

Abstract

In this short note we consider random fully connected ReLU networks of width n and depth L equipped with a mean-field weight initialization. Our purpose is to study the dependence on n and L of the maximal update (μ P) learning rate, the largest learning rate for which the mean squared change in pre-activations after one step of gradient descent remains uniformly bounded at large n, L. As in prior work on μ P [9], we find that this maximal update learning rate is independent of n for all but the first and last layer weights. However, we find that it has a non-trivial dependence of L, scaling like $L^{-3/2}$.

1 Introduction

Using a neural network requires many choices. Even after fixing an architecture, one must still specify initialization scheme, learning rate (schedule), batch size, data augmentation, regularization strength, and so on. Moreover, model performance is often highly sensitive to the setting of these hyperparameters, and yet exhaustive grid search type approaches are computationally expensive. It is therefore important to develop theoretically grounded principles for reducing the cost of hyperparameter tuning. In this short note we focus specifically on the question of how to select learning rates in a principled way. More precisely, our purpose is to generalize the *maximal update* (μ P) approach of [9] to setting learning rates to take into account network depth.

1.1 Overview of μP Approach to Learning Rates

We study learning rates in the simple setting of depth L fully connected neural networks with ReLU activations and a uniform value n for the input dimension and the hidden layers widths.¹ In such a network, by definition, each input $x \in \mathbb{R}^n$ produces an output $z^{(L+1)}(x) \in \mathbb{R}^n$ through a sequence of pre-activations $z^{(\ell)}(x) \in \mathbb{R}^n$ given by

$$z^{(\ell+1)}(x) = \begin{cases} W^{(\ell+1)}\sigma\left(z^{(\ell)}(x)\right), & \ell \ge 1\\ W^{(1)}x, & \ell = 0 \end{cases}, \quad \sigma(t) := \max\left\{0, t\right\}.$$
(1.1)

^{*}Work done while interning at Google NYC.

 $^{^{1}}$ Our computations readily generalize to the case of variable layer widths. Indeed, we carry out the proof of Theorem 1.1 in this context.

Selecting learning rates cannot be done independently of an initialization scheme. As in [9], we draw random weights for the network (1.1) from the so-called mean-field initialization

$$W_{ij}^{(\ell)} \sim \begin{cases} \mathcal{N}(0, 2/n), & \ell = 1, \dots, L\\ \mathcal{N}(0, 1/n^2), & \ell = L+1 \end{cases}$$
(1.2)

The factor of two in variance of hidden layer weights corresponds to the well-known He initialization [3], which ensures that the expected squared activations neither grow nor decay with depth:

$$\mathbb{E}\left[\left|\left|z^{(\ell)}(x)\right|\right|^{2}\right] = \left|\left|x\right|\right|^{2}, \quad \forall \ell = 1, \dots, L.$$
(1.3)

The much smaller variance of weights in the final layer distinguishes the initialization scheme (1.2) from the so-called NTK initialization [4]. The difference is twofold. First, when n is large the network output $z^{(L+1)}(x)$ is close to zero. However, crucially, the parameter gradients $\nabla_{\theta} z^{(L+1)}(x)$ are remain non-zero. Second, even in the infinite width limit $n \to \infty$ networks trained by gradient descent are capable of feature learning [6, 7, 8, 9]. This is in contrast to the setting where the final layer weight variance scales like 1/n, which corresponds to the kernel regime in which neural networks trained by SGD with a small learning rate on a mean squared error loss converge to linear models and hence cannot learn data-adaptive features [1, 4, 5].

A key contribution of [9] is that the initialization (1.2) not only leads to feature learning at large n but also allows for zero-shot learning rate transfer with respect to variable width. This means that, empirically, for a fixed depth L the learning rate at small n that leads to the smallest training loss after one epoch is close to constant as one varies n^2 . Hence, in practice, one may do logarithmic grid search for good learning rates in relatively small models (with small n) and then simply re-use the best learning rate for wider networks.

1.2 Main Result: Extending the μ P Heuristic to Deeper Networks

Instead of studying directly the training loss after one epoch [9] introduces what we will refer to here as the maximal update heuristic, which says that a good learning rate is one that corresponds to the largest change in hidden layer pre-activations after one step of GD that does not lead to a divergence at large n. More precisely, the relation (1.3) shows that *i*-th neuron pre-activation in layer ℓ corresponding to an input x that satisfies

$$\mathbb{E}\left[\left(z_{i}^{(\ell)}(x)\right)^{2}\right] = \frac{1}{n} ||x||^{2}, \qquad i = 1, \dots, n, \quad \ell = 1, \dots, L,$$

with the average being over initialization. To study the change in neuron pre-activations under GD we consider a batch $\mathcal{B} = \{(x, y)\}$ size of 1 and the associated mean-squared error

$$\mathcal{L}_{\mathcal{B}}(\theta) := \frac{1}{2} \left| \left| z^{(L+1)}(x;\theta) - y \right| \right|^2,$$

where we've emphasized the dependence of the network output $z^{(L+1)}(x;\theta)$ on the network weights θ . Let us denote by

$$\Delta^{\mathcal{B}} z_i^{(\ell)}(x) = \text{change in } z_i^{(\ell)}(x) \text{ after first step of GD on } \mathcal{L}_{\mathcal{B}}.$$

²Strictly speaking, the μ P prescription gives *n*-dependent learning rates for weights in the first and last layer and *n*-independent learning rates for weights in other layers (see Table 3 of [9]).

The maximal update heuristic then asks that we set the learning rate η so that

$$\mu P$$
 learning rate $\eta^* :=$ learning rate for which $\mathbb{E}\left[\left(\Delta^{\mathcal{B}} z_i^{(\ell)}(x)\right)^2\right] = 1,$ (1.4)

where the average is over initialization. A priori, η^* depends on both network *n* width and depth *L*. The article [9] shows that η^* does not depend on *n* and hence can be estimated accurately at small *n*. In this article, we take up the question of how η^* depends on depth. The following theorem shows that η^* is not depth-independent:

Theorem 1.1. For each $c_1 > 0$ there exists $c_2, c_3 > 0$ with the following property. Fix a network width n and depth L so that $L/n < c_1$. Then,

$$\sup_{n\geq 1} \left| \mathbb{E}\left[\left(\Delta^{\mathcal{B}} z_i^{(\ell)}(x) \right)^2 \right] - c_2 \eta^2 \ell^3 \right| \le c_3 \eta^2 \ell^2, \tag{1.5}$$

where $\mathcal{B} = \{(x, y)\}$ is any batch of size one consisting of a normalized datapoint (x, y) sampled independent of network weights and biases with:

$$\mathbb{E}\left[\frac{1}{n}\left|\left|x\right|\right|^{2}\right] = 1, \qquad \mathbb{E}\left[\left|\left|y\right|\right|^{2}\right] = 1.$$

Theorem 1.1 shows that the μP heuristic (1.4) dictates that

$$\eta^*(L) = \operatorname{const} \cdot L^{-3/2}.$$

2 Proof of Theorem 1.1

2.1 Notation and Problem Setting

We prove a slightly more general result than Theorem 1.1 in two senses. First, we allow for variable widths:

$$n_{\ell} =$$
width of layer $\ell = 0, \ldots, L+1$

Second, we will also allow for parameter-dependent learning rates:

$$\eta_{\mu} =$$
 learning rate used for parameter μ .

At the end we will restrict to the case where $\eta_{\mu} = \eta$ is independent of μ . Moreover, in order to state our proof most efficiently, we introduce some notation. Namely, we will write $x_{\alpha} \in \mathbb{R}^{n_0}$ for the network input at which we study both the forward and backward pass and will denote for brevity

$$z_{i;\alpha}^{(\ell)} := z_i^{(\ell)}(x_\alpha), \qquad z_\alpha^{(\ell)} := z^{(\ell)}(x_\alpha).$$

Thus, the batch loss $\mathcal{L}_{\mathcal{B}}$ we consider is

$$\frac{1}{2} \left\| z_{\alpha}^{(L+1)} - y_{\alpha} \right\|^2.$$

Further, we abbreviate

$$\Delta z_{i;\alpha}^{(\ell)} := \Delta^{\mathcal{B}} z_{i;\alpha}^{(\ell)}.$$

With this notation, the forward pass now takes the form

$$z_{i;\alpha}^{(\ell+1)} = \begin{cases} \sum_{j=1}^{n_0} W_{ij}^{(1)} x_{j;\alpha}, & \ell = 0\\ \sum_{j=1}^{n_{\ell-1}} W_{ij}^{(\ell)} \sigma\left(z_{j;\alpha}^{(\ell)}\right), & \ell = 1, \dots, L \end{cases}$$

and the initialization scheme is

$$W_{ij}^{(\ell+1)} \sim \begin{cases} \mathcal{N}\left(0, \frac{1}{n_L^2}\right), & \ell = L\\ \mathcal{N}\left(0, \frac{2}{n_\ell}\right), & \ell = 0, \dots, L-1 \end{cases}$$

2.2 Proof Details

We begin with the following Lemma.

Lemma 2.1. For any depth $\ell \leq L$, the pre-activation change satisfies

$$\mathbb{E}[(\Delta z_{i;\alpha}^{(\ell)})^2] = A^{(\ell)} + B^{(\ell)},$$

where

$$A^{(\ell)} := \mathbb{E}\left[\frac{1}{n_L^2} \sum_{\mu_1,\mu_2 \le \ell} \eta_{\mu_1} \eta_{\mu_2} \partial_{\mu_1} z_{1;\alpha}^{(\ell)} \partial_{\mu_2} z_{1;\alpha}^{(\ell)} \right] \times \frac{1}{n_L^2} \sum_{j_1,j_2=1}^{n_L} \left\{ \partial_{\mu_1} z_{j_1;\alpha}^{(L)} \partial_{\mu_2} z_{j_1;\alpha}^{(L)} \left(z_{j_2;\alpha}^{(L)} \right)^2 + 2z_{j_1;\alpha}^{(L)} \partial_{\mu_1} z_{j_1;\alpha}^{(L)} z_{j_2;\alpha}^{(L)} \partial_{\mu_2} z_{j_2;\alpha}^{(L)} \right\},$$

$$B^{(\ell)} := \mathbb{E}\left[\frac{1}{n_L} \sum_{\mu_1,\mu_2 \le \ell} \eta_{\mu_1} \eta_{\mu_2} \partial_{\mu_1} z_{1;\alpha}^{(\ell)} \partial_{\mu_2} z_{1;\alpha}^{(\ell)} \frac{1}{n_L} \sum_{j=1}^{n_L} \partial_{\mu_1} z_{j;\alpha}^{(L)} \partial_{\mu_2} z_{j;\alpha}^{(L)} \right].$$

$$(2.1)$$

Proof of Lemma 2.1. We first expand $\Delta z_{i;\alpha}^{(\ell)}$ by applying the chain rule:

$$\Delta z_{i;\alpha}^{(\ell)} = \sum_{\mu \le \ell} \cdot \partial_{\mu} z_{i;\alpha}^{(\ell)} \Delta \mu, \qquad (2.3)$$

where $\Delta \mu$ is the change in μ after one step of GD. The SGD update satisfies:

$$\Delta \mu = -\eta_{\mu} \partial_{\mu} \left\{ \frac{1}{2} \left\| \left| z_{\alpha}^{(L+1)} - y_{\alpha} \right\| \right|^{2} \right\} = -\eta_{\mu} \sum_{k=1}^{n_{L+1}} \partial_{\mu} z_{k;\alpha}^{(L+1)} \left(z_{k;\alpha}^{(L+1)} - y_{k;\alpha} \right).$$
(2.4)

We now combine (2.3) and (2.4) to obtain:

$$\Delta z_{i;\alpha}^{(\ell)} = \sum_{\mu \le \ell} \sum_{k=1}^{n_{L+1}} \eta_{\mu} \partial_{\mu} z_{i;\alpha}^{(\ell)} \partial_{\mu} z_{k;\alpha}^{(L+1)} \left(y_{k;\alpha} - z_{k;\alpha}^{(L+1)} \right).$$
(2.5)

Using (2.5), we obtain

$$\mathbb{E}\left[\left(\Delta z_{i;\alpha}^{(\ell)}\right)^{2}\right] = \mathbb{E}\left[\left(\sum_{\mu \leq \ell} \eta_{\mu} \partial_{\mu} z_{1;\alpha}^{(\ell)} \partial_{\mu} z_{1;\alpha}^{(L+1)} \left(z_{1;\alpha}^{(L+1)} - y_{1;\alpha}\right)\right)^{2}\right] \\ = \mathbb{E}\left[\sum_{\mu_{1},\mu_{2} \leq \ell} \eta_{\mu_{1}} \eta_{\mu_{2}} \partial_{\mu_{1}} z_{1;\alpha}^{(\ell)} \partial_{\mu_{2}} z_{1;\alpha}^{(\ell)} \partial_{\mu_{1}} z_{1;\alpha}^{(L+1)} \partial_{\mu_{2}} z_{1;\alpha}^{(L+1)} \mathbb{E}_{y}\left[\left(z_{1;\alpha}^{(L+1)} - y_{1;\alpha}\right)^{2}\right]\right].$$

$$(2.6)$$

Given the distribution of $z_{1;\alpha}^{(L+1)}$ and y, we have

$$\mathbb{E}_{y}\left[\left(z_{1;\alpha}^{(L+1)} - y\right)^{2}\right] = (z_{1;\alpha}^{(L+1)})^{2} + 1$$
(2.7)

We plug (2.7) in (2.6) and obtain

$$\mathbb{E}[(\Delta z_{i;\alpha}^{(\ell)})^2] = A^{(\ell)} + B^{(\ell)}, \qquad (2.8)$$

where

$$A^{(\ell)} = \mathbb{E}\left[\sum_{\mu_1,\mu_2 \le \ell} \eta_{\mu_1} \eta_{\mu_2} \partial_{\mu_1} z_{1;\alpha}^{(\ell)} \partial_{\mu_2} z_{1;\alpha}^{(\ell)} \partial_{\mu_1} z_{1;\alpha}^{(L+1)} \partial_{\mu_2} z_{1;\alpha}^{(L+1)} \left(z_{1;\alpha}^{(L+1)}\right)^2\right]$$
(2.9)

$$B^{(\ell)} = \mathbb{E}\left[\sum_{\mu_1,\mu_2 \le \ell} \eta_{\mu_1} \eta_{\mu_2} \partial_{\mu_1} z_{1;\alpha}^{(\ell)} \partial_{\mu_2} z_{1;\alpha}^{(\ell)} \partial_{\mu_1} z_{1;\alpha}^{(L+1)} \partial_{\mu_2} z_{1;\alpha}^{(L+1)}\right].$$
(2.10)

We integrate out the weights in layer L + 1 in (2.9) and (2.10) which yields the stated result.

Lemma 2.2. For any depth $\ell \leq L$, the constant $A^{(\ell)}$ in Lemma 2.1 satisfies $A^{(\ell)} = O(n^{-1})$. Proof of Lemma 2.2. The result is obtained essentially the same analysis at we apply to $B^{(\ell)}$ below combined with the observation that there is an extra $1/n_L$ in front of $A^{(\ell)}$ compared with $B^{(\ell)}$.

Lemma 2.2 indicates that we may neglect the contribution of $A^{(\ell)}$ in Lemma 2.1. We now focus on obtaining a recursive description for $B^{(\ell)}$.

Lemma 2.3. For any depth $\ell \leq L$, the constant $B^{(\ell)}$ in Lemma 2.1 satisfies

$$B^{(\ell)} = \mathbb{E}\left[\frac{1}{n_L}\sum_{\mu_1,\mu_2 \le \ell} \eta_{\mu_1}\eta_{\mu_2}\frac{1}{n_\ell^2}\sum_{j_1,j_2=1}^{n_\ell} \partial_{\mu_1} z_{j_1;\alpha}^{(\ell)} \partial_{\mu_2} z_{j_1;\alpha}^{(\ell)} \partial_{\mu_1} z_{j_2;\alpha}^{(\ell)} \partial_{\mu_2} z_{j_2;\alpha}^{(\ell)}\right].$$
 (2.11)

Proof of Lemma 2.3. The idea of this proof is to condition on $z_{\alpha}^{(\ell)}$ and integrate out weights in layers $\ell + 1, \ldots, L$ to obtain

$$\mathbb{E}\left[\frac{1}{n_L}\sum_{j=1}^{n_L}\partial_{\mu_1} z_{j;\alpha}^{(L)}\partial_{\mu_2} z_{j;\alpha}^{(L)} \mid z_{\alpha}^{(\ell)}\right] = \frac{1}{n_\ell}\sum_{j=1}^{n_\ell}\partial_{\mu_1} z_{j;\alpha}^{(\ell)}\partial_{\mu_2} z_{j;\alpha}^{(\ell)}.$$
(2.12)

This will yield the result once we plug (2.12) into (2.2). To see (2.12), we proceed by induction on L starting with $\ell = L$. In this case, the result is trivial. Suppose now $\ell < L$. Then we have

$$\mathbb{E}\left[\frac{1}{n_{L}}\sum_{j=1}^{n_{L}}\partial_{\mu_{1}}z_{j;\alpha}^{(L)}\partial_{\mu_{2}}z_{j;\alpha}^{(L)} \mid z_{\alpha}^{(\ell)}\right]$$

$$=\mathbb{E}\left[\frac{1}{n_{L}}\sum_{j=1}^{n_{L}}\sum_{k_{1},k_{2}=1}^{n_{L-1}}W_{jk_{1}}^{(L)}W_{jk_{2}}^{(L)}\partial_{\mu_{1}}\sigma\left(z_{k_{1};\alpha}^{(L-1)}\right)\partial_{\mu_{2}}\sigma\left(z_{k_{2};\alpha}^{(L-1)}\right) \mid z_{\alpha}^{(\ell)}\right]$$

$$=\mathbb{E}\left[\frac{1}{n_{L}}\sum_{j=1}^{n_{L}}\frac{2}{n_{L-1}}\sum_{k=1}^{n_{L-1}}\partial_{\mu_{1}}\sigma\left(z_{k;\alpha}^{(L-1)}\right)\partial_{\mu_{2}}\sigma\left(z_{k;\alpha}^{(L-1)}\right) \mid z_{\alpha}^{(\ell)}\right]$$

$$=\mathbb{E}\left[\frac{2}{n_{L-1}}\sum_{k=1}^{n_{L-1}}\left(\sigma'\left(z_{k;\alpha}^{(L-1)}\right)\right)^{2}\partial_{\mu_{1}}z_{k;\alpha}^{(L-1)}\partial_{\mu_{2}}z_{k;\alpha}^{(L-1)} \mid z_{\alpha}^{(\ell)}\right]$$

$$=\frac{1}{n_{L=1}}\sum_{k=1}^{n_{L-1}}\partial_{\mu_{1}}z_{k;\alpha}^{(L-1)}\partial_{\mu_{2}}z_{k;\alpha}^{(L-1)},$$

where in the last equality we use that $\sigma'(z_{k;\alpha}^{(\ell)})$ is distributed according to a Bernoulli 1/2 random variable and is independent of $\partial_{\mu_1} z_{k;\alpha}^{(L-1)} \partial_{\mu_2} z_{k;\alpha}^{(L-1)}$ (this can be seen by symmetrizing $W^{(L-1)} \to -W^{(L-1)}$).

Our next step is to derive a recursion for $B^{(\ell+1)}$ in terms of $B^{(\ell)}$. This is done in Lemma 2.5 below, which relies on the following result:

Proposition 2.4. Consider a random ReLU network with input dimension n_0 , L hidden layers of widths n_1, \ldots, n_L , and output dimension n_{L+1} as in (1.1). Suppose that

$$\frac{1}{n_1} + \dots + \frac{1}{n_L} \le c_1$$

for some $c_1 > 0$. For any fixed network input $x_{\alpha} \in \mathbb{R}^{n_0}$ and any $\ell = 1, \ldots, L$ we have

$$\mathbb{E}\left[\frac{1}{n_{\ell}}\sum_{j=1}^{n_{\ell}}\left(z_{j;\alpha}^{(\alpha)}\right)^{4}\right] = \Theta\left(\frac{1}{n_{0}^{2}}\left|\left|x_{\alpha}\right|\right|^{4}\right),\tag{2.13}$$

where the implicit constants depend on c_1 but are otherwise independent are n, ℓ .

Proof. This result is proved in Theorem 1 [2].

We have the following result.

Lemma 2.5. For any depth $\ell \leq L$, $B^{(\ell)}$ satisfies the following recursion:

$$B^{(\ell)} = \Theta\left(\frac{(\eta_W^{(\ell)})^2 n_{\ell-1}^2}{n_L n_\ell} \frac{1}{n_0^2} ||x_\alpha||^4\right) + \frac{\eta_W^{(\ell)} n_{\ell-1}}{n_\ell} C^{(\ell-1)} + \frac{1}{n_\ell} \widetilde{B}^{(\ell-1)} + \left(1 + \frac{1}{n_\ell}\right) B^{(\ell-1)},$$
(2.14)

where $C^{(\ell)}, \widetilde{B}^{(\ell)} > 0$ are defined as follows:

$$\widetilde{B}^{(\ell)} := \frac{1}{n_{\ell+1}} \mathbb{E} \left[\frac{1}{n_L} \sum_{\mu_1, \mu_2 \le \ell} \eta_{\mu_1} \eta_{\mu_2} \frac{1}{n_\ell^2} \sum_{j_1, j_2 = 1}^{n_\ell} \left(\partial_{\mu_1} z_{j_1; \alpha}^{(\ell)} \partial_{\mu_2} z_{j_2; \alpha}^{(\ell)} \right)^2 \right],$$
(2.15)

$$C^{(\ell)} := \mathbb{E}\left[\frac{1}{n_L} \sum_{\mu \le \ell} \eta_\mu \frac{1}{n_\ell^2} \sum_{j_1, j_2 = 1}^{n_\ell} \left(z_{j_1;\alpha}^{(\ell)} \partial_\mu z_{j_2;\alpha}^{(\ell)}\right)^2\right].$$
(2.16)

Proof of Lemma 2.5. We distinguish several cases to expand the recursion of $B^{(\ell)}$. If $\mu_1, \mu_2 \in \ell$, then the contribution to $B^{(\ell)}$ is

$$\frac{(\eta_W^{(\ell)})^2 n_{\ell-1}^2}{n_L n_\ell} \mathbb{E}\left[\frac{1}{n_{\ell-1}^2} \sum_{j_1, j_2=1}^{n_\ell - 1} \left(\sigma_{j_1}^{(\ell-1)} \sigma_{j_2}^{(\ell-1)}\right)^2\right] = \frac{(\eta_W^{(\ell)})^2 n_{\ell-1}^2}{n_L n_\ell} \Theta\left(\frac{1}{n_0^2} ||x_\alpha||^2\right)$$
(2.17)

Further, if $\mu_1 \leq \ell - 1$ and $\mu_2 \in \ell$ (or vice versa), then the contribution to $B^{(\ell)}$ is

$$2\frac{\eta_W^{(\ell)} n_{\ell-1}}{n_\ell} \mathbb{E}\left[\frac{1}{n_L} \sum_{\mu_1 \le \ell-1} \eta_{\mu_1} \frac{1}{n_{\ell-1}} \sum_{k=1}^{n_{\ell-1}} \left(\sigma_k^{(\ell-1)}\right)^2 \frac{1}{n_\ell} \sum_{j=1}^{n_\ell} \left(\partial_{\mu_1} z_j^{(\ell)}\right)^2\right] = \frac{\eta_W^{(\ell)} n_{\ell-1}}{n_\ell} C^{(\ell-1)}.$$
(2.18)

Finally, if $\mu_1, \mu_2 \leq \ell - 1$, we find the contribution to $B^{(\ell)}$ is

$$\mathbb{E}\left[\frac{1}{n_{L}}\sum_{\mu_{1},\mu_{2}\leq\ell-1}\eta_{\mu_{1}}\eta_{\mu_{2}}\left\{\frac{1}{n_{\ell}}\left(\partial_{\mu_{1}}z_{1}^{(\ell)}\partial_{\mu_{2}}z_{1}^{(\ell)}\right)^{2} + \left(1-\frac{1}{n_{\ell}}\right)\partial_{\mu_{1}}z_{1}^{(\ell)}\partial_{\mu_{2}}z_{1}^{(\ell)}\partial_{\mu_{1}}z_{2}^{(\ell)}\partial_{\mu_{2}}z_{2}^{(\ell)}\right\}\right] = \left(1+\frac{1}{n_{\ell}}\right)B^{(\ell-1)} + \frac{1}{n_{\ell}}\widetilde{B}^{(\ell-1)}.$$
(2.19)

We adding the contributions (2.17), (2.18) and (2.19) in (2.11) gives the stated result. \Box We now compute the recursion that $\widetilde{B}^{(\ell)}$ satisfies.

Lemma 2.6. For any depth $\ell \leq L$, $\widetilde{B}^{(\ell)}$ defined in (2.15) satisfies the following recursion:

$$\frac{1}{n_{\ell}}\widetilde{B}^{(\ell)} = \Theta\left(\frac{(\eta_W^{(\ell)})^2 n_{\ell-1}^2}{n_L n_{\ell}} \frac{||x_{\alpha}||^4}{n_0^2}\right) + \frac{\eta_W^{(\ell)} n_{\ell-1}}{n_{\ell}} C^{(\ell-1)} + \frac{n_{\ell-1}}{n_{\ell}} \frac{1}{n_{\ell-1}} \widetilde{B}^{(\ell-1)} + \frac{2}{n_{\ell}^2} B^{(\ell-1)}.$$
(2.20)

Proof of Lemma 2.6. We apply the same proof strategy as in Lemma 2.5 to get the result. \Box Note that (2.14) and (2.20) also depends on $C^{(\ell)}$. Its recursion is given by the following lemma.

Lemma 2.7. For any depth $\ell \leq L$, $C^{(\ell)}$ defined in (2.16) satisfies the following recursion

$$C^{(\ell)} = \Theta\left(\eta_W^{(\ell)} \frac{n_{\ell-1}}{n_L} \frac{||x_{\alpha}||^4}{n_0^2}\right) + \frac{1}{n_\ell} C^{(\ell-1)} + \left(1 + \frac{1}{n_\ell}\right) \widetilde{C}^{(\ell-1)},$$
(2.21)

where $\widetilde{C}^{(\ell)}>0$ is a sequence defined as

$$\widetilde{C}^{(\ell)} := \frac{1}{n_L} \mathbb{E} \left[\sum_{\mu \le \ell} \eta_\mu \frac{1}{n_\ell^2} \sum_{j_1, j_2 = 1}^{n_\ell} \partial_\mu z_{j_1}^{(\ell)} z_{j_1}^{(\ell)} \partial_\mu z_{j_2}^{(\ell)} z_{j_2}^{(\ell)} \right].$$
(2.22)

Proof of Lemma 2.7. We distinguish several cases to expand the recursion of $C^{(\ell)}$. If $\mu \in \ell$, the contribution to (2.16) is

$$\eta_W^{(\ell)} \frac{n_{\ell-1}}{n_L} \mathbb{E}\left[\frac{1}{n_{\ell-1}^2} \sum_{j_1, j_2=1}^{n_{\ell-1}} \left(z_{j_1}^{(\ell-1)} z_{j_2}^{(\ell-1)}\right)^2\right] = \eta_W^{(\ell)} \Theta\left(\frac{n_{\ell-1}}{n_L} \frac{||x_\alpha||^4}{n_0^2}\right)$$
(2.23)

Finally, when $\mu \leq \ell - 1$, the contribution to (2.16) is

$$\frac{1}{n_L} \mathbb{E} \left[\sum_{\mu \le \ell - 1} \eta_\mu \left\{ \frac{1}{n_\ell} \left(\partial_\mu z_1^{(\ell)} z_1^{(\ell)} \right)^2 + \left(1 - \frac{1}{n_\ell} \right) \left(\partial_\mu z_1^{(\ell)} \right)^2 \left(z_2^{(\ell)} \right)^2 \right\} \right]$$

$$= C^{(\ell - 1)} + \frac{1}{n_\ell} \widetilde{C}^{(\ell - 1)}.$$
(2.24)

Combining (2.23) and (2.24) yields the result.

We finally find the recursion of $\widetilde{C}^{(\ell)}$ that appears in (2.21).

Lemma 2.8. For any depth $\ell \leq L$, $\widetilde{C}^{(\ell)}$ satisfies the following recursion:

$$\widetilde{C}^{(\ell)} = \Theta\left(\frac{\eta_W^{(\ell)} n_{\ell-1}}{n_{\ell} n_L} \frac{||x_{\alpha}||^4}{n_0^2}\right) + \frac{1}{n_{\ell}} C^{(\ell-1)} + \left(1 + \frac{1}{n_{\ell}}\right) \widetilde{C}^{(\ell-1)}.$$
(2.25)

Proof of Lemma 2.8. We apply the same proof strategy as in Lemma 2.7 to get the result. \Box Lemma 2.9. For any depth $\ell \leq L$, we have:

$$\widetilde{C}^{(\ell)} = O(n^{-1}), \tag{2.26}$$

$$C^{(\ell)} = \Theta\left(\frac{||x_{\alpha}||^4}{2n_0^2} \sum_{\ell'=1}^{\ell} \frac{\eta_W^{(\ell')} n_{\ell'-1}}{n_L}\right)$$
(2.27)

Proof of Lemma 2.9. The first result is obtained by observing that there is extra $1/n_L$ in front of $\tilde{C}^{(\ell)}$. Regarding the recursion of $C^{(\ell)}$, we use the fact $\tilde{C}^{(\ell)}$ is small in (2.21) and then sum this equation for $\ell' = 1, \ldots, \ell$ to obtain the value of $C^{(\ell)}$.

We now specialize to the setting of uniform layer width $n_{\ell} = n$ and a global learning rate $\eta_{\mu} = \eta$ to obtain

$$C^{(\ell)} = \Theta\left(\eta\ell\right) \quad \Longrightarrow \quad \frac{1}{n}\widetilde{B}^{(\ell)} = \Theta\left(\eta^{2}\ell^{2}\right) \quad \Longrightarrow \quad B^{(\ell)} = \Theta\left(\eta^{2}L^{3}\right)\left(1 + O(L^{-1})\right),$$

completing the proof of Theorem 1.1.

3 Conclusion

In this short note we've computed how variable network depth influences the learning rate predicted by the μ P heurisdtic. We found that, unlike with respect to width, this learning rate has a non-trivial power law scaling with respect to depth (see Theorem 1.1). We leave for future work empirical validation of whether this depth dependence indeed leads to learning rate transfer in practice.

References

- [1] Simon S. Du, Xiyu Zhai, Barnabas Poczos, and Aarti Singh. Gradient descent provably optimizes over-parameterized neural networks. In *International Conference on Learning Representations*, 2019.
- [2] Boris Hanin. Which neural net architectures give rise to exploding and vanishing gradients? In Advances in Neural Information Processing Systems, 2018.
- [3] Kaiming He, Xiangyu Zhang, Shaoqing Ren, and Jian Sun. Delving deep into rectifiers: Surpassing human-level performance on imagenet classification. In *Proceedings of the IEEE international conference on computer vision*, pages 1026–1034, 2015.
- [4] Arthur Jacot, Franck Gabriel, and Clément Hongler. Neural tangent kernel: Convergence and generalization in neural networks. In Advances in neural information processing systems, pages 8571–8580, 2018.
- [5] Chaoyue Liu, Libin Zhu, and Mikhail Belkin. Loss landscapes and optimization in overparameterized non-linear systems and neural networks. Applied and Computational Harmonic Analysis, 59:85–116, 2022.
- [6] Song Mei, Andrea Montanari, and Phan-Minh Nguyen. A mean field view of the landscape of two-layer neural networks. *Proceedings of the National Academy of Sciences*, 115(33):E7665–E7671, 2018.
- [7] Phan-Minh Nguyen and Huy Tuan Pham. A rigorous framework for the mean field limit of multilayer neural networks. arXiv preprint arXiv:2001.11443, 2020.
- [8] Grant Rotskoff and Eric Vanden-Eijnden. Parameters as interacting particles: long time convergence and asymptotic error scaling of neural networks. Advances in neural information processing systems, 31, 2018.
- [9] Greg Yang, Edward J Hu, Igor Babuschkin, Szymon Sidor, Xiaodong Liu, David Farhi, Nick Ryder, Jakub Pachocki, Weizhu Chen, and Jianfeng Gao. Tensor programs v: Tuning large neural networks via zero-shot hyperparameter transfer. arXiv preprint arXiv:2203.03466, 2022.