FUNCTIONAL GHOBBER-JAMING UNCERTAINTY PRINCIPLE

K. MAHESH KRISHNA

Post Doctoral Fellow Statistics and Mathematics Unit Indian Statistical Institute, Bangalore Centre Karnataka 560 059, India Email: kmaheshak@gmail.com Date: June 5, 2023

Abstract: Let $({f_j}_{j=1}^n, {\tau_j}_{j=1}^n)$ and $({g_k}_{k=1}^n, {\omega_k}_{k=1}^n)$ be two p-orthonormal bases for a finite dimensional Banach space \mathcal{X} . Let $M, N \subseteq \{1, \ldots, n\}$ be such that

$$o(M)^{\frac{1}{q}}o(N)^{\frac{1}{p}} < \frac{1}{\max_{1 \le j,k \le n} |g_k(\tau_j)|},$$

where q is the conjugate index of p. Then for all $x \in \mathcal{X}$, we show that

(1)
$$||x|| \le \left(1 + \frac{1}{1 - o(M)^{\frac{1}{q}} o(N)^{\frac{1}{p}} \max_{1 \le j,k \le n} |g_k(\tau_j)|}\right) \left[\left(\sum_{j \in M^c} |f_j(x)|^p\right)^{\frac{1}{p}} + \left(\sum_{k \in N^c} |g_k(x)|^p\right)^{\frac{1}{p}}\right]$$

We call Inequality (1) as Functional Ghobber-Jaming Uncertainty Principle. Inequality (1) improves the uncertainty principle obtained by Ghobber and Jaming [Linear Algebra Appl., 2011].
Keywords: Uncertainty Principle, Orthonormal Basis, Hilbert space, Banach space.
Mathematics Subject Classification (2020): 42C15, 46B03, 46B04.

CONTENTS

1. Introduction	1
2. Functional Ghobber-Jaming Uncertainty Principle	2
References	7

1. INTRODUCTION

Let $d \in \mathbb{N}$ and $\widehat{}: \mathcal{L}^2(\mathbb{R}^d) \to \mathcal{L}^2(\mathbb{R}^d)$ be the unitary Fourier transform obtained by extending uniquely the bounded linear operator

$$\widehat{}: \mathcal{L}^1(\mathbb{R}^d) \cap \mathcal{L}^2(\mathbb{R}^d) \ni f \mapsto \widehat{f} \in C_0(\mathbb{R}^d); \quad \widehat{f}: \mathbb{R}^d \ni \xi \mapsto \widehat{f}(\xi) := \int_{\mathbb{R}^d} f(x) e^{-2\pi i \langle x, \xi \rangle} \, dx \in \mathbb{C}.$$

In 2007, Jaming [4] extended the uncertainty principle obtained by Nazarov for \mathbb{R} in 1993 [8] (cf. [3]). In the following theorem, Lebesgue measure on \mathbb{R}^d is denoted by m. Mean width of a measurable subset E of \mathbb{R}^d having finite measure is denoted by w(E).

Theorem 1.1. [4, 8] (Nazarov-Jaming Uncertainty Principle) For each $d \in \mathbb{N}$, there exists a universal constant C_d (depends upon d) satisfying the following: If $E, F \subseteq \mathbb{R}^d$ are measurable subsets

having finite measure, then for all $f \in \mathcal{L}^2(\mathbb{R}^d)$,

(2)
$$\int_{\mathbb{R}^d} |f(x)|^2 \, dx \le C_d e^{C_d \min\{m(E)m(F), m(E)^{\frac{1}{d}}w(F), m(F)^{\frac{1}{d}}w(E)\}} \left[\int_{E^c} |f(x)|^2 \, dx + \int_{F^c} |\widehat{f}(\xi)|^2 \, d\xi\right].$$

In particular, if f is supported on E and f is supported on F, then f = 0.

Theorem 1.1 and the milestone paper [1] of Donoho and Stark which derived finite dimensional uncertainty principles, motivated Ghobber and Jaming [2] to ask what is the exact finite dimensional analogue of Theorem 1.1? Ghobber and Jaming were able to derive the following beautiful theorem. Given a subset $M \subseteq \{1, \ldots, n\}$, the number of elements in M is denoted by o(M).

Theorem 1.2. [2] (Ghobber-Jaming Uncertainty Principle) Let $\{\tau_j\}_{j=1}^n$ and $\{\omega_j\}_{j=1}^n$ be orthonormal bases for the Hilbert space \mathbb{C}^n . If $M, N \subseteq \{1, \ldots, n\}$ are such that

(3)
$$o(M)o(N) < \frac{1}{\max_{1 \le j,k \le n} |\langle \tau_j, \omega_k \rangle|^2},$$

then for all $h \in \mathbb{C}^n$,

$$\|h\| \le \left(1 + \frac{1}{1 - \sqrt{o(M)o(N)}} \max_{1 \le j,k \le n} |\langle \tau_j, \omega_k \rangle|\right) \left[\left(\sum_{j \in M^c} |\langle h, \tau_j \rangle|^2 \right)^{\frac{1}{2}} + \left(\sum_{k \in N^c} |\langle h, \omega_k \rangle|^2 \right)^{\frac{1}{2}} \right].$$

In particular, if h is supported on M in the expansion using basis $\{\tau_j\}_{j=1}^n$ and h is supported on N in the expansion using basis $\{\omega_j\}_{j=1}^n$, then h = 0.

It is reasonable to ask whether there is a Banach space version of Ghobber-Jaming Uncertainty Principle, which when restricted to Hilbert space, reduces to Theorem 1.2? We are going to answer this question in the paper.

2. Functional Ghobber-Jaming Uncertainty Principle

In the paper, \mathbb{K} denotes \mathbb{C} or \mathbb{R} and \mathcal{X} denotes a finite dimensional Banach space over \mathbb{K} . Identity operator on \mathcal{X} is denoted by $I_{\mathcal{X}}$. Dual of \mathcal{X} is denoted by \mathcal{X}^* . Whenever 1 , <math>q denotes conjugate index of p. For $d \in \mathbb{N}$, the standard finite dimensional Banach space \mathbb{K}^d over \mathbb{K} equipped with standard $\|\cdot\|_p$ norm is denoted by $\ell^p([d])$. Canonical basis for \mathbb{K}^d is denoted by $\{\delta_j\}_{j=1}^d$ and $\{\zeta_j\}_{j=1}^d$ be the coordinate functionals associated with $\{\delta_j\}_{j=1}^d$. Motivated from the properties of orthonormal bases for Hilbert spaces, we set the following notion of p-orthonormal bases which is also motivated from the notion of p-approximate Schauder frames [7] and p-unconditional Schauder frames [6].

Definition 2.1. Let \mathcal{X} be a finite dimensional Banach space over \mathbb{K} . Let $\{\tau_j\}_{j=1}^n$ be a basis for \mathcal{X} and let $\{f_j\}_{j=1}^n$ be the coordinate functionals associated with $\{\tau_j\}_{j=1}^n$. The pair $(\{f_j\}_{j=1}^n, \{\tau_j\}_{j=1}^n)$ is said to be a **p-orthonormal basis** $(1 for <math>\mathcal{X}$ if the following conditions hold.

- (i) $||f_j|| = ||\tau_j|| = 1$ for all $1 \le j \le n$.
- (ii) For every $(a_j)_{j=1}^n \in \mathbb{K}^n$,

$$\left\|\sum_{j=1}^n a_j \tau_j\right\| = \left(\sum_{j=1}^n |a_j|^p\right)^{\frac{1}{p}}.$$

Given a p-orthonormal basis $({f_j}_{j=1}^n, {\tau_j}_{j=1}^n)$, we easily see from Definition 2.1 that

$$\|x\| = \left\|\sum_{j=1}^n f_j(x)\tau_j\right\| = \left(\sum_{j=1}^n |f_j(x)|^p\right)^{\frac{1}{p}}, \quad \forall x \in \mathcal{X}.$$

Example 2.2. The pair $(\{\zeta_j\}_{j=1}^d, \{\delta_j\}_{j=1}^d)$ is a p-orthonormal basis for $\ell^p([d])$.

Like orthonormal bases for Hilbert spaces, the following theorem characterizes all p-orthonormal bases.

Theorem 2.3. Let $(\{f_j\}_{j=1}^n, \{\tau_j\}_{j=1}^n)$ be a p-orthonormal basis for \mathcal{X} . Then a pair $(\{g_j\}_{j=1}^n, \{\omega_j\}_{j=1}^n)$ is a p-orthonormal basis for \mathcal{X} if and only if there is an invertible linear isometry $V : \mathcal{X} \to \mathcal{X}$ such that

$$g_j = f_j V^{-1}, \ \omega_j = V \tau_j, \quad \forall 1 \le j \le n.$$

Proof. (\Rightarrow) Define $V : \mathcal{X} \ni x \mapsto \sum_{j=1}^{n} f_j(x) \omega_j \in \mathcal{X}$. Since $\{\omega_j\}_{j=1}^{n}$ is a basis for \mathcal{X} , V is invertible with inverse $V^{-1} : \mathcal{X} \ni x \mapsto \sum_{j=1}^{n} g_j(x) \tau_j \in \mathcal{X}$. For $x \in \mathcal{X}$,

$$\|Vx\| = \left\|\sum_{j=1}^{n} f_j(x)\omega_j\right\| = \left(\sum_{j=1}^{n} |f_j(x)|^p\right)^{\frac{1}{p}} = \left\|\sum_{j=1}^{n} f_j(x)\tau_j\right\| = \|x\|.$$

Therefore V is isometry. Note that we clearly have $\omega_j = V\tau_j, \forall 1 \leq j \leq n$. Now let $1 \leq j \leq n$. Then

$$f_j(V^{-1}x) = f_j\left(\sum_{k=1}^n g_k(x)\tau_k\right) = \sum_{k=1}^n g_k(x)f_j(\tau_k) = g_j(x), \quad \forall x \in \mathcal{X}$$

(\Leftarrow) Since V is invertible, $\{\omega_j\}_{j=1}^n$ is a basis for \mathcal{X} . Now we see that $g_j(\omega_k) = f_j(V^{-1}V\tau_k) = f_j(\tau_k) = \delta_{j,k}$ for all $1 \leq j, k \leq n$. Therefore $\{g_j\}_{j=1}^n$ is the coordinate functionals associated with $\{\omega_j\}_{j=1}^n$. Since V is an isometry, we have $\|\omega_j\| = 1$ for all $1 \leq j \leq n$. Since V is also invertible, we have

$$\begin{aligned} \|g_j\| &= \sup_{x \in \mathcal{X}, \|x\| \le 1} |g_j(x)| = \sup_{x \in \mathcal{X}, \|x\| \le 1} |f_j(V^{-1}x)| = \sup_{Vy \in \mathcal{X}, \|Vy\| \le 1} |f_j(y)| \\ &= \sup_{Vy \in \mathcal{X}, \|y\| \le 1} |f_j(y)| = \|f_j\| = 1, \quad \forall 1 \le j \le n. \end{aligned}$$

Finally, for every $(a_j)_{j=1}^n \in \mathbb{K}^n$,

$$\left\|\sum_{j=1}^{n} a_{j}\omega_{j}\right\| = \left\|\sum_{j=1}^{n} a_{j}V\tau_{j}\right\| = \left\|V\left(\sum_{j=1}^{n} a_{j}\tau_{j}\right)\right\| = \left\|\sum_{j=1}^{n} a_{j}\tau_{j}\right\| = \left(\sum_{j=1}^{n} |a_{j}|^{p}\right)^{\frac{1}{p}}.$$

In the next result we show that Example 2.2 is prototypical as long as we consider p-orthonormal bases.

Theorem 2.4. If \mathcal{X} has a *p*-orthonormal basis $(\{f_j\}_{j=1}^n, \{\tau_j\}_{j=1}^n)$, then \mathcal{X} is isometrically isomorphic to $\ell^p([n])$.

Proof. Define $V : \mathcal{X} \ni x \mapsto \sum_{j=1}^{n} f_j(x) \delta_j \in \ell^p([n])$. By doing a similar calculation as in the direct part in the proof of Theorem 2.3, we see that V is an invertible isometry.

Now we derive main result of this paper.

Theorem 2.5. (Functional Ghobber-Jaming Uncertainty Principle) Let $(\{f_j\}_{j=1}^n, \{\tau_j\}_{j=1}^n)$ and $(\{g_k\}_{k=1}^n, \{\omega_k\}_{k=1}^n)$ be p-orthonormal bases for \mathcal{X} . If $M, N \subseteq \{1, \ldots, n\}$ are such that

$$o(M)^{\frac{1}{q}}o(N)^{\frac{1}{p}} < \frac{1}{\max_{1 \le j,k \le n} |g_k(\tau_j)|},$$

then for all $x \in \mathcal{X}$,

(4)
$$||x|| \leq \left(1 + \frac{1}{1 - o(M)^{\frac{1}{q}} o(N)^{\frac{1}{p}} \max_{1 \leq j,k \leq n} |g_k(\tau_j)|}\right) \left[\left(\sum_{j \in M^c} |f_j(x)|^p\right)^{\frac{1}{p}} + \left(\sum_{k \in N^c} |g_k(x)|^p\right)^{\frac{1}{p}}\right].$$

In particular, if x is supported on M in the expansion using basis $\{\tau_j\}_{j=1}^n$ and x is supported on N in the expansion using basis $\{\omega_k\}_{k=1}^n$, then x = 0.

Proof. Given $S \subseteq \{1, \ldots, n\}$, define

$$P_S x \coloneqq \sum_{j \in S} f_j(x) \tau_j, \quad \forall x \in \mathcal{X}, \quad \|x\|_{S,f} \coloneqq \left(\sum_{j \in S} |f_j(x)|^p\right)^{\frac{1}{p}}, \quad \|x\|_{S,g} \coloneqq \left(\sum_{j \in S} |g_j(x)|^p\right)^{\frac{1}{p}}.$$

Also define $V : \mathcal{X} \ni x \mapsto \sum_{k=1}^{n} g_k(x)\tau_k \in \mathcal{X}$. Then V is an invertible isometry. Using V we make following important calculations:

$$\|P_S x\| = \left\| \sum_{j \in S} f_j(x) \tau_j \right\| = \left(\sum_{j \in S} |f_j(x)|^p \right)^{\frac{1}{p}} = \|x\|_{S,f}, \quad \forall x \in \mathcal{X}$$

and

$$|P_S V x\| = \left\| \sum_{j \in S} f_j(V x) \tau_j \right\| = \left\| \sum_{j \in S} f_j\left(\sum_{k=1}^n g_k(x) \tau_k\right) \tau_j \right\| = \left\| \sum_{j \in S} \sum_{k=1}^n g_k(x) f_j(\tau_k) \tau_j \right\|$$
$$= \left\| \sum_{j \in S} g_j(x) \tau_j \right\| = \left(\sum_{j \in S} |g_j(x)|^p\right)^{\frac{1}{p}} = \|x\|_{S,g}, \quad \forall x \in \mathcal{X}.$$

Now let $y \in \mathcal{X}$ be such that $\{j \in \{1, ..., n\} : f_j(y) \neq 0\} \subseteq M$. Then $||P_N V y|| = ||P_N V P_M y|| \le ||P_N V P_M |||y||$ and

 $\|y\|_{N^{c},g} = \|P_{N^{c}}Vy\| = \|Vy - P_{N}Vy\| \ge \|Vy\| - \|P_{N}Vy\| = \|y\| - \|P_{N}Vy\| \ge \|y\| - \|P_{N}VP_{M}\|\|y\|.$

Therefore

(5)
$$\|y\|_{N^c,g} \ge (1 - \|P_N V P_M\|)\|y\|.$$

Let $x \in \mathcal{X}$. Note that $P_M x$ satisfies $\{j \in \{1, ..., n\} : f_j(P_M x) \neq 0\} \subseteq M$. Now using (5) we get $\|x\| = \|P_M x + P_{M^c} x\| \leq \|P_M x\| + \|P_{M^c} x\| \leq \frac{1}{1 - \|P_N V P_M\|} \|P_M x\|_{N^c,g} + \|P_{M^c} x\|$ $= \frac{1}{1 - \|P_N V P_M\|} \|P_{N^c} V P_M x\| + \|P_{M^c} x\| = \frac{1}{1 - \|P_N V P_M\|} \|P_{N^c} V (x - P_{M^c} x)\| + \|P_{M^c} x\|$ $\leq \frac{1}{1 - \|P_N V P_M\|} \|P_{N^c} V x\| + \frac{1}{1 - \|P_N V P_M\|} \|P_{M^c} x\| + \|P_{M^c} x\|$ $\leq \frac{1}{1 - \|P_N V P_M\|} \|P_{N^c} V x\| + \frac{1}{1 - \|P_N V P_M\|} \|P_{M^c} x\| + \|P_{M^c} x\|$ $= \frac{1}{1 - \|P_N V P_M\|} \|P_{N^c} V x\| + \left(1 + \frac{1}{1 - \|P_N V P_M\|}\right) \|P_{M^c} x\|$ $\leq \|P_{N^c} V x\| + \frac{1}{1 - \|P_N V P_M\|} \|P_{N^c} V x\| + \left(1 + \frac{1}{1 - \|P_N V P_M\|}\right) \|P_{M^c} x\|$ $\leq \|P_{N^c} V x\| + \frac{1}{1 - \|P_N V P_M\|} \|P_{N^c} V x\| + \left(1 + \frac{1}{1 - \|P_N V P_M\|}\right) \|P_{M^c} x\|$ $= \left(1 + \frac{1}{1 - \|P_N V P_M\|}\right) [\|P_{N^c} V x\| + \|P_{M^c} x\|] = \left(1 + \frac{1}{1 - \|P_N V P_M\|}\right) [\|x\|_{N^c,g} + \|P_{M^c} x\|]$ $= \left(1 + \frac{1}{1 - \|P_N V P_M\|}\right) \left[\left(\sum_{j \in M^c} |f_j(x)|^p\right)^{\frac{1}{p}} + \left(\sum_{k \in N^c} |g_k(x)|^p\right)^{\frac{1}{p}}\right].$

For $x \in \mathcal{X}$, we now find

$$\begin{split} \|P_{N}VP_{M}x\|^{p} &= \left\|\sum_{k\in N}f_{k}(VP_{M}x)\tau_{k}\right\|^{p} = \left(\sum_{k\in N}|f_{k}(VP_{M}x)|^{p}\right)^{\frac{1}{p}} = \sum_{k\in N}\left|(f_{k}V)\left(\sum_{j\in M}f_{j}(x)\tau_{j}\right)\right|^{p} \\ &= \sum_{k\in N}\left|\sum_{j\in M}f_{j}(x)f_{k}(V\tau_{j})\right|^{p} = \sum_{k\in N}\left|\sum_{j\in M}f_{j}(x)f_{k}\left(\sum_{r=1}^{n}g_{r}(\tau_{j})\tau_{r}\right)\right|^{p} = \sum_{k\in N}\left|\sum_{j\in M}f_{j}(x)\sum_{r=1}^{n}g_{r}(\tau_{j})f_{k}(\tau_{r})\right|^{p} \\ &= \sum_{k\in N}\left|\sum_{j\in M}f_{j}(x)g_{k}(\tau_{j})\right|^{p} \leq \sum_{k\in N}\left(\sum_{j\in M}|f_{j}(x)g_{k}(\tau_{j})|\right)^{p} \leq \left(\max_{1\leq j,k\leq n}|g_{k}(\tau_{j})|\right)^{p}\sum_{k\in N}\left(\sum_{j\in M}|f_{j}(x)|^{p}\right)^{\frac{p}{p}} \\ &= \left(\max_{1\leq j,k\leq n}|g_{k}(\tau_{j})|\right)^{p}o(N)\left(\sum_{j\in M}|f_{j}(x)|^{p}\right)^{p} \leq \left(\max_{1\leq j,k\leq n}|g_{k}(\tau_{j})|\right)^{p}o(N)\left(\sum_{j\in M}1^{q}\right)^{\frac{p}{q}} \\ &\leq \left(\max_{1\leq j,k\leq n}|g_{k}(\tau_{j})|\right)^{p}o(N)\left(\sum_{j=1}^{n}|f_{j}(x)|^{p}\right)^{\frac{p}{p}}\left(\sum_{j\in M}1^{q}\right)^{\frac{p}{q}} = \left(\max_{1\leq j,k\leq n}|g_{k}(\tau_{j})|\right)^{p}o(N)\|x\|^{p}o(M)^{\frac{p}{q}}. \end{split}$$

Therefore

$$||P_N V P_M|| \le \max_{1\le j,k\le n} |g_k(\tau_j)| o(N)^{\frac{1}{p}} o(M)^{\frac{1}{q}}$$

which gives the theorem.

Corollary 2.6. Theorem 1.2 follows from Theorem 2.5.

Proof. Let $\{\tau_j\}_{j=1}^n, \{\omega_j\}_{j=1}^n$ be two orthonormal bases for a finite dimensional Hilbert space \mathcal{H} . Define

$$f_j: \mathcal{H} \ni h \mapsto \langle h, \tau_j \rangle \in \mathbb{K}; \quad g_j: \mathcal{H} \ni h \mapsto \langle h, \omega_j \rangle \in \mathbb{K}, \quad \forall 1 \le j \le n.$$

Then p = q = 2 and $|f_j(\omega_k)| = |\langle \omega_k, \tau_j \rangle|$ for all $1 \le j, k \le n$.

By interchanging p-orthonormal bases in Theorem 2.5 we get the following theorem.

Theorem 2.7. (Functional Ghobber-Jaming Uncertainty Principle) Let $(\{f_j\}_{j=1}^n, \{\tau_j\}_{j=1}^n)$ and $(\{g_k\}_{k=1}^n, \{\omega_k\}_{k=1}^n)$ be p-orthonormal bases for \mathcal{X} . If $M, N \subseteq \{1, \ldots, n\}$ are such that

$$p(M)^{\frac{1}{q}}o(N)^{\frac{1}{p}} < \frac{1}{\max_{1 \le j,k \le n} |f_j(\omega_k)|}$$

then for all $x \in \mathcal{X}$,

$$\|x\| \le \left(1 + \frac{1}{1 - o(M)^{\frac{1}{q}} o(N)^{\frac{1}{p}} \max_{1 \le j,k \le n} |f_j(\omega_k)|}\right) \left[\left(\sum_{k \in M^c} |g_k(x)|^p \right)^{\frac{1}{p}} + \left(\sum_{j \in N^c} |f_j(x)|^p \right)^{\frac{1}{p}} \right].$$

In particular, if x is supported on M in the expansion using basis $\{\omega_k\}_{k=1}^n$ and x is supported on N in the expansion using basis $\{\tau_j\}_{j=1}^n$, then x = 0.

Observe that the constant

$$C_{d}e^{C_{d}}\min\{m(E)m(F),m(E)^{\frac{1}{d}}w(F),m(F)^{\frac{1}{d}}w(E)\}$$

in Inequality (2) is depending upon subsets E, F and not on the entire domain \mathbb{R} of functions f, \hat{f} . Thus it is natural to ask whether there is a constant sharper in Inequality (4) depending upon subsets M, N and not on $\{1, \ldots, n\}$. A careful observation in the proof of Theorem 2.5 gives following result.

Theorem 2.8. Let $(\{f_j\}_{j=1}^n, \{\tau_j\}_{j=1}^n)$ and $(\{g_k\}_{k=1}^n, \{\omega_k\}_{k=1}^n)$ be p-orthonormal bases for \mathcal{X} . If $M, N \subseteq \{1, \ldots, n\}$ are such that

$$o(M)^{\frac{1}{q}}o(N)^{\frac{1}{p}} < \frac{1}{\max_{j\in M, k\in N} |g_k(\tau_j)|},$$

then for all $x \in \mathcal{X}$,

$$\|x\| \le \left(1 + \frac{1}{1 - o(M)^{\frac{1}{q}} o(N)^{\frac{1}{p}} \max_{j \in M, k \in N} |g_k(\tau_j)|}\right) \left[\left(\sum_{j \in M^c} |f_j(x)|^p \right)^{\frac{1}{p}} + \left(\sum_{k \in N^c} |g_k(x)|^p \right)^{\frac{1}{p}} \right]$$

Similarly we have the following result from Theorem 2.7.

Theorem 2.9. Let $(\{f_j\}_{j=1}^n, \{\tau_j\}_{j=1}^n)$ and $(\{g_k\}_{k=1}^n, \{\omega_k\}_{k=1}^n)$ be p-orthonormal bases for \mathcal{X} . If $M, N \subseteq \{1, \ldots, n\}$ are such that

$$o(M)^{\frac{1}{q}}o(N)^{\frac{1}{p}} < \frac{1}{\max_{j \in N, k \in M} |f_j(\omega_k)|},$$

then for all $x \in \mathcal{X}$,

$$\|x\| \le \left(1 + \frac{1}{1 - o(M)^{\frac{1}{q}} o(N)^{\frac{1}{p}} \max_{j \in N, k \in M} |f_j(\omega_k)|}\right) \left[\left(\sum_{k \in M^c} |g_k(x)|^p\right)^{\frac{1}{p}} + \left(\sum_{j \in N^c} |f_j(x)|^p\right)^{\frac{1}{p}}\right].$$

Theorem 2.5 brings the following question.

Question 2.10. Given p and a Banach space \mathcal{X} of dimension n, for which subsets $M, N \subseteq \{1, \ldots, n\}$ and pairs of p-orthonormal bases $(\{f_j\}_{j=1}^n, \{\tau_j\}_{j=1}^n), (\{g_k\}_{k=1}^n, \{\omega_k\}_{k=1}^n)$ for \mathcal{X} , we have equality in Inequality (4)?

It is clear that we used 1 in the proof of Theorem 2.5. However, Definition 2.1 can easily be extended to include cases <math>p = 1 and $p = \infty$. This therefore leads to the following question.

Question 2.11. Whether there are Functional Ghobber-Jaming Uncertainty Principle (versions of Theorem 2.5) for 1-orthonormal bases and ∞ -orthonormal bases?

We end by mentioning that Donoho-Stark-Elad-Bruckstein-Ricaud-Torrésani Uncertainty Principle for finite dimensional Banach spaces is derived in [5] (actually, in [5] the functional uncertainty principle was derived for p-Schauder frames which is general than p-orthonormal bases. Thus it is worth to derive Theorem 2.5 or a variation of it for p-Schauder frames, which we are unable).

References

- David L. Donoho and Philip B. Stark. Uncertainty principles and signal recovery. SIAM J. Appl. Math., 49(3):906–931, 1989.
- [2] Saifallah Ghobber and Philippe Jaming. On uncertainty principles in the finite dimensional setting. Linear Algebra Appl., 435(4):751–768, 2011.
- [3] Victor Havin and Burglind Jöricke. The uncertainty principle in harmonic analysis, volume 28 of A Series of Modern Surveys in Mathematics. Springer-Verlag, Berlin, 1994.
- [4] Philippe Jaming. Nazarov's uncertainty principles in higher dimension. J. Approx. Theory, 149(1):30-41, 2007.
- [5] K. Mahesh Krishna. Functional Donoho-Stark-Elad-Bruckstein-Ricaud-Torrésani uncertainty principle. arXiv: 2304.03324v1 [math.FA] 5 April, 2023.
- [6] K. Mahesh Krishna. Group-frames for Banach spaces. arXiv: 2305.01499v1 [math.FA] 5 May, 2023.
- [7] K. Mahesh Krishna and P. Sam Johnson. Towards characterizations of approximate Schauder frame and its duals for Banach spaces. J. Pseudo-Differ. Oper. Appl., 12(1):Paper No. 9, 13, 2021.
- [8] F. L. Nazarov. Local estimates for exponential polynomials and their applications to inequalities of the uncertainty principle type. Algebra i Analiz, 5(4):3–66, 1993.