# <span id="page-0-1"></span>**PERIODIC-POINT STRUCTURES ON PARAMETRIZED SPECTRA: AN APPLICATION OF RIGIDITY**

#### CARY MALKIEWICH AND KATE PONTO

ABSTRACT. The bicategory of parameterized spectra has a remarkably rich structure. In particular, it is possible to take traces in this bicategory, which give classical invariants that count fixed points. We can also take equivariant traces, which give significant generalizations of the classical invariants that count periodic points.

Unfortunately, the existence of these traces in general depends on technical statements about the bicategory that can be difficult to verify directly. In this paper, we demonstrate the effectiveness of two tools – rigidity and deformable functors – by using them establish the structure we need to take these equivariant traces and to construct periodic-point invariants in a formal way.

### **CONTENTS**



### 1. INTRODUCTION

<span id="page-0-0"></span>The Lefschetz fixed point theorem shows that for a continuous map  $f: X \to X$ , the signed count of the fixed points of *f* can be given by an algebraic construction, namely the alternating sum of traces on rational homology. Later Dold observed in [\[Dol74\]](#page-37-1) that that this signed count is also the trace of  $\Sigma^{\infty}_{+}f$  in the stable homotopy category. In other words, the fixed points of *f* may be measured by taking an abstract trace in a symmetric monoidal category [\[DP80\]](#page-37-2).

This theorem has a generalization due to Wecken [\[Wec42\]](#page-38-0), which gives a more sensitive fixed-point invariant called the Reidemeister trace *R*(*f* ). In her thesis, the second author showed that this more sensitive invariant is also a trace, but the trace is carried out in a more sophisticated setting: the bicategory of parametrized spectra [\[Pon10\]](#page-37-3). The result is a version of *R*(*f* ) that lives in topological Hochschild homology (THH).

In [\[MP22\]](#page-37-4), the authors show how to pass to an invariant that counts periodic points in families and lives in topological restriction homology (TR). This invariant is expressed as a trace in the bicategory of *equivariant* parametrized spectra. However, in order to prove

*Date*: 7th Jun, 2023.

<sup>2020</sup> *Mathematics Subject Classification.* 18D30, 18M05, 18N10, 55P42, 55R70 .

*Key words and phrases.* parameterized spectra, coherence, rigidity, deformable functors.

<span id="page-1-1"></span>that it has the desired behavior, one needs to show that the bicategory of parametrized spectra has tensor powers that interact with the trace in a well-behaved way. In [\[MP22\]](#page-37-4), this desired structure defined explicitly and called an "*n*-Fuller structure."

In this paper, we use two tools – rigidity and deformable functors – and work in the approach to parametrized spectra from [\[Mal\]](#page-37-5) to give a detailed construction of this structure.

<span id="page-1-0"></span>**Theorem 1.1** ([§5.2\)](#page-19-0)**.** *The bicategory of parametrized spectra* Ex *has an n-Fuller structure.*

This extends earlier results from [\[MS06,](#page-37-6) [Shu08,](#page-38-1) [PS12\]](#page-37-7) showing that  $\&\mathbf{x}$  is a bicategory with shadow.

We also generalize Theorem [1.1](#page-1-0) in three directions. The first direction involves base change objects in the bicategory  $\mathcal{E}$ x that encode maps of spaces  $f: A \rightarrow B$ . Taking the product in  $\&\&x\&y$  with these base change objects has the effect of taking pullback or pushforward along *f*. We show that these objects can be chosen to be appropriately compatible with the above *n*-Fuller structure:

**Proposition 1.2** ([§5.3\)](#page-23-0)**.** *The bicategory* Ex *has a compatible system of base-change 1 cells.*

This is related to the notion of a "framed bicategory" developed in [\[Shu08\]](#page-38-1).

The second direction involves the equivariant and fiberwise versions of the bicategory Ex.

**Theorem 1.3** ([§6\)](#page-24-0)**.** *For any finite group G and base space B, the bicategories of*

- *equivariant parametrized spectra G*Ex*,*
- *fiberwise parametrized spectra* Ex*B, and*
- *equivariant fiberwise parametrized spectra*  $G\mathcal{E}_{X}$ *R,*

*also have n-Fuller structures and compatible systems of base-change 1-cells.*

Finally, we extend Theorem [1.1](#page-1-0) to encompass two change-of-group functors on equivariant parameterized spectra. Let *G* be a finite group,  $H \leq G$  a subgroup, and  $WH =$ *NH*/*H* the Weyl group of *H*.

**Theorem 1.4** ([§7\)](#page-29-0)**.** *The geometric fixed point functor and the forgetful functor induce maps of bicategories*

$$
\Phi^H : G \, \mathcal{E} \mathbf{x} \to W H \, \mathcal{E} \mathbf{x}, \qquad \iota_H^* : G \, \mathcal{E} \mathbf{x} \to H \, \mathcal{E} \mathbf{x}
$$

*and in the fiberwise case*

$$
\Phi^H: G\mathcal{E}\mathbf{x}_B \to WH\mathcal{E}\mathbf{x}_B, \qquad \iota_H^*: G\mathcal{E}\mathbf{x}_B \to H\mathcal{E}\mathbf{x}_B,
$$

*that are compatible with the shadow and the base-change objects.*

All three of these generalizations play important roles in [\[MP22\]](#page-37-4).

The rhythm of proof for all of these results is the same. We first establish a structure at the point-set level, by observing that the functors of interest have no automorphisms other than the identity. We call such functors **rigid**. Many of the functors we care about on parameterized spectra are rigid by [\[Mal,](#page-37-5) Thm 4.5.2]. We then use this fact to produce coherence isomorphisms and to verify that the relevant compatibilities between coherence isomorphisms hold.

Once we have built the desired structure at the point-set level, we show it descends to the homotopy category using the notion of a **deformable functor** from [\[DHKS04\]](#page-37-8). This is more general than the notion of a derived functor in a model category, but it has most of the same properties. The extra flexibility is very useful because the most convenient kind of fibration in the parametrized setting does not fit nicely into a model structure.

<span id="page-2-2"></span>Most importantly, deformable functors allow us to transfer the point-set coherence compatibilities to derived compatibilities in a straightforward way.

While our initial motivation was Theorem [1.1](#page-1-0) and its generalizations, we now regard the primary contribution of this paper to be the approach to the proofs presented here. It is much more general than our applications here and has wide ranging uses.

**Organization.** In Section [2](#page-2-0) we introduce rigid functors and provide the theorem that will allow us to easily verify the functors of interest in later sections are rigid. In Section [3](#page-4-0) we use the rigidity result of Section [2](#page-2-0) to show that parameterized spectra form a bicategory with a compatible shadow, *n*-Fuller structure, and base change objects.

In Section [4](#page-12-0) we recall the theory of deformable functors from [\[DHKS04\]](#page-37-8). In Section [5](#page-16-0) we combine the results of Sections [3](#page-4-0) and [4](#page-12-0) to show that the homotopy categories of parameterized spectra assemble to a bicategory with a compatible shadow, *n*-Fuller structure, and base change objects.

Section [6](#page-24-0) describes how to modify the results in Sections [3](#page-4-0) and [5](#page-16-0) to prove the analogs of Theorem [1.1](#page-1-0) for equivariant, fiberwise, and equivariant fiberwise parameterized spectra. In Section [7](#page-29-0) we apply the techniques of Sections [2](#page-2-0) and [4](#page-12-0) to functors between categories of parameterized spectra.

**Acknowledgments.** The first author was partially supported by NSF grants DMS-2005524 and DMS-2052923. The second author was partially supported by NSF grants DMS-2052905 and DMS-1810779.

### 2. RIGID MULTI-SPANS

<span id="page-2-0"></span>In the first three sections of this paper, we work in the symmetric monoidal bifibration (SMBF) [\[Shu08,](#page-38-1) 12.1] of parameterized orthogonal spectra over all base spaces. We follow the notational conventions of [\[Mal,](#page-37-5) Thm 6.1.1]. This means that we have:

- a symmetric monoidal category of parameterized orthogonal spectra over all base spaces, denoted  $\mathbb{OS}$  [\[Mal,](#page-37-5) Def 4.1.1 and 4.1.6], whose monoidal product  $\overline{\wedge}$  is called the **external product**,
- a strict symmetric monoidal functor  $\Phi$ :  $OS \rightarrow Top$  that identifies the base space of the spectrum,
- for every morphism  $f: A \rightarrow B$  in  $\mathcal{I}$ *op* and spectrum *X* over *B*, there is a spectrum  $f^*(X)$  over *A* and cartesian arrow  $f^*(X) \to X$  over *f*, and
- <span id="page-2-1"></span>• for every morphism  $f : A \to B$  in  $\mathcal{I}_{op}$  and spectrum Y over A, there is a spectrum  $f_!(Y)$  over *B* and co-cartesian arrow  $f_!(Y) \to Y$  over *f*.

The external product preserves cartesian and co-cartesian arrows and we have Beck-Chevalley isomorphisms  $j_!k^* \cong h^*f_!$  for every pullback square of topological spaces

$$
(2.1)
$$



We let  $OS(A)$  denote the fiber of  $\Phi$  over A. See [\[Shu08\]](#page-38-1) for a discussion of bifibrations.

In the applications we have in mind, the three operations  $\overline{\wedge},$   $f^*,$  and  $f_!$  tend to appear bundled together in the following form.

<span id="page-3-5"></span>**Definition 2.2.** Given a list of topological spaces *A*1, ..., *An*, *C*, a **multi-span** from the  $(A_i)$  to C is a span of the form

<span id="page-3-0"></span>(2.3) 
$$
C \longleftarrow f \longrightarrow_{A_1 \times ... \times A_n} B \underbrace{\downarrow g_1,...,g_n}_{A_1 \times ... \times A_n}.
$$

In other words, a space *B* and a choice of map from *B* to each of the spaces  $A_1, \ldots, A_n$ *C*. For each multi-span from  $(A_i)$  to *C* as in [\(2.3\)](#page-3-0) we define a functor on parametrized spectra

<span id="page-3-1"></span>(2.4) 
$$
\prod_i \mathcal{OS}(A_i) \longrightarrow \mathcal{OS}(C)
$$

$$
(X_1, ..., X_n) \longmapsto f_!(g_1, ..., g_n)^*(X_1 \overline{\wedge} ... \overline{\wedge} X_n).
$$

We refer to this as the **action** of the multi-span on the tuple of spectra  $(X_i)$ . *Remark* 2.5*.* This definition explains why we explicitly allow for a product on the right in [\(2.3\)](#page-3-0) but not on the left. (Of course nothing stops us from replacing *C* by a product.)

<span id="page-3-2"></span>**Example 2.6.** If  $n = 0$  this action takes no inputs and produces a single parametrized spectrum as output, namely  $f_! \pi^* I$ , where *I* is the unit of the symmetric monoidal category  $\varnothing$  and  $\pi$  is the map to the one-point space. Concretely,  $I$  is the sphere spectrum S over the one-point space  $\star$ , so this action produces  $\Sigma^{\infty}_{\mu}$  $_{+C}^{\infty}B$ , the fiberwise suspension spectrum over  $\overline{C}$  of  $\overline{B}_{+C} \cong B \amalg C$ .

Throughout this document we only consider the subcategory  $OS<sup>c</sup> \subseteq OS$  of spectra that are freely *f* -cofibrant [\[Mal,](#page-37-5) Def 4.2.2]. The action of a multi-span preserves this class of spectra by [\[Mal,](#page-37-5) Lem 4.3.1, Thm 4.4.6].

A key property of the functor  $(2.4)$  is that for many choices of  $g_1, \ldots, g_n, f$  it has very few automorphisms. We say that a functor is **rigid** if the only natural automorphism is the identity transformation.

**Example 2.7.** The functor

$$
\textit{Top} \times \textit{Top} \times \textit{Top} \times \textit{Top} \xrightarrow{\quad \times \quad} \textit{Top}
$$

that sends  $(A, B, C, D)$  to the product  $A \times B \times C \times D$ , is rigid.

The point of rigidity is that it makes it easy to verify coherence diagrams. For instance, once we know the above functor is rigid, we can verify the pentagon identity for the monoidal structure on *Top* (compare to Figure [3.3a\)](#page-4-1) in the following way. The composite of maps around the pentagon is a natural automorphism of  $A \times B \times C \times D$ . But the only natural isomorphism is the identity, so this composite must be the identity.

<span id="page-3-3"></span>**Theorem 2.8.** [\[Mal,](#page-37-5) Thm 4.5.2] *The functor* [\(2.4\)](#page-3-1) *is rigid if the map*

$$
B \xrightarrow{(f,g_1,\ldots,g_n)} C \times A_1 \times \ldots \times A_n
$$

*is injective.*

Motivated by this theorem, we say that the multi-span in [\(2.3\)](#page-3-0) is **rigid** if the map  $(f, g_1, \ldots, g_n)$  is injective. So if a multi-span is rigid then its action is rigid. As a consequence, any diagram of natural isomorphisms between functors isomorphic to [\(2.4\)](#page-3-1) automatically commutes.

<span id="page-3-4"></span>*Remark* 2.9. When  $n = 0$ , as in Example [2.6,](#page-3-2) the rigidity theorem simplifies to the fact that the object  $\Sigma_{+}^{\infty}$  $_{+C}^{\infty}B$  has a unique automorphism if  $B\to C$  is injective. This special case will appear frequently in Sections [3.1](#page-7-0) and [3.2.](#page-10-0)

### 3. STRUCTURES ON THE POINT-SET BICATEGORY

<span id="page-4-4"></span><span id="page-4-0"></span>In this section we use the action of multi-spans from [\(2.4\)](#page-3-1) to define an extensive formal structure on parameterized spectra. We begin by defining a bicategory of parameterized spectra. Then we add a shadow, compatibility of the external product with symmetry maps, and base change objects for maps of spaces. This is the point-set level structure whose associated homotopical structure is used in [\[MP22\]](#page-37-4) to construct periodic-point invariants.

We start with the simplest and most familiar structure, that parametrized spectra form a bicategory. This result first appears in [\[MS06\]](#page-37-6), see also [\[Shu08\]](#page-38-1). The more general result that any SMBF defines a bicategory with shadow appears in [\[PS12\]](#page-37-7). A more concise version of the following proof appears in [\[Mal,](#page-37-5) Thms 6.4.3 and 6.4.8]. We include this proof as a warm-up for the more elaborate proofs later in the section.

<span id="page-4-3"></span>**Proposition 3.1.** *There is a bicategory* OS*<sup>c</sup>* /*Top of parametrized spectra whose*

- *0-cells are spaces, and*
- *1- and 2-cells from A to B are the category*  $OS(A \times B)^c$  *of freely f-cofibrant* [\[Mal,](#page-37-5) Def 4.2.2] *parametrized orthogonal spectra over A* ×*B.*

The spans we define below make frequent use of the identity map, diagonal map, and projection map

1:  $A \rightarrow A$ ,  $\Delta: A \rightarrow A \times A$ ,  $\pi: A \rightarrow \star$ .

We concatenate maps to refer to products of these maps, for instance

$$
1\Delta\pi\colon A\times B\times C\to A\times B\times B
$$

<span id="page-4-2"></span>is the map that sends  $(a, b, c)$  to  $(a, b, b)$ .

*Remark* 3.2*.* In what follows we will often suppress isomorphisms of topological spaces that arise from associator or symmetry maps. This can be see above in the lack of parenthesizations in  $(2.3)$ . It will also notably appear in  $(3.15)$ .

When working with spans we will only be interested in the induced functors up to isomorphism and the omitted maps always induce isomorphisms.

In some cases we will include parenthesizations to help indicate the type of input spectra.

<span id="page-4-1"></span>

(A) Pentagon axiom

FIGURE 3.3. Commutative diagrams for a bicategory.

*Proof.* For this result we need to produce unit 1-cells, bicategorical composition, and associator and unitor maps that make the diagrams in Figure [3.3](#page-4-1) commute. We start with the unit and bicategorical composition.

• The unit  $U_A$  is the suspension spectrum of  $A_{+(A \times A)}$  over  $A \times A$ . Equivalently, it is unique object produced by the action of the multi-span



<span id="page-5-1"></span>• For  $X \in \mathcal{OS}^c(A \times B)$  and  $Y \in \mathcal{OS}^c(B \times C)$ , the bicategorical product  $X \circ Y$  is produced by the action of the multi-span

(3.4) 
$$
A \times C \xrightarrow{1\pi 1} A \times B \times C \xrightarrow{1\Delta 1} (A \times B) \times (B \times C).
$$

This is the first instance of parenthesization of the spaces along the bottom row by the type of 1-cell input as referenced in Remark [3.2.](#page-4-2) In this case we have a 1-cell over *A* ×*B* and one over  $B \times C$ .

We now turn to the associator and unitor maps. For these we will make use of the Beck-Chevalley isomorphisms along pullback squares in *Top*. The unit isomorphism provides a small example. The functor  $U_A \circ -$  is given by the action of the two spans along the bottom of the diagram below.

$$
A \times B
$$
  

$$
A \times A \times B
$$
  

$$
A \times A \times B
$$
  

$$
A \times A \times A \times B
$$
  

$$
A \times A \times A \times B
$$
  

$$
A \times B
$$

The square in this diagram is a pullback square of topological spaces. Therefore, the functor  $(1\Delta 1)^*(\Delta 11)$  can be replaced by  $(\Delta 1)$   $(\Delta 1)^*$ . Then composing with the projection maps gives the span with identity maps,

$$
\begin{array}{ccc}\n & & & 11 & A \times B & \xrightarrow{11} & A \times B. \\
 & & A \times B & & \xrightarrow{11} & A \times B.\n\end{array}
$$

- <span id="page-5-0"></span>• The unitors are the unique isomorphisms  $U_A \odot X \cong X \cong X \odot U_B$  given by identifying each of these expressions with the action of the identity multi-span in [\(3.5\)](#page-5-0). By Theorem [2.8](#page-3-3) this multi-span is rigid so any two functors isomorphic to this action have a unique isomorphism between them.
- The associator  $\alpha$  is the unique natural isomorphism  $(X \circ Y) \circ Z \cong X \circ (Y \circ Z)$ . It exists by identifying both sides with the action of the multi-span

$$
A \times D \xrightarrow{1 \pi \pi 1} \xrightarrow{A \times B \times C \times D} \xrightarrow{1 \Delta \Delta 1} (\overrightarrow{A} \times \overrightarrow{B}) \times (\overrightarrow{B} \times C) \times (\overrightarrow{C} \times \overrightarrow{D}).
$$

As with the unitor map, we use a Beck-Chevalley isomorphism along a pullback square of topological spaces to exchange pushforward and pullback functors.

Here we have an example of the suppressed isomorphisms referenced in Re-mark [3.2.](#page-4-2) The bottom right product is parenthesized differently for  $(X \circ Y) \circ Z$ and  $X \circ (Y \circ Z)$ , but the resulting spaces are isomorphic and we omit the map.

The axioms for a bicategory now follow similarly:

• For the pentagon axiom in Figure [3.3a,](#page-4-1) the span defining the four-fold product  $X \circ Y \circ Z \circ W$  is rigid:

(3.6) 
$$
\frac{1 \pi \pi \pi}{A \times B \times C \times D \times E} \xrightarrow{\qquad \qquad 1 \Delta \Delta \Delta 1} (A \times B) \times (B \times C) \times (C \times D) \times (D \times E).
$$

<span id="page-6-1"></span>All of the operations in Figure [3.3a](#page-4-1) are isomorphic to the action of this multispan. So by Theorem [2.8,](#page-3-3) they are uniquely isomorphic to the action of this multi-span and the diagram in Figure [3.3a](#page-4-1) commutes.

• The triangle unit axiom in Figure [3.3b](#page-4-1) follows by the same argument using the following multi-span.

$$
(3.7)
$$

(3.7) 
$$
A \times C \xrightarrow{1\pi 1} A \times B \times C \xrightarrow{1\Delta 1} (A \times B) \times (B \times C).
$$

The rhythm of this proof will be replicated for the remaining results in this section and for results in Section [6.](#page-24-0)

**Definition 3.8.** [\[Pon10\]](#page-37-3) A **shadow** on a bicategory B is a functor  $\langle \rangle$ :  $\mathcal{B}(A, A) \rightarrow$  **T** to a fixed category **T** for each 0-cell *A* and natural isomorphisms

$$
\theta: \langle X \odot Y \rangle \cong \langle Y \odot X \rangle
$$

<span id="page-6-0"></span>so the diagrams in Figure [3.9](#page-6-0) commute whenever they are defined.





# **Proposition 3.10.** *There is a shadow on the bicategory*  $\mathcal{OS}^c/\mathcal{Top}$ *.*

*Proof.* This proof is similar to the proof of Proposition [3.1](#page-4-3) in both structure and details.

• The shadow  $\langle X \rangle$  is defined by the action of the multi-span

$$
\star \stackrel{\pi}{\longleftarrow}^A \stackrel{\Delta}{\longrightarrow} (A \times A).
$$

• The rotator isomorphism  $\theta$ : « $X \circ Y$  ≥ « $Y \circ X$  exists by comparing both actions to that of the rigid span

$$
\star \stackrel{\pi\pi}{\longleftarrow} A \times B \longrightarrow A
$$
\n
$$
(A \times B) \times (B \times A).
$$

This span gives us a more interesting example of the suppressed maps refer-enced in Remark [3.2.](#page-4-2) The map  $\Delta\Delta$  has target  $A \times A \times B \times B$ . This is replaced by the isomorphic objects  $A \times B \times B \times A$  and  $B \times A \times A \times B$  to define the source and target of *θ*.

• The multi-span for the associator axiom in Figure [3.9a](#page-6-0) is

$$
\star \stackrel{\pi\pi\pi}{\longleftarrow} A \times B \times C \xrightarrow{\Delta\Delta\Delta} (A \times B) \times (B \times C) \times (C \times A).
$$

<span id="page-7-4"></span>• The multi-span for the unitor axiom in Figure [3.9b](#page-6-0) is



 $\Box$ 

We have now finished giving the more familiar structure exhibited by the bicategory OS*c* /*Top*. For readers primarily interested in the technique of proof, it may make sense to skip ahead to Section [4](#page-12-0) for the discussion of the descent of this structure to the homotopy category. The remainder of this section describes extensive additional structure that the bicategory OS*<sup>c</sup>* /*Top* enjoys. This structure is used in [\[MP22\]](#page-37-4).

<span id="page-7-0"></span>3.1. **Fuller structure.** In this subsection we describe the compatibility between the external product, permutations in the product bicategory, and the shadow on the bicategory OS*<sup>c</sup>* /*Top*.

**Definition 3.11.** Fix an integer  $n \geq 1$ . A **shadowed** *n***-Fuller structure** on a bicategory with shadow  $\mathscr B$  is the following data and conditions.

*i*. A pseudofunctor (strong functor) of bicategories

$$
\boxtimes: \underbrace{\mathscr{B} \times \ldots \times \mathscr{B}}_{n} \to \mathscr{B}.
$$

*ii*. If Γ is the pseudofunctor  $\mathcal{B} \times ... \times \mathcal{B} \to \mathcal{B} \times ... \times \mathcal{B}$  that permutes the leftmost  $\mathcal{B}$ to the right, a pseudonatural transformation

$$
\vartheta\colon\thinspace\boxtimes\circ\Gamma\to\boxtimes.
$$

*iii*. Compatibility between the 1-cells associated to Γ, the bicategorical product, and the shadow described by Figure [3.22.](#page-10-1)

<span id="page-7-3"></span>The goal of this section is the following result.

**Proposition 3.12.** *The bicategory* OS*<sup>c</sup>* /*Top has an n-Fuller structure.*

<span id="page-7-2"></span>The first step is to define the pseudofunctor.

**Lemma 3.13.** *There is a pseudofunctor (strong functor) of bicategories*

$$
\boxtimes\colon \underbrace{OS^c/\mathit{Top}\times\ldots\times OS^c/\mathit{Top}}_{n}\to OS^c/\mathit{Top}.
$$

<span id="page-7-1"></span>We call its composition isomorphism  $m_{\boxtimes}$ : ( $\boxtimes M_i$ )  $\odot$  ( $\boxtimes N_i$ )  $\cong$   $\boxtimes (M_i \odot N_i)$  and its unit  $i$  somorphism  $i$   $\boxtimes$  :  $\overset{\textstyle\cdot }{U}_{\prod_i A_i} \cong \boxtimes U_{A_i}.$ 

$$
\begin{array}{ccc}\n(\boxtimes M_i \odot \boxtimes N_i) \odot \boxtimes P_i \xrightarrow{\alpha} & \boxtimes M_i \odot (\boxtimes N_i \odot \boxtimes P_i) & U_{\prod A_i} \odot (\boxtimes M_i) \xrightarrow{\ell} & \boxtimes M_i \\
\downarrow_{\text{in} \odot m_{\boxtimes}} & \downarrow_{\text{in} \odot m_{\boxtimes}} & \downarrow_{\text{in} \odot m_{\boxtimes}} & \downarrow_{\text{in} \odot m_{\boxtimes}} & \downarrow_{\text{in} \odot m_{\boxtimes}} \\
\boxtimes (M_i \odot N_i) \odot \boxtimes P_i & \boxtimes M_i \odot \boxtimes (N_i \odot P_i) & (\boxtimes U_{A_i}) \odot (\boxtimes M_i) \xrightarrow{m_{\boxtimes}} & \boxtimes (U_{A_i} \odot M_i) \\
& \downarrow_{\text{in} \odot m_{\boxtimes}} & \downarrow_{\text{in} \odot m_{\boxtimes}} & \downarrow_{\text{in} \odot m_{\boxtimes}} & \downarrow_{\text{in} \odot m_{\boxtimes}} \\
& \downarrow_{\text{in} \odot m_{\boxtimes}} & \downarrow_{\text{in} \odot m
$$

(A) The pseudofunctor associator axiom

FIGURE 3.14. Commutative diagrams for a pseudofunctor.

*Proof.* As above, we first define the required functor, then produce the isomorphisms, and finally verify the required axioms (the diagrams in Figure [3.14\)](#page-7-1).

<span id="page-8-3"></span>• The pseudofunctor

 $(3.15)$ 

$$
\boxtimes: \underbrace{OS^c/\text{Top} \times \ldots \times OS^c/\text{Top}}_{n} \rightarrow OS^c/\text{Top}
$$

is defined on the 1-cells and 2-cells as the *n*-fold external smash product [\[Mal,](#page-37-5) Def 4.4.5], pulled back along the canonical isomorphism

$$
\prod_i (A_i \times B_i) \cong \prod_i A_i \times \prod_i B_i.
$$

• The composition isomorphism  $m_{\mathbb{R}}$ :  $(\mathbb{Z}M_i) \circ (\mathbb{Z}N_i) \cong \mathbb{Z}(M_i \circ N_i)$  arises because both sides can be identified with the action of the rigid multi-span

<span id="page-8-0"></span>
$$
\prod_i A_i \times \prod_i C_i \xleftarrow{1\pi 1} \prod_i A_i \times \prod_i B_i \times \prod_i C_i \xrightarrow{1\Delta 1} \prod_i (A_i \times B_i) \times (B_i \times C_i).
$$

This provides yet another example of the suppressed maps of Remark [3.2.](#page-4-2) The  $\text{input of } (\boxtimes M_i) \odot (\boxtimes N_i) \text{ is indexed on } \prod_i A_i \times \prod_i B_i \times \prod_i B_i \times \prod_i C_i \text{ while the result }$ is indexed on  $\prod_i A_i \times \prod_i C_i$ . The input of  $\mathbb{Z}(M_i \circ N_i)$  is indexed on  $\prod_i (A_i \times B_i) \times$  $(B_i \times C_i)$  while the output is indexed on  $\prod_i (A_i \times C_i)$ .

• Similarly, the unit isomorphism  $i_{\boxtimes}$ :  $U_{\prod_i A_i} \cong \boxtimes U_{A_i}$  comes from the rigid multispan

$$
\Pi_i A_i \times \Pi_i A_i \xrightarrow{\Delta} \Pi_i A_i \xrightarrow{\pi} \Pi_{\emptyset}.
$$

This is a manifestation of Remark [2.9](#page-3-4) where the parametrized spectrum has a unique automorphism.

• The pseudofunctor associator axiom (Figure [3.14b\)](#page-7-1) follows using the rigid multispan

$$
\prod_i A_i \times \prod_i D_i \xleftarrow{\text{ln } \pi_1 \prod_i A_i \times \prod_i B_i \times \prod_i C_i \times \prod_i D_i} \xrightarrow{\text{ln } \Delta_1} \prod_i (A_i \times B_i) \times (B_i \times C_i) \times (C_i \times D_i)
$$

• The pseudofunctor unitor axiom (Figure [3.14a\)](#page-7-1) follows using the rigid multi-span

$$
\prod_i A_i \times \prod_i B_i \longleftarrow \frac{\prod_i A_i \times \prod_i B_i}{\longrightarrow} \frac{\subseteq}{\prod_i (A_i \times B_i)}.
$$

Recall that Γ is the pseudofunctor  $\mathcal{B} \times \ldots \times \mathcal{B} \to \mathcal{B} \times \ldots \times \mathcal{B}$  that permutes the leftmost  $\mathscr{B}$  to the right.

<span id="page-8-2"></span>**Lemma 3.16.** *For the pseudofunctor defined in Lemma [3.13,](#page-7-2) there is a pseudonatural transformation*

<span id="page-8-1"></span>
$$
\vartheta\colon\thinspace\boxtimes\circ\Gamma\to\boxtimes.
$$

For each *n* tuple of objects  $(A_1,..., A_n)$  in  $\mathscr{B}$  we denote the associated object of  $\mathscr{B}(A_2 \times \ldots \times A_n \times A_1, A_1 \times \ldots \times A_n)$  by  $T_{(A_i)}$  and the natural isomorphisms by

(3.17) 
$$
\vartheta \colon T_{(A_i)} \circ (\boxtimes M_i) \xrightarrow{\cong} (\boxtimes M_{i+1}) \circ T_{(B_i)}
$$

for all  $M_i \in \mathcal{B}(A_i, B_i)$ .

*Proof.* We first define the  $T_{(A_i)}$  and  $\theta$  and then verify the required axioms (the di-agrams in Figure [3.18\)](#page-9-0) are satisfied for  $\vartheta$  and the maps defined in Proposition [3.1](#page-4-3) and Lemma [3.13.](#page-7-2)

<span id="page-9-0"></span>
$$
(T_{(A_i)} \odot \boxtimes M_i) \odot \boxtimes N_i \xrightarrow{\alpha} T_{(A_i)} \odot (\boxtimes M_i \odot \boxtimes N_i) \qquad U_{\prod A_{i+1}} \odot T_{(A_i)} \xleftarrow{\beta} T_{(A_i)} \odot U_{\prod A_i}
$$
\n
$$
\downarrow \text{id} \odot m_{\boxtimes} \qquad \qquad i_{\boxtimes} \odot \text{id} \qquad \qquad i_{\boxtimes} \odot \text{id} \qquad \qquad i_{\boxtimes} \odot \text{id} \qquad \qquad i_{\boxtimes} \odot \text{id}
$$
\n
$$
(\boxtimes M_{i+1} \odot T_{(B_i)}) \odot \boxtimes N_i \qquad T_{(A_i)} \odot \boxtimes (M_i \odot N_i) \qquad (\boxtimes U_{A_{i+1}}) \odot T_{(A_i)} \xrightarrow{\theta} T_{(A_i)} \odot (\boxtimes U_{A_i})
$$
\n
$$
\uparrow \qquad \qquad \downarrow \qquad \downarrow \qquad \qquad \downarrow \qquad \downarrow \qquad \downarrow \qquad \downarrow \qquad \qquad \downarrow \q
$$

(A) The pseudonatural composition axiom

FIGURE 3.18. Commutative diagrams for a pseudonatural transformation.

• The 1-cell  $T_{(A_i)}$  is the output of the multi-span

(3.19) 
$$
\Pi_i A_{i+1} \times \Pi_i A_i \longrightarrow \Pi_{\emptyset}.
$$

• The isomorphism  $\vartheta$  comes from comparison of both sides of the map in [\(3.17\)](#page-8-1) to the multi-span

<span id="page-9-3"></span>
$$
\Pi_i A_{i+1} \times \Pi_i B_i \xrightarrow{\Gamma_1} \Pi_i A_i \times \Pi_i B_i \xrightarrow{\cong} \Pi_i (A_i \times B_i).
$$

• The pseudonatural composition axiom (Figure [3.18a\)](#page-9-0) follows from the rigid multispan

$$
\prod_i A_{i+1} \times \prod_i C_i \xleftarrow{\Gamma \pi 1 \prod_i A_i \times \prod_i B_i \times \prod_i C_i} \xrightarrow{\phantom{a}1\Delta 1 \atop \phantom{a}1\Delta 1} \prod_i (A_i \times B_i) \times (B_i \times C_i).
$$

• The pseudonatural unit axiom (Figure [3.18b\)](#page-9-0) follows from the rigid multi-span

 $\Box$ 

$$
\Pi_i A_{i+1} \times \Pi_i A_i \longrightarrow {\Pi_i A_i} \longrightarrow {\Pi_i A_{i}}
$$

<span id="page-9-2"></span>**Lemma 3.20.** *For the 1-cells defined in the proof of Lemma [3.16](#page-8-2) there is a natural isomorphism*

 $\tau : \langle T_{(A_{i-1})} \circ \boxtimes Q_i \rangle \stackrel{\cong}{\twoheadrightarrow} \langle \!\langle Q_1 \circ \ldots \circ Q_n \rangle\!\rangle$ 

 $s$ o that the diagram in Figure [3.22](#page-10-1) commutes for all  $R_i \in \mathscr{B}(A_{i-1},B_i)$  and  $S_i \in \mathscr{B}(B_i,A_i)$ .

*Proof.* The isomorphism  $\tau$  comes from showing both source and target of  $(3.21)$  agree with the action of the span

<span id="page-9-1"></span>
$$
\star \longleftarrow {\prod_i A_i} \longrightarrow {\bigcap_{\substack{\Delta \\ \prod_i (A_{i-1} \times A_i)}}}
$$

<span id="page-10-1"></span>
$$
\langle T_{(A_{i-1})} \circ (\boxtimes R_i \circ \boxtimes S_i) \rangle \xrightarrow{\text{id} \circ m_{\boxtimes}} \langle T_{(A_{i-1})} \circ \boxtimes (R_i \circ S_i) \rangle \xrightarrow{\tau} \langle R_1 \circ S_1 \circ R_2 \circ \dots \circ R_n \circ S_n \rangle
$$
\n
$$
\downarrow^{\alpha}
$$
\n
$$
\langle (T_{(A_{i-1})} \circ \boxtimes R_i) \circ \boxtimes S_i \rangle
$$
\n
$$
\downarrow^{\beta \circ \text{id}}
$$
\n
$$
\langle (\boxtimes R_{i+1} \circ T_{(B_i)} \circ \boxtimes S_i) \rangle
$$
\n
$$
\downarrow^{\alpha}
$$
\n
$$
\langle (T_{(B_i)} \circ \boxtimes S_i) \circ \boxtimes R_{i+1} \rangle
$$
\n
$$
\downarrow^{\alpha}
$$
\n
$$
\langle T_{(B_i)} \circ (\boxtimes S_i \circ \boxtimes R_{i+1}) \rangle \xrightarrow{\text{id} \circ m_{\boxtimes}} \langle T_{(B_i)} \circ \boxtimes (S_i \circ R_{i+1}) \rangle \xrightarrow{\tau} \langle S_1 \circ R_2 \circ \dots \circ R_n \circ S_n \circ R_1 \rangle
$$



(Note the suppressed isomorphism on the pullback map.) The commuting diagram in Figure [3.22](#page-10-1) follows from the rigid multi-span

$$
\star \longleftarrow {\prod_i A_i \times \prod_i B_i \underbrace{\wedge \Delta}_{\prod_i (A_{i-1} \times B_i) \times (B_i \times A_i).} \square}
$$

<span id="page-10-0"></span>Lemmas [3.13,](#page-7-2) [3.16](#page-8-2) and [3.20](#page-9-2) together imply Proposition [3.12.](#page-7-3)

<span id="page-10-3"></span>3.2. **Base change.** In this subsection we encode the morphisms of *Top* as 1-cells in OS*c* /*Top*, and describe their compatibility with the *n*-Fuller structure.

**Definition 3.23.** If **S** is a cartesian monoidal 1-category, a **system of base-change objects for**  $\mathscr{B}$  **indexed by S** is the following data and conditions.

• A pseudofunctor  $[ \cdot \mathbf{S} \rightarrow \mathcal{B}$ .

• A vertical natural isomorphism *ν* filling the square of pseudofunctors

$$
\mathbf{S}^{\times n} \xrightarrow{\Pi} \mathbf{S} \n\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow \qquad \downarrow \n\mathcal{B}^{\times n} \xrightarrow{\boxtimes} \mathcal{B}
$$

where  $\prod$  denotes a fixed model for the *n*-fold product in **S**.

• An equality between  $T_{(B_i)}$  and the result of applying the functor  $[]$  to the isomorphism  $\prod B_i$   $\stackrel{\Gamma}{\rightarrow}$   $\prod B_{i+1}$  and compatibility between *τ* and *θ*. See Lemma [3.31.](#page-12-1)

<span id="page-10-4"></span>The goal of this section is to prove the following result.

**Proposition 3.24.** *The shadowed n-Fuller bicategory* OS*<sup>c</sup>* /*Top has base change objects indexed by the 1-category Top.*

<span id="page-10-2"></span>As in Section [3.1,](#page-7-0) we will prove this result in stages but they are shorter here. **Lemma 3.25.** *There is a pseudofunctor*  $[]$ :  $\text{Top} \rightarrow \text{OS}^c/\text{Top}$ *.* 

The 1-cell associated to a map of spaces  $B\stackrel{f}{\leftarrow}A$  will be written  $\bigl[B\stackrel{f}{\leftarrow}A \bigr]$  and the composition and unit isomorphisms will be denoted

 $m_{\text{II}}: \left[C \stackrel{g}{\leftarrow} B\right] \circ \left[B \stackrel{f}{\leftarrow} A\right] \cong \left[C \stackrel{gf}{\leftarrow} A\right], \qquad i_{\text{II}}: U_A \cong \left[A \stackrel{\text{id}}{\leftarrow} A\right].$ 

These maps need to satisfy the diagrams in Figure [3.26.](#page-11-0)

<span id="page-11-0"></span>
$$
\begin{array}{ccc}\n\left( \left[ D \stackrel{h}{\leftarrow} C \right] \circ \left[ C \stackrel{g}{\leftarrow} B \right] \right) \circ \left[ B \stackrel{f}{\leftarrow} A \right] \stackrel{\alpha}{\rightarrow} \left[ D \stackrel{h}{\leftarrow} C \right] \circ \left( \left[ C \stackrel{g}{\leftarrow} B \right] \circ \left[ B \stackrel{f}{\leftarrow} A \right] \right) & U_B \circ \left[ B \stackrel{f}{\leftarrow} A \right] \stackrel{\ell}{\longrightarrow} \left[ B \stackrel{f}{\leftarrow} A \right] \right] \\
\downarrow m_{\mathbb{U}} & \downarrow m_{\mathbb{
$$

(A) The pseudofunctor associator axiom

FIGURE 3.26. Commutative diagrams for the pseduofunctor in a system of base-change objects.

*Proof.* The base change object for a morphism  $f: A \rightarrow B$  in  $\mathcal{T}_{op}$  is the action of the multispan

 $(3.27)$ 

<span id="page-11-1"></span>
$$
B\times A \stackrel{(f,1)}{\longrightarrow} A \stackrel{\pi}{\longrightarrow} \prod_{\emptyset}.
$$

The pseudofunctor's composition isomorphism  $m_{\parallel}$  comes from the rigid span

$$
C \times A \overset{(gf,1)}{\longleftarrow} A \overset{\pi}{\longrightarrow} \prod_{\emptyset}.
$$

Its unit isomorphism  $i_{[]}$  is an identity map  $\left[ B \stackrel{=}{\leftarrow} B \right] = U_B$ .

The span in  $(3.27)$  is rigid, so base-change 1-cells have no nontrivial automorphisms. This implies that the diagrams in Figures [3.26a](#page-11-0) and [3.26b](#page-11-0) commute. (This is another manifestation of Remark [2.9.](#page-3-4))

<span id="page-11-3"></span>**Lemma 3.28.** *There is a vertical natural isomorphism ν filling the square of pseudofunctors*

(3.29)  
\n
$$
\begin{array}{ccc}\n & \text{Top}^{\times n} & \xrightarrow{\Pi} & \text{Top} \\
 & \downarrow^{\Pi} & & \downarrow^{\Pi} \\
 & & \text{(OS}^c / \text{Top})^{\times n} & \xrightarrow{\boxtimes} & \text{OS}^c / \text{Top}\n\end{array}
$$

 $where \prod denotes a \ fixed \ model for the n-fold \ product in \ Top.$ 

*Proof.* We need to show we have a natural isomorphism between the two routes in the diagram in [\(3.29\)](#page-11-2) satisfying the diagrams in Figure [3.30.](#page-12-2)

We define the canonical isomorphism

<span id="page-11-2"></span>
$$
v: \boxtimes \left[ B_i \stackrel{f_i}{\longleftarrow} A_i \right] \cong \left[ \prod B_i \stackrel{\prod f_i}{\longleftarrow} \prod A_i \right]
$$

by comparing source and target to the span

$$
\frac{\prod_i A_i}{\prod_i B_i \times \prod_i A_i} \xrightarrow{\prod_i A_i} \frac{\pi}{\prod_{\emptyset} \cdot \pi}
$$

As in the proof of Lemma [3.25,](#page-10-2) the diagrams in Figures [3.30a](#page-12-2) and [3.30b](#page-12-2) commute since the starting and ending points are isomorphic to a (rigid) base change object.  $\Box$ 

<span id="page-12-4"></span><span id="page-12-2"></span>

FIGURE 3.30. Commutative diagrams for the vertical natural isomorphism in a system of base-change objects.

In Figures [3.30a](#page-12-2) and [3.30b,](#page-12-2) all maps that happen to be equalities are notated as such. Some of these maps are not required by Definition [3.23](#page-10-3) to be equalities, but they are equalities in this construction of the bicategory.

<span id="page-12-1"></span> ${\bf Lemma \ 3.31.} \ \ Using \ the \ equality \ T_{(B_i)} = \left[\prod B_{i+1} \stackrel{\cong}{\leftarrow} \prod B_i \right] \ the \ diagram \ Figure \ 3.32 \ com {\bf Lemma \ 3.31.} \ \ Using \ the \ equality \ T_{(B_i)} = \left[\prod B_{i+1} \stackrel{\cong}{\leftarrow} \prod B_i \right] \ the \ diagram \ Figure \ 3.32 \ com {\bf Lemma \ 3.31.} \ \ Using \ the \ equality \ T_{(B_i)} = \left[\prod B_{i+1} \stackrel{\cong}{\leftarrow} \prod B_i \right] \ the \ diagram \ Figure \ 3.32 \ com$ *mutes.*

<span id="page-12-3"></span>
$$
\left[\prod B_{i+1} \stackrel{\cong}{\leftarrow} \prod B_i\right] \circ \left(\boxtimes \left[B_i \stackrel{p_i}{\leftarrow} E_i\right]\right) \stackrel{\theta}{\underset{\cong}{=}} \left(\boxtimes \left[B_{i+1} \stackrel{p_{i+1}}{\leftarrow} E_{i+1}\right]\right) \circ \left[\prod E_{i+1} \stackrel{\cong}{\leftarrow} \prod E_i\right]\right]
$$
\n
$$
\stackrel{\text{i do } v}{\underset{\text{i do } v}{=}} \left[\prod B_{i+1} \stackrel{\prod p_{i+1}}{\leftarrow} \prod E_{i+1}\right] \circ \left[\prod E_{i+1} \stackrel{\cong}{\leftarrow} \prod E_i\right]\right]
$$
\n
$$
\stackrel{\text{i do } v}{\underset{\cong}{=}} \left[\prod B_{i+1} \stackrel{\prod p_i}{\leftarrow} \prod E_i\right] \stackrel{m_{\text{II}}}{\underset{\cong}{=}} \prod E_i\right]
$$

FIGURE 3.32. The final compatibility axiom.

*Proof.* In the proof of Lemma [3.16,](#page-8-2)  $T_{(B_i)}$  is defined to be the result of the action of the span in [\(3.19\)](#page-9-3). This is the action of the span in [\(3.27\)](#page-11-1) when *f* is taken to be Γ.

As in the proof of Lemma [3.25,](#page-10-2) the diagram in Figure [3.32](#page-12-3) commutes since the starting and ending points are isomorphic to a (rigid) base change object.  $\Box$ 

<span id="page-12-0"></span>Lemmas [3.25,](#page-10-2) [3.28](#page-11-3) and [3.31](#page-12-1) together imply Proposition [3.24.](#page-10-4)

## 4. DEFORMABLE FUNCTORS

There are two bicategories of parametrized spectra: a point-set bicategory OS*<sup>c</sup>* /*Top*, in which the 2-cells are actual maps of parametrized spectra, and a homotopy bicategory  $\mathcal{E}$ x = Ho  $\mathcal{O}S^c/\mathcal{T}op$ , in which the 2-cells are morphisms in the homotopy category. To pass from the first to the second, we need to replace the product  $\odot$  by a derived product, and similarly for the other operations.

To do this we will use the framework of right-deformable functors from [\[DHKS04,](#page-37-8) 40.2], see also [\[Mal,](#page-37-5) §3.1]. In this section we recall how the framework works in general. **Definition 4.1.** A functor  $F: \mathbb{C} \to \mathbb{D}$  between categories with weak equivalences is **right-deformable** if the following data exists:

- <span id="page-13-2"></span>• a *full* subcategory **A** ⊆ **C** such that *F* preserves weak equivalences on **A**,
- a functor  $R: \mathbf{C} \to \mathbf{C}$  landing in **A**, and
- a natural transformation id  $\rightarrow R$  through weak equivalences. In other words, a natural equivalence  $X \stackrel{\sim}{\to} RX$  .

In particular, if **C** is a model category and *F* is right Quillen, we could take **A** to be the fibrant objects and *R* to be a fibrant replacement functor. To emphasize that we will *not* necessarily use fibrant replacement from a model structure, we will instead call the objects of **A** the **radiant objects**, and we say that *R* is a **radiant replacement** functor.

The **right-derived functor** of a right-deformable functor *F* is defined as

$$
\mathbb{R}F(X) := F(RX).
$$

The right-derived functor preserves weak equivalences, and therefore defines a map of homotopy categories

$$
HoC \to HoD.
$$

We say "the" right-derived functor of *F*, because up to canonical weak equivalence, R*F* does not depend on the choice of subcategory **A** or the radiant replacement functor *R*, see [\[DHKS04,](#page-37-8) 41.1, 41.2], [\[Rie17,](#page-38-2) 6.4], or [\[Mal,](#page-37-5) Prop 3.1.1]. In particular, if *F* preserves weak equivalences on all of **C**, its right-derived functor is canonically identified with *F* itself,

$$
\mathbb{R}F\simeq F.
$$

4.1. **Natural transformations.** For our applications it is not enough to have individual derived functors. We need to understand composites of derived functors and natural transformations between them.

**Definition 4.2** ([\[DHKS04,](#page-37-8) 42.3], [\[Mal,](#page-37-5) Def 3.2.9])**.** A list of composable, right-deformable functors

(4.3) 
$$
\mathbf{C} \longrightarrow \mathbf{C}_0 \xrightarrow{F_1} \mathbf{C}_1 \xrightarrow{F_2} \dots \xrightarrow{F_n} \mathbf{C}_n \longrightarrow \mathbf{D}
$$

is **coherently right-deformable** if:

- each functor  $F_i$  is right-deformable, meaning the following data exists for each  $1 \leq i \leq n$ :
	- **–** a *full* subcategory  $\mathbf{A}_{i-1}$  ⊆  $\mathbf{C}_{i-1}$  such that  $F_i$  preserves weak equivalences on **A***i*−1,
	- **–** a functor *Ri*−<sup>1</sup> : *Ci*−<sup>1</sup> → *Ci*−<sup>1</sup> whose image is contained in *Ai*−1, and
	- **–** a natural transformation id → *Ri*−<sup>1</sup> through weak equivalences,

and

•  $F_i(\mathbf{A}_{i-1}) \subset \mathbf{A}_i$ .

We take this approach to derived functors because it gives us good control over the compositions of functors and the natural transformations between them. The compatibility is described in detail in [\[Mal,](#page-37-5) Prop 3.2.4, Lem 3.2.10]; the relevant statements for us are the following results.

<span id="page-13-1"></span>**Proposition 4.4.** *For any list of coherently right deformable functors, the composite is right deformable. There is a canonical equivalence of functors (or isomorphism of functors on the homotopy category)*

$$
\mathbb{R}(F_n \circ \cdots \circ F_1) \xrightarrow{\simeq} \mathbb{R}F_n \circ \cdots \circ \mathbb{R}F_1.
$$

<span id="page-13-0"></span>**Proposition 4.5.** *Every natural transformation between right-deformable functors*

$$
\eta\colon F\Rightarrow G
$$

<span id="page-14-2"></span>*induces a canonical natural transformation of right-derived functors*

$$
\widetilde{\eta}\colon \mathbb{R} F \Rightarrow \mathbb{R} G
$$

*as functors on the homotopy category.*

This canonical natural transformation arises as the unique map making the following square commute, as functors  $HoC \rightarrow HoD$ .

$$
F \xrightarrow{\eta} G
$$
  

$$
\downarrow \qquad \qquad \downarrow
$$
  

$$
\mathbb{R}F - \xrightarrow{\tilde{\eta}} \mathbb{R}G.
$$

The uniqueness of  $\tilde{\eta}$  follows from the universal property of  $\mathbb{R}F$ , as the initial functor under *F* that preserves equivalences, see e.g. [\[DHKS04,](#page-37-8) 41.1, 41.2], [\[Rie17,](#page-38-2) 6.4].

It follows from this property that the assignment respects vertical compositions of natural transformations, and also respects horizontal compositions when the functors are coherently right-deformable:



<span id="page-14-1"></span>In other words, passage to right-derived functors is a partially-defined 2-functor on categories, functors, and natural transformations.

**Corollary 4.6.** *For coherently right-deformable functors, any isomorphism*

$$
G_m \circ \ldots \circ G_1 \cong F_n \circ \ldots \circ F_1
$$

*determines a canonical isomorphism in the homotopy category*

$$
\mathbb{R}G_m\circ\ldots\circ\mathbb{R}G_1\simeq\mathbb{R}F_n\circ\ldots\circ\mathbb{R}F_1.
$$

<span id="page-14-0"></span>**Corollary 4.7.** *Any commuting diagram between composites of coherently right-deformable functors induces a commuting diagram between the composites of right-derived functors.*

This is easiest to explain with an example. Suppose we have a commuting square of functors of the form

$$
F_2 \circ F_1 \longrightarrow G_3 \circ G_2 \circ G_1
$$
  
\n
$$
\downarrow \qquad \qquad \downarrow
$$
  
\n
$$
H_1 \longrightarrow J_2 \circ J_1
$$

and each of the lists  $\{(F_1, F_2), (G_1, G_2, G_3), (H_1), (J_1, J_2)\}$  is coherently right-deformable. Then Corollary [4.7](#page-14-0) says this induces a canonical diagram in the homotopy category



that also commutes. If, furthermore, the map  $F_2 \circ F_1 \to G_3 \circ G_2 \circ G_1$  was a composite of two maps  $F_2 \rightarrow G_3 \circ G_2$  and  $F_1 \rightarrow G_1$ , the corresponding map between right-derived functors would be a composite as well.

The upshot is the following. To verify that the functors, natural transformations, and coherences described in Section [3](#page-4-0) descend to the homotopy category, it is enough to show that every list of functors that appears is coherently right-deformable. For instance, the expression  $(X \odot Y) \odot Z$  corresponds to the list

$$
\mathcal{OS}^c(A \times B) \times \mathcal{OS}^c(B \times C) \times \mathcal{OS}^c(C \times D) \xrightarrow{\circ \times \mathrm{id}} \mathcal{OS}^c(A \times C) \times \mathcal{OS}^c(C \times D) \xrightarrow{\circ} \mathcal{OS}^c(A \times D).
$$

Once we verify that these lists are coherently right-deformable, we get the needed isomorphisms in the homotopy category from Proposition [4.5.](#page-13-0) The coherence between these isomorphisms follows from Corollary [4.7.](#page-14-0)

<span id="page-15-2"></span>*Remark* 4.8*.* The canonical isomorphisms produced by this approach admit an explicit description. If we model composites of right-derived functors by inserting copies of *R<sup>i</sup>* before every functor, then the canonical isomorphisms between composites of right-derived functors are defined by removing all of the intermediate copies of  $R_i$  from each composite, keeping the  $R_i$ s on the initial inputs, applying the point-set isomorphism, then re-inserting the intermediate copies of  $R_i$ . For an example, see Remark [5.7.](#page-17-0)

<span id="page-15-1"></span>4.2. **Extending radiant objects.** In many examples we have a "default" collection of radiant objects  $\mathbf{A}_i \subseteq \mathbf{C}_i$ , but we need to extend the collection to a slightly larger full subcategory that will contain the image of  $F(\mathbf{A}_{i-1})$ . In this section we formalize this extension procedure.

<span id="page-15-0"></span>**Lemma 4.9.** *Let*  $A \subseteq C$  *be a collection of radiant objects and*  $R: C \rightarrow C$  *a radiant replacement for a right-deformable functor*  $F: \mathbf{C} \to \mathbf{D}$ *.* 

*For any collection of objects*  $\{X_{\alpha}\}\$ ,  $X_{\alpha} \notin \mathbf{A}$ *, if each of the maps* 

$$
F(X_{\alpha}) \to F(RX_{\alpha})
$$

*is a weak equivalence, then the full subcategory* **A** ′ *whose objects are the X<sup>α</sup> and the objects of* **A***, is also a collection of radiant objects for F.*

In this case we say **A** can be **expanded** to contain the objects  ${X_a}$ .

*Proof.* It is enough to show that *F* applied to a weak equivalence between objects in **A** ′ is a weak equivalence. Let  $f: Y \to Z$  be a weak equivalence, where *Y* and *Z* are in **A**'. Then we have a commuting diagram

$$
F(Y) \longrightarrow F(RY)
$$
  
\n
$$
F(f) \qquad F(Rf)
$$
  
\n
$$
F(Z) \longrightarrow F(RZ)
$$

where the horizontal maps are induced by radiant replacement. The right vertical arrow is a weak equivalence since *RY* and *RZ* are in **A**. The top horizontal arrow is a weak equivalences, since either  $Y = X_\alpha$  for some  $\alpha$  and it is an equivalence by assumption, or  $Y ∈ A$  and then it is  $F$  of a weak equivalence in  $A$ . The bottom horizontal is a weak equivalence for the same reason. The left vertical arrow is therefore a weak equivalence by two out of three.  $\Box$ 

Our primary functor of interest is  $\circ$ , which is a functor of two variables  $F: \mathbb{C} \times \mathbb{C} \to \mathbb{D}$ . The most natural collection of radiant objects turns out to be a product category  $\mathbf{A} \times \mathbf{A}$ for some  $A \subseteq C$ , and the radiant replacement functor is a product  $R \times R$ . In this case, we apply Lemma [4.9](#page-15-0) by adding *pairs* ( $X_\alpha, Y_\alpha$ )  $\in \mathbb{C} \times \mathbb{C}$  to the subcategory  $\mathbf{A} \times \mathbf{A}$ , and checking <span id="page-16-3"></span>that the functor  $R \times R$  induces an equivalence  $F(X_\alpha, Y_\alpha) \to F(RX_\alpha, RY_\alpha)$ . Often, it is convenient to check this last condition by applying the copies of *R* one at a time:

$$
F(X_{\alpha}, Y_{\alpha}) \xrightarrow{\sim} F(RX_{\alpha}, Y_{\alpha}) \xrightarrow{\sim} F(RX_{\alpha}, RY_{\alpha}).
$$

<span id="page-16-0"></span>For instance, this happens in the proof of Lemma [5.10.](#page-18-0)

## 5. STRUCTURES ON THE HOMOTOPY BICATEGORY Ex

In this section we prove the structure we defined on the point-set bicategory OS*<sup>c</sup>* /*Top* in Section [3](#page-4-0) passes in a canonical way to the homotopy bicategory  $\mathcal{E}x = Ho(0S^c/\mathcal{I}_{op}).$ 

We define this homotopy bicategory as follows. The 0-cells of  $\mathcal{E}$ x are the 0-cells of  $OS<sup>c</sup>/Top$ , in other words, topological spaces. We form the hom categories of  $\mathcal Ex$  from those of OS<sup>c</sup>/Top by inverting the 2-cells in OS<sup>c</sup>(A × B) that are stable equivalences in the sense of [\[Mal,](#page-37-5) Def 5.1.1], or  $\pi_*$ -isomorphisms in the sense of [\[MS06,](#page-37-6) Def 12.3.4].

It remains to show that the functors of Section [3](#page-4-0) have associated derived functors on  $Ex$  that satisfy the same coherences.

The functors we studied in Section [3](#page-4-0) are composites of the external product, pushforward and pullback. We use the following result from [\[Mal,](#page-37-5) 1.0.2] to show that they descend to the homotopy category. A central role is played by the **level** *h***-fibrant spectra**. These are the spectra  $X \in \mathcal{OS}(B)$  for which each of the projection maps  $X_n \to B$  is a Hurewicz fibration [\[Mal,](#page-37-5) Definition 4.2.1].

<span id="page-16-1"></span>**Theorem 5.1.** *In each category* OS*<sup>c</sup>* (*B*) *of freely f -cofibrant parametrized orthogonal spectra*

- *the external smash product* ∧ *preserves all stable equivalences,*
- *the external smash product of level h-fibrant spectra is level h-fibrant,*
- *the pushforward f*! *preserves all stable equivalences, and level h-fibrant spectra if f is a Hurewicz fibration, and*
- *the pullback f* <sup>∗</sup> *preserves level h-fibrant spectra and stable equivalences between level h-fibrant spectra.*

*Furthermore, there is a strong symmetric monoidal level h-fibrant replacement functor P:*

$$
X \stackrel{\sim}{\rightarrow} PX, \qquad P\mathbb{S} \cong \mathbb{S}, \qquad P(X \overline{\wedge} Y) \cong PX \overline{\wedge} PY.
$$

Since  $\overline{\wedge}$  and  $f_1$  preserve all equivalences, we can take them to be their own right derived functor. The functor  $f^*$  preserves equivalences on level *h*-fibrant spectra, so it is right-deformable. We therefore get right-derived functors as in [\[Mal,](#page-37-5) Rmk 6.1.4]

$$
X\overline{\wedge}^{\mathbb{R}} Y = X \overline{\wedge} Y, \qquad (\mathbb{R}f_!)(X) = f_!(X), \qquad (\mathbb{R}f^*)(X) = f^*(PX).
$$

*Remark* 5.2*.* Theorem [5.1](#page-16-1) provides our default collection of radiant objects for this section: the freely *f* -cofibrant and level *h*-fibrant spectra. Unfortunately, not all spectra of interest are of this form. In particular, unit objects and base change objects are not level *h*-fibrant. So, when we need these 1-cells, we use the approach of Section [4.2](#page-15-1) to expand the collection of radiant objects.

The following result is the primary tool we will use to extend our collections of radiant objects. Let  $X \in \text{Top}(A \times B)$  be an arbitrary topological space over  $A \times B$ . We can add a disjoint copy of  $(A \times B)$  to *X* and take its fiberwise suspension spectrum:

$$
\Big(\Sigma^\infty_{+(A\times B)}X\Big).
$$

<span id="page-16-2"></span>The unit  $U_A$ , twisting object  $T_{A_i}$ , and more generally the base-change object  $\left[ B\stackrel{f}{\leftarrow} A \right]$ are all special cases of this construction.

<span id="page-17-5"></span>**Lemma 5.3.** *Let*  $W, X \in \textbf{Top}(A \times B), Y \in \mathcal{OS}^c(B \times C),$  and let  $f : W \to X$  be a weak equiv*alence. If the projections*  $W \rightarrow B$  *and*  $X \rightarrow B$  *are both Serre fibrations, or alternatively if each of the projections*  $Y_n \to B$  *is a Serre fibration, then the map induced by* f

$$
(\Sigma_{+(A\times B)}^{\infty}f)\odot id\colon \left(\Sigma_{+(A\times B)}^{\infty}W\right)\odot Y\to \left(\Sigma_{+(A\times B)}^{\infty}X\right)\odot Y
$$

*is an equivalence of spectra.*

*Proof.* By [\[Mal,](#page-37-5) Remark 6.4.7], the product  $\left(\sum_{i=1}^{+\infty}$  $\left(\mathbb{C}^\infty_{+(A\times B)} X\right) \odot Y$  is given at each spectrum level by the space-level circle product  $X_{+(A \times B)} \circ Y_n$ . By [\[Mal,](#page-37-5) Cor 6.5.2], this preserves equivalences provided either  $X \to B$  or  $Y_n \to B$  is a Serre fibration. The map

$$
(\Sigma_{+(A\times B)}^{\infty}f)\odot id
$$

is therefore an equivalence at every spectrum level, so it is a stable equivalence.  $\Box$ 

<span id="page-17-3"></span>5.1. **The bicategory**  $\&\mathbf{x}$ . In this section we show that  $\&\mathbf{x}$  is a bicategory. We first consider the composition product ⊙.

**Lemma 5.4.** *The functor*

$$
\odot: \mathcal{OS}^c(A \times B) \times \mathcal{OS}^c(B \times C) \longrightarrow \mathcal{OS}^c(A \times C)
$$

*preserves equivalences between pairs of level h-fibrant spectra and so is right-deformable.*

*Proof.* The first statement follows from Theorem [5.1.](#page-16-1) For the second, let the subcategory of radiant objects be pairs of level *h*-fibrant spectra. By Theorem [5.1,](#page-16-1) the functors that define ⊙ preserve weak equivalences between radiant objects.

<span id="page-17-1"></span>**Lemma 5.5.** *The lists*

$$
\circlearrowleft S^{c}(A \times B) \times \circlearrowleft S^{c}(B \times C) \times \circlearrowleft S^{c}(C \times D) \xrightarrow{\circlearrowleft \text{odd}} \circlearrowleft S^{c}(A \times C) \times \circlearrowleft S^{c}(C \times D) \xrightarrow{\circlearrowleft} \circlearrowleft S^{c}(A \times D)
$$
\n
$$
\circlearrowleft S^{c}(A \times B) \times \circlearrowleft S^{c}(B \times C) \times \circlearrowleft S^{c}(C \times D) \xrightarrow{\text{id} \times \circlearrowright} \circlearrowleft S^{c}(A \times B) \times \circlearrowleft S^{c}(B \times D) \xrightarrow{\circlearrowright} \circlearrowleft S^{c}(A \times D)
$$

*are coherently right-deformable.*

*Proof.* At each stage, we use the tuples of level *h*-fibrant spectra as our radiant objects. In the multi-span [\(3.4\)](#page-5-1) that defines  $\circ$ , the projection map  $\pi$  is a Hurewicz fibration. It follows that in the action of this span the final pushforward also preserves level *h*fibrancy. Therefore our subcategories of radiant objects are preserved by each instance of ⊙.

<span id="page-17-2"></span>**Corollary 5.6.** *The point-set associator isomorphism induces a canonical associator on the homotopy category*

 $\widetilde{\alpha}: (X \odot^{\mathbb{R}} Y) \odot^{\mathbb{R}} Z \simeq X \odot^{\mathbb{R}} (Y \odot^{\mathbb{R}} Z).$ 

*Proof.* By Lemma [5.5](#page-17-1) and Proposition [4.4](#page-13-1) the left-hand side is a right-derived functor of (*X* ⊙*Y*)⊙*Z*, and the right-hand side is a right-derived functor of *X* ⊙(*Y* ⊙*Z*). The point-set associator isomorphism therefore induces a canonical isomorphism of derived functors on the homotopy category by Corollary [4.6.](#page-14-1)  $\Box$ 

<span id="page-17-0"></span>*Remark* 5.7*.* By Remark [4.8,](#page-15-2) this derived associator is given explicitly as the composite

$$
P(PX \odot PY) \odot PZ \simeq (PX \odot PY) \odot PZ \stackrel{\alpha}{\cong} PX \odot (PY \odot PZ) \simeq PX \odot P(PY \odot PZ).
$$

<span id="page-17-4"></span>*Remark* 5.8. One might think that we are forced to make  $\mathbb{R}$ id ×  $\odot^{\mathbb{R}}$  the right-derived functor of id × ⊙ here, but that is not the case. We are perfectly free to choose id ×  $\circ^{\mathbb{R}}$ as the right-derived functor. Any two right-derived functors are canonically identified, so modifying the choice of right-derived functor does not affect the truth of any of the statements we make about it!

**Lemma 5.9.** *The diagram in Figure [3.3a](#page-4-1) commutes in the homotopy category, using the derived product* ⊙ <sup>R</sup> *and the derived associator map from Corollary [5.6:](#page-17-2)*

$$
(X \odot^{\mathbb{R}} (Y \odot^{\mathbb{R}} Z)) \odot^{\mathbb{R}} W \xrightarrow{\tilde{\alpha}} X \odot^{\mathbb{R}} ((Y \odot^{\mathbb{R}} Z) \odot^{\mathbb{R}} W)
$$
  

$$
\widetilde{\alpha} \odot^{\mathbb{R}} id \uparrow \qquad \qquad \downarrow id \odot^{\mathbb{R}} \widetilde{\alpha}
$$
  

$$
((X \odot^{\mathbb{R}} Y) \odot^{\mathbb{R}} Z) \odot^{\mathbb{R}} W \qquad X \odot^{\mathbb{R}} (Y \odot^{\mathbb{R}} (Z \odot^{\mathbb{R}} W))
$$
  

$$
\widetilde{\alpha} \downarrow \qquad \qquad \downarrow id \odot^{\mathbb{R}} \widetilde{\alpha}
$$
  

$$
(X \odot^{\mathbb{R}} Y) \odot^{\mathbb{R}} (Z \odot^{\mathbb{R}} W) \longrightarrow (X \odot^{\mathbb{R}} Y) \odot^{\mathbb{R}} (Z \odot^{\mathbb{R}} W)
$$

*Proof.* We verify the variant of Lemma [5.5](#page-17-1) in which we start with a product of four categories instead of three. The radiant objects are tuples of level *h*-fibrant spectra. This shows that every list appearing in Figure [3.3a](#page-4-1) is coherently right-deformable. By Corollary [4.7](#page-14-0) we get a commuting diagram between the right-derived functors. We identify the two upper vertical maps in this diagram as products  $\widetilde{\alpha} \circ^{\mathbb{R}}$  id by recalling that passage to the homotopy category respects horizontal composition of natural transformations.  $\Box$ 

The remaining diagrams from Section [3](#page-4-0) will follow by the same argument – the only thing to check is that every list of functors that arises is coherently right-deformable. However, to verify this for the unit isomorphism will require more care. Since  $U_A$  is not level *h*-fibrant this is our first use of the ideas in Section [4.2.](#page-15-1)

<span id="page-18-0"></span>**Lemma 5.10.** *The collection of radiant objects for* ⊙ *from Lemma [5.4](#page-17-3) can be expanded to contain pairs* (*X*,*Y*) *where one of X or Y is a unit 1-cell and the other is level h-fibrant.*

*Proof.* Recall that  $U_A = \sum_{+(A \times A)}^{\infty} A$  and  $PU_A = \sum_{+(A \times A)}^{\infty} A^I$ . The two maps

$$
A \xrightarrow{\operatorname{id}} A, \qquad A^I \xrightarrow{\operatorname{ev}_1} A,
$$

are Hurewicz fibrations. In particular, they are Serre fibrations, so by Lemma [5.3](#page-16-2) the maps

 $U_A \odot X \rightarrow PU_A \odot X$  and  $X \odot U_A \rightarrow X \odot PU_A$ 

are equivalences for any parametrized spectrum *X*.

Therefore, if *X* is level *h*-fibrant, the composites

$$
U_A \odot X \to PU_A \odot X \to PU_A \odot PX \quad \text{and} \quad X \odot U_A \to X \odot PU_A \to PX \odot PU_A
$$

are equivalences using Lemma [5.4.](#page-17-3) By Lemma [4.9,](#page-15-0) the subcategory of radiant objects can therefore be expanded to include all pairs of the form  $(U_A, X)$  and  $(X, U_A)$ .

**Corollary 5.11.** *The list*

$$
(*) \times \mathcal{OS}^{c}(A \times B) \xrightarrow{U_{A} \times id} \mathcal{OS}^{c}(A \times A) \times \mathcal{OS}^{c}(A \times B) \xrightarrow{\circ} \mathcal{OS}^{c}(A \times B)
$$

<span id="page-18-1"></span>*is coherently right-deformable.*

**Corollary 5.12.** *The point-set unitor isomorphisms induce canonical unitor isomorphisms on the homotopy category*

$$
\widetilde{\ell}: U_A \odot^{\mathbb{R}} X \simeq X, \qquad \widetilde{r}: X \odot^{\mathbb{R}} U_A \simeq X.
$$

<span id="page-18-2"></span>**Proposition 5.13.** Ex *is a bicategory with bicategorical composite given by the right derived functor of* ⊙ *from Lemma [5.4](#page-17-3) and U<sup>A</sup> as unit 1-cells.*

<span id="page-19-4"></span>*Proof.* Corollaries [5.6](#page-17-2) and [5.12](#page-18-1) give the associativity and unit isomorphisms. We verified Figure [3.3a](#page-4-1) in Lemma [5.9,](#page-17-4) so it remains to verify Figure [3.3b.](#page-4-1)

As in Lemma [5.9,](#page-17-4) this requires us to check that the list

<span id="page-19-1"></span>
$$
\mathcal{OS}^c(A \times B) \times (*) \times \mathcal{OS}^c(B \times C) \xrightarrow{\text{id} \times U_B \times \text{id}} \mathcal{OS}^c(A \times B) \times \mathcal{OS}^c(B \times B) \times \mathcal{OS}^c(B \times C)
$$
\n
$$
\xrightarrow{\odot \times \text{id}} \mathcal{OS}^c(A \times B) \times \mathcal{OS}^c(B \times C)
$$
\n
$$
\xrightarrow{\odot} \mathcal{OS}^c(A \times C)
$$

is coherently right-deformable and a similar check where we take the product on the right first. We do this by taking the radiant objects to be those spectra that are level *h*-fibrant, except in the category  $OS<sup>c</sup>(B \times B)$ , where it can be either the unit  $U_B$  or level *h*-fibrant.

As in the proof of Lemma [5.10,](#page-18-0) this is a valid subcategory of radiant objects. The radiant objects are preserved by the functors in  $(5.14)$  since  $-\circ U_B$  is isomorphic to the identity and so preserves level *h*-fibrant spectra.

By Corollary [4.7,](#page-14-0) we therefore get the commutativity of Figure [3.3b](#page-4-1) in the homotopy  $\Box$ category.

*Remark* 5.15*.* In [\[Mal,](#page-37-5) Theorem 6.4.8] several approaches are given for the construction of this bicategory. This proof is an elaboration on the second approach there. See also [\[MS06,](#page-37-6) 17.1.3] for the original construction by May and Sigurdsson.

<span id="page-19-0"></span>5.2. **Shadows and Fuller structure.** We now turn to the proof that Ex has shadows and an *n*-Fuller structure. As in the previous subsection, we separate the proof according to how we define the subcategory of radiant objects. We start with the cases where the radiant objects can be taken to be the level *h*-fibrant spectra and the unit objects *UA*:

<span id="page-19-3"></span>**Lemma 5.16.** *The functors*  $\Diamond$  *and*  $\boxtimes$  *are right deformable. There are isomorphisms*  $\theta$  *and*  $\widetilde{m}$ <sup>∞</sup> *making the diagrams in Figures* [3.9a,](#page-6-0) [3.9b](#page-6-0) and [3.14a](#page-7-1) commute for the right derived *functors of* ⊙,  $\langle \rangle$  and  $\boxtimes$ *.* 

*Proof.* We take the radiant objects for  $\langle \rangle$  to be the level *h*-fibrant spectra. Then Theo-rem [5.1](#page-16-1) implies  $\Diamond$  is right deformable. While we do not need this here, the projection map  $\pi: A \to *$  is a Hurewicz fibration and so the radiant objects are preserved.

The radiant objects for ⊠ are tuples of level *h*-fibrant spectra. Then ⊠ is right deformable and preserves radiant objects, since the external smash and pullback preserve radiant objects.

Since the radiant objects for the functors  $\odot$ ,  $\Diamond$  and  $\boxtimes$  are the same and these functors preserve radiant objects, compositions of them are coherently right-deformable. Then Proposition [4.4](#page-13-1) and Corollary [4.7](#page-14-0) provide the isomorphisms  $\theta$  and  $m_{\mathbb{N}}$  and the coherences in Figures [3.9a](#page-6-0) and [3.14a.](#page-7-1) For Figure [3.9b,](#page-6-0) the level *h*-fibrant spectra do not suffice, because they are not preserved by the functors in the following composite.

<span id="page-19-2"></span>(5.17) 
$$
\begin{aligned}\n\mathbb{OS}^c(A \times A) \times (*) & \xrightarrow{\text{id} \times U_A} \mathbb{OS}^c(A \times A) \times \mathbb{OS}^c(A \times A) \\
& \xrightarrow{\odot} \mathbb{OS}^c(A \times A) \\
& \xrightarrow{\otimes} \mathbb{OS}^c(*)\n\end{aligned}
$$

As in the proof of Proposition [5.13,](#page-18-2) we fix this by extending the collection of radiant objects in  $OS<sup>c</sup>(A \times A) \times OS<sup>c</sup>(A \times A)$  to include pairs where the first spectrum is level *h*fibrant and the second spectrum is the unit  $U_A$ . (We do not include the unit  $U_A$  as a radiant object in any of the other copies of  $OS<sup>c</sup>(A \times A)$ .) This shows that the list in [\(5.17\)](#page-19-2), and each of the other lists appearing in Figure [3.9b,](#page-6-0) is coherently right-deformable.  $\Box$ 

*Remark* 5.18. In the point set category,  $\langle PU_A \rangle$  and  $\langle U_A \rangle$  do not have the same homotopy type – one gives the suspension spectrum of the free loop space  $\Sigma_{+}^{\infty}LA$  , while the other gives only the constant loops  $\Sigma_{+}^{\infty}A$ . So, unlike ⊙, we cannot expand the radiant objects to include the unit  $U_A$  in every copy of  $OS^c(A \times A)$  above.

<span id="page-20-2"></span>Recall that the collection of radiant objects for  $\circ$  from Lemma [5.4](#page-17-3) consists of pairs of level *h*-fibrant spectra.

**Lemma 5.19.** *The collection of radiant objects for* ⊙ *from Lemma [5.4](#page-17-3) can be extended to*  $i$ nclude pairs where one or both of the spectra in the pair are base change 1-cells  $\bigl[ B \stackrel{f}{\leftarrow} A \bigr]$ .

*Proof.* This proof is similar to the proof of Lemma [5.10.](#page-18-0) We first prove that for any spectrum *Y* and any level *h*-fibrant spectrum *X* the following two maps are weak equivalences:

<span id="page-20-0"></span>(5.20) 
$$
\left[B \stackrel{f}{\leftarrow} A\right] \circ Y \longrightarrow P\left[B \stackrel{f}{\leftarrow} A\right] \circ Y
$$

<span id="page-20-1"></span>(5.21) 
$$
X \circ [B \stackrel{f}{\leftarrow} A] \longrightarrow X \circ P [B \stackrel{f}{\leftarrow} A]
$$

In each case, we tensor with the level *h*-fibrant replacement map.

As in the proof of Lemma [5.10,](#page-18-0) we recall that

$$
\left[B \stackrel{f}{\leftarrow} A\right] = \sum_{+ (B \times A)}^{\infty} A, \qquad P\left[B \stackrel{f}{\leftarrow} A\right] = \sum_{+ (B \times A)}^{\infty} (B^I \times_B A \times_A A^I).
$$

The maps  $A \to A$  and  $B^I \times_B A \times_A A^I \to A$  are both Serre fibrations, so by Lemma [5.3,](#page-16-2) the map [\(5.20\)](#page-20-0) is an equivalence.

On the other hand, if  $X \in \mathcal{OS}^c(C \times B)$  is level *h*-fibrant, then the maps  $X_n \to C \times B$ are Hurewicz fibrations, and therefore the projections  $X_n \to B$  are fibrations as well. By Lemma [5.3,](#page-16-2) the map [\(5.21\)](#page-20-1) is therefore an equivalence.

If *X* is level *h*-fibrant, the maps

$$
X \odot [B \stackrel{f}{\leftarrow} A] \stackrel{\sim}{\longrightarrow} X \odot P [B \stackrel{f}{\leftarrow} A] \stackrel{\sim}{\longrightarrow} PX \odot P [B \stackrel{f}{\leftarrow} A]
$$

are therefore both equivalences, so their composite is an equivalence. By Lemma [4.9,](#page-15-0) the subcategory of radiant objects can therefore be expanded to include all pairs of the form  $\left(X, \left[\begin{matrix} B & f \\ B & \leftarrow A \end{matrix}\right]\right)$ .

Similarly, if *Y* is level *h*-fibrant, the maps

$$
\left[B \stackrel{f}{\leftarrow} A\right] \circ Y \stackrel{\sim}{\longrightarrow} P\left[B \stackrel{f}{\leftarrow} A\right] \circ Y \stackrel{\sim}{\longrightarrow} P\left[B \stackrel{f}{\leftarrow} A\right] \circ PY
$$

are both equivalences, so the subcategory of radiant objects can be expanded to include all pairs of the form  $\left(\left[g \stackrel{f}{\leftarrow} A\right], Y\right)$  as well.

Finally, if we have two base-change 1-cells, we apply  $(5.20)$  first, then  $(5.21)$  to get the equivalence

$$
\left[C \stackrel{g}{\leftarrow} B\right] \circ \left[B \stackrel{f}{\leftarrow} A\right] \stackrel{\sim}{\longrightarrow} P\left[C \stackrel{g}{\leftarrow} B\right] \circ \left[B \stackrel{f}{\leftarrow} A\right] \stackrel{\sim}{\longrightarrow} P\left[C \stackrel{g}{\leftarrow} B\right] \circ P\left[B \stackrel{f}{\leftarrow} A\right].\right\right]
$$

Therefore the subcategory of radiant objects can be expanded to include all pairs of the  $\text{form } \left( \left[ C \stackrel{g}{\leftarrow} B \right], \left[ B \stackrel{f}{\leftarrow} A \right] \right)$  $\left| \right\rangle$ .

<span id="page-20-3"></span>*Remark* 5.22*.* We will only need Lemma [5.19](#page-20-2) for pairs in which one spectrum is level *h*-fibrant and the other is a twisting object  $T_{(A_i)}$  and for pairs in which both spectra are base change 1-cells. Since we don't gain much simplification in the proof when considering only these cases, we included the more general result.

*Remark* 5.23*.* It is important to note that we do not think of the extended collection of radiant objects in Lemma [5.19](#page-20-2) as a further extension of the collection from Lemma [5.10.](#page-18-0) Instead, we have two different expansions available to us. We are free to use either one at a given stage in a composite of right-deformable functors – we are not required to use both at the same time.

<span id="page-21-0"></span>**Lemma 5.24.** For the derived functors of  $\odot$ ,  $\Diamond$  and  $\boxtimes$  defined in Lemmas [5.4](#page-17-3) and [5.16](#page-19-3) *there are isomorphisms*  $i_{\nabla}$ ,  $\hat{\theta}$ , and  $\tilde{\tau}$  *satisfying the coherences in Figures* [3.14b,](#page-7-1) [3.18a,](#page-9-0) *[3.18b](#page-9-0) and [3.22.](#page-10-1)*

*Proof.* The proof is essentially the same as that of Lemma [5.9](#page-17-4) and Proposition [5.13.](#page-18-2) To define the isomorphism

$$
\widetilde{i}_{\boxtimes} \colon U_{\prod A_i} \simeq \boxtimes^{\mathbb{R}} U_{A_i}
$$

in the homotopy category the relevant composite of functors is

$$
(*) \xrightarrow{(U_{A_i})} \Pi \mathcal{O} \mathcal{S}^c(A_i \times A_i) \xrightarrow{\boxtimes} \mathcal{O} \mathcal{S}^c((\Pi A_i) \times (\Pi A_i)).
$$

The radiant objects consist of

- the single object of ∗ and
- the tuples in  $\prod \mathcal{OS}^c(A_i \times A_i)$  in which either
	- **–** every spectrum in the tuple is level *h*-fibrant or
	- $-$  every spectrum in the tuple is the unit  $U_{A_i}$ .

We do not allow tuples that have some level *h*-fibrant spectra and some units. (Note that it is unnecessary to choose the radiant objects in the final category  $\mathcal{O}S^c(\prod A_i \times \prod A_i)$ .) The functor  $\boxtimes = \overline{\wedge}$  preserves equivalences on these objects because it preserves  $\prod A_i$ ).) The functor  $\boxtimes = \overline{\wedge}$  preserves equivalences on these objects because it preserves all equivalences of freely *f* -cofibrant spectra. Then, as usual, Proposition [4.4](#page-13-1) defines the isomorphism  $i_{\boxtimes}$ .

We now check that each of the lists of functors in Figure [3.14b](#page-7-1) is coherently rightdeformable. Since all composites of functors are similar, we illustrate the argument with the composite in the bottom left corner:

$$
(*) \times \prod \mathcal{OS}^c(A_i \times B_i) \xrightarrow{U_{A_i} \times id} \prod \mathcal{OS}^c(A_i \times A_i) \times \prod \mathcal{OS}^c(A_i \times B_i)
$$
  

$$
\xrightarrow{\boxtimes \times \boxtimes} \mathcal{OS}^c(\prod A_i \times \prod A_i) \times \mathcal{OS}^c(\prod A_i \times \prod B_i)
$$
  

$$
\xrightarrow{\odot} \mathcal{OS}^c(\prod A_i \times B_i).
$$

Using the radiant objects and preservation properties above, we conclude by Corollary [4.7](#page-14-0) that the derived version of Figure [3.14b](#page-7-1) also commutes.

To define the isomorphism

(5.25) 
$$
\widetilde{\vartheta} \colon T_{(A_i)} \circ^{\mathbb{R}} (\boxtimes^{\mathbb{R}} M_i) \xrightarrow{\simeq} (\boxtimes^{\mathbb{R}} M_{i+1}) \circ^{\mathbb{R}} T_{(B_i)},
$$

the relevant composite of functors is

$$
(*) \times \prod \bigcirc \mathcal{S}^c(A_i \times B_i) \xrightarrow{T_{(A_i)} \times \boxtimes} \bigcirc \mathcal{S}^c(\prod A_{i+1} \times \prod A_i) \times \bigcirc \mathcal{S}^c(\prod A_i \times \prod B_i) \xrightarrow{\circ} \bigcirc \bigcirc \mathcal{S}^c(\prod A_{i+1}),
$$

and a similar composite with  $T_{(B_i)}$  on the right-hand side. In each of these categories, we take the radiant objects to be the tuples of level *h*-fibrant spectra, except in

$$
\mathcal{OS}^c(\prod A_{i+1}\times \prod A_i),
$$

where we also allow the twisting 1-cell  $T_{(A_i)}$ . Since  $T_{(A_i)}$  is a base change object for a homeomorphism it is, in particular, a base change object for a Serre fibration, so

$$
T_{(A_i)}\odot Y
$$

is level *h*-fibrant when *Y* is level *h*-fibrant. This allows us to check that radiant objects are preserved by the above functors. By Lemma [5.19,](#page-20-2) the weak equivalences between radiant objects are preserved as well, and so the above list of functors is coherently right-deformable. As before, Proposition  $4.4$  therefore defines the map  $\theta$ .

We similarly check that each of the lists of functors in Figure [3.18a](#page-9-0) is coherently right-deformable. We illustrate the argument with the entry in the top left corner:

$$
(*) \times \prod \bigcirc S^{c}(A_{i} \times B_{i}) \times \prod \bigcirc S^{c}(B_{i} \times C_{i})
$$
\n
$$
\xrightarrow{T_{(A_{i})} \times \Box \times \Box} \bigcirc S^{c}(\prod A_{i+1} \times \prod A_{i}) \times \bigcirc S^{c}(\prod A_{i} \times \prod B_{i}) \times \bigcirc S^{c}(\prod B_{i} \times \prod C_{i})
$$
\n
$$
\xrightarrow{\odot \times id} \bigcirc S^{c}(\prod A_{i+1} \times \prod B_{i}) \times \bigcirc S^{c}(\prod B_{i} \times \prod C_{i})
$$
\n
$$
\xrightarrow{\odot} \bigcirc S^{c}(\prod A_{i+1} \times \prod C_{i}).
$$

Using the radiant objects and preservation properties above, we conclude by Corollary [4.7](#page-14-0) that the derived version of Figure [3.18a](#page-9-0) also commutes.

When checking that the lists in Figure [3.18b](#page-9-0) are coherently right-deformable, we have to say a little more. We illustrate the argument with the composite in the bottom left corner:

$$
(*) \times \prod(*) \frac{T_{(A_i)} \times \prod(U_{A_i})}{\text{if } (A_i \otimes \bigcirc \bigcirc \bigcirc (1, A_{i+1} \times \prod A_i) \times \bigcirc \bigcirc (1, A_i \times A_i))}
$$
\n
$$
\xrightarrow{\text{if } \bigcirc \bigcirc \bigcirc (1, A_{i+1} \times \prod A_i) \times \bigcirc \bigcirc (1, A_i \times \prod A_i)}
$$
\n
$$
\xrightarrow{\bigcirc} \bigcirc \bigcirc^{c}(\prod A_{i+1} \times \prod A_i).
$$

The radiant objects are those tuples of spectra in which

- each spectrum is level *h*-fibrant, or
- every spectrum in the tuple is the "special" one either
	- $-$  the twisting 1-cell  $T_{(A_i)}$  in the category  $\mathcal{OS}^c(\prod A_{i+1} \times \prod A_i)$ ,
	- $-$  the unit  $U_{A_i}$  in  $\mathbb{O} \mathcal{S}^c(A_i \times A_i)$  for *every* value of *i*, or
	- $-$  the product  $\boxtimes U_{A_i}$  in  $\mathbb{O}S^c(\prod_i A_i \times \prod_i A_i)$ .

These radiant objects are all preserved by construction, and the weak equivalences between them are preserved by Lemma [5.19.](#page-20-2) We conclude by Corollary [4.7](#page-14-0) that the derived version of Figure [3.18a](#page-9-0) also commutes.

Finally, to define the isomorphism

$$
\widetilde{\tau}:\langle\!\langle T_{(A_{i-1})}\odot^{\mathbb{R}}\boxtimes^{\mathbb{R}} Q_{i}\rangle\!\rangle^{\!\mathbb{R}}\xrightarrow{\cong}\langle\!\langle Q_{1}\odot^{\mathbb{R}}\dots\odot^{\mathbb{R}} Q_{n}\rangle\!\rangle^{\!\mathbb{R}},
$$

the relevant composites of functors are

$$
(*) \times \prod \mathcal{OS}^{c}(A_{i-1} \times A_i) \xrightarrow{T_{(A_{i-1})} \times \boxtimes} \mathcal{OS}^{c}(\prod A_i \times \prod A_{i-1}) \times \mathcal{OS}^{c}(\prod A_{i-1} \times \prod A_i)
$$
  
\n
$$
\xrightarrow{\otimes} \mathcal{OS}^{c}(\prod A_i \times \prod A_i)
$$
  
\n
$$
\xrightarrow{\otimes} \mathcal{OS}^{c}(*),
$$
  
\n
$$
\prod \mathcal{OS}^{c}(A_{i-1} \times A_i) \xrightarrow{\otimes^{(n-1)}} \mathcal{OS}^{c}(A_1 \times A_1)
$$
  
\n
$$
\xrightarrow{\otimes} \mathcal{OS}^{c}(*).
$$

Here the subscripts on *A* are taken mod *n*. Again, we take the radiant objects in each category to be the level  $h$ -fibrant spectra and the twisting 1-cell  $T_{(A_i)}$  in  $\mathbb{OS}^c(\prod A_{i+1}\times\prod A_i)$ . These radiant objects and the equivalences between them are preserved, as before, by Lemma [5.19](#page-20-2) and the fact that  $T_{(A_i)} \circ -$  preserves level *h*-fibrant spectra. We see by a similar choice of radiant objects that each list of functors in Figure [3.22](#page-10-1) is coherently right-deformable. We conclude by Corollary [4.7](#page-14-0) that the derived version of Figure [3.22](#page-10-1) also commutes.  $\Box$ 

Together Lemmas [5.16](#page-19-3) and [5.24](#page-21-0) show  $\&x$  has a shadow and a compatible *n*-Fuller structure.

<span id="page-23-1"></span><span id="page-23-0"></span>5.3. **Base change objects.** It only remains to produce the natural transformations and coherences for a (derived) system of base change objects.

**Lemma 5.26.** For the derived functors of  $\circ$ ,  $\Diamond$  and  $\boxtimes$  defined in Lemmas [5.4](#page-17-3) and [5.16,](#page-19-3) *there are isomorphisms*  $\tilde{m}_{\text{[I]}}$ ,  $i_{\text{[I]}}$ , and  $\tilde{v}$  satisfying the coherences in Figures [3.26,](#page-11-0) [3.30](#page-12-2) *and [3.32.](#page-12-3)*

*Proof.* To define the isomorphism

$$
\widetilde{m}_{[]} \colon \left[ C \stackrel{g}{\leftarrow} B \right] \odot^{\mathbb{R}} \left[ B \stackrel{f}{\leftarrow} A \right] \simeq \left[ C \stackrel{gf}{\leftarrow} A \right],
$$

one of the relevant composites of functors is

$$
(*) \times (*) \xrightarrow{[g] \times [f]} \mathcal{OS}^c(C \times B) \times \mathcal{OS}^c(B \times A) \xrightarrow{\circ} \mathcal{OS}^c(C \times A).
$$

We take the radiant objects to be those pairs  $(X, Y) \in \mathcal{OS}^c(C \times B) \times \mathcal{OS}^c(B \times A)$  such that either

- *X* and *Y* are both the base change objects or
- they are both level *h*-fibrant.

The product ⊙ preserves equivalences on these pairs by Theorem [5.1](#page-16-1) and Lemma [5.19.](#page-20-2) The other composite is similar. Therefore this list is coherently right-deformable, so we get the derived isomorphism  $\widetilde{m}_{[]}$  by Proposition [4.4.](#page-13-1)

For the isomorphism

$$
\widetilde{i}_{[]} \colon U_A \simeq \left[ A \stackrel{\text{id}}{\longleftarrow} A \right],
$$

there is nothing to check, because a functor of the form  $(*) \rightarrow \mathcal{OS}^c(A \times A)$  is automatically coherently right-deformable. We therefore get the derived isomorphism  $i_{\parallel}$ , which is the same as the previous identification  $i_{\text{L}}$  on the point-set level.

We check that each of the lists of functors in Figures [3.26a](#page-11-0) and [3.26b](#page-11-0) is coherently right-deformable. We illustrate the argument with the composite in the top left corner:

$$
(*) \times (*) \times (*) \times (*) \xrightarrow{[h] \times [g] \times [f]} \mathcal{O}S^{c}(D \times C) \times \mathcal{O}S^{c}(C \times B) \times \mathcal{O}S^{c}(B \times A)
$$
  
\n
$$
\xrightarrow{\odot \times id} \mathcal{O}S^{c}(D \times B) \times \mathcal{O}S^{c}(B \times A)
$$
  
\n
$$
\xrightarrow{\odot} \mathcal{O}S^{c}(D \times A).
$$

We check this is coherently right-deformable by taking the radiant objects to be those tuples in which

- every spectrum in the tuple is the appropriate base-change object or composition of base-change objects, or
- every spectrum in the tuple is level *h*-fibrant.

The radiant objects are preserved by construction and weak equivalences are preserved by Theorem [5.1](#page-16-1) and Lemma [5.19.](#page-20-2) The other lists follow in a similar way. We conclude by Corollary [4.7](#page-14-0) that the derived versions of Figures [3.26a](#page-11-0) and [3.26b](#page-11-0) also commute.

To define the isomorphism

$$
\widetilde{v}: \boxtimes^{\mathbb{R}}\left[B_i \stackrel{f_i}{\leftarrow} A_i\right] \simeq \left[\prod B_i \stackrel{\prod f_i}{\leftarrow} \prod A_i\right]
$$

<span id="page-24-1"></span>one of the the relevant composites of functors is

$$
\Pi(*) \xrightarrow{\Pi[f_i]} \Pi \mathcal{O} \mathcal{S}^c(B_i \times A_i) \xrightarrow{\boxtimes} \mathcal{O} \mathcal{S}^c(\Pi B_i \times \Pi A_i).
$$

We take the radiant objects to be those tuples  $(X_i) \in \prod \mathcal{OS}^c(B_i \times A_i)$  such that either

- every  $X_i$  is the base change object  $\left[B_i \stackrel{f_i}{\leftarrow} A_i\right]$ , or
- every  $X_i$  is level *h*-fibrant.

The product  $\boxtimes$  preserves equivalences on these objects because it preserves all equivalences of freely *f* -cofibrant spectra. The other composite is similar. Therefore this list is coherently right-deformable, so we get the derived isomorphism  $\tilde{v}$  by Proposition [4.4.](#page-13-1)

We check that each of the lists of functors in Figures [3.30a,](#page-12-2) [3.30b](#page-12-2) and [3.32](#page-12-3) is coherently right-deformable. We illustrate the argument with the composite of functors in the top left corner of Figure [3.32:](#page-12-3)

$$
(*) \times \prod (*) \times \frac{T_{(B_i)} \times \prod [p_i]}{\text{S}^c} \text{OS}^c \left( \prod B_{i+1} \times \prod B_i \right) \times \prod \text{OS}^c (B_i \times E_i)
$$
  
\n
$$
\xrightarrow{\text{id} \times \boxtimes} \text{OS}^c \left( \prod B_{i+1} \times \prod B_i \right) \times \text{OS}^c \left( \prod B_i \times \prod E_i \right)
$$
  
\n
$$
\xrightarrow{\odot} \text{OS}^c \left( \prod B_{i+1} \times \prod E_i \right).
$$

The radiant objects are tuples of spectra in which

- every spectrum is level *h*-fibrant, or else
- every spectrum in the tuple is the "special" one either
	- $-$  the twisting 1-cell  $T_{(B_i)}$  in  $\mathcal{OS}^c(\prod B_{i+1} \times \prod B_i)$ ,
	- **−** the base change 1-cells  $\left[ B_i \stackrel{p_i}{\leftarrow} E_i \right]$  in  $\prod \theta S^c(B_i \times E_i)$ , or
	- $\bullet$  the product of base-change 1-cells  $\mathbb{E}\left[B_i \stackrel{p_i}{\leftarrow} E_i\right]$  in  $\mathbb{O}\mathcal{S}^c(\prod B_i \times \prod E_i).$

Radiant objects are preserved by construction, and equivalences between them are pre-served by Theorem [5.1](#page-16-1) and Lemma [5.19](#page-20-2) (and the fact that the product  $\boxtimes [B_i \stackrel{p_i}{\leftarrow} E_i]$  is itself isomorphic to a base change 1-cell).

The other lists follow in a similar way. We conclude by Corollary [4.7](#page-14-0) that the derived versions of Figures [3.30a,](#page-12-2) [3.30b](#page-12-2) and [3.32](#page-12-3) also commute.

<span id="page-24-0"></span>**Corollary 5.27.** Ex *is a bicategory with shadow, an n-Fuller structure, and base change objects.*

## 6. THE CASES OF  $G\mathcal{E}$ x,  $\mathcal{E}$ x<sub>B</sub>, AND  $G\mathcal{E}$ x<sub>B</sub>

The above arguments apply almost without change in  $G \&$  x, the bicategory of  $G$ equivariant parametrized spectra [\[Mal,](#page-37-5) Thm 7.4.5]. The only difference is that the isomorphisms are no longer unique as natural isomorphisms for equivariant maps – they are only unique if we insist that they are natural with respect to non-equivariant maps [\[Mal,](#page-37-5) Thm 7.2.4]. However, this is still enough to pick the isomorphisms and to confirm that they agree on the point-set level, and therefore also on the homotopy category.

**Corollary 6.1.** *G*Ex *is a bicategory with shadow, an n-Fuller structure, and base change objects.*

If we work relative to a base space *B*, more modifications are needed. Going back to the beginning, fix a base space *B* and consider the symmetric monoidal bifibration  $\mathcal{O}\mathcal{S}_{(B)}$ of parametrized orthogonal spectra over fibrations over *B* [\[Mal,](#page-37-5) Def 6.3.1]. The objects are parametrized spectra over spaces *E*, where *E* is equipped with Hurewicz fibrations  $E \rightarrow B$ . The maps are as in  $\mathcal{O}\mathcal{S}$ . The base category is the category **Fib**<sub>*B*</sub> of Hurewicz fibrations  $E \to B$  over the fixed base *B* and maps over *B*. The tensor product is the external smash product rel *B*. For a spectrum *X* over *D* and *Y* over *E*, this product  $X \overline{\wedge}_B Y$  is defined by pulling  $X \overline{\wedge} Y$  back along the canonical map  $D \times_B E \to D \times E$ .

Consider **multi-spans** in the category **Fib***B*. In other words, all the spaces have a fibration to *B* and all the maps are maps over *B*:

<span id="page-25-0"></span>(6.2) 
$$
C \longrightarrow B \longrightarrow B \longrightarrow B
$$

The definition of a **rigid** multi-span doesn't change, but the map  $E \to C \times A_1 \times ... \times A_n$ now factors through the fiber product  $C \times_B A_1 \times_B \ldots \times_B A_n$ , so we can measure rigidity by asking that the map to the fiber product be injective.

The diagonal map  $\Delta$  and projection map  $\pi$  change since our products of base spaces are taken relative to *B*:

<span id="page-25-1"></span>
$$
\Delta: A \to A \times_B A, \qquad \pi: A \to B.
$$

When we restrict to fibers over one point of *B*, these become the same maps from before. The multi-span in [\(6.2\)](#page-25-0) acts on the categories of orthogonal spectra as before:

(6.3) 
$$
\Pi_i \circ S^c(A_i) \longrightarrow \circ S^c(C)
$$

$$
(X_1, ..., X_n) \longmapsto f_!(g_1, ..., g_n)^*(X_1 \overline{\wedge}_B ... \overline{\wedge}_B X_n).
$$

Since  $\overline{\wedge}_B$  is defined by composing  $\overline{\wedge}$  with a pullback, this action is still composed of a smash product, a pullback, and a pushforward. Therefore it still preserves freely *f* cofibrant spectra and the rigidity theorem (Theorem [2.8\)](#page-3-3) still holds.

We define the point-set bicategory  $OS<sup>c</sup>/\textbf{Fib}_B$  as before:

- The 0-cells are fibrations *E* over *B*.
- The 1- and 2-cells from *A* to *C* are the category  $OS<sup>c</sup>(A \times_B C)$  of freely *f*-cofibrant parametrized orthogonal spectra over  $A \times_B C$ .
- The unit  $U_A$  is the suspension spectrum of  $A_{+(A \times_B A)}$  over  $A \times_B A$ . Equivalently, it is unique object produced by the action of the multi-span

$$
A \times_B A \xrightarrow{\Delta} A \xrightarrow{\pi} B.
$$

• For  $X \in \mathcal{OS}^c(A \times_B E)$  and  $Y \in \mathcal{OS}^c(E \times_B C)$ , the product  $X \odot_B Y$  is the action of the multi-span

$$
A \times_B C \xrightarrow{1\pi 1} A \times_B E \times_B C \xrightarrow{(1\Delta 1)} (A \times_B E) \times_B (E \times_B C).
$$

• The associator  $\alpha$  is the unique natural isomorphism  $(X \odot_B Y) \odot_B Z \cong X \odot_B (Y \odot_B Z)$ . It exists by identifying both sides with the action of the multi-span

$$
A \times_B D \xrightarrow{1 \pi \pi 1} A \times_B E \times_B C \times_B D \xrightarrow{1 \Delta \Delta 1} (A \times_B E) \times_B (E \times_B C) \times_B (C \times_B D).
$$

Since this multi-span is rigid, by Theorem [2.8,](#page-3-3) any two functors isomorphic to this action have one and only one isomorphism between them.

• Similarly, the unitors are the unique isomorphisms  $U_A \odot_B X \cong X \cong X \odot_B U_C$ .

<span id="page-26-3"></span>• The shadow  $\langle X \rangle$  is defined by the action of the multi-span

$$
B \stackrel{\pi}{\longleftarrow} A \stackrel{\Delta}{\longrightarrow} (A \times_B A).
$$

• The rotator isomorphism  $\theta$ :  $\langle X \circ_B Y \rangle \cong \langle Y \circ_B X \rangle$  exists by comparing both actions to that of the rigid span

$$
B \stackrel{\pi\pi}{\longleftarrow} A \times_B E \longrightarrow \Delta A
$$
  

$$
(A \times_B E) \times_B (E \times_B A).
$$

*Remark* 6.4*.* Note that the object *U<sup>A</sup>* is not the same as the object we denote using this symbol in Sections [3](#page-4-0) and [5,](#page-16-0) but we will rely on context to disambiguate. We will have a similar ambiguity for the base change objects  $T_{(A_i)}$  and we will also rely on context to distinguish.

**Proposition 6.5.** *These make*  $\mathbb{O}S^c$  *Fib<sub>B</sub> into a bicategory with shadow. Furthermore it has an n-Fuller structure in which*  $\boxtimes = \overline{\wedge}_B$ *, and base-change objects indexed by the category* **Fib***B.*

*Proof.* The arguments follow those in Section [3](#page-4-0) where every product is now a fiber product over  $B$ .

Before we proceed with the homotopy category, we need two lemmas. The first is a fiberwise version of Lemma [5.3.](#page-16-2) Let  $X \in \text{Top}(A \times_B C)$  be an arbitrary topological space over  $A \times_B C$ . We can add a disjoint copy of  $(A \times_B C)$  to X and take its fiberwise suspension spectrum:

$$
\Big(\Sigma_{+(A\times BC)}^\infty X\Big).
$$

As before, the unit  $U_A$ , twisting object  $T_{A_i}$ , and base-change object  $\left[C \stackrel{f}{\leftarrow} A\right]$ *B* are all special cases of this construction.

<span id="page-26-0"></span>**Lemma 6.6.** *Let*  $W, X ∈ \textbf{Top}(A ×_B C), Y ∈ \text{OS}^c(C ×_B D), and let f : W → X be a weak equiv$ *alence. If the projections*  $W \to C$  *and*  $X \to C$  *are both Serre fibrations, or alternatively if each of the projections*  $Y_n \to C$  *is a Serre fibration, then the map induced by f* 

$$
(\Sigma_{+(A\times BC)}^{\infty}f)\odot id\colon \left(\Sigma_{+(A\times BC)}^{\infty}W\right)\odot_B Y\to \left(\Sigma_{+(A\times BC)}^{\infty}X\right)\odot Y
$$

*is an equivalence of spectra.*

*Proof.* As in Lemma [5.3,](#page-16-2) the product  $\left(\sum_{i=1}^{\infty}$  $\left(\mathcal{L}_{+(A\times_B C)}X\right)\odot_B Y$  is given at each spectrum level by the space-level circle product  $X_{+(A\times BC)} \odot_B Y_n$ . By [\[Mal,](#page-37-5) Lem 6.5.1], this preserves equivalences provided either  $X \to C$  or  $Y_n \to C$  is a Serre fibration.

<span id="page-26-2"></span>The next lemma helps us check the fibration condition from Lemma [6.6.](#page-26-0) **Lemma 6.7.** *If*  $f : D \to E$  *is any map of fibrations over B, the composite* 

<span id="page-26-1"></span>(6.8) 
$$
D \times_{(E \times_B D)} (E \times_B D)^I \to (E \times_B D)^I \xrightarrow{\pi_1} D^I \xrightarrow{\text{ev}_1} D
$$

*is a Hurewicz fibration and homotopy equivalence.*

The product  $D\times_{(E\times_B D)} (E\times_B D)^I$  is formed using evaluation at  $0\in I,$  while the compos-ite in [\(6.8\)](#page-26-1) uses evaluation at  $1 \in I$ . The map  $D \to E \times_B D$  is the graph of f.

*Proof.* A homotopy inverse to the map in [\(6.8\)](#page-26-1) is the map  $d \rightarrow (d, c_{f(d),d})$  where  $c_{f(d),d}$ is the constant path at  $(f(d), d)$ . The endomorphism of *D* given by these maps is the identity. The endomorphism of  $D \times_{(E \times_B D)} (E \times_B D)^I$  given by these maps is homotopic to the identity by shrinking the path in  $E \times_B D$ .

<span id="page-27-0"></span>To see that [\(6.8\)](#page-26-1) is a fibration, we first rewrite it as the composite

(6.9) 
$$
D \times_{(E \times_B D)} (E \times_B D)^I \to (E \times_B D)^I \xrightarrow{\text{ev}_1} E \times_B D \to D.
$$

Then, suppose we are given the solid arrows in the following commutative diagram

(6.10) 
$$
X \xrightarrow{i_1 \times f_2} D \times_{(E \times_B D)} (E \times_B D)^I
$$

$$
X \times I \xrightarrow{H} \begin{cases} K_1 \times K_2 & \downarrow (6.9) \\ \downarrow & \downarrow (6.9) \end{cases}
$$

Expanding using the pullback in the top right corner, we obtain the two commutative diagrams below.

<span id="page-27-1"></span>
$$
(6.11)
$$
\n
$$
\begin{array}{ccc}\nX & \xrightarrow{f_2} & (E \times_B D)^I & X & f_1 \\
\downarrow^{k_0} & \downarrow^{k_2} & (E \times_B D) \times (E \times_B D) & X \times I - \frac{\pi}{K_2} \times (E \times_B D)^I & \downarrow^{k_2} \\
\downarrow^{k_0} & \downarrow^{k_2} & (E \times_B D) \times (E \times_B D) & X \times I - \frac{\pi}{K_2} \times (E \times_B D)^I & \downarrow^{k_2} \\
\downarrow^{k_3} & \downarrow^{k_4} & \downarrow^{k_5} & \downarrow^{k_6} \\
\downarrow^{k_6} & \downarrow^{k_7} & \downarrow^{k_8} & \downarrow^{k_9} \\
\downarrow^{k_8} & \downarrow^{k_9} & \downarrow^{k_9} & \downarrow^{k_9} & \downarrow^{k_9} \\
\downarrow^{k_9} & \downarrow^{k_8} & \downarrow^{k_9} & \downarrow^{k_9} & \downarrow^{k_9} \\
\downarrow^{k_9} & \downarrow^{k_9} & \downarrow^{k_9} & \downarrow^{k_9} & \downarrow^{k_9} & \downarrow^{k_9} \\
\downarrow^{k_9} & \downarrow^{k_9} & \downarrow^{k_9} & \downarrow^{k_9} & \downarrow^{k_9} & \downarrow^{k_9} \\
\downarrow^{k_9} & \downarrow^{k_9} & \downarrow^{k_9} & \downarrow^{k_9} & \downarrow^{k_9} & \downarrow^{k_9} & \downarrow^{k_9} \\
\downarrow^{k_9} & \downarrow^{k_9} \\
\downarrow^{k_9} & \downarrow^{k_9} \\
\downarrow^{k_9} & \downarrow^{k_9} \\
\downarrow^{k_9} & \downarrow^{k_9
$$

The product of the two evaluation maps  $(ev_0, ev_1)$  is a fibration since it is the dual of the inclusion  $\partial I \to I$ . The projection  $\pi 1: E \times_B D \to D$  is also a fibration since it is a pullback of  $E \rightarrow B$ . This gives the lift  $K_2$  in the left diagram in [\(6.11\)](#page-27-1).

Once  $K_2$  is chosen, the square on the right commutes since  $(f \times id) f_1 = ev_0(f_2)$ ev<sub>0</sub> $K_2i_0$ . In particular, the projection of ev<sub>0</sub> $K_2i_0$  onto *D* is  $f_1$ . Then the map  $K_1$  can be taken to be the composite  $f_1p: X \times I \to X \to D$ . The upper triangle in the right-hand square then commutes because  $p$  is a left inverse to  $i_0$ , and the bottom triangle commutes because  $(f \times id) f_1 p i_0 = (f \times id) f_1 = ev_0 K_2$ .

Now we proceed to the homotopy category. As in Section [5,](#page-16-0) the main task is to select collections of radiant objects that make each list of functors coherently deformable.

As in Lemma [5.4,](#page-17-3) the pairs of level *h*-fibrant spectra are a collection of radiant objects for the functor  $\odot_B$ . (For a spectrum in  $\odot S^c(A \times_B C)$ , these will be level *h*-fibrant with respect to the base space  $A \times_B C$ .)

<span id="page-27-3"></span>**Lemma 6.12.** *As in Lemmas [5.10](#page-18-0) and [5.19,](#page-20-2) this collection of radiant objects for* ⊙*<sup>B</sup> can be extended to contain either*

- *pairs in which one spectrum is a unit UA,*
- *pairs in which one spectrum is a twisting 1-cell*  $T_{(A_i)}$ , *or*
- *pairs in which both spectra are base change 1-cells*  $\left[ C \stackrel{f}{\leftarrow} A \right]$ *B .*

As in Remark [5.23,](#page-20-3) we do not need to extend the collection of radiant objects to contain all of these simultaneously.

*Proof.* For the first case where one spectrum is a unit *UA*, as in the proof of Lemma [5.10,](#page-18-0) it is enough to check that the map

<span id="page-27-2"></span>(6.13) 
$$
U_A \odot_B X \stackrel{\sim}{\rightarrow} (PU_A) \odot_B X
$$

is an equivalence. Recall that in the fiberwise setting over *B*, we have

$$
U_A = \sum_{+(A\times_B A)}^{\infty} A, \qquad PU_A = \sum_{+(A\times_B A)}^{\infty} (A\times_{(A\times_B A)}^{\infty} (A\times_B A)^I).
$$

<span id="page-28-2"></span>
$$
(6.14) \t\t A \xrightarrow{\mathrm{id}} A,
$$

(6.15) 
$$
A \times_{(A \times_B A)} (A \times_B A)^I \to (A \times_B A)^I \xrightarrow{\pi_1} A^I \xrightarrow{\text{ev}_1} A.
$$

But the first is an identity map, and the second is the special case of Lemma [6.7](#page-26-2) for the identity map id:  $A \rightarrow A$ . Therefore they are both fibrations, so [\(6.13\)](#page-27-2) is an equivalence.

For the second case where one spectrum is a twisting 1-cell  $T_{(A_i)}$ , we follow the same proof as the unit case, rather than following the proof of Lemma [5.19.](#page-20-2) It suffices to prove that the maps

$$
\left(\prod_{B} A_{i}\right) \stackrel{\text{id}}{\longrightarrow} \left(\prod_{B} A_{i}\right),
$$

(6.17)

$$
(\prod_B A_i) \times (\prod_B A_{i+1} \times_B \prod_B A_i) (\prod_B A_{i+1} \times_B \prod_B A_i)^I \to (\prod_B A_{i+1} \times_B \prod_B A_i)^I \xrightarrow{\pi_1} (\prod_B A_i)^I \xrightarrow{\text{ev}_1} (\prod_B A_i)
$$

are Hurewicz fibrations. This follows once again as a special case of Lemma [6.7.](#page-26-2)

Finally, for the case of two base-change objects it is enough to check that the map

$$
\left[D \stackrel{g}{\leftarrow} C\right]_B \odot_B \left[C \stackrel{f}{\leftarrow} A\right]_B \stackrel{\sim}{\rightarrow} P\left[D \stackrel{g}{\leftarrow} C\right]_B \odot_B P\left[C \stackrel{f}{\leftarrow} A\right]_B
$$

is an equivalence of spectra. Using [\[Mal,](#page-37-5) Remark 6.4.7] and the fact that *P* commutes with suspension spectra, this is a map of suspension spectra. Therefore, it suffices to focus attention at spectrum level zero and ignore the disjoint basepoint sections. This gives the inclusion of constant paths map

(6.18) 
$$
A \stackrel{\sim}{\rightarrow} C \times_{(D \times_B C)} (D \times_B C)^I \times_C (C \times_B A)^I \times_{(C \times_B A)} A.
$$

To see this is an equivalence, consider the composite

<span id="page-28-0"></span>
$$
C \times_{(D \times_B C)} (D \times_B C)^I \times_C (C \times_B A)^I \times_{(C \times_B A)} A \to C \times_C (C \times_B A)^I \times_{(C \times_B A)} A
$$
  
\n
$$
\cong (C \times_B A)^I \times_{(C \times_B A)} A
$$
  
\n
$$
\to A
$$

where the first piece applies the map in Lemma [6.7](#page-26-2) to the part to the left of the  $\times_c$ , the second removes the redundant *C*, and the third applies the "reflection" of the map in Lemma [6.7.](#page-26-2) These three maps are homotopy equivalences, and their composition is a left inverse to [\(6.18\)](#page-28-0). Therefore (6.18) is a homotopy equivalence as well.

<span id="page-28-1"></span>Now we define  $\mathcal{E}$ x<sub>*B*</sub> by inverting the stable equivalences in each category  $\mathcal{O}S^c(A\times_B C)$ .

**Proposition 6.19.** *Each functor in the definition of the shadowed bicategory, n-Fuller structure, and base-change objects for* OS*<sup>c</sup>* /**Fib***<sup>B</sup> has a right-derived functor. For every isomorphism and coherence axiom in these structures, the relevant lists of functors are coherently right-deformable.*

*Proof.* This proof is almost identical to that of Proposition [5.13](#page-18-2) and Lemmas [5.16,](#page-19-3) [5.24](#page-21-0) and [5.26.](#page-23-1) The role of Lemmas [5.10](#page-18-0) and [5.19](#page-20-2) is played by Lemma [6.12.](#page-27-3)

The only significant difference is that  $\boxtimes$  is replaced by  $\overline{\wedge}_B$ , which is the external smash product followed by a pullback along a diagonal map  $B \to B^n$ . This does not preserve all equivalences of freely *f* -cofibrant spectra over *A*, only those for which the composite maps  $X_n \to A \to B$  are Serre fibrations. So, we have to check that in each case it is applied, each of our radiant objects has this property.

For the level *h*-fibrant spectra *X* over  $A \rightarrow B$ , the map  $X_n \rightarrow A$  is a Hurewicz fibration. Since  $A \rightarrow B$  is a Hurewicz fibration by assumption, the composite  $X_n \rightarrow B$  is a Hurewicz <span id="page-29-2"></span>fibration. The only other cases are when *X* is either a unit or a base-change object. Since units are a special case of base-change objects, we only have to consider base-change objects.

If 
$$
X = [C \stackrel{f}{\leftarrow} A] = \sum_{+ (C \times_B A)}^{\infty} A
$$
, then at spectrum level 0, the projection to *B* is  

$$
A_{+ (C \times_B A)} \to C \times_B A \to B,
$$

which is a Hurewicz fibration because  $A \rightarrow C$  and  $C \rightarrow B$  are fibrations by assumption. At spectrum level *n*, we get the external smash product

$$
S^n \overline{\wedge} (A_{+(C \times_B A)})
$$

By [\[Mal,](#page-37-5) (2.3.4)], this is the pushout of  $A \times S^n$  and  $C \times_B A$  along A. Since all three of these spaces are *h*-fibrant over *B* and the map  $A \rightarrow (A \times S^n)$  has the homotopy extension property, this pushout is also *h*-fibrant over *B* by [\[Mal,](#page-37-5) Prop 2.1.9 and Lem 2.1.10].

In summary,  $\overline{\wedge}_B$  preserves equivalences on all of our radiant objects.

**Corollary 6.20.** Ex*<sup>B</sup> is a bicategory with shadow, an n-Fuller structure, and base change objects.*

Finally, this approach also applies to  $G\mathcal{E}x_B$ , the bicategory of G-equivariant parametrized spectra over spaces over *B*. We merely have to pick the isomorphisms that are natural with respect to all non-equivariant maps, and then they are unique and therefore coherent.

<span id="page-29-0"></span>**Corollary 6.21.** *G*Ex*<sup>B</sup> is a bicategory with shadow, an n-Fuller structure, and base change objects.*

## 7. GEOMETRIC FIXED POINTS AND REDUCTION OF GROUP ACTION

In this section we show the geometric fixed point functor and the forgetful functor from *G*-spectra to *H*-spectra for  $H \le G$  induce strong shadow functors

$$
\Phi^H : G \& \mathbf{x} \to W H \& \mathbf{x}, \qquad \iota_H^* : G \& \mathbf{x} \to H \& \mathbf{x}
$$

that commute up to vertical natural isomorphism (invertible icon in the sense of [\[Lac10\]](#page-37-9)) with the base-change objects. We then do the same for the bicategories of spectra over spaces over *B*, where *B* is a fixed space with trivial *G*-action:

> $\Phi^H$ :  $G \&g B \rightarrow W H \&g B$ ,  $\iota_E^*$  $H^*$ :  $G \, \mathcal{E} \mathbf{x}_B \rightarrow H \, \mathcal{E} \mathbf{x}_B$ .

This is the last result needed for [\[MP22,](#page-37-4) Thm 8.3].

This section will mimic Sections [2](#page-2-0) to [5.](#page-16-0) We first prove a rigidity result (Section [7.1\)](#page-29-1), then produce point set functors and natural transformations (Sections [7.2](#page-31-0) and [7.3\)](#page-32-0), and finally show this structure descends to the homotopy category (Section [7.5\)](#page-34-0).

<span id="page-29-1"></span>7.1. **Rigidity.** We start by proving a rigidity result for geometric fixed points  $\Phi^H$  that parallels Theorem [2.8.](#page-3-3)

**Definition 7.1.** [\[MM02,](#page-37-10) V§4], [\[MP22,](#page-37-4) §9] For a finite group *G* and a subgroup  $H \leq G$ , the **geometric fixed point** functor  $\Phi^H$ :  $GOS \rightarrow WHOS$  sends each parametrized *G*spectrum *X* to the coequalizer

$$
\bigvee_{V,W} F_{W^H}S^0 \overline{\wedge} \mathcal{J}_G^H(V,W) \overline{\wedge} X(V)^H \rightrightarrows \bigvee_V F_{V^H}S^0 \overline{\wedge} X(V)^H \longrightarrow \Phi^H X.
$$

Here  $V, W \subseteq \mathcal{U}$  run over finite-dimensional G-representations in some fixed complete *G*-universe U,  $\mathscr{J}_G$  is the indexing category for orthogonal *G*-spectra,  $\mathscr{J}_G^H$  denotes its  $H$ -fixed points, and  $F_{V^H}$  denotes the free spectrum at level  $V^H.$ 

<span id="page-30-7"></span>Note that if *X* is a *G*-spectrum over a space *A*, the geometric fixed points  $\Phi^H X$  is a  ${\rm spectrum~over}~A^H.$  See [\[MP22,](#page-37-4) Prop 9.3] for further discussion of geometric fixed points in the context of parameterized spectra.

<span id="page-30-0"></span>**Lemma 7.2.** [\[MP22,](#page-37-4) Prop 9.3] *The geometric fixed point functor commutes up to natural isomorphism with smash product, pullback, and pushforward on all freely f -cofibrant spectra. In particular, we have isomorphisms*

<span id="page-30-3"></span>(7.3) 
$$
\Phi^H(X \overline{\wedge} Y) \cong \Phi^H X \overline{\wedge} \Phi^H Y
$$

<span id="page-30-4"></span>
$$
\Phi^H(f^*Y) \cong (f^H)^* \Phi^H(Y)
$$

<span id="page-30-5"></span>(7.5) 
$$
\Phi^H(f_!Y) \cong (f^H)_! \Phi^H(Y).
$$

*Remark* 7.6*.* Lemma [7.2](#page-30-0) requires that we use the (CG) convention from [\[Mal,](#page-37-5) p. 6]. In other words, the levels of our spectra  $X_n$  are only assumed to be compactly generated, not compactly generated weak Hausdorff.<sup>[1](#page-30-1)</sup>

<span id="page-30-6"></span>Consider the functor

(7.7)  
\n
$$
\Pi_i G \circ S^c(A_i) \longrightarrow WH \circ S^c(C)
$$
\n
$$
(X_1, ..., X_n) \longmapsto \Phi^H f_1(g_1, ..., g_n)^* (X_1 \overline{\wedge} ... \overline{\wedge} X_n)
$$
\n
$$
\cong f_1^H (g_1^H, ..., g_n^H)^* (\Phi^H X_1 \overline{\wedge} ... \overline{\wedge} \Phi^H X_n)
$$

given by the action of a multi-span of equivariant maps

$$
C \xleftarrow{f} B \xrightarrow{(g_1,\dots,g_n)} A_1 \times \dots \times A_n
$$

<span id="page-30-2"></span>followed by the geometric fixed points functor.

**Theorem 7.8.** When  $H = G$ , this functor admits no nonidentity automorphisms. For *general H, the identity is the only automorphism that is natural when the morphisms are* extended to the H-equivariant maps, i.e. the morphisms in  $\prod_i H \mathcal{O} \mathcal{S}(\mathbf{A}_i)^c$ .

*Proof.* Without loss of generality  $H = G$ . Let  $\eta$  be any automorphism of the functor

$$
f_!^G(g_1^G,\ldots,g_n^G)^*(\Phi^G X_1 \overline{\wedge} \ldots \overline{\wedge} \Phi^G X_n).
$$

By the proof of [\[Mal,](#page-37-5) Thm 2.5.5], *η* must be the identity on any *n*-tuple of suspension  $\text{spectra } (\Sigma_{+A_1}^{\infty}(*), \ldots, \Sigma_{+A_n}^{\infty}(*))$ , on retractive *G*-spaces of the form  $(*)_{+A_i} = (*) \amalg A_i$ , where the point (\*) has trivial *G*-action and therefore maps to  $A_i^G \subseteq A_i$ .

The proof of [\[Mal17,](#page-37-11) Prop 3.17] carries over to the parametrized setting, as in [\[Mal,](#page-37-5) Thm 4.5.2], and shows that *η* must be the identity on any *n*-tuple of spectra

$$
(F_{V_1}(*)_{+A_1},\ldots,F_{V_n}(*)_{+A_n})
$$

formed by taking free spectra on nontrivial representations on the same one-point re- $\text{tractive spaces } (*)_{+A_i}.$ 

To further extend the class of spectra on which  $\eta$  is the identity, note that if  $f: X \to Y$ is a levelwise surjection of spectra, and  $\psi$  is an automorphism commuting with this map

$$
X \xrightarrow{\psi} X
$$
  
\n
$$
f \downarrow \qquad \qquad \downarrow f
$$
  
\n
$$
Y \xrightarrow{\psi} Y,
$$

<span id="page-30-1"></span><sup>&</sup>lt;sup>1</sup>It appears to also hold in the (CGWH) convention (for freely *f* -cofibrant spectra), but we do not provide a proof here.

if  $\psi$  is the identity on *X*, then  $\psi$  must also be the identity on *Y*. Now observe that  $\Phi$ <sup>*G*</sup> sends maps of spectra  $X \to Y$  for which  $X(V)^G \to Y(V)^G$  is surjective, to levelwise surjections of spectra. Furthermore,  $\overline{\wedge}$ ,  $g^*$ , and  $f_!$  all preserve levelwise surjections of spectra. So if we choose a map of tuples

$$
(X_1,\ldots,X_n)\to (Y_1,\ldots,Y_n)
$$

 $\sup X_i(V)^G \to Y_i(V)^G$  is a surjection, and if  $\eta$  is the identity on  $(X_1,\ldots,X_n),$ then *η* must also be the identity on  $(Y_1, \ldots, Y_n)$ .

If more generally  $\eta$  is the identity on a collection of such tuples  $(X_1^{\alpha},...,X_n^{\alpha})$ , and we choose maps  $X_i^{\alpha} \to Y_i$  such that the maps  $X_i^{\alpha}(V)^G \to Y_i(V)^G$  are jointly surjective, then  $t$  taking their wedge sum gives a levelwise surjective map  $\vee_a X_i^a \to Y_i$ , which then  $\Phi^G$ ,  $\overline{\wedge},$  $g^*$ , and  $f_!$  take to a levelwise surjective map of spectra. So if  $\eta$  is the identity on all of the tuples  $(X_1^{\alpha},...,X_n^{\alpha})$ , then  $\eta$  must also be the identity on  $(Y_1,...,Y_n)$ .

We now apply this principle to finish the proof. For any *n*-tuple of parametrized spectra  $(X_1,..., X_n)$ , the free-forget adjunction gives a collection of maps of spectra

$$
F_V(*)_{+A_i} \rightarrow X_i,
$$

one for each representation *V* and *G*-fixed point in  $X_i(V)$ . These maps are jointly surjective onto  $X_i(V)^G$  by construction. Therefore, by the above discussion, since  $\eta$  is the identity on every tuple of free spectra on one-point spaces, it must also be the identity on  $(X_1,..., X_n)$ .

<span id="page-31-1"></span><span id="page-31-0"></span>7.2. **Strong shadow functors.** We now use the geometric fixed points functor and Theorem [7.8](#page-30-2) to define a strong shadow functor on the point set category.

**Proposition 7.9.** *There is a functor*

$$
\Phi^H \colon G \circlearrowleft S^c / G \text{ Top} \to WH \circlearrowleft S^c / WH \text{ Top}
$$

*defined on 0-cells, hom categories, and shadow categories as follows.*

- *On objects, the image of a G-space A is the WH-space A H.*
- *On morphism categories, we take the functors*

$$
\Phi^H: G \circlearrowleft^c (A \times B) \to W H \circlearrowleft^c (A^H \times B^H).
$$

*To be precise,* Φ*<sup>H</sup> produces spectra over* (*A* × *B*) *<sup>H</sup>, and we pull back along the*  $\alpha$  *canonical identification to make these into spectra over*  $A^H\times B^H$  *.* 

• *On shadow categories, we take the functor*

$$
\Phi^H: G \circlearrowleft^c (*) \to W H \circlearrowleft^c (*).
$$

*Proof.* The natural isomorphisms

$$
m_{\Phi^H}: \Phi^H(X) \odot \Phi^H(Y) \cong \Phi^H(X \odot Y)
$$

$$
i_{\Phi^H}: U_{A^H} \cong \Phi^H(U_A)
$$

$$
s_{\Phi^H}: \langle \Phi^H(X) \rangle \cong \Phi^H \langle X \rangle
$$

on freely *f*-cofibrant spectra arise by commuting  $\Phi^H$  past the smash product, pullback, and pushforward using the isomorphisms in [\(7.3\)](#page-30-3), [\(7.4\)](#page-30-4), and [\(7.5\)](#page-30-5). In each case this is the unique isomorphism from Theorem [7.8](#page-30-2) that commutes with all *H*-equivariant maps.

To show this defines a strong shadow functor these isomorphisms must satisfy the coherences in Figure [7.10.](#page-32-1)

<span id="page-32-3"></span>• In Figure [7.10a,](#page-32-1) each of the objects is isomorphic to the action of the multi-span

$$
A \times D \xrightarrow{1\pi\pi 1} A \times B \times C \times D \xrightarrow{(A \times B) \times (B \times C) \times (C \times D)}
$$

(the one that defines the associator  $\alpha$ ), acting with geometric fixed points as in [\(7.7\)](#page-30-6). Here we again use the fact that the geometric fixed points functor commutes up to isomorphism with smash product, pullback and pushforward on freely *f* -cofibrant spectra (Lemma [7.2\)](#page-30-0). Then all functors in the diagram are isomorphic to one of the form considered in Theorem [7.8.](#page-30-2)

All of the maps are chosen to be natural with respect to *H*-equivariant maps of the inputs, and therefore the diagram commutes by Theorem [7.8.](#page-30-2)

• The coherence in Figure [7.10b](#page-32-1) follows similarly using the identity span

$$
A\times B \xrightarrow{= A\times B} A\times B.
$$

<span id="page-32-1"></span>• The coherence in Figure [7.10c](#page-32-1) follows similarly using the multi-span that defined the rotator *θ*

$$
\star \stackrel{\pi\pi}{\longleftarrow} A \times B \longrightarrow \Delta
$$
  
(*A* × *B*) × (*B* × *A*).

 $\Box$ 

$$
(\Phi^H X \circ \Phi^H Y) \circ \Phi^H Z \xrightarrow{\alpha} \Phi^H X \circ (\Phi^H Y \circ \Phi^H Z)
$$
\n
$$
m_{\Phi^H} \circ id \downarrow \qquad \qquad \downarrow id \circ m_{\Phi^H}
$$
\n
$$
\Phi^H (X \circ Y) \circ \Phi^H Z \qquad \qquad \Phi^H X \circ \Phi^H (Y \circ Z)
$$
\n
$$
m_{\Phi^H} \downarrow \qquad \qquad \downarrow
$$
\n
$$
\Phi^H ((X \circ Y) \circ Z) \xrightarrow{\alpha} \Phi^H (X \circ (Y \circ Z))
$$

(A) The shadow functor associator axiom

$$
U_{A^H} \odot \Phi^H X \longrightarrow \Phi^H X \qquad \langle \Phi^H X \odot \Phi^H Y \rangle \longrightarrow \langle \Phi^H Y \odot \Phi^H X \rangle
$$
  
\n
$$
i_{\Phi^H} \downarrow \qquad \qquad \uparrow \qquad m_{\Phi^H} \downarrow \qquad \qquad m_{\Phi^H}
$$
  
\n
$$
(\Phi^H U_A) \odot \Phi^H X \xrightarrow{m_{\Phi^H}} \Phi^H (U_A \odot X) \qquad \langle \Phi^H (X \odot Y) \rangle \qquad \langle \Phi^H (Y \odot X) \rangle
$$
  
\n(B) The shadow functor unitor axiom  
\n
$$
s_{\Phi^H} \downarrow \qquad \qquad s_{\Phi^H}
$$
  
\n
$$
\Phi^H \langle X \odot Y \rangle \longrightarrow \Phi^H \langle Y \odot X \rangle
$$

(C) The shadow functor rotator axiom

<span id="page-32-2"></span>FIGURE 7.10. Commutative diagrams for a strong shadow functor.

<span id="page-32-0"></span>7.3. **Natural transformations.** We now turn to compatibility between geometric fixed points and base change objects. We need to compare the functors

(7.11) 
$$
\Phi^H \circ [] \text{ and } [] \circ (-)^H : G \text{ Top} \to WH \text{OS}^c / WH \text{ Top}.
$$

These agree on 0-cells, so we can relate them using the less complicated concept of an icon [\[Lac10\]](#page-37-9).

<span id="page-33-4"></span><span id="page-33-0"></span>
$$
F(u) \xrightarrow{F(\mu)} F(v) \qquad 1_{F(x)} \xrightarrow{1_{G(x)}} F(u) \odot F(v) \xrightarrow{\alpha_u \odot \alpha_v} G(u) \odot G(v)
$$
\n
$$
\downarrow \alpha_u
$$
\n
$$
G(u) \xrightarrow{G(\mu)} G(v) \qquad F(1_x) \xrightarrow{\alpha_{1_x}} G(1_x)
$$
\n
$$
F(u) \odot F(v) \xrightarrow{\alpha_u \odot \alpha_v} G(u) \odot G(v)
$$
\n
$$
F(u) \odot v) \xrightarrow{\alpha_{u \odot v}} G(u) \odot v
$$
\n
$$
F(u) \odot v) \xrightarrow{\alpha_{u \odot v}} G(u) \odot v
$$
\n
$$
F(u) \odot v) \xrightarrow{\alpha_{u \odot v}} G(u) \odot v
$$
\n
$$
F(u) \odot v) \xrightarrow{\alpha_{u \odot v}} G(u) \odot v
$$
\n
$$
F(u) \odot v) \xrightarrow{\alpha_{u \odot v}} G(u) \odot v
$$
\n
$$
F(u) \odot v) \xrightarrow{\alpha_{u \odot v}} G(u) \odot v
$$
\n
$$
F(u) \odot v) \xrightarrow{\alpha_{u \odot v}} G(u) \odot v
$$

FIGURE 7.13. Commutative diagrams for a general icon.

<span id="page-33-1"></span>

(A) The pseudonatural unit axiom

(B) The pseudonatural composition axiom

FIGURE 7.16. Commutative diagrams for an icon  $\Phi^H \circ [] \to [] \circ (-)^H$ .

**Definition 7.12.** For bicategories C and D and functors  $F, G: \mathcal{C} \to \mathcal{D}$  such that  $F(x) =$ *G*(*x*) for every object *x* of C, an **icon**  $\alpha$ :  $F \rightarrow G$  consists of the following:

- For each morphism  $u: x \to y$  in C, a 2-cell  $\alpha_u: F(u) \to G(u)$ .
- For each 2-cell  $\mu: u \to v$  in C, the diagram in Figure [7.13a](#page-33-0) commutes.
- For each object  $x$  of  $C$ , the diagram in Figure [7.13b](#page-33-0) commutes.
- For each composable pair  $x \stackrel{u}{\rightarrow} y \stackrel{v}{\rightarrow} z$  in C, the diagram in Figure [7.13c](#page-33-0) commutes.

In our case [\(7.11\)](#page-32-2), the source bicategory is the 1-category *G Top*, which has no nontrivial 2-cells. This means we can dispense with Figure [7.13a](#page-33-0) and focus on Figures [7.13b](#page-33-0) and [7.13c,](#page-33-0) which become Figures [7.16a](#page-33-1) and [7.16b,](#page-33-1) respectively. Verifying these conditions is a straightforward application of Theorem [7.8.](#page-30-2)

<span id="page-33-2"></span>**Proposition 7.14.** *There is a vertical natural isomorphism (invertible icon) between the pseudofunctors in* [\(7.11\)](#page-32-2)*.*

*Proof.* These functors agree on objects, sending each *G*-space *A* to the *WH*-space *A H*. The isomorphism of spectra over  $A^H\times B^H$ 

<span id="page-33-3"></span>(7.15) 
$$
\eta : \Phi^H([B \stackrel{f}{\leftarrow} A]) \cong [B^H \stackrel{f^H}{\leftarrow} A^H]
$$

arises by commuting  $\Phi^H$  past the smash product, pullback, and pushforward, using the isomorphisms in [\(7.3\)](#page-30-3), [\(7.4\)](#page-30-4), and [\(7.5\)](#page-30-5). It is unique by the rigidity result in Theorem [7.8.](#page-30-2) The compatibilities in Figures [7.16a](#page-33-1) and [7.16b](#page-33-1) follow because these base-change objects  $\Box$  admit no nontrivial automorphisms.

*Remark* 7.17*.* The reader may have noticed that we have not included anything about the compatibility between the geometric fixed points functor and the Fuller structure. This is not because there is nothing to say, but because we have no need of it in [\[MP22\]](#page-37-4). The techniques illustrated here can be used to prove any additional compatibilities when needed.

<span id="page-34-7"></span>7.4. **Reduction of groups.** We similarly define the strong shadow functor of point-set bicategories

$$
(\iota_H)^* : G \circlearrowleft^c / G \mathit{Top} \to H \circlearrowleft^c / H \mathit{Top}
$$

and the vertical natural isomorphism (invertible icon) between the pseudofunctors

 $(\iota_H)^* \circ []$  and  $[] \circ (\iota_H)^* : G \text{ Top} \rightarrow H \circ \mathcal{S}^c / H \text{ Top}$ 

by forgetting the *G*-actions down to *H*-actions. Since this does not change the underlying set, we can pick the isomorphisms *m*, *s*, *i*, and *η* to simply be identity maps. This makes the following a less complicated version of Propositions [7.9](#page-31-1) and [7.14.](#page-33-2)

<span id="page-34-5"></span>**Proposition 7.18.** *The above data make*  $(\iota_H)^*$  *into a strong shadow functor and an invertible icon on the base-change objects.*

*Remark* 7.19*.* A rigidity theorem is not needed in this setting, but Theorem [2.8](#page-3-3) applies here. It tells us that the functor that acts by any rigid multi-span, followed by the forgetful functor, has a unique automorphism that commutes with all non-equivariant maps.

<span id="page-34-0"></span>7.5. **Descending to the homotopy category.** The process of showing that these functors descend to the homotopy category follows the same structure as in Section [5.](#page-16-0) We begin by recalling the following analog of Theorem [5.1.](#page-16-1)

<span id="page-34-4"></span>**Lemma 7.20.** [\[MP22,](#page-37-4) Lem 9.6] *The geometric fixed point functor preserves freely f cofibrant spectra and equivalences between them.*

Unfortunately, we do not have a proof that Φ*<sup>H</sup>* preserves level *h*-fibrancy for freely *f* cofibrant spectra.[2](#page-34-1) As a result, we cannot use the level *h*-fibrant spectra as our radiant objects, since they will not be preserved by  $\Phi^H$ . To get around this, we introduce a more restrictive class of "*P*-fibrant" spectra that are preserved by  $\Phi^H$ .

<span id="page-34-2"></span>**Lemma 7.21.** *For freely f -cofibrant spectra X, there is a natural isomorphism*

$$
\Phi^H P X \cong P \Phi^H X.
$$

*Proof.* The functor P is a composition of a pullback and a pushforward ([\[Mal,](#page-37-5)  $\S$ 2.7]), so this follows from the isomorphisms  $(7.4)$  and  $(7.5)$  from Lemma [7.2.](#page-30-0)

<span id="page-34-6"></span>**Definition 7.22.** A spectrum  $Y$  ∈  $GOS<sup>c</sup>(C)$  is  $P$ **-fibrant** if it is isomorphic to a spectrum of the form  $f_!g^*PX$ , where  $X \in GOS^c(A)$ ,  $g: B \to A$  is any map, and  $f: B \to C$  is an (equivariant) Hurewicz fibration.

<span id="page-34-3"></span>**Lemma 7.23.** *P-fibrant spectra are preserved by*

- *geometric fixed points* Φ*H,*
- *external smash product* ∧*,*
- *pullback g* ∗ *, and*
- *pushforward along any (equivariant) Hurewicz fibration f*! *.*

*As a result, they are also preserved by*  $\circ$  *and*  $\&$ .

*Proof.* Suppose  $Y = f_!g^*X$  is *P*-fibrant. Then  $\Phi^H Y = \Phi^H(f_!g^*PX) \cong f_!g^*P(\Phi^H X)$ , using Lemma [7.2](#page-30-0) and Lemma [7.21.](#page-34-2) The spectrum  $\Phi^H X$  is freely *f*-cofibrant, and therefore Φ*HY* is *P*-fibrant.

Since the pushforward is the last functor applied in the definition of a *P*-fibrant spectrum, a pushforward of a *P*-fibrant spectrum along a fibration is again *P*-fibrant. If we

<span id="page-34-1"></span> $2$ The issue is that the coequalizer destroys fibrancy. If we attempt to simplify the coequalizer to a colimit of pushouts of fibrant pieces, as in the proof of [\[Mal,](#page-37-5) Thm 4.4.6], we would either have to use a free presentation, in which case we can't make the input spaces fibrant, or a semifree presentation, in which case we lose control of  $\Phi^H$  because it does not have a nice formula on semifree inputs.

instead take a pullback  $h^*f_!g^*PX$ , we use the Beck-Chevalley isomorphism from [\(2.1\)](#page-2-1) to rewrite it as  $j_! k^* g^* P X$ , where  $j$  is a pullback of  $f$  and is therefore a fibration. Therefore a pullback of a *P*-fibrant spectrum is *P*-fibrant.

Theorem [5.1,](#page-16-1) along with the observation that  $OS$  is a symmetric monoidal bifibration, shows that for *P*-fibrant spectra  $Y_1$  and  $Y_2$  we have

$$
Y_1 \overline{\wedge} Y_2 \cong [(f_1)_!(g_1)^*PX_1] \overline{\wedge} [(f_2)_!(g_2)^*PX_2]
$$
  
\n
$$
\cong (f_1 \times f_2)_! ([(g_1)^*PX_1] \overline{\wedge} [(g_2)^*PX_2])
$$
  
\n
$$
\cong (f_1 \times f_2)_! (g_1 \times g_2)^* (PX_1 \overline{\wedge} PX_2)
$$
  
\n
$$
\cong (f_1 \times f_2)_! (g_1 \times g_2)^* P(X_1 \overline{\wedge} X_2).
$$

<span id="page-35-0"></span>Since  $X_1 \overline{\wedge} X_2$  is freely *f*-cofibrant,  $Y_1 \overline{\wedge} Y_2$  is *P*-fibrant. □

**Lemma 7.24.** *Every P-fibrant spectrum is level h-fibrant. Each of the operations in Lemma [7.23](#page-34-3) preserves equivalences between P-fibrant spectra.*

*Proof.* Follows directly from Theorem [5.1](#page-16-1) and Lemma [7.20.](#page-34-4) □

<span id="page-35-1"></span>We now show that the structures from Propositions [7.9](#page-31-1) and [7.14](#page-33-2) descend to the homotopy category.

**Proposition 7.25.** *The strong shadow functor* Φ*<sup>H</sup> from Proposition [7.9](#page-31-1) descends to a strong shadow functor on homotopy bicategories*

$$
\Phi^H: G \, \mathcal{E} \mathbf{x} \to W H \, \mathcal{E} \mathbf{x}.
$$

*Proof.* On 0-cells we again take each *G*-space *A* to the *WH*-space  $A^H$ . On morphism categories, we take right-derived functor

$$
\Phi^H : \text{Ho}G \circlearrowleft \otimes (A \times B)^c \to \text{Ho} \, \mathcal{W} \, H \circlearrowleft \otimes (A^H \times B^H)^c
$$

.

By Lemma [7.20,](#page-34-4)  $\Phi^H$  preserves all equivalences, so the right-derived functor is  $\Phi^H$  itself. We define the isomorphisms

$$
\widetilde{m}_{\Phi^H} : \Phi^H(X) \circ^{\mathbb{R}} \Phi^H(Y) \simeq \Phi^H(X \circ^{\mathbb{R}} Y)
$$

$$
\widetilde{i}_{\Phi^H} : U_{A^H} \simeq \Phi^H(U_A)
$$

$$
\widetilde{s}_{\Phi^H} : \langle \Phi^H(X) \rangle^{\mathbb{R}} \simeq \Phi^H \langle X \rangle^{\mathbb{R}}
$$

from their point-set versions by checking that each composite of functors that appears is coherently right-deformable. For  $\tilde{m}_{\Phi^H}$  and  $s_{\Phi^H}$ , we take the tuples of *P*-fibrant spectra as our radiant objects. For  $\tilde{i}_{\Phi^H}$ , we follow the first part of Lemma [5.24](#page-21-0) and include the unit objects  $U_A \in \mathcal{OS}^c(A \times A)$  and  $U_{A^H} \in \mathcal{OS}^c(A^H \times A^H)$  in the subcategory of radiant objects. Proposition [4.4](#page-13-1) then defines the maps  $\widetilde{m}_{\Phi^H}$ ,  $\widetilde{i}_{\Phi^H}$ , and  $\widetilde{s}_{\Phi^H}$ .

We now check that each of the lists of functors in Figure [7.10](#page-32-1) is coherently rightdeformable. We illustrate the argument with the composite in the middle left entry in Figure [7.10a:](#page-32-1)

$$
G \circlearrowleft^c (A \times B) \times G \circlearrowleft^c (B \times C) \times G \circlearrowleft^c (C \times D) \xrightarrow{\circlearrowleft \times id} G \circlearrowleft^c (A \times C) \times G \circlearrowleft^c (C \times D)
$$
\n
$$
\xrightarrow{\Phi^H \times \Phi^H} W H \circlearrowleft^c (A^H \times C^H) \times W H \circlearrowleft^c (C^H \times D^H)
$$
\n
$$
\xrightarrow{\circ} W H \circlearrowleft^c (A^H \times D^H).
$$

In this case we take the radiant objects to be all tuples of *P*-fibrant spectra. They are preserved by Lemma [7.23,](#page-34-3) and weak equivalences between them are preserved by Lemma [7.24.](#page-35-0)

<span id="page-36-2"></span>We also illustrate the argument with the bottom left entry in Figure [7.10b:](#page-32-1)

$$
(*) \times G \circlearrowleft^c (A \times B) \xrightarrow{U_A \times \text{id}} G \circlearrowleft^c (A \times A) \times G \circlearrowleft^c (A \times B)
$$
  

$$
\xrightarrow{\Phi^H \times \Phi^H} W H \circlearrowleft^c (A^H \times A^H) \times W H \circlearrowleft^c (A^H \times B^H)
$$
  

$$
\xrightarrow{\circ} W H \circlearrowleft^c (A^H \times B^H).
$$

In this case we take the radiant objects to be all P-fibrant spectra, except in  $GOS<sup>c</sup>(A \times B)$ *A*) we also take  $U_A$ , and in  $W H \mathbb{O} \mathbb{S}^c (A^H \times A^H)$  we also take  $\Phi^H U_A$ . Then the radiant objects are again preserved. To see that the weak equivalences between these radiant objects are preserved, for  $\Phi^H$  we use the fact that it preserves all equivalences of freely *f* -cofibrant spectra (Lemma [7.20\)](#page-34-4), while for ⊙ we use Lemma [5.19](#page-20-2) and the fact that  $\Phi^H U_A \cong U_{A^H}$  is isomorphic to a unit object.

We conclude by Corollary [4.7](#page-14-0) that the derived versions of the diagrams in Figure [7.10](#page-32-1) also commute, defining a strong shadow functor on the homotopy category.  $\Box$ 

*Remark* 7.26*.* As in Remark [5.7,](#page-17-0) these isomorphisms in the homotopy category can be written explicitly by inserting copies of P everywhere. This is even more true in this case since our fibrant objects are defined to be images of spectra of the form *P X*.

<span id="page-36-0"></span>**Proposition 7.27.** *The vertical natural isomorphism from Proposition [7.14](#page-33-2) descends to a vertical natural isomorphism on homotopy categories between the pseudofunctors*

$$
\Phi^H \circ []
$$
 and  $[] \circ (-)^H$ : Ho*G Top*  $\rightarrow$  *WHEx.*

*Proof.* Recall the pseudofunctor [] was defined in Lemma [3.25](#page-10-2) and its coherence isomorphisms were derived in Lemma [5.26.](#page-23-1)

We take the isomorphism  $\eta: \Phi^H([B \stackrel{f}{\leftarrow} A]) \cong [$  $B^H \stackrel{f^H}{\longleftarrow} A^H \Big\vert$  from [\(7.15\)](#page-33-3) and check that the derived versions of the diagrams in Figure [7.16](#page-33-1) commute. Again, it suffices to check that each of the lists of functors that appears is coherently right-deformable. We illustrate the argument with the entry in the top left corner of Figure [7.16b:](#page-33-1)

$$
(*) \times (*) \xrightarrow{[g] \times [f]} G \circlearrowleft^c(C \times B) \times G \circlearrowleft^c(B \times A)
$$
  
\n
$$
\xrightarrow{\Phi^H \times \Phi^H} W H \circlearrowleft^c(C^H \times B^H) \times W H \circlearrowleft^c(B^H \times A^H)
$$
  
\n
$$
\xrightarrow{\circ} W H \circlearrowleft^c(C^H \times A^H).
$$

As in Lemma [5.26,](#page-23-1) we take the radiant objects to be tuples in which

- every spectrum in the tuple is *P*-fibrant or
- every spectrum in the tuple is the appropriate base change 1-cell.

By construction, the radiant objects are preserved, and the weak equivalences between them are preserved by Lemma [7.20](#page-34-4) and Lemma [5.19.](#page-20-2)

We conclude by Corollary [4.7](#page-14-0) that the derived versions of the diagrams in Figure [7.16](#page-33-1) also commute, defining a vertical natural transformation on the homotopy category.  $\Box$ 

<span id="page-36-1"></span>**Proposition 7.28.** *The forgetful functor* (*ιH*) ∗ *from Proposition [7.18](#page-34-5) descends to a strong shadow functor on the homotopy bicategory and an invertible icon on the base-change objects.*

*Proof.* Since  $(\iota_H)^*$  preserves weak equivalences and fibrations [\[Mal,](#page-37-5) Sections 7.2 and 7.4], the proof is the same as Propositions [7.25](#page-35-1) and [7.27,](#page-36-0) but much less complicated. It is not even necessary to use *P*-fibrant spectra – the level *h*-fibrant spectra work perfectly well for this proof.  $\Box$ 

7.6. **The fiberwise case.** Fix a base space *B* with trivial *G*-action. Then the above results apply equally well to the bicategory  $G \& g$  of *G*-spectra over *G*-spaces *A* with equivariant Hurewicz fibrations  $A \rightarrow B$ . Each multi-span over *B* as in [\(6.2\)](#page-25-0) gives an action as in [\(6.3\)](#page-25-1), but with geometric fixed points  $\Phi^H$  thrown in:

(7.29) 
$$
\Pi_i G \mathcal{O} \mathcal{S}^c(A_i) \longrightarrow WH \mathcal{O} \mathcal{S}^c(C)
$$

$$
(X_1, ..., X_n) \longmapsto \Phi^H f_1(g_1, ..., g_n)^* (X_1 \overline{\wedge}_B ... \overline{\wedge}_B X_n)
$$

$$
\cong f_1^H (g_1^H, ..., g_n^H)^* (\Phi^H X_1 \overline{\wedge}_B ... \overline{\wedge}_B \Phi^H X_n).
$$

The rigidity theorem (Theorem [7.8\)](#page-30-2) applies equally well to this new functor since it is also a composition of external smash products, pullbacks, pushforwards, and  $\Phi^H$ . As in Section [6,](#page-24-0) the proofs of Propositions [7.9,](#page-31-1) [7.14](#page-33-2) and [7.18](#page-34-5) are the same, except that the products become fiber products over *B*. We conclude that at the point-set level,  $\Phi^H$ and  $(\iota_H)^*$  are strong shadow functors that commute with base-change objects up to an invertible icon.

To pass to the homotopy category, we observe that the *P*-fibrant spectra of Defini-tion [7.22](#page-34-6) are preserved by  $\overline{\wedge}_B$  because it is a composition of  $\overline{\wedge}$  and a pullback. Therefore the proofs of Propositions [7.25,](#page-35-1) [7.27](#page-36-0) and [7.28](#page-36-1) proceed as before, using Lemma [6.12](#page-27-3) to verify that ⊙*<sup>B</sup>* preserves weak equivalences between our choices of radiant objects in each case. (Note that the extra step in the proof of Proposition [6.19](#page-28-1) does not appear here, because the structures we are interested in do not use the product  $\boxtimes = \overline{\wedge_B}$  directly.) We conclude:

**Theorem 7.30.** *The geometric fixed point functor and the forgetful functor induce strong shadow functors*

> $\Phi^H$ :  $G \&g B \rightarrow WH \&g B$ ,  $\iota_L^*$  $H^*$ :  $G \&g B \rightarrow H \&g B$ ,

*that commute with base change objects up to an invertible icon.*

### <span id="page-37-0"></span>**REFERENCES**

- <span id="page-37-1"></span>[Dol74] A. Dold, *The fixed point index of fibre-preserving maps*, Invent. Math. **25** (1974), 281–297. doi[:10.1007/BF01389731](https://dx.doi.org/10.1007/BF01389731) [1](#page-0-1)
- <span id="page-37-2"></span>[DP80] A. Dold and D. Puppe, *Duality, trace, and transfer*, Proceedings of the International Conference on Geometric Topology (Warsaw, 1978), PWN, Warsaw, 1980, pp. 81–102. [1](#page-0-1)
- <span id="page-37-8"></span>[DHKS04] W. G. Dwyer, P. S. Hirschhorn, D. M. Kan, and J. H. Smith, *Homotopy limit functors on model categories and homotopical categories*, Mathematical Surveys and Monographs, vol. 113, American Mathematical Society, Providence, RI, 2004. [2,](#page-1-1) [3,](#page-2-2) [13,](#page-12-4) [14,](#page-13-2) [15](#page-14-2)
- <span id="page-37-9"></span>[Lac10] S. Lack, *Icons*, Applied Categorical Structures **18** (2010), no. 3, 289–307. arXiv[:0711.4657](https://arxiv.org/abs/0711.4657) [30,](#page-29-2) [33](#page-32-3)
- <span id="page-37-5"></span>[Mal] C. Malkiewich, *A convenient category of parameterized spectra*. arXiv[:2305.15327](https://arxiv.org/abs/2305.15327) [2,](#page-1-1) [3,](#page-2-2) [4,](#page-3-5) [5,](#page-4-4) [9,](#page-8-3) [13,](#page-12-4) [14,](#page-13-2) [17,](#page-16-3) [18,](#page-17-5) [20,](#page-19-4) [25,](#page-24-1) [27,](#page-26-3) [29,](#page-28-2) [30,](#page-29-2) [31,](#page-30-7) [35,](#page-34-7) [37](#page-36-2)
- <span id="page-37-11"></span>[Mal17] C. Malkiewich, *Cyclotomic structure in the topological Hochschild homology of DX*, Algebr. Geom. Topol. **17** (2017), no. 4, 2307–2356. arXiv[:1505.06778](https://arxiv.org/abs/1505.06778) [31](#page-30-7)
- <span id="page-37-4"></span>[MP22] C. Malkiewich and K. Ponto, *Periodic points and topological restriction homology*, Int. Math. Res. Not. IMRN (2022), no. 4, 2401–2459. doi[:10.1093/imrn/rnaa174](https://dx.doi.org/10.1093/imrn/rnaa174) arXiv[:1811.12871](https://arxiv.org/abs/1811.12871) [1,](#page-0-1) [2,](#page-1-1) [5,](#page-4-4) [8,](#page-7-4) [30,](#page-29-2) [31,](#page-30-7) [34,](#page-33-4) [35](#page-34-7)
- <span id="page-37-10"></span>[MM02] M. A. Mandell and J. P. May, *Equivariant orthogonal spectra and S-modules*, Mem. Amer. Math. Soc. **159** (2002), no. 755, x+108. doi[:10.1090/memo/0755](https://dx.doi.org/10.1090/memo/0755) [30](#page-29-2)
- <span id="page-37-6"></span>[MS06] J. P. May and J. Sigurdsson, *Parametrized homotopy theory*, Mathematical Surveys and Monographs, vol. 132, American Mathematical Society, Providence, RI, 2006. doi[:10.1090/surv/132](https://dx.doi.org/10.1090/surv/132) arXiv[:math/0411656](https://arxiv.org/abs/math/0411656) [2,](#page-1-1) [5,](#page-4-4) [17,](#page-16-3) [20](#page-19-4)
- <span id="page-37-3"></span>[Pon10] K. Ponto, *Fixed point theory and trace for bicategories*, Astérisque (2010), no. 333, xii+102. arXiv[:0807.1471](https://arxiv.org/abs/0807.1471) [1,](#page-0-1) [7](#page-6-1)
- <span id="page-37-7"></span>[PS12] K. Ponto and M. Shulman, *Duality and traces for indexed monoidal categories*, Theory Appl. Categ. **26** (2012), no. 23, 582–659. arXiv[:1211.1555](https://arxiv.org/abs/1211.1555) [2,](#page-1-1) [5](#page-4-4)
- <span id="page-38-2"></span>[Rie17] E. Riehl, *Category theory in context*, Courier Dover Publications, 2017. [14,](#page-13-2) [15](#page-14-2)
- <span id="page-38-1"></span>[Shu08] M. Shulman, *Framed bicategories and monoidal fibrations*, Theory Appl. Categ. **20** (2008), No. 18, 650–738. arXiv[:0706.1286](https://arxiv.org/abs/0706.1286) [2,](#page-1-1) [3,](#page-2-2) [5](#page-4-4)
- <span id="page-38-0"></span>[Wec42] F. Wecken, *Fixpunktklassen, I, II, III*, Math. Ann. **117, 118, 118** (1941, 1941, 1942), 659–671, 216–234, 544–477. [1](#page-0-1)

*Email address*: malkiewich@math.binghamton.edu

BINGHAMTON UNIVERSITY, PO BOX 6000, BINGHAMTON, NY 13902

*Email address*: kate.ponto@uky.edu

UNIVERSITY OF KENTUCKY, 719 PATTERSON OFFICE TOWER, LEXINGTON, KY 40506