The Weight Distributions of Two Classes of Linear **Codes From Perfect Nonlinear Functions**

Huawei Wu, Jing Yang and Keqin Feng

Abstract—In this paper, we employ general results on the value distributions of perfect nonlinear functions from \mathbb{F}_{p^m} to \mathbb{F}_p to give a unified approach to determining the weight distributions of two classes of linear codes over \mathbb{F}_p constructed from perfect nonlinear functions, where p is an odd prime and m is an odd number. When m is even, we give some mild additional conditions for similar conclusions to hold.

Index Terms-Bent functions, exponential sums, group actions, linear codes, minimum distance, perfect nonlinear, value distribution, Walsh transform, weakly regular bent, weight distribution.

I. INTRODUCTION

ET A and B be two finite abelian groups and let $f: A \to B$ be a mapping. We say that f is perfect nonlinear if for any $a \in A \setminus \{0\}$, the difference function $D_a f: A \to B$ given by $x \mapsto f(x+a) - f(x)$ is balanced, i.e., $|D_a f^{-1}(b)| = |A|/|B|$ for any $b \in B$. If |A| = |B|, perfect nonlinear mappings from A to B are also called planar functions. Perfect nonlinearity is a robust measure of nonlinearity related to differential cryptanalysis, which was first introduced by Nyberg in [14]. We refer to [2] for an intensive survey on perfect nonlinear functions (and, more generally, highly nonlinear functions).

Throughout this section, let p be an odd prime number and let $q = p^m$ with $m \in \mathbb{N}_+$, where \mathbb{N}_+ is the set of positive integers. We are mainly interested in perfect nonlinear functions from \mathbb{F}_q to itself. By the above definition, a function $f: \mathbb{F}_q \to \mathbb{F}_q$ is perfect nonlinear if for any $a \in \mathbb{F}_q^*$, the difference function $D_a f : \mathbb{F}_q \to \mathbb{F}_q$ is bijective.

It is quite difficult to construct new perfect nonlinear functions. Up to now, all known perfect nonlinear functions over \mathbb{F}_q are affine equivalent to one of the following functions:

1) Dembowski-Ostrom type:

$$\Pi_1(x) = \sum_{0 \le i \le j \le m-1} a_{ij} x^{p^i + p^j}$$

where $a_{ij} \in \mathbb{F}_q$ satisfy specific restrictions (see [1], [4], [6], [7], [17]);

2) Coulter-Matthews type: $\Pi_2(x) = x^{\frac{3^k+1}{2}}$, where p = 3, k is odd and gcd(m, k) = 1 (see [5]).

K. Feng is with the Department of Mathematical Sciences, Tsinghua University, Beijing, 100084, China (e-mail: fengkq@mail.tsinghua.edu.cn)

The work of K. Feng is supported by the National Natural Science Foundation of China under Grant 12031011.

Perfect nonlinear functions can also be used to construct linear codes with good parameters. Let $\Pi : \mathbb{F}_q \to \mathbb{F}_q$ be a perfect nonlinear function. Carlet, Ding and Yuan studied in [3] the following two classes of linear codes over \mathbb{F}_n :

$$C_{\Pi} = \{ c_{a,b} = (f_{a,b}(x))_{x \in \mathbb{F}_{a}^{*}} : a, b \in \mathbb{F}_{q} \}$$

$$f_{a,b}(x) = \operatorname{Tr}_{\mathbb{F}_a/\mathbb{F}_p}(a\Pi(x) + bx)$$

and

$$\overline{C_{\Pi}} = \{c_{a,b,c} = (f_{a,b,c}(x))_{x \in \mathbb{F}_q} : a, b, c \in \mathbb{F}_q\}$$

where

$$f_{a,b,c}(x) = \operatorname{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(a\Pi(x) + bx + c).$$

They showed that C_{Π} is a $[q-1, 2m]_p$ linear code if $\Pi(0) = 0$ and C_{Π} is a $[q, 1+2m]_p$ linear code, and developed bounds on the nonzero Hamming weights of C_{Π} and $\overline{C_{\Pi}}$. They also studied the dual codes of C_{Π} and $\overline{C_{\Pi}}$, proving results on the minimum distances.

In [16], Carlet, Ding and Yuan determined the weight distribution of C_{Π} when Π is a perfect nonlinear function of one of the following forms:

- 1) $\Pi(x) = x^{p^k+1}$, where $k \in \mathbb{N}$ and $m/\gcd(m,k)$ is an
- odd number; 2) $\Pi(x) = x^{\frac{3^k+1}{2}}$, where p = 3, k, m are odd and gcd(m,k) = 1;
- 3) $\Pi(x) = x^{10} ux^6 u^2x^2$, where p = 3, *m* is odd and $u \in \mathbb{F}_{a}^{*}$.

The authors did that on a case-by-case basis and left the Coulter-Matthews type of perfect nonlinear functions, where m is even, as an open problem. They also demonstrated that the linear codes C_{Π} constructed from the above three types of perfect nonlinear functions are either optimal or among the best codes known. Here, an [n, k, d] linear code over \mathbb{F}_p is called optimal if there does not exist an [n, k, d'] linear code over \mathbb{F}_p such that d' > d. The term "best code" can be understood in a similar manner.

In [8], Feng and Luo calculated the value distributions of certain exponential sums derived from perfect nonlinear functions and utilized their results to establish a unified method of determining the weight distribution of C_{Π} when Π is a perfect nonlinear function of Dembowski-Ostrom type or Coulter-Matthews type. Up to then, the weight distributions of the linear codes C_{Π} were completely determined for all known perfect nonlinear functions.

In [12], Li, Ling and Qu employed a unified approach to determining the weight distributions of C_{Π} and $\overline{C_{\Pi}}$ for all the

H. Wu is with the Department of Mathematical Sciences, Tsinghua University, Beijing, 100084, China (e-mail: wu-hw18@mails.tsinghua.edu.cn) J. Yang is with the Department of Mathematical Sciences, Tsinghua University, Beijing, 100084, China (e-mail: y-j@mail.tsinghua.edu.cn)

perfect nonlinear functions in the list treated in [16]. They were the first to determine the weight distributions of the linear codes $\overline{C_{\Pi}}$ and their method of determining the weight distributions of C_{Π} was new.

The main tools of the previous studies mentioned above were exponential sums and quadratic forms over finite fields. The main drawback of these studies is that they depend on the specific form of the perfect nonlinear function Π and thus they were all done on a case-by-case basis; in particular, their conclusions hold only for known perfect nonlinear functions.

However, as shown in [16], [8] and [12], the results on the weight distributions of C_{Π} and $\overline{C_{\Pi}}$ are the same for all known perfect nonlinear functions, respectively. Hence, it is natural to ask whether these results hold for an arbitrary perfect nonlinear function and whether we can find a unified approach to proving them rather than doing that on a case-by-case basis.

In [11], when p = 3, only using the assumption that Π has perfect nonlinearity, the authors adopted a new approach to determining the weight distribution of C_{Π} by determining all the possibilities for the value distribution of a perfect nonlinear function from \mathbb{F}_{3^m} to \mathbb{F}_3 . Their starting point was the following lemma concerning the value distribution of a perfect nonlinear mapping.

Lemma 1 ([2, Theorem 9]). Let (A, +) and (B, +) be abelian groups of order n and m, respectively, where m divides n, and let $f : A \to B$ be a perfect nonlinear mapping. For any $b \in B$, put $k_b = |f^{-1}(b)|$. Then we have

$$\begin{cases} \sum_{z \in B} k_z^2 = \frac{n^2 + (m-1)n}{m}, \\ \sum_{z \in B} k_z k_{z+b} = \frac{n(n-1)}{m}, \ \forall \ b \in B \setminus \{0\}, \\ \sum_{z \in B} k_z = n. \end{cases}$$
(1)

In general, the equations in the second row of (1) are not symmetric, causing difficulty in solving them. However, if $A = \mathbb{F}_{3^m}$ and $B = \mathbb{F}_3$, then the equations in (1) are simple enough (in particular, they are symmetric) for us to determine all the solutions by converting the problem into finding all the integer representations of 3^{m-1} by the binary quadratic form

$$X^2 + XY + Y^2.$$

This strategy cannot be generalized to arbitrary p since the equations in (1) are difficult to solve.

In this paper, we employ general results on the value distributions of perfect nonlinear functions from \mathbb{F}_q to \mathbb{F}_p to determine the weight distributions of C_{Π} and \overline{C}_{Π} only using the assumption that Π has perfect nonlinearity for all odd primes p when m is odd. When m is even, we give some mild additional conditions for similar conclusions to hold. Moreover, for the linear code \overline{C}_{Π} , we can say more about its codewords than just determining the weight distribution from the perspective of a group action. This paper not only provides results that go beyond the literature, but also presents some deep insights on these two classes of linear codes.

II. THE VALUE DISTRIBUTIONS OF PERFECT NONLINEAR FUNCTIONS

There is a characterization of perfect nonlinearity by means of the Fourier transform on finite abelian groups. For this, let us first recall some basic definitions and results.

Let G be a finite abelian group and let $f : G \to \mathbb{C}$ be a complex-valued function. The Fourier transform of f is defined by the complex-valued function

$$\begin{split} \widehat{f} : \widehat{G} \to \mathbb{C}, \\ \chi \mapsto \sum_{g \in G} f(g) \chi(g) \end{split}$$

where \widehat{G} is the group of characters of G, which is (in general, non-canonically) isomorphic to G. If $F: A \to B$ is a mapping between two finite abelian groups, then for any $\chi \in \widehat{B}$, we have the complex-valued function $F_{\chi} := \chi \circ F : A \to \mathbb{C}$. We define the Walsh transform $W_F: \widehat{A} \times \widehat{B} \to \mathbb{C}$ of F by

$$W_F(\phi, \chi) = \widehat{F_{\chi}}(\phi).$$

The set

$$\{W_F(\phi,\chi): \phi \in \widehat{A}, \ \chi \in \widehat{B} \setminus \{1_{\widehat{B}}\}\}$$

is called the Walsh spectrum of F, where $1_{\widehat{B}}$ is the trivial character of B sending all elements of B to $1 \in \mathbb{C}$.

The following theorem gives a characterization of perfect nonlinearity using the Walsh spectrum.

Theorem 1 ([2, Theorem 16]). Let $F : A \to B$ be a mapping between two finite abelian groups. Then F is perfect nonlinear if and only if $|W_F(\phi, \chi)| = \sqrt{|A|}$ for any $\phi \in \hat{A}$ and $\chi \in \hat{B} \setminus \{1_{\hat{B}}\}$.

With the abstract theory out of the way, let us now focus on the finite field case. Let p be a prime number and let $q = p^m$ with $m \in \mathbb{N}_+$.

If we fix a primitive *p*-th root of unity ξ_p in \mathbb{C} , then we have a natural isomorphism $\mathbb{F}_q \to \widehat{\mathbb{F}_q}$, $a \mapsto \psi_a$, where ψ_a is given by

$$\psi_a : \mathbb{F}_q \to \mathbb{C},$$
$$x \mapsto \xi_p^{\operatorname{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(ax)}$$

Therefore, if f is a complex-valued function on \mathbb{F}_q , we usually define its Fourier transform by

$$f: \mathbb{F}_q \to \mathbb{C}$$

 $a \mapsto \sum_{x \in \mathbb{F}_q} f(x) \xi_p^{\operatorname{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(ax)},$

which is a complex-valued function on \mathbb{F}_q rather than one on $\widehat{\mathbb{F}_q}$. Let *h* be a positive divisor of *m*. The Walsh transform of a function $F: \mathbb{F}_q \to \mathbb{F}_{p^h}$ is similarly defined by

$$W_F(a,b) = \sum_{x \in \mathbb{F}_q} \psi_a(x)\psi_b(F(x))$$
$$= \sum_{x \in \mathbb{F}_q} \xi_p^{\operatorname{Tr}_{\mathbb{F}_ph}/\mathbb{F}_p}(bF(x)) + \operatorname{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(ax)},$$

with $a \in \mathbb{F}_q$ and $b \in \mathbb{F}_{p^h}$. Note that, in the literature, $W_F(a, b)$ is usually defined by

$$W_F(a,b) = \sum_{x \in \mathbb{F}_q} \xi_p^{\operatorname{Tr}_{\mathbb{F}_p h}/\mathbb{F}_p(bF(x)) - \operatorname{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(ax)}$$

However, the Walsh spectra corresponding to both definitions are identical. By Theorem 1, F is perfect nonlinear if and only if $|W_F(a,b)| = \sqrt{q}$ for any $a \in \mathbb{F}_q$ and $b \in \mathbb{F}_{p^h}^*$.

Recall that a function $f : \mathbb{F}_q \to \mathbb{F}_p$ is called bent if $W_f(a) := W_f(a, 1)$ has absolute value \sqrt{q} for any $a \in \mathbb{F}_q$. The following lemma shows the equivalence between perfect nonlinearity and bentness.

Lemma 2 ([13, Theorem 2.3]). Let $f : \mathbb{F}_q \to \mathbb{F}_p$ be a function. Then f is perfect nonlinear if and only if it is bent.

This characterization of perfect nonlinear functions allows us to study their value distributions. Let $f : \mathbb{F}_q \to \mathbb{F}_p$ be a function. We say that the value distribution of f is (n_0, \dots, n_{p-1}) if $n_i = |f^{-1}(i)|$ for any $0 \le i \le p-1$. The following two theorems describe completely the value distribution of a perfect nonlinear function from \mathbb{F}_q to \mathbb{F}_p .

Theorem 2 ([13, Theorem 3.2]). Assume that m is even. Then for any bent function $f : \mathbb{F}_q \to \mathbb{F}_p$, there exists $s \in \{0, 1, \dots, p-1\}$ such that the value distribution of f is $(n_0, n_1, \dots, n_{p-1})$, where

$$n_{s} = p^{m-1} + (p-1)p^{\frac{m}{2}-1}, \qquad (2)$$

$$n_{i} = p^{m-1} - p^{\frac{m}{2}-1}, \quad i \neq s,$$

or

$$n_{s} = p^{m-1} - (p-1)p^{\frac{m}{2}-1},$$

$$n_{i} = p^{m-1} + p^{\frac{m}{2}-1}, \quad i \neq s.$$
(3)

Theorem 3 ([13, Theorem 3.4]). Assume that p and m are both odd. Then for any bent function $f : \mathbb{F}_q \to \mathbb{F}_p$, there exists $s \in \{0, 1, \dots, p-1\}$ such that the value distribution of f is $(n_0, n_1, \dots, n_{p-1})$, where

$$n_i = p^{m-1} + (\frac{i+s}{p})p^{\frac{m-1}{2}}, \quad i = 0, \dots, p-1$$

or

$$n_i = p^{m-1} - (\frac{i+s}{p})p^{\frac{m-1}{2}}, \quad i = 0, \ \cdots, \ p-1.$$

Here, $(\frac{\cdot}{p})$ is the Legendre symbol modulo p and we establish the convention that $(\frac{0}{p}) = 0$.

Remark. Note that, in [13], Theorem 3 was stated only for regular bent functions (see Definition 1). However, the proof can be easily generalized to show the validity of Theorem 3 for arbitrary bent functions.

The key points for proving Theorem 2 and Theorem 3 are the following two lemmas.

Lemma 3 ([13, Lemma before Theorem 3.4]). Assume that p is odd. If there exist $a_1, \dots, a_{p-1} \in \mathbb{Q}$ such that

$$a_{1}\xi_{p} + a_{2}\xi_{p}^{2} + \dots + a_{p-1}\xi_{p}^{p-1}$$

=
$$\begin{cases} \sqrt{p}, & \text{if } p \equiv 1 \pmod{4}, \\ i\sqrt{p}, & \text{if } p \equiv 3 \pmod{4}, \end{cases}$$

then $a_i = \left(\frac{i}{p}\right)$ for $i = 1, \dots, p-1$.

Lemma 4 ([10, Property 7, 8]). Assume that p is odd. For any bent function $f : \mathbb{F}_q \to \mathbb{F}_p$, there exists a function $f^* : \mathbb{F}_q \to \mathbb{F}_p$, which is called the dual of f, such that for any $a \in \mathbb{F}_q$,

$$W_f(a) = \begin{cases} \pm \xi_p^{f^*(a)} \sqrt{q}, & \text{if } p \equiv 1 \pmod{4}, \\ \pm i \xi_p^{f^*(a)} \sqrt{q}, & \text{if } p \equiv 3 \pmod{4} \end{cases}$$
(4)

if m is odd and

$$W_f(a) = \pm \xi_p^{f^*(a)} \sqrt{q} \tag{5}$$

if m is even. The sign ± 1 in (4) or (5) will be called the sign of f at a.

We provide a sketch of the proofs for Theorem 2 and Theorem 3 here, which will be used in the proofs of Corollary 1 and Theorem 4. Indeed, the value distribution of a bent function $f : \mathbb{F}_q \to \mathbb{F}_p$ is related to its Walsh spectrum via the following equation:

$$W_f(0) = \sum_{x \in \mathbb{F}_q} \xi_p^{f(x)} = \sum_{i=0}^{p-1} n_i \xi_p^i,$$

where $n_i = |f^{-1}(i)|$ for any $0 \le i \le p-1$. If m is even, then by Lemma 4, we have

$$W_f(0) = \epsilon \xi_p^s \sqrt{q}$$

where $\epsilon \in \{\pm 1\}$ is the sign of f at 0 and $0 \le s \le p - 1$. By employing some basic properties of cyclotomic fields, we can show that

$$n_s = p^{m-1} + \epsilon(p-1)p^{\frac{m}{2}-1}$$

and

$$n_i = n_s - \epsilon p^{\frac{m}{2}} = p^{m-1} - \epsilon p^{\frac{m}{2}-1}$$

for any $i \neq s$. In particular, the value distribution of f is of the form (2) (resp., (3)) if and only if the sign of f at 0 is +1 (resp., -1). Theorem 3 can be proved similarly, with the additional assistance of Lemma 3.

For the purpose of later exposition of the results, we introduce the definition of weakly regular bentness.

Definition 1. Let p be an odd prime number and let $f : \mathbb{F}_q \to \mathbb{F}_p$ be a bent function. If the signs ± 1 in (4) or (5) are the same for all $a \in \mathbb{F}_q$, then f is called weakly regular bent. In this case, the common sign will be called the sign of f. If $W_f(a) = \xi_p^{f^*(a)} \sqrt{q}$ for any $a \in \mathbb{F}_q$, where $f^* : \mathbb{F}_q \to \mathbb{F}_p$ is the dual of f, then f is called regular bent.

Let $\Pi : \mathbb{F}_q \to \mathbb{F}_q$ be a perfect nonlinear function. We say that Π is weakly regular perfect nonlinear if the functions $\Pi_a : \mathbb{F}_q \to \mathbb{F}_p$ given by $x \mapsto \operatorname{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(a\Pi(x))$ are weakly regular bent for all $a \in \mathbb{F}_q^*$.

Corollary 1. Assume that m is even and let $f : \mathbb{F}_q \to \mathbb{F}_p$ be a weakly regular bent function. If the value distribution of f is of the form (2) (resp., (3)), then so is that of f_b for any $b \in \mathbb{F}_q$, where

$$f_b(x) = f(x) + \operatorname{Tr}_{\mathbb{F}_a/\mathbb{F}_n}(bx)$$

Proof. As mentioned above, the form of the value distribution of a bent function from \mathbb{F}_q to \mathbb{F}_p depends only on the

sign of the function at 0. Since f is weakly regular bent, the signs of f at 0 and b are the same. Moreover, we have

$$W_f(b) = \sum_{x \in \mathbb{F}_q} \xi_p^{f(x) + \operatorname{Tr}_{\mathbb{F}_q}/\mathbb{F}_p}(bx)} = W_{f_b}(0),$$

which implies that the sign of f_b at 0 is equal to the sign of f at b. Therefore, f_b and f have the same sign at 0.

III. The Weight Distributions of C_{Π} and $\overline{C_{\Pi}}$

Throughout this section, let p be an odd prime number, let $q = p^m$ with $m \in \mathbb{N}_+$ and let $\Pi : \mathbb{F}_q \to \mathbb{F}_q$ be a perfect nonlinear function. For any $0 \le i \le q - 1$, let A_i denote the number of codewords of Hamming weight i in C_{Π} and for any $0 \le i \le q$, let $\overline{A_i}$ denote the number of codewords of Hamming weight i in $\overline{C_{\Pi}}$.

We first consider the linear code $\overline{C_{\Pi}}$, since we can say more about its codewords than just determining the weight distribution. For any $c_{a,b,c} \in \overline{C_{\Pi}}$, we say that it is a (n_0, \dots, n_{p-1}) codeword if the value distribution of $f_{a,b,c}$ is (n_0, \dots, n_{p-1}) .

If a = 0 and b = 0, then $f_{0,0,c}(x) = \operatorname{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(c)$ for any $x \in \mathbb{F}_q$. For any $0 \leq i \leq p-1$, if $\operatorname{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(c) = i$, then $c_{0,0,c} = (i, \cdots, i)$, i.e., $c_{0,0,c}$ is a $(0, \cdots, 0, p^m, 0, \cdots, 0)$ -codeword, where p^m occurs in the *i*-th entry. There is only one codeword of each of these p types in $\overline{C_{\Pi}}$.

If a = 0 but $b \neq 0$, then for any $c \in \mathbb{F}_q$, since $x \mapsto bx + c$ is a permutation of \mathbb{F}_q , $c_{0,b,c}$ is a $(p^{m-1}, \cdots, p^{m-1})$ -codeword. There are $p(q-1) = p^{m+1} - p$ such codewords.

If $a \neq 0$, then for any $b, c \in \mathbb{F}_q$, $f_{a,b,c}$ is a perfect nonlinear function from \mathbb{F}_q to \mathbb{F}_p . Put

$$\overline{\Omega} = \{c_{a,b,c} \in \overline{C_{\Pi}}: \ a \neq 0\}.$$

Since for any $a, a', b, b', c, c' \in \mathbb{F}_q$, $c_{a,b,c} = c_{a',b',c'}$ if and only if a = a', b = b' and $\operatorname{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(c - c') = 0$, the subset $\overline{\Omega}$ is well-defined, i.e., the condition that $a \neq 0$ does not depend on the way in which the codewords in \overline{C}_{Π} are represented as $c_{a,b,c}$. It is easy to see that there are $pq(q-1) = p^{m+1}(p^m-1)$ codewords in $\overline{\Omega}$.

Consider the group $G := \mathbb{F}_p^* \times \mathbb{F}_p$, whose multiplication is given by

$$(\alpha_1,\beta_1)\cdot(\alpha_2,\beta_2)=(\alpha_1\alpha_2,\alpha_1\beta_2+\beta_1).$$

The identity element of G is (1,0). For any $\alpha \in \mathbb{F}_p^*$, $\beta \in \mathbb{F}_p$ and a, b, $c \in \mathbb{F}_q$, we have

$$\alpha f_{a,b,c}(x) + \beta = \alpha \operatorname{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(a\Pi(x) + bx + c) + \beta$$
$$= \operatorname{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(\alpha a\Pi(x) + \alpha bx + \alpha c + \beta')$$

for any $x \in \mathbb{F}_q$, i.e., $\alpha f_{a,b,c} + \beta = f_{\alpha a,\alpha b,\alpha c+\beta'}$, where $\beta' \in \mathbb{F}_q$ is such that $\operatorname{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(\beta') = \beta$. Therefore, we can define a *G*-action on $\overline{\Omega}$ by

$$(\alpha,\beta) \cdot c_{a,b,c} = c_{\alpha a,\alpha b,\alpha c+\beta'}.$$

(Note that this definition does not depend on the choice of β' .)

Lemma 5. The *G*-action on $\overline{\Omega}$ is free; that is, if $(\alpha, \beta) \cdot c_{a,b,c} = c_{a,b,c}$ for some $(\alpha, \beta) \in G$ and $c_{a,b,c} \in \overline{\Omega}$, then (α, β) must be the identity element of *G*, i.e., (1, 0).

Proof. Assume that
$$(\alpha, \beta) \in G$$
 fixes $c_{a,b,c} \in \Omega$, i.e.,

$$(\alpha,\beta) \cdot c_{a,b,c} = c_{\alpha a,\alpha b,\alpha c+\beta'} = c_{a,b,c}.$$

Then

$$\alpha a = a, \ \alpha b = b, \ \operatorname{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(\alpha c + \beta' - c) = 0.$$

Since $a \neq 0$, we must have $\alpha = 1$ and thus

$$\beta = \operatorname{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(\beta') = 0.$$

Therefore, G acts freely on $\overline{\Omega}$.

In particular, there are $p^m(p^m-1)/(p-1)$ orbits of $\overline{\Omega}$ under the *G*-action and each orbit contains (p-1)p codewords. The essence of proving Theorem 4 lies in a careful examination of how the types of codewords in $\overline{\Omega}$ evolve under the *G*-action. It can be easily seen that if $c_{a,b,c}$ is a (n_0, \cdots, n_{p-1}) -codeword, then for any $(\alpha, \beta) \in G$, $(\alpha, \beta) \cdot c_{a,b,c}$ is a (n'_0, \cdots, n'_{p-1}) codeword, where $n'_i = n_{\alpha^{-1}(i-\beta)}$ for any $0 \le i \le p-1$.

For the sake of convenient exposition, we introduce the following definition.

Definition 2. Assume that m is odd. A codeword $c_{a,b,c} \in \overline{\Omega}$ is called an s^+ -codeword (resp., s^- -codeword) with $0 \le s \le p-1$ if it is a (n_0, \dots, n_{p-1}) -codeword, where

$$\begin{split} n_i &= p^{m-1} + (\frac{i+s}{p}) p^{\frac{m-1}{2}} \text{ for any } 0 \leq i \leq p-1 \\ (\text{resp., } n_i &= p^{m-1} - (\frac{i+s}{p}) p^{\frac{m-1}{2}} \text{ for any } 0 \leq i \leq p-1). \end{split}$$

Assume that m is even. A codeword $c_{a,b,c} \in \overline{\Omega}$ is called an s^+ -codeword (resp., s^- -codeword) with $0 \le s \le p-1$ if it is a (n_0, \dots, n_{p-1}) -codeword, where

$$n_{i} = \begin{cases} p^{m-1} + (p-1)p^{\frac{m}{2}-1}, & \text{if } i = s, \\ p^{m-1} - p^{\frac{m}{2}-1}, & \text{if } i \neq s, \end{cases}$$

$$\left(\text{resp., } n_{i} = \begin{cases} p^{m-1} - (p-1)p^{\frac{m}{2}-1}, & \text{if } i = s, \\ p^{m-1} + p^{\frac{m}{2}-1}, & \text{if } i \neq s, \end{cases}\right).$$

If $c_{a,b,c} \in \overline{\Omega}$ is an s^+ -codeword (resp., s^- -codeword) for some $0 \leq s \leq p-1$, then $c_{a,b,c}$ is called a positive (resp., negative) codeword and is called a strictly positive (resp., negative) codeword if furthermore $s \neq 0$.

By Theorem 2 and Theorem 3, any codeword $c_{a,b,c} \in \overline{\Omega}$ is an s^+ -codeword or an s^- -codeword for some $0 \le s \le p-1$. We can now determine the number of codewords in $\overline{C_{\Pi}}$ of each type, as follows.

Theorem 4. Let $\Pi : \mathbb{F}_q \to \mathbb{F}_q$ be a perfect nonlinear function.

1) If m is odd, then the possible types of codewords in $\overline{C_{\Pi}}$ and the corresponding numbers of codewords are listed in Table I. As a consequence, $\overline{A_i} = 0$ except for the values

$$\begin{aligned} \overline{A}_0 &= 1, \\ \overline{A}_{(p-1)p^{m-1}-p^{\frac{m-1}{2}}} &= \frac{1}{2}(p-1)p^m(p^m-1), \\ \overline{A}_{(p-1)p^{m-1}} &= (p^m-1)(p^m+p), \end{aligned}$$

$$\overline{A}_{(p-1)p^{m-1}+p^{\frac{m-1}{2}}} = \frac{1}{2}(p-1)p^m(p^m-1),$$

$$\overline{A}_{p^m} = p - 1.$$

In particular, the minimum distance of $\overline{C_{\Pi}}$ is $(p-1)p^{m-1}-p^{\frac{m-1}{2}}$ and $\overline{C_{\Pi}}$ is a 4-weight code.

- 2) If *m* is even, then the possible types of codewords in $\overline{C_{\Pi}}$ and the corresponding numbers of codewords are listed in Table I, assuming that one of the following conditions holds:
 - a) p = 3,
 - b) $|\Pi^{-1}(0)| = 1$, Π is weakly regular perfect nonlinear (see Definition 1) and the map $\mathbb{F}_q^* \to \mathbb{F}_p$ given by $a \mapsto \Pi_a^*(0)$ is not surjective, where $\Pi_a = f_{a,0,0}$ and Π_a^* is the dual of Π_a (see Lemma 4).

As a consequence, $\overline{A_i} = 0$ except for the values

$$\begin{split} A_{0} &= 1, \\ \overline{A}_{(p-1)(p^{m-1}-p^{\frac{m}{2}-1})} &= \frac{p^{m}(p^{m}-1)}{2}, \\ \overline{A}_{(p-1)p^{m-1}-p^{\frac{m}{2}-1}} &= \frac{p^{m}(p^{m}-1)(p-1)}{2}, \\ A_{(p-1)p^{m-1}} &= p^{m+1}-p, \\ \overline{A}_{(p-1)p^{m-1}+p^{\frac{m}{2}-1}} &= \frac{p^{m}(p^{m}-1)(p-1)}{2}, \\ \overline{A}_{(p-1)(p^{m-1}+p^{\frac{m}{2}-1})} &= \frac{p^{m}(p^{m}-1)}{2}, \\ \overline{A}_{p^{m}} &= p-1. \end{split}$$

In particular, the minimum distance of $\overline{C_{\Pi}}$ is $(p-1)(p^{m-1}-p^{\frac{m}{2}-1})$ and $\overline{C_{\Pi}}$ is a 6-weight code.

ſ	TABLE	ΕI

Туре	Number	
$\begin{array}{c} (0,\cdots,0,p^m,0,\cdots,0) \\ (\forall \ 0 \le i \le p-1) \end{array}$	1	
(p^{m-1},\cdots,p^{m-1})	$p^{m+1}-p$	
$s^+ \ (\forall \ 0 \le s \le p-1)$	$\frac{p^m}{2}(p^m-1)$	
$s^- (\forall \ 0 \le s \le p-1)$	$\frac{p^m}{2}(p^m-1)$	

Proof. Assume first that m is odd. Let $(\alpha, \beta) \in G$ and let $c_{a,b,c} \in \overline{\Omega}$. If $c_{a,b,c}$ is an s^+ -codeword, then $(\alpha, \beta) \cdot c_{a,b,c}$ is a $(\alpha s - \beta)^+$ -codeword if $(\frac{\alpha^{-1}}{p}) = 1$ and is a $(\alpha s - \beta)^-$ -codeword if $(\frac{\alpha^{-1}}{p}) = -1$. Similarly, if $c_{a,b,c}$ is an s^- -codeword, then $(\alpha, \beta) \cdot c_{a,b,c}$ is a $(\alpha s - \beta)^-$ -codeword if $(\frac{\alpha^{-1}}{p}) = 1$ and is a $(\alpha s - \beta)^+$ -codeword if $(\frac{\alpha^{-1}}{p}) = -1$. In particular, every orbit of $\overline{\Omega}$ contains a 0^+ -codeword.

Let *O* be an orbit of $\overline{\Omega}$ and let $c_{a,b,c}$ be a 0⁺-codeword in *O*. Then for any $(\alpha, \beta) \in G$, $(\alpha, \beta) \cdot c_{a,b,c}$ is a $(-\beta)^+$ -codeword if $(\frac{\alpha^{-1}}{p}) = 1$ and is a $(-\beta)^-$ -codeword if $(\frac{\alpha^{-1}}{p}) = -1$. In \mathbb{F}_p^* , there are (p-1)/2 elements α such that $(\frac{\alpha^{-1}}{p}) = 1$ and (p-1)/2 elements α such that $(\frac{\alpha^{-1}}{p}) = -1$. Hence, in *O*, there are $(p-1)/2 s^+$ -codewords and $(p-1)/2 s^-$ -codewords for any $0 \le s \le p-1$. This proves the assertion on the possible types of codewords in $\overline{C_{\Pi}}$.

To determine the weight distribution of $\overline{C_{\Pi}}$, we only consider the codewords in $\overline{\Omega}$, since the Hamming weights of the other codewords are easy to determine. It is clear that a codeword in $\overline{\Omega}$ has Hamming weight $(p-1)p^{m-1}$ if and only if it is a 0⁺-codeword or a 0⁻-codeword. Let $1 \le s \le p-1$. If $(\frac{s}{p}) = 1$, then any s^+ -codeword has Hamming weight $(p-1)p^{m-1} - p^{\frac{m-1}{2}}$ and any s^- -codeword has Hamming weight $(p-1)p^{m-1} + p^{\frac{m-1}{2}}$. If $(\frac{s}{p}) = -1$, the situation is reversed. Hence, in $\overline{\Omega}$ (and thus in $\overline{C_{\Pi}}$), the codewords of Hamming weight $(p-1)p^{m-1} - p^{\frac{m-1}{2}}$ are equal in number to the codewords of Hamming weight $(p-1)p^{m-1} - p^{\frac{m-1}{2}}$. This proves the assertion on the weight distribution of $\overline{C_{\Pi}}$.

Now assume that m is even. Let $(\alpha, \beta) \in G$ and let $c_{a,b,c} \in \overline{\Omega}$. If $c_{a,b,c}$ is an s^+ -codeword, then $(\alpha, \beta) \cdot c_{a,b,c}$ is a $(\alpha s + \beta)^+$ -codeword; in particular, the orbit containing $c_{a,b,c}$ contains a 0^+ -codeword. If $c_{a,b,c}$ is an s^- -codeword, then $(\alpha, \beta) \cdot c_{a,b,c}$ is a $(\alpha s + \beta)^-$ -codeword; in particular, the orbit containing $c_{a,b,c}$ contains a 0^- -codeword. Let O be an orbit of $\overline{\Omega}$. The above discussions imply that either all the codewords in O are positive or all the codewords in O are negative. An orbit of the former type will be called a positive orbit, while an orbit of the latter type will be called a negative orbit.

If O is a positive orbit of $\overline{\Omega}$ and $c_{a,b,c} \in O$ is a 0⁺codeword, then for any $(\alpha, \beta) \in G$, $(\alpha, \beta) \cdot c_{a,b,c}$ is a β^+ codeword. In particular, for any $0 \le s \le p-1$, there are p-1 s^+ -codewords in O. If $c_{a,b,c}$ is an s^+ -codeword, then

$$w_H(c_{a,b,c}) = \begin{cases} (p-1)(p^{m-1} - p^{\frac{m}{2}-1}), & \text{if } s = 0, \\ (p-1)p^{m-1} + p^{\frac{m}{2}-1}, & \text{if } s \neq 0. \end{cases}$$

Hence, there are p-1 codewords of Hamming weight $(p-1)(p^{m-1}-p^{\frac{m}{2}-1})$ and $(p-1)^2$ codewords of Hamming weight $(p-1)p^{m-1}+p^{\frac{m}{2}-1}$ in O. Similarly, if O is a negative orbit of $\overline{\Omega}$ and $c_{a,b,c} \in O$ is a 0⁻-codeword, then for any $0 \le s \le p-1$, there are p-1 s⁻-codewords in O. Moreover, there are p-1 codewords of Hamming weight $(p-1)(p^{m-1}+p^{\frac{m}{2}-1})$ and $(p-1)^2$ codewords of Hamming weight $(p-1)p^{m-1}-p^{\frac{m}{2}-1}$ in O.

Assume that there are k_+ positive orbits and k_- negative orbits in $\overline{\Omega}$. Then

$$k_{+} + k_{-} = \frac{p^{m}(p^{m} - 1)}{p - 1}.$$
(6)

To determine k_{\pm} and k_{-} , we need to consider the weight distribution of $\overline{C_{\Pi}}$ first. By the above discussions, we know that $\overline{A}_i = 0$ for $i \neq 0$, $(p-1)p^{m-1}$, p^m , $(p-1)p^{m-1} \pm p^{\frac{m}{2}-1}$ and $(p-1)(p^{m-1} \pm p^{\frac{m}{2}-1})$. Moreover, we have

$$\begin{aligned} \overline{A}_0 &= 1, \\ \overline{A}_{(p-1)(p^{m-1}-p^{\frac{m}{2}-1})} &= (p-1)k_+, \\ \overline{A}_{(p-1)p^{m-1}-p^{\frac{m}{2}-1}} &= (p-1)^2k_-, \\ \overline{A}_{(p-1)p^{m-1}} &= p^{m+1}-p, \\ \overline{A}_{(p-1)p^{m-1}+p^{\frac{m}{2}-1}} &= (p-1)^2k_+, \end{aligned}$$

$$\overline{A}_{(p-1)(p^{m-1}+p^{\frac{m}{2}-1})} = (p-1)k_{-},$$

$$\overline{A}_{p^{m}} = p-1.$$

If p = 3, then by [3, Theorem 7], the minimum distance of the dual code $\overline{C_{\Pi}}^{\perp}$ of $\overline{C_{\Pi}}$ is 5. By calculating the first four Pless power moments (see [15, p.90]), we have

$$\sum_{j=0}^{3^{m}} \overline{A}_{j} = 3^{2m+1},$$

$$\sum_{j=0}^{3^{m}} j\overline{A}_{j} = 2 \cdot 3^{3m},$$

$$\sum_{j=0}^{3^{m}} j^{2}\overline{A}_{j} = 2 \cdot 3^{3m-1}(2 \cdot 3^{m} + 1),$$

$$\sum_{j=0}^{3^{m}} j^{3}\overline{A}_{j} = 2 \cdot 3^{3m-2}(4 \cdot 3^{2m} + 2 \cdot 3^{m+1} - 1).$$

This is a system of linear equations with variables

$$\overline{A}_{2\cdot(3^{m-1}\pm 3^{\frac{m}{2}-1})}, \quad \overline{A}_{2\cdot 3^{m-1}\pm 3^{\frac{m}{2}-1}}.$$

Since the coefficient matrix of this system is a Vandermonde matrix, it has a unique solution, which is

$$\begin{split} \overline{A}_{2\cdot(3^{m-1}\pm 3^{\frac{m}{2}-1})} &= \frac{3^m(3^m-1)}{2},\\ \overline{A}_{2\cdot 3^{m-1}\pm 3^{\frac{m}{2}-1}} &= 3^m(3^m-1). \end{split}$$

It follows that

$$k_{+} = k_{-} = \frac{3^{m}(3^{m} - 1)}{4}$$

Now assume that the condition b) holds. Then the functions $\Pi_a : \mathbb{F}_q \to \mathbb{F}_p$ given by $x \mapsto \operatorname{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(a\Pi(x))$ are weakly regular bent for all $a \in \mathbb{F}_q^*$. By Corollary 1, we know that if $c_{a,0,0} \in \overline{\Omega}$ is a positive (resp., negative) codeword, then so is $c_{a,b,c}$ for any $b, c \in \mathbb{F}_q$. Hence, to prove that $k_+ = k_-$, it suffices to show that half of the codewords $c_{a,0,0}$ ($a \in \mathbb{F}_q^*$) are positive while the other half are negative. By the proof of Theorem 2, it suffices to show that half of the weakly regular bent functions Π_a ($a \in \mathbb{F}_q^*$) have sign +1 while the other half have sign -1.

By Lemma 4, for any $a \in \mathbb{F}_q^*$, we have

$$W_{\Pi_a}(0) = \epsilon_a \xi_p^{\Pi_a^*(0)} \sqrt{q},$$

where $\epsilon_a \in \{\pm 1\}$ is the sign of Π_a . Note that

$$\begin{split} \sum_{a \in \mathbb{F}_q^*} W_{\Pi_a}(0) &= \sum_{a \in \mathbb{F}_q^*} \sum_{x \in \mathbb{F}_q} \xi_p^{\mathrm{IT}_{\mathbb{F}_q/\mathbb{F}_p}(a\Pi(x))} \\ &= \sum_{x \in \mathbb{F}_q} \sum_{a \in \mathbb{F}_q^*} \xi_p^{\mathrm{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(a\Pi(x))} \\ &= \sum_{\substack{x \in \mathbb{F}_q \\ \Pi(x) = 0}} (q-1) + \sum_{\substack{x \in \mathbb{F}_q \\ \Pi(x) \neq 0}} (-1) \\ &= |\Pi^{-1}(0)|(q-1) - (q - |\Pi^{-1}(0)|) \\ &= q |\Pi^{-1}(0)| - q = 0, \end{split}$$

which implies that

$$\sum_{a \in \mathbb{F}_q^*} \epsilon_a \xi_p^{\Pi_a^*(0)} = 0.$$
⁽⁷⁾

The above equality can be reformulated as

$$\sum_{r=0}^{p-1} \Big(\sum_{\substack{a \in \mathbb{F}_q^* \\ \Pi_a^*(0)=r}} \epsilon_a\Big)\xi_p^r = 0,$$

which implies that ξ_p is a root of the polynomial

$$\sum_{r=0}^{p-1} \Big(\sum_{\substack{a \in \mathbb{F}_q^* \\ \Pi_a^*(0) = r}} \epsilon_a \Big) X^r \in \mathbb{Z}[X].$$

Since the minimal polynomial of ξ_p over \mathbb{Q} is $X^{p-1} + X^{p-2} + \cdots + 1$, the sums

$$S_r = \sum_{\substack{a \in \mathbb{F}_q^* \\ \Pi_a^*(0) = r}} \epsilon_a, \quad 0 \le r \le p - 1$$

must be all the same. Since the map $\mathbb{F}_q^* \to \mathbb{F}_p$ given by $a \mapsto \Pi_a^*(0)$ is not surjective, we have $S_i = 0$ for some $0 \leq i \leq p-1$, which implies that $S_r = 0$ for any $0 \leq r \leq p-1$. However,

$$\sum_{r=0}^{p-1} S_r = \sum_{a \in \mathbb{F}_q^*} \epsilon_a$$

= $\#\{a \in \mathbb{F}_q^* : \epsilon_a = 1\} - \#\{a \in \mathbb{F}_q^* : \epsilon_a = -1\}.$

Therefore, half of the weakly regular bent functions Π_a ($a \in \mathbb{F}_q^*$) have sign +1 while the other half have sign -1. This completes the proof.

Remark. From the proof of Theorem 4, we can observe that when m is odd, positive codewords and negative codewords in $\overline{\Omega}$ can be mutually transformed through the *G*-action, whereas when m is even, the positive and negative codeword sets are both *G*-stable. This leads to the need for additional conditions to ensure an equal number of positive and negative codewords in $\overline{\Omega}$ when m is even.

Remark. Assume that Π is of Dembowski-Ostrom type or Coulter-Matthews type. By [8, Lemma 2], $\Pi^{-1}(0) = \{0\}$. By [8, Lemma 3 ii)], Π is weakly regular perfect nonlinear and the map $\mathbb{F}_q^* \to \mathbb{F}_p$ given by $a \mapsto \Pi_a^*(0)$ is the zero constant map. Therefore, all known perfect nonlinear functions $\Pi : \mathbb{F}_q \to \mathbb{F}_q$ satisfy the condition b).

Example 1. By Theorem 4 and [9], the linear codes $\overline{C_{\Pi}}$ contain the following optimal codes:

$$[9, 5, 4; 3], [27, 7, 15; 3], [125, 7, 95; 5],$$

and the following best known codes:

[81, 9, 48; 3], [243, 11, 153; 3].

Finally, we determine the weight distribution of the linear code C_{Π} , which can also be done by considering the *G*-action and calculating the first few Pless power moments.

Theorem 5. Let $\Pi : \mathbb{F}_q \to \mathbb{F}_q$ be a perfect nonlinear function with $\Pi(0) = 0$. Assume that $\Pi(cx) \neq c\Pi(x)$ for any $x \in \mathbb{F}_q^*$ and $c \in \mathbb{F}_p \setminus \{0, 1\}$.

1) If m is odd, then $A_i = 0$ except for the values

$$A_{0} = 1,$$

$$A_{(p-1)p^{m-1}-p^{\frac{m-1}{2}}} = (p-1)(p^{m}-1)\frac{p^{m-1}+p^{\frac{m-1}{2}}}{2},$$

$$A_{(p-1)p^{m-1}} = (p^{m-1}+1)(p^{m}-1),$$

$$A_{(p-1)p^{m-1}+p^{\frac{m-1}{2}}} = (p-1)(p^{m}-1)\frac{p^{m-1}-p^{\frac{m-1}{2}}}{2}.$$

In particular, the minimum distance of the code C_{Π} is $(p-1)p^{m-1} - p^{\frac{m-1}{2}}$ and C_{Π} is a 3-weight code. If m is even then $A_{\Gamma} = 0$ except for the values

2) If m is even, then $A_i = 0$ except for the values

$$\begin{split} A_{0} &= 1, \\ A_{(p-1)(p^{m-1}-p^{\frac{m}{2}-1})} &= \frac{p^{m}-1}{2}(p^{m-1}+p^{\frac{m}{2}}-p^{\frac{m}{2}-1}), \\ A_{(p-1)p^{m-1}-p^{\frac{m}{2}-1}} &= \frac{p^{m}-1}{2}(p-1)(p^{m-1}+p^{\frac{m}{2}-1}), \\ A_{(p-1)p^{m-1}} &= p^{m}-1, \\ A_{(p-1)p^{m-1}+p^{\frac{m}{2}-1}} &= \frac{p^{m}-1}{2}(p-1)(p^{m-1}-p^{\frac{m}{2}-1}), \\ A_{(p-1)(p^{m-1}+p^{\frac{m}{2}-1})} &= \frac{p^{m}-1}{2}(p^{m-1}-p^{\frac{m}{2}}+p^{\frac{m}{2}-1}), \end{split}$$

assuming furthermore that one of the following conditions holds:

- a) p = 3,
- b) Π⁻¹(0) = {0}, Π is weakly regular perfect nonlinear and the map 𝔽^{*}_q → 𝔽_p given by a → Π^{*}_a(0) is not surjective, where Π_a = f_{a,0,0} and Π^{*}_a is the dual of Π_a.

In particular, the minimum distance of C_{Π} is $(p-1)(p^{m-1}-p^{\frac{m}{2}-1})$ and C_{Π} is a 5-weight code.

Proof. Note that $w_H(c_{a,b}) = w_H(c_{a,b,0})$ for any $a, b \in \mathbb{F}_q$. By [3, Theorem 6], the minimum distance of the dual code C_{Π}^{\perp} of C_{Π} is at least 3.

We first treat the case where m is odd. By Theorem 4, all the possible nonzero Hamming weights in C_{Π} are $(p-1)p^{m-1}$ and $(p-1)p^{m-1}\pm p^{\frac{m-1}{2}}$. Then the assertion follows by calculating the first three Pless power moments, which has been done in [16, Theorem 2].

Next we treat the case where m is even. If a = 0 and $b \neq 0$, then $w_H(c_{a,b,0}) = (p-1)p^{m-1}$. There are $p^m - 1$ such codewords. Put

$$\Omega = \{ c_{a,b,0} \in \Omega \}.$$

By Theorem 4, any codeword $c_{a,b,0} \in \Omega$ is an s^+ -codeword or an s^- -codeword, where $0 \le s \le p-1$. It is clear that there are $A_{(p-1)(p^{m-1}-p^{\frac{m}{2}-1})} 0^+$ -codewords in Ω , $A_{(p-1)(p^{m-1}+p^{\frac{m}{2}-1})} 0^-$ -codewords in Ω , $A_{(p-1)p^{m-1}+p^{\frac{m}{2}-1}}$ strictly positive codewords in Ω and $A_{(p-1)p^{m-1}-p^{\frac{m}{2}-1}}$ strictly negative codewords in Ω . If $c_{a,b,0} \in \Omega$ is a positive codeword, then in $(\{1\} \times \mathbb{F}_p) \cdot c_{a,b,0}$, there is exactly one s^+ -codeword for any $0 \leq s \leq p-1$. Similar conclusion holds for negative codewords. Since $\overline{\Omega} = (\{1\} \times \mathbb{F}_p) \cdot \Omega$, by Table I we have

$$A_{(p-1)(p^{m-1}-p^{\frac{m}{2}-1})} + A_{(p-1)p^{m-1}+p^{\frac{m}{2}-1}} = \frac{p^m(p^m-1)}{2},$$
(8)
$$A_{(p-1)(p^{m-1}+p^{\frac{m}{2}-1})} + A_{(p-1)p^{m-1}-p^{\frac{m}{2}-1}} = \frac{p^m(p^m-1)}{2}.$$

By calculating the second and the third Pless power moments, we have

$$\sum_{j=0}^{p^m-1} jA_j = p^{2m-1}(p-1)(p^m-1),$$

$$\sum_{j=0}^{p^m-1} j^2A_j = p^{2m-2}(p-1)(p^m-1)(p+(p-1)(p^m-2)).$$
(9)

Then (8) and (9) form a system of linear equations with variables

$$A_{(p-1)p^{m-1}\pm p^{\frac{m}{2}-1}}$$
 and $A_{(p-1)(p^{m-1}\pm p^{\frac{m}{2}-1})}$.

It turns out that this system has a unique solution, i.e., the one given in the theorem. This completes the proof. \Box

Remark. If Π is of Coulter-Matthews type, then Π is even, i.e., $\Pi(x) = \Pi(-x)$ for any $x \in \mathbb{F}_{3^m}$. Assume that there exists $x \in \mathbb{F}_q^*$ such that $\Pi(2x) = 2\Pi(x)$, then

$$\Pi(x) = \Pi(-x) = \Pi(2x) = 2\Pi(x),$$

which implies that $\Pi(x) = 0$. However, by [8, Lemma 2], we have $\Pi^{-1}(0) = \{0\}$, which is a contradiction. Therefore, Π satisfies the conditions in Theorem 5.

Remark. If Π is of Dembowski-Ostrom type, then

$$\Pi(x) = \sum_{0 \le i \le j \le m-1} a_{ij} x^{p^i + p^j}$$

for some $a_{ij} \in \mathbb{F}_q$. If there exists $y \in \mathbb{F}_q^*$ and $c \in \mathbb{F}_p^* \setminus \{0, 1\}$ such that $\Pi(cy) = c\Pi(y)$, then we have

$$c\Pi(y) = \Pi(cy) = \sum_{\substack{0 \le i \le j \le m-1 \\ 0 \le j \le j \le m-1 \\ 0 \le j \le j \le m-1 \\ 0 \le j \le$$

Since Π is even and $\Pi(0) = 0$, by [8, Lemma 2], we have $\Pi(y) \neq 0$. It follows that $c^2 = c$, which implies that c = 0 or 1. This is a contradiction. Therefore, Π satisfies the conditions in Theorem 5.

As a summary, in Theorem 4 and Theorem 5, we describe completely the weight distributions of the linear codes C_{Π} and $\overline{C_{\Pi}}$ only using the assumption that Π has perfect nonlinearity for all odd primes p when m is odd and give some mild additional conditions for similar conclusions to hold when m is even. As shown in the remarks, these two theorems cover all the previous relevant results (e.g., [12, Theorem 2], [12, Theorem 1, 2], [11, Theorem 4]) as special cases.

The approach employed in this paper, which studies the evolution of the codewords of a linear code under some group action is valuable and typical. It could potentially be extended to investigate other linear codes constructed from perfect nonlinear functions or bent functions (e.g., the linear codes considered in [18]).

REFERENCES

- L. Budaghyan and T. Helleseth, "New perfect nonlinear multinomials over F_{p2k} for any odd prime p," in Sequences and Their Applications-SETA 2008: 5th International Conference Lexington, KY, USA, September 14-18, 2008 Proceedings 5. Springer, 2008, pp. 403–414.
- [2] C. Carlet and C. Ding, "Highly nonlinear mappings," Journal of complexity, vol. 20, no. 2-3, pp. 205–244, 2004.
- [3] C. Carlet, C. Ding, and J. Yuan, "Linear codes from perfect nonlinear mappings and their secret sharing schemes," *IEEE Transactions on Information Theory*, vol. 51, no. 6, pp. 2089–2102, 2005.
- [4] R. Coulter, M. Henderson, L. Hu, P. Kosick, Q. Xiang, and X. Zeng, "Planar polynomials and commutative semifields two dimensional over their middle nucleus and four dimensional over their nucleus," J. Der Mathematischen Ablehnungen, vol. 14, pp. 1–5, 2007.
- [5] R. S. Coulter and R. W. Matthews, "Planar functions and planes of Lenz-Barlotti class II," *Designs, Codes and Cryptography*, vol. 10, no. 2, pp. 167–184, 1997.
- [6] P. Dembowski and T. G. Ostrom, "Planes of order n with collineation groups of order n²," *Mathematische Zeitschrift*, vol. 103, no. 3, pp. 239–258, 1968.
- [7] C. Ding and J. Yuan, "A family of skew Hadamard difference sets," *Journal of Combinatorial Theory, Series A*, vol. 113, no. 7, pp. 1526– 1535, 2006.
- [8] K. Feng and J. Luo, "Value distributions of exponential sums from perfect nonlinear functions and their applications," *IEEE transactions* on information theory, vol. 53, no. 9, pp. 3035–3041, 2007.
- [9] M. Grassl. CodeTables.de. [Online]. Available: http://www.codetables.de
- [10] P. V. Kumar, R. A. Scholtz, and L. R. Welch, "Generalized bent functions and their properties," *Journal of Combinatorial Theory, Series A*, vol. 40, no. 1, pp. 90–107, 1985.
- [11] C. Li, Q. Li, and S. Ling, "Properties and applications of preimage distributions of perfect nonlinear functions," *IEEE transactions on information theory*, vol. 55, no. 1, pp. 64–69, 2008.
- [12] C. Li, L. Qu, and S. Ling, "On the covering structures of two classes of linear codes from perfect nonlinear functions," *IEEE transactions on information theory*, vol. 55, no. 1, pp. 70–82, 2008.
- [13] K. Nyberg, "Constructions of bent functions and difference sets," in Advances in Cryptology—EUROCRYPT'90: Workshop on the Theory and Application of Cryptographic Techniques Aarhus, Denmark, May 21–24, 1990 Proceedings 9. Springer, 1991, pp. 151–160.
- [14] —, "Perfect nonlinear S-boxes," in Advances in Cryptology—EUROCRYPT'91: Workshop on the Theory and Application of Cryptographic Techniques Brighton, UK, April 8–11, 1991 Proceedings 10. Springer, 1991, pp. 378–386.
- [15] V. Pless, R. A. Brualdi, and W. C. Huffman, Handbook of Coding Theory. Elsevier Science Inc., 1998.
- [16] J. Yuan, C. Carlet, and C. Ding, "The weight distribution of a class of linear codes from perfect nonlinear functions," *IEEE transactions on information theory*, vol. 52, no. 2, pp. 712–717, 2006.
- [17] Z. Zha, G. M. Kyureghyan, and X. Wang, "Perfect nonlinear binomials and their semifields," *Finite Fields and Their Applications*, vol. 15, no. 2, pp. 125–133, 2009.
- [18] D. Zheng, X. Wang, Y. Li, and M. Yuan, "Subfield codes of linear codes from perfect nonlinear functions and their duals," *arXiv preprint* arXiv:2012.06105, 2020.