# THE DISCREPANCY OF GREATER-THAN

SRIKANTH SRINIVASAN AND AMIR YEHUDAYOFF

ABSTRACT. The discrepancy of the  $n \times n$  greater-than matrix is shown to be  $\frac{\pi}{2 \ln n}$  up to lower order terms.

The greater-than matrices appear in many areas of mathematics. They form a central example in communication complexity (see e.g. [\[10\]](#page-4-0)). They are studied in analysis as the main triangle projection (e.g. [\[6\]](#page-4-1)). They serve as a model for threshold gates in circuit complexity (e.g. [\[5\]](#page-4-2)). Understanding their properties is therefore a fundamental problem.

**Definition.** Let  $G = G_n$  be the  $n \times n$  signed greater-than matrix: for every j,  $k \in$ [n]*,*

$$
G_{j,k} = 2_{j+k \leq n} - 1.
$$

One way to capture the structure of an object is using discrepancy [\[8,](#page-4-3) [4\]](#page-4-4). On a high-level, discrepancy measures the maximum correlation with certain test functions, and it captures pseudo-randomness properties. Discrepancy of matrices plays a key role in communication complexity. It allows to lower bound randomized communication complexity (see [\[10\]](#page-4-0) and references within), and it allows to bound information complexity [\[3\]](#page-4-5). It also satisfies a direct product property [\[11,](#page-4-6) [7\]](#page-4-7).

**Definition.** The discrepancy of an  $n \times n$  real-valued matrix M with respect to a *distribution*  $\mu = \mu_{j,k}$  *on*  $[n] \times [n]$  *is* 

$$
\textsf{disc}_{\mu}(M) = \max_{x,y \in \mathbb{C}^n: \|x\|_{\infty} = \|y\|_{\infty} = 1} \Big| \sum_{j,k} M_{j,k} \mu_{j,k} x_j y_k \Big|.
$$

*The discrepancy of* M *is*

$$
\operatorname{disc}(M) = \inf_{\mu} \operatorname{disc}_{\mu}(M).
$$

Remark. *In communication complexity, the standard definition of discrepancy uses boolean vectors* x, y *instead of complex vectors. Our proof leads to a sharp bound for the complex version (which is equal to the boolean version up to constant factors).*

The main result of this note is a sharp analysis of the discrepancy of the greaterthan matrix.

Theorem.

$$
\operatorname{disc}(G_n) = (1 - o(1)) \frac{\pi}{2 \ln n}.
$$

This sharp bound improves the previous upper bound  $\operatorname{\sf disc}(G_n) \leq O(\frac{1}{\sqrt{\text{lr}}}$  $\frac{1}{\ln n}$ proved by Braverman and Weinstein [\[3\]](#page-4-5). It also improves all previous lower bounds on the two-party public-coin communication complexity of the greater-than function [\[13,](#page-4-8) [3,](#page-4-5) [9\]](#page-4-9).

To prove an upper bound on  $\textsf{disc}(G)$ , we need to choose an appropriate distribution  $\mu$ . The "natural" distributions—that were used by Viola [\[13\]](#page-4-8), by Braverman and Weinstein [\[3\]](#page-4-5), and by Ramamoorthy and Sinha [\[9\]](#page-4-9)—lead to sub-optimal discrepancy  $\Omega(\frac{1}{\sqrt{k}})$  $\frac{1}{\ln n}$ ; for more details see Section [1.1.](#page-3-0) The distribution we use is constructed via a Hilbert matrix, and it is based on ideas of Kwapien and Pelczyski [\[6\]](#page-4-1) and of Titchmarsh [\[12\]](#page-4-10) from analysis.

To prove a lower bound on  $\textsf{disc}(G)$ , we need to identify the witness vectors  $x, y$ . There is a natural and simple mechanism for locating x, y that yields an  $\Omega(\frac{1}{\ln n})$ lower bound on the discrepancy (see e.g. [\[1\]](#page-4-11)). Getting an exact bound, however, is not so simple. We use deep ideas of Bennett [\[2\]](#page-4-12) from the study of Schur multipliers. Somewhat surprisingly, the mechanism that enables to locate the witnesses  $x, y$ uses abstract machinery, like the Hahn-Banach theorem, the Riesz representation theorem and the F. and M. Riesz theorem.

### 1. The upper bound

Fix *n* for the rest of this text, and let  $H = H_n$  be the following  $n \times n$  version of the Hilbert matrix:

<span id="page-1-1"></span>(1.1) 
$$
H_{j,k} = \frac{1}{n + \frac{1}{2} - j - k}.
$$

This Hilbert matrix has three useful properties that are described in the following three claims.

<span id="page-1-0"></span>Claim 1.  $H_{j,k}G_{j,k} \geq 0$  *for all j, k.* 

<span id="page-1-2"></span>**Claim 2** (follows [\[12\]](#page-4-10)).  $|xHy| \le \pi ||x||_2 ||y||_2$  *for all*  $x, y \in \mathbb{C}^n$ .

Claim 3.  $||H||_1 \geq 2n(\ln(n) - 3)$ .

*Proof of upper bound using the three claims.* Define a distribution  $\mu^*$  on the entries by

$$
\mu_{j,k}^* = \frac{|H_{j,k}|}{\|H\|_1}.
$$

Let  $x, y \in \mathbb{C}^n$  be the vectors of  $\ell_{\infty}$ -norm one that witness the discrepancy of G with respect to  $\mu^*$ . The  $\ell_2$ -norm of x, y is at most  $\sqrt{n}$ . Think of x as a row vector and of y as a column vector. Bound

$$
\operatorname{disc}_{\mu^*}(G) = \frac{1}{\|H\|_1} |xHy| \le \frac{n}{\|H\|_1} \left| \frac{x}{\|x\|_2} H \frac{y}{\|y\|_2} \right| \le \frac{\pi}{2(\ln(n) - 3)}.
$$

*Proof of Claim [2.](#page-1-0)* Because H is symmetric, we need to upper bound its spectral norm over  $\mathbb{R}$ : for every  $x \in \mathbb{R}^n$ ,

$$
||Hx||_2 \le \pi ||x||_2.
$$

To prove this, extend  $H$  to a larger matrix. Let  $N$  be a large integer and let  $J = [-N, N] \cap \mathbb{Z}$ . Let H be the  $J \times J$  matrix defined by the formula in [\(1.1\)](#page-1-1). Let

$$
\tilde{x} \in \mathbb{R}^{J} \text{ be so that } \tilde{x}_{j} = x_{j} \text{ for every } j \in [n], \text{ and } \tilde{x}_{j} = 0 \text{ for every } j \in J \setminus [n]. \text{ Bound}
$$
\n
$$
||Hx||_{2}^{2} \le ||\tilde{H}\tilde{x}||_{2}^{2}
$$
\n
$$
= \sum_{j \in J} \left| \sum_{k \in J} \frac{1}{n + \frac{1}{2} - j - k} \tilde{x}_{k} \right|^{2}
$$
\n
$$
= \sum_{j} \left( \sum_{k} \frac{1}{(n + \frac{1}{2} - j - k)^{2}} \tilde{x}_{k}^{2} \right)
$$
\n
$$
+ \left( \sum_{k_{1} \neq k_{2}} \frac{1}{n + \frac{1}{2} - j - k_{1}} \frac{1}{n + \frac{1}{2} - j - k_{2}} \tilde{x}_{k_{1}} \tilde{x}_{k_{2}} \right)
$$
\n
$$
= \left( \sum_{k} \tilde{x}_{k}^{2} \sum_{i} \frac{1}{(n + \frac{1}{2} - j - k)^{2}} \right)
$$
\n
$$
+ \left( \sum_{k_{1} \neq k_{2}} \tilde{x}_{k_{1}} \tilde{x}_{k_{2}} \sum_{j} \frac{1}{n + \frac{1}{2} - j - k_{1}} \cdot \frac{1}{n + \frac{1}{2} - j - k_{2}} \right).
$$

Bound each of the two terms separately. For the first term, for each  $k$ ,

$$
\sum_{j\in J} \frac{1}{(n+\frac{1}{2}-j-k)^2} \le 2\sum_{\ell=0}^{\infty} \frac{1}{(\ell+\frac{1}{2})^2} = 8\sum_{\ell=0}^{\infty} \frac{1}{(2\ell+1)^2} = \pi^2;
$$

the last equality can be justified as follows. Because  $\sum_{\ell=1}^{\infty} \frac{1}{\ell^2} = \frac{\pi^2}{6}$  $\frac{r^2}{6}$ , we know that  $\sum_{\ell=1}^{\infty} \frac{1}{(2\ell)^2} = \frac{\pi^2}{24}$ , so  $\sum_{\ell=0}^{\infty} \frac{1}{(2\ell+1)^2} = \pi^2(\frac{1}{6} - \frac{1}{24})$ . For the second term, for every  $k_1 \neq k_2,$ 

$$
\left| \sum_{j \in J} \frac{1}{n + \frac{1}{2} - j - k_1} \cdot \frac{1}{n + \frac{1}{2} - j - k_2} \right|
$$
  
= 
$$
\frac{1}{|k_2 - k_1|} \cdot \left| \left( \sum_{i \in I} \frac{1}{n + \frac{1}{2} - j - k_1} \right) - \left( \sum_{i \in I} \frac{1}{n + \frac{1}{2} - j - k_2} \right) \right|
$$
  

$$
\leq \frac{1}{|k_2 - k_1|} \cdot 2|k_2 - k_1| \left| \frac{1}{N - \frac{1}{2} - n - |k_1| - |k_2|} \right|,
$$

where we used the fact that most terms cancel out. The last quantity tends to zero as  $N \to \infty$ . This completes the proof because there are only  $n(n-1)$  significant  $k_1 \neq k_2$ . □  $k_1 \neq k_2$ .

*Proof of Claim [3.](#page-1-2)* For each integer  $0 \leq \ell \leq n-1$ , there are  $n-\ell$  pairs  $(j, k)$  so that  $j + k = n + 1 - \ell$ , and there are  $n - \ell$  pairs  $(j, k)$  so that  $j + k = n + 1 + \ell$ . So,

$$
\sum_{j,k} \left| \frac{1}{n + \frac{1}{2} - j - k} \right| = \sum_{\ell=0}^{n-1} \left| \frac{n - \ell}{n + \frac{1}{2} - (n + 1 - \ell)} \right| + \left| \frac{n - \ell}{n + \frac{1}{2} - (n + 1 + \ell)} \right|
$$
  

$$
= \sum_{\ell=0}^{n-1} \frac{n - \ell}{\ell - \frac{1}{2}} + \frac{n - \ell}{\ell + \frac{1}{2}}
$$
  

$$
= n \Big( \sum_{\ell=0}^{n-1} \frac{1}{\ell - \frac{1}{2}} + \frac{1}{\ell + \frac{1}{2}} \Big) - \Big( \sum_{\ell=0}^{n-1} \frac{\ell}{\ell - \frac{1}{2}} + \frac{\ell}{\ell + \frac{1}{2}} \Big)
$$
  

$$
\ge 2n(\ln(n) - 3).
$$

<span id="page-3-0"></span>1.1. Insufficiency of previous hard distributions. The previous works [\[13,](#page-4-8) [3,](#page-4-5) [9\]](#page-4-9) used different distributions that leads to a sub-optimal bound of  $\text{disc}(G) \leq$  $O(\frac{1}{\sqrt{1}})$  $\frac{1}{\ln n}$ ; this bound was proved in [\[3\]](#page-4-5). In this section, we focus on the distribution  $\eta$ defined in [\[3\]](#page-4-5), and prove that this bound is, in fact, the best that can be obtained for this distribution.

We first recall the distribution  $\eta$ . Assume that  $n = 2^m$  for some positive integer m; otherwise work with the largest power of two smaller than n. Identify  $[n]$  with {0, 1}<sup>m</sup> by identifying j with the binary representation of j − 1 (with the first coordinate being the most significant bit and the last coordinate being the least significant). Sample  $(i, k)$  from  $\eta$  as follows. First, sample j uniformly at random from  $\{0,1\}^m$ , and independently choose a uniformly random  $i \in [m]$ . Set k to be equal to  $j$  in all coordinates less than  $i$ , to be the opposite of  $j$  in coordinate  $i$ , and chosen independently of  $j$  and uniformly in all coordinates greater than  $i$ .

#### Claim. disc $\eta(G) \ge \Omega\left(\frac{1}{\sqrt{\ln n}}\right)$  $rac{1}{\ln n}$ .

*Proof.* Define  $\underline{x} \in \mathbb{R}^n$  to be the indicator vector of all those  $j \in \{0,1\}^m$  that have at least  $\frac{m}{2} + \sqrt{m}$  many one entries, and  $y \in \mathbb{R}^n$  to be the all-ones vector. Denote by  $j_i$  the i'th coordinate of j. The event  $j \geq k$  holds exactly when  $j_i = 1$ . It follows that

$$
\begin{aligned} \text{disc}_{\eta}(G) &= \sum_{j,k} G_{j,k} \eta_{j,k} x_j \\ &= \Pr_{\eta} \left[ x_j = 1 \land j_i = 1 \right] - \Pr_{\mu} \left[ x_j = 1 \land j_i = 0 \right] \\ &= \Pr_{\eta} \left[ x_j = 1 \right] \cdot \left( 2 \Pr_{\mu} \left[ j_i = 1 \mid x_j = 1 \right] - 1 \right). \end{aligned}
$$

Standard estimates imply that  $Pr_n[x_j = 1] = \Omega(1)$ . Conditioned on any such choice for j, the random variable i is still uniform, and thus the probability that  $j_i = 1$  is at least  $\frac{1}{2} + \frac{1}{\sqrt{m}}$ .

## 2. The lower bound

In this section, we work with the matrix  $G_{j,k} = 2_{j\geq k} - 1$ . The advantage is that this G is a Toeplitz matrix  $G_{j,k} = c_{j-k}$ , where in the previous sections it was a Hankel matrix  $G_{j,k} = c_{j+k}$ . This distinction is meaningless in terms of discrepancy, but turns out to be important in analysis.

The mechanism for locating the witnesses  $x, y$  relies on the existence of a certain function  $\nu$  that encodes the interaction of G with many witnesses (via the Fourier transform). A *complex measure* is a bounded, Borel, absolutely continuous ν :  $[0, 1) \rightarrow \mathbb{C}$ . Its Fourier transform is defined to be

$$
\mathbb{Z} \ni \ell \mapsto \hat{\nu}(\ell) = \int_0^1 e^{2\pi i \ell t} \nu(t) dt \in \mathbb{C}.
$$

Its norm is defined to be

$$
||\nu|| = \sup_{f:[0,1)\to\mathbb{C}, ||f||_{\infty}=1} \Big| \int_0^1 \nu(t) f(t) dt \Big|.
$$

 $-1$ 

The following is implicit in Bennett's work.

<span id="page-4-13"></span>**Lemma 4** (implicit in [\[2\]](#page-4-12)). *There is a complex measure*  $\nu$  *so that for all*  $\ell$  *so that*  $|\ell| \leq n$ ,

(2.1) 
$$
\hat{\nu}(\ell) = \begin{cases} 1 & \ell \ge 0 \\ 0 & \ell < 0 \end{cases}
$$

*and so that*

<span id="page-4-14"></span><sup>k</sup>νk ≤ ln <sup>n</sup> π (2.2) + 2.

*Proof of Lemma [4.](#page-4-13)* The references here are in Bennett's work [\[2\]](#page-4-12). Theorem 8.1 states the existence of v satisfying  $\hat{\nu}(\ell) = M_{j+\ell,j}$  and  $\|\nu\| = \|M\|_{(\infty,1)}$  for all Toeplitz matrices that are multipliers. Let M be the  $n \times n$  Toeplitz matrix  $M_{i,k} =$  $1_{i\geq k}$ . The proof of Corollary 8.5 shows that M is a multiplier and also proves the bound on  $\|\nu\| = \|M\|_{(\infty,1)}$  stated in [\(2.2\)](#page-4-14). In the proof of Corollary 8.3, it is also proved that  $\nu$  is an absolutely continuous  $L^1$ -function. proved that  $\nu$  is an absolutely continuous  $L^1$ -function.

*Proof of lower bound.* Let  $\mu = \mu_{j,k}$  be a distribution. Without loss of generality, we can assume  $Pr_{\mu}[j \ge k] \ge \frac{1}{2}$ . By Lemma [4,](#page-4-13)

$$
\frac{1}{2} \leq \Big| \sum_{j,k} G_{j,k} \mu_{j,k} \hat{\nu}(j-k) \Big|
$$
\n
$$
= \Big| \sum_{j,k} G_{j,k} \mu_{j,k} \int \nu(t) e^{2\pi i (j-k)t} dt \Big|
$$
\n
$$
= \Big| \int \nu(t) \sum_{j,k} G_{j,k} \mu_{j,k} e^{2\pi i jt} e^{-2\pi ikt} dt \Big|
$$
\n
$$
\leq ||\nu|| \cdot \text{disc}_{\mu}(G). \qquad \Box
$$

### **REFERENCES**

- <span id="page-4-11"></span>1. Daniel Avraham and Amir Yehudayoff, *On blocky ranks of matrices*, Electronic Colloquium on Computational Complexity (ECCC), 2022.
- <span id="page-4-12"></span><span id="page-4-5"></span>2. G. Bennett, *Schur multipliers*, Duke Mathematical Journal 44 (1977), no. 3.
- 3. Mark Braverman and Omri Weinstein, *A discrepancy lower bound for information complexity*, Algorithmica 76 (2016), 846–864.
- <span id="page-4-4"></span>4. Bernard Chazelle, *The discrepancy method*, International Symposium on Algorithms and Computation, Springer, 1998, pp. 1–3.
- <span id="page-4-2"></span>5. András Hajnal, Wolfgang Maass, Pavel Pudlák, Mario Szegedy, and György Turán, Threshold *circuits of bounded depth*, Journal of Computer and System Sciences 46 (1993), no. 2, 129–154.
- <span id="page-4-1"></span>6. Stanisław Kwapień and Aleksander Pełczyński, *The main triangle projection in matrix spaces and its applications*, Studia Mathematica 34 (1970), no. 1, 43–67.
- <span id="page-4-7"></span>7. Troy Lee, Adi Shraibman, and Robert Špalek, A direct product theorem for discrepancy, CCC, 2008, pp. 71–80.
- <span id="page-4-9"></span><span id="page-4-3"></span>8. Jiri Matousek, *Geometric discrepancy: An illustrated guide*, vol. 18, Springer, 1999.
- 9. Sivaramakrishnan Natarajan Ramamoorthy and Makrand Sinha, *On the communication complexity of greater-than.*, Allerton, 2015, pp. 442–444.
- <span id="page-4-0"></span>10. Anup Rao and Amir Yehudayoff, *Communication complexity and applications*, Cambridge University Press, 2020.
- <span id="page-4-10"></span><span id="page-4-6"></span>11. Ronen Shaltiel, *Towards proving strong direct product theorems*, CCC, 2001, pp. 107–117.
- 12. Edward Charles Titchmarsh, *Reciprocal formulae involving series and integrals*, Mathematische Zeitschrift 25 (1926), no. 1, 321–347.
- <span id="page-4-8"></span>13. Emanuele Viola, *The communication complexity of addition*, Combinatorica 35 (2015), 703– 747.

Department of Computer Science, University of Copenhagen and Department of Computer Science, Aarhus University *Email address*: srsr@di.ku.dk

Department of Computer Science, University of Copenhagen and Department of Mathematics, Technion-IIT

*Email address*: amir.yehudayoff@gmail.com