

# Byzantine Multiple Access Channels — Part II: Communication With Adversary Identification

Neha Sangwan, *Student Member, IEEE*, Mayank Bakshi, *Member, IEEE*, Bikash Kumar Dey, *Member, IEEE*, and Vinod M. Prabhakaran, *Member, IEEE*

## Abstract

We introduce the problem of determining the identity of a byzantine user (internal adversary) in a communication system. We consider a two-user discrete memoryless multiple access channel where either user may deviate from the prescribed behaviour. Owing to the noisy nature of the channel, it may be overly restrictive to attempt to detect all deviations. In our formulation, we only require detecting deviations which impede the decoding of the non-deviating user's message. When neither user deviates, correct decoding is required. When one user deviates, the decoder must either output a pair of messages of which the message of the non-deviating user is correct or identify the deviating user. The users and the receiver do not share any randomness. The results include a characterization of the set of channels where communication is feasible, and an inner and outer bound on the capacity region. We also show that whenever the rate region has non-empty interior, the capacity region is same as the capacity region under randomized encoding, where each user shares independent randomness with the receiver. We also give an outer bound for this randomized coding capacity region.

## I. INTRODUCTION

In many modern wireless communication applications (e.g., the Internet of Things), devices with varying levels of security are connected over a shared communication medium. Compromised devices may allow an adversary to disrupt the communication of other devices. This motivates the question we study in this paper – is it possible to design a communication system in which malicious actions by compromised devices can be detected so that such devices can be isolated or taken offline?

We consider a two-user Multiple Access Channel (MAC) where either user may deviate from the prescribed behaviour. The deviating user (if any) is fixed for the entire duration of the transmission. We will refer to this channel as a byzantine-MAC in the rest of the paper. Owing to the noisy nature of the channel, it may be impossible or overly restrictive to attempt to detect all deviations. Indeed, it suffices to detect only such deviations which impede the correct decoding of the other user's message. We formulate a communication problem for the byzantine-MAC with the following decoding guarantee (Fig. 1): the decoder outputs either a pair of messages *or* declares one of the users to be deviating. When both users are honest, the decoder must output the correct message pair with high probability (w.h.p.); when exactly one user deviates, w.h.p., the decoder must either correctly detect the deviating user or output a message pair of which the message of the other (honest) user is correct (see Section II). No guarantees are made if both users deviate. Thus, we require that a deviating user cannot cause a decoding error for the other user without getting caught. We call this problem as that of *communication with adversary identification*. The focus of this paper is on the case where the decoder does not share any randomness with either of the encoders. We consider the average probability of error criterion, that is, an honest user sends a codeword uniformly at random. The identity of this codeword is not known to the other user.




Fig. 1. *Communication with adversary identification* in a byzantine-MAC: Reliable decoding of both the messages is required when neither user deviates. When a user (say, user B) deviates, the decoded message should either be correct for the honest user or the decoder should identify the deviating user (by outputting **b**) with high probability.

For comparison, consider the stronger guarantee of *reliable communication* where the decoder outputs a pair of messages such that the message(s) of non-deviating user(s) is correct w.h.p. [1], [2]. While achieving this clearly satisfies the requirements of the present model, it might be too demanding as we discuss in the following example.

N. Sangwan and V. Prabhakaran acknowledge support of the Department of Atomic Energy, Government of India, under project no. RTI4001. N. Sangwan's work was additionally supported by the Tata Consultancy Services (TCS) foundation through the TCS Research Scholar Program. Work of B. K. Dey was supported in part by Bharti Centre for Communication in IIT Bombay. V. Prabhakaran's work was also supported by the Science & Engineering Research Board, India through project MTR/2020/000308. The work of M. Bakshi was supported in part by the Research Grants Council of the Hong Kong Special Administrative Region, China, under Grant GRF 14300617, and in part by the National Science Foundation under Grant No. CCF-2107526.

N. Sangwan and V. M. Prabhakaran are with the School of Technology and Computer Science, Tata Institute of Fundamental Research, Mumbai 400 005 India.

M. Bakshi is with Arizona State University, Tempe, AZ, USA.

B. K. Dey is with Indian Institute of Technology Bombay, Mumbai 400 076, India.

**Example 1** (Binary erasure MAC [3, pg. 83]). *Binary erasure MAC is a deterministic MAC model with binary inputs  $X, Y$  and ternary output  $Z = X + Y$  where  $+$  is real addition and  $Z \in \{0, 1, 2\}$ . In this channel, a deviating user can run an independent copy of the honest user's encoder and inject a spurious message which will appear equally plausible to the decoder as the honest user's actual message (see Example 2 in Section V-B). Thus, it is not possible to drive the probability of error to zero for any message set of size at least two under the reliable communication guarantee. However, consider a simple scheme which uses all strings of Hamming weight 1 as codebook of user A and all strings of Hamming weight  $n - 1$  as codebook of user B. We use the following decoder: If the sum of entries of the output string is  $n + 2$  or more, user A is malicious. On the other hand, if it is  $n - 2$  or less, then user B is malicious. Note that the output will be correct with probability 1 in these cases. When the sum of entries of the output string is  $n + 1$  but there is no location with symbol 0, then the decoder declares user B to be malicious, otherwise, user A is malicious. This is because when user A is malicious and sends a string of hamming weight 2, user B, being honest, sends one of the codewords uniformly at random, the identity of which is unknown to user A. Thus, the probability that there is no 0 in the output string is  $2/n$  which vanishes with  $n$ . Similarly, when the sum of entries of the output string is  $n - 1$ , the decoder declares user A to be malicious when there is no 2 in the output string, and declares user B to be malicious otherwise. Along similar lines, we can also argue that when the sum of entries of the output string is  $n$ , with a non-vanishing probability, there is a 2 and a 0 in the output string. Such an output only corresponds to two unique input strings of users which are both valid codewords. Hence, the output would be correct. Thus, we can get a vanishing probability of error using this scheme, though with no rate. In fact, we show that for this channel, the capacity region of communication with adversary identification is the same as the (non-adversarial) capacity region of the binary erasure MAC (see Section V-A).*

Another decoding guarantee, that is weaker than the present model, allows the decoder to declare adversarial interference (in the presence of malicious user(s)) without identifying the adversary. We called this model *authenticated communication* and characterized its feasibility condition and capacity region in [4]. The feasibility condition is called *overwritability*, a notion which was introduced by Kosut and Kliever for network coding [5] and arbitrarily varying channels [6].

The present model lies between the models for reliable communication and authenticated communication in a byzantine MAC. However, obtaining results here appears to be significantly more challenging. On the one hand, for reliable communication over the two-user MAC, we may treat the channel from each user to the decoder as an arbitrarily varying channel (AVC) [7] with the other user's input as state. Hence, the users may send their messages using the corresponding AVC codes [8]. Thus, the rectangular region defined by the capacities of the two AVCs is achievable<sup>1</sup>. On the other hand, for authenticated communication over the two-user MAC, our achievable strategy in [4] involved an unauthenticated communication phase using a non-adversarial MAC code followed by separate (short) authentication phases for each user's decoded message. Failure to authenticate a user's message implies the presence of an adversary (though not its identity since the user whose message is being authenticated might have deviated to cause the authentication failure). In both the cases above, the decoder, when it accounts for the byzantine nature of the users, deals with the users one at a time. However, similar decoding strategies seem to be insufficient for adversary identification. Determining the identity of a deviating user requires dealing with the byzantine nature of both users simultaneously, thereby complicating the decoder design (see Section III).

We characterize the infeasibility of communication with adversary identification using a condition on the channel we call *spoofability* (see Fig. 2). In a spoofable channel, a deviating user may mount an attack which can be confused with an attack of the other user and which introduces a spurious message that can be confused with the actual message of the (other) honest user. When the channel is not spoofable, a deterministic code in the style of [8] can provide positive rates to both the users (Theorem 1). We give an inner and an outer bound to the deterministic coding capacity region. Our outer bound is in terms of the capacity of an arbitrarily varying multiple access channel [9] (Theorem 9). Further, in Section V-B, a comparison is made between spoofability and the feasibility conditions for the reliable communication and authenticated communication models.

In Section VI, we draw connections of the present model to the case when the users share independent randomness with the receiver. Analogous to the *dichotomy phenomenon* for arbitrarily varying channels [10] and arbitrarily varying multiple access channels [11], we show that whenever the capacity region for the deterministic case has a non-empty interior, it is the same as the capacity region for the randomized case. We also give an outer bound on the capacity region for randomized codes.

*Related works:* The model falls in the general class of adversarial channels. There is a long line of works in the information theory literature on communication in the presence of external adversaries (see [12] for a survey), for example, an arbitrarily varying channel [7], [8], [10] or an arbitrarily varying-MAC [9], [11], [13]. In these models the channel law can be arbitrarily varied by an adversary during transmission from a given set of allowed channel laws. While our model is different from these models, the technical formulation is heavily inspired. For example, similar to our deterministic coding with an average probability of error criterion, the most well studied model in arbitrarily varying channels uses a deterministic codebook with an average probability of error criterion where a user sends a codeword uniformly at random. The identity of the sent codeword(s) is hidden from the adversary. Our randomized coding model is also inspired from the corresponding randomized coding model in arbitrarily varying multiple access channels where each user shares independent randomness with the decoder which is private

<sup>1</sup>In fact, this rectangular region defined by the capacities of the two AVCs is the reliable communication capacity region since a deviating user can act exactly like the adversary in the AVC of the other user. Note that the AVCs for binary erasure MAC have zero AVC capacity.

from the other user. Our techniques also borrow from the achievability and converse techniques in the arbitrarily varying channels literature. Further, they expand and add to the existing set of available techniques for these problems. In addition to communication over arbitrarily varying channels, there has also been some recent work on authenticated communication over channels in which an external adversary may be present. In the presence of the adversary, the decoder may declare adversarial interference instead of decoding [6], [14]–[16] (In a 2-user MAC model in [16] when declaring the presence of an adversary, the decoder is required to decode at least one user’s message.).

These models are different from the present model, where a legitimate user of the channel maybe adversarial, and when declaring the presence of an adversary, we also require the decoder to output its identity. Such users are often called *byzantine* users. Communication in systems with byzantine users has also received some attention [1], [2], [4], [5], [17]–[19]. Networks coding with byzantine attacks on nodes and edges has been studied [5], [17], [18]. He and Yener [19] considered a Gaussian two-hop network with an eavesdropping and byzantine adversarial relay where the receiver is required to decode with message secrecy and detect byzantine attack. The present model, on the other hand, considers byzantine users in a multiple access channel. This model was previously considered in [1], [2], [4] but with different decoding guarantees. From a cryptographic point of view, message authentication codes where the users have pre-shared keys and communicate over noiseless channels have been extensively studied [20]–[22]. Message authentication over noisy channels has also been considered [22]–[25].

*Summary of contributions:* These are the main contributions of this work.

- We introduce the problem of communication with adversary identification in a byzantine-MAC and characterize the class of byzantine-MACs which allow positive rates using deterministic codes under the average probability of error criterion.
- We also provide inner and outer bounds to the capacity region.
- For a byzantine-MAC, we compare the feasibility condition for communication with adversary identification with the corresponding feasibility conditions in the stronger and weaker models of *reliable communication* and *authenticated communication*, and show a separation using examples.
- We show that, for communication with adversary identification, whenever positive rates can be provided to both users under deterministic coding, the capacity region of deterministic coding is the same as the randomized coding capacity region. This is like the *dichotomy phenomenon* in [10] and [11] for arbitrarily varying channels and arbitrarily varying multiple access channels respectively. We also give an outer bound on the randomized coding capacity region.

## II. SYSTEM MODEL

*Notation:* For a set  $S \subseteq \mathbb{R}^k$ , let  $\text{conv}(S)$  and  $\text{int}(S)$  denote its convex closure and interior respectively. For a set  $\mathcal{H}$ , let  $\mathcal{H}^c$  denote its complement. For a set  $\mathcal{A}$ ,  $\text{Unif}(\mathcal{A})$  denotes the uniform distribution over  $\mathcal{A}$ . We denote random variables by capital letters, like  $X, Y$  and  $\tilde{Y}$ , and their corresponding alphabets by calligraphic letters, for example,  $\mathcal{X}, \mathcal{Y}$  and  $\tilde{\mathcal{Y}}$ . Let  $\mathbf{x} \in \mathcal{X}^n$  (resp.  $\mathbf{X}$  distributed over  $\mathcal{X}^n$ ) denote the  $n$ -length vectors (resp.  $n$ -length random vectors). We denote the distribution of a random variable  $Y$  by  $P_Y$  and use the notation  $Y \sim P_Y$  to indicate this. For an alphabet  $\mathcal{X}$ , we define the set of all empirical distributions of  $n$ -length sequences in  $\mathcal{X}^n$  by  $\mathcal{P}_{\mathcal{X}}^n$ . For a distribution  $P_X \in \mathcal{P}_{\mathcal{X}}^n$ , let  $T_X^n$  denote the set of all  $n$ -length sequences  $\mathbf{x} \in \mathcal{X}^n$  with empirical distribution  $P_X$ . For a vector  $\mathbf{y} \in \mathcal{Y}^n$ , the statement  $\mathbf{y} \in T_Y^n$  is sometimes used to implicitly define  $P_Y$  as the empirical distribution of  $\mathbf{y}$  and a random variable  $Y$  distributed according to  $P_Y$ . For  $P_{XY} \in \mathcal{P}_{\mathcal{X} \times \mathcal{Y}}^n$  and  $\mathbf{x} \in T_X^n$ , we define  $T_{Y|X}^n(\mathbf{x}) \stackrel{\text{def}}{=} \{\mathbf{y} | (\mathbf{x}, \mathbf{y}) \in T_{XY}^n\}$ . For a natural number  $n$ , we denote the set  $\{1, 2, \dots, n\}$  by  $[1 : n]$ . For a real number  $a$ ,  $\exp(a)$  denotes  $2^a$  and  $\log(a)$  denotes  $\log_2(a)$ , that is,  $\exp$  and  $\log$  are with respect to base 2. For a conditional distribution  $P_{Y|X}$ , we denote its  $n$ -fold product (memoryless channel) by  $P_{Y|X}^n$ . For a vector  $\mathbf{x} \in \mathcal{X}^n$ , the term  $P_{Y|X}^n(\cdot | \mathbf{x})$  denotes the output distribution on  $\mathcal{Y}^n$  when  $\mathbf{x}$  is fed as input to the memoryless channel  $P_{Y|X}^n$ . For a 2-user multiple access channel  $W_{Z|XY}$  from input alphabets  $\mathcal{X}$  and  $\mathcal{Y}$  to output alphabet  $\mathcal{Z}$ , we will sometimes use  $W$  to simplify the notation. Its  $n$ -fold product will be denoted by  $W^n$ . For a two-user MAC  $W$ , we will use  $\mathcal{C}_{\text{MAC}}(W)$  (or simply  $\mathcal{C}_{\text{MAC}}$ ) to denote its (non-adversarial) capacity region.

Consider a two-user discrete memoryless Multiple Access Channel (MAC) as shown in Fig. 1. User A has input alphabet  $\mathcal{X}$  and user B has input alphabet  $\mathcal{Y}$ . The output alphabet of the channel is  $\mathcal{Z}$ . The sets  $\mathcal{X}$ ,  $\mathcal{Y}$  and  $\mathcal{Z}$  are finite. We study communication in a MAC where either user may deviate from the communication protocol by sending any sequence of its choice from its input alphabet. While doing so, the deviating user is unaware of other user’s input. Further, the deviating user (if any) is fixed for the entire duration of the transmission. We will refer to this channel model as a *byzantine-MAC*.

**Definition 1** (Adversary identifying code). *An  $(N_A, N_B, n)$  deterministic adversary identifying code for a byzantine-MAC consists of the following:*

- (i) Two message sets,  $\mathcal{M}_i = \{1, \dots, N_i\}$ ,  $i = A, B$ ,
- (ii) Two deterministic encoders,  $f_A : \mathcal{M}_A \rightarrow \mathcal{X}^n$  and  $f_B : \mathcal{M}_B \rightarrow \mathcal{Y}^n$ , and
- (iii) A deterministic decoder,  $\phi : \mathcal{Z}^n \rightarrow (\mathcal{M}_A \times \mathcal{M}_B) \cup \{\mathbf{a}, \mathbf{b}\}$ .

For notational convenience, let us define the decoder  $\phi_A$  for user A's message as, for  $\mathbf{z} \in \mathcal{Z}^n$ ,

$$\phi_A(\mathbf{z}) = \begin{cases} m_A & \text{if } \phi(\mathbf{z}) = (m_A, m_B) \\ \mathbf{a} & \text{if } \phi(\mathbf{z}) = \mathbf{a} \\ \mathbf{b} & \text{if } \phi(\mathbf{z}) = \mathbf{b}, \end{cases} \quad (1)$$

and the decoder  $\phi_B$  for user B's message as, for  $\mathbf{z} \in \mathcal{Z}^n$ ,

$$\phi_B(\mathbf{z}) = \begin{cases} m_B & \text{if } \phi(\mathbf{z}) = (m_A, m_B) \\ \mathbf{a} & \text{if } \phi(\mathbf{z}) = \mathbf{a} \\ \mathbf{b} & \text{if } \phi(\mathbf{z}) = \mathbf{b}. \end{cases} \quad (2)$$

The decoder outputs the symbol  $\mathbf{a}$  to declare that user A is adversarial. Similarly, an output of  $\mathbf{b}$  is used to declare that user B is adversarial. The *average probability of error*  $P_e(f_A, f_B, \phi)$  is the maximum of the average probabilities of error in the following three cases: (1) both users are honest, (2) user A is adversarial, and (3) user B is adversarial. When both users are honest, an error occurs if the decoder does not output the pair of correct messages. Let  $\mathcal{E}_{m_A, m_B} = \{\mathbf{z} : \phi(\mathbf{z}) \neq (m_A, m_B)\}$  denote the corresponding error event. The average error probability when both users are honest is

$$P_{e, \text{hon}} \stackrel{\text{def}}{=} \frac{1}{N_A \cdot N_B} \sum_{\substack{(m_A, m_B) \in \\ \mathcal{M}_A \times \mathcal{M}_B}} W^n \left( \mathcal{E}_{m_A, m_B} | f_A^{(n)}(m_A), f_B^{(n)}(m_B) \right). \quad (3)$$

When user A is adversarial, an error occurs unless the decoder's output is either the symbol  $\mathbf{a}$  or a pair of messages of which the message of user B is correct. The corresponding error event is  $\mathcal{E}_{m_B}^B \stackrel{\text{def}}{=} \{\mathbf{z} : \phi_B(\mathbf{z}) \notin \{m_B, \mathbf{a}\}\}$ . The average probability of error when user A is adversarial is

$$P_{e, \text{mal A}} \stackrel{\text{def}}{=} \max_{\mathbf{x} \in \mathcal{X}^n} \left( \frac{1}{N_B} \sum_{m_B \in \mathcal{M}_B} W^n \left( \mathcal{E}_{m_B}^B | \mathbf{x}, f_B(m_B) \right) \right). \quad (4)$$

Similarly, for  $\mathcal{E}_{m_A}^A \stackrel{\text{def}}{=} \{\mathbf{z} : \phi_A(\mathbf{z}) \notin \{m_A, \mathbf{b}\}\}$ , the average probability of error when user B is adversarial is

$$P_{e, \text{mal B}} \stackrel{\text{def}}{=} \max_{\mathbf{y} \in \mathcal{Y}^n} \left( \frac{1}{N_A} \sum_{m_A \in \mathcal{M}_A} W^n \left( \mathcal{E}_{m_A}^A | f_A(m_A), \mathbf{y} \right) \right). \quad (5)$$

We define the *average probability of error* as

$$P_e(f_A, f_B, \phi) \stackrel{\text{def}}{=} \max \{P_{e, \text{hon}}, P_{e, \text{mal A}}, P_{e, \text{mal B}}\}.$$

We note that  $\mathcal{E}_{m_A, m_B} = \mathcal{E}_{m_A}^A \cup \mathcal{E}_{m_B}^B$ . Thus,

$$\begin{aligned} P_{e, \text{hon}} &= \frac{1}{N_A \cdot N_B} \sum_{(m_A, m_B) \in \mathcal{M}_A \times \mathcal{M}_B} W^n \left( \mathcal{E}_{m_A}^A \cup \mathcal{E}_{m_B}^B | f_A(m_A), f_B(m_B) \right) \\ &\leq \frac{1}{N_A \cdot N_B} \sum_{(m_A, m_B) \in \mathcal{M}_A \times \mathcal{M}_B} \left( W^n \left( \mathcal{E}_{m_A}^A | f_A(m_A), f_B(m_B) \right) + W^n \left( \mathcal{E}_{m_B}^B | f_A(m_A), f_B(m_B) \right) \right) \\ &= \frac{1}{N_B} \sum_{m_B \in \mathcal{M}_B} \left( \frac{1}{N_A} \sum_{m_A \in \mathcal{M}_A} W^n \left( \mathcal{E}_{m_A}^A | f_A(m_A), f_B(m_B) \right) \right) \\ &\quad + \frac{1}{N_A} \sum_{m_A \in \mathcal{M}_A} \left( \frac{1}{N_B} \sum_{m_B \in \mathcal{M}_B} W^n \left( \mathcal{E}_{m_B}^B | f_A(m_A), f_B(m_B) \right) \right) \\ &\leq P_{e, \text{mal A}} + P_{e, \text{mal B}}. \end{aligned} \quad (6)$$

So, if  $P_{e, \text{mal A}}$  and  $P_{e, \text{mal B}}$  are small,  $P_{e, \text{hon}}$  is also small.

**Remark 1.** Note that the probability of error under a randomized attack is the weighted average of the probabilities of errors under the different deterministic attacks and hence maximized by a deterministic attack. Thus,  $P_{e, \text{mal B}}$  is an upper bound on the probability of error for any attack by user B, deterministic or random. Similarly,  $P_{e, \text{mal A}}$  is an upper bound for any attack by user A. Thus, the probability of error under deterministic attacks is same as that under randomized attacks.

**Definition 2** (Achievable rate pair and capacity region for communication with adversary identification).  $(R_A, R_B)$  is an achievable rate pair for communication with adversary identification if there exists a sequence of  $(\lfloor 2^{nR_A} \rfloor, \lfloor 2^{nR_B} \rfloor, n)$  adversary




Fig. 2. When (7) holds for a byzantine-MAC  $W$ , for  $(\mathbf{x}', \tilde{\mathbf{x}}, \tilde{\mathbf{y}}) \in \mathcal{X}^n \times \mathcal{X}^n \times \mathcal{Y}^n$ , the output distributions in the three cases above will be the same.




Fig. 3. A byzantine-MAC  $W$  is B-spoofable if for each  $\tilde{x}, \tilde{y}, y', z$  the conditional output distributions  $P(z|\tilde{x}, \tilde{y}, y')$  in 3(a), 3(b) and 3(c) are the same.

identifying codes  $\{f_A^{(n)}, f_B^{(n)}, \phi^{(n)}\}_{n=1}^{\infty}$  such that  $\lim_{n \rightarrow \infty} P_e(f_A^{(n)}, f_B^{(n)}, \phi^{(n)}) = 0$ . The capacity region of communication with adversary identification  $\mathcal{C}$  is the closure of the set of all such achievable rate pairs.

**Remark 2.** Note that the capacity region of a MAC where both users are honest, denoted by  $\mathcal{C}_{\text{MAC}}$ , is an outer bound on the capacity region of communication with adversary identification  $\mathcal{C}$ , that is,  $\mathcal{C} \subseteq \mathcal{C}_{\text{MAC}}$ . This is because when both users are honest, an adversary identifying code guarantees reliable decoding for both users.

### III. FEASIBILITY OF COMMUNICATION WITH ADVERSARY IDENTIFICATION

**Definition 3** (Spoofable byzantine-MAC). A byzantine-MAC  $W_{Z|XY}$  is A-spoofable if there exist conditional distributions  $Q_{Y|\tilde{X}\tilde{Y}}$  and  $Q_{X|\tilde{X}X'}$ , such that  $\forall x', \tilde{x} \in \mathcal{X}, \tilde{y} \in \mathcal{Y}, z \in \mathcal{Z}$ ,

$$\begin{aligned} & \sum_y Q_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x', y) \\ &= \sum_y Q_{Y|\tilde{X}\tilde{Y}}(y|x', \tilde{y}) W_{Z|XY}(z|\tilde{x}, y) \\ &= \sum_x Q_{X|\tilde{X}X'}(x|\tilde{x}, x') W_{Z|XY}(z|x, \tilde{y}). \end{aligned} \quad (7)$$

A byzantine-MAC  $W_{Z|XY}$  is B-spoofable (see Fig. 3.) if there exist conditional distributions  $Q_{X|\tilde{X}\tilde{Y}}$  and  $Q_{Y|\tilde{Y}Y'}$ , such that  $\forall \tilde{x} \in \mathcal{X}, \tilde{y}, y' \in \mathcal{Y}, z \in \mathcal{Z}$ ,

$$\begin{aligned} & \sum_x Q_{X|\tilde{X}\tilde{Y}}(x|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y') \\ &= \sum_x Q_{X|\tilde{X}\tilde{Y}}(x|\tilde{x}, y') W_{Z|XY}(z|x, \tilde{y}) \\ &= \sum_y Q_{Y|\tilde{Y}Y'}(y|\tilde{y}, y') W_{Z|XY}(z|\tilde{x}, y). \end{aligned} \quad (8)$$

A byzantine-MAC is spoofable if it is either A-spoofable or B-spoofable.

Our definition of spoofable channels is motivated by a scenario where the decoder cannot differentiate between two different likely transmitted codewords of user A, while at the same time it cannot blame user B for the situation since the situation appears to be possible due to an adversarial action of either user A or user B. To see this, let  $(\mathbf{x}', \tilde{\mathbf{x}}, \tilde{\mathbf{y}}) \in \mathcal{X}^n \times \mathcal{X}^n \times \mathcal{Y}^n$ . When a byzantine-MAC is A-spoofable, i.e. (7) holds, the output distributions in the following three cases are the same (see Fig. 2):

- User A sends  $\mathbf{x}'$  and an adversarial user B sends  $\mathbf{Y} \sim Q_{Y|\tilde{X}\tilde{Y}}^n(\cdot|\tilde{\mathbf{x}}, \tilde{\mathbf{y}})$ , i.e.,  $\mathbf{Y}$  is distributed as the output of the memoryless channel  $Q_{Y|\tilde{X}\tilde{Y}}$  on inputs  $\tilde{\mathbf{x}}$  and  $\tilde{\mathbf{y}}$ ;
- User A sends  $\tilde{\mathbf{x}}$  and an adversarial user B sends  $\mathbf{Y} \sim Q_{Y|\tilde{X}\tilde{Y}}^n(\cdot|\mathbf{x}', \tilde{\mathbf{y}})$ ;
- User B sends  $\tilde{\mathbf{y}}$  and an adversarial user A sends  $\mathbf{X} \sim Q_{X|\tilde{X}X'}^n(\cdot|\tilde{\mathbf{x}}, \mathbf{x}')$ .



In Lemma 3 (Sec. III-A), we use the above property of spoofable channels to show that for an A-spoofable byzantine-MAC, user-A cannot send even one bit reliably. Similarly, for a B-spoofable byzantine-MAC, user-B cannot send one bit reliably.<sup>2</sup> For an A-spoofable channel, our proof considers any given code  $(f_A, f_B, \phi)$  and independent  $M_A \sim \text{Unif}(\mathcal{M}_A)$ ,  $M'_A \sim \text{Unif}(\mathcal{M}_A)$  and  $M_B \sim \text{Unif}(\mathcal{M}_B)$ . By replacing  $(\mathbf{x}', \tilde{\mathbf{x}}, \tilde{\mathbf{y}})$  with  $(f_A(M_A), f_A(M'_A), f_B(M_B))$  in the above argument, we note that the output distributions in the following three cases are the same:

- (a) User A is honest and sends  $f_A(M_A)$  and user B is adversarial and attacks with  $\mathbf{Y} \sim Q_{Y|\tilde{X}\tilde{Y}}^n(\cdot|f_A(M'_A), f_B(M_B))$ ;
- (b) User A is honest and sends  $f_A(M'_A)$  and user B is adversarial and attacks with  $\mathbf{Y} \sim Q_{Y|\tilde{X}\tilde{Y}}^n(\cdot|f_A(M_A), f_B(M_B))$ ;
- (c) User B is honest and sends  $f_B(M_B)$  and user A is adversarial and attacks with  $\mathbf{X} \sim Q_{X|\tilde{X}X'}^n(\cdot|f_A(M_A), f_A(M'_A))$ .

Thus, the decoder cannot determine the adversarial user reliably, nor can it differentiate between  $M_A$  and  $M'_A$  as the input of user A.

Our first result states that non-spoofability characterizes the byzantine-MACs in which users can work at positive rates of communication with adversary identification.

**Theorem 1.** *If a byzantine-MAC is A-spoofable (resp. B-spoofable), communication with adversary identification from user-A (resp. user-B) is impossible. Specifically, for any  $(N_A, N_B, n)$  adversary identifying code with  $N_A \geq 2$  (resp.  $N_B \geq 2$ ), the probability of error is at least  $1/12$ . If a byzantine-MAC is neither A-spoofable nor B-spoofable, then its capacity region has a non-empty interior ( $\text{int}(\mathcal{C}) \neq \emptyset$ ), i.e., both users can communicate reliably with adversary identification at positive rates.*

**Corollary 2.**  $\text{int}(\mathcal{C}) = \emptyset$  if and only if a byzantine-MAC is spoofable.

**Remark 3.** *Theorem 1 does not cover the case when exactly one user is spoofable. A similar case is also open for Arbitrarily Varying Multiple Access MAC (AV-MAC) (see [9]).<sup>3</sup>*

Before presenting a proof of the forward direction of Theorem 1, we give an informal description of the decoder we use. For input distributions  $P_A$  and  $P_B$  on  $\mathcal{X}$  and  $\mathcal{Y}$  respectively, the decoder works by first collecting potential candidates for the messages sent by each user. A message  $m_A$  is deemed a *candidate* for user A if it is typical with some (attack) vector  $\mathbf{y}$  and the output vector  $\mathbf{z}$  according to the channel law (i.e., for some  $\eta > 0$ ,  $(f_A(m_A), \mathbf{y}, \mathbf{z}) \in T_{XYZ}^n$  such that  $D(P_{XYZ}||P_A P_Y W) \leq \eta$ ). We further prune the list of candidates by only keeping the ones which can account for all other candidates that can lead to ambiguity at the decoder. For example, for a candidate  $m_A$ , suppose there are two other candidates  $\tilde{m}_A$  and  $\tilde{m}_B$  of user A and user B respectively. The decoder is confused between  $m_A$  and  $\tilde{m}_A$ , so it cannot reliably choose an output message for user A. Neither can it adjudge one of the users to be adversarial as both users have valid message candidates. In order to get around this, we consider a message candidate  $m_A$  *viable* only if for every pair of candidates  $(\tilde{m}_A, \tilde{m}_B)$ ,  $\tilde{m}_A \neq m_A$ , such that  $(f_A(m_A), \mathbf{y}, f_A(\tilde{m}_A), f_B(\tilde{m}_B), \mathbf{z}) \in T_{XY\tilde{X}\tilde{Y}Z}^n$ , the condition  $I(\tilde{X}\tilde{Y}; XZ|Y) < \eta$  holds. Under this condition, we may infer that the channel output  $\mathbf{z}$  was likely not caused by the pair  $(\tilde{m}_A, \tilde{m}_B)$ . In fact,  $(\tilde{m}_A, \tilde{m}_B)$  is more likely to be part of the attack strategy employed by user B to produce its input vector  $\mathbf{y}$ . Similarly, if there is a pair of candidates  $(\tilde{m}_{B1}, \tilde{m}_{B2})$  of user B, the decoder can neither reliably decode user B's message, nor can it declare either user as adversarial. Then, we require that for every pair of candidates  $(\tilde{m}_{B1}, \tilde{m}_{B2})$  of user B such that  $(f_A(m_A), \mathbf{y}, f_B(\tilde{m}_{B1}), f_B(\tilde{m}_{B2}), \mathbf{z}) \in T_{XY\tilde{Y}_1\tilde{Y}_2Z}^n$ , the condition  $I(\tilde{Y}_1\tilde{Y}_2; XZ|Y) < \eta$  holds. Let  $D_A(\eta, \mathbf{z})$  be the set of all candidates of user A which pass these checks. We define  $D_B(\eta, \mathbf{z})$  analogously by interchanging the roles of users A and B. Finally, the decoder outputs as follows:

$$\phi(\mathbf{z}) \stackrel{\text{def}}{=} \begin{cases} (m_A, m_B) & \text{if } D_A(\eta, \mathbf{z}) \times D_B(\eta, \mathbf{z}) = \{(m_A, m_B)\}, \\ \mathbf{a} \text{ (blame A)} & \text{if } |D_A(\eta, \mathbf{z})| = 0, |D_B(\eta, \mathbf{z})| \neq 0, \\ \mathbf{b} \text{ (blame B)} & \text{if } |D_B(\eta, \mathbf{z})| = 0, |D_A(\eta, \mathbf{z})| \neq 0, \\ (1, 1) & \text{if } |D_A(\eta, \mathbf{z})| = |D_B(\eta, \mathbf{z})| = 0. \end{cases}$$

In the spirit of [8, Lemma 4] and [9, Lemma 1], we show that for a non-spoofable byzantine-MAC, there exists a small enough  $\eta > 0$  such that if  $|D_A(\eta, \mathbf{z})|, |D_B(\eta, \mathbf{z})| > 0$  then  $|D_A(\eta, \mathbf{z})| = |D_B(\eta, \mathbf{z})| = 1$  (Lemma 6). Thus, the decoder definition covers all the cases. We also show that  $|D_A(\eta, \mathbf{z})| = |D_B(\eta, \mathbf{z})| = 0$  is a low probability event. By analyzing the error probability of the decoder we show that for non-spoofable channels it can support positive rates for both users.

#### A. Proof of Theorem 1

We start with a proof of the converse. We prove it for the stronger version when both encoders can privately randomize.

**Lemma 3.** *If a channel is A-spoofable (resp. B-spoofable), then for any  $(N_A, N_B, n)$  adversary identifying code with  $N_A \geq 2$  (resp.  $N_B \geq 2$ ), the probability of error is at least  $1/12$ .*

<sup>2</sup>In fact, we show the result for the stronger case when users are allowed to use privately randomized encoders.

<sup>3</sup>Pereg and Steinberg [26] addressed this when encoders have private randomness, a setting we do not consider in this paper.

*Proof.* The proof uses ideas from the proof of [8, Lemma 1, page 187]. Suppose the channel is A-spoofable such that  $Q_{Y|\tilde{X}\tilde{Y}}$  and  $Q_{X|\tilde{X}X'}$  are conditional distributions satisfying (7). Let  $(F_A, F_B, \phi)$  be a given  $(N_A, N_B, n)$  code where  $F_A : [1 : N_A] \rightarrow \mathcal{X}^n$  and  $F_B : [1 : N_B] \rightarrow \mathcal{Y}^n$  are (privately) randomized maps and  $\phi : \mathcal{Z}^n \rightarrow \mathcal{M}_A \times \mathcal{M}_B \cup \{\mathbf{a}, \mathbf{b}\}$  is a deterministic map. We can define the probability of error for encoders with private randomness in a similar fashion as defined for the deterministic code in (3)-(5). Recall that  $\mathcal{E}_{m_A, m_B} = \{z : \phi(z) \neq (m_A, m_B)\}$ ,  $\mathcal{E}_{m_B}^B = \{z : \phi_B(z) \notin \{m_B, \mathbf{a}\}\}$  and  $\mathcal{E}_{m_A}^A = \{z : \phi_A(z) \notin \{m_A, \mathbf{b}\}\}$ . When both users are honest, we define

$$P_{e, \text{hon}}^{\text{pvt}} \stackrel{\text{def}}{=} \frac{1}{N_A \cdot N_B} \sum_{\substack{(m_A, m_B) \\ \in \mathcal{M}_A \times \mathcal{M}_B}} \sum_{\mathbf{x}, \mathbf{y}} \mathbb{P}(F_A(m_A) = \mathbf{x}) \mathbb{P}(F_B(m_B) = \mathbf{y}) W^n(\mathcal{E}_{m_A, m_B} | \mathbf{x}, \mathbf{y}).$$

The probability in the terms  $\mathbb{P}(F_A(m_A) = \mathbf{x})$  and  $\mathbb{P}(F_B(m_B) = \mathbf{y})$  is over the randomness of the encoders  $F_A$  and  $F_B$  respectively.

When a user is malicious, we define the probabilities of error under randomized attacks for convenience. Along the lines of Remark 1 (which considered a deterministic code), the probability of error is the same under both randomized and deterministic attacks even for privately randomized encoders. When user A is malicious, we define

$$P_{e, \text{mal A}}^{\text{pvt}} \stackrel{\text{def}}{=} \max_{P_{\mathbf{X}}} \sum_{\mathbf{x}} P_{\mathbf{X}}(\mathbf{x}) \left( \frac{1}{N_B} \sum_{m_B \in \mathcal{M}_B} \sum_{\mathbf{y}} \mathbb{P}(F_B(m_B) = \mathbf{y}) W^n(\mathcal{E}_{m_B}^B | \mathbf{x}, \mathbf{y}) \right). \quad (9)$$

Here, the maximization is over all randomized attacks  $P_{\mathbf{X}}$  distributed on  $\mathcal{X}^n$ . The probability in the term  $\mathbb{P}(F_B(m_B) = \mathbf{y})$  is over the randomness of the encoder  $F_B$ . Similarly, the average probability of error when user B is adversarial is

$$P_{e, \text{mal B}}^{\text{pvt}} \stackrel{\text{def}}{=} \max_{P_{\mathbf{Y}}} \sum_{\mathbf{y}} P_{\mathbf{Y}}(\mathbf{y}) \left( \frac{1}{N_A} \sum_{m_A \in \mathcal{M}_A} \sum_{\mathbf{x}} \mathbb{P}(F_A(m_A) = \mathbf{x}) W^n(\mathcal{E}_{m_A}^A | \mathbf{x}, \mathbf{y}) \right). \quad (10)$$

We define the *average probability of error* as

$$P_e^{\text{pvt}}(F_A, F_B, \phi) \stackrel{\text{def}}{=} \max \left\{ P_{e, \text{hon}}^{\text{pvt}}, P_{e, \text{mal A}}^{\text{pvt}}, P_{e, \text{mal B}}^{\text{pvt}} \right\}.$$

For the rest of the proof, we will use the notation  $\mathbf{z} = (z_1, z_2, \dots, z_n)$ ,  $\mathbf{x}' = (x'_1, x'_2, \dots, x'_n)$ ,  $\tilde{\mathbf{x}} = (\tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_n)$  and  $\tilde{\mathbf{y}} = (\tilde{y}_1, \tilde{y}_2, \dots, \tilde{y}_n)$  for  $n$ -length vectors  $\mathbf{z}$ ,  $\mathbf{x}'$ ,  $\tilde{\mathbf{x}}$  and  $\tilde{\mathbf{y}}$ .

Consider the following scenarios for  $i, j \in \mathcal{M}_A$ ,  $k \in \mathcal{M}_B$  and independent encoders  $F_A, F'_A$  and  $F_B$ , where  $F_A$  and  $F'_A$  are two independent copies of user A's encoder:

- (i) User A sends input to the channel according to  $F_A(i)$ . User B uses an independent copy  $F'_A$  of user A's encoder. The input of user B to the channel is produced by passing  $(F'_A(j), F_B(k))$  through  $Q_{Y|\tilde{X}\tilde{Y}}^n$ . For  $\mathbf{z} \in \mathcal{Z}^n$ , the output distribution of the channel (denoted by  $P_{i,j,k}(\mathbf{z})$ ) is given by

$$P_{i,j,k}(\mathbf{z}) \stackrel{\text{def}}{=} \sum_{\mathbf{x}', \tilde{\mathbf{x}}, \tilde{\mathbf{y}}} \mathbb{P}(F_A(i) = \mathbf{x}') \mathbb{P}(F'_A(j) = \tilde{\mathbf{x}}) \mathbb{P}(F_B(k) = \tilde{\mathbf{y}}) \left( \prod_{t=1}^n \sum_{y \in \mathcal{Y}} Q_{Y|\tilde{X}\tilde{Y}}(y | \tilde{x}_t, \tilde{y}_t) W_{Z|XY}(z_t | x'_t, y) \right). \quad (11)$$

- (ii) User B sends input according to  $F_B(k)$ . The input of user A to the channel is produced by passing  $(F_A(i), F'_A(j))$  through  $Q_{X|\tilde{X}X'}^n$ . For  $\mathbf{z} \in \mathcal{Z}^n$ , the output distribution of the channel (denoted by  $Q_{i,j,k}(\mathbf{z})$ ) is given by

$$Q_{i,j,k}(\mathbf{z}) \stackrel{\text{def}}{=} \sum_{\mathbf{x}', \tilde{\mathbf{x}}, \tilde{\mathbf{y}}} \mathbb{P}(F_A(i) = \mathbf{x}') \mathbb{P}(F'_A(j) = \tilde{\mathbf{x}}) \mathbb{P}(F_B(k) = \tilde{\mathbf{y}}) \left( \prod_{t=1}^n \sum_{x \in \mathcal{X}} Q_{X|\tilde{X}X'}(x | x'_t, \tilde{x}_t) W_{Z|XY}(z_t | x, \tilde{y}_t) \right). \quad (12)$$

By (7) (also see Fig. 2), we see that for all  $i, j \in \mathcal{M}_A, k \in \mathcal{M}_B$  and  $\mathbf{z} \in \mathcal{Z}^n$ ,

$$P_{i,j,k}(\mathbf{z}) = P_{j,i,k}(\mathbf{z}) = Q_{i,j,k}(\mathbf{z}). \quad (13)$$

In scenario (i), suppose user A chooses  $i \in \mathcal{M}_A$  uniformly at random (and is hence honest) and, independently, the adversarial user B chooses  $(j, k) \in \mathcal{M}_A \times \mathcal{M}_B$  uniformly at random. Then, from (10) and (11), we see that

$$P_{e, \text{mal B}}^{\text{pvt}} \geq \frac{1}{N_A^2 \times N_B} \sum_{i,j \in \mathcal{M}_A} \sum_{k \in \mathcal{M}_B} \sum_{\mathbf{z}: \phi_A(\mathbf{z}) \notin \{i, \mathbf{b}\}} P_{i,j,k}(\mathbf{z}).$$

Interchanging the roles of  $i$  and  $j$ , we have

$$P_{e, \text{mal A}}^{\text{pvt}} \geq \frac{1}{N_A^2 \times N_B} \sum_{i,j \in \mathcal{M}_A} \sum_{k \in \mathcal{M}_B} \sum_{\mathbf{z}: \phi_A(\mathbf{z}) \notin \{j, \mathbf{b}\}} P_{j,i,k}(\mathbf{z}).$$

In scenario (ii), suppose user B chooses  $k \in \mathcal{M}_B$  uniformly at random (and hence is honest), while, independently, the adversarial user A chooses  $(i, j) \in \mathcal{M}_A^2$  uniformly at random. Using (9) and (12), we obtain

$$P_{e, \text{mal A}}^{\text{pvt}} \geq \frac{1}{N_A^2 \times N_B} \sum_{i, j \in \mathcal{M}_A} \sum_{k \in \mathcal{M}_B} \sum_{\mathbf{z}: \phi_B(\mathbf{z}) \notin \{k, \mathbf{a}\}} Q_{i, j, k}(\mathbf{z}).$$

Thus,

$$\begin{aligned} 3P_e^{\text{pvt}}(F_A, F_B, \phi) &\geq P_{e, \text{mal B}}^{\text{pvt}} + P_{e, \text{mal B}}^{\text{pvt}} + P_{e, \text{mal A}}^{\text{pvt}} \\ &\geq \frac{1}{N_A^2 \times N_B} \sum_{i, j \in \mathcal{M}_A} \sum_{k \in \mathcal{M}_B} \left( \sum_{\mathbf{z}: \phi_A(\mathbf{z}) \notin \{i, \mathbf{b}\}} P_{i, j, k}(\mathbf{z}) + \sum_{\mathbf{z}: \phi_A(\mathbf{z}) \notin \{j, \mathbf{b}\}} P_{j, i, k}(\mathbf{z}) + \sum_{\mathbf{z}: \phi_B(\mathbf{z}) \notin \{k, \mathbf{a}\}} Q_{i, j, k}(\mathbf{z}) \right) \\ &\stackrel{(a)}{=} \frac{1}{N_A^2 \times N_B} \sum_{i, j \in \mathcal{M}_A} \sum_{k \in \mathcal{M}_B} \left( \sum_{\mathbf{z}: \phi_A(\mathbf{z}) \notin \{i, \mathbf{b}\}} P_{i, j, k}(\mathbf{z}) + \sum_{\mathbf{z}: \phi_A(\mathbf{z}) \notin \{j, \mathbf{b}\}} P_{i, j, k}(\mathbf{z}) + \sum_{\mathbf{z}: \phi_B(\mathbf{z}) \notin \{k, \mathbf{a}\}} P_{i, j, k}(\mathbf{z}) \right) \\ &\stackrel{(b)}{\geq} \frac{1}{N_A^2 \times N_B} \sum_{i, j \in \mathcal{M}_A, i \neq j} \sum_{k \in \mathcal{M}_B} \left( \sum_{\mathbf{z} \in \mathcal{Z}^n} P_{i, j, k}(\mathbf{z}) \right) \\ &= \frac{N_A(N_A - 1)N_B}{2N_A^2 \times N_B} \\ &= \frac{N_A - 1}{2N_A} \\ &\geq \frac{1}{4} \quad \text{for } N_A \geq 2, \end{aligned}$$

where (a) follows by noting from (13) that  $P_{i, j, k}(\mathbf{z}) = P_{j, i, k}(\mathbf{z}) = Q_{i, j, k}(\mathbf{z})$  and (b) follows by noting that  $\{\mathbf{z}: \phi_A(\mathbf{z}) \notin \{i, \mathbf{b}\}\} \cup \{\mathbf{z}: \phi_A(\mathbf{z}) \notin \{j, \mathbf{b}\}\} \cup \{\mathbf{z}: \phi_B(\mathbf{z}) \notin \{k, \mathbf{a}\}\} = \{\mathbf{z}: \phi_A(\mathbf{z}) \neq \mathbf{b}\} \cup \{\mathbf{z}: \phi_B(\mathbf{z}) \notin \{k, \mathbf{a}\}\} \supseteq \{\mathbf{z}: \phi_A(\mathbf{z}) \neq \mathbf{b}\} \cup \{\mathbf{z}: \phi_B(\mathbf{z}) = \mathbf{b}\} = \{\mathbf{z}: \phi_A(\mathbf{z}) \neq \mathbf{b}\} \cup \{\mathbf{z}: \phi_A(\mathbf{z}) = \mathbf{b}\} = \mathcal{Z}^n$ . Here, the second last equality follows by recalling from (1) and (2) that if  $\phi_B(\mathbf{z}) = \mathbf{b}$ , then  $\phi_A(\mathbf{z}) = \mathbf{b}$ . Thus, for any given code  $(F_A, F_B, \phi)$ , for an A-spoofable channel,  $P_e^{\text{pvt}}(F_A, F_B, \phi) \geq \frac{1}{12}$ . A similar analysis follows when the channel is B-spoofable.  $\square$

Next, for the proof of achievability of Theorem 1, we first state a *codebook lemma* which will be used to show all our achievability results. This gives a randomly generated codebook which satisfies certain properties. The technical proof of the lemma, which is along the lines of that of [8, Lemma 3], is in Appendix A. The lemma can be thought of as a generalization of [8, Lemma 3] for two users. In particular, (14) is similar to [8, Lemma 3, (3.2)], (15) and (17) are generalization of [8, Lemma 3, (3.3)] and (16) and (18) are generalization of [8, Lemma 3, (3.1)] for a pair of messages.

**Lemma 4** (Codebook lemma). *Suppose  $\mathcal{X}, \mathcal{Y}, \mathcal{Z}$  are finite. Let  $P_A \in \mathcal{P}_{\mathcal{X}}^n$  and  $P_B \in \mathcal{P}_{\mathcal{Y}}^n$ . For any  $\epsilon > 0$ , there exists  $n_0(\epsilon)$  such that for all  $n \geq n_0(\epsilon)$ ,  $N_A, N_B \geq \exp(n\epsilon)$ , there are codewords  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{N_A}$  of type  $P_A$  and  $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_{N_B}$  of type  $P_B$  such that for all  $\mathbf{x}, \mathbf{x}' \in \mathcal{X}^n$  and  $\mathbf{y}, \mathbf{y}' \in \mathcal{Y}^n$ , and joint types  $P_{X\tilde{X}\tilde{Y}Y} \in \mathcal{P}_{\mathcal{X} \times \mathcal{X} \times \mathcal{Y} \times \mathcal{Y}}^n$  and  $P_{X'\tilde{Y}_1\tilde{Y}_2Y'} \in \mathcal{P}_{\mathcal{X} \times \mathcal{Y} \times \mathcal{Y} \times \mathcal{Y}}^n$ , and for  $R_A \stackrel{\text{def}}{=} (1/n) \log N_A$  and  $R_B \stackrel{\text{def}}{=} (1/n) \log N_B$ , the following holds:<sup>4</sup>*

$$\frac{|\{m_A : (\mathbf{x}_{m_A}, \mathbf{y}) \in T_{XY}^n\}|}{N_A} \leq \exp\{-n\epsilon/2\}, \quad \text{if } I(X; Y) > \epsilon; \quad (14)$$

$$\frac{|\{m_A : (\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{m_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n \text{ for some } \tilde{m}_A \neq m_A \text{ and some } m_B\}|}{N_A} \leq \exp\{-n\epsilon/2\},$$

$$\text{if } I(X; \tilde{X}\tilde{Y}Y) - |R_A - I(\tilde{X}; \tilde{Y}Y)|^+ - |R_B - I(\tilde{Y}; Y)|^+ > \epsilon; \quad (15)$$

$$\frac{|\{(\tilde{m}_A, \tilde{m}_B) : (\mathbf{x}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n\}|}{N_A} \leq \exp\left\{n \left( |R_A - I(\tilde{X}; \tilde{Y}XY)|^+ + |R_B - I(\tilde{Y}; XY)|^+ + \epsilon \right)\right\}; \quad (16)$$

$$\frac{|\{m_A : (\mathbf{x}_{m_A}, \mathbf{y}_{\tilde{m}_{B1}}, \mathbf{y}_{\tilde{m}_{B2}}, \mathbf{y}') \in T_{X'\tilde{Y}_1\tilde{Y}_2Y'}^n \text{ for some } \tilde{m}_{B1}, \tilde{m}_{B2}\}|}{N_A} \leq \exp\{-n\epsilon/2\},$$

$$\text{if } I(X'; \tilde{Y}_1\tilde{Y}_2Y') - |R_B - I(\tilde{Y}_1; Y')|^+ - |R_B - I(\tilde{Y}_2; \tilde{Y}_1Y')|^+ > \epsilon; \quad (17)$$

$$\text{and } \frac{|\{(\tilde{m}_{B1}, \tilde{m}_{B2}) : (\mathbf{x}', \mathbf{y}_{\tilde{m}_{B1}}, \mathbf{y}_{\tilde{m}_{B2}}, \mathbf{y}') \in T_{X'\tilde{Y}_1\tilde{Y}_2Y'}^n\}|}{N_A} \leq \exp\left\{n \left( |R_B - I(\tilde{Y}_1; X'Y')|^+ + |R_B - I(\tilde{Y}_2; \tilde{Y}_1X'Y')|^+ + \epsilon \right)\right\}, \quad (18)$$

<sup>4</sup>Note that  $\exp$  and  $\log$  are with respect to base 2.



and statements analogous to (14)-(18) with the roles of users A and B are interchanged.<sup>5</sup>

With this, we are ready to prove the achievability of Theorem 1.

**Lemma 5.** *The rate region for deterministic codes is non-empty if the channel is non-spoofable.*

*Proof.* For some  $P_A$  and  $P_B$  satisfying  $\min_{x \in \mathcal{X}} P_A(x) \geq \alpha$  and  $\min_{y \in \mathcal{Y}} P_B(y) \geq \alpha$  respectively for some  $\alpha > 0$ ,  $\epsilon > 0$  to-be-defined (TBD),  $n > n_0(\epsilon)$ ,  $\delta > \epsilon$  (TBD) and  $R_A = R_B = \delta$ , the codebook is given by Lemma 4.

*Encoding.* Let  $N_A = 2^{nR_A}$ ,  $N_B = 2^{nR_B}$ ,  $\mathcal{M}_A = \{1, \dots, N_A\}$  and  $\mathcal{M}_B = \{1, \dots, N_B\}$ . For  $m_A \in \mathcal{M}_A$ ,  $f_A(m_A) = \mathbf{x}_{m_A}$  and for  $m_B \in \mathcal{M}_B$ ,  $f_B(m_B) = \mathbf{y}_{m_B}$ .

*Decoding.* For a parameter  $\eta > 0$  (TBD), let  $\mathcal{D}_\eta$  be the set of joint distributions defined as

$$D_\eta \stackrel{\text{def}}{=} \{P_{XYZ} \in \mathcal{P}_{\mathcal{X} \times \mathcal{Y} \times \mathcal{Z}}^n : D(P_{XYZ} \| P_X P_Y W) \leq \eta\}.$$

**Definition 4** ( $D_A(\eta, \mathbf{z})$ ). *For the given codebook, the parameter  $\eta > 0$  and the received channel output sequence  $\mathbf{z}$ , let  $D_A(\eta, \mathbf{z})$  be defined as the set of messages  $m_A \in \mathcal{M}_A$  such that for each  $m_A \in D_A(\eta, \mathbf{z})$ , there exists  $\mathbf{y} \in \mathcal{Y}^n$  satisfying the following conditions:*

- (i)  $(f_A(m_A), \mathbf{y}, \mathbf{z}) \in T_{XYZ}^n$  for some  $P_{XYZ} \in \mathcal{D}_\eta$ .
- (ii) For every  $(\tilde{m}_A, \tilde{m}_B) \in \mathcal{M}_A \times \mathcal{M}_B$ ,  $\tilde{m}_A \neq m_A$  and  $(\mathbf{y}', \mathbf{x}') \in \mathcal{Y}^n \times \mathcal{X}^n$  such that  $(f_A(m_A), \mathbf{y}, f_A(\tilde{m}_A), \mathbf{y}', \mathbf{x}', f_B(\tilde{m}_B), \mathbf{z}) \in T_{XY\tilde{X}Y'X'\tilde{Y}Z}^n$  with  $P_{\tilde{X}Y'Z} \in \mathcal{D}_\eta$  and  $P_{X'\tilde{Y}Z} \in \mathcal{D}_\eta$ , we have  $I(\tilde{X}\tilde{Y}; XZ|Y) < \eta$ .
- (iii) For every  $\tilde{m}_{B1}, \tilde{m}_{B2} \in \mathcal{M}_B$  where  $\tilde{m}_{B1} \neq \tilde{m}_{B2}$  and  $\mathbf{x}'_1, \mathbf{x}'_2 \in \mathcal{X}^n$  such that  $(f_A(m_A), \mathbf{y}, \mathbf{x}'_1, f_B(\tilde{m}_{B1}), \mathbf{x}'_2, f_B(\tilde{m}_{B2}), \mathbf{z}) \in T_{XYX'_1\tilde{Y}_1X'_2\tilde{Y}_2Z}^n$  with  $P_{X'_1\tilde{Y}_1Z} \in \mathcal{D}_\eta$  and  $P_{X'_2\tilde{Y}_2Z} \in \mathcal{D}_\eta$ , we have  $I(\tilde{Y}_1\tilde{Y}_2; XZ|Y) < \eta$ .

We define  $D_B(\eta, \mathbf{z})$  analogously (by interchanging the roles of user A and B). The output of the decoder (for parameter  $\eta > 0$ ) is as follows:

$$\phi(\mathbf{z}) \stackrel{\text{def}}{=} \begin{cases} (m_A, m_B), & \text{if } D_A(\eta, \mathbf{z}) \times D_B(\eta, \mathbf{z}) = \{(m_A, m_B)\} \\ \mathbf{a}, & \text{if } |D_A(\eta, \mathbf{z})| = 0, |D_B(\eta, \mathbf{z})| \neq 0 \\ \mathbf{b}, & \text{if } |D_B(\eta, \mathbf{z})| = 0, |D_A(\eta, \mathbf{z})| \neq 0 \\ (1, 1) & \text{otherwise.} \end{cases}$$

The last of the above cases (i.e.  $\phi(\mathbf{z}) = (1, 1)$ ) occurs when either of the following two events occur: (i)  $|D_B(\eta, \mathbf{z})| = |D_A(\eta, \mathbf{z})| = 0$ , (ii)  $|D_B(\eta, \mathbf{z})| \geq 1$ ,  $|D_A(\eta, \mathbf{z})| \geq 1$  and  $|D_B(\eta, \mathbf{z})| + |D_A(\eta, \mathbf{z})| \geq 3$ , that is, both  $D_B(\eta, \mathbf{z})$  and  $D_A(\eta, \mathbf{z})$  are non-empty and at least one of two sets has two or more elements. As we will see, the first event is an atypical event and hence will occur with a vanishing probability. The following lemma (proved in Appendix B) implies that the second event cannot occur for non-spoofable channels if  $\eta > 0$  is chosen sufficiently small.

**Lemma 6.** *Suppose  $\alpha > 0$ . For a channel which is not A-spoofable, for sufficiently small  $\eta > 0$ , there does not exist a distribution  $P_{XY\tilde{X}Y'X'\tilde{Y}Z} \in \mathcal{P}_{XY\tilde{X}Y'X'\tilde{Y}Z}^n$  with  $\min_x P_X(x), \min_{\tilde{x}} P_{\tilde{X}}(\tilde{x}), \min_{\tilde{y}} P_{\tilde{Y}}(\tilde{y}) \geq \alpha$  which satisfies the following:*

- (A)  $P_{XYZ} \in \mathcal{D}_\eta$ ,
- (B)  $P_{\tilde{X}Y'Z} \in \mathcal{D}_\eta$ ,
- (C)  $P_{X'\tilde{Y}Z} \in \mathcal{D}_\eta$ ,
- (D)  $I(\tilde{X}\tilde{Y}; XZ|Y) < \eta$ ,
- (E)  $I(X\tilde{Y}; \tilde{X}Z|Y') < \eta$  and
- (F)  $I(X\tilde{X}; \tilde{Y}Z|X') < \eta$ .

*Analogous condition holds for a channel which is not B-spoofable.*

For a non-spoofable channel, given  $\alpha > 0$  and small enough  $\eta > 0$ , Lemma 6 implies that if  $|D_A(\eta, \mathbf{z})|, |D_B(\eta, \mathbf{z})| \geq 1$ , then  $|D_B(\eta, \mathbf{z})| = |D_A(\eta, \mathbf{z})| = 1$ . To see this, suppose  $|D_A(\eta, \mathbf{z})| \geq 2$  and  $|D_B(\eta, \mathbf{z})| \geq 1$ . Let  $m_A, \tilde{m}_A \in D_A(\eta, \mathbf{z})$  and  $\tilde{m}_B \in D_B(\eta, \mathbf{z})$ . Then, Definition 4 implies that there exist  $\mathbf{x}, \mathbf{y}$  and  $\mathbf{y}'$  such that for  $(f_A(m_A), \mathbf{y}, f_A(\tilde{m}_A), \mathbf{y}', \mathbf{x}', f_B(\tilde{m}_B), \mathbf{z}) \in T_{XY\tilde{X}Y'X'\tilde{Y}Z}^n$ ,  $P_{XYZ} \in \mathcal{D}_\eta$ ,  $P_{\tilde{X}Y'Z} \in \mathcal{D}_\eta$ ,  $P_{X'\tilde{Y}Z} \in \mathcal{D}_\eta$ ,  $I(\tilde{X}\tilde{Y}; XZ|Y) < \eta$ ,  $I(X\tilde{Y}; \tilde{X}Z|Y') < \eta$  and  $I(X\tilde{X}; \tilde{Y}Z|X') < \eta$ . However, such joint distribution  $P_{XY\tilde{X}Y'X'\tilde{Y}Z}$  cannot exist as per Lemma 6. Analogously, for a channel which is not B-spoofable,  $|D_A(\eta, \mathbf{z})| \geq 1$  and  $|D_B(\eta, \mathbf{z})| \geq 2$  is not possible. We choose  $\eta, \epsilon$  and  $\delta (> \epsilon)$  small enough so that Lemma 6 holds and

$$\eta > 3\epsilon + 4\delta. \tag{19}$$

<sup>5</sup>See Appendix A for the full statement.

To analyze the probability of error, we first recall from (6) that  $P_{e,\text{hon}} \leq P_{e,\text{mal A}} + P_{e,\text{mal B}}$ . So, we only need to analyse the case when one of the users is malicious. We will use the ‘method of types’ for the analysis [27], [28]. Here, we restate some basic properties from [28, Chapter 2] that we need for the proof. Let  $X$  and  $Y$  be two jointly distributed random variables such that  $P_{XY} \in \mathcal{P}_{\mathcal{X} \times \mathcal{Y}}^n$ . For  $\mathbf{x} \in T_{\mathcal{X}}^n$ , a distribution  $Q$  on  $\mathcal{X}$  and a discrete memoryless channel  $U_{Y|X}$  from  $\mathcal{X}$  to  $\mathcal{Y}$ ,

$$|\mathcal{P}_{\mathcal{X}}^n| \leq (n+1)^{|\mathcal{X}|}, \quad (20)$$

$$(n+1)^{-|\mathcal{X}|} \exp(nH(X)) \leq |T_{\mathcal{X}}^n| \leq \exp(nH(X)), \quad (21)$$

$$(n+1)^{-|\mathcal{X}||\mathcal{Y}|} \exp(nH(Y|X)) \leq |T_{Y|X}^n(\mathbf{x})| \leq \exp(nH(Y|X)), \quad (22)$$

$$(n+1)^{-|\mathcal{X}|} \exp\{-nD(P_X||Q)\} \leq \sum_{\mathbf{x}' \in T_{\mathcal{X}}^n} Q^n(\mathbf{x}') \leq \exp\{-nD(P_X||Q)\} \text{ and} \\ \sum_{\mathbf{y} \in T_{Y|X}^n(\mathbf{x})} U_{Y|X}^n(\mathbf{y}|\mathbf{x}) \leq \exp\{-nD(P_{XY}||P_X U_{Y|X})\}. \quad (23)$$

We consider the case when user B is malicious. We will analyse  $P_{e,\text{mal B}}$ . Suppose a malicious user B sends  $\mathbf{y}$ . Let  $P_{e,\text{mal B}}(\mathbf{y})$  denote the probability of error when user B is malicious and sends  $\mathbf{y}$ . That is, for  $\mathcal{E}_{m_A}^A \stackrel{\text{def}}{=} \{\mathbf{z} : \phi_A(\mathbf{z}) \notin \{m_A, \mathbf{b}\}\}$ ,

$$P_{e,\text{mal B}}(\mathbf{y}) = \frac{1}{N_A} \sum_{m_A \in \mathcal{M}_A} W^n(\mathcal{E}_{m_A}^A | f_A(m_A), \mathbf{y})$$

and

$$P_{e,\text{mal B}} = \max_{\mathbf{y}} P_{e,\text{mal B}}(\mathbf{y}).$$

We will show that  $P_{e,\text{mal B}}(\mathbf{y})$  is small for each  $\mathbf{y} \in \mathcal{Y}^n$ . The analysis follows the flowchart given in Figure 4. We define the following sets.

$$\mathcal{H}_1 \stackrel{\text{def}}{=} \{m_A : (\mathbf{x}_{m_A}, \mathbf{y}) \in T_{XY}^n \text{ such that } I(X; Y) > \epsilon\}, \text{ and} \\ \mathcal{H}_2 \stackrel{\text{def}}{=} \{m_A : (\mathbf{x}_{m_A}, \mathbf{y}) \in T_{XY}^n \text{ such that } I(X; Y) \leq \epsilon\}$$

$$P_{e,\text{mal B}}(\mathbf{y}) \leq \frac{1}{N_A} |\mathcal{H}_1| + \frac{1}{N_A} \sum_{m_A \in \mathcal{H}_2} \left( \sum_{P_{XYZ} \in \mathcal{D}_{\eta}^c} \sum_{\mathbf{z} \in T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y})} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) \right) \\ + \frac{1}{N_A} \sum_{m_A \in \mathcal{H}_2} \left( \sum_{P_{XYZ} \in \mathcal{D}_{\eta}} \sum_{\substack{\mathbf{z} \in T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y}) \\ : \phi_A(\mathbf{z}) \notin \{m_A, \mathbf{b}\}}} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) \right) \\ =: P_1(\mathbf{y}) + P_2(\mathbf{y}) + P_3(\mathbf{y}). \quad (24)$$

The first term  $P_1(\mathbf{y}) = \frac{1}{N_A} |\mathcal{H}_1|$  is upper bounded by

$$|\mathcal{P}_{\mathcal{X} \times \mathcal{Y}}^n| \times \frac{|\{m_A : (\mathbf{x}_{m_A}, \mathbf{y}) \in T_{XY}^n, I(X; Y) > \epsilon\}|}{N_A}$$

which goes to zero as  $n \rightarrow \infty$  by (14) and noting that there are only polynomially many types. We now analyse the second term


$$P_2(\mathbf{y}) = \frac{1}{N_A} \sum_{m_A \in \mathcal{H}_2} \left( \sum_{P_{XYZ} \in \mathcal{D}_{\eta}^c} \sum_{\mathbf{z} \in T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y})} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) \right).$$

For any  $m_A \in \mathcal{H}_2$ ,

$$\sum_{P_{XYZ} \in \mathcal{D}_{\eta}^c} \sum_{\mathbf{z} \in T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y})} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) \stackrel{(a)}{\leq} |\mathcal{D}_{\eta}^c| \exp(-nD(P_{XYZ}||P_{XY}W)) \\ = |\mathcal{D}_{\eta}^c| \exp(-n(D(P_{XYZ}||P_X P_Y W) - I(X; Y))) \\ \leq |\mathcal{D}_{\eta}^c| \exp(-n(\eta - \epsilon)) \rightarrow 0 \text{ as } \eta > \epsilon \text{ and } |\mathcal{D}_{\eta}^c| \leq (n+1)^{|\mathcal{X}||\mathcal{Y}||\mathcal{Z}|},$$

where (a) follows from (23). We are left to analyse the last term

$$P_3(\mathbf{y}) = \frac{1}{N_A} \sum_{m_A \in \mathcal{H}_2} \left( \sum_{P_{XYZ} \in \mathcal{D}_\eta} \sum_{\substack{\mathbf{z} \in T_{ZY|XY}^n(\mathbf{x}_{m_A}, \mathbf{y}) \\ : \phi_A(\mathbf{z}) \notin \{m_A, \mathbf{b}\}}} W^n(\mathbf{z} | \mathbf{x}_{m_A}, \mathbf{y}) \right). \quad (25)$$



$P_{e, mal B}(\mathbf{y})$	the average probability of error when malicious user B sends $\mathbf{y}$
$P_1(\mathbf{y})$	the average probability that channel inputs are atypical
$P_2(\mathbf{y})$	the average probability that the channel output is atypical
$P_3(\mathbf{y})$	the average probability of error when channel inputs and output are typical
$P_{E_{m_A,1}}(\mathbf{y})$	condition (ii) in Definition 4 does not hold
$P_{E_{m_A,2}}(\mathbf{y})$	condition (iii) in Definition 4 does not hold

Fig. 4. Flowchart depicting the flow of analysis of  $P_{e, mal B}(\mathbf{y})$ , the average probability of error when user B is malicious and sends  $\mathbf{y}$ .

Recall that because of Lemma 6, whenever  $|D_A(\eta, \mathbf{z})|, |D_B(\eta, \mathbf{z})| > 0$ , we have  $|D_A(\eta, \mathbf{z})| = |D_B(\eta, \mathbf{z})| = 1$ . This implies that for  $(\mathbf{x}_{m_A}, \mathbf{y}, \mathbf{z}) \in P_{XYZ}$  such that  $P_{XYZ} \in \mathcal{D}_\eta$  and  $m_A \in \mathcal{H}_2$ , the output of  $\phi_A(\mathbf{z})$  is not in the set  $\{m_A, \mathbf{b}\}$  only if one of the following happens:

- $|D_A(\eta, \mathbf{z})| = |D_B(\eta, \mathbf{z})| = 1$ , but  $m_A \notin D_A(\eta, \mathbf{z})$ .
- $|D_A(\eta, \mathbf{z})| = 0$ .

To formalize this, we define the following sets for  $m_A \in \mathcal{M}_A$ .

$$\begin{aligned} \mathcal{G}_{m_A} &= \{\mathbf{z} : (\mathbf{x}_{m_A}, \mathbf{y}, \mathbf{z}) \in P_{XYZ}, P_{XYZ} \in \mathcal{D}_\eta, I(X; Y) \leq \epsilon\}, \\ \mathcal{G}_{m_A,0} &= \mathcal{G}_{m_A} \cap \{\mathbf{z} : \phi_A(\mathbf{z}) \notin \{m_A, \mathbf{b}\}\}, \\ \mathcal{G}_{m_A,1} &= \mathcal{G}_{m_A} \cap \{\mathbf{z} : |D_A(\eta, \mathbf{z})| = |D_B(\eta, \mathbf{z})| = 1, m_A \notin D_A(\eta, \mathbf{z})\}, \\ \mathcal{G}_{m_A,2} &= \mathcal{G}_{m_A} \cap \{\mathbf{z} : |D_A(\eta, \mathbf{z})| = 0\}, \text{ and} \\ \mathcal{G}_{m_A,3} &= \mathcal{G}_{m_A} \cap \{\mathbf{z} : m_A \notin D_A(\eta, \mathbf{z})\}. \end{aligned}$$

We are interested in  $\mathcal{G}_{m_A,0}$ . Note that  $\mathcal{G}_{m_A,0} \subseteq \mathcal{G}_{m_A,1} \cup \mathcal{G}_{m_A,2} \subseteq \mathcal{G}_{m_A,3}$ . So, it suffices to upper bound the probability of  $\mathcal{G}_{m_A,3}$  when  $\mathbf{x}_{m_A}$  is sent by user A and  $\mathbf{y}$  by user B. From the definition of  $D_A(\eta, \mathbf{z})$ , we see that  $\mathcal{G}_{m_A,3}$  is the set of  $\mathbf{z} \in \mathcal{Z}^n$  which satisfy decoding condition (i) (this is because  $\mathbf{z} \in \mathcal{G}_{m_A,3}$  implies  $\mathbf{z} \in \mathcal{G}_{m_A}$  since  $\mathcal{G}_{m_A,3} \subseteq \mathcal{G}_{m_A}$ ) but do not satisfy either decoding condition (ii) or decoding condition (iii). We capture this by defining the following sets of distributions:

$$\begin{aligned} \mathcal{P}_1 &= \{P_{\tilde{X}\tilde{Y}\tilde{Z}} \in \mathcal{P}_{\tilde{X} \times \tilde{X} \times \tilde{Y} \times \tilde{Y} \times \tilde{Z}}^n : P_{XYZ} \in \mathcal{D}_\eta, I(X; Y) \leq \epsilon, P_{\tilde{X}'\tilde{Y}'\tilde{Z}} \in \mathcal{D}_\eta \text{ for some } P_{Y'|\tilde{X}Z}, \\ &P_{X'\tilde{Y}Z} \in \mathcal{D}_\eta \text{ for some } P_{X'|\tilde{Y}Z}, P_X = P_{\tilde{X}} = P_A, P_{\tilde{Y}} = P_B \text{ and } I(\tilde{X}\tilde{Y}; XZ|Y) \geq \eta\} \end{aligned} \quad (26)$$

$$\begin{aligned} \mathcal{P}_2 = \{ & P_{X\tilde{Y}_1\tilde{Y}_2YZ} \in \mathcal{P}_{X \times Y \times Y \times Y \times Z}^n : P_{XYZ} \in \mathcal{D}_\eta, I(X; Y) \leq \epsilon, P_{X'_1\tilde{Y}_1Z} \in \mathcal{D}_\eta \text{ for some } P_{X'_1|\tilde{Y}_1Z}, \\ & P_{X'_2\tilde{Y}_2Z} \in \mathcal{D}_\eta \text{ for some } P_{X'_2|\tilde{Y}_2Z}, P_X = P_A, P_{\tilde{Y}_1} = P_{\tilde{Y}_2} = P_B \text{ and } I(\tilde{Y}_1\tilde{Y}_2; XZ|Y) \geq \eta\}. \end{aligned} \quad (27)$$

For  $P_{X\tilde{X}Y\tilde{Y}Z} \in \mathcal{P}_1$  and  $P_{X\tilde{Y}_1\tilde{Y}_2YZ} \in \mathcal{P}_2$ , let

$$\begin{aligned} \mathcal{E}_{m_A,1}(P_{X\tilde{X}Y\tilde{Y}Z}) &= \{ \mathbf{z} : \exists (\tilde{m}_A, \tilde{m}_B) \in \mathcal{M}_A \times \mathcal{M}_B, \tilde{m}_A \neq m_A, (\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}, \mathbf{y}_{\tilde{m}_B}, \mathbf{z}) \in T_{X\tilde{X}Y\tilde{Y}Z}^n \} \text{ and} \\ \mathcal{E}_{m_A,2}(P_{X\tilde{Y}_1\tilde{Y}_2YZ}) &= \{ \mathbf{z} : \exists \tilde{m}_{B1}, \tilde{m}_{B2} \in \mathcal{M}_B, \tilde{m}_{B1} \neq \tilde{m}_{B2}, (\mathbf{x}_{m_A}, \mathbf{y}_{\tilde{m}_{B1}}, \mathbf{y}_{\tilde{m}_{B2}}, \mathbf{y}, \mathbf{z}) \in T_{X\tilde{Y}_1\tilde{Y}_2YZ}^n \}. \end{aligned}$$

Note that  $\mathcal{G}_{m_A,3} = (\cup_{P_{X\tilde{X}Y\tilde{Y}Z} \in \mathcal{P}_1} \mathcal{E}_{m_A,1}(P_{X\tilde{X}Y\tilde{Y}Z})) \cup (\cup_{P_{X\tilde{Y}_1\tilde{Y}_2YZ} \in \mathcal{P}_2} \mathcal{E}_{m_A,2}(P_{X\tilde{Y}_1\tilde{Y}_2YZ}))$ .

Thus, (25) can be analyzed as below.

$$\begin{aligned} P_3(\mathbf{y}) &= \frac{1}{N_A} \sum_{m_A \in \mathcal{H}_2} \left( \sum_{P_{XYZ} \in \mathcal{D}_\eta} \sum_{\substack{\mathbf{z} \in T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y}) \\ \phi_A(\mathbf{z}) \notin \{m_A, \mathbf{b}\}}} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) \right) \\ &\leq \frac{1}{N_A} \sum_{m_A \in \mathcal{H}_2} \sum_{P_{X\tilde{X}Y\tilde{Y}Z} \in \mathcal{P}_1} W^n(\mathcal{E}_{m_A,1}(P_{X\tilde{X}Y\tilde{Y}Z})|\mathbf{x}_{m_A}, \mathbf{y}) \\ &\quad + \frac{1}{N_A} \sum_{m_A \in \mathcal{H}_2} \sum_{P_{X\tilde{Y}_1\tilde{Y}_2YZ} \in \mathcal{P}_2} W^n(\mathcal{E}_{m_A,2}(P_{X\tilde{Y}_1\tilde{Y}_2YZ})|\mathbf{x}_{m_A}, \mathbf{y}) \\ &=: P_{\mathcal{E}_{m_A,1}}(\mathbf{y}) + P_{\mathcal{E}_{m_A,2}}(\mathbf{y}). \end{aligned} \quad (28)$$

We see that  $|\mathcal{P}_1|$  and  $|\mathcal{P}_2|$  increase at most polynomially in  $n$  (see (20)) and clearly  $|\mathcal{H}_2| \leq N_A$ . So, it will suffice to uniformly upper bound  $W^n(\mathcal{E}_{m_A,1}(P_{X\tilde{X}Y\tilde{Y}Z})|\mathbf{x}_{m_A}, \mathbf{y})$  and  $W^n(\mathcal{E}_{m_A,2}(P_{X\tilde{Y}_1\tilde{Y}_2YZ})|\mathbf{x}_{m_A}, \mathbf{y})$  by a term exponentially decreasing in  $n$  for all  $P_{X\tilde{X}Y\tilde{Y}Z} \in \mathcal{P}_1$  and  $P_{X\tilde{Y}_1\tilde{Y}_2YZ} \in \mathcal{P}_2$  respectively. We start the analysis of  $P_{\mathcal{E}_{m_A,1}}(\mathbf{y})$  by upper bounding  $W^n(\mathcal{E}_{m_A,1}(P_{X\tilde{X}Y\tilde{Y}Z})|\mathbf{x}_{m_A}, \mathbf{y})$ . By using (15), we see that for  $P_{X\tilde{X}Y\tilde{Y}Z} \in \mathcal{P}_1$  such that

$$I(X; \tilde{X}\tilde{Y}Y) > |R_A - I(\tilde{X}; \tilde{Y}Y)|^+ + |R_B - I(\tilde{Y}; Y)|^+ + \epsilon,$$

we have,

$$\frac{\left| \left\{ m_A : (\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{m_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n \text{ for some } \tilde{m}_A \neq m_A \text{ and some } m_B \right\} \right|}{N_A} \leq \exp\{-n\epsilon/2\}.$$

So, for all such  $P_{X\tilde{X}Y\tilde{Y}Z}$ ,

$$\begin{aligned} & \frac{1}{N_A} \sum_{m_A \in \mathcal{H}_2} W^n(\mathcal{E}_{m_A,1}(P_{X\tilde{X}Y\tilde{Y}Z})|\mathbf{x}_{m_A}, \mathbf{y}) \\ &= \frac{1}{N_A} \sum_{\substack{m_A : (\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{m_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n \\ \tilde{m}_A \in \mathcal{M}_A, \tilde{m}_A \neq m_A, m_B \in \mathcal{M}_B}} \sum_{\mathbf{z} \in T_{Z|X\tilde{X}\tilde{Y}Y}^n(\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}, \mathbf{y}_{m_B})} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) \\ &\leq \exp\{-n\epsilon/2\}. \end{aligned}$$

Thus, it is sufficient to consider distributions  $P_{X\tilde{X}Y\tilde{Y}Z} \in \mathcal{P}_1$  for which

$$I(X; \tilde{X}\tilde{Y}Y) \leq |R_A - I(\tilde{X}; \tilde{Y}Y)|^+ + |R_B - I(\tilde{Y}; Y)|^+ + \epsilon. \quad (29)$$

For  $P_{X\tilde{X}Y\tilde{Y}Z} \in \mathcal{P}_1$  satisfying (29),

$$\begin{aligned} & \sum_{\mathbf{z} \in \mathcal{E}_{m_A,1}(P_{X\tilde{X}Y\tilde{Y}Z})} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) \\ &= \sum_{\substack{\tilde{m}_A, \tilde{m}_B : \tilde{m}_A \neq m_A, \\ (\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n}} \sum_{\mathbf{z} \in T_{Z|X\tilde{X}\tilde{Y}Y}^n(\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{y})} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) \\ &\stackrel{(a)}{\leq} \sum_{\substack{\tilde{m}_A, \tilde{m}_B : \tilde{m}_A \neq m_A, \\ (\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n}} \frac{|T_{Z|X\tilde{X}\tilde{Y}Y}^n(\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{y})|}{|T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y})|} \end{aligned}$$

$$\begin{aligned}
&\stackrel{(b)}{\leq} \sum_{\substack{\tilde{m}_A, \tilde{m}_B: \tilde{m}_A \neq m_A, \\ (\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n}} \frac{\exp\left(nH(Z|X\tilde{X}\tilde{Y}Y)\right)}{(n+1)^{-|\mathcal{X}||\mathcal{Y}||\mathcal{Z}|} \exp\left(nH(Z|XY)\right)} \\
&\stackrel{(c)}{\leq} \sum_{\substack{\tilde{m}_A, \tilde{m}_B: \\ (\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n}} \exp\left(-n\left(I(Z; \tilde{X}\tilde{Y}|XY) - \epsilon\right)\right) \\
&\stackrel{(d)}{\leq} \exp\left(n\left(|R_A - I(\tilde{X}; \tilde{Y}XY)|^+ + |R_B - I(\tilde{Y}; XY)|^+ - I(Z; \tilde{X}\tilde{Y}|XY) + 2\epsilon\right)\right), \tag{30}
\end{aligned}$$

where (a) follows by noting that whenever  $\mathbf{z}$  belongs to  $T_{Z|X\tilde{X}\tilde{Y}Y}^n(\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{y})$ ,  $\mathbf{z}$  also belongs to  $T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y})$ ; and for each  $\mathbf{z}$  in  $T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y})$ , the value of  $W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y})$  is the same and hence is upper bounded by  $1/|T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y})|$ . (b) follows from (22), (c) holds for sufficiently large  $n$  and (d) follows from (16). We see that

$$\begin{aligned}
I(Z; \tilde{X}\tilde{Y}|XY) &= I(XZ; \tilde{X}\tilde{Y}|Y) - I(X; \tilde{X}\tilde{Y}|Y) \\
&\stackrel{(a)}{\geq} \eta - I(X; \tilde{X}\tilde{Y}|Y) \\
&\stackrel{(b)}{\geq} \eta - I(X; \tilde{X}\tilde{Y}Y) \\
&\stackrel{(c)}{\geq} \eta - |R_A - I(\tilde{X}; \tilde{Y}Y)|^+ - |R_B - I(\tilde{Y}; Y)|^+ - \epsilon
\end{aligned}$$

where (a) follows from the fact that  $I(XZ; \tilde{X}\tilde{Y}|Y) \geq \eta$  for  $P_{X\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1$  (see (26)), (b) from  $I(X; \tilde{X}\tilde{Y}|Y) \leq I(X; \tilde{X}\tilde{Y}Y)$  and (c) follows from (29). Applying this to (30),

$$\begin{aligned}
&\sum_{\mathbf{z} \in \mathcal{E}_{m_A,1}(P_{X\tilde{X}\tilde{Y}YZ})} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) \\
&\leq \exp\left(n\left(|R_A - I(\tilde{X}; \tilde{Y}XY)|^+ + |R_B - I(\tilde{Y}; XY)|^+ + |R_A - I(\tilde{X}; \tilde{Y}Y)|^+ + |R_B - I(\tilde{Y}; Y)|^+ - \eta + 3\epsilon\right)\right) \\
&\stackrel{(a)}{\leq} \exp(n(4\delta - \eta + 3\epsilon)) \\
&\rightarrow 0 \text{ when } \eta > 3\epsilon + 4\delta \quad (\text{see (19)}). \tag{31}
\end{aligned}$$

Here, (a) follows by recalling that  $R_A = R_B = \delta$  and noting that  $|R_A - I(\tilde{X}; \tilde{Y}XY)|^+ + |R_B - I(\tilde{Y}; XY)|^+ + |R_A - I(\tilde{X}; \tilde{Y}Y)|^+ + |R_B - I(\tilde{Y}; Y)|^+ \leq 2R_A + 2R_B = 4\delta$ . Now, we move on to the second term  $P_{\mathcal{E}_{m_A,2}}(\mathbf{y})$  in (28). Proceeding in a similar fashion, we see that by using (17), it is sufficient to consider distributions  $P_{X\tilde{Y}_1\tilde{Y}_2YZ} \in \mathcal{P}_2$  for which

$$I\left(X; \tilde{Y}_1\tilde{Y}_2Y\right) \leq |R_B - I(\tilde{Y}_1; Y)|^+ + |R_B - I(\tilde{Y}_2; \tilde{Y}_1Y)|^+ + \epsilon. \tag{32}$$

For  $P_{X\tilde{Y}_1\tilde{Y}_2YZ} \in \mathcal{P}_2$  satisfying (32), along similar lines as the steps which led to (30) and (31),

$$\begin{aligned}
&\sum_{\mathbf{z} \in \mathcal{E}_{m_A,2}(P_{X\tilde{Y}_1\tilde{Y}_2YZ})} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) \\
&= \sum_{\substack{\tilde{m}_{B1}, \tilde{m}_{B2}: \tilde{m}_{B1} \neq \tilde{m}_{B2}, \\ (\mathbf{x}_{m_A}, \mathbf{y}_{\tilde{m}_{B1}}, \mathbf{y}_{\tilde{m}_{B2}}, \mathbf{y}) \in T_{X\tilde{Y}_1\tilde{Y}_2Y}^n}} \sum_{\mathbf{z}: (\mathbf{x}_{m_A}, \mathbf{y}_{\tilde{m}_{B1}}, \mathbf{y}_{\tilde{m}_{B2}}, \mathbf{z}) \in T_{X\tilde{Y}_1\tilde{Y}_2YZ}^n} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) \\
&\leq \sum_{\substack{\tilde{m}_{B1}, \tilde{m}_{B2}: \tilde{m}_{B1} \neq \tilde{m}_{B2}, \\ (\mathbf{x}_{m_A}, \mathbf{y}_{\tilde{m}_{B1}}, \mathbf{y}_{\tilde{m}_{B2}}, \mathbf{y}) \in T_{X\tilde{Y}_1\tilde{Y}_2Y}^n}} \frac{|T_{Z|X\tilde{Y}_1\tilde{Y}_2Y}^n(\mathbf{x}_{m_A}, \mathbf{y}_{\tilde{m}_{B1}}, \mathbf{y}_{\tilde{m}_{B2}}, \mathbf{y})|}{|T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y})|} \\
&\leq \sum_{\substack{\tilde{m}_{B1}, \tilde{m}_{B2}: \tilde{m}_{B1} \neq \tilde{m}_{B2}, \\ (\mathbf{x}_{m_A}, \mathbf{y}_{\tilde{m}_{B1}}, \mathbf{y}_{\tilde{m}_{B2}}, \mathbf{y}) \in T_{X\tilde{Y}_1\tilde{Y}_2Y}^n}} \frac{\exp\left(nH(Z|X\tilde{Y}_1\tilde{Y}_2Y)\right)}{(n+1)^{-|\mathcal{X}||\mathcal{Y}||\mathcal{Z}|} \exp\left(nH(Z|XY)\right)} \\
&\stackrel{(a)}{\leq} \sum_{\substack{\tilde{m}_{B1}, \tilde{m}_{B2}: \tilde{m}_{B1} \neq \tilde{m}_{B2}, \\ (\mathbf{x}_{m_A}, \mathbf{y}_{\tilde{m}_{B1}}, \mathbf{y}_{\tilde{m}_{B2}}, \mathbf{y}) \in T_{X\tilde{Y}_1\tilde{Y}_2Y}^n}} \exp\left(-n\left(I(Z; \tilde{Y}_1\tilde{Y}_2|XY) - \epsilon\right)\right) \\
&\stackrel{(b)}{\leq} \exp\left(n\left(|R_B - I(\tilde{Y}_1; XY)|^+ + |R_B - I(\tilde{Y}_2; \tilde{Y}_1XY)|^+ - I(Z; \tilde{Y}_1\tilde{Y}_2|XY) + 2\epsilon\right)\right) \\
&= \exp\left(n\left(|R_B - I(\tilde{Y}_1; XY)|^+ + |R_B - I(\tilde{Y}_2; \tilde{Y}_1XY)|^+ + I(X; \tilde{Y}_1\tilde{Y}_2|Y) - I(XZ; \tilde{Y}_1\tilde{Y}_2|Y) + 2\epsilon\right)\right)
\end{aligned}$$

$$\begin{aligned}
&\stackrel{(c)}{\leq} \exp\left(n\left(|R_B - I(\tilde{Y}_1; XY)|^+ + |R_B - I(\tilde{Y}_2; \tilde{Y}_1 XY)|^+ + |R_B - I(\tilde{Y}_1; Y)|^+ \right. \right. \\
&\quad \left. \left. + |R_B - I(\tilde{Y}_2; \tilde{Y}_1 Y)|^+ - \eta + 3\epsilon\right)\right) \\
&\stackrel{(d)}{\leq} \exp(n(4\delta - \eta + 3\epsilon)) \\
&\rightarrow 0 \text{ when } \eta > 3\epsilon + 4\delta \quad (\text{see (19)}).
\end{aligned}$$

where (a) holds for large  $n$ , (b) follows from (18), (c) from (32) (note that  $I(X; \tilde{Y}_1 \tilde{Y}_2 | Y) \leq I(X; \tilde{Y}_1 \tilde{Y}_2 Y)$ ) and the fact that  $I(XZ; \tilde{Y}_1 \tilde{Y}_2 | Y) \geq \eta$  since  $P_{X\tilde{Y}_1\tilde{Y}_2YZ} \in \mathcal{P}_2$  (see (27)) and (d) by recalling that  $R_B = \delta$  and hence  $|R_B - I(\tilde{Y}_1; XY)|^+ + |R_B - I(\tilde{Y}_2; \tilde{Y}_1 XY)|^+ + |R_B - I(\tilde{Y}_1; Y)|^+ + |R_B - I(\tilde{Y}_2; \tilde{Y}_1 Y)|^+ \leq 4R_B = 4\delta$ .

Thus,  $P_{e, \text{mal B}} \rightarrow 0$  as  $n \rightarrow \infty$  for the sufficiently small  $\delta > \epsilon > 0$  so that  $\eta > 3\epsilon + 4\delta$  is sufficiently small for Lemma 6 to hold. Similarly, because of symmetry, we can show that if  $\eta > 3\epsilon + 4\delta$  the probability of error goes to zero as  $n$  goes to infinity when user A is malicious.  $\square$

#### IV. CAPACITY REGION

##### A. Inner bound

For distributions  $P_A$  and  $P_B$  over  $\mathcal{X}$  and  $\mathcal{Y}$  respectively, we define  $\mathcal{P}(P_A, P_B) \stackrel{\text{def}}{=} \{P_{XY\tilde{X}\tilde{Y}Z} : P_{X\tilde{Y}Z} = P_A \times P_{\tilde{Y}} \times W \text{ for some } P_{\tilde{Y}} \text{ and } P_{\tilde{X}YZ} = P_{\tilde{X}} \times P_B \times W \text{ for some } P_{\tilde{X}}\}$ . Let  $\mathcal{R}_1(P_A, P_B)$  be the set of rate pairs  $(R_A, R_B)$  such that

$$\begin{aligned}
R_A &\leq \min_{P_{XY\tilde{X}\tilde{Y}Z} \in \mathcal{P}(P_A, P_B)} I(X; Z), \text{ and} \\
R_B &\leq \min_{P_{XY\tilde{X}\tilde{Y}Z} \in \mathcal{P}(P_A, P_B): X \perp Y} I(Y; Z|X).
\end{aligned} \tag{33}$$

Similarly, let  $\mathcal{R}_2(P_A, P_B)$  be the set of rate pairs given by

$$\begin{aligned}
R_A &\leq \min_{P_{XY\tilde{X}\tilde{Y}Z} \in \mathcal{P}(P_A, P_B): X \perp Y} I(X; Z|Y), \text{ and} \\
R_B &\leq \min_{P_{XY\tilde{X}\tilde{Y}Z} \in \mathcal{P}(P_A, P_B)} I(Y; Z).
\end{aligned} \tag{34}$$

**Theorem 7** (Inner bound). *When  $\text{int}(\mathcal{C}) \neq \emptyset$ ,*

$$\text{conv}(\cup_{P_A, P_B} (\mathcal{R}_1(P_A, P_B) \cup \mathcal{R}_2(P_A, P_B))) \subseteq \mathcal{C}.$$

The proof uses the same codebook lemma (Lemma 4) as used for the achievability of Theorem 1 but with different rates. The decoder is a slightly modified version of the decoder used in Theorem 1. The modified decoder uses similar conditions as used in the decoder for the achievability of Theorem 1, but the steps are performed in a specific order. At each step only those codewords are considered which have not been eliminated in the previous steps. Please see Appendix C for details.

**Remark 4.** *The inner bound to the capacity region given by Theorem 7 is tight for binary erasure MAC [3, pg. 83] as shown in Section V-A.*

##### B. Outer bound

The outer bound is provided in terms of the capacity region of an Arbitrarily Varying Multiple Access Channel (AV-MAC).

**Definition 5** (see [11]). *An AV-MAC  $\mathcal{W} = \{W(\cdot|\cdot, \cdot, s) | s \in \mathcal{S}\}$  is a family of MACs parameterized by the set of state symbols  $\mathcal{S}$  where MACs  $W(\cdot|\cdot, \cdot, s)$  are randomized  $\mathcal{X} \times \mathcal{Y} \rightarrow \mathcal{Z}$  maps.*

The state of an AV-MAC can vary arbitrarily during the transmission. This can also be interpreted as an adversary choosing a distribution over the state symbols (and hence a channel from the convex hull of  $\mathcal{W}$ ) for each symbol sent into the channel by the senders. Note that the adversary only knows that each sender sends a codeword uniformly at random from their codebook, and is unaware of the actual symbols sent into the channel by each sender. Similarly, the senders are unaware of the choice of adversarial state vector. We use  $\mathcal{C}_{\text{AV-MAC}}(\mathcal{W})$  (or simply  $\mathcal{C}_{\text{AV-MAC}}$ ) to denote the deterministic coding capacity region of an AV-MAC  $\mathcal{W}$ . See [9], [11] for the definition of a deterministic code and the expression of the capacity region.

**Definition 6.** *For a MAC  $W$ , let  $\tilde{\mathcal{W}}_W$  be the set of MACs  $\tilde{W}$  such that there is a pair of conditional distributions  $Q_{X'|X}$  and  $Q_{Y'|Y}$  satisfying*

$$\tilde{W}(z|x, y) = \sum_{x'} Q_{X'|X}(x'|x) W(z|x', y)$$



$$= \sum_{y'} Q_{Y'|Y}(y'|y)W(z|x, y'), \quad (35)$$

for all  $x, y, z \in \mathcal{X} \times \mathcal{Y} \times \mathcal{Z}$ .

The outer bound on the capacity region for communication with adversary identification of a byzantine-MAC  $W$  is in terms of the capacity region of the AV-MAC  $\tilde{\mathcal{W}}_W$  defined above. We first notice that  $W \in \tilde{\mathcal{W}}_W$  by choosing trivial distributions  $Q_{X'|X}(x|x) = 1$  for all  $x$  and  $Q_{Y'|Y}(y|y) = 1$  for all  $y$ . Additionally, the set  $\tilde{\mathcal{W}}_W$  is convex because for every  $(Q_{X'|X}, Q_{Y'|Y})$  and  $(Q'_{X'|X}, Q'_{Y'|Y})$  satisfying (35), the pair  $(\alpha Q_{X'|X} + (1 - \alpha)Q'_{X'|X}, \alpha Q_{Y'|Y} + (1 - \alpha)Q'_{Y'|Y})$ ,  $\alpha \in [0, 1]$  also satisfies (35). To get an outer bound, consider an adversary identifying code with a small probability of error for a MAC  $W$ . Suppose user A is malicious and attacks in the following manner: it runs its encoder on a uniformly distributed message from its message set, then passes the output of the encoder through  $\prod_{i=1}^n Q_{i, X'|X}$  where for all  $i \in [1 : n]$ ,  $(Q_{i, X'|X}, Q_{i, Y'|Y})$  satisfy (35) for some  $Q_{i, Y'|Y}$ . The output of  $\prod_{i=1}^n Q_{i, X'|X}$  is finally sent to the MAC  $W$  as input by user A. This can also be interpreted as an attack by user B using  $\prod_{i=1}^n Q_{i, Y'|Y}$ . Thus, at the receiver, it is not clear if user A attacked or user B attacked. Hence, the malicious user cannot be identified reliably. So, the decoder must output a pair of messages of which the message for the honest user is correct with high probability. This means that the decoding should be correct with high probability for  $\tilde{W}^{(n)}(\cdot|\cdot, \cdot) = \prod_{i=1}^n (\sum_{x_i \in \mathcal{X}} Q_{i, X'|X}(x_i|\cdot)W(\cdot|x_i, \cdot)) = \prod_{i=1}^n \sum_{y_i \in \mathcal{Y}} Q_{i, Y'|Y}(y_i|\cdot)W(\cdot|y_i)$ . Thus, any good adversary identifying code for the MAC  $W$  must also be a good communication code for  $\tilde{W}^{(n)}$ . In fact, we can show the following (proved later):

**Lemma 8.** Any  $(N_A, N_B, n)$  adversary identifying code  $(f_A, f_B, \phi)$  for a MAC  $W$  with  $P_e(f_A, f_B, \phi) \leq \epsilon$  is also an  $(N_A, N_B, n)$  communication code for the AV-MAC  $\tilde{\mathcal{W}}_W$  with average probability of error at most  $2\epsilon$ .

This implies that the capacity region  $\mathcal{C}$  (for communication with adversary identification) of MAC  $W$  must be a subset of the capacity region of the AV-MAC  $\tilde{\mathcal{W}}_W$  (Definition 6) parametrized by a pair of distributions  $(Q_{X'|X}, Q_{Y'|Y})$  satisfying (35). This argument is formalized below:

**Theorem 9** (Outer bound).  $\mathcal{C} \subseteq \mathcal{C}_{\text{AV-MAC}}(\tilde{\mathcal{W}}_W)$ . Moreover, there exists an AV-MAC  $\mathcal{W}_W$  such that  $\mathcal{C}_{\text{AV-MAC}}(\mathcal{W}_W) = \mathcal{C}_{\text{AV-MAC}}(\tilde{\mathcal{W}}_W)$  and  $|\mathcal{W}_W| \leq 2^{|\mathcal{X}|^2 + |\mathcal{Y}|^2}$ .<sup>6</sup>

*Proof.* Lemma 8 implies that  $\mathcal{C}$ , the capacity region of communication with adversary identification of a MAC  $W$ , is outer bounded by the capacity region  $\mathcal{C}_{\text{AV-MAC}}(\tilde{\mathcal{W}}_W)$  of the AV-MAC  $\tilde{\mathcal{W}}_W$ . Further, note that the capacity of an AV-MAC  $\mathcal{W}$  only depends on the convex hull of  $\mathcal{W}$  (see [11, Theorem 1]). So, the capacity of  $\tilde{\mathcal{W}}_W$  is the same as the capacity of another AV-MAC  $\mathcal{W}_W$  which consists of vertices of the convex polytope  $\tilde{\mathcal{W}}_W \subseteq \mathbb{R}^{|\mathcal{X}| \times |\mathcal{Y}| \times |\mathcal{Z}|}$ . The elements in the set  $\tilde{\mathcal{W}}_W$  are parameterized by  $(Q_{X'|X}, Q_{Y'|Y})$  pairs. It consists of the vertices of the polytope formed using constraints in (35) and constraints of the form: (1)  $\sum_{x'} P_{X'|X}(x'|x) = 1$  for all  $x$ , and (2)  $P_{X'|X}(x'|x) \geq 0$ . There are similar constraints for  $P_{Y'|Y}$ . Note that there are  $|\mathcal{X}|^2 + |\mathcal{Y}|^2$  inequality constraints. Every point in the resulting polytope satisfies all the equality constraints. We will get faces, edges, vertices etc. depending on the number of additional inequality constraints satisfied at that point. Thus, number of vertices  $\leq 2^{|\mathcal{X}|^2 + |\mathcal{Y}|^2}$ .  $\square$

*Proof of Lemma 8.* Consider an  $(N_A, N_B, n)$  adversary identifying code  $(f_A, f_B, \phi)$  such that  $P_e(f_A, f_B, \phi) \leq \epsilon$  For  $i \in [1 : n]$ , let  $(Q_{i, X'|X}, Q_{i, Y'|Y})$  be an arbitrary sequence of pairs of channel distributions such that each pair satisfies (35) and define  $\tilde{W}_i$  as

$$\tilde{W}_i(z|x, y) \stackrel{\text{def}}{=} \sum_{x'} Q_{i, X'|X}(x'|x)W(z|x', y) = \sum_{y'} Q_{i, Y'|Y}(y'|y)W(z|x, y')$$

for all  $x, y, z$ . Let  $Q_{X'|X} \stackrel{\text{def}}{=} \prod_{i=1}^n Q_{i, X'|X}$ ,  $Q_{Y'|Y} \stackrel{\text{def}}{=} \prod_{i=1}^n Q_{i, Y'|Y}$  and  $\tilde{W}^{(n)} \stackrel{\text{def}}{=} \prod_{i=1}^n \tilde{W}_i$ . Recall that for  $m_B \in \mathcal{M}_B$  and  $\phi_B$  as defined in (2), we define  $\mathcal{E}_{m_B}^B = \{z : \phi_B(z) \notin \{m_B, \mathbf{a}\}\}$ . Consider a malicious user-A who chooses  $M_A$  uniformly from  $[1 : N_A]$ , passes  $f_A(M_A)$  over  $Q_{X'|X}$ , and transmits the resulting vector. We may conclude that (see (4)),

$$\begin{aligned} P_{e, \text{mal A}} &= \max_{\mathbf{x} \in \mathcal{X}^n} \left( \frac{1}{N_B} \sum_{m_B} W^n \left( \mathcal{E}_{m_B}^B \mid \mathbf{x}, f_B(m_B) \right) \right) \\ &\geq \frac{1}{N_B} \sum_{m_B} \sum_{\mathbf{x}} \left( \frac{1}{N_A} \sum_{m_A} Q_{X'|X}(\mathbf{x} | f_A(m_A)) \right) W^n \left( \mathcal{E}_{m_B}^B \mid \mathbf{x}, f_B(m_B) \right) \\ &= \frac{1}{N_A \cdot N_B} \sum_{m_A, m_B} \sum_{\mathbf{x}} Q_{X'|X}(\mathbf{x} | f_A(m_A)) W^n \left( \mathcal{E}_{m_B}^B \mid \mathbf{x}, f_B(m_B) \right) \end{aligned}$$

<sup>6</sup>The capacity region of an AV-MAC only depends on its convex hull which is defined by taking convex combinations of channels under different states [9], [11]. The AV-MAC  $\mathcal{W}_W$  has the same convex hull as  $\tilde{\mathcal{W}}_W$  but with only finitely many states.

$$= \frac{1}{N_A \cdot N_B} \sum_{m_A, m_B} \tilde{W}^{(n)} \left( \mathcal{E}_{m_B}^B \mid f_A(m_A), f_B(m_B) \right). \quad (36)$$

Similarly, for  $\phi_A$  as defined in (1),  $m_A \in \mathcal{M}_A$  and  $\mathcal{E}_{m_A}^A = \{z : \phi_A(z) \notin \{m_A, \mathbf{b}\}\}$ ,

$$P_{e, \text{mal B}} \geq \frac{1}{N_A \cdot N_B} \sum_{m_A, m_B} \tilde{W}^{(n)} \left( \mathcal{E}_{m_A}^A \mid f_A(m_A), f_B(m_B) \right). \quad (37)$$

First notice that for  $(m_A, m_B) \in \mathcal{M}_A \times \mathcal{M}_B$  and  $z \in \mathcal{Z}^n$ ,

$$\begin{aligned} & \{z : \phi(z) \neq (m_A, m_B)\} \\ &= \{z : \phi_A(z) \neq m_A\} \cup \{z : \phi_B(z) \neq m_B\} \\ &\stackrel{(a)}{=} \{z : \phi_A(z) \notin \{m_A, \mathbf{b}\}\} \cup \{z : \phi_B(z) \notin \{m_B, \mathbf{a}\}\} \\ &= \mathcal{E}_{m_A}^A \cup \mathcal{E}_{m_B}^B \end{aligned}$$

where (a) holds by noting from definitions (1) and (2) that  $\phi_A = \mathbf{a}$  (resp.  $\mathbf{b}$ ) if and only if  $\phi_B = \mathbf{a}$  (resp.  $\mathbf{b}$ ). Thus,

$$\begin{aligned} & \frac{1}{N_A \cdot N_B} \sum_{m_A, m_B} \tilde{W}^{(n)} \left( \{z : \phi(z) \neq (m_A, m_B)\} \mid f_A(m_A), f_B(m_B) \right) \\ &= \frac{1}{N_A \cdot N_B} \sum_{m_A, m_B} \tilde{W}^{(n)} \left( \mathcal{E}_{m_B}^B \cup \mathcal{E}_{m_A}^A \mid f_A(m_A), f_B(m_B) \right) \\ &\stackrel{(a)}{\leq} P_{e, \text{mal A}} + P_{e, \text{mal B}} \\ &\leq 2\epsilon, \end{aligned}$$

where (a) follows from a union bound and (36)-(37). Recall that every pair  $(Q_{X'|X}, Q_{Y'|Y})$  satisfying (35) corresponds to an element in  $\tilde{\mathcal{W}}_W$ , which is a convex set (see the discussion after Definition 6). Thus, for any  $\epsilon > 0$ , an adversary identifying code for the MAC  $W$  with an average probability of error  $\epsilon$  is also a communication code for the AV-MAC  $\tilde{\mathcal{W}}_W$  with an average probability of error at most  $2\epsilon$ . So, the deterministic coding capacity region  $\mathcal{C}$  of  $W$  is contained by the deterministic coding capacity region  $\mathcal{C}_{\text{AV-MAC}}(\tilde{\mathcal{W}}_W)$  of the AV-MAC  $\tilde{\mathcal{W}}_W$ .  $\square$

We will now give an alternative proof of the first part of Theorem 1. We will use the connection of the present model with the AV-MAC given by Lemma 8 and then establish a connection between an A-spoofable byzantine-MAC and a symmetrizable- $\mathcal{X}$  [13, equation (3.2)] AV-MAC. A symmetrizable- $\mathcal{X}$  AV-MAC, introduced in [13], does not allow reliable communication by user A.

**Theorem 10** (First part of Theorem 1). *If a byzantine-MAC is A-spoofable (resp. B-spoofable), communication with adversary identification from user-A (resp. user-B) is impossible. Specifically, for any  $(N_A, N_B, n)$  adversary identifying code with  $N_A \geq 2$  (resp.  $N_B \geq 2$ ), the probability of error is at least  $1/12$ .*

*Alternate proof.* We will first show that if a byzantine-MAC is A-spoofable then the corresponding AV-MAC given by Definition 6 is symmetrizable- $\mathcal{X}$  [13, equation (3.2)].

To this end, suppose a given byzantine-MAC  $W$  is A-spoofable. This implies that (7) holds. Then, for  $\mathcal{S} = \mathcal{X}$  and by replacing  $\tilde{X}$  and with  $S$  in (7), we obtain

$$\sum_y Q_{Y|\tilde{Y}S}(y|\tilde{y}, \tilde{x}) W_{Z|XY}(z|x', y) \quad (38)$$

$$= \sum_y Q_{Y|\tilde{Y}S}(y|\tilde{y}, x') W_{Z|XY}(z|\tilde{x}, y) \quad (39)$$

$$= \sum_x Q_{X|X'S}(x|x', \tilde{x}) W_{Z|XY}(z|x, \tilde{y}) \quad \text{for all } x', \tilde{x}, \tilde{y}, z. \quad (40)$$

For every  $s \in \mathcal{S}$ , (38) and (40) imply that

$$\begin{aligned} \tilde{W}_{Z|XYS}(z|x', \tilde{y}, s) &\stackrel{\text{def}}{=} \sum_y Q_{Y|\tilde{Y}S}(y|\tilde{y}, s) W_{Z|XY}(z|x', y) \\ &= \sum_x Q_{X|X'S}(x|x', s) W_{Z|XY}(z|x, \tilde{y}) \quad \text{for all } x', \tilde{y}, s. \end{aligned}$$

Thus, the set  $\tilde{\mathcal{W}}_W$  in Definition 6 is such that

$$\left\{ \tilde{W}_{Z|XYS}(\cdot|\cdot, \cdot, s) : s \in \mathcal{S} \right\} \subseteq \tilde{\mathcal{W}}_W.$$

By (38) and (39), we note that for all  $z, \tilde{x}, \tilde{y}, x'$ ,

$$\tilde{W}_{Z|XY S}(z|x', \tilde{y}, \tilde{x}) = \tilde{W}_{Z|XY S}(z|\tilde{x}, \tilde{y}, x')$$

Thus, for  $P_{S|X}(s|x) \stackrel{\text{def}}{=} \mathbb{1}_{\{s=x\}}$ ,  $s, x \in \mathcal{X}$ , where  $\mathbb{1}$  is the indicator function,

$$\sum_s P_{S|X}(s|\tilde{x}) \tilde{W}_{Z|XY S}(z|x', \tilde{y}, s) = \sum_{s'} P_{S|X}(s'|x') \tilde{W}_{Z|XY S}(z|\tilde{x}, \tilde{y}, s)$$

for all  $x', \tilde{x}, \tilde{y}$  and  $z$ . That is, the AV-MAC  $\tilde{W}_W$  is symmetrizable- $\mathcal{X}$ . Hence, if the byzantine-MAC  $W$  is A-spoofable, then the AV-MAC  $\tilde{W}_W$  is symmetrizable- $\mathcal{X}$ .

Now, suppose that there is an  $(N_A, N_B, n)$  adversary identifying code  $(f_A, f_B, \phi)$  such that  $P_e(f_A, f_B, \phi) \leq 1/12$ . From Lemma 8, this implies that there is an  $(N_A, N_B, n)$  communication code for the AV-MAC  $\tilde{W}_W$  with average probability of error at most  $1/6$ . However, for any  $(N_A, N_B, n)$  code (with  $N_A \geq 2$ ) for a symmetrizable- $\mathcal{X}$  AV-MAC, Gubner [13, equation (3.4)] shows that the average probability of error is at least  $1/4$ . Thus, there does not exist any  $(N_A, N_B, n)$  adversary identifying code  $(f_A, f_B, \phi)$  such that  $P_e(f_A, f_B, \phi) \leq 1/12$  and  $N_A \geq 2$ .  $\square$

## V. EXAMPLES AND COMPARISON WITH OTHER MODELS

### A. Tightness of the inner bound for the Binary Erasure MAC

The binary erasure MAC [3, pg. 83] is given by  $Z = X + Y$  where  $\mathcal{X} = \mathcal{Y} = \{0, 1\}$  and  $\mathcal{Z} = \{0, 1, 2\}$ . We will show that for the binary erasure MAC, the inner bound on  $\mathcal{C}$  given by Theorem 7 is the same as its (non-adversarial) capacity region  $\mathcal{C}_{\text{MAC}}$ . Hence, it is tight.

Recall that for distributions  $P_A$  and  $P_B$  over  $\mathcal{X}$  and  $\mathcal{Y}$ ,  $\mathcal{P}(P_A, P_B) = \{P_{XY\tilde{X}\tilde{Y}Z} : P_{X\tilde{Y}Z} = P_A \times P_{\tilde{Y}} \times W \text{ for some } P_{\tilde{Y}} \text{ and } P_{\tilde{X}YZ} = P_{\tilde{X}} \times P_B \times W \text{ for some } P_{\tilde{X}}\}$ . We choose  $P_A$  and  $P_B$  arbitrarily close to the uniform distribution  $U$  on  $\{0, 1\}$  while ensuring that  $P_A \neq P_B$ . Using these distributions, we will show that for  $P_{XY\tilde{X}\tilde{Y}Z} \in \mathcal{P}(P_A, P_B)$  satisfying  $X \perp\!\!\!\perp Y$ ,  $\tilde{X} = X$  and  $\tilde{Y} = Y$ .

To this end, consider  $P_{XY\tilde{X}\tilde{Y}Z} \in \mathcal{P}(P_A, P_B)$ .

$$\mathbb{P}(Z = 0) = P_A(0)P_{\tilde{Y}}(0) = P_{\tilde{X}}(0)P_B(0). \quad (41)$$

$$\mathbb{P}(Z = 2) = (1 - P_A(0))(1 - P_{\tilde{Y}}(0)) = (1 - P_{\tilde{X}}(0))(1 - P_B(0)).$$

This implies that

$$1 + P_A(0)P_{\tilde{Y}}(0) - P_A(0) - P_{\tilde{Y}}(0) = 1 + P_{\tilde{X}}(0)P_B(0) - P_{\tilde{X}}(0) - P_B(0). \quad (42)$$

Using (41) and (42), we get  $P_A(0) + P_{\tilde{Y}}(0) = P_{\tilde{X}}(0) + P_B(0)$ . Thus,

$$P_{\tilde{X}}(0) = P_A(0) + P_{\tilde{Y}}(0) - P_B(0). \quad (43)$$

Substituting the value of  $P_{\tilde{X}}(0)$  from (43) into (41), we get  $P_A(0)P_{\tilde{Y}}(0) = P_A(0)P_B(0) + P_{\tilde{Y}}(0)P_B(0) - P_B(0)P_B(0)$ . This implies that

$$(P_A(0) - P_B(0))(P_{\tilde{Y}}(0) - P_B(0)) = 0.$$

Thus, either  $P_A(0) = P_B(0)$  or  $P_{\tilde{Y}}(0) = P_B(0)$ . Substituting this in (43), we get either  $P_A(0) = P_B(0)$  and  $P_{\tilde{X}}(0) = P_{\tilde{Y}}(0)$ , or  $P_{\tilde{Y}}(0) = P_B(0)$  and  $P_{\tilde{X}}(0) = P_A(0)$ . If we choose  $P_A$  and  $P_B$  such that  $P_A \neq P_B$ , then for every  $P_{XY\tilde{X}\tilde{Y}Z} \in \mathcal{P}(P_A, P_B)$ ,  $P_{\tilde{Y}} = P_Y = P_B$  and  $P_{\tilde{X}} = P_X = P_A$ .

We know from the definition of  $\mathcal{P}(P_A, P_B)$ , that  $X \perp\!\!\!\perp \tilde{Y}$  and  $\tilde{X} \perp\!\!\!\perp Y$ . We now analyse the case when there is a further restriction of  $X \perp\!\!\!\perp Y$  on the distributions. From the definition of  $\mathcal{P}(P_A, P_B)$ , we note that  $P_{\tilde{X}Y|X\tilde{Y}}(0, 0|0, 0) = 1$  and  $P_{\tilde{X}Y|X\tilde{Y}}(1, 1|1, 1) = 1$ . Let  $P_{\tilde{X}Y|X\tilde{Y}}(0, 1|0, 1) = \alpha$  and  $P_{\tilde{X}Y|X\tilde{Y}}(1, 0|1, 0) = 1 - \alpha$  (Note that  $P_{\tilde{X}Y|X\tilde{Y}}((0, 0)|0, 1) = P_{\tilde{X}Y|X\tilde{Y}}((1, 1)|0, 1) = 0$  by definition of  $\mathcal{P}(P_A, P_B)$ ). Similarly, let  $P_{\tilde{X}Y|X\tilde{Y}}(1, 0|1, 0) = \beta$  and  $P_{\tilde{X}Y|X\tilde{Y}}(0, 1|1, 0) = 1 - \beta$ . Thus,  $P_{XY}(0, 0) = P_{X\tilde{Y}}(0, 0)P_{\tilde{X}Y|X\tilde{Y}}(0, 0|0, 0) + P_{X\tilde{Y}}(0, 1)P_{\tilde{X}Y|X\tilde{Y}}(1, 0|0, 1) = P_X(0)P_{\tilde{Y}}(0) \cdot 1 + P_X(0)P_{\tilde{Y}}(1) \cdot (1 - \alpha)$ . Also,  $P_{XY}(0, 0) = P_X(0)P_Y(0) = P_X(0)P_{\tilde{Y}}(0)$  (The last equality follows by choosing  $P_X \neq P_Y$  (which is the same as  $P_A \neq P_B$ )). This implies that  $\alpha = 1$ . By evaluating  $P_{XY}(1, 1)$ , we can show that  $\beta = 1$ . This implies that  $\tilde{X} = X$  and  $\tilde{Y} = Y$ .

As mentioned earlier, we choose  $P_A$  and  $P_B$  arbitrarily close to uniform distributions such that  $P_A \neq P_B$ . Thus, in the limit, (33) evaluates to  $R_A \leq 0.5$  and  $R_B \leq 1$ , and (34) evaluates to  $R_A \leq 1$  and  $R_B \leq 0.5$ . Using time sharing between these two rate pairs, we obtain the entire MAC capacity region (This is the rate region  $\mathcal{C}$  in Fig. 7).

### B. Comparison with related models

In this section we contrast the present model with reliable communication and authenticated communication models.

a) *Reliable communication in a byzantine-MAC*: We consider a byzantine-MAC with a stronger decoding guarantee: the decoder, w.h.p, outputs a message pair of which the message(s) of honest user(s) is correct. In the presence of a malicious user, the channel from the honest user to the receiver can be treated as an Arbitrarily Varying Channel (AVC) [7] with the input of other user as state. Thus, the capacity region is outer bounded by the rectangular region defined by the AVC capacities of the two users' channels. Further, it is easy to see that this outer bound is achievable when both users use the corresponding AVC codes. Csiszár and Narayan show in [8] that the capacity of an AVC is zero if and only if it is *symmetrizable*. Translating this to the two-user byzantine-MAC, we define a byzantine-MAC to be *B-symmetrizable* if there exists a distribution  $P_{X|Y}$  such that

$$\sum_{x' \in \mathcal{X}} P_{X|Y}(x|y')W(z|x, y) = \sum_{x' \in \mathcal{X}} P_{X|Y}(x|y)W(z|x, y') \quad (44)$$

for all  $(x, y, z) \in \mathcal{X} \times \mathcal{Y} \times \mathcal{Z}$ . We define an A-symmetrizable byzantine-MAC analogously. A **symmetrizable** byzantine-MAC is one which is either A- or B-symmetrizable. Thus, reliable communication by both users is feasible in a byzantine-MAC if and only if it is not symmetrizable. We denote the reliable communication capacity of a byzantine-MAC by  $C_{\text{reliable}}$ .

b) *Authenticated communication in a byzantine-MAC* [4]: This model considers a byzantine-MAC with a weaker decoding guarantee: the decoder should reliably decode the messages when both users are honest. When one user is adversarial, the decoder either outputs a pair of messages of which the message of honest user is correct or it declares the presence of an adversary (without identifying it). In this case, the notion of an overwritable byzantine-MAC characterizes the class of channels with non-empty capacity region  $C_{\text{auth}}$  of authenticated communication. We say that a byzantine-MAC is *B-overwritable* [4, (1)] if there exists a distribution  $P_{X'|XY}$  such that

$$\sum_{x' \in \mathcal{X}} P_{X'|XY}(x'|x, y)W(z|x', y') = W(z|x, y) \quad (45)$$

for all  $y, y' \in \mathcal{Y}$ ,  $x \in \mathcal{X}$  and  $z \in \mathcal{Z}$ . Similarly, we can define an A-overwritable byzantine-MAC. If a byzantine-MAC is either A- or B-overwritable, we say that the byzantine-MAC is **overwritable**. Authenticated communication by both users is not feasible in an overwritable byzantine-MAC. Theorem 1 in [4] states that if the byzantine-MAC is not overwritable, then authenticated communication capacity is the same as the non adversarial capacity of the MAC, i.e.,  $C_{\text{auth}} = C_{\text{MAC}}$ .

**Proposition 11.** *All overwritable byzantine-MACs are spoofable and all spoofable byzantine-MACs are symmetrizable. Furthermore, both these inclusions are strict.*

While the inclusions in Proposition 11 are obvious from the problem definitions and the feasibility results, we nonetheless provide a direct argument. Suppose a byzantine-MAC is B-overwritable with  $P_{X'|XY}$  as the overwriting attack in (45). For any distribution  $Q_Y$  on  $\mathcal{Y}$ , let  $Q_{X|\tilde{X}\tilde{Y}}(x|\tilde{x}, \tilde{y}) \stackrel{\text{def}}{=} \sum_y Q_Y(y)P_{X'|XY}(x|\tilde{x}, y)$  for all  $x, \tilde{x}, \tilde{y}$  and  $Q_{Y|\tilde{Y}Y'}(y|\tilde{y}, y') \stackrel{\text{def}}{=} Q_Y(y)$  for all  $y, \tilde{y}, y'$ . Distributions  $Q_{X|\tilde{X}\tilde{Y}}$  and  $Q_{Y|\tilde{Y}Y'}$  as defined satisfy (8). Now, suppose a byzantine-MAC  $W_{Z|XY}$  is B-spoofable with attacks  $Q_{X|\tilde{X}\tilde{Y}}$  and  $Q_{Y|\tilde{Y}Y'}$ , satisfying (8). For all,  $x, y$ , let  $P_{X|Y}(x|y) \stackrel{\text{def}}{=} Q_{X|\tilde{X}\tilde{Y}}(x|\tilde{x}, y)$  for any  $\tilde{x} \in \mathcal{X}$ . It can be easily seen that the attack  $P_{X|Y}$  as defined satisfies (44). Examples 2 and 3 below show strict inclusion (see Fig. 5).

**Example 2** (symmetrizable, but not spoofable). *Binary erasure MAC*: It has binary inputs  $X, Y$  and outputs  $Z = X + Y$  where  $+$  is real addition, i.e.,  $\mathcal{Z} = \{0, 1, 2\}$ .

To show symmetrizability, we note that the distribution  $P_{X|Y}(x|y) = 1$  for all  $x = y$  is a symmetrizing attack in (44). Now, suppose the channel is *A-spoofable*, that is, there exist distributions  $Q_{Y|\tilde{X}\tilde{Y}}$  and  $Q_{X|\tilde{X}X'}$  such that  $\forall x', \tilde{x}, \tilde{y}, z$ ,

$$\begin{aligned} & \sum_y Q_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y})W_{Z|XY}(z|x', y) \\ &= \sum_y Q_{Y|\tilde{X}\tilde{Y}}(y|x', \tilde{y})W_{Z|XY}(z|\tilde{x}, y) \\ &= \sum_x Q_{X|\tilde{X}X'}(x|\tilde{x}, x')W_{Z|XY}(z|x, \tilde{y}). \end{aligned}$$

For  $(x', \tilde{x}, \tilde{y}, z) = (1, 0, 1, 2)$ , this gives  $Q_{Y|\tilde{X}\tilde{Y}}(1|0, 1) = 0 = Q_{X|\tilde{X}X'}(1|0, 1)$  and for  $(x', \tilde{x}, \tilde{y}, z) = (1, 0, 0, 0)$ , we get  $0 = Q_{Y|\tilde{X}\tilde{Y}}(0|1, 0) = Q_{X|\tilde{X}X'}(0|0, 1)$ . However,  $Q_{X|\tilde{X}X'}(1|0, 1) = Q_{X|\tilde{X}X'}(0|0, 1) = 0$  is not possible. Thus, the channel is not A-spoofable. Similarly, we can show that the channel is not B-spoofable.

**Example 3** (spoofable, but not overwritable). *Binary additive MAC*: It has binary inputs  $X, Y$  and outputs binary  $Z = X \oplus Y$  where  $\oplus$  is the XOR operation.

To show spoofability, note that the attacks  $Q_{X|\tilde{X}X'}(x|\tilde{x}, x') = 1/2$  for all  $x, \tilde{x}$  and  $x'$ , and  $Q_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) = 1/2$  for all  $y, \tilde{x}$  and  $\tilde{y}$ , satisfy (7) because they result in the same uniform output distribution over  $\mathcal{Z}$  in all the three cases in (7). Suppose

binary additive MAC  $Z = X \oplus Y$  is B-overwritable. Let  $P_{X'|XY}$  be the overwriting attack by user A which satisfies (45). Then, for  $(x, y, z) = (1, 1, 0)$  and all  $y'$ , (45) implies

$$P_{X'|XY}(0|1, 1)W(0|0, y') + P_{X'|XY}(1|1, 1)W(0|1, y') = W(0|1, 1) = 1.$$

For  $y' = 0$  and 1, this implies that  $P_{X'|XY}(0|1, 1) = 1$  and  $P_{X'|XY}(1|1, 1) = 1$  respectively, which is not possible simultaneously. Thus, the channel cannot be B-overwritable. Similarly, we can argue that the channel is not A-overwritable.




Fig. 5. The set of overwritable byzantine-MACs is a strict subset of the set of spoofable byzantine-MACs which, in turn, is a strict subset of the set of symmetrizable byzantine-MACs.

We also note from the problem definitions that  $\mathcal{C}_{\text{reliable}} \subseteq \mathcal{C} \subseteq \mathcal{C}_{\text{auth}} \subseteq \mathcal{C}_{\text{MAC}}$ . Next, we give an example of a channel for which  $\mathcal{C}_{\text{reliable}}$ ,  $\mathcal{C}$  and  $\mathcal{C}_{\text{auth}}$  are distinct. The example is constructed by using the byzantine-MACs in Examples 2 and 3 in parallel.

**Example 4** ( $(Z_1, Z_2) = (X_1 + Y_1, X_2 \oplus Y_2)$ ). For binary inputs  $X_1, X_2, Y_1, Y_2$ , the output is  $(Z_1, Z_2) = (X_1 + Y_1, X_2 \oplus Y_2)$ .

The channels  $Z_1 = X_1 + Y_1$  and  $Z_2 = X_2 \oplus Y_2$  are both non-overwritable and symmetrizable. Since the byzantine-MACs do not interact when used in parallel, we can show that the resultant byzantine-MAC  $(Z_1, Z_2) = (X_1 + Y_1, X_2 \oplus Y_2)$  is also non-overwritable and symmetrizable. We will first show that this channel is B-symmetrizable, that is, there exists distribution  $P_{X|Y}$  such that

$$\sum_{x \in \mathcal{X}} P_{X|Y}(x|y')W(z|x, y) = \sum_{x \in \mathcal{X}} P_{X|Y}(x|y)W(z|x, y')$$

for all  $y', y, z$ . Consider  $P_{X|Y}((x_1, x_2)|(y_1, y_2)) = 1$  when  $(x_1, x_2) = (y_1, y_2)$ . Then for  $y' = (y'_1, y'_2)$ ,  $y = (y_1, y_2)$  and  $z = (y'_1 + y_1, y'_2 \oplus y_2)$ , both the LHS and the RHS of the above equation evaluate to 1, and for every other  $z$ , they evaluate to 0. So, the channel is B-symmetrizable. Similarly, we can show that the channel is A-symmetrizable.

Next, we show that this channel is non-overwritable. Suppose the channel is B-overwritable. Let  $P_{X'|XY}$  be the overwriting attack by user A which satisfies (45). Then for  $(x, y, z) = ((1, 1), (1, 1), (2, 0))$  and all  $y' = (y'_1, y'_2)$ , (45) implies

$$\sum_{(x'_1, x'_2)} P_{X'|XY}((x'_1, x'_2)|(1, 1), (1, 1))W((2, 0)|(x'_1, x'_2), (y'_1, y'_2)) = W((2, 0)|(1, 1), (1, 1)).$$

However, for  $(y'_1, y'_2) = (0, 0)$ , the LHS evaluates to 0 whereas the RHS evaluates to 1. Hence, the channel is not B-overwritable. Similarly, we can show that the channel is not A-overwritable.

Thus,  $\mathcal{C}_{\text{reliable}} = \{(0, 0)\}$  and  $\mathcal{C}_{\text{auth}} = \mathcal{C}_{\text{MAC}}$ . To compute  $\mathcal{C}$ , we first recall Definition 6 and note that  $\tilde{W}_W$  contains a channel  $\tilde{W}$  satisfying

$$\tilde{W}(x_1 + y_1, v|(x_1, x_2), (y_1, y_2)) = 0.5,$$




Fig. 6. This figure depicts the channel in Example 4. User A has input  $(X_1, X_2)$  and user B has input  $(Y_1, Y_2)$ . The output of the channel is  $(Z_1, Z_2)$ .

for all  $x_1, x_2, y_1, y_2, v \in \{0, 1\}$ . This is obtained by using the pair  $(Q_{X'|X}, Q_{Y'|Y})$  defined by  $Q_{X'|X}((x_1, u)|(x_1, x_2)) = 0.5$  for all  $u, x_1, x_2 \in \{0, 1\}$  and  $Q_{Y'|Y}((y_1, v)|(y_1, y_2)) = 0.5$  for all  $v, y_1, y_2 \in \{0, 1\}$  in (35). We restate (35) below for completeness.

$$\begin{aligned}\tilde{W}(z|x, y) &= \sum_{x'} Q_{X'|X}(x'|x)W(z|x', y) \\ &= \sum_{y'} Q_{Y'|Y}(y'|y)W(z|x, y'),\end{aligned}$$

for all  $x, y, z \in \mathcal{X} \times \mathcal{Y} \times \mathcal{Z}$ .

Note that the channel  $\tilde{W}$  has the same first component as  $W$  (i.e., a binary erasure MAC) and a second component whose output  $Z_2$  is independent of the inputs. Using the argument following Definition 6, we can argue that  $\mathcal{C}$  is outer bounded by the (non-adversarial) capacity region of  $\tilde{W}$  which is the capacity region of the binary erasure MAC. We can show that this outer bound is tight by using an adversary identifying code for the binary erasure MAC component  $Z_1 = X_1 + Y_1$  (see Section V-A) and any arbitrary inputs for the other component. The capacity regions under these three models are plotted in Fig. 7.




Fig. 7. Capacity regions for the byzantine-MAC in Example 4:  $\mathcal{C}_{\text{reliable}} = \{0, 0\}$ ;  $\mathcal{C} = \mathcal{C}_{\text{MAC}}$  of  $Z_1 = X_1 + Y_1$ ; and  $\mathcal{C}_{\text{auth}} = \mathcal{C}_{\text{MAC}}$  of  $(Z_1, Z_2) = (X_1 + Y_1, X_2 \oplus Y_2)$ .

## VI. CONNECTIONS TO RANDOMIZED CODING CAPACITY REGION

While not the focus of this paper, in this section we comment on the relationship of the capacity region  $\mathcal{C}$  of deterministic codes with that of randomized codes. Note that we do not provide any direct achievability scheme for randomized codes in this section. Before presenting the formal definitions, we draw the reader's attention to two points about our setup for randomized codes:

- The encoders share *independent* randomness with the decoder. This is similar to the randomized code of Jahn [11] for AV-MACs.
- When a user is adversarial, we allow that user to adversarially select the realization of its randomness.<sup>7</sup>

Both these choices will prove important in making the connection between the capacity regions under deterministic and randomized codes.

**Definition 7** (Randomized adversary identifying code). An  $(N_A, N_B, L_A, L_B, n)$  randomized adversary identifying code  $(F_A, F_B, \phi_{F_A, F_B})$  for a MAC with byzantine users consists of the following:

- Two message sets,  $\mathcal{M}_i = \{1, \dots, N_i\}$ ,  $i = A, B$ ,
- Two collections of deterministic encoders,  $\Gamma_A = \{f_{A,1}, \dots, f_{A,L_A}\}$ , where  $f_{A,i} : \mathcal{M}_A \rightarrow \mathcal{X}^n$ ,  $i = 1, \dots, L_A$ , and  $\Gamma_B = \{f_{B,1}, \dots, f_{B,L_B}\}$ , where  $f_{B,i} : \mathcal{M}_B \rightarrow \mathcal{Y}^n$ ,  $i = 1, \dots, L_B$ ,
- Two independent randomized encoders  $F_A$  and  $F_B$  with distributions  $p_{F_A}$  and  $p_{F_B}$  over the sets  $\Gamma_A$  and  $\Gamma_B$ , respectively, and
- A collection of decoding maps,  $\phi_{f_A, f_B} : \mathcal{Z}^n \rightarrow (\mathcal{M}_A \times \mathcal{M}_B) \cup \{\mathbf{a}, \mathbf{b}\}$ , where  $f_A \in \Gamma_A$  and  $f_B \in \Gamma_B$ .

As before, the decoder outputs the symbol  $\mathbf{a}$  (resp.,  $\mathbf{b}$ ) to declare that user A (resp., B) is adversarial. The *average probability of error*  $P_e$  is the maximum of the average probabilities of error in the following three cases: (1) both users are honest, (2) user A is adversarial, and (3) user B is adversarial. When both users are honest, the error occurs if the decoder outputs anything other than the pair of correct messages. Let

$$\mathcal{E}_{m_A, m_B}^{f_A, f_B} = \{z : \phi_{f_A, f_B}(z) \neq (m_A, m_B)\}$$

<sup>7</sup>This is analogous to the model studied in [2] for reliable communication.



denote the error event when both users are honest. The average error probability when both users are honest is

$$P_{e,\text{hon}}^{\text{rand}} \stackrel{\text{def}}{=} \frac{1}{N_A \cdot N_B} \sum_{\substack{m_A \in \mathcal{M}_A \\ m_B \in \mathcal{M}_B}} \sum_{\substack{f_A \in \Gamma_A \\ f_B \in \Gamma_B}} p_{F_A}(f_A) p_{F_B}(f_B) W^n(\mathcal{E}_{m_A, m_B}^{f_A, f_B} | f_A(m_A), f_B(m_B)).$$

Recall that when user A is adversarial we would like the decoder's output to either be the symbol  $\mathbf{a}$  or a pair of messages of which the message of user B is correct. The error event is  $\mathcal{E}_{m_B}^{f_A, f_B} \stackrel{\text{def}}{=} \{z : \phi_{f_A, f_B}(z) \notin (\mathcal{M}_A \times \{m_B\}) \cup \{\mathbf{a}\}\}$ . The average probability of error when user A is adversarial is

$$P_{e,\text{mal A}}^{\text{rand}} \stackrel{\text{def}}{=} \max_{\substack{\mathbf{x} \in \mathcal{X}^n \\ f_A \in \Gamma_A}} \left( \frac{1}{N_B} \sum_{m_B \in \mathcal{M}_B} \sum_{f_B \in \Gamma_B} p_{F_B}(f_B) W^n(\mathcal{E}_{m_B}^{f_A, f_B} | \mathbf{x}, f_B(m_B)) \right). \quad (46)$$

Notice that the adversarial user A selects both its transmission  $\mathbf{x}$  and the code  $f_A$  (i.e., the realization of the randomness it shares with the decoder) adversarially in order to maximize the probability that the decoder (working with this code  $f_A$ ) makes an error. As in the deterministic coding case, the probability of error is maximized by deterministic attacks of the adversary. This is because for any attack distribution  $q_{\mathbf{X}, F_A}$ ,

$$\sum_{\mathbf{x}, f_A} q_{\mathbf{X}, F_A}(\mathbf{x}, f_A) \left( \frac{1}{N_B} \sum_{m_B \in \mathcal{M}_B} \sum_{f_B \in \Gamma_B} p_{F_B}(f_B) W^n(\mathcal{E}_{m_B}^{f_A, f_B} | \mathbf{x}, f_B(m_B)) \right) \leq P_{e,\text{mal A}}^{\text{rand}}.$$

Similarly, for  $\mathcal{E}_{m_A}^{f_A, f_B} \stackrel{\text{def}}{=} \{z : \phi_{f_A, f_B}(z) \notin (\{m_A\} \times \mathcal{M}_B) \cup \{\mathbf{b}\}\}$ , the average probability of error when user B is adversarial is

$$P_{e,\text{mal B}}^{\text{rand}} \stackrel{\text{def}}{=} \max_{\substack{\mathbf{y} \in \mathcal{Y}^n \\ f_B \in \Gamma_B}} \left( \frac{1}{N_A} \sum_{m_A \in \mathcal{M}_A} \sum_{f_A \in \Gamma_A} p_{F_A}(f_A) W^n(\mathcal{E}_{m_A}^{f_A, f_B} | f_A(m_A), \mathbf{y}) \right). \quad (47)$$

We define the *average probability of error* as

$$P_e^{\text{rand}} \stackrel{\text{def}}{=} \max \{ P_{e,\text{hon}}^{\text{rand}}, P_{e,\text{mal A}}^{\text{rand}}, P_{e,\text{mal B}}^{\text{rand}} \}.$$

The capacity region  $\mathcal{C}^{\text{rand}}$  of communication with adversary identification under randomized coding may be defined along the lines of Definition 2: we say that a rate  $(R_A, R_B)$  is achievable if there exists a sequence of  $(\lfloor 2^{nR_A} \rfloor, \lfloor 2^{nR_B} \rfloor, L_{A,n}, L_{B,n}, n)$  randomized adversary identifying codes (for some  $L_{A,n}, L_{B,n}$ ) such that  $P_e^{\text{rand}} \rightarrow 0$  as  $n \rightarrow \infty$ . The capacity region  $\mathcal{C}^{\text{rand}}$  is the closure of the set of all such achievable rate pairs.

#### A. Dichotomy theorem for deterministic coding capacity region

The following theorem states that when the deterministic coding capacity region  $\mathcal{C}$  has a non-empty interior, it is the same as the randomized coding capacity region  $\mathcal{C}^{\text{rand}}$ . This is analogous to similar results in the AVC literature [10, Theorem 1], [11, Theorem 1, Section IV].

**Theorem 12.**  $\mathcal{C} = \mathcal{C}^{\text{rand}}$  whenever  $(R_A, R_B) \in \mathcal{C}$  for some  $R_A, R_B > 0$ .

Since deterministic codes are a subset of randomized codes,  $\mathcal{C} \subseteq \mathcal{C}^{\text{rand}}$ . We formally show that  $\mathcal{C} \supseteq \mathcal{C}^{\text{rand}}$  in Appendix D. For this, we first prove that given any randomized adversary identifying code with a small probability of error, there exists another randomized adversary identifying code, also with a small probability of error, which requires only  $O(\log(n))$  shared random bits between each (honest) sender and the receiver (Lemma 16 in Appendix D). This *randomness reduction* argument is along the lines of Ahlswede [10] (and its extension for AV-MAC by Jahn [11]).

With this, one may construct a two-phased code for showing the achievability of Theorem 12. In the first phase of the code, a positive-rate, deterministic adversary identifying code (guaranteed to exist by the hypothesis of the theorem) is used to send  $O(\log(n))$  random bits as messages by each (honest) sender to the receiver. This establishes the  $O(\log(n))$  amount of shared randomness required for the randomized adversary identifying code. In the second phase, the shared randomness based code obtained from the randomness reduction argument is used. Note that an adversarial user can maliciously select their input in the first phase. This is why in (46) and (47) we defined the probability of errors of the randomized adversary identifying code such that the adversarial user may select their own randomness.

### B. An outer bound to the randomized coding capacity region

Let  $\mathcal{C}_{\text{AV-MAC}}^{\text{rand}}(\tilde{\mathcal{W}}_W)$  denote the randomized coding capacity for the AV-MAC  $\tilde{\mathcal{W}}_W$  (see Definition 5). Here, each user shares independent randomness with the receiver which is unknown to the adversary and the other user. This is along the lines of randomized code for AV-MACs (see [11]). Jahn [11] also gives the randomized coding capacity region expression.

**Theorem 13** (Outer bound).  $\mathcal{C}^{\text{rand}} \subseteq \mathcal{C}_{\text{AV-MAC}}^{\text{rand}}(\tilde{\mathcal{W}}_W)$ . Moreover, there exists an AV-MAC  $\mathcal{W}_W$  such that  $\mathcal{C}_{\text{AV-MAC}}^{\text{rand}}(\mathcal{W}_W) = \mathcal{C}_{\text{AV-MAC}}^{\text{rand}}(\tilde{\mathcal{W}}_W)$  and  $|\mathcal{W}_W| \leq 2^{|\mathcal{X}|^2 + |\mathcal{Y}|^2}$ <sup>8</sup>.

**Remark 5.** We will prove the outer bound for a weaker adversary who cannot choose the realization of the randomness it shares with the decoder. Thus, the outer bound holds even if the randomization is provided by the decoder (or an external agent). Note that the adversary knows the realization of its randomness and may choose its input to the channel based on this<sup>9</sup>.

*Proof.* Let  $(F_A, F_B, \phi_{F_A, F_B})$  be an  $(N_A, N_B, L_A, L_B, n)$  randomized adversary identifying code with  $P_e^{\text{rand}} \leq \epsilon$ . For  $i \in [1 : n]$ , let  $(Q_{i, X'|X}, Q_{i, Y'|Y})$  be an arbitrary sequence of pairs of conditional distributions satisfying (35) and define  $\tilde{W}_i$  as

$$\tilde{W}_i(z|x, y) \stackrel{\text{def}}{=} \sum_{x'} Q_{i, X'|X}(x'|x) W(z|x', y) = \sum_{y'} Q_{i, Y'|Y}(y'|y) W(z|x, y'), x \in \mathcal{X}, y \in \mathcal{Y}, z \in \mathcal{Z}. \quad (48)$$

Let  $Q_{X'|X} \stackrel{\text{def}}{=} \prod_{i=1}^n Q_{i, X'|X}$ ,  $Q_{Y'|Y} \stackrel{\text{def}}{=} \prod_{i=1}^n Q_{i, Y'|Y}$  and  $\tilde{W}^{(n)} \stackrel{\text{def}}{=} \prod_{i=1}^n \tilde{W}_i$ .

Let  $\mathcal{E}_{f_A, f_B}^{\text{B}, m_B} := \{z : \phi_{f_A, f_B}(z) \in \{\mathbf{b}\} \cup (\mathcal{M}_A \times (\mathcal{M}_B \setminus \{m_B\}))\}$  and  $\mathcal{E}_{f_A, f_B}^{\text{A}, m_A} := \{z : \phi_{f_A, f_B}(z) \in \{\mathbf{a}\} \cup ((\mathcal{M}_A \setminus \{m_A\}) \times \mathcal{M}_B)\}$ . Consider a malicious user-A who chooses  $M_A$  uniformly from  $\mathcal{M}_A = [1 : N_A]$ , passes  $F_A(M_A)$  over  $Q_{X'|X}$ , and transmits the resulting vector, that is, the user sends a vector distributed as  $Q_{X'|X}(\cdot | F_A(M_A))$  as input to the channel. Note that, as pointed out in Remark 5, the randomness in encoder  $F_A$  is not chosen maliciously by user-A. We may conclude from (46) and (47) that

$$P_{e, \text{mal A}}^{\text{rand}} \geq \frac{1}{N_A \cdot N_B} \sum_{m_A, m_B} \sum_{\mathbf{x}} p_{F_A}(f_A) p_{F_B}(f_B) Q_{X'|X}(\mathbf{x} | f_A(m_A)) W^n \left( \mathcal{E}_{f_A, f_B}^{\text{B}, m_B} \middle| \mathbf{x}, f_B(m_B) \right).$$

Similarly,

$$P_{e, \text{mal B}}^{\text{rand}} \geq \frac{1}{N_A \cdot N_B} \sum_{m_A, m_B} \sum_{\mathbf{y}} p_{F_A}(f_A) p_{F_B}(f_B) Q_{Y'|Y}(\mathbf{y} | f_B(m_B)) W^n \left( \mathcal{E}_{f_A, f_B}^{\text{A}, m_A} \middle| f_A(m_A), \mathbf{y} \right).$$

Using these inequalities,

$$\begin{aligned} 2\epsilon &\geq P_{e, \text{mal A}}^{\text{rand}} + P_{e, \text{mal B}}^{\text{rand}} \geq \frac{1}{N_A \cdot N_B} \sum_{m_A, m_B} \sum_{f_A, f_B} p_{F_A}(f_A) p_{F_B}(f_B) \left( \sum_{\mathbf{x}} Q_{X'|X}(\mathbf{x} | f_A(m_A)) W^n \left( \mathcal{E}_{f_A, f_B}^{\text{B}, m_B} \middle| \mathbf{x}, f_B(m_B) \right) \right. \\ &\quad \left. + \sum_{\mathbf{y}} Q_{Y'|Y}(\mathbf{y} | f_B(m_B)) W^n \left( \mathcal{E}_{f_A, f_B}^{\text{A}, m_A} \middle| f_A(m_A), \mathbf{y} \right) \right) \\ &= \frac{1}{N_A \cdot N_B} \sum_{m_A, m_B} \sum_{f_A, f_B} p_{F_A}(f_A) p_{F_B}(f_B) \left( \tilde{W}^n \left( \mathcal{E}_{f_A, f_B}^{\text{B}, m_B} \middle| f_A(m_A), f_B(m_B) \right) \right. \\ &\quad \left. + \tilde{W}^n \left( \mathcal{E}_{f_A, f_B}^{\text{A}, m_A} \middle| f_A(m_A), f_B(m_B) \right) \right) \\ &\geq \frac{1}{N_A \cdot N_B} \sum_{m_A, m_B} \sum_{f_A, f_B} p_{F_A}(f_A) p_{F_B}(f_B) \tilde{W}^{(n)} \left( \mathcal{E}_{f_A, f_B}^{\text{A}, m_A} \cup \mathcal{E}_{f_A, f_B}^{\text{B}, m_B} \middle| f_A(m_A), f_B(m_B) \right), \end{aligned} \quad (49)$$

where we use (48) and  $\tilde{W}^{(n)} = \prod_{i=1}^n \tilde{W}_i$  in the penultimate step. Notice that for  $(m_A, m_B) \in \mathcal{M}_A \times \mathcal{M}_B$ ,  $(f_A, f_B) \in \Gamma_A \times \Gamma_B$  and  $z \in \mathcal{Z}^n$ ,

$$\begin{aligned} &\{z : \phi_{f_A, f_B}(z) \neq (m_A, m_B)\} \\ &= \{z : \phi_{f_A, f_B}(z) \in \{\mathbf{a}\}\} \cup \{z : \phi_{f_A, f_B}(z) \in \{\mathbf{b}\}\} \cup \{z : \phi_{f_A, f_B}(z) \in (\mathcal{M}_A \times \mathcal{M}_B) \setminus \{(m_A, m_B)\}\} \\ &= \{z : \phi_{f_A, f_B}(z) \in \{\mathbf{b}\} \cup (\mathcal{M}_A \times (\mathcal{M}_B \setminus \{m_B\}))\} \cup \{z : \phi_{f_A, f_B}(z) \in \{\mathbf{a}\} \cup ((\mathcal{M}_A \setminus \{m_A\}) \times \mathcal{M}_B)\} \\ &= \mathcal{E}_{f_A, f_B}^{\text{B}, m_B} \cup \mathcal{E}_{f_A, f_B}^{\text{A}, m_A}. \end{aligned}$$

<sup>8</sup>The capacity region of an AV-MAC only depends on its convex hull which is defined by taking convex combinations of channels under different states [9], [11]. The AV-MAC  $\mathcal{W}_W$  has the same convex hull as  $\tilde{\mathcal{W}}_W$  but with only finitely many states.

<sup>9</sup>This is the intermediate model described in [2, Footnote 10] for the reliable communication problem.

Thus, from (49),

$$\begin{aligned} & \frac{1}{N_A \cdot N_B} \sum_{m_A, m_B} \sum_{f_A, f_B} p_{F_A}(f_A) p_{F_B}(f_B) \tilde{W}^{(n)}(\{z : \phi_{f_A, f_B}(z) \neq (m_A, m_B)\} | f_A(m_A), f_B(m_B)) \\ &= \frac{1}{N_A \cdot N_B} \sum_{m_A, m_B} \sum_{f_A, f_B} p_{F_A}(f_A) p_{F_B}(f_B) \tilde{W}^{(n)}\left(\mathcal{E}_{f_A, f_B}^{A, m_A} \cup \mathcal{E}_{f_A, f_B}^{B, m_B} \mid f_A(m_A), f_B(m_B)\right) \\ &\leq 2\epsilon. \end{aligned}$$

Recall that every pair  $(Q_{X'|X}, Q_{Y'|Y})$  satisfying (35) corresponds to an element in  $\tilde{\mathcal{W}}_W$  which is a convex set (see the discussion in Section IV-B). Thus, for all  $\epsilon > 0$ , any adversary identifying code for the byzantine-MAC  $W$  with probability of error  $\epsilon$  is also a communication code for the AV-MAC  $\tilde{\mathcal{W}}_W$  with probability of error at most  $2\epsilon$ . So, the randomized coding capacity region  $\mathcal{C}^{\text{rand}}$  of  $W$  is contained by the randomized coding capacity region  $\mathcal{C}_{\text{AV-MAC}}^{\text{rand}}(\tilde{\mathcal{W}}_W)$  of the AV-MAC  $\tilde{\mathcal{W}}_W$ .

Since the capacity region of an AV-MAC, both under randomized codes and deterministic codes, only depends on its convex hull [9], [11], we may use a similar argument as the one used in the proof of Theorem 9 to conclude that there exists an AV-MAC  $\mathcal{W}_W$  such that  $\mathcal{C}_{\text{AV-MAC}}^{\text{rand}}(\mathcal{W}_W) = \mathcal{C}_{\text{AV-MAC}}^{\text{rand}}(\tilde{\mathcal{W}}_W)$  and  $|\mathcal{W}_W| \leq 2^{|\mathcal{X}'|^2 + |\mathcal{Y}'|^2}$ .  $\square$

## REFERENCES

- [1] N. Sangwan, M. Bakshi, B. K. Dey, and V. M. Prabhakaran, "Multiple Access Channels with Byzantine Users," in *2019 IEEE Information Theory Workshop (ITW)*, pp. 1–5, 2019.
- [2] N. Sangwan, M. Bakshi, B. K. Dey, and V. M. Prabhakaran, "Byzantine Multiple Access Channels—Part I: Reliable Communication," *arXiv preprint arXiv:2211.12769*, 2022.
- [3] A. El Gamal and Y.-H. Kim, *Network information theory*. Cambridge university press, 2011.
- [4] N. Sangwan, M. Bakshi, B. K. Dey, and V. M. Prabhakaran, "Multiple access channels with adversarial users," in *2019 IEEE International Symposium on Information Theory (ISIT)*, pp. 435–439, IEEE, 2019.
- [5] O. Kosut and J. Kliewer, "Network equivalence for a joint compound-arbitrarily-varying network model," in *2016 IEEE Information Theory Workshop (ITW)*, pp. 141–145, IEEE, 2016.
- [6] O. Kosut and J. Kliewer, "Authentication capacity of adversarial channels," in *2018 IEEE Information Theory Workshop (ITW)*, pp. 1–5, IEEE, 2018.
- [7] D. Blackwell, L. Breiman, and A. Thomasian, "The capacities of certain channel classes under random coding," *The Annals of Mathematical Statistics*, vol. 31, no. 3, pp. 558–567, 1960.
- [8] I. Csiszár and P. Narayan, "The capacity of the arbitrarily varying channel revisited: Positivity, constraints," *IEEE transactions on Information Theory*, vol. 34, no. 2, pp. 181–193, 1988.
- [9] R. Ahlswede and N. Cai, "Arbitrarily varying multiple-access channels. I. Ericson's symmetrizability is adequate, Gubner's conjecture is true," *IEEE Transactions on Information Theory*, vol. 45, no. 2, pp. 742–749, 1999.
- [10] R. Ahlswede, "Elimination of correlation in random codes for arbitrarily varying channels," *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, vol. 44, no. 2, 1978.
- [11] J.-H. Jahn, "Coding of arbitrarily varying multiuser channels," *IEEE Transactions on Information Theory*, vol. 27, no. 2, pp. 212–226, 1981.
- [12] A. Lapidoth and P. Narayan, "Reliable communication under channel uncertainty," *IEEE transactions on Information Theory*, vol. 44, no. 6, pp. 2148–2177, 1998.
- [13] J. A. Gubner, "On the deterministic-code capacity of the multiple-access arbitrarily varying channel," *IEEE transactions on information theory*, vol. 36, no. 2, pp. 262–275, 1990.
- [14] A. Beemer, O. Kosut, J. Kliewer, E. Graves, and P. Yu, "Structured coding for authentication in the presence of a malicious adversary," in *2019 IEEE International Symposium on Information Theory (ISIT)*, pp. 617–621, IEEE, 2019.
- [15] E. Graves, P. Yu, and P. Spasojevic, "Keyless authentication in the presence of a simultaneously transmitting adversary," in *2016 IEEE Information Theory Workshop (ITW)*, pp. 201–205, IEEE, 2016.
- [16] A. Beemer, E. Graves, J. Kliewer, O. Kosut, and P. Yu, "Authentication and partial message correction over adversarial multiple-access channels," in *2020 IEEE Conference on Communications and Network Security (CNS)*, pp. 1–6, IEEE, 2020.
- [17] S. Jaggi, M. Langberg, S. Katti, T. Ho, D. Katabi, and M. Médard, "Resilient network coding in the presence of byzantine adversaries," in *IEEE INFOCOM 2007-26th IEEE international conference on computer communications*, pp. 616–624, IEEE, 2007.
- [18] O. Kosut, L. Tong, and N. David, "Polytope codes against adversaries in networks," *IEEE transactions on information theory*, vol. 60, no. 6, pp. 3308–3344, 2014.
- [19] X. He and A. Yener, "Strong secrecy and reliable byzantine detection in the presence of an untrusted relay," *IEEE transactions on information theory*, vol. 59, no. 1, pp. 177–192, 2012.
- [20] G. J. Simmons, "Authentication theory/coding theory," in *Workshop on the Theory and Application of Cryptographic Techniques*, pp. 411–431, Springer, 1984.
- [21] U. M. Maurer, "Authentication theory and hypothesis testing," *IEEE Transactions on Information Theory*, vol. 46, no. 4, pp. 1350–1356, 2000.
- [22] O. Gungor and C. E. Koksal, "On the basic limits of RF-fingerprint-based authentication," *IEEE transactions on information theory*, vol. 62, no. 8, pp. 4523–4543, 2016.
- [23] L. Lai, H. El Gamal, and H. V. Poor, "Authentication over noisy channels," *IEEE transactions on information theory*, vol. 55, no. 2, pp. 906–916, 2009.
- [24] S. Jiang, "Keyless authentication in a noisy model," *IEEE transactions on information forensics and security*, vol. 9, no. 6, pp. 1024–1033, 2014.
- [25] W. Tu and L. Lai, "Keyless authentication and authenticated capacity," *IEEE Transactions on Information Theory*, vol. 64, no. 5, pp. 3696–3714, 2018.
- [26] U. Pereg and Y. Steinberg, "The capacity region of the arbitrarily varying MAC: with and without constraints," in *2019 IEEE International Symposium on Information Theory (ISIT)*, pp. 445–449, IEEE, 2019.
- [27] I. Csiszár, "The method of types [information theory]," *IEEE Transactions on Information Theory*, vol. 44, no. 6, pp. 2505–2523, 1998.
- [28] I. Csiszár and J. Körner, *Information theory: coding theorems for discrete memoryless systems*. Cambridge University Press, 2011.

## APPENDIX A

### CODEBOOK FOR THEOREMS 1 AND 7

We first restate Lemma 4 by including the analogous statements when roles of users A and B are interchanged.

**Lemma.** Suppose  $\mathcal{X}, \mathcal{Y}, \mathcal{Z}$  are finite. Let  $P_A \in \mathcal{P}_{\mathcal{X}}^n$  and  $P_B \in \mathcal{P}_{\mathcal{Y}}^n$ . For any  $\epsilon > 0$ , there exists  $n_0(\epsilon)$  such that for all  $n \geq n_0(\epsilon)$ ,  $N_A, N_B \geq \exp(n\epsilon)$ , there are codewords  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{N_A}$  of type  $P_A$  and  $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_{N_B}$  of type  $P_B$  such that for all  $\mathbf{x}, \mathbf{x}', \tilde{\mathbf{x}}, \bar{\mathbf{x}} \in \mathcal{X}^n$  and  $\mathbf{y}, \mathbf{y}', \tilde{\mathbf{y}}, \bar{\mathbf{y}} \in \mathcal{Y}^n$ , and joint types  $P_{X\tilde{X}\tilde{Y}Y} \in \mathcal{P}_{\mathcal{X} \times \mathcal{X} \times \mathcal{Y} \times \mathcal{Y}}^n$ ,  $P_{X'\tilde{Y}_1\tilde{Y}_2Y'} \in \mathcal{P}_{\mathcal{X}' \times \mathcal{Y} \times \mathcal{Y} \times \mathcal{Y}}^n$ ,  $P_{\tilde{X}\tilde{X}\tilde{Y}\tilde{Y}} \in \mathcal{P}_{\mathcal{X} \times \mathcal{X} \times \mathcal{Y} \times \mathcal{Y}}^n$  and  $P_{X''\hat{X}_1\hat{X}_2Y''} \in \mathcal{P}_{\mathcal{X}'' \times \mathcal{X} \times \mathcal{X} \times \mathcal{Y}}^n$  and for  $R_A \stackrel{\text{def}}{=} (1/n) \log N_A$  and  $R_B \stackrel{\text{def}}{=} (1/n) \log N_B$ , the following holds:<sup>10</sup>

$$\frac{|\{m_A : (\mathbf{x}_{m_A}, \mathbf{y}) \in T_{X\tilde{X}Y}^n\}|}{N_A} \leq \exp\{-n\epsilon/2\}, \text{ if } I(X; Y) > \epsilon; \quad (14)$$

$$\frac{|\{m_A : (\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{m_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n \text{ for some } \tilde{m}_A \neq m_A \text{ and some } m_B\}|}{N_A} \leq \exp\{-n\epsilon/2\},$$

$$\text{if } I(X; \tilde{X}\tilde{Y}Y) - |R_A - I(\tilde{X}; \tilde{Y}Y)|^+ - |R_B - I(\tilde{Y}; Y)|^+ > \epsilon; \quad (15)$$

$$\frac{|\{(\tilde{m}_A, \tilde{m}_B) : (\mathbf{x}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n\}|}{|\{(\tilde{m}_A, \tilde{m}_B) : (\mathbf{x}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n\}|} \leq \exp\left\{n \left( |R_A - I(\tilde{X}; \tilde{Y}XY)|^+ + |R_B - I(\tilde{Y}; XY)|^+ + \epsilon \right)\right\}; \quad (16)$$

$$\frac{|\{m_A : (\mathbf{x}_{m_A}, \mathbf{y}_{\tilde{m}_{B1}}, \mathbf{y}_{\tilde{m}_{B2}}, \mathbf{y}') \in T_{X'\tilde{Y}_1\tilde{Y}_2Y'}^n \text{ for some } \tilde{m}_{B1}, \tilde{m}_{B2}\}|}{N_A} \leq \exp\{-n\epsilon/2\},$$

$$\text{if } I(X'; \tilde{Y}_1\tilde{Y}_2Y') - |R_B - I(\tilde{Y}_1; Y')|^+ - |R_B - I(\tilde{Y}_2; \tilde{Y}_1Y')|^+ > \epsilon; \quad (17)$$

$$\frac{|\{(\tilde{m}_{B1}, \tilde{m}_{B2}) : (\mathbf{x}', \mathbf{y}_{\tilde{m}_{B1}}, \mathbf{y}_{\tilde{m}_{B2}}, \mathbf{y}') \in T_{X'\tilde{Y}_1\tilde{Y}_2Y'}^n\}|}{|\{(\tilde{m}_{B1}, \tilde{m}_{B2}) : (\mathbf{x}', \mathbf{y}_{\tilde{m}_{B1}}, \mathbf{y}_{\tilde{m}_{B2}}, \mathbf{y}') \in T_{X'\tilde{Y}_1\tilde{Y}_2Y'}^n\}|} \leq \exp\left\{n \left( |R_B - I(\tilde{Y}_1; X'Y')|^+ + |R_B - I(\tilde{Y}_2; \tilde{Y}_1X'Y')|^+ + \epsilon \right)\right\}, \quad (18)$$

$$\frac{|\{m_B : (\bar{\mathbf{x}}, \mathbf{y}_{m_B}) \in T_{\tilde{X}\tilde{Y}}^n\}|}{N_A} \leq \exp\{-n\epsilon/2\}, \text{ if } I(\tilde{X}; \tilde{Y}) > \epsilon; \quad (50)$$

$$\frac{|\{m_B : (\mathbf{x}_{m_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{y}_{m_B}, \bar{\mathbf{x}}) \in T_{\tilde{X}\tilde{Y}\tilde{X}}^n \text{ for some } \tilde{m}_B \neq m_B \text{ and some } m_A\}|}{N_B} \leq \exp\{-n\epsilon/2\},$$

$$\text{if } I(\tilde{Y}; \tilde{Y}\tilde{X}\tilde{X}) - |R_B - I(\tilde{Y}; \tilde{X}\tilde{X})|^+ - |R_A - I(\tilde{X}; \tilde{X})|^+ > \epsilon; \quad (51)$$

$$\frac{|\{(\tilde{m}_A, \tilde{m}_B) : (\bar{\mathbf{y}}, \mathbf{y}_{\tilde{m}_B}, \mathbf{x}_{\tilde{m}_A}, \bar{\mathbf{x}}) \in T_{\tilde{Y}\tilde{X}\tilde{X}}^n\}|}{|\{(\tilde{m}_A, \tilde{m}_B) : (\bar{\mathbf{y}}, \mathbf{y}_{\tilde{m}_B}, \mathbf{x}_{\tilde{m}_A}, \bar{\mathbf{x}}) \in T_{\tilde{Y}\tilde{X}\tilde{X}}^n\}|} \leq \exp\left\{n \left( |R_B - I(\tilde{Y}; \tilde{X}\tilde{X}\tilde{Y})|^+ + |R_A - I(\tilde{X}; \tilde{X}\tilde{Y})|^+ + \epsilon \right)\right\}; \quad (52)$$

$$\frac{|\{m_B : (\mathbf{y}_{m_B}, \mathbf{x}_{\tilde{m}_{A1}}, \mathbf{x}_{\tilde{m}_{A2}}, \tilde{\mathbf{x}}) \in T_{Y''\hat{X}_1\hat{X}_2X''}^n \text{ for some } \tilde{m}_{A1}, \tilde{m}_{A2}\}|}{N_B} \leq \exp\{-n\epsilon/2\},$$

$$\text{if } I(Y''; \hat{X}_1\hat{X}_2X'') - |R_A - I(\hat{X}_1; X'')|^+ - |R_A - I(\hat{X}_2; \hat{X}_1X'')|^+ > \epsilon; \quad (53)$$

$$\text{and } \frac{|\{(\tilde{m}_{A1}, \tilde{m}_{A2}) : (\tilde{\mathbf{y}}, \mathbf{x}_{\tilde{m}_{A1}}, \mathbf{x}_{\tilde{m}_{A2}}, \tilde{\mathbf{x}}) \in T_{Y''\hat{X}_1\hat{X}_2X''}^n\}|}{|\{(\tilde{m}_{A1}, \tilde{m}_{A2}) : (\tilde{\mathbf{y}}, \mathbf{x}_{\tilde{m}_{A1}}, \mathbf{x}_{\tilde{m}_{A2}}, \tilde{\mathbf{x}}) \in T_{Y''\hat{X}_1\hat{X}_2X''}^n\}|} \leq \exp\left\{n \left( |R_A - I(\hat{X}_1; X''Y'')|^+ + |R_A - I(\hat{X}_2; \hat{X}_1X''Y'')|^+ + \epsilon \right)\right\}. \quad (54)$$

Here, statements (50)-(54) are analogous to (14)-(18) with the roles of users A and B are interchanged. In particular,  $(\mathbf{x}, \mathbf{x}', \mathbf{y}, \mathbf{y}')$  is replaced with  $(\bar{\mathbf{y}}, \tilde{\mathbf{y}}, \bar{\mathbf{x}}, \tilde{\mathbf{x}})$ ,  $(X, \tilde{X}, \tilde{Y}, Y)$  with  $(\tilde{Y}, \tilde{Y}, \hat{X}, \tilde{X})$ ,  $(X', \tilde{Y}_1, \tilde{Y}_2, Y')$  with  $(Y'', \hat{X}_1, \hat{X}_2, X'')$  and  $R_A$  with  $R_B$ .

*Proof.* This proof is along the lines of the proof of [8, Lemma 3]. We will generate the codebook by a random experiment. For fixed vectors  $\mathbf{x}, \mathbf{x}', \tilde{\mathbf{x}}, \bar{\mathbf{x}}, \mathbf{y}, \mathbf{y}', \tilde{\mathbf{y}}, \bar{\mathbf{y}}$  and joint types  $P_{X\tilde{X}\tilde{Y}Y}$ ,  $P_{X'\tilde{Y}_1\tilde{Y}_2Y'}$ ,  $P_{\tilde{X}\tilde{X}\tilde{Y}\tilde{Y}}$  and  $P_{X''\hat{X}_1\hat{X}_2Y''}$  satisfying the conditions of the lemma, we will show that for each of the statements (14)-(18) and (50)-(54), the probability that the statement does not hold falls doubly exponentially in  $n$ . Since,  $|\mathcal{X}^n|$ ,  $|\mathcal{Y}^n|$ ,  $|\mathcal{P}_{\mathcal{X} \times \mathcal{X} \times \mathcal{Y} \times \mathcal{Y}}^n|$ ,  $|\mathcal{P}_{\mathcal{X}' \times \mathcal{Y} \times \mathcal{Y} \times \mathcal{Y}}^n|$  and  $|\mathcal{P}_{\mathcal{X} \times \mathcal{X} \times \mathcal{Y} \times \mathcal{Y}}^n|$  grow at most exponentially in  $n$ , a union bound will imply that the probability that any of the statements (14)-(18) and (50)-(54) fail for some  $\mathbf{x}, \mathbf{x}', \tilde{\mathbf{x}}, \bar{\mathbf{x}}, \mathbf{y}, \mathbf{y}', \tilde{\mathbf{y}}, \bar{\mathbf{y}}$ ,  $P_{X\tilde{X}\tilde{Y}Y}$ ,  $P_{X'\tilde{Y}_1\tilde{Y}_2Y'}$ ,  $P_{\tilde{X}\tilde{X}\tilde{Y}\tilde{Y}}$  and  $P_{X''\hat{X}_1\hat{X}_2Y''}$  also falls doubly exponentially. This will show the existence of a codebook as required by the lemma. The proof will employ [8, Lemma A1], which is reproduced below.

<sup>10</sup>Note that exp and log are with respect to base 2.

**Lemma 14.** [8, Lemma A1] Let  $Z_1, \dots, Z_N$  be arbitrary random variables, and let  $f_i(Z_1, \dots, Z_i)$  be arbitrary with  $0 \leq f_i \leq 1$ ,  $i = 1, \dots, N$ . Then the condition

$$E[f_i(Z_1, \dots, Z_i) | Z_1, \dots, Z_{i-1}] \leq a \text{ a.s.}, \quad i = 1, \dots, N, \quad (55)$$

implies that

$$\mathbb{P} \left\{ \frac{1}{N} \sum_{i=1}^N f_i(Z_1, \dots, Z_i) > t \right\} \leq \exp \{-N(t - a \log e)\}. \quad (56)$$

We denote the type classes of  $P_A$  and  $P_B$  by  $T_A^n$  and  $T_B^n$  respectively. Let  $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_{N_A}$  be independent random vectors each uniformly distributed on  $T_A^n$  and  $\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_{N_B}$  be another set of independent random vectors (independent of  $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_{N_A}$ ) with each element uniformly distributed on  $T_B^n$ .  $(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_{N_A})$  and  $(\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_{N_B})$  are the random codebooks for user A and B respectively. Fix  $P_{X\tilde{X}\tilde{Y}Y} \in \mathcal{P}_{\mathcal{X} \times \mathcal{X} \times \mathcal{Y} \times \mathcal{Y}}^n$ ,  $P_{X'\tilde{Y}_1\tilde{Y}_2Y'} \in \mathcal{P}_{\mathcal{X} \times \mathcal{Y} \times \mathcal{Y} \times \mathcal{Y}}^n$ ,  $P_{\tilde{X}\tilde{X}\tilde{Y}\tilde{Y}} \in \mathcal{P}_{\mathcal{X} \times \mathcal{X} \times \mathcal{Y} \times \mathcal{Y}}^n$ ,  $P_{X''\hat{X}_1\hat{X}_2Y''} \in \mathcal{P}_{\mathcal{X} \times \mathcal{X} \times \mathcal{X} \times \mathcal{Y}}^n$ , such that  $P_X = P_{\tilde{X}} = P_{X'} = P_{\hat{X}} = P_{\hat{X}_1} = P_{\hat{X}_2} = P_A$  and  $P_{\tilde{Y}} = P_{\tilde{Y}_1} = P_{\tilde{Y}_2} = P_{\tilde{Y}'} = P_{\hat{Y}} = P_{Y''} = P_B$ , and  $(\mathbf{x}, \mathbf{y}) \in T_{\tilde{X}\tilde{Y}}^n$ ,  $(\mathbf{x}', \mathbf{y}') \in T_{X'Y'}^n$ ,  $(\tilde{\mathbf{x}}, \tilde{\mathbf{y}}) \in T_{\tilde{X}\tilde{Y}}^n$  and  $(\tilde{\mathbf{x}}, \tilde{\mathbf{y}}) \in T_{\tilde{X}\tilde{Y}}^n$ <sup>11</sup>. We will often use  $(\mathbf{x}_1, \dots, \mathbf{x}_{N_A})$  and  $(\mathbf{y}_1, \dots, \mathbf{y}_{N_B})$  as placeholders to denote the realizations of for  $(\mathbf{X}_1, \dots, \mathbf{X}_{N_A})$  and  $(\mathbf{Y}_1, \dots, \mathbf{Y}_{N_B})$  respectively (for example, see (57)).

Analysis of (16)

For  $i \in [1 : n]$ , define

$$g_i(\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_i) \stackrel{\text{def}}{=} \begin{cases} 1, & \text{if } \mathbf{y}_i \in T_{\tilde{Y}|XY}^n(\mathbf{x}, \mathbf{y}) \\ 0, & \text{otherwise,} \end{cases} \quad (57)$$

and for  $\tilde{\mathbf{y}} \in T_{\tilde{Y}|XY}^n(\mathbf{x}, \mathbf{y})$ ,

$$h_i^{\tilde{\mathbf{y}}}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_i) \stackrel{\text{def}}{=} \begin{cases} 1, & \text{if } \mathbf{x}_i \in T_{\tilde{X}|\tilde{Y}XY}^n(\tilde{\mathbf{y}}, \mathbf{x}, \mathbf{y}) \\ 0, & \text{otherwise.} \end{cases} \quad (58)$$

Let events  $\mathcal{E}$ ,  $\mathcal{E}_1$  and  $\mathcal{E}_2^{\tilde{\mathbf{y}}}$  be defined as

$$\begin{aligned} \mathcal{E} &\stackrel{\text{def}}{=} \left\{ \left| \{(\tilde{m}_A, \tilde{m}_B) : (\mathbf{x}, \mathbf{X}_{\tilde{m}_A}, \mathbf{Y}_{\tilde{m}_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n\} \right| \right. \\ &\quad \left. > \exp \left\{ n \left( |R_A - I(\tilde{X}; \tilde{Y}XY)|^+ + |R_B - I(\tilde{Y}; XY)|^+ + \epsilon \right) \right\} \right\}, \\ \mathcal{E}_1 &\stackrel{\text{def}}{=} \left\{ \sum_{i=1}^{N_B} g_i(\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_i) > \exp \left\{ n \left( |R_B - I(\tilde{Y}; XY)|^+ + \frac{\epsilon}{2} \right) \right\} \right\}, \text{ and} \\ \text{for } \tilde{\mathbf{y}} \in T_{\tilde{Y}|XY}^n(\mathbf{x}, \mathbf{y}), \quad \mathcal{E}_2^{\tilde{\mathbf{y}}} &\stackrel{\text{def}}{=} \left\{ \sum_{j=1}^{N_A} h_j^{\tilde{\mathbf{y}}}(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_j) > \exp \left\{ n \left( |R_A - I(\tilde{X}; \tilde{Y}XY)|^+ + \frac{\epsilon}{2} \right) \right\} \right\}. \end{aligned}$$

We note that

$$\begin{aligned} &\left| \{(\tilde{m}_A, \tilde{m}_B) : (\mathbf{x}, \mathbf{X}_{\tilde{m}_A}, \mathbf{Y}_{\tilde{m}_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n\} \right| \\ &= \sum_{i=1}^{N_B} g_i(\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_i) \left( \sum_{j=1}^{N_A} h_j^{\mathbf{Y}_i}(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_j) \right). \end{aligned}$$

Thus,

$$\mathcal{E} \subseteq \left( \bigcup_{\tilde{\mathbf{y}} \in T_{\tilde{Y}|XY}^n(\mathbf{x}, \mathbf{y})} \mathcal{E}_2^{\tilde{\mathbf{y}}} \right) \cup \mathcal{E}_1. \quad (59)$$

In order to apply Lemma 14 to (57) with  $(\mathbf{Y}_1, \dots, \mathbf{Y}_{N_B})$  as the random variables  $(Z_1, \dots, Z_N)$ , we note that

$$\begin{aligned} E[g_i(\mathbf{Y}_1, \dots, \mathbf{Y}_i) | \mathbf{Y}_1, \dots, \mathbf{Y}_{i-1}] &= \mathbb{P} \left\{ \mathbf{Y}_i \in T_{\tilde{Y}|XY}^n(\mathbf{x}, \mathbf{y}) \right\} \\ &= \frac{|T_{\tilde{Y}|XY}^n(\mathbf{x}, \mathbf{y})|}{|T_B^n|} \\ &\stackrel{(a)}{\leq} \frac{\exp \left( nH(\tilde{Y}|XY) \right)}{(n+1)^{-|\mathcal{Y}|} \exp \left( nH(\tilde{Y}) \right)} \end{aligned}$$

<sup>11</sup>Note that the lemma statements hold trivially if these conditions do not hold. For example, (14) holds trivially if  $\mathbf{y} \notin T_{\tilde{Y}}^n$ .

$$=(n+1)^{|\mathcal{Y}|} \exp\left(-nI(\tilde{Y}; XY)\right),$$

where (a) follows from (21), (22) and because  $P_B = P_{\tilde{Y}}$  (which implies  $|T_B^n| = |T_{\tilde{Y}}^n|$ ). Taking  $t = \frac{1}{N_B} \exp\left\{n\left(|R_B - I(\tilde{Y}; XY)|^+ + \frac{\epsilon}{2}\right)\right\}$  and  $n \geq n_1(\epsilon)$ , where  $n_1(\epsilon) \stackrel{\text{def}}{=} \min\left\{n : (n+1)^{|\mathcal{Y}|} \log e < \frac{1}{2} \exp\left(\frac{n\epsilon}{2}\right)\right\}$ , we see that  $N_B(t - a \log e) \geq (1/2) \exp\left(\frac{n\epsilon}{2}\right)$ . Using (56), this gives us

$$\mathbb{P}(\mathcal{E}_1) \leq \exp\left\{-\frac{1}{2} \exp\left\{\frac{n\epsilon}{2}\right\}\right\}. \quad (60)$$

Similarly, we apply Lemma 14 to (58) with  $(\mathbf{X}_1, \dots, \mathbf{X}_{N_A})$  as the random variables  $(Z_1, \dots, Z_N)$ . We can show that  $a = (n+1)^{|\mathcal{X}|} \exp\left(-nI(\tilde{X}; \tilde{Y}XY)\right)$  satisfies (55). We take  $t = \frac{1}{N_A} \exp\left\{n\left(|R_A - I(\tilde{X}; \tilde{Y}XY)|^+ + \frac{\epsilon}{2}\right)\right\}$  and  $n \geq n_2(\epsilon)$  where  $n_2(\epsilon) \stackrel{\text{def}}{=} \min\left\{n : (n+1)^{|\mathcal{X}|} \log e < \frac{1}{2} \exp\left(\frac{n\epsilon}{2}\right)\right\}$ . This gives  $N_A(t - a \log e) \geq (1/2) \exp\left(\frac{n\epsilon}{2}\right)$  which, when plugged into (56), gives

$$\mathbb{P}(\mathcal{E}_2^{\tilde{y}}) \leq \exp\left\{-\frac{1}{2} \exp\left\{\frac{n\epsilon}{2}\right\}\right\}. \quad (61)$$

Using (59), (60) and (61),

$$\mathbb{P}(\mathcal{E}) \leq \left(|T_{\tilde{Y}|XY}^n(\mathbf{x}, \mathbf{y})| + 1\right) \exp\left\{-\frac{1}{2} \exp\left\{\frac{n\epsilon}{2}\right\}\right\}. \quad (62)$$

This shows that the probability that (16) does not hold falls doubly exponentially.

#### Analysis of (14)

We will use the same arguments as used in obtaining (61). We replace  $\tilde{X}$  with  $X$  and  $(\tilde{Y}, X, Y)$  with  $Y$ , to obtain

$$\mathbb{P}\left\{|\{m_A : (\mathbf{X}_{m_A}, \mathbf{y}) \in T_{XY}^n\}| > \exp\left\{n\left(|R_A - I(X; Y)|^+ + \frac{\epsilon}{2}\right)\right\}\right\} \leq \exp\left\{-\frac{1}{2} \exp\left\{\frac{n\epsilon}{2}\right\}\right\}. \quad (63)$$

So,

$$\mathbb{P}\left\{\frac{1}{N_A} |\{m_A : (\mathbf{X}_{m_A}, \mathbf{y}) \in T_{XY}^n\}| > \exp\left\{n\left(|R_A - I(X; Y)|^+ - R_A + \frac{\epsilon}{2}\right)\right\}\right\} \leq \exp\left\{-\frac{1}{2} \exp\left\{\frac{n\epsilon}{2}\right\}\right\}.$$

We are given that  $I(X; Y) > \epsilon$ . When  $R_A > I(X; Y)$ , we have  $|R_A - I(X; Y)|^+ - R_A + \frac{\epsilon}{2} = \frac{\epsilon}{2} - I(X; Y) \leq -\frac{\epsilon}{2}$ . When  $R_A \leq I(X; Y)$ , we have  $|R_A - I(X; Y)|^+ - R_A + \frac{\epsilon}{2} = \frac{\epsilon}{2} - R_A \leq -\frac{\epsilon}{2}$  (because  $R_A \geq \epsilon$ ). Thus,  $|R_A - I(X; Y)|^+ - R_A + \frac{\epsilon}{2} \leq -\frac{\epsilon}{2}$  and

$$\begin{aligned} & \mathbb{P}\left\{\frac{1}{N_A} |\{m_A : (\mathbf{X}_{m_A}, \mathbf{y}) \in T_{XY}^n\}| > \exp\left\{\frac{-n\epsilon}{2}\right\}\right\} \\ & \leq \mathbb{P}\left\{\frac{1}{N_A} |\{m_A : (\mathbf{X}_{m_A}, \mathbf{y}) \in T_{XY}^n\}| > \exp\left\{n\left(|R_A - I(X; Y)|^+ - R_A + \frac{\epsilon}{2}\right)\right\}\right\} \\ & \leq \exp\left\{-\frac{1}{2} \exp\left\{\frac{n\epsilon}{2}\right\}\right\}. \end{aligned}$$

#### Analysis of (15)

For  $i \in [1 : N_A]$ , let  $A_i$  be the set of indices  $(j, k) \in [1 : N_A] \times [1 : N_B]$ ,  $j < i$  such that  $(\mathbf{x}_j, \mathbf{y}_k) \in T_{\tilde{X}\tilde{Y}|Y}^n(\mathbf{y})$  provided  $|A_i| \leq \exp\left\{n\left(|R_A - I(\tilde{X}; \tilde{Y}Y)|^+ + |R_B - I(\tilde{Y}; Y)|^+\right) + \frac{\epsilon}{4}\right\}$ . Otherwise,  $A_i = \emptyset$ . That is,

$$A_i \stackrel{\text{def}}{=} \begin{cases} \{(j, k) \in [1 : N_A] \times [1 : N_B] : j < i, (\mathbf{x}_j, \mathbf{y}_k) \in T_{\tilde{X}\tilde{Y}|Y}^n(\mathbf{y})\}, & \text{if } |A_i| \leq \exp\left\{n\left(|R_A - I(\tilde{X}; \tilde{Y}Y)|^+ + |R_B - I(\tilde{Y}; Y)|^+\right) + \frac{\epsilon}{4}\right\}, \\ \emptyset, & \text{otherwise.} \end{cases}$$

Let, for  $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_{N_B} \in \mathcal{Y}^n$ ,

$$f_i^{\{\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_{N_B}\}}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_i) \stackrel{\text{def}}{=} \begin{cases} 1, & \text{if } \mathbf{x}_i \in \cup_{(j,k) \in A_i} T_{\tilde{X}\tilde{Y}|Y}^n(\mathbf{x}_j, \mathbf{y}_k, \mathbf{y}) \\ 0, & \text{otherwise.} \end{cases}$$

Then,

$$\begin{aligned} & \mathbb{P}\left\{\sum_{i=1}^{N_A} f_i^{\{\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_{N_B}\}}(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_i) \neq \left|\left\{i : \mathbf{X}_i \in T_{\tilde{X}\tilde{Y}|Y}^n(\mathbf{X}_j, \mathbf{Y}_k, \mathbf{y}) \text{ for some } j < i \text{ and some } k\right\}\right|\right\} \\ & = \mathbb{P}\left\{|\{(\tilde{m}_A, \tilde{m}_B) : (\mathbf{X}_{\tilde{m}_A}, \mathbf{Y}_{\tilde{m}_B}, \mathbf{y}) \in T_{\tilde{X}\tilde{Y}|Y}^n\}| > \exp\left\{n\left(|R_A - I(\tilde{X}; \tilde{Y}Y)|^+ + |R_B - I(\tilde{Y}; Y)|^+\right) + \frac{\epsilon}{4}\right\}\right\} \end{aligned}$$



$$\leq \left( |T_{\tilde{Y}|Y}(\mathbf{y})| + 1 \right) \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon}{8} \right\} \right\}, \quad (64)$$

where the last inequality can be obtained from the definition of event  $\mathcal{E}$  and (62) where we replace  $(X, Y)$  with  $Y$ ,  $(\mathbf{x}, \mathbf{y})$  with  $\mathbf{y}$ , and  $\epsilon$  with  $\epsilon/4$ .

For  $\mathbf{y}_i \in T_{\mathbf{B}}^n$ ,  $i = 1, \dots, N_{\mathbf{B}}$ , we will now apply Lemma 14 on  $f_i^{[\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_{N_{\mathbf{B}}}]}$  with  $(\mathbf{X}_1, \dots, \mathbf{X}_{N_{\mathbf{A}}})$  as the random variables  $(Z_1, \dots, Z_N)$ . We will first compute the value of  $a$  in (55). We note that, for  $i \in [1 : N_{\mathbf{A}}]$ ,  $E \left[ f_i^{[\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_{N_{\mathbf{B}}}] }(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_i) \middle| \mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_{i-1} \right]$ , being a random function of  $(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_{i-1})$ , is a random variable. We will compute it for  $(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_{i-1}) = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{i-1})$ .

$$\begin{aligned} & E \left[ f_i^{[\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_{N_{\mathbf{B}}}] }(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_i) \middle| (\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_{i-1}) = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{i-1}) \right] \\ &= \mathbb{P} \left( \mathbf{X}_i \in \cup_{(j,k) \in A_i} T_{X|\tilde{X}\tilde{Y}Y}^n(\mathbf{x}_j, \mathbf{y}_k, \mathbf{y}) \right) \\ &\stackrel{(a)}{\leq} |A_i| \frac{\exp \left\{ nH(X|\tilde{X}\tilde{Y}Y) \right\}}{(n+1)^{-|\mathcal{X}|} \exp(nH(\tilde{X}))} \\ &= (n+1)^{|\mathcal{X}|} \exp \left\{ n \left( |R_{\mathbf{A}} - I(\tilde{X}; \tilde{Y}Y)|^+ + |R_{\mathbf{B}} - I(\tilde{Y}; Y)|^+ \right) - I(X; \tilde{X}\tilde{Y}Y) + \frac{\epsilon}{4} \right\}, \end{aligned}$$

where (a) follows from (21), (22), a union bound over  $(j, k) \in A_i$  and by noting that  $|T_{\mathbf{A}}^n| = |T_{\tilde{X}}^n|$ . For all  $i \in [1 : N_{\mathbf{A}}]$ , this upper bound holds for every realization of  $(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_{i-1})$ . Thus, in (55), we may take  $a = (n+1)^{|\mathcal{X}|} \exp \left\{ n \left( |R_{\mathbf{A}} - I(\tilde{X}; \tilde{Y}Y)|^+ + |R_{\mathbf{B}} - I(\tilde{Y}; Y)|^+ \right) - I(X; \tilde{X}\tilde{Y}Y) + \frac{\epsilon}{4} \right\}$ . If  $I(X; \tilde{X}\tilde{Y}Y) > |R_{\mathbf{A}} - I(\tilde{X}; \tilde{Y}Y)|^+ + |R_{\mathbf{B}} - I(\tilde{Y}; Y)|^+ + \epsilon$  (as postulated in (15)), (55) holds with  $a = (n+1)^{|\mathcal{X}|} \exp \left\{ -\frac{3}{4}n\epsilon \right\}$ . For  $t = \exp \left\{ \frac{-n\epsilon}{2} \right\}$  and  $n \geq n_2(\epsilon)$  with  $n_2(\epsilon) \stackrel{\text{def}}{=} \min \left\{ n : (n+1)^{|\mathcal{X}|} \log e < \frac{1}{2} \exp \left\{ \frac{n\epsilon}{4} \right\} \right\}$ , we get

$$\begin{aligned} & \mathbb{P} \left\{ \frac{1}{N_{\mathbf{A}}} \sum_{i=1}^{N_{\mathbf{A}}} f_i^{[\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_{N_{\mathbf{B}}}] }(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_i) > \exp \left\{ \frac{-n\epsilon}{2} \right\} \right\} \\ & \leq \exp \left\{ -\frac{N_{\mathbf{A}}}{2} \exp \left\{ -\frac{n\epsilon}{2} \right\} \right\} \\ & \leq \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon}{2} \right\} \right\}, \end{aligned}$$

where the last inequality uses the assumption that  $N_{\mathbf{A}} \geq \exp \{n\epsilon\}$ . Averaging over  $(\mathbf{Y}_1, \dots, \mathbf{Y}_{N_{\mathbf{B}}})$ , we get

$$\begin{aligned} & \mathbb{P} \left\{ \frac{1}{N_{\mathbf{A}}} \sum_{i=1}^{N_{\mathbf{A}}} f_i^{[\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_{N_{\mathbf{B}}}] }(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_i) > \exp \left\{ \frac{-n\epsilon}{2} \right\} \right\} \\ & \leq \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon}{2} \right\} \right\}. \end{aligned} \quad (65)$$

Let events  $\mathcal{F}_1$  and  $\mathcal{F}_2$  be defined as

$$\begin{aligned} \mathcal{F}_1 &\stackrel{\text{def}}{=} \left\{ \frac{1}{N_{\mathbf{A}}} \left| \left\{ i : \mathbf{X}_i \in T_{X|\tilde{X}\tilde{Y}Y}^n(\mathbf{X}_j, \mathbf{Y}_k, \mathbf{y}) \text{ for some } j < i \text{ and some } k \right\} \right| > \exp \left\{ \frac{-n\epsilon}{2} \right\} \right\}, \\ \mathcal{F}_2 &\stackrel{\text{def}}{=} \left\{ \sum_{i=1}^{N_{\mathbf{A}}} f_i^{[\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_{N_{\mathbf{B}}}] }(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_i) \neq \left| \left\{ i : \mathbf{X}_i \in T_{X|\tilde{X}\tilde{Y}Y}^n(\mathbf{X}_j, \mathbf{Y}_k, \mathbf{y}) \text{ for some } j < i \text{ and some } k \right\} \right| \right\}, \\ \mathcal{F}_3 &\stackrel{\text{def}}{=} \left\{ \sum_{i=1}^{N_{\mathbf{A}}} f_i^{[\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_{N_{\mathbf{B}}}] }(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_i) > \exp \left\{ \frac{-n\epsilon}{2} \right\} \right\}. \end{aligned}$$

We are interested in  $\mathbb{P}(\mathcal{F}_1)$ . We see that

$$\begin{aligned} \mathbb{P}(\mathcal{F}_1) &= \mathbb{P}(\mathcal{F}_1 \cap \mathcal{F}_2) + \mathbb{P}(\mathcal{F}_1 \cap \mathcal{F}_2^c) \\ &\leq \mathbb{P}(\mathcal{F}_2) + \mathbb{P}(\mathcal{F}_1 \cap \mathcal{F}_2^c) \\ &\leq \mathbb{P}(\mathcal{F}_2) + \mathbb{P}(\mathcal{F}_3) \\ &\stackrel{(a)}{\leq} \left( |T_{\tilde{Y}|Y}(\mathbf{y})| + 1 \right) \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon}{8} \right\} \right\} + \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon}{2} \right\} \right\} \\ &\leq \left( |T_{\tilde{Y}|Y}(\mathbf{y})| + 2 \right) \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon}{8} \right\} \right\}, \end{aligned}$$

where (a) follows from (64) and (65). Thus,

$$\begin{aligned} & \mathbb{P} \left( \frac{1}{N_A} \left| \left\{ i : \mathbf{X}_i \in T_{X|\tilde{X}\tilde{Y}Y}^n(\mathbf{X}_j, \mathbf{Y}_k, \mathbf{y}) \text{ for some } j < i \text{ and } k \right\} \right| > \exp \left\{ \frac{-n\epsilon}{2} \right\} \right) \\ & \leq \left( |T_{\tilde{Y}|Y}(\mathbf{y})| + 2 \right) \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon}{8} \right\} \right\}. \end{aligned}$$

By symmetry, we get the same upper bound when  $j > i$ . Thus,

$$\begin{aligned} & \mathbb{P} \left\{ \left| \frac{\left\{ m_A : (\mathbf{X}_{m_A}, \mathbf{X}_{\tilde{m}_A}, \mathbf{Y}_{m_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n \text{ for some } \tilde{m}_A \neq m_A \text{ and some } m_B \right\}}{N_A} \right| > \exp \{-n\epsilon/2\} \right\} \\ & < 2 \left( |T_{\tilde{Y}|Y}(\mathbf{y})| + 2 \right) \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon}{8} \right\} \right\}. \end{aligned}$$

This completes the analysis for (15).

#### Analysis of (18)

We will split the analysis in two parts as suggested by the inequalities below, where the first inequality is a union bound,

$$\begin{aligned} & \mathbb{P} \left\{ \left| \left\{ (\tilde{m}_{B1}, \tilde{m}_{B2}) : (\mathbf{x}', \mathbf{Y}_{\tilde{m}_{B1}}, \mathbf{Y}_{\tilde{m}_{B2}}, \mathbf{y}') \in T_{X'\tilde{Y}_1\tilde{Y}_2Y'}^n \right\} \right| > \exp \left\{ n \left( |R_B - I(\tilde{Y}_1; X'Y')|^+ + |R_B - I(\tilde{Y}_2; \tilde{Y}_1 X'Y')|^+ + \epsilon \right) \right\} \right\} \\ & \leq \mathbb{P} \left\{ \left| \left\{ (\tilde{m}_{B1}, \tilde{m}_{B2}) : \tilde{m}_{B1} \neq \tilde{m}_{B2}, (\mathbf{x}', \mathbf{Y}_{\tilde{m}_{B1}}, \mathbf{Y}_{\tilde{m}_{B2}}, \mathbf{y}') \in T_{X'\tilde{Y}_1\tilde{Y}_2Y'}^n \right\} \right| \right. \\ & \quad \left. > \frac{1}{2} \exp \left\{ n \left( |R_B - I(\tilde{Y}_1; X'Y')|^+ + |R_B - I(\tilde{Y}_2; \tilde{Y}_1 X'Y')|^+ + \epsilon \right) \right\} \right\} \\ & + \mathbb{P} \left\{ \left| \left\{ (\tilde{m}_{B1}, \tilde{m}_{B2}) : \tilde{m}_{B1} = \tilde{m}_{B2}, (\mathbf{x}', \mathbf{Y}_{\tilde{m}_{B1}}, \mathbf{Y}_{\tilde{m}_{B2}}, \mathbf{y}') \in T_{X'\tilde{Y}_1\tilde{Y}_2Y'}^n \right\} \right| \right. \\ & \quad \left. > \frac{1}{2} \exp \left\{ n \left( |R_B - I(\tilde{Y}_1; X'Y')|^+ + |R_B - I(\tilde{Y}_2; \tilde{Y}_1 X'Y')|^+ + \epsilon \right) \right\} \right\} \\ & \leq \mathbb{P} \left\{ \left| \left\{ (\tilde{m}_{B1}, \tilde{m}_{B2}) : \tilde{m}_{B1} \neq \tilde{m}_{B2}, (\mathbf{x}', \mathbf{Y}_{\tilde{m}_{B1}}, \mathbf{Y}_{\tilde{m}_{B2}}, \mathbf{y}') \in T_{X'\tilde{Y}_1\tilde{Y}_2Y'}^n \right\} \right| \right. \\ & \quad \left. > \exp \left\{ n \left( |R_B - I(\tilde{Y}_1; X'Y')|^+ + |R_B - I(\tilde{Y}_2; \tilde{Y}_1 X'Y')|^+ + \epsilon' \right) \right\} \right\} \\ & + \mathbb{P} \left\{ \left| \left\{ (\tilde{m}_{B1}, \tilde{m}_{B2}) : \tilde{m}_{B1} = \tilde{m}_{B2}, (\mathbf{x}', \mathbf{Y}_{\tilde{m}_{B1}}, \mathbf{Y}_{\tilde{m}_{B2}}, \mathbf{y}') \in T_{X'\tilde{Y}_1\tilde{Y}_2Y'}^n \right\} \right| \right. \\ & \quad \left. > \exp \left\{ n \left( |R_B - I(\tilde{Y}_1; X'Y')|^+ + |R_B - I(\tilde{Y}_2; \tilde{Y}_1 X'Y')|^+ + \epsilon' \right) \right\} \right\} \end{aligned}$$

for  $\epsilon' = \epsilon/2$  and sufficiently large  $n$ . We first consider the first term where  $\tilde{m}_{B1} \neq \tilde{m}_{B2}$ . We follow arguments similar to those for (16) and get an upper bound. We define

$$\tilde{g}_i(\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_i) \stackrel{\text{def}}{=} \begin{cases} 1, & \text{if } \mathbf{y}_i \in T_{\tilde{Y}_1|X'Y'}^n(\mathbf{x}', \mathbf{y}') \\ 0, & \text{otherwise.} \end{cases} \quad (66)$$

For  $\tilde{\mathbf{y}} \in T_{\tilde{Y}_1|X'Y'}^n(\mathbf{x}', \mathbf{y}')$ ,

$$\tilde{h}_i^{\tilde{\mathbf{y}}}(\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_i) \stackrel{\text{def}}{=} \begin{cases} 1, & \text{if } \mathbf{y}_i \in T_{\tilde{Y}_2|\tilde{Y}_1 X'Y'}^n(\tilde{\mathbf{y}}, \mathbf{x}', \mathbf{y}') \\ 0, & \text{otherwise.} \end{cases} \quad (67)$$

Define events  $\tilde{\mathcal{E}}$  and  $\tilde{\mathcal{E}}_1$  as

$$\begin{aligned} \tilde{\mathcal{E}} &= \left\{ \left| \left\{ (\tilde{m}_{B1}, \tilde{m}_{B2}) : \tilde{m}_{B1} \neq \tilde{m}_{B2}, (\mathbf{x}', \mathbf{Y}_{\tilde{m}_{B1}}, \mathbf{Y}_{\tilde{m}_{B2}}, \mathbf{y}') \in T_{X'\tilde{Y}_1\tilde{Y}_2Y'}^n \right\} \right| \right. \\ & \quad \left. > \exp \left\{ n \left( |R_B - I(\tilde{Y}_2; X'Y'\tilde{Y}_1)|^+ + |R_B - I(\tilde{Y}_1; X'Y')|^+ + \epsilon' \right) \right\} \right\}, \\ \tilde{\mathcal{E}}_1 &= \left\{ \sum_{i=1}^{N_B} \tilde{g}_i(\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_i) > \exp \left\{ n \left( |R_B - I(\tilde{Y}_1; X'Y')|^+ + \frac{\epsilon'}{2} \right) \right\} \right\}. \end{aligned}$$

Let  $R'_B \stackrel{\text{def}}{=} \frac{\log(N_B - 1)}{n} = \frac{\log(2^{nR_B} - 1)}{n}$ . For  $i \in [1 : N_B]$  and  $\tilde{\mathbf{y}} \in T_{\tilde{Y}_1|X'Y'}^n(\mathbf{x}', \mathbf{y}')$ , define events  $\tilde{\mathcal{E}}_2^{i, \tilde{\mathbf{y}}}$  and  $\tilde{\mathcal{E}}_{2, \dagger}^{i, \tilde{\mathbf{y}}}$  as

$$\tilde{\mathcal{E}}_2^{i, \tilde{\mathbf{y}}} = \left\{ \sum_{j=1, j \neq i}^{N_B} \tilde{h}_j^{\tilde{\mathbf{y}}}(\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_j) > \exp \left\{ n \left( |R_B - I(\tilde{Y}_2; \tilde{Y}_1 X'Y')|^+ + \frac{\epsilon'}{2} \right) \right\} \right\}$$

$$\tilde{\mathcal{E}}_{2,\dagger}^{i,\tilde{\mathbf{y}}} = \left\{ \sum_{j=1, j \neq i}^{N_B} \tilde{h}_j^{\tilde{\mathbf{y}}}(\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_j) > \exp \left\{ n \left( |R'_B - I(\tilde{Y}_2; \tilde{Y}_1 X' Y')|^+ + \frac{\epsilon'}{2} \right) \right\} \right\}$$

Note that

$$\begin{aligned} & \left| \left\{ (\tilde{m}_{B1}, \tilde{m}_{B2}) : \tilde{m}_{B1} \neq \tilde{m}_{B2} \text{ and } (\mathbf{x}', \mathbf{Y}_{\tilde{m}_{B1}}, \mathbf{Y}_{\tilde{m}_{B2}}, \mathbf{y}') \in T_{X' \tilde{Y}_1 \tilde{Y}_2 Y'}^n \right\} \right| \\ &= \sum_{i=1}^{N_B} \tilde{g}_i(\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_i) \left( \sum_{j=1, j \neq i}^{N_B} \tilde{h}_j^{\mathbf{Y}_i}(\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_j) \right). \end{aligned}$$

We see that

$$\tilde{\mathcal{E}} \subseteq \left( \cup_{i \in 2^{n R_B}} \cup_{\tilde{\mathbf{y}} \in T_{\tilde{Y}_1 | X' Y'}(\mathbf{x}', \mathbf{y}')} \tilde{\mathcal{E}}_{2,\dagger}^{i,\tilde{\mathbf{y}}} \right) \cup \tilde{\mathcal{E}}_1.$$

Notice that we need to only consider union over  $\tilde{\mathbf{y}} \in T_{\tilde{Y}_1 | X' Y'}(\mathbf{x}', \mathbf{y}')$  and not  $\tilde{\mathbf{y}} \in \mathcal{Y}^n$ . Since,  $\tilde{\mathcal{E}}_{2,\dagger}^{i,\tilde{\mathbf{y}}} \subseteq \tilde{\mathcal{E}}_{2,\dagger}^{i,\tilde{\mathbf{y}}}$  for all  $i \in [1 : N_B]$ ,

$$\tilde{\mathcal{E}} \subseteq \left( \cup_{i \in 2^{n R_B}} \cup_{\tilde{\mathbf{y}} \in T_{\tilde{Y}_1 | X' Y'}(\mathbf{x}', \mathbf{y}')} \tilde{\mathcal{E}}_{2,\dagger}^{i,\tilde{\mathbf{y}}} \right) \cup \tilde{\mathcal{E}}_1. \quad (68)$$

We apply Lemma 14 to (66) with  $(\mathbf{Y}_1, \dots, \mathbf{Y}_{N_B})$  as the random variables  $(Z_1, \dots, Z_N)$ . We can show that  $a = (n+1)^{|\mathcal{Y}|} \exp(-nI(\tilde{Y}_1; X' Y'))$  satisfies (55). We take  $t = \frac{1}{N_B} \exp \left\{ n \left( |R_B - I(\tilde{Y}_1; X' Y')|^+ + \frac{\epsilon'}{2} \right) \right\}$  and  $n \geq n_1(\epsilon')$  (recall that  $n_1(\epsilon') = \min \left\{ n : (n+1)^{|\mathcal{Y}|} \log e < \frac{1}{2} \exp(\frac{n\epsilon'}{2}) \right\}$ ). This gives  $N_B(t - a \log e) \geq (1/2) \exp(\frac{n\epsilon'}{2})$  which, when plugged into (56), gives

$$\mathbb{P}(\tilde{\mathcal{E}}_1) \leq \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon'}{2} \right\} \right\}. \quad (69)$$

Similarly, for  $i \in [1 : N_B]$ , we can apply Lemma 14 to (67) with  $(\mathbf{Y}_1, \dots, \mathbf{Y}_{i-1}, \mathbf{Y}_{i+1}, \mathbf{Y}_{N_B})$  as the random variables  $(Z_1, \dots, Z_N)$ . We can show that  $a = (n+1)^{|\mathcal{Y}|} \exp(-nI(\tilde{Y}_2; \tilde{Y}_1 X' Y'))$  satisfies (55).

Choose,  $t = \frac{1}{N_B - 1} \exp \left\{ n \left( |R'_B - I(\tilde{Y}_2; \tilde{Y}_1 X' Y')|^+ + \frac{\epsilon'}{2} \right) \right\}$  and  $n \geq n_1(\epsilon')$  to obtain

$$\mathbb{P}(\tilde{\mathcal{E}}_{2,\dagger}^{i,\tilde{\mathbf{y}}}) \leq \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon'}{2} \right\} \right\}, \text{ for } \tilde{\mathbf{y}} \in T_{\tilde{Y}_1 | X' Y'}(\mathbf{x}', \mathbf{y}'). \quad (70)$$

Using (68), (69) and (70), we see that

$$\mathbb{P}(\tilde{\mathcal{E}}) \leq \left( 2^{n R_B} |T_{\tilde{Y}_1 | X' Y'}(\mathbf{x}', \mathbf{y}')| + 1 \right) \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon'}{2} \right\} \right\}.$$

When  $\tilde{m}_{B1} = \tilde{m}_{B2}$  and  $P_{\tilde{Y}_1 \tilde{Y}_2}$  is such that  $\tilde{Y}_1 \neq \tilde{Y}_2$ ,

$$\left| \left\{ (\tilde{m}_{B1}, \tilde{m}_{B2}) : (\mathbf{x}', \mathbf{Y}_{\tilde{m}_{B1}}, \mathbf{Y}_{\tilde{m}_{B2}}, \mathbf{y}') \in T_{X' \tilde{Y}_1 \tilde{Y}_2 Y'}^n \right\} \right| = 0 \text{ w.p. } 1.$$

When  $\tilde{m}_{B1} = \tilde{m}_{B2}$  and  $P_{\tilde{Y}_1 \tilde{Y}_2}$  is such that  $\tilde{Y}_1 = \tilde{Y}_2$ ,

$$\begin{aligned} & \mathbb{P} \left\{ \left| \left\{ (\tilde{m}_{B1}, \tilde{m}_{B2}) : (\mathbf{x}', \mathbf{Y}_{\tilde{m}_{B1}}, \mathbf{Y}_{\tilde{m}_{B2}}, \mathbf{y}') \in T_{X' \tilde{Y}_1 \tilde{Y}_2 Y'}^n \right\} \right| > \exp \left\{ n \left( |R_B - I(\tilde{Y}_2; X' Y' \tilde{Y}_1)|^+ + |R_B - I(\tilde{Y}_1; X' Y')|^+ + \epsilon' \right) \right\} \right\} \\ & \leq \mathbb{P} \left\{ \left| \left\{ \tilde{m}_{B1} : (\mathbf{x}', \mathbf{Y}_{\tilde{m}_{B1}}, \mathbf{y}') \in T_{X' \tilde{Y}_1 Y'}^n \right\} \right| > \exp \left( n \left( |R_B - I(\tilde{Y}_1; X' Y')|^+ + \epsilon' \right) \right) \right\} \\ & \leq \exp \left\{ -\frac{1}{2} \exp(n\epsilon') \right\}. \end{aligned}$$

The last inequality follows from (63) where we substitute  $X$  with  $\tilde{Y}_1$ ,  $Y$  with  $(X', Y')$ ,  $\mathbf{y}$  with  $(\mathbf{x}', \mathbf{y}')$ ,  $R_A$  with  $R_B$  and  $\epsilon/2$  with  $\epsilon'$ . Thus,

$$\begin{aligned} & \mathbb{P} \left\{ \left| \left\{ (\tilde{m}_{B1}, \tilde{m}_{B2}) : (\mathbf{x}', \mathbf{Y}_{\tilde{m}_{B1}}, \mathbf{Y}_{\tilde{m}_{B2}}, \mathbf{y}') \in T_{X' \tilde{Y}_1 \tilde{Y}_2 Y'}^n \right\} \right| > \exp \left\{ n \left( |R_B - I(\tilde{Y}_1; X' Y')|^+ + |R_B - I(\tilde{Y}_2; \tilde{Y}_1 X' Y')|^+ + \epsilon' \right) \right\} \right\} \\ & \leq \left( 2^{n R_B} |T_{\tilde{Y}_1 | X' Y'}(\mathbf{x}', \mathbf{y}')| + 1 \right) \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon'}{2} \right\} \right\} + \exp \left\{ -\frac{1}{2} \exp(n\epsilon') \right\} \\ & \leq \left( 2^{n R_B} |T_{\tilde{Y}_1 | X' Y'}(\mathbf{x}', \mathbf{y}')| + 1 \right) \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon'}{4} \right\} \right\} + \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon'}{2} \right\} \right\}. \end{aligned} \quad (71)$$

This completes the analysis of (18).

Analysis of (17)

Let  $A$  be the set of indices  $(j, k) \in [1 : N_B] \times [1 : N_B]$  such that  $(\mathbf{y}_j, \mathbf{y}_k) \in T_{\tilde{Y}_1 \tilde{Y}_2 | Y'}^n(\mathbf{y}')$  provided  $|A| \leq \exp \left\{ n \left( |R_B - I(\tilde{Y}_2; \tilde{Y}_1 Y')|^+ + |R_B - I(\tilde{Y}_1; Y')|^+ \right) + \frac{\epsilon}{4} \right\}$ . Otherwise,  $A = \emptyset$ . Let

$$\begin{aligned} \tilde{f}_i^{[\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_{N_B}]}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_i) &= \begin{cases} 1, & \text{if } \mathbf{x}_i \in \cup_{(j,k) \in A} T_{X' | \tilde{Y}_1 \tilde{Y}_2 Y'}^n(\mathbf{y}_j, \mathbf{y}_k, \mathbf{y}') \\ 0, & \text{otherwise.} \end{cases} \\ \mathbb{P} \left\{ \sum_{i=1}^{N_A} \tilde{f}_i^{[\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_{N_B}]}(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_i) \neq \left| \left\{ i : \mathbf{X}_i \in T_{X' | \tilde{Y}_1 \tilde{Y}_2 Y'}^n(\mathbf{Y}_j, \mathbf{Y}_k, \mathbf{y}') \text{ for some } j \text{ and } k \right\} \right| \right\} \\ = \mathbb{P} \left\{ \left| \left\{ (\tilde{m}_{B1}, \tilde{m}_{B2}) : (\mathbf{Y}_{\tilde{m}_{B1}}, \mathbf{Y}_{\tilde{m}_{B2}}, \mathbf{y}') \in T_{\tilde{Y}_1 \tilde{Y}_2 Y'}^n \right\} \right| \right. \\ \left. > \exp \left\{ n \left( |R_B - I(\tilde{Y}_1; \tilde{Y}_2 Y')|^+ + |R_B - I(\tilde{Y}_1; Y')|^+ \right) + \frac{\epsilon}{4} \right\} \right\} \\ \leq \left( 2^{n R_B} |T_{\tilde{Y} | Y'}^n(\mathbf{y}')| + 1 \right) \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon}{8} \right\} \right\} + \exp \left\{ -\frac{1}{2} \exp \{n\epsilon\} \right\}, \end{aligned} \quad (72)$$

where last inequality follows from (71) by replacing  $(\mathbf{x}', \mathbf{y}')$  with  $\mathbf{y}'$ ,  $(\mathbf{X}', \mathbf{Y}')$  with  $\mathbf{Y}'$  and  $\frac{\epsilon}{2}$  (or  $\epsilon'$ ) with  $\frac{\epsilon}{4}$ .

For  $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_{N_B} \in \mathcal{Y}^n$ , we will apply Lemma 14 on  $\tilde{f}_i^{[\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_{N_B}]}$  with  $(\mathbf{X}_1, \dots, \mathbf{X}_{N_A})$  as the random variables  $(Z_1, \dots, Z_N)$ . We will first compute the value of  $a$  in (55).

$$\begin{aligned} E \left[ \tilde{f}_i^{[\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_{N_B}]}(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_i) \mid (\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_{i-1}) \right] \\ = \mathbb{P} \left( \mathbf{X}_i \in \cup_{(j,k) \in A} T_{X' | \tilde{Y}_1 \tilde{Y}_2 Y'}^n(\mathbf{y}_j, \mathbf{y}_k, \mathbf{y}') \right) \\ \stackrel{(a)}{\leq} |A| \frac{\exp \left\{ nH(X' | \tilde{Y}_1 \tilde{Y}_2 Y') \right\}}{(n+1)^{-|\mathcal{X}|} \exp(nH(X'))} \\ \leq (n+1)^{|\mathcal{X}|} \exp \left\{ n \left( |R_B - I(\tilde{Y}_2; \tilde{Y}_1 Y')|^+ + |R_B - I(\tilde{Y}_1; Y')|^+ \right) - I(X'; \tilde{Y}_1 \tilde{Y}_2 Y') + \frac{\epsilon}{4} \right\}. \end{aligned}$$

where (a) follows from (21), (22), a union bound over  $(j, k) \in A$  and by noting that  $|T_A^n| = |T_{X'}^n|$ . If  $I(X'; \tilde{Y}_1 \tilde{Y}_2 Y') > |R_B - I(\tilde{Y}_2; \tilde{Y}_1 Y')|^+ + |R_B - I(\tilde{Y}_1; Y')|^+ + \epsilon$  (which (17) postulates), (55) holds with  $a = (n+1)^{|\mathcal{X}|} \exp \left\{ -\frac{3}{4} n\epsilon \right\}$ . For  $t = \exp \left\{ \frac{-n\epsilon}{2} \right\}$  and  $n \geq n_2(\epsilon)$  (recall that  $n_2(\epsilon) = \min \{ n : (n+1)^{|\mathcal{X}|} \log e < \frac{1}{2} \exp \left\{ \frac{n\epsilon}{4} \right\} \}$ ), we get

$$\begin{aligned} \mathbb{P} \left\{ \frac{1}{N_A} \sum_{i=1}^{N_A} \tilde{f}_i^{[\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_{N_B}]}(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_i) > \exp \left\{ \frac{-n\epsilon}{2} \right\} \right\} \\ \leq \exp \left\{ -\frac{N_A}{2} \exp \left\{ -\frac{n\epsilon}{2} \right\} \right\} \\ \leq \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon}{2} \right\} \right\} \end{aligned}$$

where the last inequality uses the assumption that  $N_A \geq \exp \{n\epsilon\}$ . Averaging over  $(\mathbf{Y}_1, \dots, \mathbf{Y}_B)$ , we get

$$\begin{aligned} \mathbb{P} \left\{ \frac{1}{N_A} \sum_{i=1}^{N_A} \tilde{f}_i^{[\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_{N_B}]}(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_i) > \exp \left\{ \frac{-n\epsilon}{2} \right\} \right\} \\ \leq \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon}{2} \right\} \right\}. \end{aligned} \quad (73)$$

Let events  $\tilde{\mathcal{F}}_1$  and  $\tilde{\mathcal{F}}_2$  be defined as

$$\begin{aligned} \tilde{\mathcal{F}}_1 &= \left\{ \frac{1}{N_A} \left| \left\{ i : \mathbf{X}_i \in T_{X' | \tilde{Y}_1 \tilde{Y}_2 Y'}^n(\mathbf{Y}_j, \mathbf{Y}_k, \mathbf{y}') \text{ for some } j \text{ and } k \right\} \right| > \exp \left\{ \frac{-n\epsilon}{2} \right\} \right\}, \\ \tilde{\mathcal{F}}_2 &= \left\{ \sum_{i=1}^{N_A} \tilde{f}_i^{[\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_{N_B}]}(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_i) \neq \left| \left\{ i : \mathbf{X}_i \in T_{X' | \tilde{Y}_1 \tilde{Y}_2 Y'}^n(\mathbf{Y}_j, \mathbf{Y}_k, \mathbf{y}') \text{ for some } j \text{ and } k \right\} \right| \right\}, \end{aligned}$$

$$\tilde{\mathcal{F}}_3 = \left\{ \sum_{i=1}^{N_A} \tilde{f}_i^{[Y_1, Y_2, \dots, Y_{N_B}]}(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_i) > \exp \left\{ \frac{-n\epsilon}{2} \right\} \right\}.$$

We are interested in  $\mathbb{P}(\tilde{\mathcal{F}}_1)$ . We see that

$$\begin{aligned} \mathbb{P}(\tilde{\mathcal{F}}_1) &= \mathbb{P}(\tilde{\mathcal{F}}_1 \cap \tilde{\mathcal{F}}_2) + \mathbb{P}(\tilde{\mathcal{F}}_1 \cap \tilde{\mathcal{F}}_2^c) \\ &\leq \mathbb{P}(\tilde{\mathcal{F}}_2) + \mathbb{P}(\tilde{\mathcal{F}}_1 \cap \tilde{\mathcal{F}}_2^c) \\ &\leq \mathbb{P}(\tilde{\mathcal{F}}_2) + \mathbb{P}(\tilde{\mathcal{F}}_3) \\ &\stackrel{(a)}{\leq} \left( 2^{nR_B} |T_{\tilde{Y}|Y'}(\mathbf{y}')| + 1 \right) \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon}{8} \right\} \right\} + \exp \left\{ -\frac{1}{2} \exp \{n\epsilon\} \right\} + \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon}{2} \right\} \right\} \\ &= \left( 2^{nR_B} |T_{\tilde{Y}|Y'}(\mathbf{y}')| + 3 \right) \exp \left\{ -\frac{1}{2} \exp \left\{ \frac{n\epsilon}{8} \right\} \right\}, \end{aligned}$$

where (a) follows from (72) and (73). Statements (50)-(54) are analogous to (14)-(18) with the roles of users A and B are interchanged. In particular,  $(\mathbf{x}, \mathbf{x}', \mathbf{y}, \mathbf{y}')$  is replaced with  $(\tilde{\mathbf{y}}, \tilde{\mathbf{y}}, \tilde{\mathbf{x}}, \tilde{\mathbf{x}})$ ,  $(X, \tilde{X}, \tilde{Y}, Y)$  with  $(\tilde{Y}, \tilde{Y}, \tilde{X}, \tilde{X})$ ,  $(X', \tilde{Y}_1, \tilde{Y}_2, Y')$  with  $(Y'', \tilde{X}_1, \tilde{X}_2, X'')$  and  $R_A$  with  $R_B$ . Thus, we can follow a similar analysis and show that the probability that any of the statements (50), (51), (52), (53) or (54) do not hold for some  $\tilde{\mathbf{x}}, \tilde{\mathbf{x}}, \tilde{\mathbf{y}}, \tilde{\mathbf{y}}$ ,  $P_{\tilde{X}\tilde{X}\tilde{Y}\tilde{Y}}$  and  $P_{X''\tilde{X}_1\tilde{X}_2Y''}$  also falls doubly exponentially.  $\square$

## APPENDIX B PROOF OF LEMMA 6

**Lemma.** Suppose  $\alpha > 0$ . For a channel which is not A-spoofable, for sufficiently small  $\eta > 0$ , there does not exist a distribution  $P_{XY\tilde{X}Y'X'\tilde{Y}Z} \in \mathcal{P}_{XY\tilde{X}Y'X'\tilde{Y}Z}^n$  with  $\min_x P_X(x), \min_{\tilde{x}} P_{\tilde{X}}(\tilde{x}), \min_{\tilde{y}} P_{\tilde{Y}}(\tilde{y}) \geq \alpha$  which satisfies the following:

- (A)  $P_{XYZ} \in D_\eta$ ,
- (B)  $P_{\tilde{X}Y'Z} \in D_\eta$ ,
- (C)  $P_{X'\tilde{Y}Z} \in D_\eta$ ,
- (D)  $I(\tilde{X}\tilde{Y}; XZ|Y) < \eta$ ,
- (E)  $I(X\tilde{Y}; \tilde{X}Z|Y') < \eta$  and
- (F)  $I(X\tilde{X}; \tilde{Y}Z|X') < \eta$ .

Similarly, for a channel which is not B-spoofable, there does not exist a distribution  $P_{X'\tilde{Y}_1X'_2\tilde{Y}_2XYZ} \in \mathcal{P}_{X'\tilde{Y}_1X'_2\tilde{Y}_2XYZ}^n$  with  $\min_x P_X(x), \min_{\tilde{y}_1} P_{\tilde{Y}_1}(\tilde{y}_1), \min_{\tilde{y}_2} P_{\tilde{Y}_2}(\tilde{y}_2) \geq \alpha > 0$  which satisfies the following:

- (G)  $P_{XYZ} \in D_\eta$ ,
- (H)  $P_{X'\tilde{Y}_1Z} \in D_\eta$ ,
- (I)  $P_{X'_2\tilde{Y}_2Z} \in D_\eta$ ,
- (J)  $I(\tilde{Y}_1\tilde{Y}_2; XZ|Y) < \eta$ ,
- (K)  $I(X\tilde{Y}_2; \tilde{Y}_1Z|X'_1) < \eta$  and
- (L)  $I(X\tilde{Y}_1; \tilde{Y}_2Z|X'_2) < \eta$ .

*Proof.* Suppose for a channel which is not A-spoofable, there exists  $P_{XY\tilde{X}Y'X'\tilde{Y}Z} \in \mathcal{P}_{XY\tilde{X}Y'X'\tilde{Y}Z}^n$  which satisfies (A)-(F). Using (A) and (D), we obtain that

$$\begin{aligned} 2\eta &> D(P_{XYZ} || P_X P_Y W_{Z|XY}) + I(\tilde{X}\tilde{Y}; XZ|Y) \\ &= D(P_{XYZ} || P_X P_Y W_{Z|XY}) + D(P_{XY\tilde{X}\tilde{Y}Z} || P_Y P_{\tilde{X}\tilde{Y}|Y} P_{XZ|Y}) \\ &= \sum_{x,y,\tilde{x},\tilde{y},z} P_{XY\tilde{X}\tilde{Y}Z}(x,y,\tilde{x},\tilde{y},z) \left( \log \left\{ \frac{P_{XYZ}(x,y,z)}{P_X(x)P_Y(y)W_{Z|XY}(z|x,y)} \right\} + \log \left\{ \frac{P_{XY\tilde{X}\tilde{Y}Z}(x,y,\tilde{x},\tilde{y},z)}{P_Y(y)P_{\tilde{X}\tilde{Y}|Y}(\tilde{x},\tilde{y}|y)P_{XZ|Y}(x,z|y)} \right\} \right) \\ &= \sum_{x,y,\tilde{x},\tilde{y},z} P_{XY\tilde{X}\tilde{Y}Z}(x,y,\tilde{x},\tilde{y},z) \left( \log \left\{ \frac{P_{XYZ}(x,y,z) \times P_{XY\tilde{X}\tilde{Y}Z}(x,y,\tilde{x},\tilde{y},z)}{P_X(x)P_Y(y)W_{Z|XY}(z|x,y) \times P_Y(y)P_{\tilde{X}\tilde{Y}|Y}(\tilde{x},\tilde{y}|y)P_{XZ|Y}(x,z|y)} \right\} \right) \\ &= \sum_{x,y,\tilde{x},\tilde{y},z} P_{XY\tilde{X}\tilde{Y}Z}(x,y,\tilde{x},\tilde{y},z) \left( \log \left\{ \frac{P_{XY\tilde{X}\tilde{Y}Z}(x,y,\tilde{x},\tilde{y},z)}{P_X(x)P_Y(y)W_{Z|XY}(z|x,y)P_{\tilde{X}\tilde{Y}|Y}(\tilde{x},\tilde{y}|y)} \right\} \right) \\ &= \sum_{x,y,\tilde{x},\tilde{y},z} P_{XY\tilde{X}\tilde{Y}Z}(x,y,\tilde{x},\tilde{y},z) \left( \log \left\{ \frac{P_{XY\tilde{X}\tilde{Y}Z}(x,y,\tilde{x},\tilde{y},z)}{P_X(x)W_{Z|XY}(z|x,y)P_{\tilde{X}\tilde{Y}}(y,\tilde{x},\tilde{y})} \right\} \right) \end{aligned}$$

$$\begin{aligned}
&= D(P_{XY\tilde{X}\tilde{Y}Z} \| P_X P_{\tilde{X}\tilde{Y}} P_{Y|\tilde{X}\tilde{Y}} W_{Z|XY}) \\
&\stackrel{(a)}{\geq} D(P_{X\tilde{X}\tilde{Y}Z} \| P_X P_{\tilde{X}\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^1),
\end{aligned} \tag{74}$$

where  $V_{Z|X\tilde{X}\tilde{Y}}^{(1)}(z|x, \tilde{x}, \tilde{y}) \stackrel{\text{def}}{=} \sum_y P_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y)$  and (a) follows from the log sum inequality.

Before we proceed, recall that the total variation distance between distributions  $P_X$  and  $Q_X$  distributed on an alphabet  $\mathcal{X}$ ,  $d_{TV}(P_X, Q_X)$  is defined as

$$d_{TV}(P_X, Q_X) \stackrel{\text{def}}{=} \frac{1}{2} \sum_{x \in \mathcal{X}} |P_X(x) - Q_X(x)|.$$

The total variation distance and KL divergence between two distributions are related by the Pinsker's inequality as stated below.

$$D(P_X \| Q_X) \geq \frac{1}{\ln(2)} (d_{TV}(P_X, Q_X))^2. \tag{75}$$

With this, using Pinsker's inequality (75) on (74), we obtain

$$d_{TV}(P_{X\tilde{X}\tilde{Y}Z}, P_X P_{\tilde{X}\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(1)}) < c\sqrt{\eta}. \tag{76}$$

where  $c = \sqrt{2 \ln 2}$ . Similarly, using (B) and (E), we obtain

$$d_{TV}(P_{\tilde{X}X\tilde{Y}Z}, P_{\tilde{X}} P_X \tilde{Y} V_{Z|X\tilde{X}\tilde{Y}}^{(2)}) < c\sqrt{\eta} \tag{77}$$

where  $V_{Z|X\tilde{X}\tilde{Y}}^{(2)}(z|x, \tilde{x}, \tilde{y}) \stackrel{\text{def}}{=} \sum_{y'} P_{Y'|X\tilde{Y}}(y'|x, \tilde{y}) W_{Z|XY}(z|\tilde{x}, y')$ . Finally, using (C) and (F), we get

$$d_{TV}(P_{X\tilde{X}\tilde{Y}Z}, P_X \tilde{X} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(3)}) < c\sqrt{\eta} \tag{78}$$

where  $V_{Z|X\tilde{X}\tilde{Y}}^{(3)}(z|x, \tilde{x}, \tilde{y}) \stackrel{\text{def}}{=} \sum_{x'} P_{X'|X\tilde{X}}(x'|x, \tilde{x}) W_{Z|XY}(z|x', \tilde{y})$ . Next, we note that

$$\begin{aligned}
&2d_{TV}(P_X P_{\tilde{X}\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(1)}, P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(1)}) \\
&= \sum_{x, \tilde{x}, \tilde{y}, z} \left| P_X(x) P_{\tilde{X}\tilde{Y}}(\tilde{x}, \tilde{y}) V_{Z|X\tilde{X}\tilde{Y}}^{(1)}(z|x, \tilde{x}, \tilde{y}) - P_X(x) P_{\tilde{X}}(\tilde{x}) P_{\tilde{Y}}(\tilde{y}) V_{Z|X\tilde{X}\tilde{Y}}^{(1)}(z|x, \tilde{x}, \tilde{y}) \right| \\
&= \left( \sum_x P_X(x) \right) \left( \sum_z V_{Z|X\tilde{X}\tilde{Y}}^{(1)}(z|x, \tilde{x}, \tilde{y}) \right) \sum_{\tilde{x}, \tilde{y}} |P_{\tilde{X}\tilde{Y}}(\tilde{x}, \tilde{y}) - P_{\tilde{X}}(\tilde{x}) P_{\tilde{Y}}(\tilde{y})| \\
&= 2d_{TV}(P_{\tilde{X}\tilde{Y}}, P_{\tilde{X}} P_{\tilde{Y}}) < 2c\sqrt{\eta} \text{ by (77)}.
\end{aligned}$$

Using this and (76),

$$\begin{aligned}
2c\sqrt{\eta} &> d_{TV}(P_{X\tilde{X}\tilde{Y}Z}, P_X P_{\tilde{X}\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(1)}) + d_{TV}(P_X P_{\tilde{X}\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(1)}, P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(1)}) \\
&\stackrel{(a)}{\geq} d_{TV}(P_{X\tilde{X}\tilde{Y}Z}, P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(1)}),
\end{aligned}$$

where (a) is the triangle inequality. Thus,

$$d_{TV}(P_{X\tilde{X}\tilde{Y}Z}, P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(1)}) < 2c\sqrt{\eta}. \tag{79}$$

Similarly, using (76) to show that  $d_{TV}(P_{\tilde{X}} P_X P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(2)}, P_{\tilde{X}} P_X \tilde{Y} V_{Z|X\tilde{X}\tilde{Y}}^{(2)}) < c\sqrt{\eta}$  and (77), we obtain

$$d_{TV}(P_{X\tilde{X}\tilde{Y}Z}, P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(2)}) < 2c\sqrt{\eta}, \tag{80}$$

and using (76) to show that  $d_{TV}(P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(3)}, P_X \tilde{X} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(3)}) < c\sqrt{\eta}$  and (78), we obtain

$$d_{TV}(P_{X\tilde{X}\tilde{Y}Z}, P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(3)}) < 2c\sqrt{\eta}. \tag{81}$$

We use (79) and (81) to write the following:

$$\begin{aligned}
&d_{TV}(P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(1)}, P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(3)}) \\
&\leq d_{TV}(P_{X\tilde{X}\tilde{Y}Z}, P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(1)}) + d_{TV}(P_{X\tilde{X}\tilde{Y}Z}, P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(3)})
\end{aligned}$$



$$< 4c\sqrt{\eta}. \quad (82)$$

Similarly, using (80) and (81), we may write the following:

$$d_{TV} \left( P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(2)}, P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(3)} \right) < 4c\sqrt{\eta}, \quad (83)$$

and using (79) and (80), we may write the following:

$$d_{TV} \left( P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(1)}, P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(2)} \right) < 4c\sqrt{\eta}. \quad (84)$$

Let  $Q_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) \stackrel{\text{def}}{=} \frac{P_{Y'|X\tilde{Y}}(y|\tilde{x}, \tilde{y}) + P_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y})}{2}$  for all  $\tilde{x} \in \mathcal{X}$ ,  $y, \tilde{y} \in \mathcal{Y}$  and  $Q_{X'|X\tilde{X}}(x'|x, \tilde{x}) = \frac{P_{X'|X\tilde{X}}(x'|\tilde{x}, x) + P_{X'|X\tilde{X}}(x'|x, \tilde{x})}{2}$  for all  $x', x, \tilde{x} \in \mathcal{X}$ . With this,

$$\begin{aligned} & \max_{x, \tilde{x}, \tilde{y}, z} 2\alpha^3 \left| \sum_y Q_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - \sum_{x'} Q_{X'|X\tilde{X}}(x'|x, \tilde{x}) W_{Z|XY}(z|x', \tilde{y}) \right| \\ &= \max_{x, \tilde{x}, \tilde{y}, z} \alpha^3 \left| \left( \sum_y P_{Y'|X\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - \sum_{x'} P_{X'|X\tilde{X}}(x'|\tilde{x}, x) W_{Z|XY}(z|x', \tilde{y}) \right) \right. \\ & \quad \left. + \left( \sum_y P_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - \sum_{x'} P_{X'|X\tilde{X}}(x'|x, \tilde{x}) W_{Z|XY}(z|x', \tilde{y}) \right) \right| \\ &\leq \max_{x, \tilde{x}, \tilde{y}, z} \alpha^3 \left| \left( \sum_y P_{Y'|X\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - \sum_{x'} P_{X'|X\tilde{X}}(x'|\tilde{x}, x) W_{Z|XY}(z|x', \tilde{y}) \right) \right| \\ & \quad + \max_{x, \tilde{x}, \tilde{y}, z} \alpha^3 \left| \left( \sum_y P_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - \sum_{x'} P_{X'|X\tilde{X}}(x'|x, \tilde{x}) W_{Z|XY}(z|x', \tilde{y}) \right) \right| \\ &\stackrel{(a)}{=} \max_{x, \tilde{x}, \tilde{y}, z} \alpha^3 \left| \left( \sum_y P_{Y'|X\tilde{Y}}(y|x, \tilde{y}) W_{Z|XY}(z|\tilde{x}, y) - \sum_{x'} P_{X'|X\tilde{X}}(x'|x, \tilde{x}) W_{Z|XY}(z|x', \tilde{y}) \right) \right| \\ & \quad + \max_{x, \tilde{x}, \tilde{y}, z} \alpha^3 \left| \left( \sum_y P_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - \sum_{x'} P_{X'|X\tilde{X}}(x'|x, \tilde{x}) W_{Z|XY}(z|x', \tilde{y}) \right) \right| \\ &\stackrel{(b)}{\leq} \max_{x, \tilde{x}, \tilde{y}, z} P_X(x) P_{\tilde{X}}(\tilde{x}) P_{\tilde{Y}}(\tilde{y}) \left| \left( \sum_y P_{Y'|X\tilde{Y}}(y|x, \tilde{y}) W_{Z|XY}(z|\tilde{x}, y) - \sum_{x'} P_{X'|X\tilde{X}}(x'|x, \tilde{x}) W_{Z|XY}(z|x', \tilde{y}) \right) \right| \\ & \quad + \max_{x, \tilde{x}, \tilde{y}, z} P_X(x) P_{\tilde{X}}(\tilde{x}) P_{\tilde{Y}}(\tilde{y}) \left| \left( \sum_y P_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - \sum_{x'} P_{X'|X\tilde{X}}(x'|x, \tilde{x}) W_{Z|XY}(z|x', \tilde{y}) \right) \right| \\ &\leq \max_{x, \tilde{x}, \tilde{y}, z} \left| \left( P_X(x) P_{\tilde{X}}(\tilde{x}) P_{\tilde{Y}}(\tilde{y}) \sum_y P_{Y'|X\tilde{Y}}(y|x, \tilde{y}) W_{Z|XY}(z|\tilde{x}, y) - P_X(x) P_{\tilde{X}}(\tilde{x}) P_{\tilde{Y}}(\tilde{y}) \sum_{x'} P_{X'|X\tilde{X}}(x'|x, \tilde{x}) W_{Z|XY}(z|x', \tilde{y}) \right) \right| \\ & \quad + \max_{x, \tilde{x}, \tilde{y}, z} \left| \left( P_X(x) P_{\tilde{X}}(\tilde{x}) P_{\tilde{Y}}(\tilde{y}) \sum_y P_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - P_X(x) P_{\tilde{X}}(\tilde{x}) P_{\tilde{Y}}(\tilde{y}) \sum_{x'} P_{X'|X\tilde{X}}(x'|x, \tilde{x}) W_{Z|XY}(z|x', \tilde{y}) \right) \right| \\ &\leq \sum_{x, \tilde{x}, \tilde{y}, z} \left| \left( P_X(x) P_{\tilde{X}}(\tilde{x}) P_{\tilde{Y}}(\tilde{y}) \sum_y P_{Y'|X\tilde{Y}}(y|x, \tilde{y}) W_{Z|XY}(z|\tilde{x}, y) - P_X(x) P_{\tilde{X}}(\tilde{x}) P_{\tilde{Y}}(\tilde{y}) \sum_{x'} P_{X'|X\tilde{X}}(x'|x, \tilde{x}) W_{Z|XY}(z|x', \tilde{y}) \right) \right| \\ & \quad + \sum_{x, \tilde{x}, \tilde{y}, z} \left| \left( P_X(x) P_{\tilde{X}}(\tilde{x}) P_{\tilde{Y}}(\tilde{y}) \sum_y P_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - P_X(x) P_{\tilde{X}}(\tilde{x}) P_{\tilde{Y}}(\tilde{y}) \sum_{x'} P_{X'|X\tilde{X}}(x'|x, \tilde{x}) W_{Z|XY}(z|x', \tilde{y}) \right) \right| \\ &= 2d_{TV} \left( P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(2)}, P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(3)} \right) + 2d_{TV} \left( P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(1)}, P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(2)} \right) \\ &\stackrel{(c)}{<} 16c\sqrt{\eta}, \end{aligned}$$

where (a) is obtained by exchanging  $x$  and  $\tilde{x}$  in the first term, (b) follows by recalling that  $\min_{\tilde{x}} P_{\tilde{X}}(\tilde{x}), \min_{\tilde{y}} P_{\tilde{Y}}(\tilde{y}) \geq \alpha$  and (c) follows from (82) and (83). Thus, we have shown that

$$\max_{x, \tilde{x}, \tilde{y}, z} \left| \sum_y Q_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - \sum_{x'} Q_{X'|X\tilde{X}}(x'|x, \tilde{x}) W_{Z|XY}(z|x', \tilde{y}) \right| < \frac{8c\sqrt{\eta}}{\alpha^3}. \quad (85)$$

Next, we will show the following similar statement:

$$\max_{x, \tilde{x}, \tilde{y}, z} \left| \sum_y Q_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - \sum_{y'} Q_{Y|\tilde{X}\tilde{Y}}(y'|x, \tilde{y}) W_{Z|XY}(z|\tilde{x}, y') \right| < \frac{8c\sqrt{\eta}}{\alpha^3}. \quad (86)$$

Consider,

$$\begin{aligned} & 2\alpha^3 \max_{x, \tilde{x}, \tilde{y}, z} \left| \sum_y Q_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - \sum_{y'} Q_{Y|\tilde{X}\tilde{Y}}(y'|x, \tilde{y}) W_{Z|XY}(z|\tilde{x}, y') \right| \\ &= \max_{x, \tilde{x}, \tilde{y}, z} \alpha^3 \left| \sum_y \left( P_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) + P_{Y'|X\tilde{Y}}(y|\tilde{x}, \tilde{y}) \right) W_{Z|XY}(z|x, y) - \sum_{y'} \left( P_{Y'|X\tilde{Y}}(y'|x, \tilde{y}) + P_{Y|\tilde{X}\tilde{Y}}(y'|x, \tilde{y}) \right) W_{Z|XY}(z|\tilde{x}, y') \right| \\ &\leq \max_{x, \tilde{x}, \tilde{y}, z} \alpha^3 \left| \sum_y P_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - \sum_{y'} P_{Y'|X\tilde{Y}}(y'|x, \tilde{y}) W_{Z|XY}(z|\tilde{x}, y') \right| \\ &\quad + \max_{x, \tilde{x}, \tilde{y}, z} \alpha^3 \left| \sum_y P_{Y'|X\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - \sum_{y'} P_{Y|\tilde{X}\tilde{Y}}(y'|x, \tilde{y}) W_{Z|XY}(z|\tilde{x}, y') \right| \\ &= 2 \max_{x, \tilde{x}, \tilde{y}, z} \alpha^3 \left| \sum_y P_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - \sum_{y'} P_{Y'|X\tilde{Y}}(y'|x, \tilde{y}) W_{Z|XY}(z|\tilde{x}, y') \right| \\ &\leq 2 \max_{x, \tilde{x}, \tilde{y}, z} P_X(x) P_{\tilde{X}}(\tilde{x}) P_{\tilde{Y}}(\tilde{y}) \left| \sum_y P_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - \sum_{y'} P_{Y'|X\tilde{Y}}(y'|x, \tilde{y}) W_{Z|XY}(z|\tilde{x}, y') \right| \\ &\leq 2 \sum_{x, \tilde{x}, \tilde{y}, z} \left| P_X(x) P_{\tilde{X}}(\tilde{x}) P_{\tilde{Y}}(\tilde{y}) \sum_y P_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - P_X(x) P_{\tilde{X}}(\tilde{x}) P_{\tilde{Y}}(\tilde{y}) \sum_{y'} P_{Y'|X\tilde{Y}}(y'|x, \tilde{y}) W_{Z|XY}(z|\tilde{x}, y') \right| \\ &= 4d_{TV} \left( P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(1)}, P_X P_{\tilde{X}} P_{\tilde{Y}} V_{Z|X\tilde{X}\tilde{Y}}^{(2)} \right) \\ &\stackrel{(a)}{<} 16c\sqrt{\eta}, \end{aligned}$$

where (a) follows from (84). With this, we have shown (86). Using (85) and (86), we have shown that for the given channel there exist conditional distributions  $Q_{Y|\tilde{X}\tilde{Y}}$  and  $Q_{X'|X\tilde{X}}$  such that

$$\max \left\{ \max_{x, \tilde{x}, \tilde{y}, z} \left| \sum_y Q_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - \sum_{x'} Q_{X'|X\tilde{X}}(x'|x, \tilde{x}) W_{Z|XY}(z|x', \tilde{y}) \right|, \right. \\ \left. \max_{x, \tilde{x}, \tilde{y}, z} \left| \sum_y Q_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - \sum_{y'} Q_{Y|\tilde{X}\tilde{Y}}(y'|x, \tilde{y}) W_{Z|XY}(z|\tilde{x}, y') \right| \right\} < \frac{8c\sqrt{\eta}}{\alpha^3}. \quad (87)$$

Suppose the channel is not A-spoofable (i.e. (7) does not hold), then for every pair  $(Q_{Y|\tilde{X}\tilde{Y}}, Q_{X'|X\tilde{X}})$ , there exists  $\zeta(Q_{Y|\tilde{X}\tilde{Y}}, Q_{X'|X\tilde{X}}) > 0$  such that the following holds:

$$\max \left\{ \max_{x, \tilde{x}, \tilde{y}, z} \left| \sum_y Q_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - \sum_{x'} Q_{X'|X\tilde{X}}(x'|x, \tilde{x}) W_{Z|XY}(z|x', \tilde{y}) \right|, \right. \\ \left. \max_{x, \tilde{x}, \tilde{y}, z} \left| \sum_y Q_{Y|\tilde{X}\tilde{Y}}(y|\tilde{x}, \tilde{y}) W_{Z|XY}(z|x, y) - \sum_{y'} Q_{Y|\tilde{X}\tilde{Y}}(y'|x, \tilde{y}) W_{Z|XY}(z|\tilde{x}, y') \right| \right\} > \zeta(Q_{Y|\tilde{X}\tilde{Y}}, Q_{X'|X\tilde{X}}).$$

This contradicts (87) for  $\eta$  small enough such that  $\zeta(Q_{Y|\tilde{X}\tilde{Y}}, Q_{X'|X\tilde{X}}) > 8c\sqrt{\eta}/\alpha^3$ . This completes the proof for a channel which is not A spoofable. The proof for a channel which is not B spoofable is along the similar lines. It can be obtained

by interchanging the roles of users A and B and making the following replacements in the above proof:  $\tilde{X} \rightarrow \tilde{Y}_1$ ,  $Y' \rightarrow X'_1$ ,  $X' \rightarrow X'_2$ , and  $\tilde{Y} \rightarrow \tilde{Y}_2$ .  $\square$

## APPENDIX C

### Proof of Theorem 7

*Proof.* For some  $P_A$  and  $P_B$  satisfying  $\min_{x \in \mathcal{X}} P_A(x), \min_{y \in \mathcal{Y}} P_B(y) \geq \alpha$  for some  $\alpha > 0$ ,  $\epsilon > 0$  (TBD),  $n \geq n_0(\epsilon)$  and a rate pair  $(R_A, R_B) \in \mathcal{R}_2(P_A, P_B)$  given by (34) with  $R_A, R_B > \epsilon$ , consider the codebook given by Lemma 4.

*Encoding.* Let  $N_A = 2^{nR_A}$ ,  $N_B = 2^{nR_B}$ ,  $\mathcal{M}_A = \{1, \dots, N_A\}$  and  $\mathcal{M}_B = \{1, \dots, N_B\}$ . For  $m_A \in \mathcal{M}_A$ ,  $f_A(m_A) = \mathbf{x}_{m_A}$  and for  $m_B \in \mathcal{M}_B$ ,  $f_B(m_B) = \mathbf{y}_{m_B}$ .

*Decoding.* For a parameter  $\eta > 0$  (TBD), let  $\mathcal{D}_\eta$  be the set of joint distributions defined as

$$D_\eta \stackrel{\text{def}}{=} \{P_{XYZ} \in \mathcal{P}_{\mathcal{X} \times \mathcal{Y} \times \mathcal{Z}}^n : D(P_{XYZ} \| P_X P_Y W) \leq \eta\}.$$

For a given codebook, the parameter  $\eta$  and the received channel output sequence  $\mathbf{z}$ , decoding happens in five steps. In the first step, we populate sets  $A_1$  and  $B_1$  containing candidate messages for user A and B respectively. In steps 2-5, we prune these sets by sequentially removing the candidates which do not satisfy certain conditions.

**Step 1:** Let  $A_1 = \{m_A \in \mathcal{M}_A : (f_A(m_A), \bar{\mathbf{y}}, \mathbf{z}) \in T_{XYZ}^n \text{ for some } \bar{\mathbf{y}} \in \mathcal{Y}^n \text{ such that } P_{XYZ} \in \mathcal{D}_\eta\}$  and

$$B_1 = \{m_B \in \mathcal{M}_B : (\bar{\mathbf{x}}, f_B(m_B), \mathbf{z}) \in T_{XYZ}^n \text{ for some } \bar{\mathbf{x}} \in \mathcal{X}^n \text{ such that } P_{XYZ} \in \mathcal{D}_\eta\}.$$

**Step 2:** Let  $C_1 = \{m_A \in A_1 : \text{for every } \tilde{m}_{B1}, \tilde{m}_{B2} \in B_1 \text{ and } \bar{\mathbf{y}} \in \mathcal{Y}^n \text{ with } (f_A(m_A), \bar{\mathbf{y}}, f_B(\tilde{m}_{B1}), f_B(\tilde{m}_{B2}), \mathbf{z}) \in T_{XY\tilde{Y}_1\tilde{Y}_2}^n$ <sup>12</sup> that satisfies  $P_{XYZ} \in \mathcal{D}_\eta$ , we have  $I(\tilde{Y}_1\tilde{Y}_2; XZ|Y) > \eta\}$ . Let  $A_2 = A_1 \setminus C_1$ .

**Step 3:** Let  $C_2 = \{m_B \in B_1 : \text{for every } \tilde{m}_{A1}, \tilde{m}_{A2} \in A_2 \text{ and } \bar{\mathbf{x}} \in \mathcal{X}^n \text{ with } (\bar{\mathbf{x}}, f_B(m_B), f_A(\tilde{m}_{A1}), f_A(\tilde{m}_{A2}), \mathbf{z}) \in T_{XY\tilde{X}_1\tilde{X}_2}^n$  that satisfies  $P_{XYZ} \in \mathcal{D}_\eta$ , we have  $I(\tilde{X}_1\tilde{X}_2; YZ|X) > \eta\}$ . Let  $B_2 = B_1 \setminus C_2$ .

**Step 4:** Let  $C_3 = \{m_A \in A_2 : \text{for every } (\tilde{m}_A, \tilde{m}_B) \in A_2 \times B_2, \tilde{m}_A \neq m_A \text{ and } \bar{\mathbf{y}} \in \mathcal{Y}^n \text{ with } (f_A(m_A), \bar{\mathbf{y}}, f_A(\tilde{m}_A), f_B(\tilde{m}_B), \mathbf{z}) \in T_{XY\tilde{X}\tilde{Y}}^n$  such that  $P_{XYZ} \in \mathcal{D}_\eta$ , we have  $I(\tilde{X}\tilde{Y}; XZ|Y) > \eta\}$ . Let  $A_3 = A_2 \setminus C_3$ .

**Step 5:** Let  $C_4 = \{m_B \in B_2 : \text{for every } (\tilde{m}_A, \tilde{m}_B) \in A_3 \times B_2, \tilde{m}_B \neq m_B \text{ and } \bar{\mathbf{x}} \in \mathcal{X}^n \text{ with } (\bar{\mathbf{x}}, f_B(m_B), f_A(\tilde{m}_A), f_B(\tilde{m}_B), \mathbf{z}) \in T_{XY\tilde{X}\tilde{Y}}^n$  such that  $P_{XYZ} \in \mathcal{D}_\eta$ , we have  $I(\tilde{X}\tilde{Y}; YZ|X) > \eta\}$ . Let  $B_3 = B_2 \setminus C_4$ .

After steps 1-5, the decoded output is as follows.

$$\phi(\mathbf{z}) = \begin{cases} (m_A, m_B) & \text{if } A_3 \times B_3 = \{(m_A, m_B)\}, \\ \mathbf{a} & \text{if } |A_3| = 0, |B_3| \neq 0, \\ \mathbf{b} & \text{if } |A_3| \neq 0, |B_3| = 0 \text{ and} \\ (1, 1) & \text{otherwise.} \end{cases}$$

Similar to the decoder for achievability of Theorem 1, the last of the above cases (*i.e.*  $\phi(\mathbf{z}) = (1, 1)$ ) occurs when either of the following two events occur: (i)  $|A_3| = |B_3| = 0$  and (ii)  $|A_3|, |B_3| \geq 1$  and  $|A_3| + |B_3| \geq 3$ . The first event will be shown to be an atypical event and hence will occur with vanishing probability. For small enough choices of  $\eta > 0$ , Lemma 6 implies that the second event cannot happen in a non-spoofable channel. We consider such an  $\eta$ .

As noted previously (see (6))  $P_{e, \text{hon}} \leq P_{e, \text{mal A}} + P_{e, \text{mal B}}$ . Thus, it is sufficient to analyze the case when one of the users is adversarial. We first consider the case when user B is malicious while user A is honest. Recall that for  $\mathcal{E}_{m_A}^A = \{\mathbf{z} : \phi_A(\mathbf{z}) \notin \{m_A, \mathbf{b}\}\}$ , the probability of error is

$$P_{e, \text{mal B}} = \max_{\mathbf{y} \in \mathcal{Y}^n} \frac{1}{N_A} \sum_{m_A \in \mathcal{M}_A} W^n(\mathcal{E}_{m_A}^A | f_A(m_A), \mathbf{y}).$$

For each  $\mathbf{y} \in \mathcal{Y}^n$ , we will obtain a uniform upper bound on  $P_{e, \text{mal B}}$  which goes to zero as  $n \rightarrow \infty$ . So, let us suppose that user B attacks with  $\mathbf{y} \in \mathcal{Y}^n$  and analyze

$$P := \frac{1}{N_A} \sum_{m_A \in \mathcal{M}_A} W^n(\mathcal{E}_{m_A}^A | f_A(m_A), \mathbf{y}).$$

For some  $\epsilon$  (TBD) satisfying  $0 < \epsilon < \eta/3$ , let

$$\mathcal{H} \stackrel{\text{def}}{=} \{m_A : (\mathbf{x}_{m_A}, \mathbf{y}) \in T_{XY}^n \text{ such that } I(X; Y) > \epsilon\}.$$

Then,

$$P \leq \frac{1}{N_A} |\mathcal{H}| + \frac{1}{N_A} \sum_{m_A \in \mathcal{H}^c} W^n(\mathcal{E}_{m_A}^A | f_A(m_A), \mathbf{y})$$

<sup>12</sup>Note that the statement  $(f_A(m_A), \bar{\mathbf{y}}, f_B(\tilde{m}_{B1}), f_B(\tilde{m}_{B2}), \mathbf{z}) \in T_{XY\tilde{Y}_1\tilde{Y}_2}^n$  is used to define the distribution  $P_{XY\tilde{Y}_1\tilde{Y}_2}$ .

$$=: P_1 + P_2.$$

Here,  $\mathcal{H}^c$  denotes the complement of  $\mathcal{H}$ . The first term on the RHS,

$$P_1 \leq |\mathcal{P}_{\mathcal{X} \times \mathcal{Y}}^n| \times \frac{|\{m_A : (\mathbf{x}_{m_A}, \mathbf{y}) \in T_{XY}^n, I(X; Y) > \epsilon\}|}{N_A}$$

which goes to zero as  $n \rightarrow \infty$  by (14) and noting that there are only polynomially many types. In order to analyze the second term  $P_2$ , for  $\mathbf{y}' \in \mathcal{Y}^n$ , let  $\mathcal{E}_1(\mathbf{y}')$  be defined as

$$\mathcal{E}_1(\mathbf{y}') = \{\mathbf{z} : (\mathbf{x}_{m_A}, \mathbf{y}', \mathbf{z}) \in T_{XYZ}^n \text{ for some } P_{XYZ} \in \mathcal{D}_\eta\}.$$

Then,

$$\begin{aligned} P_2 &= \frac{1}{N_A} \sum_{m_A \in \mathcal{H}^c} W^n(\mathcal{E}_{m_A}^A | f_A(m_A), \mathbf{y}) \\ &= \frac{1}{N_A} \sum_{m_A \in \mathcal{H}^c} W^n((\mathcal{E}_1(\mathbf{y})^c \cap \mathcal{E}_{m_A}^A) \cup (\mathcal{E}_1(\mathbf{y}) \cap \mathcal{E}_{m_A}^A) | f_A(m_A), \mathbf{y}) \\ &\leq \frac{1}{N_A} \sum_{m_A \in \mathcal{H}^c} W^n((\mathcal{E}_1(\mathbf{y})^c) | f_A(m_A), \mathbf{y}) + W^n((\mathcal{E}_1(\mathbf{y}) \cap \mathcal{E}_{m_A}^A) | f_A(m_A), \mathbf{y}) \\ &= \frac{1}{N_A} \sum_{m_A \in \mathcal{H}^c} \left( \sum_{P_{XYZ} \in \mathcal{D}_\eta^c} \sum_{\mathbf{z} \in T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y})} W^n(\mathbf{z} | \mathbf{x}_{m_A}, \mathbf{y}) \right) + \frac{1}{N_A} \sum_{m_A \in \mathcal{H}^c} \left( \sum_{P_{XYZ} \in \mathcal{D}_\eta} \sum_{\mathbf{z} \in T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y}) \cap \mathcal{E}_{m_A}^A} W^n(\mathbf{z} | \mathbf{x}_{m_A}, \mathbf{y}) \right) \\ &=: P_{2a} + P_{2b}. \end{aligned}$$

To analyze  $P_{2a}$ , consider any  $m_A \in \mathcal{H}^c$ ,

$$\begin{aligned} \sum_{\substack{P_{XYZ} \in \mathcal{D}_\eta^c, \\ I(X; Y) \leq \epsilon}} \sum_{\mathbf{z} \in T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y})} W^n(\mathbf{z} | \mathbf{x}_{m_A}, \mathbf{y}) &\leq \sum_{\substack{P_{XYZ} \in \mathcal{D}_\eta^c, \\ I(X; Y) \leq \epsilon}} \exp(-nD(P_{XYZ} || P_{XY}W)) \\ &= \sum_{\substack{P_{XYZ} \in \mathcal{D}_\eta^c, \\ I(X; Y) \leq \epsilon}} \exp(-n(D(P_{XYZ} || P_X P_Y W) - I(X; Y))) \\ &= \sum_{\substack{P_{XYZ} \in \mathcal{D}_\eta^c, \\ I(X; Y) \leq \epsilon}} \exp(-n(\eta - \epsilon)) \\ &\leq |\mathcal{D}_\eta^c| \exp(-n(\eta - \epsilon)). \end{aligned}$$

Thus,

$$\begin{aligned} P_{2a} &\leq \frac{|\mathcal{H}^c|}{N_A} |\mathcal{D}_\eta^c| \exp(-n(\eta - \epsilon)) \\ &\rightarrow 0 \text{ as } \epsilon < \eta/3 \text{ and } |\mathcal{D}_\eta^c| \leq (n+1)^{|\mathcal{X}||\mathcal{Y}||\mathcal{Z}|}, \end{aligned}$$

We are left to analyze

$$P_{2b} = \frac{1}{N_A} \sum_{m_A \in \mathcal{H}^c} \left( \sum_{P_{XYZ} \in \mathcal{D}_\eta} \sum_{\mathbf{z} \in T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y}) \cap \mathcal{E}_{m_A}^A} W^n(\mathbf{z} | \mathbf{x}_{m_A}, \mathbf{y}) \right).$$

In order to proceed, we define the following sets,

$$\begin{aligned} \mathcal{P}_1^\eta &= \{P_{X\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_{\mathcal{X} \times \mathcal{X} \times \mathcal{Y} \times \mathcal{Y} \times \mathcal{Z}}^n : P_{XYZ} \in \mathcal{D}_\eta, P_{\tilde{X}Y'Z} \in \mathcal{D}_\eta \text{ for some } P_{Y'| \tilde{X}Z}, P_{X'\tilde{Y}Z} \in \mathcal{D}_\eta \\ &\quad \text{for some } P_{X'|\tilde{Y}Z}, P_X = P_{\tilde{X}} = P_A, P_{\tilde{Y}} = P_B, I(\tilde{Y}; X) \leq 2\eta, I(\tilde{Y}; \tilde{X}) \leq 2\eta \text{ and } I(\tilde{X}\tilde{Y}; XZ|Y) \geq \eta\} \\ \mathcal{P}_2^\eta &= \{P_{X\tilde{Y}_1\tilde{Y}_2YZ} \in \mathcal{P}_{\mathcal{X} \times \mathcal{Y} \times \mathcal{Y} \times \mathcal{Y} \times \mathcal{Z}}^n : P_{XYZ} \in \mathcal{D}_\eta, P_{X_1\tilde{Y}_1Z} \in \mathcal{D}_\eta \text{ for some } P_{X_1'|\tilde{Y}_1Z}, P_{X_2'\tilde{Y}_2Z} \in \mathcal{D}_\eta \\ &\quad \text{for some } P_{X_2'|\tilde{Y}_2Z}, P_X = P_A, P_{\tilde{Y}_1} = P_{\tilde{Y}_2} = P_B \text{ and } I(\tilde{Y}_1\tilde{Y}_2; XZ|Y) \geq \eta\}. \end{aligned}$$

For  $P_{X\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1^\eta$  and  $P_{X\tilde{Y}_1\tilde{Y}_2YZ} \in \mathcal{P}_2^\eta$ , let

$$\begin{aligned} \mathcal{E}_{m_A,1}(P_{X\tilde{X}\tilde{Y}YZ}) &= \{\mathbf{z} : \exists (\tilde{m}_A, \tilde{m}_B) \in \mathcal{M}_A \times \mathcal{M}_B, \tilde{m}_A \neq m_A, (\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}, \mathbf{y}_{\tilde{m}_B}, \mathbf{z}) \in T_{X\tilde{X}\tilde{Y}YZ}^n\} \text{ and} \\ \mathcal{E}_{m_A,2}(P_{X\tilde{Y}_1\tilde{Y}_2YZ}) &= \{\mathbf{z} : \exists \tilde{m}_{B1}, \tilde{m}_{B2} \in \mathcal{M}_B, (\mathbf{x}_{m_A}, \mathbf{y}_{\tilde{m}_{B1}}, \mathbf{y}_{\tilde{m}_{B2}}, \mathbf{y}, \mathbf{z}) \in T_{X\tilde{Y}_1\tilde{Y}_2YZ}^n\}. \end{aligned}$$

With these definitions,

$$\begin{aligned} \left\{ \mathbf{z} \in T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y}) \cap \mathcal{E}_{m_A}^A \text{ where } P_{XYZ} \in \mathcal{D}_\eta \right\} &\stackrel{(a)}{\subseteq} \{ \mathbf{z} : m_A \in A_1 \cap C_1 \} \cup \{ \mathbf{z} : m_A \in A_2 \cap C_3 \} \\ &\stackrel{(b)}{\subseteq} \left( \bigcup_{P_{X\tilde{Y}_1\tilde{Y}_2YZ} \in \mathcal{P}_2^\eta} \mathcal{E}_{m_A,2}(P_{X\tilde{Y}_1\tilde{Y}_2YZ}) \right) \cup \left( \bigcup_{P_{X\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1^\eta} \mathcal{E}_{m_A,1}(P_{X\tilde{X}\tilde{Y}YZ}) \right) \end{aligned} \quad (88)$$

where (a) follows from the decoder definition (Note that  $\mathbf{z} \in T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y})$  for  $P_{XYZ} \in \mathcal{D}_\eta$  implies that  $\mathbf{z} \in A_1$  and  $\mathbf{z} \in \mathcal{E}_{m_A}^A$  further implies that it was eliminated in **Step 2** or **Step 4**, i.e.,  $\mathbf{z} \in A_1 \cap C_1$  or  $\mathbf{z} \in A_2 \cap C_3$ ). To see (b), first notice that  $\{ \mathbf{z} : m_A \in A_1 \cap C_1 \} \subseteq \left( \bigcup_{P_{X\tilde{Y}_1\tilde{Y}_2YZ} \in \mathcal{P}_2^\eta} \mathcal{E}_{m_A,2}(P_{X\tilde{Y}_1\tilde{Y}_2YZ}) \right)$ . Further, for  $m_A \in C_3$ , we only consider pairs  $(\tilde{m}_A, \tilde{m}_B)$  belonging to  $A_2 \times B_2$  defined in **Step 2** and **Step 3**. Thus, for  $(\mathbf{x}_{m_A}, \mathbf{y}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{z}) \in T_{XY\tilde{X}\tilde{Y}Z}^n$ , we have  $P_{XYZ} \in \mathcal{D}_\eta$ ,  $P_{\tilde{X}\tilde{Y}'Z} \in \mathcal{D}_\eta$  for some  $P_{Y'\tilde{X}Z}$ ,  $P_{X'\tilde{Y}Z} \in \mathcal{D}_\eta$  for some  $P_{X'|\tilde{Y}Z}$  and  $I(X\tilde{X}; \tilde{Y}Z|X') \leq \eta$  (see **Step 3**). For such distributions, the following lemma implies that  $\{ \mathbf{z} : m_A \in A_2 \cap C_3 \} \subseteq \left( \bigcup_{P_{X\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1^\eta} \mathcal{E}_{m_A,1}(P_{X\tilde{X}\tilde{Y}YZ}) \right)$ .

**Lemma 15.** For a distribution  $P_{XY\tilde{X}Y'X'\tilde{Y}Z} \in \mathcal{P}_{\mathcal{X} \times \mathcal{Y} \times \mathcal{X}' \times \mathcal{Y}' \times \mathcal{X} \times \mathcal{Y} \times \mathcal{Z}}^n$  satisfying

- (A)  $P_{XYZ} \in \mathcal{D}_\eta$
- (B)  $P_{\tilde{X}\tilde{Y}'Z} \in \mathcal{D}_\eta$
- (C)  $P_{X'\tilde{Y}Z} \in \mathcal{D}_\eta$ , and
- (D)  $I(X\tilde{X}; \tilde{Y}Z|X') < \eta$ ,

we have  $I(\tilde{Y}; X) \leq 2\eta$  and  $I(\tilde{Y}; \tilde{X}) \leq 2\eta$ .

The proof of this lemma is given after the current proof. It follows by adding (C) and (D) and an application of log-sum inequality. Continuing the analysis of  $P_{2b}$ , we use (88) to write

$$\begin{aligned} P_{2b} &= \frac{1}{N_A} \sum_{m_A \in \mathcal{H}^c} \left( \sum_{P_{XYZ} \in \mathcal{D}_\eta} \sum_{\mathbf{z} \in T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y}) \cap \mathcal{E}_{m_A}^A} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) \right) \\ &\leq \frac{1}{N_A} \sum_{m_A \in \mathcal{M}_A} \sum_{P_{X\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1^\eta} W^n(\mathcal{E}_{m_A,1}(P_{X\tilde{X}\tilde{Y}YZ})|\mathbf{x}_{m_A}, \mathbf{y}) \\ &\quad + \frac{1}{N_A} \sum_{m_A \in \mathcal{M}_A} \sum_{P_{X\tilde{Y}_1\tilde{Y}_2YZ} \in \mathcal{P}_2^\eta} W^n(\mathcal{E}_{m_A,2}(P_{X\tilde{Y}_1\tilde{Y}_2YZ})|\mathbf{x}_{m_A}, \mathbf{y}). \end{aligned} \quad (89)$$

We see that  $|\mathcal{P}_1^\eta|$  and  $|\mathcal{P}_2^\eta|$  grow at most polynomial in  $n$ . So, it will suffice to uniformly upper bound  $W^n(\mathcal{E}_{m_A,1}(P_{X\tilde{X}\tilde{Y}YZ})|\mathbf{x}_{m_A}, \mathbf{y})$  and  $W^n(\mathcal{E}_{m_A,2}(P_{X\tilde{Y}_1\tilde{Y}_2YZ})|\mathbf{x}_{m_A}, \mathbf{y})$  by a term exponentially decreasing in  $n$  for all  $P_{X\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1^\eta$ ,  $P_{X\tilde{Y}_1\tilde{Y}_2YZ} \in \mathcal{P}_2^\eta$  and  $m_A \in \mathcal{M}_A$ . We start with  $W^n(\mathcal{E}_{m_A,1}(P_{X\tilde{X}\tilde{Y}YZ})|\mathbf{x}_{m_A}, \mathbf{y})$ . By using (15), we see that for  $P_{X\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1^\eta$  such that

$$I(X; \tilde{X}\tilde{Y}Y) > |R_A - I(\tilde{X}; \tilde{Y}Y)|^+ + |R_B - I(\tilde{Y}; Y)|^+ + \epsilon, \quad (90)$$

we have,

$$\left| \frac{\left\{ m_A : (\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{m_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n \text{ for some } \tilde{m}_A \neq m_A \text{ and some } m_B \right\}}{N_A} \right| \leq \exp\{-n\epsilon/2\}.$$

So, for all  $P_{X\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1^\eta$  satisfying (90),

$$\begin{aligned} &\frac{1}{N_A} \sum_{m_A \in \mathcal{H}^c} W^n(\mathcal{E}_{m_A,1}(P_{X\tilde{X}\tilde{Y}YZ})|\mathbf{x}_{m_A}, \mathbf{y}) \\ &= \frac{1}{N_A} \sum_{\substack{m_A: (\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{m_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n \\ \tilde{m}_A \in \mathcal{M}_A, \tilde{m}_A \neq m_A, m_B \in \mathcal{M}_B}} \sum_{\mathbf{z} \in T_{Z|X\tilde{X}\tilde{Y}Y}^n(\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{m_B})} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) \\ &\leq \exp\{-n\epsilon/2\}. \end{aligned}$$

Thus, it is sufficient to consider distributions  $P_{X\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1^\eta$  for which

$$I(X; \tilde{X}\tilde{Y}Y) \leq |R_A - I(\tilde{X}; \tilde{Y}Y)|^+ + |R_B - I(\tilde{Y}; Y)|^+ + \epsilon. \quad (91)$$

For  $P_{X\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1^\eta$  satisfying (91),

$$\sum_{\mathbf{z} \in \mathcal{E}_{m_A,1}(P_{X\tilde{X}\tilde{Y}YZ})} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y})$$

$$\begin{aligned}
&= \sum_{\substack{\tilde{m}_A, \tilde{m}_B: \tilde{m}_A \neq m_A \\ (\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n}} \sum_{\mathbf{z}: (\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{y}, \mathbf{z}) \in T_{X\tilde{X}\tilde{Y}YZ}^n} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) \\
&\stackrel{(a)}{\leq} \sum_{\substack{\tilde{m}_A, \tilde{m}_B: \tilde{m}_A \neq m_A \\ (\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n}} \frac{|T_{Z|X\tilde{X}\tilde{Y}Y}^n(\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{y})|}{|T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y})|} \\
&\stackrel{(b)}{\leq} \sum_{\substack{\tilde{m}_A, \tilde{m}_B: \tilde{m}_A \neq m_A \\ (\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n}} \frac{\exp\left(nH(Z|X\tilde{X}\tilde{Y}Y)\right)}{(n+1)^{-|\mathcal{X}||\mathcal{Y}||\mathcal{Z}|} \exp\left(nH(Z|XY)\right)} \\
&\stackrel{(c)}{\leq} \sum_{\substack{\tilde{m}_A, \tilde{m}_B: \\ (\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{y}) \in T_{X\tilde{X}\tilde{Y}Y}^n}} \exp\left(-n\left(I(Z; \tilde{X}\tilde{Y}|XY) - \epsilon\right)\right) \\
&\stackrel{(d)}{\leq} \exp\left(n\left(|R_A - I(\tilde{X}; \tilde{Y}|XY)|^+ + |R_B - I(\tilde{Y}; XY)|^+ - I(Z; \tilde{X}\tilde{Y}|XY) + 2\epsilon\right)\right) \tag{92}
\end{aligned}$$

where (a) follows by noting that whenever  $\mathbf{z}$  belongs to  $T_{Z|X\tilde{X}\tilde{Y}Y}^n(\mathbf{x}_{m_A}, \mathbf{x}_{\tilde{m}_A}, \mathbf{y}_{\tilde{m}_B}, \mathbf{y})$ ,  $\mathbf{z}$  also belongs to  $T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y})$ ; and for each  $\mathbf{z} \in T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y})$ , the value of  $W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y})$  is the same and is upper bounded by  $1/|T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y})|$ . (b) follows from (22), (c) holds for large  $n$  and (d) follows from (16). To analyze (92), we will separately consider the following cases which together cover all the possibilities.

- 1)  $R_A \leq I(\tilde{X}; \tilde{Y}|Y)$  and  $R_B \leq I(\tilde{Y}; Y)$
- 2)  $I(\tilde{X}; \tilde{Y}|Y) < R_A$  and  $R_B \leq I(\tilde{Y}; XY)$
- 3)  $R_A \leq I(\tilde{X}; \tilde{Y}|XY)$  and  $I(\tilde{Y}; Y) < R_B$
- 4)  $I(\tilde{X}; \tilde{Y}|XY) < R_A$  and  $I(\tilde{Y}; XY) < R_B$

Case 1:  $R_A \leq I(\tilde{X}; \tilde{Y}|Y)$  and  $R_B \leq I(\tilde{Y}; Y)$

In this case, (91) implies that  $I(\tilde{X}; \tilde{Y}|Y) \leq \epsilon$  which implies that  $I(X; \tilde{X}\tilde{Y}|Y) \leq \epsilon$ . Thus, using the condition  $I(XZ; \tilde{X}\tilde{Y}|Y) \geq \eta$  from definition of  $\mathcal{P}_1^\eta$ , we see that

$$\begin{aligned}
I(Z; \tilde{X}\tilde{Y}|XY) &= I(XZ; \tilde{X}\tilde{Y}|Y) - I(X; \tilde{X}\tilde{Y}|Y) \\
&\geq \eta - \epsilon.
\end{aligned}$$

Using (92), this implies that

$$\begin{aligned}
\sum_{\mathbf{z} \in \mathcal{E}_{m_A, 1}(P_{X\tilde{X}\tilde{Y}YZ})} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) &\leq \exp(-n(\eta - 3\epsilon)) \\
&\rightarrow 0 \text{ because } \eta > 3\epsilon.
\end{aligned}$$

Case 2:  $I(\tilde{X}; \tilde{Y}|Y) < R_A$  and  $R_B \leq I(\tilde{Y}; XY)$

Using (91), we have

$$\begin{aligned}
-|R_B - I(\tilde{Y}; Y)|^+ &\leq R_A - I(\tilde{X}; \tilde{Y}|Y) - I(X; \tilde{X}\tilde{Y}|Y) + \epsilon \\
&= R_A - I(\tilde{X}; \tilde{Y}|Y) - I(X; \tilde{Y}|Y) - I(X; \tilde{X}|\tilde{Y}|Y) + \epsilon \\
&= R_A - I(\tilde{X}; \tilde{Y}|XY) - I(X; \tilde{Y}|Y) + \epsilon.
\end{aligned}$$

This implies that

$$R_A - I(\tilde{X}; \tilde{Y}|XY) + \epsilon \geq I(X; \tilde{Y}|Y) - |R_B - I(\tilde{Y}; Y)|^+.$$

We will argue that the RHS of the above inequality is non-negative. When  $R_B \leq I(\tilde{Y}; Y)$ , the RHS is  $I(X; \tilde{Y}|Y)$  which is non-negative. Otherwise, when  $I(\tilde{Y}; Y) < R_B \leq I(\tilde{Y}; XY)$ ,

$$\begin{aligned}
I(X; \tilde{Y}|Y) - |R_B - I(\tilde{Y}; Y)|^+ &= I(X; \tilde{Y}|Y) - R_B + I(\tilde{Y}; Y) \\
&= I(X; Y) + I(X; \tilde{Y}|Y) - R_B + I(\tilde{Y}; Y) \\
&= I(\tilde{Y}; XY) - R_B + I(X; Y) \geq 0.
\end{aligned}$$

So, again the RHS is non-negative. This implies that  $R_A \geq I(\tilde{X}; \tilde{Y}|XY) - \epsilon$ . Hence  $|R_A - I(\tilde{X}; \tilde{Y}|XY)|^+ \leq R_A - I(\tilde{X}; \tilde{Y}|XY) + \epsilon$ . Thus, from (92),

$$\sum_{\mathbf{z} \in \mathcal{E}_{m_A, 1}} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) \leq \exp\left(n\left(R_A - I(\tilde{X}; \tilde{Y}|XY) - I(Z; \tilde{X}\tilde{Y}|XY) + 3\epsilon\right)\right)$$



$$\begin{aligned}
&= \exp\left(n\left(R_A - I(\tilde{X}; \tilde{Y}XY) - I(Z; \tilde{Y}|XY) - I(Z; \tilde{X}|\tilde{Y}XY) + 3\epsilon\right)\right) \\
&= \exp\left(n\left(R_A - I(\tilde{X}; Z\tilde{Y}XY) - I(Z; \tilde{Y}|XY) + 3\epsilon\right)\right) \\
&\leq \exp\left(n\left(R_A - I(\tilde{X}; Z\tilde{Y}) + 3\epsilon\right)\right) \\
&\leq \exp\left(n\left(R_A - I(\tilde{X}; Z|\tilde{Y}) + 3\epsilon\right)\right) \\
&\rightarrow 0 \text{ for } R_A \leq \min_{P_{X\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1^\eta} I(\tilde{X}; Z|\tilde{Y}) - 3\epsilon.
\end{aligned} \tag{93}$$

Taking limit  $\mathcal{P}_1^\eta \rightarrow \mathcal{P}_1^0$  (recall that  $\epsilon < \eta/3$ ), we get the following rate bound

$$R_A \leq \min_{P_{X\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1^0} I(\tilde{X}; Z|\tilde{Y}). \tag{94}$$

Case 3:  $R_A \leq I(\tilde{X}; \tilde{Y}XY)$  and  $I(\tilde{Y}; Y) < R_B$

Using (91), we obtain that

$$\begin{aligned}
-|R_A - I(\tilde{X}; \tilde{Y}Y)|^+ &\leq R_B - I(\tilde{Y}; Y) - I(X; \tilde{X}\tilde{Y}Y) + \epsilon \\
&= R_B - I(\tilde{Y}; Y) - I(X; Y) - I(X; \tilde{Y}|Y) - I(X; \tilde{X}|\tilde{Y}Y) + \epsilon \\
&= R_B - I(\tilde{Y}; XY) - I(X; Y) - I(X; \tilde{X}|\tilde{Y}Y) + \epsilon
\end{aligned}$$

This implies that

$$R_B - I(\tilde{Y}; XY) + \epsilon \geq I(X; Y) + I(X; \tilde{X}|\tilde{Y}Y) - |R_A - I(\tilde{X}; \tilde{Y}Y)|^+.$$

Similar to the previous case, we will argue that RHS of the above inequality is non-negative. When  $R_A \leq I(\tilde{X}; \tilde{Y}Y)$ , it is clearly true. Otherwise, when  $I(\tilde{X}; \tilde{Y}Y) < R_A \leq I(\tilde{X}; \tilde{Y}XY)$ , then

$$\begin{aligned}
I(X; \tilde{X}|\tilde{Y}Y) - |R_A - I(\tilde{X}; \tilde{Y}Y)|^+ &= I(X; \tilde{X}|\tilde{Y}Y) - R_A + I(\tilde{X}; \tilde{Y}Y) \\
&= I(\tilde{X}; \tilde{Y}XY) - R_A \geq 0.
\end{aligned}$$

Thus, for  $R_A \leq I(\tilde{X}; \tilde{Y}XY)$  and  $I(\tilde{Y}; Y) < R_B$ , we have  $R_B - I(\tilde{Y}; XY) + \epsilon \geq 0$ . This implies that  $|R_B - I(\tilde{Y}; XY)|^+ \leq R_B - I(\tilde{Y}; XY) + \epsilon$ . So, from (92),

$$\begin{aligned}
\sum_{\mathbf{z} \in \mathcal{E}_{m_A, 1}(P_{X\tilde{X}\tilde{Y}YZ})} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) &\leq \exp\left(n\left(R_B - I(\tilde{Y}; XY) - I(Z; \tilde{X}\tilde{Y}|XY) + 3\epsilon\right)\right) \\
&= \exp\left(n\left(R_B - I(\tilde{Y}; XY) - I(Z; \tilde{Y}|XY) - I(Z; \tilde{X}|XY\tilde{Y}) + 3\epsilon\right)\right) \\
&= \exp\left(n\left(R_B - I(\tilde{Y}; XYZ) - I(Z; \tilde{X}|XY\tilde{Y}) + 3\epsilon\right)\right) \\
&\leq \exp\left(n\left(R_B - I(\tilde{Y}; XZ) + 3\epsilon\right)\right) \\
&\leq \exp\left(n\left(R_B - I(\tilde{Y}; Z|X) + 3\epsilon\right)\right) \\
&\rightarrow 0 \text{ if } R_B < \min_{P_{X\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1^\eta} I(\tilde{Y}; Z|X) - 3\epsilon.
\end{aligned} \tag{95}$$

Taking limit  $\mathcal{P}_1^\eta \rightarrow \mathcal{P}_1^0$ , we get the rate bound,

$$R_B \leq \min_{P_{X\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1^0} I(\tilde{Y}; Z|X). \tag{96}$$

Case 4:  $I(\tilde{X}; \tilde{Y}XY) < R_A$  and  $I(\tilde{Y}; XY) < R_B$

From (92), we have

$$\begin{aligned}
\sum_{\mathbf{z} \in \mathcal{E}_{m_A, 1}(P_{X\tilde{X}\tilde{Y}YZ})} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) &\leq \exp\left(n\left(R_A - I(\tilde{X}; \tilde{Y}XY) + R_B - I(\tilde{Y}; XY) - I(Z; \tilde{X}\tilde{Y}|XY) + 2\epsilon\right)\right) \\
&\leq \exp\left(n\left(R_A - I(\tilde{X}; XY|\tilde{Y}) + R_B - I(\tilde{Y}; XY) - I(Z; \tilde{X}\tilde{Y}|XY) + 2\epsilon\right)\right) \\
&= \exp\left(n\left(R_A + R_B - I(\tilde{X}\tilde{Y}; XYZ) + 3\epsilon\right)\right) \\
&\leq \exp\left(n\left(R_A + R_B - I(\tilde{X}\tilde{Y}; Z) + 3\epsilon\right)\right) \\
&\rightarrow 0 \text{ if } R_A + R_B < \min_{P_{X\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1^\eta} I(\tilde{X}\tilde{Y}; Z) - 3\epsilon.
\end{aligned} \tag{97}$$

Taking limit  $\mathcal{P}_1^\eta \rightarrow \mathcal{P}_1^0$ , we get the rate bound,

$$R_A + R_B \leq \min_{P_{\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1^0} I(\tilde{X}\tilde{Y}; Z) \quad (98)$$

We define

$$\mathcal{P}_1^+ \stackrel{\text{def}}{=} \{P_{\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_{\tilde{X} \times \tilde{Y} \times \mathcal{X} \times \mathcal{Y} \times Z}^n : P_{XYZ} \in \mathcal{D}_0, P_{\tilde{X}Y'Z} \in \mathcal{D}_0 \text{ for some } P_{Y'|XZ}, P_{X'YZ} \in \mathcal{D}_0 \\ \text{for some } P_{X'|\tilde{Y}Z}, P_X = P_{\tilde{X}} = P_A, P_{\tilde{Y}} = P_B \text{ and } I(\tilde{X}; \tilde{Y}) = 0, I(X; \tilde{Y}) = 0\}$$

We see that  $\mathcal{P}_1^0 \subseteq \mathcal{P}_1^+$ . Using this and collecting (94), (96) and (98), we obtain,

$$R_A \leq \min_{P_{\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1^+} I(\tilde{X}; Z|\tilde{Y}) \quad (99)$$

$$R_B \leq \min_{P_{\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1^+} I(\tilde{Y}; Z|X) \quad (100)$$

$$R_A + R_B \leq \min_{P_{\tilde{X}\tilde{Y}YZ} \in \mathcal{P}_1^+} I(\tilde{X}\tilde{Y}; Z) \quad (101)$$

Now, we move to the analysis of  $W^n(\mathcal{E}_{m_A,2}(P_{X\tilde{Y}_1\tilde{Y}_2YZ})|\mathbf{x}_{m_A}, \mathbf{y})$  (see the second term in the RHS of (89)). We see that by using (17), it is sufficient to consider distribution  $P_{X\tilde{Y}_1\tilde{Y}_2YZ} \in \mathcal{P}_2^\eta$  for which

$$I(X; \tilde{Y}_1\tilde{Y}_2Y) \leq |R_B - I(\tilde{Y}_1; Y)|^+ + |R_B - I(\tilde{Y}_2; \tilde{Y}_1Y)|^+ + \epsilon. \quad (102)$$

For  $P_{X\tilde{Y}_1\tilde{Y}_2YZ} \in \mathcal{P}_2^\eta$  satisfying (102),

$$\begin{aligned} & \sum_{\mathbf{z} \in \mathcal{E}_{m_A,2}(P_{X\tilde{Y}_1\tilde{Y}_2YZ})} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) \\ & \leq \sum_{\substack{\tilde{m}_{B1}, \tilde{m}_{B2}: \tilde{m}_{B1} \neq \tilde{m}_{B2}, \\ (\mathbf{x}_{m_A}, \mathbf{y}^{\tilde{m}_{B1}}, \mathbf{y}^{\tilde{m}_{B2}}) \in T_{X\tilde{Y}_1\tilde{Y}_2Y}^n}} \sum_{\mathbf{z}: (\mathbf{x}_{m_A}, \mathbf{y}^{\tilde{m}_{B1}}, \mathbf{y}^{\tilde{m}_{B2}}, \mathbf{y}, \mathbf{z}) \in T_{X\tilde{Y}_1\tilde{Y}_2YZ}^n} W^n(\mathbf{z}|\mathbf{x}_{m_A}, \mathbf{y}) \\ & \leq \sum_{\substack{\tilde{m}_{B1}, \tilde{m}_{B2}: \tilde{m}_{B1} \neq \tilde{m}_{B2}, \\ (\mathbf{x}_{m_A}, \mathbf{y}^{\tilde{m}_{B1}}, \mathbf{y}^{\tilde{m}_{B2}}) \in T_{X\tilde{Y}_1\tilde{Y}_2Y}^n}} \frac{|T_{Z|X\tilde{Y}_1\tilde{Y}_2Y}^n(\mathbf{x}_{m_A}, \mathbf{y}^{\tilde{m}_{B1}}, \mathbf{y}^{\tilde{m}_{B2}}, \mathbf{y})|}{|T_{Z|XY}^n(\mathbf{x}_{m_A}, \mathbf{y})|} \\ & \leq \sum_{\substack{\tilde{m}_{B1}, \tilde{m}_{B2}: \tilde{m}_{B1} \neq \tilde{m}_{B2}, \\ (\mathbf{x}_{m_A}, \mathbf{y}^{\tilde{m}_{B1}}, \mathbf{y}^{\tilde{m}_{B2}}) \in T_{X\tilde{Y}_1\tilde{Y}_2Y}^n}} \frac{\exp(nH(Z|X\tilde{Y}_1\tilde{Y}_2Y))}{(n+1)^{-|\mathcal{X}||\mathcal{Y}||\mathcal{Z}|} \exp(nH(Z|XY))} \\ & \stackrel{(a)}{\leq} \sum_{\substack{\tilde{m}_{B1}, \tilde{m}_{B2}: \tilde{m}_{B1} \neq \tilde{m}_{B2}, \\ (\mathbf{x}_{m_A}, \mathbf{y}^{\tilde{m}_{B1}}, \mathbf{y}^{\tilde{m}_{B2}}) \in T_{X\tilde{Y}_1\tilde{Y}_2Y}^n}} \exp\left(-n\left(I(Z; \tilde{Y}_1\tilde{Y}_2|XY) - \epsilon\right)\right) \\ & \stackrel{(b)}{\leq} \exp\left(n\left(|R_B - I(\tilde{Y}_1; XY)|^+ + |R_B - I(\tilde{Y}_2; \tilde{Y}_1XY)|^+ - I(Z; \tilde{Y}_1\tilde{Y}_2|XY) + 2\epsilon\right)\right), \end{aligned} \quad (103)$$

where (a) holds for large  $n$  and (b) follows from (18).

Note that, in the analysis of  $W^n(\mathcal{E}_{m_A,1}(P_{X\tilde{X}\tilde{Y}YZ})|\mathbf{x}_{m_A}, \mathbf{y})$  (see the steps leading to (92)), if we replace  $R_A$  with  $R_B$ ,  $\tilde{Y}$  with  $\tilde{Y}_1$  and  $\tilde{X}$  with  $\tilde{Y}_2$ , (92) changes to (103) and the condition (91) on the distribution changes to (102). With these replacements, we see that (103) goes to zero when the following hold (cf. (93),(95),(97)):

$$R_B < I(\tilde{Y}_2; Z\tilde{Y}_1) - 3\epsilon \quad (104)$$

$$R_B < I(\tilde{Y}_1; XZ) - 3\epsilon \quad (105)$$

$$2R_B < I(\tilde{Y}_2; \tilde{Y}_1XYZ) + I(\tilde{Y}_1; XYZ) - 3\epsilon. \quad (106)$$

Here, (106) is obtained by noting that  $I(\tilde{Y}_2; \tilde{Y}_1XY) + I(\tilde{Y}_1; XY) + I(Z; \tilde{Y}_1\tilde{Y}_2|XY) = I(\tilde{Y}_2; \tilde{Y}_1XY) + I(\tilde{Y}_1; XY) + I(Z; \tilde{Y}_1|XY) + I(Z; \tilde{Y}_2|XY\tilde{Y}_1)$ . We first note that the rate bounds in (104), (105) and (106) still hold when

$$R_B < I(\tilde{Y}_2; Z) - 3\epsilon$$

$$R_B < I(\tilde{Y}_1; Z) - 3\epsilon$$

$$2R_B < I(\tilde{Y}_2; Z) + I(\tilde{Y}_1; Z) - 3\epsilon.$$

With this, we proceed similar to the previous case and define

$$\mathcal{P}_2^+ \stackrel{\text{def}}{=} \{P_{X\tilde{Y}_1\tilde{Y}_2YZ} \in \mathcal{P}_{\mathcal{X} \times \mathcal{Y} \times \mathcal{Y} \times \mathcal{Z}}^n : P_{XYZ} \in \mathcal{D}_0, P_{X'_1\tilde{Y}_1Z} \in \mathcal{D}_0 \text{ for some } P_{X'_1|\tilde{Y}_1Z}, \\ P_{X'_2\tilde{Y}_2Z} \in \mathcal{D}_0 \text{ for some } P_{X'_2|\tilde{Y}_2Z}, P_X = P_A, P_{\tilde{Y}_1} = P_{\tilde{Y}_2} = P_B\}.$$

Note that  $\mathcal{P}_2^0 \subseteq \mathcal{P}_2^+$ . Using this and taking limit  $\mathcal{P}_2^\eta \rightarrow \mathcal{P}_2^0$  we get the following rate bounds,

$$R_B \leq \min_{P_{XY\tilde{Y}_1\tilde{Y}_2Z} \in \mathcal{P}_2^0} I(\tilde{Y}_2; Z) \quad (107)$$

$$R_B \leq \min_{P_{XY\tilde{Y}_1\tilde{Y}_2Z} \in \mathcal{P}_2^0} I(\tilde{Y}_1; Z) \quad (108)$$

$$2R_B \leq \min_{P_{XY\tilde{Y}_1\tilde{Y}_2Z} \in \mathcal{P}_2^0} I(\tilde{Y}_2; Z) + I(\tilde{Y}_1; Z) \quad (109)$$

When user A is malicious, error will occur either in *Step 1* or *Step 3* or *Step 5*. Similar to the current case, we can show that error will not happen in *Step 1* w.h.p. because of typicality. For *Step 3* and *Step 5*, we will get bounds of the form (99), (100) and (101). This is because we only consider the candidates which have passed *Step 2*. Hence, we get independence conditions (as we got from Lemma 15 here). Thus, combining (99), (100), (101), (107), (108),(109) and bounds from the case when user A is malicious, we get the following rate region:

$$R_A \leq \min_{P_{XY'X'YZ} \in \mathcal{P}} I(X; Z|Y) \\ R_B \leq \min_{P_{XY'X'YZ} \in \mathcal{P}} I(Y; Z)$$

where  $\mathcal{P}$  is the set of distributions

$$\mathcal{P} \stackrel{\text{def}}{=} \{P_{XY'X'YZ} : P_{XY'Z} = P_A P_{Y'W}, P_{X'YZ} = P_{X'} P_B W, X \perp\!\!\!\perp Y\}.$$

This gives us one corner point (given by (34)) of the rate region, we get the other corner point (given by (33)) by changing the order of decoding by performing *Step 3* before *Step 2*. □

*Proof of Lemma 15.* Adding (C) and (D),

$$\begin{aligned} 2\eta &> D(P_{X'\tilde{Y}Z} \| P_{X'} P_{\tilde{Y}} W) + I(X\tilde{X}; \tilde{Y}Z | X') \\ &= D(P_{X'\tilde{Y}Z} \| P_{X'} P_{\tilde{Y}} W) + D(P_{X'\tilde{Y}X\tilde{X}Z} \| P_{X'} P_{X\tilde{X}|X'} P_{\tilde{Y}Z|X'}) \\ &= \sum_{x', \tilde{y}, x, \tilde{x}, z} P_{X'\tilde{Y}X\tilde{X}Z}(x', \tilde{y}, x, \tilde{x}, z) \left( \log \left\{ \frac{P_{X'\tilde{Y}Z}(x', \tilde{y}, z)}{P_{X'}(x') P_{\tilde{Y}}(\tilde{y}) W(z|x', \tilde{y})} \right\} + \log \left\{ \frac{P_{X'\tilde{Y}X\tilde{X}Z}(x', \tilde{y}, x, \tilde{x}, z)}{P_{X'}(x') P_{X\tilde{X}|X'}(x, \tilde{x}|x') P_{\tilde{Y}Z|X'}(\tilde{y}, z|x')} \right\} \right) \\ &= \sum_{x', \tilde{y}, x, \tilde{x}, z} P_{X'\tilde{Y}X\tilde{X}Z}(x', \tilde{y}, x, \tilde{x}, z) \left( \log \left\{ \frac{P_{X'\tilde{Y}Z}(x', \tilde{y}, z) \times P_{X'\tilde{Y}X\tilde{X}Z}(x', \tilde{y}, x, \tilde{x}, z)}{P_{X'}(x') P_{\tilde{Y}}(\tilde{y}) W(z|x', \tilde{y}) \times P_{X'}(x') P_{X\tilde{X}|X'}(x, \tilde{x}|x') P_{\tilde{Y}Z|X'}(\tilde{y}, z|x')} \right\} \right) \\ &= \sum_{x', \tilde{y}, x, \tilde{x}, z} P_{X'\tilde{Y}X\tilde{X}Z}(x', \tilde{y}, x, \tilde{x}, z) \left( \log \left\{ \frac{P_{X'\tilde{Y}X\tilde{X}Z}(x', \tilde{y}, x, \tilde{x}, z)}{P_{X'}(x') P_{\tilde{Y}}(\tilde{y}) W(z|x', \tilde{y}) P_{X\tilde{X}|X'}(x, \tilde{x}|x')} \right\} \right) \\ &= \sum_{x', \tilde{y}, x, \tilde{x}, z} P_{X'\tilde{Y}X\tilde{X}Z}(x', \tilde{y}, x, \tilde{x}, z) \left( \log \left\{ \frac{P_{X'\tilde{Y}X\tilde{X}Z}(x', \tilde{y}, x, \tilde{x}, z)}{P_{\tilde{Y}}(\tilde{y}) P_{X\tilde{X}}(x', x, \tilde{x}) W(z|x', \tilde{y})} \right\} \right) \\ &= D(P_{X'\tilde{Y}X\tilde{X}Z} \| P_{\tilde{Y}} P_{X\tilde{X}} P_{X'|X\tilde{X}} W) \\ &\stackrel{(a)}{\geq} D(P_{\tilde{Y}X\tilde{X}} \| P_{\tilde{Y}} P_{X\tilde{X}}) \\ &= I(\tilde{Y}; X\tilde{X}), \end{aligned}$$

where (a) follows from the log-sum inequality. Note that  $I(\tilde{Y}; X\tilde{X}) = I(\tilde{Y}; X) + I(\tilde{Y}; \tilde{X}|X) = I(\tilde{Y}; \tilde{X}) + I(\tilde{Y}; X|\tilde{X})$ . Thus,  $I(\tilde{Y}; X\tilde{X}) \leq 2\eta$  implies  $I(\tilde{Y}; \tilde{X}) \leq 2\eta$  and  $I(\tilde{Y}; X) \leq 2\eta$  as mutual information is always non-negative. □

#### APPENDIX D PROOF OF THEOREM 12

We first prove Lemma 16 which gives the randomness reduction argument.

**Lemma 16** (Randomness reduction). *There exists  $n_0(\cdot) : \mathbb{R}^+ \rightarrow \mathbb{N}$  such that given any  $(N_A, N_B, L_A, L_B, n)$  adversary identifying code  $(F_A, F_B, \phi_{F_A, F_B})$  with  $P_e^{\text{rand}}$  denoting its average probability of error and  $\epsilon > 0$ , if*

$$\epsilon > 2 \log(1 + P_e^{\text{rand}}),$$

*and  $n \geq n_0(\epsilon)$  there exists an  $(N_A, N_B, n^2, n^2, n)$  adversary identifying code  $(F'_A, F'_B, \phi_{F'_A, F'_B})$  where the distributions  $p_{F'_A}$  and  $p_{F'_B}$  are the uniform distributions over encoder sets  $\Gamma'_A \subseteq \Gamma_A$  and  $\Gamma'_B \subseteq \Gamma_B$  (with  $|\Gamma'_A| = |\Gamma'_B| = n^2$ ) respectively, and the average probability of error is at most  $\epsilon$ . That is,*

$$\frac{1}{N_A \cdot N_B} \sum_{\substack{m_A \in \mathcal{M}_A \\ m_B \in \mathcal{M}_B}} \sum_{\substack{f'_A \in \Gamma'_A \\ f'_B \in \Gamma'_B}} \frac{1}{n^2 \times n^2} \sum_{\substack{z: \phi_{f'_A, f'_B}(z) \\ \notin \{(m_A, m_B)\}}} W^n(z | f'_A(m_A), f'_B(m_B)) < \epsilon, \quad (110)$$

$$\max_{\substack{\mathbf{x} \in \mathcal{X}^n \\ f'_A \in \Gamma'_A}} \left( \frac{1}{N_B} \sum_{m_B \in \mathcal{M}_B} \sum_{f'_B \in \Gamma'_B} \frac{1}{n^2} \sum_{\substack{z: \phi_{f'_A, f'_B}(z) \\ \notin \{(\mathcal{M}_A \times \{m_B\}) \cup \{\mathbf{a}\}\}}} W^n(z | \mathbf{x}, f'_B(m_B)) \right) < \epsilon \text{ and} \quad (111)$$

$$\max_{\substack{\mathbf{y} \in \mathcal{Y}^n \\ f'_B \in \Gamma'_B}} \left( \frac{1}{N_A} \sum_{m_A \in \mathcal{M}_A} \sum_{f'_A \in \Gamma'_A} \frac{1}{n^2} \sum_{\substack{z: \phi_{f'_A, f'_B}(z) \\ \notin \{(\{m_A\} \times \mathcal{M}_B) \cup \{\mathbf{b}\}\}}} W^n(z | f'_A(m_A), \mathbf{y}) \right) < \epsilon. \quad (112)$$

*Proof.* The proof is along the lines of [28, Lemma 12.8] and Jahn [11, Theorem 1]. Let  $\{F_{A,i}\}, i = 1, \dots, n^2$  be i.i.d. according to  $p_{F_A}$ . Similarly, let  $\{F_{B,j}\}, j = 1, \dots, n^2$  be i.i.d. according to  $p_{F_B}$ . Further, let  $\{F_{A,i}\}_{i=1}^{n^2}$  and  $\{F_{B,j}\}_{j=1}^{n^2}$  be independent.

Define  $e_A(f_A, f_B, \mathbf{x})$  to be the error probability for fixed encoding maps  $f_A$  for user A and  $f_B$  for user B, and the channel inputs chosen by the adversarial user A as  $\mathbf{x}$ , i.e.,

$$e_A(f_A, f_B, \mathbf{x}) \stackrel{\text{def}}{=} \frac{1}{N_B} \sum_{m_B \in \mathcal{M}_B} \sum_{\substack{z: \phi_{f_A, f_B}(z) \\ \notin (\mathcal{M}_A \times \{m_B\}) \cup \{\mathbf{a}\}}} W^n(z | \mathbf{x}, f_B(m_B)).$$

Similarly, for adversarial user B with input  $\mathbf{y} \in \mathcal{Y}^n$ , let  $e_B(f_A, f_B, \mathbf{y})$  be defined as

$$e_B(f_A, f_B, \mathbf{y}) \stackrel{\text{def}}{=} \frac{1}{N_A} \sum_{m_A \in \mathcal{M}_A} \sum_{\substack{z: \phi_{f_A, f_B}(z) \\ \notin (\{m_A\} \times \mathcal{M}_B) \cup \{\mathbf{b}\}}} W^n(z | f_A(m_A), \mathbf{y}).$$

When both users are honest, we define

$$e(f_A, f_B) \stackrel{\text{def}}{=} \frac{1}{N_A N_B} \sum_{(m_A, m_B) \in \mathcal{M}_A \times \mathcal{M}_B} \sum_{\substack{z: \phi_{f_A, f_B}(z) \\ \neq (m_A, m_B)}} W^n(z | f_A(m_A), f_B(m_B)).$$

For any  $j, j' \in [1 : n^2]$ , note that  $e_A(F_{A,j}, F_{B,j'}, \mathbf{x})$ ,  $e_B(F_{A,j}, F_{B,j'}, \mathbf{y})$  and  $e(F_{A,j}, F_{B,j'})$ , as functions of  $F_{A,j}$  and  $F_{B,j'}$ , are random variables. We will show that

$$\mathbb{P} \left\{ \left( \frac{1}{n^2 \times n^2} \sum_{j, j' \in [1 : n^2]} e(F_{A,j}, F_{B,j'}) \geq \epsilon \right) \cup \left( \bigcup_{\mathbf{x} \in \mathcal{X}^n, j \in [1 : n^2]} \left( \frac{1}{n^2} \sum_{j' \in [1 : n^2]} e_A(F_{A,j}, F_{B,j'}, \mathbf{x}) \geq \epsilon \right) \right) \cup \left( \bigcup_{\mathbf{y} \in \mathcal{Y}^n, j' \in [1 : n^2]} \left( \frac{1}{n^2} \sum_{j \in [1 : n^2]} e_B(F_{A,j}, F_{B,j'}, \mathbf{y}) \geq \epsilon \right) \right) \right\} \quad (113)$$

is less than 1. This will imply the existence of  $n^2$  deterministic encoders  $\Gamma'_A = \{f_{A,j} : j \in [1 : n^2]\}$  and  $\Gamma'_B = \{f_{B,j} : j \in [1 : n^2]\}$  satisfying (110), (111) and (112).

For  $j' \in n^2$ , a fixed encoder  $f_A$  for user A and an input vector  $\mathbf{x}$  for malicious user A, let  $e_A^*(f_A, p_{F_B}, \mathbf{x}) \stackrel{\text{def}}{=} \mathbb{E}_{F_{B,j'}} e_A(f_A, F_{B,j'}, \mathbf{x})$ . Note that for the given code  $(F_A, F_B, \phi_{F_A, F_B})$ , the average probability of error when user A is malicious,

$$P_{e, \text{mal}}^{\text{rand}} = \max_{\substack{\mathbf{x} \in \mathcal{X}^n \\ f_A \in \Gamma_A}} e_A^*(f_A, p_{F_B}, \mathbf{x}).$$

For any  $j \in [1 : n^2]$ ,

$$\begin{aligned} \mathbb{P} \left( \frac{1}{n^2} \sum_{j' \in [1:n^2]} e_A(F_{A,j}, F_{B,j'}, \mathbf{x}) \geq \epsilon \right) &= \mathbb{P} \left( \exp \left( \sum_{j' \in [1:n^2]} e_A(F_{A,j}, F_{B,j'}, \mathbf{x}) \right) \geq \exp(n^2 \epsilon) \right) \\ &\leq \exp(-n^2 \epsilon) \mathbb{E} \left[ \exp \left( \sum_{j' \in [1:n^2]} e_A(F_{A,j}, F_{B,j'}, \mathbf{x}) \right) \right] \\ &= \exp(-n^2 \epsilon) \mathbb{E} \left[ \prod_{j' \in [1:n^2]} \exp(e_A(F_{A,j}, F_{B,j'}, \mathbf{x})) \right] \end{aligned}$$

But,

$$\begin{aligned} \mathbb{E} \left[ \prod_{j' \in [1:n^2]} \exp(e_A(F_{A,j}, F_{B,j'}, \mathbf{x})) \right] &= \mathbb{E}_{F_{A,j}} \left[ \mathbb{E}_{(F_{B,1}, \dots, F_{B,n^2}) | F_{A,j}} \left[ \prod_{j' \in [1:n^2]} \exp(e_A(F_{A,j}, F_{B,j'}, \mathbf{x})) \right] \right] \\ &= \sum_{f_A \in \Gamma_A} p_{F_A}(f_A) \mathbb{E}_{(F_{B,1}, \dots, F_{B,n^2}) | F_{A,j}=f_A} \left[ \prod_{j' \in [1:n^2]} \exp(e_A(F_{A,j}, F_{B,j'}, \mathbf{x})) \right] \\ &\stackrel{(a)}{=} \sum_{f_A \in \Gamma_A} p_{F_A}(f_A) \prod_{j' \in [1:n^2]} \mathbb{E}_{F_{B,j'}} [\exp(e_A(f_A, F_{B,j'}, \mathbf{x}))] \\ &\stackrel{(b)}{=} \sum_{f_A \in \Gamma_A} p_{F_A}(f_A) (\mathbb{E}_{F_{B,1}} [\exp(e_A(f_A, F_{B,1}, \mathbf{x}))])^{n^2} \\ &\stackrel{(c)}{\leq} \sum_{f_A \in \Gamma_A} p_{F_A}(f_A) (\mathbb{E}_{F_{B,1}} [1 + e_A(f_A, F_{B,1}, \mathbf{x})])^{n^2} \\ &= \sum_{f_A \in \Gamma_A} p_{F_A}(f_A) (1 + \mathbb{E}_{F_{B,1}} [e_A(f_A, F_{B,1}, \mathbf{x})])^{n^2} \\ &= \sum_{f_A \in \Gamma_A} p_{F_A}(f_A) (1 + e_A^*(f_A, p_{F_B}, \mathbf{x}))^{n^2} \\ &\leq \sum_{f_A \in \Gamma_A} p_{F_A}(f_A) (1 + P_{e, \text{mal A}}^{\text{rand}})^{n^2} \\ &\leq (1 + P_e^{\text{rand}})^{n^2}, \end{aligned}$$

where (a) holds because  $F_{A,j} \perp\!\!\!\perp (F_{B,1}, \dots, F_{B,n^2})$  and  $\{\exp(e_A(f_A, F_{B,j'}, \mathbf{x}))\}_{j'=1}^{n^2}$  are i.i.d. random variables. (b) also follows from the i.i.d. nature of  $\{\exp(e_A(f_A, F_{B,j'}, \mathbf{x}))\}_{j'=1}^{n^2}$ . To see inequality (c), recall that  $\exp(\alpha)$  stands for  $2^\alpha$  which is upper bounded by  $(1 + \alpha)$  for  $0 \leq \alpha \leq 1$ . Thus,

$$\begin{aligned} \mathbb{P} \left( \frac{1}{n^2} \sum_{j' \in [1:n^2]} e_A(F_{A,j}, F_{B,j'}, \mathbf{x}) \geq \epsilon \text{ for any } \mathbf{x} \in \mathcal{X}^n \text{ and } j \in [1 : n^2] \right) \\ \leq |\mathcal{X}|^n \cdot n^2 \cdot \exp(-n^2 \epsilon) (1 + P_e^{\text{rand}})^{n^2} \\ = \exp \left\{ -n^2 \left( \epsilon - \frac{\log |\mathcal{X}|}{n} - \frac{2 \log n}{n^2} - \log(1 + P_e^{\text{rand}}) \right) \right\}. \quad (114) \end{aligned}$$

Similarly, we have

$$\begin{aligned} \mathbb{P} \left( \frac{1}{n^2} \sum_{j \in [1:n^2]} e_B(F_{A,j}, F_{B,j'}, \mathbf{y}) \geq \epsilon \text{ for any } \mathbf{y} \in \mathcal{Y}^n \text{ and } j' \in [1 : n^2] \right) \\ \leq |\mathcal{Y}|^n \cdot n^2 \cdot \exp(-n^2 (\epsilon - \log(1 + P_e^{\text{rand}}))) \\ = \exp \left\{ -n^2 \left( \epsilon - \frac{\log |\mathcal{Y}|}{n} - \frac{2 \log n}{n^2} - \log(1 + P_e^{\text{rand}}) \right) \right\}. \quad (115) \end{aligned}$$

Next, we will compute an upper bound on  $\mathbb{P}\left(\frac{1}{n^2 \times n^2} \sum_{j \in [1:n^2]} \sum_{j' \in [1:n^2]} e(F_{A,j}, F_{B,j'}) \geq \epsilon\right)$ . To this end, let  $\Sigma_{n^2} \stackrel{\text{def}}{=} \{\tau_i : i \in [0 : n^2 - 1]\}$  be a set of permutations (in fact cyclic shifts) of  $(1, 2, \dots, n^2)$  such that

$$\tau_i(j) = (i + j) \bmod n^2 \text{ for all } j \in [1 : n^2].$$

With this,

$$\mathbb{P}\left(\frac{1}{n^2 \times n^2} \sum_{j \in [1:n^2]} \sum_{j' \in [1:n^2]} e(F_{A,j}, F_{B,j'}) \geq \epsilon\right) = \mathbb{P}\left(\frac{1}{n^2} \sum_{\sigma \in \Sigma_{n^2}} \left(\frac{1}{n^2} \sum_{j \in [1:n^2]} e(F_{A,j}, F_{B,\sigma(j)})\right) \geq \epsilon\right).$$

For  $\sigma \in \Sigma_{n^2}$ , let  $P_\sigma \stackrel{\text{def}}{=} \frac{1}{n^2} \sum_{j \in [1:n^2]} e(F_{A,j}, F_{B,\sigma(j)})$ .

$$\begin{aligned} \mathbb{P}\left(\sum_{\sigma \in \Sigma_{n^2}} P_\sigma \geq n^2 \epsilon\right) &\leq \mathbb{P}\left(\cup_{\sigma \in \Sigma_{n^2}} (P_\sigma \geq \epsilon)\right) \\ &\leq \sum_{\sigma \in \Sigma_{n^2}} \mathbb{P}(P_\sigma \geq \epsilon). \end{aligned}$$

Note that for  $\sigma \in \Sigma_{n^2}$ ,  $P_\sigma$  are identically distributed random variables. Thus,

$$\mathbb{P}\left(\frac{1}{n^2 \times n^2} \sum_{j \in [1:n^2]} \sum_{j' \in [1:n^2]} e(F_{A,j}, F_{B,j'}) \geq \epsilon\right) \leq n^2 \mathbb{P}(P_{\tau_0} \geq \epsilon).$$

But,

$$\begin{aligned} \mathbb{P}(P_{\tau_0} \geq \epsilon) &= \mathbb{P}\left(\frac{1}{n^2} \sum_{j \in [1:n^2]} e(F_{A,j}, F_{B,\tau_0(j)}) \geq \epsilon\right) \\ &= \mathbb{P}\left(\frac{1}{n^2} \sum_{j \in [1:n^2]} e(F_{A,j}, F_{B,j}) \geq \epsilon\right) \\ &= \mathbb{P}\left(\sum_{j \in [1:n^2]} e(F_{A,j}, F_{B,j}) \geq n^2 \epsilon\right) \\ &= \mathbb{P}\left(\exp\left(\sum_{j \in [1:n^2]} e(F_{A,j}, F_{B,j})\right) \geq \exp(n^2 \epsilon)\right) \\ &\leq \exp(-n^2 \epsilon) \mathbb{E}\left[\exp\left(\sum_{j \in [1:n^2]} e(F_{A,j}, F_{B,j})\right)\right]. \end{aligned}$$

Note that  $\{e(F_{A,j}, F_{B,j})\}_{j=1}^{n^2}$  are i.i.d. random variables. Hence,

$$\begin{aligned} \mathbb{E}\left[\exp\left(\sum_{j \in [1:n^2]} e(F_{A,j}, F_{B,j})\right)\right] &= \mathbb{E}\left[\prod_{j \in [1:n^2]} \exp(e(F_{A,j}, F_{B,j}))\right] \\ &= \prod_{j \in [1:n^2]} \mathbb{E}[\exp(e(F_{A,j}, F_{B,j}))] \\ &= (\mathbb{E}[\exp(e(F_{A,1}, F_{B,1}))])^{n^2} \\ &\stackrel{(a)}{\leq} (1 + \mathbb{E}(e(F_{A,1}, F_{B,1})))^{n^2} \\ &\stackrel{(b)}{\leq} (1 + P_e^{\text{rand}})^{n^2} \end{aligned}$$

where (a) holds because for  $0 \leq \alpha \leq 1$ ,  $\exp(\alpha) = 2^\alpha \leq 1 + \alpha$  and (b) holds because for the given code  $(F_A, F_B, \phi_{F_A, F_B})$ ,  $P_{e, \text{hon}}^{\text{rand}} = \mathbb{E}(e(F_{A,1}, F_{B,1}))$ . Thus,

$$\mathbb{P}\left(\frac{1}{n^2 \times n^2} \sum_{j \in [1:n^2]} \sum_{j' \in [1:n^2]} e(F_{A,j}, F_{B,j'}) \geq \epsilon\right) \leq n^2 \exp(-n^2 \epsilon) (1 + P_e^{\text{rand}})^{n^2}$$



$$\begin{aligned}
&= \exp(2 \log n - n^2 \epsilon + n^2 \log(1 + P_e^{\text{rand}})) \\
&= \exp\left(-n^2 \left(\epsilon - \frac{2 \log n}{n^2} - \log(1 + P_e^{\text{rand}})\right)\right). \tag{116}
\end{aligned}$$

From (114), (115) and (116), we note that by using a union bound, (113) is upper bounded by

$$\begin{aligned}
&\exp\left\{-n^2 \left(\epsilon - \frac{\log |\mathcal{X}|}{n} - \frac{2 \log n}{n^2} - \log(1 + P_e^{\text{rand}})\right)\right\} + \exp\left\{-n^2 \left(\epsilon - \frac{\log |\mathcal{Y}|}{n} - \frac{2 \log n}{n^2} - \log(1 + P_e^{\text{rand}})\right)\right\} \\
&+ \exp\left(-n^2 \left(\epsilon - \frac{2 \log n}{n^2} - \log(1 + P_e^{\text{rand}})\right)\right),
\end{aligned}$$

which is less than 1 for large enough  $n(=n_0(\epsilon))$  which depends only on the input alphabet sizes and  $\epsilon$ .  $\square$

*Proof of Theorem 12.* We first restate the theorem below.

**Theorem.**  $\mathcal{C} = \mathcal{C}^{\text{rand}}$  whenever  $(R_A, R_B) \in \mathcal{C}$  for some  $R_A, R_B > 0$ .

Since deterministic codes are a subset of randomized codes,  $\mathcal{C} \subseteq \mathcal{C}^{\text{rand}}$ . It only remains to show that  $\mathcal{C} \supseteq \mathcal{C}^{\text{rand}}$  whenever  $(R'_A, R'_B) \in \mathcal{C}$  for some  $R'_A, R'_B > 0$ .

Let  $\epsilon > 0$  (TBD) and large enough  $n$  satisfying  $n \geq n_0(\epsilon)$  (given by Lemma 16). Consider an achievable rate pair  $(R_A, R_B) \in \mathcal{C}^{\text{rand}}$ . This implies that for  $(N_A, N_B) = (2^{nR_A}, 2^{nR_B})$ , there exists an  $(N_A, N_B, L_A, L_B, n)$  adversary identifying code  $(F_A, F_B, \phi_{F_A, F_B})$  with probability of error  $P_e^{\text{rand}}$  which vanishes with  $n$ . Choose  $\epsilon > 0$  such that

$$\epsilon > 2 \log(1 + P_e^{\text{rand}}).$$

By Lemma 16, there exists an  $(N_A, N_B, n^2, n^2, n)$  adversary identifying code  $(F'_A, F'_B, \phi_{F'_A, F'_B})$  where the distributions  $p_{F'_A}$  and  $p_{F'_B}$  on the encoder sets  $\Gamma'_A$  and  $\Gamma'_B$  (with  $|\Gamma'_A| = |\Gamma'_B| = n^2$ ) respectively, are uniform, and the average probability of error is at most  $\epsilon$ . That is,

$$\frac{1}{N_A \cdot N_B} \sum_{\substack{m_A \in \mathcal{M}_A \\ m_B \in \mathcal{M}_B}} \sum_{\substack{l_A \in n^2 \\ l_B \in n^2}} \frac{1}{n^2 \times n^2} \sum_{\substack{z: \phi_{f_{A, l_A}, f_{B, l_B}}(z) \\ \notin \{(m_A, m_B)\}}} W^n(z | f_{A, l_A}(m_A), f_{B, l_B}(m_B)) < \epsilon, \tag{117}$$

$$\max_{\substack{\mathbf{x} \in \mathcal{X}^n \\ l_A \in n^2}} \left( \frac{1}{N_B} \sum_{m_B \in \mathcal{M}_B} \sum_{l_B \in n^2} \frac{1}{n^2} \sum_{\substack{z: \phi_{f_{A, l_A}, f_{B, l_B}}(z) \\ \notin \{(\mathcal{M}_A \times \{m_B\}) \cup \{\mathbf{a}\}\}}} W^n(z | \mathbf{x}, f_{B, l_B}) \right) < \epsilon \text{ and} \tag{118}$$

$$\max_{\substack{\mathbf{y} \in \mathcal{Y}^n \\ l_B \in n^2}} \left( \frac{1}{N_A} \sum_{m_A \in \mathcal{M}_A} \sum_{l_A \in n^2} \frac{1}{n^2} \sum_{\substack{z: \phi_{f_{A, l_A}, f_{B, l_B}}(z) \\ \notin \{(\{m_A\} \times \mathcal{M}_B) \cup \{\mathbf{b}\}\}}} W^n(z | f_{A, l_A}(m_A), \mathbf{y}) \right) < \epsilon. \tag{119}$$

Further, since  $(R'_A, R'_B) \in \mathcal{C}$  for  $R'_A, R'_B > 0$  (i.e.  $(R'_A, R'_B)$  is an achievable rate pair), there exists an  $(n^2, n^2, k_n)$  code  $(\hat{f}_A, \hat{f}_B, \hat{\phi})$  where  $k_n/n \rightarrow 0$  and

$$P_e(\hat{f}_A, \hat{f}_B, \hat{\phi}) \leq \epsilon \tag{120}$$

for large enough  $n$ . We choose sufficiently large  $n$  such that (117), (118), (119) and (120) hold. For a vector sequence  $\tilde{\mathbf{s}} \in \mathcal{S}^{k_n+n}$  for any alphabet  $\mathcal{S}$ , we write  $\tilde{\mathbf{s}} = (\hat{\mathbf{s}}, \mathbf{s})$ , where  $\hat{\mathbf{s}}$  denotes the first  $k_n$ -length part of  $\tilde{\mathbf{s}}$  and  $\mathbf{s}$  denotes the last  $n$ -length part of the  $\tilde{\mathbf{s}}$ . Let  $(\tilde{f}_A, \tilde{f}_B, \tilde{\phi})$  be a new  $(\tilde{N}_A, \tilde{N}_B, \tilde{n})$  code where  $\tilde{n} := k_n + n$ , message set for user- $i \in \{A, B\}$ ,  $\tilde{\mathcal{M}}_i = [1 : \tilde{N}_i] := \{1, 2, \dots, n^2\} \times [1 : N_i]$ . Further, for  $l_A \in [1 : n^2]$ ,  $m_A \in [1 : N_A]$ , let  $\tilde{m}_A := (l_A, m_A)$ . We define  $\tilde{f}_A(\tilde{m}_A) = \tilde{f}_A(l_A, m_A) := (\hat{f}_A(l_A), f_{A, l_A}(m_A))$ . For  $\tilde{\mathbf{z}} = (\hat{\mathbf{z}}, \mathbf{z})$ , if  $\hat{\phi}(\hat{\mathbf{z}}) = (\hat{l}_A, \hat{l}_B)$ , we define  $\tilde{\phi}(\tilde{\mathbf{z}}) = \phi_{f_{A, \hat{l}_A}, f_{B, \hat{l}_B}}(\mathbf{z})$ . Otherwise, if  $\hat{\phi}(\hat{\mathbf{z}}) \in \{\mathbf{a}, \mathbf{b}\}$ ,  $\tilde{\phi}(\tilde{\mathbf{z}}) = \hat{\phi}(\hat{\mathbf{z}})$ . Then, for the new code,

$$\begin{aligned}
P_{e, \text{mal B}} &= \max_{\tilde{\mathbf{y}} \in \mathcal{Y}^{\tilde{n}}} \frac{1}{\tilde{N}_A} \sum_{\tilde{m}_A \in \tilde{\mathcal{M}}_A} \mathbb{P}\left(\left(\tilde{\phi}(\tilde{\mathbf{Z}}) \notin (\{\tilde{m}_A\} \times \tilde{\mathcal{M}}_B) \cup \{\mathbf{b}\}\right) \middle| \tilde{\mathbf{X}} = \tilde{f}_A(\tilde{m}_A), \tilde{\mathbf{Y}} = \tilde{\mathbf{y}}\right) \\
&= \max_{(\tilde{\mathbf{y}} \times \mathbf{y}) \in \mathcal{Y}^{\tilde{n}}} \left( \frac{1}{n^2 N_A} \sum_{(l_A, m_A) \in \tilde{\mathcal{M}}_A} \mathbb{P}\left(\left(\hat{\phi}(\hat{\mathbf{Z}}) \notin (\{\{l_A\} \times [1 : n^2]\} \cup \{\mathbf{b}\})\right) \cup \right.
\end{aligned}$$

$$\begin{aligned}
& \left( \hat{\phi}(\hat{\mathbf{Z}}) = (l_A, l_B) \text{ for some } l_B, \phi_{f_{l_A}, f_{l_B}}(\mathbf{Z}) \notin (\{m_A\} \times \mathcal{M}_B) \cup \{\mathbf{b}\} \mid \tilde{\mathbf{X}} = \tilde{f}_A(l_A, m_A), \tilde{\mathbf{Y}} = (\hat{\mathbf{y}} \times \mathbf{y}) \right) \\
& \leq \max_{\hat{\mathbf{y}} \in \mathcal{Y}^{k_n}} \frac{1}{n^2} \sum_{l_A \in [1:n^2]} \mathbb{P} \left( \left( \hat{\phi}(\hat{\mathbf{Z}}) \notin \{(\{l_A\} \times [1:n^2])\} \cup \{\mathbf{b}\} \right) \mid \hat{\mathbf{X}} = \hat{f}_A(l_A), \hat{\mathbf{Y}} = \hat{\mathbf{y}} \right) \\
& \quad + \max_{\substack{\mathbf{y} \in \mathcal{Y}^n \\ l_B \in [1:n^2]}} \frac{1}{n^2 N_A} \sum_{\substack{l_A \in [1:n^2] \\ m_A \in \mathcal{M}_A}} \mathbb{P} \left( \phi_{f_{l_A}, f_{l_B}}(\mathbf{Z}) \notin (\{m_A\} \times \mathcal{M}_B) \cup \{\mathbf{b}\} \mid \mathbf{X} = f_{A, l_A}(m_A), \mathbf{Y} = \mathbf{y} \right) \\
& \leq 2\epsilon
\end{aligned}$$

where the last inequality follows from (119) and (120). Similarly, we can argue that  $P_{e, \text{mal A}} \leq 2\epsilon$ . Next, we note from (6) that  $P_{e, \text{hon}} \leq P_{e, \text{mal A}} + P_{e, \text{mal B}} \leq 4\epsilon$ .  $\square$