

LIPSCHITZ ESTIMATES AND AN APPLICATION TO TRACE FORMULAE

TIRTHANKAR BHATTACHARYYA, ARUP CHATTOPADHYAY, SAIKAT GIRI,
AND CHANDAN PRADHAN

ABSTRACT. In this note, we provide an elementary proof for the expression of $f(U) - f(V)$ in the form of a double operator integral for every Lipschitz function f on the unit circle \mathbb{T} and for a pair of unitary operators (U, V) with $U - V \in \mathcal{S}_2(\mathcal{H})$ (the Hilbert-Schmidt class). As a consequence, we obtain the Schatten 2-Lipschitz estimate $\|f(U) - f(V)\|_2 \leq \|f\|_{\text{Lip}(\mathbb{T})} \|U - V\|_2$ for all Lipschitz functions $f : \mathbb{T} \rightarrow \mathbb{C}$. Moreover, we develop an approach to the operator Lipschitz estimate for a pair of contractions with the assumption that one of them is a strict contraction, which significantly extends the class of functions from results known earlier. More specifically, for each $p \in (1, \infty)$ and for every pair of contractions (T_0, T_1) with $\|T_0\| < 1$, there exists a constant $d_{f,p,T_0} > 0$ such that $\|f(T_1) - f(T_0)\|_p \leq d_{f,p,T_0} \|T_1 - T_0\|_p$ for all Lipschitz functions on \mathbb{T} . Using our Lipschitz estimates, we establish a modified Krein trace formula applicable to a specific category of pairs of contractions featuring Hilbert-Schmidt perturbations.

1. INTRODUCTION

Given a complex separable Hilbert space \mathcal{H} , the algebra of bounded linear operators acting on \mathcal{H} will be denoted by $\mathcal{B}(\mathcal{H})$. For $p \in [1, \infty)$, the Schatten p -class $\mathcal{S}_p(\mathcal{H})$ is

$$\mathcal{S}_p(\mathcal{H}) = \left\{ A \in \mathcal{B}(\mathcal{H}) : \|A\|_p \stackrel{\text{def}}{=} \left(\sum_{n \in \mathbb{N}} s_n(A)^p \right)^{1/p} < \infty \right\},$$

where $s_n(A)$ are the singular values of A . The unsuffixed $\|A\|$ will always mean the operator norm.

An intriguing question in perturbation theory is to decide to what extent a function $f(A)$ of an operator A can change when the operator A undergoes a small perturbation.

Definition 1.1. *Let I be an interval in \mathbb{R} . Let $f : I \rightarrow \mathbb{C}$ be a continuous function. Then*

- (1) *f is called operator Lipschitz if there is a constant C_f such that*

$$\|f(A) - f(B)\| \leq C_f \|A - B\| \tag{1.1}$$

for all self-adjoint operators A, B with $\sigma(A) \cup \sigma(B) \subset I$ and $A - B \in \mathcal{B}(\mathcal{H})$.

- (2) *f is called \mathcal{S}_p -Lipschitz for $1 \leq p < \infty$ if there exists a constant $C_{f,p}$ such that*

$$\|f(A) - f(B)\|_p \leq C_{f,p} \|A - B\|_p, \tag{1.2}$$

for all self-adjoint operators A, B with $\sigma(A) \cup \sigma(B) \subset I$ and $A - B \in \mathcal{S}_p(\mathcal{H})$.

2010 *Mathematics Subject Classification.* 47A20, 47A55, 47A56, 47B10, 42B30, 30H10.

Key words and phrases. Unitary operators, Contractions, Unitary dilations, Lipschitz functions.

It is a natural guess that the \mathcal{S}_p -Lipschitz functions depend on p . A surprising result of Potapov and Sukochev [35] shows that for $p \in (1, \infty)$, these classes of functions are same and coincide with the class of scalar Lipschitz functions on \mathbb{R} . Indeed, they proved the following theorem.

Theorem 1.2. [35, Theorem 1] *Let f be a Lipschitz function on \mathbb{R} . For every $1 < p < \infty$ there is a constant c_p such that*

$$\|f(A) - f(B)\|_p \leq c_p \|f\|_{\text{Lip}(\mathbb{R})} \|A - B\|_p, \quad (1.3)$$

where A and B are self-adjoint (possibly unbounded) linear operators such that $A - B \in \mathcal{S}_p(\mathcal{H})$.

The above result resolved a number of conjectures of Krein [22], Farforovskaya [16] and Peller [33]. However, for $p = 1$ (due to E. B. Davies [13]) and for $p = \infty$ (due to Farforovskaya [14], [15] and Kato [17]), the classes of \mathcal{S}_p -Lipschitz functions are smaller than the class of Lipschitz functions on \mathbb{R} . For more on the history of this subject we refer the reader to the recent survey [3] by Aleksandrov and Peller.

Recently, in [6] an analogue of Theorem 1.2 was obtained in the unitary setting, as follows.

Theorem 1.3. [6, Theorem 2] *For each $p \in (1, \infty)$, there is a constant $d_p > 0$ such that*

$$\|f(U_1) - f(U_0)\|_p \leq d_p \|f\|_{\mathcal{L}\text{ip}} \|U_1 - U_0\|_p, \text{ where } \|f\|_{\mathcal{L}\text{ip}} = \sup_{\substack{\lambda, \mu \in \mathbb{R} \\ \lambda \neq \mu}} \frac{|f(e^{i\lambda}) - f(e^{i\mu})|}{|\lambda - \mu|}, \quad (1.4)$$

for all Lipschitz functions $f : \mathbb{T} \rightarrow \mathbb{C}$ and all unitary operators U_0 and U_1 such that $U - U_0 \in \mathcal{S}_p(\mathcal{H})$. Further $d_p \leq 32(c_p + 9)$, where c_p is given by Theorem 1.2.

It was shown in [10] that $c_p = O(p^2/p - 1)$, which is the best constant known.

Going forward, we denote the open unit disk of \mathbb{C} by \mathbb{D} , the unit circle by \mathbb{T} and the disk algebra by $A(\mathbb{D})$. It is natural to identify the disk algebra $A(\mathbb{D})$ with the Banach subalgebra $A(\mathbb{T})$ of all continuous functions on the unit circle \mathbb{T} with vanishing negative Fourier coefficients and the natural norm of $C(\mathbb{T})$.

The $\mathcal{S}_p(\mathcal{H})$ -Lipschitz property has been studied for unitaries (see [6]), for contractions (with analytic functions, see [18], [19]), dissipative operators (see [1], [5]) and for normal operators (see [2], [19], [31]).

An operator $T \in \mathcal{B}(\mathcal{H})$ is a contraction if $\|T\| \leq 1$ and is a strict contraction if the inequality is strict. For every contraction T we have the defect operators $D_T = (I - T^*T)^{1/2}$ and $D_{T^*} = (I - TT^*)^{1/2}$.

Definition 1.4. *For $\alpha > 0$ and a domain Ω in \mathbb{C} , a function $f : \Omega \rightarrow \mathbb{C}$ is said to be α -Lipschitz if there is a constant $C > 0$ such that*

$$|f(z) - f(w)| \leq C|z - w|^\alpha, \quad (1.5)$$

for arbitrary $z, w \in \Omega$ (we use the notation $\text{Lip}_\alpha(\Omega)$ for the class of α -Lipschitz functions on Ω , and $\text{Lip}(\Omega)$ for $\text{Lip}_1(\Omega)$). The Lipschitz norm $\|f\|_{\text{Lip}(\Omega)}$ of f is the smallest positive value of C .

There are three main objectives of this paper:

- (1) to obtain \mathcal{S}_2 -Lipschitz estimate for pair of unitaries. This improves the constant d_2 over the existing estimates in [19, Corollary 6.1] and Theorem 1.3.
- (2) to translate the result of Theorem 1.3 from unitaries to contractions. This extends the class of functions from earlier known results.

(3) to establish a modified Krein trace formula for pairs of contractions.

The celebrated expression of $f(U) - f(V)$ for every f with $f' \in \text{Lip}_\alpha(\mathbb{T})$ in terms of double operator integrals (see Equation (2.8)) established by Birman and Solomyak in [7], holds true for a large class of scalar-valued functions f , viz., for any $f \in \text{Lip}(\mathbb{T})$. Indeed, $f' \in \text{Lip}_\alpha(\mathbb{T})$ implies $f \in C^1(\mathbb{T})$, which further implies $f \in \text{Lip}(\mathbb{T})$, but $f \in \text{Lip}(\mathbb{T})$ does not necessarily imply $f' \in \text{Lip}_\alpha(\mathbb{T})$. We give an elementary proof of the expression of $f(U) - f(V)$ in terms of a double operator integral in Theorem 2.4. Note that the original proof of Birman and Solomyak cannot be adapted unless $f' \in \text{Lip}_\alpha(\mathbb{T})$. As a consequence of the expression, we obtain the \mathcal{S}_2 -Lipschitz estimate for a pair of unitaries.

We would like to highlight a novel approach adopted here, viz., the Schäffer dilation for contractions [40]. The \mathcal{S}_p -Lipschitz property for pairs of contractions has been studied in literature (cf. [18, Theorem 4.2], [19, Theorem 6.4]). The existent results are for Lipschitz functions in the disk algebra $\mathcal{A}(\mathbb{T})$. We extend this to arbitrary Lipschitz functions on the unit circle when one of the contractions is strict. Our estimates yields a modified version of the celebrated Krein's trace formula (see Theorem 4.2) for certain pair of contractions.

The article is structured as follows. In Section 2, we will give a brief discussion on the double operator integral. En route, we establish the operator Lipschitz estimate for pairs of unitaries via the double operator integral technique. Section 3 deals with Lipschitz estimates for a pair of contractions when one of them is strict. In Section 4, we establish the modified Krein trace formula. We end this note by proposing an open problem in Section 5.

2. THE DOUBLE OPERATOR INTEGRAL

This section is devoted to obtain an operator Lipschitz estimate for a pair of unitaries on the Hilbert-Schmidt ideal $\mathcal{S}_2(\mathcal{H})$ via the double operator integral technique. We start with a brief discussion on the construction and its algebraic properties which will be used throughout this section. Such an approach was first suggested by Birman and Solomyak in [7] (see also [8], [9]).

Let $E(\cdot)$ and $F(\cdot)$ be two spectral measures given on the Borel σ -algebra $\mathcal{B}([0, 2\pi])$. We construct a new spectral measure on the product σ -algebra $\mathcal{B}([0, 2\pi]) \times \mathcal{B}([0, 2\pi])$, Define it on product sets by

$$G(\delta \times \sigma)(X) := E(\delta)XF(\sigma), \quad \forall X \in \mathcal{S}_2(\mathcal{H}) \quad (2.1)$$

and extend using standard techniques.

Definition 2.1. Let $\Phi \in L^\infty([0, 2\pi] \times [0, 2\pi], G)$, where G is defined as in (2.1). The Birman-Solomyak double operator integral $T_\Phi^G : \mathcal{S}_2(\mathcal{H}) \rightarrow \mathcal{S}_2(\mathcal{H})$ is defined as the integral of the symbol Φ with respect to the spectral measure $G(\cdot)$, that is,

$$T_\Phi^G(X) := \int_0^{2\pi} \int_0^{2\pi} \Phi(\lambda_1, \lambda_2) dG(\lambda_1, \lambda_2)(X), \quad X \in \mathcal{S}_2(\mathcal{H}).$$

The notation

$$T_\Phi^G(X) := \int_0^{2\pi} \int_0^{2\pi} \Phi(\lambda_1, \lambda_2) dE(\lambda_1) X dF(\lambda_2), \quad X \in \mathcal{S}_2(\mathcal{H}), \quad (2.2)$$

will be used frequently. We note that T_Φ^G is bounded on $\mathcal{S}_2(\mathcal{H})$ and $\|T_\Phi^G\| = \|\Phi\|_\infty$.

The following properties of T_Φ^G on $\mathcal{S}_2(\mathcal{H})$ are useful:

- (a) $T_{\alpha\phi+\beta\psi} = \alpha T_\phi + \beta T_\psi$, $\alpha, \beta \in \mathbb{C}$;

(b) $T_{\phi\psi} = T_\phi T_\psi$;

(c) If $\Phi(\lambda_1, \lambda_2) = \phi_1(\lambda_1) \cdot \phi_2(\lambda_2)$ for some $\Phi, \phi_1, \phi_2 \in L^\infty([0, 2\pi] \times [0, 2\pi], G)$, then

$$T_\Phi^G(X) := \left(\int_0^{2\pi} \phi_1(\lambda_1) dE(\lambda_1) \right) \cdot X \cdot \left(\int_0^{2\pi} \phi_2(\lambda_2) dF(\lambda_2) \right)$$

holds for all $X \in \mathcal{S}_2(\mathcal{H})$.

The first main result in this section can now be stated.

Theorem 2.2. *Let $f : \mathbb{T} \rightarrow \mathbb{C}$ be a Lipschitz function on \mathbb{T} . Then for any pair of unitaries U, V and for $X \in \mathcal{S}_2(\mathcal{H})$, we have*

$$\begin{aligned} f(U)X - Xf(V) &= T_{h^{[1]}}^G(UX - XV), \text{ and} \\ \|f(U)X - Xf(V)\|_2 &\leq \|f\|_{\text{Lip}(\mathbb{T})} \|UX - XV\|_2, \end{aligned} \quad (2.3)$$

where

$$h^{[1]}(\lambda, \mu) = \begin{cases} \frac{f(e^{i\lambda}) - f(e^{i\mu})}{e^{i\lambda} - e^{i\mu}} & \text{if } \lambda \neq \mu \\ 0 & \text{if } \lambda = \mu. \end{cases} \quad (2.4)$$

Proof. By the spectral theorem we have

$$U = \int_0^{2\pi} e^{i\lambda} dE(\lambda) \quad \text{and} \quad V = \int_0^{2\pi} e^{i\mu} dF(\mu), \quad (2.5)$$

where $E(\cdot)$ and $F(\cdot)$ are the spectral measures associated with the unitaries U and V respectively. Let $p_1(\lambda, \mu) = e^{i\lambda}$ and $p_2(\lambda, \mu) = e^{i\mu}$. Therefore $T_{p_1}^G(X) = UX$, $T_{p_2}^G(X) = XV$ and $T_{f \circ p_1}^G(X) = f(U)X$, $T_{f \circ p_2}^G(X) = Xf(V)$, which are the immediate implications of (c).

Again (a) and (b) imply

$$\begin{aligned} f(U)X - Xf(V) &= T_{f \circ p_1}^G(X) - T_{f \circ p_2}^G(X) \\ &= T_{(f \circ p_1) - (f \circ p_2)}^G(X) \\ &= T_{(p_1 - p_2)h^{[1]}}^G(X) \\ &= T_{h^{[1]}}^G(T_{p_1}^G(X) - T_{p_2}^G(X)) = T_{h^{[1]}}^G(UX - XV), \end{aligned} \quad (2.6)$$

where $h^{[1]}(\cdot, \cdot)$ is as given in (2.4). Therefore, (2.6) further implies that

$$\|f(U)X - Xf(V)\|_2 \leq \|h^{[1]}\|_\infty \|UX - XV\|_2. \quad (2.7)$$

This completes the proof of the theorem as $\|h^{[1]}\|_\infty = \|f\|_{\text{Lip}(\mathbb{T})}$. \square

We shall need the following theorem which is a special case of [11, Theorem 2.2]. The original idea goes back to Voiculescu [42].

Theorem 2.3. *Let T be a normal contraction on \mathcal{H} and $V \in \mathcal{S}_2(\mathcal{H})$. Then there exists a sequence $\{P_n\}$ of finite rank projections, such that $\{P_n\} \uparrow I$ and each of the following terms*

$$(i) \|P_n^\perp T P_n\|_2, \quad (ii) \|P_n T P_n^\perp\|_2, \quad (iii) \|P_n^\perp V\|_2, \quad (iv) \|P_n^\perp V^*\|_2$$

converge to 0 as $n \rightarrow \infty$.

Taking E and F in equation (2.2) to be the spectral measures of U and V respectively, we know from [7, Theorem 11] that $T_{h^{[1]}}^G(U - V)$ is a Hilbert-Schmidt operator and, in fact, equals $f(U) - f(V)$ for $f' \in \text{Lip}_\alpha(\mathbb{T})$ for $\alpha > 0$.

Theorem 2.4. *Let U and V be two unitary operators on \mathcal{H} such that $U - V \in \mathcal{S}_2(\mathcal{H})$. Let $f : \mathbb{T} \rightarrow \mathbb{C}$ be a Lipschitz function on \mathbb{T} , then*

$$f(U) - f(V) = \int_0^{2\pi} \int_0^{2\pi} \frac{f(e^{i\lambda}) - f(e^{i\mu})}{e^{i\lambda} - e^{i\mu}} dE(\lambda) (U - V) dF(\mu). \quad (2.8)$$

Furthermore,

$$\|f(U) - f(V)\|_2 \leq \|f\|_{\text{Lip}(\mathbb{T})} \|U - V\|_2. \quad (2.9)$$

Proof. Applying Theorem 2.2 to $(f(U)P_n - P_nf(V))$, where P_n are as in Theorem 2.3, we get

$$\|f(U)P_n - P_nf(V)\|_2 \leq \|f\|_{\text{Lip}(\mathbb{T})} \|UP_n - P_nV\|_2, \quad (2.10)$$

Again, note that

$$\begin{aligned} \|(UP_n - P_nV) - (U - V)\|_2 &= \|UP_n^\perp - P_n^\perp V\|_2 \\ &= \|P_nUP_n^\perp + P_n^\perp UP_n^\perp - P_n^\perp VP_n - P_n^\perp VP_n^\perp\|_2 \\ &\leq \|P_nUP_n^\perp\|_2 + \|P_n^\perp VP_n\|_2 + \|P_n^\perp(U - V)\|_2. \end{aligned} \quad (2.11)$$

Therefore, from (2.10) and (2.11), we have $\|(UP_n - P_nV) - (U - V)\|_2 \rightarrow 0$ as $n \rightarrow \infty$. As a result,

$$\begin{aligned} \lim_{n \rightarrow \infty} \|T_{h^{[1]}}^G(UP_n - P_nV) - T_{h^{[1]}}^G(U - V)\|_2 &= 0 \quad \text{and} \\ \lim_{n \rightarrow \infty} \|(f(U)P_n - P_nf(V)) - T_{h^{[1]}}^G(U - V)\| &= 0, \end{aligned}$$

where $h^{[1]}$ and $G(\cdot)$ are the same as in (2.4) and (2.1) respectively. Hence

$$\begin{aligned} f(U) - f(V) &= T_{h^{[1]}}^G(U - V) \\ &= \int_0^{2\pi} \int_0^{2\pi} \frac{f(e^{i\lambda}) - f(e^{i\mu})}{e^{i\lambda} - e^{i\mu}} dE(\lambda) (U - V) dF(\mu), \end{aligned}$$

which implies

$$\|f(U) - f(V)\|_2 \leq \|f\|_{\text{Lip}(\mathbb{T})} \|U - V\|_2.$$

□

Remark 2.5. *Note that $\|f\|_{\mathcal{L}\text{ip}} \leq \|f\|_{\text{Lip}(\mathbb{T})}$ for every Lipschitz function $f : \mathbb{T} \rightarrow \mathbb{C}$. It is worth mentioning that $|\lambda - \mu| \leq \pi$ implies $|e^{i\lambda} - e^{i\mu}| \geq \frac{2}{\pi}|\lambda - \mu|$, and this further implies that $\|f\|_{\text{Lip}(\mathbb{T})} \leq \frac{\pi}{2}\|f\|_{\mathcal{L}\text{ip}}$. Therefore, for two unitary operators U and V , if $|\lambda - \mu| \leq \pi$ for any $e^{i\lambda} \in \sigma(U)$ and $e^{i\mu} \in \sigma(V)$, then (2.9) establishes a better estimate compared to the previously known estimate in Theorem 1.3.*

3. OPERATOR LIPSCHITZ ESTIMATES FOR PAIR OF CONTRACTIONS

The goal of this section is prove one of the central results of the paper. This concerns the operator Lipschitz estimates for pair of contractions. The proof is based on the Lipschitz properties of Schäffer unitary dilations of contractions. We start by reviewing the construction of a functional calculus $f \mapsto f(T)$ for a contraction T , which will be used in the course of the investigation.

Let $N \in \mathbb{N}$. For a trigonometric polynomial $p_N(z) = \sum_{n=-N}^N a_n z^n$, $a_n \in \mathbb{C}$, define $p_N(T)$ as follows

$$p_N(T) = \sum_{n=-N}^{-1} a_n T^{*|n|} + \sum_{n=0}^N a_n T^n, \quad (3.1)$$

for a contraction T .

Definition 3.1. (Power unitary dilation) *Let T be a contraction on \mathcal{H} . We say that T has a power unitary dilation if there exists another Hilbert space \mathcal{K} containing \mathcal{H} as a subspace, and a unitary operator $U_T : \mathcal{K} \rightarrow \mathcal{K}$ such that*

$$T^n = P_{\mathcal{H}} U_T^n|_{\mathcal{H}} \quad \text{for } n \in \mathbb{Z}, \quad (3.2)$$

where $T^{-|n|} = (T^*)^{|n|}$, and $P_{\mathcal{H}}$ is the orthogonal projection from \mathcal{K} onto \mathcal{H} . Hence, for any trigonometric polynomial p ,

$$p(T) = P_{\mathcal{H}} p(U_T)|_{\mathcal{H}}. \quad (3.3)$$

Now the following lemma extends the definition (3.3) to every $f \in C(\mathbb{T})$.

Lemma 3.2. *Let $f \in C(\mathbb{T})$ and $T \in \mathcal{B}(\mathcal{H})$ be a contraction. Let U_T be a power unitary dilation of T on a bigger Hilbert space \mathcal{K} . Define*

$$f(T) = P_{\mathcal{H}} f(U_T)|_{\mathcal{H}}. \quad (3.4)$$

Here $P_{\mathcal{H}}$ denotes the orthogonal projection from \mathcal{K} onto \mathcal{H} . Then $f(T)$ is a bounded operator satisfying the von Neumann inequality

$$\|f(T)\| \leq \|f\|_{\infty}, \quad (3.5)$$

and $f(T)$ is independent of the choice of the power unitary dilation U_T . This functional calculus is linear but not multiplicative.

Proof. Let U_T and V_T be two power unitary dilations of T on Hilbert spaces $\mathcal{K}_1 \supset \mathcal{H}$ and $\mathcal{K}_2 \supset \mathcal{H}$ respectively. Now, for a trigonometric polynomial p , consider the following inequality

$$\|P_{\mathcal{H}} f(U_T)|_{\mathcal{H}} - P_{\mathcal{H}} f(V_T)|_{\mathcal{H}}\| \leq \|P_{\mathcal{H}} f(U_T)|_{\mathcal{H}} - P_{\mathcal{H}} p(U_T)|_{\mathcal{H}}\| + \|P_{\mathcal{H}} f(V_T)|_{\mathcal{H}} - P_{\mathcal{H}} p(V_T)|_{\mathcal{H}}\|. \quad (3.6)$$

Finally, from (3.6) using (3.5) and the denseness property of trigonometric polynomials in $C(\mathbb{T})$, we conclude the proof of the lemma. \square

As is well-known, there is a trade off between the conditions on the operators and the conditions on the functions when proving perturbation estimates. The following result has fewer conditions on the operators whereas Theorem 3.9 has fewer conditions on the functions involved.

Proposition 3.3. *Let T_0 and T_1 be two contractions on \mathcal{H} such that $T_1 - T_0 \in \mathcal{S}_p(\mathcal{H})$, $1 \leq p \leq \infty$. Let f be a function on the unit circle \mathbb{T} such that $\sum_{n \in \mathbb{Z}} |n \hat{f}(n)| < \infty$. Then*

$$\|f(T_1) - f(T_0)\|_p \leq \left(\sum_{n \in \mathbb{Z} \setminus \{0\}} |n \hat{f}(n)| \right) \|T_1 - T_0\|_p. \quad (3.7)$$

Proof. Let T_0, T_1 be two contractions. Then

$$f(T_1) - f(T_0) = \sum_{n=1}^{\infty} \hat{f}(-n) (T_1^{*n} - T_0^{*n}) + \sum_{n=1}^{\infty} \hat{f}(n) (T_1^n - T_0^n),$$

which implies

$$\|f(T_1) - f(T_0)\|_p \leq \left(\sum_{n \in \mathbb{Z} \setminus \{0\}} |n \hat{f}(n)| \right) \|T_1 - T_0\|_p.$$

□

In our journey towards getting results with fewer conditions on the function, we first obtain a preliminary result by restricting the class of operators,

Lemma 3.4. *If $f \in \text{Lip}(\mathbb{T})$ satisfies $\sum_{n \in \mathbb{Z}} |n \hat{f}(n)| < \infty$, and if both T_0 and T_1 are strict contractions such that $T_1 - T_0 \in \mathcal{S}_p(\mathcal{H})$, then*

$$\|f(T_1) - f(T_0)\|_p \leq \sqrt{2} \|f\|_{\text{Lip}(\mathbb{T})} (1 - \max\{\|T_1\|, \|T_0\|\}^2)^{-1/2} \|T_1 - T_0\|_p. \quad (3.8)$$

Proof. Suppose $0 \leq \|T_0\| \leq \|T_1\| < 1$. Then

$$\begin{aligned} \|f(T_1) - f(T_0)\|_p &\leq \left(\sum_{n \in \mathbb{Z} \setminus \{0\}} |n \hat{f}(n)| \|T_1\|^{|n|-1} \right) \|T_1 - T_0\|_p \\ &\leq \sqrt{2} \left[\sum_{n \in \mathbb{Z} \setminus \{0\}} |n \hat{f}(n)|^2 \right]^{1/2} \left[\sum_{n=0}^{\infty} \|T_1\|^{2n} \right]^{1/2} \|T_1 - T_0\|_p \\ &= \sqrt{2} \|f'\|_2 (1 - \|T_1\|^2)^{-1/2} \|T_1 - T_0\|_p \\ &\leq \sqrt{2} \|f\|_{\text{Lip}(\mathbb{T})} (1 - \|T_1\|^2)^{-1/2} \|T_1 - T_0\|_p. \end{aligned}$$

□

We shall need the following lemma from [12] for our use.

Lemma 3.5. [12, Lemma 2.1] *Let A and B be two positive contractions on \mathcal{H} . If the null space of B , $\mathcal{N}(B) = \{0\}$, then for every $f \in \mathcal{H}$*

$$(A - B)f = \int_0^{+\infty} e^{-tA} (A^2 - B^2) e^{-tB} f \, dt, \quad (3.9)$$

where the right hand side of (3.9) exists as an improper strong Riemann-Bochner integral.

Based on the lemma above, we obtain the following crucial result.

Lemma 3.6. *Let A and B be two positive contractions on \mathcal{H} . If $B \geq \delta I > 0$ for some positive δ and if for any $p \in [1, \infty]$, $(A^2 - B^2) \in \mathcal{S}_p(\mathcal{H})$, then $(A - B) \in \mathcal{S}_p(\mathcal{H})$, and*

$$\|A - B\|_p \leq \frac{1}{\delta} \|A^2 - B^2\|_p. \quad (3.10)$$

Proof. By Lemma 3.5, we have the following identity in $\mathcal{B}(\mathcal{H})$:

$$A - B = \int_0^{+\infty} e^{-tA} (A^2 - B^2) e^{-tB} dt, \quad (3.11)$$

since the hypothesis in Lemma 3.6 implies $\mathcal{N}(B) = \{0\}$. Moreover, the integral in (3.11) converges in $\mathcal{S}_p(\mathcal{H})$ -topology since $\|e^{-tA}\| \leq 1$, $\|e^{-tB}\| \leq e^{-\delta t}$. Hence

$$\|A - B\|_p \leq \int_0^{+\infty} \|A^2 - B^2\|_p e^{-t\delta} dt = \frac{1}{\delta} \|A^2 - B^2\|_p.$$

This completes the proof of the lemma. \square

The following corollary is an immediate consequence of Lemma 3.6.

Corollary 3.7. *Let T_0 and T_1 be two contractions on \mathcal{H} such that one of them is strict. Then $T_1 - T_0 \in \mathcal{S}_p(\mathcal{H})$ implies $D_{T_1} - D_{T_0}$ and $D_{T_1^*} - D_{T_0^*}$ both are in $\mathcal{S}_p(\mathcal{H})$.*

Schäffer [40] found an explicit power unitary dilation of a contraction T on a Hilbert space \mathcal{H} . Let $\ell_2(\mathcal{H}) = \mathcal{H} \oplus \mathcal{H} \oplus \dots$. The dilation space is $\ell_2(\mathcal{H}) \oplus \mathcal{H} \oplus \ell_2(\mathcal{H})$ and the unitary operator is

$$U_T = \begin{bmatrix} \ell_2(\mathcal{H}) & \mathcal{H} & \ell_2(\mathcal{H}) \\ S^* & 0 & 0 \\ D_{T^*} P_{\mathcal{H}} & T & 0 \\ -T^* P_{\mathcal{H}} & D_T & S \end{bmatrix} \begin{matrix} \ell_2(\mathcal{H}) \\ \mathcal{H} \\ \ell_2(\mathcal{H}) \end{matrix}, \quad (3.12)$$

where S is the unilateral shift on $\ell_2(\mathcal{H})$ of multiplicity $\dim(\mathcal{H})$ and $P_{\mathcal{H}}$ is the orthogonal projection from $\ell_2(\mathcal{H})$ onto $\mathcal{H} \oplus 0 \oplus 0 \oplus \dots$.

We have arrived at the main result of this section.

Theorem 3.8. *Let $1 \leq p < \infty$ and T_0, T_1 be two contractions on \mathcal{H} having the following properties:*

- (i) *one of the contractions T_0, T_1 is strict.*
- (ii) *$T_1 - T_0 \in \mathcal{S}_p(\mathcal{H})$.*

Let U_{T_0} and U_{T_1} on $\ell_2(\mathcal{H}) \oplus \mathcal{H} \oplus \ell_2(\mathcal{H})$ be Schäffer matrix dilations for T_0 and T_1 respectively. Then

$$\|U_{T_1} - U_{T_0}\|_p \leq 2^{\frac{1}{p}} \left[\frac{2 + (1 - \|T_0\|^2)^{1/2}}{(1 - \|T_0\|^2)^{1/2}} \right] \|T_1 - T_0\|_p.$$

Proof. Let T_0 and T_1 be two contractions such that $\|T_0\| < 1$ and $T_1 - T_0 \in \mathcal{S}_p(\mathcal{H})$. Let U_{T_0} and U_{T_1} be Schäffer matrix dilation for T_0 and T_1 respectively as above. The difference of U_{T_0} and U_{T_1} is given by

$$\begin{aligned} U_{T_1} - U_{T_0} &= \begin{bmatrix} 0 & 0 & 0 \\ (D_{T_1^*} - D_{T_0^*})P_{\mathcal{H}} & T_1 - T_0 & 0 \\ (T_0^* - T_1^*)P_{\mathcal{H}} & D_{T_1} - D_{T_0} & 0 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 & 0 \\ 0 & T_1 - T_0 & 0 \\ (T_0^* - T_1^*)P_{\mathcal{H}} & 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 & 0 \\ (D_{T_1^*} - D_{T_0^*})P_{\mathcal{H}} & 0 & 0 \\ 0 & D_{T_1} - D_{T_0} & 0 \end{bmatrix}. \end{aligned} \quad (3.13)$$

From the identity (3.13) and the Corollary 3.7, we arrive at

$$\|U_{T_1} - U_{T_0}\|_p \leq 2^{\frac{1}{p}} \|T_1 - T_0\|_p + \left[\|D_{T_1^*} - D_{T_0^*}\|_p^p + \|D_{T_1} - D_{T_0}\|_p^p \right]^{\frac{1}{p}}. \quad (3.14)$$

Since T_0 is a strict contraction so $T_0^*T_0$ is also a strict contraction and the spectrum of $T_0^*T_0$, $\sigma(T_0^*T_0)$ is contained in $[0, \|T_0^*T_0\|] = [0, \|T_0\|^2] \subseteq [0, 1)$. Let $\delta = (1 - \|T_0\|^2)^{1/2} > 0$. Then from the spectral theorem of $T_0^*T_0$, we have $D_{T_0} \geq \delta I > 0$. Applying Lemma 3.6 for the pair $(A = D_{T_1}, B = D_{T_0})$ we obtain

$$\|D_{T_1} - D_{T_0}\|_p \leq \frac{2}{(1 - \|T_0\|^2)^{1/2}} \|T_1 - T_0\|_p. \quad (3.15)$$

By a similar argument we also have

$$\|D_{T_1^*} - D_{T_0^*}\|_p \leq \frac{2}{(1 - \|T_0\|^2)^{1/2}} \|T_1 - T_0\|_p. \quad (3.16)$$

Therefore, using (3.15) and (3.16) in (3.14) we derive

$$\|U_{T_1} - U_{T_0}\|_p \leq 2^{\frac{1}{p}} \left[\frac{2 + (1 - \|T_0\|^2)^{1/2}}{(1 - \|T_0\|^2)^{1/2}} \right] \|T_1 - T_0\|_p.$$

This completes the proof. \square

The following theorem is now easy.

Theorem 3.9. *Let $f : \mathbb{T} \rightarrow \mathbb{C}$ be a Lipschitz function on \mathbb{T} and $1 < p < \infty$. Then for any pair of contractions (T_0, T_1) satisfying $\|T_0\| < 1$ and the hypothesis (ii), has the following estimate:*

$$\|f(T_1) - f(T_0)\|_p \leq 2^{\frac{1}{p}} \left[\frac{2 + (1 - \|T_0\|^2)^{1/2}}{(1 - \|T_0\|^2)^{1/2}} \right] d_p \|f\|_{\mathcal{L}_{\text{ip}}} \|T_1 - T_0\|_p, \quad (3.17)$$

where d_p is same as in Theorem 1.3, $\|T_0\| < 1$ and $\|f\|_{\mathcal{L}_{\text{ip}}}$ as in (1.4).

Proof. Let T_0, T_1 be two contractions and U_{T_0}, U_{T_1} be two Schäffer unitary dilations for T_0 and T_1 respectively. Since f is a Lipschitz function on \mathbb{T} , so by Theorem 1.3 and Theorem 3.8, we arrive at

$$\|f(T_1) - f(T_0)\|_p \leq \|f(U_{T_1}) - f(U_{T_0})\|_p \leq 2^{\frac{1}{p}} \left[\frac{2 + (1 - \|T_0\|^2)^{1/2}}{(1 - \|T_0\|^2)^{1/2}} \right] d_p \|f\|_{\mathcal{L}_{\text{ip}}} \|T_1 - T_0\|_p.$$

This completes the proof. \square

4. APPLICATION: THE MODIFIED KREIN TRACE FORMULA

In this section, we apply Theorem 3.9 to prove the existence of the spectral shift function and the associated modified Krein trace formula for a pair of contractions differing by a Hilbert-Schmidt operator.

The spectral shift function introduced by Lifshitz in [25] and generalized to a general situation by Krein guarantees the existence of a unique integrable function ξ on $\Omega = \mathbb{R}$ for arbitrary pairs of self-adjoint operators T_0 and T_1 with $T_1 - T_0$ in the trace class ideal such that the trace formula

$$\text{Tr}(f(T_1) - f(T_0)) = \int_{\Omega} f'(z) \xi(z) dz \quad (4.1)$$

holds if the Fourier transform of f' is in $L^1(\mathbb{R})$ (see [20]). The function ξ is called the *spectral shift function* and the formula (4.1) is known as the *Krein trace formula*.

A similar result was obtained in [21] for pairs of unitary operators (T_0, T_1) with $T_1 - T_0 \in \mathcal{S}_1(\mathcal{H})$ and $\Omega = \mathbb{T}$ such that (4.1) holds whenever f' has an absolutely convergent Fourier series. More than five decades later, a better sufficient condition on f was found by Aleksandrov and Peller in [4, Theorem 4.1], which states that (4.1) holds for any operator Lipschitz function f on \mathbb{T} . This brings to mind the paper by Mohapatra and Sinha [29, 30], where the authors gave an alternative proof of Krein trace formula (4.1) for pairs of self-adjoint operators as well as for pairs of unitary operators, which is purely functional analytic in nature.

Note that the existence of a complex-valued integrable spectral shift function ξ for a pair of contractions (T_0, T_1) with trace class difference $T_1 - T_0 \in \mathcal{S}_1(\mathcal{H})$ corresponding to all $f \in A(\mathbb{T})$, which are operator Lipschitz on \mathbb{T} , was proved in [26] (under an additional assumption $\rho(T_0) \cap \mathbb{D} \neq \emptyset$, where $\rho(T_0)$ is the resolvent set of T_0) and in [28] (in full generality). This settles the longstanding problem of obtaining a trace formula for arbitrary pairs of contractions with trace class differences as well as proving the existence of an integrable spectral shift function for such contractions. For more on the Krein trace formula, we refer to [23, 24, 36, 37, 38, 39].

In this present section, we treat the case of a pair of contractions (T_0, T_1) to establish a formula for $\text{Tr}[(f(T_1) - f(T_0))X]$ whenever $f \in C^1(\mathbb{T})$ and $T_1 - T_0, X \in \mathcal{S}_2(\mathcal{H})$. To this aim, the following theorem is crucial which establishes a formula for $\text{Tr}[(f(U) - f(V))Z]$ for two unitaries U and V such that $U - V \in \mathcal{S}_2(\mathcal{H})$ and $Z \in \mathcal{S}_2(\mathcal{H})$. We omit the proof because Skripka obtained a higher-order version of the following trace formula in [41, Theorem 4.5] and a similar reasoning will prove it.

Theorem 4.1. *Let U and V be two unitary operators acting on \mathcal{H} and let*

$$f \in \mathcal{F}_1(\mathbb{T}) := \left\{ f \in C^1(\mathbb{T}) : \sum_{n \in \mathbb{Z}} |n \hat{f}(n)| < \infty \right\}.$$

Let $U - V \in \mathcal{S}_2(\mathcal{H})$. Then for all $Z \in \mathcal{S}_2(\mathcal{H})$, there exists a function $\eta = \eta_{U, V, Z} \in L^1([0, 2\pi])$ satisfying

$$\text{Tr} \left[(f(U) - f(V))Z \right] = \int_0^{2\pi} f'(e^{it}) \eta(t) dt, \quad (4.2)$$

for every $f \in \mathcal{F}_1(\mathbb{T})$.

Now we are in a stage to state and prove our main result in this section.

Theorem 4.2. *Let T_0 and T_1 be two contractions and U_{T_0}, U_{T_1} be the unitary dilations of T_0 and T_1 , given by (3.12) respectively. If $U_{T_1} - U_{T_0} \in \mathcal{S}_2(\mathcal{H})$, then for each $f \in \text{Lip}(\mathbb{T})$,*

$$\left[(f(T_1) - f(T_0))X \right] \in \mathcal{S}_1(\mathcal{H}), \quad (4.3)$$

for every $X \in \mathcal{S}_2(\mathcal{H})$. Moreover, for each $X \in \mathcal{S}_2(\mathcal{H})$, there exists a function $\eta = \eta_{T_0, T_1, X} \in L^1([0, 2\pi])$ (unique upto an additive constant) satisfying

$$\text{Tr} \left[(f(T_1) - f(T_0))X \right] = \int_0^{2\pi} \frac{d}{dt} f(e^{it}) \eta(t) dt, \quad (4.4)$$

for every $f \in C^1(\mathbb{T})$.

Proof. Theorem 2.4 and Hölder's inequality implies

$$\begin{aligned} \left\| \left[(f(T_1) - f(T_0))X \right] \right\|_1 &\leq \|f(T_1) - f(T_0)\|_2 \|X\|_2 \\ &\leq \|f(U_{T_1}) - f(U_{T_0})\|_2 \|X\|_2 \\ &\leq \|f\|_{\text{Lip}(\mathbb{T})} \|U_{T_1} - U_{T_0}\|_2 \|X\|_2. \end{aligned} \quad (4.5)$$

Therefore (4.3) holds true. Now for $f \in C^1(\mathbb{T})$, from (4.5), we have

$$\left| \text{Tr} \left[(f(T_1) - f(T_0))X \right] \right| \leq \|f'\|_\infty \|U_{T_1} - U_{T_0}\|_2 \|X\|_2. \quad (4.6)$$

Hence there exists a complex Borel measure μ on \mathbb{T} such that

$$\text{Tr} \left[(f(T_1) - f(T_0))X \right] = \int_{\mathbb{T}} f'(z) d\mu(z), \quad (4.7)$$

for all $f \in C^1(\mathbb{T})$. Now by change of variable in the integration of (4.7) we arrive at

$$\int_{\mathbb{T}} f'(z) d\mu(z) = \int_0^{2\pi} \frac{d}{dt} f(e^{it}) d\nu(t), \quad (4.8)$$

where ν is the push forward measure for μ on $[0, 2\pi]$. On the other hand, for $f \in \mathcal{F}_1(\mathbb{T}) \subset C^1(\mathbb{T})$, Theorem 4.1 and the Schäffer matrix representations of U_{T_1} and U_{T_0} and the cyclicity property of trace together give

$$\begin{aligned} \text{Tr} \left[(f(T_1) - f(T_0))X \right] &= \text{Tr} \left[\hat{P}_{\mathcal{H}} (f(U_{T_1}) - f(U_{T_0})) \hat{P}_{\mathcal{H}} \left(\hat{P}_{\mathcal{H}} X \hat{P}_{\mathcal{H}} \right) \right] \\ &= \text{Tr} \left[(f(U_{T_1}) - f(U_{T_0})) \left(\hat{P}_{\mathcal{H}} X \hat{P}_{\mathcal{H}} \right) \right] \\ &= \int_0^{2\pi} \frac{d}{dt} f(e^{it}) \eta(t) dt, \end{aligned} \quad (4.9)$$

where $\hat{P}_{\mathcal{H}}$ is the orthogonal projection from $\ell_2(\mathcal{H}) \oplus \mathcal{H} \oplus \ell_2(\mathcal{H})$ onto $0 \oplus \mathcal{H} \oplus 0$, and $\eta = \eta_{U_{T_0}, U_{T_1}, \hat{P}_{\mathcal{H}} X \hat{P}_{\mathcal{H}}} \in L^1[0, 2\pi]$. Therefore, from (4.7), (4.8) and (4.9) we arrive at

$$\int_0^{2\pi} \frac{d}{dt} f(e^{it}) \eta(t) dt = \int_0^{2\pi} \frac{d}{dt} f(e^{it}) d\nu(t), \quad \text{for all } f \in \mathcal{F}_1(\mathbb{T}),$$

which implies,

$$\int_0^{2\pi} e^{int} \eta(t) dt = \int_0^{2\pi} e^{int} d\nu(t), \quad \text{for all } n \in \mathbb{Z} \setminus \{0\}, \quad (4.10)$$

which further implies that $d\nu(t) = (\eta(t) + c)dt$, for some constant c . This establishes (4.4). Let η_1 and η_2 be two $L^1[0, 2\pi]$ functions satisfying (4.4), then

$$\int_0^{2\pi} e^{int} \eta_1(t) dt = \int_0^{2\pi} e^{int} \eta_2(t) dt, \quad \text{for all } n \in \mathbb{Z} \setminus \{0\}. \quad (4.11)$$

This shows that η_1 and η_2 are unique upto an additive constant. This completes the proof of the theorem. \square

Corollary 4.3. *Thanks to Theorems 3.8 and 3.9, the above Theorem 4.2 holds for all pairs of contractions, with one of them being strict, when their difference is a Hilbert-Schmidt operator.*

5. AN OPEN PROBLEM

In this section, we state an open problem for functions of commuting contractions. To be more precise, if (T_1, R_1) and (T_2, R_2) are two pairs of doubly commuting contractions such that $T_1 - T_2, R_1 - R_2 \in \mathcal{S}_p(\mathcal{H})$, $1 < p < \infty$ and $f \in \text{Lip}(\mathbb{T}^2)$ then is it possible to give a Schatten von Neumann estimate for the difference $(f(T_1, R_1) - f(T_2, R_2))$?

This question has been partially considered by V. Peller in [34]. Indeed, he proved the following:

Theorem 5.1. *Let $1 \leq p < \infty$ and let (T_1, R_1) and (T_2, R_2) be pairs of commuting contractions such that $T_1 - T_2, R_1 - R_2 \in \mathcal{S}_p(\mathcal{H})$. Suppose that f is a function of class $B_{\infty,1}^1(\mathbb{T}^2)_+$. Then $(f(T_1, R_1) - f(T_2, R_2)) \in \mathcal{S}_p(\mathcal{H})$ and*

$$\|f(T_1, R_1) - f(T_2, R_2)\|_p \leq \text{const} \|f\|_{B_{\infty,1}^1(\mathbb{T}^2)} \max\{\|T_1 - T_2\|_p, \|R_1 - R_2\|_p\}, \quad (5.1)$$

where $B_{\infty,1}^1(\mathbb{T}^2)_+$ is the analytic subspace of the Besov space $B_{\infty,1}^1(\mathbb{T}^2)$.

For a detailed exposition on Besov spaces and its properties the reader is directed to [34, Section 2]. Note that we intend not only to remove the restrictive analyticity condition on the function f but also to extend the function space to the whole of $\text{Lip}(\mathbb{T}^2)$. To accomplish this, we set up the following concepts.

Let (T_1, R_1) and (T_2, R_2) be two pairs of doubly commuting contractions acting on \mathcal{H} , i.e., $T_i R_i = R_i T_i$ and $T_i^* R_i = R_i^* T_i^*$ for $i = 1, 2$. Let (U_1, V_1) and (U_2, V_2) be the commuting unitary dilations of the pairs (T_1, R_1) and (T_2, R_2) respectively on the Hilbert spaces $\mathcal{K}_1 \supset \mathcal{H}$ and $\mathcal{K}_2 \supset \mathcal{H}$. If $P_{\mathcal{H}}, Q_{\mathcal{H}}$ are the projections from \mathcal{K}_1 onto \mathcal{H} and \mathcal{K}_2 onto \mathcal{H} respectively, then for any trigonometric polynomial $p(z_1, z_2)$, we have

$$p(T_1, R_1) = P_{\mathcal{H}} p(U_1, V_1)|_{\mathcal{H}} \quad \text{and} \quad p(T_2, R_2) = Q_{\mathcal{H}} p(U_2, V_2)|_{\mathcal{H}} \quad (5.2)$$

and by a verbatim repetition of the argument employed in Lemma 3.2, one can extend (5.2) to every $f \in C(\mathbb{T}^2)$. We close the paper with the following open problem.

Problem 5.2. *Let $1 < p < \infty$ and let (T_1, R_1) and (T_2, R_2) be two pairs of doubly commuting contractions on \mathcal{H} satisfying the following conditions:*

- (i) T_2 and R_2 are strict contractions.
- (ii) $T_1 - T_2 \in \mathcal{S}_p(\mathcal{H}), R_1 - R_2 \in \mathcal{S}_p(\mathcal{H})$.

Suppose that f is a function of class $\text{Lip}(\mathbb{T}^2)$. Then is it true that

$$(f(T_1, R_1) - f(T_2, R_2)) \in \mathcal{S}_p(\mathcal{H})?$$

If yes, does there exist a constant $d_{p,T_2,R_2} > 0$ such that

$$\|f(T_1, R_1) - f(T_2, R_2)\|_p \leq d_{p,T_2,R_2} \|f\|_{\text{Lip}(\mathbb{T}^2)} \max\{\|T_1 - T_2\|_p, \|R_1 - R_2\|_p\} \quad (5.3)$$

holds, where

$$\|f\|_{\text{Lip}(\mathbb{T}^2)} = \inf \left\{ K > 0 : |f(z_1, w_1) - f(z_2, w_2)| \leq K \max\{|z_1 - z_2|, |w_1 - w_2|\} \right\} \quad (5.4)$$

Acknowledgment: T. Bhattacharyya is supported by a J C Bose Fellowship JCB/2021/000041 of SERB. A. Chattopadhyay is supported by the Core Research Grant (CRG), File No: CRG/2023/004826, of SERB. S. Giri acknowledges the support by the Prime Minister's Research Fellowship (PMRF), Government of India. C. Pradhan acknowledges support from JCB/2021/000041 as well as IoE post-doctoral fellowship. This research is supported by the DST FIST program-2021 [TPN-700661].

Data Availability Statement: Data sharing is not applicable to this article as no data sets were generated or analyzed during the current study.

Conflicts of interests: The authors declare that they have no conflict of interest.

REFERENCES

- [1] A. B. Aleksandrov and V. V. Peller, *Functions of perturbed dissipative operators*, Algebra i Analiz., **23** (2011), no. 2, 9–51.
- [2] A. B. Aleksandrov, V. V. Peller, D. S. Potapov and F. A. Sukochev, *Functions of normal operators under perturbations*, Adv. Math., **226** (2011), no. 6, 5216–5251.
- [3] A. B. Aleksandrov and V. V. Peller, *Operator Lipschitz functions*, Uspekhi Mat. Nauk, **71** (2016), no. 4(430), 3–106 (in Russian). Translation: Russ. Math. Surv., **71** (2016), no.4, 605–702.
- [4] A. B. Aleksandrov and V. V. Peller, *Krein’s trace formula for unitary operators and operator Lipschitz functions*, Funktsional. Anal. i Prilozhen., **50** (2016), no. 3, 1–11.
- [5] A. B. Aleksandrov and V. V. Peller, *Dissipative operators and operator Lipschitz functions*, Proc. Amer. Math. Soc., **147** (2019), no. 5, 2081–2093.
- [6] P. J. Ayre, M. G. Cowling and F. A. Sukochev, *Operator Lipschitz estimates in the unitary setting*, Proc. Amer. Math. Soc., **144** (2016), no. 3, 1053–1057.
- [7] M. S. Birman and M. Z. Solomyak, *Double Stieltjes operator integrals*, Problems of Mathematical Physics, Izdat (Leningrad University, Leningrad, 1966, in Russian), pp. 33–67. English translation in: Topics in Mathematical Physics, vol. **1** (Consultants Bureau Plenum Publishing Corporation, New York, 1968), pp. 25–54.
- [8] M. S. Birman and M. Z. Solomyak, *Double Stieltjes operator integrals II*, Problems of Mathematical Physics, Izdat, no. 2 (Leningrad University, Leningrad, 1967, in Russian), pp. 26–60. English translation in: Topics in Mathematical Physics, vol. **2** (Consultants Bureau, New York, 1968), pp. 19–46.
- [9] M. S. Birman and M. Z. Solomyak, *Double Stieltjes operator integrals III*, Problems of Mathematical Physics, Izdat vol. **6** (Leningrad University, Leningrad, 1973, in Russian), pp. 27–53.
- [10] M. Caspers, S. Montgomery-Smith, D. Potapov and F. Sukochev, *The best constants for operator Lipschitz functions on Schatten classes*, J. Funct. Anal., **267** (2014), 3557–3579.
- [11] A. Chattopadhyay and K. B. Sinha, *Trace formula for Two Variables*, [arXiv:1402.0792](https://arxiv.org/abs/1402.0792) (2014), Preprint.
- [12] A. Chattopadhyay and K. B. Sinha, *Trace formula for contractions and its representation in \mathbb{D}* , J. Operator Theory, **88** (2022), no. 2, 275–288.
- [13] E. B. Davies, *Lipschitz continuity of functions of operators in the Schatten classes*, J. London Math. Soc. (2), **37** (1988), no. 1, 148–157.
- [14] Yu. B. Farforovskaya, *Estimates of the closeness of spectral decompositions of self-adjoint operators in the Kantorovich-Rubinshtein metric*, Vestnik Leningrad. Univ., **22** (1967), no. 19, 155–156 (Russian).
- [15] Yu. B. Farforovskaya, *The connection of the Kantorovich-Rubinshtein metric for spectral resolutions of selfadjoint operators with functions of operators*, Vestnik Leningrad. Univ., **23** (1968), no. 19, 94–97 (Russian).
- [16] Yu. B. Farforovskaya, *On the estimation of the difference $f(B) - f(A)$ in the classes \mathcal{S}_p* , Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI), **74** (1974), 194–195 (Russian).
- [17] T. Kato, *Continuity of the map $S \mapsto |S|$ for linear operators*, Proc. Japan Acad., **49** (1973), 157–160.
- [18] E. Kissin and V. S. Shulman, *On fully operator Lipschitz functions*, J. Funct. Anal., **253** (2007), no. 2, 711–728.
- [19] E. Kissin, D. Potapov, V. Shulman and F. Sukochev, *Operator smoothness in Schatten norms for functions of several variables: Lipschitz conditions, differentiability and unbounded derivations*, Proc. Lond. Math. Soc. (3), **105** (2012), no. 4, 661–702.
- [20] M. G. Krein, *On a trace formula in perturbation theory*, Mat. Sb., **33** (1953), no. 3, 597–626.
- [21] M. G. Krein, *On perturbation determinants and a trace formula for unitary and self-adjoint operators*, Dokl. Akad. Nauk SSSR, **144** (1962), 268–271 (in Russian).
- [22] M. G. Krein, *Some new studies in the theory of perturbations of self-adjoint operators* (Russian), First Math. Summer School, Part I (Russian), Izdat. Naukova Dumka, Kiev., (1964), 103–187.

- [23] M. G. Krein, *Perturbation determinants and a trace formula for some classes of pairs of operators*, J. Operator Theory, **17** (1987), no. 1, 129–187.
- [24] H. Langer, *Eine Erweiterung der Spurformel der Störungstheorie*, Math. Nachr., **30** (1965), 123–135.
- [25] I. M. Lifshitz, *On a problem of the theory of perturbations connected with quantum statistics* Russian, Uspehi Matem. Nauk (N.S.), **7** (1952), no. 1(47), 171–180.
- [26] M. M. Malamud and H. Neidhardt, *Trace formulas for additive and non-additive perturbations*, Adv. Math., **274** (2015), 736–832.
- [27] M. M. Malamud, H. Neidhardt and V. V. Peller, *A trace formula for functions of contractions and analytic operator Lipschitz functions*, C. R. Math. Acad. Sci. Paris, **355** (2017), no. 7, 806–811.
- [28] M. M. Malamud, H. Neidhardt and V. V. Peller, *Absolutely continuity of spectral shift*, J. Funct. Anal., **276** (2019), no. 5, 1575–1621.
- [29] A. Mohapatra and K. B. Sinha, *Spectral shift function and trace formula. Spectral and inverse spectral theory*, Proc. Indian Acad. Sci. Math. Sci., **104** (1994), no. 4, 819–853.
- [30] A. Mohapatra and K. B. Sinha, *Spectral shift function and trace formula for unitaries—new proof*, Integral Equations Operator Theory, **24** (1996), no. 3, 285–297.
- [31] F. L. Nazarov and V. V. Peller, *Functions of perturbed n -tuples of commuting self-adjoint operators*, J. Funct. Anal., **266** (2014), no. 8, 5398–5428.
- [32] V. V. Peller, *Hankel operators in the perturbation theory of unitary and self-adjoint operators*, Funkt. Anal. i Priloz., **19** (1985), no. 2, 37–51; English transl.: Functional Anal. Appl., **19** (1985), no. 2, 111–123.
- [33] V. V. Peller, *For which f does $A - B \in S_p$ imply that $f(A) - f(B) \in S_p$?*, in Operators in indefinite metric spaces, scattering theory and other topics (Bucharest, 1985), Oper. Theory Adv. Appl., **24** (1987), 289–294. Birkhäuser, Basel.
- [34] V. V. Peller, *Functions of commuting contractions under perturbation*, Math. Nachr., **292** (2019), no. 5, 1151–1160.
- [35] D. Potapov and F. Sukochev, *Operator-Lipschitz functions in Schatten-von Neumann classes*, Acta Math., **207** (2011), no. 2, 375–389.
- [36] A. V. Rybkin, *A trace formula for a contractive and a unitary operator*, Funkt. Anal. i Priloz., **21** (1987), no. 4, 85–87.
- [37] A. V. Rybkin, *The discrete and the singular spectrum in the trace formula for a contractive and a unitary operator*, Funkt. Anal. i Priloz., **23** (1989), no. 3, 84–85.
- [38] A. V. Rybkin, *The spectral shift function, the characteristic function of a contraction and a generalized integral*, Mat. Sb., **185** (1994), no. 10, 91–144.
- [39] A. V. Rybkin, *On A -integrability of the spectral shift function of unitary operators arising in the Lax-Phillips scattering theory*, Duke Math. J., **83** (1996), no. 3, 683–699.
- [40] J. J. Schäffer, *On unitary dilations of contractions*, Proc. Amer. Math. Soc., **6** (1955), 322.
- [41] A. Skripka, *Estimates and trace formulas for unitary and resolvent comparable perturbations*, Adv. Math., **311** (2017), 481–509.
- [42] D. Voiculescu, *Some results on norm-ideal perturbations of Hilbert space operators*, J. Operator Theory, **2** (1979), no. 1, 3–37.

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE, BANGALORE 560012, INDIA
Email address: tirtha@iisc.ac.in

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY GUWAHATI, GUWAHATI, 781039, INDIA
Email address: arupchatt@iitg.ac.in, 2003arupchattopadhyay@gmail.com

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY GUWAHATI, GUWAHATI, 781039, INDIA
Email address: saikat.giri@iitg.ac.in, saikatgiri90@gmail.com

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE, BANGALORE 560012, INDIA
Email address: chandanp@iisc.ac.in; chandan.pradhan2108@gmail.com