

Anatomical basis of human sex differences in ECG identified by automated torso-cardiac three-dimensional reconstruction

Hannah J. Smith¹, Blanca Rodriguez¹, Yuling Sang², Marcel Beetz², Robin P. Choudhury³, Vicente Grau², and Abhirup Banerjee^{2,3}

¹ Computational Cardiovascular Science Group, Department of Computer Science, University of Oxford, Oxford OX1 3QD, United Kingdom

² Institute of Biomedical Engineering, Department of Engineering Science, University of Oxford, Oxford OX3 7DQ, United Kingdom

³ Division of Cardiovascular Medicine, Radcliffe Department of Medicine, University of Oxford, Oxford OX3 9DU, United Kingdom

***Address for correspondence.**

Abhirup Banerjee, Institute of Biomedical Engineering, Department of Engineering Science, University of Oxford, Oxford OX3 7DQ, United Kingdom, abhirup.banerjee@eng.ox.ac.uk

Author contributions. HJS, BR, VG, and AB conceptualised the study and designed the research. BR, VG, RPC, and AB supervised the research. HJS and AB acquired and curated the dataset. AB and HJS developed the reconstruction pipeline. HJS, AB, YS, and MB pre-processed and cleaned the data. HJS and AB conducted the study and analysed the data and the results. HJS, AB, BR, and VG prepared the original draft and reviewed and edited the manuscript. All authors discussed the results, provided comments regarding the manuscript, and agreed on the final draft.

Abstract

Background and Aims

The electrocardiogram (ECG) is routinely used for diagnosis and risk stratification following myocardial infarction (MI), though its interpretation is confounded by anatomical variability and sex differences. Women have a higher incidence of missed MI diagnosis and poorer outcomes following infarction. Sex differences in ECG biomarkers and torso-ventricular anatomy have not been well characterised, largely due to the absence of high-throughput torso reconstruction methods.

Methods

This work presents quantification of sex differences in ECG versus anatomical biomarkers in healthy and post-MI subjects, enabled by a novel, end-to-end automated pipeline for torso-ventricular anatomical reconstruction from clinically standard cardiac magnetic resonance imaging. Personalised 3D torso-ventricular reconstructions were generated for 425 post-MI subjects and 1051 healthy controls from the UK Biobank. Regression models were created relating the extracted torso-ventricular and ECG parameters.

Results

Half the sex difference in QRS durations is explained by smaller ventricles in women both in healthy (3.4 ± 1.3 ms of 6.0 ± 1.5 ms) and post-MI (4.5 ± 1.4 ms of 8.3 ± 2.5 ms) subjects. Lower baseline ST_T amplitude in women is also associated with smaller ventricles, and more superior and posterior cardiac position. Post-MI T wave amplitude and R axis deviations are more strongly associated with a more posterior and horizontal cardiac position in women rather than electrophysiology as in men.

Conclusion

A novel computational pipeline enables the three-dimensional reconstruction of 1476 torso-cardiac geometries of healthy and post-myocardial infarction subjects, quantification of sex and BMI-related differences and association with ECG biomarkers. Any ECG-based tool should be reviewed considering anatomical sex differences to avoid sex-biased outcomes.

What's new

- A novel end-to-end automated pipeline for torso-cardiac anatomical reconstruction from standard clinical imaging developed and validated, widely applicable in clinical contexts for the delivery of personalised medicine.
- Sex-differences in torso-ventricular anatomical features quantified and their impact on electrocardiographic biomarkers assessed in 1476 healthy and post-myocardial infarction subjects.
- Lower female ST-segment amplitude is associated with differences in cardiac size and position, and body mass index plays an important modulating effect, which must be corrected for in diagnostic tools for myocardial infarction.
- Significant sex-differences in the contribution of anatomical versus electrophysiological changes in the electrocardiogram following myocardial infarction are identified, which are relevant to patient risk stratification.
- Torso-cardiac anatomical reconstructions open an avenue for personalised modelling and simulation within the Digital Twin vision.

Keywords. Torso reconstruction, magnetic resonance imaging, myocardial infarction, sex differences, risk stratification, machine learning

Introduction

The electrocardiogram (ECG) is a vital tool for routine clinical assessment of cardiac electrical abnormalities. However, the relative position and orientation of the heart with respect to the electrode locations can affect ECG complex morphology. Computational and experimental studies have demonstrated that torso-ventricular orientation can substantially affect the ECG, potentially confounding disease diagnosis, and that ECG variability due to anatomy exceeds that of electrophysiology in healthy subjects [1-5]. The intersectional relationship between the anatomy and demographic features indicates that the accuracy of ECG interpretation will be skewed by characteristics such as sex and body mass index (BMI) if these factors are not considered [6-8].

Following MI, risk stratification tools are used to identify patients susceptible to sudden cardiac death (SCD), primarily from ventricular arrhythmias [9]. Current risk stratification guidelines that primarily focus on the left ventricular ejection fraction (LVEF) fail to identify the majority of SCD cases, prompting calls for supplemental ECG biomarkers to be used [10]. These include measures of slowed conduction, such as prolonged QRS duration [11, 12] and of increased repolarisation heterogeneity, which are affected by the duration, amplitude, and axis angles of the QRS and T waves [13, 14]. Acutely, ischemic ECG changes are a fundamental consideration in diagnosis and categorisation of MI events [15]. Changes to the ST segment are particularly important, with ST-elevation MI (STEMI) mandating immediate reperfusion therapy [16, 17]. However, as variation from anatomical factors has been shown to exceed thresholds for diagnosing pathology [2, 3], without correction for anatomy, their use is fundamentally problematic.

There is a higher incidence of missed diagnosis of acute MI in women, and hospital care can be less aggressive [18-22]. This is a potential contributor to the higher female mortality following MI [23, 24], alongside factors such as differing demographic risk characteristics and comorbidities [22, 25, 26].

Women's ECGs typically exhibit shorter QRS durations and lower ST junction (STj) and T wave amplitudes, but longer QT intervals, than male [27]. The origin of these differences is multifactorial, with both electrophysiological and structural factors contributing. Testosterone increases repolarising potassium currents and decreases L-type calcium currents, both acting to speed up repolarisation [6]. Additionally, left ventricular mass is smaller in women than men [28], a factor known to affect QRS duration and STj and T wave amplitudes [29, 30]. Decreases in torso volume are known to increase QRS and T wave amplitudes [1, 2], making the smaller female torso a potential contributor to ECG sex differences, but the impact on STj amplitude is not known. Sex differences in cardiac position have been noted, particularly in the superior-inferior axis [31]. In women, electrophysiological remodelling post-infarction may be less substantial than in men, partially from the protective effects of estrogen [32], suggesting that the effects of anatomical differences may be proportionally more significant than in men. Therefore, it is important to investigate the effect of demographic differences on parameters considered for integration into risk stratification tools following MI. However, differences in the post-myocardial infarction (MI) population and the quantification of the effect of anatomical differences on key ECG biomarkers from clinical ECGs is largely unknown. The investigation of population-wide trends requires automated torso reconstruction from clinically standard cardiac magnetic resonance (CMR) imaging to facilitate high-throughput analysis of the ECG-anatomy relationship.

Accordingly, our current work aims to (1) develop and validate an automated image processing pipeline to reconstruct 3D torso models using clinically standard CMR images, (2) quantify differences in torso-ventricular anatomy and ECG biomarkers, and (3) identify the key anatomical

determinants of ECG biomarkers for healthy and chronic-stage post-MI subjects of both sexes. We hypothesise that whereas the smaller female heart and torso could explain baseline sex differences in some ECG parameters, sex differences in healthy versus post-MI subjects will also be affected by cardiac orientation in the torso as well as non-anatomical electrophysiological factors. Anatomical effects may be proportionally more significant in women than men due to their predicted smaller electrophysiological changes. In order to explore this, we have developed a novel automated image processing pipeline to quantify anatomical biomarkers from personalised 3D torso and heart models reconstructed from clinical CMR images of 1051 healthy and 425 post-MI subjects, and to investigate sex differences in the relationship between torso-ventricular anatomy and key ECG biomarkers. These insights will lay the groundwork for a process to correct for the effect of torso-ventricular anatomy on the ECG, enabling personalised, automated ECG interpretation.

Methods

Computational pipeline

Figure 1A illustrates the fully automated pipeline used to create personalised 3D torso reconstructions from clinically standard CMR imaging. This integrates and reformulates state-of-the-art image processing techniques in order to give the first fully automated, and extensively validated, torso-ventricular reconstruction pipeline of its kind. In short, the pipeline combines machine learning-based contour extraction, statistical shape modelling and automated mesh deformation in the following steps:

Three-stage torso contour extraction from CMR imaging using machine learning to form sparse torso representation: Approximately 60 scout images per subject were first segmented using a convolutional neural network; then their outline was extracted with automated post-processing, and finally refined using a second network. U-Net [33] was used for both network architectures, as described in Smith et al. [34].

Fitting sparse representation contours to a statistical shape model of the human body to give an initial fitted torso: To generate torso meshes from sparse torso contours, a statistical shape model (SSM) of human body shapes made from a large optical scan dataset was fitted [35, 36]. As described in Supplementary Appendix 1.2.1, the principal components of the SSM were iteratively adjusted, alongside rigid transformations of the SSM in order to minimise distance between the surface and contours.

Automated deformation of the initial torso towards the extracted contours using thin plate splines to give a final dense torso mesh: As the fitted surface is inherently limited by variability in the SSM, a final deformation step was applied in order to better capture inter-subject variation. A thin plate splines method was used to pull the surface towards the contours, whilst retaining smoothness and regularity [37], as described in Supplementary Appendix 1.2.2. Laplacian smoothing, decimation, and affine transformation were applied to ensure the local geometric and topological characteristics of the reconstructed mesh.

Integration with a sparse cardiac mesh and subsequent extraction of torso-ventricular anatomical parameters: Cardiac surfaces were segmented using the method in Banerjee et al. [38] and torso-ventricular anatomical parameters were extracted, as detailed in Supplementary Appendix 1.3. The torso volume and two metrics of cardiac size were estimated. The locations of the ECG electrodes were identified on the mean SSM torso, and their positions were transformed with the mesh such

that the resulting torso had electrodes located in equivalent locations. The position of the heart centre relative to the electrodes was estimated in the x (lateral), y (posterior) and z (superior) directions. The orientation of the heart was quantified by the Euler angles of the cardiac axes with respect to the torso axes. This includes the spin of the cardiac short axis plane around the torso vertical axis, the verticality of the cardiac long axis and the tilt of the cardiac short axis plane.

Validation of the torso reconstruction pipeline

The CMR images of 30 subjects, chosen to be representative of the underlying dataset, were manually annotated to form an independent test set for validation. Final results of the three-stage torso contour extraction process were compared with simply using a single U-Net architecture to process raw images directly to refined contours in order to assess the accuracy of the torso-air boundary placement. Separate reconstructions of the 30 test subjects were made using the fully automated pipeline and using manually annotated contours. For each subject, the surface-to-surface distance between the two reconstructions was calculated to evaluate error propagation from contouring to reconstruction, as in Supplementary Appendix 1.4. The mean surface-to-contour distance was calculated for each test subject to evaluate the quality of the reconstruction.

Dataset

The UK Biobank (UKB) dataset [39] was chosen for its high quality CMR imaging and 12-lead ECGs. 1646 control subjects were randomly selected from the UKB cohort alongside 479 subjects with a history of MI, as defined in Supplementary Appendix 1.1. The healthy cohort was made by excluding control subjects with disease diagnoses, and subjects in either population were excluded if their ECG or imaging data was not complete, as described in Supplementary Appendix 1.1. The final population for reconstruction and statistical analysis comprised 1051 healthy and 425 post-MI subjects (total 1476), as shown in Tables 1 and 2. 92.7% of subjects identified as White British. The post-MI population was 6.1 ± 0.4 years older and had 2.4 ± 0.2 kg/m² higher BMI, with no statistically significant differences in ethnicity.

Statistical analysis

ECG parameters were obtained from the UK Biobank database, which automatically extracted them with the proprietary software CardioSoft [40]. The distribution of each anatomical and ECG parameter were compared between populations as in Supplementary Appendix 1.5. STj amplitude refers to the amplitude of the ST segment measured at the junction point (the end of the QRS complex). The contribution of each anatomical parameter to the sex difference in ECG biomarkers, and the remaining estimated effect of electrophysiology, was calculated by regression analysis, as in Supplementary Appendix 1.5. Similarly, their contribution to the difference between healthy and post-MI ECG biomarkers was estimated.

Results

Reconstructed torso surfaces matched input contours to within 2.19mm:

Figure 1B showcases the capability of the computational image processing pipeline to extract torso contours and reconstruct them into 3D meshes. Figure 1B(i) demonstrates that the three-stage segmentation, automated post-processing, and refinement procedure improved the Dice coefficient of torso contour extraction, compared with a single-stage contouring network, across all image views. The median surface-to-contour distance, showing the quality of the reconstructed torso surface, as depicted in Figure 1B(ii), for the test subjects was 2.19mm (inter-quartile range - IQR: 1.98-2.37mm). The median surface-to-surface distance between torso meshes reconstructed using

the manually annotated contours and the machine learning pipeline for the 30 test subjects was 0.82mm (IQR: 0.69-1.10mm). Figure 1B(iii) shows the reconstructed surfaces for the test subjects with the smallest and largest median reconstruction errors. Both reconstructed surfaces are smooth and anatomically realistic, and, across most of the torso, errors are within the millimetre range.

Approximately 50% of QRS shortening in women versus men is explained by small cavity volumes:

Figures 2A and 2B show that for both healthy and post-MI subjects, QRS duration was shorter in women versus men in all leads. Figures 2C and 2D illustrate the contribution of each anatomical factor, and the remaining effect of electrophysiology, to the difference between female and male QRS duration. This is calculated from the product of the regression coefficients and anatomical sex differences, as further described in Supplementary Appendix 1.5. In healthy subjects, this shows that sex differences in cavity volume (enclosed by the endocardial surfaces of both ventricles) contributed a mean value across all leads of 3.4 ± 1.3 ms, approximately half of the 6.0 ± 1.5 ms mean total sex difference in QRS duration. Similarly for post-MI subjects, cavity volume contributed a mean of 4.5 ± 1.4 ms of a total sex difference of 8.3 ± 2.5 ms. The effect of electrophysiology consistently contributed between 1.0 ± 1.0 ms and 2.3 ± 1.2 ms to the sex difference in the precordial leads QRS for healthy subjects; but in post-MI subjects, the electrophysiological contribution was considerably more variable, increasing markedly from 0.2 ± 2.8 ms in V1 to 6.3 ± 3.0 ms in V6.

There were several anatomical differences for post-MI patients compared with healthy subjects. Torso volume was larger than in healthy subjects in women and men (by 3.7 ± 0.9 dm³ (10.2 \pm 3.3%) and 3.5 ± 0.5 dm³ (8.2 \pm 1.9%) respectively, Supplementary Figure 3). Additionally, for women, in post-MI compared with healthy subjects the heart was positioned more posteriorly by 3.1 ± 0.9 mm. The heart was also oriented with its long axis more horizontally by $6.0 \pm 1.2^\circ$ and $4.3 \pm 0.5^\circ$ in post-MI versus healthy women and men respectively. These changes were related to the increased BMI of post-MI subjects, as shown in Supplementary Figure 3K. Age was also considered, as shown in Supplementary Figure 3J, but its effects are considerably less substantial.

Figures 2E and 2F demonstrate that the QRS prolongation in MI compared with healthy population was again less substantial in women than men with mean percentage reduction in QRS duration, across all leads, of $3.4 \pm 2.3\%$ and $5.8 \pm 1.5\%$ respectively. The contribution of each anatomical factor to the difference between healthy and post-MI subjects' QRS durations was calculated similarly by the product of the corresponding regression coefficients with the anatomical difference between healthy and post-MI subjects. Further details are given in Supplementary Appendix 1.5. This is illustrated in Figures 2G and 2H. Comparing Figures 2G and 2H demonstrates that the reduced amount of QRS duration prolongation following MI in women was associated with smaller electrophysiological contributions than in men. There was also an increased amount of compensating shortening in QRS duration (shown as negative contributions) from anatomical factors, primarily torso volume, in women. For both women and men, the more horizontal cardiac long axis in post-MI compared with healthy subjects was associated with a substantial proportion of their QRS prolongation for all precordial, and most limb, leads.

ST_j amplitude is lower in women than men in all precordial leads for healthy and post-MI subjects, explained by a smaller left ventricular (LV) mass and more superior and posterior position:

Figure 3A and S6A illustrate sex differences in ST_j amplitude, measured at the end of the QRS complex, in healthy and post-MI subjects. They show that for both populations, ST_j amplitude is significantly lower in the precordial leads in women versus men, particularly in the septal and

anterior leads. Figure 3B and Supplementary Figure 6B show the estimated anatomical contributions to this sex difference. They demonstrate that this can be partly explained by women's decreased LV mass (by $29.3 \pm 5.4\%$ and $27.1 \pm 5.7\%$ in healthy and post-MI subjects respectively), $9.6 \pm 0.9\text{mm}$ more superior position for healthy subjects, and $4.3 \pm 1.0\text{mm}$ more posterior position for post-MI subjects. While the smaller female LV mass generally added to the electrophysiological sex difference, their smaller torso volume had a negative contribution, as a smaller torso volume was generally associated with a higher STj amplitude. Baseline STj amplitude in lead V4 was $0.028 \pm 0.003\text{mV}$ lower in healthy women than healthy men. As BMI positively correlates with torso volume, this meant that for healthy women with $\text{BMI} > 25\text{kgm}^{-2}$ STj amplitude was $0.047 \pm 0.004\text{mV}$ lower than for healthy men with $\text{BMI} < 25\text{kgm}^{-2}$.

T wave amplitude (TWA) following MI is more strongly associated with a more posterior cardiac position and more horizontal orientation in women than men:

Figures 3C and 3D show the change in TWA between healthy and post-MI subjects, split by sex. They illustrate that in women, TWA was only substantially reduced following MI in precordial leads V4-6, alongside most limb leads, whereas in men this occurred in all leads. The anatomical contributions of this reduction are illustrated in Figures 3E and 3F. They demonstrate that, for both women and men, the reduced cardiac long axis verticality and larger torso volume of post-MI subjects compared with healthy controls contribute to their reduced TWA, but in women the more posterior position also contributes. The more posterior post-MI cardiac position and more horizontal orientation play an important role in TWA in women compared with men. The estimated electrophysiological component was consistently associated with reduced TWA in the precordial leads for men, primarily in the anterior leads (V3-V4). However, in women this was actually associated with increased TWA in the septal leads (V1-V2), with significant reductions only in the lateral leads (V5-V6). These increases are however masked by the negative contribution of anatomical factors, such as posterior position and torso volume, leading to little difference in the TWA for the septal and anterior leads in women.

R axis deviations following MI are associated with more horizontal cardiac orientation in women, but disease-related electrophysiological abnormalities in men:

Whereas the previous ECG parameters can be assessed in individual leads, ECG-derived axis angles are a composite metric, positively correlated with amplitudes in leads such as lead aVF, and negatively correlated with lead I [41]. Figure 4A breaks down the anatomical contributions to R and T axis shifts in post-MI compared with healthy subjects. It illustrates that for men the electrophysiological contribution dominates their $12.0 \pm 2.8^\circ$ left deviated R axis following MI. However, Figure 4B reveals that for women their $5.3 \pm 4.0^\circ$ left deviation in the R axis is mainly associated with a more horizontal cardiac long axis, with electrophysiological factors playing very little role. This is partly explained by female R axes being more sensitive to changes in verticality – the regression coefficient between R axis and verticality, shown in Figure 4C, is higher for women than men. Figures 4A and 4B also demonstrate that the female T axis was more right deviated than male, largely associated with electrophysiology.

Figure 4E shows the regression coefficient of R axis against torso volume. It reveals torso volume significantly affected R axis for both sexes in healthy, but not post-MI subjects. For example, in healthy women, every 10dm^3 increase in torso volume was associated with a decrease in R axis of $7.2 \pm 3.0^\circ$. This was related to increased distance to the LL electrode (inversely related to the R axis) without as substantial an increase in distance to the LA electrode (positively related to the R axis). For healthy women, every 10dm^3 increase in torso volume was associated with an increased distance to the heart centre of $19.4 \pm 0.7\text{mm}$ and $5.9 \pm 0.6\text{mm}$ for the LL and LA electrodes,

respectively. However, in post-MI women, the gap between these differences was smaller, with corresponding increased distances of 16.6 ± 1.8 mm and 8.3 ± 1.7 mm.

Discussion

This study provides a comprehensive evaluation into sex differences of the interplay between anatomy and key ECG biomarkers in healthy and post-MI subjects. The development and validation of a novel computational image processing pipeline for automated 3D torso reconstruction enables high-throughput investigations into the effects of anatomical variability on the study of a wide range of cardiac diseases. The adaptation of the pipeline to standard clinical cardiac imaging allows for the exploitation of extensive databases, such as the UK Biobank, and sets the groundwork for clinical tools to personalise ECG interpretation considering patient anatomy.

Our analysis provides the following insights in how sex-related anatomical differences affected the impact of myocardial infarction on key ECG biomarkers. 1) Smaller cavity volumes are associated with approximately half of the shorter QRS durations in women than men in both healthy and post-MI subjects. 2) Female ST_T amplitude is lower than male in all precordial leads for healthy and post-MI subjects, associated with a smaller LV mass and more superior and posterior position. 3) Female TWA changes following MI are more associated with anatomical factors than male. 4) R axis is more left deviated in MI than healthy populations; for women this is associated with a more horizontal cardiac long axis, but for men with electrophysiology.

The high Dice coefficient for the contour extraction and low surface-to-contour distances show that the extracted contours are near the ground truth torso-air boundary on the clinical imaging, as well as to the final reconstructed surfaces. As detailed in Smith et al. [34], the three-step segmentation, automated post-processing, and refinement process was particularly effective at removing errors inside the torso at locations of sharp gradient between dark and light, for example due to air in the lungs. Whilst reconstructed 3D torso surfaces were largely smooth and realistic, there were small irregularities, generally on the shoulder and waist regions. These were not typically associated with incomplete CMR artifact removal and may instead result from images being taken on different breath-holds or the image sparsity. The low surface-to-surface distance between reconstructions made using the automatically and manually derived contours demonstrates that the small errors in contour extraction did not lead to significant errors in electrode placement on the torso meshes. Surfaces were generally more accurate on the upper left quadrant of the torso, which is where most electrodes are placed. The validity of the torso-ventricular reconstruction pipeline is further supported as known trends in torso and cardiac size, position, and orientation were qualitatively reproduced, and furthermore were quantitatively characterised. This includes the more superior female cardiac position [31, 42] and the horizontal shift in the cardiac long axis for subjects with a larger BMI [7, 43, 44]. As this pipeline uses clinically standard CMR imaging, it can be more easily translated to a clinical tool to correct for the effect of anatomical variation on the ECG in a variety of medical contexts.

QRS duration and its dispersion have been a key component of proposed electrocardiographic risk stratification for SCD following MI. However, findings of its effectiveness are mixed, with some trials suggesting it is a predictor of arrhythmic risk following MI [11, 12], but others finding that it is more likely a predictor of general mortality than arrhythmic inducibility [45, 46]. Our finding of QRS prolongation in post-MI subjects is consistent with previous studies finding increased QRS duration in the acute stages post-MI, particularly cooccurring with ST changes in leads exploring the infarcted region [47]. It is associated with slowed conduction velocity in the infarcted region itself, involvement of the conduction system such as the bundle branches and Purkinje, and in the chronic

case, with dilation and hypertrophy of non-infarcted myocardium [45, 48-51]. Both clinically and moderately prolonged QRS durations have been associated with increased risk following MI [52, 53]. We found that even small differences in the QRS duration are significant in the context of risk stratification, as in lead aVL, mean QRS duration would have to increase by 27% more for women than men to exceed 120ms and by 68% more to exceed 100ms. The combination of increased TWA and QRS prolongation in the septal leads from electrophysiology for women may specifically point to hypertrophy or remodelling of the calcium channels in the septum. However, this may be missed if structure were not accounted for, particularly as structural changes mask the septal TWA increase. Our finding that QRS duration is consistently related to cardiac orientation, particularly for healthy subjects, is broadly in line with simulation studies [2], but goes further to characterise trends across large populations and between demographic groups. The increased sensitivity of QRS duration to cardiac orientation for healthy over post-MI subjects may be explained by the deformation of the depolarisation wave's path through the myocardium for infarcted patients [48]. Our finding that QRS duration is positively associated with cardiac size is in line with both previous simulation studies [1], and with the biophysical understanding that larger distances require longer to traverse [54]. However, the estimation of the proportion of sex differences associated with structural variation is striking. This suggests that if demographic characteristics that impact anatomy, particularly sex, are taken into account as proxies for those anatomical differences, corrected QRS duration may be more representative of the underlying electrophysiological changes that predispose post-MI patients to SCD. This may therefore improve its accuracy in risk stratification, without the need for CMR imaging.

The finding that STj amplitude was substantially lower in women than men in all septal and anterior leads (V1-V4) is striking. The ECG data was not from subjects undergoing myocardial infarctions, so is instructive in sex differences in the normal baseline of STj amplitude, from which any deviations can be measured. The Fourth Universal Definition of Myocardial Infarction only has sex-specific thresholds in leads V2 and V3 for identifying ST-elevation, a key component in the diagnosis of ST-elevated myocardial infarctions (STEMIs) [15]. The dual effects of cardiac and torso volume necessitate the interpretation of ST-elevation through the lens of the anatomical context of the patient. For example, in lead V4, the clinical threshold for both female and male ST-elevation is 0.1mV [15]. However, due to their lower baseline values, overweight healthy women would need a 63% larger increase in STj amplitude to be classified as ST-elevated than normal weight healthy men. This is a critical observation in the context of higher incidence of missed diagnosis and increased female MI mortality following infarction [20, 23, 24]. Previous studies have suggested STj amplitude in acute ischemia is affected by cardiac rotations and translations, but this was severely limited by the use of a single, canine heart [55]. Therefore, this work represents a significant step forward in the characterisation of anatomical influences of its baseline level, from which deviations can then be more accurately identified in the context of anatomical variation. It also highlights the important intersectional effects of demographic characteristics such as sex and BMI, suggesting that both of these factors should be accounted for in assessing ST-elevation to improve accuracy of MI diagnosis and classification.

Changes in TWA are particularly clinically relevant in MI. Alterations to the polarity and amplitude of the T wave are used in diagnosis of acute MI [15] and TWA affects proposed risk stratification tools, particularly markers of repolarisation abnormalities [9, 56]. Acutely, hyperkalaemia can lead to hyperacute T waves [57, 58], and repolarisation heterogeneities to T wave inversion [59]. Conversely, in the progression to chronic MI, reductions in L-type calcium and sodium currents [60, 61] can both cause TWA reductions [62, 63]. However, particularly in the female case, a substantial proportion of TWA changes are associated with anatomical differences, such as the decreased

verticality of the post-MI heart. This implies that without anatomical corrections, such as adjusting for BMI, the female TWA may be less representative of its underlying electrophysiology, and therefore less representative of any pathology. This increased effect of anatomical parameters in women is partly due to their larger difference in key anatomical factors such as posterior position and long axis orientation from healthy to post-MI subjects. This reflects their larger increase in BMI, associated with a more horizontal cardiac long axis, as described in previous works suggesting this was related to an upwards shift in the diaphragm [7, 44]. Furthermore, the female TWA is more sensitive to changes in anatomical parameters. This could be explained by sex differences in the directions of the T wave vectors and nonlinearities in the structure-function relationships. Our finding that TWA is affected by cardiac position and orientation is in line with previous simulation studies [2], and with the physical understanding that amplitude increases with decreased distance to the electrode and better alignment between the lead axes and electrical vectors. However, while their small number of anatomies (N=5) limited their conclusions to these relationships being significant but patient dependent, the vast increase in anatomies that our work considered (N=1476) facilitated more specific characterisations of population and demographic trends. This demonstrates the power of automated, high-throughput anatomical reconstruction to ensure that electrophysiological parameters are equally representative of underlying pathology in women and men.

Electrical axes are fundamental to parameters proposed to stratify arrhythmic risk post-MI, particularly markers of electrical heterogeneity such as QRS-T angle [64]. However, the relationship between electrical and anatomical axes is complex and poorly understood, with previous works on smaller populations even disagreeing over whether there was any significant relationship between the two [7, 65, 66]. Our finding that the verticality of the cardiac long axis is particularly significant in determining R axis angle, especially for women, represents a step forward in understanding this relationship. Moreover, our work finds that while the left-deviation of the R axis may look similar in both sexes, in men this is representative of an electrophysiological change, presumably alterations to the direction of the QRS vector caused by both hypertrophy and areas of electrical inactivity. However, for women this change is almost entirely associated with anatomical differences. This suggests that it is particularly important to correct R axis angles for anatomical factors in women to avoid misattribution of deviations to electrophysiological pathology. The significant relationship between torso volume and R axis angle for healthy subjects indicates that BMI may be a critical consideration when using ECG based biomarkers that are sensitive to axis angles. The fact that increased torso volume moved the LL electrode further away from the heart centre than the LA electrode suggests that this relationship is caused by a stretching of the anatomical plane in which these angles are viewed. Therefore, this represents another area in which a clinical tool that accounts for anatomical variation in interpreting ECG biomarkers would reduce differential accuracy in risk stratification between demographic groups.

Due to the demographics of the UK Biobank [39], this study only included subjects aged 45-80 years old; however, the incidence of MI is very low in people younger than this range [24]. The ethnic background of the dataset also had limited diversity; however, this is roughly representative of the UK Biobank population and thus the ethnic composition of the UK at the time of recruitment [67]. As these databases become more diverse, the mediating effect of ethnic background on the cardiac structure-function relationship should be investigated. Whilst the healthy subjects had a roughly equal sex balance, the post-MI subjects had a male-female ratio of approximately 4:1, in line with sex differences in diagnosed MI events [68]. The statistical shape model used for initial torso reconstruction did not have separate male and female models, which may have led to an underestimation in the sex differences in the torsos of male and female anatomies. However, the

deformation step of the reconstruction was designed to address this issue by modelling personalised torso variations. The exact electrode positions of the obtained ECGs were not available in the UK Biobank dataset, so these were estimated by placing the electrodes on standard positions on the personalised torso. However, this estimation process allows for use of many similar clinically standard datasets, avoiding complex acquisition protocols. As the anatomical factors cannot represent a perfectly complete and independent basis set, the electrophysiological contribution can only be estimated. However, as there is more anatomical variation than can be represented using the given factors, such as the smaller scale geometry of the heart, the role of the anatomy versus electrophysiology was likely under- rather than overestimated. Whilst the study benefits from the electrophysiological and anatomical variation that using clinical data facilitates, it is impossible to fully separate the effects of these factors on ECG biomarkers, as characteristics such as sex affect both categories, making them interdependent. This means that this work complements computational studies, which have less interpersonal variability, but isolate the effect of positional and rotational changes. Future work should focus on the characterisation of ECG biomarkers in myocardial infarction and other disease conditions. The automated torso reconstruction pipeline this work proposes enables the expansion of the number of anatomies such future computational studies would investigate.

To conclude, this work presents a novel automated pipeline for personalised torso reconstruction and demonstrates its power in exploiting large clinical databases by relating torso-ventricular anatomy and ECG biomarkers for 1051 healthy and 425 post-MI subjects. Results show the considerable influence of anatomical factors in demographic differences of ECG biomarkers. Women and individuals with higher BMI may be disproportionately affected. This underlines the importance of, and lays the foundations for, personalised ECG interpretation that considers a subject's individual torso-ventricular anatomy, facilitating improvement in diagnosis and risk stratification tools.

Acknowledgements

This research has been conducted using the UK Biobank Resource under Application Number '40161'. The authors would like to thank Dr Ernesto Zacur for providing valuable suggestions on the 3D anatomical mesh reconstruction.

Funding

HJS is supported by a Wellcome Trust Studentship under Grant No. 102161/Z/13/Z. AB is a Royal Society University Research Fellow and is supported by the Royal Society Grant No. URF\R1\221314. The works of AB, RPC, and VG are supported by the British Heart Foundation (BHF) Project under Grant PG/20/21/35082. This work was funded by a Wellcome Trust Fellowship in Basic Biomedical Sciences to BR (214290/Z/18/Z) and the CompBioMed 2 Centre of Excellence in Computational Biomedicine (European Commission Horizon 2020 research and innovation programme, grant agreements No. 823712) to BR and VG.

Conflict of interest: none declared.

Data availability statement

The UK Biobank data is publicly available and can be accessed by application upon approval here: <https://www.ukbiobank.ac.uk/enable-your-research/register>. The reconstruction pipeline code will be made publicly available upon acceptance.

References

- 1 Mincholé, A., Zacur, E., Ariga, R., Grau, V., and Rodriguez, B.: 'MRI-Based Computational Torso/Biventricular Multiscale Models to Investigate the Impact of Anatomical Variability on the ECG QRS Complex', *Front Physiol*, 2019, 10
- 2 Nguyễn, U.C., Potse, M., Regoli, F., Caputo, M.L., Conte, G., Murzilli, R., Muzzarelli, S., Moccetti, T., Caiani, E.G., Prinzen, F.W., Krause, R., and Auricchio, A.: 'An in-silico analysis of the effect of heart position and orientation on the ECG morphology and vectorcardiogram parameters in patients with heart failure and intraventricular conduction defects', *Journal of Electrocardiology*, 2015, 48, (4), pp. 617-625
- 3 MacLeod, R.S., Ni, Q., Punske, B., Ershler, P.R., Yilmaz, B., and Taccardi, B.: 'Effects of heart position on the body-surface electrocardiogram', *Journal of Electrocardiology*, 2000, 33, pp. 229-237
- 4 Bradley, C.P., Pullan, A.J., and Hunter, P.J.: 'Effects of Material Properties and Geometry on Electrocardiographic Forward Simulations', *Annals of Biomedical Engineering*, 2000, 28, (7), pp. 721-741
- 5 Hoekema, R., Uijen, G.J.H., and Oosterom, A.v.: 'Geometrical aspects of the interindividual variability of multilead ECG recordings', *IEEE Transactions on Biomedical Engineering*, 2001, 48, (5), pp. 551-559
- 6 Prajapati, C., Koivumäki, J., Pekkanen-Mattila, M., and Aalto-Setälä, K.: 'Sex differences in heart: from basics to clinics', *European Journal of Medical Research*, 2022, 27, (1), pp. 241
- 7 Sathananthan, G., Aggarwal, G., Zahid, S., Byth, K., Chik, W., Friedman, D., and Thiagalingam, A.: 'Computed tomography-guided in vivo cardiac orientation and correlation with ECG in individuals without structural heart disease and in age-matched obese and older individuals', *Clinical Anatomy*, 2015, 28, (4), pp. 487-493
- 8 Sang, Y., Beetz, M., and Grau, V.: 'Generation of 12-Lead Electrocardiogram with Subject-Specific, Image-Derived Characteristics Using a Conditional Variational Autoencoder', in Editor (Ed.)^(Eds.): 'Book Generation of 12-Lead Electrocardiogram with Subject-Specific, Image-Derived Characteristics Using a Conditional Variational Autoencoder' (2022, edn.), pp. 1-5
- 9 Bui, A.H., and Waks, J.W.: 'Risk Stratification of Sudden Cardiac Death After Acute Myocardial Infarction', *J Innov Card Rhythm Manag*, 2018, 9, (2), pp. 3035-3049
- 10 Dagres, N., and Hindricks, G.: 'Risk stratification after myocardial infarction: is left ventricular ejection fraction enough to prevent sudden cardiac death?', *European Heart Journal*, 2013, 34, (26), pp. 1964-1971
- 11 Dhar, R., Alsheikh-Ali, A.A., Estes, N.A.M., Moss, A.J., Zareba, W., Daubert, J.P., Greenberg, H., Case, R.B., and Kent, D.M.: 'Association of prolonged QRS duration with ventricular tachyarrhythmias and sudden cardiac death in the Multicenter Automatic Defibrillator Implantation Trial II (MADIT-II)', *Heart Rhythm*, 2008, 5, (6), pp. 807-813
- 12 Zimetbaum, P.J., Buxton, A.E., Batsford, W., Fisher, J.D., Hafley, G.E., Lee, K.L., O'Toole, M.F., Page, R.L., Reynolds, M., and Josephson, M.E.: 'Electrocardiographic Predictors of Arrhythmic Death and Total Mortality in the Multicenter Unsustained Tachycardia Trial', *Circulation*, 2004, 110, (7), pp. 766-769
- 13 Waks, J.W., and Tereshchenko, L.G.: 'Global electrical heterogeneity: A review of the spatial ventricular gradient', *Journal of Electrocardiology*, 2016, 49, (6), pp. 824-830
- 14 Oehler, A., Feldman, T., Henrikson, C.A., and Tereshchenko, L.G.: 'QRS-T Angle: A Review', *Annals of Noninvasive Electrocardiology*, 2014, 19, (6), pp. 534-542
- 15 Thygesen, K., Alpert, J.S., Jaffe, A.S., Chaitman, B.R., Bax, J.J., Morrow, D.A., White, H.D., and Group, E.S.C.S.D.: 'Fourth universal definition of myocardial infarction (2018)', *European Heart Journal*, 2019, 40, (3), pp. 237-269
- 16 Collet, J.-P., Thiele, H., Barbato, E., Barthélémy, O., Bauersachs, J., Bhatt, D.L., Dendale, P., Dorobantu, M., Edvardsen, T., Folliguet, T., Gale, C.P., Gilard, M., Jobs, A., Jüni, P., Lambrinou, E., Lewis, B.S., Mehilli, J., Meliga, E., Merkely, B., Mueller, C., Roffi, M., Rutten, F.H., Sibbing, D., Siontis, G.C.M., Chettibi, M., Hayrapetyan, H.G., Metzler, B., Najafov, R., Stelmashok, V.I., Claeys, M., Kušljugić, Z., Gatzov, P.M., Skoric, B., Panayi, G., Mates, M., Sorensen, R., Shokry, K., Marandi, T., Kajander, O.A., Commeau, P., Aladashvili, A., Massberg, S., Nikas, D., Becker, D., Guðmundsdóttir, I.J., Peace, A.J., Beigel, R., Indolfi, C., Aidargaliyeva, N., Elezi, S., Beishenkulov, M., Maca, A., Gustiene, O., Degrell, P., Cassar Maempel, A., Ivanov, V., Damman, P., Kedev, S., Steigen, T.K., Legutko, J., Morais, J., Vinereanu, D., Duplyakov, D., Zavatta, M., Pavlović, M., Orban, M., Bunc, M., Ibañez, B., Hofmann, R., Gaemperli, O., Marjeh, Y.B., Addad, F., Tutar, E., Parkhomenko, A., Karia, N., and Group, E.S.C.S.D.: '2020 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: The Task Force for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC)', *European Heart Journal*, 2021, 42, (14), pp. 1289-1367
- 17 Ibanez, B., James, S., Agewall, S., Antunes, M.J., Bucciarelli-Ducci, C., Bueno, H., Caforio, A.L.P., Crea, F., Goudevenos, J.A., Halvorsen, S., Hindricks, G., Kastrati, A., Lenzen, M.J., Prescott, E., Roffi, M., Valgimigli, M., Varenhorst, C., Vranckx, P., Widimský, P., and Group, E.S.C.S.D.: '2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation: The Task Force for the management of acute myocardial infarction in patients presenting with ST-segment elevation of the European Society of Cardiology (ESC)', *European Heart Journal*, 2018, 39, (2), pp. 119-177

- 18 Bugiardini, R., and Cenko, E.: 'Sex differences in myocardial infarction deaths', *The Lancet*, 2020, 396, (10244), pp. 72-73
- 19 Brush, J.E., Krumholz, H.M., Greene, E.J., and Dreyer, R.P.: 'Sex Differences in Symptom Phenotypes Among Patients With Acute Myocardial Infarction', *Circulation: Cardiovascular Quality and Outcomes*, 2020, 13, (2), pp. e005948
- 20 Pope, J.H., Aufderheide, T.P., Ruthazer, R., Woolard, R.H., Feldman, J.A., Beshansky, J.R., Griffith, J.L., and Selker, H.P.: 'Missed diagnoses of acute cardiac ischemia in the emergency department', *New England Journal of Medicine*, 2000, 342, (16), pp. 1163-1170
- 21 Chandra, N.C., Ziegelstein, R.C., Rogers, W.J., Tiefenbrunn, A.J., Gore, J.M., French, W.J., Rubison, M., and for the National Registry of Myocardial Infarction, I.: 'Observations of the Treatment of Women in the United States With Myocardial Infarction: A Report From the National Registry of Myocardial Infarction-I', *Archives of Internal Medicine*, 1998, 158, (9), pp. 981-988
- 22 Bairey Merz, C.N., Shaw Leslee, J., Reis Steven, E., Bittner, V., Kelsey Sheryl, F., Olson, M., Johnson, B.D., Pepine Carl, J., Mankad, S., Sharaf Barry, L., Rogers William, J., Pohost Gerald, M., Lerman, A., Quyyumi Arshed, A., Sopko, G., and null, n.: 'Insights From the NHLBI-Sponsored Women's Ischemia Syndrome Evaluation (WISE) Study', *Journal of the American College of Cardiology*, 2006, 47, (3_Supplement), pp. S21-S29
- 23 Fang, J., Alderman, M.H., Keenan, N.L., and Ayala, C.: 'Acute Myocardial Infarction Hospitalization in the United States, 1979 to 2005', *The American Journal of Medicine*, 2010, 123, (3), pp. 259-266
- 24 Benjamin, E.J., Muntner, P., Alonso, A., Bittencourt, M.S., Callaway, C.W., Carson, A.P., Chamberlain, A.M., Chang, A.R., Cheng, S., Das, S.R., Delling, F.N., Djousse, L., Elkind, M.S.V., Ferguson, J.F., Fornage, M., Jordan, L.C., Khan, S.S., Kissela, B.M., Knutson, K.L., Kwan, T.W., Lackland, D.T., Lewis, T.T., Lichtman, J.H., Longenecker, C.T., Loop, M.S., Lutsey, P.L., Martin, S.S., Matsushita, K., Moran, A.E., Mussolino, M.E., O'Flaherty, M., Pandey, A., Perak, A.M., Rosamond, W.D., Roth, G.A., Sampson, U.K.A., Satou, G.M., Schroeder, E.B., Shah, S.H., Spartano, N.L., Stokes, A., Tirschwell, D.L., Tsao, C.W., Turakhia, M.P., VanWagner, L.B., Wilkins, J.T., Wong, S.S., Virani, S.S., and null, n.: 'Heart Disease and Stroke Statistics—2019 Update: A Report From the American Heart Association', *Circulation*, 2019, 139, (10), pp. e56-e528
- 25 Coventry, L.L., Finn, J., and Bremner, A.P.: 'Sex differences in symptom presentation in acute myocardial infarction: A systematic review and meta-analysis', *Heart & Lung*, 2011, 40, (6), pp. 477-491
- 26 Shaw Leslee, J., Bairey Merz, C.N., Pepine Carl, J., Reis Steven, E., Bittner, V., Kelsey Sheryl, F., Olson, M., Johnson, B.D., Mankad, S., Sharaf Barry, L., Rogers William, J., Wessel Timothy, R., Arant Christopher, B., Pohost Gerald, M., Lerman, A., Quyyumi Arshed, A., Sopko, G., and null, n.: 'Insights From the NHLBI-Sponsored Women's Ischemia Syndrome Evaluation (WISE) Study', *Journal of the American College of Cardiology*, 2006, 47, (3_Supplement), pp. S4-S20
- 27 Kittnar, O.: 'Sex Related Differences in Electrocardiography', *Physiological Research*, 2023, 72
- 28 Daniels, S.R., Kimball, T.R., Morrison, J.A., Khoury, P., Witt, S., and Meyer, R.A.: 'Effect of Lean Body Mass, Fat Mass, Blood Pressure, and Sexual Maturation on Left Ventricular Mass in Children and Adolescents', *Circulation*, 1995, 92, (11), pp. 3249-3254
- 29 Kansal, S., Roitman, D., and Sheffield, L.: 'A quantitative relationship of electrocardiographic criteria of left ventricular hypertrophy with echocardiographic left ventricular mass: a multivariate approach', *Clin Cardiol*, 1983, 6, (9), pp. 456-463
- 30 Gambill, C.L., Wilkins, M.L., Haisty, W.K., Anderson, S.T., Maynard, C., Wagner, N.B., Startt Selvester, R.H., and Wagner, G.S.: 'T wave amplitudes in normal populations: Variation with ECG lead, sex, and age', *Journal of Electrocardiology*, 1995, 28, (3), pp. 191-197
- 31 Güneş Bolatlı, N.Ü., Mustafa Koplay, Zeliha Fazlıoğulları, Ahmet Kağan Karabulut: 'Examination of the levels of structures in the thorax in multidetector computerized tomography images', *Journal of Surgery and Medicine*, 2020, 4, (9), pp. 784-789
- 32 Piro, M., Della Bona, R., Abbate, A., Biasucci Luigi, M., and Crea, F.: 'Sex-Related Differences in Myocardial Remodeling', *Journal of the American College of Cardiology*, 2010, 55, (11), pp. 1057-1065
- 33 Ronneberger, O., Fischer, P., and Brox, T.: 'U-Net: Convolutional Networks for Biomedical Image Segmentation', in Editor (Ed.)^(Eds.): 'Book U-Net: Convolutional Networks for Biomedical Image Segmentation' (Springer International Publishing, 2015, edn.), pp. 234-241
- 34 Smith, H.J., Banerjee, A., Choudhury, R.P., and Grau, V.: 'Automated Torso Contour Extraction from Clinical Cardiac MR Slices for 3D Torso Reconstruction', in Editor (Ed.)^(Eds.): 'Book Automated Torso Contour Extraction from Clinical Cardiac MR Slices for 3D Torso Reconstruction' (2022, edn.), pp. 3809-3813
- 35 Robinette, K.M., Daanen, H., and Paquet, E.: 'The CAESAR project: a 3-D surface anthropometry survey', in Editor (Ed.)^(Eds.): 'Book The CAESAR project: a 3-D surface anthropometry survey' (1999, edn.), pp. 380-386
- 36 Pishchulin, L., Wuhrer, S., Helten, T., Theobalt, C., and Schiele, B.: 'Building statistical shape spaces for 3D human modeling', *Pattern Recognition*, 2017, 67, pp. 276-286
- 37 Rohr, K., Stiehl, H.S., Sprengel, R., Buzug, T.M., Weese, J., and Kuhn, M.H.: 'Landmark-based elastic registration using approximating thin-plate splines', *IEEE Transactions on Medical Imaging*, 2001, 20, (6), pp. 526-534
- 38 Banerjee, A., Camps, J., Zacur, E., Andrews, C.M., Rudy, Y., Choudhury, R.P., Rodriguez, B., and Grau, V.: 'A completely automated pipeline for 3D reconstruction of human heart from 2D cine magnetic resonance slices', *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 2021, 379, (2212), pp. 20200257
- 39 Petersen, S.E., Matthews, P.M., Bamberg, F., Bluemke, D.A., Francis, J.M., Friedrich, M.G., Leeson, P., Nagel, E., Plein, S., Rademakers, F.E., Young, A.A., Garratt, S., Peakman, T., Sellors, J., Collins, R., and Neubauer, S.: 'Imaging in

- population science: cardiovascular magnetic resonance in 100,000 participants of UK Biobank - rationale, challenges and approaches', *Journal of Cardiovascular Magnetic Resonance*, 2013, 15, (1), pp. 46
- 40 Schlegel, T.T., DePalma, J.L., and Moradi, S.: 'Real-time, high frequency QRS electrocardiograph', in Editor (Ed.) (Eds.): 'Book Real-time, high frequency QRS electrocardiograph' (Google Patents, 2006, edn.), pp.
- 41 Dougherty, J.D.: 'The relation of the frontal QRS axis to the anatomic position of the heart', *Journal of Electrocardiology*, 1970, 3, (3), pp. 267-284
- 42 Mirjalili, S.A., Hale, S.J.M., Buckenham, T., Wilson, B., and Stringer, M.D.: 'A reappraisal of adult thoracic surface anatomy', *Clinical Anatomy*, 2012, 25, (7), pp. 827-834
- 43 Rautaharju, P.M., Zhou, S.H., and Calhoun, H.P.: 'Ethnic differences in ECG amplitudes in North American white, black, and Hispanic men and women: Effect of obesity and age', *Journal of Electrocardiology*, 1994, 27, pp. 20-31
- 44 Engblom, H., Hedström, E., Palmer, J., Wagner, G.S., and Arheden, H.: 'Determination of the left ventricular long-axis orientation from a single short-axis MR image: relation to BMI and age', *Clinical Physiology and Functional Imaging*, 2004, 24, (5), pp. 310-315
- 45 Yerra, L., Anavekar, N., Skali, H., Zelenkofske, S., Velazquez, E., McMurray, J., Pfeffer, M., and Solomon, S.D.: 'Association of QRS duration and outcomes after myocardial infarction: the VALIANT trial', *Heart Rhythm*, 2006, 3, (3), pp. 313-316
- 46 Buxton Alfred, E., Sweeney Michael, O., Wathen Mark, S., Josephson Mark, E., Otterness Mary, F., Hogan-Miller, E., Stark Alice, J., DeGroot Paul, J., and null, n.: 'QRS Duration Does Not Predict Occurrence of Ventricular Tachyarrhythmias in Patients With Implanted Cardioverter-Defibrillators', *Journal of the American College of Cardiology*, 2005, 46, (2), pp. 310-316
- 47 Chávez-González, E., Jiménez, A.R., and Moreno-Martínez, F.: 'QRS duration and dispersion for predicting ventricular arrhythmias in early stage of acute myocardial infarction', *Medicina Intensiva (English Edition)*, 2017, 41, (6), pp. 347-355
- 48 Bacharova, L., Bang, L.E., Szathmary, V., and Mateasik, A.: 'Imaging QRS complex and ST segment in myocardial infarction', *Journal of Electrocardiology*, 2014, 47, (4), pp. 438-447
- 49 Sheridan, D.J., Culling, W., and Penny, W.J.: 'Electrophysiological Disturbances Associated with Acute Myocardial Infarction', *European Heart Journal*, 1986, 7, (suppl_B), pp. 11-18
- 50 Brembilla-Perrot, B., Houriez, P., Claudon, O., Preiss, J.-P., and De La Chaise, A.T.: 'Evolution of QRS Duration after Myocardial Infarction: Clinical Consequences', *Pacing and Clinical Electrophysiology*, 1999, 22, (10), pp. 1466-1475
- 51 Strauss, D.G., and Selvester, R.H.: 'The QRS complex—a biomarker that “images” the heart: QRS scores to quantify myocardial scar in the presence of normal and abnormal ventricular conduction', *Journal of Electrocardiology*, 2009, 42, (1), pp. 85-96
- 52 Surawicz, B., Childers, R., Deal, B.J., and Gettes, L.S.: 'AHA/ACCF/HRS Recommendations for the Standardization and Interpretation of the Electrocardiogram', *Circulation*, 2009, 119, (10), pp. e235-e240
- 53 Pudil, R., Feinberg, M.S., Hod, H., Boyko, V., Mandelzweig, L., and Behar, S.: 'The prognostic significance of intermediate QRS prolongation in acute myocardial infarction', *International Journal of Cardiology*, 2001, 78, (3), pp. 233-239
- 54 van Oosterom, A., Hoekema, R., and Uijen, G.J.H.: 'Geometrical factors affecting the interindividual variability of the ECG and the VCG', *Journal of Electrocardiology*, 2000, 33, pp. 219-227
- 55 MacLeod, R.S., Lux, R.L., and Taccardi, B.: 'A possible mechanism for electrocardiographically silent changes in cardiac repolarization', *Journal of Electrocardiology*, 1998, 30, pp. 114-121
- 56 Tereshchenko, L.G., McNitt, S., Han, L., Berger, R.D., and Zareba, W.: 'ECG Marker of Adverse Electrical Remodeling Post-Myocardial Infarction Predicts Outcomes in MADIT II Study', *PLOS ONE*, 2012, 7, (12), pp. e51812
- 57 Somers, M.P., Brady, W.J., Perron, A.D., and Mattu, A.: 'The prominent T wave: Electrocardiographic differential diagnosis', *The American Journal of Emergency Medicine*, 2002, 20, (3), pp. 243-251
- 58 Montague, B.T., Ouellette, J.R., and Buller, G.K.: 'Retrospective review of the frequency of ECG changes in hyperkalemia', *Clin J Am Soc Nephrol*, 2008, 3, (2), pp. 324-330
- 59 Mandel, W.J., Burgess, M.J., Neville, J., Jr., and Abildskov, J.A.: 'Analysis of T-wave abnormalities associated with myocardial infarction using a theoretic model', *Circulation*, 1968, 38, (1), pp. 178-188
- 60 Hund, T.J., Decker, K.F., Kanter, E., Mohler, P.J., Boyden, P.A., Schuessler, R.B., Yamada, K.A., and Rudy, Y.: 'Role of activated CaMKII in abnormal calcium homeostasis and INa remodeling after myocardial infarction: Insights from mathematical modeling', *Journal of Molecular and Cellular Cardiology*, 2008, 45, (3), pp. 420-428
- 61 Decker, K.F., and Rudy, Y.: 'Ionic mechanisms of electrophysiological heterogeneity and conduction block in the infarct border zone', *Am J Physiol Heart Circ Physiol*, 2010, 299, (5), pp. H1588-1597
- 62 Moss, A.J., Zareba, W., Benhorin, J., Locati, E.H., Hall, W.J., Robinson, J.L., Schwartz, P.J., Towbin, J.A., Vincent, G.M., Lehmann, M.H., Keating, M.T., MacCluer, J.W., and Timothy, K.W.: 'ECG T-Wave Patterns in Genetically Distinct Forms of the Hereditary Long QT Syndrome', *Circulation*, 1995, 92, (10), pp. 2929-2934
- 63 Zemzemi, N., and Rodriguez, B.: 'Effects of L-type calcium channel and human ether-a-go-go related gene blockers on the electrical activity of the human heart: a simulation study', *EP Europace*, 2015, 17, (2), pp. 326-333
- 64 Borleffs, C.J.W., Scherptong, R.W.C., Man, S.-C., van Welsenes, G.H., Bax, J.J., van Erven, L., Swenne, C.A., and Schalij, M.J.: 'Predicting Ventricular Arrhythmias in Patients With Ischemic Heart Disease', *Circulation: Arrhythmia and Electrophysiology*, 2009, 2, (5), pp. 548-554

- 65 Engblom, H., Foster, J.E., Martin, T.N., Groenning, B., Pahlm, O., Dargie, H.J., Wagner, G.S., and Arheden, H.: 'The relationship between electrical axis by 12-lead electrocardiogram and anatomical axis of the heart by cardiac magnetic resonance in healthy subjects', *American Heart Journal*, 2005, 150, (3), pp. 507-512
- 66 Gnalini Sathananthan, S.Z., Gunjan Aggarwal, William Chik, Daniel Friedman, Aravinda Thiagalingam: 'Cardiac orientation: is there a correlation between the anatomical and the electrical axis of the heart?', *The British Journal of Cardiology*, 2015, 22, (2)
- 67 Fry, A., Littlejohns, T.J., Sudlow, C., Doherty, N., Adamska, L., Sprosen, T., Collins, R., and Allen, N.E.: 'Comparison of Sociodemographic and Health-Related Characteristics of UK Biobank Participants With Those of the General Population', *American Journal of Epidemiology*, 2017, 186, (9), pp. 1026-1034
- 68 Bhatnagar, P., Wickramasinghe, K., Williams, J., Rayner, M., and Townsend, N.: 'The epidemiology of cardiovascular disease in the UK 2014', *Heart*, 2015, 101, (15), pp. 1182




Figure 1. A: Proposed end-to-end automated 3D torso reconstruction pipeline from 2D standard clinical cardiac magnetic resonance scans. The torso contours are first extracted from the images. A SSM is fitted to the contours, which is then optimally deformed. SSM: statistical shape model, α : spin of the cardiac short axis plane around the torso vertical axis, β : verticality of the cardiac long axis, γ : tilt of the cardiac short axis plane. **B: (i)** comparison of Dice coefficient between single stage contouring (blue) and 3 stage segmentation, postprocessing and refinement (red). **(ii)** surface-to-contour distance for an example case between the automatically reconstructed surface and the automatically generated torso contours. **(iii)** surface-to-surface distance between the torso mesh created using the automated pipeline and the manually annotated contours for the subjects with the smallest (left) and largest (right) electrode error respectively.




Figure 2. **A, B:** QRS duration across subjects for each ECG lead in healthy and post-MI subjects respectively with women shown as pale red circles and men as cyan triangles. **C:** contribution of each parameter to the difference in QRS duration (QRSd) from women to men for healthy subjects, calculated by multiplying the regression coefficient for each factor by its mean difference between male and female populations. **D:** correspondingly from women to men for post-MI subjects. **E, F:** mean QRS durations across subjects for each ECG lead in men and women respectively. Healthy subjects are shown as black squares, post-MI subjects as red diamonds. **G:** contribution of each parameter to the change in QRS duration from healthy to post-MI subjects in men, and **H:** correspondingly from healthy to post-MI subjects in women, calculated by multiplying the regression coefficient for each factor by its mean difference between healthy and post-MI populations. Female y-axis limits have been adjusted by the difference in healthy QRS duration between sexes for ease of comparison.




Figure 3. A: Mean ST amplitudes measured at the J point (STj), across subjects for each ECG lead in healthy subjects with women shown as pale red circles and men as cyan triangles. **B:** contribution of each parameter to the difference in STj amplitude from women to men for healthy subjects, calculated by multiplying the regression coefficient for each factor by its mean difference between male and female populations. Corresponding figures for post-MI subjects are shown in Supplementary Figure 6. **C, D:** mean T wave amplitudes (TWA) across subjects for each ECG lead in men and women respectively. Healthy subjects are shown as black squares, post-MI subjects as red diamonds. **E:** contribution of each parameter to the change in T wave amplitude from healthy to post-MI subjects in men, and **F:** correspondingly from healthy to post-MI subjects in women, calculated by multiplying the regression coefficient for each factor by its mean difference between healthy and post-MI populations. Female y-axis limits have been adjusted by the difference in healthy T wave amplitude between sexes for ease of comparison.




Figure 4. A, B: Contribution of each parameter to the change in axis angle from healthy to post-MI subjects in men and women respectively, calculated by multiplying the regression coefficient for each factor by its mean difference between healthy and post-MI populations. C, D: normalised regression coefficients showing the association between the verticality of the cardiac long axis and the R and T axes respectively for the healthy and post-MI populations. Bars shown in red represent regression coefficients significantly different from 0, at a significance level of 0.05. E, F: normalised regression coefficients showing the association between torso volume and the R and T axes respectively for the healthy and post-MI populations.

Table 1. Demographic characteristics of the 1476 subjects from the UK Biobank cohort.

Characteristics	Healthy (N = 1051)	MI (N = 425)	p-value
Age (years)	61.1 ± 7.5	67.3 ± 6.2	< 0.0001
Sex: Female, n (%)	581 (54.3)	84 (19.8)	< 0.0001
BMI (kg/m ²)	25.9 ± 4.1	28.1 ± 4.4	< 0.0001
Systolic blood pressure	133±17	142±19	< 0.0001
Diastolic blood pressure	80.3±10.0	81.4±10.7	0.08
Ethnicity, n (%)			0.5
White	1021 (97.1)	413 (97.2)	1
Mixed	3 (0.3)	1 (0.2)	0.9
Asian or Asian British	8 (0.8)	6 (1.4)	0.2
Black or Black British	7 (0.7)	1 (0.2)	0.3
Chinese	5 (0.5)	0 (0.0)	0.2
Other ethnic group	5 (0.5)	2 (0.5)	1
Prefer not to answer	2 (0.2)	2 (0.5)	0.3

Continuous variables are shown as mean ± standard deviation and categorical variables as number of subjects (percentage in bracket). The p-value refers to the null hypothesis that the mean, or proportion in a category, is equal in the healthy and MI subpopulations. Distributions of continuous variables were tested for normality and equal variance, and the appropriate statistical test was selected, as outlined in the statistical methods Supplementary Appendix 1.5. χ^2 -tests were performed to assess the proportions of subjects for categorical variables. MI: myocardial infarction, BMI: body mass index.

Table 2. Disease and treatment characteristics of 425 post-MI subjects from the UK Biobank cohort.

Characteristics	Male (N = 341)	Female (N = 84)	p-value
Disease prevalence (%)			
Primary (essential) hypertension	241 (70.7)	50 (59.5)	0.07
Atrial fibrillation or flutter	60 (17.6)	10 (11.9)	0.3
Heart failure	49 (14.4)	10 (11.9)	0.1
Renal failure	34 (10.0)	11 (13.1)	0.5
Diabetes	29 (8.5)	6 (7.1)	0.9
Surgical treatment (%)			
Replacement of coronary artery	52 (15.2)	1 (1.2)	0.0009
Bypass of coronary artery	57 (16.7)	2 (2.4)	0.001
Balloon angioplasty of coronary artery without stent	41 (12.0)	5 (6.0)	0.2
Balloon angioplasty of coronary artery with stent	135 (39.6)	21 (25.0)	0.02

Categorical variables as number of subjects (percentage in bracket). χ^2 -tests were performed to assess the proportions of subjects with each disease or treatment. The p-value refers to the null hypothesis that the proportion in a category is equal in the male and female subpopulations. Surgical treatment information is sourced from hospital records, whereas disease information is obtained from multiple sources including hospital records, primary care and self-report.

Supplementary Appendix 1: Methods

1.1 Dataset. Myocardial infarction (MI) was defined as any history of MI including ST- and non-ST-elevation MI, as obtained from patients' self-report in the baseline questionnaire and nurse-led interview, and linked hospital admission data [1]. The exclusion criteria for the subjects used in the larger statistical analysis are shown in Supplemental Figure 1. From the 1646 controls, 3 cases were excluded for incomplete image sets, and 592 were excluded for having at least one of 82 disease diagnoses in their UKB records, including circulatory system, renal, genitourinary, and endocrine disorders. The final set included 1051 'healthy' controls (581 female and 470 male), as shown in the left panel of S1. From the 479 post-MI subjects, 6 were excluded for incomplete image sets, 13 for missing ECG data, and 35 were excluded due to their imaging occurring before the MI event, thus retaining 425 post-MI subjects (84 female and 341 male), as shown in the right panel of S1. To select the test set used to evaluate the automated pipeline, both male and female subjects in the control population were split into deciles by their BMI. Then each decile was evenly sampled to create a test set that spanned the range of BMI for both sexes. The range of ages of the test subjects was then calculated to ensure that they represented the underlying population.

Note that the UKB protocol for imaging and ECG involved both being performed in a supine position.

1.2 Reconstruction. To generate the initial 3D torso mesh, a statistical shape model (SSM) was applied over the automatically extracted sparse torso contours in 3D space, which was then optimally deformed by minimising the distance to the extracted contours to produce the final 3D torso mesh.

1.2.1 Fitting the SSM. In order to generate the torso meshes from the sparse torso contours, a high-resolution statistical shape model (SSM) of human body shapes, was employed for the initial 3D torso reconstruction. The SSM was generated from the 3D optical body surface scanning of 4308 subjects in the CAESAR study, the largest commercially available scan database to date [2]. Detailed information on the SSM generation is provided in [3]. As the initial condition, the mean SSM was translated into the subject's coordinate system by positioning the centre of heart from the mean SSM onto the centre of the subject's heart. The latter centre was approximated by the closest point to the intersection lines between the cine short axis (SAX) and long axis (LAX) slices. Over the automatically extracted sparse torso contours in 3D space, the SSM was fitted by the optimal estimation of principal components, followed by rigid transformation, such that:

$$\min_{q,R} \sum_i d(C_i, R(\bar{X} + \Phi q))^2$$

where C_i is the i^{th} point on sparse torso representation, \bar{X} the mean shape model, $\Phi = (\phi_1, \phi_2, \dots)$ the SSM modes of variations, q the estimated set of parameters along the principal modes, and R the rigid transformation. The fitting of the reconstructions was restricted to the first 40 principal components of the SSM.

1.2.2 Deforming for patient-specific torso reconstruction. The initial generated torso mesh is inherently limited by the variability in the SSM. Hence, in order to accurately capture subject-specific variations, a final deformation step was performed via approximate thin plate splines (TPS) [4] to produce the final 3D torso mesh, minimising the distance to the extracted contours. In case the automated torso contour extraction generated sparse outlier points, any torso points over the 95th percentile were removed.

The initial deformation field was computed by considering the closest point on the SSM-generated mesh from the points on the sparse representation. A series of small, smooth deformations using approximate TPS was then applied to push the 3D mesh closer to the torso points, while preserving the smoothness and local topological properties [5]. Let F be the deformation field such that:

$$\arg \min_F \lambda \sum_i (||F(P_i) - Q_i'||)^2 + J_2^3(F) \text{ and } Q_i' = P_i + \beta \frac{(Q_i - P_i)}{||Q_i - P_i||}$$

where $\{P_i\}$ is the set of closest points on the surface M_t to the torso points $\{Q_i\}$ on the sparse model, and J_2^3 is the TPS functional using derivatives of order 2 and image dimensions 3. The deformation $M_{t+1} = F(M_t)$ was iteratively applied in a diffeomorphic manner, resulting in a composition of several smooth approximations approaching initial mesh M towards the sparse representation. Laplacian smoothing, decimation, and affine transformation were applied in the end to ensure the local geometric and topological characteristics of the reconstructed mesh. Finally, the torso surface was remeshed with a restricted Frontal-Delaunay algorithm using the mesh generator JIGSAW [6] with a specified element size of 1cm.

1.3 Definition of anatomical parameters. The torso volume was calculated using PyVista [7], as the volume enclosed by the reconstructed torso surface. The cardiac volume was estimated from the extracted cardiac contours [8]. Volume of the cavity was obtained using Simpson's rule over the segmented endocardial surfaces – as described in Beetz et al. [9] this provides good estimates of cardiac volumes. Left ventricular (LV) mass was measured from the volume difference between LV epicardial and endocardial surfaces multiplied by 1.05 [10].

The x position (lateral) of the heart relative to the ECG electrodes was determined by setting the location of the most rightward electrode on the body as $x = 0$ and the most leftward as $x = 1$ and noting the position of the centre of the heart along this scale. y (posterior) and z (superior) positions were found similarly by setting the most frontward electrode as $y = 0$, backward as $y = 1$, downward as $z = 0$, and upward as $z = 1$.

For estimating the heart position, the intersection point of the LAX four-chamber plane, LAX two-chamber plane, and the middle slice of the SAX stack was identified. For estimating the heart orientation, the X-axis was first measured in the cardiac system as the intersecting line between the LAX four-chamber plane and the mid SAX slice, while the Z-axis was identified as the perpendicular line to the X-axis that passed through the LAX four-chamber plane. The Y-axis was estimated as normal to the XZ-plane. For calculating the Euler angles (α , β , γ), as shown in Figure 1A, the intersecting line, denoted as x' , between the XY-plane in the cardiac system and the xy-plane in the real (torso) coordinate system was first identified. α was measured as the angle between the x-axis in the real system and the intersecting line x' , essentially presenting the spin of the XY-plane around the z-axis. The γ angle was calculated as the angle between the X-axis in the cardiac system and the intersecting line, i.e., the tilt of the X-axis with respect to the xy-plane. β was measured as the angle between the Z-axis in the cardiac system and the z-axis in the real (torso) system, i.e., the verticality of the cardiac long axis. The calculations of the coordinate systems and angles are presented in the lower-left corner of Figure 1A.

1.4 Reconstruction validation. In order to evaluate the performance of the contour extraction procedure, reconstructions of 30 subjects made using the fully automated pipeline were compared with those made using manually annotated contours. A ray tracing method [11] was applied to obtain the distance between 3D torso surfaces. For each point on one surface, the normal to the surface was found, and the nearest intersection between that line and the second surface (in either direction) was obtained. Since the normal is not guaranteed to intersect the surface or may intersect at a point on the opposing side of the torso, the nearest neighboring point on the second surface was also found for each point. The minimum between these two distances was taken for each point as the surface-to-surface distance. The distance between each electrode location on the different torsos was also taken as a measure of similarity for the reconstructions. The mean surface-to-contour distance was calculated for each test subject to evaluate the quality of the reconstruction.

1.5 Statistical methods. *Choice of ECG parameters.* ECG parameters were chosen due to their availability in the UKB dataset (particularly their availability in all leads, which excludes parameters such as QT interval) and their relevance to diagnosis and risk stratification of myocardial infarction. A range of durations, amplitudes and axis angles were chosen in order to investigate the differing effect of anatomical parameters on each category.

Distribution of anatomical and ECG parameters for all subpopulations. Split violin plots were created to show the distribution of each anatomical and electrophysiological parameter separately for healthy males, healthy females, post-MI males and post-MI females. Outliers were removed, as defined by values that were more than three times the inter quartile range above the third quartile or similarly below the first quartile. Mean values of all biomarkers were compared between healthy and post-MI subjects of either sex, and male and female subjects of either MI status. The Shapiro-Wilk test was used to ascertain whether both subpopulations had a normal distribution. If there was significant evidence of non-normality, the Wilcoxon test was used. The Levene test was used to determine whether the subpopulations had an equal variance. If there was sufficient evidence of nonequal variance, the Welch's t-test was used, otherwise, the standard t-test was used. For populations with statistically significant differences between them, arrows were drawn from the lower mean population to the higher mean population with stars denoting the statistical significance of the relationship. * denotes relationships with p-values between 0.05 and 0.01, ** between 0.01 and 0.001, *** between 0.001 and 0.0001 and **** less than 0.0001. The mean and standard deviation of each anatomical and ECG biomarker was recorded, alongside the p-values for differences between subpopulations.

Age and body mass index (BMI) association. Each anatomical parameter was separately regressed against age for the healthy male, healthy female, post-MI male and post-MI female populations. This was repeated against BMI. This was recorded in a heat map and associated table, with the confidence intervals and two-sided p-values for the regression coefficient to be non-zero.

Normalised correlation coefficients. As the two measures of cardiac size (total cavity volume and left ventricular mass) were highly colinear, one was selected for each ECG biomarker depending on the minimisation of mean Akaike information criterion (AIC) across the 12 leads. For each lead, the ECG parameter was regressed against the chosen cardiac size parameter, torso volume, the relative heart centre positions in the x, y and z directions and the three cardiac orientation parameters (spin, verticality and tilt). In order to compare the regression coefficients for anatomical and ECG parameters on different scales, the regression coefficients were normalised by multiplying by the standard deviation of the anatomical parameter and then dividing by the standard deviation of the ECG biomarker, before being plotted. This was done separately for the healthy male, healthy female, post-MI male and post-MI female populations.

Sex differences. Separately for the post-MI and healthy populations, the difference in the mean ECG biomarker between the male and female cohorts was plotted for each lead, discarding any missing values. For each lead the ECG parameter was regressed against the chosen cardiac size parameter, torso volume, the relative heart centre positions in the x, y and z directions and the three cardiac orientation parameters (spin, verticality and tilt), and also the sex categorical parameter. For the continuous variables (all anatomical parameters), the individual raw regression coefficients were then multiplied by the difference in the mean value between the male and female populations to give their estimated contribution to the sex difference in the ECG parameter. The coefficient for sex is the remaining sex difference when all of the chosen anatomical parameters were controlled for, so is taken as an estimate for the electrophysiological contribution. This was done separately for the healthy and post-MI populations.

MI differences. Separately for the male and female cohorts, the difference in the mean ECG biomarker between the post-MI and healthy populations was plotted for each lead, discarding any missing values. For each lead, the ECG parameter was regressed against the chosen cardiac size parameter, torso volume, the relative heart centre positions in the x, y and z directions and the three cardiac orientation parameters (spin, verticality and tilt) and also the healthy vs post-MI categorical parameter. For the continuous variables (all anatomical parameters), the individual regression

coefficients were then multiplied by the difference in the mean value between the healthy and post-MI populations to give their estimated contribution to the difference in the ECG parameter between healthy and post-MI. The coefficient for MI status is the remaining healthy versus post-MI difference when all of the chosen anatomical parameters were controlled for, so is taken as an estimate for the electrophysiological contribution.


Supplementary Appendix 2: Results

2.1 Anatomy-QRS duration relationship. Mean AIC across leads, a measure of prediction error, was lower with cavity volume than LV mass (12311 and 12316 respectively). Therefore, cavity volume was chosen as the representative of cardiac size for QRS duration.


2.2 Anatomy-STj amplitude relationship. Mean AIC across leads was lower with LV mass than cavity volume (-4841 and -4834 respectively). Therefore, LV mass was chosen as the representative of cardiac size for STj amplitude.

2.3 Anatomy-TWA amplitude relationship. Mean AIC across leads was lower with cavity volume than LV mass (-965 and -963 respectively). Therefore, cavity volume was chosen as the representative of cardiac size for TWA.


2.4 Anatomy-axis angles relationship. Mean AIC across R and T axes was lower with cavity volume than LV mass (1459 and 14660 respectively). Therefore, cavity volume was chosen as the representative of cardiac size for both axis angles.




Supplementary Figure 1. Exclusion pipeline for reconstruction dataset. Both healthy and post-myocardial infarction (MI) subjects were required to have sufficient images across the view subtypes to reconstruct the torso and cardiac geometries. Subjects also were excluded if the ECG was missing or invalid (all healthy subjects had completed ECG). In the healthy dataset, subjects with at least 1 of 82 disease diagnoses were omitted, including circulatory system disorders, renal disease, genitourinary disorders, and endocrine disorders. The most common reason for exclusion was primary hypertension. Post-MI subjects were excluded if the recorded date of the MI event was after that of the imaging visit where the CMR images and ECG were taken.



Supplementary Figure 2. Depiction of the relative coordinate system in which the heart position was measured. A heart centre equidistant in the z direction to the highest (most superior) and lowest (most inferior) electrodes would have a relative z coordinate of 0.5.




Supplementary Figure 3. A-I Distributions of geometrical biomarkers. Horizontal lines show statistically significant differences between subpopulation means with arrows pointing to the larger mean. Heat map of correlation coefficients for each geometrical parameter with age (**J**) and BMI (**K**). Correlations with $p > 0.05$ are shaded grey. CVol: total cavity volume, LVm: left ventricular mass, TVol: torso volume, x: lateral heart centre position, y: posterior position, z: superior position, Vert: verticality of the cardiac long axis. HM: healthy male, HF: healthy female, PM: post-MI male, and PF: post-MI female.




Supplementary Figure 4. Distributions of ECG biomarkers for healthy and post-MI populations and proportions of subjects with AV block and pathological Q waves. Horizontal lines show statistically significant differences between subpopulation means with arrows pointing to the larger mean. Many distributions show features of non-normality, such as skew, heavy tails or bimodality. Post-MI subjects have longer PQ and QT intervals and P wave duration, and a lower T amplitude for both sexes. Post-MI males also have a slower heart rate, lower P amplitude, longer QRS duration, left-deviated R axis, and longer QTc. Post-MI subjects are substantially more likely to have AV block and pathological Q waves. The T axis shows a higher variability in post-MI individuals for both sexes. AV: atrioventricular, QTc: corrected QT interval.

Healthy




Post-MI




Supplementary Figure 5. Normalised regression coefficients with 95% confidence intervals for QRS duration in all ECG leads for healthy and post-MI subjects. CVol: total cavity volume, TVol: torso volume, x, y, z: x (medial), y (posterior), z (superior) coordinate of the heart centre relative to the electrodes, Vert: verticality of the cardiac long axis. Bars shown in red represent regression coefficients that are significantly different from 0, at a significance level of 0.05. QRS duration is shortened by a more vertical cardiac orientation (less negative β) for at least one sex in a majority of

leads. Increasing cavity volume prolongs the QRS duration. QRS duration is largely unaffected by adjustments in the cardiac location. Unlike healthy subjects, for post-MI individuals QRS duration is largely unaffected by cardiac orientation. QRS duration again usually does not significantly vary with cardiac location. In most leads increasing the cavity volume prolongs the QRS duration.




Supplementary Figure 6. A mean ST amplitudes measured at the junction (end of the QRS complex) across subjects for each ECG lead in post-MI subjects with females shown as pale red circles and males as cyan triangles. **B** contribution of each parameter to the difference in STj amplitude from females to males for post-MI subjects, calculated by multiplying the regression coefficient for each factor by its mean difference between male and female populations.




Supplementary Figure 7. Normalised regression coefficients with 95% confidence intervals for STj amplitude in all ECG leads for healthy subjects. LVm: left ventricular mass, TVol: torso volume, x, y, z: x (medial), y (posterior), z (superior) coordinate of the heart centre relative to the electrodes, Vert: verticality of the cardiac long axis. Bars shown in red represent regression coefficients that are significantly different from 0, at a significance level of 0.05. Significant positional correlations generally follow the pattern that the closer the heart centre was to the position of the exploring electrode, the

higher the STj amplitude. Increases in left ventricular mass are associated with increased STj amplitude in some of the precordial leads.



Supplementary Figure 8. Normalised regression coefficients with 95% confidence intervals for T wave amplitude in all ECG leads for healthy subjects. CVol: total cavity volume, TVol: torso volume, x, y, z: x (medial), y (posterior), z (superior) coordinate of the heart centre relative to the electrodes, Vert: verticality of the cardiac long axis. Bars shown in red represent regression coefficients that are significantly different from 0, at a significance level of 0.05. T wave amplitude in many leads increases as the heart is oriented more vertically (less negative β). Significant positional correlations generally

follow the pattern that the closer the heart centre was to the position of the exploring electrode, the higher the T wave amplitude.



Supplementary Figure 9. Normalised regression coefficients with 95% confidence intervals for R axis (top) and T axis (bottom), in healthy (left) and post-MI (right) subjects. For healthy but not post-MI subjects, R axis is negatively associated with torso volume. R axis is significantly affected by cardiac orientation, but not T axis. CVol: total cavity volume, TVol: torso volume, x, y, z: x (medial), y (posterior), z (superior) coordinate of the heart centre relative to the electrodes, Vert: verticality of the cardiac long axis. Bars shown in red represent regression coefficients significantly different from 0, at a significance level of 0.05.

Supplementary Table 1. Mean and standard deviation values of anatomical and ECG biomarkers in each subpopulation.

Anatomical biomarkers								
Biomarker	Healthy			MI			Male	Female
	Male N=470	Female N=581	p-value	Male N=341	Female N=84	p-value	p-value	p-value
Total cavity volume (ml)	326 (62)	249 (42)	< 0.0001	320 (63)	245 (47)	< 0.0001	0.09	0.4
LV mass (g)	94.3 (17.0)	66.7 (11.0)	< 0.0001	97.3 (19.3)	70.9 (13.3)	< 0.0001	0.06	0.003
Torso volume (dm ³)	43.4 (7.5)	36.5 (7.6)	< 0.0001	47.0 (7.7)	40.3 (8.0)	< 0.0001	< 0.0001	< 0.0001
Lateral (x) position	0.642 (0.034)	0.643 (0.036)	0.7	0.635 (0.035)	0.634 (0.041)	0.8	0.002	0.02
Posterior (y) position	0.727 (0.069)	0.743 (0.071)	0.0006	0.736 (0.070)	0.770 (0.069)	< 0.0001	0.06	0.0004
Superior (z) position	0.544 (0.037)	0.569 (0.033)	< 0.0001	0.549 (0.036)	0.565 (0.034)	0.0002	0.09	0.2
Spin α (°)	41.4 (7.6)	44.0 (8.4)	< 0.0001	43.0 (8.4)	44.8 (9.9)	0.1	0.005	0.5
Verticality β (°)	-63.4 (7.9)	-59.5 (9.0)	< 0.0001	-67.7 (7.6)	-65.4 (10.0)	0.05	< 0.0001	< 0.0001
Tilt γ (°)	-21.2 (7.5)	-23.6 (8.4)	< 0.0001	-19.2 (7.1)	-21.5 (9.1)	0.03	0.0002	0.02
ECG biomarkers								
Biomarker	Healthy	MI	Male	Female				
	Male N=470	Female N=581	p-value	Male N=341	Female N=84	p-value	p-value	p-value
Heart rate (l) (bpm)	59.9 (10.4)	62.4 (10.7)	< 0.0001	58.5 (11.2)	61.2 (10.7)	0.02	0.007	0.1
P duration (l) (ms)	98.7 (15.9)	95.1 (14.7)	< 0.0001	100.3 (19.1)	98.9 (14.8)	0.2	0.04	0.03
Mean P amplitude (mV)	0.0755 (0.0260)	0.0747 (0.0384)	0.3	0.0707 (0.0186)	0.0708 (0.0170)	0.8	0.005	0.3
P axis (°)	50.9 (22.1)	49.6 (22.4)	0.5	48.6 (25.6)	49.1 (21.6)	0.9	0.2	0.6
PQ interval (l) (ms)	165 (25)	158 (23)	< 0.0001	178 (32)	164 (28)	0.001	< 0.0001	0.02
Mean QRS duration (ms)	85.8 (10.8)	79.8 (9.1)	< 0.0001	90.7 (15.1)	82.5 (14.5)	< 0.0001	< 0.0001	0.1
R axis (°)	27.6 (35.9)	31.2 (34.9)	0.1	15.6 (41.5)	25.8 (34.1)	0.003	< 0.0001	0.3
QT interval (l) (ms)	415 (31)	421 (33)	0.003	428 (31)	429 (32)	0.8	< 0.0001	0.02
QTc (l) (ms)	411 (24)	426 (29)	< 0.0001	418 (26)	429 (27)	0.002	0.0009	0.6
Mean STj amplitude (mV)	0.0106 (0.0225)	-0.0008 (0.0190)	< 0.0001	0.0088 (0.0231)	-0.0002 (0.0176)	0.002	0.09	0.8
Mean T amplitude (mV)	0.248 (0.109)	0.160 (0.084)	< 0.0001	0.189 (0.108)	0.144 (0.077)	< 0.0001	< 0.0001	0.03
T axis (°)	38.9 (30.4)	41.0 (31.6)	0.08	39.2 (43.8)	45.6 (37.6)	0.06	0.7	0.1

p-values are measured based on appropriate two-sided test (as discussed in statistical methods) for: first two columns, the mean male and female values being different and last two columns, the mean healthy and post-MI values being different. p-values are shaded red if below 0.05, i.e., statistically significant. LV: left ventricular. Post-MI hearts were found to be placed in a more medial position (lower x) and less vertical orientation (more negative β) than those of healthy subjects. For females the cardiac position was also found to be more posterior (larger y) in post-MI than healthy subjects. Torso volume was considerably larger in post-MI subjects. Females had a more posterior (larger y) and superior (larger z) cardiac position than males and their orientation was more vertical (less negative β). Cavity volume, LV mass and torso volume were substantially lower in females than males. Post-MI subjects have longer P durations and PQ and QT intervals, and a lower T amplitude for both sexes. Post-MI males also have a slower heart rate, lower P amplitude, longer QRS duration, left-deviated R axis, and longer QTc. AV: atrioventricular, QTc: corrected QT interval.

Supplementary Table 2. Associations between age and BMI and anatomical biomarkers.

Biomarker	Age			
	Healthy		MI	
	Male N=470	Female N=581	Male N=341	Female N=84
Total cavity volume (ml)	-0.277[-0.364, -0.189], p < 0.0001	-0.251[-0.330, -0.172], p < 0.0001	-0.172[-0.277, -0.067], p = 0.001	-0.042[-0.261, 0.178], p = 0.7
LV mass (g)	-0.238[-0.326, -0.150], p < 0.0001	-0.107[-0.188, -0.026], p = 0.01	-0.117[-0.223, -0.011], p = 0.03	-0.064[-0.283, 0.155], p = 0.6
Torso volume (dm ³)	-0.021[-0.111, 0.070], p = 0.7	-0.014[-0.096, 0.068], p = 0.7	-0.024[-0.131, 0.083], p = 0.7	-0.195[-0.410, 0.020], p = 0.08
Lateral (x) position	0.062[-0.029, 0.153], p = 0.2	0.032[-0.050, 0.113], p = 0.4	-0.056[-0.163, 0.051], p = 0.3	0.039[-0.181, 0.258], p = 0.7
Posterior (y) position	0.063[-0.027, 0.154], p = 0.2	0.039[-0.042, 0.121], p = 0.3	-0.041[-0.148, 0.066], p = 0.4	-0.201[-0.416, 0.014], p = 0.07
Superior (z) position	-0.070[-0.161, 0.021], p = 0.1	-0.127[-0.208, -0.046], p = 0.002	-0.123[-0.229, -0.017], p = 0.02	0.057[-0.162, 0.276], p = 0.6
Spin α (°)	-0.089[-0.180, 0.001], p = 0.05	0.027[-0.054, 0.109], p = 0.5	-0.103[-0.210, 0.003], p = 0.06	0.060[-0.160, 0.279], p = 0.6
Verticality β (°)	-0.113[-0.203, -0.022], p = 0.01	-0.218[-0.298, -0.139], p < 0.0001	-0.100[-0.206, 0.007], p = 0.07	-0.215[-0.429, 0.000], p = 0.05
Tilt γ (°)	0.068[-0.023, 0.158], p = 0.1	0.088[0.007, 0.170], p = 0.03	0.052[-0.055, 0.158], p = 0.3	-0.080[-0.299, 0.139], p = 0.5
BMI				
Biomarker	Healthy		MI	
	Male N=470	Female N=581	Male N=341	Female N=84
Total cavity volume (ml)	0.000[-0.093, 0.093], p = 1	0.065[-0.017, 0.148], p = 0.1	-0.047[-0.155, 0.062], p = 0.4	0.028[-0.203, 0.259], p = 0.8
LV mass (g)	0.277[0.188, 0.367], p < 0.0001	0.247[0.167, 0.327], p < 0.0001	0.235[0.129, 0.340], p < 0.0001	0.198[-0.029, 0.425], p = 0.09
Torso volume (dm ³)	0.878[0.846, 0.909], p < 0.0001	0.924[0.894, 0.955], p < 0.0001	0.922[0.879, 0.965], p < 0.0001	0.936[0.859, 1.012], p < 0.0001
Lateral (x) position	-0.425[-0.509, -0.341], p < 0.0001	-0.478[-0.551, -0.405], p < 0.0001	-0.486[-0.582, -0.391], p < 0.0001	-0.544[-0.738, -0.349], p < 0.0001
Posterior (y) position	0.440[0.358, 0.522], p < 0.0001	0.479[0.406, 0.552], p < 0.0001	0.320[0.217, 0.423], p < 0.0001	0.431[0.230, 0.631], p < 0.0001
Superior (z) position	0.284[0.196, 0.372], p < 0.0001	0.303[0.225, 0.381], p < 0.0001	0.228[0.123, 0.333], p < 0.0001	0.075[-0.146, 0.297], p = 0.5
Spin α (°)	0.032[-0.060, 0.124], p = 0.5	-0.039[-0.121, 0.044], p = 0.4	-0.003[-0.111, 0.106], p = 1	0.161[-0.065, 0.387], p = 0.2
Verticality β (°)	-0.538[-0.616, -0.461], p < 0.0001	-0.533[-0.604, -0.463], p < 0.0001	-0.450[-0.547, -0.354], p < 0.0001	-0.423[-0.630, -0.215], p < 0.0001
Tilt γ (°)	0.389[0.302, 0.475], p < 0.0001	0.379[0.303, 0.455], p < 0.0001	0.351[0.249, 0.452], p < 0.0001	0.214[-0.010, 0.437], p = 0.06

The values represent regression coefficient [95% confidence interval] (two-sided p-value) of each parameter with age. Regression coefficients are normalised by the standard deviation of both the dependent and independent variables. Cells are shaded red if the p-value is below 0.05, i.e., statistically significant. LV: left ventricular. Age primarily affects the size of the heart, with increasing age reducing both cavity volume and LV mass in most subpopulations. Increasing age additionally moves the heart to a more inferior (lower z) position and makes the orientation less vertical (more negative β) in some subpopulations. Most geometrical parameters are more strongly associated with BMI than age. As BMI increases, the heart centre moves to a more medial (lower x), posterior (higher

y), and superior (higher z) position. The cardiac orientation also becomes more horizontal (more negative β) with increasing BMI. BMI does not significantly affect the cavity volume, but does make the LV mass larger, suggesting that the heart wall becomes thicker.

Supplementary References

1. Littlejohns, T.J., Sudlow, C., Allen, N.E., and Collins, R.: 'UK Biobank: opportunities for cardiovascular research', *European Heart Journal*, 2019, 40, (14), pp. 1158-1166
2. Robinette, K.M., Daanen, H., and Paquet, E.: 'The CAESAR project: a 3-D surface anthropometry survey', in Editor (Ed.)^(Eds.): 'Book The CAESAR project: a 3-D surface anthropometry survey' (1999, edn.), pp. 380-386
3. Pishchulin, L., Wuhrer, S., Helten, T., Theobalt, C., and Schiele, B.: 'Building statistical shape spaces for 3D human modeling', *Pattern Recognition*, 2017, 67, pp. 276-286
4. Rohr, K., Stiehl, H.S., Sprengel, R., Buzug, T.M., Weese, J., and Kuhn, M.H.: 'Landmark-based elastic registration using approximating thin-plate splines', *IEEE Transactions on Medical Imaging*, 2001, 20, (6), pp. 526-534
5. Villard, B., Grau, V., and Zacur, E.: 'Surface Mesh Reconstruction from Cardiac MRI Contours', in Editor (Ed.)^(Eds.): 'Book Surface Mesh Reconstruction from Cardiac MRI Contours' (2018, edn.), pp.
6. Engwirda, D.: 'Generalised primal-dual grids for unstructured co-volume schemes', *Journal of Computational Physics*, 2018, 375, pp. 155-176
7. Kaszynski, B.S.a.A.: 'PyVista: 3D plotting and mesh analysis through a streamlined interface for the Visualization Toolkit (VTK)', *Journal of Open Source Software*, 2019, 4, (37), pp. 1450
8. Bai, W., Sinclair, M., Tarroni, G., Oktay, O., Rajchl, M., Vaillant, G., Lee, A.M., Aung, N., Lukaschuk, E., Sanghvi, M.M., Zemrak, F., Fung, K., Paiva, J.M., Carapella, V., Kim, Y.J., Suzuki, H., Kainz, B., Matthews, P.M., Petersen, S.E., Piechnik, S.K., Neubauer, S., Glocker, B., and Rueckert, D.: 'Automated cardiovascular magnetic resonance image analysis with fully convolutional networks', *Journal of Cardiovascular Magnetic Resonance*, 2018, 20, (1), pp. 65
9. Beetz, M., Banerjee, A., Ossenberg-Engels, J., and Grau, V.: 'Multi-class point cloud completion networks for 3D cardiac anatomy reconstruction from cine magnetic resonance images', *Medical Image Analysis*, 2023, 90, pp. 102975
10. Shapiro, E.P., Rogers, W.J., Beyar, R., Soulen, R.L., Zerhouni, E.A., Lima, J.A., and Weiss, J.L.: 'Determination of left ventricular mass by magnetic resonance imaging in hearts deformed by acute infarction', *Circulation*, 1989, 79, (3), pp. 706-711
11. <https://docs.pyvista.org/examples/01-filter/distance-between-surfaces.html#distance-between-two-surfaces>, accessed 31 January 2023