
ADAPTIVE EXPERIMENTAL DESIGN FOR POLICY LEARNING

A PREPRINT

Masahiro Kato¹, Kyohei Okumura², Takuya Ishihara³, and Toru Kitagawa⁴

¹Data Analytics Team, Mizuho–DL Financial Technology, Co., Ltd.

²Department of Economics, Northwestern University

³Graduate School of Economics and Management, Tohoku University

⁴Department of Economics, Brown University

February 9, 2024

ABSTRACT

Evidence-based targeting has been a topic of growing interest among the practitioners of policy and business. Formulating decision-maker’s policy learning as a fixed-budget best arm identification (BAI) problem with contextual information, we study an optimal adaptive experimental design for policy learning with multiple treatment arms. In the sampling stage, the planner assigns treatment arms adaptively over sequentially arriving experimental units upon observing their contextual information (covariates). After the experiment, the planner recommends an individualized assignment rule to the population. Setting the worst-case *expected regret* as the performance criterion of adaptive sampling and recommended policies, we derive its asymptotic lower bounds, and propose a strategy, *Adaptive Sampling-Policy Learning strategy* (PLAS), whose leading factor of the regret upper bound aligns with the lower bound as the size of experimental units increases.

1 Introduction

This study designs an adaptive experiment for decision-making given multiple treatment arms, such as arms in slot machines, diverse therapies, and distinct unemployment assistance programs. The primary objective is to identify the best treatment arm for individuals given covariates, often referred to as a context, at the end of an experiment. Our problem is termed contextual fixed-budget best arm identification (BAI), an instance of the stochastic multi-armed bandit (MAB) problem (Thompson, 1933; Lai and Robbins, 1985). Our setting is a generalization of the *fixed-budget BAI* problem to minimize the expected simple regret at the end of a fixed number of rounds of an adaptive experiment, called a *budget* or *sample size* (Bubeck et al., 2009, 2011; Audibert et al., 2010).

In our setting, at each round of an adaptive experiment, a decision-maker sequentially assigns one of the treatment arms to a research subject based on past observations and contextual information observed before the treatment assignment. At the end of the experiment, the experimenter recommends an estimated best treatment arm for future experimental subjects. We measure the performance of a strategy of the decision-maker using the expected simple regret, which is a difference between the maximum expected outcome that could be achieved given full knowledge of distributions of the treatment arms and an expected outcome of a recommended treatment arm by the strategy.

The decision-maker follows a strategy which consists of a sampling rule and a recommendation rule. Specifically, the decision-maker assigns units following the sampling rule and recommends a treatment arm following the recommendation rule. A strategy of the decision-maker effectiveness is evaluated using the expected simple regret, which is a difference between the maximum context-conditional expected outcome that could be achieved given full knowledge of distributions of the treatment arms and a context-conditional expected outcome of a treatment arm estimated as the best treatment arm by the strategy. Our goal is to develop a strategy that minimizes the expected simple regret.

The focus of BAI is on an optimal strategy consisting of a sampling and recommendation rule. While the sampling rule dictates treatment arm assignment during each adaptive experiment round, the recommendation rule determines which treatment arm to be recommended at the end of the experiment. Typically, we first develop a lower bound (theoretical

limit) for the expected simple regret. Subsequently, we evaluate the upper bound (performance) of our suggested strategy to discuss its optimality.

The challenge in this issue arises from the need for context-specific recommendations. Unlike prior studies that did not consider contextual information, developing a model capturing the relationship between context and outcomes becomes imperative. To address this issue, we define a function suggesting a treatment arm given a context as a *policy*. In our approach, the recommendation rule of a strategy employs a policy trained by observations obtained from an adaptive experiment.

Specifically, we propose a strategy that assigns treatment arms following the Adaptive Sampling (AS) and recommends a treatment arm using Policy Learning (PL). In the AS rule, a decision-maker assigns a treatment arm to an experimental unit following a probability depending on variances of experiment units' outcomes. Because the variances are unknown, the decision-maker estimates them during the experiment and continues updating the assignment probability. At the end of the experiment, the decision-maker trains a policy using observations obtained in the experiment and recommends a treatment arm using the trained policy. We refer to our strategy as the PLAS strategy.

Given the inherent uncertainties, we use the minimax criterion for performance assessment, which evaluates the worst-case scenario among a set of bandit models (Bubeck et al., 2011). Our research identifies the leading factor in the lower bound as the variances of potential outcomes, also providing a variance-dependent sampling rule. We subsequently show that the Adaptive Sampling (AS)-Policy Learning (PL) strategy is asymptotically minimax optimal, as its foremost factor of the worst-case expected simple regret aligns with the lower bound.

In summary, our contributions include: (i) a lower bound for the worst-case expected simple regret; (ii) the PLAS strategy with a closed-form target assignment ratio, characterized by the variances of outcomes, (iii) an asymptotic minimax optimality of the PLAS strategy, . These findings contribute to a variety of subjects, including statistical decision theory and causal inference, in addition to BAI.

1.1 Related Work

This section provides a literature review of related work.

BAI and ordinal optimization. The MAB problem has been explored as an instance of the sequential decision-making problem (Thompson, 1933; Robbins, 1952; Lai and Robbins, 1985), where BAI is a paradigm within this context (Even-Dar et al., 2006; Audibert et al., 2010; Bubeck et al., 2011). BAI research can be traced back to sequential testing, ranking, and selection problems in the 1940s (Wald, 1947; Bechhofer et al., 1968).

Subsequent studies in operations research, particularly in the realm of ordinal optimization, have garnered considerable attention (Chen et al., 2000; Glynn and Juneja, 2004). These studies predominantly focus on devising optimal strategies under the assumption of known distributional parameters.

The machine learning community has reframed the problem of sequential decision-making as the BAI problem, placing a specific emphasis on estimating unknown distributions (Jennison et al., 1982; Even-Dar et al., 2006; Audibert et al., 2010; Bubeck et al., 2011). Audibert et al. (2010) propose the UCB-E and Successive Rejects (SR) strategies. Bubeck et al. (2011) demonstrates minimax optimal strategies for expected simple regret in a non-asymptotic setting by extending the minimax lower bound of Auer et al. (2002).

Carpentier and Locatelli (2016) further enhances the minimax lower bound, showing the optimality of Audibert et al. (2010)'s methods in terms of leading factors in expected simple regret. Based on their lower bound, Yang and Tan (2022) proposes minimax optimal linear fixed-budget BAI.

In addition to minimax evaluation, Komiyama et al. (2023) develop an optimal strategy whose upper bound for simple Bayesian regret lower bound aligns with their derived lower bound. Atsidakou et al. (2023) propose a Bayes optimal strategy for minimizing the probability of misidentification, revealing a surprising result that a $1/\sqrt{T}$ -factor dominates the evaluation.

Russo (2016), Qin et al. (2017), and Shang et al. (2020) propose Bayesian BAI strategies that are optimal in terms of posterior convergence rate. However, Kasy and Sautmann (2021) and Ariu et al. (2021) discuss that such optimality does not necessarily guarantee asymptotic optimality for the probability of misidentification in fixed-budget BAI.

In contrast to the approaches of Bubeck et al. (2011) and Carpentier and Locatelli (2016), regarding asymptotic optimality, Kaufmann et al. (2016) derives distribution-dependent lower bounds for BAI with fixed confidence and a fixed budget, based on change-of-measure arguments and building upon the work of Lai and Robbins (1985). Following their work, Garivier and Kaufmann (2016) proposes an optimal strategy for BAI with fixed confidence; however, in the fixed-budget setting, there is currently a lack of strategies whose upper bound matches the lower bound established by

Kaufmann et al. (2016). This issue has been discussed by Kaufmann (2020), Ariu et al. (2021), Qin (2022), Degenne (2023), Kato (2023b), and Wang et al. (2023).

The problem of BAI with contextual information is still under investigation. For example, Tekin and van der Schaar (2015), Guan and Jiang (2018), Deshmukh et al. (2018), Kato and Ariu (2021), and Qin and Russo (2022) consider this problem, but their analyses and settings differ from those employed in this study.

Our proposed strategy assigns treatment arms with a probability depending on their variances. Variance-dependent BAI has been explored by Chen et al. (2000), Glynn and Juneja (2004), Kaufmann et al. (2016), Sauro (2020), Jourdan et al. (2023), Kato et al. (2023), Kato (2023b,a), and Lalitha et al. (2023). Our choice of treatment-assignment probability is also inspired by van der Laan (2008) and Hahn et al. (2011) in adaptive experimental design for efficient treatment effect estimation.

Decision theory and treatment choice. Beyond BAI, our study is further related to statistical decision theory (Wald, 1949). Manski (2000, 2002, 2004) extend this decision theory and introduce the treatment choice problem from a decision theory perspective, independent of BAI. They focus on recommending the best treatment arm using non-experimental, independently and identically distributed (i.i.d.) observations without adaptive experimental design (Schlag, 2007; Stoye, 2009, 2012; Manski and Tetenov, 2016; Dominiczak and Manski, 2017, 2022). Hirano and Porter (2009) employs the limit experiment framework (Le Cam, 1972, 1986; Lehmann and Casella, 1998; van der Vaart, 1991, 1998) for discussing the problem of treatment choice, where the class of alternative hypotheses comprises local models, with parameters of interest converging to the true parameters at a rate of $1/\sqrt{T}$. Armstrong (2022) and Hirano and Porter (2023) apply this framework to adaptive experimental design. Adusumilli (2021, 2022) present an alternative minimax evaluation of bandit strategies for both regret minimization and BAI, based on a formulation utilizing a diffusion process proposed by Wager and Xu (2021) and the limit experiment framework (Le Cam, 1960, 1972, 1986; van der Vaart, 1991, 1998).

Policy learning. Inspired by supervised learning and statistical decision theory, various off-policy learning methods have been proposed. Swaminathan and Joachims (2015) proposes counterfactual risk minimization, an extension of empirical risk minimization for policy learning. Kitagawa and Tetenov (2018) extends counterfactual risk minimization by linking it to the viewpoint of treatment choice and proposes welfare maximization. Athey and Wager (2021) refines the method proposed by Kitagawa and Tetenov (2018). Zhou et al. (2023) shows a tight upper bound for general policy learning methods. Zhan et al. (2022) develops a policy learning method from adaptively collected observations.

For the expected simple regret, Bubeck et al. (2011) shows that the Uniform-Empirical Best Arm (EBA) strategy is minimax optimal for bandit models with bounded supports. Kock et al. (2023) extends the results to cases where parameters of interest are functionals of the distribution and finds that optimal sampling rules are not uniform. Adusumilli (2022, 2021) consider a different minimax evaluation of bandit strategies for both regret minimization and BAI problems, based on a formulation using a diffusion process, as proposed by Wager and Xu (2021).

Other related work. Efficient estimation of ATE via adaptive experiments constitutes another area of related literature. van der Laan (2008) and Hahn et al. (2011) propose experimental design methods to estimate ATE more efficiently by using covariate information in treatment assignments. Karlan and Wood (2014) examine donors' responses to new information by applying the method of Hahn et al. (2011). Subsequently, Tabord-Meehan (2022) and Kato et al. (2020) attempt to improve these studies, and more recently, Gupta et al. (2021) proposes the use of instrumental variables in this context. Viviano (2022) explores experimental designs for network inference.

Kock et al. (2023) generalizes the results of Bubeck et al. (2011) for the case where the parameter of interest is a functional of the distribution and finds that, in contrast to the results Bubeck et al. (2011), the target allocation ratio is not uniform.

We employ the Augmented Inverse Probability Weighting (AIPW) estimator in policy learning. The AIPW estimator has been extensively used in the fields of causal inference and semiparametric inference (Tsiatis, 2007; Bang and Robins, 2005; Chernozhukov et al., 2018). More recently, it has also been utilized in other MAB problems, as seen in Kim et al. (2021), Ito et al. (2022), Zimmert and Seldin (2021), and Masoudian and Seldin (2021).

1.2 Organization

The structure of this paper is as follows: Section 2 defines our problem. Section 3 develops lower bounds for the worst-case expected simple regret and the target assignment ratio. Section 4 introduces the PLAS strategy, and Section 5 presents upper bounds for the proposed strategy and its asymptotic minimax optimality.

2 Problem Setting

This study considers an adaptive experiment with a fixed budget (sample size) $T \in \mathbb{N}$, a set of treatment arms $[K] := \{1, 2, \dots, K\}$, and a decision-maker who aims to identify the best treatment arm. In each round $t \in [T]$, experimental units sequentially visit, and the decision-maker can assign treatment arms to them. At the end of the experiment, the decision-maker recommends an estimated best treatment arm. We refer to this problem as *fixed-budget BAI with contextual information*.

2.1 Potential Outcomes

Following the Neyman-Rubin causal model (Neyman, 1923; Rubin, 1974), let $Y^a \in \mathcal{Y}$ be a potential random outcome of treatment arm $a \in [K]$, where $\mathcal{Y} \subset \mathbb{R}$ denotes a set of possible outcomes. We also define $X \in \mathcal{X}$ as a context, also called covariates, that characterizes an experimental unit, where $\mathcal{X} \subset \mathbb{R}^d$ denote a set of possible d -dimensional covariates. We define a tuple (Y^1, \dots, Y^K, X) as a $(\mathcal{Y}^K \times \mathcal{X})$ -valued random variable following a probability distribution $P \in \mathcal{M}(\mathcal{Y}^K \times \mathcal{X})$, where $\mathcal{M}(\mathcal{T})$ denotes the set of all Borel probability measures on a topological space \mathcal{T} . In our setting, a distribution P of (Y^1, \dots, Y^K, X) belongs to a bandit model \mathcal{P} defined below. We refer to the bandit model as a location-shift bandit model. Let \mathbb{E}_P and Var_P be the expectation and variance operator under P , respectively. For each $P \in \mathcal{P}$, let us denote the first and second moment of the potential outcome as $\mu^a(P)(x) := \mathbb{E}_P[Y^a | X = x]$ and $\nu^a(P)(x) := \mathbb{E}_P[(Y^a)^2 | X = x]$, respectively.

Definition 2.1 (Location-shift bandit model). *Let $(\sigma^a)^2: \mathcal{X} \rightarrow [\underline{C}, \bar{C}]$ be a function that is exogenously given but unknown to the decision-maker, where \underline{C} and \bar{C} are universal constants such that $0 < \underline{C} < \bar{C} < \infty$ and known to the decision-maker. Let ζ be a distribution of X that is exogenously given but unknown to the decision-maker, which is supported on \mathcal{X} . Then, a location-shift bandit model \mathcal{P} is defined as follows:*

$$\mathcal{P} := \mathcal{P}_\zeta := \left\{ P \in \mathcal{M}(\mathcal{Y}^K \times \mathcal{X}) : \forall a \in [K] \quad \forall x \in \mathcal{X}, \quad \text{Var}_P(Y^a | X = x) = \sigma^a(x), \right. \\ \left. |\mu^a(P)(x)| < \bar{C}, \quad |\nu^a(P)(x)| < \bar{C}, \quad \text{marg}_{\mathcal{X}}(P) = \zeta \right\},$$

where $\text{marg}_{\mathcal{X}}(P)$ denotes the marginal distribution of P on \mathcal{X}

Our lower and upper bounds are characterized by $\sigma^a(x)$, given independently of the experiment. Location-shift models are a commonly employed assumption in statistical analysis (Lehmann and Casella, 1998). A key example is a Gaussian distribution, where for all P , the variances are fixed and only mean parameters shift. Note that we can omit the condition $|\nu^a(P)(x)| < \bar{C}$ from the boundedness of the variance. However, we introduce it to simplify the definition of our strategies.

2.2 Adaptive Experiment

We consider a decision-maker who aims to identify the best treatment arm

$$a^*(P)(x) \in \arg \max_{a \in [K]} \mathbb{E}_P[Y^a | X = x]$$

for each context $x \in \mathcal{X}$ through an adaptive experiment. A fixed number of rounds T , called the budget or sample size, is exogenously given. At each round $t \in [T]$, the decision-maker uses the following procedure:

1. A potential outcome $(Y_t^1, Y_t^2, \dots, Y_t^K, X_t)$ is drawn from P .
2. The decision-maker observes a context X_t .
3. The decision-maker assigns treatment arm A_t to an experimental unit based on past observations $(X_s, A_s, Y_s)_{s=1}^{t-1}$ and the context X_t .
4. The decision-maker observes an outcome of the assigned treatment arm, $Y_t = \sum_{a \in [K]} \mathbb{1}[A_t = a] Y_t^a$.

This process is referred to as the *exploration phase*. Note that outcomes from unassigned treatment arms remain unobserved. This setting is called the bandit feedback or Rubin causal model (Neyman, 1923; Rubin, 1974). Through the exploration phase, the decision-maker obtains observations $(Y_t, A_t, X_t)_{t=1}^T$. After round T , the decision-maker recommends an estimated best treatment arm $\hat{a}_T(x)$ for each context x given the observations $(X_t, A_t, Y_t)_{t=1}^T$.

2.3 Strategy with Policy Learning

A strategy of the decision-maker defines which treatment arm to assign in each period during the exploration phase and which treatment arm to recommend as an estimated best arm for each context. More precisely, the strategy consists of a pair $((A_t)_{t \in [T]}, \hat{a}_T)$ in which:

- the *sampling rule* determines which treatment arm A_t to be assigned in round t based on the past observations. In other words, A_t is \mathcal{G}_t -measurable, where $\mathcal{G}_t := \sigma(X_1, A_1, Y_1, \dots, X_{t-1}, A_{t-1}, Y_{t-1}, X_t)$ for each $t \in [T]$.
- the *recommendation rule* returns an estimated best treatment arm for each context \hat{a}_T based on the observations collected during the exploration phase. In other words, for each $x \in \mathcal{X}$, $\hat{a}_T(x)$ is \mathcal{F}_T -measurable, where $\mathcal{F}_T := \sigma(X_1, A_1, Y_1, \dots, X_T, A_T, Y_T)$.

Policy. In this study, we impose a restriction on a class of recommendation rules that the decision-maker can use. We assume that there is an *exogenously* given class of policies Π , whose typical element is a policy $\pi: [K] \times \mathcal{X} \rightarrow [0, 1]$ that is measurable and satisfies $\sum_a \pi(a, x) = 1$ for each x . Here, $\pi(a, x)$ denotes a probability that the decision-maker recommends treatment arm $a \in [K]$ as an estimated best treatment arm for context x . With a slight abuse of notation, we write $\pi(a | x)$ instead of $\pi(a, x)$. We require the decision-maker to obtain an estimator $\hat{a}_T(x)$ of $a^*(P)(x)$ as follows: first, the decision-maker constructs a policy $\pi \in \Pi$ based on the observations collected during the exploration phase; then, $\hat{a}_T(x)$ is drawn from $\pi(\cdot | x)$ for each x .

Optimal policy. We evaluate the performance of policies via a simple regret. The *value* of policy $\pi \in \Pi$ under P is the expected outcome when the decision-maker uses a policy π , which is defined as

$$Q(P)(\pi) := \mathbb{E}_P \left[\sum_{a \in [K]} \pi(a | X) \mu^a(P)(X); \pi \right],$$

and the optimal policy within class Π is defined accordingly:

$$\pi^*(P) := \arg \max_{\pi \in \Pi} Q(P)(\pi),$$

Then, given a strategy with a policy $\hat{\pi} \in \Pi$ of the decision-maker, we define a simple regret for each context $x \in \mathcal{X}$ under $P \in \mathcal{P}$ is defined as

$$r_T(P)(\hat{\pi})(x) := \sum_{a \in [K]} \pi^*(P)(a | x) \mu^a(P)(x) - \mu^{\hat{a}_T(x)}(P)(x),$$

and the marginalized simple regret is defined as

$$R_T(P)(\hat{\pi}) := \mathbb{E}_{X \sim \zeta} [r_T(P)(\hat{\pi})(X)] = Q(P)(\pi^*(P)) - Q(P)(\hat{\pi}).$$

Then, we define the *expected simple regret* as

$$\mathbb{E}_P [R_T(P)(\hat{\pi})],$$

where the expectation is taken over $\hat{\pi}$. This expected simple regret is our performance measure of interest. We also refer to the expected simple regret $\mathbb{E}_P [R_T(P)(\hat{\pi})]$ as the *policy regret*. The decision-maker aims to identify the best treatment arm with a smaller expected simple regret.

3 Lower Bounds for the Worst-case Expected Simple Regret

In this section, we derive lower bounds for the worst-case expected simple regret.

3.1 Restrictions to Strategies

To derive lower bounds, we often utilize an alternative hypothesis. We consider a bandit model whose expected outcomes are the same among the K treatment arms. We refer to it as the null bandit model.

Definition 3.1 (Null bandit models). *A bandit model $P \in \mathcal{P}$ is called a null bandit model if the expected outcomes are equal: $\mu^1(P)(x) = \mu^2(P)(x) = \dots = \mu^K(P)(x)$ for any $x \in \mathcal{X}$.*

We consider a restricted class of strategies such that under null bandit models, any strategy in this class recommends one of the arms with an equal probability ($1/K$) in the limit.

Definition 3.2 (Null consistent strategy). *We say a strategy is null consistent if for any null bandit model $P \in \mathcal{P}$, we have*

$$\left| \mathbb{P}_P(\hat{a}_T(X) = a \mid X = x) - \mathbb{P}_P(\hat{a}_T(X) = b \mid X = x) \right| \rightarrow 0 \quad (T \rightarrow \infty).$$

for all $x \in \mathcal{X}$ and for any $a, b \in [K]$.

Under null consistent strategies, $|\mathbb{P}_P(\hat{a}_T(X) = a \mid X = x) - 1/K| = o(1)$ holds for each $a \in [K]$ as $T \rightarrow \infty$.

3.2 Lower Bounds for Location-Shift Bandit Models

Next, we introduce the Natarajan dimension, a metric that measures the complexity of a policy class Π (Natarajan, 1989). Our lower bounds are characterized by the Natarajan dimension.

Definition 3.3 (Natarajan dimension). *We say that Π shatters M points $\{s_1, s_2, \dots, s_M\} \subseteq \mathcal{X}$ if there exist $f_1, f_{-1} : \{s_1, s_2, \dots, s_M\} \rightarrow [K]$ such that*

1. for any $j \in [M]$, $f_{-1}(s_j) \neq f_1(s_j)$ holds;
2. for any $\sigma := \{\sigma_1, \sigma_2, \dots, \sigma_M\} \in \{\pm 1\}^M$, there exists a policy $\pi \in \Pi$ such that for any $j \in [M]$,

$$\pi(s_j) = \begin{cases} f_1(s_j) & \text{if } \sigma_j = 1 \\ f_{-1}(s_j) & \text{if } \sigma_j = -1 \end{cases}.$$

The Natarajan dimension of Π , denoted by $d_N(\Pi)$, is the maximum cardinality of a set shattered by Π .

Let $d_{VC}(\Pi)$ be the Vapnik-Chervonenkis (VC) dimension of Π . Note that when $K = 2$, the Natarajan dimension is equivalent to the VC dimension; that is, when $K = 2$, $d_{VC}(\Pi) = d_N(\Pi)$ holds.

We derive the following lower bounds of the expected simple regret, which hold for any null consistent strategies and depend on the complexity of a policy class Π , measured by the Natarajan dimension. The proof is shown in Appendix A.

Theorem 3.4. *There exists a distribution ζ on \mathcal{X} such that for any $K \geq 2$, any null consistent strategy π with a policy class Π such that $d_N(\Pi) = M$ satisfies*

$$\sup_{P \in \mathcal{P}_\zeta} \sqrt{T} \mathbb{E}_P [R(P)(\pi)] \geq \frac{1}{8} \mathbb{E}_{X \sim \zeta} \left[\sqrt{M \sum_{a \in [K]} (\sigma^a(X))^2} \right],$$

where $\mathbb{E}_{X \sim \zeta}$ denotes the expectation of a random variable X under the probability distribution ζ .

Our lower bounds also depend on the variances of outcomes Y^a .

When $K = 2$, we can obtain a tighter lower bound than the one in Theorem 3.4. The proof is provided in Section A.4.

Theorem 3.5. *There exists a distribution ζ on \mathcal{X} such that for $K = 2$, any null consistent strategy π with policy class Π such that $d_{VC}(\Pi) = M$ satisfies*

$$\sup_{P \in \mathcal{P}_\zeta} \sqrt{T} \mathbb{E}_P [R(P)(\pi)] \geq \frac{1}{8} \mathbb{E}_{X \sim \zeta} \left[\sqrt{M (\sigma^1(X) + \sigma^2(X))^2} \right].$$

Here, note that

$$\sum_{a \in \{1,2\}} (\sigma^a(x))^2 \leq (\sigma^1(x) + \sigma^2(x))^2$$

holds for each $x \in \mathcal{X}$. Therefore, when $K = 2$, we use the lower bound in Theorem 3.5 and when $K \geq 3$, we use the one in Theorem 3.4.

In the course of proving Theorem 3.4, we address the following optimization problem with respect to w :

$$\min_{w \in \mathcal{W}} \left\{ \sum_{s \in \mathcal{S}} \max_{a \in [K]} \left\{ \sqrt{\frac{(\sigma^a(s))^2}{w(a|s)}} \right\} \right\},$$

where $\mathcal{S} := \{s_1, s_2, \dots, s_M\} \subseteq \mathcal{X}$ and \mathcal{W} is the set of all measurable functions $w : \mathcal{X} \times [K] \rightarrow (0, 1)$ such that $\sum_{a \in [K]} w(a | x) = 1$ for each $x \in \mathcal{X}$.

The solutions w^* , whose explicit forms appear in (1) and (2) in Section 4.1, works as a conjecture for $\frac{1}{T} \sum_{t=1}^T \mathbb{E}_P [\mathbb{1}[A_t = a] | X_t = x]$ of an optimal strategy that we derive in the following sections. Here, $\frac{1}{T} \sum_{t=1}^T \mathbb{E}_P [\mathbb{1}[A_t = a] | X_t = x]$ represents a fraction of units assigned to treatment arm a during the exploration phase. We refer to w^* as the *target assignment ratio*.¹

Remark (Efficiency gain). For each $a \in [K]$, let $(\sigma^a)^2$ be the unconditional variance of Y_t^a , and $w : [K] \rightarrow [0, 1]$ be an assignment ratio such that $\sum_{a \in [K]} w(a) = 1$ when we cannot utilize the contextual information. Then, the following inequality holds:

$$\begin{aligned} \sup_{P \in \mathcal{P}} \sqrt{T} \mathbb{E}_P [R(P) (\hat{\pi})] &\geq \frac{1}{8} \inf_{w \in \mathcal{W}} \max_{a \in [K]} \sqrt{M (\sigma^a)^2 / w(a)} \\ &\geq \frac{1}{8} \inf_{w \in \mathcal{W}} \max_{a \in [K]} \sqrt{M \mathbb{E}_{X \sim \zeta} [(\sigma^a(X))^2 / w(a | X)]} \\ &\geq \frac{1}{8} \inf_{w \in \mathcal{W}} \max_{a \in [K]} \mathbb{E}_{X \sim \zeta} \left[\sqrt{M (\sigma^a(X))^2 / w(a | X)} \right]. \end{aligned}$$

This result implies that we can minimize a lower bound by using contextual information; that is, strategies utilizing contextual information are more efficient than ones not utilizing contextual information.

4 PLAS Strategy

Our strategy consists of the following sampling and recommendation rules. At each round $t = 1, 2, \dots, T$, our sampling rule randomly assigns a unit to a treatment arm with a probability identical to an estimated target assignment ratio. After the final round T , our recommendation rule recommends a treatment arm with the highest value of a policy trained by maximizing empirical policy value. We refer to our strategy as the *adaptive-sampling (AS) and policy learning (PL) strategy*. We show a pseudo-code in Algorithm 1.

4.1 Target Assignment Ratio

As introduced in Section 3.2, the target assignment ratio is the expected value of the sample average of A_t of a strategy under which a leading factor of its expected simple regret aligns with that of our derived lower bound. The target sample assignment ratios are used to define a sampling rule in our proposed strategy.

From the arguments in Section 3, when $K = 2$, for each $a \in [K] = \{1, 2\}$, we define the target assignment ratio as

$$w^*(a | x) = \frac{\sigma^a(x)}{\sigma^1(x) + \sigma^2(x)}; \quad (1)$$

when $K \geq 3$, for each $a \in [K]$, we define the target assignment ratio as

$$w^*(a | x) = \frac{(\sigma^a(x))^2}{\sum_{b \in [K]} (\sigma^b(x))^2}. \quad (2)$$

These target assignment ratios are ex-ante unknown to the decision-maker since the variance $(\sigma^a(x))^2$ is unknown. Therefore, the decision-maker needs to estimate it during an adaptive experiment and employ the estimator as a probability of assigning a treatment arm.

4.2 Sampling Rule with Adaptive Sampling (AS)

For $a \in [K]$ and $t \in [T]$, let $\hat{w}_t(a | x)$ be an estimated target assignment ratio at round t .

As an initialization, for each $t = 1, 2, \dots, K$, we assign treatment arm $A_t = t$ and set $\hat{w}_t(a | X_t) = 1/K$ and $\hat{\mu}_t^a(X_t) = 0$ for all $a \in [K]$.

¹Target assignment ratios play important roles in the literature of BAI. Also see [Garivier and Kaufmann \(2016\)](#).

In each round $t \in [T]$, we obtain ξ_t from the uniform distribution on $[0, 1]$. In each round $t > K$, we assign treatment arm $A_t = 1$ if $\xi_t \leq \widehat{w}_t(1 | X_t)$ and $A_t = a$ for $a \geq 2$ if $\xi_t \in (\sum_{b=1}^{a-1} \widehat{w}_t(b | X_t), \sum_{b=1}^a \widehat{w}_t(b | X_t)]$; that is, we assign treatment arm a with probability $\widehat{w}_t(a | X_t)$. Note that we do not use ξ_t in each $t = 1, 2, \dots, K$, but it is required for defining an estimator of the expected outcomes.

In round $t > K$, for all $a \in [K]$, we estimate the target assignment ratio w^* using past observations \mathcal{F}_{t-1} , such that for all $a \in [K]$ and $x \in \mathcal{X}$, $\widehat{w}_t(a | x) > 0$ and $\sum_{a \in [K]} \widehat{w}_t(a | x) = 1$. We first estimate $\mu^a(P)(x)$ and $\nu^a(P)(x)$ using only past samples \mathcal{F}_{t-1} , denoted by $\widehat{\mu}_t^a(x)$ and $\widehat{\nu}_t^a(x)$, respectively. Let $(\widehat{\sigma}_t^{\dagger a}(x))^2 = \widehat{\nu}_t^a(x) - (\widehat{\mu}_t^a(x))^2$ for all $a \in [K]$ and $x \in \mathcal{X}$. Then, we estimate the variance $(\sigma^a(x))^2$ for all $a \in [K]$ and $x \in \mathcal{X}$ as $(\widehat{\sigma}_t^a(x))^2 = \text{thre}((\widehat{\sigma}_t^{\dagger a}(x))^2, \overline{C}, \underline{C})$, where $\text{thre}(A, a, b) = \max\{\min\{A, a\}, b\}$. Lastly, we define \widehat{w}_t by replacing the variances in w^* with corresponding estimators; that is, when $K = 2$, for each $a \in \{1, 2\}$, $\widehat{w}_t(a | X_t) = \frac{\widehat{\sigma}_t^1(X_t)}{\widehat{\sigma}_t^1(X_t) + \widehat{\sigma}_t^2(X_t)}$; when $K \geq 3$, for each $a \in [K]$, $\widehat{w}_t(a | X_t) = \frac{(\widehat{\sigma}_t^a(X_t))^2}{\sum_{b \in [K]} (\widehat{\sigma}_t^b(X_t))^2}$.

To show upper bounds, we use $\widehat{\mu}_t^a(x)$ and $\widehat{\nu}_t^a(x)$ such that $\widehat{\mu}_t^a(x)$ and $\widehat{\nu}_t^a(x)$ converge to the true functions for each a and x almost surely, and $|\widehat{\mu}_t^a(x)| < \overline{C}$ holds. We can use nonparametric estimators, such as the nearest neighbor regression estimator and kernel regression estimator, which have been proven to converge to the true function almost surely under a bounded sampling probability \widehat{w}_t by [Yang and Zhu \(2002\)](#) and [Qian and Yang \(2016\)](#). To estimate $\mu^a(P)(x)$ and $\nu^a(P)(x)$, we can use any estimators if $\widehat{\mu}_t^a(x)$ and $\widehat{\nu}_t^a(x)$ converge to the true functions almost surely, and $|\widehat{\mu}_t^a(x)| < \overline{C}$ for each a and x . It should be noted that we do not assume specific convergence rates for estimators for $\mu^a(P)(x)$ and w^* as the asymptotic optimality of the AIPW estimator can be demonstrated without them ([van der Laan, 2008](#); [Kato et al., 2020, 2021](#)).

4.3 Recommendation Rule with Policy Learning

The following part presents our recommendation rule. To recommend the conditionally best treatment arm $a^*(P)(x)$, we train a policy $\pi : [K] \times \mathcal{X} \rightarrow [0, 1]$ by maximizing the empirically approximated policy value function, which we will describe below.

At the end of an experiment, we estimate the policy value $Q(\pi)$ by using the augmented doubly robust estimator, which is defined as follows:

$$\widehat{Q}_T(\pi) := \frac{1}{T} \sum_{t=1}^T \sum_{a \in [K]} \pi(a | X_t) \widehat{\Gamma}_t^a, \quad (3)$$

where

$$\widehat{\Gamma}_t^a := \frac{\mathbb{1}[A_t = a](c_T(Y_t) - \widehat{\mu}_t^a(X_t))}{\widehat{w}_t(a | X_t)} + \widehat{\mu}_t^a(X_t), \quad c_T(Y_t) = \text{thre}(Y_t, U_T, -U_T),$$

and U_T is a positive value that approaches infinity as $T \rightarrow \infty$. Then, we train a policy by

$$\widehat{\pi}_T^{\text{PLAS}} := \arg \max_{\pi \in \Pi} \widehat{Q}_T(\pi)$$

By using this trained policy function, given $x \in \mathcal{X}$, we recommend $\widehat{a}_T(x) \in [K]$ as the best treatment arm with probability $\widehat{\pi}_T^{\text{PLAS}}(\widehat{a}_T(x) | x)$.

The AIPW estimator debiases the sample selection bias resulting from treatment assignment based on contextual information. Additionally, the AIPW estimator possesses the following properties: (i) its components $\{\widehat{\Gamma}_t^a - \mu^a(P)(x)\}_{t=1}^T$ are a martingale difference sequence, allowing us to employ the martingale limit theorems in derivation of the upper bound; (ii) it has the minimal asymptotic variance among the possible estimators. For example, other estimators with a martingale property, such as the inverse probability weighting (IPW) estimator, may be employed, yet their asymptotic variance would be greater than that of the AIPW estimator. The t -th element of the sum in the AIPW estimator utilizes nuisance parameters $(\mu^a(P)(x)$ and w^*) estimated from past observations up to round $t - 1$ for constructing a martingale difference sequence ([van der Laan, 2008](#); [Hadad et al., 2021](#); [Kato et al., 2020, 2021](#)). For those reasons, this estimator is often used in the context of adaptive experimental design.

5 Policy Upper Bound

This section provides upper bounds for the expected simple regret of the PLAS strategy. First, we assume the following convergence rate for estimators of $\mu^a(P)(x)$ and $w^*(a | x)$.

Algorithm 1 PLAS strategy

Parameter: Positive constants C_μ and C_{σ^2} .

Initialization:

for $t = 1$ to K **do**

Assign $A_t = t$. For each $a \in [K]$, set $\hat{w}_t(a | x) = 1/K$.

end for

for $t = K + 1$ to T **do**

Observe X_t .

Construct \hat{w}_t by using the estimators of the variances.

Draw ξ_t from the uniform distribution on $[0, 1]$.

$A_t = 1$ if $\xi_t \leq \hat{w}_t(1 | X_t)$ and $A_t = a$ for $a \geq 2$ if $\xi_t \in \left(\sum_{b=1}^{a-1} \hat{w}_t(b | X_t), \sum_{b=1}^a \hat{w}_t(b | X_t) \right]$.

end for

Construct $\hat{Q}(\pi)$ following (3).

Train a policy $\hat{\pi}_T^{\text{PLAS}}$ as $\hat{\pi}_T^{\text{PLAS}} = \arg \max_{\pi \in \Pi} \hat{Q}(\pi)$.

Recommend \hat{a}_T following $\hat{\pi}_T^{\text{PLAS}}$.

Assumption 5.1. For any ζ , any $P \in \mathcal{P}_\zeta$, and all $a \in [K]$,

$$\sup_{x \in \mathcal{X}} \left| \hat{w}_T(a | x) - w^*(a | x) \right| \xrightarrow{\text{a.s.}} 0, \quad \sup_{x \in \mathcal{X}} \left| \hat{\mu}_T^a(x) - \mu^a(P)(x) \right| \xrightarrow{\text{a.s.}} 0.$$

holds as $T \rightarrow \infty$.

Next, we define an entropy integral of a policy class.

Definition 5.2. Given the feature domain \mathcal{X} , a policy class Π , a set of n points $\{x_1, \dots, x_n\} \subset \mathcal{X}$, define:

1. Hamming distance between any two policies π_1 and π_2 in Π : $H(\pi_1, \pi_2) = \frac{1}{n} \sum_{j=1}^n \mathbb{1}[\pi_1(x_j) \neq \pi_2(x_j)]$.
2. ϵ -Hamming covering number of the set $\{x_1, \dots, x_n\}$:
 $\mathbb{N}_H(\epsilon, \Pi, \{x_1, \dots, x_n\})$ is the smallest number K of policies $\{\pi_1, \dots, \pi_K\}$ in Π , such that $\forall \pi \in \Pi, \exists \pi_i, H(\pi, \pi_i) \leq \epsilon$.
3. ϵ -Hamming covering number of Π : $\mathbb{N}_H(\epsilon, \Pi) = \sup \{ \mathbb{N}_H(\epsilon, \Pi, \{x_1, \dots, x_m\}) \mid m \geq 1, x_1, \dots, x_m \in \mathcal{X} \}$.
4. Entropy integral: $\kappa(\Pi) = \int_0^1 \sqrt{\log \mathbb{N}_H(\epsilon^2, \Pi)} d\epsilon$.

The entropy represents the complexity of a policy class, as well as the Natarajan dimension. Between the entropy integral $\kappa(\Pi)$ and the Natarajan dimension $d_N(\Pi)$, $\kappa(\Pi) \leq C \sqrt{\log(d) d_N(\Pi)}$ holds for some universal constant $C > 0$ (Jin, 2023; Zhan et al., 2022) when $K \geq 3$. When $K = 2$, $d_N(\Pi)$ is equal to the VC dimension, and $\kappa(\Pi) \leq 2.5 \sqrt{d_N(\Pi)}$ holds (Haussler, 1995).

Furthermore, we make the following assumption for the ϵ -Hamming covering number.

Assumption 5.3. For all $\epsilon \in (0, 1)$, $\mathbb{N}_H(\epsilon, \Pi) \leq C \exp(D(\frac{1}{\epsilon})^\omega)$ for some constants $C, D > 0, 0 < \omega < 0.5$.

For analysis, let us define the following quantities:

$$\begin{aligned} \mu_c^a(P)(X) &:= \mathbb{E}_P[c_T(Y^a)], \\ Q_c(P)(\pi) &:= \mathbb{E}_{X \sim \zeta} \left[\sum_{a \in [K]} \pi(a | X) \mu_c^a(P)(X) \right], \\ \pi_c^*(P) &:= \arg \max_{\pi \in \Pi} Q_c(P)(\pi), \\ R_c(P)(\pi) &:= Q_c(P)(\pi_c^*) - Q_c(P)(\pi). \end{aligned}$$

These quantities are used in the conditions in the main theorem and the following proof. For Q and Q_c , we make the following assumption.

Assumption 5.4. For any ζ , any $P \in \mathcal{P}_\zeta$, and any $\pi \in \Pi$, $\sqrt{T}Q(P)(\pi) - \sqrt{T}Q_c(P)(\pi) \rightarrow 0$ holds as $T \rightarrow \infty$. Additionally, for any $P \in \mathcal{P}$, $\max_{\pi \in \Pi} \sqrt{T}Q(P)(\pi) - \max_{\pi \in \Pi} \sqrt{T}Q_c(P)(\pi) \rightarrow 0$ holds as $T \rightarrow \infty$.

Because we set U_T in $c_T(\cdot)$ as $U_T \rightarrow \infty$ when $T \rightarrow \infty$, this assumption implies that when a clipping $c_T(Y_t) = \text{thre}(Y_t, U_T, -U_T)$ asymptotically vanishes as $T \rightarrow \infty$, π_c^* approaches π^* , a optimal policy under outcomes without the clipping.

Then, we obtain the following upper bound for the expected simple regret of the PLAS strategy.

Theorem 5.5 (Upper bound). *Suppose that Assumptions 5.1–5.4 hold. Then, for any ζ and any $P \in \mathcal{P}_\zeta$, the following upper bounds hold:*

- When $K \geq 3$, under the PLAS strategy,

$$\sqrt{T}\mathbb{E}_P \left[R(P) (\hat{\pi}_T^{\text{PLAS}}) \right] \leq (108.8\kappa(\Pi) + 870.4) \mathbb{E}_{X \sim \zeta} \left[\sqrt{\sum_{a=1}^K (\sigma^a(X))^2} \right] + o(1)$$

holds as $T \rightarrow \infty$.

- When $K = 2$,

$$\sqrt{T}\mathbb{E}_P \left[R(P) (\hat{\pi}_T^{\text{PLAS}}) \right] \leq (108.8\kappa(\Pi) + 870.4) \mathbb{E}_{X \sim \zeta} \left[\sqrt{(\sigma^a(X) + \sigma^2(X))^2} \right] + o(1)$$

holds as $T \rightarrow \infty$.

Additionally, for any ζ and any $P \in \mathcal{P}_\zeta$, the following upper bound hold:

- When $K \geq 3$, under the PLAS strategy, if $d_N(\Pi) = M$, then

$$\sqrt{T}\mathbb{E}_P \left[R(P) (\hat{\pi}_T^{\text{PLAS}}) \right] \leq (108.8C\sqrt{\log(d)M} + 870.4) \mathbb{E}_{X \sim \zeta} \left[\sqrt{\sum_{a=1}^K (\sigma^a(X))^2} \right] + o(1)$$

holds as $T \rightarrow \infty$, where $C > 0$ is a universal constant.

- When $K = 2$, under the PLAS strategy, if $d_{VC}(\Pi) = M$, then

$$\sqrt{T}\mathbb{E}_P \left[R(P) (\hat{\pi}_T^{\text{PLAS}}) \right] = (272\sqrt{M} + 870.4) \mathbb{E}_{X \sim \zeta} \left[\sqrt{\sum_{a=1}^K (\sigma^a(X))^2} \right] + o(1)$$

holds as $T \rightarrow \infty$.

From Lemma 5.5 and the relationship between the entropy integral and the Natarajan dimension, the following theorem holds.

We note that the leading factor in these upper bounds are $\mathbb{E}_{X \sim \zeta} \left[\sqrt{\log(d)M \sum_{a=1}^K (\sigma^a(X))^2} \right]$ and $\mathbb{E}_{X \sim \zeta} \left[\sqrt{M (\sigma^a(X) + \sigma^2(X))^2} \right]$, a product of the policy complexity (Natarajan dimension $d_N(\Pi)$) and outcome variances. This theorem implies that the leading factors align with the lower bounds with high probability. Here, the terms $\sqrt{\sum_{a=1}^K \mathbb{E}_{X \sim \zeta} [(\sigma^a(X))^2]}$ and $\sqrt{\mathbb{E}_{X \sim \zeta} [(\sigma^a(X) + \sigma^2(X))^2]}$ remain, which are relatively smaller than the leading factors if M is sufficiently large. We might tight these terms as $\mathbb{E}_{X \sim \zeta} \left[\sqrt{\sum_{a=1}^K (\sigma^a(X))^2} \right]$ and $\mathbb{E}_{X \sim \zeta} \left[\sqrt{(\sigma^a(X) + \sigma^2(X))^2} \right]$, which is an open issue.

In the following sections, we prove the upper bound. To show the upper bound, we aim to use the result of Zhou et al. (2023), which provides an upper bound for expected simple regret in the problem of policy learning with multiple treatment arms. Here, note that we cannot directly apply the results of Zhou et al. (2023) because they assume that the observations are i.i.d. in their study, but observations are non-i.i.d. in our study. They also assume that their outcomes are bounded, but our problem satisfies the requirement by clipping the outcomes using $c_T(\cdot)$. To circumvent the issue of non-i.i.d. observations, we show an asymptotic equivalence between our empirical policy value $\hat{Q}_T(\pi)$ and

a hypothetical empirical policy value constructed from hypothetical i.i.d. observations in Section 5.1. Then, using the results of Zhou et al. (2023), we upper bound a regret for $Q_c(P)(\pi_c^*) - Q_c(P)(\hat{\pi})$ with the hypothetical policy value in Section 5.2. Finally, in Section 5.3, we derive the upper bounds for $R(\pi^*) = Q(P)(\pi^*) - Q(P)(\hat{\pi})$ from the upper bounds for $Q_c(P)(\pi_c^*) - Q_c(P)(\hat{\pi})$.

5.1 Asymptotic Equivalence

Let $w_t(0 | x)$ be 0. We write $\hat{\Gamma}_t^a$ as

$$\hat{\Gamma}_t^a = \hat{\Gamma}^a((Y_t^a)_{a \in [K]}, \xi_t, X_t) = \frac{\mathbb{1} \left[\sum_{b=0}^{a-1} \hat{w}_t(b | X_t) \leq \xi_t \leq \sum_{b=0}^a \hat{w}_t(b | X_t) \right] (c_T(Y_t^a) - \hat{\mu}_t^a(X_t))}{\hat{w}_t(a | X_t)} + \hat{\mu}_t^a(X_t).$$

This expression equals the original definition of $\hat{\Gamma}_t^a$. Let $w^*(0 | x)$ be 0. Then, we define

$$\Gamma_t^{*a}((Y_t^a)_{a \in [K]}, \xi_t, X_t) := \frac{\mathbb{1} \left[\sum_{b=0}^{a-1} w^*(b | X_t) \leq \xi_t \leq \sum_{b=0}^a w^*(b | X_t) \right] (c_T(Y_t^a) - \mu^a(P)(X_t))}{w^*(a | X_t)} + \mu^a(P)(X_t).$$

Here, we show that $\hat{\Gamma}_t^a$ and Γ_t^{*a} are asymptotically equivalent. The proof is shown in Appendix B.

Lemma 5.6. *Suppose that Assumption 5.1 holds. Then,*

$$\sqrt{T} \hat{Q}_T(\hat{\pi}_T^{\text{PLAS}}) = \sqrt{T} \sum_{t=1}^T \sum_{a \in [K]} \pi(a | X_t) \Gamma_c^{*a}((Y_t^a)_{a \in [K]}, \xi_t, X_t) + o_P(1).$$

holds as $T \rightarrow \infty$.

Denote $\sqrt{T} \sum_{t=1}^T \sum_{a \in [K]} \pi(a | X_t) \Gamma_c^{*a}$ by $\hat{Q}_T(\hat{\pi}_T^{\text{PLAS}})$. Note that $\hat{Q}_T(\hat{\pi}_T^{\text{PLAS}})$ consists of only i.i.d. observations; therefore, we can directly apply the results of Zhou et al. (2023) to derive the upper bound for the policy regret.

This technique of the asymptotic equivalence is inspired by Hahn et al. (2011) and is important because, for dependent observations (even if they are martingales), we cannot apply the tools for upper-bounding regrets or risks, such as the Rademacher complexity. For instance, Zhan et al. (2022) addresses this issue and establishes an off-policy learning method from adaptively collected observations by utilizing the Rademacher complexity for martingales developed by Rakhlin et al. (2015). However, our interest lies in developing upper bounds depending on variances because our lower bounds also depend on variances, and such upper bounds are considered to be tight. Although Zhou et al. (2023) derives such upper bounds for policy learning from i.i.d. observations using the local Rademacher complexity, it is unclear whether we can use the results of Zhou et al. (2023) with the Rademacher complexity for martingales developed by Rakhlin et al. (2015). In contrast, in this study, if we restrict the problem to BAI and the evaluation metric to the worst-case expected simple regret, we show that we can apply the results of Zhou et al. (2023) and avoid the use of the Rademacher complexity for martingales by bypassing a hypothetical policy value that only depends on i.i.d. observations.

5.2 Upper Bound under I.I.D. Observations

Because $\hat{Q}_T(\hat{\pi}_T^{\text{PLAS}})$ is asymptotically equivalent to a policy value that consists of i.i.d. observations, we can apply the results of policy learning with i.i.d. observations to bound the policy regret. Specifically, we modify a regret upper bound shown by Zhou et al. (2023), given as the following lemma.

Lemma 5.7 (Modified upper bound). *Suppose that Assumptions 5.1 and 5.3 hold. Then, for any ζ and any $P \in \mathcal{P}_\zeta$,*

$$\mathbb{E}_P \left[R_c(P)(\hat{\pi}_T^{\text{PLAS}}) \right] = \mathbb{E}_P \left[Q_c(P)(\pi_c^*) - Q_c(P)(\hat{\pi}_T^{\text{PLAS}}) \right] = \mathbb{E}_P \left[(54.4\kappa(\Pi) + 435.2) \frac{\Upsilon_*}{\sqrt{T}} + O\left(\frac{\sqrt{U_T}}{T^{3/4}}\right) \right]$$

holds, where

$$\Upsilon_* = \mathbb{E} \left[\sqrt{\sup_{\pi_1, \pi_2 \in \Pi} \sum_{t \in [T]} \left\{ \sum_{a \in [K]} (\pi_1(a | Z_t) - \pi_2(a | Z_t)) \Gamma^{*a}(Y_t, \xi_t, X_t) \right\}^2} \right].$$

The proof is shown in Appendix C.

5.3 Asymptotic Optimality

Finally, we derive the upper bounds of $R(P)(\hat{\pi}_T^{\text{PLAS}}) = Q(P)(\pi^*) - Q(P)(\hat{\pi}_T^{\text{PLAS}})$ from the upper bounds of $R_c(P)(\hat{\pi}_T^{\text{PLAS}}) = Q_c(P)(\pi_c^*) - Q_c(P)(\hat{\pi}_T^{\text{PLAS}})$.

By using Assumption 5.4, we can evaluate the regret $R(\pi^*)$ as follows:

$$\begin{aligned}
R(P)(\hat{\pi}_T^{\text{PLAS}}) &= R(P)(\hat{\pi}_T^{\text{PLAS}}) - R_c(P)(\hat{\pi}_T^{\text{PLAS}}) + R_c(P)(\hat{\pi}_T^{\text{PLAS}}) \\
&= Q(P)(\pi^*) - Q(P)(\hat{\pi}_T^{\text{PLAS}}) - \left\{ Q_c(P)(\pi_c^*) - Q_c(P)(\hat{\pi}_T^{\text{PLAS}}) \right\} + \left\{ Q_c(P)(\pi_c^*) - Q_c(P)(\hat{\pi}_T^{\text{PLAS}}) \right\} \\
&= \left\{ Q(P)(\pi^*) - Q(P)(\pi_c^*) \right\} + \left(\left\{ Q(P)(\pi_c^*) - Q(P)(\hat{\pi}_T^{\text{PLAS}}) \right\} - \left\{ Q_c(P)(\pi_c^*) - Q_c(P)(\hat{\pi}_T^{\text{PLAS}}) \right\} \right) \\
&\quad + \left\{ Q_c(P)(\pi_c^*) - Q_c(P)(\hat{\pi}_T^{\text{PLAS}}) \right\} \\
&= \left\{ Q_c(P)(\pi_c^*) - Q_c(P)(\hat{\pi}_T^{\text{PLAS}}) \right\} + o(1/\sqrt{T}) = R_c(P)(\hat{\pi}_T^{\text{PLAS}}) + o(1/\sqrt{T}).
\end{aligned}$$

Let $c_T = T^{1/10}$. From the Cauchy-Schwarz inequality, we have

$$\begin{aligned}
\Upsilon_* &= \mathbb{E} \left[\sqrt{\sup_{\pi_1, \pi_2 \in \Pi} \sum_{t \in [T]} \left\{ \sum_{a \in [K]} (\pi_1(a | X_t) - \pi_2(a | X_t)) \Gamma_c^{*a}(Y_t, \xi_t, X_t) \right\}^2} \right] \\
&\leq \mathbb{E} \left[\sqrt{\sup_{\pi_1, \pi_2 \in \Pi} \sum_{t \in [T]} \left\{ \sum_{a \in [K]} (\pi_1(a | X_t) - \pi_2(a | X_t)) \right\}^2 \left\{ \sum_{a \in [K]} \Gamma_c^{*a}(Y_t, \xi_t, X_t) \right\}^2} \right] \\
&\leq \mathbb{E} \left[\sqrt{\sup_{\pi_1, \pi_2 \in \Pi} \sum_{t \in [T]} 4 \left\{ \sum_{a \in [K]} \Gamma_c^{*a}(Y_t, \xi_t, X_t) \right\}^2} \right] \\
&= 2 \mathbb{E} \left[\sqrt{\sum_{t \in [T]} \left\{ \sum_{a \in [K]} \Gamma_c^{*a}(Y_t, \xi_t, X_t) \right\}^2} \right].
\end{aligned}$$

From the law of iterated expectations and the Jensen inequality, we have

$$\begin{aligned}
\Upsilon_* &\leq 2 \mathbb{E} \left[\sqrt{\sum_{t \in [T]} \left\{ \sum_{a \in [K]} \Gamma_c^{*a}(Y_t, \xi_t, X_t) \right\}^2} \right] \\
&= 2 \mathbb{E}_{X \sim \zeta} \left[\sqrt{\mathbb{E} \left[\sum_{t \in [T]} \left\{ \sum_{a \in [K]} \Gamma_c^{*a}(Y_t, \xi_t, X_t) \right\}^2 \right]} \right] = 2 \mathbb{E}_{X \sim \zeta} \left[\sqrt{T \mathbb{E} \left[\left\{ \sum_{a \in [K]} \Gamma_c^{*a}(Y_t, \xi_t, X_t) \right\}^2 \right]} \right].
\end{aligned}$$

In conclusion, we obtain

$$\begin{aligned}
\Upsilon_*/\sqrt{T} &\leq 2 \mathbb{E}_{X \sim \zeta} \left[\sqrt{\mathbb{E} \left[\left\{ \sum_{a \in [K]} \Gamma_c^{*a}(Y_t, \xi_t, X_t) \right\}^2 \right]} \right] \\
&= 2 \mathbb{E}_{X \sim \zeta} \left[\sqrt{\mathbb{E} \left[\left\{ \sum_{a \in [K]} \Gamma^{*a}(Y_t, \xi_t, X_t) \right\}^2 \right]} \right] + o(1) = 2 \mathbb{E}_{X \sim \zeta} \left[\sqrt{\sum_{a \in [K]} \frac{(\sigma^a(X))^2}{w^*(a | X)}} \right] + o(1),
\end{aligned}$$

as $T \rightarrow \infty$ ($c_T \rightarrow \infty$).

Similarly, we obtain

$$\tilde{\Upsilon}_*/\sqrt{T} \leq 2 \sqrt{\sum_{a \in [K]} \mathbb{E}_{X \sim \zeta} \left[\frac{(\sigma^a(X))^2}{w^*(a|X)} \right]} + o(1).$$

By substituting w^* for each case with $K = 2$ and $K \geq 3$, we obtain Theorem 5.5.

6 Conclusion

In this study, we presented an adaptive experiment with policy learning. Our main contributions are the lower bound of strategies, the PLAS strategy, and its regret upper bound. First, by employing the lower bounds developed by Kaufmann et al. (2016), we derived lower bounds for the expected simple regret. Our derived lower bounds depend on the variances of outcomes. Then, based on the lower bounds, we developed the PLAS strategy, which trains a policy at the end of the experiment. Lastly, we showed upper bounds for the PLAS strategy.

From a technical viewpoint, we showed how we use Rademacher complexity for i.i.d. samples in an adaptive experiment with non-i.i.d. samples. We did not employ complexity measures for non-i.i.d. samples presented by Rakhlin et al. (2015) and Foster et al. (2023). Our technique relies on an approach used in Hahn et al. (2011), and we extend this approach by using sample splitting, also known as double machine learning (van der Laan, 2008; Zheng and van der Laan, 2011; Chernozhukov et al., 2018; Hadad et al., 2021; Kato et al., 2020, 2021).

We also contributed to the literature on policy learning by providing a variance-dependent lower bound, which works in observational studies with i.i.d. samples, and by discussing matching upper bounds. Our derived lower bound differs from that in Athey and Wager (2021) and more tightly depends on variances, which also requires the refinement of existing upper bounds.

Our next step is to tighten the lower and upper bounds. When $K = 2$, we showed that assigning each treatment arm with a proportion of standard deviations is optimal, which is consistent with existing work such as Neyman (1934), Glynn and Juneja (2004), and Kaufmann et al. (2016). In contrast, we showed that assigning each treatment arm with a proportion of variances is optimal when $K \geq 3$. However, other studies for fixed-budget BAI without contextual information, such as Glynn and Juneja (2004), Kaufmann et al. (2016), and Kato (2023a), show that strategies with different sampling rules are optimal. Filling the gap between our study and those existing studies is an open issue.

References

- Adusumilli, K. (2021), “Risk and optimal policies in bandit experiments,” arXiv:2112.06363.
- (2022), “Minimax policies for best arm identification with two arms,” arXiv:2204.05527.
- Ariu, K., Kato, M., Komiyama, J., McAlinn, K., and Qin, C. (2021), “Policy Choice and Best Arm Identification: Asymptotic Analysis of Exploration Sampling,” arXiv:2109.08229.
- Armstrong, T. B. (2022), “Asymptotic Efficiency Bounds for a Class of Experimental Designs,” arXiv:2205.02726.
- Athey, S. and Wager, S. (2021), “Policy Learning With Observational Data,” *Econometrica*, 89, 133–161.
- Atsidakou, A., Katariya, S., Sanghavi, S., and Kveton, B. (2023), “Bayesian Fixed-Budget Best-Arm Identification,” arXiv:2211.08572.
- Audibert, J.-Y., Bubeck, S., and Munos, R. (2010), “Best Arm Identification in Multi-Armed Bandits,” in *Conference on Learning Theory*, pp. 41–53.
- Auer, P., Cesa-Bianchi, N., Freund, Y., and Schapire, R. E. (2002), “The Nonstochastic Multiarmed Bandit Problem,” *SIAM Journal on Computing*, 32, 48–77.
- Bang, H. and Robins, J. M. (2005), “Doubly Robust Estimation in Missing Data and Causal Inference Models,” *Biometrics*, 61, 962–973.
- Bartlett, P. L. and Mendelson, S. (2003), “Rademacher and gaussian complexities: risk bounds and structural results,” *Journal of Machine Learning Research*, 3, 463–482.
- Bechhofer, R., Kiefer, J., and Sobel, M. (1968), *Sequential Identification and Ranking Procedures: With Special Reference to Koopman-Darmois Populations*, University of Chicago Press.

- Bubeck, S., Munos, R., and Stoltz, G. (2009), “Pure Exploration in Multi-armed Bandits Problems,” in *Algorithmic Learning Theory (ALT)*.
- (2011), “Pure exploration in finitely-armed and continuous-armed bandits,” *Theoretical Computer Science*.
- Carpentier, A. and Locatelli, A. (2016), “Tight (Lower) Bounds for the Fixed Budget Best Arm Identification Bandit Problem,” in *Conference on Learning Theory (COLT)*.
- Chen, C.-H., Lin, J., Yücesan, E., and Chick, S. E. (2000), “Simulation Budget Allocation for Further Enhancing TheEfficiency of Ordinal Optimization,” *Discrete Event Dynamic Systems*, 10, 251–270.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C., Newey, W., and Robins, J. (2018), “Double/debiased machine learning for treatment and structural parameters,” *The Econometrics Journal*.
- Degenne, R. (2023), “On the Existence of a Complexity in Fixed Budget Bandit Identification,” .
- Deshmukh, A. A., Sharma, S., Cutler, J. W., Moldwin, M., and Scott, C. (2018), “Simple Regret Minimization for Contextual Bandits,” arXiv:1810.07371.
- Dominitz, J. and Manski, C. F. (2022), “Minimax-regret sample design in anticipation of missing data, with application to panel data,” *Journal of Econometrics*, 226, 104–114, annals Issue in Honor of Gary Chamberlain.
- Dominitz, J. and Manski, F. C. (2017), “More Data or Better Data? A Statistical Decision Problem,” *The Review of Economic Studies*, 84, 1583–1605.
- Even-Dar, E., Mannor, S., Mansour, Y., and Mahadevan, S. (2006), “Action Elimination and Stopping Conditions for the Multi-Armed Bandit and Reinforcement Learning Problems.” *Journal of machine learning research*.
- Foster, D., Foster, D. J., Golowich, N., and Rakhlin, A. (2023), “On the Complexity of Multi-Agent Decision Making: From Learning in Games to Partial Monitoring,” in *Conference on Learning Theory (COLT)*.
- Garivier, A. and Kaufmann, E. (2016), “Optimal Best Arm Identification with Fixed Confidence,” in *Conference on Learning Theory*.
- Glynn, P. and Juneja, S. (2004), “A large deviations perspective on ordinal optimization,” in *Proceedings of the 2004 Winter Simulation Conference*, IEEE, vol. 1.
- Guan, M. and Jiang, H. (2018), “Nonparametric Stochastic Contextual Bandits,” *AAAI Conference on Artificial Intelligence*.
- Gupta, S., Lipton, Z. C., and Childers, D. (2021), “Efficient Online Estimation of Causal Effects by Deciding What to Observe,” in *Advances in Neural Information Processing Systems (NeurIPS)*.
- Hadad, V., Hirshberg, D. A., Zhan, R., Wager, S., and Athey, S. (2021), “Confidence intervals for policy evaluation in adaptive experiments,” *Proceedings of the National Academy of Sciences*, 118.
- Hahn, J., Hirano, K., and Karlan, D. (2011), “Adaptive experimental design using the propensity score,” *Journal of Business and Economic Statistics*.
- Haussler, D. (1995), “Sphere packing numbers for subsets of the Boolean n-cube with bounded Vapnik-Chervonenkis dimension,” *Journal of Combinatorial Theory, Series A*, 69, 217–232.
- Hirano, K. and Porter, J. R. (2009), “Asymptotics for Statistical Treatment Rules,” *Econometrica*, 77, 1683–1701.
- (2023), “Asymptotic Representations for Sequential Decisions, Adaptive Experiments, and Batched Bandits,” arXiv:2302.03117.
- Ito, S., Tsuchiya, T., and Honda, J. (2022), “Adversarially Robust Multi-Armed Bandit Algorithm with Variance-Dependent Regret Bounds,” in *Conference on Learning Theory*.
- Jennison, C., Johnstone, I. M., and Turnbull, B. W. (1982), “Asymptotically Optimal Procedures for Sequential Adaptive Selection of the Best of Several Normal Means,” in *Statistical Decision Theory and Related Topics III*, Academic Press, pp. 55–86.
- Jin, Y. (2023), “Upper bounds on the Natarajan dimensions of some function classes,” arXiv:2209.07015.
- Jourdan, M., Rémy, D., and Emilie, K. (2023), “Dealing with Unknown Variances in Best-Arm Identification,” in *Proceedings of The 34th International Conference on Algorithmic Learning Theory*, vol. 201, pp. 776–849.
- Karlan, D. and Wood, D. H. (2014), “The Effect of Effectiveness: Donor Response to Aid Effectiveness in a Direct Mail Fundraising Experiment,” Working Paper 20047, National Bureau of Economic Research.
- Kasy, M. and Sautmann, A. (2021), “Adaptive Treatment Assignment in Experiments for Policy Choice,” *Econometrica*, 89, 113–132.

- Kato, M. (2023a), “Locally Optimal Fixed-Budget Best Arm Identification in Two-Armed Gaussian Bandits with Unknown Variances,” arXiv:2312.12741.
- (2023b), “Worst-Case Optimal Multi-Armed Gaussian Best Arm Identification with a Fixed Budget,” arXiv:2310.19788.
- Kato, M. and Ariu, K. (2021), “The Role of Contextual Information in Best Arm Identification,” arXiv:2106.14077.
- Kato, M., Imaizumi, M., Ishihara, T., and Kitagawa, T. (2023), “Asymptotically Optimal Fixed-Budget Best Arm Identification with Variance-Dependent Bounds,” arXiv:2302.02988v2.
- Kato, M., Ishihara, T., Honda, J., and Narita, Y. (2020), “Adaptive Experimental Design for Efficient Treatment Effect Estimation: Randomized Allocation via Contextual Bandit Algorithm,” arXiv:2002.05308.
- Kato, M., McAlinn, K., and Yasui, S. (2021), “The Adaptive Doubly Robust Estimator and a Paradox Concerning Logging Policy,” in *Advances in Neural Information Processing Systems (NeurIPS)*.
- Kaufmann, E. (2020), *Contributions to the Optimal Solution of Several Bandits Problems*, Habilitation à Diriger des Recherches, Université de Lille.
- Kaufmann, E., Cappé, O., and Garivier, A. (2016), “On the Complexity of Best-Arm Identification in Multi-Armed Bandit Models,” *Journal of Machine Learning Research*, 17, 1–42.
- Kim, W., Kim, G.-S., and Paik, M. C. (2021), “Doubly Robust Thompson Sampling with Linear Payoffs,” in *Advances in Neural Information Processing Systems (NeurIPS)*.
- Kitagawa, T. and Tetenov, A. (2018), “Who Should Be Treated? Empirical Welfare Maximization Methods for Treatment Choice,” *Econometrica*, 86, 591–616.
- Kock, A. B., Preinerstorfer, D., and Veliyev, B. (2023), “Treatment recommendation with distributional targets,” *Journal of Econometrics*, 234, 624–646.
- Komiyama, J., Ariu, K., Kato, M., and Qin, C. (2023), “Rate-Optimal Bayesian Simple Regret in Best Arm Identification,” *Mathematics of Operations Research*.
- Lai, T. and Robbins, H. (1985), “Asymptotically efficient adaptive allocation rules,” *Advances in Applied Mathematics*.
- Lalitha, A., Kalantari, K., Ma, Y., Deoras, A., and Kveton, B. (2023), “Fixed-Budget Best-Arm Identification with Heterogeneous Reward Variances,” in *Conference on Uncertainty in Artificial Intelligence*.
- Le Cam, L. (1960), *Locally Asymptotically Normal Families of Distributions. Certain Approximations to Families of Distributions and Their Use in the Theory of Estimation and Testing Hypotheses*, University of California Publications in Statistics. vol. 3. no. 2, Berkeley & Los Angeles.
- (1972), “Limits of experiments,” in *Theory of Statistics*, University of California Press, pp. 245–282.
- (1986), *Asymptotic Methods in Statistical Decision Theory (Springer Series in Statistics)*, Springer.
- Lehmann, E. L. and Casella, G. (1998), *Theory of Point Estimation*, Springer-Verlag.
- Manski, C. (2000), “Identification problems and decisions under ambiguity: Empirical analysis of treatment response and normative analysis of treatment choice,” *Journal of Econometrics*, 95, 415–442.
- Manski, C. F. (2002), “Treatment choice under ambiguity induced by inferential problems,” *Journal of Statistical Planning and Inference*, 105, 67–82.
- (2004), “Statistical Treatment Rules for Heterogeneous Populations,” *Econometrica*, 72, 1221–1246.
- Manski, C. F. and Tetenov, A. (2016), “Sufficient trial size to inform clinical practice,” *Proceedings of the National Academy of Sciences*, 113, 10518–10523.
- Masoudian, S. and Seldin, Y. (2021), “Improved Analysis of the Tsallis-INF Algorithm in Stochastically Constrained Adversarial Bandits and Stochastic Bandits with Adversarial Corruptions,” in *Conference on Learning Theory (COLT)*.
- Natarajan, B. K. (1989), “On learning sets and functions,” *Machine Learning*, 4, 67–97.
- Neyman, J. (1923), “Sur les applications de la theorie des probabilites aux experiences agricoles: Essai des principes,” *Statistical Science*, 5, 463–472.
- (1934), “On the Two Different Aspects of the Representative Method: the Method of Stratified Sampling and the Method of Purposive Selection,” *Journal of the Royal Statistical Society*, 97, 123–150.
- Qian, W. and Yang, Y. (2016), “Kernel Estimation and Model Combination in A Bandit Problem with Covariates,” *Journal of Machine Learning Research*.

- Qin, C. (2022), “Open Problem: Optimal Best Arm Identification with Fixed-Budget,” in *Conference on Learning Theory*.
- Qin, C., Klabjan, D., and Russo, D. (2017), “Improving the Expected Improvement Algorithm,” in *Advances in Neural Information Processing Systems (NeurIPS)*.
- Qin, C. and Russo, D. (2022), “Adaptivity and Confounding in Multi-Armed Bandit Experiments,” arXiv:2202.09036.
- Rakhlin, A., Sridharan, K., and Tewari, A. (2015), “Sequential complexities and uniform martingale laws of large numbers,” *Probability Theory and Related Fields*, 161, 111–153.
- Robbins, H. (1952), “Some aspects of the sequential design of experiments,” *Bulletin of the American Mathematical Society*.
- Rubin, D. B. (1974), “Estimating causal effects of treatments in randomized and nonrandomized studies,” *Journal of Educational Psychology*.
- Russo, D. (2016), “Simple Bayesian Algorithms for Best Arm Identification,” arXiv:1602.08448.
- Sauro, L. (2020), “Rapidly finding the Best Arm using Variance,” in *European Conference on Artificial Intelligence*.
- Schlag, K. H. (2007), “Eleven% Designing Randomized Experiments under Minimax Regret,” *Unpublished manuscript, European University Institute, Florence*.
- Shang, X., de Heide, R., Menard, P., Kaufmann, E., and Valko, M. (2020), “Fixed-confidence guarantees for Bayesian best-arm identification,” in *International Conference on Artificial Intelligence and Statistics*, vol. 108, pp. 1823–1832.
- Stoye, J. (2009), “Minimax regret treatment choice with finite samples,” *Journal of Econometrics*, 151, 70–81.
- (2012), “Minimax regret treatment choice with covariates or with limited validity of experiments,” *Journal of Econometrics*, 166, 138–156.
- Swaminathan, A. and Joachims, T. (2015), “Counterfactual Risk Minimization,” in *Proceedings of the 24th International Conference on World Wide Web*, Association for Computing Machinery, p. 939–941.
- Tabord-Meehan, M. (2022), “Stratification Trees for Adaptive Randomization in Randomized Controlled Trials,” *The Review of Economic Studies*.
- Tekin, C. and van der Schaar, M. (2015), “RELEAF: An Algorithm for Learning and Exploiting Relevance,” *IEEE Journal of Selected Topics in Signal Processing*.
- Thompson, W. R. (1933), “On the likelihood that one unknown probability exceeds another in view of the evidence of two samples,” *Biometrika*.
- Tsiatis, A. (2007), *Semiparametric Theory and Missing Data*, Springer Series in Statistics, Springer New York.
- van der Laan, M. J. (2008), “The Construction and Analysis of Adaptive Group Sequential Designs,” <https://biostats.bepress.com/ucbbiostat/paper232>.
- van der Vaart, A. (1991), “An Asymptotic Representation Theorem,” *International Statistical Review / Revue Internationale de Statistique*, 59, 97–121.
- (1998), *Asymptotic Statistics*, Cambridge Series in Statistical and Probabilistic Mathematics, Cambridge University Press.
- Viviano, D. (2022), “Experimental Design under Network Interference,” arXiv:2003.08421.
- Wager, S. and Xu, K. (2021), “Diffusion Asymptotics for Sequential Experiments,” arXiv:2101.09855.
- Wald, A. (1947), “Foundations of a General Theory of Sequential Decision Functions,” *Econometrica*, 15, 279–313.
- (1949), “Statistical Decision Functions,” *The Annals of Mathematical Statistics*, 20, 165 – 205.
- Wang, P.-A., Ariu, K., and Proutiere, A. (2023), “On Uniformly Optimal Algorithms for Best Arm Identification in Two-Armed Bandits with Fixed Budget,” arXiv:2308.12000.
- Yang, J. and Tan, V. (2022), “Minimax Optimal Fixed-Budget Best Arm Identification in Linear Bandits,” in *Advances in Neural Information Processing Systems (NeurIPS)*.
- Yang, Y. and Zhu, D. (2002), “Randomized Allocation with nonparametric estimation for a multi-armed bandit problem with covariates,” *Annals of Statistics*, 30, 100–121.
- Zhan, R., Ren, Z., Athey, S., and Zhou, Z. (2022), “Policy Learning with Adaptively Collected Data,” arXiv:2105.02344.
- Zheng, W. and van der Laan, M. J. (2011), “Cross-Validated Targeted Minimum-Loss-Based Estimation,” in *Targeted Learning: Causal Inference for Observational and Experimental Data*, Springer-Verlag New York, Springer Series in Statistics.

- Zhou, Z., Athey, S., and Wager, S. (2023), “Offline Multi-Action Policy Learning: Generalization and Optimization,” *Operations Research*, 71, 148–183.
- Zimmert, J. and Seldin, Y. (2021), “Tsallis-INF: An Optimal Algorithm for Stochastic and Adversarial Bandits,” *Journal of Machine Learning Research*, 22.

A Proof of the Minimax Lower Bounds (Theorems 3.4 and 3.5)

In this section, we establish the proofs for Theorems 3.4 and 3.5. Initially, we focus on proving Theorem 3.4, which presents a lower bound applicable for cases where $K \geq 2$. Following this, Section A.4 introduces a refined lower bound specifically for $K = 2$. While the initial lower bound from Theorem 3.4 is valid for $K = 2$, Theorem 3.5 offers a tighter lower bound.

A.1 Transportation Lemma

Let $f_P^a(y^a | s)$ be a density of Y^a conditional on $X = s$ under P . Let $\zeta_P(s)$ be a density of X under P .

Kaufmann et al. (2016) derives the following result based on a change-of-measure argument, which is the principal tool in our lower bound. Let us define a density of $(Y^1, Y^2, \dots, Y^K, X)$ under a bandit model $P \in \mathcal{P}$ as

$$p(y^1, y^2, \dots, y^K, s) = \prod_{a \in [K]} f_P^a(y^a | s) \zeta_P(s).$$

Between two bandit models $P, Q \in \mathcal{P}$, following the proof of Lemma 1 in Kaufmann et al. (2016), referred to as the *transportation lemma*, we define the log-likelihood ratio of a sequence of observations $(X_t, Y_t, A_t)_{t=1}^T$ under a given strategy as

$$L_T(P, Q) = \sum_{t=1}^T \sum_{a \in [K]} \mathbb{1}[A_t = a] \log \left(\frac{f_P^a(Y_t^a | X_t)}{f_Q^a(Y_t^a | X_t)} \right).$$

As discussed by Kaufmann et al. (2016), the transportation lemma immediately yields the following lemma.

Lemma A.1 (Lemma 1 and Remark 2 in Kaufmann et al. (2016)). *Suppose that for any two bandit model $P, Q \in \mathcal{P}$ with K treatment arms and for all $a \in [K]$, the distributions P^a and Q^a are mutually absolutely continuous, where P^a and Q^a are distributions of (Y^a, X) under P and Q , respectively. Then, for any $a \in [K]$ and $x \in \mathcal{X}$, any strategy satisfies*

$$|\mathbb{P}_P(\hat{a}(x) = a) - \mathbb{P}_Q(\hat{a}(x) = a)| \leq \sqrt{\frac{\mathbb{E}_P[L_T(P, Q)]}{2}}$$

A.2 Restricted Bandit Models

Fix any policy class Π with Natarajan dimension M . We can find a set of M points $\mathcal{S} := \{s_1, s_2, \dots, s_M\}$ that are shattered by Π .

Gaussian bandit models. We choose a specific ζ , a marginal distribution on X , and define a specific subclass \mathcal{P}^\dagger of a class of location-shift bandit models to derive a lower bound. Let ζ be a distribution on \mathcal{X} such that $\text{supp } \zeta = \mathcal{S}$ and $\Pr_\zeta(X = s) = 1/M$ for any $s \in \mathcal{S}$. We focus on Gaussian bandit models, where outcomes follow Gaussian distributions conditional on contexts. A subclass $\mathcal{P}^\dagger \subseteq \mathcal{P}$ is defined as follows:

$$\mathcal{P}^\dagger := \left\{ P \in \mathcal{P} : \forall a \in [K] \forall s \in \mathcal{S} \left[Y^a | X = s \sim \mathcal{N} \left(\mu^a(s), (\sigma^a(s))^2 \right), \mu^a(s) \in \mathbb{R} \right], \text{marg}_{\mathcal{X}}(P) = \zeta \right\},$$

where $\sigma^a : \mathcal{X} \rightarrow \mathbb{R}_+$ is given (introduced in Definition 2.1).

Best treatment arm. Suppose for now that, for each context $s \in \mathcal{S}$, the best treatment arm $d(s) \in \arg \max_{a \in [K]} \mathbb{E}_P[Y^a | X = s]$ is uniquely determined for each $s \in \mathcal{S}$ under the true distribution P .² Let $\mathbf{d} = (d(s))_{s \in \mathcal{S}}$ be a vector of best treatment arms for each context.

Let $\Delta(\mathbf{d}) = (\Delta_{d(s_1)}(s_1), \Delta_{d(s_2)}(s_2), \dots, \Delta_{d(s_M)}(s_M))$, where $\Delta_{d(s)}(s)$ is a K -dimensional vector such that its $d(s)$ -th element is $\Delta^{d(s)}(s)$ and the other elements are zero; that is,

$$\Delta_{d(s)}(s) = \left(\underbrace{0 \ 0 \ \dots \ 0}_{d(s)-1} \ \Delta^{d(s)}(s) \ \underbrace{0 \ \dots \ 0 \ 0}_{K-d(s)} \right)^\top$$

²Later, we focus on the situation in which nature always chooses a distribution that satisfies this property.

Null instance. Given constants $(m^s)_{s \in \mathcal{S}} \in \mathbb{R}^M$,³ for each $s \in \mathcal{S}$, let $P^{\sharp, s} \in \mathcal{P}^\dagger$ be an instance such that

$$\mu^1(P^{\sharp, s})(s) = \dots = \mu^K(P^{\sharp, s})(s) = m^s$$

holds.

Alternative hypothesis. The set of alternative hypotheses $\mathcal{Q}^\dagger \subseteq \mathcal{P}^\dagger$ is defined as follows:

$$\mathcal{Q}^\dagger := \left\{ P \in \mathcal{P}^\dagger : \begin{aligned} &\forall s \in \mathcal{S} \quad \mu^{d(s)}(P)(s) = m^s + \Delta^{d(s)}(s), \\ &\forall s \in \mathcal{S} \quad \forall b \in [K] \setminus \{d(s)\} \quad f_{P^{\sharp, s}}^b(y | s) = f_P^b(y | s), \\ &\forall s \in \mathcal{S} \quad \forall s' \in \mathcal{S} \setminus \{s\} \quad \forall a \in [K] \quad f_{P^{\sharp, s}}^a(y | s') = f_P^a(y | s'), \\ &\forall s \in \mathcal{S} \quad \Delta^{d(s)}(s) > 0, \quad d(s) \in [K], \quad m^s \in \mathbb{R} \end{aligned} \right\},$$

where $f_P(y | s)$ is a density of Y^a conditioned on $X = s$ under an instance P . We denote an element of \mathcal{Q}^\dagger by $Q_{\Delta(d)}$.⁴ Note that for any $Q_{\Delta(d)} \in \mathcal{Q}^\dagger$, arm $d(s)$ is indeed the optimal arm for context s , and we have $\mu^b(P)(s) = m^s$ for any $s \in \mathcal{S}$ and $b \in [K] \setminus \{d(s)\}$ by definition.

Change of measure. For any $Q_{\Delta(d)} \in \mathcal{Q}^\dagger$ and for each $s \in \mathcal{S}$, the following equation holds:

$$\begin{aligned} L_T(P^{\sharp, s}, Q_{\Delta(d)}) &= \sum_{t=1}^T \sum_{a \in [K]} \left\{ \mathbb{1}[A_t = a] \log \left(\frac{f_{P^{\sharp, s}}^a(Y_t^a | X_t)}{f_{Q_{\Delta(d)}}^a(Y_t^a | X_t)} \right) \right\} \\ &= \sum_{t=1}^T \sum_{a \in [K]} \sum_{s' \in \mathcal{S}} \left\{ \mathbb{1}[A_t = a] \log \left(\frac{f_{P^{\sharp, s'}}^a(Y_t^a | s')}{f_{Q_{\Delta(d)}}^a(Y_t^a | s')} \right) \right\} \mathbb{1}[X_t = s'] \\ &\stackrel{(i)}{=} \sum_{t=1}^T \sum_{a \in [K]} \left\{ \mathbb{1}[A_t = a] \log \left(\frac{f_{P^{\sharp, s}}^a(Y_t^a | s)}{f_{Q_{\Delta(d)}}^a(Y_t^a | s)} \right) \right\} \mathbb{1}[X_t = s] \\ &\stackrel{(ii)}{=} \sum_{t=1}^T \left\{ \mathbb{1}[A_t = d(s)] \log \left(\frac{f_{P^{\sharp, s}}^{d(s)}(Y_t^{d(s)} | s)}{f_{Q_{\Delta(d)}}^{d(s)}(Y_t^{d(s)} | s)} \right) \right\} \mathbb{1}[X_t = s]. \end{aligned}$$

In $\stackrel{(i)}{=}$, we used $\frac{f_{P^{\sharp, s'}}^a(Y_t^a | s')}{f_{Q_{\Delta(d)}}^a(Y_t^a | s')} = 1$ for $s' \neq s$ from the definition of $P^{\sharp, s}$. In $\stackrel{(ii)}{=}$, we used the assumption that for each $b \in [K] \setminus \{d(s)\}$, it holds that $f_{P^{\sharp, s}}^b(y | s) = f_{Q_{\Delta(d)}}^b(y | s)$.

Given a strategy, its *assignment ratio* $w : [K] \times \mathcal{X} \rightarrow [0, 1]$ under P is defined as follows:

$$w(a | x) := \mathbb{E}_P \left[\sum_{t=1}^T \mathbb{1}\{A_t = a\} \mid X = x \right].$$

For submodel \mathcal{Q}^\dagger , the following lemma holds.

Lemma A.2. For any $P^{\sharp, s} \in \mathcal{P}^\dagger$, $Q_{\Delta(d)} \in \mathcal{Q}^\dagger$, $s \in \mathcal{S}$, and $T \in \mathbb{N}$, the following equality holds:

$$\mathbb{E}_{P^{\sharp, s}} [L_T(P^{\sharp, s}, Q_{\Delta(d)})] = \frac{T}{2M} \frac{(\Delta^{d(s)}(s))^2}{\frac{(\sigma^{d(s)}(s))^2}{w(d(s)|s)}}.$$

Proof. We have

$$\mathbb{E}_{P^{\sharp, s}} [L_T(P^{\sharp, s}, Q_{\Delta(d)})] = \mathbb{E}_{P^{\sharp, s}} \left[\sum_{t=1}^T \left\{ \mathbb{1}[A_t = d(s)] \log \left(\frac{f_{P^{\sharp, s}}^{d(s)}(Y_t^{d(s)} | s)}{f_{Q_{\Delta(d)}}^{d(s)}(Y_t^{d(s)} | s)} \right) \right\} \mathbb{1}[X_t = s] \right]$$

³When $K \geq 3$, we can use any arbitrary $(m^s)_{s \in \mathcal{S}} \in \mathbb{R}^M$. When $K = 2$, we should choose $(m^s)_{s \in \mathcal{S}} \in \mathbb{R}^M$ carefully.

⁴Note that $Q_{\Delta(d)}$ also depends on $(m_s)_s$, but we omit it for notational convenience.

$$\begin{aligned}
&= \sum_{t=1}^T \mathbb{E}_{P^{\sharp,s}} \left[\mathbb{E}_{P^{\sharp,s}} \left[\left\{ \mathbb{1}[A_t = d(s)] \log \left(\frac{f_{P^{\sharp,s}}^{d(s)}(Y_t^{d(s)} | s)}{f_{Q_{\Delta(d)}}^{d(s)}(Y_t^{d(s)} | s)} \right) \right\} \mid X_t = s \right] \mathbb{1}[X_t = s] \right] \\
&= \sum_{t=1}^T \mathbb{E}_{P^{\sharp,s}} \left[w(d(s) | s) \mathbb{E}_{P^{\sharp,s}} \left[\log \left(\frac{f_{P^{\sharp,s}}^{d(s)}(Y_t^{d(s)} | s)}{f_{Q_{\Delta(d)}}^{d(s)}(Y_t^{d(s)} | s)} \right) \mid X_t = s \right] \mathbb{1}[X_t = s] \right] \\
&= Tw(d(s) | s) \frac{(\mu^{d(s)}(Q_{\Delta(d)})(s) - \mu^{d(s)}(P^{\sharp,s})(s))^2}{2(\sigma^{d(s)}(s))^2} / M \\
&= \frac{T}{2M} \frac{(\Delta^{d(s)}(s))^2}{(\sigma^{d(s)}(s))^2 / w(d(s) | s)}.
\end{aligned}$$

To show the second last equality, we use $\mathbb{E}_{P^{\sharp,s}} \left[\log \left(\frac{f_{P^{\sharp,s}}^{d(s)}(Y_t^{d(s)} | s)}{f_{Q_{\Delta(d)}}^{d(s)}(Y_t^{d(s)} | s)} \right) \mid X_t = s \right] = \frac{(\mu^{d(s)}(Q_{\Delta(d)})(s) - \mu^{d(s)}(P^{\sharp,s})(s))^2}{2(\sigma^{d(s)}(s))^2}$, which corresponds to a KL divergence between two Gaussian distributions. \square

A.3 Proof of Theorem 3.4: General Minimax Lower Bounds

We derive the lower bound the expected simple regret.

Proof of Theorem 3.4. Our proof below is built on the following line of reasoning: First, suppose that nature selects a true distribution in a two-step process: Initially, nature determines $(e^a(s))_{s,a} \in \mathbb{R}_+^{M \times K}$, ensuring that $\sum_a e^a(s) = 1$ for each state $s \in \mathcal{S}$. It then selects the optimal arms $d(s)$ with a probability of $e^{d(s)}(s)$ for each state s . Subsequently, nature chooses $Q_{\Delta(d)} \in \mathcal{Q}^\dagger$ for some $\Delta_{(d)}$ and $(m_s)_{s \in \mathcal{S}}$, which represents the true distribution.⁵ We then focus on the expected simple regret that the decision-maker could encounter under these strategies of nature, regardless of the strategy employed by the decision-maker. By construction, there must exist at least one distribution $Q_{\Delta(d)} \in \mathcal{Q}^\dagger \subseteq \mathcal{P}$ that attains this regret. Hence, this value serves as a lower bound for the regret.

Fix any strategy of the decision-maker. First, observe that, for each $P \in \mathcal{P}^\dagger$, the expected simple regret can be simplified as follows:

$$\begin{aligned}
R(P)(\pi) &= \mathbb{E}_P \left[\sum_{a \in [K]} \left(\mu^{a^*(P)(X)}(P)(X) - \mu^a(P)(X) \right) \mathbb{P}_P(\widehat{a}_T(X) = a \mid X) \right] \\
&= \sum_{s \in \mathcal{S}} \sum_{a \in [K]} \left(\mu^{a^*(P)(s)}(P)(s) - \mu^a(P)(s) \right) \mathbb{P}_P(\widehat{a}_T(X) = a \mid X = s) \mathbb{P}_P(X = s) \\
&= \sum_{s \in \mathcal{S}} \sum_{a \in [K]} \left(\mu^{a^*(P)(s)}(P)(s) - \mu^a(P)(s) \right) \mathbb{P}_P(\widehat{a}_T(s) = a) / M.
\end{aligned}$$

Next, we decompose the expected simple regret by using the definition of \mathcal{P}^\dagger : fix any $(e^a(s))_{s,a}$ and $\Delta_{(d)}$. Then, the decision-maker's expected simple regret experiences is:

$$\begin{aligned}
&\sum_{s \in \mathcal{S}} \sum_{d(s) \in [K]} e^{d(s)}(s) \sum_{b \in [K] \setminus \{d(s)\}} \left(\mu^{d(s)}(Q_{\Delta(d)})(s) - \mu^b(Q_{\Delta(d)})(s) \right) \mathbb{P}_{Q_{\Delta(d)}}(\widehat{a}_T(s) = b) / M \\
&= \sum_{s \in \mathcal{S}} \sum_{d(s) \in [K]} e^{d(s)}(s) \sum_{b \in [K] \setminus \{d(s)\}} \left(\mu^{d(s)}(Q_{\Delta(d)})(s) - \mu^{P^{\sharp,s}}(s) \right) \mathbb{P}_{Q_{\Delta(d)}}(\widehat{a}_T(s) = b) / M \\
&= \sum_{s \in \mathcal{S}} \sum_{d(s) \in [K]} e^{d(s)}(s) \left\{ \sum_{b \in [K] \setminus \{d(s)\}} \Delta^{d(s)}(s) \mathbb{P}_{Q_{\Delta(d)}}(\widehat{a}_T(s) = b) \right\} / M \\
&= \sum_{s \in \mathcal{S}} \sum_{d(s) \in [K]} e^{d(s)}(s) \left\{ \Delta^{d(s)}(s) \mathbb{P}_{Q_{\Delta(d)}}(\widehat{a}_T(s) \neq d(s)) \right\} / M
\end{aligned}$$

⁵Note that the choice of $(m_s)_{s \in \mathcal{S}}$ does not affect any objects in the proof.

$$= \sum_{s \in \mathcal{S}} \sum_{d(s) \in [K]} e^{d(s)}(s) \left\{ \Delta^{d(s)}(s) \left(1 - \mathbb{P}_{Q_{\Delta^{(d)}}}(\widehat{a}_T(s) = d(s)) \right) \right\} / M, \quad (4)$$

Note that, as we saw in the beginning of the proof, the regret lower bound is bounded from below by (4).

From Lemma A.1. and the definition of null consistent strategies, we have

$$\begin{aligned} (4) &= \sum_{s \in \mathcal{S}} \sum_{d(s) \in [K]} e^{d(s)}(s) \left\{ \Delta^{d(s)} \left(1 - \mathbb{P}_{P^{\sharp, s}}(\widehat{a}_T(s) = d(s)) + \mathbb{P}_{P^{\sharp, s}}(\widehat{a}_T(s) = d(s)) - \mathbb{P}_{Q_{\Delta^{(d)}}}(\widehat{a}_T(s) = d(s)) \right) \right\} / M \\ &\geq \sum_{s \in \mathcal{S}} \sum_{d(s) \in [K]} e^{d(s)}(s) \left\{ \Delta^{d(s)}(s) \left\{ 1 - \mathbb{P}_{P^{\sharp, s}}(\widehat{a}_T(s) = d(s)) - \sqrt{\frac{\mathbb{E}_{P^{\sharp, s}} \left[L_T^{d(s)}(P^{\sharp, s}, Q_{\Delta^{(d)}}) \right]}{2}} \right\} \right\} / M. \end{aligned} \quad (5)$$

From $\mathbb{P}_{P^{\sharp, s}}(\widehat{a}_T(s) = d(s)) = 1/K$ and Lemma A.2, we obtain

$$(5) = \sum_{s \in \mathcal{S}} \sum_{d(s) \in [K]} e^{d(s)}(s) \Delta^{d(s)}(s) \left\{ 1 - \frac{1}{K} - \sqrt{\frac{T(\Delta^{d(s)}(s))^2}{4M \frac{(\sigma^{d(s)}(s))^2}{w(d(s)|s)}}}} \right\} / M.$$

Let

$$\mathcal{E} := \left\{ (e^d(s))_{d,s} \in \mathbb{R}^{[K] \times \mathcal{S}} : e^d(s) \in [0, 1], \forall s \in \mathcal{S}, \sum_{d \in [K]} e^d(s) = 1 \right\},$$

and denote its typical element by e . By substituting $\Delta^{d(s)}(s) = \sqrt{M \frac{(\sigma^{d(s)}(s))^2}{w(d(s)|s)}} / 2T$, taking the infimum with respect to w , and taking the supremum with respect to $(e^a(s))_{a,s}$, we obtain the following regret lower bound:

$$\frac{1}{8} \frac{1}{\sqrt{MT}} \sum_{s \in \mathcal{S}} \sup_{e \in \mathcal{E}} \inf_{w \in \mathcal{W}} \sum_{d(s) \in [K]} e^{d(s)}(s) \left\{ \sqrt{\frac{(\sigma^{d(s)}(s))^2}{w(d(s)|s)}} \right\}. \quad (6)$$

Fix any $s \in \mathcal{S}$. Let $h(e, w) := \sum_{d \in [K]} e^d(s) \sqrt{\frac{(\sigma^{d(s)}(s))^2}{w(d(s)|s)}}$ and consider $\sup_{e \in \mathcal{E}} \inf_{w \in \mathcal{W}} h(e, w)$. Note that h is concave in e , convex in w , and continuous in (e, w) . Denote the closure of \mathcal{W} by $\overline{\mathcal{W}}$. First, observe that

$$\sup_{e \in \mathcal{E}} \inf_{w \in \mathcal{W}} h(e, w) \geq \sup_{e \in \mathcal{E}} \inf_{w \in \overline{\mathcal{W}}} h(e, w).$$

Since $\overline{\mathcal{W}}$ is compact and $h(e, \cdot)$ is continuous for each e , we have

$$\sup_{e \in \mathcal{E}} \inf_{w \in \overline{\mathcal{W}}} h(e, w) = \sup_{e \in \mathcal{E}} \min_{w \in \overline{\mathcal{W}}} h(e, w).$$

By Berge's Maximum Theorem, $\min_{w \in \overline{\mathcal{W}}} h(e, w)$ is continuous in w . Since \mathcal{E} is compact, we have

$$\sup_{e \in \mathcal{E}} \min_{w \in \overline{\mathcal{W}}} h(e, w) = \max_{e \in \mathcal{E}} \min_{w \in \overline{\mathcal{W}}} h(e, w).$$

Since \mathcal{E} and $\overline{\mathcal{W}}$ are both compact and convex and h is concave-convex, by the minimax theorem, we have

$$\max_{e \in \mathcal{E}} \min_{w \in \overline{\mathcal{W}}} h(e, w) = \min_{w \in \overline{\mathcal{W}}} \max_{e \in \mathcal{E}} h(e, w).$$

Combining these results, we have

$$\sup_{e \in \mathcal{E}} \inf_{w \in \mathcal{W}} h(e, w) \geq \min_{w \in \overline{\mathcal{W}}} \max_{e \in \mathcal{E}} h(e, w).$$

Let us consider $\max_{e \in \mathcal{E}} h(e, \mathbf{w})$ for a fixed \mathbf{w} . At the optimum, we have $e^d(s) = 1$ iff $d \in \arg \max_{d \in [K]} \sqrt{\frac{(\sigma^d(s))^2}{w(d|s)}}$. Thus, we have

$$\min_{\mathbf{w} \in \overline{\mathcal{W}}} \max_{e \in \mathcal{E}} h(e, \mathbf{w}) = \min_{\mathbf{w} \in \overline{\mathcal{W}}} \max_{d \in [K]} \sqrt{\frac{(\sigma^d(s))^2}{w(d|s)}}. \quad (7)$$

For each s , we consider the following constrained optimization:

$$\begin{aligned} \min_{R \geq 0, \mathbf{w} \in \overline{\mathcal{W}}} \quad & R \\ \text{s.t.} \quad & R \geq \sqrt{\frac{(\sigma^d(s))^2}{w(d|s)}} \quad \forall d \in [K] \end{aligned}$$

Note that if (R^*, \mathbf{w}^*) is the optimal solution to this problem, \mathbf{w}^* is also the optimal solution to (7). After some algebra, we can show that

$$w^*(d|s) = \frac{(\sigma^d(s))^2}{\sum_{a \in [K]} (\sigma^a(s))^2}$$

for each $d \in [K]$.

Thus, we have

$$(7) = \sqrt{\sum_{a \in [K]} (\sigma^a(s))^2},$$

and hence

$$\begin{aligned} (6) &\geq \frac{1}{8} \frac{1}{\sqrt{TM}} \sum_{s \in \mathcal{S}} \sqrt{\sum_{a \in [K]} (\sigma^a(s))^2} = \frac{1}{8} \frac{1}{\sqrt{T}} \sum_{s \in \mathcal{S}} \sqrt{M \sum_{a \in [K]} (\sigma^a(s))^2} \frac{1}{M} \\ &= \frac{1}{8} \frac{1}{\sqrt{T}} \mathbb{E}_{X \sim \zeta} \left[\sqrt{M \sum_{a \in [K]} (\sigma^a(s))^2} \right]. \end{aligned} \quad (8)$$

The last equality follows since we choose ξ so that ξ puts equal probabilities on \mathcal{S} . This implies that, for any strategy of the decision-maker, there exists a distribution in \mathcal{P} under which the simple regret is lower bounded by (8). \square

Although this lower bound is applicable to a case with $K = 2$, we can tighten the lower bound by changing the definition of the parametric submodel.

A.4 Refined Minimax Lower Bounds for Two-armed Bandits (Proof of Theorem 3.5)

When $K = 2$, we can derive a tighter lower bound. As in previous sections, we choose $\xi \in \mathcal{M}(\mathcal{X})$ such that $\Pr_\xi(X = s) = 1/M$ for each $s \in \mathcal{S}$.

Define \mathbf{d} as before (see Section A.2.) Let $\mathbf{\Delta} = (\Delta(s_1), \Delta(s_2), \dots, \Delta(s_M))$, where $\Delta(s) \in (0, \infty)$ is a scalar.

Null instance. Given constants $(m^s)_{s \in \mathcal{S}} \in \mathbb{R}^M$, for each $s \in \mathcal{S}$, let $P^{\#,s} \in \mathcal{P}^\dagger$ be an instance such that

$$\mu^1(P^{\#,s})(s) = \mu^2(P^{\#,s})(s) = m^s.$$

Note that, unlike the $K \geq 2$ case where the value of $(m_s)_s$ does not affect the derived lower bound, we will optimize $(m^s)_{s \in \mathcal{S}}$ to derive a refined lower bound for the $K = 2$ case.

Alternative hypothesis. Define the set of alternative hypotheses $\mathcal{Q}^\dagger \subseteq \mathcal{P}^\dagger \subseteq \mathcal{P}$ as follows:

$$\mathcal{Q}^\dagger := \left\{ P \in \mathcal{P}^\dagger: \begin{aligned} &\forall s \in \mathcal{S} \quad \mu^{d(s)}(P)(s) = m^s + \Delta(s) \\ &\forall s \in \mathcal{S} \quad \forall b \in [K] \setminus \{d(s)\} \quad f_{P^{\sharp,s}}^b(y | s) = f_P^b(y | s) \\ &\forall s \in \mathcal{S} \quad \forall s' \in \mathcal{S} \setminus \{s\} \quad \forall a \in [K] \quad f_{P^{\sharp,s}}^a(y | s') = f_P^a(y | s'), \\ &\forall s \in \mathcal{S} \quad \Delta(s) > 0, d(s) \in [K], m_s \in \mathbb{R} \end{aligned} \right\}$$

where $f_P(y | s)$ is a density of Y^a conditioned on $X = s$ under an instance P . We denote a typical element of \mathcal{Q}^\dagger by Q_Δ .⁶ Note that for $Q_\Delta \in \mathcal{Q}^\dagger$, $\forall s \in \mathcal{S} \quad \forall b \in [K] \setminus \{d(s)\} \quad \mu^b(P)(s) = m^s$ holds by definition.

Change of measure. For any $Q_{\Delta(a)} \in \mathcal{Q}^\dagger$ and $s \in \mathcal{S}$, the following equation holds:

$$\begin{aligned} L_T(P^{\sharp,s}, Q_\Delta) &= \sum_{t=1}^T \sum_{a \in [K]} \left\{ \mathbb{1}[A_t = a] \log \left(\frac{f_{P^{\sharp,s}}^a(Y_t^a | X_t)}{f_{Q_\Delta}^a(Y_t^a | X_t)} \right) \right\} \\ &= \sum_{t=1}^T \sum_{a \in [K]} \sum_{s' \in \mathcal{S}} \left\{ \mathbb{1}[A_t = a] \log \left(\frac{f_{P^{\sharp,s'}}^a(Y_t^a | s')}{f_{Q_\Delta}^a(Y_t^a | s')} \right) \right\} \mathbb{1}[X_t = s'] \\ &\stackrel{(i)}{=} \sum_{t=1}^T \sum_{a \in [K]} \left\{ \mathbb{1}[A_t = a] \log \left(\frac{f_{P^{\sharp,s}}^a(Y_t^a | s)}{f_{Q_\Delta}^a(Y_t^a | s)} \right) \right\} \mathbb{1}[X_t = s]. \end{aligned}$$

In ⁽ⁱ⁾, we used $\frac{f_{P^{\sharp,s'}}^a(Y_t^a | s')}{f_{Q_\Delta}^a(Y_t^a | s')} = 1$ for $s' \neq s$ from the definition of $P^{\sharp,s}$.

Proof of Theorem 3.5. By the same argument as in Section A.3, we have the following lower bound for the simple regret (equation (5)):

$$\sum_{s \in \mathcal{S}} \sum_{d(s) \in [K]} e^{d(s)}(s) \left\{ \Delta^{d(s)}(s) \left\{ 1 - \mathbb{P}_{P^{\sharp,s}}(\widehat{a}_T(s) = d(s)) - \sqrt{\frac{\mathbb{E}_{P^{\sharp,s}} [L_T^{d(s)}(P^{\sharp,s}, Q_{\Delta(a)})]}{2}} \right\} \right\} / M. \quad (9)$$

By the same argument as in the proof of Lemma A.2, we have

$$\mathbb{E}_{P^{\sharp,s}} [L_T(P^{\sharp,s}, Q_\Delta)] = \frac{T}{2M} \left\{ w(1 | s) \frac{(\mu^1(Q_\Delta)(s) - m^s)^2}{(\sigma^1(s))^2} + w(2 | s) \frac{(\mu^2(Q_\Delta)(s) - m^s)^2}{(\sigma^2(s))^2} \right\}.$$

We consider the following optimization problem with respect to m^s :

$$\min_{m_s \in \mathbb{R}} \left\{ w(1 | s) \frac{(\mu^1(Q_\Delta)(s) - m^s)^2}{(\sigma^1(s))^2} + w(2 | s) \frac{(\mu^2(Q_\Delta)(s) - m^s)^2}{(\sigma^2(s))^2} \right\}.$$

The solution is

$$m^s = \frac{c^1 \mu^1(Q_{\Delta_1})(s) + c^2 \mu^2(Q_{\Delta_2})(s)}{c^1 + c^2},$$

where

$$c^a = \frac{w(a | s)}{(\sigma^a(s))^2},$$

⁶Note that Q_Δ also depends on $(m_s)_s$, but we omit it for notational convenience.

and, the optimal value is

$$(\Delta(s))^2 \left[\frac{(\sigma^1(s))^2}{w(1|s)} + \frac{(\sigma^2(s))^2}{w(2|s)} \right]^{-1}. \quad (10)$$

Since $\mathbb{P}_{P^{\#,s}}(\hat{a}_T(s) = d(s)) = 1/K$, by (9) and (10), we have the following regret lower bound:

$$\inf_{w \in \mathcal{W}} \sup_{\substack{\Delta(s) \in (0, \infty) \\ \text{for } s \in \mathcal{S}}} \sum_{s \in \mathcal{S}} \sum_{d(s) \in [K]} e^{d(s)} \Delta(s) \left\{ 1 - \frac{1}{K} - \sqrt{\frac{T \Delta^2(s)}{4M \left\{ \frac{(\sigma^1(s))^2}{w(1|s)} + \frac{(\sigma^2(s))^2}{w(2|s)} \right\}}} \right\} / M.$$

By substituting $\Delta(s) = \sqrt{M \left\{ \frac{(\sigma^1(s))^2}{w(1|s)} + \frac{(\sigma^2(s))^2}{w(2|s)} \right\}} / 2T$, we obtain the following regret lower bound:

$$\begin{aligned} & \frac{1}{8} \frac{1}{\sqrt{MT}} \sum_{s \in \mathcal{S}} \sup_{e \in \mathcal{E}} \inf_{w \in \mathcal{W}} \sum_{d(s) \in [K]} e^{d(s)} \left\{ \sqrt{\frac{(\sigma^1(s))^2}{w(1|s)} + \frac{(\sigma^2(s))^2}{w(2|s)}} \right\} \\ &= \frac{1}{8} \frac{1}{\sqrt{MT}} \sum_{s \in \mathcal{S}} \sup_{e \in \mathcal{E}} \inf_{w \in \mathcal{W}} \left\{ \sqrt{\frac{(\sigma^1(s))^2}{w(1|s)} + \frac{(\sigma^2(s))^2}{w(2|s)}} \right\} \sum_{d(s) \in [K]} e^{d(s)} \\ &= \frac{1}{8} \frac{1}{\sqrt{MT}} \sum_{s \in \mathcal{S}} \inf_{w \in \mathcal{W}} \left\{ \sqrt{\frac{(\sigma^1(s))^2}{w(1|s)} + \frac{(\sigma^2(s))^2}{w(2|s)}} \right\}. \end{aligned}$$

Consider the following optimization problem:

$$\min_{w \in \mathcal{W}} \sqrt{\frac{(\sigma^1(s))^2}{w(1|s)} + \frac{(\sigma^2(s))^2}{w(2|s)}}.$$

The solution is

$$w(a|s) = \frac{\sigma^a(s)}{\sigma^1(s) + \sigma^2(s)},$$

and the optimal value is

$$\sqrt{(\sigma^1(s) + \sigma^2(s))^2}.$$

Therefore, we obtain the following lower bound:

$$\frac{1}{8} \frac{1}{\sqrt{MT}} \sum_{s \in \mathcal{S}} \sqrt{(\sigma^1(s) + \sigma^2(s))^2} = \frac{1}{8} \frac{1}{\sqrt{T}} \mathbb{E}_{X \sim \xi} \sqrt{M(\sigma^1(X) + \sigma^2(X))^2}.$$

This completes the proof. □

B Proof of Lemma 5.6

In this section, we show

$$\sqrt{T} \hat{Q}_T(\pi) = \sqrt{T} \frac{1}{T} \sum_{t=1}^T \sum_{a \in [K]} \pi(a|X_t) \hat{\Gamma}_t^a = \sqrt{T} \frac{1}{T} \sum_{t=1}^T \sum_{a \in [K]} \pi(a|X_t) \Gamma^{*a}(Y_t, \xi_t, X_t) + o_P(1). \quad (11)$$

Here, note that

$$\begin{aligned}
& \frac{1}{\sqrt{T}} \sum_{t=1}^T \pi(a | X_t) \widehat{\Gamma}_t^a(Y_t^a, \xi_t, X_t) \\
&= \frac{1}{\sqrt{T}} \sum_{t=1}^T \pi(a | X_t) \left\{ \frac{\mathbb{1} \left[\sum_{b=0}^{a-1} \widehat{w}_t(b | X_t) \leq \xi_t \leq \sum_{b=0}^a \widehat{w}_t(b | X_t) \right] (Y_t^a - \widehat{\mu}_t^a(X_t))}{\widehat{w}_t(a | X_t)} + \widehat{\mu}_t^a(X_t) \right\} \\
&= \frac{1}{\sqrt{T}} \sum_{t=1}^T \pi(a | X_t) \left\{ \frac{\mathbb{1} \left[\sum_{b=0}^{a-1} w^*(b | X_t) \leq \xi_t \leq \sum_{b=0}^a w^*(b | X_t) \right] (Y_t^a - \mu^a(P)(X_t))}{w^*(a | X_t)} + \mu^a(P)(X_t) \right\} + o_P(1)
\end{aligned} \tag{12}$$

holds. Therefore, to show (11), we show that the equality from the second to the third lines holds.

Proof. Let us define

$$\begin{aligned}
& G_t(Y_t^a, X_t, \xi_t; \{\widehat{w}_t(b | X_t)\}_{b \in [K]}, \widehat{\mu}_T^a(X_t)) \\
&:= \pi(a | X_t) \left\{ \frac{\mathbb{1} \left[\sum_{b=0}^{a-1} \widehat{w}_t(b | X_t) \leq \xi_t \leq \sum_{b=0}^a \widehat{w}_t(b | X_t) \right] (Y_t^a - \widehat{\mu}_t^a(X_t))}{\widehat{w}_t(a | X_t)} + \widehat{\mu}_t^a(X_t) \right\}.
\end{aligned}$$

Then, we obtain

$$\begin{aligned}
& G_t(Y_t^a, X_t, \xi_t; \{\widehat{w}_t(b | X_t)\}_{b \in [K]}, \widehat{\mu}_T^a(X_t)) \\
&= G_t(Y_t^a, X_t, \xi_t; \{w^*(b | X_t)\}_{b \in [K]}, \mu^a(P)(X_t)) \\
&\quad - G_t(Y_t^a, X_t, \xi_t; \{w^*(b | X_t)\}_{b \in [K]}, \mu^a(P)(X_t)) + G_t(Y_t^a, X_t, \xi_t; \{\widehat{w}_t(b | X_t)\}_{b \in [K]}, \widehat{\mu}_T^a(X_t)) \\
&= G_t(Y_t^a, X_t, \xi_t; \{w^*(b | X_t)\}_{b \in [K]}, \mu^a(P)(X_t)) + B_t,
\end{aligned}$$

where

$$B_t := G_t(Y_t^a, X_t, \xi_t; \{\widehat{w}_t(b | X_t)\}_{b \in [K]}, \widehat{\mu}_T^a(X_t)) - G_t(Y_t^a, X_t, \xi_t; \{w^*(b | X_t)\}_{b \in [K]}, \mu^a(P)(X_t)).$$

To show (12), we consider showing $\frac{1}{\sqrt{T}} \sum_{t=1}^T B_t \rightarrow 0$ as $T \rightarrow \infty$ in probability.

We show $\frac{1}{\sqrt{T}} \sum_{t=1}^T B_t \rightarrow 0$ as $T \rightarrow \infty$ in probability by using the properties of martingales. First, we have

$$\begin{aligned}
& \mathbb{E}[B_t | X_t, \mathcal{F}_{t-1}] \\
&= \mathbb{E} \left[\pi(a | X_t) \left\{ \frac{\mathbb{1} \left[\sum_{b=0}^{a-1} \widehat{w}_t(b | X_t) \leq \xi_t \leq \sum_{b=0}^a \widehat{w}_t(b | X_t) \right] (Y_t^a - \widehat{\mu}_t^a(X_t))}{\widehat{w}_t(a | X_t)} + \widehat{\mu}_t^a(X_t) \right\} \middle| X_t, \mathcal{F}_{t-1} \right] \\
&\quad - \mathbb{E} \left[\pi(a | X_t) \left\{ \frac{\mathbb{1} \left[\sum_{b=0}^{a-1} w^*(b | X_t) \leq \xi_t \leq \sum_{b=0}^a w^*(b | X_t) \right] (Y_t^a - \mu^a(P)(X_t))}{w^*(a | X_t)} + \mu^a(P)(X_t) \right\} \middle| X_t, \mathcal{F}_{t-1} \right] \\
&= \mathbb{E} \left[\pi(a | X_t) \left\{ \frac{\widehat{w}_t(a | X_t)(Y_t^a - \widehat{\mu}_t^a(X_t))}{\widehat{w}_t(a | X_t)} + \widehat{\mu}_t^a(X_t) \right\} \middle| X_t, \mathcal{F}_{t-1} \right] \\
&\quad - \mathbb{E} \left[\pi(a | X_t) \left\{ \frac{w^*(a | X_t)(Y_t^a - \mu^a(P)(X_t))}{w^*(a | X_t)} + \mu^a(P)(X_t) \right\} \middle| X_t, \mathcal{F}_{t-1} \right] \\
&= \pi(a | X_t) \mu^a(P)(X_t) - \pi(a | X_t) \mu^a(P)(X_t) \\
&= 0.
\end{aligned}$$

This result implies that $\{B_t\}_{t=1}^T$ is a martingale difference sequence (MDS) because $\mathbb{E}[B_t | X_t, \mathcal{F}_{t-1}] = \mathbb{E}[\mathbb{E}[B_t | X_t, \mathcal{F}_{t-1}]] = 0$ holds.

Besides, from $\tilde{w}_t(a | X_t) - w^*(a | X_t) \xrightarrow{\text{a.s.}} 0$ and $\hat{\mu}_t^a(X_t) - \mu^a(P)(X_t) \xrightarrow{\text{a.s.}} 0$, $\mathbb{E}[B_t^2 | X_t, \mathcal{F}_{t-1}]$ also converges to zero almost surely; that is,

$$\begin{aligned} & \mathbb{E}[B_t^2 | X_t, \mathcal{F}_{t-1}] \\ &= \mathbb{E} \left[\pi^2(a | X_t) \left(\left\{ \frac{\mathbb{1} \left[\sum_{b=0}^{a-1} \hat{w}_t(b | X_t) \leq \xi_t \leq \sum_{b=0}^a \hat{w}_t(b | X_t) \right] (Y_t^a - \hat{\mu}_t^a(X_t))}{\hat{w}_t(a | X_t)} + \hat{\mu}_t^a(X_t) \right\} \right. \right. \\ & \quad \left. \left. - \left\{ \frac{\mathbb{1} \left[\sum_{b=0}^{a-1} w^*(b | X_t) \leq \xi_t \leq \sum_{b=0}^a w^*(b | X_t) \right] (Y_t^a - \mu^a(P)(X_t))}{w^*(a | X_t)} + \mu^a(P)(X_t) \right\} \right)^2 \middle| X_t, \mathcal{F}_{t-1} \right] \\ & \xrightarrow{\text{a.s.}} 0. \end{aligned}$$

This is because from $\hat{\mu}_t^a(X_t) - \mu^a(P)(X_t) \xrightarrow{\text{a.s.}} 0$,

$$\begin{aligned} & \mathbb{E} \left[\pi^2(a | X_t) \left(\left\{ \frac{\mathbb{1} \left[\sum_{b=0}^{a-1} \hat{w}_t(b | X_t) \leq \xi_t \leq \sum_{b=0}^a \hat{w}_t(b | X_t) \right] (Y_t^a - \hat{\mu}_t^a(X_t))}{\hat{w}_t(a | X_t)} + \hat{\mu}_t^a(X_t) \right\} \right. \right. \\ & \quad \left. \left. - \left\{ \frac{\mathbb{1} \left[\sum_{b=0}^{a-1} \hat{w}_t(b | X_t) \leq \xi_t \leq \sum_{b=0}^a \hat{w}_t(b | X_t) \right] (Y_t^a - \mu^a(P)(X_t))}{w^*(a | X_t)} + \mu^a(P)(X_t) \right\} \right)^2 \middle| X_t, \mathcal{F}_{t-1} \right] \\ & \xrightarrow{\text{a.s.}} 0. \end{aligned}$$

holds. Additionally, from $\tilde{w}_t(a | X_t) - w^*(a | X_t) \xrightarrow{\text{a.s.}} 0$, for any $\varepsilon > 0$, there exists $T(\varepsilon) > 0$ such that for any $t > T(\varepsilon)$, $|\tilde{w}_t(a | x) - w^*(a | x)| < \varepsilon$ holds for all $a \in [K]$ with probability one; that is,

$$\begin{aligned} & \mathbb{E} \left[\pi^2(a | X_t) \left(\left\{ \frac{\mathbb{1} \left[\sum_{b=0}^{a-1} \hat{w}_t(b | X_t) \leq \xi_t \leq \sum_{b=0}^a \hat{w}_t(b | X_t) \right] (Y_t^a - \mu^a(P)(X_t))}{\hat{w}_t(a | X_t)} + \mu^a(P)(X_t) \right\} \right. \right. \\ & \quad \left. \left. - \left\{ \frac{\mathbb{1} \left[\sum_{b=0}^{a-1} w^*(b | X_t) \leq \xi_t \leq \sum_{b=0}^a w^*(b | X_t) \right] (Y_t^a - \mu^a(P)(X_t))}{\hat{w}_t(a | X_t)} + \mu^a(P)(X_t) \right\} \right)^2 \middle| X_t, \mathcal{F}_{t-1} \right] \\ & \leq \mathbb{E} \left[\left\{ \frac{\mathbb{1} \left[\min \left\{ \sum_{b=0}^{a-1} \hat{w}_t(b | X_t), \sum_{b=0}^{a-1} w^*(b | X_t) \right\} \leq \xi_t \leq \max \left\{ \sum_{b=0}^{a-1} \hat{w}_t(b | X_t), \sum_{b=0}^{a-1} w^*(b | X_t) \right\} \right]}{\hat{w}_t^2(a | X_t)} \right\} \right. \\ & \quad \left. \times \pi^2(a | X_t) (Y_t^a - \mu^a(P)(X_t))^2 \middle| X_t, \mathcal{F}_{t-1} \right] \\ & + \mathbb{E} \left[\left\{ \frac{\mathbb{1} \left[\min \left\{ \sum_{b=0}^a \hat{w}_t(b | X_t), \sum_{b=0}^a w^*(b | X_t) \right\} \leq \xi_t \leq \max \left\{ \sum_{b=0}^a \hat{w}_t(b | X_t), \sum_{b=0}^a w^*(b | X_t) \right\} \right]}{\hat{w}_t^2(a | X_t)} \right\} \right. \\ & \quad \left. \times \pi^2(a | X_t) (Y_t^a - \mu^a(P)(X_t))^2 \middle| X_t, \mathcal{F}_{t-1} \right] \\ & \leq \mathbb{E} \left[\left\{ \frac{\mathbb{1} \left[\sum_{b=0}^{a-1} \left\{ w^*(b | X_t) - \varepsilon \right\} \leq \xi_t \leq \sum_{b=0}^{a-1} \left\{ w^*(b | X_t) + \varepsilon \right\} \right]}{\hat{w}_t^2(a | X_t)} \right\} \pi^2(a | X_t) (Y_t^a - \mu^a(P)(X_t))^2 \middle| X_t, \mathcal{F}_{t-1} \right] \\ & + \mathbb{E} \left[\left\{ \frac{\mathbb{1} \left[\sum_{b=0}^a \left\{ w^*(b | X_t) - \varepsilon \right\} \leq \xi_t \leq \sum_{b=0}^a \left\{ w^*(b | X_t) + \varepsilon \right\} \right]}{\hat{w}_t^2(a | X_t)} \right\} \pi^2(a | X_t) (Y_t^a - \mu^a(P)(X_t))^2 \middle| X_t, \mathcal{F}_{t-1} \right] \end{aligned}$$

$$\leq \frac{2(a-1)\varepsilon}{\widehat{w}_t^2(a|X_t)} \mathbb{E} \left[\pi^2(a|X_t) \left(Y_t^a - \mu^a(P)(X_t) \right)^2 \mid \mathcal{F}_{t-1} \right] + \frac{2a\varepsilon}{\widehat{w}_t^2(a|X_t)} \mathbb{E} \left[\pi^2(a|X_t) \left(Y_t^a - \mu^a(P)(X_t) \right)^2 \mid X_t, \mathcal{F}_{t-1} \right]$$

$$\leq C\varepsilon$$

holds with probability one, where $C > 0$ is a constant independent from t . This implies that for any $\varepsilon' > 0$, there exists $T'(\varepsilon') > 0$ such that for any $t > T'(\varepsilon')$,

$$\left| \mathbb{E} \left[\pi^2(a|X_t) \left(\frac{\mathbb{1} \left[\sum_{b=0}^{a-1} \widehat{w}_t(b|X_t) \leq \xi_t \leq \sum_{b=0}^a \widehat{w}_t(b|X_t) \right] (Y_t^a - \mu^a(P)(X_t))}{\widehat{w}_t(a|X_t)} + \mu^a(P)(X_t) \right) \right. \right. \\ \left. \left. - \left(\frac{\mathbb{1} \left[\sum_{b=0}^{a-1} w^*(b|X_t) \leq \xi_t \leq \sum_{b=0}^a w^*(b|X_t) \right] (Y_t^a - \mu^a(P)(X_t))}{\widehat{w}_t(a|X_t)} + \mu^a(P)(X_t) \right) \right)^2 \mid X_t, \mathcal{F}_{t-1} \right] \right| < \varepsilon'$$

with probability one. We also have

$$\mathbb{E} \left[\pi^2(a|X_t) \left(\frac{\mathbb{1} \left[\sum_{b=0}^{a-1} w^*(b|X_t) \leq \xi_t \leq \sum_{b=0}^a w^*(b|X_t) \right] (Y_t^a - \mu^a(P)(X_t))}{\widehat{w}_t(a|X_t)} + \mu^a(P)(X_t) \right) \right. \\ \left. - \left(\frac{\mathbb{1} \left[\sum_{b=0}^{a-1} w^*(b|X_t) \leq \xi_t \leq \sum_{b=0}^a w^*(b|X_t) \right] (Y_t^a - \mu^a(P)(X_t))}{w^*(a|X_t)} + \mu^a(P)(X_t) \right) \right)^2 \mid X_t, \mathcal{F}_{t-1} \right] \\ \xrightarrow{\text{a.s.}} 0.$$

Thus, $\mathbb{E}[B_t^2 \mid X_t, \mathcal{F}_{t-1}] \xrightarrow{\text{a.s.}} 0$ holds.

Based on these results, from Chebyshev's inequality, $\frac{1}{\sqrt{T}} \sum_{t=1}^T B_t$ converges to zero in probability. This is because for any $v > 0$,

$$\mathbb{P}_P \left(\left| \frac{1}{\sqrt{T}} \sum_{t=1}^T B_t - \mathbb{E} \left[\frac{1}{\sqrt{T}} \sum_{t=1}^T B_t \right] \right| \geq v \right) = \mathbb{P}_P \left(\left| \frac{1}{\sqrt{T}} \sum_{t=1}^T B_t \right| \geq v \right) \leq \frac{\text{Var}_P \left(\frac{1}{\sqrt{T}} \sum_{t=1}^T B_t \right)}{v^2} \quad (13)$$

Because B_t is an MDS, the covariance between B_t and B_s for $t \neq s$ is zero; that is, if $s < t$, $\text{Cov}(B_t, B_s) = \mathbb{E}[B_t B_s] = \mathbb{E}[B_s \mathbb{E}[B_t \mid \mathcal{F}_{t-1}]] = 0$.

Therefore, we can show $\frac{1}{\sqrt{T}} \sum_{t=1}^T B_t \xrightarrow{P} 0$ by showing

$$\text{Var}_P \left(\frac{1}{\sqrt{T}} \sum_{t=1}^T B_t \right) = \frac{1}{T} \sum_{t=1}^T \text{Var}(B_t) \rightarrow 0, \quad (14)$$

as $T \rightarrow \infty$, where we used that the covariance between B_t and B_s for $t \neq s$ is zero.

To show (14), we show

$$\frac{1}{T} \sum_{t=1}^T \text{Var}_P(B_t \mid X_t, \mathcal{F}_{t-1}) = \frac{1}{T} \sum_{t=1}^T \mathbb{E}_P[B_t^2 \mid X_t, \mathcal{F}_{t-1}] \xrightarrow{P} 0,$$

by using $\mathbb{E}[B_t^2 \mid X_t, \mathcal{F}_{t-1}] \xrightarrow{\text{a.s.}} 0$.

Let u_t be $u_t = \mathbb{E}_P[B_t^2 \mid X_t, \mathcal{F}_{t-1}]$. Fix some positive $\epsilon > 0$ and $\delta > 0$. Almost-sure convergence of u_t to zero as $t \rightarrow \infty$ implies that we can find a large enough $t(\epsilon)$ such that $|u_t| < \epsilon$ for all $t \geq t(\epsilon)$ with probability at least $1 - \delta$. Let $\mathcal{E}(\epsilon)$ denote the event in which this happens; that is, $\mathcal{E}(\epsilon) = \{|u_t| < \epsilon \quad \forall t \geq t(\epsilon)\}$. Under this event, for $T > t(\epsilon)$,

$$\sum_{t=1}^T |u_t| \leq \sum_{t=1}^{t(\epsilon)} C + \sum_{t=t(\epsilon)+1}^T \epsilon = t(\epsilon)C + T\epsilon.$$

Therefore, we obtain

$$\begin{aligned} \mathbb{P}\left(\frac{1}{T}\sum_{t=1}^T|u_t| > 2\epsilon\right) &= \mathbb{P}\left(\left\{\frac{1}{T}\sum_{t=1}^T|u_t| > 2\epsilon\right\} \cap \mathcal{E}(\epsilon)\right) + \mathbb{P}\left(\left\{\frac{1}{T}\sum_{t=1}^T|u_t| > 2\epsilon\right\} \cap \mathcal{E}^c(\epsilon)\right) \\ &\leq \mathbb{P}\left(\frac{t(\epsilon)}{T}C + \epsilon > 2\epsilon\right) + \mathbb{P}(\mathcal{E}^c(\epsilon)) = \mathbb{P}\left(\frac{t(\epsilon)}{T}C > \epsilon\right) + \mathbb{P}(\mathcal{E}^c(\epsilon)). \end{aligned}$$

Letting $T \rightarrow \infty$, for arbitrarily small $\delta > 0$, $\frac{1}{T}\sum_{t=1}^T \mathbb{E}_P[B_t^2 | X_t, \mathcal{F}_{t-1}] \xrightarrow{P} 0$ as $T \rightarrow \infty$ holds.

Then, from the dominated convergence theorem and the boundedness of B_t^2 , $\frac{1}{T}\sum_{t=1}^T \text{Var}_P(B_t) \xrightarrow{P} 0$ as $T \rightarrow \infty$ holds⁷.

Therefore, from (13), $\frac{1}{\sqrt{T}}\sum_{t=1}^T B_t \xrightarrow{P} 0$ holds, which implies

$$\begin{aligned} &\frac{1}{\sqrt{T}}\sum_{t=1}^T \pi(a | X_t) \widehat{\Gamma}_t^a(Y_t^a, \xi_t, X_t) \\ &= \frac{1}{\sqrt{T}}\sum_{t=1}^T \pi(a | X_t) \left\{ \frac{\mathbb{1}\left[\sum_{b=0}^{a-1} w^*(b | X_t) \leq \xi_t \leq \sum_{b=0}^a w^*(a | X_t)\right] (Y_t^a - \mu^a(P)(X_t))}{w^*(a | X_t)} + \mu^a(P)(X_t) \right\} + o_P(1) \end{aligned}$$

as $T \rightarrow \infty$. □

C Proof of Lemma 5.7

This section provides the proof of Lemma 5.7. In Appendix C.1, we introduce an upper bound of the Rademacher complexity shown by Zhou et al. (2023). Then, in Appendix C.2, we prove a new lemma (Lemma C.3), which directly yields Lemma 5.7.

C.1 Upper bound of the Rademacher complexity.

Let us define a policy class

$$\Pi^D := \left\{ h : \mathcal{X} \times \prod_{a=1}^K \mathbb{R} \rightarrow \mathbb{R} \mid h(x, (\Gamma^a)_{a \in [K]}) := \sum_{a \in [K]} \left(\pi_1(a | x) - \pi_2(a | x) \right) \Gamma^a, \pi_1, \pi_2 \in \Pi \right\}$$

Then, let us define the Rademacher complexity as follows.

Definition C.1. Let $\{Z_t\}_{t=1}^T$ be a sequence of i.i.d. Rademacher random variables $Z_t \in \{-1, +1\}$: $\mathbb{P}[Z_t = +1] = \mathbb{P}[Z_t = -1] = \frac{1}{2}$.

- The empirical Rademacher complexity $\mathfrak{R}_T(\Pi^D; \{X_t, \Gamma_t^{*a}\}_{t=1}^T)$ of a function class Π^D is defined as

$$\begin{aligned} &\mathfrak{R}_T(\Pi^D; \{X_t, \Gamma_t^{*a}\}_{t=1}^T) \\ &= \mathbb{E} \left[\sup_{\pi_1, \pi_2 \in \Pi} \frac{1}{T} \left| \sum_{t=1}^T Z_t \sum_{a \in [K]} \left(\pi_1(a | X_t) - \pi_2(a | X_t) \right) \Gamma_t^{*a} \right| \mid \{X_t, \Gamma_t^{*a}\}_{t=1}^T \right], \end{aligned}$$

where the expectation is taken with respect to Z_1, \dots, Z_T .

- The Rademacher complexity $\mathfrak{R}_T(\Pi^D)$ of the function class Π^D is the expected value taken with respect to the observations $\{X_t, \Gamma_t^{*a}\}_{t=1}^T$ of the empirical Rademacher complexity:

$$\mathfrak{R}_T(\Pi^D) := \mathbb{E} \left[\mathfrak{R}_T(\Pi^D; \{X_t, \Gamma_t^{*a}\}_{t=1}^T) \right].$$

⁷Our proof for this part refers to the proof of Lemma 10 in Hadad et al. (2021).

Our proof starts from the following result about the Rademacher complexity shown by [Zhou et al. \(2023\)](#).

Lemma C.2 (From the inequality above (C.14) and the inequality in (C.19) of [Zhou et al. \(2023\)](#)). *Suppose that Assumptions 5.1 and 5.3 hold. Then, for any ζ and $P \in \mathcal{P}_\zeta$,*

$$\mathfrak{R}_T(\Pi^D) \leq 13.6\sqrt{2} \{\kappa(\Pi) + 8\} \frac{\Upsilon_*}{\sqrt{T}} + o\left(\frac{\sqrt{U_T}}{T^{3/4}}\right).$$

holds, where

$$\Upsilon_* = \mathbb{E} \left[\sqrt{\sup_{\pi_1, \pi_2 \in \Pi} \sum_{t \in [T]} \left\{ \sum_{a \in [K]} \left(\pi_1(a | Z_t) - \pi_2(a | Z_t) \right) \Gamma^{*a}(Y_t, \xi_t, X_t) \right\}^2} \right].$$

Following this upper bound, [Zhou et al. \(2023\)](#) applies the Jensen inequality to bound Υ_* by

$$\Upsilon_* \leq \sqrt{\mathbb{E} \left[\sup_{\pi_1, \pi_2 \in \Pi} \left\{ \sum_{a \in [K]} \left(\pi_1(a | Z_t) - \pi_2(a | Z_t) \right) \Gamma^{*a}(Y_t, \xi_t, X_t) \right\}^2 \right]}.$$

Then, they apply the Talagrand inequality to bound the regret as

$$\Upsilon_* \leq \sqrt{\sup_{\pi_1, \pi_2 \in \Pi} \mathbb{E} \left[\left\{ \sum_{a \in [K]} \left(\pi_1(a | Z_t) - \pi_2(a | Z_t) \right) \Gamma^{*a}(Y_t, \xi_t, X_t) \right\}^2 \right]}.$$

However, the use of the Jensen inequality yields a loose upper bound, which results in a mismatch between the upper bound and our derived lower bound. Therefore, in our proof, we consider bounding Υ_* without using Jensen's inequality.

C.2 Proof of Lemma 5.7

Based on this proof strategy, we show the following lemma.

Lemma C.3 (From the equation above (C.14) in [Zhou et al. \(2023\)](#)). *Suppose that Assumptions 5.1 and 5.3 hold. Then, for each $P \in \mathcal{P}$,*

$$\mathbb{E}_P [R_c(P) (\hat{\pi}_T^{\text{PLAS}})] \leq 54.4\sqrt{2} \{\kappa(\Pi) + 8\} \frac{\Upsilon_*}{\sqrt{T}} + o\left(\frac{\sqrt{U_T}}{T^{3/4}}\right).$$

holds, where Υ_* is defined in Lemma C.2.

Lemma 5.7 directly follows from Lemma C.3 by multiplying both sides by \sqrt{T} and letting $T \rightarrow \infty$.

Proof of Lemma C.3. Recall that

$$\hat{Q}_T(\pi) := \frac{1}{T} \sum_{t=1}^T \sum_{a \in [K]} \pi(a | X_t) \hat{\Gamma}_t^a,$$

Let us define the following quantities:

$$\begin{aligned} \tilde{Q}_T(\pi) &:= \frac{1}{T} \sum_{t=1}^T \sum_{a \in [K]} \pi(a | X_t) \Gamma_c^{*a}((Y_t^a)_{a \in [K]}, \xi_t, X_t), \\ \hat{\Delta}(\pi_1, \pi_2) &:= \hat{Q}_T(\pi_1) - \hat{Q}_T(\pi_2), \\ \tilde{\Delta}(\pi_1, \pi_2) &:= \tilde{Q}_T(\pi_1) - \tilde{Q}_T(\pi_2), \\ \Delta(\pi_1, \pi_2) &:= Q_c(P)(\pi_1) - Q_c(P)(\pi_2). \end{aligned}$$

Then, we have

$$\begin{aligned}
& R_c(P) (\widehat{\pi}_T^{\text{PLAS}}) \\
&= Q_c(P) (\pi_c^*) - Q_c(P) (\widehat{\pi}_T^{\text{PLAS}}) \\
&= \widehat{Q}_T (\pi_c^*) - \widehat{Q}_T (\pi^{\text{PLAS}}) + \Delta (\pi_c^*, \pi^{\text{PLAS}}) - \widehat{\Delta} (\pi_c^*, \pi^{\text{PLAS}}) \\
&\leq \left| \Delta (\pi_c^*, \pi^{\text{PLAS}}) - \widehat{\Delta} (\pi_c^*, \pi^{\text{PLAS}}) \right| \\
&\leq \sup_{\pi_1, \pi_2 \in \Pi} \left| \Delta (\pi_1, \pi_2) - \widehat{\Delta} (\pi_1, \pi_2) \right| \\
&\leq \sup_{\pi_1, \pi_2 \in \Pi} \left| \widehat{\Delta} (\pi_1, \pi_2) - \widetilde{\Delta} (\pi_1, \pi_2) \right| + \sup_{\pi_1, \pi_2 \in \Pi} \left| \Delta (\pi_1, \pi_2) - \widetilde{\Delta} (\pi_1, \pi_2) \right|.
\end{aligned}$$

From Lemma 5.6, we have

$$\sqrt{T} R_c(P) \leq \sup_{\pi_1, \pi_2 \in \Pi} \left| \Delta (\pi_1, \pi_2) - \widetilde{\Delta} (\pi_1, \pi_2) \right| + o_P(1).$$

Therefore, we obtain

$$\mathbb{E}_P [R_c(P) (\widehat{\pi}_T^{\text{PLAS}})] \leq \mathbb{E}_P \left[\sup_{\pi_1, \pi_2 \in \Pi} \left| \Delta (\pi_1, \pi_2) - \widetilde{\Delta} (\pi_1, \pi_2) \right| \right],$$

where we used that $\sup_{\pi_1, \pi_2 \in \Pi} \left| \widehat{\Delta} (\pi_1, \pi_2) - \widetilde{\Delta} (\pi_1, \pi_2) \right|$ is a bounded random variable.

Then, from the property of the Rademacher complexity (Bartlett and Mendelson, 2003) and Lemma C.2, we have

$$\begin{aligned}
& \mathbb{E}_P \left[\sup_{\pi_1, \pi_2 \in \Pi} \left| \Delta (\pi_1, \pi_2) - \widetilde{\Delta} (\pi_1, \pi_2) \right| \right] \\
&\leq 4\mathfrak{R}_T(\Pi) \\
&\leq 54.4\sqrt{2} \{ \kappa(\Pi) + 8 \} \frac{\Upsilon^*}{\sqrt{T}} + o\left(\frac{\sqrt{U_T}}{T^{3/4}}\right).
\end{aligned}$$

This completes the proof. □