Uniform Recovery Guarantees for Quantized Corrupted Sensing Using Structured or Generative Priors

Junren Chen*, Zhaoqiang Liu[†], Meng Ding[‡], and Michael K. Ng[§]

Abstract. This paper studies quantized corrupted sensing where the measurements are contaminated by unknown corruption and then quantized by a dithered uniform quantizer. We establish uniform guarantees for Lasso that ensure the accurate recovery of all signals and corruptions using a single draw of the sub-Gaussian sensing matrix and uniform dither. For signal and corruption with structured priors (e.g., sparsity, low-rankness), our uniform error rate for constrained Lasso typically coincides with the non-uniform one up to logarithmic factors, indicating that the uniformity costs very little. By contrast, our uniform error rate for unconstrained Lasso exhibits worse dependence on the structured parameters due to regularization parameters larger than the ones for non-uniform recovery. These results complement the non-uniform ones recently obtained in [Sun, Cui and Liu, 2022] and provide more insights for understanding actual applications where the sensing ensemble is typically fixed and the corruption may be adversarial. For signal and corruption living in the ranges of some Lipschitz continuous generative models (referred to as generative priors), we achieve uniform recovery via constrained Lasso with a measurement number proportional to the latent dimensions of the generative models. We present experimental results to corroborate our theories. From the technical side, our treatments to the two kinds of priors are (nearly) unified and share the common key ingredients of (global) quantized product embedding (QPE) property, which states that the dithered uniform quantization (universally) preserves inner product. As a by-product, our QPE result refines the one in [Xu and Jacques, 2020] under sub-Gaussian random matrix, and in this specific instance we are able to sharpen the uniform error decaying rate (for the projected-back projection estimator with signals in some convex symmetric set) presented therein from $O(m^{-1/16})$ to $O(m^{-1/8})$.

1 Introduction In corrupted sensing, our goal is to recover the signal $x^{\star} \in \mathbb{R}^n$ and corruption $v^{\star} \in \mathbb{R}^m$ from relatively few measurements

(1.1)
$$\boldsymbol{y} = \boldsymbol{\Phi}\boldsymbol{x}^{\star} + \sqrt{m}\boldsymbol{v}^{\star} + \boldsymbol{\epsilon},$$

where $\Phi \in \mathbb{R}^{m \times n}$ is the sensing matrix, $\epsilon \in \mathbb{R}^m$ represents the noise vector, \boldsymbol{y} denotes the measurements from which we seek to recover \boldsymbol{x}^* and \boldsymbol{v}^* . When the corruption \boldsymbol{v}^* does not appear, (1.1) reduces to the classical compressed sensing problem [12,28,32]; hence, corrupted sensing is a more challenging generalization of compressed sensing. While the corrupted sensing problem is ill-posed in general, faithful recovery can be achieved even in a high-dimensional

^{*}J. Chen is with Department of Mathematics, The University of Hong Kong (chenjr58@connect.hku.hk). J. Chen was supported by a Hong Kong Ph.D. Fellowship from the Hong Kong Research Grant Council (HKRGC).

[†]Z. Liu is with the School of Computer Science and Engineering, University of Electronic Science and Technology of China (zqliu12@gmail.com). Z. Liu was supported by the Shenzhen Fundamental Research Program (No. JCYJ20220530164812027), Sichuan Science and Technology Program (No. 2021YFS0374), and Sichuan Science and Technology Program (No. 2022YFS0600).

[‡]M. Ding is with the School of Mathematics, Southwest Jiaotong University (dingmeng56@163.com). M. Ding was supported by the National Natural Science Foundation of China under Grant 12201522 and the Fundamental Research Funds for the Central Universities under Grant 2682023CX069.

[§]M. K. Ng is with Department of Mathematics, Hong Kong Baptist University (michael-ng@hkbu.edu.hk). M. K. Ng was supported by HKRGC GRF 12300218, 12300519, 17201020, 17300021, C1013-21GF, C7004-21GF and Joint NSFC-RGC N-HKU76921.

regime (i.e., $m \ll n$) by utilizing additional structures of (x^{\star}, v^{\star}) , such as sparsity and lowrankness. In the literature, recovery methods with theoretical guarantees have been well developed in a long line of works, first for some specific instances of (1.1) like sparse signal recovery or low-rank matrix sensing under sparse corruption [21, 49, 61, 62, 82], and then for the more general cases where x^{\star} and v^{\star} exhibit some structures that are often captured by Gaussian width [14, 33, 57]. We note that (1.1) captures a series of applications in imaging problems. Specifically, associated with various operators Φ , the reduced model $y = \Phi x^{\star} + \epsilon$ (without corruption \boldsymbol{v}^{\star}) already models most problems in computational imaging [2], from classical tasks like deblurring, inpainting and super-resolution, to a wide range of tomographic imaging applications such as magnetic resonance imaging and X-ray computed tomography (e.g., [29, 31]). These inverse problems of recovering x^{\star} from y are collectively referred to as *image reconstruction*, where prior knowledge on the image signal is often available (e.g., sparsity in some dictionary or basis, (approximately) low-rankness, smoothness [39,84]). However, the flexibility of including the corruption v^{\star} , as per (1.1), becomes necessary in certain imaging problems, in which the recovery of the corruption often provides useful information. We provide two specific examples:

- The *face recognition* example [10,80] can be modeled by (1.1), where the columns of the dictionary Φ are the training face images, the prior on the present face image y is that it can be represented as a *sparse* linear combination of the training faces. However, it is unreasitic to assume that the prior exactly holds true (e.g., due to undesired parts in y such as glasses and shadows), and a useful remedy is to complement our prior via a sparse corruption v^{*}.
- In some *image reconstruction* problems our goal is to recover the image and the impulsive signal v^* , with Φ being certain dictionary that generates the image or the sensing matrix that produces the compressive measurements of the signal x^* . See the star-galaxy separation example described in [75] for instance.

Additionally, the corrupted sensing model (1.1) has found applications in sensor network analysis [41], subspace clustering [30], latent variable modeling [13], among others.

Note that the data is inevitably quantized to finite precision in digital signal processing [37, 79], and working with coarsely quantized data has proven effective in many large scale machine learning or signal processing systems [40, 83, 85]. Consequently, recent years have witnessed rapidly increasing literature on quantized compressed sensing. These works proposed various quantization schemes that are associated with accurate (post-quantization) recovery methods, including 1-bit quantization [7,19,27,42,46,65,66,77], uniform (multi-bit) quantization [17,46, 77,81], and other adaptive quantization methods as surveyed in [24]. The recent work [76] even extended the theoretical foundation of 1-bit compressed sensing [42] to learning a signal set. However, nearly all of them are restricted to classical compressed sensing without accounting for the additional corruption v^* . The single exception is a recent work [75], in which the authors analyzed corrupted sensing under a dithered uniform quantizer $\mathcal{Q}_{\delta}(\cdot + \tau)$:

(1.2)
$$\dot{\boldsymbol{y}} := \mathcal{Q}_{\delta}(\boldsymbol{y} + \boldsymbol{\tau}) = \mathcal{Q}_{\delta}(\boldsymbol{\Phi}\boldsymbol{x}^{\star} + \sqrt{m}\boldsymbol{v}^{\star} + \boldsymbol{\epsilon} + \boldsymbol{\tau}),$$

where $\boldsymbol{y} = \boldsymbol{\Phi}\boldsymbol{x}^{\star} + \sqrt{m}\boldsymbol{v}^{\star} + \boldsymbol{\epsilon}$ is the unquantized measurements in corrupted sensing as per (1.1), $\boldsymbol{\tau} \in \mathbb{R}^m$ is the uniform dither, $\mathcal{Q}_{\delta}(\cdot)$ is the uniform quantizer with resolution δ , see

subsection 2.4. Under sub-Gaussian Φ , they showed that accurate recovery of $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star})$ can be achieved by Lasso, thus confirming the compatibility of the dithered uniform quantization and the recovery of the additional structured corruption \boldsymbol{v}^{\star} . Nonetheless, their recovery guarantees are *non-uniform* and only ensure the recovery of a single pair of $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star})$ fixed before drawing (Φ, ϵ, τ) , with the implication that a new realization of (Φ, ϵ, τ) is needed for the sensing and recovery of a different pair of $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star})$. Thus, the possibility of uniform recovery in quantized corrupted sensing (1.2) remains unaddressed:

Is it possible to recover all signals and corruptions using a fixed sensing ensemble (Φ, ϵ, τ) ? If yes, what is the cost of uniformity compared to non-uniform recovery?

Importance of Uniformity: Compressed sensing theories are developed to promote the understanding of its many real-world applications, and the above uniform recovery question is important for theory of (1.2) for the following reasons:

- Uniform recovery is a highly sought-after notion in compressed/corrupted sensing theory since the sensing matrix Φ is typically fixed in applications (e.g., think of the above face recognition example), and one expects that the single fixed sensing ensemble works for all possible signals (and corruptions) that may arise. In fact, uniformity is a defining property for achieving *compression* in some applications, and non-uniform recovery with new sensing ensemble for new signal could be unrealistic (if not impossible) since the memory of (Φ, τ) is already heavier than the signal itself.
- In view of the corruption v^* in (1.2), a uniform guarantee also offers stronger robustness than a non-uniform guarantee. Specifically, a uniform guarantee tolerates v^* generated in an *adversarial* manner according to the knowledge of $(\Phi, \epsilon, \tau, x^*)$, and the error bound remains valid as long as v^* satisfies certain structured assumption like sparsity. In contrast, a non-uniform guarantee only works for a fixed v^* oblivious to (Φ, ϵ, τ) .

Generative Prior: Beyond the classical structured priors, it was proposed in [5] to assume that the desired signal in compressed sensing lies in the range of a pre-trained generative model, known as a generative prior. This new perspective has led to successful numerical results such as a significant reduction of the required number of measurements for accurate recovery, as well as attracted much research attention with various extensions such as nonlinear models [52, 55, 69], MRI applications [43, 70], and information-theoretic bounds [47, 56], among others. Note that generative prior has now been widely applied to imaging-related inverse problems; we refer interested readers to [63]. For the specific quantized corrupted sensing problem (1.2), the results in [75] are restricted to classical structured priors promoted by certain norms, and it is unclear whether their theory extends to generative priors. Without considering quantization, the linear case of corrupted sensing with generative priors has been studied in [3,4] under the name of "generative demixing" (here, [3] is the extended version of the published conference paper [4]).

1.1 Main Results In this paper, we establish *uniform* recovery guarantees for (1.2) with structured priors on (x^*, v^*) using (un)constrained Lasso. Our uniform guarantees state that a single realization of (Φ, ϵ, τ) can be used for simultaneously recovering all (x^*, v^*)

in the structured sets,¹ thus answering the above question in affirmative. We also provide careful comparison with [75] to unveil the cost of uniformity. Noticing the recent trend of using generative prior for compressed sensing, we also present a uniform recovery guarantee for quantized corrupted sensing where x^* and v^* are equipped with generative priors (see Assumption 5 for details). We highlight and summarize our major results as follows:

- Structured Priors via Constrained Lasso: We establish uniform recovery guarantees for quantized corrupted sensing (1.2) with structured priors on $(\boldsymbol{x}^*, \boldsymbol{v}^*)$ via Lasso. In constrained Lasso, our uniform error rate in Theorem 3.2 exhibits a decaying rate of $O(m^{-1/2})$ and typically coincides with the non-uniform one in [75] up to logarithmic factors (see Remark 3.3 and Corollary 3.5, Corollary 3.7), indicating that the uniformity costs very little. To our best knowledge, even going back to compressed sensing (without corruption) associated with the dithered uniform quantizer, our Theorem 3.2 provides the sharpest uniform error rate for a computationally feasible decoder (see Remark 3.9).
- Structured Priors via Unconstrained Lasso: In unconstrained Lasso, our result in Theorem 3.12 decays in m with a rate of $O(m^{-1/2})$ but exhibits a worse dependence on the structured parameter. To our best knowledge, this is the first uniform guarantee for quantized compressed sensing via unconstrained Lasso (see Remark 3.13 and Corollary 3.14, Corollary 3.15). The results in this and the previous dot point strengthen the non-uniform ones in [75] and shed more light on the understanding of actual applications with fixed sensing ensemble and possibly adversarial corruption.
- Generative Priors: We present the first result in Theorem 3.16 for quantized corrupted sensing using generative priors, which assume that the signal and corruption lie in the ranges of some Lipschitz continuous generative models with latent dimensions k and k'. Our result guarantees that roughly $\tilde{O}\left(\frac{(1+\delta^2)(k+k')}{\mu^2}\right)$ measurements (up to logarithmic factors) suffice for achieving a uniform ℓ_2 -norm recovery error of μ via constrained Lasso. Note that this also implies a decaying rate of $O(m^{-1/2})$; see Remark 3.17.

1.2 Technical Contributions and Technically Related Works Considering the theoretical nature of our work, we provide an introduction from a more technical aspect.

A Sketch of our Techniques: While most existing works focused on either structured prior or generative prior, we provide near unified treatments for the two kinds of priors, with the common key ingredient for achieving uniformity being the global quantized product embedding (QPE) property, which states that the dithered quantization uniformly preserves inner product, i.e., $|\langle Q_{\delta}(a + \tau) - a, b \rangle|$ is uniformly small over all (a, b) in certain constraint sets. The most general version of QPE is presented in Theorem B.3 in Appendix B.2, and a version sufficient for proving our main theorems is given in Corollary B.4 in Appendix B.3. The proofs of these QPE results, despite following the conceptually simple covering strategy, appear to be the most technical and tedious part of this work. To avoid being overly technical in the main body, we collect the statements and proofs of QPE in Appendix B. Equipped with QPE and a set of useful concentration inequalities (see Appendix A.1), the major differences in proving our three main theorems (Theorem 3.2, 3.12, 3.16) lie in estimating the Gaussian

 $^{^{1}}$ This is a generalization of traditional signal structures such as sparsity and low-rank, see Definition 3.1 and Assumption 3.

UNIFORM GUARANTEES FOR QUANTIZED CORRUPTED SENSING

width and Kolmogorov entropy of various constrain sets, which will be settled in Appendix A.2.

We provide comparisons with the works most relevant to this paper in techniques (Readers less interested in proof techniques could safely skip the discussions below):

- Comparing with [81] on QPE: Under the name of *limited projection distortion* (LPD) property, Xu and Jacques [81] utilized QPE for analyzing the projected-back projection (PBP) estimator. In this work, we show that global QPE also serves as the key ingredient in analyzing uniform recovery via Lasso. Compared to the global QPE in [81], our Theorem B.3 is a generalization and provides (instance-wise) improvement under sub-Gaussian sensing matrix; see Corollary D.1 and Corollary D.3 in Appendix D.1. As an interesting by-product of Corollary D.3, under sub-Gaussian sensing matrix, we are able to improve the uniform error rate of PBP over a symmetric convex signal set from $O(m^{-1/16})$ to $O(m^{-1/8})$; see Proposition D.5 in Appendix D.2.
- Comparing with [18,36] on the Approach to Uniformity: The recent work [36] due to Genzel and Stollenwerk developed a unified approach to proving uniform recovery guarantees for constrained Lasso in non-linear compressed sensing $y_i = f_i(\Phi_i^{\top} x^{\star})$, where the possibly random $f_i(\cdot)$ captures some non-linearity that can be unknown and/or discontinuous. However, under discontinuous $f_i(\cdot)$, their general strategy leads to a uniform decaying rate $O(m^{-1/4})$ inferior to our $O(m^{-1/2})$. More recently, Chen et al. [18] extended the scope of [36] to non-linear compressed sensing with generative prior. They observed that using a different concentration inequality yields tighter bound for the generative case, thus they managed to prove a uniform decaying rate of $O(m^{-1/2})$ for discontinuous $f_i(\cdot)$ (e.g., various quantization models). However, as we will discuss in Remark 3.11 and Remark 3.18, some hurdle arises if we follow the general strategy in [18,36] to prove our main theorems, thus our techniques are not implied by these two works. Indeed, our QPE-based analysis suggests a possible strategy to improve the rate $O(m^{-1/4})$ in [36] to $O(m^{-1/2})$ under discontinuous $f_i(\cdot)$.

1.3 Paper Outline In section 2, we provide preliminaries and set up notations. In section 3, we present our main theorems. Experimental results are reported in section 4. We provide some remarks to conclude the paper in section 5. We provide technical lemmas and useful propositions in Appendix A to support our analysis. We develop a general global QPE result in Appendix B.2 and present a version sufficient for proving our main theorems in Appendix B.3. The proofs of results in the main body, if missing, are deferred to Appendix C. In Appendix D, we present more implications of our QPE result (Appendix D.1) and then obtain a by-product for PBP estimator (Appendix D.2). In Appendix E, we provide a list of recurring notation (Table 1) to improve the readability of this paper.

2 Preliminaries We first collect some generic notations. We represent matrices and vectors by boldface letters, scalars by regular letters. For positive integer m we write $[m] := \{1, ..., m\}$. We use |S| to denote the cardinality of any finite set S. For a vector $\boldsymbol{x} = [x_i] \in \mathbb{R}^d$, we work with the ℓ_p -norm $\|\boldsymbol{x}\|_p = (\sum_i |x_i|^p)^{1/p}$ $(p \ge 1)$, max norm $\|\boldsymbol{x}\|_{\infty} = \max_i |x_i|$, and zero norm $\|\boldsymbol{x}\|_0$ that counts the number of non-zero entries. We write the standard Euclidean sphere in *n*-dimensional space as $\mathbb{S}^{n-1} = \{\boldsymbol{x} \in \mathbb{R}^n : \|\boldsymbol{x}\|_2 = 1\}$, the set of *s*-sparse vectors as $\Sigma_s^n = \{\boldsymbol{x} \in \mathbb{R}^n : \|\boldsymbol{x}\|_0 \le s\}$. The inner product of $\boldsymbol{x}, \boldsymbol{y} \in \mathbb{R}^n$ is $\langle \boldsymbol{x}, \boldsymbol{y} \rangle = \boldsymbol{x}^\top \boldsymbol{y}$. Given a matrix $\boldsymbol{A} \in \mathbb{R}^{p \times q}$, we denote its operator norm (that equals the maximal singular value), Frobenius

norm, nuclear norm (i.e., sum of singular values) by $\|\boldsymbol{A}\|_{\text{op}}$, $\|\boldsymbol{A}\|_{\text{F}}$, $\|\boldsymbol{A}\|_{\text{nu}}$, respectively. The inner product between matrices $\boldsymbol{A}, \boldsymbol{B}$ is $\langle \boldsymbol{A}, \boldsymbol{B} \rangle = \text{Tr}(\boldsymbol{A}^{\top}\boldsymbol{B})$. The set of matrices with rank not exceeding r is denoted by $M_r^{p,q} = \{\boldsymbol{A} \in \mathbb{R}^{p \times q} : \text{rank}(\boldsymbol{A}) \leq r\}$.

Given a norm $f(\cdot)$ in \mathbb{R}^n (resp. $\mathbb{R}^{p \times q}$), we denote the corresponding ball with radius r by $\mathbb{B}_f^n(r) = \{ \boldsymbol{x} \in \mathbb{R}^n : f(\boldsymbol{x}) \leq r \}$ (resp. $\mathbb{B}_f^{p,q}(r) = \{ \boldsymbol{A} \in \mathbb{R}^{p \times q} : f(\boldsymbol{A}) \leq r \}$). We let $\mathbb{B}_f^n := \mathbb{B}_f^n(1)$, $\mathbb{B}_f^{p,q} := \mathbb{B}_f^{p,q}(1)$ be the unit ball. For instance, \mathbb{B}_2^n , \mathbb{B}_1^n are respectively the ℓ_2 -ball, ℓ_1 -ball in \mathbb{R}^n , $\mathbb{B}_F^{p,q}$, $\mathbb{B}_{nu}^{p,q}$ are respectively the Frobenius norm ball, nuclear norm ball in $\mathbb{R}^{p \times q}$. The dual norm of $f(\cdot)$ is defined as $f^*(\boldsymbol{x}) := \sup_{\boldsymbol{y} \in \mathbb{B}_f} \langle \boldsymbol{x}, \boldsymbol{y} \rangle$, and we note the Hölder's inequality $\langle \boldsymbol{x}, \boldsymbol{y} \rangle \leq f(\boldsymbol{x}) \cdot f^*(\boldsymbol{y})$. The descent cone of f at a point \boldsymbol{x} and its normalized counterpart are defined as

(2.1)
$$\mathcal{D}_f(\boldsymbol{x}) := \{ \boldsymbol{u} : \exists t > 0, \text{ s.t. } f(\boldsymbol{x} + t\boldsymbol{u}) \le f(\boldsymbol{x}) \}, \ \mathcal{D}_f^*(\boldsymbol{x}) := \mathcal{D}_f(\boldsymbol{x}^*) \cap \mathbb{S}^{n-1} \}$$

Throughout this paper, $\mathbb{P}(\cdot)$, $\mathbb{E}(\cdot)$, $\mathbb{I}(\cdot)$ stand for probability, expectation, indicator function, respectively. We make no attempt to refine multiplicative constants, and we use C, C_i, c, c_i to denote absolute constants whose value may vary from line to line. For some quantities I_1, I_2 , We write $I_1 = O(I_2)$ or $I_1 \leq I_2$ if $I_1 \leq CI_2$ holds for some absolute constant C; Conversely, we write $I_1 = \Omega(I_2)$ or $I_1 \gtrsim I_2$ if $I_1 \geq cI_2$ for some c; we refer to (C, c) behind \leq or \geq as the implied constant. For some quantity I, we may write "certain event holds with probability exceeding $1 - \exp(-\Omega(I))$ " to state that this event holds with probability at least $1 - \exp(cI)$ for some absolute constant c > 0. We will write $I_1 \simeq I_2$ to state that $I_1 = O(I_2)$ and $I_1 = \Omega(I_2)$ simultaneously hold. Given $W \subset \mathbb{R}^d$ we use $\mathscr{U}(W)$ to denote the uniform distribution over W. Also, $\mathcal{N}(\mu, \Sigma)$ represents Gaussian variable/vector with mean μ and covariance Σ .

2.1 Sub-Gaussian Random Variable The sub-Gaussian norm of a random variable X is defined as $||X||_{\psi_2} = \inf\{t > 0 : \mathbb{E}(X^2/t^2) \le 2\}$, and we have $||X||_{\psi_2} \asymp \sup_{p \ge 1} p^{-1/2} (\mathbb{E}|X|^p)^{1/p}$ and hence $||X||_{\psi_2} = O(K)$ if $|X| \le K$. X is said to be sub-Gaussian if $||X||_{\psi_2} < \infty$, and sub-Gaussian X has probability tail resembling that of a Gaussian variable:

(2.2)
$$\mathbb{P}(|X| \ge t) \le 2 \exp\left(-\frac{ct^2}{\|X\|_{\psi_2}^2}\right)$$

holds for any t > 0. Moreover, for independent *zero-mean* random variables X_i 's we have (see [78, Prop. 2.6.1])

(2.3)
$$\left\|\sum_{i} X_{i}\right\|_{\psi_{2}}^{2} \leq C \sum_{i} \|X_{i}\|_{\psi_{2}}^{2}.$$

A random vector $\boldsymbol{X} \in \mathbb{R}^n$ is sub-Gaussian if it has finite sub-Gaussian norm, which is defined as $\|\boldsymbol{X}\|_{\psi_2} = \sup_{\boldsymbol{v} \in \mathbb{S}^{n-1}} \|\boldsymbol{v}^\top \boldsymbol{X}\|_{\psi_2}$. Assume \boldsymbol{X} has independent *zero-mean* entries X_i 's satisfying $\|X_i\|_{\psi_2} \leq K$, then $\|\boldsymbol{X}\|_{\psi_2} = O(K)$ [78, Lem. 3.4.2]. Readers may refer to [78, Sections 2–3] for more details.

2.2 Covering Number and Kolmogorov Entropy Given $\mathcal{K} \subset \mathbb{R}^n$, a subset $\mathcal{G} \subset \mathcal{K}$ is said to be an ε -net (with respect to Euclidean distance) of \mathcal{K} , if for any $\boldsymbol{x} \in \mathcal{K}$ there exists $\boldsymbol{x}' \in \mathcal{G}$

satisfying $\|\boldsymbol{x} - \boldsymbol{x}'\|_2 \leq \varepsilon$, i.e., $\mathcal{K} \subset \bigcup_{\boldsymbol{x} \in \mathcal{G}} (\boldsymbol{x} + \mathbb{B}_2^n(\varepsilon))$. The covering number of \mathcal{K} under radius ε , denoted $\mathscr{N}(\mathcal{K}, \varepsilon)$, is defined to be the smallest possible cardinality of an ε -net of \mathcal{K} . We note the following monotonicity of covering number [78, Exercise 4.2.10]:

(2.4)
$$\mathcal{N}(\mathcal{K},\varepsilon) \leq \mathcal{N}(\mathcal{K}',\varepsilon/2), \text{ if } \mathcal{K} \subset \mathcal{K}'.$$

We will more frequently work with the equivalent notion called Kolmogorov entropy:

(2.5)
$$\mathscr{H}(\mathcal{K},\varepsilon) = \log \mathscr{N}(\mathcal{K},\varepsilon).$$

2.3 Gaussian Width and Gaussian Complexity Suppose that \boldsymbol{g} has i.i.d. $\mathcal{N}(0,1)$ entries, then the Gaussian width of $\mathcal{K} \subset \mathbb{R}^n$ is defined as $\omega(\mathcal{K}) = \mathbb{E} \sup_{\boldsymbol{x} \in \mathcal{K}} \langle \boldsymbol{g}, \boldsymbol{x} \rangle$, which is a geometric quantity that precisely and stably captures the intrinsic dimension of \mathcal{K} . Gaussian width is closely related to Kolmogorov entropy. Specifically, we can bound the Kolmogorov entropy by Gaussian width via Sudakov's inequality [78, Thm. 8.1.13]

(2.6)
$$\mathscr{H}(\mathcal{K},\varepsilon) \leq \frac{\omega^2(\mathcal{K})}{\varepsilon^2}.$$

We can also estimate $\omega(\mathcal{K})$ by $\mathscr{H}(\mathcal{K}, \cdot)$ via Dudley's inequality [78, Thm. 8.1.10]

(2.7)
$$\omega(\mathcal{K}) \le C \int_0^\infty \sqrt{\mathscr{H}(\mathcal{K},\varepsilon)} \, \mathrm{d}\varepsilon.$$

A slightly different notion is the Gaussian complexity defined as $\gamma(\mathcal{K}) = \mathbb{E} \sup_{\boldsymbol{x} \in \mathcal{K}} |\langle \boldsymbol{g}, \boldsymbol{x} \rangle|$. In many cases $\omega(\mathcal{K})$ and $\gamma(\mathcal{K})$ are of similar scaling. For instance, $\omega(\mathcal{K}) \simeq \gamma(\mathcal{K})$ holds if $0 \in \mathcal{K}$. More generally, we note the following relation from [78, Exercise 7.6.9]

(2.8)
$$\frac{1}{3} \Big(\omega(\mathcal{K}) + \|\boldsymbol{x}\|_2 \Big) \le \gamma(\mathcal{K}) \le 2 \Big(\omega(\mathcal{K}) + \|\boldsymbol{x}\|_2 \Big)$$

that holds for any $\boldsymbol{x} \in \mathcal{K}$. Given $\mathcal{K} \subset \mathbb{R}^n$ and some $\rho > 0$, we will work with the localized version of \mathcal{K} defined as

(2.9)
$$\mathcal{K}_{\text{loc}}^{(\rho)} = (\mathcal{K} - \mathcal{K}) \cap \mathbb{B}_2^n(\rho).$$

We denote the radius of \mathcal{K} by $\operatorname{rad}(\mathcal{K}) = \sup_{a \in \mathcal{K}} \|a\|_2$.

2.4 Dithered Uniform Quantization For some resolution $\delta > 0$, the uniform quantizer $\mathcal{Q}_{\delta}(\cdot)$ quantizes a scalar *a* to

(2.10)
$$\mathcal{Q}_{\delta}(a) = \delta\left(\left\lfloor \frac{a}{\delta} \right\rfloor + \frac{1}{2}\right).$$

Note that $\mathcal{Q}_{\delta}(\cdot)$ enjoys the bounded distortion property, i.e.,

(2.11)
$$|\mathcal{Q}_{\delta}(a) - a| \le \frac{\delta}{2}$$

holds for any a. In this paper, we use a dithered uniform quantizer that involves a uniform dither $\tau \sim \mathscr{U}\left[-\frac{\delta}{2}, \frac{\delta}{2}\right]$. Specifically, we quantize a to $\mathcal{Q}_{\delta}(a+\tau)$, and we refer to $\xi := \mathcal{Q}_{\delta}(a+\tau)-a$ as quantization noise, which is always bounded because

(2.12)
$$|\xi| \le |\mathcal{Q}_{\delta}(a+\tau) - (a+\tau)| + |\tau| \le \frac{\delta}{2} + \frac{\delta}{2} = \delta.$$

With dithering, the nice property is that the quantization noise is zero-mean:

(2.13)
$$\mathbb{E}(\xi) = \mathbb{E}[\mathcal{Q}_{\delta}(a+\tau)] - a = 0,$$

see [17, 37, 38, 77] for instance.² To quantize a vector $\boldsymbol{a} \in \mathbb{R}^m$, we apply the dithered uniform quantizer to each entry in a *memoryless* manner. That is, we draw a random uniform dither $\boldsymbol{\tau} \sim \mathscr{U}([-\frac{\delta}{2}, \frac{\delta}{2}]^m)$ and then quantize \boldsymbol{a} to $\mathcal{Q}_{\delta}(\boldsymbol{a}+\boldsymbol{\tau})$. Let $\boldsymbol{\xi} = \mathcal{Q}_{\delta}(\boldsymbol{a}+\boldsymbol{\tau})-\boldsymbol{a}$ be the quantization noise. It follows that, for a fixed $\boldsymbol{a} \in \mathbb{R}^m$, entries of $\boldsymbol{\xi}$ are independent (since entries of $\boldsymbol{\tau}$ are independent), zero-mean (due to (2.13)), and bounded by δ (see (2.12)), and hence the sub-Gaussian norm of each entry also scales as $O(\delta)$. Taken collectively, we arrive at (see subsection 2.1)

(2.14)
$$\|\boldsymbol{\xi}\|_{\psi_2} = O(\delta).$$

3 Main Results Recall that the corrupted sensing problem can be formulated as

(3.1)
$$\boldsymbol{y} = \boldsymbol{\Phi}\boldsymbol{x}^{\star} + \sqrt{m}\boldsymbol{v}^{\star} + \boldsymbol{\epsilon}$$

where v^{\star} is the corruption mixed with the clean measurements Φx^{\star} of the signal x^{\star} , and ϵ represents measurement noise. In this work, we study a more challenging nonlinear model that involves quantization of y, adopting a dithered uniform quantizer following prior works [17,46,75,77,81]. For some quantization level $\delta > 0$,³ we acquire the quantized measurements as

(3.2)
$$\dot{\boldsymbol{y}} := \mathcal{Q}_{\delta}(\boldsymbol{y} + \boldsymbol{\tau}) = \mathcal{Q}_{\delta}(\boldsymbol{\Phi}\boldsymbol{x}^{\star} + \sqrt{m}\boldsymbol{v}^{\star} + \boldsymbol{\epsilon} + \boldsymbol{\tau}),$$

where $\boldsymbol{\tau} \sim \mathscr{U}([-\frac{\delta}{2}, \frac{\delta}{2}]^m)$ is the uniform dither independent of $\boldsymbol{\Phi}$ and $\boldsymbol{\epsilon}$. We denote the quantization noise (that depends on $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star})$) by

(3.3)
$$\boldsymbol{\xi}_{\boldsymbol{x}^{\star},\boldsymbol{v}^{\star}} := \dot{\boldsymbol{y}} - \boldsymbol{y} = \mathcal{Q}_{\delta}(\boldsymbol{\Phi}\boldsymbol{x}^{\star} + \sqrt{m}\boldsymbol{v}^{\star} + \boldsymbol{\epsilon} + \boldsymbol{\tau}) - (\boldsymbol{\Phi}\boldsymbol{x}^{\star} + \sqrt{m}\boldsymbol{v}^{\star} + \boldsymbol{\epsilon}),$$

with the k-th entry denoted by $(\boldsymbol{\xi}_{\boldsymbol{x}^{\star},\boldsymbol{v}^{\star}})_k$. Then we can also write (3.2) as

(3.4)
$$\dot{\boldsymbol{y}} = \boldsymbol{\Phi}\boldsymbol{x}^{\star} + \sqrt{m}\boldsymbol{v}^{\star} + \boldsymbol{\epsilon} + \boldsymbol{\xi}_{\boldsymbol{x}^{\star},\boldsymbol{v}^{\star}}.$$

²It is revealed by (2.13) that the benefit of dithering is to whiten the quantization noise. As a brief introduction, we mention that the use of dithering (prior to quantization) dates back to early engineering works [44,51] and theoretical analysis [73], while in the past few years it has regained a surge of research interest in various estimation/recovery problems, including compressed sensing [27, 46, 75, 77, 81], matrix completion [9, 17, 19, 22], and more recently covariance estimation [15, 25, 26] and reduced-rank regression [20].

³Smaller δ corresponds to higher resolution. Specifically, letting $\delta \to 0$ returns the unquantized (full-data) setting.

Moreover, let Φ_i^{\top} be the *i*-th row of Φ , v_i^{\star} and ϵ_i be respectively the *i*-th entry of v^{\star} and ϵ , then the *i*-th quantized measurement \dot{y}_i is given by

(3.5)
$$\dot{y}_i = \mathcal{Q}_{\delta}(\boldsymbol{\Phi}_i^{\top} \boldsymbol{x}^{\star} + \sqrt{m} \boldsymbol{v}_i^{\star} + \boldsymbol{\epsilon}_i + \boldsymbol{\tau}_i) = \boldsymbol{\Phi}_i^{\top} \boldsymbol{x}^{\star} + \sqrt{m} \boldsymbol{v}_i^{\star} + \boldsymbol{\epsilon}_i + (\boldsymbol{\xi}_{\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star}})_i.$$

Note that the recovery behaviour under a sub-Gaussian sensing matrix serves as an important benchmark in compressed sensing. In this work, we make the following assumption.

Assumption 1 (Random Sensing Ensemble). $\Phi_1, ..., \Phi_m$ are independent, zero-mean, isotropic (i.e., $\mathbb{E}(\Phi_i \Phi_i^{\top}) = I_n$), sub-Gaussian sensing vectors satisfying $\|\Phi_i\|_{\psi_2} \leq K$ for all *i* and for some absolute constant K; $\epsilon_1, ..., \epsilon_m$ are independent of each other and of Φ , and sub-Gaussian satisfying $\|\epsilon_i\|_{\psi_2} \leq E$; $\tau_1, ..., \tau_m$ are independent of each other and of (Φ, ϵ) , and $\tau_i \sim \mathscr{U}[-\frac{\delta}{2}, \frac{\delta}{2}]$.

To handle the high-dimensional regime where $m \ll n$, it is standard and necessary to utilize the low-complexity structures of $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star})$. Following [14] we assume that the structure can be promoted by some norm, for instance, ℓ_1 -norm for sparsity, ℓ_1/ℓ_2 -norm for group sparsity, and nuclear norm for low-rankness.

Assumption 2 (Structures Promoted by Norms). For some low-complexity sets $\mathcal{K}_{\boldsymbol{x}} \subset \mathbb{R}^n$ and $\mathcal{K}_{\boldsymbol{v}} \subset \mathbb{R}^m$, we assume that $\boldsymbol{x}^{\star} \in \mathcal{K}_{\boldsymbol{x}}$ and $\boldsymbol{v}^{\star} \in \mathcal{K}_{\boldsymbol{v}}$. The structures of \boldsymbol{x}^{\star} and \boldsymbol{v}^{\star} can be promoted by the norms $f(\boldsymbol{x})$ and $g(\boldsymbol{v})$, respectively.

We will investigate the recovery performance of two types of Lasso: the constrained Lasso⁴

(3.6)
$$(\hat{\boldsymbol{x}}, \hat{\boldsymbol{v}}) = \arg\min_{\substack{\boldsymbol{x} \in \mathbb{R}^n \\ \boldsymbol{v} \in \mathbb{R}^m}} \| \dot{\boldsymbol{y}} - \boldsymbol{\Phi} \boldsymbol{x} - \sqrt{m} \boldsymbol{v} \|_2, \text{ s.t. } f(\boldsymbol{x}) \le f(\boldsymbol{x}^{\star}), \ g(\boldsymbol{v}) \le g(\boldsymbol{v}^{\star}),$$

and the unconstrained Lasso

(3.7)
$$(\hat{\boldsymbol{x}}, \hat{\boldsymbol{v}}) = \arg\min_{\substack{\boldsymbol{x} \in \mathbb{R}^n \\ \boldsymbol{v} \in \mathbb{R}^m}} \| \dot{\boldsymbol{y}} - \boldsymbol{\Phi} \boldsymbol{x} - \sqrt{m} \boldsymbol{v} \|_2^2 + \lambda_1 \cdot f(\boldsymbol{x}) + \lambda_2 \cdot g(\boldsymbol{v}).$$

Note that the loss function $\mathcal{L}(\boldsymbol{x}, \boldsymbol{v}) := \| \boldsymbol{\dot{y}} - \boldsymbol{\Phi}\boldsymbol{x} - \sqrt{m}\boldsymbol{v} \|_2^2$ is simply the regular ℓ_2 -loss with full observations \boldsymbol{y} substituted by the quantized ones $\boldsymbol{\dot{y}}$.

The non-uniform guarantees in [75] state that for any fixed $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star})$, (3.6) and (3.7) deliver comparably accurate recovery using a random realization of $(\boldsymbol{\Phi}, \boldsymbol{\dot{y}})$ according to Assumption 1. By contrast, the primary goal of this paper is to establish uniform recovery guarantees that ensure the accurate recovery of *all* possible pairs of $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star})$ using $(\boldsymbol{\Phi}, \boldsymbol{\dot{y}})$, where the quantized measurements $\boldsymbol{\dot{y}}$ are produced by a single draw of $(\boldsymbol{\Phi}, \boldsymbol{\epsilon}, \boldsymbol{\tau})$. Compared to non-uniform guarantees, our uniform ones provide more insights in understanding the actual applications with fixed sensing ensemble and possibly adversarial corruption, as we explained in section 1.

3.1 Structured Priors with Constrained Lasso Our first set of results are for (3.2) where x^* and v^* lie in some structured sets, which we follow [81, Section 3.1] and define as follows.

⁴Though it is possible the pursue a relaxation (e.g., [36, 67]), we follow prior works such as [14, 20, 75, 77] to consider the constrained Lasso with the best possible constraint to allow for an descent-cone-based analysis.

Definition 3.1 (Structured Set). If $\mathcal{K} \subset \mathbb{R}^p$ is a cone satisfying

(3.8)
$$\mathscr{H}(\mathcal{K} \cap \mathbb{B}_2^p, \eta) \le C \cdot \omega^2(\mathcal{K} \cap \mathbb{B}_2^p) \log\left(1 + \frac{1}{\eta}\right)$$

for all $\eta > 0$ and for some absolute constant C, then we say \mathcal{K} is a structured set.

Structured set is a generalization of various prototypical structures utilized in compressed sensing, e.g., (group) sparse vectors, low-rank matrices (*cf.* Proposition A.6), subspaces, union of subspaces, among others. In Definition 3.1, the distinguishing feature of a structured set is that its Kolmogorov entropy is only logarithmically dependent on its covering radius, in contrast to Sudakov's inequality (2.6) that holds for arbitrary subset. We note that notions analogous to structured sets have been widely adopted in the literature [16, 23, 64].

To derive a uniform error bound in quantized compressed sensing, it is standard to concentrate on $\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star}$ with bounded ℓ_2 -norm (e.g., [46, Thm. 3], [81, Thm. 4.1]). Without loss of generality, we assume that $\boldsymbol{x}^{\star} \in \mathbb{B}_2^n$ and $\boldsymbol{v}^{\star} \in \mathbb{B}_2^m$ and state our structured set assumption as follows.

Assumption 3 (Structured Priors). Given a pair of structured sets $\mathcal{K}^0_{\boldsymbol{x}}$ and $\mathcal{K}^0_{\boldsymbol{v}}$ as per Definition 3.1, we let $\mathcal{K}_{\boldsymbol{x}} = \mathcal{K}^0_{\boldsymbol{x}} \cap \mathbb{B}^n_2$ and $\mathcal{K}_{\boldsymbol{v}} = \mathcal{K}^0_{\boldsymbol{v}} \cap \mathbb{B}^m_2$. According to Definition 3.1, for any $\eta > 0$ we have

(3.9)
$$\mathscr{H}(\mathcal{K}_{\boldsymbol{x}},\eta) \lesssim \omega^2(\mathcal{K}_{\boldsymbol{x}}) \log\left(1+\frac{1}{\eta}\right), \ \mathscr{H}(\mathcal{K}_{\boldsymbol{v}},\eta) \lesssim \omega^2(\mathcal{K}_{\boldsymbol{v}}) \log\left(1+\frac{1}{\eta}\right).$$

We first consider constrained Lasso (3.6) and present an upper bound that holds uniformly for all $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star}) \in \mathcal{K}_{\boldsymbol{x}} \times \mathcal{K}_{\boldsymbol{v}}$.

Theorem 3.2 (Uniform Recovery via Constrained Lasso). Under Assumption 1–Assumption 3, we define the constraint sets

(3.10)
$$\mathcal{D}_{\boldsymbol{x}} := \cup_{\boldsymbol{x}^{\star} \in \mathcal{K}_{\boldsymbol{x}}} \mathcal{D}_f(\boldsymbol{x}^{\star}), \ \mathcal{D}_{\boldsymbol{v}} := \cup_{\boldsymbol{v}^{\star} \in \mathcal{K}_{\boldsymbol{v}}} \mathcal{D}_g(\boldsymbol{v}^{\star})$$

(3.11)
$$\mathcal{D}_{\boldsymbol{x}}^* := \mathcal{D}_{\boldsymbol{x}} \cap \mathbb{S}^{n-1} = \cup_{\boldsymbol{x}^\star \in \mathcal{K}_{\boldsymbol{x}}} \mathcal{D}_f^*(\boldsymbol{x}^\star), \ \mathcal{D}_{\boldsymbol{v}}^* := \mathcal{D}_{\boldsymbol{v}} \cap \mathbb{S}^{m-1} = \cup_{\boldsymbol{v}^\star \in \mathcal{K}_{\boldsymbol{v}}} \mathcal{D}_g^*(\boldsymbol{v}^\star).$$

Suppose that the positive scalars (ζ, ρ_1, ρ_2) and the sample size m satisfy

(3.12)
$$\zeta = \frac{4\delta(\mathscr{H}(\mathcal{K}_{\boldsymbol{x}},\rho_1) + \mathscr{H}(\mathcal{K}_{\boldsymbol{v}},\rho_2))}{m},$$

(3.13)
$$\rho_1 \le \frac{c_1 \zeta}{(\log \frac{\delta}{\zeta})^{1/2}}, \ \omega \left((\mathcal{K}_{\boldsymbol{x}})_{\text{loc}}^{(\rho_1)} \right) \le c_2 \zeta \sqrt{\frac{m\zeta}{\delta}}, \ \rho_2 \le c_3 \zeta \sqrt{\frac{\zeta}{\delta}}$$

for small enough (c_1, c_2, c_3) . If for large enough C_4 it holds that

(3.14)
$$m \ge C_4 \Big(\gamma^2(\mathcal{D}_{\boldsymbol{x}}^*) + \gamma^2(\mathcal{D}_{\boldsymbol{v}}^*) + \mathscr{H}(\mathcal{K}_{\boldsymbol{x}}, \rho_1) + \mathscr{H}(\mathcal{K}_{\boldsymbol{v}}, \rho_2) \Big)$$

then with probability exceeding

$$(3.15) \qquad 1 - 12 \exp\left(-\Omega(\mathscr{H}(\mathcal{K}_{\boldsymbol{x}}, \rho_1) + \mathscr{H}(\mathcal{K}_{\boldsymbol{v}}, \rho_2))\right) - 6 \exp\left(-\Omega(\gamma^2(\mathcal{D}_{\boldsymbol{x}}^*) + \gamma^2(\mathcal{D}_{\boldsymbol{v}}^*))\right)$$

on a single draw of (Φ, ϵ, τ) , the following uniform error bound holds true for all $(x^{\star}, v^{\star}) \in \mathcal{K}_{x} \times \mathcal{K}_{v}$:

(3.16)
$$\sqrt{\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{2}^{2} + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2}} \lesssim \frac{(E+\delta)\big(\gamma(\mathcal{D}_{\boldsymbol{x}}^{*}) + \gamma(\mathcal{D}_{\boldsymbol{v}}^{*})\big) + \delta(\mathscr{H}(\mathcal{K}_{\boldsymbol{x}},\rho_{1}) + \mathscr{H}(\mathcal{K}_{\boldsymbol{v}},\rho_{2}))^{1/2}}{\sqrt{m}},$$

where $\Delta_x = \hat{x} - x^*$ and $\Delta_v = \hat{v} - v^*$, with (\hat{x}, \hat{v}) being the solution to (3.6)

Proof. We present the proof in three steps.

Step 1: Problem Reduction Our first step is to reduce the problem to bounding some random processes. We assume that Δ_x and Δ_v are non-zero with no loss of generality.

Using Optimality: Starting with the optimality of (\hat{x}, \hat{v}) that implies

(3.17)
$$\|\dot{\boldsymbol{y}} - \boldsymbol{\Phi}\hat{\boldsymbol{x}} - \sqrt{m}\hat{\boldsymbol{v}}\|_2^2 \le \|\dot{\boldsymbol{y}} - \boldsymbol{\Phi}\boldsymbol{x}^{\star} - \sqrt{m}\boldsymbol{v}^{\star}\|_2^2,$$

we substitute $\hat{x} = x^* + \Delta_x$ and $\hat{v} = v^* + \Delta_v$ into the left-hand side and then expand the square to obtain

(3.18)
$$\|\boldsymbol{\Phi}\boldsymbol{\Delta}_{\boldsymbol{x}} + \sqrt{m}\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2} \leq 2\langle \boldsymbol{\dot{y}} - (\boldsymbol{\Phi}\boldsymbol{x}^{\star} + \sqrt{m}\boldsymbol{v}^{\star}), \boldsymbol{\Phi}\boldsymbol{\Delta}_{\boldsymbol{x}} + \sqrt{m}\boldsymbol{\Delta}_{\boldsymbol{v}} \rangle$$

(3.19)
$$= 2\langle \boldsymbol{\epsilon}, \boldsymbol{\Phi} \boldsymbol{\Delta}_{\boldsymbol{x}} + \sqrt{m} \boldsymbol{\Delta}_{\boldsymbol{v}} \rangle + 2\langle \boldsymbol{\xi}_{\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star}}, \boldsymbol{\Phi} \boldsymbol{\Delta}_{\boldsymbol{x}} + \sqrt{m} \boldsymbol{\Delta}_{\boldsymbol{v}} \rangle,$$

where in (3.19) we substitute (3.4), and recall that $\boldsymbol{\xi}_{\boldsymbol{x}^{\star},\boldsymbol{v}^{\star}}$ is the quantization noise associated with $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star})$ defined in (3.3). In pursuit of uniform error bound, we must ensure that each step proceeds universally for all $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star}) \in \mathcal{K}_{\boldsymbol{x}} \times \mathcal{K}_{\boldsymbol{v}}$; for clarity, we will take supremum/infimum at this early stage, which requires us to identify contraint sets that accommodate the estimation errors.

Identifying Constraint Sets: Because $f(\hat{x}) \leq f(x^*)$, $g(\hat{v}) \leq g(v^*)$, we have $\Delta_x \in \mathcal{D}_f(x^*)$ and $\Delta_v \in \mathcal{D}_g(v^*)$. Combining with \mathcal{D}_x and \mathcal{D}_v defined in (3.10), we have

(3.20)
$$(\boldsymbol{\Delta}_{\boldsymbol{x}}, \boldsymbol{\Delta}_{\boldsymbol{v}}) \in \mathcal{D}_{\boldsymbol{x}, \boldsymbol{v}} := \mathcal{D}_{\boldsymbol{x}} \times \mathcal{D}_{\boldsymbol{v}}, \ \forall (\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star}) \in \mathcal{K}_{\boldsymbol{x}} \times \mathcal{K}_{\boldsymbol{v}}.$$

Besides $\mathcal{D}_{\boldsymbol{x},\boldsymbol{v}}$ in (3.20), we further define its localized version $\mathcal{D}_{\boldsymbol{x},\boldsymbol{v}}^* = \mathcal{D}_{\boldsymbol{x},\boldsymbol{v}} \cap \mathbb{S}^{m+n-1}$; along with (3.11), we have that

(3.21)
$$\frac{\boldsymbol{\Delta}_{\boldsymbol{x}}}{\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{2}} \in \mathcal{D}_{\boldsymbol{x}}^{*}, \ \frac{\boldsymbol{\Delta}_{\boldsymbol{v}}}{\|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}} \in \mathcal{D}_{\boldsymbol{v}}^{*}, \ \frac{(\boldsymbol{\Delta}_{\boldsymbol{x}}, \boldsymbol{\Delta}_{\boldsymbol{v}})}{(\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{2}^{2} + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2})^{1/2}} \in \mathcal{D}_{\boldsymbol{x},\boldsymbol{v}}^{*}$$

holds universally for all $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star}) \in \mathcal{K}_{\boldsymbol{x}, \boldsymbol{v}}$.

Bounding Both Sides of (3.18): Uniformly for all (x^*, v^*) , the left-hand side of (3.18) is lower bounded by

(3.22)
$$\left(\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{2}^{2} + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2} \right) \cdot \inf_{(\boldsymbol{a},\boldsymbol{b})\in\mathcal{D}_{\boldsymbol{x},\boldsymbol{v}}^{*}} \left\| \boldsymbol{\Phi}\boldsymbol{a} + \sqrt{m}\boldsymbol{b} \right\|_{2}^{2} := \left(\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{2}^{2} + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2} \right) \cdot I_{1},$$

and (3.19) is (upper) bounded by

(3.23)
$$2\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{2} \cdot \sup_{\boldsymbol{c}\in\mathcal{D}_{\boldsymbol{x}}^{*}} \langle \boldsymbol{\epsilon}, \boldsymbol{\Phi}\boldsymbol{c} \rangle + 2\|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2} \cdot \sup_{\boldsymbol{d}\in\mathcal{D}_{\boldsymbol{v}}^{*}} \langle \boldsymbol{\epsilon}, \sqrt{m}\boldsymbol{d} \rangle$$

(3.24)
$$+ 2\left(\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{2}^{2} + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2}\right)^{1/2} \cdot \sup_{\boldsymbol{a} \in \mathcal{K}_{\boldsymbol{x}}} \sup_{\boldsymbol{b} \in \mathcal{K}_{\boldsymbol{v}}} \sup_{(\boldsymbol{c}, \boldsymbol{d}) \in \mathcal{D}_{\boldsymbol{x}, \boldsymbol{v}}^{*}} \langle \boldsymbol{\xi}_{\boldsymbol{a}, \boldsymbol{b}}, \boldsymbol{\Phi} \boldsymbol{c} + \sqrt{m} \boldsymbol{d} \rangle$$

(3.25) := 2
$$\| \boldsymbol{\Delta}_{\boldsymbol{x}} \|_{2} \cdot I_{2} + 2 \| \boldsymbol{\Delta}_{\boldsymbol{v}} \|_{2} \cdot I_{3} + 2 (\| \boldsymbol{\Delta}_{\boldsymbol{x}} \|_{2}^{2} + \| \boldsymbol{\Delta}_{\boldsymbol{v}} \|_{2}^{2})^{1/2} \cdot I_{4},$$

This manuscript is for review purposes only.

where in (3.24) we introduce a generic notation for quantization noise similar to (3.3):

(3.26)
$$\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}} := \mathcal{Q}_{\delta}(\boldsymbol{\Phi}\boldsymbol{a} + \sqrt{m}\boldsymbol{b} + \boldsymbol{\epsilon} + \boldsymbol{\tau}) - (\boldsymbol{\Phi}\boldsymbol{a} + \sqrt{m}\boldsymbol{b} + \boldsymbol{\epsilon}),$$

and for convenience we denote the random processes that arose by I_1, I_2, I_3, I_4 . Therefore, we obtain that

(3.27)
$$(\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{2}^{2} + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2}) \cdot I_{1} \leq 2 \Big(\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{2} \cdot I_{2} + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2} \cdot I_{3} + \big(\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{2}^{2} + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2}\big)^{1/2} \cdot I_{4}\Big)$$

holds uniformly for all $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star}) \in \mathcal{K}_{\boldsymbol{x}} \times \mathcal{K}_{\boldsymbol{v}}$.

Step 2: Bounding I_1, I_2, I_3, I_4 With (3.27), all that remains is to bound I_1, I_2, I_3, I_4 from the correct side. We provide a sketch of our techniques in this step:

- We apply the extended matrix deviation inequality (Proposition A.1) to get a lower bound on I_1 ;
- We apply Proposition A.4 with the randomness of Φ to bound I_2 ;
- We apply Proposition A.2 with the randomness of ϵ to bound I_3 ;
- We apply the global QPE property for structured sets (Corollary B.4) to bound I_4 .

Bounding I_1 : For any $t \ge 0$, Proposition A.1 yields that the event

(3.28)
$$\sup_{(\boldsymbol{a},\boldsymbol{b})\in\mathcal{D}^*_{\boldsymbol{x},\boldsymbol{v}}} \left| \|\boldsymbol{\Phi}\boldsymbol{a} + \sqrt{m}\boldsymbol{b}\|_2 - \sqrt{m} \right| \le C \left(\gamma(\mathcal{D}^*_{\boldsymbol{x},\boldsymbol{v}}) + t \right)$$

holds with probability exceeding $1 - \exp(-t^2)$. Note that Proposition A.7 gives $\gamma(\mathcal{D}_{\boldsymbol{x},\boldsymbol{v}}^*) \approx \gamma(\mathcal{D}_{\boldsymbol{x}}^*) + \gamma(\mathcal{D}_{\boldsymbol{v}}^*)$, so the sample complexity (3.14) implies $m \gtrsim \gamma^2(\mathcal{D}_{\boldsymbol{x},\boldsymbol{v}}^*)$, and we can set $t = \gamma(\mathcal{D}_{\boldsymbol{x},\boldsymbol{v}}^*)$ in (3.28) and obtain that the event

(3.29)
$$\sup_{(\boldsymbol{a},\boldsymbol{b})\in\mathcal{D}^*_{\boldsymbol{x},\boldsymbol{v}}} \left| \|\boldsymbol{\Phi}\boldsymbol{a} + \sqrt{m}\boldsymbol{b}\|_2 - \sqrt{m} \right| \leq \frac{\sqrt{m}}{2}$$

holds with probability exceeding $1 - \exp(-\gamma^2(\mathcal{D}^*_{\boldsymbol{x},\boldsymbol{v}}))$. Combining with triangle inequality, (3.29) gives

(3.30)
$$\sqrt{I_1} = \inf_{(\boldsymbol{a},\boldsymbol{b})\in\mathcal{D}^*_{\boldsymbol{x},\boldsymbol{v}}} \left\|\boldsymbol{\Phi}\boldsymbol{a} + \sqrt{m}\boldsymbol{b}\right\|_2$$

(3.31)
$$\geq \sqrt{m} - \sup_{(\boldsymbol{a},\boldsymbol{b})\in\mathcal{D}^*_{\boldsymbol{x},\boldsymbol{v}}} \left| \|\boldsymbol{\Phi}\boldsymbol{a} + \sqrt{m}\boldsymbol{b}\|_2 - \sqrt{m} \right| \geq \frac{\sqrt{m}}{2},$$

thus yielding the desired lower bound on I_1 : $I_1 \geq \frac{m}{4}$.

Bounding I_2 : Conditioning on ϵ , we invoke Proposition A.4 to obtain that for any $t \ge 0$, the event

(3.32)
$$I_2 \le C_1 \|\boldsymbol{\epsilon}\|_2 \big(\omega(\mathcal{D}_{\boldsymbol{x}}^*) + t \big)$$

holds with probability exceeding $1 - 2\exp(-t^2)$. We set $t = \gamma(\mathcal{D}^*_{\boldsymbol{x},\boldsymbol{v}})$ to obtain that the event $I_2 \leq C_1 \|\boldsymbol{\epsilon}\|_2 \cdot \omega(\mathcal{D}^*_{\boldsymbol{x},\boldsymbol{v}})$ holds with probability exceeding $1 - 2\exp(-\gamma^2(\mathcal{D}^*_{\boldsymbol{x},\boldsymbol{v}}))$. Then we deal

with the randomness of ϵ . By $\|\epsilon\|_{\psi_2} = O(E)$ from Assumption 1, we can use [78, Exercise 6.3.5] to bound $\|\epsilon\|_2$ and obtain

(3.33)
$$\mathbb{P}\left(\|\boldsymbol{\epsilon}\|_{2} \geq C_{2}E\sqrt{m} + t\right) \leq \exp\left(-\frac{c_{3}t^{2}}{E^{2}}\right)$$

for any $t \ge 0$. We set $t \simeq E\sqrt{m}$ to obtain that $\|\boldsymbol{\epsilon}\|_2 \le E\sqrt{m}$ holds with probability exceeding $1 - \exp(-\Omega(m))$. Therefore, we arrive at the desired bound

$$(3.34) I_2 \lesssim E\sqrt{m} \cdot \gamma(\mathcal{D}^*_{\boldsymbol{x},\boldsymbol{v}})$$

that holds with probability exceeding $1 - 2\exp(-\gamma^2(\mathcal{D}^*_{\boldsymbol{x},\boldsymbol{v}})) - \exp(-\Omega(m))$.

Bounding I_3 : For any $d_1, d_2 \in \mathcal{D}^*_{\boldsymbol{v}} \cup \{0\}$ we have

(3.35)
$$\|\langle \boldsymbol{\epsilon}, \sqrt{m}\boldsymbol{d}_1 \rangle - \langle \boldsymbol{\epsilon}, \sqrt{m}\boldsymbol{d}_2 \rangle\|_{\psi_2} = \sqrt{m} \|\langle \boldsymbol{\epsilon}, \boldsymbol{d}_1 - \boldsymbol{d}_2 \rangle\|_{\psi_2} \le E\sqrt{m} \|\boldsymbol{d}_1 - \boldsymbol{d}_2\|_2.$$

Thus, Proposition A.2 implies that for any $t \ge 0$, the event $I_3 \le C_4 E \sqrt{m}(\omega(\mathcal{D}_{\boldsymbol{v}}^*) + t)$ holds with probability exceeding $1 - 2\exp(-t^2)$. Setting $t = \gamma(\mathcal{D}_{\boldsymbol{x},\boldsymbol{v}}^*)$ gives

$$(3.36) I_3 \lesssim E\sqrt{m} \cdot \gamma(\mathcal{D}^*_{\boldsymbol{x},\boldsymbol{v}})$$

that holds with probability at least $1 - 2 \exp(-\gamma^2(\mathcal{D}^*_{\boldsymbol{x},\boldsymbol{v}}))$.

Bounding I_4 : This is the most challenging part in our analysis, but we leave the development of QPE to Appendix B to allow for a clean analysis in the main body. With (3.12), (3.13) and (3.14) we can apply Corollary B.4 to obtain that

(3.37)
$$I_4 \le C_5 \delta \sqrt{m} \Big(\omega(\mathcal{D}^*_{\boldsymbol{x}, \boldsymbol{v}}) + \sqrt{\mathscr{H}(\mathcal{K}_{\boldsymbol{x}}, \rho_1) + \mathscr{H}(\mathcal{K}_{\boldsymbol{v}}, \rho_2)} \Big)$$

holds with probability at least $1 - 12 \exp(-\Omega(\mathscr{H}(\mathcal{K}_{\boldsymbol{x}}, \rho_1) + \mathscr{H}(\mathcal{K}_{\boldsymbol{v}}, \rho_2)))$.

Step 3: Combining Everything We are in the position to combine everything together to conclude the proof. Substituting the bounds (3.31) on I_1 , (3.34), (3.36) and (3.37) on (I_2, I_3, I_4) into (3.27) yields

(3.38)
$$m \left(\| \boldsymbol{\Delta}_{\boldsymbol{x}} \|_{2}^{2} + \| \boldsymbol{\Delta}_{\boldsymbol{v}} \|_{2}^{2} \right)$$
$$\lesssim \sqrt{m} \left(\| \boldsymbol{\Delta}_{\boldsymbol{x}} \|_{2}^{2} + \| \boldsymbol{\Delta}_{\boldsymbol{v}} \|_{2}^{2} \right)^{1/2} \left((E + \delta) \cdot \gamma(\mathcal{D}_{\boldsymbol{x},\boldsymbol{v}}^{*}) + \delta \sqrt{\mathscr{H}(\mathcal{K}_{\boldsymbol{x}},\rho_{1}) + \mathscr{H}(\mathcal{K}_{\boldsymbol{v}},\rho_{2})} \right)$$

that holds uniformly for all $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star}) \in \mathcal{K}_{\boldsymbol{x}} \times \mathcal{K}_{\boldsymbol{v}}$. Rearranging, along with $\gamma(\mathcal{D}_{\boldsymbol{x},\boldsymbol{v}}^{\star}) \asymp \gamma(\mathcal{D}_{\boldsymbol{x}}^{\star}) + \gamma(\mathcal{D}_{\boldsymbol{v}}^{\star})$ from Proposition A.7, yields the desired bound (3.16). All that remains is to count the probability terms: We rule out probability terms of $\exp(-\gamma^2(\mathcal{D}_{\boldsymbol{x},\boldsymbol{v}}^{\star}))$ to ensure (3.31), $2\exp(-\gamma^2(\mathcal{D}_{\boldsymbol{x},\boldsymbol{v}}^{\star})) + \exp(-\Omega(m))$ for (3.34), $2\exp(-\gamma^2(\mathcal{D}_{\boldsymbol{x},\boldsymbol{v}}^{\star}))$ for (3.36), $12\exp(-\Omega(\mathcal{H}(\mathcal{K}_{\boldsymbol{x}},\rho_1) + \mathcal{H}(\mathcal{K}_{\boldsymbol{v}},\rho_2)))$ for (3.37). Combining with $\gamma^2(\mathcal{D}_{\boldsymbol{x},\boldsymbol{v}}^{\star}) \asymp \gamma^2(\mathcal{D}_{\boldsymbol{x}}^{\star}) + \gamma^2(\mathcal{D}_{\boldsymbol{v}}^{\star})$ and (3.14), we can promise that the uniform error bound holds with the probability stated in (3.15).

Remark 3.3 (The Cost of Uniformity: Constrained Lasso with Structured Priors). To see the implication of Theorem 3.2 on structured priors (Assumption 3), we substitute (3.9) into (3.16) to obtain the uniform ℓ_2 -norm error bound

(3.39)
$$\tilde{O}\left(\frac{(E+\delta)[\gamma(\mathcal{D}_{\boldsymbol{x}}^*)+\gamma(\mathcal{D}_{\boldsymbol{v}}^*)]+\delta\cdot[\omega(\mathcal{K}_{\boldsymbol{x}})+\omega(\mathcal{K}_{\boldsymbol{v}})]}{\sqrt{m}}\right),$$

where we use $O(\cdot)$ to omit some logarithmic factors on (ρ_1, ρ_2) . We compare (3.39) with the non-uniform bound [75, Thm. 1]

(3.40)
$$O\left(\frac{(E+\delta)\left[\omega\left(\mathcal{D}_{f}^{*}(\boldsymbol{x}^{\star})\right)+\gamma\left(\mathcal{D}_{g}^{*}(\boldsymbol{v}^{\star})\right)\right]}{\sqrt{m}}\right)$$

and elaborate the cost of uniformity by noting two differences:

- First, the term $\omega(\mathcal{D}_{f}^{*}(\boldsymbol{x}^{\star})) + \gamma(\mathcal{D}_{g}^{*}(\boldsymbol{v}^{\star}))$ regarding some fixed $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star})$ in (3.40) is substituted with $\gamma(\mathcal{D}_{\boldsymbol{x}}^{*}) + \gamma(\mathcal{D}_{\boldsymbol{v}}^{*})$ in (3.39), where $\mathcal{D}_{\boldsymbol{x}}^{*}$ and $\mathcal{D}_{\boldsymbol{v}}^{*}$ are defined in (3.11). This appears rather natural to us in uniform recovery, the local complexity quantity (e.g., $\omega(\mathcal{D}_{f}^{*}(\boldsymbol{x}^{\star})))$ upgrades to a global one (e.g., $\omega(\mathcal{D}_{\boldsymbol{x}}^{*}))$ concerning all signals and corruptions.
- Second, our uniform bound also presents the additional term $\tilde{O}\left(\frac{\delta}{\sqrt{m}}\left[\omega(\mathcal{K}_{\boldsymbol{x}}) + \omega(\mathcal{K}_{\boldsymbol{v}})\right]\right)$. Note that this term vanishes in non-uniform recovery where $\mathcal{K}_{\boldsymbol{x}}$ and $\mathcal{K}_{\boldsymbol{v}}$ only contain one point. Thus, we can think of that (3.40) also implicitly includes this term, and from this viewpoint the two bounds (3.39) and (3.40) stay consistent.

We further note that similar phenomena were also observed from the results of [36] (e.g., Theorem 1 therein). As we shall see, in the most interesting cases of structured priors (such as sparsity and low-rank), (3.39) and (3.40) are typically of the same scaling up to logarithmic factors, indicating that the uniformity costs very little.

Remark 3.4 (The Role of Quantization Resolution δ). Due to random dithering, the quantization resolution δ appears in (3.16) as multiplicative factors, agreeing with similar findings in [17, 20, 75, 77, 81] and confirming the intuition that the recovery worsens under coarser quantization (i.e., larger δ).

To illustrate the implications of Theorem 3.2, we provide two concrete examples in Corollary 3.5 and Corollary 3.7. The proofs of these corollaries can be found in Appendix C. Equipped with the general Theorem 3.2, the proofs can be done by selecting (ρ_1, ρ_2) and estimating the geometric quantities. Due to the feature of structured signal sets as per (3.9), using some extremely small (ρ_1, ρ_2) to render (3.13) only leads to logarithmic degradation. That being said, we still (slightly) refine the choice of (ρ_1, ρ_2) to lessen the logarithmic factor (our delicate QPE result Corollary B.4 allows us to do so); see Remark 3.6, Remark 3.8 below.

Corollary 3.5 (Sparse Signal and Sparse Corruption). Under Assumption 1–Assumption 3, we assume that \mathbf{x}^{\star} is s-sparse and \mathbf{v}^{\star} is k-sparse, i.e., $\mathcal{K}_{\mathbf{x}} = \Sigma_s^n \cap \mathbb{B}_2^n$ and $\mathcal{K}_{\mathbf{v}} = \Sigma_k^m \cap \mathbb{B}_2^m$ in Assumption 3, and accordingly we use $f(\mathbf{x}) = \|\mathbf{x}\|_1$ and $g(\mathbf{v}) = \|\mathbf{v}\|_1$ in Assumption 2. If $m \geq C_1 s \log(\frac{nm^{3/2}}{s^{5/2}\delta}) + C_1 k \log(\frac{m^{5/2}}{k^{5/2}\delta})$ holds with large enough C_1 , then with probability exceeding $1 - C_2 \exp(-\Omega(s \log \frac{en}{s} + k \log \frac{em}{k}))$ on a single draw of $(\mathbf{\Phi}, \boldsymbol{\epsilon}, \boldsymbol{\tau})$, the following uniform error bound holds true for all $(\mathbf{x}^{\star}, \mathbf{v}^{\star}) \in \mathcal{K}_{\mathbf{x}} \times \mathcal{K}_{\mathbf{v}}$:

(3.41)
$$\sqrt{\|\mathbf{\Delta}_{x}\|_{2}^{2} + \|\mathbf{\Delta}_{v}\|_{2}^{2}} \lesssim \frac{E\sqrt{s\log\frac{en}{s} + k\log\frac{em}{k}} + \delta\sqrt{s\log\frac{nm^{3/2}}{s^{5/2}\delta} + k\log\frac{m^{5/2}}{k^{5/2}\delta}}}{\sqrt{m}}$$

where $\Delta_x = \hat{x} - x^*$, $\Delta_v = \hat{v} - v^*$, with (\hat{x}, \hat{v}) being the solution to (3.6).

Remark 3.6 (Elimination of Logarithmic Factors). Provided the additional scaling conditions $m \leq n$ and $\delta \geq \left(\frac{\min\{s,k\}}{m}\right)^N$ for some positive integer N (note that they are very mild and

cover most interesting settings), we have $\log(\frac{nm^{3/2}}{s^{5/2}\delta}) \lesssim \log(\frac{n}{s})$ and $\log(\frac{m^{5/2}}{k^{5/2}\delta}) \lesssim \log(\frac{m}{k})$. Thus notably, our Corollary 3.5 coincides with its non-uniform counterpart in [75, Coro. 1] without suffering from logarithmic degradation (rather, the cost is at most a larger multiplicative constant which both works do not aim to refine).

Corollary 3.7 (Low-Rank Signal and Sparse Corruption). Under Assumption 1–Assumption 3, we assume that $\mathbf{x}^{\star} \in \mathbb{R}^{p \times q}$ is of rank no greater than r, 5 and \mathbf{v}^{\star} is k-sparse, i.e., $\mathcal{K}_{\mathbf{x}} = M_r^{p,q} \cap \mathbb{B}_F^{p,q}$ and $\mathcal{K}_{\mathbf{v}} = \Sigma_k^m \cap \mathbb{B}_2^m$ in Assumption 3, and accordingly we use $f(\mathbf{x}) = \|\mathbf{x}\|_{\mathrm{nu}}$ and $g(\mathbf{v}) = \|\mathbf{v}\|_1$ in Assumption 2. If $m \geq C_1 r(p+q) \log(\frac{m^{3/2}}{\delta(r(p+q))^{3/2}}) + C_1 k \log(\frac{m^{5/2}}{k^{5/2}\delta})$ for some large enough C_1 , then with probability exceeding $1 - C_2 \exp(-\Omega(r(p+q) + k \log \frac{em}{k})))$ on a single draw of $(\mathbf{\Phi}, \boldsymbol{\epsilon}, \boldsymbol{\tau})$, the following uniform error bound holds true for all $(\mathbf{x}^{\star}, \mathbf{v}^{\star}) \in \mathcal{K}_{\mathbf{x}} \times \mathcal{K}_{\mathbf{v}}$: (3.42)

$$\sqrt{\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{\mathrm{F}}^{2} + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2}} \lesssim \frac{E\sqrt{r(p+q) + k\log\frac{em}{k}} + \delta\sqrt{r(p+q)\log(\frac{m^{3/2}}{\delta(r(p+q))^{3/2}}) + k\log(\frac{m^{5/2}}{k^{5/2}\delta})}{\sqrt{m}}$$

where $\Delta_x = \hat{x} - x^*$, $\Delta_v = \hat{v} - v^*$, with (\hat{x}, \hat{v}) being the solution to (3.6).

Remark 3.8. Analogously to Remark 3.6, provided the additional scaling conditions of $\delta \gtrsim (\frac{r(p+q)}{m})^N$ and $\delta \gtrsim (\frac{k}{m})^N$ for some positive integer N, one can further simplify (3.42) to

(3.43)
$$\frac{\delta\sqrt{r(p+q)\log(\frac{m}{r(p+q)})} + E\sqrt{r(p+q)} + (\delta+E)\sqrt{k\log\frac{em}{k}}}{\sqrt{m}}$$

This only exhibits an additional factor of $\log^{1/2}(\frac{m}{r(p+q)})$ compared to the non-uniform counterpart in [75, Coro. 2].

We close this subsection by comparing with relevant results and claiming our contributions.

Remark 3.9 (Related Works and the Novelty of Our Results). Restricted to Lasso, the prior developments are as follows:

- Non-uniform guarantees were presented in [77, Thm. III.1] for compressed sensing and in [75, Thm. 1] for corrupted sensing.
- The only existing uniform guarantee for constrained Lasso was obtained in [36, Coro. 4], but only applies to classical compressed sensing (without the need of recovering an additional structured corruption) and typically yields an error rate of $O(\sqrt{\delta}[\frac{\omega^2(\mathcal{K}_x)}{m}]^{1/4})$, which is slower than our (3.39) under structured priors (Assumption 3). We will further note that it may not be sensible (if not impossible) to follow the proof technique in [36] to prove Theorem 3.2; see Remark 3.11.

Note that constrained Lasso is a general recipe for nonlinear compressed sensing models [35, 55, 67], but there have also been uniform guarantees for other recovery methods (see also the less extensive discussion in [36, Sec. 4.3]):

• Jung et al. devised and theoretically analyzed a more specialized recovery method for quantized compressed sensing [46, Thm. 3]. However, under the dithered uniform quantizer, their uniform error rate translates into $O\left(\left[\frac{\delta\omega^2(\mathcal{K}_x)}{m}\right]^{1/3}\right)$ in the worst case, and it

⁵When substituted into (3.1), we view \boldsymbol{x}^{\star} as a (n := pq)-dimensional vector by vectorization.

is unknown whether their result can be sharpened for structured sets like Σ_s^n since their statement requires \mathcal{K}_x to be convex;

• Xu and Jacques analyzed the projected-back projection (PBP) estimator in [81, Sec. 7.3A], providing a rate of $\tilde{O}((1+\delta)\frac{\omega(\mathcal{K}_x)}{\sqrt{m}})$ (logarithmic factors omitted) for structured sets. Although the rate is comparable to ours, one downside of the PBP estimator is that it does not achieve exact reconstruction in a noiseless unquantized case [81, Sec. 7.3C]. (In contrast, Lasso achieves exact reconstruction in a noiseless unquantized case; see [14, 33] for instance.)

In a nutshell, in compressed/corrupted sensing associated with the dithered uniform quantizer, our Theorem 3.2 presents the *sharpest* uniform error rate over structured set (Definition 3.1), and note that this is achieved by constrained Lasso which returns exact recovery in a noisyless unquantized setting.

Remark 3.10 (Optimality). We note that the rates in Corollary 3.5 and Corollary 3.7 are near minimax optimal when the sub-Gaussian noise ϵ is severer than the quantization noise (i.e., $E \gtrsim \delta$). Such optimality is implied by adding together the lower bounds (e.g., from [68, Thm. 4.2]) for the two estimation problems $y_1 = \Phi x^* + \epsilon$ and $y_2 = \sqrt{m}v^* + \epsilon$.⁶ Nonetheless, for the noiseless case with quantization (i.e., $E = 0, \delta > 0$), the information theoretic limit exhibits a decaying rate of $O(m^{-1})$ that is faster than our $O(m^{-1/2})$ (e.g., [8]), and we suspect that such faster rate cannot be achieved by Lasso due to some fundamental performance limit.⁷

Remark 3.11 (Technical Comparison with [36]). Genzel and Stollenwerk [36] developed a general strategy to achieve uniform recovery, which consists of two ingredients: (i) Constructing Lipschitz approximation for handling the discontinuity of $f_i(\cdot)$ (if any), and (ii) Applying the concentration inequality [36, Thm. 8] (due to Mendelson [58]) to bounding the product processes arising in the analysis. However, it might not be sensible (if not impossible) to follow their techniques to prove Theorem 3.2 for two reasons. First, when $f_i(\cdot)$ contains some discontinuity (as with our (1.2)), their general strategy leads to a uniform decaying rate of $O(m^{-1/4})$ (see [36, Sec. 4]) that is inferior to our $O(m^{-1/2})$, and it is unclear how to get faster uniform rate without incorporating existing embedding result available in the literature.⁸ Second, the appearance of the corruption v^* poses additional hurdle to the approach in [36], since v^* leads to random processes beyond the scope of [36, Thm. 8]. Conversely, our work of getting $O(m^{-1/2})$ decaying rate based on QPE suggests the possibility of improving the slow rate of $O(m^{1/4})$ in [36] under more general discontinuous $f_i(\cdot)$ — one may deal with discontinuity of $f_i(\cdot)$ by proper product embedding property (or limited projection distortion as termed by [81]) rather than constructing Lipschitz approximation.

⁶The minimax lower bound for $y_1 = \Phi x^* + \epsilon$ applies to the estimation of x^* from (3.2), since the additional corruption and quantization can only decrease our ability to estimate x^* . Similarly, the minimax lower bound for $y_2 = \sqrt{m}v^* + \epsilon$ stands when estimating v^* from (3.2).

⁷Though we are not aware of a rigorous analysis, to our best knowledge, all proved rates for *Lasso* in quantized compressed sensing with memorylesss quantizer are no faster than $O(m^{-1/2})$; see similar discussion in [36, P. 34].

⁸This is a workaround proposed in [36, Sec. 5]. Nonetheless, the needed embedding result may not exist in the literature for the problem at hand, and this is the case for our quantized corrupted sensing problem.

3.2 Structured Priors with Unconstrained Lasso We now turn our attention to unconstrained Lasso (3.7), which is more practical than the constrained Lasso in (3.6) in the sense that it does not require prior estimates of $(f(\boldsymbol{x}^*), g(\boldsymbol{v}^*))$. Rather, as we shall see, a fixed large enough choice of the regularization parameters (λ_1, λ_2) works uniformly for all pairs of signal and corruption. To proceed, we first define the restricted compatibility constant between $f(\cdot)$ and ℓ_2 -norm over some constraint set $\mathcal{X} \subset \mathbb{R}^n$ as

(3.44)
$$\alpha_f(\mathcal{X}) = \sup_{\boldsymbol{x}\in\mathcal{X}} \frac{f(\boldsymbol{x})}{\|\boldsymbol{x}\|_2}.$$

Compared to constrained Lasso, the analysis of unconstrained Lasso is more technical in the following senses:

- The derivation of a low-complexity constraint set that contains the estimation error becomes non-trivial, as contrasted to the straightforward $\Delta_x \in \mathcal{D}_f(x^*)$ and $\Delta_v \in \mathcal{D}_g(v^*)$ for constrained Lasso;
- Some other additional efforts are needed, e.g., bounding the compatibility constant that is in general technically challenging.

Without pursuing full generality, we make the following decomposable assumption on $f(\cdot)$ and $g(\cdot)$ to facilitate the estimation of compatibility constant.

Assumption 4 (Decomposable Norm). Regarding the sets $\mathcal{K}_{\boldsymbol{x}}, \mathcal{K}_{\boldsymbol{v}}$ and associated norms $f(\cdot), g(\cdot)$ for promoting certain structure (Assumption 2), we assume that:

• (Decomposibility) Given any $a \in \mathcal{K}_{x}$, there exists a pair of linear subspaces $(\mathcal{X}_{a}, \overline{\mathcal{X}}_{a})$ (possibly depending on a) with $\mathcal{X}_{a} \subset \overline{\mathcal{X}}_{a}$, such that $a \in \mathcal{X}_{a}$, and $f(\cdot)$ is decomposable over $(\mathcal{X}_{a}, \overline{\mathcal{X}}_{a}^{\perp})$.⁹

(3.45)
$$f(\boldsymbol{x}_1 + \boldsymbol{x}_2) = f(\boldsymbol{x}_1) + f(\boldsymbol{x}_2), \quad \forall \ \boldsymbol{x}_1 \in \mathcal{X}_{\boldsymbol{a}}, \boldsymbol{x}_2 \in \overline{\mathcal{X}}_{\boldsymbol{a}}^{\perp}.$$

Similarly, given any $\mathbf{b} \in \mathcal{K}_{\mathbf{v}}$ there exists a pair of linear subspaces $(\mathcal{V}_{\mathbf{b}}, \overline{\mathcal{V}}_{\mathbf{b}})$ (possibly depending on \mathbf{b}) with $\mathcal{V}_{\mathbf{b}} \subset \overline{\mathcal{V}}_{\mathbf{b}}$, such that $\mathbf{b} \in \mathcal{V}_{\mathbf{b}}$, and the decomposibility $g(\mathbf{v}_1 + \mathbf{v}_2) = g(\mathbf{v}_1) + g(\mathbf{v}_2)$ holds for any $\mathbf{v}_1 \in \mathcal{V}_{\mathbf{a}}$, $\mathbf{v}_2 \in \overline{\mathcal{V}}_{\mathbf{b}}^{\perp}$.

• (Uniform Bound on Compatibility Constant) There exist $\alpha_{\boldsymbol{x}}$ and $\alpha_{\boldsymbol{v}}$ such that $\alpha_f(\overline{\mathcal{X}}_{\boldsymbol{a}}) \leq \alpha_{\boldsymbol{x}}$ holds uniformly for all $\boldsymbol{a} \in \mathcal{K}_{\boldsymbol{x}}$, and that $\alpha_g(\overline{\mathcal{V}}_{\boldsymbol{b}}) \leq \alpha_{\boldsymbol{v}}$ holds uniformly for all $\boldsymbol{b} \in \mathcal{K}_{\boldsymbol{v}}$, where $(\overline{\mathcal{X}}_{\boldsymbol{a}}, \overline{\mathcal{V}}_{\boldsymbol{b}})$ are the linear subspaces identified for a specific $(\boldsymbol{a}, \boldsymbol{b}) \in \mathcal{K}_{\boldsymbol{x}} \times \mathcal{K}_{\boldsymbol{v}}$ in the preceding dot point, $\alpha_f(\overline{\mathcal{X}}_{\boldsymbol{a}})$ and $\alpha_g(\overline{\mathcal{V}}_{\boldsymbol{b}})$ are the compatibility constants defined as per (3.44).

It is well-known that this decomposibility assumption is satisfied $\|\cdot\|_1$, $\|\cdot\|_{nu}$, $\|\cdot\|_{\ell_1/\ell_2}$ and so on [59,60], thus covering the most interesting cases of structured priors. As a canonical example, for s-sparse structured prior together with $f(\boldsymbol{x}) = \|\boldsymbol{x}\|_1$, we will let $\mathcal{X}_{\boldsymbol{a}} = \overline{\mathcal{X}}_{\boldsymbol{a}} =$ $\{\boldsymbol{v} \in \mathbb{R}^n : \operatorname{supp}(\boldsymbol{v}) \subset \operatorname{supp}(\boldsymbol{a})\}$ for any $\boldsymbol{a} \in \mathcal{K}_{\boldsymbol{x}} = \Sigma_s^n \cap \mathbb{B}_2^n$, under which it is evident that $\alpha_{\boldsymbol{x}} = \sqrt{s}$ is a uniform bound on $\alpha_f(\overline{\mathcal{X}}_{\boldsymbol{a}})$ (see more details in the proof of Corollary 3.14).

⁹For a given linear subapce \mathcal{X} , we denote its orthogonal complement by \mathcal{X}^{\perp} .

Theorem 3.12 (Uniform Recovery via Unconstrained Lasso). Under Assumption 1–Assumption 4, suppose that the positive scalars (ζ, ρ_1, ρ_2) and the sample size m satisfy

(3.46)
$$\zeta = \frac{4\delta(\mathscr{H}(\mathcal{K}_{\boldsymbol{x}},\rho_1) + \mathscr{H}(\mathcal{K}_{\boldsymbol{v}},\rho_2))}{m},$$

(3.47)
$$\rho_1 \le \frac{c_1 \zeta}{(\log \frac{\delta}{\zeta})^{1/2}}, \ \omega \left((\mathcal{K}_{\boldsymbol{x}})_{\text{loc}}^{(\rho_1)} \right) \le c_2 \zeta \sqrt{\frac{m\zeta}{\delta}}, \ \rho_2 \le c_3 \zeta \sqrt{\frac{\zeta}{\delta}}$$

for small enough (c_1, c_2, c_3) , and we also suppose that $f(\boldsymbol{x}) \geq \|\boldsymbol{x}\|_2$ holds for any $\boldsymbol{x} \in \mathbb{R}^n$, $g(\boldsymbol{v}) \geq \|\boldsymbol{v}\|_2$ holds for any $\boldsymbol{v} \in \mathbb{R}^m$.¹⁰ We set

(3.48)
$$\lambda_1 = C_4(E+\delta)\sqrt{m} \cdot \omega(\mathbb{B}_f^n) + C_4\delta\sqrt{m} \cdot \sqrt{\mathscr{H}(\mathcal{K}_x,\rho_1) + \mathscr{H}(\mathcal{K}_v,\rho_2)},$$

(3.49)
$$\lambda_2 = C_5(E+\delta)\sqrt{m} \cdot \omega(\mathbb{B}_g^m) + C_5\delta\sqrt{m} \cdot \sqrt{\mathscr{H}(\mathcal{K}_{\boldsymbol{x}},\rho_1)} + \mathscr{H}(\mathcal{K}_{\boldsymbol{v}},\rho_2),$$

for some large enough C_4, C_5 . If for some sufficiently large implied constant, it holds that

$$(3.50) \quad m \gtrsim \left(\alpha_{\boldsymbol{x}} + \frac{\lambda_2 \alpha_{\boldsymbol{v}}}{\lambda_1}\right)^2 \omega^2(\mathbb{B}_f^n) + \left(\alpha_{\boldsymbol{v}} + \frac{\lambda_1 \alpha_{\boldsymbol{x}}}{\lambda_2}\right)^2 \omega^2(\mathbb{B}_g^m) + \mathscr{H}(\mathcal{K}_{\boldsymbol{x}}, \rho_1) + \mathscr{H}(\mathcal{K}_{\boldsymbol{v}}, \rho_2),$$

then with probability exceeding

$$(3.51) \quad 1 - C_6 \exp\left(-c_7 \min\left\{\omega^2(\mathbb{B}_f^n), \omega^2(\mathbb{B}_g^m)\right\}\right) - 24 \exp\left(-\Omega(\mathscr{H}(\mathcal{K}_x, \rho_1) + \mathscr{H}(\mathcal{K}_v, \rho_2))\right)$$

on a single draw of (Φ, ϵ, τ) , the following uniform error bound holds true for all $(x^*, v^*) \in \mathcal{K}_x \times \mathcal{K}_v$:

(3.52)
$$\sqrt{\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{2}^{2} + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2}} \lesssim \frac{\lambda_{1}\alpha_{\boldsymbol{x}} + \lambda_{2}\alpha_{\boldsymbol{v}}}{m},$$

where $\Delta_x = \hat{x} - x^*$ and $\Delta_v = \hat{v} - v^*$, with (\hat{x}, \hat{v}) being the solution to (3.7)

Proof. We assume that Δ_x and Δ_v are non-zero without loss of generality. Note that the optimality of (\hat{x}, \hat{v}) gives

$$(3.53) \|\dot{\boldsymbol{y}} - \boldsymbol{\Phi}\hat{\boldsymbol{x}} - \sqrt{m}\hat{\boldsymbol{v}}\|_{2}^{2} + \lambda_{1}f(\hat{\boldsymbol{x}}) + \lambda_{2}g(\hat{\boldsymbol{v}}) \leq \|\dot{\boldsymbol{y}} - \boldsymbol{\Phi}\boldsymbol{x}^{\star} - \sqrt{m}\boldsymbol{v}^{\star}\|_{2}^{2} + \lambda_{1}f(\boldsymbol{x}^{\star}) + \lambda_{2}g(\boldsymbol{v}^{\star}).$$

We perform some calculation and reformulate the inequality as follows:

$$(3.56) \quad \leq 2 \Big(f(\boldsymbol{\Delta}_{\boldsymbol{x}}) \cdot \sup_{\boldsymbol{c} \in \mathbb{B}_{f}^{n}} \langle \boldsymbol{\epsilon} + \boldsymbol{\xi}_{\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star}}, \boldsymbol{\Phi} \boldsymbol{c} \rangle + g(\boldsymbol{\Delta}_{\boldsymbol{v}}) \cdot \sup_{\boldsymbol{d} \in \mathbb{B}_{g}^{m}} \langle \boldsymbol{\epsilon} + \boldsymbol{\xi}_{\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star}}, \sqrt{m} \boldsymbol{d} \rangle \Big)$$

(3.57)
$$+ \lambda_1 \big(f(\boldsymbol{x}^{\star}) - f(\hat{\boldsymbol{x}}) \big) + \lambda_2 \big(g(\boldsymbol{v}^{\star}) - g(\hat{\boldsymbol{v}}) \big)$$

$$(3.58) \leq 2\Big(f(\boldsymbol{\Delta}_{\boldsymbol{x}}) \cdot \big[I_1 + I_2\big] + g(\boldsymbol{\Delta}_{\boldsymbol{v}}) \cdot \big[I_3 + I_4\big]\Big) + \lambda_1\big(f(\boldsymbol{x}^{\star}) - f(\hat{\boldsymbol{x}})\big) + \lambda_2\big(g(\boldsymbol{v}^{\star}) - g(\hat{\boldsymbol{v}})\big)$$

¹⁰This is a very mild condition because $f(\cdot)$ and $g(\cdot)$ are norms that promote low-complexity structure, thus naturally dominating ℓ_2 -norm; see, e.g., [72].

where the first inequality (3.54)–(3.55) is obtained from (3.53) by substituting $\hat{\boldsymbol{x}} = \boldsymbol{x}^{\star} + \Delta_{\boldsymbol{x}}$ and $\hat{\boldsymbol{v}} = \boldsymbol{v}^{\star} + \Delta_{\boldsymbol{v}}$ into $\|\boldsymbol{\dot{y}} - \boldsymbol{\Phi}\hat{\boldsymbol{x}} - \sqrt{m}\hat{\boldsymbol{v}}\|_2^2$ and then expanding the square; then, in (3.56) we substitute $\boldsymbol{\dot{y}} - \boldsymbol{\Phi}\boldsymbol{x}^{\star} - \sqrt{m}\boldsymbol{v}^{\star} = \boldsymbol{\epsilon} + \boldsymbol{\xi}_{\boldsymbol{x}^{\star},\boldsymbol{v}^{\star}}$ (3.4) and then take the supremum over $\boldsymbol{c} = \frac{\Delta_{\boldsymbol{x}}}{f(\Delta_{\boldsymbol{x}})} \in \mathbb{B}_f^n$ and $\boldsymbol{d} = \frac{\Delta_{\boldsymbol{v}}}{g(\Delta_{\boldsymbol{v}})} \in \mathbb{B}_g^m$; moreover, in (3.58) we further take the supremum with respect to $\boldsymbol{\xi}_{\boldsymbol{x}^{\star},\boldsymbol{v}^{\star}}$ over $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star}) \in \mathcal{K}_{\boldsymbol{x}} \times \mathcal{K}_{\boldsymbol{v}}$ and introduce the shorthand

(3.59)
$$I_1 := \sup_{\boldsymbol{c} \in \mathbb{B}_f^n} \langle \boldsymbol{\epsilon}, \boldsymbol{\Phi} \boldsymbol{c} \rangle, \ I_2 := \sup_{\boldsymbol{a} \in \mathcal{K}_{\boldsymbol{x}}} \sup_{\boldsymbol{b} \in \mathcal{K}_{\boldsymbol{v}}} \sup_{\boldsymbol{c} \in \mathbb{B}_f^n} \langle \boldsymbol{\xi}_{\boldsymbol{a}, \boldsymbol{b}}, \boldsymbol{\Phi} \boldsymbol{c} \rangle,$$

(3.60)
$$I_3 := \sup_{\boldsymbol{d} \in \mathbb{B}_g^m} \langle \boldsymbol{\epsilon}, \sqrt{m} \boldsymbol{d} \rangle, \ I_4 := \sup_{\boldsymbol{a} \in \mathcal{K}_{\boldsymbol{x}}} \sup_{\boldsymbol{b} \in \mathcal{K}_{\boldsymbol{v}}} \sup_{\boldsymbol{d} \in \mathbb{B}_g^m} \langle \boldsymbol{\xi}_{\boldsymbol{a}, \boldsymbol{b}}, \sqrt{m} \boldsymbol{d} \rangle,$$

where $\xi_{a,b}$ is the quantization noise as per (3.26). We pause to provide an outline of the remainder of this proof:

- Step 1: Bounding I_1, I_2, I_3, I_4 in (3.59)–(3.60) by the techniques similar to those in the proof of Theorem 3.2. The high-probability bounds imply $\lambda_1 \ge 4(I_1+I_2), \lambda_2 \ge 4(I_3+I_4)$;
- Step 2: Based on Assumption 4, identifying the constraint set where (Δ_x, Δ_v) lives and establishing a uniform lower bound on $\|\Phi \Delta_x + \sqrt{m} \Delta_v\|_2^2$ via Proposition A.1;
- Step 3: Combining everything to conclude the proof.

Step 1: Bounding I_1, I_2, I_3, I_4 Similarly to the proof of Theorem 3.2, we bound the error terms associated with ϵ (i.e., I_1, I_3) via Proposition A.2 and Proposition A.4, and bound the terms associated with quantization noise (i.e., I_2, I_4) via QPE property specialized to structured sets (Corollary B.4).

Bounding I_1 : Note that $\mathbb{B}_f^n \subset \mathbb{B}_2^n$ since $f(\boldsymbol{x}) \geq \|\boldsymbol{x}\|_2$ holds for all $\boldsymbol{x} \in \mathbb{R}^n$. Conditioning on $\boldsymbol{\epsilon}$, Proposition A.4 provides that $I_1 \leq C_1 \|\boldsymbol{\epsilon}\|_2 \cdot \omega(\mathbb{B}_f^n)$ holds with probability exceeding $1 - \exp(-\omega^2(\mathbb{B}_f^n))$. Moreover, we can still bound $\|\boldsymbol{\epsilon}\|_2$ as in (3.33), which implies $\|\boldsymbol{\epsilon}\|_2 = O(E\sqrt{m})$ with probability exceeding $1 - \exp(-\Omega(m))$. Thus, the bound on I_1

(3.61)
$$I_1 \lesssim E\sqrt{m} \cdot \omega(\mathbb{B}_f^n)$$

holds with probability exceeding $1 - \exp(-\omega^2(\mathbb{B}^n_f)) - \exp(-\Omega(m))$.

Bounding I_2 : With the assumptions (3.46)–(3.47) and (3.50), we can invoke Corollary B.4 with $(\mathcal{A}, \mathcal{B}, \mathcal{E}) = (\mathcal{K}_{\boldsymbol{x}}, \mathcal{K}_{\boldsymbol{v}}, \mathbb{B}_f^n \times \{0\})$, yielding that the event

(3.62)
$$I_2 \lesssim \delta \sqrt{m} \cdot \left(\omega(\mathbb{B}_f^n) + \sqrt{\mathscr{H}(\mathcal{K}_{\boldsymbol{x}}, \rho_1) + \mathscr{H}(\mathcal{K}_{\boldsymbol{v}}, \rho_2)} \right)$$

holds with probability exceeding $1 - 12 \exp(-\Omega(\mathscr{H}(\mathcal{K}_{\boldsymbol{x}}, \rho_1) + \mathscr{H}(\mathcal{K}_{\boldsymbol{v}}, \rho_2))))$.

Bounding I_3 : We note that $\mathbb{B}_g^m \subset \mathbb{B}_2^m$ due to $g(\boldsymbol{v}) \geq \|\boldsymbol{v}\|_2$ holds for any $\boldsymbol{v} \in \mathbb{R}^m$. Analogously to "Bounding I_3 " in the proof of Theorem 3.2, we can apply Proposition A.2 to obtain to get $I_3 \leq E\sqrt{m}(\omega(\mathbb{B}_g^m) + t)$ with probability exceeding $1 - 2\exp(-t^2)$. Setting $t = \omega(\mathbb{B}_q^m)$ yields the bound on I_3

$$(3.63) I_3 \lesssim E\sqrt{m} \cdot \omega(\mathbb{B}^m_a)$$

with probability exceeding $1 - \exp(-\omega^2(\mathbb{B}_q^m))$.

Bounding I_4 : With the assumptions (3.46)–(3.47) and (3.50), we can invoke Corollary B.4 with $(\mathcal{A}, \mathcal{B}, \mathcal{E}) = (\mathcal{K}_x, \mathcal{K}_v, \{0\} \times \mathbb{B}_q^m)$. This yields that the event

(3.64)
$$I_4 \lesssim \delta \sqrt{m} \cdot \left(\omega(\mathbb{B}_g^m) + \sqrt{\mathscr{H}(\mathcal{K}_x, \rho_1) + \mathscr{H}(\mathcal{K}_v, \rho_2)} \right)$$

holds with probability exceeding $1 - 12 \exp(-\Omega(\mathscr{H}(\mathcal{K}_{\boldsymbol{x}}, \rho_1) + \mathscr{H}(\mathcal{K}_{\boldsymbol{v}}, \rho_2))))$. Compared to our choice of (λ_1, λ_2) in (3.48)–(3.49), we arrive at

(3.65)
$$\lambda_1 \ge 4(I_1 + I_2), \ \lambda_2 \ge 4(I_3 + I_4)$$

that hold with the promised probability.

Step 2: Constraining Estimation Errors Substituting (3.65) into (3.58) yields

(3.66)
$$\|\boldsymbol{\Phi}\boldsymbol{\Delta}_{\boldsymbol{x}} + \sqrt{m}\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2} \leq \frac{\lambda_{1}}{2}f(\boldsymbol{\Delta}_{\boldsymbol{x}}) + \frac{\lambda_{2}}{2}g(\boldsymbol{\Delta}_{\boldsymbol{v}}) + \lambda_{1}(f(\boldsymbol{x}^{\star}) - f(\hat{\boldsymbol{x}})) + \lambda_{2}(g(\boldsymbol{v}^{\star}) - g(\hat{\boldsymbol{v}})).$$

Since $\| \boldsymbol{\Phi} \boldsymbol{\Delta}_{\boldsymbol{x}} + \sqrt{m} \boldsymbol{\Delta}_{\boldsymbol{v}} \|_2^2 \ge 0$, we obtain

(3.67)
$$\lambda_1 (f(\hat{\boldsymbol{x}}) - f(\boldsymbol{x}^{\star})) + \lambda_2 (g(\hat{\boldsymbol{v}}) - g(\boldsymbol{v}^{\star})) \leq \frac{\lambda_1}{2} f(\boldsymbol{\Delta}_{\boldsymbol{x}}) + \frac{\lambda_2}{2} g(\boldsymbol{\Delta}_{\boldsymbol{v}}).$$

For a linear subspace \mathcal{X} , we use $\mathcal{P}_{\mathcal{X}}(\cdot)$ to denote the projection onto \mathcal{X} under ℓ_2 -norm. Given any $\mathbf{x}^{\star} \in \mathcal{K}_{\mathbf{x}}$, we can pick a pair of linear subspaces $(\mathcal{X}, \overline{\mathcal{X}}) := (\mathcal{X}_{\mathbf{x}^{\star}}, \overline{\mathcal{X}}_{\mathbf{x}^{\star}})$ as in Assumption 4 (they depend on \mathbf{x}^{\star} , while we will proceed with $(\mathcal{X}, \overline{\mathcal{X}})$ to avoid cumbersome notation) such that $f(\cdot)$ is decomposable over $(\mathcal{X}, \overline{\mathcal{X}}^{\perp})$. Then we can proceed as

(3.68)
$$f(\hat{\boldsymbol{x}}) = f(\boldsymbol{x}^{\star} + \boldsymbol{\Delta}_{\boldsymbol{x}}) = f(\boldsymbol{x}^{\star} + \mathcal{P}_{\overline{\mathcal{X}}}\boldsymbol{\Delta}_{\boldsymbol{x}} + \mathcal{P}_{\overline{\mathcal{X}}^{\perp}}\boldsymbol{\Delta}_{\boldsymbol{x}})$$

$$(3.69) \geq f(\boldsymbol{x}^{\star} + \mathcal{P}_{\overline{\mathcal{X}}^{\perp}} \boldsymbol{\Delta}_{\boldsymbol{x}}) - f(\mathcal{P}_{\overline{\mathcal{X}}} \boldsymbol{\Delta}_{\boldsymbol{x}})$$

(3.70)
$$= f(\boldsymbol{x}^{\star}) + f(\mathcal{P}_{\overline{\mathcal{X}}^{\perp}} \boldsymbol{\Delta}_{\boldsymbol{x}}) - f(\mathcal{P}_{\overline{\mathcal{X}}} \boldsymbol{\Delta}_{\boldsymbol{x}}),$$

where in (3.69) we use triangle inequality and in (3.70) we use the decomposibility of $f(\cdot)$ as per (3.45). Thus, we arrive at

(3.71)
$$f(\hat{\boldsymbol{x}}) - f(\boldsymbol{x}^{\star}) \ge f(\mathcal{P}_{\overline{\mathcal{X}}^{\perp}} \boldsymbol{\Delta}_{\boldsymbol{x}}) - f(\mathcal{P}_{\overline{\mathcal{X}}} \boldsymbol{\Delta}_{\boldsymbol{x}}).$$

Regarding the corruption, there exists $(\mathcal{V}, \overline{\mathcal{V}}) := (\mathcal{V}_{v^{\star}}, \overline{\mathcal{V}}_{v^{\star}})$ for a given $v^{\star} \in \mathcal{K}_{v}$ as in Assumption 4 such that $g(\cdot)$ is decomposable over $(\mathcal{V}, \overline{\mathcal{V}}^{\perp})$, and we can similarly show

(3.72)
$$g(\hat{\boldsymbol{v}}) - g(\boldsymbol{v}^{\star}) \ge g(\mathcal{P}_{\overline{\mathcal{V}}^{\perp}} \boldsymbol{\Delta}_{\boldsymbol{v}}) - g(\mathcal{P}_{\overline{\mathcal{V}}} \boldsymbol{\Delta}_{\boldsymbol{v}}).$$

Substituting (3.71)–(3.72) into the left-hand side of (3.67), and then use triangle inequality $f(\mathbf{\Delta}_{x}) \leq f(\mathcal{P}_{\overline{\chi}^{\perp}}\mathbf{\Delta}_{x}) + f(\mathcal{P}_{\overline{\chi}}\mathbf{\Delta}_{x})$ and $g(\mathbf{\Delta}_{v}) \leq g(\mathcal{P}_{\overline{\mathcal{V}}^{\perp}}\mathbf{\Delta}_{v}) + g(\mathcal{P}_{\overline{\mathcal{V}}}\mathbf{\Delta}_{v})$ in the right-hand side, we obtain

$$(3.73) \qquad \lambda_1 f(\mathcal{P}_{\overline{\mathcal{X}}^{\perp}} \mathbf{\Delta}_{\boldsymbol{x}}) + \lambda_2 g(\mathcal{P}_{\overline{\mathcal{V}}^{\perp}} \mathbf{\Delta}_{\boldsymbol{v}}) \le 3\lambda_1 f(\mathcal{P}_{\overline{\mathcal{X}}} \mathbf{\Delta}_{\boldsymbol{x}}) + 3\lambda_2 g(\mathcal{P}_{\overline{\mathcal{V}}} \mathbf{\Delta}_{\boldsymbol{v}}).$$

Based on this, we can derive the following

(3.74)
$$\lambda_1 f(\boldsymbol{\Delta}_{\boldsymbol{x}}) + \lambda_2 g(\boldsymbol{\Delta}_{\boldsymbol{v}})$$

$$(3.75) \qquad \leq \lambda_1 f(\mathcal{P}_{\overline{\mathcal{X}}} \mathbf{\Delta}_{\mathbf{x}}) + \lambda_1 f(\mathcal{P}_{\overline{\mathcal{X}}^{\perp}} \mathbf{\Delta}_{\mathbf{x}}) + \lambda_2 g(\mathcal{P}_{\overline{\mathcal{V}}} \mathbf{\Delta}_{\mathbf{v}}) + \lambda_2 g(\mathcal{P}_{\overline{\mathcal{V}}^{\perp}} \mathbf{\Delta}_{\mathbf{v}})$$

(3.76)
$$\leq 4\lambda_1 f(\mathcal{P}_{\overline{\mathcal{X}}} \boldsymbol{\Delta}_{\boldsymbol{x}}) + 4\lambda_2 g(\mathcal{P}_{\overline{\mathcal{V}}} \boldsymbol{\Delta}_{\boldsymbol{v}})$$

(3.77)
$$\leq 4\lambda_1 \alpha_{\boldsymbol{x}} \| \boldsymbol{\Delta}_{\boldsymbol{x}} \|_2 + 4\lambda_2 \alpha_{\boldsymbol{v}} \| \boldsymbol{\Delta}_{\boldsymbol{v}} \|_2,$$

where (3.75) follows from triangle inequality, in (3.76) we substitute (3.73), and moreover, (3.77) is due to $\alpha_f(\overline{\mathcal{X}}) \leq \alpha_x$ and $\alpha_g(\overline{\mathcal{V}}) \leq \alpha_v$ that hold uniformly for all $(\boldsymbol{x}^*, \boldsymbol{v}^*)$ (Assumption 4); in more detail, e.g., $f(\mathcal{P}_{\overline{\mathcal{X}}} \boldsymbol{\Delta}_{\boldsymbol{x}}) \leq \alpha_f(\overline{\mathcal{X}}) \| \mathcal{P}_{\overline{\mathcal{X}}} \boldsymbol{\Delta}_{\boldsymbol{x}} \|_2 \leq \alpha_{\boldsymbol{x}} \| \boldsymbol{\Delta}_{\boldsymbol{x}} \|_2$. Therefore, we arrive at

(3.78)
$$(\boldsymbol{\Delta}_{\boldsymbol{x}}, \boldsymbol{\Delta}_{\boldsymbol{v}}) \in \mathcal{C}(\lambda_1, \lambda_2), \ \forall (\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star}) \in \mathcal{K}_{\boldsymbol{x}} \times \mathcal{K}_{\boldsymbol{v}},$$

(it is easy to check that each step is uniform for all pairs of (x^*, v^*) ,) where the constraint set is given by

(3.79)
$$\mathcal{C}(\lambda_1,\lambda_2) = \left\{ (\boldsymbol{c},\boldsymbol{d}) \in \mathbb{R}^n \times \mathbb{R}^m : \lambda_1 f(\boldsymbol{c}) + \lambda_2 g(\boldsymbol{d}) \le 4\lambda_1 \alpha_{\boldsymbol{x}} \|\boldsymbol{c}\|_2 + 4\lambda_2 \alpha_{\boldsymbol{v}} \|\boldsymbol{d}\|_2 \right\}.$$

Uniform Lower Bound on $\|\Phi \Delta_x + \sqrt{m} \Delta_v\|_2^2$: Equipped with (3.78), we are now able to establish a uniform lower bound on $\|\Phi \Delta_x + \sqrt{m} \Delta_v\|_2^2$. Note that $\mathcal{C}(\lambda_1, \lambda_2)$ is a cone, and we localize it as $\mathcal{C}^* = \mathcal{C}(\lambda_1, \lambda_2) \cap \mathbb{S}^{n+m-1}$. Then we invoke Proposition A.1 with $\mathcal{T} = \mathcal{C}^*$ and $t = \omega(\mathbb{B}_f^n)$ to obtain that the event¹¹

(3.80)
$$\sup_{(\boldsymbol{c},\boldsymbol{d})\in\mathcal{C}^*} \left| \|\boldsymbol{\Phi}\boldsymbol{c} + \sqrt{m}\boldsymbol{d}\|_2 - \sqrt{m} \right| \leq C_1 \left(\omega(\mathcal{C}^*) + \omega(\mathbb{B}_f^n) \right)$$

holds with probability exceeding $1 - \exp(-\omega^2(\mathbb{B}_f^n))$. Note that Proposition A.8 provides a bound on $\omega(\mathcal{C}^*)$, which indicates that (3.50) implies $m \gtrsim \omega^2(\mathcal{C}^*) + \omega^2(\mathbb{B}_f^n)$. Thus, we can assume that the right-hand side of (3.80) is bounded by $\frac{1}{2}\sqrt{m}$ and achieve

(3.81)
$$\inf_{(\boldsymbol{c},\boldsymbol{d})\in\mathcal{C}^*} \|\boldsymbol{\Phi}\boldsymbol{c} + \sqrt{m}\boldsymbol{d}\|_2 \ge \sqrt{m} - \sup_{(\boldsymbol{c},\boldsymbol{d})\in\mathcal{C}^*} \|\boldsymbol{\Phi}\boldsymbol{c} + \sqrt{m}\boldsymbol{d}\|_2 - \sqrt{m}$$

$$(3.82) \geq \sqrt{m} - \frac{\sqrt{m}}{2} = \frac{\sqrt{m}}{2}$$

Combining with (3.78), we obtain

(3.83)
$$\|\boldsymbol{\Phi}\boldsymbol{\Delta}_{\boldsymbol{x}} + \sqrt{m}\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2} \ge \left(\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{2}^{2} + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2}\right) \inf_{(\boldsymbol{c},\boldsymbol{d})\in\mathcal{C}^{*}} \|\boldsymbol{\Phi}\boldsymbol{c} + \sqrt{m}\boldsymbol{d}\|_{2}^{2}$$

$$(3.84) \geq \frac{m}{4} \left(\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_2^2 + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_2^2 \right).$$

¹¹By Proposition A.1 we can bound the left-hand side of (3.80) by $O(\gamma(\mathcal{C}^*) + \omega(\mathbb{B}^n_f))$. Note that $\boldsymbol{v} \in \mathcal{C}^*$ implies $-\boldsymbol{v} \in \mathcal{C}^*$, hence let $\boldsymbol{g} \sim \mathcal{N}(0, \boldsymbol{I}_{n+m})$ we have $\omega(\mathcal{C}^*) = \mathbb{E}\sup_{\boldsymbol{v} \in \mathcal{C}^*} (\boldsymbol{g}^\top \boldsymbol{v}) \geq \mathbb{E}\sup_{\boldsymbol{v} \in \mathcal{C}^*} \max\{\boldsymbol{g}^\top \boldsymbol{v}, \boldsymbol{g}^\top(-\boldsymbol{v})\} = \mathbb{E}\sup_{\boldsymbol{v} \in \mathcal{C}^*} |\boldsymbol{g}^\top \boldsymbol{v}| = \gamma(\mathcal{C}^*)$. Thus, we arrive at the bound given in the right-hand side of (3.80).

Step 3: Combining Everything We derive the desired bound by bounding both sides of (3.66): By (3.84), the left-hand side is uniformly lower bounded by $\frac{m}{4}(\|\Delta_{\boldsymbol{x}}\|_2^2 + \|\Delta_{\boldsymbol{v}}\|_2^2)$; Using triangle inequality and (3.77) in (3.86), along with simple algebra in (3.87), the right-hand side can be uniformly bounded by

(3.85)
$$\frac{\lambda_1}{2}f(\boldsymbol{\Delta}_{\boldsymbol{x}}) + \frac{\lambda_2}{2}g(\boldsymbol{\Delta}_{\boldsymbol{v}}) + \lambda_1 \big(f(\boldsymbol{x}^{\star}) - f(\hat{\boldsymbol{x}})\big) + \lambda_2 \big(g(\boldsymbol{v}^{\star}) - g(\hat{\boldsymbol{v}})\big)$$

(3.86)
$$\leq \frac{3\lambda_1}{2} f(\boldsymbol{\Delta}_{\boldsymbol{x}}) + \frac{3\lambda_2}{2} g(\boldsymbol{\Delta}_{\boldsymbol{v}}) \leq 6 \left(\lambda_1 \alpha_{\boldsymbol{x}} \| \boldsymbol{\Delta}_{\boldsymbol{x}} \|_2 + \lambda_2 \alpha_{\boldsymbol{v}} \| \boldsymbol{\Delta}_{\boldsymbol{v}} \|_2 \right)$$

(3.87)
$$\leq 6(\lambda_1 \alpha_{\boldsymbol{x}} + \lambda_2 \alpha_{\boldsymbol{v}}) \left(\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_2^2 + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_2^2 \right)^{1/2}$$

Rearranging immediately yields the claim that holds with the promised probability as per (3.51).

Substituting (3.48)–(3.49) into (3.52) yields the following more explicit uniform bound on $(\|\Delta_x\|_2^2 + \|\Delta_v\|_2^2)^{1/2}$

$$(3.88) \ O\left(\frac{(E+\delta)\big(\alpha_{\boldsymbol{x}}\cdot\boldsymbol{\omega}(\mathbb{B}_{f}^{n})+\alpha_{\boldsymbol{v}}\cdot\boldsymbol{\omega}(\mathbb{B}_{g}^{m})\big)+\delta(\alpha_{\boldsymbol{x}}+\alpha_{\boldsymbol{v}})\big(\mathscr{H}(\mathcal{K}_{\boldsymbol{x}},\rho_{1})+\mathscr{H}(\mathcal{K}_{\boldsymbol{v}},\rho_{2})\big)^{1/2}}{\sqrt{m}}\right).$$

To see the implication of Theorem 3.12 on structured priors (Assumption 3), we substitute (3.9) into (3.88) and omit some logarithmic factors on (ρ_1, ρ_2) , then we obtain the uniform bound

(3.89)
$$\tilde{O}\left(\frac{(E+\delta)\left(\alpha_{\boldsymbol{x}}\cdot\boldsymbol{\omega}(\mathbb{B}_{f}^{n})+\alpha_{\boldsymbol{v}}\cdot\boldsymbol{\omega}(\mathbb{B}_{g}^{m})\right)+\delta(\alpha_{\boldsymbol{x}}+\alpha_{\boldsymbol{v}})\left(\boldsymbol{\omega}(\mathcal{K}_{\boldsymbol{x}})+\boldsymbol{\omega}(\mathcal{K}_{\boldsymbol{v}})\right)}{\sqrt{m}}\right).$$

Remark 3.13 (The Cost of Uniformity: Unconstrained Lasso with Structured Priors). The non-uniform error rate for (3.7) in [75, Thm. 2] also reads as $O(\frac{\lambda_1 \alpha_x + \lambda_2 \alpha_v}{m})$ (when adjusted to our notation), with the parameters $\lambda_1 \simeq (E + \delta)\sqrt{m} \cdot \omega(\mathbb{B}_f^n)$ and $\lambda_2 \simeq (E + \delta)\sqrt{m} \cdot \omega(\mathbb{B}_g^m)$ to guarantee that

(3.90)
$$\lambda_1 \gtrsim \sup_{\boldsymbol{c} \in \mathbb{B}_f^n} \langle \boldsymbol{\epsilon} + \boldsymbol{\xi}_{\boldsymbol{x}^\star, \boldsymbol{v}^\star}, \boldsymbol{\Phi} \boldsymbol{c} \rangle, \ \lambda_2 \gtrsim \sup_{\boldsymbol{d} \in \mathbb{B}_g^m} \langle \boldsymbol{\epsilon} + \boldsymbol{\xi}_{\boldsymbol{x}^\star, \boldsymbol{v}^\star}, \sqrt{m} \boldsymbol{d} \rangle$$

holds for a fixed (x^*, v^*) ; see Remark 5 therein. Thus, their non-uniform error rate translates into

(3.91)
$$O\left(\frac{(E+\delta)\left(\alpha_{\boldsymbol{x}}\cdot\boldsymbol{\omega}(\mathbb{B}_{f}^{n})+\alpha_{\boldsymbol{v}}\cdot\boldsymbol{\omega}(\mathbb{B}_{g}^{m})\right)}{\sqrt{m}}\right).$$

However, to achieve uniformity, we must use sufficiently large λ_1 and λ_2 to ensure that (3.90) holds for all $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star})$, i.e.,

(3.92)
$$\lambda_1 \gtrsim \sup_{\boldsymbol{a} \in \mathcal{K}_{\boldsymbol{x}}} \sup_{\boldsymbol{b} \in \mathcal{K}_{\boldsymbol{v}}} \sup_{\boldsymbol{c} \in \mathbb{B}_f^n} \langle \boldsymbol{\epsilon} + \boldsymbol{\xi}_{\boldsymbol{a}, \boldsymbol{b}}, \boldsymbol{\Phi} \boldsymbol{c} \rangle, \ \lambda_2 \gtrsim \sup_{\boldsymbol{a} \in \mathcal{K}_{\boldsymbol{x}}} \sup_{\boldsymbol{b} \in \mathcal{K}_{\boldsymbol{v}}} \sup_{\boldsymbol{d} \in \mathbb{B}_g^m} \langle \boldsymbol{\epsilon} + \boldsymbol{\xi}_{\boldsymbol{a}, \boldsymbol{b}}, \sqrt{m} \boldsymbol{d} \rangle.$$

For this purpose, the second term of λ_1 in (3.48) (or λ_2 in (3.49)) is additional compared to the λ_1 and λ_2 in [75]. This leads to the additional term

(3.93)
$$\tilde{O}\left(\frac{\delta(\alpha_{\boldsymbol{x}} + \alpha_{\boldsymbol{v}})(\omega(\mathcal{K}_{\boldsymbol{x}}) + \omega(\mathcal{K}_{\boldsymbol{v}}))}{\sqrt{m}}\right)$$

in our uniform error rate (3.89), which we further discuss as follows:

- Under the regular scaling of $E \simeq \delta$, the additional term (3.93) is typically dominant, and as a result our uniform rate (3.89) often exhibits worse dependence on structured parameters than its non-uniform counterpart (3.91). For instance, in recovery of *s*sparse \mathbf{x}^{\star} we have $\mathcal{K}_{\mathbf{x}} = \sum_{s}^{n} \cap \mathbb{B}_{2}^{n}$ and $f(\mathbf{x}) = \|\mathbf{x}\|_{1}$, and thus $\omega(\mathcal{K}_{\mathbf{x}}) \simeq \sqrt{s \log \frac{en}{s}}$ and $\omega(\mathbb{B}_{f}^{n}) \simeq \sqrt{\log n}$, then the term $\frac{\alpha_{\mathbf{x}} \cdot \omega(\mathcal{K}_{\mathbf{x}})}{\sqrt{m}}$ in (3.93) already loses a factor of \sqrt{s} compared to $\frac{\alpha_{\mathbf{x}} \cdot \omega(\mathbb{B}_{f}^{n})}{\sqrt{m}}$ in (3.91). The readers can clearly see such degradation by comparing [75, Coros 3–4] and our Corollary 3.14–Corollary 3.15 below.
- Nevertheless, to the best of our knowledge, there exists no uniform result for unconstrained Lasso in quantized compressed/corrupted sensing (see Remark 3.9 for a review), thereby pointing to the open question on the possibility of improvement.
- Moreover, we observe that the additional term (3.93) presents a multiplicative factor δ , reflecting that the gap between uniform recovery and non-uniform recovery closes when δ is extremely small or in an unquantized setting where $\delta = 0$.

We present some concrete outcomes of Theorem 3.12, with proofs deferred to Appendix C.

Corollary 3.14 (Sparse Signal and Sparse Corruption). We consider the same settings as in Corollary 3.5 (i.e., $\mathbf{x}^{\star} \in \mathcal{K}_{\mathbf{x}} = \Sigma_s^n \cap \mathbb{B}_2^n$ with $f(\mathbf{x}) = \|\mathbf{x}\|_1$, $\mathbf{v}^{\star} \in \mathcal{K}_{\mathbf{v}} = \Sigma_k^m \cap \mathbb{B}_2^m$ with $g(\mathbf{v}) = \|\mathbf{v}\|_1$, but use (3.7) as recovery program. We set

$$\lambda_1 = C_1 \delta \sqrt{ms \log\left(\frac{nm^{3/2}}{s^{5/2}\delta}\right) + mk \log\left(\frac{m^{5/2}}{k^{5/2}\delta}\right)} + C_1(E+\delta)\sqrt{m\log n},$$
$$\lambda_2 = C_2 \delta \sqrt{ms \log\left(\frac{nm^{3/2}}{s^{5/2}\delta}\right) + mk \log\left(\frac{m^{5/2}}{k^{5/2}\delta}\right)} + C_2(E+\delta)\sqrt{m\log m}$$

with sufficiently large C_1, C_2 . If $m \ge C_3(s+k)\log(mn) + C_3s\log(\frac{nm^{3/2}}{s^{5/2}\delta}) + C_3k\log(\frac{m^{5/2}}{k^{5/2}\delta})$ for large enough C_3 , then with probability exceeding $1 - n^{-c_4} - m^{-c_5}$ on a single draw of $(\mathbf{\Phi}, \boldsymbol{\epsilon}, \boldsymbol{\tau})$, the following uniform error bound holds true for all $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star}) \in \mathcal{K}_{\boldsymbol{x}} \times \mathcal{K}_{\boldsymbol{v}}$:

$$\sqrt{\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{2}^{2} + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2}} \lesssim \frac{(E+\delta)\sqrt{s\log n + k\log m} + \delta\sqrt{(s+k)\left(s\log(\frac{nm^{3/2}}{s^{5/2}\delta}) + k\log(\frac{m^{5/2}}{k^{5/2}\delta})\right)}}{\sqrt{m}}$$

where $\Delta_x = \hat{x} - x^*, \Delta_v = \hat{v} - v^*, (\hat{x}, \hat{v})$ is the solution to (3.7).

Corollary 3.15 (Low-Rank Signal and Sparse Corruption). We consider the same settings as in Corollary 3.7 (i.e., $\mathbf{x}^{\star} \in \mathcal{K}_{\mathbf{x}} = M_r^{p,q} \cap \mathbb{B}_{\mathrm{F}}^{p,q}$ with $f(\mathbf{x}) = \|\mathbf{x}\|_{\mathrm{nu}}$, \mathbf{x}^{\star} is vectorized as a (pq)-dimensional vector when substituted into (3.1), $\mathbf{v}^{\star} \in \mathcal{K}_{\mathbf{v}} = \Sigma_k^m \cap \mathbb{B}_2^m$ with $g(\mathbf{v}) = \|\mathbf{v}\|_1$, but use (3.7) as recovery program. We set

$$\lambda_{1} = C_{1}\delta \cdot \sqrt{mr(p+q)\log\left(\frac{m^{3/2}}{\delta(r(p+q))^{3/2}}\right) + mk\log\left(\frac{m^{5/2}}{k^{5/2}\delta}\right) + C_{1}(E+\delta)\sqrt{m(p+q)}},$$

$$\lambda_{2} = C_{2}\delta \cdot \sqrt{mr(p+q)\log\left(\frac{m^{3/2}}{\delta(r(p+q))^{3/2}}\right) + mk\log\left(\frac{m^{5/2}}{k^{5/2}\delta}\right)} + C_{2}(E+\delta)\sqrt{m\log m}$$

with sufficiently large C_1, C_2 . If $m \ge C_3(r+k)(p+q+\log m) + C_3r(p+q)\log(\frac{m^{3/2}}{\delta(r(p+q))^{3/2}}) + C_3k\log\frac{m^{5/2}}{k^{5/2}\delta}$ for large enough C_3 , then with probability exceeding $1 - C_4\exp(-c_5(p+q)) - C_6m^{-c_7}$ on a single draw of (Φ, ϵ, τ) , the following uniform error bound holds true for all $(\boldsymbol{x^*}, \boldsymbol{v^*}) \in \mathcal{K}_{\boldsymbol{x}} \times \mathcal{K}_{\boldsymbol{v}}$:

$$\begin{split} \sqrt{\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{\mathrm{F}}^{2} + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2}} \lesssim \frac{(E+\delta)\sqrt{r(p+q) + k\log m}}{\sqrt{m}} \\ + \frac{\delta\sqrt{(r+k)\left(r(p+q)\log(\frac{m^{3/2}}{\delta(r(p+q))^{3/2}}) + k\log(\frac{m^{5/2}}{\delta k^{5/2}})\right)}}{\sqrt{m}} \end{split}$$

where $\Delta_x = \hat{x} - x^{\star}, \Delta_v = \hat{v} - v^{\star}, (\hat{x}, \hat{v})$ is the solution to (3.7).

3.3 Generative Priors To handle the case where traditional structured priors fail to precisely characterize the underlying signal, it was recently proposed to use a generative prior for compressed sensing, i.e., assuming that the desired signal lies in the range of a generative model [5]. This new perspective for compressed sensing has led to numerical success and attracted much research interest. The goal of this subsection is to establish uniform recovery guarantee for quantized corrupted sensing using generative priors. Following the long list of prior works on this field (e.g., [4, 5, 18, 55]), we formulate the generative priors on signal and corruption as follows.

Assumption 5 (Generative Priors). For some r, r' > 0, let $G : \mathbb{B}_2^k(r) \to \mathbb{R}^n$, $H : \mathbb{B}_2^{k'}(r') \to \mathbb{R}^m$ be some generative models. We assume that $G(\cdot)$ is L-Lipschitz continuous, $H(\cdot)$ is L'-Lipschitz continuous:

(3.94)
$$\|G(\boldsymbol{a}_1) - G(\boldsymbol{a}_2)\|_2 \le L \|\boldsymbol{a}_1 - \boldsymbol{a}_2\|_2, \ \forall \boldsymbol{a}_1, \boldsymbol{a}_2 \in \mathbb{R}^k,$$

(3.95)
$$\|H(\boldsymbol{b}_1) - H(\boldsymbol{b}_2)\|_2 \le L' \|\boldsymbol{b}_1 - \boldsymbol{b}_2\|_2, \ \forall \boldsymbol{b}_1, \boldsymbol{b}_2 \in \mathbb{R}^{k'}$$

and assume that the signal and corruption lie in the range of the two generative models:

(3.96)
$$\boldsymbol{x}^{\star} \in \mathcal{K}_{\boldsymbol{x}} := G\big(\mathbb{B}_{2}^{k}(r)\big), \boldsymbol{v}^{\star} \in \mathcal{K}_{\boldsymbol{v}} := H\big(\mathbb{B}_{2}^{k'}(r')\big).$$

Recall that $\dot{\boldsymbol{y}}$ are the quantized measurements as per (3.2). We naturally extend the scope of constrained Lasso (3.6) by substituting the norm constraints with the generative priors on $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star})$:

(3.97)
$$(\hat{\boldsymbol{x}}, \hat{\boldsymbol{v}}) = \arg\min_{\substack{\boldsymbol{x} \in \mathbb{R}^n \\ \boldsymbol{v} \in \mathbb{R}^m}} \| \dot{\boldsymbol{y}} - \boldsymbol{\Phi} \boldsymbol{x} - \sqrt{m} \boldsymbol{v} \|_2, \text{ s.t. } \boldsymbol{x} \in G(\mathbb{B}_2^k(r)), \ \boldsymbol{v} \in H(\mathbb{B}_2^{k'}(r')).$$

We note that it is in general hard to exactly optimize (3.97) due to the highly non-convex constraint, while fortunately there have been some practical approaches to *approximately* solve this program [53,71,74]. Note that all prior recovery methods in this area exhibit optimization issue of this type but have proven effective in practice.

We present a uniform recovery guarantee of (3.97).

Theorem 3.16 (Quantized Corrupted Sensing with Generative Priors). Under Assumption 1 and Assumption 5, we let $\mu \in (0,1)$ be some given recovery accuracy and suppose that m = O(n). If for some sufficiently large C_1 we have

$$(3.98) \ m \ge \frac{C_1 E^2}{\mu^2} \Big(k \log \frac{Lr}{\mu} + k' \log \frac{L'r'}{\mu} \Big) + C_1 \Big(1 + \frac{\delta^2}{\mu^2} \Big) \Big(k \log \Big(\frac{Lr n^{3/2}}{\mu \delta k^{3/2}} \Big) + k' \log \Big(\frac{L'r' m^{3/2}}{\mu \delta (k')^{3/2}} \Big) \Big),$$

then with probability exceeding $1 - C_2 \exp(-\Omega(k \log(Lr) + k' \log(L'r')))$ on a single draw of $(\mathbf{\Phi}, \boldsymbol{\epsilon}, \boldsymbol{\tau})$, the uniform error bound $\sqrt{\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_2^2 + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_2^2} \leq \mu$ holds for all $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star}) \in \mathcal{K}_{\boldsymbol{x}} \times \mathcal{K}_{\boldsymbol{v}}$, where $\boldsymbol{\Delta}_{\boldsymbol{x}} = \hat{\boldsymbol{x}} - \boldsymbol{x}^{\star}$ and $\boldsymbol{\Delta}_{\boldsymbol{v}} = \hat{\boldsymbol{v}} - \boldsymbol{v}^{\star}$, with $(\hat{\boldsymbol{x}}, \hat{\boldsymbol{v}})$ being the solution to (3.97).

A Sketch of the Proof. The proof of Theorem 3.16 is analogous to that of Theorem 3.2, with the major differences lying in constraining the range of the estimation error (Δ_x, Δ_v) , which also appeals to separate treatments in estimating the Gaussian width and Kolmogorov entropy (see Proposition A.9). The full proof can be found in Appendix C.

Remark 3.17 (Uniform Error Bound for Theorem 3.16). Provided that $m \gtrsim k \log(\frac{Lrn^{3/2}}{\mu \delta k^{3/2}}) + k' \log(\frac{L'r'm^{3/2}}{\mu \delta (k')^{3/2}})$ for large enough implied constant, the sample complexity (3.98) sufficient for achieving a uniform ℓ_2 -error of μ implies a uniform error bound

(3.99)
$$\begin{split} \sqrt{\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{2}^{2} + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2}} \lesssim E\sqrt{\frac{k\log(\frac{Lr\sqrt{m}}{E\sqrt{k}}) + k'\log(\frac{L'r'\sqrt{m}}{E\sqrt{k'}})}{m}}{m}} \\ + \delta\sqrt{\frac{k\log(\frac{Lrn^{2}}{\delta^{2}k^{2}}) + k'\log(\frac{L'r'm^{2}}{\delta^{2}(k')^{2}})}{m}}, \end{split}$$

which can be verified by some algebra.

Remark 3.18 (Technical Comparison with [18]). Restricted to compressed sensing with generative prior, most recovery guarantees for non-linear models are non-uniform. As a followup of [36], Chen et al. [18] built a unified framework for proving uniform recovery guarantee in non-linear compressed sensing with generative prior. Specifically, they handled potential discontinuity of $f_i(\cdot)$ by constructing Lipschitz approximation as in [36], but used [18, Thm. 2] (rather than [36, Thm. 8]) to bound the product processes. Their key observation is that the replacement of concentration inequality yields tighter bound for the generative case [18, Remark 8], thus they proved a uniform decaying rate of $O(m^{1/2})$ even when $f_i(\cdot)$ contains some discontinuity. (In contrast, [36] only achieves a uniform decaying rate of $O(m^{-1/4})$ under discontinuous $f_i(\cdot)$, see Remark 3.11.) However, as with Remark 3.11, it is unclear whether their approach applies to analyzing corrupted sensing, since the corruption leads to random process out of the scope of [18, Thm. 2].¹² Moreover, while the present paper provides unified analysis of two priors, it is unclear whether the techniques in [18] can be adapted to structured prior.

4 Experimental Results In this section we provide experimental results to corroborate and demonstrate our uniform recovery guarantees. Due to the theoretical nature of our work, an extensive set of experiments will not pursued.

4.1 Structured Priors First, we consider using structured priors for corruption sensing, for which our main theoretical results are Theorem 3.2 and Theorem 3.12. The aim of our first set of experiments is to show the recovery performance of (un)constrained Lasso under two settings, namely sparse signal recovery from sparse corruption (Corollary 3.5 and Corollary 3.14) and low-rank matrix recovery from sparse corruption (Corollary 3.7 and Corollary 3.15). All simulations in this subsection are performed using MATLAB R2018b on a desktop with a 3.70 GHz Intel Core i7-8700M CPU and 32 GB RAM.

We use a realization of the ensemble (Φ, τ, ϵ) to recover a fixed (x^*, v^*) for simulating non-uniform recovery. In contrast, with a single realization of the sensing ensemble, the error rates in our theorems holds uniformly for all $(x^*, v^*) \in \mathcal{K}_x \times \mathcal{K}_v$, or equivalently interpreted, they are upper bounds on the following quantity:

(4.1)
$$\sup_{\boldsymbol{x}^{\star} \in \mathcal{K}_{\boldsymbol{x}}} \sup_{\boldsymbol{v}^{\star} \in \mathcal{K}_{\boldsymbol{v}}} \left(\| \hat{\boldsymbol{x}} - \boldsymbol{x}^{\star} \|_{2}^{2} + \| \hat{\boldsymbol{v}} - \boldsymbol{v}^{\star} \|_{2}^{2} \right)^{1/2}.$$

Nonetheless, under a fixed (Φ, τ, ϵ) , it is in general impossible to track (4.1) since $\mathcal{K}_{\boldsymbol{x}} \times \mathcal{K}_{\boldsymbol{v}}$ is typically infinite set (e.g., $\mathcal{K}_{\boldsymbol{x}} = \Sigma_s^n \cap \mathbb{B}_2^n$). To provide some clues to demonstrate our theories, we instead utilize a fixed (Φ, τ, ϵ) to recover multiple signal-and-corruption pairs in a testing set \mathcal{X}_{test} , and we will track the maximum recovery error

(4.2)
$$\sup_{(\boldsymbol{x}^{\star},\boldsymbol{v}^{\star})\in\mathcal{X}_{test}} \left(\|\boldsymbol{\hat{x}}-\boldsymbol{x}^{\star}\|_{2}^{2}+\|\boldsymbol{\hat{v}}-\boldsymbol{v}^{\star}\|_{2}^{2}\right)^{1/2}$$

as an approximation of (4.1). We adopt the following general principles in our simulations:

- Construction of \mathcal{X}_{test} : Given the cardinality $|\mathcal{X}_{test}|$, the (x^*, v^*) in \mathcal{X}_{test} are independently, randomly created by the construction of sparse vector and low-rank matrix below.
- Calculation of (4.2): In a single trial that simulates the uniform recovery over some \mathcal{X}_{test} , we calculate (4.2) under a single draw of (Φ, τ, ϵ) . We will report (4.2) as its mean value in 10 independent trials.
- Construction of a Vector in Σ_s^N : We let the support of the vector uniformly distributed over all $\binom{N}{s}$ possibilities, and then draw the non-zero entries from $\mathcal{N}(0, 1)$.
- Construction of a Matrix in M^{p,q}_r: We use UV[⊤] where where U ∈ ℝ^{p×r} and V ∈ ℝ^{q×r} are independent random matrices with orthonormal columns, generated by the Matlab code "[U,S,U1]=svd(randn(p,r),r); [V,S,V1]=svd(randn(q,r),r)".

¹²As we reviewed in section 1, under generative prior, the only existing result that accommodates a generative corruption \boldsymbol{v}^{\star} was presented in [3,4] and restricted to the linear case.

• Tuning Parameters: In unconstrained Lasso we provide the best possible constraint for each pair $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star}) \in \mathcal{X}_{test}$, namely $f(\boldsymbol{x}) \leq f(\boldsymbol{x}^{\star})$ and $g(\boldsymbol{v}) \leq g(\boldsymbol{v}^{\star})$. In unconstrained Lasso, we properly choose a large enough (λ_1, λ_2) and then use it for all $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star}) \in \mathcal{K}_{test}$.

4.1.1 Non-Uniformity v.s. Uniformity We first compare non-uniform recovery and uniform recovery (over \mathcal{X}_{test}) to demonstrate the major theoretical achievements of this work.

Constrained Lasso: we simulate Corollary 3.5 with $(n, s, k, \delta, E) = (256, 2, 2, 0.1, 0)$. We construct four test sets with different cardinalities: $|\mathcal{X}_{test}| = 1, 10, 100, 300$. Then we test these four cases under a properly chosen range of the measurement number m. Note that $|\mathcal{X}_{test}| = 1$ reduces to non-uniform recovery, while $|\mathcal{X}_{test}| = 300$ represents the highest level of uniformity that we simulate. Over these four test sets, we report (4.2) as its mean value in 10 independent trials under properly chosen measurement number, and then plot the loglog curves in Figure 1(left). We note the following observations from Figure 1(left) that are consistent with our Corollary 3.5:

- All curves decay in a rate of $O(m^{-1/2})$;
- To achieve the same recovery error over a "larger" \mathcal{X}_{test} (which corresponds to a higher level of uniformity) requires more measurements.

Since (4.1) is always larger than (4.2), the log-log curve of (4.1) will further shift to the right compared to the green curve (corresponding to $|\mathcal{X}_{test}| = 300$) in Figure 1(left), whereas regarding this curve corresponding to the "actual" uniform recovery (that we cannot experimentally track), our Corollary 3.5 guarantees the following: (i) Fixing other parameters, this curve still decays in a rate of $O(m^{-1/2})$; (ii) The measurement number needed to achieve the same recovery error as the curves in Figure 1(left) is still of the scaling law $\tilde{O}(s+k)$ (logarithmic factors omitted).

Unconstrained Lasso: We provide similar experiment results for unconstrained Lasso to demonstrate our theory. We simulate Corollary 3.14 with $(n, s, k, \delta, E) = (256, 2, 2, 0.1, 0)$ and track the quantity (4.2) with $|\mathcal{X}_{test}| = 1, 10, 100, 300$. The results are displayed in Figure 1(right), and we note the similar observations that all curves are decaying roughly in a rate of $O(m^{-1/2})$, and that larger \mathcal{X}_{test} requires more measurements to ensure that (4.2) is smaller than some desired accuracy.

Naturally, the log-log curve of (4.1) will further shift to the right compared to the green curve (under $|\mathcal{X}_{test}| = 300$) in Figure 1(right), while regarding this curve corresponding to the "actual" uniform recovery (that we cannot experimentally track), our Corollary 3.14 promises the following: (i) This curve still decays i a rate of $O(m^{-1/2})$ when other parameters are fixed; (ii) The measurement number needed to achieve the same recovery error as the curves in Figure 1(right) is of scaling $\tilde{O}(s^2 + k^2)$ (logarithmic factors omitted, and note that we do not know whether this is sharp).

4.1.2 Constrained Lasso In order to demonstrate the role of different parameters in our uniform bound, we proceed to more simulations for constrained Lasso.

Sparse Recovery from Sparse Corruption: We simulate Corollary 3.5 with $(n, \delta, E) = (256, 0.1, 0)$ and vary the measurement number m between 150 and 500. We consider both Gaussian design $\Phi \sim \mathcal{N}^{m \times n}(0, 1)$ and Bernoulli design that has i.i.d. zero-mean $\{-1, 1\}$ -



Figure 1. From Non-Uniformity to Uniformity: (Left) Sparse Recovery via Constrained Lasso (Corollary 3.5); (Right) Sparse Recovery via Unconstrained Lasso (Corollary 3.14)



Figure 2. Log-log error curves for the constrained Lasso under Gaussian or Bernoulli measurements. valued entries.¹³ We simulate the uniform recovery with size-100 \mathcal{X}_{test} in the cases of "s =

¹³Bernoulli design is an example that demonstrates the benefit of using dithering. Without the random dither

k = 5", "s = k = 10" and "s = k = 15", and we report (4.2) as log-log curves in Figure 2(a) and Figure 2(d) for Gaussian design and Bernoulli design, respectively. Clearly, the results under two designs are similar, and note the two observations that are consistent with our theory. First, all curves decrease with m in the theoretical rate $O(m^{-1/2})$. Second, the errors increase under larger (s, k), for which the intuition is that weaker sparse priors on $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star})$ correspond to a harder high-dimensional estimation problem.

Low-Rank Recovery from Sparse Corruption: We simulate Corollary 3.7 with $(p,q,\delta,E) = (16,16,0.1,0)$ and vary measurements m between 200 and 1200. Under a realization of (Φ, τ) , we track (4.2) with size-100 \mathcal{X}_{test} in the cases of "(r,k) = (1,5)", "(r,k) = (2,10)" and "(r,k) = (3,15)". The log-log curves corresponding to Gaussian and Bernoulli designs are shown in Figure 2(b) and Figure 2(e) respectively, which are consistent with our theoretical uniform bound.



Figure 3. Log-log error curves for the unconstrained Lasso under Gaussian or Bernoulli measurements.

The Role of δ and E: We also use the setting of Corollary 3.5 to illustrate the role played by the quantization resolution δ and the robustness to noise ϵ . We test Gaussian noise

 τ , the identifiability issue arises under Bernoulli design even in compressed sensing without the corruption v^* (e.g., [17,75]). On the other hand, by using dithering, recovery can be ensured under general sub-Gaussian Φ .

 $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \sigma^2 \boldsymbol{I}_m)$ (note that the noise level σ can be simply understood as E in Assumption 1) in the cases of " $(\sigma, \delta) = (0.02, 0.1)$ ", " $(\sigma, \delta) = (0.02, 0.2)$ ", " $(\sigma, \delta) = (0.04, 0.1)$ " and " $(\sigma, \delta) = (0.04, 0.2)$ ". The log-log curves are displayed in Figure 2(c) and Figure 2(f). As predicted by our uniform bound (3.41), larger δ and severer sub-Gaussian noise (i.e., larger E) lift the curves higher but do not affect the decaying rate of $O(m^{-1/2})$.

4.1.3 Unconstrained Lasso We conduct parallel experiments using unconstrained Lasso (3.7). Note that unconstrained Lasso (3.7) might be more practical than its constrained counterpart, in the sense that we can use a program with fixed large enough (λ_1, λ_2) for all 100 pairs of $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star})$ in \mathcal{X}_{test} . We show the results in Figure 3, which are consistent with the uniform error bounds in Corollary 3.14–Corollary 3.15 in terms of decaying rate, qualitative dependence on structured parameters, δ and E.

4.2 Generative Priors In this subsection, we present proof-of-concept experimental results for the case of using generative priors. In particular, we consider the case that the signal is close to the range of a generative model, and the corruption vector is also close to the range of another generative model. All the experiments were conducted using the Python 3.10.6 and PyTorch 2.0.0 framework on an NVIDIA RTX 3060 Laptop 6GB GPU. We modify (4.2) and track the following two quantities for signal and corruption:

(4.3)
$$\sup_{(\boldsymbol{x}^{\star},\boldsymbol{v}^{\star})\in\mathcal{X}_{test}}\frac{\|\hat{\boldsymbol{x}}-\boldsymbol{x}^{\star}\|_{2}}{\|\boldsymbol{x}^{\star}\|_{2}}, \quad \sup_{(\boldsymbol{x}^{\star},\boldsymbol{v}^{\star})\in\mathcal{X}_{test}}\frac{\|\hat{\boldsymbol{v}}-\boldsymbol{v}^{\star}\|_{2}}{\|\boldsymbol{v}^{\star}\|_{2}},$$

which are just the maximum relative error over a test set of (x^{\star}, v^{\star}) denoted by \mathcal{X}_{test} .

Demixing "8" from "1" in MNIST: First, we follow [3, 4] to train two variational auto-encoders (VAEs) for the training images of digits 8 and 1 in the MNIST dataset [48] respectively. The decoders of these two VAEs were composed of a fully connected neural network with ReLU activation functions. The VAEs had an input dimension k = k' = 20 and an output dimension of $m = n = 28 \times 28 = 784$, with two hidden layers consisting of 500 neurons each. We used the Adam optimizer with a mini-batch size of 100 and a learning rate of 0.001 to train these VAEs.

We take the images of digit 8 as the signal and those of digit 1 as the corruption vectors. We use $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \sigma^2 \boldsymbol{I}_m)$ to simulate sub-Gaussian noise. To demonstrate uniform recovery, we use a single realization of $(\boldsymbol{\Phi}, \boldsymbol{\epsilon}, \boldsymbol{\tau})$ to track the maximum relative error in (4.3), where \mathcal{X}_{test} contains 20 test images of digits 8 and 1 from the testing set of MNIST. As before, we report the quantities in (4.3) as its mean value in 10 independent random trials.

We use the constrained Lasso (3.97) to reconstruct $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star})$. Similarly to the algorithm proposed in [5], we employ the gradient descent algorithm to minimize the following objective function over $\mathbb{R}^k \times \mathbb{R}^{k'}$:

$$\mathcal{L}(\boldsymbol{z}, \boldsymbol{z}') := \| \dot{\boldsymbol{y}} - \boldsymbol{\Phi} G(\boldsymbol{z}) - \sqrt{m} H(\boldsymbol{z'}) \|_2,$$

where $G : \mathbb{B}_2^k(r) \to \mathbb{R}^n$, $H : \mathbb{B}_2^{k'}(r') \to \mathbb{R}^m$ are the generative models as per Assumption 5.¹⁴ Our algorithm is referred to as QCS_Gen. We follow the settings in [5] and perform 10 random

¹⁴Since r and r' can typically scale as large as $n^{\Theta(d)}$ with d being the number of layers [5], we do not impose constraints of $\|\boldsymbol{z}\|_2 \leq r$ and $\|\boldsymbol{z}'\|_2 \leq r'$ in our experiments.

S	C	8	8	8	8	8	8	8	8
S Gen	8	8	8	8	8	8	8	8	8
Original	1	j	ł	١	١	ł	}	١	1
dcs_Gen	/	J	١	١	١	l	}	١	1
Original	8	8	8	B	8	8	\$	8	45
dcs_Gen	8	8	8	8	8	8	8	8	S
Original	ŧ	ł	١	1	1	1	1	1	}
dcs_Ger	1	١	١	1	l	/	1	1	}

Figure 4. Reconstructed images for digits 8 and 1 of MNIST under Gaussian measurements with $\sigma = 2$ and $\delta = 10$.



Figure 5. Reconstructed images for digits 8 and 1 of MNIST under Bernoulli measurements with $\sigma = 10$ and $\delta = 20$.

restarts with 1000 gradient descent steps per restart. The optimal reconstruction is chosen based on the lowest measurement error.

We test the sensing matrix $\mathbf{\Phi} \in \mathbb{R}^{n \times n}$ under both Gaussian and Bernoulli designs. Examples of reconstructed signals and corruptions are presented in Figure 4 and Figure 5, with the quantitative results showcased in Figure 6 and Figure 7. The ℓ_2 -norm of the test images is about 10, and the reconstructed images have impressive accuracy, even in a coarsely quantized and highly noisy setting with $\delta = 20$ and $\sigma = 10$.

Demixing Images in CelebA from Digits in MNIST: Since the image vectors of digits 8 and 1 have the same data dimension, the above experiment with x^* and v^* both from MNIST dataset is a demixing task without compression. In order to achieve compression, we additionally performed experiments for the case in which the signals are selected from the test images of the CelebA dataset [54] with data dimension $n = 3 \times 64 \times 64 = 12288$ and



Figure 6. Quantitative results of the performance of QCS_Gen under Gaussian measurements for digits 8 and 1 of MNIST.



Figure 7. Quantitative results of the performance of QCS_Gen under Bernoulli measurements for digits 8 and 1 of MNIST.

the corruptions correspond to the test images of the MNIST dataset with data dimension m = 784. Since the number of measurements m is much smaller than n, the measurement matrix $\mathbf{\Phi}$ of dimension $m \times n = 784 \times 12288$ is used for our simulations of corrupted sensing with compression (as $m \ll n$).

We train a VAE on the training set of the MNIST dataset, which comprises 60,000 images for digits 0 to 9. The decoder of the VAE is a fully connected neural network with two hidden layers and 500 neurons each, ReLU activations in the layers, and an input dimension of k = 20and output dimension of n = 784. We also employ the Adam optimizer with a mini-batch size of 100 and a learning rate of 0.001 for training.

The CelebA database contains more than 200,000 face images of celebrities, on which we



Figure 8. Reconstructed images for CelebA and MNIST under Gaussian measurements and $\sigma = 0.1$, $\delta = 1.0$.

train a deep convolutional generative adversarial network (DCGAN) following the settings in https://pytorch.org/tutorials/beginner/dcgan_faces_tutorial.html. The latent dimension of the generator for this model is 100 and the number of epochs for training is 20. We select 20 images from the test set of CelebA as our signals and 20 test images of MNIST as our corruptions, and we conduct 5 random trials. As the images of CelebA and MNIST differ significantly in their ℓ_2 -norm, we normalize both of them to have unit ℓ_2 -norm prior to generating the quantized observations. All other settings remain the same as those applied for the case of using two VAEs for digits 8 and 1 of MNIST.

Examples of reconstructed signals and corruptions can be seen in Figure 8 and Figure 9, and the relative error for each is quantified in Figure 10 and Figure 11. We observe that when m is much smaller than n, accurate reconstructions of the multiple test images can be achieved using a single draw of (Φ, ϵ, τ), as theoretically supported by our uniform recovery guarantee. Consistent with (3.98) in Theorem 3.16, under fixed δ , the error increases under larger σ . Also, while fixing the noise level σ , larger δ (that represents coarser quantization) corresponds to larger error, indicating that a trade-off between quantization resolution and recovery accuracy is important in practice.

5 Concluding Remarks In this work, we established uniform recovery guarantees for the problem of quantized corrupted sensing, using a dithered uniform quantizer, as well as structured or generative priors on the signal and corruption. Unlike the non-uniform results in [75], our results allow one to use a fixed realization of the sensing ensemble to accurately recover all signals and corruptions of interest. Our main techniques to prove uniformity are certain quantized embedding properties obtained from covering arguments; based on them, interestingly, our treatments for structured priors and generative priors are nearly unified. All the uniform error bounds exhibit decaying rates of $O(\frac{1}{\sqrt{m}})$. Specifically, the bounds for constrained Lasso typically coincide with the corresponding non-uniform ones up to logarithmic factors, while those for unconstrained Lasso usually display worse dependence on the structured parameters like sparsity level or matrix rank, creating a gap between constrained Lasso and unconstrained Lasso whose closing presents an open question. To demonstrate uniform recovery, in the ex-



Figure 9. Reconstructed images for CelebA and MNIST under Bernoulli measurements and $\sigma = 0.2$, $\delta = 0.2$.



Figure 10. Quantitative results of the performance of QCS_Gen under Gaussian measurements for CelebA and MNIST.

periments we use a fixed realization of (Φ, ϵ, τ) to accurately recover all (x^*, v^*) in some testing sets, and the numerical results are consistent with our uniform bounds. For future research, besides the aforementioned open question concerning using unconstrained Lasso for uniform recovery, corrupted sensing under 1-bit quantization is also worth investigation.

Appendix A. Auxiliary Facts. We collect auxiliary results in this Appendix, including some known concentration inequalities and technical lemmas that support our proofs.

A.1 Concentration Inequalities First of all, we provide the extended matrix deviation inequality that is well tailored to suit the analysis of corrupted sensing. We comment that setting $\mathcal{T} = \mathcal{T}_0 \times \{0\}$ for some $\mathcal{T}_0 \subset \mathbb{R}^n$ returns the regular matrix deviation inequality in [45, 50, 78]

Proposition A.1 (Extended matrix deviation inequality, Theorem 1 in [14]). Let $\Phi \in \mathbb{R}^{m \times n}$ be the sub-Gaussian sensing matrix described in Assumption 1, we let $\mathcal{T} \subset \mathbb{R}^n \times \mathbb{R}^m$ be a



Figure 11. Quantitative results of the performance of QCS_Gen under Bernoulli measurements for CelebA and MNIST.

bounded subset, then for any $t \geq 0$, the event

$$\sup_{(\boldsymbol{a},\boldsymbol{b})\in\mathcal{T}} \left| \|\boldsymbol{\Phi}\boldsymbol{a} + \sqrt{m}\boldsymbol{b}\|_2 - \sqrt{m} \left(\|\boldsymbol{a}\|_2^2 + \|\boldsymbol{b}\|_2^2 \right)^{1/2} \right| \le C \left(\gamma(\mathcal{T}) + t \cdot \operatorname{rad}(\mathcal{T}) \right)$$

holds with probability at least $1 - \exp(-t^2)$.

We will utilize the following result to prove a local version of the quantized product embedding property (see Lemma B.2).

Proposition A.2 (Exercise 8.6.5 in [78]). Consider a random process $(Z_a)_{a \in \mathcal{T}}$ indexed by points a in a bounded subset $\mathcal{T} \subset \mathbb{R}^n$. Assuming that $Z_0 = 0$, and for all $a, b \in \mathcal{T} \cup \{0\}$ we have

(A.1)
$$||Z_{a} - Z_{b}||_{\psi_{2}} \leq M ||a - b||_{2}.$$

Then for any $t \ge 0$, the event

(A.2)
$$\sup_{\boldsymbol{a}\in\mathcal{T}} \left| \boldsymbol{Z}_{\boldsymbol{a}} \right| \le CM \cdot \left[\boldsymbol{\omega}(\mathcal{T}) + t \cdot \operatorname{rad}(\mathcal{T}) \right]$$

holds with probability exceeding $1 - 2\exp(-t^2)$.

Next, we present a result that precisely characterizes the range of a low-complexity set under the sub-Gaussian map Φ . In particular, it provides uniform bound on the *l*-th largest measurement (since this is evidently no larger than the left-hand side of (A.3) below), which proves an effective tool in bounding the number of large perturbations (see (B.31) in the proof of Theorem B.3). We note that Proposition A.3 is the most crucial ingredient for getting our improvement on [81], which is to be presented in Appendix D.2.

Proposition A.3 (Adapted from Theorem 2.10 in [27]). Let $\Phi_1, ..., \Phi_m$ be independent, isotropic sub-Gaussian sensing vectors satisfying $\max_i \|\Phi_i\|_{\psi_2} = O(1)$, we consider some $\mathcal{T} \subset$

 \mathbb{R}^n . If $1 \leq l \leq m$, then for some absolute constants C_1, C_2 , the event¹⁵

(A.3)
$$\sup_{\boldsymbol{a}\in\mathcal{T}}\max_{\substack{I\subset[m]\\|I|\leq l}} \left(\frac{1}{l}\sum_{i\in I}|\langle \boldsymbol{\Phi}_i,\boldsymbol{a}\rangle|^2\right)^{1/2} \leq C_1\left(\frac{\omega(\mathcal{T})}{\sqrt{l}} + \operatorname{rad}(\mathcal{T})\sqrt{\log\frac{em}{l}}\right)$$

holds with probability at least $1 - 2\exp(-C_2 l \log \frac{em}{l})$.

Finally, we present a Chevet-type inequality that is indeed a simple outcome of Proposition A.2.

Proposition A.4 (Chevet-type inequality, Lemma 4 in [14]). Let Φ be the sub-Gaussian sensing matrix described in Assumption 1, $w \in \mathbb{R}^m$ be fixed, and \mathcal{T} be a bounded subset of \mathbb{R}^n . Then for any $t \geq 0$, the event

(A.4)
$$\sup_{\boldsymbol{a}\in\mathcal{T}} |\langle \boldsymbol{w}, \boldsymbol{\Phi}\boldsymbol{a}\rangle| \leq C \|\boldsymbol{w}\|_2 \big(\omega(\mathcal{T}) + t \cdot \operatorname{rad}(\mathcal{T})\big),$$

holds with probability exceeding $1 - 2\exp(-t^2)$.

A.2 Estimates of Geometric Quantities We collect some useful estimates on geometric quantities, specifically on Gaussian width, Gaussian complexity or Kolmogorov entropy (a notion equivalent to covering number). We begin with the well-known bounds on Gaussian width and Kolmogorov entropy for the structured priors of sparsity and low-rankness.

Proposition A.5 (See, e.g., [65]). We can estimate the Gaussian width of $\Sigma_s^n \cap \mathbb{S}^{n-1}$, $\mathbb{B}_2^n \cap \sqrt{s}\mathbb{B}_1^n$ as follows:

$$\max\left\{\omega(\Sigma_s^n \cap \mathbb{S}^{n-1}), \omega(\mathbb{B}_2^n \cap \sqrt{s}\mathbb{B}_1^n)\right\} \asymp \sqrt{s\log\frac{en}{s}}.$$

Proposition A.6 (See, e.g., [1, 11, 66]). Given some $\epsilon > 0$, recall that Σ_s^n is the set of all s-sparse signals in \mathbb{R}^n , and $M_r^{p,q}$ is the set of $p \times q$ matrices with rank not exceeding r, then we have

$$\mathscr{H}(\Sigma_s^n \cap \mathbb{B}_2^n, \epsilon) \le s \log\left(\frac{9n}{\epsilon s}\right) \text{ and } \mathscr{H}(M_r^{p,q} \cap \mathbb{B}_{\mathrm{F}}^{p,q}, \epsilon) \le 2r(p+q)\log\left(\frac{9}{\epsilon}\right).$$

Then, we give two results on relations between complexity quantities. With structured priors, Proposition A.7 will be used in the analysis of constrained Lasso, while Proposition A.8 is for analyzing unconstrained Lasso.

Proposition A.7. Given two cones $\mathcal{A} \subset \mathbb{R}^n$ and $\mathcal{B} \subset \mathbb{R}^m$, let $\mathcal{C} = \mathcal{A} \times \mathcal{B}$ and write $\mathcal{A}^* = \mathcal{A} \cap \mathbb{S}^{n-1}$, $\mathcal{B}^* = \mathcal{B} \cap \mathbb{S}^{m-1}$, $\mathcal{C}^* = \mathcal{C} \cap \mathbb{S}^{m+n-1}$. Then we have

$$\max\{\gamma(\mathcal{A}^*), \gamma(\mathcal{B}^*)\} \le \gamma(\mathcal{C}^*) \le \gamma(\mathcal{A}^*) + \gamma(\mathcal{B}^*).$$

¹⁵In the original statement of [27, Thm. 2.10], $\omega(\mathcal{T})$ in the right-hand side of (A.3) should be $\gamma(\mathcal{T})$, while we can safely use $\omega(\mathcal{T})$ here because (2.8) gives $\gamma(\mathcal{K}) \leq 2\omega(\mathcal{K}) + 2\operatorname{rad}(\mathcal{T})$, and we observe that $\frac{\operatorname{rad}(\mathcal{T})}{l^{1/2}} \leq \operatorname{rad}(\mathcal{T})(\log \frac{em}{l})^{1/2}$ holds for any $l \in [1, m]$.

Proof. Because $0 \in \mathcal{A}$ and $0 \in \mathcal{B}$, it is easy to see that $\mathcal{A}^* \times \{0\} \subset \mathcal{C}^*$ and $\{0\} \times \mathcal{B}^* \subset \mathcal{C}^*$, which implies $\gamma(\mathcal{A}^*) = \gamma(\mathcal{A}^* \times \{0\}) \leq \gamma(\mathcal{C}^*)$, and similarly $\gamma(\mathcal{B}^*) \leq \gamma(\mathcal{C}^*)$. To prove the second inequality, we first observe that $\mathcal{C}^* \subset \bigcup_{c \in [0,1]} (c\mathcal{A}^* \times \sqrt{1-c^2}\mathcal{B}^*)$, and we let g_i be standard Gaussian vector with self-evident dimension. Then following the definition of Gaussian complexity we have

$$\begin{split} \gamma(\mathcal{C}^*) &\leq \mathbb{E} \sup_{\substack{c \in [0,1] \\ \boldsymbol{a} \in \mathcal{A}^*, \boldsymbol{b} \in \mathcal{B}^*}} \left| \langle \boldsymbol{g}_1, c \cdot \boldsymbol{a} \rangle + \langle \boldsymbol{g}_2, \sqrt{1 - c^2} \cdot \boldsymbol{b} \rangle \right| \\ &\leq \mathbb{E} \sup_{\boldsymbol{a} \in \mathcal{A}^*} \left| \langle \boldsymbol{g}_1, \boldsymbol{a} \rangle \right| + \mathbb{E} \sup_{\boldsymbol{b} \in \mathcal{B}^*} \left| \langle \boldsymbol{g}_2, \boldsymbol{b} \rangle \right| = \gamma(\mathcal{A}^*) + \gamma(\mathcal{B}^*), \end{split}$$

which completes the proof.

Proposition A.8. Let $f(\cdot)$ be a norm in \mathbb{R}^n , $g(\cdot)$ be a norm in \mathbb{R}^m , $\lambda_1, \lambda_2, \alpha_x, \alpha_v$ be some positive parameters, C be some absolute constant. Define the cone

(A.5)
$$C(\lambda_1, \lambda_2) = \{(\boldsymbol{a}, \boldsymbol{b}) \in \mathbb{R}^n \times \mathbb{R}^m : \lambda_1 f(\boldsymbol{a}) + \lambda_2 g(\boldsymbol{b}) \le C\lambda_1 \alpha_{\boldsymbol{x}} \|\boldsymbol{a}\|_2 + C\lambda_2 \alpha_{\boldsymbol{v}} \|\boldsymbol{b}\|_2 \}$$

and let $\mathcal{C}^* = \mathcal{C}(\lambda_1, \lambda_2) \cap \mathbb{S}^{n+m-1}$, then we have

(A.6)
$$\omega(\mathcal{C}^*) \lesssim \left(\alpha_{\boldsymbol{x}} + \frac{\lambda_2 \alpha_{\boldsymbol{v}}}{\lambda_1}\right) \cdot \omega(\mathbb{B}_f^n) + \left(\alpha_{\boldsymbol{v}} + \frac{\lambda_1 \alpha_{\boldsymbol{x}}}{\lambda_2}\right) \cdot \omega(\mathbb{B}_g^m)$$

Proof. Let $h_1 \sim \mathcal{N}(0, I_n), h_2 \sim \mathcal{N}(0, I_m)$, then we proceed as

(A.7)
$$\omega(\mathcal{C}^*) = \mathbb{E} \sup_{(\boldsymbol{a}, \boldsymbol{b}) \in \mathcal{C}^*} \langle \boldsymbol{h}_1, \boldsymbol{a} \rangle + \langle \boldsymbol{h}_2, \boldsymbol{b} \rangle$$

(A.8)
$$\leq \mathbb{E} \sup_{(\boldsymbol{a},\boldsymbol{b})\in\mathcal{C}^*} \left[\lambda_1 f(\boldsymbol{a}) \cdot f^*\left(\frac{\boldsymbol{h}_1}{\lambda_1}\right) + \lambda_2 g(\boldsymbol{b}) \cdot g^*\left(\frac{\boldsymbol{h}_2}{\lambda_2}\right)\right]$$

(A.9)
$$\leq \mathbb{E} \sup_{(\boldsymbol{a},\boldsymbol{b})\in\mathcal{C}^*} \left[\left(\lambda_1 f(\boldsymbol{a}) + \lambda_2 g(\boldsymbol{b}) \right) \cdot \left(\sup_{\boldsymbol{a}\in\mathbb{B}_f^n} \left\langle \frac{\boldsymbol{h}_1}{\lambda_1}, \boldsymbol{a} \right\rangle + \sup_{\boldsymbol{b}\in\mathbb{B}_g^m} \left\langle \frac{\boldsymbol{h}_2}{\lambda_2}, \boldsymbol{b} \right\rangle \right] \right]$$

(A.10)
$$\lesssim \mathbb{E}\left[(\lambda_1 \alpha_{\boldsymbol{x}} + \lambda_2 \alpha_{\boldsymbol{v}}) \cdot \left(\sup_{\boldsymbol{a} \in \mathbb{B}_f^n} \left\langle \frac{\boldsymbol{h}_1}{\lambda_1}, \boldsymbol{a} \right\rangle + \sup_{\boldsymbol{b} \in \mathbb{B}_g^m} \left\langle \frac{\boldsymbol{h}_2}{\lambda_2}, \boldsymbol{b} \right\rangle \right] \right]$$

(A.11)
$$\lesssim \left(\alpha_{\boldsymbol{x}} + \frac{\lambda_2 \alpha_{\boldsymbol{v}}}{\lambda_1}\right) \cdot \omega(\mathbb{B}_f^n) + \left(\alpha_{\boldsymbol{v}} + \frac{\lambda_1 \alpha_{\boldsymbol{x}}}{\lambda_2}\right) \cdot \omega(\mathbb{B}_g^m),$$

where in (A.8) we apply Hölder's inequality with proper re-scaling, (A.9) follows from the definitions of $f^*(\cdot), g^*(\cdot)$ and some simple relaxation, (A.10) is due to (A.5) and $\mathcal{C}^* \subset \mathbb{S}^{n+m-1}$, and in the last inequality we use the definition of Gaussian width. The proof is complete.

We close this subsection with another set of geometric quantities estimates specialized to the analysis of generative priors.

Proposition A.9. Let $\mathcal{K}_{\boldsymbol{x}}$ and $\mathcal{K}_{\boldsymbol{v}}$ be described in Assumption 5, then the following statements hold true for any $\eta_1 \in (0, Lr), \ \eta_2 \in (0, L'r')$: (a) The Kolmogorov entropy of $\mathcal{K}_{\boldsymbol{x}}$ and $\mathcal{K}_{\boldsymbol{v}}$ is bounded as follows:

(A.12)
$$\mathscr{H}(\mathcal{K}_{\boldsymbol{x}},\eta_1) \le k \log \frac{3Lr}{\eta_1}, \quad \mathscr{H}(\mathcal{K}_{\boldsymbol{v}},\eta_2) \le k' \log \frac{3L'r'}{\eta_2}$$

(b) Let $\mathcal{K}_{\boldsymbol{x}}^- = \mathcal{K}_{\boldsymbol{x}} - \mathcal{K}_{\boldsymbol{x}}, \ \mathcal{K}_{\boldsymbol{v}}^- = \mathcal{K}_{\boldsymbol{v}} - \mathcal{K}_{\boldsymbol{v}}$, then we have

(A.13)
$$\mathscr{H}(\mathcal{K}_{\boldsymbol{x}}^{-},\eta_{1}) \leq 2k \log \frac{6Lr}{\eta_{1}}, \quad \mathscr{H}(\mathcal{K}_{\boldsymbol{v}}^{-},\eta_{2}) \leq 2k' \log \frac{6L'r'}{\eta_{2}}.$$

Moreover, for any $0 < \eta < \min\{Lr, L'r'\}$ it holds that

(A.14)
$$\mathscr{H}(\mathcal{K}_{\boldsymbol{x}}^{-} \times \mathcal{K}_{\boldsymbol{v}}^{-}, \eta) \leq 2k \log \frac{12Lr}{\eta} + 2k' \log \frac{12L'r'}{\eta}$$

(c) Given any $\mu \in (0, 1)$, we define \mathcal{E} and \mathcal{E}^* as

(A.15)
$$\mathcal{E} = \left\{ (\boldsymbol{c}, \boldsymbol{d}) : \boldsymbol{c} \in \mathcal{K}_{\boldsymbol{x}}^{-}, \ \boldsymbol{d} \in \mathcal{K}_{\boldsymbol{v}}^{-}, \ (\|\boldsymbol{c}\|_{2}^{2} + \|\boldsymbol{d}\|_{2}^{2})^{1/2} \ge 2\mu \right\}$$

(A.16)
$$\mathcal{E}^* = \left\{ (\boldsymbol{c}, \boldsymbol{d}) / (\|\boldsymbol{c}\|_2^2 + \|\boldsymbol{d}\|_2^2)^{1/2} : (\boldsymbol{c}, \boldsymbol{d}) \in \mathcal{E} \right\},$$

then for any $0 < \eta < \min\{Lr, L'r'\}$ we have

(A.17)
$$\mathscr{H}(\mathcal{E}^*,\eta) \le 2k \log \frac{24Lr}{\mu\eta} + 2k' \log \frac{24L'r'}{\mu\eta}$$

Moreover, we have

(A.18)
$$\omega(\mathcal{E}^*) \lesssim \left(k \log \frac{Lr}{\mu} + k' \log \frac{L'r'}{\mu}\right)^{1/2}.$$

Proof. We present the proofs of (a), (b) and (c) separately.

The Proof of (a) By the covering number of the ℓ_2 -ball [78, Coro. 4.2.13], we can construct \mathcal{G} as a $\left(\frac{\eta_1}{L}\right)$ -net of $\mathbb{B}_2^k(r)$ with cardinality not exceeding $\left(\frac{2Lr}{\eta_1}+1\right)^k$, and hence not exceeding $\left(\frac{3Lr}{\eta_1}\right)^k$ because $\eta_1 < Lr$. Since $G(\cdot)$ is L-Lipschitz, $G(\mathcal{G})$ is an η_1 -net of $\mathcal{K}_{\boldsymbol{x}} = G(\mathbb{B}_2^k(r))$, thus implying $\mathscr{H}(\mathcal{K}_{\boldsymbol{x}},\eta_1) \leq k \log \frac{3Lr}{\eta_1}$. One can similarly derive the bound for $\mathscr{H}(\mathcal{K}_{\boldsymbol{v}},\eta_2)$.

The Proof of (b) From the result (a) that we just proved, we can construct \mathcal{G}_1 as an $\left(\frac{\eta_1}{2}\right)$ -net of $\mathcal{K}_{\boldsymbol{x}}$ such that $\log |\mathcal{G}_1| \leq k \log \frac{6Lr}{\eta_1}$. It is not hard to show $\mathcal{G}_1 - \mathcal{G}_1$ is a η_1 -net of $\mathcal{K}_{\boldsymbol{x}}^-$, and note that $\log |\mathcal{G}_1 - \mathcal{G}_1| \leq \log |\mathcal{G}_1|^2 \leq 2k \log \frac{6Lr}{\eta_1}$. We can similarly prove $\mathscr{H}(\mathcal{K}_{\boldsymbol{v}}^-, \eta_2) \leq 2k' \log \frac{6L'r'}{\eta_2}$, hence (A.13) follows. To prove (A.14), it is sufficient to note the following simple fact: if \mathcal{G}_2 is an $\frac{\eta}{2}$ -net of $\mathcal{K}_{\boldsymbol{x}}^-$, \mathcal{G}_3 is an $\frac{\eta}{2}$ -net of $\mathcal{K}_{\boldsymbol{v}}^-$, then $\mathcal{G}_2 \times \mathcal{G}_3$ is an η -net of $\mathcal{K}_{\boldsymbol{x}}^- \times \mathcal{K}_{\boldsymbol{v}}^-$.

The Proof of (c) Because $\mathcal{E} \subset \mathcal{K}_{\boldsymbol{x}} \times \mathcal{K}_{\boldsymbol{v}}^-$, by (2.4) and (A.14), we can let \mathcal{G}_4 be an $(\mu\eta)$ -net of \mathcal{E} satisfying $\log |\mathcal{G}_4| \leq 2k \log \frac{24Lr}{\mu\eta} + 2k' \log \frac{24L'r'}{\mu\eta}$. Then it suffices to show that

$$\mathcal{G}_4^* = \Big\{ rac{(m{a},m{b})}{(\|m{a}\|_2^2 + \|m{b}\|_2^2)^{1/2}} : (m{a},m{b}) \in \mathcal{G}_4 \Big\}$$

is an η -net of \mathcal{E}^* , which we prove in the following two dot points:

- By (A.16), any $(a_1, b_1) \in \mathcal{E}^*$ can be written as $c_2/\|c_2\|_2 =: (a_2, b_2)/(\|a_2\|_2^2 + \|b_2\|_2^2)^{1/2}$ for some $c_2 := (a_2, b_2) \in \mathcal{E}$. Since \mathcal{G}_4 is a $(\mu\eta)$ -net of \mathcal{E} , we can pick $c_3 := (a_3, b_3) \in \mathcal{G}_4$ such that $\|c_3 - c_2\|_2 \le \mu\eta$.
- Note that $c_3/\|c_3\|_2 \in \mathcal{G}_4^*$, and we have

$$\begin{split} \left\| \frac{\boldsymbol{c}_2}{\|\boldsymbol{c}_2\|_2} - \frac{\boldsymbol{c}_3}{\|\boldsymbol{c}_3\|_2} \right\|_2 &\leq \left\| \frac{\boldsymbol{c}_2}{\|\boldsymbol{c}_2\|_2} - \frac{\boldsymbol{c}_3}{\|\boldsymbol{c}_2\|_2} \right\|_2 + \left\| \frac{\boldsymbol{c}_3}{\|\boldsymbol{c}_2\|_2} - \frac{\boldsymbol{c}_3}{\|\boldsymbol{c}_3\|_3} \right\|_2 \\ &\leq \frac{\|\boldsymbol{c}_2 - \boldsymbol{c}_3\|_2}{\|\boldsymbol{c}_2\|_2} + \frac{\|\|\boldsymbol{c}_3\|_2 - \|\boldsymbol{c}_2\|_2\|}{\|\boldsymbol{c}_2\|_2} \\ &\leq \frac{2\|\boldsymbol{c}_2 - \boldsymbol{c}_3\|}{\|\boldsymbol{c}_2\|_2} \leq \frac{2\eta\mu}{2\mu} = \eta, \end{split}$$

where the last inequality holds because $c_2 \in \mathcal{E}$ satisfies $||c_2||_2 \ge 2\mu$. Therefore, \mathcal{G}_4^* is an η -net of \mathcal{E}^* .

Therefore, $\mathscr{H}(\mathcal{E}^*, \eta) \leq \log |\mathcal{G}_4|$, the bound in (A.17) follows. It remains to prove (A.18), and our strategy is to estimate $\omega(\mathcal{E}^*)$ from $\mathscr{H}(\mathcal{E}^*, \eta)$ via Dudley's inequality (2.7). Note that $\mathcal{E}^* \subset \mathbb{S}^{n+m-1}$, we thus have

$$\begin{split} \omega(\mathcal{E}^*) &\lesssim \int_0^\infty \sqrt{\mathscr{H}(\mathcal{E}^*, \eta)} \, \mathrm{d}\eta = \int_0^2 \sqrt{\mathscr{H}(\mathcal{E}^*, \eta)} \, \mathrm{d}\eta \\ &\leq \sqrt{2k} \int_0^2 \sqrt{\log \frac{24Lr}{\mu\eta}} \, \mathrm{d}\eta + \sqrt{2k'} \int_0^2 \sqrt{\log \frac{24L'r'}{\mu\eta}} \, \mathrm{d}\eta \\ &\lesssim \left(k \log \frac{Lr}{\mu} + k' \log \frac{L'r'}{\mu}\right)^{1/2}, \end{split}$$

where we apply (A.17) in the second inequality. The proof is complete.

Appendix B. Quantized Product Embedding. In this appendix, we develop quantized product embedding (QPE) property for analyzing the uniform recovery of Lasso in quantized corrupted sensing. In brief, QPE states that the dithered uniform quantization universally preserves inner product. More precisely, with random dithering given by $\tau \sim \mathscr{U}([-\frac{\tau}{2}, \frac{\tau}{2}]^m)$, QPE ensures that $\langle Q_{\delta}(\boldsymbol{a}+\boldsymbol{\tau}), \boldsymbol{b} \rangle$ is close to $\langle \boldsymbol{a}, \boldsymbol{b} \rangle$, with the closeness holding uniformly for all $(\boldsymbol{a}, \boldsymbol{b})$ in some constraint sets of interest.

B.1 An Observation and Local QPE We start from a simple observation due to Xu and Jacques [81, Lem. 6.1], which bounds the number of measurements exhibiting some "discontinuity" due to the quantizer $Q_{\delta}(\cdot)$.

Lemma B.1. Given some $\delta, \zeta > 0$ satisfying $\zeta \in (0, \frac{\delta}{2})$, and a fixed $\boldsymbol{a} = [a_i] \in \mathbb{R}^m$. Associated with $\boldsymbol{\tau} = [\tau_i] \sim \mathscr{U}[-\frac{\delta}{2}, \frac{\delta}{2}]^m$, we define the random set

(B.1)
$$\mathcal{Z} = \left\{ i \in [m] : \mathcal{Q}_{\delta}(a_i + \tau_i + t) \text{ is discontinuous in } [-\zeta, \zeta] \right\}.$$

Then, for any $t \geq 0$, we have

(B.2)
$$\mathbb{P}\left(|\mathcal{Z}| \ge \frac{5m\zeta}{\delta}\right) \le \exp\left(-\frac{m\zeta}{\delta}\right),$$

where $|\mathcal{Z}|$ is the random variable denoting the cardinality of \mathcal{Z} .

Proof. First, we note that $\mathcal{Q}_{\delta}(a_i + \tau_i + t)$ is discontinuous in $[-\zeta, \zeta]$ if and only if $\mathcal{Q}_{\delta}(t)$ is discontinuous in $[a_i + \tau_i - \zeta, a_i + \tau_i + \zeta]$, and further, the latter statement is evidently equivalent to the event

$$E_1 = \big\{ [a_i + \tau_i - \zeta, a_i + \tau_i + \zeta] \cap (\delta \mathbb{Z}) = \emptyset \big\}.$$

Moreover, under our assumptions of $\tau_i \sim \mathscr{U}([-\frac{\delta}{2}, \frac{\delta}{2}])$ and $\zeta \in (0, \frac{\delta}{2})$, it is not hard to see that, $\mathbb{P}(E_1) = \frac{2\zeta}{\delta}$ holds true independent of the location of $[a_i - \zeta, a_i + \zeta]$. Therefore, $|\mathcal{Z}| \sim \text{Bin}(m, \frac{2\zeta}{\delta})$ (i.e., $|\mathcal{Z}|$ follows a binomial distribution with m trials and probability of success $\frac{2\zeta}{\delta}$ for each), hence it can be written as $|\mathcal{Z}| = \sum_{k=1}^m Z_k$ with i.i.d. $Z_k \sim \text{Bernoulli}(\frac{2\zeta}{\delta})$. Because for any integer $q \geq 3$ we have $\sum_{k=1}^m \mathbb{E}|Z_k|^q = \frac{2m\zeta}{\delta}$, we can invoke Bernstein's inequality [6, Thm. 2.10] to obtain that, for any t > 0, the event $\sum_{k=1}^m (Z_k - \mathbb{E}Z_k) \leq 2(\frac{m\zeta t}{\delta})^{1/2} + t$ holds with probability exceeding $1 - \exp(-t)$. Setting $t = \frac{m\zeta}{\delta}$ and substituting $\mathbb{E}|\mathcal{Z}| = \sum_{k=1}^m \mathbb{E}Z_k = \frac{2m\zeta}{\delta}$ yield the desired claim.

We then establish the QPE with quantized measurements generated by a fixed signalcorruption pair (a, b). This "local" QPE will be sufficient for establishing non-uniform guarantee, and indeed, the lemma below readily follows from [75, Lem. 2]. We provide a proof for completeness.

Lemma B.2 (Local QPE). Given some bounded sets $\mathcal{A} \subset \mathbb{R}^n$, $\mathcal{B} \subset \mathbb{R}^m$, $\mathcal{E} \subset \mathbb{R}^{n+m}$ and some $\delta > 0$, we fix $\mathbf{a} \in \mathcal{A}$ and $\mathbf{b} \in \mathcal{B}$, and assume that the sub-Gaussian matrix $\mathbf{\Phi} \in \mathbb{R}^{m \times n}$, sub-Gaussian noise $\boldsymbol{\epsilon}$ and the random dither $\boldsymbol{\tau} \sim \mathscr{U}[-\frac{\delta}{2}, \frac{\delta}{2}]^m$ are as described in Assumption 1, then for $\mathbf{c} \in \mathbb{R}^n$ and $\mathbf{d} \in \mathbb{R}^m$, and some absolute constant C, the event

(B.3)
$$\sup_{(\boldsymbol{c},\boldsymbol{d})\in\mathcal{E}} \left| \left\langle \boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}, \boldsymbol{\Phi}\boldsymbol{c} + \sqrt{m}\boldsymbol{d} \right\rangle \right| \leq C\sqrt{m}\delta\left(\omega(\mathcal{E}) + u \cdot \operatorname{rad}(\mathcal{E})\right)$$

holds with probability at least $1 - 2\exp(-u^2)$, where we denote the quantization noise of $\Phi a + \sqrt{m}b$ by the shorthand

(B.4)
$$\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}} := \mathcal{Q}_{\delta}(\boldsymbol{\Phi}\boldsymbol{a} + \sqrt{m}\boldsymbol{b} + \boldsymbol{\epsilon} + \boldsymbol{\tau}) - (\boldsymbol{\Phi}\boldsymbol{a} + \sqrt{m}\boldsymbol{b} + \boldsymbol{\epsilon}).$$

Proof. We note that it suffices to consider $\epsilon = 0$; for the general setting, since ϵ is independent of (Φ, τ) , we can condition on ϵ and write

(B.5)
$$\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}} = \mathcal{Q}_{\delta}(\boldsymbol{\Phi}\boldsymbol{a} + \sqrt{m}\boldsymbol{\tilde{b}} + \boldsymbol{\tau}) - (\boldsymbol{\Phi}\boldsymbol{a} + \sqrt{m}\boldsymbol{\tilde{b}})$$

with $\tilde{\boldsymbol{b}} := \boldsymbol{b} + \frac{\boldsymbol{\epsilon}}{\sqrt{m}}$, then the result can be obtained from the case of $\boldsymbol{\epsilon} = 0$.

To get the desired bound on $\sup_{(c,d)\in\mathcal{E}} |\langle \boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}, \boldsymbol{\Phi}\boldsymbol{c} + \sqrt{m}\boldsymbol{d} \rangle|$, we view $\langle \boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}, \boldsymbol{\Phi}\boldsymbol{c} + \sqrt{m}\boldsymbol{d} \rangle$ as a random process indexed by $(\boldsymbol{c},\boldsymbol{d}) \in \mathcal{E}$ and seek to apply Proposition A.2. Given any $(\boldsymbol{c}_1,\boldsymbol{d}_1), (\boldsymbol{c}_2,\boldsymbol{d}_2) \in \mathcal{E} \cup \{0\}$, we first verify (A.1) as follows:

(B.6)
$$\left\|\left\langle \boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}, \boldsymbol{\Phi}\boldsymbol{c}_{1} + \sqrt{m}\boldsymbol{d}_{1} \right\rangle - \left\langle \boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}, \boldsymbol{\Phi}\boldsymbol{c}_{2} + \sqrt{m}\boldsymbol{d}_{2} \right\rangle \right\|_{\psi_{2}}$$

(B.7)
$$\leq \left\| \langle \boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}, \boldsymbol{\Phi}(\boldsymbol{c}_1 - \boldsymbol{c}_2) \rangle \right\|_{\psi_2} + \left\| \langle \boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}, \sqrt{m}(\boldsymbol{d}_1 - \boldsymbol{d}_2) \rangle \right\|_{\psi_2}$$

(B.8) $\lesssim \sqrt{m\delta} \cdot \|\boldsymbol{c}_1 - \boldsymbol{c}_2\|_2 + \sqrt{m\delta} \cdot \|\boldsymbol{d}_1 - \boldsymbol{d}_2\|_2$

(B.9)
$$\lesssim \sqrt{m\delta} \cdot \|(\boldsymbol{c}_1, \boldsymbol{d}_1) - (\boldsymbol{c}_2, \boldsymbol{d}_2)\|_2$$

where (B.8) is due to

(B.10)
$$\|\langle \boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}, \sqrt{m}(\boldsymbol{d}_1 - \boldsymbol{d}_2) \rangle\|_{\psi_2} \lesssim \sqrt{m}\delta \cdot \|\boldsymbol{d}_1 - \boldsymbol{d}_2\|_2$$

(B.11)
$$\|\langle \boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}, \boldsymbol{\Phi}(\boldsymbol{c}_1 - \boldsymbol{c}_2) \rangle\|_{\psi_2} \lesssim \sqrt{m}\delta \cdot \|\boldsymbol{c}_1 - \boldsymbol{c}_2\|_2.$$

Note that (B.10) follows from $\|\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}\|_{\psi_2} = O(\delta)$ (see (2.14)). To explain (B.11), by $\|\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}\|_{\infty} \leq \delta$ (see (2.12)), and with respect to the randomness of τ_k we have $\mathbb{E}_{\tau_k}[(\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}})_k] = 0$ (see (2.13)) that implies

(B.12)
$$\mathbb{E}\left((\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}})_{k}\boldsymbol{\Phi}_{k}^{\top}(\boldsymbol{c}_{1}-\boldsymbol{c}_{2})\right)=\mathbb{E}\left(\mathbb{E}_{\tau_{k}}[(\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}})_{k}]\boldsymbol{\Phi}_{k}^{\top}(\boldsymbol{c}_{1}-\boldsymbol{c}_{2})\right)=0,$$

we can use (2.3) to obtain

(B.13)
$$\left\| \langle \boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}, \boldsymbol{\Phi}(\boldsymbol{c}_1 - \boldsymbol{c}_2) \rangle \right\|_{\psi_2}^2 \leq \sum_{k=1}^m \left\| (\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}})_k \boldsymbol{\Phi}_k^\top (\boldsymbol{c}_1 - \boldsymbol{c}_2) \right\|_{\psi_2}^2 \lesssim m \delta^2 \|\boldsymbol{c}_1 - \boldsymbol{c}_2\|_2^2.$$

Now we invoke Proposition A.2 to obtain that for some absolute constant C, the event

(B.14)
$$\sup_{(\boldsymbol{c},\boldsymbol{d})\in\mathcal{E}} \left| \langle \boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}, \boldsymbol{\Phi}\boldsymbol{c} + \sqrt{m}\boldsymbol{d} \rangle \right| \le C\sqrt{m}\delta\big(\omega(\mathcal{E}) + u \cdot \operatorname{rad}(\mathcal{E})\big)$$

holds with probability exceeding $1 - \exp(-u^2)$, as desired.

B.2 Global QPE: The General Version In pursuit of a uniform guarantee, the local QPE in Lemma B.2 is no longer enough. Rather, we will need a global QPE property that holds universally for all $\xi_{a,b}$ with $(a, b) \in \mathcal{A} \times \mathcal{B}$ (see (B.4) for $\xi_{a,b}$). To achieve the desired uniformity, we follow the standard approach of the covering argument that comprises two steps: (i) establish the desired property over the discrete nets \mathcal{G}_1 and \mathcal{G}_2 that respectively approximate \mathcal{A} and \mathcal{B} , (ii) extend the desired property from $(a, b) \in \mathcal{G}_1 \times \mathcal{G}_2$ to $(a, b) \in \mathcal{A} \times \mathcal{B}$. While (i) can be done by applying Lemma B.2 along with a union bound, the key challenge lies in (ii) due to the discontinuity of $\mathcal{Q}_{\delta}(\cdot)$ (note that the extension in (ii) typically relies on certain notion of continuity). We overcome the difficulty by a strategy similar to [81, Prop. 6.1] (see (B.26) below that characterizes certain continuity of the dithered quantizer), but with the sub-Gaussianity of Φ we manage to provide refinements by using Proposition A.3 (see more discussions in Appendix D).

Theorem B.3 (Global QPE). Given some bounded sets $\mathcal{A} \subset \mathbb{R}^n$, $\mathcal{B} \subset \mathbb{R}^m$, $\mathcal{E} \subset \mathbb{R}^{n+m}$ and some $\delta > 0$, we further define $\mathcal{C} \subset \mathbb{R}^n$ and $\mathcal{D} \subset \mathbb{R}^m$ as

(B.15)
$$\mathcal{C} = \left\{ \boldsymbol{c} \in \mathbb{R}^n : \exists \boldsymbol{d} \in \mathbb{R}^m, \text{ such that } (\boldsymbol{c}, \boldsymbol{d}) \in \mathcal{E} \right\},\$$

(B.16)
$$\mathcal{D} = \left\{ \boldsymbol{d} \in \mathbb{R}^m : \exists \boldsymbol{c} \in \mathbb{R}^n, \text{ such that } (\boldsymbol{c}, \boldsymbol{d}) \in \mathcal{E} \right\},\$$

and assume that the sub-Gaussian matrix $\mathbf{\Phi} \in \mathbb{R}^{m \times n}$, sub-Gaussian noise $\boldsymbol{\epsilon}$ and the random dither $\boldsymbol{\tau} \sim \mathscr{U}[-\frac{\delta}{2}, \frac{\delta}{2}]^m$ are as described in Assumption 1. Suppose that (ζ, ρ_1, ρ_2) are positive scalars satisfying

(B.17)
$$\zeta \le c_1 \delta, \ \rho_1 \le \frac{c_1 \zeta}{(\log \frac{\delta}{\zeta})^{1/2}}, \ \rho_2 \le c_1 \zeta$$

for some sufficiently small c_1 . If for some large enough absolute constant $C_2 > 0$ we have

(B.18)
$$m \ge \frac{2\delta \cdot \mathscr{H}(\mathcal{A}, \rho_1) + 2\delta \cdot \mathscr{H}(\mathcal{B}, \rho_2)}{\zeta} + \frac{C_2 \cdot \omega^2(\mathcal{A}_{\text{loc}}^{(\rho_1)})}{\zeta^2}$$

then with the quantization noise $\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}$ being given in (B.4), the event

(B.19)
$$\sup_{\boldsymbol{a}\in\mathcal{A}}\sup_{\boldsymbol{b}\in\mathcal{B}}\sup_{(\boldsymbol{c},\boldsymbol{d})\in\mathcal{E}}\left|\langle\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}},\boldsymbol{\Phi}\boldsymbol{c}+\sqrt{m}\boldsymbol{d}\rangle\right| \lesssim \delta\sqrt{m}\cdot\omega(\mathcal{E})+\delta\sqrt{m\hat{U}}\cdot\mathrm{rad}(\mathcal{D})$$

(B.20)
$$+ \delta \sqrt{m} \cdot \operatorname{rad}(\mathcal{C}) \cdot \left(\sqrt{\mathscr{H}(\mathcal{A}, \rho_1)} + \sqrt{\mathscr{H}(\mathcal{B}, \rho_2)} + \frac{\hat{U}}{\sqrt{m}} \left[\log \frac{\delta}{\zeta} \right]^{1/2} \right)$$

holds with probability exceeding $1 - 12 \exp(-c_3 \mathscr{H}(\mathcal{A}, \rho_1) - c_3 \mathscr{H}(\mathcal{B}, \rho_2))$ on a single draw of (Φ, τ) . In (B.20), \hat{U} is a quantity scaling as

(B.21)
$$\hat{U} \approx \frac{m\zeta}{\delta} + \frac{m\rho_2^2}{\zeta^2} + \frac{\omega^2(\mathcal{A}_{\text{loc}}^{(\rho_1)})}{\zeta^2}.$$

Proof. Again, we can simply concentrate on the case where $\boldsymbol{\epsilon} = 0$; we can reduce the general case to the case with $\boldsymbol{\epsilon} = 0$ by conditioning on $\boldsymbol{\epsilon}$ and writing (B.5) with $\tilde{\boldsymbol{b}} = \boldsymbol{b} + \frac{\boldsymbol{\epsilon}}{\sqrt{m}} \in \tilde{\mathcal{B}} := \mathcal{B} + \frac{\boldsymbol{\epsilon}}{\sqrt{m}}$; since $\mathscr{H}(\mathcal{B}, \cdot) = \mathscr{H}(\tilde{\mathcal{B}}, \cdot)$ always holds, the result can be readily obtained from the case of $\boldsymbol{\epsilon} = 0$. We seek to bound

(B.22)
$$\sup_{\boldsymbol{a}\in\mathcal{A}}\sup_{\boldsymbol{b}\in\mathcal{B}}\sup_{(\boldsymbol{c},\boldsymbol{d})\in\mathcal{E}}\left|\langle\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}},\boldsymbol{\Phi}\boldsymbol{c}+\sqrt{m}\boldsymbol{d}\rangle\right|.$$

We pause to provide an outline for this most technical proof in this work (see Table 1 in Appendix E for a table of the recurring notation):

- In Step 1 we bound the random process over nets of \mathcal{A} and \mathcal{B} , which is done by applying local QPE (Lemma B.2) and then a union bound.
- In Step 2 we build three useful events E₁, E₂, E₃: E₁ in (B.26) bounds the number of measurements exhibiting some discontinuity due to the quantizer (those in Z_{a,b} as per (B.24)); E₂ in (B.34) bounds the number of measurements exhibiting large perturbations associated with a ∈ A (those in J_a^A as per (B.29)); and E₃ in (B.36) bounds the number of measurements with large perturbations associated with b ∈ B (those in J_b^B as per (B.30)).

- In Step 3 we strengthen the bound from Step 1 to a uniform bound over $\mathcal{A} \times \mathcal{B}$. We provide different treatments to two classes of measurements. The first class collects the "bad" ones in $\mathcal{Z}_{a,b} \cup \mathcal{J}_a^{\mathcal{A}} \cup \mathcal{J}_b^{\mathcal{B}}$ (B.39) that only account for a few measurements (see (B.41)), the second class collects the remaining benign measurements that enjoy some nice property (see (B.51)).
- In **Step 4** we choose parameters and use (B.17) and (B.18) to get the bound and probability in the theorem statement.

Step 1: Uniform Bound Over Nets For some $\rho_1, \rho_2 > 0$ to be chosen, we let \mathcal{G}_1 be the minimal ρ_1 -net of \mathcal{A} with $\log |\mathcal{G}_1| = \mathscr{H}(\mathcal{A}, \rho_1)$, \mathcal{G}_2 be the minimal ρ_2 -net of \mathcal{B} with $\log |\mathcal{G}_2| = \mathscr{H}(\mathcal{B}, \rho_2)$. Then, for any $t_1 > 0$, we apply the non-uniform bound in Lemma B.2 along with a union bound over $(\boldsymbol{a}, \boldsymbol{b}) \in \mathcal{G}_1 \times \mathcal{G}_2$, yielding that the event

(B.23)
$$\sup_{\boldsymbol{a}\in\mathcal{G}_1} \sup_{\boldsymbol{b}\in\mathcal{G}_2} \sup_{(\boldsymbol{c},\boldsymbol{d})\in\mathcal{E}} \left| \langle \boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}, \boldsymbol{\Phi}\boldsymbol{c} + \sqrt{m}\boldsymbol{d} \rangle \right| \le C\sqrt{m}\delta\big(\omega(\mathcal{E}) + t_1 \cdot \operatorname{rad}(\mathcal{E})\big)$$

holds with probability at least $1 - 2\exp(\mathscr{H}(\mathcal{A},\rho_1) + \mathscr{H}(\mathcal{B},\rho_2) - t_1^2)$.

Step 2: Some Useful Events Recall that we need to further strengthen (B.23) from $(a, b) \in \mathcal{G}_1 \times \mathcal{G}_2$ to $(a, b) \in \mathcal{A} \times \mathcal{B}$. To this end, we pause to build some useful high-probability events that aid the subsequent analysis. For given $(a, b) \in \mathcal{A} \times \mathcal{B}$ and $\zeta > 0$, we define the random set

(B.24)
$$\mathcal{Z}_{\boldsymbol{a},\boldsymbol{b}} = \left\{ i \in [m] : \mathcal{Q}_{\delta}(\boldsymbol{\Phi}_{i}^{\top}\boldsymbol{a} + \sqrt{m} \cdot b_{i} + \tau_{i} + t) \text{ is discontinuous in } t \in [-\zeta,\zeta] \right\}$$

and let its cardinality be $|\mathcal{Z}_{a,b}|$.

Bounding $|\mathcal{Z}_{a,b}|$ over nets: For some $\zeta \in (0, \frac{\delta}{2})$ to be chosen later, given (a, b) and conditioning on Φ , we utilize Lemma B.1 (with respect to the randomness of τ) to obtain

(B.25)
$$\mathbb{P}\left(|\mathcal{Z}_{\boldsymbol{a},\boldsymbol{b}}| \geq \frac{5m\zeta}{\delta}\right) \leq \exp\left(-\frac{m\zeta}{\delta}\right).$$

Further taking a union bound over $(a, b) \in \mathcal{G}_1 \times \mathcal{G}_2$, we obtain that the event

(B.26)
$$E_1 = \left\{ \sup_{\mathbf{a} \in \mathcal{G}_1} \sup_{\mathbf{b} \in \mathcal{G}_2} |\mathcal{Z}_{\mathbf{a},\mathbf{b}}| \le \frac{5m\zeta}{\delta} \right\}$$

holds with probability exceeding $1 - \exp\left(\mathscr{H}(\mathcal{A}, \rho_1) + \mathscr{H}(\mathcal{B}, \rho_2) - \frac{m\zeta}{\delta}\right)$. Note that for $i \notin \mathbb{Z}_{a,b}$, $\mathcal{Q}_{\delta}(\mathbf{\Phi}_i^{\top} \mathbf{a} + \sqrt{m} \cdot \mathbf{b}_i + \tau_i + t)$ is continuous in $t \in [-\zeta, \zeta]$, which along with the definition of $\mathcal{Q}_{\delta}(\cdot)$ means that $\mathcal{Q}_{\delta}(\mathbf{\Phi}_i^{\top} \mathbf{a} + \sqrt{m} \cdot \mathbf{b}_i + \tau_i + t)$ remains constant in $t \in [-\zeta, \zeta]$. In essence, E_1 bounds $|\mathcal{Z}_{a,b}|$ to be no larger than $\frac{5m\zeta}{\delta}$, universally over the nets $(\mathbf{a}, \mathbf{b}) \in \mathcal{G}_1 \times \mathcal{G}_2$, and thus indicates certain continuity of the quantization in the following sense: for measurement in $[m] \setminus \mathcal{Z}_{a,b}$ (that is the majority under small enough ζ) a perturbation smaller than ζ will not change the quantized value.

For any $(\boldsymbol{a}, \boldsymbol{b}) \in \mathcal{A} \times \mathcal{B}$ there exists $(\boldsymbol{a}', \boldsymbol{b}') \in \mathcal{G}_1 \times \mathcal{G}_2$ satisfying $\|\boldsymbol{a} - \boldsymbol{a}'\|_2 \leq \rho_1$ and $\|\boldsymbol{b} - \boldsymbol{b}'\|_2 \leq \rho_2$, and we suppose that such pair of $(\boldsymbol{a}', \boldsymbol{b}')$ has been chosen for every $(\boldsymbol{a}, \boldsymbol{b}) \in \mathcal{A} \times \mathcal{B}$ $(\boldsymbol{a}' \text{ and } \boldsymbol{b}' \text{ evidently depend on } (\boldsymbol{a}, \boldsymbol{b})$, but we omit such dependence to avoid cumbersome notation). To get uniform bound over $\mathcal{A} \times \mathcal{B}$ from a bound over nets (see (B.23)), we will need to compare $\mathcal{Q}_{\delta}(\mathbf{\Phi}_i^{\top} \mathbf{a} + \sqrt{m} \cdot b_i + \tau_i)$ with $\mathcal{Q}_{\delta}(\mathbf{\Phi}_i^{\top} \mathbf{a}' + \sqrt{m} \cdot b_i' + \tau_i)$, and we note that the former can be written as

(B.27)
$$\mathcal{Q}_{\delta} \left(\boldsymbol{\Phi}_{i}^{\top} \boldsymbol{a}' + \sqrt{m} \cdot \boldsymbol{b}_{i}' + \tau_{i} + \boldsymbol{\Phi}_{i}^{\top} (\boldsymbol{a} - \boldsymbol{a}') + \sqrt{m} \cdot (\boldsymbol{b}_{i} - \boldsymbol{b}_{i}') \right),$$

thus we need to study how the perturbations $\mathbf{\Phi}_i^{\top}(\mathbf{a}-\mathbf{a}')$ and $\sqrt{m} \cdot (b_i - b'_i)$ affect the quantized value. As explained above, on the event E_1 , for any $i \notin \mathcal{Z}_{\mathbf{a}',\mathbf{b}'}$, if $|\mathbf{\Phi}_i^{\top}(\mathbf{a}-\mathbf{a}') + \sqrt{m} \cdot (b_i - b'_i)| \leq \zeta$, then it holds that

(B.28)
$$\mathcal{Q}_{\delta}(\boldsymbol{\Phi}_{i}^{\top}\boldsymbol{a} + \sqrt{m}\cdot\boldsymbol{b}_{i} + \tau_{i}) = \mathcal{Q}_{\delta}(\boldsymbol{\Phi}_{i}^{\top}\boldsymbol{a}' + \sqrt{m}\cdot\boldsymbol{b}_{i}' + \tau_{i}).$$

In order to utilize this property, we proceed to bound the number of large perturbations $\boldsymbol{\Phi}_i^{\top}(\boldsymbol{a}-\boldsymbol{a}')$ associated with \boldsymbol{a} , and similarly large perturbations $\sqrt{m} \cdot (b_i - b_i')$ associated with \boldsymbol{b} . More precisely, given $\boldsymbol{a} \in \mathcal{A}$ and $\boldsymbol{b} \in \mathcal{B}$, we define the index sets for large perturbations as

(B.29)
$$\mathcal{J}_{\boldsymbol{a}}^{\mathcal{A}} = \left\{ i \in [m] : |\boldsymbol{\Phi}_{i}^{\top}(\boldsymbol{a} - \boldsymbol{a}')| \geq \frac{\zeta}{2} \right\},$$

(B.30)
$$\mathcal{J}_{\boldsymbol{b}}^{\mathcal{B}} = \left\{ i \in [m] : \sqrt{m} \cdot \left| b_i - b'_i \right| \ge \frac{\zeta}{2} \right\},$$

and denote their cardinalities by $|\mathcal{J}_a^{\mathcal{A}}|$ and $|\mathcal{J}_b^{\mathcal{B}}|$, respectively.

Bounding $|\mathcal{J}_{\boldsymbol{a}}^{\mathcal{A}}|$ over \mathcal{A} : For some $l \in [1, m]$ to be chosen, we apply Proposition A.3 with $\mathcal{T} = \mathcal{A}_{\text{loc}}^{(\rho_1)} := (\mathcal{A} - \mathcal{A}) \cap \mathbb{B}_2^n(\rho)$ to obtain that the event

(B.31)
$$\sup_{\boldsymbol{v}\in\mathcal{A}_{\text{loc}}^{(\rho_1)}}\max_{\substack{I\subset[m]\\|I|\leq l}} \left(\frac{1}{l}\sum_{i\in I}|\langle\boldsymbol{\Phi}_i,\boldsymbol{v}\rangle|^2\right)^{1/2} \leq \frac{\zeta}{3}$$

holds with probability at least $1 - 2\exp(C_1 l \log \frac{em}{l})$, as long as

(B.32)
$$\frac{\omega(\mathcal{A}_{\text{loc}}^{(\rho_1)})}{\sqrt{l}} + \rho_1 \sqrt{\log \frac{em}{l}} \le c_2 \zeta$$

holds with sufficiently small c_2 , as dictated by the right-hand side of (A.3).¹⁶ We suppose that we are on the event (B.31) and will choose (ρ_1, l, ζ) satisfying (B.32) later. Then, given $\boldsymbol{a} \in \mathcal{A}$ and the corresponding $\boldsymbol{a}' \in \mathcal{G}_1$ satisfying $\|\boldsymbol{a} - \boldsymbol{a}'\|_2 \leq \rho_1$, we have $\boldsymbol{a} - \boldsymbol{a}' \in \mathcal{A}_{\text{loc}}^{(\rho_1)}$ and thus (B.31) yields

(B.33)
$$\max_{\substack{I \subset [m] \\ |I| \le l}} \left(\frac{1}{l} \sum_{i \in I} \left| \langle \boldsymbol{\Phi}_i, \boldsymbol{a} - \boldsymbol{a}' \rangle \right|^2 \right)^{1/2} \le \frac{\zeta}{3}.$$

¹⁶The right-hand side of (A.3) dictates that, to ensure (B.31) holding with high probability, it suffices to have $\frac{\omega(\mathcal{A}_{loc}^{(\rho_1)})}{l^{1/2}} + \rho_1 (\log \frac{em}{l})^{1/2} \leq c_2 \zeta$ with sufficiently small C_2 .

Observe that the left-hand side of (B.33) is an upper bound on the *l*-th largest elements in $\{|\boldsymbol{\Phi}_{i}^{\top}(\boldsymbol{a}-\boldsymbol{a}')|: i=1,...,m\}$, we thus obtain $|\mathcal{J}_{\boldsymbol{a}}^{\mathcal{A}}| \leq l$. Since this argument applies to any $\boldsymbol{a} \in \mathcal{A}$ (and the corresponding $\boldsymbol{a}' \in \mathcal{G}_{1}$), (B.31) implies the event

(B.34)
$$E_2 = \left\{ \sup_{\boldsymbol{a} \in \mathcal{A}} |\mathcal{J}_{\boldsymbol{a}}^{\mathcal{A}}| \le l \right\}.$$

Bounding $|\mathcal{J}_{\boldsymbol{b}}^{\mathcal{B}}|$ over \mathcal{B} : We consider $\boldsymbol{b} \in \mathcal{B}$ and the corresponding $\boldsymbol{b}' \in \mathcal{G}_2$ satisfying $\|\boldsymbol{b} - \boldsymbol{b}'\|_2 \leq \rho_2$. Without the modulation of $\boldsymbol{\Phi}$, we will have less available information on $\{\sqrt{m} \cdot |b_i - b'_i| : i \in [m]\}$ but merely $\|\boldsymbol{b} - \boldsymbol{b}'\|_2 \leq \rho_2$. To still get a bound on $|\mathcal{J}_{\boldsymbol{b}}^{\mathcal{B}}|$, we observe that

(B.35)
$$\rho_2^2 \ge \|\boldsymbol{b} - \boldsymbol{b}'\|_2^2 \ge |\mathcal{J}_{\boldsymbol{b}}^{\mathcal{B}}| \cdot \left(\frac{\zeta}{2\sqrt{m}}\right)^2 = \frac{\zeta^2 |\mathcal{J}_{\boldsymbol{b}}^{\mathcal{B}}|}{4m},$$

which implies $|\mathcal{J}_{\boldsymbol{b}}^{\mathcal{B}}| \leq \frac{4m\rho_2^2}{\zeta^2}$. Note that this holds deterministically for all $\boldsymbol{b} \in \mathcal{B}$ (and the corresponding $\boldsymbol{b}' \in \mathcal{G}_2$), and hence the event

(B.36)
$$E_3 = \left\{ \sup_{\boldsymbol{b} \in \mathcal{B}} |\mathcal{J}_{\boldsymbol{b}}^{\mathcal{B}}| \le \frac{4m\rho_2^2}{\zeta^2} \right\}$$

holds deterministically.

Step 3: Extension to the Whole Sets Equipped with the high-probability events E_1 , E_2 and E_3 , we are in a position to strengthen the bound over $\mathcal{G}_1 \times \mathcal{G}_2$ (see (B.23)) to $(\boldsymbol{a}, \boldsymbol{b}) \in \mathcal{A} \times \mathcal{B}$. For any $(\boldsymbol{a}, \boldsymbol{b}) \in \mathcal{A} \times \mathcal{B}$, recall that we have chosen $(\boldsymbol{a}', \boldsymbol{b}') \in \mathcal{G}_1 \times \mathcal{G}_2$ satisfying $\|\boldsymbol{a} - \boldsymbol{a}'\|_2 \leq \rho_1$ and $\|\boldsymbol{b} - \boldsymbol{b}'\|_2 \leq \rho_2$, and we begin with

$$(B.37) \sup_{\substack{(c,d)\in\mathcal{E}\\(c,d)\in\mathcal{E}}} \left| \langle \boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}, \boldsymbol{\Phi}\boldsymbol{c} + \sqrt{m}\boldsymbol{d} \rangle \right| \leq \sup_{\substack{(c,d)\in\mathcal{E}\\(c,d)\in\mathcal{E}}} \left| \langle \boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}, \boldsymbol{\Phi}\boldsymbol{c} + \sqrt{m}\boldsymbol{d} \rangle \right| + \sup_{\substack{(c,d)\in\mathcal{E}\\(c,d)\in\mathcal{E}}} \left| \langle \boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}, \boldsymbol{\Phi}\boldsymbol{c} + \sqrt{m}\boldsymbol{d} \rangle \right| \\ (B.38) \leq \sup_{\substack{c\in\mathcal{C}\\\vdots=I_1}} \left| \langle \boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}, \boldsymbol{\Phi}\boldsymbol{c} \rangle \right| + \underbrace{\sqrt{m} \cdot \sup_{\boldsymbol{d}\in\mathcal{D}} \left| \langle \boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}, \boldsymbol{\Phi}\boldsymbol{c} \rangle \right|}_{:=I_2} + C\sqrt{m}\delta \cdot \left(\omega(\mathcal{E}) + t_1 \cdot \operatorname{rad}(\mathcal{E}) \right),$$

where in (B.38) we apply (B.23). We will need to separately bound I_1 and I_2 , while we discuss two kinds of measurements before proceeding.

Bad Measurements: We define for any $(a, b) \in \mathcal{A} \times \mathcal{B}$ (and the corresponding $(a', b') \in \mathcal{G}_1 \times \mathcal{G}_2$) the index set

(B.39)
$$\mathcal{U}_{\boldsymbol{a},\boldsymbol{b}} = \mathcal{Z}_{\boldsymbol{a}',\boldsymbol{b}'} \cup \mathcal{J}_{\boldsymbol{a}}^{\mathcal{A}} \cup \mathcal{J}_{\boldsymbol{b}}^{\mathcal{B}},$$

which collect the "bad" measurements that either lack certain continuity regarding the quantizer (i.e., measurements in $\mathcal{Z}_{a,b}$) or present large perturbations regarding a or b (i.e., measurements in $\mathcal{J}_a^{\mathcal{A}} \cup \mathcal{J}_b^{\mathcal{B}}$). Fortunately, the "bad" measurements are not that many, since on the events E_1 (B.26), E_2 (B.34), E_3 (B.36) we have

(B.40)
$$\sup_{\boldsymbol{a}\in\mathcal{A}}\sup_{\boldsymbol{b}\in\mathcal{B}}|\mathcal{U}_{\boldsymbol{a},\boldsymbol{b}}| \leq \sup_{\boldsymbol{a}'\in\mathcal{G}_1}\sup_{\boldsymbol{b}'\in\mathcal{G}_2}|\mathcal{Z}_{\boldsymbol{a}',\boldsymbol{b}'}| + \sup_{\boldsymbol{a}\in\mathcal{A}}|\mathcal{J}_{\boldsymbol{a}}^{\mathcal{A}}| + \sup_{\boldsymbol{b}\in\mathcal{B}}|\mathcal{J}_{\boldsymbol{b}}^{\mathcal{B}}|$$

(B.41)
$$\leq \frac{5m\zeta}{\delta} + \frac{4m\rho_2^2}{\zeta^2} + l := U_0.$$

By (B.17) $\frac{5m\zeta}{\delta} + \frac{4m\rho_2^2}{\zeta^2} \leq m$ with small enough implied constant, and we will choose ℓ in (B.77) below satisfying $\ell \leq m$ for small enough constant (see (B.78) for the value of U_0 after choosing l). Thus, by rounding that has minimal impact on our analysis, we can assume that U_0 is an integer in [1, m]. To further control the impact of bad measurements in the worst case, we note the following deterministic bound that holds for any $i \in [m]$:

(B.42)
$$|(\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}})_{i} - (\boldsymbol{\xi}_{\boldsymbol{a}',\boldsymbol{b}'})_{i}| = \left| \left[\mathcal{Q}_{\delta}(\boldsymbol{\Phi}_{i}^{\top}\boldsymbol{a} + \sqrt{m}b_{i} + \tau_{i}) - (\boldsymbol{\Phi}_{i}^{\top}\boldsymbol{a} + \sqrt{m}b_{i}) \right] - \left[\mathcal{Q}_{\delta}(\boldsymbol{\Phi}_{i}^{\top}\boldsymbol{a}' + \sqrt{m}b_{i}' + \tau_{i}) - (\boldsymbol{\Phi}_{i}^{\top}\boldsymbol{a}' + \sqrt{m}b_{i}') \right] \right|$$

(B.44)
$$= \left| \left[\mathcal{Q}_{\delta}(\boldsymbol{\Phi}_{i}^{\top}\boldsymbol{a} + \sqrt{m}b_{i} + \tau_{i}) - (\boldsymbol{\Phi}_{i}^{\top}\boldsymbol{a} + \sqrt{m}b_{i} + \tau_{i}) \right] \right|$$

(B.45)
$$-\left[\mathcal{Q}_{\delta}(\boldsymbol{\Phi}_{i}^{\top}\boldsymbol{a}'+\sqrt{m}\boldsymbol{b}_{i}'+\tau_{i})-(\boldsymbol{\Phi}_{i}^{\top}\boldsymbol{a}'+\sqrt{m}\boldsymbol{b}_{i}'+\tau_{i})\right]\right|$$

(B.46)
$$\leq 2 \cdot \sup_{x \in \mathbb{R}} |\mathcal{Q}_{\delta}(a) - a| \leq \delta$$

Benign Measurements: By contrast, measurements not in $\mathcal{U}_{a,b}$ enjoy some nice property; In particular, for $i \notin \mathcal{U}_{a,b}$ we have $i \notin \mathcal{J}_a^{\mathcal{A}} \cup \mathcal{J}_b^{\mathcal{B}}$ and hence

(B.47)
$$|\boldsymbol{\Phi}_i^{\top}(\boldsymbol{a} - \boldsymbol{a}') + \sqrt{m} \cdot (b_i - b_i')|$$

(B.48)
$$\leq |\boldsymbol{\Phi}_i^{\top}(\boldsymbol{a} - \boldsymbol{a}')| + |\sqrt{m} \cdot (b_i - b_i')| \leq \frac{\zeta}{2} + \frac{\zeta}{2} = \zeta,$$

thus (B.28) holds true, which allows us to simplify the *i*-th entry of $\xi_{a,b} - \xi_{a',b'}$ as

(B.49)
$$(\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}})_i - (\boldsymbol{\xi}_{\boldsymbol{a}',\boldsymbol{b}'})_i = \mathcal{Q}_{\delta}(\boldsymbol{\Phi}_i^{\top}\boldsymbol{a} + \sqrt{m}b_i + \tau_i) - (\boldsymbol{\Phi}_i^{\top}\boldsymbol{a} + \sqrt{m}b_i)$$

(B.50)
$$-\mathcal{Q}_{\delta}(\boldsymbol{\Phi}_{i}^{\top}\boldsymbol{a}'+\sqrt{m}\boldsymbol{b}_{i}'+\tau_{i})+(\boldsymbol{\Phi}_{i}^{\top}\boldsymbol{a}'+\sqrt{m}\boldsymbol{b}_{i}')$$

(B.51)
$$= \mathbf{\Phi}_i^{\top} (\mathbf{a}' - \mathbf{a}) + \sqrt{m} \cdot (b_i' - b_i).$$

Decomposition: According to $\mathcal{U}_{a,b}$ we can always decompose $\xi_{a,b} - \xi_{a',b'}$ into

(B.52)
$$\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}} - \boldsymbol{\xi}_{\boldsymbol{a}',\boldsymbol{b}'} = \boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(1)} + \boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(2)},$$

with $h_{a,b}^{(1)}$ and $h_{a,b}^{(2)}$ respectively accommodating the entries in $\mathcal{U}_{a,b}$ and $[m] \setminus \mathcal{U}_{a,b}$, i.e.,

(B.53)
$$(\boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(1)})_i = (\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}})_i - (\boldsymbol{\xi}_{\boldsymbol{a}',\boldsymbol{b}'})_i, \ (\boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(2)})_i = 0$$

when $i \in \mathcal{U}_{\boldsymbol{a},\boldsymbol{b}}$; otherwise, $(\boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(1)})_i = 0$ and

(B.54)
$$(\boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(2)})_i = (\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}})_i - (\boldsymbol{\xi}_{\boldsymbol{a}',\boldsymbol{b}'})_i = \boldsymbol{\Phi}_i^\top (\boldsymbol{a}' - \boldsymbol{a}) + \sqrt{m} \cdot (b_i' - b_i)$$

UNIFORM GUARANTEES FOR QUANTIZED CORRUPTED SENSING

when $i \notin \mathcal{U}_{a,b}$, with the second equality following from (B.51).

Bounding I_1 : By substituting (B.52) we can start with

(B.55)
$$I_1 \leq \sup_{\boldsymbol{c} \in \mathcal{C}} \left| \langle \boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(1)}, \boldsymbol{\Phi} \boldsymbol{c} \rangle \right| + \sup_{\boldsymbol{c} \in \mathcal{C}} \left| \langle \boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(2)}, \boldsymbol{\Phi} \boldsymbol{c} \rangle \right|.$$

Recall from (B.41) and (B.46) that $\|\boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(1)}\|_0 \leq U_0$ and $\|\boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(1)}\|_{\infty} \leq \delta$ hold uniformly for all $(\boldsymbol{a},\boldsymbol{b}) \in \mathcal{A} \times \mathcal{B}$. Thus, to bound the first term in (B.55), we can restrict our attention to entries in the support of $\boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(1)}$ and apply Cauchy-Schwarz inequality to obtain

(B.56)
$$\sup_{\boldsymbol{c}\in\mathcal{C}} \left| \langle \boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(1)}, \boldsymbol{\Phi}\boldsymbol{c} \rangle \right| \le \|\boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(1)}\|_2 \cdot \sup_{\boldsymbol{c}\in\mathcal{C}} \left(\sum_{i\in \operatorname{supp}(\boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(1)})} \left|\boldsymbol{\Phi}_i^{\top}\boldsymbol{c}\right|^2 \right)^{1/2}$$

(B.57)
$$\leq \delta \sqrt{U_0} \cdot \sup_{\boldsymbol{c} \in \mathcal{C}} \max_{\substack{I \subset [m] \\ |I| \leq U_0}} \left(\sum_{i \in I} |\boldsymbol{\Phi}_i^{\top} \boldsymbol{c}|^2 \right)^{1/2}$$

(B.58)
$$\leq C_3 \delta \sqrt{U_0} \cdot \left(\omega(\mathcal{C}) + \operatorname{rad}(\mathcal{C}) \sqrt{U_0 \log \frac{em}{U_0}} \right),$$

where (B.58) holds with probability at least $1 - 2 \exp(-C_4 U_0 \log \frac{em}{U_0})$ due to a straightforward application of Proposition A.3. Next, we seek to bound the second term in (B.55). By (B.54) we can proceed as

(B.59)
$$\sup_{\boldsymbol{c}\in\mathcal{C}} \left| \langle \boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(2)}, \boldsymbol{\Phi}\boldsymbol{c} \rangle \right| \leq \left\| \boldsymbol{\Phi}(\boldsymbol{a}'-\boldsymbol{a}) + \sqrt{m}(\boldsymbol{b}'-\boldsymbol{b}) \right\|_{2} \cdot \sup_{\boldsymbol{c}\in\mathcal{C}} \|\boldsymbol{\Phi}\boldsymbol{c}\|_{2}$$

(B.60)
$$\leq \left(\sup_{\boldsymbol{v}\in\mathcal{A}_{\text{loc}}^{(\rho_1)}} \|\boldsymbol{\Phi}\boldsymbol{v}\|_2 + \sqrt{m}\rho_2\right) \cdot \sup_{\boldsymbol{c}\in\mathcal{C}} \|\boldsymbol{\Phi}\boldsymbol{c}\|_2.$$

Now we apply Proposition A.3 to achieve the following two bounds (for some absolute constants C_5, C_6):¹⁷

• Proposition A.3 with $\mathcal{T} = \mathcal{A}_{loc}^{(\rho_1)}$ and l = m yields that the event

(B.61)
$$\sup_{\boldsymbol{v}\in\mathcal{A}_{\text{loc}}^{(\rho_1)}} \|\boldsymbol{\Phi}\boldsymbol{v}\|_2 \le C_5 \Big(\omega\big(\mathcal{A}_{\text{loc}}^{(\rho_1)}\big) + \sqrt{m} \cdot \rho_1\Big)$$

holds with probability exceeding $1 - 2 \exp(-C_6 m)$.

• Proposition A.3 with $\mathcal{T} = \mathcal{C}$ and l = m yields that the event

(B.62)
$$\sup_{\boldsymbol{c}\in\mathcal{C}} \|\boldsymbol{\Phi}\boldsymbol{c}\|_2 \le C_5 \Big(\omega(\mathcal{C}) + \sqrt{m}\cdot \operatorname{rad}(\mathcal{C})\Big)$$

holds with probability exceeding $1 - 2 \exp(-C_6 m)$.

¹⁷Alternatively, one can achieve this by using matrix deviation inequality; see Proposition A.1 with $\mathcal{T} = \mathcal{T}_0 \times \{0\}$ for some $\mathcal{T}_0 \subset \mathbb{R}^n$.

Substituting (B.61) and (B.62) into (B.60) we obtain

(B.63)
$$\sup_{\boldsymbol{c}\in\mathcal{C}} \left| \langle \boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(2)}, \boldsymbol{\Phi}\boldsymbol{c} \rangle \right| \leq C_7 \big(\omega(\mathcal{C}) + \sqrt{m} \cdot \operatorname{rad}(\mathcal{C}) \big) \Big(\omega \big(\mathcal{A}_{\operatorname{loc}}^{(\rho_1)} \big) + \sqrt{m}\rho_1 + \sqrt{m}\rho_2 \Big).$$

Note that all arguments in bounding I_1 hold universally for all $(a, b) \in \mathcal{A} \times \mathcal{B}$. Thus, combining (B.55), (B.58) and (B.63) immediately yields

(B.64)

$$\sup_{\boldsymbol{a}\in\mathcal{A}}\sup_{\boldsymbol{b}\in\mathcal{B}}I_{1} \leq C_{3}\delta\sqrt{U_{0}}\cdot\left(\omega(\mathcal{C})+\operatorname{rad}(\mathcal{C})\sqrt{U_{0}\log\frac{em}{U_{0}}}\right) + C_{7}\left(\omega(\mathcal{C})+\sqrt{m}\cdot\operatorname{rad}(\mathcal{C})\right)\left(\omega\left(\mathcal{A}_{\operatorname{loc}}^{(\rho_{1})}\right)+\sqrt{m}\rho_{1}+\sqrt{m}\rho_{2}\right)$$

where U_0 is given in (B.41).

Bounding I_2 : By substituting (B.52), we proceed as

(B.65)
$$I_2 \leq \sqrt{m} \cdot \operatorname{rad}(\mathcal{D}) \cdot \|\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}} - \boldsymbol{\xi}_{\boldsymbol{a}',\boldsymbol{b}'}\|_2$$

(B.66)
$$\leq \sqrt{m} \cdot \operatorname{rad}(\mathcal{D}) \cdot \left(\|\boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(1)}\|_{2} + \|\boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(2)}\|_{2} \right)$$

Recall from (B.41) and (B.46) that $\|\boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(1)}\|_0 \leq U_0$ and $\|\boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(1)}\|_{\infty} \leq \delta$ hold uniformly for all $(\boldsymbol{a},\boldsymbol{b}) \in \mathcal{A} \times \mathcal{B}$, and hence we have $\|\boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(1)}\|_2 \leq \delta \sqrt{U_0}$. Then, (B.54) gives

(B.67)
$$\|\boldsymbol{h}_{\boldsymbol{a},\boldsymbol{b}}^{(2)}\|_{2} \leq \|\boldsymbol{\Phi}(\boldsymbol{a}-\boldsymbol{a}')\|_{2} + \sqrt{m}\|\boldsymbol{b}-\boldsymbol{b}'\|_{2}$$

(B.68)
$$\leq \sup_{\boldsymbol{v} \in \mathcal{A}_{\text{loc}}^{(\rho_1)}} \|\boldsymbol{\Phi}\boldsymbol{v}\|_2 + \sqrt{m} \cdot \rho_2$$

(B.69)
$$\leq C_5 \Big(\omega \big(\mathcal{A}_{\text{loc}}^{(\rho_1)} \big) + \sqrt{m} \rho_1 + \sqrt{m} \rho_2 \Big),$$

where we use (B.61) in (B.69). We note that all arguments in bounding I_2 hold universally for all $(\boldsymbol{a}, \boldsymbol{b}) \in \mathcal{A} \times \mathcal{B}$. Substituting the bounds $\|\boldsymbol{h}_{\boldsymbol{a}, \boldsymbol{b}}^{(1)}\|_2 \leq \delta \sqrt{U_0}$ and (B.69) into (B.66), we thus obtain

(B.70)
$$\sup_{\boldsymbol{a}\in\mathcal{A}}\sup_{\boldsymbol{b}\in\mathcal{B}}I_2 \leq \sqrt{m}\cdot \operatorname{rad}(\mathcal{D})\Big(\delta\sqrt{U_0} + C_5\Big(\omega\big(\mathcal{A}_{\operatorname{loc}}^{(\rho_1)}\big) + \sqrt{m}\rho_1 + \sqrt{m}\rho_2\Big)\Big)$$

where U_0 is given in (B.41).

Step 4: Combining and Parameters Selection We are in a position to combine everything together. Taking supremum over $(\boldsymbol{a}, \boldsymbol{b}) \in \mathcal{A} \times \mathcal{B}$ in (B.38) and then substituting (B.64) and (B.70), we obtain that as long as $\rho_1 > 0, l \in [1, m], \zeta \in (0, \frac{\delta}{2})$ are chosen such that (B.32) holds, then the event

(B.71) $\sup_{\boldsymbol{a}\in\mathcal{A}} \sup_{\boldsymbol{b}\in\mathcal{B}} \sup_{(\boldsymbol{c},\boldsymbol{d})\in\mathcal{E}} \left| \langle \boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}}, \boldsymbol{\Phi}\boldsymbol{c} + \sqrt{m}\boldsymbol{d} \rangle \right|$

$$(B.72) \lesssim \sqrt{m\delta} \cdot \left(\omega(\mathcal{E}) + t_1 \cdot \operatorname{rad}(\mathcal{E})\right) + \delta \sqrt{U_0} \left(\omega(\mathcal{C}) + \sqrt{U_0 \log \frac{em}{U_0}} \cdot \operatorname{rad}(\mathcal{C}) + \sqrt{m} \cdot \operatorname{rad}(\mathcal{D})\right)$$

$$(B.73) + \left(\omega \left(\mathcal{A}_{\operatorname{loc}}^{(\rho_1)}\right) + \sqrt{m\rho_1} + \sqrt{m\rho_2}\right) \left(\omega(\mathcal{C}) + \sqrt{m} \cdot \operatorname{rad}(\mathcal{C}) + \sqrt{m} \cdot \operatorname{rad}(\mathcal{D})\right)$$

holds with probability exceeding

(B.74)
$$1 - 2\exp\left(\mathscr{H}(\mathcal{A},\rho_1) + \mathscr{H}(\mathcal{B},\rho_2) - t_1^2\right) - \exp\left(\mathscr{H}(\mathcal{A},\rho_1) + \mathscr{H}(\mathcal{B},\rho_2) - \frac{m_{\zeta}}{\delta}\right)$$

(B.75)
$$-2\exp\left(-C_1 l \log \frac{cm}{l}\right) - 2\exp\left(-C_4 U_0 \log \frac{cm}{U_0}\right) - 4\exp(-C_6 m),$$

where the terms in (B.74) stem from (B.23) and the event E_1 (B.26), the terms in (B.75) are from (B.31), (B.58), and (B.61)–(B.62).

Choosing Parameters: We specify the parameter whose (near) optimal choice is clear at this stage, while we still leave other parameters generic since their optimal values may depend on $(\mathcal{A}, \mathcal{B}, \mathcal{E})$. Specifically, we set (we suppose that l below is chosen as an integer in [1, m] without loss of generality, since we can we just round otherwise)

(B.76)
$$t_1 = 2\sqrt{\mathscr{H}(\mathcal{A},\rho_1)} + 2\sqrt{\mathscr{H}(\mathcal{B},\rho_2)},$$

(B.77)
$$l = \frac{m\zeta}{\delta} + \frac{C_2 \cdot \omega^2(\mathcal{A}_{\text{loc}}^{(p_1)})}{\zeta^2}$$

with sufficiently large C_2 such that $\frac{\omega(\mathcal{A}_{loc}^{(\rho_1)})}{\sqrt{l}} \leq \frac{c_2\zeta}{2}$. We show that our choice (B.77) satisfies (B.32) that is needed to ensure (B.31). Specifically, $l \geq \frac{C_2 \cdot \omega^2(\mathcal{A}_{loc}^{(\rho_1)})}{\zeta^2}$ implies $\frac{\omega(\mathcal{A}_{loc}^{(\rho_1)})}{\sqrt{l}} \lesssim \zeta$ with small enough implied constant, and $l \geq \frac{m\zeta}{\delta}$ along with $\zeta \lesssim \delta$ from (B.17) implies $\rho_1 \sqrt{\log \frac{em}{l}} \leq \rho_1 \sqrt{\log \frac{e\delta}{\zeta}} \lesssim \zeta$ with small enough implied constant. We recall the value of U_0 given in (B.41), which together with (B.77) reads as

(B.78)
$$U_0 = \frac{6m\zeta}{\delta} + \frac{4m\rho_2^2}{\zeta^2} + \frac{C_2 \cdot \omega^2(\mathcal{A}_{\text{loc}}^{(\rho_1)})}{\zeta^2}.$$

Simplifying (B.74)–(B.75): Under (B.76) and (B.18) that implies $\frac{m\zeta}{\delta} \geq 2\mathscr{H}(\mathcal{A}, \rho_1) + 2\mathscr{H}(\mathcal{B}, \rho_2)$, we can relax the probability terms in (B.74) to

(B.79)
$$2\exp\left(\mathscr{H}(\mathcal{A},\rho_1) + \mathscr{H}(\mathcal{B},\rho_2) - t_1^2\right) + \exp\left(\mathscr{H}(\mathcal{A},\rho_1) + \mathscr{H}(\mathcal{B},\rho_2) - \frac{m\zeta}{\delta}\right)$$

(B.80)
$$\leq 3 \exp\left(-\mathcal{H}(\mathcal{A},\rho_1) - \mathcal{H}(\mathcal{B},\rho_2)\right)$$

Besides, since (B.41) and (B.77) give $U_0 \ge l \ge \frac{m\zeta}{\delta}$, so we have

$$(B.81) \quad 2\exp(-C_1 l\log\frac{em}{l}) + 2\exp(-C_4 U_0\log\frac{em}{U_0}) \le 4\exp(-c_8 \mathscr{H}(\mathcal{A},\rho_1) - c_8 \mathscr{H}(\mathcal{B},\rho_2))$$

for some absolute constant $c_8 > 0$. Moreover, (B.18) and $\zeta \leq \frac{\delta}{2}$ imply $m \geq 2 \mathscr{H}(\mathcal{A}, \rho_1) + 2 \mathscr{H}(\mathcal{B}, \rho_2)$, and so we have

(B.82)
$$4\exp(-C_6m) \le 4\exp(-c_8 \mathscr{H}(\mathcal{A},\rho_1) - c_8 \mathscr{H}(\mathcal{B},\rho_2))$$

This manuscript is for review purposes only.

provided that c_8 is chosen sufficiently small. Overall, from (B.74)-(B.75) we can promise that (B.71)-(B.73) holds with probability exceeding

(B.83)
$$1 - 12 \exp(-c_9 \mathscr{H}(\mathcal{A}, \rho_1) - c_9 \mathscr{H}(\mathcal{B}, \rho_2))$$

with some $c_9 > 0$.

Simplifying (B.71)–(B.73): We enforce some typical scaling such that the terms in (B.73) is dominated by those in (B.72), up to multiplicative factors. For clarity, we collect the developments as follows:

- From (B.78) and $\zeta \in (0, \frac{\delta}{2})$ we have $\delta \sqrt{U_0} \gtrsim \frac{\delta}{\zeta} \omega(\mathcal{A}_{\text{loc}}^{(\rho_1)}) \gtrsim \omega(\mathcal{A}_{\text{loc}}^{(\rho_1)})$, hence the terms $\omega(\mathcal{A}_{\text{loc}}^{(\rho_1)}) \cdot (\omega(\mathcal{C}) + \sqrt{m} \cdot \text{rad}(\mathcal{D})) \text{ from (B.73) are dominated by } \delta\sqrt{U_0} \cdot (\omega(\mathcal{C}) + \sqrt{m} \cdot \text{rad}(\mathcal{D})) \text{ from (B.72);}$
- Note that (B.17) provides $\rho_1 + \rho_2 \lesssim \zeta$ with small enough implied constant. Then, from (B.78) and $\zeta \in (0, \frac{\delta}{2})$ we have $\delta \sqrt{U_0} \geq \sqrt{m\zeta\delta} \geq \sqrt{m\zeta} \gtrsim \sqrt{m(\rho_1 + \rho_2)}$, and hence the terms $\sqrt{m}(\rho_1 + \rho_2) \cdot (\omega(\mathcal{C}) + \sqrt{m} \cdot \operatorname{rad}(\mathcal{D}))$ from (B.73) are dominated by $\delta \sqrt{U_0} \cdot (\omega(\mathcal{C}) + \sqrt{m} \cdot \operatorname{rad}(\mathcal{D}))$ from (B.72);
- We further show that the remaining terms in (B.73), namely $(\sqrt{m} \cdot \omega(\mathcal{A}_{\text{loc}}^{(\rho_1)}) + m(\rho_1 + \omega))$ (ρ_2)) · rad(\mathcal{C}), are dominated by $\delta U_0 \sqrt{\log \frac{em}{U_0}}$ · rad(\mathcal{C}) from (B.72). It suffices to show

(B.84)
$$\sqrt{m} \cdot \omega(\mathcal{A}_{\text{loc}}^{(\rho_1)}) + m(\rho_1 + \rho_2) \lesssim \delta U_0.$$

By (B.78) we have $\delta U_0 \ge 6m\zeta$. On the other hand, (B.17) and (B.18) evidently imply $\sqrt{m} \cdot \omega(\mathcal{A}_{\text{loc}}^{(\rho_1)}) + m(\rho_1 + \rho_2) \lesssim m\zeta.$ (B.84) hence follows.

By the above discussions, under the scaling conditions stated in our theorem statement, the terms in (B.72) dominate the ones in (B.73). Further substituting t_1 in (B.76) yields the simplified bound:

 $\sup_{\boldsymbol{a}\in\mathcal{A}}\sup_{\boldsymbol{b}\in\mathcal{B}}\sup_{(\boldsymbol{c},\boldsymbol{d})\in\mathcal{E}}\left|\langle\boldsymbol{\xi}_{\boldsymbol{a},\boldsymbol{b}},\boldsymbol{\Phi}\boldsymbol{c}+\sqrt{m}\boldsymbol{d}\rangle\right|$ (B.85)

(B.86)
$$\leq \sqrt{m\delta} \cdot \omega(\mathcal{E}) + \sqrt{m\delta} \cdot \operatorname{rad}(\mathcal{E}) \cdot \left(\sqrt{\mathscr{H}(\mathcal{A},\rho_1)} + \sqrt{\mathscr{H}(\mathcal{B},\rho_2)}\right)$$

(B.87)
$$+ \delta \sqrt{U_0} \Big(\omega(\mathcal{C}) + \sqrt{U_0 \log \frac{em}{U_0}} \cdot \operatorname{rad}(\mathcal{C}) + \sqrt{m} \cdot \operatorname{rad}(\mathcal{D}) \Big).$$

We denote U_0 by \hat{U} as in the theorem statement (B.21). To complete the proof, it remains to make some final simplification:

- By (B.78) we have $\log \frac{em}{U_0} \lesssim \log \frac{\delta}{\zeta}$ and hence we can relax $\delta U_0 \sqrt{\log \frac{em}{U_0}} \cdot \operatorname{rad}(\mathcal{C})$ to the term $\delta \hat{U} \sqrt{\log \frac{\delta}{\zeta}} \cdot \operatorname{rad}(\mathcal{C})$ in (B.20);
- By $U_0 \leq m$ we have $\delta \sqrt{U_0} \cdot \omega(\mathcal{C}) \leq \delta \sqrt{m} \cdot \omega(\mathcal{E})$, and we can only retain $\delta \sqrt{m} \cdot \omega(\mathcal{E})$ as in (B.19):
- By $\operatorname{rad}(\mathcal{E}) \leq \operatorname{rad}(\mathcal{C}) + \operatorname{rad}(\mathcal{D})$ we can bound $\sqrt{m\delta} \cdot \operatorname{rad}(\mathcal{E}) \cdot (\sqrt{\mathscr{H}(\mathcal{A},\rho_1)} + \sqrt{\mathscr{H}(\mathcal{B},\rho_2)})$ by $\sqrt{m\delta} \cdot \operatorname{rad}(\mathcal{C}) \cdot (\sqrt{\mathscr{H}(\mathcal{A},\rho_1)} + \sqrt{\mathscr{H}(\mathcal{B},\rho_2)}) + \sqrt{m\delta} \cdot \operatorname{rad}(\mathcal{D}) \cdot (\sqrt{\mathscr{H}(\mathcal{A},\rho_1)} + \sqrt{\mathscr{H}(\mathcal{B},\rho_2)}).$ Observe that $\sqrt{m\delta} \cdot \operatorname{rad}(\mathcal{D}) \cdot (\sqrt{\mathscr{H}(\mathcal{A}, \rho_1)} + \sqrt{\mathscr{H}(\mathcal{B}, \rho_2)})$ is dominated by $\delta \sqrt{m\hat{U}} \cdot \operatorname{rad}(\mathcal{D})$

UNIFORM GUARANTEES FOR QUANTIZED CORRUPTED SENSING

(due to (B.21) and (B.18)), we can thus simply retain $\sqrt{m\delta} \cdot \operatorname{rad}(\mathcal{C}) \cdot (\sqrt{\mathscr{H}(\mathcal{A},\rho_1)} + \sqrt{\mathscr{H}(\mathcal{B},\rho_2)})$ as in (B.20).

The proof is now complete.

B.3 Global QPE for Structured Sets We consider the setting where \mathcal{A} and \mathcal{B} in Theorem B.3 are structured sets with Kolmogorov entropy depending on the covering radius in a logarithmic manner; see Definition 3.1. With properly chosen parameters (ζ , ρ_1 , ρ_2), Theorem B.3 specializes to the following. We will explain in Remark B.5 that the QPE below is sufficient for the proofs of our main theorems (Theorem 3.2, 3.12, 3.16).

Corollary B.4 (Global QPE for Structured Sets). Given some bounded sets $\mathcal{A} \subset \mathbb{R}^n$, $\mathcal{B} \subset \mathbb{R}^m$, $\mathcal{E} \subset \mathbb{R}^{n+m}$ and some $\delta > 0$, we assume that the sub-Gaussian matrix $\mathbf{\Phi} \in \mathbb{R}^{m \times n}$ and the random dither $\boldsymbol{\tau} \sim \mathscr{U}[-\frac{\delta}{2}, \frac{\delta}{2}]^m$ are as described in Assumption 1. Suppose that (ζ, ρ_1, ρ_2) are positive scalars satisfying

(B.88)
$$\zeta = \frac{4\delta(\mathscr{H}(\mathcal{A},\rho_1) + \mathscr{H}(\mathcal{B},\rho_2))}{m}$$

(B.89)
$$\rho_1 \le \frac{c_1 \zeta}{(\log \frac{\delta}{\zeta})^{1/2}}, \ \omega \left(\mathcal{A}_{\text{loc}}^{(\rho_1)} \right) \le c_2 \zeta \sqrt{\frac{m\zeta}{\delta}}, \ \rho_2 \le c_3 \zeta \sqrt{\frac{\zeta}{\delta}}$$

for some sufficiently small absolute constants (c_1, c_2, c_3) , and suppose that

(B.90)
$$m \ge C_4 \left(\mathscr{H}(\mathcal{A}, \rho_1) + \mathscr{H}(\mathcal{B}, \rho_2) \right)$$

for large enough C_4 . Then, with the quantization noise $\xi_{a,b}$ being given in (B.4), for some absolute constant C_5 the event

(B.91) $\sup_{\boldsymbol{a} \in \mathcal{A}} \sup_{\boldsymbol{b} \in \mathcal{B}} \sup_{(\boldsymbol{c}, \boldsymbol{d}) \in \mathcal{E}} \left| \langle \boldsymbol{\xi}_{\boldsymbol{a}, \boldsymbol{b}}, \boldsymbol{\Phi} \boldsymbol{c} + \sqrt{m} \boldsymbol{d} \rangle \right|$

(B.92)
$$\leq C_5 \delta \sqrt{m} \Big(\omega(\mathcal{E}) + \operatorname{rad}(\mathcal{E}) \sqrt{\mathscr{H}(\mathcal{A}, \rho_1) + \mathscr{H}(\mathcal{B}, \rho_2)} \Big)$$

holds with probability exceeding $1 - 12 \exp(-c_6 \mathscr{H}(\mathcal{A}, \rho_1) - c_6 \mathscr{H}(\mathcal{B}, \rho_2))$ on a single draw of (Φ, τ) .

Proof. We prove the statement using the general global QPE property presented in Theorem B.3. Given \mathcal{E} , recall that \mathcal{C} and \mathcal{D} are defined in (B.15) and (B.16).

Verifying (B.17)–(B.18) First, we verify (B.17). Under (B.90), ζ in (B.88) evidently satisfies $\zeta \leq c_1 \delta$ with small enough c_1 , verifying the first condition in (B.17). Then, (B.89) gives $\rho_2 \leq c_3 \zeta \sqrt{\frac{\zeta}{\delta}} \leq c_3 \zeta$, which provides the third condition in (B.17). Combining with $\rho_1 \leq \frac{c_1 \zeta}{(\log \frac{\zeta}{\delta})^{1/2}}$, we know that the scaling conditions in (B.17) are satisfied by (B.88)–(B.89). Next, we verify (B.18). By substituting (B.88) we find that it suffices to verify $m \geq \frac{C_2}{\zeta^2} \omega^2 (\mathcal{A}_{\text{loc}}^{(\rho_1)})$ for some large enough C_2 , and note that this is guaranteed by the second condition in (B.89) that provides $m \geq \frac{\omega^2(\mathcal{A}_{\text{loc}}^{(\rho_1)})}{c_2^2 \zeta^2} \cdot \frac{\delta}{\zeta}$ for small enough c_2 (since $\delta \geq \zeta$).

This manuscript is for review purposes only.

Simplifying (B.19)–(B.20) Because the last two conditions in (B.89) imply

(B.93)
$$\frac{m\rho_2^2}{\zeta^2} + \frac{\omega^2(\mathcal{A}_{\text{loc}}^{(\rho_1)})}{\zeta^2} \le (c_2^2 + c_3)\frac{m\zeta}{\delta},$$

 \hat{U} given in (B.21) simplifies to $\hat{U} \simeq \frac{m\zeta}{\delta} = 4(\mathscr{H}(\mathcal{A}, \rho_1) + \mathscr{H}(\mathcal{B}, \rho_2))$, with the equality following from (B.88). Thus, we have

(B.94)
$$\frac{\hat{U}}{\sqrt{m}} \left[\log \frac{\delta}{\zeta} \right]^{1/2} \lesssim \frac{\mathscr{H}(\mathcal{A}, \rho_1) + \mathscr{H}(\mathcal{B}, \rho_2)}{\sqrt{m}} \sqrt{\log \left(\frac{m}{\mathscr{H}(\mathcal{A}, \rho_1) + \mathscr{H}(\mathcal{B}, \rho_2)} \right)}$$

(B.95)
$$\lesssim \sqrt{\mathscr{H}(\mathcal{A},\rho_1)} + \sqrt{\mathscr{H}(\mathcal{B},\rho_2)},$$

where the second inequality follows from (B.90). Therefore, the bound in (B.19)-(B.20) simplifies to

(B.96)
$$O\left(\delta\sqrt{m}\cdot\omega(\mathcal{E})+\delta\sqrt{m\hat{U}}\cdot\mathrm{rad}(\mathcal{D})+\delta\sqrt{m}\cdot\mathrm{rad}(\mathcal{C})\sqrt{\mathscr{H}(\mathcal{A},\rho_1)+\mathscr{H}(\mathcal{B},\rho_2)}\right)$$

(B.97)
$$= O\Big(\delta\sqrt{m}\big(\omega(\mathcal{E}) + \operatorname{rad}(\mathcal{E})\sqrt{\mathscr{H}(\mathcal{A},\rho_1) + \mathscr{H}(\mathcal{B},\rho_2)}\big)\Big),$$

where in the second line we use $\hat{U} \leq \mathscr{H}(\mathcal{A}, \rho_1) + \mathscr{H}(\mathcal{B}, \rho_2)$ and $\max\{\operatorname{rad}(\mathcal{C}), \operatorname{rad}(\mathcal{D})\} \leq \operatorname{rad}(\mathcal{E})$. We have arrived at the desired bound in Corollary B.4, and note that the promised probability directly follows from Theorem B.3. The proof is complete.

Remark B.5. Corollary B.4 is tailored to fit the case where \mathcal{A} and \mathcal{B} are structured sets as per Definition 3.1, but more generally put, it works well for \mathcal{A}, \mathcal{B} with Kolmogorov entropy logarithmically depending on the covering radius. By Proposition A.9(a), this is also the case when \mathcal{A} and \mathcal{B} are the ranges of some Lipschitz generative models (as per Assumption 5), thus Corollary B.4 applies to the analysis of generative prior. Therefore, Corollary B.4 is a version of QPE sufficient for proving our main theorems, and we will further present other implications of Theorem B.3 for the case where \mathcal{A}, \mathcal{B} are arbitrary sets in Appendix D.1.

Appendix C. Deferred Proofs. We collect the proofs of Corollary 3.5 and Corollary 3.7 (concrete outcomes of Theorem 3.2), Corollary 3.14 and Corollary 3.15 (concrete outcomes of Theorem 3.12), and Theorem 3.16 (for generative prior) in this appendix.

In the first two proofs we will invoke Theorem 3.2, with the two major steps being: (i) bounding the geometric complexity quantities $\gamma(\mathcal{D}_{\boldsymbol{x}}^*)$ and $\gamma(\mathcal{D}_{\boldsymbol{v}}^*)$; (ii) selecting (ρ_1, ρ_2) to render (3.12)–(3.13).

C.1 The Proof of Corollary 3.5 (Recovering Sparse Signal and Sparse Corruption via Constrained Lasso)

Proof.

Step 1: Bounding $\gamma(\mathcal{D}_{\boldsymbol{x}}^*)$ and $\gamma(\mathcal{D}_{\boldsymbol{v}}^*)$ For any $\boldsymbol{v} \in \mathcal{D}_{\boldsymbol{x}}$, there exists some *s*-sparse \boldsymbol{x} such that $\|\boldsymbol{x} + t\boldsymbol{v}\|_1 \leq \|\boldsymbol{x}\|_1$ holds for some t > 0. We let $\mathcal{S} = \operatorname{supp}(\boldsymbol{x})$. Because $\mathcal{S} \subset [n]$ and $|\mathcal{S}| \leq s$, we have (given $\boldsymbol{v} \in \mathbb{R}^n$ and $\mathcal{S} \subset [n]$ we obtain $\boldsymbol{v}_{\mathcal{S}}$ from \boldsymbol{v} by only retaining entries in \mathcal{S} while setting others zero)

(C.1)
$$\|\boldsymbol{x}\|_{1} \ge \|\boldsymbol{x} + t\boldsymbol{v}_{\mathcal{S}^{c}}\|_{1} = \|\boldsymbol{x} + t\boldsymbol{v}_{\mathcal{S}}\|_{1} + t\|\boldsymbol{v}_{\mathcal{S}^{c}}\|_{1} \ge \|\boldsymbol{x}\|_{1} - t\|\boldsymbol{v}_{\mathcal{S}}\|_{1} + t\|\boldsymbol{v}_{\mathcal{S}^{c}}\|_{1},$$

which provides $\|\boldsymbol{v}_{\mathcal{S}^c}\|_1 \leq \|\boldsymbol{v}_{\mathcal{S}}\|_1$. Thus, we obtain

(C.2)
$$\|v\|_1 \le 2\|v_{\mathcal{S}}\|_1 \le 2\sqrt{s}\|v_{\mathcal{S}}\|_2 \le 2\sqrt{s}\|v\|_2.$$

Because $\mathcal{D}_{\boldsymbol{x}}^* = \mathcal{D}_{\boldsymbol{x}} \cap \mathbb{S}^{n-1}$, we have $\mathcal{D}_{\boldsymbol{x}}^* \subset \mathbb{B}_2^n \cap \mathbb{B}_1^n(2\sqrt{s})$, hence Proposition A.5 gives $\omega(\mathcal{D}_{\boldsymbol{x}}^*) \lesssim \sqrt{s \log \frac{en}{s}}$. By (2.8), this also bounds $\gamma(\mathcal{D}_{\boldsymbol{x}}^*)$ (up to multiplicative constant). Similarly, we have $\gamma(\mathcal{D}_{\boldsymbol{v}}^*) \lesssim \sqrt{k \log \frac{em}{k}}$.

Step 2: Selecting (ρ_1, ρ_2) Recall that $\zeta = \frac{4\delta}{m}(\mathscr{H}(\mathcal{K}_x, \rho_1) + \mathscr{H}(\mathcal{K}_v, \rho_2))$ as per (3.12), and we claim that setting

(C.3)
$$\rho_1 = c\delta\left(\frac{s}{m}\right)^{3/2}, \ \rho_2 = c\delta\left(\frac{k}{m}\right)^{3/2}$$

with small enough c satisfies (3.13). The reasoning is as follows:

- In view of Proposition A.6, $m \gtrsim s \log(\frac{nm^{3/2}}{s^{5/2}\delta}) + k \log(\frac{m^{5/2}}{k^{5/2}\delta})$ in Corollary 3.5 implies (3.14) needed in Theorem 3.2;
- Verifying $\rho_1 \lesssim \frac{\zeta}{(\log \frac{\delta}{\zeta})^{1/2}}$, $\rho_2 \lesssim \zeta \sqrt{\frac{\zeta}{\delta}}$: By (3.14) we have $m \gtrsim \mathscr{H}(\mathcal{K}_{\boldsymbol{x}}, \rho_1) + \mathscr{H}(\mathcal{K}_{\boldsymbol{v}}, \rho_2)$ that implies $\zeta \lesssim \delta$. Substituting $\zeta = \frac{4\delta}{m} (\mathscr{H}(\mathcal{K}_{\boldsymbol{x}}, \rho_1) + \mathscr{H}(\mathcal{K}_{\boldsymbol{v}}, \rho_2))$ finds

$$\rho_1 + \rho_2 \lesssim \delta\left(\frac{s}{m}\right)^{3/2} + \delta\left(\frac{k}{m}\right)^{3/2} \lesssim \zeta \sqrt{\frac{\zeta}{\delta}}.$$

Because $\frac{\delta}{\zeta} \ge C_1$ holds for some large C_1 , and so $\rho_1 + \rho_2 \le c_2 \zeta \sqrt{\frac{\zeta}{\delta}}$ with small enough c_2 suffices to ensure $\rho_1 \lesssim \frac{\zeta}{(\log \frac{\delta}{\zeta})^{1/2}}$ and $\rho_2 \lesssim \zeta \sqrt{\frac{\zeta}{\delta}}$.

• Verifying $\omega(\mathcal{A}_{\text{loc}}^{(\rho_1)}) \lesssim \zeta \sqrt{\frac{m\zeta}{\delta}}$: Observe that $\mathcal{A}_{\text{loc}}^{(\rho_1)} \subset \Sigma_{2s}^n \cap \rho_1 \mathbb{B}_2^n$, and so $\omega(\mathcal{A}_{\text{loc}}^{(\rho_1)}) \lesssim \rho_1 s \log \frac{en}{s}$. Further, by $\rho_1 \lesssim \zeta$ and $s \log \frac{em}{s} \lesssim \sqrt{\mathscr{H}(\mathcal{A}, \rho_1)} \lesssim \sqrt{\frac{m\zeta}{\delta}}$ we arrive at $\omega(\mathcal{A}_{\text{loc}}^{(\rho_1)}) \lesssim \zeta \sqrt{\frac{m\zeta}{\delta}}$.

Note that we have derived explicit bounds on the geometric quantities and chosen ρ_1, ρ_2 to satisfy the conditions in Theorem 3.2. Now we can simply invoke Theorem 3.2 to prove the desired claim.

C.2 The Proof of Corollary 3.7 (Recovering Low-Rank Signal and Sparse Corruption via Constrained Lasso)

Proof.

Step 1: Bounding $\gamma(\mathcal{D}_{\boldsymbol{x}}^*)$ and $\gamma(\mathcal{D}_{\boldsymbol{v}}^*)$ [34, Coro. 2.1] gives $\gamma(\mathcal{D}_{\boldsymbol{x}}^*) \lesssim \sqrt{r(p+q)}$. As shown in the proof of Corollary 3.5, we have $\gamma(\mathcal{D}_{\boldsymbol{v}}^*) \lesssim \sqrt{k \log \frac{em}{k}}$.

Step 2: Selecting (ρ_1, ρ_2) This is similar to Step 2 in the proof of Corollary 3.5. We recall $\zeta = \frac{4\delta}{m}(\mathscr{H}(\mathcal{K}_{\boldsymbol{x}}, \rho_1) + \mathscr{H}(\mathcal{K}_{\boldsymbol{v}}, \rho_2))$ as per (3.12) and claim that the choice

(C.4)
$$\rho_1 = c\delta \left(\frac{r(p+q)}{m}\right)^{3/2}, \ \rho_2 = c\delta \left(\frac{k}{m}\right)^{3/2}$$

with small enough c satisfies the (3.12)–(3.13) in Theorem 3.2. We note the following dot points to explain this:

- In view of Proposition A.6, $m \gtrsim r(p+q)\log(\frac{m^{3/2}}{\delta(r(p+q))^{3/2}}) + k\log(\frac{m^{5/2}}{k^{5/2}\delta})$ in Corollary 3.7 implies (3.14) needed in Theorem 3.2;
- $\rho_1 \lesssim \frac{\zeta}{(\log \frac{\delta}{\zeta})^{1/2}}$, $\rho_2 \lesssim \zeta \sqrt{\frac{\zeta}{\delta}}$: (3.14) implies $m \gtrsim \mathscr{H}(\mathcal{K}_{\boldsymbol{x}}, \rho_1) + \mathscr{H}(\mathcal{K}_{\boldsymbol{v}}, \rho_2)$ and hence $\zeta \lesssim \delta$. By $\zeta = \frac{4\delta}{m}(\mathscr{H}(\mathcal{K}_{\boldsymbol{x}}, \rho_1) + \mathscr{H}(\mathcal{K}_{\boldsymbol{v}}, \rho_2))$ it is easy to verify

$$\rho_1 + \rho_2 \lesssim \delta \left(\frac{r(p+q)}{m} \right)^{3/2} + \delta \left(\frac{k}{m} \right)^{3/2} \lesssim \zeta \sqrt{\frac{\zeta}{\delta}}.$$

Because $\frac{\delta}{\zeta} \geq C_1$ holds for some large C_1 , $\rho_1 + \rho_2 \lesssim \zeta \sqrt{\frac{\zeta}{\delta}}$ ensures $\rho_1 \lesssim \frac{\zeta}{(\log \frac{\delta}{\zeta})^{1/2}}$ and $\rho_2 \lesssim \zeta \sqrt{\frac{\zeta}{\delta}}$.

• $\omega(\mathcal{A}_{\text{loc}}^{(\rho_1)}) \lesssim \zeta \sqrt{\frac{m\zeta}{\delta}}$: Observe that $\mathcal{A}_{\text{loc}}^{(\rho_1)} \subset M_{2r}^{p,q} \cap \mathbb{B}_{\mathrm{F}}^{p,q}(\rho_1)$, and so $\omega(\mathcal{A}_{\text{loc}}^{(\rho_1)}) \lesssim \rho_1 r(p+q)$. Further, $\rho_1 \lesssim \zeta$ and $r(p+q) \lesssim \sqrt{\mathscr{H}(\mathcal{A},\rho_1)} \lesssim \sqrt{\frac{m\zeta}{\delta}}$ imply $\omega(\mathcal{A}_{\text{loc}}^{(\rho_1)}) \lesssim \zeta \sqrt{\frac{m\zeta}{\delta}}$.

Applying Theorem 3.2 yields the desired Corollary 3.7.

Next, we prove the recovery guarantees for unconstrained Lasso. We apply Theorem 3.12 by several steps: (1) Verifying Assumption 4, (2) Selecting (ρ_1, ρ_2) to render (3.46)–(3.47), and (3) Estimating the geometric quantities.

C.3 The Proof of Corollary 3.14 (Recovering Sparse Signal and Sparse Corruption via Unconstrained Lasso)

Proof. We present the proof in three steps.

Step 1: Verifying Assumption 4 Given any $a \in \mathcal{K}_{x}$, we take $\mathcal{X}_{a} = \overline{\mathcal{X}}_{a} = \{w \in \mathbb{R}^{n} : \operatorname{supp}(w) \subset \operatorname{supp}(a)\}$. Then we have $\overline{\mathcal{X}}_{a}^{\perp} = \{w \in \mathbb{R}^{n} : \operatorname{supp}(w) \subset ([n] \setminus \operatorname{supp}(a))\}$, and note that decomposibility (3.45) immediately follows since $||w_{1} + w_{2}||_{1} = ||w_{1}||_{1} + ||w_{2}||_{1}$ holds for any $w_{1}, w_{2} \in \mathbb{R}^{n}$ with $\operatorname{supp}(w_{1}) \subset \operatorname{supp}(a)$ and $\operatorname{supp}(w_{2}) \subset ([n] \setminus \operatorname{supp}(a))$. Moreover, for any $w \in \overline{\mathcal{X}}_{a}$ we have $||w||_{1} \leq \sqrt{s}||w||_{2}$, and hence $\alpha_{\|\cdot\|_{1}}(\overline{\mathcal{X}}_{a}) \leq \sqrt{s}$; This holds uniformly for all $a \in \mathcal{K}_{x}$ and hence we can take $\alpha_{x} = \sqrt{s}$. Similarly, regarding the k-sparse corruption, Assumption 4 is also satisfied with $\alpha_{v} = \sqrt{k}$.

Step 2: Selecting (ρ_1, ρ_2) As shown in Step 2 of the proof of Corollary 3.5, setting $\rho_1 = c\delta(\frac{s}{m})^{3/2}$ and $\rho_2 = c\delta(\frac{k}{m})^{3/2}$ with small enough c satisfies (3.46)–(3.47).

Step 3: Estimating Geometric Quantities For the estimations of $\mathscr{H}(\mathcal{K}_{\boldsymbol{x}}, \cdot)$ and $\mathscr{H}(\mathcal{K}_{\boldsymbol{v}}, \cdot)$ we use Proposition A.6. For the ℓ_1 -ball we have $\omega(\mathbb{B}_f^n) \approx \sqrt{\log n}$ and $\omega(\mathbb{B}_g^m) \approx \sqrt{\log m}$ [78, Example 7.5.9]. Now we can easily see that (λ_1, λ_2) in Corollary 3.14 satisfies (3.48)–(3.49), and the sample complexity stated in Corollary 3.14 satisfies (3.50).

With the above preparations, the result immediately follows from Theorem 3.12.

C.4 The Proof of Corollary 3.15 (Recovering Low-Rank Signal and Sparse Corruption via Unconstrained Lasso)

Proof. We present the proof in three steps.

Step 1: Verifying Assumption 4 Given any $a \in M_r^{p,q}$, we let its singular value decomposition be

(C.5)
$$\boldsymbol{a} = \begin{bmatrix} \boldsymbol{U}_1 & \boldsymbol{U}_2 \end{bmatrix} \begin{bmatrix} \boldsymbol{\Sigma} & \boldsymbol{0} \\ \boldsymbol{0} & \boldsymbol{0} \end{bmatrix} \begin{bmatrix} \boldsymbol{V}_1^\top \\ \boldsymbol{V}_2^\top \end{bmatrix}.$$

Then we define $\mathcal{X}_{\boldsymbol{a}} = \{\boldsymbol{U}_{1}\boldsymbol{A}\boldsymbol{V}_{1}^{\top}: \boldsymbol{A} \in \mathbb{R}^{r \times r}\}, \ \overline{\mathcal{X}}_{\boldsymbol{a}} = \{\boldsymbol{U}_{1}\boldsymbol{A}\boldsymbol{V}_{1}^{\top} + \boldsymbol{U}_{1}\boldsymbol{B}\boldsymbol{V}_{2}^{\top} + \boldsymbol{U}_{2}\boldsymbol{C}\boldsymbol{V}_{1}^{\top}: \boldsymbol{A} \in \mathbb{R}^{r \times r}, \boldsymbol{B} \in \mathbb{R}^{r \times (q-r)}, \boldsymbol{C} \in \mathbb{R}^{(p-r) \times r}\}.$ Note that $\overline{\mathcal{X}}_{\boldsymbol{a}}^{\perp} = \{\boldsymbol{U}_{2}\boldsymbol{A}\boldsymbol{V}_{2}^{\top}: \boldsymbol{A} \in \mathbb{R}^{(p-r) \times (q-r)}\}$, then it is not hard to verify the decomposibility (3.45), since $\|\boldsymbol{w}_{1} + \boldsymbol{w}_{2}\|_{\mathrm{nu}} = \|\boldsymbol{w}_{1}\|_{\mathrm{nu}} + \|\boldsymbol{w}_{2}\|_{\mathrm{nu}}$ holds for any $\boldsymbol{w}_{1} \in \mathcal{X}_{\boldsymbol{a}}$ and $\boldsymbol{w}_{2} \in \overline{\mathcal{X}}_{\boldsymbol{a}}^{\perp}$. Moreover, any $\boldsymbol{w} \in \overline{\mathcal{X}}_{\boldsymbol{a}}$ has rank not exceeding 2r, and hence we have $\|\boldsymbol{x}\|_{\mathrm{nu}} \leq \sqrt{2r}\|\boldsymbol{x}\|_{\mathrm{F}}$. Note that $\|\cdot\|_{\mathrm{F}}$ is just the ℓ_{2} -norm when \boldsymbol{x} is viewed as vector, thus we have $\alpha_{\|\cdot\|_{\mathrm{nu}}}(\mathcal{X}_{\boldsymbol{a}}) \leq \sqrt{2r}$. This holds uniformly for all $\boldsymbol{a} \in M_{r}^{p,q}$, thus we can take $\alpha_{\boldsymbol{x}} = \sqrt{2r}$ in Assumption 4. It has been shown in the proof of Corollary 3.14 that the k-sparse corruption satisfies Assumption 4 with $\alpha_{\boldsymbol{v}} = \sqrt{k}$.

Step 2: Selecting (ρ_1, ρ_2) As shown in Step 2 in the proof of Corollary 3.7, setting $\rho_1 = c\delta(\frac{r(p+q)}{m})^{3/2}$ and $\rho_2 = c\delta(\frac{k}{m})^{3/2}$ with small enough c, along with the sample complexity stated in Corollary 3.15, satisfies (3.46)–(3.13).

Step 3: Estimating Geometric Quantities For estimations of $\mathscr{H}(\mathcal{K}_{\boldsymbol{x}}, \cdot)$ and $\mathscr{H}(\mathcal{K}_{\boldsymbol{v}}, \cdot)$ we use Proposition A.6. [78, Example 7.5.9] gives $\omega(\mathbb{B}_{g}^{m}) \simeq \sqrt{\log m}$. Moreover, we show $\omega(\mathbb{B}_{f}^{n}) = \omega(\mathbb{B}_{nu}^{p,q}) \simeq \sqrt{p+q}$ in the following. First, note that a matrix that has only one non-zero row in \mathbb{B}_{2}^{q} (or only one non-zero column in \mathbb{B}_{2}^{p}) belongs to $\mathbb{B}_{nu}^{p,q}$, which implies $\omega(\mathbb{B}_{nu}^{p,q}) \geq \max\{\omega(\mathbb{B}_{2}^{p}), \omega(\mathbb{B}_{2}^{q})\} = \Omega(\sqrt{p+q})$. Second, let $\boldsymbol{G} \sim \mathcal{N}^{p \times q}(0,1)$ we have $\omega(\mathbb{B}_{nu}^{p,q}) =$ $\mathbb{E}\sup_{\|\boldsymbol{A}\|_{nu}=1}\langle \boldsymbol{G}, \boldsymbol{A} \rangle \leq \mathbb{E}\|\boldsymbol{G}\|_{op} \lesssim \sqrt{p+q}$ [78, Exercise 4.4.6].

With the above preparations, we are ready to invoke Theorem 3.12 to obtain the desired claim. $\hfill\blacksquare$

C.5 The Proof of Theorem 3.16 (Uniform Recovery Guarantee under Generative Priors)

Proof. By writing $\mathcal{K}_{\boldsymbol{x}}^- = \mathcal{K}_{\boldsymbol{x}} - \mathcal{K}_{\boldsymbol{x}}$ and $\mathcal{K}_{\boldsymbol{v}}^- = \mathcal{K}_{\boldsymbol{v}} - \mathcal{K}_{\boldsymbol{v}}$, it is immediate from the constraint of (3.97) that

(C.6)
$$\boldsymbol{\Delta}_{\boldsymbol{x}} = \hat{\boldsymbol{x}} - \boldsymbol{x}^{\star} \in \mathcal{K}_{\boldsymbol{x}}^{-}, \ \boldsymbol{\Delta}_{\boldsymbol{v}} = \hat{\boldsymbol{v}} - \boldsymbol{v}^{\star} \in \mathcal{K}_{\boldsymbol{v}}^{-}.$$

We may omit some details because the techniques are analogous to those for proving Theorem 3.2. We present the proofs in three steps.

Step 1: Problem Reduction We first reduce the proof to bounding several random processes.

Identifying Constraint Sets: Note that we want to prove $\sqrt{\|\Delta_{\boldsymbol{x}}\|_2^2 + \|\Delta_{\boldsymbol{v}}\|_2^2} \leq \mu$ for all $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star}) \in \mathcal{K}_{\boldsymbol{x}} \times \mathcal{K}_{\boldsymbol{v}}$ and for some given accuracy $\mu \in (0, 1)$. Up to rescaling it suffices to prove $\|\boldsymbol{\Delta}\|_2 = \sqrt{\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_2^2 + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_2^2} \leq 3\mu$. Hence, we can assume

(C.7)
$$\sqrt{\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_2^2 + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_2^2} \ge 2\mu$$

since the bound holds trivially when $(\|\Delta_x\|_2^2 + \|\Delta_v\|_2^2)^{1/2} < 2\mu$. Therefore, we can proceed with the constraint

(C.8)
$$(\boldsymbol{\Delta}_{\boldsymbol{x}}, \boldsymbol{\Delta}_{\boldsymbol{v}}) \in \mathcal{E} := \{ (\boldsymbol{c}, \boldsymbol{d}) : \boldsymbol{c} \in \mathcal{K}_{\boldsymbol{x}}^{-}, \ \boldsymbol{d} \in \mathcal{K}_{\boldsymbol{v}}^{-}, \ (\|\boldsymbol{c}\|_{2}^{2} + \|\boldsymbol{d}\|_{2}^{2})^{1/2} \ge 2\mu \},$$

where the constraint set \mathcal{E} is defined as per (A.15). To accmmodate the normalized error, we introduce \mathcal{E}^*

(C.9)
$$\frac{(\boldsymbol{\Delta}_{\boldsymbol{x}}, \boldsymbol{\Delta}_{\boldsymbol{v}})}{(\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{2}^{2} + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2})^{1/2}} \in \mathcal{E}^{*} = \left\{ (\boldsymbol{c}, \boldsymbol{d}) / (\|\boldsymbol{c}\|_{2}^{2} + \|\boldsymbol{d}\|_{2}^{2})^{1/2} : (\boldsymbol{c}, \boldsymbol{d}) \in \mathcal{E} \right\}$$

as per (A.16). Besides, we further define

(C.10)
$$\mathcal{C} = \left\{ \boldsymbol{c} \in \mathbb{R}^n : (\boldsymbol{c}, \boldsymbol{d}) \in \mathcal{E}^* \text{ for some } \boldsymbol{d} \in \mathbb{R}^m \right\}$$

(C.11)
$$\mathcal{D} = \{ \boldsymbol{d} \in \mathbb{R}^m : (\boldsymbol{c}, \boldsymbol{d}) \in \mathcal{E}^* \text{ for some } \boldsymbol{c} \in \mathbb{R}^n \},\$$

then we note the relation

(C.12)
$$\max\{\omega(\mathcal{C}), \omega(\mathcal{D})\} \le \omega(\mathcal{E}^*) \lesssim \left(k \log \frac{Lr}{\mu} + k' \log \frac{L'r'}{\mu}\right)^{1/2},$$

where the first inequality can be seen by [78, Exercise 7.5.4], the second inequality follows from Proposition A.9(c). In the sequel, we will proceed with the constraints

(C.13)
$$\boldsymbol{\Delta}_{\boldsymbol{x}}/(\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_2^2 + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_2^2)^{1/2} \in \mathcal{C},$$

(C.14)
$$\boldsymbol{\Delta}_{\boldsymbol{v}}/(\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_2^2 + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_2^2)^{1/2} \in \mathcal{D}.$$

Note that all above constraints hold universally for all $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star})$ that need further consideration; those $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star})$ that fail to satisfy these constraints must satisfy $(\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_2^2 + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_2^2)^{1/2} < 2\mu$ and are already done.

Using Optimality: From $\|\dot{\boldsymbol{y}} - \boldsymbol{\Phi}\hat{\boldsymbol{x}} - \sqrt{m}\hat{\boldsymbol{v}}\|_2 \leq \|\dot{\boldsymbol{y}} - \boldsymbol{\Phi}\boldsymbol{x}^{\star} - \sqrt{m}\boldsymbol{v}^{\star}\|_2$, we substitute $\hat{\boldsymbol{x}} = \boldsymbol{x}^{\star} + \boldsymbol{\Delta}_{\boldsymbol{x}}$ and $\hat{\boldsymbol{v}} = \boldsymbol{v}^{\star} + \boldsymbol{\Delta}_{\boldsymbol{v}}$, expand the square, and then substitute (3.4) to obtain

(C.15)
$$\|\boldsymbol{\Phi}\boldsymbol{\Delta}_{\boldsymbol{x}} + \sqrt{m}\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2} \leq 2\langle\boldsymbol{\epsilon} + \boldsymbol{\xi}_{\boldsymbol{x}^{\star},\boldsymbol{v}^{\star}}, \boldsymbol{\Phi}\boldsymbol{\Delta}_{\boldsymbol{x}} + \sqrt{m}\boldsymbol{\Delta}_{\boldsymbol{v}}\rangle,$$

see (3.3) and (3.26) for $\boldsymbol{\xi}_{\boldsymbol{x}^{\star},\boldsymbol{v}^{\star}}$. Combining with the constraints in (C.9), (C.10) and (C.11), we bound both sides of (C.15) to arrive at

(C.16)
$$(\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{2}^{2} + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2}) \cdot I_{1} \leq 2(\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{2}^{2} + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2})^{1/2} \cdot (I_{2} + I_{3} + I_{4})$$

where the random terms (to be bounded) are given by

(C.17)
$$I_1 := \inf_{(\boldsymbol{c},\boldsymbol{d})\in\mathcal{E}^*} \|\boldsymbol{\Phi}\boldsymbol{c} + \sqrt{m}\boldsymbol{d}\|_2^2, \ I_2 := \sup_{\boldsymbol{c}\in\mathcal{C}} \langle \boldsymbol{\epsilon}, \boldsymbol{\Phi}\boldsymbol{c} \rangle$$

(C.18)
$$I_3 := \sup_{\boldsymbol{d} \in \mathcal{D}} \langle \boldsymbol{\epsilon}, \sqrt{m} \boldsymbol{d} \rangle, \ I_4 := \sup_{\boldsymbol{a} \in \mathcal{K}_{\boldsymbol{v}}} \sup_{\boldsymbol{b} \in \mathcal{K}_{\boldsymbol{v}}} \sup_{(\boldsymbol{c}, \boldsymbol{d}) \in \mathcal{E}^*} \langle \boldsymbol{\xi}_{\boldsymbol{a}, \boldsymbol{b}}, \boldsymbol{\Phi} \boldsymbol{c} + \sqrt{m} \boldsymbol{d} \rangle$$

Step 2: Bounding I_1, I_2, I_3, I_4 Parallel to the proof of Theorem 3.2, our techniques to bound I_1, I_2, I_3, I_4 are Proposition A.1, Proposition A.4, Proposition A.2 and Corollary B.4, respectively. The additional technicalities are the estimates on Gaussian width and Kolmogorov entropy developed in Proposition A.9.

Bounding I_1 : Because $\mathcal{E}^* \subset \mathbb{S}^{n+m-1}$, Proposition A.1 yields that the event

(C.19)
$$\sup_{(\boldsymbol{c},\boldsymbol{d})\in\mathcal{E}^*} \left| \|\boldsymbol{\Phi}\boldsymbol{c} + \sqrt{m}\boldsymbol{d}\|_2 - \sqrt{m} \right| \leq C_1 \big(\gamma(\mathcal{E}^*) + t\big)$$

holds with probability exceeding $1 - \exp(-t^2)$. Notice that the bound on $\omega(\mathcal{E}^*)$ in (C.12) remains valid for $\gamma(\mathcal{E}^*)$ due to (2.8). Since (3.98) implies $m \gtrsim k \log \frac{Lr}{\mu} + k' \log \frac{L'r'}{\mu}$, we can set $t = \sqrt{k \log \frac{Lr}{\mu} + k' \log \frac{L'r'}{\mu}}$ and still assume that the right-hand side of (C.19) is bounded by $\frac{\sqrt{m}}{2}$, thus obtaining that the event

(C.20)
$$\sqrt{I_1} = \inf_{(\boldsymbol{c},\boldsymbol{d})\in\mathcal{E}^*} \|\boldsymbol{\Phi}\boldsymbol{c} + \sqrt{m}\boldsymbol{d}\|_2$$

(C.21)
$$\geq \sqrt{m} - \sup_{(\boldsymbol{c},\boldsymbol{d})\in\mathcal{E}^*} \left| \|\boldsymbol{\Phi}\boldsymbol{c} + \sqrt{m}\boldsymbol{d}\|_2 - \sqrt{m} \right| \geq \frac{\sqrt{m}}{2}$$

holds with probability exceeding $1 - \exp(-k \log \frac{Lr}{\mu} - k' \log \frac{L'r'}{\mu})$. Bounding I_2 : We derive the bound following similar courses as in the corresponding part in the proof of Theorem 3.2. Conditioning on ϵ , for any $t \geq 0$, Proposition A.4 gives that the event

(C.22)
$$I_2 \le C_2 \|\boldsymbol{\epsilon}\|_2 (\omega(\mathcal{C}) + t)$$

holds with probability exceeding $1 - 2\exp(-t^2)$. By repeating the argument in (3.33) we can show that $\|\boldsymbol{\epsilon}\|_2 \lesssim E\sqrt{m}$ holds with probability exceeding $1 - \exp(-\Omega(m))$, and hence also exceeding $1 - \exp(-\Omega(k \log \frac{Lr}{\mu} + k' \log \frac{L'r'}{\mu}))$ by (3.98). Combining with (C.12), we set $t = (k \log \frac{Lr}{\mu} + k' \log \frac{L'r'}{\mu})^{1/2}$ in (C.22) to obtain the bound on I_2

(C.23)
$$I_2 \lesssim E\sqrt{m} \cdot \left(k \log \frac{Lr}{\mu} + k' \log \frac{L'r'}{\mu}\right)^{1/2},$$

with probability exceeding $1 - 3 \exp(-\Omega(k \log \frac{Lr}{\mu} + k' \log \frac{L'r'}{\mu}))$. **Bounding** I_3 : Due to (3.35), for any $t \ge 0$, Proposition A.2 yields that the event $I_3 \lesssim$ $E\sqrt{m}(\omega(\mathcal{D})+t)$ holds with probability exceeding $1-2\exp(-t^2)$. Combining with (C.12), we set $t = (k \log \frac{Lr}{\mu} + k' \log \frac{L'r'}{\mu})^{1/2}$ to obtain that the bound on I_3

(C.24)
$$I_3 \lesssim E\sqrt{m} \cdot \left(k \log \frac{Lr}{\mu} + k' \log \frac{L'r'}{\mu}\right)^{1/2}$$

holds with probability exceeding $1 - 2 \exp(-k \log \frac{Lr}{\mu} - k' \log \frac{L'r'}{\mu})$.

Bounding I_4 : We apply Corollary B.4 to bound I_4 , so the major work lies in selecting (ρ_1, ρ_2) such that (B.88)–(B.89) hold. We claim that setting

(C.25)
$$\rho_1 = c\delta\left(\frac{k}{n}\right)^{3/2}, \ \rho_2 = c\delta\left(\frac{k'}{m}\right)^{3/2}$$

works, with the reasoning provided below:¹⁸

- In general, we do not have lower bound on the Kolmogorov entropy of \mathcal{K}_{x} and \mathcal{K}_{v} in (3.96), but we note that the $\mathscr{H}(\mathcal{A}, \rho_{1})$ and the $\mathscr{H}(\mathcal{B}, \rho_{2})$ appearing in the statement of Corollary B.4 can be replaced by their upper bounds, and we will simply use (A.12).
- That being mentioned, from (B.88) we have $\zeta \gtrsim \frac{\delta(k+k')}{m}$, and so $\rho_2 = c\delta(\frac{k'}{m})^{3/2} \lesssim \zeta \sqrt{\frac{\zeta}{\delta}}$. Since (3.98) implies $m \gtrsim k+k'$, we can assume that $\frac{\delta}{\zeta}$ is sufficiently large. Hence, to ensure $\rho_1 \lesssim \frac{\zeta}{(\log \frac{\delta}{\zeta})^{1/2}}$, it suffices to ensure $\rho_1 \lesssim \zeta \sqrt{\frac{\zeta}{\delta}}$. Under m = O(n), $\rho_1 = c\delta(\frac{k}{n})^{3/2}$ with small enough c evidently satisfies this.
- It remains to verify the second condition in (B.89), namely $\omega(\mathcal{A}_{\text{loc}}^{(\rho_1)}) \lesssim \zeta \sqrt{\frac{m\zeta}{\delta}}$, and by $\zeta \gtrsim \frac{\delta k}{m}$ it suffices to ensure $\omega(\mathcal{A}_{\text{loc}}^{(\rho_1)}) \lesssim \frac{\delta k^{3/2}}{m}$. This can be justified by $\omega(\mathcal{A}_{\text{loc}}^{(\rho_1)}) \leq \rho_1 \omega(\mathbb{B}_2^n) \leq \rho_1 \sqrt{n} = c \delta \frac{k^{3/2}}{n}$, where we use $\omega(\mathbb{B}_2^n) \leq \sqrt{n}$ from [78, Prop. 7.5.2(f)].

Moreover, note that (3.98) implies $m \gtrsim k \log(\frac{Lrn^{3/2}}{\delta k^{3/2}}) + k' \log(\frac{L'r'm^{3/2}}{\delta (k')^{3/2}})$, and thus we can apply Corollary B.4 to obtain that the bound

(C.26)
$$I_4 \lesssim \delta \sqrt{mk \log\left(\frac{Lrn^{3/2}}{\mu\delta k^{3/2}}\right) + mk' \log\left(\frac{L'r'm^{3/2}}{\mu\delta(k')^{3/2}}\right)}$$

that holds with probability exceeding $1 - 12 \exp(-\Omega(k \log(\frac{Lrn^{3/2}}{\mu \delta k^{3/2}}) + k' \log(\frac{L'r'm^{3/2}}{\mu \delta (k')^{3/2}})))$.

Step 3: Combining Everything We substitute the bounds (C.21), (C.23), (C.24), (C.26) into (C.16) and perform simple rearrangement. This yields the bound (universally for all $(\boldsymbol{x}^{\star}, \boldsymbol{v}^{\star})$ that may violate $(\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_2^2 + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_2^2)^{1/2} \leq 2\mu$)

$$\left(\|\boldsymbol{\Delta}_{\boldsymbol{x}}\|_{2}^{2} + \|\boldsymbol{\Delta}_{\boldsymbol{v}}\|_{2}^{2} \right)^{1/2} \lesssim \frac{E\sqrt{k\log(\frac{Lr}{\mu}) + k'\log(\frac{L'r'}{\mu})} + \delta\sqrt{k\log(\frac{Lrn^{3/2}}{\mu\delta k^{3/2}}) + k'\log(\frac{L'r'm^{3/2}}{\mu\delta(k')^{3/2}})}{\sqrt{m}}$$

with probability exceeding $1 - C_3 \exp(-\Omega(k \log(Lr) + k' \log(L'r')))$. Therefore, under the sample size given in (3.98), we again obtain $(\|\Delta_x\|_2^2 + \|\Delta_v\|_2^2)^{1/2} \leq 3\mu$, completing the proof.

Appendix D. Technical By-Product. We demonstrate that our global QPE property Theorem B.3 is a generalization and instance-wise improvement (under Gaussian sensing matrix Φ) of the one developed in [81]. Then, as an interesting enough technical by-product, we improve the uniform error rate of the *projected back-projection* (PBP) estimator over bounded convex signal set in [81] from $O(m^{-1/16})$ to $O(m^{-1/8})$.

¹⁸Unlike in the case of structured priors, we do not aim to carefully choose ρ_1 but simply set it small enough to justify (B.89). The reason is that other parameters (L, r, L', r') appearing in the logarithm typically dominate (m, n). (As a result, most works in generative compressed sensing do not refine logarithmic factor.)

D.1 Implications of Theorem B.3 Recall that we have specialized Theorem B.3 to the case where \mathcal{A} and \mathcal{B} are *structured sets* (see Definition 3.1) in Corollary B.4. Here, for *arbitrary* sets $\mathcal{A} \subset \mathbb{R}^n$ and $\mathcal{B} \subset \mathbb{R}^m$, we further present some direct outcomes of our general Theorem B.3 and compare them with [81].

The key ingredient in [81] for achieving global QPE is their Proposition 6.1, which can be recovered from our Theorem B.3 by setting $\mathcal{A} = \mathcal{C} = \{0\}$ (see Remark D.2).

Corollary D.1 (QPE (Almost) Coincident with Proposition 6.1 in [81]). Given $\mathcal{B}, \mathcal{D} \subset \mathbb{B}_2^m$, we consider the uniform quantizer $\mathcal{Q}_{\delta}(\cdot)$ associated with uniform dither $\boldsymbol{\tau} \sim \mathscr{U}([-\frac{\delta}{2}, \frac{\delta}{2}]^m)$. Given any small enough $\epsilon > 0$, if

(D.1)
$$m \ge C_1(\frac{\mathscr{H}(\mathcal{B}, \delta\epsilon^3)}{\epsilon^2} + \frac{\omega^2(\mathcal{D})}{\epsilon^2})$$

holds for some sufficiently large C_1 , then the event

(D.2)
$$\sup_{\boldsymbol{b}\in\mathcal{B}}\sup_{\boldsymbol{d}\in\mathcal{D}}\left|\langle \mathcal{Q}_{\delta}(\sqrt{m}\boldsymbol{b}+\boldsymbol{\tau})-\sqrt{m}\boldsymbol{b},\sqrt{m}\boldsymbol{d}\rangle\right|\leq C_{2}\delta m\epsilon$$

holds with probability exceeding $1 - 12 \exp(-c_3 \mathscr{H}(\mathcal{B}, \delta \epsilon^3))$.

Proof. Applying Theorem B.3: We invoke Theorem B.3 with $\mathcal{A} = \mathcal{C} = \{0\}, \mathcal{E} = \{0\} \times \mathcal{D}, \epsilon = 0$. Using arbitrarily small ρ_1 that renders the second condition in (B.17), we always have $\mathscr{H}(\mathcal{A}, \rho_1) = 0$ and $\omega^2(\mathcal{A}_{\text{loc}}^{(\rho_1)}) = 0$ due to $\mathcal{A} = \{0\}$. Combining with $\mathcal{B}, \mathcal{D} \subset \mathbb{B}_2^m$, Theorem B.3 gives that, if $\zeta \in (0, \frac{\delta}{2})$ and $\rho_2 \lesssim \zeta, m \gtrsim \frac{2\delta \cdot \mathscr{H}(\mathcal{B}, \rho_2)}{\zeta}$, then the event

(D.3)
$$\sup_{\boldsymbol{b}\in\mathcal{B}}\sup_{\boldsymbol{d}\in\mathcal{D}}\left|\langle \mathcal{Q}_{\delta}(\sqrt{m}\boldsymbol{b}+\boldsymbol{\tau})-\sqrt{m}\boldsymbol{b},\sqrt{m}\boldsymbol{d}\rangle\right| \lesssim \delta m\left(\frac{\omega(\mathcal{D})}{\sqrt{m}}+\sqrt{\frac{\zeta}{\delta}}+\frac{\rho_{2}}{\zeta}\right)$$

holds with probability exceeding $1 - 12 \exp(-\Omega(\mathscr{H}(\mathcal{B}, \rho_2))))$.

Choosing Parameters: Given sufficiently small $\epsilon > 0$, we set

 $\zeta = \delta \epsilon^2, \ \rho_2 = \epsilon \zeta = \delta \epsilon^3$

that satisfy the required conditions $\zeta \in (0, \frac{\delta}{2})$ and $\rho_2 \lesssim \zeta$. Under such choice, the required sample size of $m \gtrsim \frac{2\delta \cdot \mathscr{H}(\mathcal{B}, \rho_2)}{\zeta}$ reads as $m \gtrsim \frac{\mathscr{H}(\mathcal{B}, \delta\epsilon^3)}{\epsilon^2}$, which is satisfied due to (D.1), and the right-hand side of (D.3) becomes $\delta m(\frac{\omega(\mathcal{D})}{\sqrt{m}} + 2\epsilon)$. Combining with $m \gtrsim \frac{\omega^2(\mathcal{D})}{\epsilon^2}$, we arrive at the desired bound of $O(\delta m\epsilon)$. The promised probability is directly dictated from Theorem B.3.

Remark D.2. There is no essential difference between our Corollary D.1 and [81, Prop. 6.1], and we simply note the specific two points: (i) [81, Prop. 6.1] is stated for a fixed $\boldsymbol{d} = \boldsymbol{d}_0$, which corresponds to the special case of Corollary D.1 with $\mathcal{D} = \{\boldsymbol{d}_0\}$; (ii) [81, Prop. 6.1] is stated for $\tilde{\boldsymbol{b}} := \sqrt{m}\boldsymbol{b} \in \tilde{\mathcal{B}} := \sqrt{m}\mathcal{B}$, so the sample complexity is consistent since $\mathscr{H}(\mathcal{B}, \delta\epsilon^3) = \mathscr{H}(\tilde{\mathcal{B}}, \sqrt{m}\delta\epsilon^3)$ always holds.

Next, we show that improvement can be obtained if (\mathbf{b}, \mathbf{d}) in (D.2) is modulated by a sub-Gaussian sensing matrix (rather than being simply re-scaled by a factor of \sqrt{m}), as will be discussed in Remark D.4. To get the improved QPE property, we invoke Theorem B.3 with $\mathcal{B} = \mathcal{D} = \{0\}$.

Corollary D.3 (Improved QPE under Sub-Gaussian Matrix). Given $\mathcal{A}, \mathcal{C} \subset \mathbb{B}_2^n$, we assume that the sub-Gaussian sensing matrix Φ and uniform dither τ are as described in Assumption 1. Given any small enough $\epsilon > 0$, we let $\rho = \frac{c_0 \delta \epsilon}{\sqrt{\log \epsilon^{-1}}}$ for sufficiently small absolute constant c_0 . If

(D.4)
$$m \ge C_1 \left(\frac{\mathscr{H}(\mathcal{A}, \rho)}{\epsilon^2 \log(\epsilon^{-1})} + \frac{\omega^2(\mathcal{A}_{\text{loc}}^{(\rho)})}{\delta^2 \epsilon^3} + \frac{\omega^2(\mathcal{C})}{\epsilon^2 \log(\epsilon^{-1})} \right)$$

holds for some sufficiently large C_1 , then the event

(D.5)
$$\sup_{\boldsymbol{a}\in\mathcal{A}}\sup_{\boldsymbol{c}\in\mathcal{C}}\left|\langle\mathcal{Q}_{\delta}(\boldsymbol{\Phi}\boldsymbol{a}+\boldsymbol{\tau})-\boldsymbol{\Phi}\boldsymbol{a},\boldsymbol{\Phi}\boldsymbol{c}\rangle\right|\leq C_{2}\delta m\epsilon\sqrt{\log\epsilon^{-1}}$$

holds with probability exceeding $1 - 12 \exp(-c_3 \mathscr{H}(\mathcal{A}, \rho))$.

Proof. Applying Theorem B.3: We invoke Theorem B.3 with $\mathcal{B} = \mathcal{D} = \{0\}$, $\mathcal{E} = \mathcal{C} \times \{0\}$, $\boldsymbol{\epsilon} = 0$. Using arbitrarily small ρ_2 to render $\rho_2 \lesssim \zeta$ needed in (B.17), we always have $\mathscr{H}(\mathcal{B}, \rho_2) = 0$ since $\mathscr{B} = \{0\}$. Combining with $\mathcal{A}, \mathcal{C} \subset \mathbb{B}_2^n$, Theorem B.3 gives that, if for some positive scalars $\rho = \rho_1, \zeta$ we have

(D.6)
$$\zeta \in \left(0, \frac{\delta}{2}\right), \ \rho \lesssim \frac{\zeta}{(\log \frac{\delta}{\zeta})^{1/2}}, \ m \gtrsim \frac{\delta \cdot \mathscr{H}(\mathcal{A}, \rho)}{\zeta} + \frac{\omega^2(\mathcal{A}_{\text{loc}}^{(\rho)})}{\zeta^2},$$

then the event

(D.7)
$$\sup_{\boldsymbol{a}\in\mathcal{A}}\sup_{\boldsymbol{c}\in\mathcal{C}}\left|\langle\mathcal{Q}_{\delta}(\boldsymbol{\Phi}\boldsymbol{a}+\boldsymbol{\tau})-\boldsymbol{\Phi}\boldsymbol{a},\boldsymbol{\Phi}\boldsymbol{c}\rangle\right|$$

(D.8)
$$\lesssim \delta m \left(\frac{\omega(\mathcal{C})}{\sqrt{m}} + \frac{\sqrt{\mathscr{H}(\mathcal{A},\rho)}}{\sqrt{m}} + \frac{\zeta}{\delta} \sqrt{\log \frac{\delta}{\zeta}} + \frac{\omega^2(\mathcal{A}_{\rm loc}^{(\rho)})}{m\zeta^2} \sqrt{\log \frac{\delta}{\zeta}} \right)$$

holds with probability exceeding $1 - 12 \exp(-\Omega(\mathscr{H}(\mathcal{A}, \rho)))$.

Choosing Parameters: We proceed with the parametrization $\zeta = \delta \epsilon$ with the given small enough ϵ , which along with $\rho = \frac{c_0 \delta \epsilon}{\sqrt{\log \epsilon^{-1}}}$ ensures the first two conditions in (D.6). Note that (D.4), additionally implies

$$m \gtrsim \frac{\mathscr{H}(\mathcal{A}, \rho)}{\epsilon} + \frac{\omega^2(\mathcal{A}_{\mathrm{loc}}^{(\rho)})}{\delta^2 \epsilon^2},$$

which is just the third condition in (D.6) due to $\zeta = \delta \epsilon$. Therefore, the bound (D.8) on (D.7) holds with the promised probability, and by substituting $\zeta = \delta \epsilon$ it reads as

(D.9)
$$O\left(\delta m \left(\frac{\omega(\mathcal{C})}{\sqrt{m}} + \frac{\sqrt{\mathscr{H}(\mathcal{A},\rho_1)}}{\sqrt{m}} + \epsilon \sqrt{\log \epsilon^{-1}} + \frac{\omega^2(\mathcal{A}_{\text{loc}}^{(\rho)})\sqrt{\log \epsilon^{-1}}}{m\delta^2\epsilon^2}\right)\right)$$

The desired claim (D.5) thus follows under the sample complexity in (D.4) since under the assumed sample complexity (D.4), (D.9) scales as $O(\delta m \epsilon \sqrt{\log \epsilon^{-1}})$ — specifically, $m \gtrsim \frac{\mathscr{H}(\mathcal{A},\rho_1)}{\epsilon^2 \log(\epsilon^{-1})}$ ensures $\frac{\sqrt{\mathscr{H}(\mathcal{A},\rho)}}{\sqrt{m}} \lesssim \epsilon \sqrt{\log \epsilon^{-1}}$, $m \gtrsim \frac{\omega^2(\mathcal{A}_{\mathrm{loc}}^{(\rho)})}{\delta^2 \epsilon^3}$ ensures $\frac{\omega^2(\mathcal{A}_{\mathrm{loc}}^{(\rho)})\sqrt{\log \epsilon^{-1}}}{m\delta^2 \epsilon^2} \lesssim \epsilon \sqrt{\log \epsilon^{-1}}$, and $m \gtrsim \frac{\omega^2(\mathcal{C})}{\epsilon^2 \log(\epsilon^{-1})}$ implies $\frac{\omega(\mathcal{C})}{\sqrt{m}} \lesssim \epsilon \sqrt{\log \epsilon^{-1}}$. The proof is complete.

Remark D.4 (Comparing Corollary D.1 and Corollary D.3). Note that the random process in (D.5) reduces to the one in (D.2) when $\Phi = \sqrt{m}I$. In this remark, we show that distortion in (D.5) exhibits a decaying rate in *m* faster than (D.2), due to the modulation of the sub-Gaussian matrix Φ . Recall that Corollary D.1 and Corollary D.3 aim to handle arbitrary signal sets \mathcal{A} and \mathcal{B} , thus Sudakov's inequality (2.6) is tight. Moreover, we use the simple bound (that follows from $\rho \leq 1$ and [78, Prop. 7.5.2(e)])

(D.10)
$$\omega^2(\mathcal{A}_{\text{loc}}^{(\rho)}) \le \omega^2(\mathcal{A} - \mathcal{A}) = 4\omega^2(\mathcal{A}).$$

Now we present Corollary D.1 and Corollary D.3 in the form of error rate:

• The Decaying Rate of Corollary D.1: By Sudakov's inequality, $m \gtrsim \frac{\omega^2(\mathcal{B})}{\delta^2 \epsilon^8} + \frac{\omega^2(\mathcal{D})}{\epsilon^2}$ (with large enough implied constant, implicitly below), or equivalently

$$\delta\epsilon\gtrsim\delta^{3/4}\Bigl(rac{\omega^2(\mathcal{B})}{m}\Bigr)^{1/8}+\delta\Bigl(rac{\omega^2(\mathcal{D})}{m}\Bigr)^{1/2},$$

suffices for ensuring (D.1). Therefore, provided that $m \gtrsim \frac{\omega^2(\mathcal{B})}{\delta^2} + \omega^2(\mathcal{D})$, (D.2) in Corollary D.1 implies the following bound on the QPE distortion:

(D.11)
$$\sup_{\boldsymbol{b}\in\mathcal{B}}\sup_{\boldsymbol{d}\in\mathcal{D}}\frac{1}{m}\left|\left\langle \mathcal{Q}_{\delta}(\sqrt{m}\boldsymbol{b}+\boldsymbol{\tau})-\sqrt{m}\boldsymbol{b},\sqrt{m}\boldsymbol{d}\right\rangle\right| \lesssim \delta^{3/4}\left(\frac{\omega^{2}(\mathcal{B})}{m}\right)^{1/8}+\delta\left(\frac{\omega^{2}(\mathcal{D})}{m}\right)^{1/2}.$$

• The Decaying Rate of Corollary D.3: By Sudakov's inequality and (D.10), the condition

(D.12)
$$m \gtrsim \frac{\omega^2(\mathcal{A})}{\rho^2 \epsilon^2 \log(\epsilon^{-1})} + \frac{\omega^2(\mathcal{A})}{\delta^2 \epsilon^3} + \frac{\omega^2(\mathcal{C})}{\epsilon^2 \log(\epsilon^{-1})}$$

suffices for ensuring (D.4). By substituting $\rho = \frac{c_0 \delta \epsilon}{\sqrt{\log \epsilon^{-1}}}$ and $\epsilon < 1$, we can write (D.12) as $m \gtrsim \frac{\omega^2(\mathcal{A})}{\delta^2 \epsilon^4} + \frac{\omega^2(\mathcal{C})}{\epsilon^2 \log(\epsilon^{-1})}$, and further note that this can be guaranteed by two conditions:

(D.13)
$$\epsilon \gtrsim \left(\frac{\omega^2(\mathcal{A})}{\delta^2 m}\right)^{1/4} \text{ and } \delta \epsilon \sqrt{\log \epsilon^{-1}} \gtrsim \delta \left(\frac{\omega^2(\mathcal{C})}{m}\right)^{1/2}$$

Moreover, since $\epsilon \sqrt{\log \epsilon^{-1}}$ is monotonically increasing with ϵ when ϵ is sufficiently small, the first condition in (D.13) is equivalent to

(D.14)
$$\delta\epsilon\sqrt{\log\epsilon^{-1}} \gtrsim \sqrt{\delta} \left(\frac{\omega^2(\mathcal{A})}{m}\right)^{1/4} \left(\log\frac{\delta^2 m}{\omega^2(\mathcal{A})}\right)^{1/2}.$$

Overall, the above analysis shows that

(D.15)
$$\delta\epsilon\sqrt{\log\epsilon^{-1}} \gtrsim \sqrt{\delta} \left(\frac{\omega^2(\mathcal{A})}{m}\log^2\left(\frac{\delta^2 m}{\omega^2(\mathcal{A})}\right)\right)^{1/4} + \delta\left(\frac{\omega^2(\mathcal{C})}{m}\right)^{1/2}$$

with sufficiently large implied constant can imply (D.4). Therefore, provided that $m \gtrsim \frac{\omega^2(\mathcal{A})}{\delta^2} + \omega^2(\mathcal{C})$, (D.5) in Corollary D.3 implies the following bound on the QPE distortion:

(D.16)
$$\sup_{\boldsymbol{a}\in\mathcal{A}} \sup_{\boldsymbol{c}\in\mathcal{C}} \frac{1}{m} |\langle \mathcal{Q}_{\delta}(\boldsymbol{\Phi}\boldsymbol{a}+\boldsymbol{\tau}) - \boldsymbol{\Phi}\boldsymbol{a}, \boldsymbol{\Phi}\boldsymbol{c} \rangle| \\ \lesssim \sqrt{\delta} \Big(\frac{\omega^2(\mathcal{A})}{m} \log^2 \Big(\frac{\delta^2 m}{\omega^2(\mathcal{A})} \Big) \Big)^{1/4} + \delta \Big(\frac{\omega^2(\mathcal{C})}{m} \Big)^{1/2}.$$

Comparing (D.11) and (D.16), it shall be clear that Corollary D.3 provides decaying rate of the QPE distortion faster than Corollary D.1.

D.2 Improving Uniform Error Decaying Rate for PBP We present an interesting byproduct as our final technical development: with Corollary D.3, under sub-Gaussian measurement matrix, we are able to improve the uniform recovery guarantee for the projected-back projection (PBP) estimator over signals from a convex and symmetric set \mathcal{K} in [81, Sec. 7.3B].

PBP Estimator and Uniform Guarantee in [81] We first review the PBP estimator and the related result in [81]. Suppose that the signal $x^* \in \mathbb{R}^n$ lies in some convex and symmetric set \mathcal{K} with $\operatorname{rad}(\mathcal{K}) \leq 1$, and the reader may think of a typical example given by the set of effectively sparse signals (e.g., [65, 66])

(D.17)
$$\mathcal{K} = \mathbb{B}_1^n(\sqrt{s}) \cap \mathbb{B}_2^n = \{ \boldsymbol{x} \in \mathbb{R}^n : \|\boldsymbol{x}\|_1 \le \sqrt{s}, \|\boldsymbol{x}\|_2 \le 1 \},$$

which is essentially larger than the set of exactly s-sparse signals $\Sigma_s^n \cap \mathbb{B}_2^n$. Under the sensing matrix $\mathbf{\Phi} \in \mathbb{R}^{m \times n}$ and uniform dither $\boldsymbol{\tau} \sim \mathscr{U}([-\frac{\delta}{2}, \frac{\delta}{2}]^m)$, we observe the quantized measurements $\dot{\boldsymbol{y}} = \mathcal{Q}_{\delta}(\boldsymbol{\Phi}\boldsymbol{x}^{\star} + \boldsymbol{\tau})$. Let $\mathcal{P}_{\mathcal{K}}(\cdot)$ be the projection operator onto \mathcal{K} under ℓ_2 -norm, then the PBP estimator is given by (e.g., [68, 81])

(D.18)
$$\hat{\boldsymbol{x}}_{\text{PBP}} = \mathcal{P}_{\mathcal{K}} \Big(\frac{1}{m} \boldsymbol{\Phi}^{\top} \boldsymbol{\dot{y}} \Big).$$

Given a general RIP matrix Φ (one that satisfies restricted isometry property (RIP)), it was shown in [81, Sec. 7.3B] that PBP achieves uniform recovery over all $x^* \in \mathcal{K}$ with the following error rate (up to logarithmic factors)

(D.19)
$$\|\hat{\boldsymbol{x}}_{\text{PBP}} - \boldsymbol{x}^{\star}\|_{2} = \tilde{O}\left((1+\delta)^{\frac{1}{2}} \left(\frac{\omega^{2}(\mathcal{K})}{m}\right)^{\frac{1}{16}}\right).$$

We also recap some argument from the proofs of [81, Thm. 4.3, Coro. 3.1] for bounding the PBP estimation error. Let $\boldsymbol{a}_{\boldsymbol{x}^{\star}} = \frac{1}{m} \boldsymbol{\Phi}^{\top} \boldsymbol{\dot{y}}$ (note that this depends on \boldsymbol{x}^{\star}) be the intermediate estimator, then the PBP estimator can be written as $\hat{\boldsymbol{x}}_{\text{PBP}} = \mathcal{P}_{\mathcal{K}}(\boldsymbol{a}_{\boldsymbol{x}^{\star}})$, and we can proceed

as

(D.20)
$$\|\hat{x}_{PBP} - x^{\star}\|_{2}^{2}$$

(D.21)
$$= \|\mathcal{P}_{\mathcal{K}}(\boldsymbol{a}_{\boldsymbol{x}^{\star}}) - \mathcal{P}_{\mathcal{K}}(\boldsymbol{x}^{\star})\|_{2}^{2}$$

$$(D.22) \leq \langle \boldsymbol{x}^* - \boldsymbol{a}_{\boldsymbol{x}^*}, \mathcal{P}_{\mathcal{K}}(\boldsymbol{x}^*) - \mathcal{P}_{\mathcal{K}}(\boldsymbol{a}_{\boldsymbol{x}^*}) \rangle$$

(D.23)
$$\leq |\langle \boldsymbol{x}^{\star} - \boldsymbol{a}_{\boldsymbol{x}^{\star}}, \mathcal{P}_{\mathcal{K}}(\boldsymbol{x}^{\star})\rangle| + |\langle \boldsymbol{x}^{\star} - \boldsymbol{a}_{\boldsymbol{x}^{\star}}, \mathcal{P}_{\mathcal{K}}(\boldsymbol{a}_{\boldsymbol{x}^{\star}})\rangle|$$

(D.24)
$$\leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{x}^{\star} - \frac{1}{m} \boldsymbol{\Phi}^{\top} \cdot \mathcal{Q}_{\delta}(\boldsymbol{\Phi} \boldsymbol{x}^{\star} + \boldsymbol{\tau}), \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{x}^{\star} - \frac{1}{m} \boldsymbol{\Phi}^{\top} \cdot \mathcal{Q}_{\delta}(\boldsymbol{\Phi} \boldsymbol{x}^{\star} + \boldsymbol{\tau}), \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{x}^{\star} - \frac{1}{m} \boldsymbol{\Phi}^{\top} \cdot \mathcal{Q}_{\delta}(\boldsymbol{\Phi} \boldsymbol{x}^{\star} + \boldsymbol{\tau}), \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \left\langle \boldsymbol{u} \right\rangle \right| \leq$$

(D.25)
$$\leq 2 \sup_{\boldsymbol{v} \in \mathcal{K}} \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \langle \boldsymbol{v}, \boldsymbol{u} \rangle - \frac{1}{m} \langle \mathcal{Q}_{\delta}(\boldsymbol{\Phi}\boldsymbol{v} + \boldsymbol{\tau}), \boldsymbol{\Phi}\boldsymbol{u} \rangle \right|$$

(D.26)
$$\leq 2 \sup_{\boldsymbol{u}, \boldsymbol{v} \in \mathcal{K}} \left| \langle \boldsymbol{v}, \boldsymbol{u} \rangle - \frac{1}{m} \langle \boldsymbol{\Phi} \boldsymbol{v}, \boldsymbol{\Phi} \boldsymbol{u} \rangle \right| + 2 \sup_{\boldsymbol{u}, \boldsymbol{v} \in \mathcal{K}} \frac{1}{m} \left| \langle \mathcal{Q}_{\delta}(\boldsymbol{\Phi} \boldsymbol{v} + \boldsymbol{\tau}) - \boldsymbol{\Phi} \boldsymbol{v}, \boldsymbol{\Phi} \boldsymbol{u} \rangle \right|$$
$$:= I_{1}$$

where (D.22) follows from the non-expansivity of the projector onto \mathcal{K} , in (D.24) we take the supremum over $\mathcal{P}_{\mathcal{K}}(\boldsymbol{x}^{\star}), \mathcal{P}_{\mathcal{K}}(\boldsymbol{a}_{\boldsymbol{x}^{\star}}) \in \mathcal{K}$ and substitute $\boldsymbol{a}_{\boldsymbol{x}^{\star}} = \frac{1}{m} \boldsymbol{\Phi}^{\top} \dot{\boldsymbol{y}} = \frac{1}{m} \boldsymbol{\Phi}^{\top} \cdot \mathcal{Q}_{\delta}(\boldsymbol{\Phi} \boldsymbol{x}^{\star} + \boldsymbol{\tau}),$ in (D.25) we take the supremum over $x^{\star} \in \mathcal{K}$, and (D.26) follows from triangle inequality. Then, the critical observation made by [81] can be summarized as follows:

I_1 can be bounded by RIP, I_2 can be bounded by QPE.

To see how one can bound I_1 via RIP, suppose that $\mathcal{K} \subset \mathbb{B}_2^n$, the RIP over the convex symmetric \mathcal{K} with distortion β is formulated as

(D.27)
$$\sup_{\boldsymbol{w}\in\mathcal{K}} \left|\frac{1}{m} \|\boldsymbol{\Phi}\boldsymbol{w}\|_{2}^{2} - \|\boldsymbol{w}\|_{2}^{2}\right| \leq \beta,$$

and one can bound I_1 by (D.27) since

(D.28)
$$I_{1} = \sup_{\boldsymbol{u}, \boldsymbol{v} \in \mathcal{K}} \left| \frac{1}{m} \left\| \boldsymbol{\Phi} \left(\frac{\boldsymbol{u} + \boldsymbol{v}}{2} \right) \right\|_{2}^{2} - \frac{1}{m} \left\| \boldsymbol{\Phi} \left(\frac{\boldsymbol{u} - \boldsymbol{v}}{2} \right) \right\|_{2}^{2} - \left\| \frac{\boldsymbol{u} + \boldsymbol{v}}{2} \right\|_{2}^{2} + \left\| \frac{\boldsymbol{u} - \boldsymbol{v}}{2} \right\|_{2}^{2} \right\|_{2}^{2}$$

(D.29)
$$\leq 2 \sup_{\boldsymbol{w} \in \mathcal{K}} \left| \frac{1}{m} \| \boldsymbol{\Phi} \boldsymbol{w} \|_2^2 - \| \boldsymbol{w} \|_2^2 \right| \leq \beta \leq 2\beta,$$

where (D.29) holds because $\frac{u+v}{2}, \frac{u-v}{2} \in \mathcal{K}$ (recall that \mathcal{K} is convex and symmetric).

Improved Rate under Sub-Gaussian Φ In the specific instance of sub-Gaussian Φ , we are able to improve (D.19) by bounding I_2 via our Corollary D.3. We formally present this as the following statement.

Proposition D.5 (Improved PBP Uniform Rate under sub-Gaussian Matrix). Let the sub-Gaussian matrix $\mathbf{\Phi} \in \mathbb{R}^{m \times n}$ and the uniform dither $\boldsymbol{\tau} \sim \mathscr{U}([-\frac{\delta}{2}, \frac{\delta}{2}]^m)$ be as described in Assumption 1, $x^{\star} \in \mathcal{K}$ for some convex symmetric $\mathcal{K} \subset \mathbb{B}_2^n$, and from the quantized observations $\dot{\boldsymbol{y}} = \mathcal{Q}_{\delta}(\boldsymbol{\Phi}\boldsymbol{x}^{\star} + \boldsymbol{\tau})$ we recover \boldsymbol{x}^{\star} by PBP as per (D.18). If $m \gtrsim (1 + \delta^{-2})\omega^2(\mathcal{K})$ with large enough implied constant, then for some small enough absolute constants c_0, c_1 , with probability exceeding $1 - \exp(-\omega^2(\mathcal{K})) - 12 \exp(-c_1 \mathscr{H}(\mathcal{K}, c_0 \delta))$ on a single draw of (Φ, τ) , the error rate

(D.30)
$$\|\hat{\boldsymbol{x}}_{\text{PBP}} - \boldsymbol{x}^{\star}\|_{2} \lesssim \delta^{\frac{1}{4}} \left(\frac{\omega^{2}(\mathcal{K})}{m} \log^{2}\left(\frac{\delta^{2}m}{\omega^{2}(\mathcal{K})}\right)\right)^{\frac{1}{8}} + (1+\delta)^{\frac{1}{2}} \left(\frac{\omega^{2}(\mathcal{K})}{m}\right)^{\frac{1}{4}}$$

holds uniformly for all $x^{\star} \in \mathcal{K}$.

Proof. We bound I_1 and I_2 in (D.26) separately.

Bounding I_1 Recall from (D.29) that $I_1 \leq 2\beta$ as long as the RIP in (D.27) holds. Thus, we only need to identify the value of β by bounding $\sup_{\boldsymbol{w}\in\mathcal{K}} |\frac{1}{m} \|\boldsymbol{\Phi}\boldsymbol{w}\|_2^2 - \|\boldsymbol{w}\|_2^2$. To achieve this, we set $\mathcal{T} = \mathcal{K} \times \{0\}$ to obtain that for any $t \geq 0$, the event $\sup_{\boldsymbol{w}\in\mathcal{K}} |\frac{\|\boldsymbol{\Phi}\boldsymbol{w}\|_2}{\sqrt{m}} - \|\boldsymbol{w}\|_2| \lesssim \frac{\omega(\mathcal{K})+t}{\sqrt{m}}$ holds with probability exceeding $1 - \exp(-t^2)$ (note that for symmetric \mathcal{K} we have $\omega(\mathcal{K}) = \gamma(\mathcal{K})$). Then, we set $t = \omega(\mathcal{K})$ to obtain

(D.31)
$$\sup_{\boldsymbol{w}\in\mathcal{K}} \left| \frac{\|\boldsymbol{\Phi}\boldsymbol{w}\|_2}{\sqrt{m}} - \|\boldsymbol{w}\|_2 \right| \lesssim \frac{\omega(\mathcal{K})}{\sqrt{m}}$$

with probability exceeding $1 - \exp(-\omega^2(\mathcal{K}))$. Note that $\frac{\omega(\mathcal{K})}{\sqrt{m}} = O(1)$, and hence we have $\sup_{\boldsymbol{w}\in\mathcal{K}} \frac{\|\boldsymbol{\Phi}\boldsymbol{w}\|_2}{\sqrt{m}} \leq \sup_{\boldsymbol{w}\in\mathcal{K}} |\frac{\|\boldsymbol{\Phi}\boldsymbol{w}\|_2}{\sqrt{m}} - \|\boldsymbol{w}\|_2 + \sup_{\boldsymbol{w}\in\mathcal{K}} \|\boldsymbol{w}\|_2 = O(1)$. Thus we have

(D.32)
$$\sup_{\boldsymbol{w}\in\mathcal{K}} \left| \frac{\|\boldsymbol{\Phi}\boldsymbol{w}\|_{2}^{2}}{m} - \|\boldsymbol{w}\|_{2}^{2} \right| \leq \sup_{\boldsymbol{w}\in\mathcal{K}} \left| \frac{\|\boldsymbol{\Phi}\boldsymbol{w}\|_{2}}{\sqrt{m}} - \|\boldsymbol{w}\|_{2} \right| \cdot \sup_{\boldsymbol{w}\in\mathcal{K}} \left| \frac{\|\boldsymbol{\Phi}\boldsymbol{w}\|_{2}}{\sqrt{m}} + \|\boldsymbol{w}\|_{2} \right| \lesssim \frac{\omega(\mathcal{K})}{\sqrt{m}}.$$

Combining with (D.29), we obtain that $I_1 \lesssim \frac{\omega(\mathcal{K})}{\sqrt{m}}$ with probability exceeding $1 - \exp(-\omega^2(\mathcal{K}))$.

Bounding I_2 We bound I_2 by Corollary D.3. As reformulated in Remark D.4, provided that $m \gtrsim (1 + \delta^{-2})\omega^2(\mathcal{K})$, Corollary D.3 implies that

(D.33)
$$I_2 \lesssim \sqrt{\delta} \left(\frac{\omega^2(\mathcal{K})}{m} \log^2\left(\frac{\delta^2 m}{\omega^2(\mathcal{K})}\right)\right)^{1/4} + \delta\left(\frac{\omega^2(\mathcal{K})}{m}\right)^{1/2}$$

holds with probability exceeding $1 - 12 \exp(-c_1 \mathscr{H}(\mathcal{K}, c_0 \delta))$ for some small enough c_0 (recall that $\rho \approx \frac{\delta \epsilon}{\sqrt{\log \epsilon^{-1}}}$ for small enough ϵ in Corollary D.3). Substituting $I_1 \lesssim \frac{\omega(\mathcal{K})}{\sqrt{m}}$ and (D.33) into (D.26) yields the claim.

Remark D.6. Under the regular scaling of $\delta \simeq 1$, our Proposition D.5 provides a uniform error rate of $\tilde{O}((\frac{\omega^2(\mathcal{K})}{m})^{1/8})$, which improves on $\tilde{O}((\frac{\omega^2(\mathcal{K})}{m})^{1/16})$ in [81] in the specific instance of sub-Gaussian Φ .

Remark D.7 (Improving the Non-Uniform Error Rate). While for general RIP matrix $\mathbf{\Phi}$ the non-uniform error rate (for recovering a fixed $\mathbf{x}^{\star} \in \mathcal{K}$) reads as $\tilde{O}((1+\delta)^{1/2}(\frac{\omega^2(\mathcal{K})}{m})^{1/8})$ [81, Sec. 7.3B], we note that a faster decaying rate of $O(m^{-1/4})$ can also be obtained if $\mathbf{\Phi}$ is sub-Gaussian:

UNIFORM GUARANTEES FOR QUANTIZED CORRUPTED SENSING

• First note that (D.20)–(D.26) bounds the recovery error for a fixed x^{\star} as

(D.34)
$$\|\hat{\boldsymbol{x}}_{\text{PBP}} - \boldsymbol{x}^{\star}\|_{2}^{2} \leq 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \left| \langle \boldsymbol{x}^{\star}, \boldsymbol{u} \rangle - \frac{1}{m} \langle \boldsymbol{\Phi} \boldsymbol{x}^{\star}, \boldsymbol{\Phi} \boldsymbol{u} \rangle \right|$$

(D.35)
$$+ 2 \sup_{\boldsymbol{u} \in \mathcal{K}} \frac{1}{m} \left| \langle \mathcal{Q}_{\delta}(\boldsymbol{\Phi} \boldsymbol{x}^{\star} + \boldsymbol{\tau}) - \boldsymbol{\Phi} \boldsymbol{x}^{\star}, \boldsymbol{\Phi} \boldsymbol{u} \rangle \right| := I_{3} + I_{4};$$

• Then, by (D.29) and (D.32) we obtain $I_3 \lesssim \frac{\omega(\mathcal{K})}{\sqrt{m}}$, and Lemma B.2 (local QPE, with $\boldsymbol{a} = \boldsymbol{x}^{\star}, \boldsymbol{b} = 0, \boldsymbol{\epsilon} = 0, \boldsymbol{\mathcal{E}} = \mathcal{K} \times \{0\}$) gives $I_4 \lesssim \frac{\delta \cdot \omega(\mathcal{K})}{\sqrt{m}}$, thus yielding the non-uniform rate $O((1+\delta)^{1/2}(\frac{\omega^2(\mathcal{K})}{m})^{1/4}).$

Comparing with the uniform rate $O(m^{-1/8})$ (D.30) provides such implication: when estimating signals living in a convex symmetric set, the cost of uniform recovery is essential. Moreover, we note that under a general non-linear model with Gaussian Φ , the non-uniform recovery of PBP has been systematically studied in [68], and and their rate also reads as $O(m^{-1/4})$ for the set of effectively sparse signals [68, Sec. 2.6] (this is the canonical example of bounded convex signal set).

Appendix E. A Table of Recurring Notation.

REFERENCES

- R. G. BARANIUK, M. A. DAVENPORT, R. A. DEVORE, AND M. B. WAKIN, A simple proof of the restricted isometry property for random matrices, Constructive Approximation, (2007).
- [2] H. H. BARRETT AND K. J. MYERS, Foundations of image science, John Wiley & Sons, 2013.
- [3] A. BERK, Deep generative demixing: Recovering Lipschitz signals from noisy subgaussian mixtures, arXiv preprint arXiv:2010.06652, (2020).
- [4] A. BERK, Deep generative demixing: Error bounds for demixing subgaussian mixtures of Lipschitz signals, in IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP), IEEE, 2021, pp. 4010–4014.
- [5] A. BORA, A. JALAL, E. PRICE, AND A. G. DIMAKIS, Compressed sensing using generative models, in International Conference on Machine Learning, PMLR, 2017, pp. 537–546.
- [6] S. BOUCHERON, G. LUGOSI, AND P. MASSART, Concentration inequalities: A nonasymptotic theory of independence, Oxford university press, 2013.
- [7] P. T. BOUFOUNOS AND R. G. BARANIUK, 1-bit compressive sensing, in Annual Conference on Information Sciences and Systems, IEEE, 2008, pp. 16–21.
- [8] P. T. BOUFOUNOS, L. JACQUES, F. KRAHMER, AND R. SAAB, Quantization and compressive sensing, in Compressed Sensing and its Applications: MATHEON Workshop 2013, Springer, 2015, pp. 193–237.
- [9] T. CAI AND W.-X. ZHOU, A max-norm constrained minimization approach to 1-bit matrix completion., J. Mach. Learn. Res., 14 (2013), pp. 3619–3647.
- [10] E. J. CANDÈS, X. LI, Y. MA, AND J. WRIGHT, Robust principal component analysis?, Journal of the ACM (JACM), 58 (2011), pp. 1–37.
- [11] E. J. CANDES AND Y. PLAN, Tight oracle inequalities for low-rank matrix recovery from a minimal number of noisy random measurements, IEEE Transactions on Information Theory, 57 (2011), pp. 2342–2359.
- [12] E. J. CANDÈS, J. ROMBERG, AND T. TAO, Robust uncertainty principles: Exact signal reconstruction from highly incomplete frequency information, IEEE Transactions on Information Theory, 52 (2006), pp. 489–509.
- [13] V. CHANDRASEKARAN, S. SANGHAVI, P. A. PARRILO, AND A. S. WILLSKY, Rank-sparsity incoherence for matrix decomposition, SIAM Journal on Optimization, 21 (2011), pp. 572–596.
- [14] J. CHEN AND Y. LIU, Stable recovery of structured signals from corrupted sub-gaussian measurements, IEEE Transactions on Information Theory, 65 (2018), pp. 2976–2994.

- [15] J. CHEN AND M. K. NG, A parameter-free two-bit covariance estimator with improved operator norm error rate, arXiv preprint arXiv:2308.16059, (2023).
- [16] J. CHEN AND M. K. NG, Uniform exact reconstruction of sparse signals and low-rank matrices from phase-only measurements, IEEE Transactions on Information Theory, 69 (2023), pp. 6739–6764.
- [17] J. CHEN, M. K. NG, AND D. WANG, Quantizing heavy-tailed data in statistical estimation: (near) minimax rates, covariate quantization, and uniform recovery, IEEE Transactions on Information Theory, (2023).
- [18] J. CHEN, J. SCARLETT, M. NG, AND Z. LIU, A unified framework for uniform signal recovery in nonlinear generative compressed sensing, in Thirty-seventh Conference on Neural Information Processing Systems, 2023, https://openreview.net/forum?id=vUXNNLatFv.
- [19] J. CHEN, C.-L. WANG, M. K. NG, AND D. WANG, High dimensional statistical estimation under uniformly dithered one-bit quantization, IEEE Transactions on Information Theory, 69 (2023), pp. 5151– 5187.
- [20] J. CHEN, Y. WANG, AND M. K. NG, Quantized low-rank multivariate regression with random dithering, IEEE Transactions on Signal Processing, 71 (2023), pp. 3913–3928.
- [21] Y. CHEN, A. JALALI, S. SANGHAVI, AND C. CARAMANIS, Low-rank matrix recovery from errors and erasures, IEEE Transactions on Information Theory, 59 (2013), pp. 4324–4337.
- [22] M. A. DAVENPORT, Y. PLAN, E. VAN DEN BERG, AND M. WOOTTERS, 1-bit matrix completion, Information and Inference: A Journal of the IMA, 3 (2014), pp. 189–223.
- [23] S. DIRKSEN, Dimensionality reduction with subgaussian matrices: a unified theory, Foundations of Computational Mathematics, 16 (2016), pp. 1367–1396.
- [24] S. DIRKSEN, Quantized compressed sensing: a survey, in Compressed Sensing and Its Applications: Third International MATHEON Conference 2017, Springer, 2019, pp. 67–95.
- [25] S. DIRKSEN AND J. MALY, Tuning-free one-bit covariance estimation using data-driven dithering, arXiv preprint arXiv:2307.12613, (2023).
- [26] S. DIRKSEN, J. MALY, AND H. RAUHUT, Covariance estimation under one-bit quantization, The Annals of Statistics, 50 (2022), pp. 3538–3562.
- [27] S. DIRKSEN AND S. MENDELSON, Non-gaussian hyperplane tessellations and robust one-bit compressed sensing, Journal of the European Mathematical Society, 23 (2021), pp. 2913–2947.
- [28] D. L. DONOHO, Compressed sensing, IEEE Transactions on Information Theory, 52 (2006), pp. 1289– 1306.
- [29] I. A. ELBAKRI AND J. A. FESSLER, Statistical image reconstruction for polyenergetic x-ray computed tomography, IEEE transactions on medical imaging, 21 (2002), pp. 89–99.
- [30] E. ELHAMIFAR AND R. VIDAL, Sparse subspace clustering: Algorithm, theory, and applications, IEEE Transactions on Pattern Analysis and Machine Intelligence, 35 (2013), pp. 2765–2781.
- [31] J. A. FESSLER, Model-based image reconstruction for mri, IEEE signal processing magazine, 27 (2010), pp. 81–89.
- [32] S. FOUCART, H. RAUHUT, S. FOUCART, AND H. RAUHUT, An invitation to compressive sensing, Springer, 2013.
- [33] R. FOYGEL AND L. MACKEY, Corrupted sensing: Novel guarantees for separating structured signals, IEEE Transactions on Information Theory, 60 (2014), pp. 1223–1247.
- [34] T. FUCHS, D. GROSS, P. JUNG, F. KRAHMER, R. KUENG, AND D. STÖGER, Proof methods for robust low-rank matrix recovery, in Compressed Sensing in Information Processing, Springer, 2022, pp. 37– 75.
- [35] M. GENZEL, High-dimensional estimation of structured signals from non-linear observations with general convex loss functions, IEEE Transactions on Information Theory, 63 (2016), pp. 1601–1619.
- [36] M. GENZEL AND A. STOLLENWERK, A unified approach to uniform signal recovery from nonlinear observations, Foundations of Computational Mathematics, (2022), pp. 1–74.
- [37] R. M. GRAY AND D. L. NEUHOFF, Quantization, IEEE transactions on information theory, 44 (1998), pp. 2325–2383.
- [38] R. M. GRAY AND T. G. STOCKHAM, Dithered quantizers, IEEE Transactions on Information Theory, 39 (1993), pp. 805–812.
- [39] S. GU, Q. XIE, D. MENG, W. ZUO, X. FENG, AND L. ZHANG, Weighted nuclear norm minimization and its applications to low level vision, International journal of computer vision, 121 (2017), pp. 183–208.

UNIFORM GUARANTEES FOR QUANTIZED CORRUPTED SENSING

- [40] O. A. HANNA, Y. H. EZZELDIN, C. FRAGOULI, AND S. DIGGAVI, Quantization of distributed data for learning, IEEE Journal on Selected Areas in Information Theory, 2 (2021), pp. 987–1001.
- [41] J. HAUPT, W. U. BAJWA, M. RABBAT, AND R. NOWAK, Compressed sensing for networked data, IEEE Signal Processing Magazine, 25 (2008), pp. 92–101.
- [42] L. JACQUES, J. N. LASKA, P. T. BOUFOUNOS, AND R. G. BARANIUK, Robust 1-bit compressive sensing via binary stable embeddings of sparse vectors, IEEE transactions on information theory, 59 (2013), pp. 2082–2102.
- [43] A. JALAL, M. ARVINTE, G. DARAS, E. PRICE, A. G. DIMAKIS, AND J. TAMIR, Robust compressed sensing MRI with deep generative priors, Advances in Neural Information Processing Systems, 34 (2021), pp. 14938–14954.
- [44] N. JAYANT AND L. RABINER, The application of dither to the quantization of speech signals, Bell System Technical Journal, 51 (1972), pp. 1293–1304.
- [45] H. JEONG, X. LI, Y. PLAN, AND O. YILMAZ, Sub-gaussian matrices on sets: Optimal tail dependence and applications, Communications on Pure and Applied Mathematics, 75 (2022), pp. 1713–1754.
- [46] H. C. JUNG, J. MALY, L. PALZER, AND A. STOLLENWERK, Quantized compressed sensing by rectified linear units, IEEE Transactions on Information Theory, 67 (2021), pp. 4125–4149.
- [47] A. KAMATH, E. PRICE, AND S. KARMALKAR, On the power of compressed sensing with generative models, in International Conference on Machine Learning, PMLR, 2020, pp. 5101–5109.
- [48] Y. LECUN, L. BOTTOU, Y. BENGIO, AND P. HAFFNER, Gradient-based learning applied to document recognition, Proceedings of the IEEE, 86 (1998), pp. 2278–2324.
- [49] X. LI, Compressed sensing and matrix completion with constant proportion of corruptions, Constructive Approximation, 37 (2013), pp. 73–99.
- [50] C. LIAW, A. MEHRABIAN, Y. PLAN, AND R. VERSHYNIN, A simple tool for bounding the deviation of random matrices on geometric sets, in Geometric Aspects of Functional Analysis, Springer, 2017, pp. 277–299.
- [51] J. LIMB, Design of dither waveforms for quantized visual signals, The Bell System Technical Journal, 48 (1969), pp. 2555–2582.
- [52] Z. LIU, S. GOMES, A. TIWARI, AND J. SCARLETT, Sample complexity bounds for 1-bit compressive sensing and binary stable embeddings with generative priors, in International Conference on Machine Learning, PMLR, 2020, pp. 6216–6225.
- [53] Z. LIU, J. LIU, S. GHOSH, J. HAN, AND J. SCARLETT, Generative principal component analysis, in ICLR, 2022.
- [54] Z. LIU, P. LUO, X. WANG, AND X. TANG, Deep learning face attributes in the wild, in Proceedings of the IEEE International Conference on Computer Vision, 2015, pp. 3730–3738.
- [55] Z. LIU AND J. SCARLETT, The generalized Lasso with nonlinear observations and generative priors, Advances in Neural Information Processing Systems, 33 (2020), pp. 19125–19136.
- [56] Z. LIU AND J. SCARLETT, Information-theoretic lower bounds for compressive sensing with generative models, IEEE Journal on Selected Areas in Information Theory, 1 (2020), pp. 292–303.
- [57] M. B. MCCOY AND J. A. TROPP, Sharp recovery bounds for convex demixing, with applications, Foundations of Computational Mathematics, 14 (2014), pp. 503–567.
- [58] S. MENDELSON, Upper bounds on product and multiplier empirical processes, Stochastic Processes and their Applications, 126 (2016), pp. 3652–3680.
- [59] S. NEGAHBAN AND M. J. WAINWRIGHT, Restricted strong convexity and weighted matrix completion: Optimal bounds with noise, The Journal of Machine Learning Research, 13 (2012), pp. 1665–1697.
- [60] S. N. NEGAHBAN, P. RAVIKUMAR, M. J. WAINWRIGHT, AND B. YU, A unified framework for highdimensional analysis of m-estimators with decomposable regularizers, Statistical science, 27 (2012), pp. 538–557.
- [61] N. H. NGUYEN AND T. D. TRAN, Robust Lasso with missing and grossly corrupted observations, IEEE Transactions on Information Theory, 59 (2012), pp. 2036–2058.
- [62] N. H. NGUYEN AND T. D. TRAN, Exact recoverability from dense corrupted observations via l₁minimization, IEEE Transactions on Information Theory, 59 (2013), pp. 2017–2035.
- [63] G. ONGIE, A. JALAL, C. A. METZLER, R. G. BARANIUK, A. G. DIMAKIS, AND R. WILLETT, Deep learning techniques for inverse problems in imaging, IEEE Journal on Selected Areas in Information Theory, 1 (2020), pp. 39–56.

- [64] S. OYMAK AND B. RECHT, Near-optimal bounds for binary embeddings of arbitrary sets, arXiv preprint arXiv:1512.04433, (2015).
- [65] Y. PLAN AND R. VERSHYNIN, Robust 1-bit compressed sensing and sparse logistic regression: A convex programming approach, IEEE Transactions on Information Theory, 59 (2012), pp. 482–494.
- [66] Y. PLAN AND R. VERSHYNIN, One-bit compressed sensing by linear programming, Communications on Pure and Applied Mathematics, 66 (2013), pp. 1275–1297.
- [67] Y. PLAN AND R. VERSHYNIN, The generalized Lasso with non-linear observations, IEEE Transactions on information theory, 62 (2016), pp. 1528–1537.
- [68] Y. PLAN, R. VERSHYNIN, AND E. YUDOVINA, High-dimensional estimation with geometric constraints, Information and Inference: A Journal of the IMA, 6 (2017), pp. 1–40.
- [69] S. QIU, X. WEI, AND Z. YANG, Robust one-bit recovery via ReLU generative networks: Near-optimal statistical rate and global landscape analysis, in International Conference on Machine Learning, PMLR, 2020, pp. 7857–7866.
- [70] T. M. QUAN, T. NGUYEN-DUC, AND W.-K. JEONG, Compressed sensing MRI reconstruction using a generative adversarial network with a cyclic loss, IEEE Transactions on Medical Imaging, 37 (2018), pp. 1488–1497.
- [71] A. RAJ, Y. LI, AND Y. BRESLER, Gan-based projector for faster recovery with convergence guarantees in linear inverse problems, in ICCV, 2019, pp. 5602–5611.
- [72] G. RASKUTTI, M. YUAN, AND H. CHEN, Convex regularization for high-dimensional multiresponse tensor regression, The Annals of Statistics, 47 (2019), pp. 1554–1584.
- [73] L. SCHUCHMAN, Dither signals and their effect on quantization noise, IEEE Transactions on Communication Technology, 12 (1964), pp. 162–165.
- [74] V. SHAH AND C. HEGDE, Solving linear inverse problems using GAN priors: An algorithm with provable guarantees, in ICASSP, IEEE, 2018, pp. 4609–4613.
- [75] Z. SUN, W. CUI, AND Y. LIU, Quantized corrupted sensing with random dithering, IEEE Transactions on Signal Processing, 70 (2022), pp. 600–615.
- [76] J. TACHELLA AND L. JACQUES, Learning to reconstruct signals from binary measurements alone, Transactions on Machine Learning Research, (2023), https://openreview.net/forum?id=ioFIAQOBOS. Featured Certification.
- [77] C. THRAMPOULIDIS AND A. S. RAWAT, The generalized Lasso for sub-gaussian measurements with dithered quantization, IEEE Transactions on Information Theory, 66 (2020), pp. 2487–2500.
- [78] R. VERSHYNIN, High-dimensional probability: An introduction with applications in data science, vol. 47, Cambridge university press, 2018.
- [79] B. WIDROW AND I. KOLLÁR, Quantization noise: roundoff error in digital computation, signal processing, control, and communications, Cambridge University Press, 2008.
- [80] J. WRIGHT, A. Y. YANG, A. GANESH, S. S. SASTRY, AND Y. MA, Robust face recognition via sparse representation, IEEE Transactions on Pattern Analysis and Machine Intelligence, 31 (2008), pp. 210– 227.
- [81] C. XU AND L. JACQUES, Quantized compressive sensing with RIP matrices: The benefit of dithering, Information and Inference: A Journal of the IMA, 9 (2020), pp. 543–586.
- [82] H. XU, C. CARAMANIS, AND S. MANNOR, Outlier-robust PCA: The high-dimensional case, IEEE Transactions on Information Theory, 59 (2012), pp. 546–572.
- [83] T. YANG, J. MALY, S. DIRKSEN, AND G. CAIRE, Plug-in channel estimation with dithered quantized signals in spatially non-stationary massive mimo systems, arXiv preprint arXiv:2301.04641, (2023).
- [84] G. YU, G. SAPIRO, AND S. MALLAT, Solving inverse problems with piecewise linear estimators: From gaussian mixture models to structured sparsity, IEEE Transactions on Image Processing, 21 (2011), pp. 2481–2499.
- [85] H. ZHANG, J. LI, K. KARA, D. ALISTARH, J. LIU, AND C. ZHANG, Zipml: Training linear models with end-to-end low precision, and a little bit of deep learning, in International Conference on Machine Learning, PMLR, 2017, pp. 4035–4043.

Table 1	
$Table \ of \ recurring \ notation.$	

Introduced in main text							
x^{\star}, n	Underlying signal and its dimension						
$\boldsymbol{v^{\star}},m$	Underlying corruption and its dimension						
$oldsymbol{\Phi}, oldsymbol{\Phi}_i^ op$	Sub-Gaussian sensing matrix and its i -th sensing vector (row)						
ϵ, E	Independent sub-Gaussian noise with sub-Gaussian norm bounded by E						
$\delta, oldsymbol{ au}, \mathcal{Q}_{\delta}(\cdot)$	Quantization resolution, uniform dither, uniform quantizer						
$oldsymbol{y}, oldsymbol{\dot{y}}$	Unquantized noisy measurements (3.1) , quantized measurements (3.2)						
$\xi_{a,b}$	Quantization noise associated with $(\boldsymbol{a}, \boldsymbol{b})$ (3.26)						
$f(\cdot), g(\cdot)$	Norms for promoting the structures of x^* and v^* (Assumption 2)						
$\mathcal{D}_f(oldsymbol{x}), \mathcal{D}_f^*(oldsymbol{x})$	Descent cone of f at \boldsymbol{x} and its normalized counterpart (2.1)						
$\mathscr{N}(\mathcal{K},arepsilon)$	The covering number of \mathcal{K} with radius ε under Euclidean distance						
$\mathscr{H}(\mathcal{K},arepsilon)$	The Kolmogorov entropy defined as $\mathscr{H}(\mathcal{K},\varepsilon) = \log \mathscr{N}(\mathcal{K},\varepsilon)$						
$\omega(\mathcal{K})$	The Gaussian width of $\mathcal{K} \subset \mathbb{R}^n$: $\omega(\mathcal{K}) = \mathbb{E}_{\boldsymbol{g} \sim \mathcal{N}(0, \boldsymbol{I}_n)} \sup_{\boldsymbol{x} \in \mathcal{K}} \boldsymbol{g}^\top \boldsymbol{x}$						
$\gamma(\mathcal{K})$	The Gaussian complexity of $\mathcal{K} \subset \mathbb{R}^n$: $\gamma(\mathcal{K}) = \mathbb{E}_{\boldsymbol{g} \sim \mathcal{N}(0, \boldsymbol{I}_n)} \sup_{\boldsymbol{x} \in \mathcal{K}} \boldsymbol{g}^\top \boldsymbol{x} $						
$\operatorname{rad}(\mathcal{K})$	The radius of \mathcal{K} : $\operatorname{rad}(\mathcal{K}) = \sup_{\boldsymbol{x} \in \mathcal{K}} \ \boldsymbol{x}\ _2$						
$\mathcal{K}_{loc}^{(ho)}$	The localized version of \mathcal{K} : $\mathcal{K}_{loc}^{(\rho)} = (\mathcal{K} - \mathcal{K}) \cap \mathbb{B}_2^n(\rho)$						
$\mathcal{D}_{oldsymbol{x}},\mathcal{D}^*_{oldsymbol{x}},\mathcal{D}_{oldsymbol{v}},\mathcal{D}^*_{oldsymbol{v}}$	Constraint sets for analyzing constrained Lasso $(3.10)-(3.11)$						
$\alpha_f(\mathcal{X})$	Compatibility constant between f and ℓ_2 -norm over set \mathcal{X} (3.44)						
Δ_x, Δ_v	Reconstruction error of the signal and the corruption						
$\alpha_{\boldsymbol{x}}, \alpha_{\boldsymbol{v}}$	Uniform bounds on $\alpha_f(\overline{\mathcal{X}}_{\boldsymbol{x}^{\star}})$ and $\alpha_g(\overline{\mathcal{V}}_{\boldsymbol{v}^{\star}})$ (Assumption 4)						
ρ_1, ρ_2, ζ	Parameters to be chosen in Theorem 3.2, Theorem 3.12						
$I_1, I_2, I_3,$ etc.	Random processes that we need to bound						
$\mathcal{C}(\lambda_1,\lambda_2)$	Constraint set for analyzing unconstrained Lasso (3.79)						
k, L, r, k', L', r'	Parameters for formulating generative priors (Assumption 5)						
Σ^n_s	The set of s -sparse n -dimensional vectors						
$M_r^{p,q}$	$M_r^{p,q}$ The set of $p \times q$ matrices with rank not exceeding r						
Introduced in appendices							
$\mathcal{K}_{oldsymbol{x}}^{-},\mathcal{K}_{oldsymbol{v}}^{-},\mathcal{E},\mathcal{E}^{*}$	Constraint sets for analyzing generative case (Proposition A.9)						
ρ_1, ρ_2	Covering radius for the covering arguments in the proof of Theorem B.3						
ζ	A parameter in $(0, \frac{\delta}{2})$ introduced in the proof of Theorem B.3						
$\mathcal{Z}_{a,b}$	"Bad" measurements suffering from discontinuity $(B.24)$						
$[\mathcal{J}_{oldsymbol{a}}^{\mathcal{A}},\mathcal{J}_{oldsymbol{b}}^{\mathcal{B}}]$	"Bad" measurements suffering from large perturbations (B.29)–(B.30)						
E_1, E_2, E_3	Events that aid the proof of Theorem B.3: (B.26), (B.34), (B.36)						
$\mathcal{U}_{\boldsymbol{a},\boldsymbol{b}}$	"Bad" measurements associated with $(\boldsymbol{a}, \boldsymbol{b})$ (B.39)						
U_0	Uniform upper bound on the cardinality of $\mathcal{U}_{\boldsymbol{a},\boldsymbol{b}}$ (B.41)						