

On Sparse Covers of Minor Free Graphs, Low Dimensional Metric Embeddings, and other applications

Arnold Filtser *
Bar-Ilan University¹

¹Bar-Ilan University

Abstract

Given a metric space (X, d_X) , a (β, s, Δ) -sparse cover is a collection of clusters $\mathcal{C} \subseteq P(X)$ with diameter at most Δ , such that for every point $x \in X$, the ball $B_X(x, \frac{\Delta}{\beta})$ is fully contained in some cluster $C \in \mathcal{C}$, and x belongs to at most s clusters in \mathcal{C} . Our main contribution is to show that the shortest path metric of every K_r -minor free graphs admits $(O(r), O(r^2), \Delta)$ -sparse cover, and for every $\epsilon > 0$, $(4 + \epsilon, O(\frac{1}{\epsilon})^r, \Delta)$ -sparse cover (for arbitrary $\Delta > 0$). We then use this sparse cover to show that every K_r -minor free graph embeds into $\ell_\infty^{\tilde{O}(\frac{1}{\epsilon})^{r+1} \cdot \log n}$ with distortion $3 + \epsilon$ (resp. into $\ell_\infty^{\tilde{O}(r^2) \cdot \log n}$ with distortion $O(r)$). Further, we provide applications of these sparse covers into padded decompositions, sparse partitions, universal TSP / Steiner tree, oblivious buy at bulk, name independent routing, and path reporting distance oracles.

Contents

1	Introduction	1	5.2	Embedding into ℓ_∞ from sparse partition cover scheme	17
1.1	Low dimensional metric embeddings into ℓ_∞	3	5.3	Removing the dependence on the aspect ratio	21
1.2	Padded Decompositions	4	5.4	Improved distortion for minor free graphs	23
1.3	Further Applications	4	6	From sparse covers to padded decompositions	25
1.4	Related Work	6	7	Further applications	28
2	Preliminaries	6	7.1	Sparse Partitions	28
3	Technical Overview	7	7.2	Universal TSP and Universal Steiner Tree	28
4	Sparse Covers for Minor Free Graphs	11	7.3	Oblivious Buy-at-Bulk	29
4.1	Buffered cop decomposition.	11	7.4	Name-Independent Routing	31
4.2	Additional properties of buffered cop decomposition	12	7.5	Path-Reporting Distance Oracle	32
4.3	Sparse Covers Construction	14	8	Conclusion and Open Problems	33
5	Metric Embedding into ℓ_∞ from Sparse Partition Cover	16	A	Proof of Lemma 5	41
5.1	Prefix Free Codes	17			

*Email: arnold.filtser@biu.ac.il. This research was supported by the ISRAEL SCIENCE FOUNDATION (grant No. 1042/22).

1 Introduction

Given a metric space (X, d_X) a (β, s, Δ) -sparse cover is a collection of clusters $\mathcal{C} \subseteq P(X)$ all with diameter at most Δ , such that every point belongs to at most s clusters, and every ball $B_X(x, \frac{\Delta}{\beta})$ is fully contained in some cluster $C \in \mathcal{C}$. Sparse covers are very useful for algorithmic design, and in particular for divide and conquer. Since their introduction by Awerbuch and Peleg [AP90b], sparse covers found numerous applications. A partial list of which includes: compact routing schemes [PU99, Pel00, TZ01, AGM⁺08, AGM05, AGMW10, BLT14], distant-dependent distributed directories [AP91, Pel93, Pel00, BLT14], network synchronizers [AW04, AP90a, Lyn96, Pel00, BLT14], distributed deadlock prevention [AKP94], construction of spanners and ultrametric covers [HIS13, FN22, LS23, HMO23, FL22a, FGN23], metric embeddings [Rao99, KLMN05], universal TSP and Steiner tree constructions [JLN⁺05, BDR⁺12, Fil20b, BCF⁺23], and Oblivious buy-at-bulk [SBI11].

We will study sparse covers in weighted graphs, where we distinguish between two different types of diameter. The *weak* diameter of a cluster $A \subseteq V$ in a weighted graph $G = (V, E, w)$ is the maximum pairwise distance $\max_{u, v \in A} d_G(u, v)$ w.r.t. the original shortest path distance, while the *strong* diameter is the maximum pairwise distance $\max_{u, v \in A} d_{G[A]}(u, v)$ in the induced subgraph. We continue with a formal definition:

Definition 1 (Sparse Cover). *Given a weighted graph $G = (V, E, w)$, a collection of clusters $\mathcal{C} = \{C_1, \dots, C_t\}$ is called a weak/strong (β, s, Δ) sparse cover if the following conditions hold.*

1. *Bounded diameter: The weak/strong diameter of every cluster $C_i \in \mathcal{C}$ is bounded by Δ .*
2. *Padding: For each $v \in V$, there exists a cluster $C_i \in \mathcal{C}$ such that $B_G(v, \frac{\Delta}{\beta}) \subseteq C_i$.*
3. *Sparsity: For each $v \in V$, there are at most s clusters in \mathcal{C} containing v .*

If the clusters \mathcal{C} can be partitioned into s partitions $\mathcal{P}_1, \dots, \mathcal{P}_s$ s.t. $\mathcal{C} = \cup_{i=1}^s \mathcal{P}_i$, then $\{\mathcal{P}_1, \dots, \mathcal{P}_s\}$ is called a weak/strong (β, s, Δ) sparse partition cover. We say that a graph G admits a weak/strong (β, s) sparse (partition) cover scheme, if for every parameter $\Delta > 0$ it admits a weak/strong (β, s, Δ) sparse (partition) cover that can be constructed in expected polynomial time. Sparse partition cover scheme is abbreviated SPCS.

The notion of sparse cover scheme is the more common in the literature. However, SPCS provides additional structure that is crucial for different applications. ¹ Obtaining strong diameter guarantee is also considerably more challenging, however it is frequently required in “subgraph based” applications. ²

Sparse covers were introduced by Awerbuch and Peleg [AP90b] who showed that for every $k \in \mathbb{N}$, every n vertex graph admits a strong $(4k - 2, 2k \cdot n^{\frac{1}{k}})$ -SPCS. Klein, Plotkin, and Rao [KPR93] constructed a celebrated weak padded decomposition ³ for K_r -minor free graphs with padding parameter $O(r^3)$ (later improved to $O(r^2)$ [FT03]). It is folklore that padded decomposition with padding parameter ρ implies a $(\rho, O(\log n))$ -SPCS (by taking the union of $O(\log n)$ independent partitions). Thus [KPR93, FT03] implied a weak $(O(r^2), O(\log n))$ -SPCS for K_r -minor free graphs.

¹In this paper we require the partition property of SPCS for metric embeddings into ℓ_∞ , and for the oblivious buy-at-bulk. This property is also crucial in the construction of ultrametric covers [FL22a, Fil23, FGN23].

²In this paper we use the strong diameter guarantee for routing, buy-at-bulk, and path reporting distance oracle.

³Roughly, padded decomposition with padding parameter ρ is a random partition into clusters of diameter at most Δ such that every ball of radius $\frac{\Delta}{\rho}$ is fully contained in a single cluster with probability at least $\frac{1}{2}$. See Definition 2.

Family	Padding	Sparsity	SPCS?	Diameter	Ref
General	$4k - 2$	$2k \cdot n^{\frac{1}{k}}$	yes	strong	[AP90b]
Planar	32	18	yes (implicit)	strong	[BLT14]
K_r -minor free	$O(r^3)$	$O(\log n)$	yes	weak	[KPR93]
	$O(r^2)$	$O(\log n)$	yes	weak	[FT03]
	$O(r)$	$O(\log n)$	yes	weak	[AGG ⁺ 19]
	$O(r)$	$O(\log n)$	yes	strong	[Fil19a]
	8	$f(r) \cdot \log n$ ⁽⁴⁾	yes (implicit)	strong	[BLT14]
	$O(r^2)$	2^r	yes	weak	[FT03, KLMN05]
	$O(r^2)$	$2^{O(r)} \cdot r!$	no	strong	[AGMW10]
	$O(r)$	$O(r^2)$	yes	strong	Theorem 1
$4 + \varepsilon$	$O(\frac{1}{\varepsilon})^r$	yes	strong		

Table 1: Summary of new and previous work on sparse covers.

Krauthgamer *et al.* [KLMN05] observed that one can use [KPR93, FT03] padded decomposition to construct a weak $(O(r^2), 2^r)$ -SPCS for K_r -minor free graphs (see [Fil20b] for an explicit proof). This was the first sparse cover where all the parameters are independent from the cardinality of the vertex set n . Busch *et al.* [BLT14] used shortest path separators to obtain a strong $(8, f(r) \cdot \log n)$ -SPCS. ⁴ Abraham *et al.* [AGMW10] constructed a strong sparse cover. Specifically, they made some adaptations to [KPR93] to obtain a strong $(O(r^2), 2^{O(r)} \cdot r!)$ -sparse cover scheme (not a SPCS). The final major piece in our story is a weak padded decomposition for K_r -minor free graphs with padding parameter $O(r)$ by Abraham *et al.* [AGG⁺19]. Which was later improved to strong padded decomposition with the same parameter by Filtser [Fil19a]. In particular, this implies a strong $(O(r), O(\log n))$ -sparse cover scheme. Interestingly, it was unknown how to turn this padded decomposition into a sparse cover with sparsity independent of n . Indeed, Filtser explicitly asked whether it is possible to construct $(O(r), g(r))$ -sparse cover scheme [Fil19a]. The main result of this paper is an affirmative answer to this question. A decade after the publication of [AGG⁺14] we finally managed to turn this padded decomposition into a sparse cover.

Theorem 1 (Cover for Minor Free Graphs). *Every K_r -minor free graph admits the following:*

- Strong $(O(r), O(r^2))$ -SPCS.
- For $\varepsilon \in (0, \frac{1}{2})$, strong $(4 + \varepsilon, O(\frac{1}{\varepsilon})^r)$ -SPCS.

The first bullet in Theorem 1 improves over the previous SPCS state of the art [FT03, KLMN05, Fil20b] in a threefold manner: (1) the padding is improved quadratically from $O(r^2)$ to $O(r)$, (2) the sparsity is improved exponentially from 2^r to $O(r^2)$, (3) the diameter guarantee is now strong. Alternately, the second bullet improves the stretch from $O(r^2)$ to $O(1)$ while keeping a similar sparsity. See Table 1 for a comparison of ours and previous work.

⁴ $f(r)$ is the enormous constant hiding in the Robertson Seymour [RS03] structure theorem. Johnson [Joh87] estimated $f(r) \geq 2 \uparrow (2 \uparrow (2 \uparrow \frac{r}{2}) + 3)$ where $2 \uparrow t$ is the exponential tower function ($2 \uparrow 0 = 1$ and $2 \uparrow t = 2^{2 \uparrow (t-1)}$).

1.1 Low dimensional metric embeddings into ℓ_∞

Metric embedding is a map between two metric spaces that approximately preserves pairwise distances. Given a (finite) metric space (X, d_X) , a map $\phi: V \rightarrow \mathbb{R}^k$, and a norm $\|\cdot\|$, the *contraction* and *expansion* of the map ϕ are the smallest $\xi, \rho \geq 1$, respectively, such that for every pair $x, y \in X$,

$$\frac{1}{\xi} \cdot d(x, y) \leq \|\phi(x) - \phi(y)\| \leq \rho \cdot d(x, y) \quad .$$

The *distortion* of the map is then $\xi \cdot \rho$. The expansion ρ is also called the Lipschitz constant of the embedding ϕ . Metric embeddings into norm spaces were thoroughly studied [Bou85, LLR95, Mat96, ABN11], and have a plethora of applications.

There is a special interest for metric embeddings into ℓ_∞ . From an algorithmic viewpoint, there is a significant advantage in the additional structure the norm space is providing. One example being nearest neighbor search (NNS) [Ind98, BG19], where ℓ_∞ enjoys succinct data structures. NNS for many other spaces works by first embedding the space into ℓ_∞ , and then using the ℓ_∞ NNS data structure to answer queries in the original metric space (see e.g. [FI99, Ind02]). From a geometric view point, ℓ_∞ is a special norm as it is a universal host metric space. That is, every finite metric spaces embeds isometrically (that is with distortion 1) into ℓ_∞ (the so called Fréchet embedding). However, there are n -point metric spaces of which every isometric embedding into ℓ_∞ requires $\Omega(n)$ dimensions [LLR95]. It is desirable to have low dimensional metric embeddings, as these are much more useful for algorithmic design. Matoušek [Mat96] showed that for every integer $t \geq 2$, every metric space embeds with distortion $2t - 1$ into ℓ_∞ of dimension $O(n^{1/t} \cdot t \cdot \log n)$ (which is almost tight assuming the Erdős girth conjecture [Erd64]). For distortion $O(\log n)$, Abraham *et al.* [ABN11] later improved the dimension to $O(\log n)$.

For more restrictive metric spaces better results are known. Linial *et al.* [LLR95] showed that every n -point tree metric embeds isometrically into $\ell_\infty^{O(\log n)}$, while Neiman [Nei16] showed that every metric space with doubling dimension d embeds into $\ell_\infty^{\varepsilon^{-O(d)} \cdot \log n}$ with distortion $1 + \varepsilon$. Krauthgamer *et al.* [KLMN05] showed that every n -point K_r -minor free graph G embeds into $\ell_\infty^{\tilde{O}(3^r \cdot \log n)}$ with distortion $O(r^2)$. Their embedding follows from a SPCS based on [KPR93, FT03]. However, their construction uses additional properties of that cover, and we cannot simply plug in our SPCS to get an improved embedding. We show that under very general conditions,⁵ one can use SPCS in a black box manner to obtain a metric embedding into ℓ_∞ (see Theorem 6). As a corollary, we improve both the distortion and dimension compared to [KLMN05]. We then slightly tailor the embedding for minor free graphs to push the distortion down all the way to $3 + \varepsilon$. See Table 2 for a comparison of new and old results.

Corollary 1. *Every n vertex K_r -minor free graph embeds into $\ell_\infty^{O(r^2 \cdot \log r \cdot \log n)}$ with distortion $O(r)$.*

Theorem 2. *For every $\varepsilon \in (0, \frac{1}{2})$, every n vertex K_r -minor free graph G can be embedded into $\ell_\infty^{O(\frac{1}{\varepsilon})^{r+1} \cdot \log \frac{1}{\varepsilon} \cdot \log \frac{n}{\varepsilon}}$ with distortion $3 + \varepsilon$.*

⁵In fact the reduction hold in general with no condition, while the dimension slightly increases. See Lemma 8.

Family	Distortion	Dimension	Ref
General Metric	1	$n - 1$	Fréchet
	$2k - 1$	$O(k \cdot n^{1/k} \cdot \log n)$	[Mat96]
	$O(\log n)$	$O(\log n)$	[ABN11]
Tree	1	$\Theta(\log n)$	[LLR95]
K_r -Minor Free	$O(r^2)$	$\tilde{O}(3^r) \cdot \log n$	[KLMN05]
	$O(r)$	$\tilde{O}(r^2) \cdot \log n$	Corollary 1
	$3 + \varepsilon$	$\tilde{O}(\frac{1}{\varepsilon})^{r+1} \cdot \log n$	Theorem 2

Table 2: Summary of new and previous work on metric embeddings into ℓ_∞ .

1.2 Padded Decompositions

Padded decomposition is a random partition of the vertices of a graph such that nearby vertices are likely to be clustered together. We begin with a formal definition:

Definition 2 (Padded Decomposition). *A distribution \mathcal{D} over partitions of a graph $G = (V, E, w)$ is strong (resp. weak) (β, δ, Δ) -padded decomposition if every $\mathcal{P} \in \text{supp}(\mathcal{D})$ is strongly (resp. weakly) Δ -bounded and for any $0 \leq \gamma \leq \delta$, and $z \in V$, $\Pr[B_G(z, \gamma\Delta) \subseteq P(z)] \geq e^{-\beta\gamma}$. We say that G admits (β, δ) -padded decomposition scheme if for every $\Delta > 0$, it admits a (β, δ, Δ) -padded decomposition.*

One can think of padded decompositions as bring “dual” to sparse partitions. Indeed, given a (β, δ, Δ) -padded decomposition, for $\gamma < \delta$, by taking the union of $e^{-\beta\gamma} \cdot O(\log n)$ independent samples from the padded decomposition we will obtain a $(\gamma, e^{-\beta\gamma} \cdot O(\log n))$ -SPCS. However, no similar statement in the other direction was ever made. Here we show that in general, given a sparse cover one can construct a padded decomposition (alas only with a weak diameter guarantee).

Theorem 3 (From sparse covers to padded decompositions). *Consider a weighted graph $G = (V, E, w)$ that admits a (β, s, Δ) -sparse cover. Then G admits weak $(O(\beta \cdot \log s), \frac{1}{4\beta}, \Delta)$ -padded decomposition.*

By plugging in our sparse cover from [Theorem 1](#) (with constant ε) we recover the state of the art result [\[AGG⁺19\]](#).

Corollary 2. *Every K_r -minor free graph admits a weak $(O(r), \Omega(1))$ -padded decomposition scheme.*

The proof in [\[AGG⁺19\]](#) is non-intuitive, and uses complicated and sophisticated probabilistic analysis based on a potential function. From the other hand, the proof of [Theorem 3](#) is short and elementary. Thus we obtain a simplified proof of this core-stone result. The only known lower bound on the padding parameter is $\Omega(\log r)$ which holds due to the fact that expanders are K_n -minor free and have padding parameter $\Omega(\log n)$ [\[Bar96\]](#). Abraham *et al.* [\[AGG⁺19\]](#) conjectured that K_r -minor free graphs admit padded decomposition with padding parameter $O(\log r)$. [Theorem 3](#) now provides a natural avenue for solving this conjecture: construct better covers!

1.3 Further Applications

Sparse partition and universal TSP / Steiner tree Given a weighted graph $G = (V, E, w)$, a (α, τ, Δ) -sparse partition is a partition \mathcal{C} of V into clusters with weak diameter at most Δ , such

that every ball of radius $\frac{\Delta}{\alpha}$ intersects at most τ clusters from \mathcal{C} . G admits a (α, τ) -sparse partition scheme if it admits (α, τ, Δ) -sparse partition for every $\Delta > 0$. Using our [Theorem 1](#), we construct $(O(r), O(r^2))$ -sparse partition scheme for K_r minor free graphs, improving over the previous state of the art of $(O(r^2), 2^r)$ -sparse partition scheme [[JLN⁺05](#), [Fil20b](#)]. See [Corollary 3](#) in [Section 7.1](#) for further details.

In the universal TSP problem, we are given a metric space (X, d_X) and the goal is to choose a single permutation π (a universal TSP) of X , such that given a subset $S \subseteq X$, we visit the points in S w.r.t. the order in π . This is the induced TSP tour by π . The permutation π has stretch ρ , if the length of the induced tour for every subset S is at most ρ times larger than the optimal tour for S . There is a general reduction from sparse partition scheme to universal TSP. Using this reduction and our sparse partitions ([Corollary 3](#)), given a shortest path metric of an n point K_r minor free graph, we construct universal TSP with stretch $O(r^4) \cdot \log n$ ([Corollary 4](#)), exponentially improving the dependence on r compared with previous results $(O(1)^r \cdot \log n)$. The same phenomena occurs also for the universal Steiner tree problem. See [Section 7.2](#) for further details.

Oblivious buy-at-bulk Given a weighted graph $G = (V, E, w)$ and a *canonical fusion function* $f : \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$ (see [Section 7.3](#) for definition), in the oblivious buy-at-bulk problem, the goal is to pick a route P_i for every possible demand pair $\delta_i = (s_i, t_i) \in \binom{V}{2}$. Then, given a specific set of demands $A = \{\delta_1, \dots, \delta_k\}$, the cost of our oblivious solution $\mathcal{P} = \{P_1, \dots, P_k\}$ is $\text{cost}(\mathcal{P}) = \sum_{e \in E} f(\varphi_e) \cdot w(e)$, where φ_e is the number of paths in \mathcal{P} using e . The solution is said to have approximation ratio ρ if for every subset of demands, the induced cost of the oblivious solution is at most ρ times the optimal solution. Due to the concavity of the canonical fusion function f , it is advantageous for the chosen paths to intersect as much as possible. Srinivasagopalan *et al.* [[SBI11](#)], implicitly showed that if a graph G admits a strong (β, s) -sparse cover scheme, where each cover is k -colorable (see [Definition 6](#)), then one can efficiently compute a solution for the oblivious buy-at-bulk problem with approximation ratio $O(s \cdot \beta^2 \cdot k \cdot \log n)$. We observe that our sparse cover scheme from [Theorem 1](#) is $O(r^2)$ -colorable, and conclude that K_r minor free graphs admit a solution with approximation ratio $O(r^6 \cdot \log n)$ (see [Corollary 5](#)), answering an open problem from [[SBI11](#)]. Note that nothing better for minor free graphs compared with general graphs was previously known. This is a previous covers were either not strong [[KPR93](#), [FT03](#)], or not colorable [[AGMW10](#), [AGG⁺19](#), [Fil19a](#)].

Name Independent Routing Here we are given an unweighted graph $G = (V, E)$ where the names (and ports) of all nodes are fixed. The goal is to design a compact routing scheme that will allow sending packages in the network, where routing decisions are made using small local routing tables, and the resulting routing paths are approximate shortest paths. This regime is considered more challenging and practical from the regime where nodes names and ports could be chosen by the routing scheme designer. Given a hereditary graph family that has α -orientation (see [Section 7.4](#)), where each graph possess strong (τ, β) -sparse cover scheme, Abraham *et al.* [[AGMW10](#)] constructed name independent compact routing scheme with stretch $O(\beta)$, $O(\log n + \log \tau)$ -bit headers and tables of $O(\frac{\log^3 n}{\log \log n} + \alpha \cdot \log n) \cdot \tau \cdot \log D$ bits. We use our strong sparse covers ([Theorem 1](#)) to construct name independent compact routing scheme significantly improving over previous work. See [Corollary 6](#), and [Table 3](#) for a comparison.

Path reporting distance oracle A path reporting distance oracle (PRDO) for a weighted graph $G = (V, E, w)$ is a succinct data structure that given a query $\{s, t\}$, efficiently returns an

approximate s - t shortest path P . We say that a PRDO has stretch k and query time t , if for every query (s, t) , the oracle returns a path P of weight at most $k \cdot d_G(s, t)$ in $O(|P|) + t$ time. PRDO were first studied for general graphs. Later, Elkin *et al.* [ENW16] constructed a PRDO for K_r -minor free graphs based on strong sparse covers. We plug in our strong sparse cover from Theorem 1 to obtain improvements in both space and stretch, see Corollary 7 (and Section 7.5).

1.4 Related Work

We provided background on sparse covers in the introduction. We refer to the cited papers for additional background on sparse partitions, UTSP, UST, routing and distance oracles. Here we provide additional background on metric embeddings in order to put our results (Corollary 1, Theorem 2) in a wider context. We begin with metric embeddings into ℓ_p spaces. Every n point metric space embeds into $\ell_2^{O(\log n)}$ with distortion $O(\log n)$ [Bou85], which is also tight [LLR95]. Planar graphs, and more generally fixed minor free graphs, embed into $\ell_2^{O(\log n)}$ with distortion $O(\sqrt{\log n})$ [Rao99, AFGN22], which is also tight [NR02]. The big open question here is regarding the embedding of such graphs into ℓ_1 . The upper bounds are the same as for ℓ_2 , while the only lower bound is 2 [LR10]. A long standing conjecture by Gupta *et al.* [GNRS04] states that every graph family excluding a fixed minor, and in particular planar graphs, can be embedded into ℓ_1 with constant distortion. Some partial progress for planar graph was made in the cases where we care only about vertices laying on a small number of faces [KLR19, Fil20a], or only about vertex pairs laying on the same face [OS81, Kum22].

Refined notion of distortion in metric embeddings were studied, such as scaling distortion [ABN11, BFN19b], and terminal/prioritized distortion [EFN17, EFN18]. In particular, there been study of prioritized low dimensional embeddings into ℓ_∞ [FGK23, EN22]. Online metric embeddings into normed spaces were also studied [IMSZ10, NR20, BFT24].

A different venue of research is metric embeddings into trees, or more generally low treewidth graphs. Every n -point metric space stochastically embeds into trees with expected distortion $O(\log n)$ [Bar96, FRT04]. This result is tight even when the metric is the shortest path metric of a planar graph [AKPW95]. Every planar graph with diameter Δ can be (deterministically) embedded into a graph with treewidth $\tilde{O}(\varepsilon^{-3})$ with additive distortion $\varepsilon \cdot \Delta$ [FKS19, FL22b, CCL⁺23]. Every K_r -minor free graphs stochastically embeds into graphs with treewidth $f(r) \cdot O(\frac{\log \log n}{\varepsilon})^2$ with expected additive distortion $\varepsilon \cdot \Delta$ [CFKL20, FL22b]. Clan embeddings and Ramsey-type embeddings of K_r -minor free graphs were also studied [FL22a]. Finally, recently it was shown that every K_r -minor free graphs stochastically embeds into graphs with treewidth $f(r) \cdot \tilde{O}(\frac{1}{\varepsilon}) \cdot \text{poly}(\log(n \cdot \Phi))$ with multiplicative expected distortion $1 + \varepsilon$ [CLPP23] (here Φ is the aspect ratio).

2 Preliminaries

\tilde{O} notation hides poly-logarithmic factors, that is $\tilde{O}(g) = O(g) \cdot \text{polylog}(g)$. All logarithms are at base 2 (unless specified otherwise), \ln stand for the natural logarithm. Given a set A , $\binom{A}{2} = \{\{x, y\} \mid x, y \in A, x \neq y\}$ denotes all the subsets of size 2. For a number $k \in \mathbb{N}$, $[k] = \{1, 2, \dots, k\}$.

We consider connected undirected graphs $G = (V, E, w)$ with edge weights $w : E \rightarrow \mathbb{R}_{\geq 0}$. We say that vertices v, u are neighbors if $\{v, u\} \in E$. Let d_G denote the shortest path metric in G . $B_G(v, r) = \{u \in V \mid d_G(v, u) \leq r\}$ is the closed ball of radius r around v . For a vertex $v \in V$ and a subset $A \subseteq V$, let $d_G(x, A) := \min_{a \in A} d_G(x, a)$, where $d_G(x, \emptyset) = \infty$. For a subset of vertices $A \subseteq V$,

$G[A]$ denotes the induced graph on A , and $G \setminus A := G[V \setminus A]$. The *diameter* of a graph G is $\text{diam}(G) = \max_{v,u \in V} d_G(v,u)$, i.e. the maximal distance between a pair of vertices. Given a subset $A \subseteq V$, the *weak-diameter* of A is $\text{diam}_G(A) = \max_{v,u \in A} d_G(v,u)$, i.e. the maximal distance between a pair of vertices in A , w.r.t. to d_G . The *strong-diameter* of A is $\text{diam}(G[A])$, the diameter of the graph induced by A .

The *aspect ratio* of a metric space (X, d_X) is usually defined as the ratio between the maximum and minimum distances. However, here we mainly work with the shortest path distance in graphs, where we allow 0-weights. Formally, here the shortest path metric is actually a pseudometric. Accordingly, we will define aspect ratio of a graph $G = (V, E, w)$ as the ratio between the maximum distance to the minimum non zero distance: $\Phi(G) = \frac{\max_{u,v \in V} d_G(u,v)}{\min_{u,v \in V \text{ s.t. } d_G(u,v) > 0} d_G(u,v)}$.

A graph H is a *minor* of a graph G if we can obtain H from G by edge deletions/contractions, and isolated vertex deletions. A graph family \mathcal{G} is *H-minor-free* if no graph $G \in \mathcal{G}$ has H as a minor. Some examples of minor free graphs are planar graphs (K_5 and $K_{3,3}$ minor-free), outer-planar graphs (K_4 and $K_{3,2}$ minor-free), series-parallel graphs (K_4 minor-free) and trees (K_3 minor-free).

The ℓ_∞ -norm of a vector $x = (x_1, \dots, x_k) \in \mathbb{R}^k$ is $\|x\|_\infty := \max_{i \in [k]} |x_i|$. An embedding from a metric space (X, d_X) into ℓ_∞ is a function $f : X \rightarrow \mathbb{R}^k$. The embedding f has distortion $c \cdot t$ if for every $x, y \in X$, $\frac{1}{c} \cdot d_X(x, y) \leq \|f(x) - f(y)\|_\infty \leq t \cdot d_X(x, y)$. t is the *expansion* (also known as a *Lipschitz constant*) of f , while c is that *contraction* of f . An embedding with distortion 1 (where $c = t = 1$) is called *isometric*. Embedding $f : X \rightarrow \ell_\infty^k$ can be viewed as a collection of embeddings $\{f_i\}_{i=1}^k$ into the line \mathbb{R} , where $f_i(x)$ equals to the i 'th coordinate of $f(x)$. We will also denote $(f(x))_i = f_i(x)$. Using this notation, f has expansion t if for every $x, y \in X$ and $i \in [k]$, $|f_i(x) - f_i(y)| \leq t \cdot d_X(x, y)$. Similarly, f has contraction c if for every $x, y \in X$ there is some $i \in [k]$ such that $|f_i(x) - f_i(y)| \geq \frac{1}{c} \cdot d_X(x, y)$. In this case, we will say that the pair x, y is *satisfied* by the coordinate i .

3 Technical Overview

Cop Decomposition. Abraham *et al.* [AGG⁺19] constructed a padded decomposition for K_r -minor free graphs based on the cops-and-robbers game [And86]. Fix the scale parameter $\Delta > 0$. The process works as follows: pick arbitrary x_1 , and let the ball $B_G(x_1, r_1)$ be the first cluster η_1 , where $r_1 \in [0, \Delta]$ is sampled using truncated exponential distribution. To construct the second cluster, pick an arbitrary connected component C_2 of $G \setminus \eta_1$, and arbitrary $x_2 \in C_2$. Let T_2 be a shortest path from x_2 to some vertex y with a neighbor in η_1 . Then the second cluster $\eta_2 = B_{G[C_2]}(T_2, r_2)$ is a ball around T_2 in the graph induced by the connected component, where the radius $r_2 \in [0, \Delta]$ is sampled using truncated exponential distribution. In general, suppose that we already constructed clusters $\eta_1, \dots, \eta_{k-1}$. Let C_k be an arbitrary connected component of $G \setminus \cup_{i < k} \eta_i$, and $x_k \in C_k$ arbitrary vertex. Let $\mathcal{K}_{C_k} \subseteq \{\eta_1, \dots, \eta_{k-1}\}$ be all the previously created clusters η_i , such that there is an edge from η_i to C_k . Let T_k be a shortest path tree in C_k , with x_k as a root, and such that for every $\eta_i \in \mathcal{K}_{C_k}$, there is an edge from a vertex in T_k to η_i . In particular, T_k will have at most $|\mathcal{K}_{C_k}|$ leaves. The k 'th cluster $\eta_k = B_{G[C_k]}(T_k, r_k)$ is a ball around T_k in the induced graph $G[C_k]$, where $r_k \in [0, \Delta]$ is sampled using truncated exponential distribution. See Figure 1 for illustration.

One can run the cop decomposition on general graphs without getting any interesting structure. What makes it particularly interesting for K_r -minor free graphs is the fact that the size of the set \mathcal{K}_{C_k} of neighboring previously created clusters is always bounded by $r - 2$. Indeed, one can argue that if $|\mathcal{K}_{C_k}| \geq r - 1$, than one can contract all the internal edges inside each cluster in \mathcal{K}_{C_k} , and

the connected component C_k , and obtain K_r as a minor. Thus T_k is a shortest path tree (w.r.t. $G[C_k]$) with at most $r - 2$ leaves, and the cluster C_k is a ball of radius at most Δ around this tree. We will call each such cluster a *supernode*, and the shortest path tree T_k its *skeleton*. In addition, we construct a tree \mathcal{T} over the supernodes. Here each supernode η_k will be the child in \mathcal{T} of the last created supernode $\eta_i \in \mathcal{K}_{C_k}$. Note that the only outgoing edges from η_k are either to its descendants or ancestors in \mathcal{T} . Furthermore, η_k have at most $r - 2$ “neighbor” ancestors. We will call the connected component C_k where we constructed the supernode η_k (with skeleton T_k) the *domain* of η_k , denoted $\text{dom}(\eta_k)$. Note that the vertices in $\text{dom}(\eta_k)$ will either belong to η_k , or to the descendants of η_k w.r.t. \mathcal{T} . See [Figure 1](#) for illustration.

Due to the truncated exponential distribution, Abraham *et al.* [[AGG⁺19](#)] showed that a small ball $B_G(x, \gamma\Delta)$ is likely to be fully contained in a single supernode.⁶ Unfortunately, the supernodes $\{\eta_i\}_{i \geq 1}$ do not have a bounded diameter. Nevertheless, following [[Fil19a](#)], using the skeleton T_k , one can partition each supernode η_k into clusters of diameter $O(\Delta)$, while cutting each small ball only with a small probability. Combining these two processes together, one obtains a strong $(O(r), \Omega(\frac{1}{r}), O(\Delta))$ -padded decomposition. However, it is unclear if it is possible to use the cop decomposition to create a sparse cover. Indeed, the “dependence tree” \mathcal{T} does not have a bounded depth, and the entire process looks very chaotic. Indeed, every small change in the sampling of the radii leads to a completely different outcome. In contrast, the [[KPR93](#)] (as well as [[AGMW10](#)]) clustering process had only a depth of r , and thus [[KLMN05](#)] enumerated all the possible choices in the process leading to a sparse cover of exponential sparsity.

Buffered Cop Decomposition. In a recent work, Chang *et al.* [[CCL⁺24](#)] obtained a new “separation”-property in the cop-decomposition. Instead of growing a ball with a random radius around the skeleton T_k , Chang *et al.* constructed the supernode T_K deterministically. The new separation property is the following, consider a supernode η , and a vertex $v \in \text{dom}(\eta)$ such that v belongs to a supernode η' , which is descendant of η , but there is no edge from η to η' . Then $d_{G[\text{dom}(\eta)]}(\eta, v) > \frac{\Delta}{r} = \gamma$. In other words, for every descendant η' of η , either they are neighbors, or every path in $\text{dom}(\eta)$ from η to η' is of length at least γ .⁷ Chang *et al.* called the new partition *Buffered Cop Decomposition*, because there is now a buffer between non-neighboring clusters. They used the new buffered cop decomposition to construct a *shortcut-partition* (which is a generalization of scattering partition [[Fil20b](#)]). Roughly, a shortcut-partition is a partition of the vertices into clusters of diameter at most $\varepsilon \cdot D$, such that for every pair of vertices u, v at distance at most D , there is an approximate shortest path going through at most $O_r(\frac{1}{\varepsilon})$ clusters. Chang *et al.* used their shortcut-partitions to construct tree covers [[BFN19a](#), [CCL⁺23](#)], distance oracles [[Tho04b](#), [Kle02](#), [AG06](#)], to solve the Steiner point removal problem [[Fil19b](#), [Fil20b](#), [KKN15](#), [Che18](#), [FKT19](#)], and to construct additive embedding of apex minor free graphs into low treewidth graphs [[FKS19](#), [CFKL20](#), [FL21](#), [FL22b](#), [CCL⁺23](#)].

Sparse Covers. The starting point of this paper is the buffered cop decomposition of [[CCL⁺24](#)]. We begin by observing some additional properties. Let \vec{G}_C be a digraph where the supernodes are

⁶Specifically, using a sophisticated argument, based on a potential function, Abraham *et al.* [[AGG⁺19](#)] showed that with probability at least $e^{-O(r)\gamma}$, the ball $B_G(x, \gamma\Delta)$ is fully contained in a single supernode.

⁷Roughly speaking, [[CCL⁺24](#)] begin with the supernode η_k being equal to the skeleton T_k . Then, as the algorithm progresses, each time the buffer property is violated we add the violating vertices to a previously created supernode. One can argue that the depth of this process is bounded by r (once for each neighboring ancestor supernode), and thus the cluster vertices are all within Δ distance from the skeleton.

the vertices, and there is a directed edge from a supernode η to its ancestor η' (w.r.t. \mathcal{T}) iff there is an edge between vertices in η and η' . \vec{G}_C is a DAG (directed acyclic graph) with maximum out-degree $r - 2$, and it has additional crucial property: if v has outgoing edges towards u, z than there has to be an edge between u and z (in one way or another). We call a graph with this property a *transitive DAG* (see Definition 4). In transitive DAG's, neighboring vertices tend to share many of their neighbors. Denote by $B_{\vec{G}_C}(\eta, q)$ the set of supernodes towards which there is a directed path from η of length at most q . We use the transitive DAG property to show that the size of $B_{\vec{G}_C}(\eta, q)$ is bounded by $\binom{r+q}{r}$ (Lemma 1). Note that crucially, for $q \geq r$, $\binom{r+q}{r} \approx O(q)^r$ the growth rate is sub-exponential in q . Next, we generalize the buffer property to argue that for every ancestor supernode η' of η such that $\eta' \notin B_{\vec{G}_C}(\eta, 2q+1)$ it holds that the distance from every vertex $v \in \eta$ to η' in $\text{dom}(\eta')$ is at least $(q+1) \cdot \frac{\Delta}{r}$ (Lemma 2). Combining these two properties together, it follows that for every vertex v there are at most $\binom{r+q}{r}$ ancestor supernodes at distance $q \cdot \frac{\Delta}{r}$.

Similar to the process of padded decomposition [AGG⁺19, Fil19a], our sparse cover is constructed in two steps. First we cover the vertices using *enlarged* supernodes, and then we separately cover each enlarged supernode. Fix $q \geq 0$, and for every supernode η , let $\hat{\eta} = B_{G[\text{dom}(\eta)]}(\eta, q \cdot \frac{\Delta}{r})$ be all the vertices at distance at most $q \cdot \frac{\Delta}{r}$ from η in $\text{dom}(\eta)$. In particular, a vertex $v \in \eta$ can join only to the enlarged supernodes which are ancestors of η at distance at most $q \cdot \frac{\Delta}{r}$. Consider the ball $B = B_G(v, \frac{q}{2} \cdot \frac{\Delta}{r})$, and let η be the first supernode containing some vertex from B . By the minimality of η , and the triangle inequality, it will follow that the ball $B \subseteq \hat{\eta}$ is contained in the enlarged supernode. From the other hand, due to the properties discussed above, each vertex v will belong to at most $\binom{r+q}{r}$ enlarged supernodes. Thus we get both the sparsity and the padding properties we wanted. The only missing property at this point is the bounded diameter. Next we cover each enlarged supernode $\hat{\eta}$ using the skeleton T_η . T_η consist of at most r shortest path. We go over these shortest paths, and choose a Δ -net N . Specifically, a set such that every two net points are at distance at least Δ , and every point has a net point at distance at most Δ . Fix $R = 2\Delta + q \cdot \frac{\Delta}{r}$. Due to the properties of shortest paths, every point has at most $O(r+q)$ net points at distance $2R$. Now taking all the balls of radius R around net points provides us the desired sparse cover for the enlarged supernode. Taking the union of all the covers for all the enlarged supernodes we obtain the sparse cover for the graph. Fixing $q = 1$ we obtain padding $O(r)$ and sparsity $O(r^2)$, while by taking $q = \Theta(\frac{r}{\varepsilon})$ we obtain padding $4 + \varepsilon$ and sparsity $O(\frac{1}{\varepsilon})^r$.

Metric embedding into ℓ_∞ . Our metric embedding into ℓ_∞ is based on our SPCS. The first to construct a sparse cover based metric embedding was Rao [Rao99], who used the [KPR93] padded decomposition to embed K_r minor free graphs into ℓ_2 with distortion $O(r^3 \cdot \sqrt{\log n})$. Given a partition \mathcal{P} , for a vertex v belonging to cluster $v \in C_v \in \mathcal{P}$, let $\partial_{\mathcal{P}}(v) = d_X(v, V \setminus C_v)$ be the distance between v to the boundary of the cluster C_v . Note that if v is padded $B_G(v, \frac{\Delta}{\beta}) \subseteq C_v$, then $\partial_{\mathcal{P}}(v) > \frac{\Delta}{\beta}$. For each cluster $C \in \mathcal{P}$, sample $\alpha_C \in \{\pm 1\}$ u.a.r. . For every vertex v , send v to $\alpha_{C_v} \cdot \partial_{\mathcal{P}}(v)$. By the triangle inequality, it follows that the expansion is at most 2. But for which vertex pairs can we guarantee small contraction?

Consider a pair u, v at distance $d_G(v, u) \in (\Delta, 2\Delta]$, and suppose that \mathcal{P} has diameter at most Δ , and v is padded in \mathcal{P} . Then u and v must belong to different clusters C_u, C_v . If it so happened that $\alpha_{C_u} \neq \alpha_{C_v}$, then $|\alpha_{C_v} \cdot \partial_{\mathcal{P}}(v) - \alpha_{C_u} \cdot \partial_{\mathcal{P}}(u)| = \partial_{\mathcal{P}}(v) + \partial_{\mathcal{P}}(u) \geq \frac{\Delta}{\beta} \geq \frac{d_G(u, v)}{2\beta}$, and we obtain a bound on the contraction! Rao took $O(\log n)$ independent samples of the coefficients $\{\alpha_C\}_{C \in \mathcal{P}}$, and gets the contraction guarantee in a constant fraction of the samples. Next, Rao also took

$O(\log n)$ independent partitions to get that v is padded in a constant fraction of them. Finally, Rao sampled partitions for all possible distance scales, concatenated the resulting embeddings of them all, and obtained the desired $O(r^3 \cdot \sqrt{\log n})$ distortion. Assuming all the distances are in $[1, \text{poly}(n)]$, there are $O(\log n)$ different distances scales, and the resulting dimension is $O(\log^3 n)$: one log for the number of scales, one log to sample many partitions for each scale, and one log to sample the coefficients $\{\alpha_C\}_{C \in \mathcal{P}}$. Nevertheless, in ℓ_2 , using dimension reduction [JL84], one can reduce the dimension to $O(\log n)$ without significantly increasing the distortion.

The focus of our paper is embeddings into ℓ_∞ . One can repeat Rao's embedding exactly as is, and get embedding into $\ell_\infty^{O(\log^3 n)}$ with distortion $O(r^3)$ (or $O(r)$ using the improved padding parameter from [AGG⁺19]). Unfortunately, there is no general dimension reduction in ℓ_∞ (ala [JL84]). Nevertheless, the dimension still can be dramatically improved. First, observe that when embedding into ℓ_∞ , we don't need to succeed on a constant fraction of the partitions (or the α coefficients), it is enough to be successful only once! Thus, instead taking $O(\log n)$ independent samples from a padded decomposition, one can use a SPCS. Indeed, Krauthgamer *et al.* [KLMN05] constructed an $(O(r^2), 3^r)$ -SPCS for K_r minor free graphs. Using this SPCS immediately leads to an embedding into $\ell_\infty^{\tilde{O}(3^r \cdot \log^2 n)}$ with distortion $O(r^2)$. To remove additional $\log n$ factor, [KLMN05] used an additional property of the [KPR93] based SPCS that does not hold in general: it is possible to create 3^r partitions such that for every pair u, v , there will be a single partition where both u and v will be padded **simultaneously**. [KLMN05] heavily relied on this property, while our SPCS (and actually all the others as well) lacking it. Hence we cannot apply [KLMN05] as is.

Our solution follow similar lines to [KLMN05], but avoids using the additional special structure of [KPR93]. Consider a graph G with a (β, τ) -SPCS. First, for every scale $\Delta_i = \rho^i$, for $\rho = O(\frac{\beta}{\varepsilon})$, create τ partitions $\mathcal{P}_i^1, \mathcal{P}_i^2, \dots, \mathcal{P}_i^\tau$, all with diameter Δ_i , and such that every vertex is β -padded in one of them (that is $\forall v, B_G(v, \frac{\Delta}{\beta})$ is contained in some cluster). Next, create, laminar partitions that closely resemble the original partitions. Specifically, we create $\{\tilde{\mathcal{P}}_i^1\}_{i \in \mathbb{Z}}, \{\tilde{\mathcal{P}}_i^2\}_{i \in \mathbb{Z}}, \dots, \{\tilde{\mathcal{P}}_i^\tau\}_{i \in \mathbb{Z}}$, where for every i, j , $\tilde{\mathcal{P}}_i^j$ refines \mathcal{P}_{i+1}^j , $\tilde{\mathcal{P}}_i^j$ has radius at most $(1+\varepsilon) \cdot \Delta_i$, and for every $v \in V$, and $i \in \mathbb{Z}$, $B_G(v, \frac{\Delta}{(1+\varepsilon)\beta})$ is fully contained in a cluster of one of $\tilde{\mathcal{P}}_i^1, \dots, \tilde{\mathcal{P}}_i^\tau$. In other words, for every $v \in V$, and $i \in \mathbb{Z}$, $\max_{j \in [\tau]} \partial_{\tilde{\mathcal{P}}_i^j}(v) > \frac{\Delta_i}{(1+\varepsilon)\beta}$. Similar laminar partitions were also created in [KLMN05].

Next, our goal is to embed w.r.t. each laminar partition $\{\tilde{\mathcal{P}}_i^j\}_{i \in \mathbb{Z}}$ independently, such that for every pair u, v which is separated in partition $\tilde{\mathcal{P}}_i^j$ it will hold that $\|f(u) - f(v)\|_\infty \geq \partial_{\tilde{\mathcal{P}}_i^j}(v) + \partial_{\tilde{\mathcal{P}}_i^j}(u)$.⁸ Note that given such embedding for all the laminar partitions, we can concatenate them all to obtain the desired distortion. Indeed, constant expansion follows by triangle inequality, while for every u, v , $\|f(u) - f(v)\|_\infty \geq \max_{i,j} \left(\partial_{\tilde{\mathcal{P}}_i^j}(v) + \partial_{\tilde{\mathcal{P}}_i^j}(u) \right) = \Omega\left(\frac{d_G(u,v)}{\beta}\right)$. To create the embedding w.r.t. to the laminar partition $\{\tilde{\mathcal{P}}_i^j\}_{i \in \mathbb{Z}}$, let $i_0 \in \mathbb{Z}$ be a the maximum index such that $\tilde{\mathcal{P}}_{i_0}^j$ is partition into singletons, and let $k \in \mathbb{N}$ be the maximum such that $\tilde{\mathcal{P}}_{i_0+k}^j$ is not the trivial partition into a single cluster $\{V\}$. Clearly it is enough to embed only w.r.t. the laminar partition $\{\tilde{\mathcal{P}}_i^j\}_{i=i_0}^{i_0+k}$. Let $\tilde{\mathcal{P}}_{i_0+k}^j = \{A_1, \dots, A_m\}$ be the clusters in the top partition in our hierarchy. We create a prefix free code $\alpha: \tilde{\mathcal{P}}_{i_0+k}^j \rightarrow \{\pm 1\}^*$. Specifically, each cluster A_q is assigned a string of ± 1 of length at most $2 \cdot \left\lceil \log \frac{|V|}{|A_q|} \right\rceil$. The strings of different clusters might be of different length. However, for every $A_q, A_{q'}$

⁸For comparison, [KLMN05] used the simultaneous padding property of [KPR93] and only guaranteed $\|f(u) - f(v)\|_\infty \geq \min\{\partial_{\tilde{\mathcal{P}}_i^j}(v), \partial_{\tilde{\mathcal{P}}_i^j}(u)\}$.

there is an index s such that $\alpha_s(A_q), \alpha_s(A_{q'})$ exist and differ. Then the embedding of each vertex $v \in A_q$ defined by concatenating the coordinates $\left(\alpha_1(A_q) \cdot \partial_{\tilde{\mathcal{P}}_{i_0+k}^j}(v), \alpha_2(A_q) \cdot \partial_{\tilde{\mathcal{P}}_{i_0+k}^j}(v), \dots\right)$ with an embedding created inductively for A_q w.r.t. the induced partition by $\{\tilde{\mathcal{P}}_i^j\}_{i=i_0}^{i_0+k-1}$. To bound the contraction, consider a pair u, v and suppose that they are first separated in level $k' \in [i_0, i_0 + k]$. Then the embeddings of u and v are “aligned” in scales $[k'+1, i_0+k]$, while in scale k' they belong to respective clusters $A_v, A_u \in \tilde{\mathcal{P}}_{k'}^j$. The respective codes $\alpha(A_v), \alpha(A_u)$ will differ in some coordinate s and thus $\|f(u) - f(v)\|_\infty \geq \left|\alpha_s(A_u) \cdot \partial_{\tilde{\mathcal{P}}_{k'}^j}(u) - \alpha_s(A_v) \cdot \partial_{\tilde{\mathcal{P}}_{k'}^j}(v)\right| = \partial_{\tilde{\mathcal{P}}_{k'}^j}(u) + \partial_{\tilde{\mathcal{P}}_{k'}^j}(v)$. To conclude a bound on the contraction it remains to observe that for every partition $\tilde{\mathcal{P}}_l^j$ that refines $\tilde{\mathcal{P}}_{k'}^j$ it holds that $\partial_{\tilde{\mathcal{P}}_{k'}^j}(v) + \partial_{\tilde{\mathcal{P}}_{k'}^j}(u) \geq \partial_{\tilde{\mathcal{P}}_l^j}(v) + \partial_{\tilde{\mathcal{P}}_l^j}(u)$.

The overall number of coordinates used for the embedding of the laminar partition $\{\tilde{\mathcal{P}}_i^j\}_{i=i_0}^{i_0+k}$ is $2 \cdot \lceil \log n \rceil + 2(k+1)$, where k have to be bounded by a logarithm of the aspect ratio $O(\log \Phi)$. If we started from a (β, τ) -SPCS, there are τ laminar partitions and thus the overall dimension is $O(\tau \cdot \log(n \cdot \Phi))$. We remove the dependence on the aspect ratio using fairly standard techniques. Specifically, by observing that for far enough scales, we can use the same coordinate. For this to hold, when treating scale ρ^i , we need to “contract” all vertex pairs at distance at most $\frac{\rho^i}{n^2}$ (as otherwise a pair can accumulate error in unbounded number of scales). Hence we can remove the dependence on the aspect ratio only if the contracted graph still admits a SPCS.

4 Sparse Covers for Minor Free Graphs

This section is devoted to proving a meta-theorem (Theorem 5) that given a buffered cop decomposition, constructs a sparse cover in a black box manner. Afterwards, our main Theorem 1 will follow as a corollary of this meta theorem, and the buffered cop decomposition of K_r minor free graphs from [CCL⁺24]. We begin in Section 4.1 with recalling the definition of buffered cop decomposition from [CCL⁺24]. Then, in Section 4.2 we prove some properties of buffered cop decomposition (mainly bounding the possible number of supernodes at a certain distance, see Lemmas 1 and 2). Finally, in Section 4.3, we construct a sparse cover from a buffered cop decomposition.

4.1 Buffered cop decomposition.

Let $G = (V, E, w)$ be a weighted graph. A *supernode* $\eta \subseteq V$ with *skeleton* T_η and *radius* Δ is an induced subgraph $G[\eta]$ of G containing a tree T_η where every vertex in η is within distance Δ of T_η w.r.t. induced shortest path metric, that is $\eta = B_{G[\eta]}(T_\eta, \Delta)$. A *buffered cop decomposition* for G is a partition of G into vertex-disjoint supernodes, together with a tree \mathcal{T} called the *partition tree*, whose nodes are the supernodes of G . For any supernode η , the *domain* $\text{dom}(\eta)$ denotes the subgraph induced by the union of all vertices in supernodes in the subtree of \mathcal{T} rooted at η . See Figure 1 for an illustration of supernodes, their domain, and a buffered cop decomposition.

Definition 3. A (Δ, γ, w) -buffered cop decomposition for G is a buffered cop decomposition \mathcal{T} that satisfies the following properties:

- [Supernode radius.] *Every supernode η has radius at most Δ .*
- [Shortest-path skeleton.] *For every supernode η , let \mathcal{A}_η be the set of ancestor supernodes η' of η such that there is an edge from $\text{dom}(\eta)$ to η' . Then $|\mathcal{A}_\eta| \leq w$. The skeleton T_η is an*




Figure 1: Illustration of an (Δ, γ, w) -buffered cop decomposition of the unweighted grid graph (left) together with the associated tree \mathcal{T} . There are 9 different supernodes η_1, \dots, η_9 , all colored with different colors. Each supernode η contains a shortest path tree T_η (the bold lines) with at most 3 leaves, where all the vertices $x \in \eta$ in the super node are at distance at most $\Delta = 6$ from T_η . The domain of each supernode consist of all the supernodes in its subtree. For example $\text{dom}(\eta_5) = G[\eta_5 \cup \eta_7 \cup \eta_8 \cup \eta_9]$, and $\text{dom}(\eta_3) = G[V \setminus (\eta_1 \cup \eta_2)]$. As η_3 and η_9 are not adjacent, the distance from η_3 to any vertex in η_9 (w.r.t. $\text{dom}(\eta_3)$) is at least γ .

*SSSP tree in $\text{dom}(\eta)$, with at most w leaves (the root is not counted). In particular, for every $\eta' \in \mathcal{A}_\eta$, there is an edge from T_η to η' .*⁹


- [Supernode buffer.] *Let η be a supernode, and let η' be another supernode that is an ancestor of η in the partition tree \mathcal{T} . Then either η and η' are adjacent in G , or for every vertex v in $\text{dom}(\eta)$, we have $d_{\text{dom}(\eta')}(v, \eta') > \gamma$.*

Chang *et al.* [CCL⁺24] constructed a buffered cop decomposition for K_r -minor free graphs based on the classic cop decomposition (see [And86, AGG⁺19, Fil19a]).

Theorem 4 (Buffered cop decomposition). *Let G be a K_r -minor-free graph, and let Δ be a positive number. Then G admits a $(\Delta, \Delta/r, r-1)$ -buffered cop decomposition (efficiently computable).*

4.2 Additional properties of buffered cop decomposition

Consider a (Δ, γ, w) -buffered cop decomposition \mathcal{T} of a graph G . In this subsection we will prove some additional properties of the buffered cop decomposition, that will later be useful in our construction of the sparse covers. Let $\vec{G}_\mathcal{C}$ be a directed graph with the supernodes \mathcal{C} as vertices. For a supernode η , we add a directed edge from η to any adjacent ancestor supernode η' . See illustration on the right for the digraph $\vec{G}_\mathcal{C}$ w.r.t. the decomposition from Figure 1. Note that $\vec{G}_\mathcal{C}$ is a DAG as all edges are directed towards the ancestors w.r.t. \mathcal{T} . $\vec{G}_\mathcal{C}$ has another crucial property: suppose that there are outgoing edges from η to η_1 and η_2 , where η_1 is an ancestor of η_2 in \mathcal{T} . Then there is an outgoing edge from η_2 to η_1 . Indeed, as η and η_1 are neighbors, there is an edge from $\text{dom}(\eta)$



⁹In the original definition of buffered cop decomposition in [CCL⁺24] there were no explicit requirement for edges between T_η to \mathcal{A}_η . However it holds in their construction.

to η_1 . As $\text{dom}(\eta) \subseteq \text{dom}(\eta_2)$, there is an edge from $\text{dom}(\eta_2)$ to η_1 . In particular, $\eta_1 \in \mathcal{A}_{\eta_2}$ and hence there is an edge from T_{η_2} (and in particular from η_2) to η_1 . We call such a graph a *Transitive DAG*.

Definition 4 (Transitive DAG). A digraph $\vec{G} = (V, \vec{E})$ is called *transitive DAG* if it contains no cycles, and for every $x, y, z \in V$ such that $(x, y), (x, z) \in \vec{E}$ it holds that either $(y, z) \in \vec{E}$ or $(z, y) \in \vec{E}$.

Another important property of \vec{G}_C is that the maximum out degree of a vertex is w . Indeed, this just follows from the definition of buffered cop decomposition. Give a digraph \vec{G} , and a vertex $v \in V$, the directed ball $B_{\vec{G}}(v, q)$ is the set of vertices u such that there is a directed path from v to u in \vec{G} of length at most q . v is called the center of the directed ball, while q is it's radius. We next bound the size of directed balls in a transitive DAG of bounded out degree.

Lemma 1. Let $\vec{G} = (V, \vec{E})$ be a transitive DAG of maximum out degree w . Then for every $v \in V$, and $q \in \mathbb{N}$, $|B_{\vec{G}}(v, q)| \leq \binom{w+q}{w}$.

Proof. The proof is by induction on q . The base case is when $q = 0$, and indeed every directed ball of radius 0 is of size $\binom{w+0}{w} = 1$ (containing only the center). Let $v_0, v_1, v_2, \dots, v_n$ be a topological ordering of \vec{G} vertices. That is for every edge $(v_i, v_j) \in \vec{E}$ it holds that $i < j$. Denote $v = v_{i_0}$. We can assume that v_{i_0} has exactly w outgoing edges, as otherwise we can just add artificial neighbors to the right of v_n , which will only increase the size of the directed ball. Denote by $v_{i_1}, v_{i_2}, \dots, v_{i_w}$ the endpoints of the w edges going out of v . For $s \in [0, w-1]$, let $I_s = \{v_{i_s}, v_{i_{s+1}}, \dots, v_{i_{s+1}-1}\}$ be the ‘‘interval’’ of vertices starting at v_{i_s} and ending just before $v_{i_{s+1}}$. Denote also $I_w = \{v_{i_w}, \dots, v_n\}$.

For $s \in [0, w]$, denote by $B_s = I_s \cap B_{\vec{G}}(v_{i_0}, q)$ the intersection of I_s with $B_{\vec{G}}(v_{i_0}, q)$. Clearly $B_s = B_{\vec{G}[I_s]}(v_{i_s}, q-1)$. That is, v_{i_s} is the only neighbor of v in I_s , and as \vec{G} is a DAG, all the paths from v_{i_0} to vertices in I_s must go through v_{i_s} . Next, we argue that for every vertex $u_j \in B_s$, u_j has outgoing edges towards $v_{i_{s+1}}, v_{i_{s+2}}, v_{i_w}$. The proof is by induction w.r.t. the topological order. The base case is v_{i_s} , and it holds by the transitive DAG rule as v_{i_0} has outgoing edges towards v_{i_s}, \dots, v_{i_w} . By induction, consider $u_j \in B_s$ and suppose that the last edge on its shortest paths from v_{i_s} to u_j is $(u_{j'}, u_j)$. By the inductive hypothesis, $u_{j'}$ has outgoing edges towards $v_{i_{s+1}}, \dots, v_{i_w}$. According to the rule of the transitive DAG, u_j also has outgoing edges towards $v_{i_{s+1}}, \dots, v_{i_w}$, as required. We conclude that the induced digraph $\vec{G}[B_s]$ has maximum outgoing degree of s (as the maximum out degree in \vec{G} is w and each such vertex has already $w-s$ outgoing edges out of B_s). By the inductive hypothesis, we conclude $|B_s| \leq \binom{s+(q-1)}{s}$.

For convenience denote $B_0 = \{v_{i_0}\}$ (one can think of it as the ball in the first interval, that contains only v_{i_0} and has out degree 0). Then $|B_0| = 1 = \binom{q-1}{0}$. We conclude

$$|B_{\vec{G}}(v, q)| = \sum_{s=0}^w |B_s| \leq \sum_{s=0}^w \binom{s+q-1}{s} \stackrel{(*)}{=} \binom{w+q}{w}.$$

To argue why equality $(*)$ holds we will use a story. Denote by A all the subsets of $[w+q]$ of size w . Clearly $|A| = \binom{w+q}{w}$. In addition, for $s \in [0, w]$ note by A_s all the subsets of $[w+q]$ where the maximum number not chosen is $q+s$. That is, every subset in A_s have to contain all the number in $\{q+s+1, w+s+2, \dots, w+q\}$, and in addition s number from $\{1, \dots, q+s-1\}$. That is $|A_s| = \binom{s+q-1}{s}$.

As $A = \cup_{s=0}^w A_s$, and all these sets are disjoint, it holds that $\binom{w+q}{w} = |A| = \sum_{s=0}^w |A_s| = \sum_{s=0}^w \binom{s+q-1}{s}$, proving quality $(*)$, and thus the induction and the lemma follows. \square

Next, we extend the supernode buffer property as a function of the distance in $\vec{G}_{\mathcal{C}}$.

Lemma 2. *Consider a vertex $v \in V$ that belongs to a supernode $v \in \eta \in \mathcal{C}$, and consider an ancestor supernode $\eta' \in \mathcal{C}$ of η such that $\eta' \notin B_{\vec{G}_{\mathcal{C}}}(\eta, 2q+1)$ for some $q \geq 0$. Then $d_{\text{dom}(\eta')}(v, \eta') > (q+1) \cdot \gamma$.*

Proof. The proof is by induction on q . The base case is when $q = 0$, As η' is an ancestor of η , $v \in \text{dom}(\eta')$. However, as $\eta' \notin B_{\vec{G}_{\mathcal{C}}}(\eta, 1)$, η and η' are not adjacent in G . By the definition of buffered cop decomposition, it follows that $d_{\text{dom}(\eta')}(v, \eta') > \gamma$.

For the inductive step, let P be the shortest path in $G[\text{dom}(\eta')]$ from v to η' . Let η_1, \dots, η_k be the supernodes in $B_{\vec{G}_{\mathcal{C}}}(\eta, 1)$. Let u be the first vertex along P out of $\text{dom}(\eta) \cup \cup_{i=1}^k \eta_i$. u belong to some supernode $\tilde{\eta}$. Note that $\tilde{\eta}$ is an ancestor of η . However, as $u \in \text{dom}(\eta')$ it holds that η' is an ancestor of $\tilde{\eta}$. Clearly, $\tilde{\eta} \notin B_{\vec{G}_{\mathcal{C}}}(\eta, 1)$. However, as there is an edge from a vertex in $\text{dom}(\eta) \cup \cup_{i=1}^k \eta_i$ to $\tilde{\eta}$, it holds that $d_{\vec{G}_{\mathcal{C}}}(\eta, \tilde{\eta}) = 2$, and hence $\tilde{\eta} \notin B_{\vec{G}_{\mathcal{C}}}(\eta', 2(q-1)+1)$. Using the induction hypothesis, it holds that $d_{\text{dom}(\eta')}(u, \eta') > q \cdot \gamma$. Using the base of the induction, $d_{\text{dom}(\tilde{\eta})}(v, u) > \gamma$. We conclude that $w(P) > q \cdot (\gamma + \gamma = q+1) \cdot \gamma$, as required. \square

4.3 Sparse Covers Construction

This subsection is devoted to proving a the following meta theorem, which given a buffered cop decomposition constructs a sparse cover. As a corollary, we will obtain our main [Theorem 4](#).

Theorem 5 (From buffered cop decomposition to sparse covers). *Consider a graph $G = (V, E, w)$ that admits a (Δ, γ, w) -buffered cop decomposition where $\gamma \leq \Delta$. Then for every $q \geq 1$, G admits a strong $\left(4 \cdot \left(\frac{2\Delta}{q\gamma} + 1\right), \binom{w+2q-1}{w} \cdot 4 \cdot w \cdot \left(2 + \frac{q\gamma}{\Delta}\right), 2 \cdot (2\Delta + q \cdot \gamma)\right)$ -sparse partition cover.*

Proof. Consider a (Δ, γ, w) -buffered cop decomposition \mathcal{T} of a graph G . Let q be the input parameter, and let $k = \binom{w+2q-1}{w}$. Consider two supernodes η, η' , where η' is an ancestor of η . We say that η and η' are *nearby* if $d_{G_{\mathcal{C}}}(\eta, \eta') \leq 2q - 1$. Note that by [Lemma 1](#), each supernode $\eta \in \mathcal{C}$ has at most k nearby nodes (including itself). We first partition the supernodes into sets $\mathcal{S}_1, \mathcal{S}_2, \dots, \mathcal{S}_k$ such that no two nearby supernodes belong to the same set. This can be done greedily w.r.t the order induced by \mathcal{T} . Specifically, when we construct \mathcal{S}_1 all the supernodes are active. The root η joins \mathcal{S}_1 . We mark η and all its nearby supernodes as inactive. Then all the maximal supernode w.r.t. the order induced by \mathcal{T} among the remaining active supernodes join \mathcal{S}_1 . We also mark the nearby supernodes of the newly joining supernodes as inactive. We continue with this process until no active supernodes remain. In general, to construct \mathcal{S}_i , we begin by marking all the supernodes in $\cup_{j=1}^{i-1} \mathcal{S}_j$ as inactive, and continue with the same process. That is, all the maximal supernodes η' (w.r.t. order induced by \mathcal{T} among the remaining active supernodes) join \mathcal{S}_i , we then mark all their nearby supernodes as inactive, and so on.

We argue that every supernode η joins one of $\mathcal{S}_1, \mathcal{S}_2, \dots, \mathcal{S}_k$. Indeed, the only way η might become inactive without joining \mathcal{S}_i is if one of its nearby ancestor supernodes joins \mathcal{S}_i . However, as η has only $k-1$ nearby ancestor clusters (not counting itself), if η was prevented from joining $\mathcal{S}_1, \mathcal{S}_2, \dots, \mathcal{S}_{k-1}$, it will necessarily join \mathcal{S}_k . Consider \mathcal{S}_i . For every supernode $\eta \in \mathcal{S}_i$, let $\hat{\eta} = B_{G[\text{dom}(\eta)]}(\eta, q \cdot \gamma)$ be the ball of radius $q \cdot \gamma$ around η in $\text{dom}(\eta)$. We argue the following:

Lemma 3. Consider the collection $\{\hat{\eta}\}_{\eta \in \mathcal{S}_1}, \dots, \{\hat{\eta}\}_{\eta \in \mathcal{S}_k}$. It holds that:

1. For every $i \in [k]$, $\{\hat{\eta}\}_{\eta \in \mathcal{S}_i}$ is a partial partition of V . That is for every $\eta_1, \eta_2 \in \mathcal{S}_i$, $\hat{\eta}_1 \cap \hat{\eta}_2 = \emptyset$.
2. For every $i \in [k]$ and $\eta \in \mathcal{S}_i$, $\hat{\eta}$ contains a skeleton T_η which is a shortest path tree in $G[\hat{\eta}]$ with at most w leafs.
3. For every vertex $v \in V$, there is some index $i \in [k]$ and supernode $\eta \in \mathcal{S}_i$ such that $d_{G[\hat{\eta}]}(v, T_\eta) \leq \Delta + \frac{q\gamma}{2}$, and $B_G(v, \frac{q\gamma}{2}) \subseteq \hat{\eta}$.

Proof. We begin by proving the first point. Consider $\eta_1, \eta_2 \in \mathcal{S}_i$. If η_1, η_2 does not have ancestry relationship, then $\text{dom}(\eta_1), \text{dom}(\eta_2)$ are disjoint, which implies $\hat{\eta}_1 \cap \hat{\eta}_2 = \emptyset$. We thus can assume w.l.o.g. that η_1 is an ancestor of η_2 . Suppose for the sake of contradiction that there is a vertex $v \in \hat{\eta}_1 \cap \hat{\eta}_2$. It holds that $v \in \text{dom}(\eta_1)$, and $d_{G[\text{dom}(\eta_1)]}(\eta_1, v) \leq q \cdot \gamma$. From the other hand, as $\eta_1, \eta_2 \in \mathcal{S}_i$ they are not nearby. In particular, $d_{G_C}(\eta_1, \eta_2) > 2q - 1$ which implies $\eta_1 \notin B_{\vec{G}}(\eta_2, 2q - 1)$. But by Lemma 2, $d_{G[\text{dom}(\eta_1)]}(\eta_1, v) \geq d_{G[\text{dom}(\eta_1)]}(\eta_1, \eta_2) > q \cdot \gamma$, a contradiction.

The second point follows directly from the construction. For the third point, let η be the first supernode w.r.t. the order induced by \mathcal{T} that intersects the ball $B_G(v, \frac{q\gamma}{2})$. Suppose that $\eta \in \mathcal{S}_i$. By the minimality of η , it holds that $B_G(v, \frac{q\gamma}{2}) \subseteq \text{dom}(\eta)$. In particular, as η intersects $B_G(v, \frac{q\gamma}{2})$, by triangle inequality it follows that $d_{G[\hat{\eta}]}(v, T_\eta) \leq \Delta + \frac{q\gamma}{2}$, and $B_G(v, \frac{q\gamma}{2}) \subseteq \hat{\eta}$. \square

Next we argue that each enlarged supernode can be efficiently clustered.

Lemma 4. Consider an enlarged supernode $\hat{\eta}$. There is set of partitions $\mathcal{C}_1^{\hat{\eta}}, \dots, \mathcal{C}_q^{\hat{\eta}}$ of size $s \leq 4w \cdot (2 + \frac{q\gamma}{\Delta})$ with strong diameter $2 \cdot (2\Delta + q \cdot \gamma)$ such that for every vertex $v \in \hat{\eta}$ at distance at most $\Delta + \frac{q\gamma}{2}$ from T_η , the ball $B_{G[\hat{\eta}]}(v, \frac{q\gamma}{2})$ is fully contained in some cluster in some partition.

Proof. Let $N \subseteq T_\eta$ be an Δ -net. That is, a set of points in T_η at pairwise distance greater than Δ , and such that for every point $v \in T_\eta$, there is some net point $u \in N$ at distance at most $d_{G[T_\eta]}(u, v) \leq \Delta$. Such a set can be constructed greedily. We say that a pair of net points $u, v \in N$ is nearby if $d_{G[\hat{\eta}]}(v, u) \leq 2 \cdot (2\Delta + q \cdot \gamma)$.

Claim 1. Every net point $v \in N$ has at most $s = 4w \cdot (2 + \frac{q\gamma}{\Delta})$ nearby net points (including v).

Proof. T_η is a shortest path tree with at most w leafs. In particular, T_η consist of the shortest paths P_1, \dots, P_k for $k \leq w$. Fix some P_i . Let u_1, u_2, \dots, u_t be the net points along P_i which are nearby v , sorted w.r.t. their position along P_i . As P_i is a shortest path, and the pairwise distance between every pair of consecutive net points is at least Δ , we have that $d_{G[\hat{\eta}]}(u_1, u_t) > (t - 1) \cdot \Delta$. From the other hand, as both u_1, u_t are nearby v , by the triangle inequality it holds that $d_{G[\hat{\eta}]}(u_1, u_t) \leq d_{G[\hat{\eta}]}(u_1, v) + d_{G[\hat{\eta}]}(v, u_t) \leq 4 \cdot (2\Delta + q \cdot \gamma)$. It follows that $(t - 1) \cdot \Delta < 4 \cdot (2\Delta + q \cdot \gamma)$ and thus $t \leq 4 \cdot (2 + \frac{q\gamma}{\Delta})$. The claim now follows as there are at most w such paths. \square

Partition N into subsets N_1, \dots, N_s such that no two nearby points will belong to the same set N_i (this can be done in a greedy manner). For every subset N_i , consider the partition with the clusters $\{B_{G[\hat{\eta}]}(v, 2\Delta + q \cdot \gamma) \mid v \in N_i\}$, and the rest of $\hat{\eta}$ vertices being singletons. Note that this is indeed a partition as N_i does not contain nearby net points and hence all this balls are disjoint. Clearly the strong diameter of each cluster is at most $2 \cdot (2\Delta + q \cdot \gamma)$. Finally, consider a vertex $v \in \hat{\eta}$ such that there is $x \in T_\eta$ with $d_{G[\hat{\eta}]}(v, x) \leq \Delta + \frac{q\gamma}{2}$. Let $y \in N$ be the closest net point to x . By the triangle inequality $d_{G[\hat{\eta}]}(v, y) \leq d_{G[\hat{\eta}]}(v, x) + d_{G[\hat{\eta}]}(x, y) \leq 2\Delta + \frac{q\gamma}{2}$. Let i be an index such that

$y \in N_i$. It follows that the ball $B_{G[\hat{\eta}]}(y, 2\Delta + q \cdot \gamma)$, which belongs to the respective partition of N_i contains $B_{G[\hat{\eta}]}(v, \frac{q\gamma}{2})$ as required. \square

Our final set of partitions is constructed naturally. We first use [Lemma 3](#) to obtain the collections $\{\hat{\eta}\}_{\eta \in \mathcal{S}_1}, \dots, \{\hat{\eta}\}_{\eta \in \mathcal{S}_k}$ of enlarged supernodes. Then for every $i \in [k]$, and every $\hat{\eta} \in \mathcal{S}_i$, we use [Lemma 4](#) to obtain partitions $\mathcal{C}_1^{\hat{\eta}}, \dots, \mathcal{C}_s^{\hat{\eta}}$. Finally, for every $i \in [k]$, and $j \in [s]$, we create the partition $\left\{ \mathcal{C}_j^{\hat{\eta}} \right\}_{\eta \in \mathcal{S}_i}$ of G , and add all the non clustered vertices as singletons. Note that we obtained $k \cdot s = \binom{w+2q-1}{w} \cdot 4 \cdot w \cdot \left(2 + \frac{q\gamma}{\Delta}\right)$ partitions. Clearly, each cluster in every partition has strong diameter at most $2 \cdot (2\Delta + q \cdot \gamma)$. Finally, we prove the cover property. Consider a vertex $v \in V$. By [Lemma 3](#) there is some index $i \in [k]$ and supernode $\eta \in \mathcal{S}_i$ such that $d_{G[\hat{\eta}]}(v, T_\eta) \leq \Delta + \frac{q\gamma}{2}$, and $B_G(v, \frac{q\gamma}{2}) \subseteq \hat{\eta}$. By [Lemma 4](#), the ball $B_{G[\hat{\eta}]}(v, \frac{q\gamma}{2}) = B_G(v, \frac{q\gamma}{2})$ is fully contained in some cluster in some partition. Thus our padding parameter is $\frac{2 \cdot (2\Delta + q\gamma)}{\frac{q\gamma}{2}} = 4 \cdot \left(\frac{2\Delta}{q\gamma} + 1\right)$, as required. \square

We are now ready to prove our main [Theorem 1](#) (restated for convenience).

Theorem 1 (Cover for Minor Free Graphs). *Every K_r -minor free graph admits the following:*

- *Strong $(O(r), O(r^2))$ -SPCS.*
- *For $\varepsilon \in (0, \frac{1}{2})$, strong $(4 + \varepsilon, O(\frac{1}{\varepsilon})^r)$ -SPCS.*

Proof. According to [[CCL⁺24](#)] ([Theorem 4](#)), every K_r minor free graph G admits $(\Delta, \Delta/r, r-1)$ -buffered cop decomposition for every $\Delta > 0$. Using [Theorem 5](#) it follows that G admits strong $\left(4 \cdot \left(\frac{2r}{q} + 1\right), \binom{r+2q-2}{r-1} \cdot 4 \cdot r \cdot \left(2 + \frac{q}{r}\right), 2 \cdot \left(2 + \frac{q}{r}\right) \cdot \Delta\right)$ -sparse partition cover, for arbitrary $\Delta > 0$. It follows that every K_r minor free graph admits strong $\left(4 \cdot \left(\frac{2r}{q} + 1\right), \binom{r+2q-2}{r-1} \cdot 4 \cdot r \cdot \left(2 + \frac{q}{r}\right)\right)$ -SPCS.

Fixing $q = 1$ we obtain that every K_r minor free graph admits strong $\left(4 \cdot (2r + 1), \binom{r}{r-1} \cdot 4 \cdot r \cdot \left(2 + \frac{1}{r}\right)\right) = (O(r), O(r^2))$ -SPCS. Next, fix $q = \frac{8r}{\varepsilon}$, using the identity $\binom{n}{k} \leq \left(\frac{n \cdot e}{k}\right)^k$, we have $\binom{r + \frac{16r}{\varepsilon} - 2}{r-1} \cdot 4 \cdot r \cdot \left(2 + \frac{8}{\varepsilon}\right) \leq \left(\frac{(r + \frac{16r}{\varepsilon} - 2) \cdot e}{r-1}\right)^{r-1} \cdot O\left(\frac{r}{\varepsilon}\right) = O\left(\frac{1}{\varepsilon}\right)^r$, and thus G admits a strong $(4 + \varepsilon, O(\frac{1}{\varepsilon})^r)$ -SPCS. The theorem follows by scaling ε accordingly. \square

5 Metric Embedding into ℓ_∞ from Sparse Partition Cover

The main result of this section is a meta theorem ([Theorem 6](#)) that transforms a SPCS into a low distortion/dimension metric embedding into ℓ_∞ . This reduction is applicable for graph families which are closed under re-weighting (abbreviated CURW). A graph family \mathcal{F} is CURW if for every weighted graph $G = (V, E, w) \in \mathcal{F}$ in the family, it holds that for every weight function $w' : E \rightarrow \mathbb{R}_{\geq 0}$, the reweighted graph $G' = (V, E, w') \in \mathcal{F}$ is also in the family. Many graph families are CURW, some examples being general graphs, planar graphs, graphs with bounded treewidth \setminus pathwidth \setminus genus, and most importantly for us, H minor free graphs. There are also well studied families which are not CURW, such as graphs with bounded doubling dimension, or bounded highway dimension.

Theorem 6 (From SPCS to Metric embedding). *Consider a CURW graph family \mathcal{F} , such that every $G \in \mathcal{F}$ admits a (β, τ) -SPCS. Then for every $\varepsilon > 0$, every n -point graph $G = (V, E, w) \in \mathcal{F}$ admits an efficiently computable embedding $f : V \rightarrow \ell_\infty^k$ with distortion $(1 + \varepsilon) \cdot 2\beta$ and dimension $k = O\left(\frac{\tau}{\varepsilon} \cdot \log \frac{\beta}{\varepsilon} \cdot \log\left(\frac{n \cdot \beta}{\varepsilon}\right)\right)$.*

Using our [Theorem 1](#) with $\varepsilon = 1$, [Corollary 1](#) follows. We restate it for convenience:

Corollary 1. *Every n vertex K_r -minor free graph embeds into $\ell_\infty^{O(r^2 \cdot \log r \cdot \log n)}$ with distortion $O(r)$.*

One can also use our $(4 + \varepsilon, O(\frac{1}{\varepsilon})^r)$ -SPCS to obtain embedding into $\ell_\infty^{O(\frac{1}{\varepsilon})^{r+1} \cdot \log \frac{1}{\varepsilon} \cdot \log \frac{n}{\varepsilon}}$ with distortion $8 + \varepsilon$. However, for the case of minor-free graphs, we open the black box of [Theorem 6](#) and obtain an improved embedding with distortion $3 + \varepsilon$ (see [Theorem 2](#)).

The plan for the section is as follows: first in [Section 5.1](#) we construct a prefix free code, which will later be used in the construction of the embedding. Then, in [Section 5.2](#) we construct a reduction from SPCS into metric embedding (see [Lemma 8](#)). Note that there is no assumption of a CURW graph family here. The resulting dimension is $O\left(\frac{\tau}{\varepsilon} \cdot \log \frac{\beta}{\varepsilon} \cdot \log(n\Phi)\right)$, where $\Phi = \frac{\max_{u,v} d_G(u,v)}{\min_{u,v} d_G(u,v)}$ is the aspect ratio. Then in [Section 5.3](#) we use [Lemma 8](#) to obtain our main [Theorem 6](#). Finally, in [Section 5.4](#) we improve the result for minor free graph to obtain distortion $3 + \varepsilon$.

5.1 Prefix Free Codes

This sub-section is devoted to the construction of prefix free codes. $\{\pm 1\}^*$ stands for all finite sequences of $\{\pm 1\}$.

Definition 5. *A code for elements X is a function $h : X \rightarrow \{\pm 1\}^*$. The code is prefix free if for every $x, y \in X$, $h(x)$ and $h(y)$ are not the prefixes of each other. That is, there is no $L \in \{\pm 1\}^*$ such that either $h(x) \circ L = h(y)$, or $h(y) \circ L = h(x)$.*

In the following lemma we construct a prefix free code where the length of the code word of x is bounded by $2 \cdot \left\lceil \log \frac{\mu(X)}{\mu(x)} \right\rceil$. This statement is not new, and even better prefix free codes are known (see e.g. [\[KN76\]](#)). Nevertheless, we provide a proof of [Lemma 5](#) in [Appendix A](#) for the sake of completeness.

Lemma 5. *Consider a set X with a weight function $\mu : X \rightarrow \mathbb{R}_{>0}$. Denote $\mu(X) = \sum_{x \in X} \mu(x)$. Then there is a prefix free code h , such that for every $x \in X$, $|h(x)| \leq 2 \cdot \left\lceil \log \frac{\mu(X)}{\mu(x)} \right\rceil$.*

5.2 Embedding into ℓ_∞ from sparse partition cover scheme

In this subsection, given a SPCS, we construct a metric embedding into ℓ_∞ , where the dimension depends on the aspect ratio ([Lemma 8](#)). We begin this subsection by showing that we can turn a collection of unrelated partitions at different scales into laminar partitions ([Lemma 6](#)). This statement was implicitly proved in [\[KLMN05\]](#), as well as in other constructions. The following lemma is an adaptation from [\[FL22a\]](#). We provide a proof for the sake of completeness.

Lemma 6. *Consider a finite metric space (X, d_X) that admits a (β, τ) -sparse partition cover scheme. Then for every $a > 0$, and $\varepsilon \in (0, 1)$ there is a collection of hierarchical partitions of X : $\{\mathcal{P}_q^i\}_{q \in \mathbb{Z}, i \in [\tau]}$ with the following properties:*

1. For each j , $\{\mathcal{P}_i^j\}_{i \in \mathbb{Z}}$ is a hierarchical partition. That is $\forall i, j$, \mathcal{P}_i^j is a refinement of \mathcal{P}_{i+1}^j .
2. Fix $\Delta_i = a \cdot (\frac{4\beta}{\varepsilon})^i$. For every j and $i \in \mathbb{Z}$, \mathcal{P}_i^j has diameter $(1 + \varepsilon) \cdot \Delta_i$.
3. For every $x \in X$ and $i \in \mathbb{Z}$, there is some $j \in [\tau]$ such that the ball $B_X(x, \frac{\Delta_i}{\beta(1+\varepsilon)})$ is fully contained in some cluster of \mathcal{P}_i^j .

Proof. Assume w.l.o.g. that the minimal pairwise distance in X is 1 (otherwise one can scale accordingly), while the maximal pairwise distance is Φ . Set $I = \lceil \log_{4\beta/\epsilon} \Phi/a \rceil$. For $i \in [0, I]$, let $\mathbb{P}_i = \{\mathcal{P}_i^1, \dots, \mathcal{P}_i^\tau\}$ be a (β, τ, Δ_i) -sparse partition cover (we assume that \mathcal{P}_i has exactly τ partitions, we can enforce this assumption by duplicating partitions if necessary). Fix some j . For $i < 0$, let \mathcal{P}_i^j be the partition where each vertex is a singleton. For $i > I$ let \mathcal{P}_i^j be the trivial partition with a single cluster $\{X\}$. Consider $\{\mathcal{P}_i^j\}_{i \geq 0}^I$. We will inductively define a new set of partitions $\{\tilde{\mathcal{P}}_i^j\}_{i \geq 0}^I$, enforcing it to be a laminar system, while still closely resembling the original partitions.

Levels below 0 and above I stay as-is. Inductively, for any $i \geq 0$, after constructing $\tilde{\mathcal{P}}_{i-1}^j$ from \mathcal{P}_{i-1}^j , we will construct $\tilde{\mathcal{P}}_i^j$ from \mathcal{P}_i^j using $\tilde{\mathcal{P}}_{i-1}^j$. Let $\mathcal{P}_i^j = \{C_1, \dots, C_\phi\}$ be the clusters in the partition \mathcal{P}_i^j . For each $q \in [1, \phi]$, let $Y_q = X \setminus \cup_{a < q} \tilde{C}_a$ be the set of unclustered points (w.r.t. level i , before iteration q). Let $C'_q = C_q \cap Y_q$ be the cluster C_q restricted to vertices in Y_q , and let $S_{C'_q} = \{C \in \tilde{\mathcal{P}}_{i-1}^j \mid C \cap C'_q \neq \emptyset\}$ be the set of new level- $(i-1)$ clusters with non empty intersection with C'_q . We set the new cluster $\tilde{C}_q = \cup S_{C'_q}$ to be the union of these clusters, and continue iteratively. Clearly, $\tilde{\mathcal{P}}_{i-1}^j$ is a refinement of $\tilde{\mathcal{P}}_i^j$. We conclude that $\{\tilde{\mathcal{P}}_i^j\}_{i \geq -1}$ is a laminar hierarchical set of partitions that refine each other.

We next argue by induction that $\tilde{\mathcal{P}}_i^j$ has diameter $(1 + \epsilon) \cdot \Delta_i$. Consider $\tilde{C}_q \in \tilde{\mathcal{P}}_i^j$. It consists of $C'_q \subseteq C_q \in \mathcal{P}_i^j$ and of clusters in $\tilde{\mathcal{P}}_{i-1}^j$ intersecting C'_q . As the diameter of C'_q is bounded by $\text{diam}(C_q) \leq \Delta_i$, and by the induction hypothesis, the diameter of each cluster $C \in \tilde{\mathcal{P}}_{i-1}^j$ is bounded by $(1 + \epsilon) \cdot \Delta_{i-1}$, we conclude that the diameter of \tilde{C}_q is bounded by

$$\Delta_i + 2 \cdot (1 + \epsilon) \Delta_{i-1} = \Delta_i \cdot \left(1 + \frac{2(1 + \epsilon)}{4\beta} \cdot \epsilon \right) \leq (1 + \epsilon) \cdot \Delta_i ,$$

since $\beta \geq 1$ and $\epsilon < 1$.

Finally, we argue that every ball is fully contained in some cluster. Fix, $x \in X$ and $i \in \mathbb{Z}$. If $i < 0$ or $i > I$ then there is nothing to prove. We will thus assume $i \in [0, I]$. Set

$$R = \frac{\Delta_i}{\beta} - (1 + \epsilon) \cdot \Delta_{i-1} = \frac{\Delta_i}{\beta} \cdot \left(1 - (1 + \epsilon) \cdot \frac{\epsilon}{4} \right) \geq \frac{\Delta_i}{(1 + \epsilon) \cdot \beta} .$$

By construction, there is some index j such that the ball $B_X(x, \frac{\Delta_i}{\beta})$ is fully contained in a single cluster C_q in the partition \mathcal{P}_i^j . It will be enough to show that the ball $B_X(x, R)$ is fully contained in the cluster \tilde{C}_q of $\tilde{\mathcal{P}}_i^j$. Suppose for the sake of contradiction otherwise, then there have to be a vertex $y \in B_X(x, R) \setminus Y_q$. In particular, y belongs to a cluster $Q \in \tilde{\mathcal{P}}_j^{i-1}$ that intersected $C_{q'} \in \mathcal{P}_i^j$ for $q' < q$. Let $z \in C_{q'} \cap Q$. As the maximum diameter of $\tilde{\mathcal{P}}_j^{i-1}$ cluster is $(1 + \epsilon) \cdot \Delta_{i-1}$, it follows that

$$d_X(z, x) \leq d_X(z, y) + d_X(y, x) \leq (1 + \epsilon) \cdot \Delta_{i-1} + R = \frac{\Delta_i}{\beta} .$$

But this implies that $z \in C_q$ and not $C_{q'}$, a contradiction. \square

Consider a partition \mathcal{P} , and a point $x \in X$. Let $C_x \in \mathcal{P}$ be the cluster containing x . We denote by $\partial_{\mathcal{P}}(x) = d_X(x, X \setminus C_x)$ the distance between x and the boundary of its cluster C_x . The following embedding will be the core stone of our construction:

Lemma 7. Consider a finite metric space (X, d_X) , and a hierarchy of laminar partitions $\mathcal{P}_1, \dots, \mathcal{P}_k$ of X (that is \mathcal{P}_i is a refinement of \mathcal{P}_{i+1}). Then there is an embedding $f : X \rightarrow \ell_\infty^{2 \log |X| + 2k}$ such that:

1. Lipschitz: $\forall x, y \in X, \|f(x) - f(y)\|_\infty \leq 2 \cdot d_X(x, y)$.
2. For every pair of points $x, y \in X$ and partition \mathcal{P}_i where x, y belong to different clusters, it holds that $\|f(x) - f(y)\|_\infty \geq \partial_{\mathcal{P}_i}(x) + \partial_{\mathcal{P}_i}(y)$.

Proof. The construction and proof is by induction on $|X|$, the number of metric points. The base case where $|X| = 1$ is trivial, as the empty embedding will do. Suppose that the partition \mathcal{P}_k is into the clusters X_1, X_2, \dots, X_m . If $m = 1$ (that is the trivial partition), then we can simply ignore this partition (as we don't guarantee anything w.r.t. this partition). If all the partitions in the hierarchy are trivial, then again there is nothing to prove (as no pair is ever separated) and the empty embedding will do. We thus can assume that $m \geq 2$.

Using the induction hypothesis, we construct functions $f_j : X_j \rightarrow \ell_\infty^{2 \log |X_j| + 2(k-1)}$ fulfilling the conditions of the lemma. Let Y be a set with the elements being X_1, \dots, X_m and weight function $\mu(X_j) = |X_j|$. Note that $\mu(Y) = |X|$. Using Lemma 5 we obtain a prefix free code $h : \{X_1, \dots, X_m\} \rightarrow \{\pm 1\}^*$ such that for every $j \in [1, m]$, $|h(X_j)| = 2 \cdot \left\lceil \log \frac{\mu(Y)}{\mu(X_j)} \right\rceil = 2 \cdot \left\lceil \log \frac{|X|}{|X_j|} \right\rceil$ (we can obtain equality instead of weak inequality by padding with arbitrary ± 1). We define a new embedding $f : X \rightarrow \ell_\infty$, such that for a point $x \in X_j$, for every $q \leq 2 \cdot \left\lceil \log \frac{\mu(Y)}{\mu(X_j)} \right\rceil$, the q 'th coordinate equal to $(f(x))_q = (h(X_j))_q \cdot \partial_{\mathcal{P}_k}(x)$. The rest of the coordinates will be just a concatenation of $f_j(x)$. For every point $x \in X_j$, the total number of coordinates is bounded by

$$2 \log |X_j| + 2(k-1) + 2 \cdot \left\lceil \log \frac{|X|}{|X_j|} \right\rceil \leq 2 \log |X_j| + 2k + 2 \cdot \log \frac{|X|}{|X_j|} = 2k + 2 \log |X| .$$

We can pad with 0 coordinates to get exactly $2k + 2 \log |X|$ coordinates.

Next we prove that our embeddings is expanding by at most a factor of 2. The proof follows the construction, and hence it is also by induction on $|X|$. Consider a pair $x, y \in X$. Suppose first that x, y belong to the same cluster X_j in the partition \mathcal{P}_k . For every coordinate q among the first $2 \cdot \left\lceil \log \frac{|X|}{|X_j|} \right\rceil$ coordinates, by the triangle inequality it holds that

$$\begin{aligned} |(f(x))_q - (f(y))_q| &= |h_q(X_j) \cdot \partial_{\mathcal{P}_k}(x) - h_q(X_j) \cdot \partial_{\mathcal{P}_k}(y)| \\ &= |d_X(x, X \setminus X_j) - d_X(y, X \setminus X_j)| \leq d_X(x, y) . \end{aligned} \quad (1)$$

The rest of the coordinates are a concatenation of $f_j(x), f_j(y)$ and the upper bound follows by the induction hypothesis.

Next, suppose that $x \in X_j, y \in X_{j'}$ for $j \neq j'$. Consider a coordinate q . Note that $(f(x))_q$ is either a 0, or equals to $\sigma_x \cdot \partial_{\mathcal{P}_{k'}}(x) = \sigma_x \cdot d_X(x, X \setminus Y_j)$ where $\sigma_x \in \{\pm 1\}$, $k' \in [1, k]$, and Y_j is the cluster containing x in $\mathcal{P}_{k'}$. In particular, $Y_j \subset X_j$. Similarly $(f(y))_q$ is either a 0 or $\sigma_y \cdot d_X(y, X \setminus Y_{j'})$ for $\sigma_y \in \{\pm 1\}$ and $Y_{j'} \subseteq X_{j'}$. It holds that

$$\begin{aligned} |(f(x))_q - (f(y))_q| &\leq |(f(x))_q| + |(f(y))_q| \leq |d_X(x, X \setminus Y_j)| + |d_X(y, X \setminus Y_{j'})| \\ &\leq |d_X(x, y)| + |d_X(y, x)| = 2 \cdot d_X(y, x) , \end{aligned} \quad (2)$$

where the third inequality holds as $y \notin Y_j$ and $x \notin Y_{j'}$.

It remains to prove the lower bound. Consider a pair $x, y \in X$ that belong to different clusters in \mathcal{P}_i . Let $X_j, X_{j'} \in \mathcal{P}_i$ be the clusters containing x and y respectively. Let $i' \in [i, k]$ be the maximal index such that x, y belong to different clusters in $\mathcal{P}_{i'}$. Let $Y_j, Y_{j'} \in \mathcal{P}_{i'}$ be the clusters containing x and y respectively. As $i \leq i'$, \mathcal{P}_i is a refinement of $\mathcal{P}_{i'}$. In particular, $X_j \subseteq Y_j$ and $X_{j'} \subseteq Y_{j'}$. As x and y belong to the same clusters in the partitions $\mathcal{P}_{j'+1}, \dots, \mathcal{P}_k$, their first coordinates of $f(x), f(y)$ correspond to these partitions and are “aligned”. Suppose that all these embeddings fill the first s coordinates. Then we have the partition w.r.t. $\mathcal{P}_{i'}$. In particular, we used a prefix free code h , and hence there is some q such that both $h_q(Y_j), h_q(Y_{j'})$ are defined and different. We conclude

$$\begin{aligned} \|f(x) - f(y)\|_\infty &\geq |(f(x))_{s+q} - (f(y))_{s+q}| \\ &= |h_q(Y_j) \cdot \partial_{\mathcal{P}_{j'}}(x) - h_q(Y_{j'}) \cdot \partial_{\mathcal{P}_{j'}}(y)| \\ &= |\partial_{\mathcal{P}_{j'}}(x)| + |\partial_{\mathcal{P}_{j'}}(y)| \geq |\partial_{\mathcal{P}_j}(x)| + |\partial_{\mathcal{P}_j}(y)|, \end{aligned}$$

where the last inequality holds as $\partial_{\mathcal{P}_{j'}}(x) = d_X(x, X \setminus Y_j) \geq d_X(x, X \setminus X_j) = \partial_{\mathcal{P}_j}(x)$ because $X_j \subseteq Y_j$. Similarly for y . □

Lemma 8. *Consider an n -point metric space (X, d_X) that admits a (β, τ) -sparse partition cover scheme, and such that all the pairwise distances in X are between 1 and Φ . Then for every $\varepsilon \in (0, 1)$, there is an (efficiently computable) embedding $f : X \rightarrow \ell_\infty^k$ with distortion $(1 + \varepsilon) \cdot 2\beta$, for $k = O\left(\frac{\tau}{\varepsilon} \cdot \log \frac{\beta}{\varepsilon} \cdot \log(n\Phi)\right)$.*

Proof. We will construct embedding with distortion $(1 + O(\varepsilon)) \cdot \beta$, afterwards one can obtain the lemma by scaling ε accordingly. For every $q \in \left\{1, 2, \dots, \left\lceil \log_{1+\varepsilon} \frac{4\beta}{\varepsilon} \right\rceil\right\}$ let $a_q = (1 + \varepsilon)^q$. Using Lemma 6 with parameters a_q and ε , construct hierarchical partitions $\{\mathcal{P}_i^{1,q}\}_{i \in \mathbb{Z}}, \dots, \{\mathcal{P}_i^{\tau,q}\}_{i \in \mathbb{Z}}$. Fix $I = \lceil \log_{4\beta/\varepsilon} \Phi \rceil$. For every $q \in \left\lceil \log_{1+\varepsilon} \frac{4\beta}{\varepsilon} \right\rceil$ and $j \in [\tau]$, consider the hierarchical partition $\{\mathcal{P}_i^{j,q}\}_{i=-1}^I$. Using Lemma 7, we construct an embedding $f_{j,q} : X \rightarrow \ell_\infty^{2 \log n + 2(I+2)}$. Our final embedding is simply a concatenation of all this embeddings: $\circ_{j \in [\tau], q \in \left\lceil \log_{1+\varepsilon} \frac{4\beta}{\varepsilon} \right\rceil} f_{j,q}$. The overall dimension is $\tau \cdot \lceil \log_{1+\varepsilon} \frac{4\beta}{\varepsilon} \rceil \cdot (2 \log n + 2(I+1)) = O\left(\frac{\tau}{\varepsilon} \cdot \log \frac{\beta}{\varepsilon} \cdot \log(n\Phi)\right)$.

Next we bound the distortion. Consider a pair $x, y \in X$. Let $q \in [0, \lceil \log_{1+\varepsilon} \frac{4\beta}{\varepsilon} \rceil]$, and $i \in [-1, I]$ be indices such that $(1 + \varepsilon)^q \left(\frac{4\beta}{\varepsilon}\right)^i < \frac{d_X(x,y)}{1+\varepsilon} \leq (1 + \varepsilon)^{q+1} \left(\frac{4\beta}{\varepsilon}\right)^i$. Fix $\Delta_i = a_q \cdot \left(\frac{4\beta}{\varepsilon}\right)^i$, and note that $(1 + \varepsilon) \cdot \Delta_i = (1 + \varepsilon)^{q+1} \cdot \left(\frac{4\beta}{\varepsilon}\right)^i < d_X(x, y)$. There is some $j \in [\tau]$ such that in the hierarchical partition $\{\mathcal{P}_{i'}^{j,q}\}_{i'=-1}^I$, the ball $B_X(x, \frac{\Delta_i}{\beta(1+\varepsilon)})$ is fully contained in some cluster of $\mathcal{P}_i^{j,q}$. In particular, $\partial_{\mathcal{P}_i^{j,q}}(x) > \frac{\Delta_i}{\beta(1+\varepsilon)}$. According to Lemma 6, $\mathcal{P}_i^{j,q}$ is a $(1 + \varepsilon) \cdot \Delta_i$ bounded partition. It follows that x and y have to belong to different clusters in $\mathcal{P}_i^{j,q}$. Using Lemma 7 we conclude

$$\begin{aligned} \|f(x) - f(y)\|_\infty &\geq \|f_{j,q}(x) - f_{j,q}(y)\|_\infty \geq \partial_{\mathcal{P}_i^{j,q}}(x) + \partial_{\mathcal{P}_i^{j,q}}(y) \\ &\geq \frac{\Delta_i}{\beta(1 + \varepsilon)} \geq \frac{d_X(x, y)}{\beta(1 + \varepsilon)^3} = \frac{d_X(x, y)}{\beta(1 + O(\varepsilon))}. \end{aligned} \tag{3}$$

Furthermore, as all the embeddings created by Lemma 7 are 2-Lipschitz, so is f . The lemma now follows. □

5.3 Removing the dependence on the aspect ratio

In this section we prove [Theorem 6](#). The following is the main lemma of the subsection, where we show that in a metric embeddings from a CURW family to ℓ_∞ , one can remove the dependence on the aspect ratio. This is a general phenomena, independent of SPCS.

Lemma 9. *Consider a CURW graph family \mathcal{F} , and suppose that every n - point graph $G \in \mathcal{F}$ with aspect ratio Φ can be embedded into ℓ_∞^k with expansion ρ , contraction β , and dimension $k = \varphi(n, \Phi, \rho, \beta)$, where $\varphi : \mathbb{N} \times \mathbb{R}_{\geq 1}^3 \rightarrow \mathbb{N}$ is some coordinate-wise monotone function. Then for every $\epsilon \in (0, \frac{1}{2})$, every n - point graph $G \in \mathcal{F}$ can be embedded into $\ell_\infty^{k'}$ with expansion $(1 + \delta) \cdot \rho$, contraction $(1 + \epsilon) \cdot \beta$, and dimension $3 \cdot \varphi(n, \left(\frac{8 \cdot \rho \cdot \beta}{\epsilon}\right)^2 \cdot n^3, \rho, \beta)$.*

Combining [Lemmas 8](#) and [9](#) we get our main meta [Theorem 6](#). The rest of this section is devoted to the proof of [Lemma 9](#).

Proof of Lemma 9. Fix $s = \frac{8 \cdot \rho \cdot \beta \cdot n}{\epsilon}$. For a parameter $\alpha > 0$, let $G^\alpha = (V, E, w_\alpha)$ be the graph with the following weight function for every $e \in E$,

$$w^\alpha(e) = \begin{cases} \alpha & w(e) \geq \alpha \\ w(e) & \frac{\alpha}{s^2} < w(e) < \alpha \\ 0 & \frac{\alpha}{s^2} \geq w(e) \end{cases} .$$

That is we reduce all edge weights to be at most α , and we nullified (essentially contract) all edges of weight at most $\frac{\alpha}{s^2}$. Note that as \mathcal{F} is CURW, $G^\alpha \in \mathcal{F}$. We next analyze the properties of G^α .

Claim 2. *The graph G^α has the following properties:*

1. For every $x, y \in V$, $d_{G^\alpha}(x, y) \leq \min \{d_G(x, y), n \cdot \alpha\}$.
2. For every $x, y \in X$ such that $d_G(x, y) \in [\frac{\alpha}{s}, \alpha]$, $d_{G^\alpha}(x, y) \geq (1 - \frac{n}{s}) \cdot d_G(x, y)$.
3. The induced shortest path metric has aspect ratio $n \cdot s^2$.

Proof. (1). The edge weights only decreased, so clearly $d_{G^\alpha}(x, y) \leq d_G(x, y)$. In addition, the shortest path from x to y in G^α consist of at most $n - 1$ edges, all of weight at most α , so it has weight at most $n \cdot \alpha$. (2). Consider a pair x, y such that $d_G(x, y) \in [\frac{\alpha}{s}, \alpha]$. Let P be the shortest path from x to y in G^α . As $d_G(x, y) < \alpha$, we can assume that P does not contain any edges of weight at least α . It holds that

$$d_{G^\alpha}(x, y) = w^\alpha(P) > w(P) - n \cdot \frac{\alpha}{s^2} \geq d_G(x, y) - \frac{n}{s} \cdot d_G(x, y) . \quad (4)$$

(3). Recall that our definition of aspect ratio ignores 0 distances: $\Phi(G) = \frac{\max_{u, v \in V} d_G(u, v)}{\min_{u, v \in V \text{ s.t. } d_G(u, v) > 0} d_G(u, v)}$ (see [Section 2](#)). It follows that the minimal distance in G^α is $\frac{\alpha}{s^2}$, and hence by point (1), the aspect ratio is bounded by $\frac{n \cdot \alpha}{\frac{\alpha}{s^2}} = n \cdot s^2$. \square

Let $\alpha_i = \Phi \cdot s^{-i}$. For every $i \geq 0$, construct an embedding $f_i : X \rightarrow \ell_\infty$ w.r.t. $d_{G^{\alpha_i}}$ with expansion ρ and contraction β and dimension $\varphi(n, n \cdot s^2, \rho, \beta)$. We will then reuse coordinates such that all the embeddings $\{f_{3i}\}_{i \geq 0}$ all use the same coordinates. Similarly for $\{f_{3i+1}\}_{i \geq 0}$ and $\{f_{3i+2}\}_{i \geq 0}$. We will




Figure 2: Illustration of the contraction and expansion proof in Lemma 9.

denote these embeddings by g_0, g_1, g_2 respectively. This finishes the definition of our embedding f . Clearly the overall number of coordinates is bounded by $3 \cdot \varphi(n, n \cdot s^2, \rho, \beta) = 3 \cdot \varphi(n, \left(\frac{8 \cdot \rho \cdot \beta}{\epsilon}\right)^2 \cdot n^3, \rho, \beta)$. Next we bound expansion and contraction (see Figure 2 for illustration).

Consider $x, y \in X$, and assume w.l.o.g. that there is i such that $\alpha_{3i+1} < d_G(x, y) \leq \alpha_{3i}$ (the other cases is where there is an i such that either $\alpha_{3i+2} < d_G(x, y) \leq \alpha_{3i+1}$ or $\alpha_{3i+3} < d_G(x, y) \leq \alpha_{3i+2}$ are treated symmetrically). For every $j \leq 3i - 2$ (if any), it holds that $d_G(x, y) \leq \alpha_{3i} = \alpha_j \cdot s^{j-3i} \leq \alpha_j \cdot s^{-2}$. Hence $d_G^{\alpha_j}(x, y) = 0$, which implies $\|f_j(x) - f_j(y)\|_\infty = 0$. For every $j \geq 3i + 2$,

$$d_G^{\alpha_j}(x, y) \leq n \cdot \alpha_j = n \cdot s^{(3i+1)-j} \cdot \alpha_{3i+1} \leq n \cdot s^{(3i+1)-j} \cdot d_G(x, y) .$$

We conclude

$$\begin{aligned} \|g_0(x) - g_0(y)\|_\infty &= \left\| \sum_{q \geq 0} f_{3q}(x) - \sum_{q \geq 0} f_{3q}(y) \right\|_\infty \\ &\leq \|f_{3i}(x) - f_{3i}(y)\|_\infty + \sum_{q \geq i+1} \|f_{3q}(x) - f_{3q}(y)\|_\infty \\ &\leq \rho \cdot d_G(x, y) + \rho \cdot \sum_{q \geq i+1} n \cdot s^{(3i+1)-3q} \cdot d_G(x, y) \\ &= \rho \cdot d_G(x, y) + \rho \cdot \frac{n}{s^2} \cdot \frac{1}{1-s^{-3}} \cdot d_G(x, y) < \left(1 + \frac{2n}{s^2}\right) \cdot \rho \cdot d_G(x, y) . \end{aligned}$$

$$\begin{aligned} \|g_1(x) - g_1(y)\|_\infty &\leq \|f_{3i+1}(x) - f_{3i+1}(y)\|_\infty + \sum_{q \geq i+1} \|f_{3q+1}(x) - f_{3q+1}(y)\|_\infty \\ &\leq \rho \cdot d_G(x, y) + \rho \cdot \sum_{q \geq i+1} n \cdot s^{(3i+1)-(3q+1)} \cdot d_G(x, y) \\ &= \left(1 + \frac{n}{s^3} \cdot \frac{1}{1-s^{-3}}\right) \cdot \rho \cdot d_G(x, y) \leq \left(1 + \frac{2n}{s^3}\right) \cdot \rho \cdot d_G(x, y) \end{aligned}$$

$$\begin{aligned} \|g_2(x) - g_2(y)\|_\infty &\leq \|f_{3i-1}(x) - f_{3i-1}(y)\|_\infty + \sum_{q \geq i} \|f_{3q+2}(x) - f_{3q+2}(y)\|_\infty \\ &\leq \rho \cdot d_G(x, y) + \rho \cdot \sum_{q \geq i} n \cdot s^{(3i+1)-(3q+2)} \cdot d_G(x, y) \\ &= \left(1 + \frac{n}{s} \cdot \frac{1}{1-s^{-3}}\right) \cdot \rho \cdot d_G(x, y) \leq \left(1 + \frac{2n}{s}\right) \cdot \rho \cdot d_G(x, y) \end{aligned}$$

From the other hand, according to [Claim 2](#), $d_{G^{\alpha_{3i}}}(x, y) \geq (1 - \frac{n}{s}) \cdot d_G(x, y)$. It holds that

$$\begin{aligned}
\|g_0(x) - g_0(y)\|_\infty &\geq \|f_{3i}(x) - f_{3i}(y)\|_\infty - \sum_{q \geq i+1} \|f_{3q}(x) - f_{3q}(y)\|_\infty \\
&\geq \frac{1}{\beta} \cdot d_{G^{\alpha_{3i}}}(x, y) - \rho \cdot \sum_{q \geq i+1} n \cdot s^{(3i+1)-3q} \cdot d_G(x, y) \\
&= \frac{1}{\beta} \cdot (1 - \frac{n}{s}) \cdot d_G(x, y) - \rho \cdot \frac{n}{s^2} \cdot \frac{1}{1 - s^{-3}} \cdot d_G(x, y) \\
&> \left(1 - \frac{n}{s} - \rho \cdot \beta \cdot \frac{2n}{s^2}\right) \cdot \frac{1}{\beta} \cdot d_G(x, y) .
\end{aligned}$$

As $\|f(x) - f(y)\|_\infty = \max_{j \in \{0, 1, 2\}} \|g_j(x) - g_j(y)\|_\infty$, and according to our choice of s we conclude that

$$\begin{aligned}
\|f(x) - f(y)\|_\infty &\leq \left(1 + \frac{2n}{s}\right) \cdot \rho \cdot d_G(x, y) \leq (1 + \epsilon) \cdot \rho \cdot d_G(x, y) \\
\|f(x) - f(y)\|_\infty &\geq \left(1 - \frac{n}{s} - \rho \cdot \beta \cdot \frac{2n}{s^2}\right) \cdot \frac{1}{\beta} \cdot d_G(x, y) \geq \frac{1}{(1 + \epsilon) \cdot \beta} \cdot d_G(x, y)
\end{aligned}$$

□

5.4 Improved distortion for minor free graphs

This subsection is devoted to proving [Theorem 2](#), which improves the distortion of the ℓ_∞ embedding of K_r -minor free graphs to $3 + \epsilon$. The following is our key lemma, which is parallel to [Lemma 8](#).

Lemma 10. *Fix $\epsilon \in (0, \frac{1}{2})$ and $\Phi \geq 1$, and consider a weighted n vertex K_r -minor free graph $G = (V, E, w)$ such that all the pairwise distances in X are between 1 and Φ . Then G can be embedded into $\ell_\infty^{O(\epsilon)^{r+1} \cdot \log \frac{1}{\epsilon} \cdot \log(n\Phi)}$ with distortion $3 + \epsilon$.*

Proof. The embedding is essentially the same as in the construction of [Lemma 8](#), and the main changes are in the analysis. Roughly speaking, first we subdivide all the edges of our graph, this will reduce the expansion from 2 to $1 + \epsilon$. Then we will dive into the specific details of the sparse cover, and show that the contraction is only $3 + O(\epsilon)$.

Consider our K_r minor free graph $G = (V, E, w)$. Let $\tilde{G} = (\tilde{V}, \tilde{E}, \tilde{w})$ be a graph, where we replace every edge $e = (v, u) \in E$ with a path consisting of $\frac{1}{\epsilon}$ edges of length $\epsilon \cdot w(e)$. Note that all the pairwise distances between original vertices remained exactly the same, and that \tilde{G} is still K_r -minor free, thus it also admits the sparse covers from [Theorem 1](#). We will embed only the original vertices V into ℓ_∞ , however the partitions will be created w.r.t. \tilde{G} . This small change will improve our upper bound in [Lemma 7](#) from expansion 2 to expansion $1 + \epsilon$. Indeed, consider a pair $x, y \in V$, and recall the proof of [Lemma 7](#). The proof is by induction on $|V|$. The case where x, y belong to the same cluster in \mathcal{P}_k is treated in the same way (as [eq. \(1\)](#) guarantees $|(f(x))_q - (f(y))_q| \leq d_G(x, y)$). In the second case, x and y correspond so different clusters, where according to [eq. \(2\)](#), $|(f(x))_q - (f(y))_q| \leq d_{\tilde{G}}(x, \tilde{V} \setminus Y_j) + d_{\tilde{G}}(y, \tilde{V} \setminus Y_{j'})$ where $x \in Y_j$, $y \in Y_{j'}$, and $Y_j \cap Y_{j'} = \emptyset$ are disjoint. In \tilde{G} , the shortest path P from x to y in \tilde{G} must go through $x' \notin Y_j$, and $y' \notin Y_{j'}$ such that x', y' are consecutive vertices of a subdivided edge e (see two illustrations bellow). In particular, $d_{\tilde{G}}(x', y') = \epsilon \cdot w(e) \leq \epsilon \cdot w(P)$. It follows that

$$\begin{aligned}
|(f(x))_q - (f(y))_q| &\leq d_{\tilde{G}}(x, \tilde{V} \setminus Y_j) + d_{\tilde{G}}(y, \tilde{V} \setminus Y_{j'}) \\
&\leq d_{\tilde{G}}(x, x') + d_{\tilde{G}}(y, y') \leq d_G(x, y) + d_G(x', y') < (1 + \varepsilon) \cdot d_G(x, y) .
\end{aligned}$$



It follows that the embedding produced by [Theorem 6](#) has expansion $1 + \varepsilon$.

Next we improve the contraction guarantee. The basis of our construction is the $(4 \cdot (1 + \varepsilon), O(\frac{1}{\varepsilon})^r)$ -SPCS of [Theorem 1](#). We begin by observing that the padding guarantee in [Theorem 5](#) is somewhat stronger than what is required by the definition of SPCS. Indeed, we follow the construction (and notation) of [Theorem 5](#). Note that here $\gamma = \frac{\Delta}{r}$, and for the construction of our strong $(4 \cdot (1 + \varepsilon), O(\frac{1}{\varepsilon})^r)$ -SPCS we used $q = \frac{2r}{\varepsilon}$. In [Lemmas 3](#) and [4](#) it was shown that the cluster satisfying a point x consist of a ball $B_{G[\hat{\eta}]}(v, 2\Delta + q \cdot \gamma) = B_{G[\hat{\eta}]}(v, (1 + \varepsilon) \cdot \frac{2\Delta}{\varepsilon})$ where $B_G(x, \frac{q\gamma}{2}) = B_G(x, \frac{\Delta}{\varepsilon}) \subseteq \hat{\eta}$ and $d_{G[\hat{\eta}]}(v, x) \leq 2\Delta + \frac{q\gamma}{2} = (1 + 2\varepsilon) \cdot \frac{\Delta}{\varepsilon}$. Note that the diameter of the corresponding partition is $(1 + \varepsilon) \cdot \frac{4\Delta}{\varepsilon}$, and that $\partial_{\mathcal{P}}(x) \geq \frac{\Delta}{\varepsilon}$. In other words, fix $\Psi = \frac{4\Delta}{\varepsilon}$, then there is a cluster $C_x \in \mathcal{P}$ such that $B_G(x, \Psi) \subseteq C \subseteq B_G(v, (1 + \varepsilon) \cdot 2\Psi)$, and $d_G(x, v) \leq (1 + 2\varepsilon) \cdot \Psi$. Next, in [Lemma 6](#), the partition \mathcal{P} was slightly changed to into $\tilde{\mathcal{P}}$ in order to become a part of an hierarchy. In particular, it holds that there is a cluster $\tilde{C} \in \tilde{\mathcal{P}}$ such that $B_G(x, \frac{\Psi}{1 + \varepsilon}) \subseteq \tilde{C} \subseteq B_G(v, (1 + \varepsilon)^2 \cdot 2\Psi)$. Furthermore, for a vertex $y \in V$ such that $d_G(x, y) > (3 + 8\varepsilon) \cdot \Psi$ it holds that

$$d_G(v, y) \geq d_G(x, y) - d_G(v, x) > (3 + 8\varepsilon) \cdot \Psi - (1 + 2\varepsilon) \cdot \Psi = (1 + 3\varepsilon) \cdot 2\Psi > (1 + \varepsilon)^2 \cdot 2\Psi ,$$

and thus y does not belong to \tilde{C} .

Fix a pair $x, y \in V$. In [Lemma 8](#) we constructed partition for every possible scale. Consider the scale (i, q) such that $(1 + \varepsilon)^q (\frac{4\beta}{\varepsilon})^i < \frac{4}{3 + 8\varepsilon} \cdot d_G(x, y) \leq (1 + \varepsilon)^{q+1} (\frac{4\beta}{\varepsilon})^i$. Fix $\Psi = \frac{1}{4} \cdot (1 + \varepsilon)^q (\frac{4\beta}{\varepsilon})^i$. Following the discussion above, for this scale, there was some partition $\mathcal{P}_i^{j,q}$ with diameter at most $(1 + \varepsilon)^2 \cdot 4\Psi$ such that the ball $B_G(x, \frac{\Psi}{1 + \varepsilon})$ was contained in a single cluster, and every vertex at distance greater than $(3 + 8\varepsilon) \cdot \Psi$ from x belonged to a different cluster. In particular, as $d_G(x, y) > \frac{3 + 8\varepsilon}{4} \cdot (1 + \varepsilon)^q (\frac{4\beta}{\varepsilon})^i = (3 + 8\varepsilon) \cdot \Psi$, y belongs to a different cluster. Using [Lemma 7](#) we conclude

$$\begin{aligned}
\|f(x) - f(y)\|_\infty &\geq \|f_{j,q}(x) - f_{j,q}(y)\|_\infty \geq \partial_{\mathcal{P}_i^{j,q}}(x) + \partial_{\mathcal{P}_i^{j,q}}(y) \\
&\geq \frac{\Psi}{1 + \varepsilon} = \frac{(3 + 8\varepsilon) \cdot \Psi}{(1 + \varepsilon) \cdot (3 + 8\varepsilon)} > \frac{d_G(x, y)}{3 \cdot (1 + O(\varepsilon))} .
\end{aligned}$$

Thus we indeed obtained distortion $3 + O(\varepsilon)$ as promised. Similarly to [Lemma 8](#), the overall dimension is bounded by $O\left(\frac{r}{\varepsilon} \cdot \log \frac{\beta}{\varepsilon} \cdot \log(n\Phi)\right) = O\left(\frac{1}{\varepsilon}\right)^{r+1} \cdot \log \frac{1}{\varepsilon} \cdot \log(n\Phi)$. \square

Finally, we apply [Lemma 9](#) on [Lemma 10](#) to get [Theorem 2](#), restated bellow for convenience.

Theorem 2. *For every $\varepsilon \in (0, \frac{1}{2})$, every n vertex K_r -minor free graph G can be embedded into $\ell_\infty^{O(\frac{1}{\varepsilon})^{r+1} \cdot \log \frac{1}{\varepsilon} \cdot \log \frac{n}{\varepsilon}}$ with distortion $3 + \varepsilon$.*

6 From sparse covers to padded decompositions

This section is devoted to proving our meta [Theorem 3](#) (restated below for convenience) that reduces sparse covers into padded decomposition. As a corollary, we recover the state of the art padded decomposition for K_τ minor-free graphs ([Corollary 2](#)) via a much simpler proof (compared with [\[AGG⁺19\]](#)). More importantly, every further improvement on sparse covers will now immediately lead to an improved padded decomposition!

Theorem 3 (From sparse covers to padded decompositions). *Consider a weighted graph $G = (V, E, w)$ that admits a (β, s, Δ) -sparse cover. Then G admits weak $(O(\beta \cdot \log s), \frac{1}{4\beta}, \Delta)$ -padded decomposition.*

Proof. Given a cover \mathcal{C} , we will sample a partition \mathcal{P} with the desired padding properties. To avoid confusion, the sets of \mathcal{C} (resp. \mathcal{P}) will be called \mathcal{C} -clusters (resp. \mathcal{P} -clusters). Our decomposition is morally [\[MPX13\]](#)-based, and closely follows [\[Fil19a\]](#). Filtser [\[Fil19a\]](#) proved that the existence of a “sparse net” implies padded decomposition. Specifically, if there is a net N such that every vertex v has a net point at distance at most $d_G(v, N) \leq \Delta$, and the number of net points in the ball $B_G(v, 3\Delta)$ is at most τ , then G admits a strong $(O(\ln \tau), \frac{1}{16}, 4\Delta)$ -padded decomposition. One can notice that by taking balls of radius 2Δ around the sparse net N , we will get a $(\frac{1}{4}, \tau, 4\Delta)$ -sparse cover, and thus our [Theorem 3](#) is in a sense generalization of [\[Fil19a\]](#) to arbitrary sparse covers (alas only with a weak diameter guarantee).

For a \mathcal{C} -cluster $C \subseteq V$, and a vertex $v \in V$. Denote by $\partial_C(v) = d_G(v, V \setminus C)$ the distance between v and the boundary of the \mathcal{C} -cluster C . Note that if $v \notin C$, $\partial_C(v) = 0$, while $B_G(v, r) \subseteq C$ implies $\partial_C(v) > r$. Furthermore, for every pair of vertices $u, v \in V$, $|\partial_C(u) - \partial_C(v)| = |d_G(u, V \setminus C) - d_G(v, V \setminus C)| \leq d_G(u, v)$.

To create a padded decompositions, following previous works, we will use truncated exponential distribution. That is, exponential distribution conditioned on the event that the outcome lays in a certain interval. The $[0, 1]$ -truncated exponential distribution with parameter λ is denoted by $\text{Texp}(\lambda)$, the density function is then $g(y) = \frac{\lambda e^{-\lambda y}}{1 - e^{-\lambda}}$, for $y \in [0, 1]$.

Consider a (β, s, Δ) -sparse cover \mathcal{C} . For every \mathcal{C} -cluster C we sample $\delta_C \sim \text{Texp}(\lambda)$ using truncated exponential distribution with parameter $\lambda = 2 + 2 \ln s$. For a \mathcal{C} -cluster C , we define a function $f_C : V \rightarrow \mathbb{R}_{\geq 0}$, as follows: $f_C(v) = \delta_C \cdot \frac{\Delta}{\beta} + \partial_C(v)$. We next create a partition $\mathcal{P} = \{P_C\}_{C \in \mathcal{C}}$ where each vertex $v \in V$ joins the \mathcal{P} -cluster P_C associated with the \mathcal{C} -cluster C that maximizes $f_C(v)$.

Fix a vertex $v \in V$. As \mathcal{C} is a sparse cover, there is some \mathcal{C} -cluster $C_v \in \mathcal{C}$ such that $B_G(v, \frac{\Delta}{\beta}) \subseteq C_v$, implying $f_{C_v}(v) > \frac{\Delta}{\beta}$. From the other hand, for every \mathcal{C} -cluster C which does not contain v , it holds that $f_C(v) \leq \delta_C \cdot \frac{\Delta}{\beta} + 0 \leq \frac{\Delta}{\beta}$. It follows that $f_C(v) < f_{C_v}(v)$. Hence v can only join a \mathcal{P} -cluster in P_C that associated with a \mathcal{C} -cluster C that contains it. In particular, for every $C \in \mathcal{C}$, $P_C \subseteq C$. As every \mathcal{C} -cluster has diameter at most Δ , we conclude that \mathcal{P} has diameter at most Δ . Note that the diameter guarantee in \mathcal{P} is weak, regardless of the diameter guarantee in \mathcal{C} .

Claim 3. *Consider a vertex v , and let $C_{(1)}, C_{(2)}, \dots$ be an ordering of the \mathcal{C} -clusters w.r.t. the values $\{f_C(v)\}_{C \in \mathcal{C}}$. That is $f_{C_{(1)}}(v) \geq f_{C_{(2)}}(v) \geq \dots$. Set $\Upsilon = f_{C_{(1)}}(v) - f_{C_{(2)}}(v)$. Then for every $r < \frac{\Upsilon}{2}$, it holds that $B_G(v, r) \subseteq P_{C_{(1)}}$.*

Proof. Fix $r < \frac{\Upsilon}{2}$. For every $u \in B_G(v, r)$, and every center $C \neq C_{(1)}$ it holds that,

$$\begin{aligned}
f_{C_{(1)}}(u) &= \delta_{C_{(1)}} \cdot \frac{\Delta}{\beta} + \partial_{C_{(1)}}(u) > \delta_{C_{(1)}} \cdot \frac{\Delta}{\beta} + \partial_{C_{(1)}}(v) - d_G(u, v) \\
&= f_{C_{(1)}}(v) - d_G(u, v) \geq f_C(v) + \Upsilon - d_G(u, v) \\
&= \delta_C \cdot \frac{\Delta}{\beta} + \partial_C(v) + \Upsilon - d_G(u, v) \\
&\geq \delta_C \cdot \frac{\Delta}{\beta} + \partial_C(u) + \Upsilon - 2 \cdot d_G(u, v) > f_C(u).
\end{aligned}$$

In particular, $u \in C_{(1)}$, and thus $B_G(v, r) \subseteq P_{C_{(1)}}$ as required. \square

Next we analyzing the padding probability. Consider some vertex $v \in V$, and parameter $\gamma \leq \frac{1}{4}$. We will first argue that the ball $B = B_G(v, \gamma \cdot \frac{\Delta}{\beta})$ is fully contained in a single \mathcal{P} -cluster with probability at least $e^{-4\gamma\lambda}$. The theorem will then follow due to scaling. Let $\mathcal{C} = \{C_1, C_2, \dots\}$ by an arbitrarily ordering of the \mathcal{C} -clusters. Denote by \mathcal{F}_i the event that v joins the cluster associated with C_i , i.e. $v \in P_{C_i}$. Denote by \mathcal{Q}_i the event that $v \in P_{C_i}$, but not all of the vertices in B joined P_{C_i} , that is $v \in P_{C_i} \cap B \neq B$. Let $\mathcal{C}_v \subseteq \mathcal{C}$ be the clusters containing v . Note that for every $C_i \notin \mathcal{C}_v$, $\Pr[\mathcal{F}_i] = \Pr[\mathcal{Q}_i] = 0$. To prove our assertion, it is enough to show that $\Pr[\cup_i \mathcal{Q}_i] \leq 1 - e^{-4\gamma\lambda}$. Set $\alpha = e^{-2\gamma\lambda}$.

Claim 4. For every i , $\Pr[\mathcal{Q}_i] \leq (1 - \alpha) \left(\Pr[\mathcal{F}_i] + \frac{1}{e^{\lambda} - 1} \right)$.

Proof. As the order in \mathcal{C} is arbitrary, assume w.l.o.g. that $i = |\mathcal{C}|$ and denote $C = C_{|\mathcal{C}|}$, $\mathcal{Q} = \mathcal{Q}_i$, $\mathcal{F} = \mathcal{F}_i$, and $\delta = \delta_{C_i}$. Let $X \in [0, 1]^{|\mathcal{C}|-1}$ be the vector where the j 'th coordinate equals δ_{x_j} . Set $\rho_X = \max \left\{ 0, \frac{\beta}{\Delta} \cdot \left(\max_{C' \neq C} \left\{ \delta_{C'} \cdot \frac{\Delta}{\beta} + \partial_{C'}(v) \right\} - \partial_C(v) \right) \right\}$. Note that ρ_X is the minimal value of δ such that if $\delta > \rho_X$, then C has the maximal value $f_C(v)$, and therefore v will join P_C . Note that it is possible that $\rho_X > 1$. Conditioning on all the other samples having values X , and assuming first that $\rho_X \leq 1$, it holds that

$$\Pr[\mathcal{F} \mid X] = \Pr[\delta > \rho_X] = \int_{\rho_X}^1 \frac{\lambda \cdot e^{-\lambda y}}{1 - e^{-\lambda}} dy = \frac{e^{-\rho_X \cdot \lambda} - e^{-\lambda}}{1 - e^{-\lambda}}.$$

If $\delta > \rho_X + 2\gamma$, then

$$\begin{aligned}
f_C(v) &= \delta_C \cdot \frac{\Delta}{\beta} + \partial_C(v) > (\rho_X + 2\gamma) \cdot \frac{\Delta}{\beta} + \partial_C(v) \\
&\geq 2\gamma \cdot \frac{\Delta}{\beta} + \max_{C' \neq C} \left\{ \delta_{C'} \cdot \frac{\Delta}{\beta} + \partial_{C'}(u) \right\} = 2\gamma \cdot \frac{\Delta}{\beta} + \max_{C' \neq C} f_{C'}(v).
\end{aligned}$$

In particular, by [Claim 3](#) the ball B will be contained in C . We conclude

$$\begin{aligned}
\Pr[\mathcal{Q} \mid X] &\leq \Pr[\rho_X \leq \delta \leq \rho_X + 2\gamma] \\
&= \int_{\rho_X}^{\max\{1, \rho_X + 2\gamma\}} \frac{\lambda \cdot e^{-\lambda y}}{1 - e^{-\lambda}} dy \\
&\leq \frac{e^{-\rho_X \cdot \lambda} - e^{-(\rho_X + 2\gamma) \cdot \lambda}}{1 - e^{-\lambda}} \\
&= (1 - e^{-2\gamma \cdot \lambda}) \cdot \frac{e^{-\rho_X \cdot \lambda}}{1 - e^{-\lambda}} \\
&= (1 - \alpha) \cdot \left(\Pr[\mathcal{F} \mid X] + \frac{1}{e^\lambda - 1} \right).
\end{aligned}$$

Note that if $\rho_X > 1$ then $\Pr[\mathcal{Q} \mid X] = 0 \leq (1 - \alpha) \cdot \left(\Pr[\mathcal{F} \mid X] + \frac{1}{e^\lambda - 1} \right)$ as well. Denote by h the density function of the distribution over all possible values of X . Using the law of total probability, we can bound the probability of \mathcal{Q} :

$$\begin{aligned}
\Pr[\mathcal{Q}] &= \int_X \Pr[\mathcal{Q} \mid X] \cdot h(X) dX \\
&\leq (1 - \alpha) \cdot \int_X \left(\Pr[\mathcal{F} \mid X] + \frac{1}{e^\lambda - 1} \right) \cdot h(X) dX \\
&= (1 - \alpha) \cdot \left(\Pr[\mathcal{F}] + \frac{1}{e^\lambda - 1} \right)
\end{aligned}$$

□

Next, we bound the probability that the ball B is cut.

$$\begin{aligned}
\Pr[\cup_i \mathcal{Q}_i] &= \sum_{C_i \in \mathcal{C}_v} \Pr[\mathcal{Q}_i] \leq (1 - \alpha) \cdot \sum_{C_i \in \mathcal{C}_v} \left(\Pr[\mathcal{F}_i] + \frac{1}{e^\lambda - 1} \right) \\
&\leq (1 - e^{-2\gamma \cdot \lambda}) \cdot \left(1 + \frac{s}{e^\lambda - 1} \right) \\
&\leq (1 - e^{-2\gamma \cdot \lambda}) \cdot (1 + e^{-2\gamma \cdot \lambda}) = 1 - e^{-4\gamma \cdot \lambda},
\end{aligned}$$

where the last inequality follows as $e^{-2\gamma \cdot \lambda} = \frac{e^{-2\gamma \cdot \lambda} (e^\lambda - 1)}{e^\lambda - 1} \geq \frac{e^{-2\gamma \cdot \lambda} \cdot e^{\lambda - 1}}{e^\lambda - 1} \geq \frac{e^{\frac{\lambda}{2} - 1}}{e^\lambda - 1} = \frac{s}{e^\lambda - 1}$, where we used that $\gamma \in (0, \frac{1}{4}]$, and $\lambda = 2 + 2 \ln s$.

We conclude that our distribution indeed produces a weak $(4\beta \cdot \lambda, \frac{1}{4\beta}, \Delta)$ -padded decomposition. Indeed, we already established that the diameter is at most Δ . Next, fix $\gamma \leq \frac{1}{4\beta}$, and $v \in V$. Denote by $P(v)$ the \mathcal{P} -cluster containing v . Then

$$\Pr_{\mathcal{P}}[B_G(v, \gamma \cdot \Delta) \subseteq P(v)] = \Pr_{\mathcal{P}}\left[B_G(v, \beta \cdot \gamma \cdot \frac{\Delta}{\beta}) \subseteq P(v)\right] \geq e^{-4\beta \cdot \gamma \cdot \lambda},$$

where we used ?? w.r.t. $\beta \cdot \gamma$, and the fact that $\beta \cdot \gamma \leq \frac{1}{4}$. The theorem now follows.

□

7 Further applications

7.1 Sparse Partitions

Given a weighted graph $G = (V, E, w)$, a weak/strong (α, τ, Δ) -sparse partition is a partition \mathcal{C} of V such that:

- **Low Diameter:** $\forall C \in \mathcal{C}$, the C has a weak/strong diameter at most Δ ;
- **Ball Preservation:** $\forall v \in V$, the ball $B_G(v, \frac{\Delta}{\alpha})$ intersects at most τ clusters from \mathcal{C} .

We say that the graph $G = (V, E, w)$ admits a weak/strong (α, τ) -sparse partition scheme if for every $\Delta > 0$, G admits a weak/strong (α, τ, Δ) -sparse partition.

Sparse partition have been used to construct universal TSP and universal Steiner tree [JLN⁺05, BDR⁺12, Fil20a, BCF⁺23] (see Section 7.2). Recently they been also used to obtain an $O(\frac{d}{\log d})$ approximation for the facility location problem in \mathbb{R}^d in the geometric streaming model [CFJ⁺22].

Jia *et al.* [JLN⁺05] implicitly proved (see [Fil20a] for an explicit proof) that if a space admits a weak (β, s) -sparse cover scheme, then it admits a weak (β, s) -sparse partition scheme. Using Theorem 1 we conclude:

Corollary 3. *Every K_r -minor free graph G admits a weak $(O(r), O(r^2))$ -sparse partition scheme. In addition, for $\varepsilon \in (0, \frac{1}{2})$, G admits a weak $(4 + \varepsilon, O(\frac{1}{\varepsilon})^r)$ -sparse partition.*

The previous state of the art follows directly from the previously best know weak sparse cover scheme, that is weak $(O(r^2), O(2^r))$ -sparse partition scheme (see [Fil20a], and also [KPR93, FT03]).

7.2 Universal TSP and Universal Steiner Tree

Consider a postman providing post service for a set X of clients with n different locations (with distance measure d_X). Each morning the postman receives a subset $S \subseteq X$ of the required deliveries for the day. In order to minimize the total tour length, one solution may be to compute each morning an (approximation of an) optimal TSP tour for the set S . An alternative solution will be to compute a *Universal TSP* (UTSP) tour. This is a universal tour R containing all the points X . Given a subset S , $R(S)$ is the tour visiting all the points in S w.r.t. the order induced by R . Given a tour T denote its length by $|T|$. The *stretch* of R is the maximum ratio among all subsets $S \subseteq X$ between the length of $R(S)$ and the length of the optimal TSP tour on S , $\max_{S \subseteq X} \frac{|R(S)|}{|\text{Opt}(S)|}$.

A closely related problem to UTSP is the *Universal Steiner tree* (UST). Consider the problem of designing a network that allows a server to broadcast a message to a single set of clients. If sending a message over a link incurs some cost then designing the best broadcast network is classically modeled as the Steiner tree problem [Hwa76]. However, if the server have to solve this problem repeatedly, with different client sets, it is desirable to construct a single network that will optimize the cost of the broadcast for every possible subset of clients.

Given a metric space (X, d_X) and root $r \in X$, a ρ -approximate universal Steiner tree (UST) is a weighted tree T over X such that for every $S \subseteq X$ containing r , we have

$$w(T\{S\}) \leq \rho \cdot \text{OPT}_S$$

where $T\{S\} \subseteq T$ is the minimal subtree of T connecting S , and OPT_S is the minimum weight Steiner tree connecting S in X . If the tree T is not required to be a subgraph of G , we will call the problem metric UST.

Jia *et al.* [JLN⁺05] showed that for every n -point metric space that admits weak (σ, τ) -sparse partition scheme, there is a polynomial time algorithm that given a root $\text{rt} \in V$ computes a UTSP with stretch $O(\tau\sigma^2 \log_\tau n)$, and a metric UST with stretch $O(\tau\sigma^2 \log_\tau n)$. Using our Corollary 3, we conclude:

Corollary 4. *Consider an n -point K_r -minor free weighted graph $G = (V, E, w)$. Then G admits a solution to both UTSP and metric UST with stretch $O(\frac{r^4}{\log r}) \cdot \log n$.*

The previous state of the art had stretch $O(2^r \cdot r) \cdot \log n$, thus we obtain an exponential improvement in the dependence on r . Interestingly, there is an $\Omega(\frac{\log n}{\log \log n})$ lower bound for the UST problem on the $n \times n$ grid, which is K_5 -minor free, thus essentially, in the context of K_r -minor free graphs, the dependence on r is the only parameter left to optimize.

It is important to note that the UST returned by [JLN⁺05] (and thus by Corollary 4) is not a subgraph of G . The (non-metric) UST problem was also studied and there is a similar reduction [BDR⁺12]. Formally, a strong (τ, σ, ρ) -sparse partition hierarchy is a set of laminar partitions $\mathcal{P}_1, \mathcal{P}_2, \dots$, such that \mathcal{P}_i refines \mathcal{P}_{i+1} , and each \mathcal{P}_i is a strong (τ, σ, ρ^i) -sparse partition. Busch *et al.* [BDR⁺12] showed that if a graph admits strong (τ, σ, ρ) -sparse partition then there is a subgraph solution to the UST problem with stretch $O(\tau^2 \cdot \sigma^2 \cdot \rho \cdot \log n)$. This reduction was recently used by Busch *et al.* [BCF⁺23] to obtain a solution for the UST problem on general graphs with stretch $O(\log^7 n)$. [BCF⁺23] also showed solution with stretch $O(\log n)$ for graph with constant doubling dimension or pathwidth. However, nothing better than general graphs is known for K_r -minor free graphs, or even planar graphs. One reason being that even trees do not admit better than $(\tilde{\Omega}(\log n), \tilde{\Omega}(\log n))$ -strong sparse partition scheme [Fil20b]. Improving the stretch factor for K_r -minor free graphs is a fascinating open problem.

7.3 Oblivious Buy-at-Bulk

In the *buy-at-bulk* problem we are given a weighted graph $G = (V, E, w)$, the goal is to satisfy a set of demands $A \subseteq \binom{V}{2}$, by routing these demands over the graph while minimizing the cost. In more details, $\delta_i = (s_i, t_i)$ is a unit of demand that induces an unsplittable unit of flow from source node $s_i \in V$ to a destination $t_i \in V$. Given a set of demands $A = \{\delta_1, \delta_2, \dots, \delta_k\}$, a valid solution is a set of paths $\mathcal{P} = \{P_1, \dots, P_k\}$, where P_i is a path from s_i to t_i . The paths can overlap. For an edge $e \in E$, denote by φ_e the number of paths in \mathcal{P} that use e . A function $f : \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$ is called *canonical fusion function* if it is (1) concave, (2) non-decreasing, (3) $f(0) = 0$, and (4) sub-additive (that is $\forall x, y \in \mathbb{N}, f(x + y) \leq f(x) + f(y)$). In the buy-at-bulk problem we are given a canonical fusion function f . The cost of a solution \mathcal{P} is $\text{cost}(\mathcal{P}) = \sum_{e \in E} f(\varphi_e) \cdot w(e)$. The goal is to find a valid solution of minimum cost. The canonical fusion function provides the following intuition: there is a discount as you use more and more of the same edge. So it is generally beneficial for the paths in \mathcal{P} to overlap. Note that the buy-at-bulk problem is NP-hard, as the Steiner tree problem is a special case (where the canonical fusion function is $f(0) = 0$ and $f(i) = 1$ for $i \geq 1$).

In the *oblivious buy-at-bulk* problem we have to choose a collection of paths \mathcal{P} without knowing the demands. That is, for every possible demand $\delta_i \in \binom{V}{2}$, we have to add to \mathcal{P} choose a path $P(\delta_i)$ between its endpoints. Then given a set of demands $A \subseteq \binom{V}{2}$, $\text{cost}(\mathcal{P}, A) = \sum_{e \in E} f(\varphi_e(\mathcal{P}, A)) \cdot w(e)$,

where $\varphi_e(\mathcal{P}, A) = |\{P(\delta_i) \mid e \in P(\delta_i) \ \& \ \delta_i \in A\}|$ is the number of paths associated with the demands in A that use e . The approximation ratio of \mathcal{P} , is the ratio between the induced cost, to the optimal cost for the worst possible set of demands A :

$$\text{Approximatio ratio}(\mathcal{P}) = \max_{A \subseteq \binom{V}{2}} \frac{\text{cost}(\mathcal{P}, A)}{\text{opt}(A)} .$$

Gupta *et al.* [GHR06] provided an algorithm achieving approximation ratio $O(\log^2 n)$. Interestingly, their algorithm is oblivious not only to the demand pairs A , but also to the canonical fusion function f . A lower bound of $\Omega(\log n)$ is known already for the case where the graph G is planar [IW91]. Later, Srinivasagopalan *et al.* [SBI11] improved the approximation ratio for the case of planar graphs to $O(\log n)$. Their solution is also oblivious to both demands and function. Srinivasagopalan *et al.* [SBI11] left it as an explicit open problem to “obtain efficient solutions to other related network topologies, such as minor-free graphs.” More than a decade later, and compared with general graphs, nothing better for K_r -minor free graphs is known.

Our contribution here is to answer the question from [SBI11] affirmatively. Specifically, that every K_r -minor free graph admits a solution with approximation ratio $\text{poly}(r) \cdot \log n$. This is tight up to the dependence on r .

Srinivasagopalan *et al.* [SBI11] implicitly proved that if a graph admits a “colorable” strong sparse cover scheme, than it also admits an efficiently computable solution with small approximation ratio for the oblivious buy-at-bulk problem.

Definition 6. Consider a strong (β, s, Δ) -sparse cover \mathcal{C} of $G = (V, E, w)$. We say that cluster $C \in \mathcal{C}$ β -satisfies a vertex $x \in V$ if the ball $B_G(x, \frac{\Delta}{\beta}) \subseteq C$ is contained in C . Two clusters $C_1, C_2 \in \mathcal{C}$ are neighbors if there is a pair of vertices $x \in C_1$ and $y \in C_2$ at distance at most $d_G(x, y) \leq \frac{\Delta}{\beta}$ such that x, y are β -satisfied by X_1, X_2 respectively. A k -coloring is a function $\chi: \mathcal{C} \rightarrow [k]$ such that no two neighboring clusters are colored using the same color.

Srinivasagopalan *et al.* [SBI11] showed that the strong sparse cover of [BLT14] for planar graphs is 18-colorable. Note that given a strong (β, s, Δ) -sparse partition cover, it is clearly s -colorable. Indeed, we can give all the clusters in each partition the same color. As the clusters in each partitions are all disjoint, this is a valid coloring. We conclude that our sparse covers from Theorem 1 are $O(r^2)$ -colorable.

Srinivasagopalan *et al.* [SBI11] showed that if a graph G admits a strong (β, s) -sparse cover scheme, where each cover is k -colorable, than one can efficiently compute a solution for the oblivious buy-at-bulk problem with approximation ratio $O(s \cdot \beta^2 \cdot k \cdot \log n)$. Interestingly, no colorable strong sparse covers for K_r -minor free graphs were known before. Hence the question from [SBI11] remained open until now. Using our Theorem 1 we conclude:

Corollary 5. For every n -vertex weighted K_r -minor free graph $G = (V, E, w)$ admits an efficiently computable solution to the oblivious buy-at-bulk problem with approximation ratio $O(r^6 \cdot \log n)$. Furthermore, the solution is also oblivious to the concave faction f .

Remark 1. The Universal Steiner tree, and Oblivious buy-at-bulk problem sound very similar. Indeed, one might be tempted to think that the Universal Steiner tree problem is a special case where there is only a single source, and the canonical function gets only the values $\{0, 1\}$. However, in the universal Steiner tree we have additional demand- the solution has to be a tree! Indeed, at present, there are significant gaps between the best solutions for this problems on both general, and K_r -minor free graphs.

7.4 Name-Independent Routing

A *compact routing scheme* in a network is a mechanism that allows packets to be delivered from any node to any other node. The network is represented as a undirected graph, and each node can forward incoming data by using local information stored at the node, called a *routing table*, and the (short) packet's *header*. The *stretch* of a routing scheme is the worst-case ratio between the length of a path on which a packet is routed to the shortest possible path. When designing a compact routing scheme our goal is to optimize the trade-off between the stretch and the size of the routing table and header. Here we focus on the name-independent model where each node is assigned an arbitrary unique network identifier, which cannot be chosen by the routing scheme designer. In addition, each edge will have an arbitrary port (that also cannot be changed). Note that there is a different regime of labeled routing, where the designer can choose the node labels and edge ports, see e.g. [TZ01, Tho04a, AG06, Che13, ACE+20, Fil21, FL21]. The labeled routing regime admits much better compact routing schemes. Indeed for general graphs, for every $k \in \mathbb{N}$ one can construct labeled routing scheme with stretch $3.68k$, routing table size $O(k \cdot n^{\frac{1}{k}})$, and label size $O(k \cdot \log n)$ (see also [ACE+20, FL21] for different trade-off's). In K_r -minor free graphs one can even construct labeled routing scheme with stretch $1 + \varepsilon$ and label and table size of $\varepsilon^{-1} \cdot f(r) \cdot \text{polylog}(n)$ ⁴ [AG06].

The name-independent regime is considered more practical, as it does not assumes that the sender knows the artificially chosen label of the destination. Further, it can cope much better with changes in the network. There are also applications that pose constrains on nodes addresses that are not easily satisfied by the artificial smartly chosen labels (such as distributed hash tables). However, the name-independent regime is much more challenging. Indeed, any name-independent routing scheme on unweighted stars (trees of depth 1) requires table size of $\Omega(n \cdot \log n)$ bits to get stretch bellow 3 [AGM+08]. The situation in weighted stars is even more dire, as every name-independent routing scheme with stretch bellow $2k + 1$ (for $k \geq 1$) requires routing tables of at least $\Omega((n \cdot \log n)^{\frac{1}{k}})$ bits [AGM06]. We will thus focus on unweighted graphs.

Abraham *et al.* [AGMW10] used their strong sparse covers, as well as name-independent routing scheme for trees [AGM04], to design a name-independent routing scheme for K_r -minor free graphs. We can use our new strong sparse cover to improve the various parameters in this routing scheme. In the following we elaborate on that.

A hereditary graph family is a family \mathcal{F} such that for every $G \in \mathcal{F}$, every subgraph H of G also belongs to $H \in \mathcal{F}$. A graph posses an α -orientation if it is possible to direct all edges such that every vertex has out-degree at most α . A family \mathcal{F} has an α -orientation if every $G \in \mathcal{F}$ has α -orientation. [AGMW10] implicitly proved the following meta theorem: consider a hereditary graph family \mathcal{F} that has an α -orientation, and let $G \in \mathcal{F}$ be an unweighted graph with diameter D that admits a strong (τ, β) -sparse cover scheme. Then there is a polynomial time constructible name-independent routing seheme, in the fixed port model with stretch $O(\beta)$, and using $O(\log n + \log \tau)$ -bit headers, in which every node requires tables of $O(\frac{\log^3 n}{\log \log n} + \alpha \cdot \log n) \cdot \tau \cdot \log D$ bits.

It is well known that the hereditary graph family of K_r minor free graphs has an $O(r\sqrt{\log r})$ -orientation (see e.g. [AGMW10]). Combining it with our [Theorem 1](#) we conclude

Corollary 6. *For every n vertex unweighted K_r -minor free graph G with diameter D , there is a polynomial time constructible name-independent routing scheme, in the fixed port model, with:*

- *Stretch $O(r)$, $O(\log n)$ bit headers, and $O(\frac{\log^3 n}{\log \log n} + r\sqrt{\log r} \cdot \log n) \cdot r^2 \cdot \log D$ bit table.*

	Stretch	Header	Table	Ref
(1)	$O(r^2)$	$O(\log n + r \log r)$	$O(1)^r \cdot r! \cdot \frac{\log^3 n}{\log \log n} \cdot \log D$	[AGMW10]
(2)	$O(1)$		$O(f(r) \cdot \log^3 n)$	[BLT14]
(3)	$O(r)$	$O(\log n)$	$O(\frac{\log^3 n}{\log \log n} + r\sqrt{\log r \cdot \log n}) \cdot r^2 \cdot \log D$	Corollary 6
(4)	$O(1)$	$O(r + \log n)$	$O(1)^r \cdot \frac{\log^3 n}{\log \log n} \cdot \log D$	Corollary 6

Table 3: Summary of new and previous work on name-independent routing schemes in K_r -minor free graphs in the fixed parameter model. The input graph are unweighted, and D denotes its diameter. $f(r)$ is an extremely fast growing function ⁴. In [BLT14], the header size is not explicitly stated, and the bound on the table is only in expectation. Comparing our result (3) with [AGMW10] we quadratically improved the stretch, and exponentially improved the dependence on r in the routing table size. Comparing [AGMW10] with (4) we improved the stretch from $O(r^2)$ to constant, and somewhat improved the dependence on r in both table size and header. Comparing (4) with [BLT14], we very significantly improved the table size, and improved the size bound to hold in the worst case.

- *Stretch $O(1)$, $O(r + \log n)$ bit headers, and $O(1)^r \cdot \frac{\log^3 n}{\log \log n} \cdot \log D$ bit table.*

See Table 3 for comparison of Corollary 6 with previous results.

7.5 Path-Reporting Distance Oracle

Given a weighted graph $G = (V, E, w)$, a path-reporting distance oracle (PRDO) is a succinct data structure that given a query $\{u, v\} \in \binom{V}{2}$ quickly returns a $u - v$ path which is an approximate $u - v$ shortest path. Formally we say that a distance oracle has stretch k if on query $\{u, v\}$, it returns a $u - v$ path P of weight at most $k \cdot d_G(u, v)$. We say that the query time is t , if the time it takes the distance oracle to return a path P is at most $O(|P|) + t$. In the study of PRDO's [EP16, ENW16, ES23], the goal is to optimize the trade-off between stretch, query time, and space.

Elkin, Neiman and Wulff-Nilsen [ENW16] constructed a PRDO using strong sparse covers with very small query time. Given a weighted graph $G = (V, E, w)$ with aspect ratio Φ that admits a strong (β, s) -sparse cover scheme, [ENW16] implicitly constructed a PRDO with stretch $O(\beta)$, space $O(s \cdot \log \Phi)$ (in machine words), and query time $O(\log \log \Phi)$. In fact, by decreasing the gap between the different scales in their construction to $1 + \varepsilon$, their proof will lead to a PRDO with stretch $(1 + \varepsilon) \cdot \beta$, space $O(\frac{s}{\varepsilon} \cdot \log \Phi)$, and query time $O(\log \log \Phi + \log \frac{1}{\varepsilon})$.

Elkin *et al.* [ENW16] used the strong $(O(r^2), O(\log n))$ -sparse cover scheme that follows from the padded decomposition of [AGG⁺19] to construct their PRDO for K_r -minor free graphs with stretch $O(r^2)$, size $O(n \cdot \log n \cdot \log \Phi)$ and query time $O(\log \log \Phi)$. Later, Filtser [Fil19a] used his improved strong padded decomposition to obtain a strong $(O(r), O(\log n))$ -sparse cover scheme, and plugged it into the PRDO construction of [ENW16], improving the stretch parameter from $O(r^2)$ to $O(r)$. Using our strong $(O(r), O(r^2))$ -sparse cover scheme (Theorem 1), we can get a further improvement, reducing the space to $O(n \cdot r^2 \cdot \log \Phi)$. Furthermore, using our strong $(4 + \varepsilon, O(\frac{1}{\varepsilon})^r)$ -sparse cover scheme (Theorem 1), and the observations made in the proof of the contraction argument in Lemma 10, we can obtain a PRDO with stretch $3 + \varepsilon$, size $n \cdot O(\frac{1}{\varepsilon})^{r+1} \cdot \log \Phi$ and query time $O(\log \log \Phi + \log \frac{1}{\varepsilon})$.

Corollary 7. *Every n -vertex K_r -minor free weighted graph G with aspect ratio Φ , admits a path-reporting distance oracle with:*

- Stretch $O(r)$, size $O(n \cdot r^2 \cdot \log \Phi)$ and query time $O(\log \log \Phi)$.
- Stretch $3 + \varepsilon$, size $n \cdot O(\frac{1}{\varepsilon})^{r+1} \cdot \log \Phi$ and query time $O(\log \log \Phi + \log \frac{1}{\varepsilon})$.

8 Conclusion and Open Problems

This paper is mainly concerned with sparse covers for minor free graphs. Our first contribution is to transform from the recently introduced notion of buffered cop decomposition [CCL⁺24] to sparse covers. As a result, we obtain a strong $(O(r), O(r^2))$ -SPCS, and strong $(4 + \varepsilon, O(\frac{1}{\varepsilon})^r)$ -SPCS for K_r -minor free graphs (Theorem 1). This significantly improves both sparsity and padding parameters compared to previous work, obtains the strong diameter guarantee, and the “partition based” property. We then use this new covers to construct low distortion and dimension ℓ_∞ embedding of K_r minor free graphs (Corollary 1, and Theorem 2), improving significantly over the previous work [KLMN05] in both dimension and distortion. Next, three decades after the notions of sparse covers and padded decompositions were introduced [AP91, KPR93], we observe that sparse covers imply padded decomposition (Theorem 3), and recover the state of the art padded decomposition for minor free graphs [AGG⁺19]. Finally, we show several applications of our new sparse covers. Specifically, we obtained better sparse partitions, and as a result get an exponential improvement in the dependence on r for both universal TSP, and metric universal Steiner tree problems. We also get a similar exponential improvement for the oblivious buy-at-bulk problem. For the name independent routing scheme in K_r -minor free graphs, we get quadratic improvement in the stretch and exponential improvement in the space (in the dependence on r) compared with previous work [AGMW10] (or just ginormous improvement in the space compared with [BLT14]). We also obtain similar improvements for path reporting distance oracles.

Our work leaves several open questions:

1. The main open question is to obtain improve sparse covers for K_r -minor free graphs. Note that we don’t have any lower bound beyond what follows from general graphs. In particular, it might be possible to obtain $(O(\log r), O(\log r))$ -sparse cover scheme, or $(O(1), \text{poly}(r))$ -sparse cover scheme. Due to our Theorem 3, such sparse covers will imply padded decomposition with parameter $\tilde{O}(\log r)$, solving the conjecture from [AGG⁺19].
2. Metric embeddings into ℓ_∞ . In Theorem 2 we got distortion $3 + \varepsilon$ which looks like a natural barrier for sparse cover based techniques. It is very interesting to see if we can get below 3, even for planar graphs. Optimally, we would like to see an embedding with distortion $1 + \varepsilon$ and $g(r, \varepsilon) \cdot \log n$ coordinated, for an arbitrary function g .
3. Universal Steiner tree. In this paper, using the reduction from [JLN⁺05] we obtain a solution for the metric UST problem in K_r -minor free graphs with stretch $O(\frac{r^4}{\log r}) \cdot \log n$. However, for the usual (non-metric) UST problem, the state of the art for K_r -minor free (or even planar) graphs is $O(\log^7 n)$ [BCF⁺23], the same as for general graphs. It is interesting to see if we can use the topological structure to get an improved stretch.

Acknowledgments

The author would like to thank James R Lee for helpful discussions regarding the embedding of planar graphs into ℓ_∞ from [KLMN05]. The author would also like to thank Shmuel Tomi Klein

for the reference to [KN76].

References

- [ABN11] I. Abraham, Y. Bartal, and O. Neiman. Advances in metric embedding theory. *Advances in Mathematics*, 228(6):3026–3126, 2011, doi:<https://doi.org/10.1016/j.aim.2011.08.003>. 3, 4, 6
- [ACE⁺20] I. Abraham, S. Chechik, M. Elkin, A. Filtser, and O. Neiman. Ramsey spanning trees and their applications. *ACM Trans. Algorithms*, 16(2):19:1–19:21, 2020. preliminary version published in SODA 2018, doi:[10.1145/3371039](https://doi.org/10.1145/3371039). 31
- [AFGN22] I. Abraham, A. Filtser, A. Gupta, and O. Neiman. Metric embedding via shortest path decompositions. *SIAM J. Comput.*, 51(2):290–314, 2022. a preliminary version appeared in the proceedings of STOC 18, doi:[10.1137/19m1296021](https://doi.org/10.1137/19m1296021). 6
- [AG06] I. Abraham and C. Gavoille. Object location using path separators. In *Proceedings of the Twenty-fifth Annual ACM Symposium on Principles of Distributed Computing*, PODC '06, pages 188–197, 2006. Full version: <https://www.cse.huji.ac.il/~ittaia/papers/AG-TR.pdf>, doi:[10.1145/1146381.1146411](https://doi.org/10.1145/1146381.1146411). 8, 31
- [AGG⁺14] I. Abraham, C. Gavoille, A. Gupta, O. Neiman, and K. Talwar. Cops, robbers, and threatening skeletons: Padded decomposition for minor-free graphs. In *Proceedings of the 46th Annual ACM Symposium on Theory of Computing*, STOC '14, page 79–88, 2014, doi:[10.1145/2591796.2591849](https://doi.org/10.1145/2591796.2591849). 2
- [AGG⁺19] I. Abraham, C. Gavoille, A. Gupta, O. Neiman, and K. Talwar. Cops, robbers, and threatening skeletons: Padded decomposition for minor-free graphs. *SIAM J. Comput.*, 48(3):1120–1145, 2019. preliminary version published in STOC 2014, doi:[10.1137/17M1112406](https://doi.org/10.1137/17M1112406). 2, 4, 5, 7, 8, 9, 10, 12, 25, 32, 33
- [AGM04] I. Abraham, C. Gavoille, and D. Malkhi. Routing with improved communication-space trade-off. In R. Guerraoui, editor, *Distributed Computing, 18th International Conference, DISC 2004, Amsterdam, The Netherlands, October 4-7, 2004, Proceedings*, volume 3274 of *Lecture Notes in Computer Science*, pages 305–319. Springer, 2004, doi:[10.1007/978-3-540-30186-8_22](https://doi.org/10.1007/978-3-540-30186-8_22). 31
- [AGM05] I. Abraham, C. Gavoille, and D. Malkhi. Compact routing for graphs excluding a fixed minor. In P. Fraigniaud, editor, *Distributed Computing, 19th International Conference, DISC 2005, Cracow, Poland, September 26-29, 2005, Proceedings*, volume 3724 of *Lecture Notes in Computer Science*, pages 442–456. Springer, 2005, doi:[10.1007/11561927_32](https://doi.org/10.1007/11561927_32). 1
- [AGM06] I. Abraham, C. Gavoille, and D. Malkhi. On space-stretch trade-offs: lower bounds. In P. B. Gibbons and U. Vishkin, editors, *SPAA 2006: Proceedings of the 18th Annual ACM Symposium on Parallelism in Algorithms and Architectures, Cambridge, Massachusetts, USA, July 30 - August 2, 2006*, pages 207–216. ACM, 2006, doi:[10.1145/1148109.1148143](https://doi.org/10.1145/1148109.1148143). 31
- [AGM⁺08] I. Abraham, C. Gavoille, D. Malkhi, N. Nisan, and M. Thorup. Compact name-independent routing with minimum stretch. *ACM Trans. Algorithms*, 4(3):37:1–37:12, 2008, doi:[10.1145/1367064.1367077](https://doi.org/10.1145/1367064.1367077). 1, 31
- [AGMW10] I. Abraham, C. Gavoille, D. Malkhi, and U. Wieder. Strong-diameter decompositions of minor free graphs. *Theory of Computing Systems*, 47(4):837–855, 2010, doi:[10.1007/s00224-010-9283-6](https://doi.org/10.1007/s00224-010-9283-6). 1, 2, 5, 8, 31, 32, 33
- [AKP94] B. Awerbuch, S. Kutten, and D. Peleg. On buffer-economical store-and-forward deadlock prevention. *IEEE Trans. Commun.*, 42(11):2934–2937, 1994, doi:[10.1109/26.328973](https://doi.org/10.1109/26.328973). 1

- [AKPW95] N. Alon, R. M. Karp, D. Peleg, and D. B. West. A graph-theoretic game and its application to the k-server problem. *SIAM J. Comput.*, 24(1):78–100, 1995. preliminary version published in On-Line Algorithms 1991, doi:10.1137/S0097539792224474. 6
- [And86] T. Andreae. On a pursuit game played on graphs for which a minor is excluded. *J. Comb. Theory, Ser. B*, 41(1):37–47, 1986, doi:10.1016/0095-8956(86)90026-2. 7, 12
- [AP90a] B. Awerbuch and D. Peleg. Network synchronization with polylogarithmic overhead. In *Proc. 31st IEEE Symp. on Foundations of Computer Science*, pages 514–522, 1990. 1
- [AP90b] B. Awerbuch and D. Peleg. Sparse partitions. In *Proceedings of the 31st IEEE Symposium on Foundations of Computer Science (FOCS)*, pages 503–513, 1990, doi:10.1109/FSCS.1990.89571. 1, 2
- [AP91] B. Awerbuch and D. Peleg. Concurrent online tracking of mobile users. In L. Chapin, editor, *Proceedings of the Conference on Communications Architecture & Protocols, SIGCOMM 1991, Zürich, Switzerland, September 3-6, 1991*, pages 221–233. ACM, 1991, doi:10.1145/115992.116013. 1, 33
- [AW04] H. Attiya and J. L. Welch. *Distributed computing - fundamentals, simulations, and advanced topics (2. ed.)*. Wiley series on parallel and distributed computing. Wiley, 2004. 1
- [Bar96] Y. Bartal. Probabilistic approximations of metric spaces and its algorithmic applications. In *37th Annual Symposium on Foundations of Computer Science, FOCS '96, Burlington, Vermont, USA, 14-16 October, 1996*, pages 184–193, 1996, doi:10.1109/SFCS.1996.548477. 4, 6
- [BCF⁺23] O. Busch, D. Q. Chen, A. Filtser, D. Hathcock, D. E. Hershkowitz, and R. Rajaraman. One tree to rule them all: Poly-logarithmic universal steiner tree. In *2023 IEEE 64th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 60–76, 2023, doi:10.1109/FOCS57990.2023.00012. 1, 28, 29, 33
- [BDR⁺12] C. Busch, C. Dutta, J. Radhakrishnan, R. Rajaraman, and S. Srivathsan. Split and join: Strong partitions and universal steiner trees for graphs. In *53rd Annual IEEE Symposium on Foundations of Computer Science, FOCS 2012, New Brunswick, NJ, USA, October 20-23, 2012*, pages 81–90. IEEE Computer Society, 2012, doi:10.1109/FOCS.2012.45. 1, 28, 29
- [BFN19a] Y. Bartal, N. Fandina, and O. Neiman. Covering metric spaces by few trees. In *46th International Colloquium on Automata, Languages, and Programming, ICALP 2019, July 9-12, 2019, Patras, Greece*, pages 20:1–20:16, 2019, doi:10.4230/LIPIcs.ICALP.2019.20. 8
- [BFN19b] Y. Bartal, A. Filtser, and O. Neiman. On notions of distortion and an almost minimum spanning tree with constant average distortion. *J. Comput. Syst. Sci.*, 105:116–129, 2019. preliminary version published in SODA 2016, doi:10.1016/j.jcss.2019.04.006. 6
- [BFT24] S. Bhore, A. Filtser, and C. D. Toth. Online duet between metric embeddings and minimum-weight perfect matchings. In *Proceedings of the 2024 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 4564–4579, 2024, arXiv:https://epubs.siam.org/doi/pdf/10.1137/1.9781611977912.162, doi:10.1137/1.9781611977912.162. 6
- [BG19] Y. Bartal and L. Gottlieb. Approximate nearest neighbor search for ℓ_p -spaces ($2 < p < \infty$). *Theor. Comput. Sci.*, 757:27–35, 2019, doi:10.1016/J.TCS.2018.07.011. 3
- [BLT14] C. Busch, R. LaFortune, and S. Tirthapura. Sparse covers for planar graphs and graphs that exclude a fixed minor. *Algorithmica*, 69(3):658–684, 2014, doi:10.1007/S00453-013-9757-4. 1, 2, 30, 32, 33
- [Bou85] J. Bourgain. On lipschitz embedding of finite metric spaces in hilbert space. *Israel Journal of Mathematics*, 52(1-2):46–52, 1985, doi:10.1007/BF02776078. 3, 6

- [CCL⁺23] H.-C. Chang, J. Conroy, H. Le, L. Milenkovic, S. Solomon, and C. Than. Covering planar metrics (and beyond): $O(1)$ trees suffice. In *2023 IEEE 64th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 2231–2261, 2023, doi:[10.1109/FOCS57990.2023.00139](https://doi.org/10.1109/FOCS57990.2023.00139). 6, 8
- [CCL⁺24] H.-C. Chang, J. Conroy, H. Le, L. Milenković, S. Solomon, and C. Than. Shortcut partitions in minor-free graphs: Steiner point removal, distance oracles, tree covers, and more. In *Proceedings of the 2024 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 5300–5331, 2024, arXiv:<https://epubs.siam.org/doi/pdf/10.1137/1.9781611977912.191>, doi:[10.1137/1.9781611977912.191](https://doi.org/10.1137/1.9781611977912.191). 8, 11, 12, 16, 33
- [CFJ⁺22] A. Czumaj, A. Filtser, S. H. Jiang, R. Krauthgamer, P. Veselý, and M. Yang. Streaming facility location in high dimension via new geometric hashing. *CoRR*, abs/2204.02095, 2022. conference version published in STOC 2022, arXiv:[2204.02095](https://arxiv.org/abs/2204.02095), doi:[10.48550/ARXIV.2204.02095](https://doi.org/10.48550/ARXIV.2204.02095). 28
- [CFKL20] V. Cohen-Addad, A. Filtser, P. N. Klein, and H. Le. On light spanners, low-treewidth embeddings and efficient traversing in minor-free graphs. *CoRR*, abs/2009.05039, 2020. To appear in FOCS 2020, <https://arxiv.org/abs/2009.05039>, arXiv:[2009.05039](https://arxiv.org/abs/2009.05039). 6, 8
- [Che13] S. Chechik. Compact routing schemes with improved stretch. In P. Fatourou and G. Taubenfeld, editors, *ACM Symposium on Principles of Distributed Computing, PODC '13, Montreal, QC, Canada, July 22-24, 2013*, pages 33–41. ACM, 2013, doi:[10.1145/2484239.2484268](https://doi.org/10.1145/2484239.2484268). 31
- [Che18] Y. K. Cheung. Steiner point removal - distant terminals don't (really) bother. In A. Czumaj, editor, *Proceedings of the Twenty-Ninth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2018, New Orleans, LA, USA, January 7-10, 2018*, pages 1353–1360. SIAM, 2018, doi:[10.1137/1.9781611975031.89](https://doi.org/10.1137/1.9781611975031.89). 8
- [CLPP23] V. Cohen-Addad, H. Le, M. Pilipczuk, and M. Pilipczuk. Planar and minor-free metrics embed into metrics of polylogarithmic treewidth with expected multiplicative distortion arbitrarily close to 1^* . In *2023 IEEE 64th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 2262–2277, 2023, doi:[10.1109/FOCS57990.2023.00140](https://doi.org/10.1109/FOCS57990.2023.00140). 6
- [EFN17] M. Elkin, A. Filtser, and O. Neiman. Terminal embeddings. *Theoretical Computer Science*, 697:1 – 36, 2017, doi:<https://doi.org/10.1016/j.tcs.2017.06.021>. 6
- [EFN18] M. Elkin, A. Filtser, and O. Neiman. Prioritized metric structures and embedding. *SIAM J. Comput.*, 47(3):829–858, 2018, doi:[10.1137/17M1118749](https://doi.org/10.1137/17M1118749). 6
- [EN22] M. Elkin and O. Neiman. Lossless prioritized embeddings. *SIAM J. Discret. Math.*, 36(3):1529–1550, 2022, doi:[10.1137/21M1436221](https://doi.org/10.1137/21M1436221). 6
- [ENW16] M. Elkin, O. Neiman, and C. Wulff-Nilsen. Space-efficient path-reporting approximate distance oracles. *Theor. Comput. Sci.*, 651:1–10, 2016, doi:[10.1016/j.tcs.2016.07.038](https://doi.org/10.1016/j.tcs.2016.07.038). 6, 32
- [EP16] M. Elkin and S. Pettie. A linear-size logarithmic stretch path-reporting distance oracle for general graphs. *ACM Trans. Algorithms*, 12(4):50:1–50:31, 2016, doi:[10.1145/2888397](https://doi.org/10.1145/2888397). 32
- [Erd64] P. Erdős. Extremal problems in graph theory. In *in "Theory of Graphs and Its Applications," Proc. Sympos. Smolenice*, pages 29–36, 1964. 3
- [ES23] M. Elkin and I. Shabat. Path-reporting distance oracles with logarithmic stretch and size $o(n \log \log n)$. In *2023 IEEE 64th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 2278–2311, 2023, doi:[10.1109/FOCS57990.2023.00141](https://doi.org/10.1109/FOCS57990.2023.00141). 32
- [FGK23] A. Filtser, L. Gottlieb, and R. Krauthgamer. Labelings vs. embeddings: On distributed and prioritized representations of distances. *Discret. Comput. Geom.*, 2023, doi:[10.1007/s00454-023-00565-2](https://doi.org/10.1007/s00454-023-00565-2). 6

- [FGN23] A. Filtser, Y. Gitlitz, and O. Neiman. Light, reliable spanners. *CoRR*, abs/2307.16612, 2023, [arXiv:2307.16612](https://arxiv.org/abs/2307.16612), [doi:10.48550/ARXIV.2307.16612](https://doi.org/10.48550/ARXIV.2307.16612). 1
- [FI99] M. Farach-Colton and P. Indyk. Approximate nearest neighbor algorithms for hausdorff metrics via embeddings. In *40th Annual Symposium on Foundations of Computer Science, FOCS '99, 17-18 October, 1999, New York, NY, USA*, pages 171–180. IEEE Computer Society, 1999, [doi:10.1109/SFFCS.1999.814589](https://doi.org/10.1109/SFFCS.1999.814589). 3
- [Fil19a] A. Filtser. On strong diameter padded decompositions. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques, APPROX/RANDOM 2019, September 20-22, 2019, Massachusetts Institute of Technology, Cambridge, MA, USA*, pages 6:1–6:21, 2019, [doi:10.4230/LIPIcs.APPROX-RANDOM.2019.6.2](https://doi.org/10.4230/LIPIcs.APPROX-RANDOM.2019.6.2), 5, 8, 9, 12, 25, 32
- [Fil19b] A. Filtser. Steiner point removal with distortion $O(\log k)$ using the relaxed-voronoi algorithm. *SIAM J. Comput.*, 48(2):249–278, 2019, [doi:10.1137/18M1184400](https://doi.org/10.1137/18M1184400). 8
- [Fil20a] A. Filtser. A face cover perspective to ℓ_1 embeddings of planar graphs. In S. Chawla, editor, *Proceedings of the 2020 ACM-SIAM Symposium on Discrete Algorithms, SODA 2020, Salt Lake City, UT, USA, January 5-8, 2020*, pages 1945–1954. SIAM, 2020, [doi:10.1137/1.9781611975994.120.6](https://doi.org/10.1137/1.9781611975994.120.6), 28
- [Fil20b] A. Filtser. Scattering and sparse partitions, and their applications. In A. Czumaj, A. Dawar, and E. Merelli, editors, *47th International Colloquium on Automata, Languages, and Programming, ICALP 2020, July 8-11, 2020, Saarbrücken, Germany (Virtual Conference)*, volume 168 of *LIPIcs*, pages 47:1–47:20. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020, [doi:10.4230/LIPIcs.ICALP.2020.47.1](https://doi.org/10.4230/LIPIcs.ICALP.2020.47.1), 2, 5, 8, 29
- [Fil21] A. Filtser. Hop-constrained metric embeddings and their applications. In *62nd IEEE Annual Symposium on Foundations of Computer Science, FOCS 2021, Denver, CO, USA, February 7-10, 2022*, pages 492–503. IEEE, 2021, [doi:10.1109/FOCS52979.2021.00056](https://doi.org/10.1109/FOCS52979.2021.00056). 31
- [Fil23] A. Filtser. Labeled nearest neighbor search and metric spanners via locality sensitive orderings. In E. W. Chambers and J. Gudmundsson, editors, *39th International Symposium on Computational Geometry, SoCG 2023, June 12-15, 2023, Dallas, Texas, USA*, volume 258 of *LIPIcs*, pages 33:1–33:18. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2023, [doi:10.4230/LIPIcs.SoCG.2023.33.1](https://doi.org/10.4230/LIPIcs.SoCG.2023.33.1)
- [FKS19] E. Fox-Epstein, P. N. Klein, and A. Schild. Embedding planar graphs into low-treewidth graphs with applications to efficient approximation schemes for metric problems. In *Proceedings of the 30th Annual ACM-SIAM Symposium on Discrete Algorithms, SODA '19*, page 1069–1088, 2019, [doi:10.1137/1.9781611975482.66.6](https://doi.org/10.1137/1.9781611975482.66.6), 8
- [FKT19] A. Filtser, R. Krauthgamer, and O. Trabelsi. Relaxed voronoi: A simple framework for terminal-clustering problems. In J. T. Fineman and M. Mitzenmacher, editors, *2nd Symposium on Simplicity in Algorithms, SOSA 2019, January 8-9, 2019, San Diego, CA, USA*, volume 69 of *OASICS*, pages 10:1–10:14. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2019, [doi:10.4230/OASICS.SOSA.2019.10.8](https://doi.org/10.4230/OASICS.SOSA.2019.10.8)
- [FL21] A. Filtser and H. Le. Clan embeddings into trees, and low treewidth graphs. In S. Khuller and V. V. Williams, editors, *STOC '21: 53rd Annual ACM SIGACT Symposium on Theory of Computing, Virtual Event, Italy, June 21-25, 2021*, pages 342–355. ACM, 2021, [doi:10.1145/3406325.3451043](https://doi.org/10.1145/3406325.3451043). 8, 31
- [FL22a] A. Filtser and H. Le. Locality-sensitive orderings and applications to reliable spanners. In S. Leonardi and A. Gupta, editors, *STOC '22: 54th Annual ACM SIGACT Symposium on Theory of Computing, Rome, Italy, June 20 - 24, 2022*, pages 1066–1079. ACM, 2022, [doi:10.1145/3519935.3520042](https://doi.org/10.1145/3519935.3520042). 1, 6, 17

- [FL22b] A. Filtser and H. Le. Low treewidth embeddings of planar and minor-free metrics. In *63rd IEEE Annual Symposium on Foundations of Computer Science, FOCS 2022, Denver, CO, USA, October 31 - November 3, 2022*, pages 1081–1092. IEEE, 2022, doi:[10.1109/FOCS54457.2022.00105](https://doi.org/10.1109/FOCS54457.2022.00105). 6, 8
- [FN22] A. Filtser and O. Neiman. Light spanners for high dimensional norms via stochastic decompositions. *Algorithmica*, 2022, doi:[10.1007/s00453-022-00994-0](https://doi.org/10.1007/s00453-022-00994-0). 1
- [FRT04] J. Fakcharoenphol, S. Rao, and K. Talwar. A tight bound on approximating arbitrary metrics by tree metrics. *J. Comput. Syst. Sci.*, 69(3):485–497, November 2004. preliminary version published in STOC 2003, doi:[10.1016/j.jcss.2004.04.011](https://doi.org/10.1016/j.jcss.2004.04.011). 6
- [FT03] J. Fakcharoenphol and K. Talwar. An improved decomposition theorem for graphs excluding a fixed minor. In *RANDOM-APPROX*, pages 36–46, 2003. 1, 2, 3, 5, 28
- [GHR06] A. Gupta, M. T. Hajiaghayi, and H. Räcke. Oblivious network design. In *Proceedings of the Seventeenth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2006, Miami, Florida, USA, January 22-26, 2006*, pages 970–979. ACM Press, 2006. 30
- [GNRS04] A. Gupta, I. Newman, Y. Rabinovich, and A. Sinclair. Cuts, trees and l_1 -embeddings of graphs. *Comb.*, 24(2):233–269, 2004, doi:[10.1007/S00493-004-0015-X](https://doi.org/10.1007/S00493-004-0015-X). 6
- [HIS13] S. Har-Peled, P. Indyk, and A. Sidiropoulos. Euclidean spanners in high dimensions. In *Proceedings of the Twenty-Fourth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2013, New Orleans, Louisiana, USA, January 6-8, 2013*, pages 804–809, 2013, doi:[10.1137/1.9781611973105.57](https://doi.org/10.1137/1.9781611973105.57). 1
- [HMO23] S. Har-Peled, M. Mendel, and D. Oláh. Reliable spanners for metric spaces. *ACM Trans. Algorithms*, 19(1):7:1–7:27, 2023, doi:[10.1145/3563356](https://doi.org/10.1145/3563356). 1
- [Huf52] D. A. Huffman. A method for the construction of minimum-redundancy codes. *Proceedings of the IRE*, 40(9):1098–1101, 1952, doi:[10.1109/JRPROC.1952.273898](https://doi.org/10.1109/JRPROC.1952.273898). 41
- [Hwa76] F. K. Hwang. On steiner minimal trees with rectilinear distance. *SIAM Journal on Applied Mathematics*, 30(1):104–114, 1976, arXiv:<https://doi.org/10.1137/0130013>, doi:[10.1137/0130013](https://doi.org/10.1137/0130013). 28
- [IMSZ10] P. Indyk, A. Magen, A. Sidiropoulos, and A. Zouzias. Online embeddings. In M. J. Serna, R. Shaltiel, K. Jansen, and J. D. P. Rolim, editors, *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques, 13th International Workshop, APPROX 2010, and 14th International Workshop, RANDOM 2010, Barcelona, Spain, September 1-3, 2010. Proceedings*, volume 6302 of *Lecture Notes in Computer Science*, pages 246–259. Springer, 2010, doi:[10.1007/978-3-642-15369-3_19](https://doi.org/10.1007/978-3-642-15369-3_19). 6
- [Ind98] P. Indyk. On approximate nearest neighbors in non-euclidean spaces. In *39th Annual Symposium on Foundations of Computer Science, FOCS '98, November 8-11, 1998, Palo Alto, California, USA*, pages 148–155. IEEE Computer Society, 1998, doi:[10.1109/SFCS.1998.743438](https://doi.org/10.1109/SFCS.1998.743438). 3
- [Ind02] P. Indyk. Approximate nearest neighbor algorithms for Fréchet distance via product metrics. In *Proceedings of the 8th Symposium on Computational Geometry*, pages 102–106, Barcelona, Spain, June 2002. ACM Press, doi:[10.1145/513400.513414](https://doi.org/10.1145/513400.513414). 3
- [IW91] M. Imase and B. M. Waxman. Dynamic steiner tree problem. *SIAM J. Discret. Math.*, 4(3):369–384, 1991, doi:[10.1137/0404033](https://doi.org/10.1137/0404033). 30
- [JL84] W. Johnson and J. Lindenstrauss. Extensions of Lipschitz mappings into a Hilbert space. *Contemporary Mathematics*, 26:189–206, 1984. see [here](#). 10

- [JLN⁺05] L. Jia, G. Lin, G. Noubir, R. Rajaraman, and R. Sundaram. Universal approximations for tsp, steiner tree, and set cover. In H. N. Gabow and R. Fagin, editors, *Proceedings of the 37th Annual ACM Symposium on Theory of Computing, Baltimore, MD, USA, May 22-24, 2005*, pages 386–395. ACM, 2005, doi:[10.1145/1060590.1060649](https://doi.org/10.1145/1060590.1060649). 1, 5, 28, 29, 33
- [Joh87] D. S. Johnson. The NP-completeness column: An ongoing guide (column 19). *Journal of Algorithms*, 8(3):438–448, 1987. 2
- [KKN15] L. Kamma, R. Krauthgamer, and H. L. Nguyen. Cutting corners cheaply, or how to remove steiner points. *SIAM J. Comput.*, 44(4):975–995, 2015, doi:[10.1137/140951382](https://doi.org/10.1137/140951382). 8
- [Kle02] P. N. Klein. Preprocessing an undirected planar network to enable fast approximate distance queries. In *Proceedings of the Thirteenth Annual ACM-SIAM Symposium on Discrete Algorithms, January 6-8, 2002, San Francisco, CA, USA*, pages 820–827, 2002. see [here](#). 8
- [KLMN05] R. Krauthgamer, J. R. Lee, M. Mendel, and A. Naor. Measured descent: a new embedding method for finite metrics. *Geometric and Functional Analysis*, 15(4):839–858, 2005. preliminary version published in FOCS 2004, doi:[10.1007/s00039-005-0527-6](https://doi.org/10.1007/s00039-005-0527-6). 1, 2, 3, 4, 8, 10, 17, 33
- [KLR19] R. Krauthgamer, J. R. Lee, and H. Rika. Flow-cut gaps and face covers in planar graphs. In T. M. Chan, editor, *Proceedings of the Thirtieth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2019, San Diego, California, USA, January 6-9, 2019*, pages 525–534. SIAM, 2019, doi:[10.1137/1.9781611975482.33](https://doi.org/10.1137/1.9781611975482.33). 6
- [KN76] G. O. H. Katona and T. O. H. Nemetz. Huffman codes and self-information. *IEEE Trans. Inf. Theory*, 22(3):337–340, 1976, doi:[10.1109/TIT.1976.1055554](https://doi.org/10.1109/TIT.1976.1055554). 17, 34
- [KPR93] P. Klein, S. A. Plotkin, and S. Rao. Excluded minors, network decomposition, and multicommodity flow. In *Proceedings of the twenty-fifth annual ACM symposium on Theory of computing, STOC '93*, pages 682–690, New York, NY, USA, 1993. ACM, doi:<http://doi.acm.org/10.1145/167088.167261>. 1, 2, 3, 5, 8, 9, 10, 28, 33
- [Kum22] N. Kumar. An approximate generalization of the okamura-seymour theorem. In *63rd IEEE Annual Symposium on Foundations of Computer Science, FOCS 2022, Denver, CO, USA, October 31 - November 3, 2022*, pages 1093–1101. IEEE, 2022, doi:[10.1109/FOCS54457.2022.00106](https://doi.org/10.1109/FOCS54457.2022.00106). 6
- [LLR95] N. Linial, E. London, and Y. Rabinovich. The geometry of graphs and some of its algorithmic applications. *Comb.*, 15(2):215–245, 1995. preliminary version published in FOCS 1994, doi:[10.1007/BF01200757](https://doi.org/10.1007/BF01200757). 3, 4, 6
- [LR10] J. R. Lee and P. Raghavendra. Coarse differentiation and multi-flows in planar graphs. *Discret. Comput. Geom.*, 43(2):346–362, 2010, doi:[10.1007/S00454-009-9172-4](https://doi.org/10.1007/S00454-009-9172-4). 6
- [LS23] H. Le and S. Solomon. A unified framework for light spanners. In B. Saha and R. A. Servedio, editors, *Proceedings of the 55th Annual ACM Symposium on Theory of Computing, STOC 2023, Orlando, FL, USA, June 20-23, 2023*, pages 295–308. ACM, 2023, doi:[10.1145/3564246.3585185](https://doi.org/10.1145/3564246.3585185). 1
- [Lyn96] N. A. Lynch. *Distributed Algorithms*. Morgan Kaufmann, 1996. 1
- [Mat96] J. Matoušek. On the distortion required for embedding finite metric spaces into normed spaces. *Israel Journal of Mathematics*, 93(1):333–344, 1996, doi:[10.1007/BF02761110](https://doi.org/10.1007/BF02761110). 3, 4
- [MPX13] G. L. Miller, R. Peng, and S. C. Xu. Parallel graph decompositions using random shifts. In G. E. Blelloch and B. Vöcking, editors, *25th ACM Symposium on Parallelism in Algorithms and Architectures, SPAA '13, Montreal, QC, Canada - July 23 - 25, 2013*, pages 196–203. ACM, 2013, doi:[10.1145/2486159.2486180](https://doi.org/10.1145/2486159.2486180). 25

- [Nei16] O. Neiman. Low dimensional embeddings of doubling metrics. *Theory Comput. Syst.*, 58(1):133–152, 2016, doi:[10.1007/S00224-014-9567-3](https://doi.org/10.1007/S00224-014-9567-3). 3
- [NR02] I. Newman and Y. Rabinovich. A lower bound on the distortion of embedding planar metrics into euclidean space. In *SCG '02: Proceedings of the eighteenth annual symposium on Computational geometry*, pages 94–96. ACM Press, 2002, doi:<http://doi.acm.org/10.1145/513400.513412>. 6
- [NR20] I. Newman and Y. Rabinovich. Online embedding of metrics. In S. Albers, editor, *17th Scandinavian Symposium and Workshops on Algorithm Theory, SWAT 2020, June 22-24, 2020, Tórshavn, Faroe Islands*, volume 162 of *LIPICs*, pages 32:1–32:13. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020, doi:[10.4230/LIPICs.SWAT.2020.32](https://doi.org/10.4230/LIPICs.SWAT.2020.32). 6
- [OS81] H. Okamura and P. Seymour. Multicommodity flows in planar graphs. *Journal of Combinatorial Theory, Series B*, 31(1):75 – 81, 1981, doi:[http://dx.doi.org/10.1016/S0095-8956\(81\)80012-3](http://dx.doi.org/10.1016/S0095-8956(81)80012-3). 6
- [Pel93] D. Peleg. Distance-dependent distributed directories. *Inf. Comput.*, 103(2):270–298, 1993, doi:[10.1006/INCO.1993.1020](https://doi.org/10.1006/INCO.1993.1020). 1
- [Pel00] D. Peleg. *Distributed Computing: A Locality-Sensitive Approach*. SIAM, Philadelphia, PA, 2000, doi:[10.1137/1.9780898719772](https://doi.org/10.1137/1.9780898719772). 1
- [PU89] D. Peleg and E. Upfal. A trade-off between space and efficiency for routing tables. *J. ACM*, 36(3):510–530, 1989, doi:[10.1145/65950.65953](https://doi.org/10.1145/65950.65953). 1
- [Rao99] S. Rao. Small distortion and volume preserving embeddings for planar and Euclidean metrics. In *Proceedings of the Fifteenth Annual Symposium on Computational Geometry, Miami Beach, Florida, USA, June 13-16, 1999*, pages 300–306, 1999, doi:[10.1145/304893.304983](https://doi.org/10.1145/304893.304983). 1, 6, 9
- [RS03] N. Robertson and P. D. Seymour. Graph minors. XVI. Excluding a non-planar graph. *Journal of Combinatorial Theory Series B*, 89(1):43–76, 2003, doi:[10.1016/S0095-8956\(03\)00042-X](https://doi.org/10.1016/S0095-8956(03)00042-X). 2
- [SBI11] S. Srivathsan, C. Busch, and S. S. Iyengar. Oblivious buy-at-bulk in planar graphs. In N. Katoh and A. Kumar, editors, *WALCOM: Algorithms and Computation - 5th International Workshop, WALCOM 2011, New Delhi, India, February 18-20, 2011. Proceedings*, volume 6552 of *Lecture Notes in Computer Science*, pages 33–44. Springer, 2011, doi:[10.1007/978-3-642-19094-0_6](https://doi.org/10.1007/978-3-642-19094-0_6). 1, 5, 30
- [Tho04a] M. Thorup. Compact oracles for reachability and approximate distances in planar digraphs. *Journal of the ACM*, 51(6):993–1024, 2004, doi:[10.1145/1039488.1039493](https://doi.org/10.1145/1039488.1039493). 31
- [Tho04b] M. Thorup. Compact oracles for reachability and approximate distances in planar digraphs. *J. ACM*, 51(6):993–1024, November 2004, doi:[10.1145/1039488.1039493](https://doi.org/10.1145/1039488.1039493). 8
- [TZ01] M. Thorup and U. Zwick. Compact routing schemes. In *Proceedings of the Thirteenth Annual ACM Symposium on Parallel Algorithms and Architectures, SPAA 2001, Heraklion, Crete Island, Greece, July 4-6, 2001*, pages 1–10, 2001, doi:[10.1145/378580.378581](https://doi.org/10.1145/378580.378581). 1, 31

A Proof of Lemma 5

We restate Lemma 5 for convenience.

Lemma 5. *Consider a set X with a weight function $\mu : X \rightarrow \mathbb{R}_{>0}$. Denote $\mu(X) = \sum_{x \in X} \mu(x)$. Then there is a prefix free code h , such that for every $x \in X$, $|h(x)| \leq 2 \cdot \left\lceil \log \frac{\mu(X)}{\mu(x)} \right\rceil$.*

Proof. Recall Huffman prefix free code [Huf52]. Huffman constructs a binary tree such that each edge is associated with ± 1 , and the elements X are the leafs. Then $h(x)$ is simply the string obtain by traversing the unique path from the root to x . It is clearly prefix free. The tree constructed as follows: pick the two elements $x, y \in X$ with minimal weight. Let $z \notin X$, and define a new set $X' = X \setminus \{x, y\} \cup \{z\}$ with the same weight function μ on $X \setminus \{x, y\}$, and let $\mu(z) = \mu(x) + \mu(y)$. Note that $\mu(X') = \mu(X)$. Construct a binary tree recursively on X' . Finally add back x, y as the two children of z , assigning $+1$ (resp. -1) to the edge $\{z, x\}$ (resp. $\{z, y\}$). As a result we obtain a binary tree. Huffman [Huf52] proved that this code minimizes the average length: $\sum_{x \in X} \mu(x) \cdot |h(x)|$. In contrast, here we are interested in bounding the length of each code word separately.

To prove the lemma, it is enough to argue for every x , that the root to x unique path has length at most $2 \cdot \left\lceil \log \frac{\mu(X)}{\mu(x)} \right\rceil$. The proof is by induction on $|X|$. The base case, where $|X| = 1$ clearly holds as the path is of length 0. Next we assume $|X| \geq 2$. Consider an arbitrary point $x \in X$. If $\mu(x) \geq \frac{1}{2}\mu(X)$, then x will be “contracted” exactly once, and thus again the lemma holds. Otherwise, if in the entire process x is contracted at most twice, than again we are done (as $2 \cdot \left\lceil \log \frac{\mu(X)}{\mu(x)} \right\rceil \geq 2$). Thus we can assume that x was contracted at least three times.

Consider the the first time when x is combined with some element y into a point z . At this time the set was X_i , and x, y had the minimal weight. In particular for every $q \in X_i \setminus \{x, y\}$, $\mu(x) \leq \mu(q)$. Consider the first time the point z is contracted with a point q into z' . q is combined from elements in $X_i \setminus \{x, y\}$, and hence $\mu(x) \leq \mu(q)$. It follows that $\mu(z') = \mu(y) + \mu(x) + \mu(q) > 2 \cdot \mu(x)$. Note that in the created tree, the path from x to the root goes though z' . By the induction hypothesis, there will be a path from z' to the root of length at most $\left\lceil 2 \cdot \log \frac{\mu(X)}{\mu(z')} \right\rceil$. It follows that there will be a path from x to the root of length at most

$$2 + \left\lceil 2 \cdot \log \frac{\mu(X)}{\mu(z')} \right\rceil = \left\lceil 2 + 2 \cdot \log \frac{\mu(X)}{\mu(z')} \right\rceil = \left\lceil 2 \cdot \left(1 + \log \frac{\mu(X)}{\mu(z')} \right) \right\rceil = \left\lceil 2 \cdot \log \frac{2\mu(X)}{\mu(z')} \right\rceil \leq \left\lceil 2 \cdot \log \frac{\mu(X)}{\mu(x)} \right\rceil,$$

as required. □