

Hypothesis Spaces for Deep Learning

Rui Wang*, Yuesheng Xu[†] and Mingsong Yan[‡]

Abstract

This paper introduces a hypothesis space for deep learning that employs deep neural networks (DNNs). By treating a DNN as a function of two variables, the physical variable and parameter variable, we consider the primitive set of the DNNs for the parameter variable located in a set of the weight matrices and biases determined by a prescribed depth and widths of the DNNs. We then complete the linear span of the primitive DNN set in a weak* topology to construct a Banach space of functions of the physical variable. We prove that the Banach space so constructed is a reproducing kernel Banach space (RKBS) and construct its reproducing kernel. We investigate two learning models, regularized learning and minimum interpolation problem in the resulting RKBS, by establishing representer theorems for solutions of the learning models. The representer theorems unfold that solutions of these learning models can be expressed as linear combination of a finite number of kernel sessions determined by given data and the reproducing kernel.

Key words: Reproducing kernel Banach space, deep learning, deep neural network, representer theorem for deep learning

1 Introduction

Deep learning has been a huge success in applications. Mathematically, its success is due to the use of deep neural networks (DNNs), neural networks of multiple layers, to describe decision functions. Various mathematical aspects of DNNs as an approximation tool were investigated recently in a number of studies [9, 11, 13, 16, 20, 27, 28, 31]. As pointed out in [8], learning processes do not take place in a vacuum. Classical learning methods took place in a reproducing kernel Hilbert space (RKHS) [1], which leads to representation of learning solutions in terms of a combination of a finite number of kernel sessions [19] of a universal kernel [17]. Reproducing kernel Hilbert spaces as appropriate hypothesis spaces for classical learning methods provide a foundation for mathematical analysis of the learning methods. A natural and imperative question is what are appropriate hypothesis spaces for deep learning. Although hypothesis spaces for learning with shallow neural networks (networks of one hidden layer) were investigated recently in a number of studies, (e.g. [2, 6, 18, 21]), appropriate hypothesis spaces for deep learning are still absent. The goal of the present study is to understand this imperative theoretical issue.

The road-map of constructing the hypothesis space for deep learning may be described as follows. We treat a DNN as a function of two variables, one being the physical variable and the

*School of Mathematics, Jilin University, Changchun 130012, P. R. China. E-mail address: rwang11@jlu.edu.cn.

[†]Department of Mathematics and Statistics, Old Dominion University, Norfolk, VA 23529, USA. This author is also a Professor Emeritus of Mathematics, Syracuse University, Syracuse, NY 13244, USA. E-mail address: y1xu@odu.edu. All correspondence should be sent to this author.

[‡]Department of Mathematics and Statistics, Old Dominion University, Norfolk, VA 23529, USA. E-mail address: myan007@odu.edu.

other being the parameter variable. We then consider the set of the DNNs as functions of the physical variable for the parameter variable taking all elements of the set of the weight matrices and biases determined by a prescribed depth and widths of the DNNs. Upon completing the linear span of the DNN set in a weak* topology, we construct a Banach space of functions of the physical variable. We establish that the resulting Banach space is a reproducing kernel Banach space (RKBS), on which point-evaluation functionals are continuous, and construct an asymmetric reproducing kernel, for the space, which is a function of the two variables, the physical variable and the parameter variable. We regard the constructed RKBS as the hypothesis space for deep learning. We remark that when deep neural networks reduce to shallow network (having only one hidden layer), our hypothesis space coincides the space for shallow learning studied in [2].

Upon introducing the hypothesis space for deep learning, we investigate two learning models, the regularized learning and minimum interpolation problem in the resulting RKBS. We establish representer theorems for solutions of the learning models by employing theory of the reproducing kernel Banach space developed in [25, 26, 29] and representer theorems for solutions of learning in a general RKBS established in [4, 23, 24]. Like the representer theorems for the classical learning in RKHSs, the resulting representer theorems for the two deep learning models in the RKBS reveal that although the learning models are of infinite dimension, their solutions lay in finite dimensional manifolds. More specifically, they can be expressed as a linear combination of a finite number of kernel sessions, the reproducing kernel evaluated the parameter variable at points determined by given data. The representer theorems established in this paper is data-dependent. Even when deep neural networks reduce to a shallow network, the corresponding representer theorem is still new to our best acknowledge. The hypothesis space and the representer theorems for the two deep learning models in it provide us prosperous insights of deep learning and supply deep learning a sound mathematical foundation for further investigation.

We organize this paper in six sections. We describe in Section 2 an innate deep learning model with DNNs. Aiming at formulating reproducing kernel Banach spaces as hypothesis spaces for deep learning, in Section 3 we elucidate the notion of vector-valued reproducing kernel Banach spaces. Section 4 is entirely devoted to the development of the hypothesis space for deep learning. We specifically show that the completion of the linear span of the primitive DNN set, pertaining to the innate learning model, in a weak* topology is an RKBS, which constitutes the hypothesis space for deep learning. In Section 5, we study learning models in the RKBS, establishing representer theorems for solutions of two learning models (regularized learning and minimum norm interpolation) in the hypothesis space. We conclude this paper in Section 6 with remarks on advantages of learning in the proposed hypothesis space.

2 Learning with Deep Neural Networks

We describe in this section an innate learning model with DNNs, considered wildly in the machine learning community.

We first recall the notation of DNNs. Let s and t be positive integers. A DNN is a vector-valued function from \mathbb{R}^s to \mathbb{R}^t formed by compositions of functions, each of which is defined by an activation function applied to an affine map. Specifically, for a given univariate function $\sigma : \mathbb{R} \rightarrow \mathbb{R}$, we define a vector-valued function by

$$\sigma(x) := [\sigma(x_1), \dots, \sigma(x_s)]^\top, \quad \text{for } x := [x_1, x_2, \dots, x_s]^\top \in \mathbb{R}^s.$$

For each $n \in \mathbb{N}$, let $\mathbb{N}_n := \{1, 2, \dots, n\}$. For k vector-valued functions f_j , $j \in \mathbb{N}_k$, where the range of f_j is contained in the domain of f_{j+1} , for $j \in \mathbb{N}_{k-1}$, we denote the consecutive composition of

$f_j, j \in \mathbb{N}_k$, by

$$\bigodot_{j=1}^k f_j := f_k \circ f_{k-1} \circ \cdots \circ f_2 \circ f_1,$$

whose domain is that of f_1 . Suppose that $D \in \mathbb{N}$ is prescribed and fixed. Throughout this paper, we always let $m_0 := s$ and $m_D := t$. We specify positive integers $m_j, j \in \mathbb{N}_{D-1}$. For $\mathbf{W}_j \in \mathbb{R}^{m_j \times m_{j-1}}$ and $\mathbf{b}_j \in \mathbb{R}^{m_j}, j \in \mathbb{N}_D$, a DNN is a function defined by

$$\mathcal{N}^D(x) := \left(\mathbf{W}_D \bigodot_{j=1}^{D-1} \sigma(\mathbf{W}_j \cdot + \mathbf{b}_j) + \mathbf{b}_D \right) (x), \quad x \in \mathbb{R}^s. \quad (1)$$

Note that x is the input vector and \mathcal{N}^D has $D - 1$ hidden layers and an output layer, which is the D -th layer.

A DNN may be represented in a recursive manner. From definition (1), a DNN can be defined recursively by

$$\mathcal{N}^1(x) := \mathbf{W}_1 x + \mathbf{b}_1, \quad x \in \mathbb{R}^s$$

and

$$\mathcal{N}^{j+1}(x) := \mathbf{W}_{j+1} \sigma(\mathcal{N}^j(x)) + \mathbf{b}_{j+1}, \quad x \in \mathbb{R}^s, \quad \text{for all } j \in \mathbb{N}_{D-1}.$$

We write \mathcal{N}^D as $\mathcal{N}^D(\cdot, \{\mathbf{W}_j, \mathbf{b}_j\}_{j=1}^D)$ when it is necessary to indicate the dependence of DNNs on the parameters. In this paper, when we write the set $\{\mathbf{W}_j, \mathbf{b}_j\}_{j=1}^D$ associated with the neural network \mathcal{N}^D , we implicitly give it the order inherited from the definition of \mathcal{N}^D . Throughout this paper, we assume that the activation function σ is continuous.

It is advantageous to consider the DNN \mathcal{N}^D defined above as a function of two variables, one being the physical variable $x \in \mathbb{R}^s$ and the other being the parameter variable $\theta := \{\mathbf{W}_j, \mathbf{b}_j\}_{j=1}^D$. Given positive integers $m_j, j \in \mathbb{N}_{D-1}$, we let

$$\mathbb{W} := \{m_j : j \in \mathbb{N}_{D-1}\} \quad (2)$$

denote the width set and define the primitive set of DNNs of D layers by

$$\mathcal{A}_{\mathbb{W}} := \{\mathcal{N}^D(\cdot, \{\mathbf{W}_j, \mathbf{b}_j\}_{j=1}^D) : \mathbf{W}_j \in \mathbb{R}^{m_j \times m_{j-1}}, \mathbf{b}_j \in \mathbb{R}^{m_j}, j \in \mathbb{N}_D\}. \quad (3)$$

Clearly, the set $\mathcal{A}_{\mathbb{W}}$ defined by (3) depends not only on \mathbb{W} but also on D . For the sake of simplicity, we will not indicate the dependence on D in our notation when ambiguity is not caused. For example, we will use \mathcal{N} for \mathcal{N}^D . Moreover, an element of $\mathcal{A}_{\mathbb{W}}$ is a vector-valued function mapping from \mathbb{R}^s to \mathbb{R}^t . We shall understand the set $\mathcal{A}_{\mathbb{W}}$. To this end, we define the parameter space Θ by letting

$$\Theta = \Theta_{\mathbb{W}} := \bigotimes_{j \in \mathbb{N}_D} (\mathbb{R}^{m_j \times m_{j-1}} \otimes \mathbb{R}^{m_j}). \quad (4)$$

Note that Θ is measurable. For $x \in \mathbb{R}^s$ and $\theta \in \Theta$, we define

$$\mathcal{N}(x, \theta) := \mathcal{N}^D(x, \{\mathbf{W}_j, \mathbf{b}_j\}_{j=1}^D). \quad (5)$$

For $x \in \mathbb{R}^s$ and $\theta \in \Theta$, there holds $\mathcal{N}(x, \theta) \in \mathbb{R}^t$. In this notation, set $\mathcal{A}_{\mathbb{W}}$ may be written as

$$\mathcal{A}_{\mathbb{W}} = \{\mathcal{N}(\cdot, \theta) : \theta \in \Theta_{\mathbb{W}}\}.$$

We now describe the innate learning model with DNNs. Suppose that a training dataset

$$\mathbb{D}_m := \{(x_j, y_j) \in \mathbb{R}^s \times \mathbb{R}^t : j \in \mathbb{N}_m\} \quad (6)$$

is given and we would like to train a neural network from the dataset. We denote by $\mathcal{L}(\mathcal{N}, \mathbb{D}_m) : \Theta \rightarrow \mathbb{R}$ a loss function determined by the dataset \mathbb{D}_m . For example, a loss function may take the form

$$\mathcal{L}(\mathcal{N}, \mathbb{D}_m)(\theta) := \sum_{j \in \mathbb{N}_m} \|\mathcal{N}(x_j, \theta) - y_j\|,$$

where $\|\cdot\|$ is a norm of \mathbb{R}^t . Given a loss function, a typical deep learning model is to train the parameters $\theta \in \Theta_{\mathbb{W}}$ from the training dataset \mathbb{D}_m by solving the optimization problem

$$\min\{\mathcal{L}(\mathcal{N}, \mathbb{D}_m)(\theta) : \theta \in \Theta_{\mathbb{W}}\}, \quad (7)$$

where \mathcal{N} has the form in equation (5). Equivalently, optimization problem (7) may be written as

$$\min\{\mathcal{L}(\mathcal{N}, \mathbb{D}_m) : \mathcal{N} \in \mathcal{A}_{\mathbb{W}}\}. \quad (8)$$

Model (8) is an innate learning model considered wildly in the machine learning community. Note that the set $\mathcal{A}_{\mathbb{W}}$ lacks either algebraic or topological structures. It is difficult to conduct mathematical analysis for learning model (8). Even the existence of its solution is not guaranteed.

We introduce a vector space that contains $\mathcal{A}_{\mathbb{W}}$ and consider learning in the vector space. For this purpose, given a set \mathbb{W} of weight widths defined by (2), we define the set

$$\mathcal{B}_{\mathbb{W}} := \left\{ \sum_{l=1}^n c_l \mathcal{N}(\cdot, \theta_l) : c_l \in \mathbb{R}, \theta_l \in \Theta_{\mathbb{W}}, l \in \mathbb{N}_n, n \in \mathbb{N} \right\}. \quad (9)$$

In the next proposition, we present properties of $\mathcal{B}_{\mathbb{W}}$.

Proposition 1. *If \mathbb{W} is the width set defined by (2), then*

- (i) $\mathcal{B}_{\mathbb{W}}$ defined by (9) is the smallest vector space on \mathbb{R} that contains the set $\mathcal{A}_{\mathbb{W}}$,
- (ii) $\mathcal{B}_{\mathbb{W}}$ is of infinite dimension,
- (iii) $\mathcal{B}_{\mathbb{W}} \subset \bigcup_{n \in \mathbb{N}} \mathcal{A}_{n\mathbb{W}}$.

Proof. It is clear that $\mathcal{B}_{\mathbb{W}}$ may be identified as the linear span of $\mathcal{A}_{\mathbb{W}}$, that is,

$$\mathcal{B}_{\mathbb{W}} = \text{span} \{\mathcal{N}(\cdot, \theta) : \theta \in \Theta_{\mathbb{W}}\}.$$

Thus, $\mathcal{B}_{\mathbb{W}}$ is the smallest vector space containing $\mathcal{A}_{\mathbb{W}}$. Item (ii) follows directly from the definition (9) of $\mathcal{B}_{\mathbb{W}}$.

It remains to prove Item (iii). To this end, we let $f \in \mathcal{B}_{\mathbb{W}}$. By the definition (9) of $\mathcal{B}_{\mathbb{W}}$, there exist $n' \in \mathbb{N}$, $c_l \in \mathbb{R}$, $\theta_l \in \Theta_{\mathbb{W}}$, for $l \in \mathbb{N}_{n'}$ such that

$$f(\cdot) = \sum_{l=1}^{n'} c_l \mathcal{N}(\cdot, \theta_l).$$

It suffices to show that $f \in \mathcal{A}_{n'\mathbb{W}}$. Noting that $\theta_l := \{\mathbf{W}_j^l, \mathbf{b}_j^l\}_{j=1}^D$, for $l \in \mathbb{N}_{n'}$, we set

$$\widetilde{\mathbf{W}}_1 := \begin{bmatrix} \mathbf{W}_1^1 \\ \mathbf{W}_1^2 \\ \vdots \\ \mathbf{W}_1^{n'} \end{bmatrix}, \quad \widetilde{\mathbf{W}}_j := \begin{bmatrix} \mathbf{W}_j^1 & \mathbf{0} & \cdots & \mathbf{0} \\ \mathbf{0} & \mathbf{W}_j^2 & \cdots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \cdots & \mathbf{W}_j^{n'} \end{bmatrix}, \quad j \in \mathbb{N}_{D-1} \setminus \{1\}, \quad \widetilde{\mathbf{b}}_j := \begin{bmatrix} \mathbf{b}_j^1 \\ \mathbf{b}_j^2 \\ \vdots \\ \mathbf{b}_j^{n'} \end{bmatrix}, \quad j \in \mathbb{N}_{D-1},$$

and

$$\widetilde{\mathbf{W}}_D := [c_1 \mathbf{W}_D^1 \quad c_2 \mathbf{W}_D^2 \quad \cdots \quad c_{n'} \mathbf{W}_D^{n'}], \quad \widetilde{\mathbf{b}}_D := \sum_{j=1}^{n'} c_j \mathbf{b}_D^j.$$

Clearly, we have that $\widetilde{\mathbf{W}}_1 \in \mathbb{R}^{(n'm_1) \times m_0}$, $\widetilde{\mathbf{b}}_j \in \mathbb{R}^{n'm_j}$, $j \in \mathbb{N}_{D-1}$, $\widetilde{\mathbf{W}}_j \in \mathbb{R}^{(n'm_j) \times (n'm_{j-1})}$, $j \in \mathbb{N}_{D-1} \setminus \{1\}$, $\widetilde{\mathbf{W}}_D \in \mathbb{R}^{m_D \times (n'm_{D-1})}$, and $\widetilde{\mathbf{b}}_D \in \mathbb{R}^{m_D}$. Direct computation confirms that $f(\cdot) = \mathcal{N}(\cdot, \widetilde{\theta})$ with $\widetilde{\theta} := \{\widetilde{\mathbf{W}}_j, \widetilde{\mathbf{b}}_j\}_{j=1}^D$. By definition (3), $f \in \mathcal{A}_{n'\mathbb{W}}$. \square

Proposition 1 reveals that $\mathcal{B}_{\mathbb{W}}$ is the smallest vector space that contains $\mathcal{A}_{\mathbb{W}}$. Hence, it is a reasonable substitute of $\mathcal{A}_{\mathbb{W}}$. Motivated by Proposition 1, we propose the following alternative learning model

$$\inf\{\mathcal{L}(\mathcal{N}, \mathbb{D}_m) : \mathcal{N} \in \mathcal{B}_{\mathbb{W}}\}. \quad (10)$$

For a given width set \mathbb{W} , unlike learning model (8) which searches a minimizer in set $\mathcal{A}_{\mathbb{W}}$, learning model (10) seeks a minimizer in the vector space $\mathcal{B}_{\mathbb{W}}$, which contains $\mathcal{A}_{\mathbb{W}}$ and is contained in $\mathcal{A} := \bigcup_{n \in \mathbb{N}} \mathcal{A}_{n\mathbb{W}}$. According to Proposition 1, learning model (10) is “semi-equivalent” to learning model (8) in the sense that

$$\mathcal{L}(\mathcal{N}_{\mathcal{A}}, \mathbb{D}_m) \leq \mathcal{L}(\mathcal{N}_{\mathcal{B}_{\mathbb{W}}}, \mathbb{D}_m) \leq \mathcal{L}(\mathcal{N}_{\mathcal{A}_{\mathbb{W}}}, \mathbb{D}_m), \quad (11)$$

where $\mathcal{N}_{\mathcal{B}_{\mathbb{W}}}$ is a minimizer of model (10), $\mathcal{N}_{\mathcal{A}_{\mathbb{W}}}$ and $\mathcal{N}_{\mathcal{A}}$ are the minimizers of model (8) and model (8) with the set $\mathcal{A}_{\mathbb{W}}$ replaced by \mathcal{A} , respectively. One might argue that since model (8) is a finite dimension optimization problem while model (10) is an infinite dimensional one, the alternative model (10) may add unnecessary complexity to the original model. Although model (10) is of infinite dimension, the algebraic structure of the vector space $\mathcal{B}_{\mathbb{W}}$ and its topological structure that will be equipped later provide us with great advantages for mathematical analysis of learning on the space. As a matter of fact, the vector-valued RKBS to be obtained by completing the vector space $\mathcal{B}_{\mathbb{W}}$ in a weak* topology will lead to the miraculous representer theorem, of the learned solution, which reduces the infinite dimensional optimization problem to a finite dimension one. This addresses the challenges caused by the infinite dimension of the space $\mathcal{B}_{\mathbb{W}}$.

3 Vector-Valued Reproducing Kernel Banach Space

It was proved in the last section that for a given width set \mathbb{W} , the set $\mathcal{B}_{\mathbb{W}}$ defined by (9) is the smallest vector space that contains the primitive set $\mathcal{A}_{\mathbb{W}}$. One of the aims of this paper is to establish that the vector space $\mathcal{B}_{\mathbb{W}}$ is dense in a weak* topology in a vector-valued RKBS. For this purpose, in this section we describe the notion of vector-valued RKBSs.

A Banach space \mathcal{B} with the norm $\|\cdot\|_{\mathcal{B}}$ is called a space of vector-valued functions on a prescribed set X if \mathcal{B} is composed of vector-valued functions defined on X and for each $f \in \mathcal{B}$, $\|f\|_{\mathcal{B}} = 0$ implies that $f(x) = \mathbf{0}$ for all $x \in X$. For each $x \in X$, we define the point evaluation operator $\delta_x : \mathcal{B} \rightarrow \mathbb{R}^n$ as

$$\delta_x(f) := f(x), \quad f \in \mathcal{B}.$$

We provide the definition of vector-valued RKBSs below.

Definition 2. A Banach space \mathcal{B} of vector-valued functions from X to \mathbb{R}^n is called a vector-valued RKBS if there exists a norm $\|\cdot\|$ of \mathbb{R}^n such that for each $x \in X$, the point evaluation operator δ_x is continuous with respect to the norm $\|\cdot\|$ of \mathbb{R}^n on \mathcal{B} , that is, for each $x \in X$, there exists a constant $C_x > 0$ such that

$$\|\delta_x(f)\| \leq C_x \|f\|_{\mathcal{B}}, \quad \text{for all } f \in \mathcal{B}.$$

Note that since all norms of \mathbb{R}^n are equivalent, if a Banach space \mathcal{B} of vector-valued functions from X to \mathbb{R}^n is a vector-valued RKBS with respect to a norm of \mathbb{R}^n , then it must be a vector-valued RKBS with respect to any other norm of \mathbb{R}^n . Thus, the property of point evaluation operators being continuous on space \mathcal{B} is independent of the choice of the norm of the output space \mathbb{R}^n .

The notion of RKBSs was originally introduced in [29], to guarantee the stability of sampling process and to serve as a hypothesis space for sparse machine learning. Vector-valued RKBSs were studied in [14, 30], in which the definition of the vector-valued RKBS involves an abstract Banach space, with a specific norm, as the output space of functions. In Definition 2, we limit the output space to the Euclidean space \mathbb{R}^n without specifying a norm, due to the special property that norms on \mathbb{R}^n are all equivalent.

We reveal in the next proposition that point evaluation operators are continuous if and only if component-wise point evaluation functionals are continuous. To this end, for a vector-valued function $f : X \rightarrow \mathbb{R}^n$, for each $j \in \mathbb{N}_n$, we denote by $f_j : X \rightarrow \mathbb{R}$ the j -th component of f , that is,

$$f(x) := [f_j(x) : j \in \mathbb{N}_n]^\top, \quad x \in X.$$

Moreover, for each $x \in X$, $k \in \mathbb{N}_n$, we introduce a linear functional $\delta_{x,k} : \mathcal{B} \rightarrow \mathbb{R}$ by

$$\delta_{x,k}(f) := f_k(x), \text{ for } f := [f_k : k \in \mathbb{N}_n]^\top \in \mathcal{B}.$$

Proposition 3. *A Banach space \mathcal{B} of vector-valued functions from X to \mathbb{R}^n is a vector-valued RKBS if and only if for each $x \in X$, $k \in \mathbb{N}_n$, there exists a constant $C_{x,k} > 0$ such that*

$$|\delta_{x,k}(f)| \leq C_{x,k} \|f\|_{\mathcal{B}}, \text{ for all } f \in \mathcal{B}. \quad (12)$$

Proof. Suppose that \mathcal{B} is a vector-valued RKBS. Definition 2 together with the norm equivalence of the Euclidean space \mathbb{R}^n ensures that for each $x \in X$, point evaluation operator δ_x is continuous with respect to the maximum norm $\|\cdot\|_\infty$ of \mathbb{R}^n . That is, for each $x \in X$, there exists a positive constant C_x such that $\|f(x)\|_\infty \leq C_x \|f\|_{\mathcal{B}}$ for all $f \in \mathcal{B}$. Hence, for each $x \in X$, $k \in \mathbb{N}_n$, there holds that $|\delta_{x,k}(f)| = |f_k(x)| \leq \|f(x)\|_\infty \leq C_x \|f\|_{\mathcal{B}}$ for all $f \in \mathcal{B}$. That is, inequality (12) holds true with $C_{x,k} := C_x$.

Conversely, we assume that for each $x \in X$, $k \in \mathbb{N}_n$, there exists $C_{x,k} > 0$ such that (12) holds. Note that for each $x \in X$, there holds $\|\delta_x(f)\|_\infty = \max_{k \in \mathbb{N}_n} |f_k(x)|$ for all $f \in \mathcal{B}$. By setting $C_x := \max_{k \in \mathbb{N}_n} C_{x,k}$, we get from inequality (12) that $\|\delta_x(f)\|_\infty \leq C_x \|f\|_{\mathcal{B}}$ for all $f \in \mathcal{B}$. That is, for each $x \in X$, the point evaluation operator δ_x is continuous with respect to the maximum norm $\|\cdot\|_\infty$ of \mathbb{R}^n , and thus, by Definition 2, \mathcal{B} is a vector-valued RKBS. \square

We next identify a reproducing kernel for a vector-valued RKBS. We need the notion of the δ -dual space of a vector-valued RKBS. For a Banach space B with a norm $\|\cdot\|_B$, we denote by B^* the dual space of B , which is composed of all continuous linear functionals on B endowed with the norm

$$\|\nu\|_{B^*} := \sup_{\|f\|_B \leq 1} |\nu(f)|, \text{ for all } \nu \in B^*.$$

The dual bilinear form $\langle \cdot, \cdot \rangle_B$ on $B^* \times B$ is defined by

$$\langle \nu, f \rangle_B := \nu(f), \text{ for all } \nu \in B^* \text{ and } f \in B.$$

Suppose that \mathcal{B} is a vector-valued RKBS of functions from X to \mathbb{R}^n , with the dual space \mathcal{B}^* . We set

$$\Delta := \text{span}\{\delta_{x,j} : x \in X, j \in \mathbb{N}_n\}. \quad (13)$$

Proposition 3 reveals that for each $x \in X$ and $j \in \mathbb{N}_n$, $\delta_{x,j}$ is a continuous linear functional on \mathcal{B} . As a result, Δ is a subset of \mathcal{B}^* . We denote by \mathcal{B}' the closure of Δ in the norm topology on \mathcal{B}^* and call it the δ -dual space of \mathcal{B} . Clearly, we have that $\mathcal{B}' \subseteq \mathcal{B}^*$ and \mathcal{B}' is the smallest Banach space that contains all point-evaluation functionals on \mathcal{B} . We remark that the δ -dual space of a scalar-valued RKBS was originally introduced in [25].

We identify in the next proposition a reproducing kernel for the vector-valued RKBS \mathcal{B} under the assumption that the δ -dual space \mathcal{B}' is isometrically isomorphic to a Banach space of functions from a set X' to \mathbb{R} .

Proposition 4. *Suppose that \mathcal{B} is a vector-valued RKBS of functions from X to \mathbb{R}^n and its δ -dual space \mathcal{B}' is isometrically isomorphic to a Banach space of functions from X' to \mathbb{R} , then there exists a unique vector-valued function $K : X \times X' \rightarrow \mathbb{R}^n$ such that $K_j(x, \cdot) \in \mathcal{B}'$ for all $x \in X$, $j \in \mathbb{N}_n$, and*

$$f_j(x) = \langle K_j(x, \cdot), f \rangle_{\mathcal{B}}, \text{ for all } f = [f_j : j \in \mathbb{N}_n] \in \mathcal{B} \text{ and all } x \in X, j \in \mathbb{N}_n. \quad (14)$$

Proof. Since \mathcal{B} is a vector-valued RKBS, Proposition 3 ensures that for each $x \in X$, $j \in \mathbb{N}_n$, $\delta_{x,j}$ is a continuous linear functional on \mathcal{B} . By noting that \mathcal{B}' is isometrically isomorphic to a Banach space of functions from X' to \mathbb{R} , there exists $k_{x,j} \in \mathcal{B}'$ such that

$$\delta_{x,j}(f) = \langle k_{x,j}, f \rangle_{\mathcal{B}}, \text{ for all } f \in \mathcal{B}. \quad (15)$$

By defining the j -th component of the vector-valued function $K : X \times X' \rightarrow \mathbb{R}^n$ by

$$K_j(x, x') := k_{x,j}(x'), \quad x \in X, x' \in X', j \in \mathbb{N}_n, \quad (16)$$

we have that $K_j(x, \cdot) \in \mathcal{B}'$ for all $x \in X$ and $j \in \mathbb{N}_n$. Substituting equation (16) into the right-hand side of equation (15) with noting that $\delta_{x,j}(f) = f_j(x)$, we obtain equation (14).

It remains to verify that the function K on $X \times X'$ satisfying the above properties is unique. Assume that there exists another $\tilde{K} : X \times X' \rightarrow \mathbb{R}^n$ such that $\tilde{K}_j(x, \cdot) \in \mathcal{B}'$ for all $x \in X$, $j \in \mathbb{N}_n$, and $f_j(x) = \langle \tilde{K}_j(x, \cdot), f \rangle_{\mathcal{B}}$, for all $f \in \mathcal{B}$, $x \in X$, and $j \in \mathbb{N}_n$. It follows from the above equation and equation (14) that $\langle K_j(x, \cdot) - \tilde{K}_j(x, \cdot), f \rangle_{\mathcal{B}} = 0$, for all $f \in \mathcal{B}$ and all $x \in X$. That is, for all $x \in X$, $j \in \mathbb{N}_n$, $K_j(x, \cdot) - \tilde{K}_j(x, \cdot) = 0$. Noting that \mathcal{B}' is isometrically isomorphic to a Banach space of functions on X' , we conclude that $K_j(x, x') - \tilde{K}_j(x, x') = 0$ for all $x \in X$, $x' \in X'$, and $j \in \mathbb{N}_n$, which is equivalent to $K = \tilde{K}$. \square

We call the vector-valued function $K : X \times X' \rightarrow \mathbb{R}^n$ that satisfies $K_j(x, \cdot) \in \mathcal{B}'$ for all $x \in X$, $j \in \mathbb{N}_n$ and equation (14) the reproducing kernel for the vector-valued RKBS \mathcal{B} . Moreover, equation (14) is called the reproducing property. Clearly, we have that $K(x, \cdot) \in (\mathcal{B}')^n$ for all $x \in X$. The notion of the vector-valued RKBS and its reproducing kernel will serve as a basis for us to understand the hypothesis space for deep learning in the next section.

It is worth of pointing out that although \mathcal{B} is a space of vector-valued functions, the δ -dual space \mathcal{B}' defined here is a space of scalar-valued functions. This is determined by the form of the point evaluation functionals in set Δ defined by (13). The way of defining the δ -dual space of the vector-valued RKBS \mathcal{B} is not unique. One can also define a δ -dual space of the vector-valued RKBS \mathcal{B} as a space of vector-valued functions. In this paper, we adopt the current form of \mathcal{B}' since it is simple and sufficient to serve our purposes. Other forms of the δ -dual space will be investigated in a different occasion.

4 Hypothesis Space

In this section, we return to understanding the vector space $\mathcal{B}_{\mathbb{W}}$ introduced in section 2 from the RKBS viewpoint. Specifically, our goal is to introduce a vector-valued RKBS in which the vector space $\mathcal{B}_{\mathbb{W}}$ is weakly* dense. The resulting vector-valued RKBS will serve as the hypothesis space for deep learning.

We first construct the vector-valued RKBS. Recalling the parameter space Θ defined by equation (4), we use $C_0(\Theta)$ to denote the space of the continuous *scalar-valued* functions vanishing at infinity on Θ . We equip the maximum norm on $C_0(\Theta)$, namely, $\|f\|_{\infty} := \sup_{\theta \in \Theta} |f(\theta)|$, for all $f \in C_0(\Theta)$. For the function $\mathcal{N}(x, \theta)$, $x \in \mathbb{R}^s$, $\theta \in \Theta$, defined by equation (5), we denote by $\mathcal{N}_k(x, \theta)$ the k -th component of $\mathcal{N}(x, \theta)$, for $k \in \mathbb{N}_t$. We require that all components $\mathcal{N}_k(x, \cdot)$ with a weight belong to $C_0(\Theta)$ for all $x \in \mathbb{R}^s$. Specifically, we assume that there exists a continuous weight function $\rho : \Theta \rightarrow \mathbb{R}$ such that the functions

$$\mathcal{N}_k(x, \cdot)\rho(\cdot) \in C_0(\Theta), \quad \text{for all } x \in \mathbb{R}^s, k \in \mathbb{N}_t.$$

An example of such a weight function is given by the rapidly decreasing function

$$\rho(\theta) := \exp(-\|\theta\|_2^2), \quad \theta \in \Theta. \quad (17)$$

We need a measure on the set Θ . A Radon measure [10] on Θ is a Borel measure on Θ that is finite on all compact sets of Θ , outer regular on all Borel sets of Θ , and inner regular on all open sets of Θ . Let $\mathcal{M}(\Theta)$ denote the space of finite Radon measures $\mu : \Theta \rightarrow \mathbb{R}$, equipped with the total variation norm

$$\|\mu\|_{\text{TV}} := \sup \left\{ \sum_{k=1}^{\infty} |\mu(E_k)| : \Theta = \bigcup_{k=1}^{\infty} E_k, E_i \cap E_j = \emptyset \text{ whenever } i \neq j \right\}, \quad (18)$$

where E_k are required to be measurable. Note that $\mathcal{M}(\Theta)$ is the dual space of $C_0(\Theta)$ (see, for example, [7]). Moreover, the dual bilinear form on $\mathcal{M}(\Theta) \times C_0(\Theta)$ is given by

$$\langle \mu, g \rangle_{C_0(\Theta)} := \int_{\Theta} g(\theta) d\mu(\theta), \quad \text{for } \mu \in \mathcal{M}(\Theta), g \in C_0(\Theta). \quad (19)$$

For $\mu \in \mathcal{M}(\Theta)$, we let

$$f_{\mu}^k(\cdot) := \int_{\Theta} \mathcal{N}_k(\cdot, \theta) \rho(\theta) d\mu(\theta), \quad k \in \mathbb{N}_t, \quad (20)$$

and

$$f_{\mu}(\cdot) := \left[f_{\mu}^k(\cdot) : k \in \mathbb{N}_t \right]^{\top}.$$

We introduce the vector space

$$\mathcal{B}_{\mathcal{N}} := \{ f_{\mu} : \mu \in \mathcal{M}(\Theta) \}, \quad (21)$$

with norm

$$\|f_{\mu}\|_{\mathcal{B}_{\mathcal{N}}} := \inf \{ \|\nu\|_{\text{TV}} : f_{\nu} = f_{\mu}, \nu \in \mathcal{M}(\Theta) \}, \quad (22)$$

where f_{μ}^k , $k \in \mathbb{N}_t$, are defined by equation (20) and $\|\cdot\|_{\text{TV}}$ is defined as (18). Note that in definition (22) of the norm $\|f_{\mu}\|_{\mathcal{B}_{\mathcal{N}}}$, the infimum is taken over all the measures $\nu \in \mathcal{M}(\Theta)$ that satisfy t equality constraints

$$\int_{\Theta} \mathcal{N}_k(\cdot, \theta) \rho(\theta) d\mu(\theta) = \int_{\Theta} \mathcal{N}_k(\cdot, \theta) \rho(\theta) d\nu(\theta), \quad k \in \mathbb{N}_t.$$

In particular, in the case $t = 1$, where f_μ reduces to a neural network of a scalar-valued output, the norm $\|f_\mu\|_{\mathcal{B}_\mathcal{N}}$ is taken over the measures $\nu \in \mathcal{M}(\Theta)$ that satisfies only a single equality constraint. The bigger t is, the larger the norm $\|f_\mu\|_{\mathcal{B}_\mathcal{N}}$ will be. We remark that the special case of $\mathcal{B}_\mathcal{N}$ with \mathcal{N} being a scalar-valued neural network of a *single* hidden layer was recently studied in [2].

We next show that the space $\mathcal{B}_\mathcal{N}$ defined by (21) with norm (22) is a Banach space having a pre-dual space. This is done by showing that $\mathcal{B}_\mathcal{N}$ is isometrically isomorphic to a quotient space. We recall the concept of the quotient space. Let B be a Banach space with its dual space B^* and M be a closed subspace of B . For each f in B , the translate $f + M$ which contains f is called the coset of M . The quotient space B/M is defined by $B/M := \{f + M : f \in B\}$, with the quotient norm

$$\|f + M\|_{B/M} := \inf \{\|f + g\| : g \in M\}, \quad f \in B.$$

It is known [15] that the quotient space B/M is a Banach space. We say that a Banach space B has a pre-dual space if there exists a Banach space B_* such that $(B_*)^* = B$ and we call the space B_* a pre-dual space of B . We also need the notion of annihilators. Let M and M' be subsets of B and B^* , respectively. The annihilator of M in B^* is defined by

$$M^\perp := \{\nu \in B^* : \langle \nu, f \rangle_B = 0, \text{ for all } f \in M\}.$$

The annihilator of M' in B is defined by

$${}^\perp M' := \{f \in B : \langle \nu, f \rangle_B = 0, \text{ for all } \nu \in M'\}.$$

We review a result about the dual space of a closed subspace of a Banach space. Specifically, let M be a closed subspace of a Banach space B . For each $\nu \in B^*$, we denote by $\nu|_M$ the restriction of ν to M . It is clear that $\nu|_M \in M^*$ and $\|\nu|_M\|_{M^*} \leq \|\nu\|_{B^*}$. The dual space M^* may be identified as B^*/M^\perp . In fact, by Theorem 10.1 in Chapter III of [7], the map $\tau : B^*/M^\perp \rightarrow M^*$ defined by

$$\tau(\nu + M^\perp) := \nu|_M, \text{ for } \nu \in B^*,$$

is an isometric isomorphism between B^*/M^\perp and M^* .

For the purpose of proving that $\mathcal{B}_\mathcal{N}$ is a Banach space, we identify the quotient space which is isometrically isomorphic to $\mathcal{B}_\mathcal{N}$. To this end, we introduce a closed subspace of $C_0(\Theta)$ as

$$\mathcal{S} := \overline{\text{span}}\{\mathcal{N}_k(x, \cdot)\rho(\cdot) : x \in \mathbb{R}^s, k \in \mathbb{N}_t\}, \quad (23)$$

where the closure is taken in the maximum norm. From definition (23), it is clear that \mathcal{S} is a Banach space of functions defined on the parameter space Θ .

Proposition 5. *Let Θ be the parameter space defined by (4). If for each $x \in \mathbb{R}^s$ and $k \in \mathbb{N}_t$, the function $\mathcal{N}_k(x, \cdot)\rho(\cdot) \in C_0(\Theta)$, then the space $\mathcal{B}_\mathcal{N}$ defined by (21) endowed with the norm (22) is a Banach space with a pre-dual space \mathcal{S} defined by (23).*

Proof. It is clear that $\mathcal{B}_\mathcal{N}$ is a normed space endowed with the norm $\|\cdot\|_{\mathcal{B}_\mathcal{N}}$. It suffices to verify that $\mathcal{B}_\mathcal{N}$ is complete. This is done by showing that $\mathcal{B}_\mathcal{N}$ is isometrically isomorphic to the quotient space $\mathcal{M}(\Theta)/\mathcal{S}^\perp$. We first characterize the annihilator of \mathcal{S} in $\mathcal{M}(\Theta)$. It follows from definition (23) of \mathcal{S} that $\mu \in \mathcal{S}^\perp$ if and only if $\langle \mu, \mathcal{N}_k(x, \cdot)\rho(\cdot) \rangle_{C_0(\Theta)} = 0$ for all $x \in \mathbb{R}^s$ and $k \in \mathbb{N}_t$. The latter is equivalent to $f_\mu^k(x) = 0$ for all $x \in \mathbb{R}^s$ and $k \in \mathbb{N}_t$, due to definition (20) of f_μ^k and definition (19) of the dual bilinear form on $\mathcal{M}(\Theta) \times C_0(\Theta)$. That is, $f_\mu = 0$. Consequently, we conclude that

$$\mathcal{S}^\perp = \{\mu \in \mathcal{M}(\Theta) : f_\mu = 0\}. \quad (24)$$

We next let φ be the map from $\mathcal{B}_{\mathcal{N}}$ to $\mathcal{M}(\Theta)/\mathcal{S}^\perp$ defined for $f_\mu \in \mathcal{B}_{\mathcal{N}}$ by

$$\varphi(f_\mu) := \mu + \mathcal{S}^\perp, \quad \mu \in \mathcal{M}(\Theta), \quad (25)$$

and show that φ is an isometric isomorphism.

We first show that φ is an isometry. For any $f_\mu \in \mathcal{B}_{\mathcal{N}}$ with $\mu \in \mathcal{M}(\Theta)$, we have that $\nu \in \mathcal{M}(\Theta)$ satisfies $f_\nu = f_\mu$ if and only if $f_{\nu-\mu} = 0$. Due to representation (24) of \mathcal{S}^\perp , the latter is equivalent to $\nu - \mu \in \mathcal{S}^\perp$. That is, $\nu = \mu + \mu'$ for some $\mu' \in \mathcal{S}^\perp$. Hence, we get by definition (22) that

$$\|f_\mu\|_{\mathcal{B}_{\mathcal{N}}} = \inf \left\{ \|\mu + \mu'\|_{\text{TV}} : \mu' \in \mathcal{S}^\perp \right\}.$$

This together with the definition of the quotient norm yields that

$$\|f_\mu\|_{\mathcal{B}_{\mathcal{N}}} = \|\mu + \mathcal{S}^\perp\|_{\mathcal{M}(\Theta)/\mathcal{S}^\perp},$$

which with (25) leads to $\|\varphi(f_\mu)\|_{\mathcal{M}(\Theta)/\mathcal{S}^\perp} = \|f_\mu\|_{\mathcal{B}_{\mathcal{N}}}$. In other words, φ is an isometry. Due to its isometry property, φ is injective. Clearly, φ is surjective. Hence, it is bijective. Consequently, φ is an isometric isomorphism from $\mathcal{B}_{\mathcal{N}}$ to the Banach space $\mathcal{M}(\Theta)/\mathcal{S}^\perp$, and thus, $\mathcal{B}_{\mathcal{N}}$ is complete.

We now show that $\mathcal{B}_{\mathcal{N}}$ is isometrically isomorphic to the dual space of \mathcal{S} . Note that \mathcal{S} is a closed subspace of $C_0(\Theta)$ and $(C_0(\Theta))^* = \mathcal{M}(\Theta)$. By Theorem 10.1 in [7] with $B := C_0(\Theta)$ and $M := \mathcal{S}$, we have that the map $\tau : \mathcal{M}(\Theta)/\mathcal{S}^\perp \rightarrow \mathcal{S}^*$ defined by

$$\tau(\mu + \mathcal{S}^\perp) := \mu|_{\mathcal{S}}, \quad \mu \in \mathcal{M}(\Theta),$$

is an isometric isomorphism. As has been shown earlier, the map φ defined by (25) is an isometric isomorphism from $\mathcal{B}_{\mathcal{N}}$ to $\mathcal{M}(\Theta)/\mathcal{S}^\perp$. As a result, $\tau \circ \varphi$ provides an isometric isomorphism from $\mathcal{B}_{\mathcal{N}}$ to \mathcal{S}^* . \square

Proposition 5 and the theorems that follow require that for each $x \in \mathbb{R}^s$ and $k \in \mathbb{N}_t$, the function $\mathcal{N}_k(x, \cdot)\rho(\cdot)$ belongs to $C_0(\Theta)$. This requirement in fact imposes a hypothesis to the activation function σ : (i) σ is continuous and (ii) when the weight function ρ is chosen as (17), we need to select the activation function σ having a growth rate no greater than polynomials. We remark that many commonly used activation functions satisfy this requirement. They include the ReLU function

$$\sigma(x) := \max\{0, x\}, \quad x \in \mathbb{R},$$

and the sigmoid function

$$\sigma(x) := \frac{1}{1 + e^{-x}}, \quad x \in \mathbb{R}.$$

Now that the space $\mathcal{B}_{\mathcal{N}}$ with the norm $\|\cdot\|_{\mathcal{B}_{\mathcal{N}}}$, guaranteed by Proposition 5, is a Banach space, we denote by $\mathcal{B}_{\mathcal{N}}^*$ the dual space of $\mathcal{B}_{\mathcal{N}}$ endowed with the norm

$$\|\ell\|_{\mathcal{B}_{\mathcal{N}}^*} = \sup\{|\langle \ell, f_\mu \rangle_{\mathcal{B}_{\mathcal{N}}}| : \|f_\mu\|_{\mathcal{B}_{\mathcal{N}}} = 1, f_\mu \in \mathcal{B}_{\mathcal{N}}\}, \quad \text{for } \ell \in \mathcal{B}_{\mathcal{N}}^*. \quad (26)$$

The dual space $\mathcal{B}_{\mathcal{N}}^*$ is again a Banach space. Moreover, it follows from Proposition 5 that the space \mathcal{S} is a pre-dual space of $\mathcal{B}_{\mathcal{N}}$, that is, $(\mathcal{B}_{\mathcal{N}})^* = \mathcal{S}$. We remark that the dual bilinear form on $\mathcal{B}_{\mathcal{N}} \times \mathcal{S}$ is given by

$$\langle f_\mu, g \rangle_{\mathcal{S}} = \langle \mu, g \rangle_{C_0(\Theta)}, \quad \text{for } f_\mu \in \mathcal{B}_{\mathcal{N}}, g \in \mathcal{S}. \quad (27)$$

According to Proposition 5, the space \mathcal{S} is the pre-dual space of $\mathcal{B}_{\mathcal{N}}$, that is, $\mathcal{S}^* = \mathcal{B}_{\mathcal{N}}$. Thus, we obtain that $\mathcal{S}^{**} = \mathcal{B}_{\mathcal{N}}^*$. It is well-known (for example, see [7]) that $\mathcal{S} \subseteq \mathcal{S}^{**}$ in the sense of isometric embedding. Hence, $\mathcal{S} \subseteq \mathcal{B}_{\mathcal{N}}^*$ and there holds

$$\langle g, f_\mu \rangle_{\mathcal{B}_{\mathcal{N}}^*} = \langle f_\mu, g \rangle_{\mathcal{S}}, \quad \text{for all } f_\mu \in \mathcal{B}_{\mathcal{N}} \text{ and all } g \in \mathcal{S}. \quad (28)$$

We now turn to establishing that $\mathcal{B}_{\mathcal{N}}$ is a vector-valued RKBS on \mathbb{R}^s .

Theorem 6. Let Θ be the parameter space defined by (4). If for each $x \in \mathbb{R}^s$ and $k \in \mathbb{N}_t$, the function $\mathcal{N}_k(x, \cdot)\rho(\cdot)$ belongs to $C_0(\Theta)$, then the Banach space $\mathcal{B}_{\mathcal{N}}$ defined by (21) endowed with the norm (22) is a vector-valued RKBS on \mathbb{R}^s .

Proof. According to Proposition 3 with $X := \mathbb{R}^s$, it suffices to prove that for each $x \in \mathbb{R}^s$, $k \in \mathbb{N}_t$, there exists a positive constant $C_{x,k}$ such that

$$|f_{\mu}^k(x)| \leq C_{x,k} \|f_{\mu}\|_{\mathcal{B}_{\mathcal{N}}}, \text{ for all } f_{\mu} \in \mathcal{B}_{\mathcal{N}}. \quad (29)$$

To this end, for any $f_{\mu} \in \mathcal{B}_{\mathcal{N}}$, we obtain from definition (20) of f_{μ}^k that

$$|f_{\mu}^k(x)| \leq \|\mathcal{N}_k(x, \cdot)\rho(\cdot)\|_{\infty} \|\nu\|_{\text{TV}}, \quad (30)$$

for any $\nu \in \mathcal{M}(\Theta)$ satisfying $f_{\nu} = f_{\mu}$. By taking infimum of both sides of inequality (30) over $\nu \in \mathcal{M}(\Theta)$ satisfying $f_{\nu} = f_{\mu}$ and employing definition (22), we obtain that

$$|f_{\mu}^k(x)| \leq \|\mathcal{N}_k(x, \cdot)\rho(\cdot)\|_{\infty} \|f_{\mu}\|_{\mathcal{B}_{\mathcal{N}}}.$$

Letting $C_{x,k} := \|\mathcal{N}_k(x, \cdot)\rho(\cdot)\|_{\infty}$, we get inequality (29). \square

Next, we identify the reproducing kernel of the vector-valued RKBS $\mathcal{B}_{\mathcal{N}}$. According to Proposition 4, the existence of the reproducing kernel requires to characterize the δ -dual space of $\mathcal{B}_{\mathcal{N}}$. We note that the δ -dual space $\mathcal{B}'_{\mathcal{N}}$ is the closure of

$$\Delta := \text{span}\{\delta_{x,k} : x \in \mathbb{R}^s, k \in \mathbb{N}_t\},$$

in the norm topology (26) of $\mathcal{B}_{\mathcal{N}}^*$. We will show that Δ is isometrically isomorphic to

$$\mathbb{S} := \text{span}\{\mathcal{N}_k(x, \cdot)\rho(\cdot) : x \in \mathbb{R}^s, k \in \mathbb{N}_t\},$$

a subspace of \mathcal{S} . To this end, we introduce a mapping $\Psi : \Delta \rightarrow \mathbb{S}$ by

$$\Psi \left(\sum_{j \in \mathbb{N}_m} \alpha_j \delta_{x_j, k_j} \right) := \sum_{j \in \mathbb{N}_m} \alpha_j \mathcal{N}_{k_j}(x_j, \cdot)\rho(\cdot), \quad (31)$$

for all $m \in \mathbb{N}$, $\alpha_j \in \mathbb{R}$, $x_j \in \mathbb{R}^s$, $k_j \in \mathbb{N}_t$, and $j \in \mathbb{N}_m$.

Lemma 7. The map Ψ defined by (31) is an isometric isomorphism between Δ and \mathbb{S} .

Proof. We first prove that Ψ is an isometry, that is, $\|\ell\|_{\mathcal{B}_{\mathcal{N}}^*} = \|\Psi(\ell)\|_{\infty}$, for all $\ell \in \Delta$. Let ℓ be an arbitrary element of Δ . Then there exist $m \in \mathbb{N}$, $\alpha_j \in \mathbb{R}$, $x_j \in \mathbb{R}^s$, $k_j \in \mathbb{N}_t$, and $j \in \mathbb{N}_m$ such that $\ell = \sum_{j \in \mathbb{N}_m} \alpha_j \delta_{x_j, k_j}$. By definition (26) and the definition of the functionals δ_{x_j, k_j} , $j \in \mathbb{N}_m$, we have that

$$\|\ell\|_{\mathcal{B}_{\mathcal{N}}^*} = \sup \left\{ \left| \sum_{j \in \mathbb{N}_m} \alpha_j f_{\mu}^{k_j}(x_j) \right| : \|f_{\mu}\|_{\mathcal{B}_{\mathcal{N}}} = 1, f_{\mu} \in \mathcal{B}_{\mathcal{N}} \right\}. \quad (32)$$

We next compute $\|\Psi(\ell)\|_{\infty}$. By noting that $\Psi(\ell) \in \mathbb{S}$ and $\mathbb{S}^* = \mathcal{B}_{\mathcal{N}}$, we have that

$$\|\Psi(\ell)\|_{\infty} = \sup \left\{ |\langle f_{\mu}, \Psi(\ell) \rangle_{\mathcal{S}}| : \|f_{\mu}\|_{\mathcal{B}_{\mathcal{N}}} = 1, f_{\mu} \in \mathcal{B}_{\mathcal{N}} \right\}.$$

Substituting equation (27) with $g := \Psi(\ell)$ into the right hand side of the above equation, we get that

$$\|\Psi(\ell)\|_\infty = \sup \left\{ |\langle \mu, \Psi(\ell) \rangle_{C_0(\Theta)}| : \|f_\mu\|_{\mathcal{B}_\mathcal{N}} = 1, f_\mu \in \mathcal{B}_\mathcal{N} \right\}. \quad (33)$$

According to definition (31) of Ψ , there holds for any $f_\mu \in \mathcal{B}_\mathcal{N}$ that

$$\langle \mu, \Psi(\ell) \rangle_{C_0(\Theta)} = \sum_{j \in \mathbb{N}_m} \alpha_j \langle \mu, \mathcal{N}_{k_j}(x_j, \cdot) \rho(\cdot) \rangle_{C_0(\Theta)}.$$

This together with definition (20) yields that $\langle \mu, \Psi(\ell) \rangle_{C_0(\Theta)} = \sum_{j \in \mathbb{N}_m} \alpha_j f_\mu^{k_j}(x_j)$. Involving this equation in the right-hand side of (33) leads to

$$\|\Psi(\ell)\|_\infty = \sup \left\{ \left| \sum_{j \in \mathbb{N}_m} \alpha_j f_\mu^{k_j}(x_j) \right| : \|f_\mu\|_{\mathcal{B}_\mathcal{N}} = 1, f_\mu \in \mathcal{B}_\mathcal{N} \right\}. \quad (34)$$

Comparing (32) and (34), we obtain that $\|\ell\|_{\mathcal{B}_\mathcal{N}^*} = \|\Psi(\ell)\|_\infty$ and hence, Ψ is an isometry between Δ and \mathbb{S} . The isometry of Ψ further implies its injectivity. Moreover, Ψ is linear and surjective. Thus, Ψ is bijective. Therefore, Ψ is an isometric isomorphism between Δ and \mathbb{S} . \square

The isometrically isomorphic relation between Δ and \mathbb{S} is preserved after completing them. We state this result in the following lemma without proof.

Lemma 8. *Suppose that A and B are Banach spaces with norms $\|\cdot\|_A$ and $\|\cdot\|_B$, respectively. Let A_0 and B_0 be dense subsets of A and B , respectively. If A_0 is isometrically isomorphic to B_0 , then A is isometrically isomorphic to B .*

Lemma 8 may be obtained by applying Theorem 1.6-2 in [12]. With the help of Lemmas 7 and 8, we identify in the following theorem the reproducing kernel for the RKBS $\mathcal{B}_\mathcal{N}$.

Theorem 9. *Let Θ be the parameter space defined by (4). Suppose that for each $x \in \mathbb{R}^s$ and $k \in \mathbb{N}_t$, the function $\mathcal{N}_k(x, \cdot) \rho(\cdot)$ belongs to $C_0(\Theta)$. If the vector-valued RKBS $\mathcal{B}_\mathcal{N}$ is defined by (21) with the norm (22), then the vector-valued function*

$$\mathcal{K}(x, \theta) := \mathcal{N}(x, \theta) \rho(\theta), \quad \text{for } (x, \theta) \in \mathbb{R}^s \times \Theta, \quad (35)$$

is the reproducing kernel for space $\mathcal{B}_\mathcal{N}$.

Proof. We employ Proposition 4 with $X := \mathbb{R}^s$ and $X' := \Theta$ to establish that the function \mathcal{K} defined by (35) is the reproducing kernel of space $\mathcal{B}_\mathcal{N}$. According to Lemma 7, Δ is isometrically isomorphic to \mathbb{S} . Since $\mathcal{B}'_\mathcal{N}$ and \mathcal{S} are the completion of Δ and \mathbb{S} , respectively, by Lemma 8, we conclude that the δ -dual space $\mathcal{B}'_\mathcal{N}$ of $\mathcal{B}_\mathcal{N}$ is isometrically isomorphic to \mathcal{S} , which is a Banach space of functions from Θ to \mathbb{R} . Hence, Proposition 4 ensures that there exists a unique reproducing kernel for $\mathcal{B}_\mathcal{N}$.

We next verify that the vector-valued function \mathcal{K} defined by (35) is the reproducing kernel for $\mathcal{B}_\mathcal{N}$. By noting that the δ -dual space $\mathcal{B}'_\mathcal{N}$ is isometrically isomorphic to \mathcal{S} , we have for each $x \in \mathbb{R}^s$ and each $k \in \mathbb{N}_t$ that $\mathcal{K}_k(x, \cdot) := \mathcal{N}_k(x, \cdot) \rho(\cdot) \in \mathcal{B}'_\mathcal{N}$. The space \mathcal{S} , guaranteed by Proposition 5, is a pre-dual space of $\mathcal{B}_\mathcal{N}$. Hence, by equation (28) with $g := \mathcal{K}_k(x, \cdot)$, we obtain for each $x \in \mathbb{R}^s$, $k \in \mathbb{N}_t$ that

$$\langle \mathcal{K}_k(x, \cdot), f_\mu \rangle_{\mathcal{B}_\mathcal{N}} = \langle f_\mu, \mathcal{K}_k(x, \cdot) \rangle_{\mathcal{S}}, \quad \text{for all } f_\mu \in \mathcal{B}_\mathcal{N}.$$

Substituting equation (27) with $g := \mathcal{K}_k(x, \cdot)$ into the right-hand side of the above equation leads to

$$\langle \mathcal{K}_k(x, \cdot), f_\mu \rangle_{\mathcal{B}_\mathcal{N}} = \langle \mu, \mathcal{K}_k(x, \cdot) \rangle_{C_0(\Theta)}, \text{ for all } f_\mu \in \mathcal{B}_\mathcal{N}.$$

This together with definitions (19), (35) and (20) implies the reproducing property

$$\langle \mathcal{K}_k(x, \cdot), f_\mu \rangle_{\mathcal{B}_\mathcal{N}} = f_\mu^k(x), \text{ for all } f_\mu \in \mathcal{B}_\mathcal{N}. \quad (36)$$

Consequently, \mathcal{K} is the reproducing kernel of $\mathcal{B}_\mathcal{N}$. \square

The reproducing kernel defined by (35) in Theorem 9 is an asymmetric kernel, unlike a reproducing kernel in a reproducing kernel Hilbert space, which is always symmetric. It is the asymmetry of the “kernel” that allows us to encode one variable of the kernel function as the physical variable and one as the parameter variable. Theorem 9 restricted to the shallow network is still new to our best knowledge. We will show in the next section a solution of a deep learning model may be expressed as a combination of a finite number of kernel sessions, a kernel with the parameter variable evaluated at a point of the parameter space determined by given data.

We are ready to prove that the vector space $\mathcal{B}_\mathbb{W}$, defined by equation (9), is weakly* dense in the vector-valued RKBS $\mathcal{B}_\mathcal{N}$. For this purpose, we recall the concept of the weak* topology. Let B be a Banach space. The weak* topology of the dual space B^* is the smallest topology for B^* such that, for each $f \in B$, the linear functional $\nu \rightarrow \langle \nu, f \rangle_B$ on B^* is continuous with respect to the topology. For a subset M' of B^* , we denote by $\overline{M'}^{w^*}$ the closure of M' in the weak* topology of B^* . We remark that the fact that the Banach space $\mathcal{B}_\mathcal{N}$ has a pre-dual space \mathcal{S} makes it valid for $\mathcal{B}_\mathcal{N}$ to be equipped with the weak* topology, the topology of \mathcal{S}^* .

Theorem 10. *Let Θ be the parameter space defined by (4) and \mathbb{W} the width set defined by (2). If for each $x \in \mathbb{R}^s$ and $k \in \mathbb{N}_t$, the function $\mathcal{N}_k(x, \cdot)\rho(\cdot)$ belongs to $C_0(\Theta)$, then $\mathcal{B}_\mathbb{W}$ is a subspace of $\mathcal{B}_\mathcal{N}$ and*

$$\overline{\mathcal{B}_\mathbb{W}}^{w^*} = \mathcal{B}_\mathcal{N}. \quad (37)$$

Proof. It has been shown in Proposition 1 that $\mathcal{B}_\mathbb{W}$ is a vector space. We now show that $\mathcal{B}_\mathbb{W}$ is a subspace of $\mathcal{B}_\mathcal{N}$. For any $f \in \mathcal{B}_\mathbb{W}$, there exist $n \in \mathbb{N}$, $c_l \in \mathbb{R}$, $\theta_l \in \Theta$, $l \in \mathbb{N}_n$ such that $f = \sum_{l=1}^n c_l \mathcal{N}(\cdot, \theta_l)\rho(\theta_l)$. By choosing $\mu := \sum_{l=1}^n c_l \delta_{\theta_l}$, we have that $\mu \in \mathcal{M}(\Theta)$. We then obtain from definition (20) that

$$f_\mu^k(x) = \sum_{l=1}^n c_l \mathcal{N}_k(x, \theta_l)\rho(\theta_l), \text{ for all } x \in \mathbb{R}^s, k \in \mathbb{N}_t.$$

This together with the representation of f yields that $f = f_\mu$ and thus, $f \in \mathcal{B}_\mathcal{N}$. Consequently, we have that $\mathcal{B}_\mathbb{W} \subseteq \mathcal{B}_\mathcal{N}$.

It remains to prove equation (37). Proposition 5 ensures that $\mathcal{S}^* = \mathcal{B}_\mathcal{N}$, in the sense of being isometrically isomorphic. Hence, $\mathcal{B}_\mathbb{W}$ is a subspace of the dual space of \mathcal{S} . It follows from Proposition 2.6.6 of [15] that $({}^\perp \mathcal{B}_\mathbb{W})^\perp = \overline{\mathcal{B}_\mathbb{W}}^{w^*}$. It suffices to verify that $({}^\perp \mathcal{B}_\mathbb{W})^\perp = \mathcal{B}_\mathcal{N}$. Due to definition (9) of $\mathcal{B}_\mathbb{W}$, $g \in {}^\perp \mathcal{B}_\mathbb{W}$ if and only if

$$\langle \mathcal{N}(\cdot, \theta)\rho(\theta), g \rangle_{\mathcal{S}} = 0, \text{ for all } \theta \in \Theta. \quad (38)$$

By equation (27) with $f_\mu := \mathcal{N}(\cdot, \theta)\rho(\theta)$ with $\mu = \delta_\theta$, equation (38) is equivalent to $\langle \delta_\theta, g \rangle_{C_0(\Theta)} = 0$, for all $\theta \in \Theta$, which leads to $g(\theta) = 0$ for all $\theta \in \Theta$. That is, $g = 0$. Therefore, $({}^\perp \mathcal{B}_\mathbb{W})^\perp = \{0\}$. This together with the definition of annihilators leads to $({}^\perp \mathcal{B}_\mathbb{W})^\perp = \mathcal{B}_\mathcal{N}$, which completes the proof of this theorem. \square

To close this section, we summarize the properties of the space $\mathcal{B}_{\mathcal{N}}$ established in Theorems 6, 9 and 10 as follows:

- (i) The space $\mathcal{B}_{\mathcal{N}}$ is a vector-valued RKBS.
- (ii) The vector-valued function \mathcal{K} defined by (35) is the reproducing kernel for space $\mathcal{B}_{\mathcal{N}}$.
- (iii) The space $\mathcal{B}_{\mathcal{N}}$ is the weak* completion of the vector space $\mathcal{B}_{\mathbb{W}}$.

These favorable properties of the space $\mathcal{B}_{\mathcal{N}}$ motivate us to take it as the hypothesis space for deep learning. Thus, we consider the following learning model

$$\inf\{\mathcal{L}(f_{\mu}, \mathbb{D}_m) : f_{\mu} \in \mathcal{B}_{\mathcal{N}}\}. \quad (39)$$

If we denote by $\mathcal{N}_{\mathcal{B}_{\mathcal{N}}}$ the neural network learned from the model (39), then, according to (11), we have that

$$\mathcal{L}(\mathcal{N}_{\mathcal{B}_{\mathcal{N}}}, \mathbb{D}_m) \leq \mathcal{L}(\mathcal{N}_{\mathcal{B}_{\mathbb{W}}}, \mathbb{D}_m) \leq \mathcal{L}(\mathcal{N}_{\mathcal{A}_{\mathbb{W}}}, \mathbb{D}_m).$$

Even though learning model (39) is like model (10), which is of infinite dimension (unlike model (8), which is of finite dimension), we will show in the next section that a solution of learning model (39) lays in a finite dimensional manifold determined by the kernel \mathcal{K} and a given data set.

5 Representer Theorems for Learning Solutions

In this section, we consider learning a target function in $\mathcal{B}_{\mathcal{N}}$ from the sampled dataset \mathbb{D}_m defined by (6). Learning such a function is an ill-posed problem, whose solutions often suffer from overfitting. For this reason, instead of solving the learning model (39) directly, we consider a related regularization problem and MNI problem in the RKBS $\mathcal{B}_{\mathcal{N}}$. The goal of this section is to establish representer theorems for solutions of these two learning models.

We start with describing the regularized learning problem in the vector-valued RKBS $\mathcal{B}_{\mathcal{N}}$. For the dataset \mathbb{D}_m defined by (6), we define the set $\mathcal{X} := \{x_j : j \in \mathbb{N}_m\}$ and the matrix $\mathbf{Y} := [y_j^k : k \in \mathbb{N}_t, j \in \mathbb{N}_m] \in \mathbb{R}^{t \times m}$, where for each $j \in \mathbb{N}_m$, y_j^k , $k \in \mathbb{N}_t$, are the components of vector y_j . We introduce an operator $\mathbf{I}_{\mathcal{X}} : \mathcal{B}_{\mathcal{N}} \rightarrow \mathbb{R}^{t \times m}$ by

$$\mathbf{I}_{\mathcal{X}}(f_{\mu}) := \left[f_{\mu}^k(x_j) : k \in \mathbb{N}_t, j \in \mathbb{N}_m \right], \text{ for } f_{\mu} \in \mathcal{B}_{\mathcal{N}}. \quad (40)$$

We choose a loss function $\mathcal{Q} : \mathbb{R}^{t \times m} \rightarrow \mathbb{R}_+ := [0, +\infty)$ and define

$$\mathcal{L}(f_{\mu}, \mathbb{D}_m) := \mathcal{Q}(\mathbf{I}_{\mathcal{X}}(f_{\mu}) - \mathbf{Y}), \text{ for } f_{\mu} \in \mathcal{B}_{\mathcal{N}}. \quad (41)$$

Examples of loss functions $\mathcal{Q}(\mathbf{M})$ may be chosen as a norm of the matrix \mathbf{M} . The proposed regularization problem is formed by adding a regularization term $\lambda \|f_{\mu}\|_{\mathcal{B}_{\mathcal{N}}}$ to the data fidelity term $\mathcal{Q}(\mathbf{I}_{\mathcal{X}}(f_{\mu}) - \mathbf{Y})$. That is,

$$\inf \{ \mathcal{Q}(\mathbf{I}_{\mathcal{X}}(f_{\mu}) - \mathbf{Y}) + \lambda \|f_{\mu}\|_{\mathcal{B}_{\mathcal{N}}} : f_{\mu} \in \mathcal{B}_{\mathcal{N}} \}, \quad (42)$$

where λ is a positive regularization parameter. The learning model (42) allows us to learn a function f_{μ} in the space $\mathcal{B}_{\mathcal{N}}$ by solving the optimization problem (42).

We first comment on the existence of a solution to the regularization problem (42). The next proposition follows directly from Proposition 40 of [23].

Proposition 11. *Suppose that m distinct points $x_j \in \mathbb{R}^s$, $j \in \mathbb{N}_m$, and $\mathbf{Y} \in \mathbb{R}^{t \times m}$ are given. If $\lambda > 0$ and the loss function \mathcal{Q} is lower semi-continuous on $\mathbb{R}^{t \times m}$, then the regularization problem (42) has at least one solution.*

Proof. We have shown in Proposition 5 that the Banach space $\mathcal{B}_{\mathcal{N}}$ has the pre-dual space \mathcal{S} . We specify the function $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ appearing in Proposition 40 of [23] as the identity function $\varphi(x) = x$, $x \in \mathbb{R}_+$, which is lower semi-continuous, increasing and coercive. Since the loss function \mathcal{Q} is lower semi-continuous on $\mathbb{R}^{t \times m}$, the assumptions in Proposition 40 of [23] are all satisfied. Thus, we conclude from Proposition 40 of [23] that the regularization problem (42) has at least one solution. \square

It is known that regularization problems are closely related to MNI problems (see, for example, [23]). The MNI problem aims at finding a vector-valued function f_μ in $\mathcal{B}_{\mathcal{N}}$, having the smallest norm and satisfying the interpolation condition $f_\mu(x_j) = y_j$, $j \in \mathbb{N}_m$. In other words, the MNI problem has the form

$$\inf \{ \|f_\mu\|_{\mathcal{B}_{\mathcal{N}}} : f_\mu(x_j) = y_j, f_\mu \in \mathcal{B}_{\mathcal{N}}, j \in \mathbb{N}_m \}. \quad (43)$$

Associated with the set \mathcal{X} and matrix \mathbf{Y} , we introduce a subset $\mathcal{M}_{\mathcal{X}, \mathbf{Y}}$ of $\mathcal{B}_{\mathcal{N}}$ as

$$\mathcal{M}_{\mathcal{X}, \mathbf{Y}} := \{ f_\mu \in \mathcal{B}_{\mathcal{N}} : \mathbf{I}_{\mathcal{X}}(f_\mu) = \mathbf{Y} \}. \quad (44)$$

We then reformulate the MNI problem (43) in an equivalent form as

$$\inf \left\{ \|f_\mu\|_{\mathcal{B}_{\mathcal{N}}} : f_\mu \in \mathcal{M}_{\mathcal{X}, \mathbf{Y}} \right\}. \quad (45)$$

The MNI problem (45) has a solution if and only if the set $\mathcal{M}_{\mathcal{X}, \mathbf{Y}}$ is nonempty. The non-emptiness of the set $\mathcal{M}_{\mathcal{X}, \mathbf{Y}}$ pertains to the existence of interpolation of a function in $\mathcal{B}_{\mathcal{N}}$ to any given data \mathbb{D}_m . For the sake of keeping focus on the main issues of this paper, this issue will be postponed to a different occasion. In this paper, we will always assume that the set $\mathcal{M}_{\mathcal{X}, \mathbf{Y}}$ is nonempty.

Recall that the vector-valued function \mathcal{K} defined by (35) is the reproducing kernel for $\mathcal{B}_{\mathcal{N}}$. By using the reproducing property (36), we represent the operator $\mathbf{I}_{\mathcal{X}}$ defined by (40) as

$$\mathbf{I}_{\mathcal{X}}(f_\mu) = [\langle \mathcal{K}_k(x_j, \cdot), f_\mu \rangle_{\mathcal{B}_{\mathcal{N}}} : k \in \mathbb{N}_t, j \in \mathbb{N}_m], \text{ for } f_\mu \in \mathcal{B}_{\mathcal{N}}.$$

Clearly, the MNI problem (45) includes tm interpolation conditions which are produced by the linear functionals in the set

$$\mathbb{K}_{\mathcal{X}} := \{ \mathcal{K}_k(x_j, \cdot) : j \in \mathbb{N}_m, k \in \mathbb{N}_t \}.$$

It follows from Proposition 1 of [23] that the existence of a solution of the MNI problem (45) is guaranteed if the functionals in $\mathbb{K}_{\mathcal{X}}$ are linearly independent elements in \mathcal{S} . In fact, the linear independence of the functionals in $\mathbb{K}_{\mathcal{X}}$ is a sufficient condition that ensures the non-emptiness of the subset $\mathcal{M}_{\mathcal{X}, \mathbf{Y}}$ defined by (44) for any given $\mathbf{Y} \in \mathbb{R}^{t \times m}$. If the subset $\mathcal{M}_{\mathcal{X}, \mathbf{Y}}$ is nonempty but the functionals in $\mathbb{K}_{\mathcal{X}}$ are linearly dependent, one may replace $\mathbb{K}_{\mathcal{X}}$ by its maximal linearly independent subset. Hence, without loss of generality, we will assume that the functionals in $\mathbb{K}_{\mathcal{X}}$ are linearly independent throughout the rest of this paper.

A solution of the regularization problem (42) may be identified as a solution of an MNI problem in the form of (45) with different data. In fact, according to [23], every solution $\hat{f}_\mu \in \mathcal{B}_{\mathcal{N}}$ of the regularization problem (42) is also a solution of the MNI problem (45) with $\mathbf{Y} := \mathbf{I}_{\mathcal{X}}(\hat{f}_\mu)$. In addition, if $\hat{f}_\mu \in \mathcal{B}_{\mathcal{N}}$ is a solution of the MNI problem (45), then \hat{f}_μ is a solution of the regularization problem

$$\inf \{ \mathcal{Q}(\mathbf{I}_{\mathcal{X}}(f_\mu) - \mathbf{Y}) + \lambda \|f_\mu\|_{\mathcal{B}_{\mathcal{N}}} : f_\mu \in \mathcal{M}_{\mathcal{X}, \mathbf{Y}} \},$$

and there holds the relation

$$C^* \leq \mathcal{Q}(\mathbf{I}_{\mathcal{X}}(\hat{f}_\mu) - \mathbf{Y}) + \lambda \|\hat{f}_\mu\|_{\mathcal{B}_{\mathcal{N}}},$$

where C^* is the infimum of the regularization problem (42), since $\mathcal{M}_{\mathcal{X}, \mathbf{Y}}$ is a subset of $\mathcal{B}_{\mathcal{N}}$.

We now establish a representer theorem for a solution of the MNI problem (45). This can be accomplished by applying the explicit and data-dependent representer theorem for the MNI problem in a general Banach space setting, established in our recent paper [24] to the current setting. To this end, we review some necessary notions of Convex Analysis and recall a result of [24]. Let \mathbb{X} be a Hausdorff locally convex topological vector space. A subset A of \mathbb{X} is called a convex set if $tx + (1-t)y \in A$ for all $x, y \in A$ and all $t \in [0, 1]$. The convex hull of a subset A of \mathbb{X} , denoted by $\text{co}(A)$, is the smallest convex set that contains A . The closed convex hull of A , denoted by $\overline{\text{co}}(A)$, is the smallest closed convex set that contains A , where the closure is taken under the topology of \mathbb{X} . Suppose that A is a nonempty closed convex subset of \mathbb{X} . An element $z \in A$ is said to be an extreme point of A if $x, y \in A$ and $tx + (1-t)y = z$ for some $t \in (0, 1)$ implies that $x = y = z$. By $\text{ext}(A)$ we denote the set of extreme points of A . The celebrated Krein-Milman theorem [15] states that if A is a nonempty compact convex subset of \mathbb{X} , then A is the closed convex hull of its set of extreme points, that is, $A = \overline{\text{co}}(\text{ext}(A))$. Let B be a Banach space endowed with norm $\|\cdot\|_B$. Clearly, the norm $\|\cdot\|_B$ is a convex function on B . The subdifferential of the norm function $\|\cdot\|_B$ at each $f \in B \setminus \{0\}$ is defined by

$$\partial\|\cdot\|_B(f) := \{\nu \in B^* : \|\nu\|_{B^*} = 1, \langle \nu, f \rangle_B = \|f\|_B\}.$$

Notice that the dual space B^* of B equipped with the weak* topology is a Hausdorff locally convex topological vector space. Moreover, for any $f \in B \setminus \{0\}$, the subdifferential set $\partial\|\cdot\|_B(f)$ is a convex and weakly* compact subset of B^* . Hence, the Krein-Milman theorem ensures that for any $f \in B \setminus \{0\}$,

$$\partial\|\cdot\|_B(f) = \overline{\text{co}}(\text{ext}(\partial\|\cdot\|_B(f))),$$

where the closed convex hull is taken under the weak* topology of B^* .

We now review the representer theorem for the MNI problem in a general Banach space having a pre-dual space established in [24]. Suppose that B is a Banach space having a pre-dual space B_* . Let $\nu_j, j \in \mathbb{N}_n$, be linearly independent elements in B_* and $\mathbf{z} := [z_j : j \in \mathbb{N}_n] \in \mathbb{R}^n$ be a given vector. Set $\mathcal{V} := \text{span}\{\nu_j : j \in \mathbb{N}_n\}$. We define an operator $\mathcal{L} : B \rightarrow \mathbb{R}^n$ by $\mathcal{L}(f) := [\langle \nu_j, f \rangle_B : j \in \mathbb{N}_n]$, for all $f \in B$, and introduce a subset of B as $M_{\mathbf{z}} := \{f \in B : \mathcal{L}(f) = \mathbf{z}\}$. The MNI problem with the given data $\{(\nu_j, y_j) : j \in \mathbb{N}_n\}$ considered in [24] has the form

$$\inf \{\|f\|_B : f \in M_{\mathbf{z}}\}. \quad (46)$$

The representer theorem established in Proposition 7 of [24] provides a representation of any extreme point of the solution set of the MNI problem (46) with $\mathbf{z} \in \mathbb{R}^n$. We describe this result in the next lemma.

Lemma 12. *Suppose that B is a Banach space having a pre-dual space B_* . Let $\nu_j \in B_*$, $j \in \mathbb{N}_n$, be linearly independent and $\mathbf{z} \in \mathbb{R}^n \setminus \{\mathbf{0}\}$. If \mathcal{V} and $M_{\mathbf{z}}$ are defined as above and $\hat{\nu} \in \mathcal{V}$ satisfies*

$$(\|\hat{\nu}\|_{B_*} \partial\|\cdot\|_{B_*}(\hat{\nu})) \cap M_{\mathbf{z}} \neq \emptyset, \quad (47)$$

then for any extreme point \hat{f} of the solution set of the MNI problem (46), there exist $\gamma_j \in \mathbb{R}$, $j \in \mathbb{N}_n$, with $\sum_{j \in \mathbb{N}_n} \gamma_j = \|\hat{\nu}\|_{B_}$ and $u_j \in \text{ext}(\partial\|\cdot\|_{B_*}(\hat{\nu}))$, $j \in \mathbb{N}_n$, such that*

$$\hat{f} = \sum_{j \in \mathbb{N}_n} \gamma_j u_j.$$

It was pointed out in [24] that the element $\hat{\nu}$ satisfying (47) can be obtained through solving a dual problem of (46). Moreover, we remark that the solution set is a nonempty, convex and weakly* compact subset of B . Hence, by the Krein-Milman theorem, the set of extreme points of the solution set is nonempty and moreover, any solution of problem (46) can be expressed as the weak* limit of a sequence in the convex hull of the set of extreme points.

We now present a representer theorem for a solution of the MNI problem (45), which is a direct consequence of Lemma 12. We introduce a subspace of \mathcal{S} , which is defined by (23) and has been proved to be a pre-dual space of $\mathcal{B}_{\mathcal{N}}$, by

$$\mathcal{V}_{\mathcal{N}} := \text{span } \mathbb{K}_{\mathcal{X}}, \quad (48)$$

and denote by $\mathbb{S}_{\mathcal{X}, \mathbf{Y}}$ the solution set of the MNI problem (45).

We prepare applying Lemma 12 to the MNI problem (45). To this end, we introduce the dual problem of problem (45) as

$$\sup \left\{ \sum_{k \in \mathbb{N}_t} \sum_{j \in \mathbb{N}_m} c_{kj} y_j^k : \left\| \sum_{k \in \mathbb{N}_t} \sum_{j \in \mathbb{N}_m} c_{kj} \mathcal{K}_k(x_j, \cdot) \right\|_{\infty} = 1 \right\}. \quad (49)$$

Note that the dual problem is a finite dimensional optimization problem which has the same optimal value, denoted by C^* , as the MNI problem (45). It has been proved in [5] that there exists at least one solution for the dual problem of the MNI problem in $\ell_1(\mathbb{N})$. By a similar argument, we can show the existence of a solution of the dual problem (49). Suppose that $\hat{\mathbf{c}} := [\hat{c}_{kj} : k \in \mathbb{N}_t, j \in \mathbb{N}_m] \in \mathbb{R}^{t \times m}$ is a solution of the dual problem (49). We let

$$\hat{g}(\cdot) := C^* \sum_{k \in \mathbb{N}_t} \sum_{j \in \mathbb{N}_m} \hat{c}_{kj} \mathcal{K}_k(x_j, \cdot). \quad (50)$$

Theorem 13. *Suppose that m distinct points $x_j \in \mathbb{R}^s$, $j \in \mathbb{N}_m$, and $\mathbf{Y} \in \mathbb{R}^{t \times m} \setminus \{\mathbf{0}\}$ are given, and the functionals in $\mathbb{K}_{\mathcal{X}}$ are linearly independent. Let \hat{g} be the function defined by (50). Then for any $\hat{f} \in \text{ext}(\mathbb{S}_{\mathcal{X}, \mathbf{Y}})$, there exist $\gamma_{\ell} \in \mathbb{R}$, $\ell \in \mathbb{N}_{tm}$, with $\sum_{\ell \in \mathbb{N}_{tm}} \gamma_{\ell} = \|\hat{g}\|_{\infty}$ and $h_{\ell} \in \text{ext}(\|\cdot\|_{\infty}(\hat{g}))$, $\ell \in \mathbb{N}_{tm}$, such that*

$$\hat{f}(x) = \sum_{\ell \in \mathbb{N}_{tm}} \gamma_{\ell} h_{\ell}(x), \quad x \in \mathbb{R}^s. \quad (51)$$

Proof. Proposition 5 ensures that the vector-valued RKBS $\mathcal{B}_{\mathcal{N}}$ has the pre-dual space \mathcal{S} . Note that the functionals in $\mathbb{K}_{\mathcal{X}}$ belong to the pre-dual space \mathcal{S} and are linearly independent. Moreover, since \hat{g} is the function defined by (50), we have that $\hat{g} \in \mathcal{V}_{\mathcal{N}}$, and according to Proposition 37 of [24], \hat{g} satisfies the condition

$$(\|\hat{g}\|_{\infty} \|\partial\| \cdot \|\cdot\|_{\infty}(\hat{g})) \cap \mathcal{M}_{\mathcal{X}, \mathbf{Y}} \neq \emptyset. \quad (52)$$

Hence, the hypothesis of Lemma 12 is satisfied. Then by Lemma 12, we can represent any extreme point \hat{f} of the solution set $\mathbb{S}_{\mathcal{X}, \mathbf{Y}}$ of the MNI problem (45) as in equation (51) for some $\gamma_{\ell} \in \mathbb{R}$, $\ell \in \mathbb{N}_{tm}$, with $\sum_{\ell \in \mathbb{N}_{tm}} \gamma_{\ell} = \|\hat{g}\|_{\infty}$ and $h_{\ell} \in \text{ext}(\|\partial\| \cdot \|\cdot\|_{\infty}(\hat{g}))$, $\ell \in \mathbb{N}_{tm}$. \square

Theorem 13 provides for each extreme point of the solution set of problem (45) an explicit and data-dependent representation by using the elements in $\text{ext}(\|\partial\| \cdot \|\cdot\|_{\infty}(\hat{g}))$. Even more significantly, the essence of Theorem 13 is that although the MNI problem (45) is of infinite dimension, every extreme point of its solution set lays in a *finite* dimensional manifold spanned by tm elements $h_{\ell} \in \text{ext}(\|\partial\| \cdot \|\cdot\|_{\infty}(\hat{g}))$.

As we have demonstrated earlier, the element \hat{g} satisfying (52) can be obtained by solving the dual problem (49) of (45). Since \hat{g} is an element in \mathcal{S} , the subdifferential $\partial\|\cdot\|_\infty(\hat{g})$ is a subset of the space $\mathcal{B}_\mathcal{N}$, which is the dual space of \mathcal{S} . Notice that the subdifferential set $\partial\|\cdot\|_\infty(\hat{g})$ may not be included in space $\mathcal{B}_\mathbb{W}$ defined by (9) which is spanned by the kernel sessions $\mathcal{K}(\cdot, \theta)$, $\theta \in \Theta$. However, a learning solution in the vector-valued RKBS $\mathcal{B}_\mathcal{N}$ is expected to be represented by the kernel sessions $\mathcal{K}(\cdot, \theta)$, $\theta \in \Theta$. For the purpose of obtaining a kernel representation for a solution of problem (45), alternatively to problem (45), we consider a closely related MNI problem in the measure space $\mathcal{M}(\Theta)$ and apply the representer theorem established in [24] to it. We then translate the resulting representer theorem for the MNI problem in $\mathcal{M}(\Theta)$ to that for problem (45), by using the relation between the solutions of these two problems.

We now introduce the MNI problem in the measure space $\mathcal{M}(\Theta)$ with respect to the the sampled dataset \mathbb{D}_m and show the relation between its solution and a solution of problem (45). By defining an operator $\tilde{\mathbf{I}}_\mathcal{X} : \mathcal{M}(\Theta) \rightarrow \mathbb{R}^{t \times m}$ by

$$\tilde{\mathbf{I}}_\mathcal{X}(\mu) := [\langle \mathcal{K}_k(x_j, \cdot), \mu \rangle_{\mathcal{M}(\Theta)} : k \in \mathbb{N}_t, j \in \mathbb{N}_m], \text{ for all } \mu \in \mathcal{M}(\Theta),$$

and introducing a subset $\tilde{\mathcal{M}}_{\mathcal{X}, \mathbf{Y}}$ of $\mathcal{M}(\Theta)$ as

$$\tilde{\mathcal{M}}_{\mathcal{X}, \mathbf{Y}} := \left\{ \mu \in \mathcal{M}(\Theta) : \tilde{\mathbf{I}}_\mathcal{X}(\mu) = \mathbf{Y} \right\}, \quad (53)$$

we formulate the MNI problem in $\mathcal{M}(\Theta)$ as

$$\inf \left\{ \|\mu\|_{\text{TV}} : \mu \in \tilde{\mathcal{M}}_{\mathcal{X}, \mathbf{Y}} \right\}. \quad (54)$$

The next proposition reveals the relation between the solutions of the MNI problems (45) and (54).

Proposition 14. *Suppose that m distinct points $x_j \in \mathbb{R}^s$, $j \in \mathbb{N}_m$, and $\mathbf{Y} \in \mathbb{R}^{t \times m} \setminus \{\mathbf{0}\}$ are given, and the functionals in $\mathbb{K}_\mathcal{X}$ are linearly independent. If $\hat{\mu}$ is a solution of the MNI problem (54), then $f_{\hat{\mu}}(x) := \left[f_{\hat{\mu}}^k(x) : k \in \mathbb{N}_t \right]^\top$, $x \in \mathbb{R}^s$, with $f_{\hat{\mu}}^k$, $k \in \mathbb{N}_t$, defined as in (20) with μ replaced by $\hat{\mu}$, is a solution of the MNI problem (45) and $\|f_{\hat{\mu}}\|_{\mathcal{B}_\mathcal{N}} = \|\hat{\mu}\|_{\text{TV}}$.*

Proof. Note that the measure space $\mathcal{M}(\Theta)$ has the pre-dual space $C_0(\Theta)$. Hence, it follows from Proposition 1 of [23] that the linear independence of the functionals in $\mathbb{K}_\mathcal{X}$ ensures the existence of a solution of problem (54). Assume that $\hat{\mu}$ is a solution of problem (54). We then obtain that $\hat{\mu} \in \tilde{\mathcal{M}}_{\mathcal{X}, \mathbf{Y}}$ and

$$\|\hat{\mu}\|_{\text{TV}} \leq \|\mu\|_{\text{TV}}, \text{ for all } \mu \in \tilde{\mathcal{M}}_{\mathcal{X}, \mathbf{Y}}. \quad (55)$$

By equations (27) and (28) with $g := \mathcal{K}_k(x_j, \cdot)$, we have for each $k \in \mathbb{N}_t$ and each $j \in \mathbb{N}_m$ that

$$\langle \mathcal{K}_k(x_j, \cdot), f_\mu \rangle_{\mathcal{B}_\mathcal{N}} = \langle \mu, \mathcal{K}_k(x_j, \cdot) \rangle_{C_0(\Theta)}, \text{ for all } \mu \in \mathcal{M}(\Theta). \quad (56)$$

Note that $\mathcal{K}_k(x_j, \cdot) \in C_0(\Theta)$ can be viewed as a bounded linear functional on $\mathcal{M}(\Theta)$ and

$$\langle \mu, \mathcal{K}_k(x_j, \cdot) \rangle_{C_0(\Theta)} = \langle \mathcal{K}_k(x_j, \cdot), \mu \rangle_{\mathcal{M}(\Theta)}.$$

Substituting the above equation into the right hand of equation (56) leads to

$$\langle \mathcal{K}_k(x_j, \cdot), f_\mu \rangle_{\mathcal{B}_\mathcal{N}} = \langle \mathcal{K}_k(x_j, \cdot), \mu \rangle_{\mathcal{M}(\Theta)}, \text{ for all } \mu \in \mathcal{M}(\Theta).$$

This implies that $\mathbf{I}_{\mathcal{X}}(f_{\mu}) = \widetilde{\mathbf{I}}_{\mathcal{X}}(\mu)$ for all $\mu \in \mathcal{M}(\Theta)$. As a result, $\mu \in \widetilde{\mathcal{M}}_{\mathcal{X}, \mathbf{Y}}$ if and only if $f_{\mu} \in \mathcal{M}_{\mathcal{X}, \mathbf{Y}}$. Since $\hat{\mu} \in \widetilde{\mathcal{M}}_{\mathcal{X}, \mathbf{Y}}$, we get that $f_{\hat{\mu}} \in \mathcal{M}_{\mathcal{X}, \mathbf{Y}}$. It suffices to verify that

$$\|f_{\hat{\mu}}\|_{\mathcal{B}_{\mathcal{N}}} \leq \|f_{\mu}\|_{\mathcal{B}_{\mathcal{N}}}, \quad \text{for all } f_{\mu} \in \mathcal{M}_{\mathcal{X}, \mathbf{Y}}.$$

Let f_{μ} be an arbitrary element in $\mathcal{M}_{\mathcal{X}, \mathbf{Y}}$. For any $\nu \in \mathcal{M}(\Theta)$ satisfying $f_{\mu} = f_{\nu}$, there holds $f_{\nu} \in \mathcal{M}_{\mathcal{X}, \mathbf{Y}}$. Thus, $\nu \in \widetilde{\mathcal{M}}_{\mathcal{X}, \mathbf{Y}}$. It follows from inequality (55) that

$$\|\hat{\mu}\|_{\text{TV}} \leq \|\nu\|_{\text{TV}}. \quad (57)$$

By taking infimum of both sides of the inequality (57) over $\nu \in \mathcal{M}(\Theta)$ satisfying $f_{\mu} = f_{\nu}$ and noting the definition (22) of the norm $\|f_{\mu}\|_{\mathcal{B}_{\mathcal{N}}}$, we get that

$$\|\hat{\mu}\|_{\text{TV}} \leq \|f_{\mu}\|_{\mathcal{B}_{\mathcal{N}}}. \quad (58)$$

Again by the definition (22) of the norm $\|f_{\mu}\|_{\mathcal{B}_{\mathcal{N}}}$, we obtain that

$$\|f_{\hat{\mu}}\|_{\mathcal{B}_{\mathcal{N}}} \leq \|\hat{\mu}\|_{\text{TV}}. \quad (59)$$

Combining inequalities (58) with (59), we conclude that $\|f_{\hat{\mu}}\|_{\mathcal{B}_{\mathcal{N}}} \leq \|f_{\mu}\|_{\mathcal{B}_{\mathcal{N}}}$. Therefore, $f_{\hat{\mu}}$ is a solution of the MNI problem (45). Moreover, by taking $\mu = \hat{\mu}$ in (58), we get that $\|\hat{\mu}\|_{\text{TV}} \leq \|f_{\hat{\mu}}\|_{\mathcal{B}_{\mathcal{N}}}$. This together with inequality (59) leads to $\|f_{\hat{\mu}}\|_{\mathcal{B}_{\mathcal{N}}} = \|\hat{\mu}\|_{\text{TV}}$. \square

We next derive a representer theorem for a solution of problem (54) by employing Lemma 12. Applying Lemma 12 to problem (54) requires the representation of the extreme points of the subdifferential set $\partial\|\cdot\|_{\infty}(g)$ for any nonzero $g \in C_0(\Theta)$. Here, the subdifferential set $\partial\|\cdot\|_{\infty}(g)$ is a subset of the measure space $\mathcal{M}(\Theta)$. For each $g \in C_0(\Theta)$, let $\Theta(g)$ denote the subset of Θ where the function g attains its maximum norm $\|g\|_{\infty}$, that is,

$$\Theta(g) := \{\theta \in \Theta : |g(\theta)| = \|g\|_{\infty}\}.$$

For each $g \in C_0(\Theta)$, we introduce a subset of $\mathcal{M}(\Theta)$ by

$$\Omega(g) := \{\text{sign}(g(\theta))\delta_{\theta} : \theta \in \Theta(g)\}. \quad (60)$$

Lemma 26 in [24] essentially states that if $g \in C_0(\Theta) \setminus \{0\}$, then

$$\text{ext}(\partial\|\cdot\|_{\infty}(g)) = \Omega(g). \quad (61)$$

We denote by $\widetilde{\mathcal{S}}_{\mathcal{X}, \mathbf{Y}}$ the solution set of the MNI problem (54). We note that the MNI problem (54) shares the same dual problem (49) with the MNI problem (45).

Proposition 15. *Suppose that m distinct points $x_j \in \mathbb{R}^s$, $j \in \mathbb{N}_m$, and $\mathbf{Y} \in \mathbb{R}^{t \times m} \setminus \{\mathbf{0}\}$ are given, and the functionals in $\mathbb{K}_{\mathcal{X}}$ are linearly independent. Let \hat{g} be the function defined by (50). Then for any $\hat{\mu} \in \text{ext}(\widetilde{\mathcal{S}}_{\mathcal{X}, \mathbf{Y}})$, there exist $\gamma_{\ell} \in \mathbb{R}$, $\ell \in \mathbb{N}_{tm}$, with $\sum_{\ell \in \mathbb{N}_{tm}} \gamma_{\ell} = \|\hat{g}\|_{\infty}$ and $\theta_{\ell} \in \Theta(\hat{g})$, $\ell \in \mathbb{N}_{tm}$, such that*

$$\hat{\mu} = \sum_{\ell \in \mathbb{N}_{tm}} \gamma_{\ell} \text{sign}(\hat{g}(\theta_{\ell})) \delta_{\theta_{\ell}}. \quad (62)$$

Proof. Note that the measure space $\mathcal{M}(\Theta)$ has the pre-dual space $C_0(\Theta)$ and the functionals $\mathcal{K}_k(x_j, \cdot)$, $k \in \mathbb{N}_t$, $j \in \mathbb{N}_m$, which belong to the pre-dual space $C_0(\Theta)$, are linearly independent. By Proposition 37 in [24], the function \hat{g} defined by (50) satisfies $\hat{g} \in \mathcal{V}_{\mathcal{N}}$ and

$$(\|\hat{g}\|_{\infty} \partial\|\cdot\|_{\infty}(\hat{g})) \cap \widetilde{\mathcal{M}}_{\mathcal{X}, \mathbf{Y}} \neq \emptyset,$$

in which the subdifferential set $\partial\|\cdot\|_{\infty}(\hat{g})$ is a subset of the measure space $\mathcal{M}(\Theta)$. As a result, the hypothesis of Lemma 12 is satisfied. According to Lemma 12, for any $\hat{\mu} \in \text{ext}(\widetilde{\mathcal{S}}_{\mathcal{X}, \mathbf{Y}})$, there exist $\gamma_{\ell} \in \mathbb{R}$, $\ell \in \mathbb{N}_{tm}$, with $\sum_{\ell \in \mathbb{N}_{tm}} \gamma_{\ell} = \|\hat{g}\|_{\infty}$ and $u_{\ell} \in \text{ext}(\partial\|\cdot\|_{\infty}(\hat{g}))$, $\ell \in \mathbb{N}_{tm}$, such that

$$\hat{\mu} = \sum_{\ell \in \mathbb{N}_{tm}} \gamma_{\ell} u_{\ell}. \quad (63)$$

It follows from equation (61) that for each $\ell \in \mathbb{N}_{tm}$, we have that $u_{\ell} \in \Omega(\hat{g})$. By definition (60) of the set $\Omega(\hat{g})$, for each $\ell \in \mathbb{N}_{tm}$, there exists $\theta_{\ell} \in \Theta(\hat{g})$ such that $u_{\ell} = \text{sign}(\hat{g}(\theta_{\ell}))\delta_{\theta_{\ell}}$. Therefore, we may rewrite the representation (63) of $\hat{\mu}$ as (62). \square

Proposition 15 provides a representation for an extreme point of the solution set of the MNI problem (54). This solution can be converted via Proposition 14 to a solution of the MNI problem (45). We present this result in the next theorem.

Theorem 16. *Suppose that m distinct points $x_j \in \mathbb{R}^s$, $j \in \mathbb{N}_m$, and $\mathbf{Y} \in \mathbb{R}^{t \times m} \setminus \{\mathbf{0}\}$ are given, and the functionals in $\mathbb{K}_{\mathcal{X}}$ are linearly independent. Let $\mathcal{V}_{\mathcal{N}}$ and $\widetilde{\mathcal{M}}_{\mathcal{X}, \mathbf{Y}}$ be defined by (48) and (53), respectively, and let \hat{g} be the function defined by (50). Then the MNI problem (45) has a solution \hat{f} in the form*

$$\hat{f}(x) = \sum_{\ell \in \mathbb{N}_{tm}} \gamma_{\ell} \text{sign}(\hat{g}(\theta_{\ell})) \mathcal{K}(x, \theta_{\ell}), \quad x \in \mathbb{R}^s, \quad (64)$$

for some $\gamma_{\ell} \in \mathbb{R}$, $\ell \in \mathbb{N}_{tm}$, with $\sum_{\ell \in \mathbb{N}_{tm}} \gamma_{\ell} = \|\hat{g}\|_{\infty}$ and $\theta_{\ell} \in \Theta(\hat{g})$, $\ell \in \mathbb{N}_{tm}$.

Proof. By Proposition 1 of [23], the MNI problem (54) has at least one solution. That is, $\widetilde{\mathcal{S}}_{\mathcal{X}, \mathbf{Y}}$ is nonempty and moreover, $\text{ext}(\widetilde{\mathcal{S}}_{\mathcal{X}, \mathbf{Y}})$ is nonempty. We choose $\hat{\mu} \in \text{ext}(\widetilde{\mathcal{S}}_{\mathcal{X}, \mathbf{Y}})$. Proposition 15 ensures that there exist $\gamma_{\ell} \in \mathbb{R}$, $\ell \in \mathbb{N}_{tm}$, with $\sum_{\ell \in \mathbb{N}_{tm}} \gamma_{\ell} = \|\hat{g}\|_{\infty}$ and $\theta_{\ell} \in \Theta(\hat{g})$, $\ell \in \mathbb{N}_{tm}$, such that $\hat{\mu}$ may be expressed as in equation (62). Since $\hat{\mu}$ is a solution of problem (54), we get by Proposition 14 that $f_{\hat{\mu}} = [f_{\hat{\mu}}^k : k \in \mathbb{N}_t]$ is a solution of the MNI problem (45). By definition (20) of $f_{\hat{\mu}}^k$, $k \in \mathbb{N}_t$, we have that

$$f_{\hat{\mu}}^k(x) = \int_{\Theta} \mathcal{K}_k(x, \theta) d\hat{\mu}(\theta), \quad \text{for } x \in \mathbb{R}^s, k \in \mathbb{N}_t. \quad (65)$$

Substituting representation (62) of $\hat{\mu}$ into the right-hand side of equation (65) yields that

$$f_{\hat{\mu}}^k(x) = \sum_{\ell \in \mathbb{N}_{tm}} \gamma_{\ell} \text{sign}(\hat{g}(\theta_{\ell})) \mathcal{K}_k(x, \theta_{\ell}), \quad \text{for } x \in \mathbb{R}^s, k \in \mathbb{N}_t.$$

By letting $\hat{f} := f_{\hat{\mu}}$, we conclude that the MNI problem (45) has a solution \hat{f} in the form of (64). \square

We now return to considering the regularization problem (42). Our goal is to establish representation theorems for a solution of the regularization problem. To ensure the existence of a solution of the regularization problem (42), we always assume that the loss function \mathcal{Q} is lower semi-continuous

on $\mathbb{R}^{t \times m}$. Before establishing the representer theorems for a solution of problem (42), we point out the relation between the solutions of this problem and the MNI problem (45). We denote by $\mathbb{R}_{\mathcal{X}, \mathbf{Y}}$ the solution set of problem (42). By Proposition 11, $\mathbb{R}_{\mathcal{X}, \mathbf{Y}}$ is nonempty. We then introduce a subset $\mathcal{D}_{\mathcal{X}, \mathbf{Y}}$ of $\mathbb{R}^{t \times m}$ by

$$\mathcal{D}_{\mathcal{X}, \mathbf{Y}} := \mathbf{I}_{\mathcal{X}}(\mathbb{R}_{\mathcal{X}, \mathbf{Y}}). \quad (66)$$

Now, recalling that $\mathbb{S}_{\mathcal{X}, \mathbf{Z}}$ denotes the solution set of the MNI problem (45), it follows from Proposition 41 of [23] that

$$\bigcup_{\mathbf{Z} \in \mathcal{D}_{\mathcal{X}, \mathbf{Y}}} \mathbb{S}_{\mathcal{X}, \mathbf{Z}} = \mathbb{R}_{\mathcal{X}, \mathbf{Y}}. \quad (67)$$

Moreover, by Lemma 11 of [24], if the loss function \mathcal{Q} is convex, then

$$\text{ext}(\mathbb{R}_{\mathcal{X}, \mathbf{Y}}) \subset \bigcup_{\mathbf{Z} \in \mathcal{D}_{\mathcal{X}, \mathbf{Y}}} \text{ext}(\mathbb{S}_{\mathcal{X}, \mathbf{Z}}). \quad (68)$$

Below, we convert the representer theorem for a solution of problem (45) stated in Theorem 13 to that for the regularization problem (42) by making use of the relation between the solutions of problems (67) and (68).

Theorem 17. *Suppose that m distinct points $x_j \in \mathbb{R}^s$, $j \in \mathbb{N}_m$, and $\mathbf{Y} \in \mathbb{R}^{t \times m}$ are given, $\lambda > 0$. Let $\mathcal{V}_{\mathcal{N}}$ be defined by (48) and $\mathcal{D}_{\mathcal{X}, \mathbf{Y}}$ be defined by (66).*

1. *If $\mathcal{D}_{\mathcal{X}, \mathbf{Y}} \neq \{\mathbf{0}\}$, then there exists $\hat{f} \in \mathbb{R}_{\mathcal{X}, \mathbf{Y}}$ such that*

$$\hat{f}(x) = \sum_{\ell \in \mathbb{N}_{tm}} \alpha_{\ell} h_{\ell}(x), \quad x \in \mathbb{R}^s, \quad (69)$$

for some $\hat{g} \in \mathcal{V}_{\mathcal{N}}$, $\gamma_{\ell} \in \mathbb{R}$, $\ell \in \mathbb{N}_{tm}$, with $\sum_{\ell \in \mathbb{N}_{tm}} \gamma_{\ell} = \|\hat{g}\|_{\infty}$ and $h_{\ell} \in \text{ext}(\partial\|\cdot\|_{\infty}(\hat{g}))$, $\ell \in \mathbb{N}_{tm}$.

2. *If the loss function \mathcal{Q} is convex, then every nonzero extreme point \hat{f} of $\mathbb{R}_{\mathcal{X}, \mathbf{Y}}$ has the form of (69) for some $\hat{g} \in \mathcal{V}_{\mathcal{N}}$, $\gamma_{\ell} \in \mathbb{R}$, $\ell \in \mathbb{N}_{tm}$, with $\sum_{\ell \in \mathbb{N}_{tm}} \gamma_{\ell} = \|\hat{g}\|_{\infty}$ and $h_{\ell} \in \text{ext}(\partial\|\cdot\|_{\infty}(\hat{g}))$, $\ell \in \mathbb{N}_{tm}$.*

Proof. We first prove Item 1. Note that $\mathbb{R}_{\mathcal{X}, \mathbf{Y}}$ is a nonempty set. It follows from the hypothesis $\mathcal{D}_{\mathcal{X}, \mathbf{Y}} \neq \{\mathbf{0}\}$ that there exists $\hat{h} \in \mathbb{R}_{\mathcal{X}, \mathbf{Y}}$ such that $\hat{\mathbf{Z}} := \mathbf{I}_{\mathcal{X}}(\hat{h}) \neq \mathbf{0}$. According to equation (67) with noting that $\hat{\mathbf{Z}} \in \mathcal{D}_{\mathcal{X}, \mathbf{Y}}$, we have that $\mathbb{S}_{\mathcal{X}, \hat{\mathbf{Z}}} \subset \mathbb{R}_{\mathcal{X}, \mathbf{Y}}$ and thus, $\text{ext}(\mathbb{S}_{\mathcal{X}, \hat{\mathbf{Z}}}) \subset \mathbb{R}_{\mathcal{X}, \mathbf{Y}}$. We choose $\hat{f} \in \text{ext}(\mathbb{S}_{\mathcal{X}, \hat{\mathbf{Z}}})$ and verify that \hat{f} can be represented as in (69). To this end, we choose $\hat{\mathbf{c}} := [\hat{c}_{kj} : k \in \mathbb{N}_t, j \in \mathbb{N}_m] \in \mathbb{R}^{t \times m}$ to be a solution of the dual problem (49) with \mathbf{Y} replaced by $\hat{\mathbf{Z}}$. Let \hat{g} be the function defined by (50) with $\hat{\mathbf{c}}$. Theorem 13 ensures that \hat{f} , as an extreme point of the solution set $\mathbb{S}_{\mathcal{X}, \hat{\mathbf{Z}}}$, can be represented as in (69) for some $\gamma_{\ell} \in \mathbb{R}$, $\ell \in \mathbb{N}_{tm}$, with $\sum_{\ell \in \mathbb{N}_{tm}} \gamma_{\ell} = \|\hat{g}\|_{\infty}$ and $h_{\ell} \in \text{ext}(\partial\|\cdot\|_{\infty}(\hat{g}))$, $\ell \in \mathbb{N}_{tm}$.

We next show Item 2. Assume that \hat{f} is an arbitrary nonzero extreme point of $\mathbb{R}_{\mathcal{X}, \mathbf{Y}}$, that is $\hat{f} \in \text{ext}(\mathbb{R}_{\mathcal{X}, \mathbf{Y}}) \setminus \{0\}$. Because the loss function \mathcal{Q} is convex, the inclusion relation (68) is satisfied. By (68), there exists $\hat{\mathbf{Z}} \in \mathcal{D}_{\mathcal{X}, \mathbf{Y}}$ such that $\hat{f} \in \text{ext}(\mathbb{S}_{\mathcal{X}, \hat{\mathbf{Z}}})$. Clearly, $\hat{\mathbf{Z}} \neq \mathbf{0}$. Assume to the contrary that $\hat{\mathbf{Z}} = \mathbf{0}$. We then must have that $\mathbb{S}_{\mathcal{X}, \hat{\mathbf{Z}}} = \{0\}$. As a result, $\hat{f} = 0$, which is a contradiction. Again, let \hat{g} be defined by (50) with $\hat{\mathbf{c}}$ being a solution of problem (49) with \mathbf{Y} being replaced by $\hat{\mathbf{Z}}$. By Theorem 13, we can represent \hat{f} as in (69) for some $\gamma_{\ell} \in \mathbb{R}$, $\ell \in \mathbb{N}_{tm}$, with $\sum_{\ell \in \mathbb{N}_{tm}} \gamma_{\ell} = \|\hat{g}\|_{\infty}$ and $h_{\ell} \in \text{ext}(\partial\|\cdot\|_{\infty}(\hat{g}))$, $\ell \in \mathbb{N}_{tm}$. \square

Similarly to Theorem 13 for a solution of the MNI problem, Theorem 17 ensures that each extreme point of the solution set of the regularization problem (42) lays in a *finite* dimensional manifold spanned by tm elements $h_\ell \in \text{ext}(\partial\|\cdot\|_\infty(\hat{g}))$ and it has an explicit and data-dependent representation.

We further show that there exists a solution of the regularization problem (42) that can be represented as a linear combination of a finite number of kernel sessions.

Theorem 18. *Suppose that m distinct points $x_j \in \mathbb{R}^s$, $j \in \mathbb{N}_m$, and $\mathbf{Y} \in \mathbb{R}^{t \times m}$ are given, $\lambda > 0$. Let $\mathcal{V}_\mathcal{N}$ be defined by (48) and $\mathcal{D}_{\mathcal{X}, \mathbf{Y}}$ be defined by (66). Then there exists $\hat{f} \in \mathbb{R}_{\mathcal{X}, \mathbf{Y}}$ such that*

$$\hat{f}(x) = \sum_{\ell \in \mathbb{N}_{tm}} \gamma_\ell \text{sign}(\hat{g}(\theta_\ell)) \mathcal{K}(x, \theta_\ell), \quad x \in \mathbb{R}^s, \quad (70)$$

for some $\hat{g} \in \mathcal{V}_\mathcal{N}$, $\gamma_\ell \in \mathbb{R}$, $\ell \in \mathbb{N}_{tm}$, with $\sum_{\ell \in \mathbb{N}_{tm}} \gamma_\ell = \|\hat{g}\|_\infty$ and $\theta_\ell \in \Theta(\hat{g})$, $\ell \in \mathbb{N}_{tm}$.

Proof. Since $\mathbb{R}_{\mathcal{X}, \mathbf{Y}}$ is nonempty and $\mathcal{D}_{\lambda, \mathbf{Y}} \neq \{\mathbf{0}\}$, there exists $\hat{h} \in \mathbb{R}_{\mathcal{X}, \mathbf{Y}}$ such that $\hat{\mathbf{Z}} := \mathbf{I}_\mathcal{X}(\hat{h}) \neq \mathbf{0}$. We choose \hat{g} in the form of (50), where $\hat{\mathbf{c}}$ is a solution of problem (49) with \mathbf{Y} being replaced by $\hat{\mathbf{Z}}$. According to Theorem 16, the MNI problem (45) with $\mathbf{Y} := \hat{\mathbf{Z}}$ has a solution \hat{f} in the form of (70), for some $\gamma_\ell \in \mathbb{R}$, $\ell \in \mathbb{N}_{tm}$, with $\sum_{\ell \in \mathbb{N}_{tm}} \gamma_\ell = \|\hat{g}\|_\infty$ and $\theta_\ell \in \Theta(\hat{g})$, $\ell \in \mathbb{N}_{tm}$. In other words, $\hat{f} \in \mathbb{S}_{\mathcal{X}, \hat{\mathbf{Z}}}$ and it has the form of (70). It follows from relation (67) that $\mathbb{S}_{\mathcal{X}, \hat{\mathbf{Z}}} \subset \mathbb{R}_{\mathcal{X}, \mathbf{Y}}$, which implies that $\hat{f} \in \mathbb{R}_{\mathcal{X}, \mathbf{Y}}$. \square

To close this section, we remark on relevant existing work on representer theorems for deep learning solution. Representer theorems for deep learning have been investigated in [3, 22]. Bohn et al. studied in [3] the composition of reproducing kernel Hilbert spaces as the hypothesis space, and derived a representer theorem for deep kernel learning, which led to a solution in the form of composition of a finite number of kernel sessions. In [22], Unser proposed to optimize the activation functions in the deep neural network by a second-order total-variation regularization, and the resulting representer theorem ensures that the activation function for each node in the optimal network is a linear spline with adaptive knots.

6 Concluding Remarks

We have introduced the hypothesis space $\mathcal{B}_\mathcal{N}$ for deep learning. The hypothesis space that we have come up with is a vector-valued RKBS which has a unique vector-valued reproducing kernel \mathcal{K} and the weak* completion of the vector space $\mathcal{B}_\mathbb{W}$ which is the linear span of the primitive set $\mathcal{A}_\mathbb{W}$ for deep learning. The hypothesis space allows us to understand mathematical insights of deep learning. Specifically, by exploiting the hypothesis space, we have developed representer theorems for solutions of two deep learning models the minimum norm interpolation and regularization problem with deep neural networks.

In the remaining part of this section, we discuss relations among several learning models considered in this paper and comment on advantages of learning in the proposed hypothesis space $\mathcal{B}_\mathcal{N}$.

First, we remark on relations among several learning models studied in this paper. Suppose that the loss function $\mathcal{L}(f_\mu, \mathbb{D}_m)$ takes the form of (41) with a function $\mathcal{Q} : \mathbb{R}^{t \times m} \rightarrow \mathbb{R}_+$ satisfying the condition $\mathcal{Q}(0) = 0$. If $\hat{f}_\mu \in \mathcal{B}_\mathcal{N}$ is a solution of (43), then we have that $\mathcal{L}(\hat{f}_\mu, \mathbb{D}_m) = 0$ and thus, \hat{f}_μ is a solution of the learning model (39). Furthermore, let $f_{\mu, \lambda}$ be a solution of the regularized learning problem (42), and denote by $f_{\mu, 0}$ the limit of $f_{\mu, \lambda}$ as $\lambda \rightarrow 0$ if the limit exists. Then, we

clearly have that $f_{\mu,0}$ is a solution of the learning model (39). Therefore, learning models (42) and (43) are favorable stable substitutes of learning model (39), which may suffer from instability.

Learning in the RKBS space $\mathcal{B}_{\mathcal{N}}$ has several advantages. First of all, the primitive learning model (8), a model commonly considered in the machine learning community, is not guaranteed to have a solution since the primitive set $\mathcal{A}_{\mathbb{W}}$ has neither algebraic nor topological structures. Since the space $\mathcal{B}_{\mathcal{N}}$, which has the pre-dual space \mathcal{S} , is the completion of the linear span $\mathcal{B}_{\mathbb{W}}$ (of $\mathcal{A}_{\mathbb{W}}$) in the weak* topology, it is the smallest RKBS that contains $\mathcal{A}_{\mathbb{W}}$. On one hand, it embraces the intrinsic features of the set $\mathcal{A}_{\mathbb{W}}$ and on the other hand, it has desired algebraic and topological structures which allow us to conduct mathematical analysis of learning on it. Hence, it is natural to consider the RKBS $\mathcal{B}_{\mathcal{N}}$ as a hypothesis space for deep learning problems. Due to the algebraic and topological structures of $\mathcal{B}_{\mathcal{N}}$, unlike model (8), the learning model (42) is guaranteed to have a solution under a mild condition. Last but not least, the reproducing kernel of the RKBS $\mathcal{B}_{\mathcal{N}}$ furnishes a learning solution in the RKBS $\mathcal{B}_{\mathcal{N}}$ a representation in terms of the reproducing kernel, which leads to a representer theorem of deep learning. The resulting representer theorems reveal that although the learning models on the proposed hypothesis space are of infinite dimension, their solutions lay in finite dimensional manifolds and can be expressed as a linear combination of a finite number of kernel sessions determined by given data and the reproducing kernel. In summary, introducing the hypothesis space $\mathcal{B}_{\mathcal{N}}$ to deep learning enables us to understand the insights of deep learning and provide a foundation for its mathematical analysis.

Acknowledgement

R. Wang is supported in part by the Natural Science Foundation of China under grant 12171202 and by the National Key Research and Development Program of China under grants 2020YFA0714101; Y. Xu is supported in part by the US National Science Foundation under grant DMS-2208386, and by the US National Institutes of Health under grant R21CA263876. M. Yan is supported in part by Cheng Fund from College of Science at Old Dominion University. All correspondence should be sent to Y. Xu.

References

- [1] N. ARONSZAJN, *Theory of reproducing kernels*, Transactions of the American mathematical society, 68 (1950), pp. 337–404.
- [2] F. BARTOLUCCI, E. D. VITO, L. ROSASCO, AND S. VIGOGNA, *Understanding neural networks with reproducing kernel Banach spaces*, Applied and Computational Harmonic Analysis, 62 (2023), pp. 194–236.
- [3] B. BOHN, M. GRIEBEL, AND C. RIEGER, *A representer theorem for deep kernel learning*, Journal of Machine Learning Research, 20 (2019), pp. 1–32.
- [4] R. CHENG, R. WANG, AND Y. XU, *A duality approach to regularized learning problems in Banach spaces*, Journal of Complexity, 81 (2024), p. 101818.
- [5] R. CHENG AND Y. XU, *Minimum norm interpolation in the $\ell_1(\mathbb{N})$ space*, Analysis and Applications, 19 (2021), pp. 21–42.
- [6] S.-Y. CHUNG AND Q. SUN, *Barron space for graph convolution neural networks*, arXiv preprint arXiv:2311.02838, (2023).

- [7] J. B. CONWAY, *A Course in Functional Analysis*. 2nd Edition, Springer-Verlag, New York, 1990.
- [8] F. CUCKER AND S. SMALE, *On the mathematical foundations of learning*, Bulletin of the American Mathematical Society, 39 (2002), pp. 1–49.
- [9] I. DAUBECHIES, R. DEVORE, S. FOUCART, B. HANIN, AND G. PETROVA, *Nonlinear approximation and (deep) ReLU networks*, Constructive Approximation, 55 (2022), pp. 127–172.
- [10] G. B. FOLLAND, *Real analysis: modern techniques and their applications.*, vol. 40, John Wiley & Sons, 1999.
- [11] J. HUANG, Y. JIAO, Z. LI, S. LIU, Y. WANG, AND Y. YANG, *An error analysis of generative adversarial networks for learning distributions*, The Journal of Machine Learning Research, 23 (2022), pp. 5047–5089.
- [12] E. KREYSZIG, *Introductory Functional Analysis with Applications*, John Wiley & Sons, New York, 1978.
- [13] Y. LI, S. LU, P. MATHÉ, AND S. V. PEREVERZEV, *Two-layer networks with the $ReLU^k$ activation function: Barron spaces and derivative approximation*, Numerische Mathematik, 156 (2024), pp. 319–344.
- [14] R. LIN, G. SONG, AND H. ZHANG, *Multi-task learning in vector-valued reproducing kernel Banach spaces with the ℓ_1 norm*, Journal of Complexity, 63 (2021), p. 101514.
- [15] R. E. MEGGINSON, *An Introduction to Banach Space Theory*. Graduate Texts in Mathematics, vol. 183, Springer-Verlag, New York, 1998.
- [16] H. N. MHASKAR AND T. POGGIO, *Deep vs. shallow networks: An approximation theory perspective*, Analysis and Applications, 14 (2016), pp. 829–848.
- [17] C. A. MICCHELLI, Y. XU, AND H. ZHANG, *Universal kernels*, Journal of Machine Learning Research, 7 (2006), pp. 2651–2667.
- [18] R. PARHI AND R. D. NOWAK, *Banach space representer theorems for neural networks and ridge splines*, Journal of Machine Learning Research, 22 (2021), pp. 1960–1999.
- [19] B. SCHÖLKOPF, R. HERBRICH, AND A. J. SMOLA, *A generalized representer theorem*, in Proceeding of the Fourteenth Annual Conference on Computational Learning Theory and the Fifth European Conference on Computational Learning Theory, Springer-Verlag, London, 2001, pp. 416–426.
- [20] Z. SHEN, H. YANG, AND S. ZHANG, *Optimal approximation rate of ReLU networks in terms of width and depth*, Journal de Mathématiques Pures et Appliquées, 157 (2022), pp. 101–135.
- [21] J. SHENOUDA, R. PARHI, K. LEE, AND R. D. NOWAK, *Vector-valued variation spaces and width bounds for DNNs: insights on weight decay regularization*, arXiv preprint arXiv:2305.16534, (2023).
- [22] M. UNSER, *A representer theorem for deep neural networks*, Journal of Machine Learning Research, 20 (2019), pp. 1–30.

- [23] R. WANG AND Y. XU, *Representer theorems in Banach spaces: minimum norm interpolation, regularized learning and semi-discrete inverse problems*, Journal of Machine Learning Research, 22 (2021), pp. 1–65.
- [24] R. WANG, Y. XU, AND M. YAN, *Sparse representer theorems for learning in reproducing kernel Banach spaces*, Journal of Machine Learning Research, 25 (2024), pp. 1–45.
- [25] Y. XU, *Sparse machine learning in Banach spaces*, Applied Numerical Mathematics, 187 (2023), pp. 138–157.
- [26] Y. XU AND Q. YE, *Generalized Mercer kernels and reproducing kernel Banach spaces*, Memoirs of the American Mathematical Society, 258 (2019), p. 1243.
- [27] Y. XU AND H. ZHANG, *Convergence of deep convolutional neural networks*, Neural Networks, 153 (2022), pp. 553–563.
- [28] Y. XU AND H. ZHANG, *Convergence of deep ReLU networks*, Neurocomputing, 571 (2024), p. 127174.
- [29] H. ZHANG, Y. XU, AND J. ZHANG, *Reproducing kernel Banach spaces for machine learning.*, Journal of Machine Learning Research, 10 (2009), pp. 2741–2775.
- [30] H. ZHANG AND J. ZHANG, *Vector-valued reproducing kernel Banach spaces with applications to multi-task learning*, Journal of Complexity, 29 (2013), pp. 195–215.
- [31] D.-X. ZHOU, *Universality of deep convolutional neural networks*, Applied and computational harmonic analysis, 48 (2020), pp. 787–794.