

A rate-distortion framework for MCMC algorithms: geometry and factorization of multivariate Markov chains

Michael C.H. Choi^{*1}, Youjia Wang^{†2}, and Geoffrey Wolfer^{‡3}

¹Department of Statistics and Data Science and Yale-NUS College, National
University of Singapore, Singapore

²Department of Statistics and Data Science, National University of
Singapore, Singapore

³Center for AI Project, RIKEN, Tokyo, Japan[§]

April 22, 2024

Abstract

We introduce a framework rooted in a rate distortion problem for Markov chains, and show how a suite of commonly used Markov Chain Monte Carlo (MCMC) algorithms are specific instances within it, where the target stationary distribution is controlled by the distortion function. Our approach offers a unified variational view on the optimality of algorithms such as Metropolis-Hastings, Glauber dynamics, the swapping algorithm and Feynman-Kac path models. Along the way, we analyze factorizability and geometry of multivariate Markov chains. Specifically, we demonstrate that induced chains on factors of a product space can be regarded as information projections with respect to a particular divergence. This perspective yields Han–Shearer type inequalities for Markov chains as well as applications in the context of large deviations and mixing time comparison.

Keywords: Markov chains, f -divergence, mutual information, Markov chain Monte Carlo, large deviations, rate distortion, matrix nearness problem, Metropolis-Hastings,

*Email: mchchoi@nus.edu.sg, corresponding author

†Email: e1124868@u.nus.edu

‡Email: geoffrey.wolfer@riken.jp

§Research conducted while at RIKEN AIP. Current address: Center for Data Science, Waseda University, Tokyo, Japan. email: geo.wolfer@aoni.waseda.jp

simulated annealing, Feynman-Kac path models, Han's inequality, Shearer's lemma, information geometry

AMS 2020 subject classification: 60F10, 60J10, 60J22, 94A15, 94A17

Contents

1	Introduction	3
2	Distance to independence and factorizability of Markov chains	7
2.1	Distance to independence and the closest product chain	12
2.1.1	The closest product chain with prescribed marginals and a large deviation principle of Markov chains	16
2.1.2	A coordinate descent algorithm for finding the closest product chain	19
2.2	Leave-one-out transition matrices and the Han's inequality for KL divergence of Markov chains	19
2.3	Leave- S -out transition matrices and the Shearer's lemma for Markov chains	22
2.4	$(S_i)_{i=1}^n$ -factorizable transition matrices and the distance to $(S_i)_{i=1}^n$ -factorizability	26
2.5	$(C_i)_{i=1}^n$ -factorizable transition matrices with respect to a graph	28
2.6	Comparisons of mixing and hitting time parameters between P and its information projections	31
3	Unified view of key MCMC algorithms via rate distortion theory	34
3.1	Variational interpretation of the probability law of MCMC algorithms	39
3.1.1	Example: Metropolis-Hastings type Markov chains with general acceptance function	40
3.1.2	Example: Glauber dynamics, its marginal transition matrices and the distance to independence	42
3.1.3	Example: swapping algorithm, its marginal transition matrices and consequences of the contraction principle	44
3.1.4	Example: Feynman-Kac path models	46
3.1.5	Example: simulated annealing	47
4	Geometry of multivariate Markov chains induced from the information divergence rate	48
4.1	Information geometry of Markov chains	48

4.2	The exponential family of product chains	49
4.3	The m-family of multivariate chains with prescribed marginal edge measures	53
4.4	Resulting geodesic foliation	54
4.5	Partial projections and the Han–Shearer entropy gap	55

1 Introduction

Consider two random variables X, Y on a common finite state space Ω . We denote by $\mathcal{P}(\Omega)$ to be the set of probability masses on Ω . We write $p_{(X,Y)} \in \mathcal{P}(\Omega^2)$ to be the joint probability mass of (X, Y) while p_X (resp. p_Y) $\in \mathcal{P}(\Omega)$ denotes the marginal probability mass of X (resp. Y) with respect to $p_{(X,Y)}$. Recall that the Kullback-Leibler (KL) divergence from ν to μ is given by

$$\tilde{D}_{KL}(\mu\|\nu) := \sum_{x \in \Omega} \mu(x) \ln \left(\frac{\mu(x)}{\nu(x)} \right),$$

where the usual convention of $0 \ln(0/0) := 0$ applies and $\mu, \nu \in \mathcal{P}(\Omega)$. With the above setup in mind, the classical notion of mutual information between X, Y is defined to be

$$I(X; Y) := \tilde{D}_{KL}(p_{(X,Y)}\|p_X \otimes p_Y),$$

where the symbol $(\mu \otimes \nu)(x, y) := \mu(x)\nu(y)$ for all $x, y \in \Omega$ denotes the product distribution of $\mu, \nu \in \mathcal{P}(\Omega)$. Note that $\mu \otimes \nu \in \mathcal{P}(\Omega^2)$. It can be shown, via the chain rule of the KL divergence (Cover and Thomas, 2006, Theorem 2.5.3), that $p_X \otimes p_Y$ is the unique closest product distribution to $p_{(X,Y)}$, that is,

$$I(X; Y) = \min_{\mu, \nu \in \mathcal{P}(\Omega)} \tilde{D}_{KL}(p_{(X,Y)}\|\mu \otimes \nu). \tag{1}$$

In other words, the mutual information $I(X; Y)$ can broadly be interpreted as an entropic “distance” to the closest product distribution, that is, an entropic distance to independence. From (1), we immediately see that $I(X; Y)$ is non-negative, and vanishes if and only if X, Y are independent (Cover and Thomas, 2006, equation (2.90)).

Among various applications and usages of mutual information in information theory, we are particularly interested in the rate distortion optimization problem with respect to a so-called distortion cost function $c : \Omega \times \Omega \rightarrow \mathbb{R}$. Let $p_{Y|X}$ be the law of Y given X , and suppose that p_Y is known. Then the (Lagrangian relaxation of the) rate distortion optimization ((Cover and Thomas, 2006, Section 10.7), (Polyanskiy and Wu, 2022, Equation (5.30)), (Shafieipoorfard et al., 2016, Section 4)) amounts to the minimization problem

$$\min_{p_{Y|X}} I(X; Y) + \mathbb{E}[c(X, Y)] = \min_{p_{Y|X}} \tilde{D}_{KL}(p_{(X,Y)}\|p_X \otimes p_Y) + \mathbb{E}[c(X, Y)], \tag{2}$$

in which a minimizer $p_{Y|X}^* \in \arg \min_{p_{Y|X}} I(X; Y) + \mathbb{E}[c(X, Y)]$ is given by, for $x, y \in \Omega$,

$$p_{Y|X}^*(y|x) = \frac{p_Y(y)e^{-c(x,y)}}{\sum_{z \in \Omega} p_Y(z)e^{-c(x,z)}}. \quad (3)$$

For a proof of this fact we point out (Cover and Thomas, 2006, Equation (10.124)) and (Polyanskiy and Wu, 2022, Paragraph below (5.30)). We refer interested readers to these cited sources and the references therein for further exposition on rate distortion and its relation with other classical information theory problems such as computation of the channel capacity.

In this paper, we propose to similarly define a rate distortion problem in the context of Markov chains. We will motivate our definition in Section 3, by demonstrating that numerous established MCMC algorithms coincide with solutions of specific instances of this rate distortion problem. More precisely, we consider two transition matrices M, L on a common finite state space \mathcal{X} and we write the set of all transition matrices on \mathcal{X} to be $\mathcal{L}(\mathcal{X})$. Let $f : \mathbb{R}_+ \rightarrow \mathbb{R}$ be a convex function with $f(1) = 0$, and $\pi \in \mathcal{P}(\mathcal{X})$. Analogous to f -divergence between infinitesimal generators of continuous-time Markov chains in (Diaconis and Miclo, 2009, Proposition 1.5), we define the f -divergence from L to M with respect to π to be

$$D_f^\pi(M \| L) := \sum_{x \in \mathcal{X}} \pi(x) \sum_{y \in \mathcal{X}} L(x, y) f\left(\frac{M(x, y)}{L(x, y)}\right),$$

where several standard conventions apply in this definition, see Definition 2.1 below. Note that π is arbitrary and M, L may or not admit π as their respective stationary distribution. In the context of MCMC, we can naturally choose π as the target distribution and consider the f -divergence of two MCMC samplers from L to M with respect to this chosen π . In the special case of $f(t) = t \ln t$ that generates the KL divergence, we shall write D_{KL}^π , and this coincides with the KL divergence rate from L to M when these two admit π as the stationary distribution, which will be the topic of Section 4. For $M \in \mathcal{L}(\mathcal{X}^{(1)})$ and $L \in \mathcal{L}(\mathcal{X}^{(2)})$, their tensor product $M \otimes L \in \mathcal{L}(\mathcal{X}^{(1)} \times \mathcal{X}^{(2)})$ is defined to be

$$(M \otimes L)((x^1, x^2), (y^1, y^2)) := M(x^1, y^1)L(x^2, y^2),$$

where $x^i, y^i \in \mathcal{X}^{(i)}$ for $i = 1, 2$. Suppose now the state space $\mathcal{X} = \mathcal{X}^{(1)} \times \dots \times \mathcal{X}^{(d)}$ takes on a product form for $d \in \mathbb{N}$. Given a $P \in \mathcal{L}(\mathcal{X})$, what is the closest product Markov chain? To put this question in concrete applications, we can think of an interacting particle system with d particles or agents, such as the voter model, that is described by a transition matrix P . What is the dynamics of the closest independent system to P in which each particle or agent evolves independently of each other? That is, we are interested in seeking a minimizer of

$$\mathbb{I}_f^\pi(P) := \min_{L_i \in \mathcal{L}(\mathcal{X}^{(i)}), \forall i \in [d]} D_f^\pi(P \| \otimes_{i=1}^d L_i),$$

which is analogous to the classical mutual information as in (1). Note that we write $\llbracket d \rrbracket := \{1, \dots, d\}$ for $d \in \mathbb{N}$. We will first verify that \mathbb{I}_f^π shares similar geometric properties with the distance to product distributions (i.e. mutual information between two random variables). As a result, $\mathbb{I}_f^\pi(P)$ can be interpreted as a distance to independence of a given P .

Orthogonality considerations allow us to identify and determine the closest product chain under KL divergence, which in turn allows us to compare the worst-case KL divergence mixing time of the original chain with that of the closest product chain. This can be seen as a Markov chain version of the matrix nearness problem as in Higham (1989); Li and Lim (2023), as we are seeking the closest product chain from a given P . In the dual case when f generates the reverse KL divergence, we present a large deviation principle of Markov chains where $\mathbb{I}_f^\pi(P)$ plays a role in the exponent of large deviation probability. These results are presented in Section 2.1 and 2.1.1 below.

We proceed to generalize these notions further. We introduce the leave-one-out, or more generally leave- S -out, transition matrix, and investigate the factorizability of transition matrix with respect to partition or cliques of a given graph in Section 2.2 to 2.5. Observing that leave- S -out transition matrices are instances of Markov chain decomposition (Jerrum et al., 2004) or induced chains Aldous and Fill (2002), we deduce comparison methods for hitting and mixing time parameters such as spectral gap and log-Sobolev constant between P and its information projections.

Note that in Lacker (2023) a similar problem has been investigated in the context of diffusion processes, where the author studies the closest independent diffusion process of a given multivariate diffusion process and identifies the associated Wasserstein gradient flow and consequences for the McKean-Vlasov equation. On the other hand in the present manuscript, we focus on the closest independent Markov chain problem and the underlying geometry induced by information divergences between transition matrices.

We proceed to tackle the rate distortion optimization problem of a source Markov chain $M \in \mathcal{L}(\mathcal{X})$ with respect to π and cost function c . Analogous to (2), we consider

$$\min_{P \in \mathcal{L}(\mathcal{X})} D_{KL}^\pi(P||M) + \mathbb{E}_{X \sim \pi, Y|X=x \sim P(x, \cdot)}[c(X, Y)], \quad (4)$$

in which a minimizer is given by, for $x, y \in \mathcal{X}$,


$$\Psi_c(M)(x, y) := \frac{M(x, y)e^{-c(x, y)}}{\sum_{z \in \mathcal{X}} M(x, z)e^{-c(x, z)}}.$$

The proof and detailed discussions can be found in Section 3 below. We also note that similar optimization problem has been investigated in the literature of risk-sensitive control of Markov chains (Fleming and Hernández-Hernández, 1997, Lemma 2.12). Thus, one can intuitively understand that the source chain M is distorted by the cost function c to yield $\Psi_c(M)$, and this resulting distortion is optimal in the sense of attaining (4).


We shall demonstrate in Section 3 that with different choices of the source Markov chain M and cost function c , the minimizer $\Psi_c(M)$ corresponds to different Markov chain Monte

Carlo (MCMC) algorithms such as Metropolis-Hastings, Glauber dynamics, Feynman-Kac path models, swapping algorithm and simulated annealing. Thus, the transition matrices of these algorithms are optimal in the sense that they are respectively the minimizer of a corresponding rate distortion optimization problem (4). This suggests a unifying variational representation of these common MCMC algorithms. In this vein, we mention that our work is in line with the body of literature on optimal MCMC where one seeks to find a transition matrix that optimizes a mixing criteria, such as minimizing the worst-case or average asymptotic variance, minimizing the second largest eigenvalue in modulus or minimizing the average hitting time, see for instance [Borkar and Miclo \(2020\)](#); [Boyd et al. \(2004, 2009\)](#); [Chen et al. \(2012\)](#); [Huang et al. \(2018\)](#); [Wu and Chu \(2015\)](#).

In [Billera and Diaconis \(2001\)](#); [Diaconis and Miclo \(2009\)](#) and [Choi and Wolfer \(2024+\)](#), one is searching for the closest π -reversible Markov generator by projecting a proposal generator onto the π -reversible space. This yields generators such as Metropolis-Hastings or Barker proposal, and these resulting algorithms are all π -reversible. However, it is not clear at the moment whether this geometric projection approach can recover more complicated MCMC algorithms such as the swapping algorithm. In the rate distortion optimization problem, we are searching for the closest, not necessarily reversible, Markov chain subject to a distortion cost constraint, where the constraint may enforce π -reversibility or π -stationarity. Using this rate distortion approach, we manage to recover non-reversible MCMC algorithms such as non-reversible Metropolis-Hastings in Section 3.1.1 and more complicated algorithms such as the swapping algorithm in Section 3.1.3. Another unified framework, known as involutive MCMC, has been proposed and analyzed in [Neklyudov et al. \(2020\)](#), where different MCMC algorithms can be considered as special cases in the framework by choosing different involution functions. On the other hand, our proposed approach naturally links with variational principles in information theory via mutual information and rate-distortion optimization.



(a). Encoder-decoder in rate distortion. An example of x is an input image such as the cameraman image [Schreiber \(1978\)](#), while y is the resulting distorted image.



(b). Selection-mutation in MCMC algorithms. An example of x is an initial configuration of the Ising model, and y is the output configuration of an MCMC algorithm.

Figure 1: Analogy between encoder-decoder and selection-mutation

We illustrate this connection between rate distortion and MCMC algorithms in Figure 1. In rate distortion theory, we write x to be an input signal or data source such as a video or an image. It is then fed into the encoder to generate an encoded signal x' for transmission, and then passed into the decoder to generate the output y . If the objective is to minimize (2), then an optimal conditional distribution $p_{Y|X}$ is given as in (3). On the other hand, the evolution of many MCMC algorithms can be described as a two-stage selection-mutation procedure. In this context, the input x is considered as the initial configuration of the MCMC algorithm, such as a configuration of an Ising model. A source Markov chain or proposal chain M is utilized for selection: think of the Glauber dynamics in which a random site is selected with its spin value flipped, which generates a candidate configuration x' . x' is then allowed to mutate, such as by feeding it into a Metropolis filter, to generate the output y . If the objective is to minimize (4), then the optimal transition matrix from x to y is given by $\Psi_c(M)$.

Finally, in Section 4, we inspect the geometry of multivariate Markov chains induced from the conventional information divergence rate. Specifically, we demonstrate that product transition matrices forms an exponential family while multivariate transition matrices with prescribed marginal edge measures form a mixture family. This leads to the establishment of an m-foliation on the set of multivariate transition matrices.

The rest of this paper is organized as follows. In Section 2, we first recall the notion of f -divergences between transition matrices and probability measures. We derive a few important properties of these divergences, which allow us to define an entropic distance to independence of a given P on a finite product state space in Section 2.1. In Section 2.1.1, we determine and identify the closest product chain under KL divergence, and present a large deviation principle in this context. We investigate the factorizability of P with respect to partition or cliques of a given graph in Section 2.2 to 2.6. These considerations eventually lead us to study a rate distortion optimization problem of a source Markov chain in Section 3, where we demonstrate that many common MCMC algorithms can be shown to be an optimal chain under suitable source chain and cost function. In Section 4, following Nagaoka (2005), we analyze the geometric structure of irreducible multivariate Markov chains.

2 Distance to independence and factorizability of Markov chains

On a finite state space \mathcal{X} , we define $\mathcal{L} = \mathcal{L}(\mathcal{X})$ as the set of transition matrices of discrete-time homogeneous Markov chains. We denote by $\mathcal{P}(\mathcal{X})$ to be the set of probability masses on \mathcal{X} . Let $\pi \in \mathcal{P}(\mathcal{X})$ be any given positive probability distribution, and denote $\mathcal{L}(\pi) \subset \mathcal{L}$ as the set of π -reversible transition matrices on \mathcal{X} , where a transition matrix $P \in \mathcal{L}$ is said to be π -reversible if $\pi(x)P(x, y) = \pi(y)P(y, x)$ for all $x, y \in \mathcal{X}$. We also say that $P \in \mathcal{L}$ is π -stationary if it satisfies $\pi P = \pi$. Suppose that P is π -stationary, then the π -dual or the time reversal of P , P^* , is defined to be $P^*(x, y) := \frac{\pi(y)}{\pi(x)}P(y, x)$, for all $x, y \in \mathcal{X}$.

First, we give the definition of f -divergence of Markov chains and recall that of probability measures.

Definition 2.1 (f -divergence of Markov chains and of probability measures). *Let $f : \mathbb{R}^+ \rightarrow \mathbb{R}$ be a convex function and $f(1) = 0$. For given $\pi \in \mathcal{P}(\mathcal{X})$ and transition matrices $M, L \in \mathcal{L}$, we define the f -divergence from L to M with respect to π as*

$$D_f^\pi(M\|L) := \sum_{x \in \mathcal{X}} \pi(x) \sum_{y \in \mathcal{X}} L(x, y) f\left(\frac{M(x, y)}{L(x, y)}\right). \quad (5)$$

For two probability measures $\mu, \nu \in \mathcal{P}(\Omega)$ with Ω finite, the f -divergence from ν to μ is defined to be

$$\tilde{D}_f(\mu\|\nu) := \sum_{x \in \Omega} \nu(x) f\left(\frac{\mu(x)}{\nu(x)}\right), \quad (6)$$

where we apply the usual convention that $0f(\frac{0}{0}) := 0$ and $0f(\frac{a}{0}) := af'(+\infty)$ with $f'(+\infty) := \lim_{x \rightarrow 0^+} xf(\frac{1}{x})$ for $a > 0$ in the two definitions above. We also adapt the convention that $0 \cdot \infty := 0$.

In the special case of taking $f(t) = t \ln t$, we recover the Kullback-Leibler (KL) divergence. In this case we shall write D_{KL}^π and \tilde{D}_{KL} respectively. In particular, when M is assumed to be π -stationary, we write $D(M\|L) := D_{KL}^\pi(M\|L)$ can be interpreted as the KL divergence rate from L to M , see [Rached et al. \(2004\)](#). This divergence $D(M\|L)$ shall be investigated in Section 4 in studying the information geometry of multivariate chains, while other sections concern a general D_{KL}^π with respect to a fixed π . Notably f -divergences of Markov chains have also been proposed for estimating the transition matrix from samples generated from Markov chain, for instance in [Hao et al. \(2018\)](#).

In the sequel, a majority of our focus is devoted to state space that takes on a product form, that is, $\mathcal{X} = \mathcal{X}^{(1)} \times \dots \times \mathcal{X}^{(d)} =: \times_{i=1}^d \mathcal{X}^{(i)}$ for $d \in \mathbb{N}$. A transition matrix $P \in \mathcal{L}(\times_{i=1}^d \mathcal{X}^{(i)})$ is said to be of product form if there exists $M_i \in \mathcal{L}(\mathcal{X}^{(i)})$ for $i \in \llbracket d \rrbracket$ such that P can be expressed as a tensor product of the form

$$P = \otimes_{i=1}^d M_i.$$

This notion of product chain has appeared in ([Levin and Peres, 2017](#), Exercise 12.7) and differs from another slightly different “product-type chain” in [Mathé \(1998\)](#) or “product chain” introduced in ([Levin and Peres, 2017](#), Section 12.4, 20.4). Analogously, a probability mass $\mu \in \mathcal{P}(\times_{i=1}^d \Omega^{(i)})$ is said to be of product form if there exists $\nu_i \in \mathcal{P}(\Omega^{(i)})$ for $i \in \llbracket d \rrbracket$ such that

$$\mu = \otimes_{i=1}^d \nu_i.$$

With the above notations in mind, we are now ready to introduce f -divergences from a product chain (resp. product distribution) to a given M (resp. μ). Let $f : \mathbb{R}^+ \rightarrow \mathbb{R}$ be a convex function and $f(1) = 0$. For given $\pi \in \mathcal{P}(\times_{i=1}^d \mathcal{X}^{(i)})$, $M \in \mathcal{L}(\times_{i=1}^d \mathcal{X}^{(i)})$ and a product

transition matrix $\otimes_{i=1}^d L_i$ with $L_i \in \mathcal{L}(\mathcal{X}^{(i)})$ for $i \in \llbracket d \rrbracket$, the f -divergence from $\otimes_{i=1}^d L_i$ to M with respect to π is

$$D_f^\pi(M \parallel \otimes_{i=1}^d L_i).$$

Similarly, for $\mu \in \mathcal{P}(\times_{i=1}^d \Omega^{(i)})$ and a product distribution $\otimes_{i=1}^d \nu_i$ with $\nu_i \in \mathcal{P}(\Omega^{(i)})$ for $i \in \llbracket d \rrbracket$, the f -divergence from $\otimes_{i=1}^d \nu_i$ to μ is

$$\tilde{D}_f(\mu \parallel \otimes_{i=1}^d \nu_i).$$

In particular, when we choose $f(t) = t \ln t$ which generates the KL divergence, we write

$$D_{KL}^\pi(M \parallel \otimes_{i=1}^d L_i), \quad \tilde{D}_{KL}(\mu \parallel \otimes_{i=1}^d \nu_i).$$

Remark 2.1 (On mutual information and interaction information). *We remark that there exists a body of literature on various generalizations of mutual information to the case of $d > 2$ random variables, see for example [Ghassami and Kiyavash \(2017\)](#) and the references therein.*

Observe that in the special case when $\pi = \delta_x$ with $x = (x^1, \dots, x^d)$, the Dirac point mass at x , we have $D_f^\pi(M \parallel \otimes_{i=1}^d L_i) = \tilde{D}_f(M((x^1, \dots, x^d), \cdot) \parallel (\otimes_{i=1}^d L_i)((x^1, \dots, x^d), \cdot))$. As such, we can interpret D_f^π as a generalization of \tilde{D}_f . We also note that D_f^π can be written as a π -weighted average of \tilde{D}_f , since we have

$$D_f^\pi(M \parallel \otimes_{i=1}^d L_i) = \sum_{x \in \mathcal{X}} \pi(x) \tilde{D}_f(M(x, \cdot) \parallel (\otimes_{i=1}^d L_i)(x, \cdot)).$$

Our next proposition summarizes some fundamental yet useful properties of D_f^π :

Proposition 2.1. *Denote the state space to be $\mathcal{X} = \times_{i=1}^d \mathcal{X}^{(i)}$. Let $\pi \in \mathcal{P}(\mathcal{X})$, $M \in \mathcal{L}(\mathcal{X})$ and $M_i, L_i \in \mathcal{L}(\mathcal{X}^{(i)})$ for $i \in \llbracket d \rrbracket$.*

1. (Non-negativity)

$$D_f^\pi(M \parallel \otimes_{i=1}^d L_i) \geq 0.$$

Suppose that π is a positive probability mass, that is, $\min_{x \in \mathcal{X}} \pi(x) > 0$. Then the equality holds if and only if $M = \otimes_{i=1}^d L_i$.

2. (Convexity) Fix $L_i \in \mathcal{L}(\mathcal{X}^{(i)})$ for $i \in \llbracket d \rrbracket$. The mapping

$$\mathcal{L}(\mathcal{X}) \ni M \mapsto D_f^\pi(M \parallel \otimes_{i=1}^d L_i)$$

is convex in M .

3. (Chain rule of KL divergence) Let $\pi = \otimes_{i=1}^d \pi^{(i)}$ be a product distribution with $\pi^{(i)} \in \mathcal{P}(\mathcal{X}^{(i)})$. Then we have

$$D_{KL}^\pi(\otimes_{i=1}^d M_i \| \otimes_{i=1}^d L_i) = \sum_{i=1}^d D_{KL}^{\pi^{(i)}}(M_i \| L_i),$$

where each $D_{KL}^{\pi^{(i)}}(M_i \| L_i)$ is weighted by $\pi^{(i)}$ for $i \in \llbracket d \rrbracket$.

4. (Bounds of squared Hellinger distance) Let $f(t) = (\sqrt{t} - 1)^2$ that generates the squared Hellinger distance and $\pi = \otimes_{i=1}^d \pi^{(i)}$ be a product distribution with $\pi^{(i)} \in \mathcal{P}(\mathcal{X}^{(i)})$. We have

$$\max_{i \in \llbracket d \rrbracket} D_f^{\pi^{(i)}}(M_i \| L_i) \leq D_f^\pi(\otimes_{i=1}^d M_i \| \otimes_{i=1}^d L_i) \leq \sum_{i=1}^d D_f^{\pi^{(i)}}(M_i \| L_i).$$

5. (Bisection property) Suppose that M is π -stationary and L_i is $\pi^{(i)}$ -stationary, where $\pi = \otimes_{i=1}^d \pi^{(i)}$ is a product distribution and $\pi^{(i)} \in \mathcal{P}(\mathcal{X}^{(i)})$ for $i \in \llbracket d \rrbracket$. Then we have

$$D_f^\pi(M \| \otimes_{i=1}^d L_i) = D_f^\pi(M^* \| \otimes_{i=1}^d L_i^*).$$

In particular, if $L_i \in \mathcal{L}(\pi^{(i)})$, then the above leads to

$$D_f^\pi(M \| \otimes_{i=1}^d L_i) = D_f^\pi(M^* \| \otimes_{i=1}^d L_i).$$

Remark 2.2. The Hellinger distance is commonly used to assess the convergence to equilibrium of product Markov chains, see for example [Chen and Kumagai \(2018\)](#); [Levin and Peres \(2017\)](#).

Remark 2.3. Note that Proposition 2.1 item (1), (2) and (5) also hold when the second argument is not a product transition matrix.

Proof. For brevity, throughout this proof we write $x = (x^1, \dots, x^d)$ and $y = (y^1, \dots, y^d)$. We first prove item (1). Since D_f^π is a f -divergence from $\otimes_{i=1}^d L_i$ to M with respect to π , it is non-negative according to [Wang and Choi \(2023\)](#). Since π is a positive probability mass, equality holds if and only if for all $x \in \mathcal{X}$ we have $\tilde{D}_f(M(x, \cdot) \| (\otimes_{i=1}^d L_i)(x, \cdot)) = 0$ if and only if $M = \otimes_{i=1}^d L_i$ (see for instance [Polyanskiy and Wu \(2022\)](#)).

Next, we prove item (2). We see that

$$D_f^\pi(M \| \otimes_{i=1}^d L_i) = \sum_{x \in \mathcal{X}} \pi(x) \tilde{D}_f(M(x, \cdot) \| (\otimes_{i=1}^d L_i)(x, \cdot)).$$

Since $M(x, \cdot) \mapsto \tilde{D}_f(M(x, \cdot) \| (\otimes_{i=1}^d L_i)(x, \cdot))$ is convex and $D_f^\pi(M \| \otimes_{i=1}^d L_i)$ is a π -weighted sum of convex functions, it is convex.

We proceed to prove item (3). First, we consider the case where $L_i(x^i, y^i) > 0$ for all i or $M_i(x^i, y^i) = 0$ whenever $L_i(x^i, y^i) = 0$, i.e. $M_i(x^i, \cdot) \ll L_i(x^i, \cdot)$ for all i . We see that

$$\begin{aligned}
D_{KL}^\pi(\otimes_{i=1}^d M_i \| \otimes_{i=1}^d L_i) &= \sum_{x, y \in \mathcal{X}} \pi(x) \prod_{j=1}^d M_j(x^j, y^j) \ln \left(\prod_{i=1}^d \frac{M_i(x^i, y^i)}{L_i(x^i, y^i)} \right) \\
&= \sum_{x, y \in \mathcal{X}} \pi(x) \prod_{j=1}^d M_j(x^j, y^j) \sum_{i=1}^d \ln \left(\frac{M_i(x^i, y^i)}{L_i(x^i, y^i)} \right) \\
&= \sum_{i=1}^d \sum_{x, y \in \mathcal{X}} \pi(x) \prod_{j=1}^d M_j(x^j, y^j) \ln \left(\frac{M_i(x^i, y^i)}{L_i(x^i, y^i)} \right) \\
&= \sum_{i=1}^d \sum_{x^i, y^i} \pi^{(i)}(x^i) M_i(x^i, y^i) \ln \left(\frac{M_i(x^i, y^i)}{L_i(x^i, y^i)} \right) \\
&= \sum_{i=1}^d D_{KL}^{\pi^{(i)}}(M_i \| L_i).
\end{aligned}$$

Next, we consider the case where there exists i such that $M_i(x^i, y^i) > 0$ yet $L_i(x^i, y^i) = 0$. Since we take $f(t) = t \ln t$ in KL divergence, we have $f'(\infty) = \infty$, and hence both sides are ∞ in item (3).

Now, we prove item (4). For the upper bound, we note that

$$\begin{aligned}
D_f^\pi(\otimes_{i=1}^d M_i \| \otimes_{i=1}^d L_i) &= \sum_{x \in \mathcal{X}} \pi(x) \tilde{D}_f((\otimes_{i=1}^d M_i)(x, \cdot) \| (\otimes_{i=1}^d L_i)(x, \cdot)) \\
&\leq \sum_{x \in \mathcal{X}} \pi(x) \sum_{i=1}^d \tilde{D}_f(M_i(x^i, \cdot) \| L_i(x^i, \cdot)) \\
&= \sum_{i=1}^d \sum_{x^i \in \mathcal{X}^{(i)}} \pi^{(i)}(x^i) \tilde{D}_f(M_i(x^i, \cdot) \| L_i(x^i, \cdot)) \\
&= \sum_{i=1}^d D_f^{\pi^{(i)}}(M_i \| L_i),
\end{aligned}$$

where we apply (Levin and Peres, 2017, Lemma 20.9) in the inequality. On the other hand, the lower bound can be seen via

$$\begin{aligned}
D_f^\pi(\otimes_{i=1}^d M_i \| \otimes_{i=1}^d L_i) &= \sum_{x \in \mathcal{X}} \pi(x) \tilde{D}_f((\otimes_{i=1}^d M_i)(x, \cdot) \| (\otimes_{i=1}^d L_i)(x, \cdot)) \\
&\geq \sum_{x \in \mathcal{X}} \pi(x) \max_{i \in [d]} \tilde{D}_f(M_i(x^i, \cdot) \| L_i(x^i, \cdot)) \\
&\geq \sum_{x \in \mathcal{X}} \pi(x) \tilde{D}_f(M_i(x^i, \cdot) \| L_i(x^i, \cdot))
\end{aligned}$$

$$= D_f^{\pi^{(i)}}(M_i \| L_i),$$

where the first inequality follows from (Chen and Kumagai, 2018, Proposition 2.3). The desired result follows by taking maximum over $i \in \llbracket d \rrbracket$.

Finally, we prove item (5). We compute that

$$\begin{aligned} D_f^\pi(M \| \otimes_{i=1}^d L_i) &= \sum_{x, y \in \mathcal{X}} \pi(x) \prod_{i=1}^d L_i(x^i, y^i) f\left(\frac{M(x, y)}{\prod_{i=1}^d L_i(x^i, y^i)}\right) \\ &= \sum_{L_i(x^i, y^i) > 0 \ \forall i \in \llbracket d \rrbracket} \pi(x) \prod_{i=1}^d L_i(x^i, y^i) f\left(\frac{M(x, y)}{\prod_{i=1}^d L_i(x^i, y^i)}\right) \\ &\quad + \sum_{M(x, y) > 0, \ L_i(x^i, y^i) = 0 \text{ for some } i \in \llbracket d \rrbracket} \pi(x) \prod_{i=1}^d L_i(x^i, y^i) f\left(\frac{M(x, y)}{\prod_{i=1}^d L_i(x^i, y^i)}\right) \\ &= \sum_{L_i^*(y^i, x^i) > 0 \ \forall i \in \llbracket d \rrbracket} \pi(y) \prod_{i=1}^d L_i^*(y^i, x^i) f\left(\frac{M^*(y, x)}{\prod_{i=1}^d L_i^*(y^i, x^i)}\right) \\ &\quad + \sum_{M(x, y) > 0, \ L_i(x^i, y^i) = 0 \text{ for some } i \in \llbracket d \rrbracket} \pi(x) M(x, y) f'(\infty) \\ &= \sum_{L_i^*(y^i, x^i) > 0 \ \forall i \in \llbracket d \rrbracket} \pi(y) \prod_{i=1}^d L_i^*(y^i, x^i) f\left(\frac{M^*(y, x)}{\prod_{i=1}^d L_i^*(y^i, x^i)}\right) \\ &\quad + \sum_{M^*(y, x) > 0, \ L_i^*(y^i, x^i) = 0 \text{ for some } i \in \llbracket d \rrbracket} \pi(y) M^*(y, x) f'(\infty) \\ &= D_f^\pi(M^* \| \otimes_{i=1}^d L_i^*). \end{aligned}$$

□

2.1 Distance to independence and the closest product chain

Given a Markov chain with transition matrix P on a finite product state space, how far away is it from being a product chain? In other words, what is the (information-theoretic) “distance”, in a broad sense, to independence? One possible way to measure this distance is by means of projection. Throughout this section, unless otherwise specified we shall consider a product state space of the form $\mathcal{X} = \times_{i=1}^d \mathcal{X}^{(i)}$. We now define

Definition 2.2 (Distance to independence of P with respect to D_f^π). *Given $P \in \mathcal{L}(\mathcal{X})$, we define the distance to independence of P with respect to D_f^π to be*

$$\mathbb{I}_f^\pi(P) := \min_{L_i \in \mathcal{L}(\mathcal{X}^{(i)}), \forall i \in \llbracket d \rrbracket} D_f^\pi(P \| \otimes_{i=1}^d L_i). \quad (7)$$

In particular, when we take $f(t) = t \ln t$ that generates the KL divergence, we write $\mathbb{I}^\pi(P) := \mathbb{I}_f^\pi(P)$ in this case. If P is π -stationary, then we also write $\mathbb{I}(P) = \mathbb{I}^\pi(P)$.

Note that since $L \mapsto D_f^\pi(M||L)$ is continuous and the set $\times_{i=1}^d \mathcal{L}(\mathcal{X}^{(i)})$ is compact, the minimization problem (7) is always attained. The next result states that this distance to independence of P is zero if and only if P is a product chain under suitable assumptions on π . This is analogous to the property that if two random variables are independent, then their correlation is zero.

Proposition 2.2. *Assume the same setting as in Proposition 2.1. Let $\pi \in \mathcal{P}(\mathcal{X})$ and $P \in \mathcal{L}(\mathcal{X})$. We have*

$$\mathbb{I}_f^\pi(P) \geq 0.$$

Suppose that π is a positive probability mass, that is, $\min_{x \in \mathcal{X}} \pi(x) > 0$. Then the equality holds if and only if P is a product chain.

Proof. The non-negativity is clear from Proposition 2.1. If P is a product chain, then clearly $\mathbb{I}_f(P) = 0$. For the other direction, if $\mathbb{I}_f(P) = 0$, then $D_f^\pi(P||\otimes_{i=1}^d L_i) = 0$ for some L_i since the minimization in (7) is exactly attained. By Proposition 2.1, $P = \otimes_{i=1}^d L_i$. \square

Let us now recall the notion of edge measure of a Markov chain (see e.g. (Levin and Peres, 2017, equation (7.5))). Let $\pi \in \mathcal{P}(\mathcal{X})$ and $P \in \mathcal{L}(\mathcal{X})$. The edge measure $\pi \boxtimes P \in \mathcal{P}(\mathcal{X} \times \mathcal{X})$ is defined to be, for $x, y \in \mathcal{X}$,

$$(\pi \boxtimes P)(x, y) := \pi(x)P(x, y).$$

Note that this edge measure encodes the probability of observing a consecutive pair generated from the chain with transition matrix P starting from the distribution π .

We proceed to define the i th marginal transition matrix of P with respect to π :

Definition 2.3 ($P_\pi^{(i)}$: the i th marginal transition matrix of P with respect to π). *Assume the same setting as in Proposition 2.1. Let $\pi \in \mathcal{P}(\mathcal{X})$ be a positive probability mass, $P \in \mathcal{L}(\mathcal{X})$ and $i \in \llbracket d \rrbracket$. For any $(x^i, y^i) \in \mathcal{X}^{(i)} \times \mathcal{X}^{(i)}$, we define*

$$\begin{aligned} P_\pi^{(i)}(x^i, y^i) &:= \frac{\sum_{(x^j, y^j); j \neq i} \pi(x^1, \dots, x^d) P((x^1, \dots, x^d), (y^1, \dots, y^d))}{\sum_{x^j; j \neq i} \pi(x^1, \dots, x^d)} \\ &= \frac{\sum_{(x^j, y^j); j \neq i} (\pi \boxtimes P)((x^1, \dots, x^d), (y^1, \dots, y^d))}{\pi^{(i)}(x^i)}, \end{aligned}$$

where $\pi^{(i)}$ is the i th marginal probability mass of π . Note that $P_\pi^{(i)} \in \mathcal{L}(\mathcal{X}^{(i)})$. When P is π -stationary, we omit the subscript and write $P^{(i)}$.

First, we note that $P^{(i)}$ can be understood as a special case of the keep- S -in transition matrix to be introduced in Definition 2.5 below, which is a further special case of various notions of “projection chains” investigated in Aldous and Fill (2002); Chen et al. (1999); Jerrum et al. (2004). Second, we see that if P is π -stationary, then $P_\pi^{(i)}$ is $\pi^{(i)}$ -stationary. Similarly, if P is π -reversible, then $P_\pi^{(i)}$ is $\pi^{(i)}$ -reversible. As a result, $\otimes_{i=1}^d P_\pi^{(i)}$ is thus $\otimes_{i=1}^d \pi^{(i)}$ -stationary, and hence in general $\otimes_{i=1}^d P_\pi^{(i)}$ is not π -stationary. In the case where $\pi = \otimes_{i=1}^d \pi^{(i)}$ is a product stationary distribution, then $\otimes_{i=1}^d P_\pi^{(i)}$ is π -stationary. A generalization of the above discussions can be found in Proposition 2.3 below.

For a concrete example of $P_\pi^{(i)}$, we point to the examples of Glauber dynamics and swapping algorithm where we calculate explicitly their respective marginal transition matrices in Section 3.1.2 and Section 3.1.3.

In our next result, we state that under the KL divergence and positivity of π , a Pythagorean identity holds and it implies that the product chain with transition matrix $\otimes_{i=1}^d P_\pi^{(i)}$ is the unique closest product chain to P :

Theorem 2.1. *Assume the same setting as in Proposition 2.1. Let $\pi \in \mathcal{P}(\mathcal{X})$, $P \in \mathcal{L}(\mathcal{X})$ and $L_i \in \mathcal{L}(\mathcal{X}^{(i)})$ for $i \in \llbracket d \rrbracket$. We then have*

1. (Pythagorean identity of D_{KL}^π) *Let π be a positive probability mass. We have*

$$\begin{aligned} D_{KL}^\pi(P \| \otimes_{i=1}^d L_i) &= D_{KL}^\pi(P \| \otimes_{i=1}^d P_\pi^{(i)}) + D_{KL}^\pi(\otimes_{i=1}^d P_\pi^{(i)} \| \otimes_{i=1}^d L_i) \\ &= D_{KL}^\pi(P \| \otimes_{i=1}^d P_\pi^{(i)}) + \sum_{i=1}^d D_{KL}^{\pi^{(i)}}(P_\pi^{(i)} \| L_i), \end{aligned}$$

where each $D_{KL}^{\pi^{(i)}}(P_\pi^{(i)} \| L_i)$ is weighted by $\pi^{(i)}$, the i th marginal distribution of π . In particular, the unique minimizer that solves (7) is given by $\otimes_{i=1}^d P_\pi^{(i)}$, that is,

$$\mathbb{I}^\pi(P) = D_{KL}^\pi(P \| \otimes_{i=1}^d P_\pi^{(i)}).$$

2. (Bisection property) *Suppose that P is π -stationary, where $\pi = \otimes_{i=1}^d \pi^{(i)}$ is a product distribution and $\pi^{(i)} \in \mathcal{P}(\mathcal{X}^{(i)})$ for $i \in \llbracket d \rrbracket$. We have*

$$\mathbb{I}^\pi(P) = \mathbb{I}^\pi(P^*).$$

In other words, the distance to independence of P with respect to D_{KL}^π and that of its time-reversal P^* is the same.

Remark 2.4 (Distance to independence as KL divergence rate between P and its closest product chain). *Suppose that P is π -stationary, and $(\mathbf{X}_n)_{n \in \mathbb{N}} = (X_n^1, X_n^2, \dots, X_n^d)_{n \in \mathbb{N}}$ (resp. $(\mathbf{Y}_n)_{n \in \mathbb{N}} = (Y_n^1, Y_n^2, \dots, Y_n^d)_{n \in \mathbb{N}}$) is the discrete-time homogeneous Markov chain with transition matrix P (resp. $\otimes_{i=1}^d P_\pi^{(i)}$). Then, the distance to independence of P can be written as*

$$\mathbb{I}(P) = D(P \| \otimes_{i=1}^d P_\pi^{(i)}) = \lim_{n \rightarrow \infty} \frac{1}{n} \tilde{D}_{KL}(\mu_n \| \nu_n),$$

where $\mathbf{X}_n \sim \mu_n, \mathbf{Y}_n \sim \nu_n$ and the right hand side can be interpreted as the KL divergence rate from the closest product chain to P . The rightmost expression in the equality above is also known as the mutual information rate [Chen et al. \(2018\)](#); [Holliday et al. \(2002\)](#).

Proof. We first prove item (1). We first consider the case where there exists $x, y \in \mathcal{X}$ such that $P(x, y) > 0$ yet $(\otimes_{i=1}^d L_i)(x, y) = 0$. This implies that $P_\pi^{(i)}(x^i, y^i) > 0$ for all i . Thus, both $D_{KL}^\pi(P \parallel \otimes_{i=1}^d L_i) = D_{KL}^\pi(P_\pi^{(i)} \parallel L_i) = +\infty$ and the identity holds. Next, we consider the case with $P(x, \cdot) \ll (\otimes_{i=1}^d L^{(i)})(x, \cdot)$ for all x . We see that

$$\begin{aligned} D_{KL}^\pi(P \parallel \otimes_{i=1}^d L_i) &= D_{KL}^\pi(P \parallel \otimes_{i=1}^d P_\pi^{(i)}) + \sum_{x,y} \pi(x) P(x, y) \ln \left(\frac{(\otimes_{i=1}^d P_\pi^{(i)})(x, y)}{(\otimes_{i=1}^d L_i)(x, y)} \right) \\ &= D_{KL}^\pi(P \parallel \otimes_{i=1}^d P_\pi^{(i)}) + \sum_{i=1}^d \sum_{x^i, y^i} \pi^{(i)}(x^i) \sum_{x^j, y^j; j \neq i} \frac{\pi(x) P(x, y)}{\pi^{(i)}(x^i)} \ln \left(\frac{P_\pi^{(i)}(x^i, y^i)}{L_i(x^i, y^i)} \right) \\ &= D_{KL}^\pi(P \parallel \otimes_{i=1}^d P_\pi^{(i)}) + \sum_{i=1}^d \sum_{x^i, y^i} \pi^{(i)}(x^i) P_\pi^{(i)}(x^i, y^i) \ln \left(\frac{P_\pi^{(i)}(x^i, y^i)}{L_i(x^i, y^i)} \right) \\ &= D_{KL}^\pi(P \parallel \otimes_{i=1}^d P_\pi^{(i)}) + \sum_{i=1}^d D_{KL}^\pi(P_\pi^{(i)} \parallel L_i). \end{aligned}$$

In view of [Proposition 2.1](#),

$$D_{KL}^\pi(P \parallel \otimes_{i=1}^d L_i) \geq D_{KL}^\pi(P \parallel \otimes_{i=1}^d P_\pi^{(i)})$$

and equality holds if and only if $D_{KL}^\pi(P_\pi^{(i)} \parallel L_i) = 0$ for all i if and only if $P_\pi^{(i)} = L_i$ for all i .

Next, we prove item (2). First we see that $P_\pi^{(i)}$ is $\pi^{(i)}$ -stationary with $P_\pi^{(i)*} = P_\pi^{*(i)}$. Thus, by item (1) and [Proposition 2.1](#) item (5), we have

$$\mathbb{I}^\pi(P) = D_{KL}^\pi(P \parallel \otimes_{i=1}^d P_\pi^{(i)}) = D_{KL}^\pi(P^* \parallel \otimes_{i=1}^d P_\pi^{(i)*}) = D_{KL}^\pi(P^* \parallel \otimes_{i=1}^d P_\pi^{*(i)}) = \mathbb{I}^\pi(P^*).$$

□

One possible application of the Pythagorean identity lies in assessing the convergence to equilibrium of P . This is in part motivated by [\(Hsu et al., 2019, Section 10\)](#) which suggests looking into ‘‘Markov chains with factored transition kernels with a few factors’’. Suppose that P is ergodic (i.e. irreducible and aperiodic) with a product form stationary distribution $\pi = \otimes_{i=1}^d \pi^{(i)}$. Let $\Pi \in \mathcal{L}(\mathcal{X})$ be the matrix where each row is π , and $\Pi^{(i)} \in \mathcal{L}(\mathcal{X}^{(i)})$ be a matrix where each row is $\pi^{(i)}$ for all i . We thus see that

$$D_{KL}^\pi(P^n \parallel \Pi) \leq \max_{x \in \mathcal{X}} \tilde{D}_{KL}(P^n(x, \cdot) \parallel \pi).$$

On the other hand, we can lower bound $D_{KL}^\pi(P^n \|\Pi)$ via the KL divergence from $\Pi^{(i)}$ to $P_\pi^{n(i)}$ using the Pythagorean identity in Theorem 2.1:

$$D_{KL}^\pi(P^n \|\Pi) \geq \max_{i \in \llbracket d \rrbracket} D_{KL}^{\pi^{(i)}}(P_\pi^{n(i)} \|\Pi^{(i)}).$$

As a result this yields, for $\varepsilon > 0$,

$$t_{mix}(\varepsilon) \geq \max_{i \in \llbracket d \rrbracket} t_{mix}^{(i)}(\varepsilon),$$

where $t_{mix}(\varepsilon) := \inf\{n \in \mathbb{N}; \max_{x \in \mathcal{X}} \tilde{D}_{KL}(P^n(x, \cdot) \|\pi) < \varepsilon\}$ is the worst-case KL divergence mixing time of P and $t_{mix}^{(i)}(\varepsilon) := \inf\{n \in \mathbb{N}; D_{KL}^{\pi^{(i)}}(P_\pi^{n(i)} \|\Pi^{(i)}) < \varepsilon\}$ is an average-case KL divergence mixing time of the i th marginal transition matrix of P^n , namely $P_\pi^{n(i)}$. The interpretation is that the first time for P to be ε close to π in the sense of KL divergence is at least larger than the average-case marginal transition matrix KL divergence mixing time. In Section 2.6, we shall compare ergodicity constants, such as the spectral gap and log-Sobolev constant, between P and its information projections.

2.1.1 The closest product chain with prescribed marginals and a large deviation principle of Markov chains

Fix $i \in \llbracket d \rrbracket$ and suppose we are prescribed with $L_j \in \mathcal{L}(\mathcal{X}^{(j)})$ for all $j \in \llbracket d \rrbracket$, $j \neq i$ and a transition matrix $P \in \mathcal{L}(\mathcal{X})$. We consider the problem of finding the closest product chain of the form $(\otimes_{j=1}^{i-1} L_j) \otimes L \otimes (\otimes_{j=i+1}^d L_j)$. In other words, we are interested in seeking a minimizer of

$$L_*^{(i)} = L_*^{(i)}(P, L_1, \dots, L_{i-1}, L_{i+1}, \dots, L_d, f, \pi) \in \arg \min_{L \in \mathcal{L}(\mathcal{X}^{(i)})} D_f^\pi(P \| (\otimes_{j=1}^{i-1} L_j) \otimes L \otimes (\otimes_{j=i+1}^d L_j)).$$

In view of the previous section, in the special case where $L_j = P_\pi^{(j)}$ for all $j \neq i$, the j th marginal transition matrix of P as introduced in Definition 2.3, it seems natural to guess that $L_*^{(i)}$ is $P_\pi^{(i)}$. Our next result shows that, depending on the choice of f , $L_*^{(i)}$ can in fact be weighted averages of L_j and P in a broad sense. Therefore, the seemingly natural product chain with transition matrix $\otimes_{i=1}^d P_\pi^{(i)}$ is not necessarily the closest product chain under some information divergences.

Theorem 2.2. *Fix $i \in \llbracket d \rrbracket$ and suppose we are prescribed with $L_j \in \mathcal{L}(\mathcal{X}^{(j)})$ for all $j \in \llbracket d \rrbracket$, $j \neq i$ and a transition matrix $P \in \mathcal{L}(\mathcal{X})$. Let π be a positive probability mass.*

1. (Reverse KL divergence) Let $f(t) = -\ln t$ that generates the reverse KL divergence. The unique $L_*^{(i)}$ is given by, for $x^i, y^i \in \mathcal{X}^{(i)}$,

$$L_*^{(i)}(x^i, y^i) \propto \prod_{x^{(-i)}, y^{(-i)}} P(x, y) \frac{\pi(x) \mathbf{z}(x^{(-i)}, y^{(-i)})}{Z(x^i, y^i)}, \quad (8)$$

where $x^{(-i)} := (x^1, \dots, x^{i-1}, x^{i+1}, \dots, x^d)$, $\mathbf{Z}(x^{(-i)}, y^{(-i)}) := \prod_{j=1; j \neq i}^n L_j(x^j, y^j)$ and $Z(x^i, y^i) = \sum_{x^{(-i)}, y^{(-i)}} \pi(x) \mathbf{Z}(x^{(-i)}, y^{(-i)})$.

2. (α -divergence) Let $f(t) = \frac{1}{\alpha-1}(t^\alpha - 1)$ that generates the α -divergence with $\alpha \in (0, 1) \cup (1, \infty)$. The unique $L_*^{(i)}$ is given by, for $x^i, y^i \in \mathcal{X}^{(i)}$,

$$L_*^{(i)}(x^i, y^i) \propto \left(\sum_{x^{(-i)}, y^{(-i)}} \pi(x) \left(\prod_{j=1; j \neq i}^d L_j(x^j, y^j) \right)^{1-\alpha} P(x, y)^\alpha \right)^{1/\alpha}.$$

3. (KL divergence) Let $f(t) = t \ln t$ that generates the KL divergence. The unique $L_*^{(i)}$ is given by

$$L_*^{(i)} = P_\pi^{(i)}.$$

Note that $L_*^{(i)}$ depends on P and π and does not depend on L_j with $j \neq i$.

Proof. We first prove item (1). We write down

$$\begin{aligned} D_f^\pi(P \| (\otimes_{j=1}^{i-1} L_j) \otimes L \otimes (\otimes_{j=i+1}^d L_j)) \\ = \sum_{x, y} \pi(x) L(x^i, y^i) \prod_{j=1; j \neq i}^d L_j(x^j, y^j) \ln \left(\frac{L(x^i, y^i) \prod_{j=1; j \neq i}^d L_j(x^j, y^j)}{P(x, y)} \right). \end{aligned}$$

Differentiating the above with respect to $L(x^i, y^i)$ and noting that z^i is chosen such that $L(x^i, z^i) = 1 - \sum_{y^i \in \mathcal{X}^i; y^i \neq z^i} L(x^i, y^i)$, we set the derivative to be zero to give

$$\sum_{x^{(-i)}, y^{(-i)}} \pi(x) \prod_{j=1; j \neq i}^d L_j(x^j, y^j) \ln \left(\frac{L(x^i, y^i) P(x, (y^1, \dots, z^i, \dots, y^d))}{L(x^i, z^i) P(x, y)} \right) = 0.$$

Using $\mathbf{Z}(x^{(-i)}, y^{(-i)}) = \prod_{j=1; j \neq i}^d L_j(x^j, y^j)$ and $Z(x^i, y^i) = \sum_{x^{(-i)}, y^{(-i)}} \pi(x) \mathbf{Z}(x^{(-i)}, y^{(-i)})$, we then see that

$$L_*^{(i)}(x^i, y^i) \propto \prod_{x^{(-i)}, y^{(-i)}} P(x, y)^{\frac{\pi(x) \mathbf{Z}(x^{(-i)}, y^{(-i)})}{Z(x^i, y^i)}}.$$

Next, we prove item (2). In this case we have

$$\begin{aligned} D_f^\pi(P \| (\otimes_{j=1}^{i-1} L_j) \otimes L \otimes (\otimes_{j=i+1}^d L_j)) \\ = \frac{1}{\alpha-1} \sum_{x, y} \pi(x) \left(L(x^i, y^i) \prod_{j=1; j \neq i}^d L_j(x^j, y^j) \right)^{1-\alpha} P(x, y)^\alpha - 1. \end{aligned}$$

We then differentiate the above with respect to $L(x^i, y^i)$ and note that z^i satisfies $L(x^i, z^i) = 1 - \sum_{y^i \in \mathcal{X}^i; y^i \neq z^i} L(x^i, y^i)$. Setting the derivative to be zero leads to

$$L_*^{(i)}(x^i, y^i) \propto \left(\sum_{x^{(-i)}, y^{(-i)}} \pi(x) \left(\prod_{j=1; j \neq i}^d L_j(x^j, y^j) \right)^{1-\alpha} P(x, y)^\alpha \right)^{1/\alpha}.$$

Finally, we prove item (3). We first note that

$$\begin{aligned} D_f^\pi(P \| (\otimes_{j=1}^{i-1} L_j) \otimes L \otimes (\otimes_{j=i+1}^d L_j)) \\ = \sum_{x, y} \pi(x) P(x, y) \ln \left(\frac{P(x, y)}{L(x^i, y^i) \prod_{j=1; j \neq i}^d L_j(x^j, y^j)} \right). \end{aligned}$$

Differentiating the above with respect to $L(x^i, y^i)$ and noting that z^i is chosen such that $L(x^i, z^i) = 1 - \sum_{y^i \in \mathcal{X}^i; y^i \neq z^i} L(x^i, y^i)$, we set the derivative to be zero to give

$$L_*^{(i)}(x^i, y^i) \propto \sum_{x^{(-i)}, y^{(-i)}} \pi(x) P(x, y),$$

that is, $L_*^{(i)} = P_\pi^{(i)}$. □

One application of Theorem 2.2 lies in the large deviation analysis and Sanov's theorem of Markov chains, in which we apply the results obtained in [Fayolle and de La Fortelle \(2002\)](#). We refer readers to [Dembo and Zeitouni \(2010\)](#); [Natarajan \(1985\)](#); [Vidyasagar \(2014\)](#) for related literature on large deviations of Markov chains.

Let $X = (X_n)_{n \in \mathbb{N}_0}$ be the Markov chain with transition matrix P . Define the pair empirical measure of X to be

$$E_n := \frac{1}{n} \left(\sum_{i=1}^{n-1} \delta_{(X_i, X_{i+1})} + \delta_{(X_n, X_1)} \right). \quad (9)$$

Theorem 2.3 (A Sanov's theorem of Markov chains). *Fix $i \in \llbracket d \rrbracket$. Let $\pi = \otimes_{l=1}^d \pi^{(l)}$. Suppose we are prescribed with $\pi^{(j)}$ -stationary $L_j \in \mathcal{L}(\mathcal{X}^{(j)})$ for all $j \in \llbracket d \rrbracket$, $j \neq i$ and a π -stationary $P \in \mathcal{L}(\mathcal{X})$. Let K be the set*

$$\begin{aligned} K_i &= K_i(L_1, \dots, L_{i-1}, L_{i+1}, \dots, L_d) \\ &:= \{(\otimes_{j=1}^{i-1} L_j) \otimes M \otimes (\otimes_{j=i+1}^d L_j); \pi^{(i)}\text{-stationary } M \in \mathcal{L}(\mathcal{X}^{(i)})\}, \end{aligned}$$

and $f(t) = -\ln t$ that generates the reverse KL divergence. We have

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P}(E_n \in K_i) \leq -D_f^\pi(P \| (\otimes_{j=1}^{i-1} L_j) \otimes L_*^{(i)} \otimes (\otimes_{j=i+1}^d L_j)),$$

where we recall that E_n is the pair empirical measure as introduced in (9) and $L_*^{(i)}$ is given in (8). Note that the above result holds without any restriction on the initial distribution of the chain X .

Proof. The plan is to invoke Theorem 1.1 in [Fayolle and de La Fortelle \(2002\)](#). Let us first assume that K is a subset of the set of balanced measures. Then by [Fayolle and de La Fortelle \(2002\)](#) and Theorem 2.2, these yield

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \ln \mathbb{P}(E_n \in K_i) \leq - \inf_{L \in K} D_f^\pi(P \| L) = -D_f^\pi(P \| (\otimes_{j=1}^{i-1} L_j) \otimes L_*^{(i)} \otimes (\otimes_{j=i+1}^d L_j)).$$

It remains to verify that K_i is a subset of the set of balanced measures, in which we readily see that

$$(\pi \boxtimes (\otimes_{j=1}^{i-1} L_j) \otimes M \otimes (\otimes_{j=i+1}^d L_j))(\mathcal{X}, \cdot) = \pi(\cdot) = (\pi \boxtimes (\otimes_{j=1}^{i-1} L_j) \otimes M \otimes (\otimes_{j=i+1}^d L_j))(\cdot, \mathcal{X}).$$

□

2.1.2 A coordinate descent algorithm for finding the closest product chain

Let us recall that in Theorem 2.1, we have shown $\otimes_{i=1}^d P_\pi^{(i)}$ is the unique closest product chain to a given P . In other choices of f -divergences such as the reverse KL divergence or the α -divergence, we did not manage to derive a closed form formula for the closest product chain. In these cases, if we have the closed form of the closest product chain with prescribed marginals as in Theorem 2.2, we can derive a coordinate descent algorithm to find approximately the closest product chain.

Suppose that the algorithm is initiated with $L_i^0 \in \mathcal{L}(\mathcal{X}^{(i)})$ for $i \in \llbracket d \rrbracket$. At iteration $l \in \mathbb{N}$ and for each $i \in \llbracket d \rrbracket$, we compute that

$$L_i^l = L_i^{(i)}(P, L_1^l, \dots, L_{i-1}^l, L_{i+1}^{l-1}, \dots, L_d^{l-1}, f, \pi) \in \arg \min_{L \in \mathcal{L}(\mathcal{X}^{(i)})} D_f^\pi(P \| (\otimes_{j=1}^{i-1} L_j^l) \otimes L \otimes (\otimes_{j=i+1}^d L_j^{l-1})).$$

In the case of reverse KL divergence or α -divergence, these are computed in Theorem 2.2. The sequence $(\otimes_{i=1}^d L_i^l)_{l \in \mathbb{N}}$ satisfies

$$D_f^\pi(P \| \otimes_{i=1}^d L_i^l) \geq D_f^\pi(P \| \otimes_{i=1}^d L_i^{l+1})$$

for $l \geq 0$.

2.2 Leave-one-out transition matrices and the Han's inequality for KL divergence of Markov chains

Fix $i \in \llbracket d \rrbracket$, and consider a product state space $\mathcal{X} = \times_{j=1}^d \mathcal{X}^{(j)}$. We define the leave- i -out state space to be $\mathcal{X}^{(-i)} = \times_{j=1; j \neq i}^d \mathcal{X}^{(j)}$. Let $x^j \in \mathcal{X}^{(j)}$ for all $j \in \llbracket d \rrbracket$, and recall that we write $x^{(-i)} = (x^1, \dots, x^{i-1}, x^{i+1}, \dots, x^d) \in \mathcal{X}^{(-i)}$ as in Theorem 2.2. Let $\mu, \otimes_{j=1}^d \nu_j \in \mathcal{P}(\mathcal{X})$. The leave- i -out distribution of μ is given by

$$\mu^{(-i)}(x^{(-i)}) := \sum_{x^i} \mu(x^1, \dots, x^d).$$

In particular, this yields

$$(\otimes_{j=1}^d \nu_j)^{(-i)} = \otimes_{j=1; j \neq i}^d \nu_j.$$

We remark that these leave-one-out distributions are widely used in forming the jackknife estimator in mathematical statistics and cross-validation in the training of machine learning algorithms.

The Han's inequality for KL divergence between discrete probability masses (Boucheron et al., 2013, Theorem 4.9) is given by

$$\tilde{D}_{KL}(\mu \| \otimes_{j=1}^d \nu_j) \leq \sum_{i=1}^d \tilde{D}_{KL}(\mu \| \otimes_{j=1}^d \nu_j) - \tilde{D}_{KL}(\mu^{(-i)} \| \otimes_{j=1; j \neq i}^d \nu_j). \quad (10)$$

In this subsection, we aim at understanding generalization of the Han's inequality to KL divergence between transition matrices of Markov chains. Let us now define the leave- i -out transition matrix of a given P by suitably marginalizing the i th coordinate:

Definition 2.4 ($P_\pi^{(-i)}$): the leave- i -out transition matrix of P with respect to π . Let $\pi \in \mathcal{P}(\mathcal{X})$ be a positive probability mass, $P \in \mathcal{L}(\mathcal{X})$ and $i \in \llbracket d \rrbracket$. For any $(x^{(-i)}, y^{(-i)}) \in \mathcal{X}^{(-i)} \times \mathcal{X}^{(-i)}$, we define

$$\begin{aligned} P_\pi^{(-i)}(x^{(-i)}, y^{(-i)}) &:= \frac{\sum_{(x^i, y^i)} \pi(x^1, \dots, x^d) P((x^1, \dots, x^d), (y^1, \dots, y^d))}{\sum_{x^i} \pi(x^1, \dots, x^d)} \\ &= \frac{(\pi \boxtimes P)^{(-i)}(x^{(-i)}, y^{(-i)})}{\pi^{(-i)}(x^{(-i)})}. \end{aligned}$$

Note that $P_\pi^{(-i)} \in \mathcal{L}(\mathcal{X}^{(-i)})$. When P is π -stationary, we omit the subscript and write $P^{(-i)}$.

Note that if P is π -stationary, then $P_\pi^{(-i)}$ is $\pi^{(-i)}$ -stationary, and hence $P_\pi^{(i)} \otimes P_\pi^{(-i)}$ is $\pi^{(i)} \otimes \pi^{(-i)}$ -stationary. Thus, in general $P_\pi^{(S)} \otimes P_\pi^{(-S)}$ is not π -stationary. In the case where $\pi = \otimes_{i=1}^d \pi^{(i)}$ is a product stationary distribution, then $P_\pi^{(i)} \otimes P_\pi^{(-i)}$ is π -stationary. If P is π -reversible, then $P_\pi^{(-i)}$ is $\pi^{(-i)}$ -reversible. A generalization of these results and their consequences can be found in Proposition 2.3 below.

We now state the Han's inequality for KL divergence of Markov chains:

Theorem 2.4 (Han's inequality for KL divergence of Markov chains). Let $\pi = \otimes_{j=1}^d \pi^{(j)} \in \mathcal{P}(\mathcal{X})$ be a positive product distribution, $P \in \mathcal{L}(\mathcal{X})$ and $L_j \in \mathcal{L}(\mathcal{X}^{(j)})$ for $j \in \llbracket d \rrbracket$. For $d \geq 2$, we have

$$D_{KL}^\pi(P \| \otimes_{j=1}^d L_j) \leq \sum_{i=1}^d D_{KL}^\pi(P \| \otimes_{j=1}^d L_j) - D_{KL}^{\pi^{(-i)}}(P^{(-i)} \| \otimes_{j=1; j \neq i}^d L_j).$$

Proof. Using the Han's inequality for KL divergence of probability masses (10), we see that

$$D_{KL}^\pi(P \| \otimes_{j=1}^d L_j) = \tilde{D}_{KL}(\pi \boxtimes P \| \otimes_{j=1}^d (\pi^{(j)} \boxtimes L_j))$$

$$\begin{aligned}
&\leq \sum_{i=1}^d \tilde{D}_{KL}(\pi \boxtimes P \| \otimes_{j=1}^d (\pi^{(j)} \boxtimes L_j)) - \tilde{D}_{KL}((\pi \boxtimes P)^{(-i)} \| \otimes_{j=1; j \neq i}^d (\pi^{(j)} \boxtimes L_j)) \\
&= \sum_{i=1}^d D_{KL}^\pi(P \| \otimes_{j=1}^d L_j) - D_{KL}^{\pi^{(-i)}}(P_\pi^{(-i)} \| \otimes_{j=1; j \neq i}^d L_j).
\end{aligned}$$

□

For $j \neq i$ and $d \geq 3$, if we consider the j th marginal transition matrix of $P^{(-i)}$ with respect to π , we compute that to be

$$(P_\pi^{(-i)})_{\pi^{(-i)}}^{(j)} = P_\pi^{(j)}. \quad (11)$$

Thanks to Theorem 2.1, this yields

$$\begin{aligned}
\mathbb{I}^\pi(P) &= D_{KL}^\pi(P \| \otimes_{j=1}^d P_\pi^{(j)}), \\
\mathbb{I}^{\pi^{(-i)}}(P_\pi^{(-i)}) &= D_{KL}^{\pi^{(-i)}}(P_\pi^{(-i)} \| \otimes_{j=1; j \neq i}^d (P_\pi^{(-i)})_{\pi^{(-i)}}^{(j)}) = D_{KL}^{\pi^{(-i)}}(P_\pi^{(-i)} \| \otimes_{j=1; j \neq i}^d P_\pi^{(j)}).
\end{aligned}$$

Applying these two identities to Theorem 2.4 by taking $L_j = P_\pi^{(j)}$ therein leads to

Corollary 2.1 (Han's inequality for distance to independence of P with respect to π). *Let $\pi = \otimes_{j=1}^d \pi^{(j)} \in \mathcal{P}(\mathcal{X})$ be a positive product distribution and $P \in \mathcal{L}(\mathcal{X})$. For $d \geq 2$, we have*

$$\mathbb{I}^\pi(P) \leq \sum_{i=1}^d \mathbb{I}^\pi(P) - \mathbb{I}^{\pi^{(-i)}}(P_\pi^{(-i)}),$$

or equivalently,

$$\mathbb{I}^\pi(P) \geq \frac{1}{d-1} \sum_{i=1}^d \mathbb{I}^{\pi^{(-i)}}(P_\pi^{(-i)}).$$

Remark 2.5. *Corollary 2.1 can be recursively applied to obtain further lower bounds, say for $d \geq 3$, we can apply the Han's inequality to $\mathbb{I}^{\pi^{(-i)}}(P_\pi^{(-i)})$ and so on.*

Note that the case of $d = 2$ trivially holds since $\mathbb{I}^{\pi^{(-1)}}(P_\pi^{(-1)}) = \mathbb{I}^{\pi^{(-2)}}(P_\pi^{(-2)}) = 0$ while $\mathbb{I}^\pi(P) \geq 0$ by the non-negativity of KL divergence. Also, when P is itself a product chain, equality holds obviously since the distance to independence of P or $P_\pi^{(-i)}$ are all zeros. Intuitively, we can understand that the distance to independence of P with respect to π is at least greater than the ‘‘average’’ distances to independence of the leave-one-out transition matrices.

We now consider the information projection of P onto the space $\{L_i \otimes M; L_i \in \mathcal{L}(\mathcal{X}^{(i)}), M \in \mathcal{L}(\mathcal{X}^{(-i)})\}$. That is, it is the space of transition matrices on \mathcal{X} where the i th coordinate is independent while the remaining coordinates form a block. We develop a Pythagorean identity in this case:

Theorem 2.5. (*Pythagorean identity*) Let $\pi \in \mathcal{P}(\mathcal{X})$ be a positive probability mass, $P \in \mathcal{L}(\mathcal{X})$, $L^{(i)} \in \mathcal{L}(\mathcal{X}^{(i)})$ and $M \in \mathcal{L}(\mathcal{X}^{(-i)})$ for $i \in \llbracket d \rrbracket$. We then have

$$\begin{aligned} D_{KL}^\pi(P \| L_i \otimes M) &= D_{KL}^\pi(P \| P_\pi^{(i)} \otimes P_\pi^{(-i)}) + D_{KL}^\pi(P_\pi^{(i)} \otimes P_\pi^{(-i)} \| L_i \otimes M) \\ &= D_{KL}^\pi(P \| P_\pi^{(i)} \otimes P_\pi^{(-i)}) + D_{KL}^{\pi^{(i)}}(P_\pi^{(i)} \| L_i) + D_{KL}^{\pi^{(-i)}}(P_\pi^{(-i)} \| M), \end{aligned}$$

where $D_{KL}^{\pi^{(i)}}(P_\pi^{(i)} \| L_i)$ is weighted by $\pi^{(i)}$, the i th marginal distribution of π . In other words, $P_\pi^{(i)} \otimes P_\pi^{(-i)}$, the tensor product of the i th marginal transition matrix and the leave- i -out transition matrix, is the unique minimizer of

$$\min_{L_i \in \mathcal{L}(\mathcal{X}^{(i)}), M \in \mathcal{L}(\mathcal{X}^{(-i)})} D_{KL}^\pi(P \| L_i \otimes M) = D_{KL}^\pi(P \| P_\pi^{(i)} \otimes P_\pi^{(-i)}).$$

Proof.

$$\begin{aligned} D_{KL}^\pi(P \| L_i \otimes M) &= D_{KL}^\pi(P \| P_\pi^{(i)} \otimes P_\pi^{(-i)}) + \sum_{x,y} \pi(x) P(x,y) \ln \left(\frac{(P_\pi^{(i)} \otimes P_\pi^{(-i)})(x,y)}{(L_i \otimes M)(x,y)} \right) \\ &= D_{KL}^\pi(P \| P_\pi^{(i)} \otimes P_\pi^{(-i)}) + \sum_{x^i, y^i} \pi^{(i)}(x^i) P_\pi^{(i)}(x^i, y^i) \ln \left(\frac{P_\pi^{(i)}(x^i, y^i)}{L_i(x^i, y^i)} \right) \\ &\quad + \sum_{x^{(-i)}, y^{(-i)}} \pi^{(-i)}(x^{(-i)}) P_\pi^{(-i)}(x^{(-i)}, y^{(-i)}) \ln \left(\frac{P_\pi^{(-i)}(x^{(-i)}, y^{(-i)})}{M(x^{(-i)}, y^{(-i)})} \right) \\ &= D_{KL}^\pi(P \| P_\pi^{(i)} \otimes P_\pi^{(-i)}) + D_{KL}^{\pi^{(i)}}(P_\pi^{(i)} \| L_i) + D_{KL}^{\pi^{(-i)}}(P_\pi^{(-i)} \| M). \end{aligned}$$

□

2.3 Leave- S -out transition matrices and the Shearer's lemma for Markov chains

It should hopefully be apparent to the readers that the results in Section 2.2 can be generalized to consider transition matrices where we marginalize $k \in \llbracket d-1 \rrbracket$ coordinates of a given P . In the special case of $k = 1$, we obtain the leave-one-out transition matrices. In the other special case of $k = d-1$, we recover $P_\pi^{(i)}$ for $i \in \llbracket d \rrbracket$, the i th marginal transition matrix. Generally, we can marginalize a subset $S \subseteq \llbracket d \rrbracket$ of the d coordinates.

Fix $S \subseteq \llbracket d \rrbracket$ and consider a product state space $\mathcal{X} = \times_{j=1}^d \mathcal{X}^{(j)}$. The leave- S -out state space is defined to be $\mathcal{X}^{(-S)} := \times_{j=1; j \notin S}^n \mathcal{X}^{(j)}$, while the keep- S -in state space is defined to be $\mathcal{X}^{(S)} := \times_{j=1; j \in S}^n \mathcal{X}^{(j)}$. Let $x^j \in \mathcal{X}^{(j)}$ for all $j \in \llbracket d \rrbracket$, and we write $x^{(-S)} := (x^j)_{j \notin S} \in \mathcal{X}^{(-S)}$, $x^{(S)} := (x^j)_{j \in S} \in \mathcal{X}^{(S)}$. Let $\mu, \otimes_{j=1}^d \nu_j \in \mathcal{P}(\mathcal{X})$. The leave- S -out distribution of μ is given by

$$\mu^{(-S)}(x^{(-S)}) := \sum_{i \in S} \sum_{x^i} \mu(x^1, \dots, x^d).$$

In particular, this yields

$$(\otimes_{j=1}^d \nu_j)^{(-S)} = \otimes_{j=1; j \notin S}^d \nu_j.$$

The keep- S -in distribution of μ is the leave- $\llbracket d \rrbracket \setminus S$ -out distribution of μ , that is,

$$\mu^{(S)}(x^{(S)}) := \mu^{(-\llbracket d \rrbracket \setminus S)}(x^{(-\llbracket d \rrbracket \setminus S)}) = \sum_{i \notin S} \sum_{x^i} \mu(x^1, \dots, x^d).$$

This leads to

$$(\otimes_{j=1}^d \nu_j)^{(S)} = \otimes_{j=1; j \in S}^d \nu_j.$$

To demonstrate the consistency of the notations, we note that, for $j \in \llbracket d \rrbracket$,

$$\begin{aligned} \mathcal{X}^{(-j)} &= \mathcal{X}^{(-\{j\})}, & \mathcal{X}^{(j)} &= \mathcal{X}^{\{j\}}, \\ \mu^{(-j)} &= \mu^{(-\{j\})}, & \mu^{(j)} &= \mu^{\{j\}}. \end{aligned}$$

Given a sequence $(S_i)_{i=1}^n$ with $S_i \subseteq \llbracket d \rrbracket$, where each $j \in \llbracket d \rrbracket$ belongs to at least r of S_i . The Shearer's lemma of entropy (Polyanskiy and Wu, 2022, Theorem 1.8) is given by

$$H(X_1, \dots, X_d) \leq \frac{1}{r} \sum_{i=1}^n H((X_l)_{l \in S_i}), \quad (12)$$

where $(X_l)_{l \in S_i} \sim \mu^{(S_i)}$ and $H((X_l)_{l \in S_i}) := -\sum_{x^{(S_i)}} \mu^{(S_i)}(x^{(S_i)}) \ln \mu^{(S_i)}(x^{(S_i)})$ is the entropy of $\mu^{(S_i)}$, while the KL divergence version of the Shearer's lemma (Gavinsky et al., 2015, Corollary 2.8) is stated as, for $\mu, \nu = \otimes_{j=1}^d \nu_j \in \mathcal{P}(\mathcal{X})$,

$$\tilde{D}_{KL}(\mu \parallel \nu) \geq \frac{1}{r} \sum_{i=1}^n \tilde{D}_{KL}(\mu^{(S_i)} \parallel \nu^{(S_i)}). \quad (13)$$

Next, we introduce the leave- S -out transition matrix:

Definition 2.5 ($P_\pi^{(-S)}$ and $P_\pi^{(S)}$): the leave- S -out and keep- S -in transition matrix of P with respect to π . Let $\pi \in \mathcal{P}(\mathcal{X})$ be a positive probability mass, $P \in \mathcal{L}(\mathcal{X})$ and $S \subseteq \llbracket d \rrbracket$. For any $(x^{(-S)}, y^{(-S)}) \in \mathcal{X}^{(-S)} \times \mathcal{X}^{(-S)}$, we define

$$\begin{aligned} P_\pi^{(-S)}(x^{(-S)}, y^{(-S)}) &:= \frac{\sum_{(x^i, y^i) \in \mathcal{X}^{(S)} \times \mathcal{X}^{(S)}} \pi(x^1, \dots, x^d) P((x^1, \dots, x^d), (y^1, \dots, y^d))}{\sum_{x^i \in \mathcal{X}^{(S)}} \pi(x^1, \dots, x^d)} \\ &= \frac{(\pi \boxtimes P)^{(-S)}(x^{(-S)}, y^{(-S)})}{\pi^{(-S)}(x^{(-S)})}. \end{aligned}$$

Note that $P_\pi^{(-S)} \in \mathcal{L}(\mathcal{X}^{(-S)})$. The keep- S -in transition matrix of P with respect to π is

$$P_\pi^{(S)} := P_\pi^{(-\llbracket d \rrbracket \setminus S)} \in \mathcal{L}(\mathcal{X}^{(S)}).$$

In the special case of $S = \{i\}$ for $i \in \llbracket d \rrbracket$, we write

$$P_\pi^{(-i)} = P_\pi^{(-\{i\})}, \quad P_\pi^{(i)} = P_\pi^{\{i\}}.$$

When P is π -stationary, we omit the subscript and write $P^{(-S)}, P^{(S)}$.

We note that when P is π -stationary, the keep- S -in transition matrix $P^{(S)}$ can be viewed as a special case of the “projection chain” of P in [Jerrum et al. \(2004\)](#), or as an “induced chain” of P in ([Aldous and Fill, 2002](#), Section 4.6). The latter is also known as a “collapsed chain” in [Chen et al. \(1999\)](#), which can be understood as the opposite of the lifting procedure in MCMC.

Let $\Omega_{x^{(S)}} := \{x^{(S)}\} \times \llbracket d \rrbracket \setminus S$. We then see that $(\Omega_{x^{(S)}})_{x^{(S)} \in \mathcal{X}^{(S)}}$ is a partition of the state space, that is, $\mathcal{X} = \times_{i=1}^d \mathcal{X}^{(i)} = \cup_{x^{(S)} \in \mathcal{X}^{(S)}} \Omega_{x^{(S)}}$. Let \bar{P} be the “projection chain” of P with respect to $(\Omega_{x^{(S)}})_{x^{(S)} \in \mathcal{X}^{(S)}}$ in the sense of [Jerrum et al. \(2004\)](#), which is defined to be

$$\bar{P}(x^{(S)}, y^{(S)}) := \frac{\sum_{x \in \Omega_{x^{(S)}}, y \in \Omega_{y^{(S)}}} \pi(x) P(x, y)}{\sum_{x \in \Omega_{x^{(S)}}} \pi(x)}.$$

This yields $\bar{P} = P^{(S)}$.

Recall that (X_n) is the Markov chain with transition matrix P on \mathcal{X} . Let $\phi : \mathcal{X} \rightarrow \mathcal{X}^{(S)}$ be the projection map onto $\mathcal{X}^{(S)}$ defined to be $\phi(x) := x^{(S)}$. Note that $(\phi(X_n))$ is not a Markov chain in general and is a Markov chain in the lumpable case, see [Kemeny and Snell \(1983\)](#); [Wolfer and Watanabe \(2024+\)](#).

We summarize the ergodicity properties of $P^{(S)}$ appeared in [Aldous and Fill \(2002\)](#); [Jerrum et al. \(2004\)](#). We can understand that these properties are inherited from the counterpart properties of the original P :

Proposition 2.3. *Let $P \in \mathcal{L}(\mathcal{X})$, $S \subseteq \llbracket d \rrbracket$ and $\pi \in \mathcal{P}(\mathcal{X})$ be a positive probability mass. We have*

1. *If P is π -stationary, then $P_\pi^{(S)}$ is $\pi^{(S)}$ -stationary.*
2. *If P is π -reversible, then $P_\pi^{(S)}$ is $\pi^{(S)}$ -reversible.*
3. *If P is π -stationary, then $P_\pi^{(S)} \otimes P_\pi^{(-S)}$ is $\pi^{(S)} \otimes \pi^{(-S)}$ -stationary. In particular, if $\pi = \otimes_{i=1}^d \pi^{(i)}$ is a product stationary distribution, then $P_\pi^{(S)} \otimes P_\pi^{(-S)}$ is π -stationary.*
4. *If P is lazy, that is, $P(x, x) \geq 1/2$ for all $x \in \mathcal{X}$, then $P_\pi^{(S)}$ is lazy.*
5. *If P is ergodic (i.e. irreducible and aperiodic), then $P_\pi^{(S)}$ is ergodic.*

In the context of MCMC, there are often situations in which we are only interested in sampling from a subset S out of the d coordinates of the stationary distribution π of a sampler. Thus, $P_\pi^{(S)}$ offers a natural projection sampler to approximately sample from $\pi^{(S)}$. As a concrete example, we can consider the run-and-tumble Markov chains [van Ginkel et al. \(2021\)](#) where the algorithm maintains both the positions and directions of d particles. In such setting, we are often interested in sampling from the stationary distribution of the positions of the particles only and discard the samples from the directions. Another concrete example would be the swapping algorithm that we shall discuss in [Section 3.1.3](#), where one maintains

a system of Markov chains over a range of temperatures. The swapping algorithm is designed to sample from the Boltzmann-Gibbs distribution at the lowest temperature of the algorithm and samples that correspond to higher temperatures are often discarded. A third example is the auxiliary MCMC methods [Higdon \(1998\)](#), where one artificially add auxiliary variables in a MCMC algorithm to improve convergence, and at the end of simulation the samples of these auxiliary variables are discarded. In these algorithms, it is therefore natural to consider $P_\pi^{(S)}$ as a candidate sampler for $\pi^{(S)}$. We shall compare hitting and mixing time parameters between P and $P_\pi^{(S)}$ in [Section 2.6](#).

We now state the Shearer's lemma for KL divergence of Markov chains:

Theorem 2.6 (Shearer's lemma for KL divergence of Markov chains). *Let $\pi = \otimes_{j=1}^d \pi^{(j)} \in \mathcal{P}(\mathcal{X})$ be a positive product distribution, $P, L = \otimes_{j=1}^d L_j \in \mathcal{L}(\mathcal{X})$ and $L_j \in \mathcal{L}(\mathcal{X}^{(j)})$ for $j \in \llbracket d \rrbracket$. Given a sequence $(S_i)_{i=1}^n$ with $S_i \subseteq \llbracket d \rrbracket$, where each $j \in \llbracket d \rrbracket$ belongs to at least r of S_i . We have*

$$D_{KL}^\pi(P \| L) \geq \frac{1}{r} \sum_{i=1}^n D_{KL}^{\pi^{(S_i)}}(P_\pi^{(S_i)} \| L^{(S_i)}).$$

Note that $L^{(S_i)} = \otimes_{j \in S_i} L_j$. In particular, if we take $S_i = \llbracket d \rrbracket \setminus \{i\}$ and $n = d$ so that $r = d-1$, we recover the Han's inequality in [Theorem 2.4](#).

Proof. Using the Shearer's lemma for KL divergence of probability masses in [\(13\)](#), we see that

$$\begin{aligned} D_{KL}^\pi(P \| \otimes_{j=1}^d L_j) &= \tilde{D}_{KL}(\pi \boxtimes P \| \otimes_{j=1}^d (\pi^{(j)} \boxtimes L_j)) \\ &\geq \frac{1}{r} \sum_{i=1}^n \tilde{D}_{KL}((\pi \boxtimes P)^{(S_i)} \| \otimes_{j \in S_i} (\pi^{(j)} \boxtimes L_j)) \\ &= \frac{1}{r} \sum_{i=1}^n D_{KL}^{\pi^{(S_i)}}(P_\pi^{(S_i)} \| \otimes_{j \in S_i} L_j). \end{aligned}$$

□

For $j \in S_i$, if we consider the j th marginal transition matrix of $P_\pi^{(S_i)}$ with respect to $\pi^{(S_i)}$, we compute that to be

$$(P_\pi^{(S_i)})_{\pi^{(S_i)}}^{(j)} = P_\pi^{(j)}. \tag{14}$$

Using again [Theorem 2.1](#) leads to

$$\begin{aligned} \mathbb{I}^\pi(P) &= D_{KL}^\pi(P \| \otimes_{j=1}^d P_\pi^{(j)}), \\ \mathbb{I}^{\pi^{(S_i)}}(P_\pi^{(S_i)}) &= D_{KL}^{\pi^{(S_i)}}(P_\pi^{(S_i)} \| \otimes_{j \in S_i} (P_\pi^{(S_i)})_{\pi^{(S_i)}}^{(j)}) = D_{KL}^{\pi^{(S_i)}}(P_\pi^{(S_i)} \| \otimes_{j \in S_i} P_\pi^{(j)}). \end{aligned}$$

Applying these two equalities to [Theorem 2.6](#) by taking $L_j = P_\pi^{(j)}$ therein leads to

Corollary 2.2 (Shearer's lemma for distance to independence of P with respect to π). *Let $\pi = \otimes_{j=1}^d \pi^{(j)} \in \mathcal{P}(\mathcal{X})$ be a positive product distribution, $P, L = \otimes_{j=1}^d L_j \in \mathcal{L}(\mathcal{X})$ and $L_j \in \mathcal{L}(\mathcal{X}^{(j)})$ for $j \in \llbracket d \rrbracket$. Let $(S_i)_{i=1}^n$ be a sequence with $S_i \subseteq \llbracket d \rrbracket$, where each $j \in \llbracket d \rrbracket$ belongs to at least r of S_i . We have*

$$\mathbb{I}^\pi(P) \geq \frac{1}{r} \sum_{i=1}^n \mathbb{I}^{\pi^{(S_i)}}(P_\pi^{(S_i)}).$$

In particular, if we take $S_i = \llbracket d \rrbracket \setminus \{i\}$ and $n = d$ so that $r = d - 1$, we recover the Han's inequality in Corollary 2.1. Equality holds if P is itself a product chain where both sides equal to zero.

2.4 $(S_i)_{i=1}^n$ -factorizable transition matrices and the distance to $(S_i)_{i=1}^n$ -factorizability

Throughout this subsection, we consider a mutually exclusive partition of $\llbracket d \rrbracket$ that we denote by $(S_i)_{i=1}^n$. We also assume that $|S_i| \geq 1$ for all i , and hence $n \in \llbracket d \rrbracket$. Intuitively, we can understand that $\llbracket d \rrbracket$ is partitioned into n blocks with each block being S_i . The aim of this subsection is to find the closest $(S_i)_{i=1}^n$ -factorizable transition matrix to a given multivariate P and hence to compute the distance to $(S_i)_{i=1}^n$ -factorizability of P . In other words, we are looking for transition matrices which are independent across different blocks but possibly dependent within each block of coordinates.

We now define the set of $(S_i)_{i=1}^n$ -factorizable transition matrices.

Definition 2.6 ($(S_i)_{i=1}^n$ -factorizable transition matrices). *Consider a mutually exclusive partition $(S_i)_{i=1}^n$ of $\llbracket d \rrbracket$ with $|S_i| \geq 1$. A transition matrix $P \in \mathcal{L}(\mathcal{X})$ is said to be $(S_i)_{i=1}^n$ -factorizable if there exists $L_i \in \mathcal{L}(\mathcal{X}^{(S_i)})$ such that P can be written as*

$$P = \otimes_{i=1}^n L_i.$$

We write $\mathcal{L}_{\otimes_{i=1}^n S_i}(\mathcal{X})$ to be the set of all $(S_i)_{i=1}^n$ -factorizable transition matrices. In particular, we write $\mathcal{L}_{\otimes}(\mathcal{X}) := \mathcal{L}_{\otimes_{i=1}^d \{i\}}(\mathcal{X})$ to be the set of product transition matrices on \mathcal{X} .

It is obvious to see that $\mathcal{L}_{\otimes}(\mathcal{X}) \subseteq \mathcal{L}_{\otimes_{i=1}^n S_i}(\mathcal{X})$ for any choice of the partition $(S_i)_{i=1}^n$.

Next, we consider the information projection of P onto the space $\mathcal{L}_{\otimes_{i=1}^n S_i}(\mathcal{X})$. We develop a Pythagorean identity in this case:

Theorem 2.7. (*Pythagorean identity*) *Consider a mutually exclusive partition $(S_i)_{i=1}^n$ of $\llbracket d \rrbracket$ with $|S_i| \geq 1$. Let $\pi \in \mathcal{P}(\mathcal{X})$ be a positive probability mass, $P \in \mathcal{L}(\mathcal{X})$, $L_i \in \mathcal{L}(\mathcal{X}^{(S_i)})$ for $i \in \llbracket n \rrbracket$. We then have*

$$D_{KL}^\pi(P \| \otimes_{i=1}^n L_i) = D_{KL}^\pi(P \| \otimes_{i=1}^n P_\pi^{(S_i)}) + D_{KL}^\pi(\otimes_{i=1}^n P_\pi^{(S_i)} \| \otimes_{i=1}^n L_i)$$

$$= D_{KL}^\pi(P \| \otimes_{i=1}^n P_\pi^{(S_i)}) + \sum_{i=1}^n D_{KL}^{\pi^{(S_i)}}(P_\pi^{(S_i)} \| L_i),$$

where we recall that $P_\pi^{(S_i)}$ is the keep- S_i -in transition matrices of P with respect to π in Definition 2.5, while $D_{KL}^{\pi^{(S_i)}}(P_\pi^{(S_i)} \| L_i)$ is weighted by $\pi^{(S_i)}$, the keep- S_i -in marginal distribution of π . In other words, $\otimes_{i=1}^n P_\pi^{(S_i)}$, the tensor product of the keep- S_i -in transition matrices, is the unique minimizer of

$$\min_{L_i \in \mathcal{L}(\mathcal{X}^{(S_i)})} D_{KL}^\pi(P \| \otimes_{i=1}^n L_i) = D_{KL}^\pi(P \| \otimes_{i=1}^n P_\pi^{(S_i)}).$$

Remark 2.6. In fact we have already seen two special cases. In the first special case, we take $n = d$ and $S_i = \{i\}$, in which we recover the Pythagorean identity in Theorem 2.1. In the second special case, for $d \geq 2$, we take $n = 2$ and $S_1 = \{i\}, S_2 = \llbracket d \rrbracket \setminus \{i\}$, in which we retrieve the Pythagorean identity in Theorem 2.5.

Proof.

$$\begin{aligned} D_{KL}^\pi(P \| \otimes_{i=1}^n L_i) &= D_{KL}^\pi(P \| \otimes_{i=1}^n P_\pi^{(S_i)}) + \sum_{x,y} \pi(x) P(x,y) \ln \left(\frac{(\otimes_{i=1}^n P_\pi^{(S_i)})(x,y)}{(\otimes_{i=1}^n L_i)(x,y)} \right) \\ &= D_{KL}^\pi(P \| \otimes_{i=1}^n P_\pi^{(S_i)}) + \sum_{i=1}^n \sum_{x^{(S_i)}, y^{(S_i)}} \pi^{(S_i)}(x^{(S_i)}) P_\pi^{(S_i)}(x^{(S_i)}, y^{(S_i)}) \ln \left(\frac{P_\pi^{(S_i)}(x^{(S_i)}, y^{(S_i)})}{L_i(x^{(S_i)}, y^{(S_i)})} \right) \\ &= D_{KL}^\pi(P \| \otimes_{i=1}^n P_\pi^{(S_i)}) + \sum_{i=1}^n D_{KL}^{\pi^{(S_i)}}(P_\pi^{(S_i)} \| L_i). \end{aligned}$$

□

Using Theorem 2.7 we now introduce a distance to the space $\mathcal{L}_{\otimes_{i=1}^n S_i}(\mathcal{X})$ of a given multivariate P :

Definition 2.7 (Distance to $(S_i)_{i=1}^n$ -factorizability of P with respect to π). Consider a mutually exclusive partition $(S_i)_{i=1}^n$ of $\llbracket d \rrbracket$ with $|S_i| \geq 1$. Let $\pi \in \mathcal{P}(\mathcal{X})$ be a positive probability mass and $P \in \mathcal{L}(\mathcal{X})$. We define

$$\mathbb{I}^\pi(P, \mathcal{L}_{\otimes_{i=1}^n S_i}(\mathcal{X})) := \min_{L_i \in \mathcal{L}(\mathcal{X}^{(S_i)})} D_{KL}^\pi(P \| \otimes_{i=1}^n L_i) = D_{KL}^\pi(P \| \otimes_{i=1}^n P_\pi^{(S_i)}).$$

In particular, if we take $S_i = \{i\}$ and $n = d$, we recover the distance to independence of P with respect to π as introduced in (7):

$$\mathbb{I}^\pi(P) = \mathbb{I}^\pi(P, \mathcal{L}_\otimes(\mathcal{X})) = D_{KL}^\pi(P \| \otimes_{i=1}^d P_\pi^{(i)}).$$

Finally, the above results give an interesting decomposition of the distance to independence of P in terms of the distance to $(S_i)_{i=1}^n$ -factorizability of P .

Corollary 2.3 (Decomposition of the distance to independence of P). *Consider a mutually exclusive partition $(S_i)_{i=1}^n$ of $\llbracket d \rrbracket$ with $|S_i| \geq 1$. Let $\pi \in \mathcal{P}(\mathcal{X})$ be a positive probability mass and $P \in \mathcal{L}(\mathcal{X})$. We have*

$$\begin{aligned} \underbrace{\mathbb{I}^\pi(P)}_{\text{distance to independence of } P} &= \underbrace{\mathbb{I}^\pi(P, \mathcal{L}_{\otimes_{i=1}^n S_i}(\mathcal{X}))}_{\text{distance to } (S_i)_{i=1}^n\text{-factorizability of } P} + \underbrace{\mathbb{I}^\pi(\otimes_{i=1}^n P_\pi^{(S_i)})}_{\text{distance to independence of } \otimes_{i=1}^n P_\pi^{(S_i)}} \\ &= \underbrace{\mathbb{I}^\pi(P, \mathcal{L}_{\otimes_{i=1}^n S_i}(\mathcal{X}))}_{\text{distance to } (S_i)_{i=1}^n\text{-factorizability of } P} + \sum_{i=1}^n \underbrace{\mathbb{I}^{\pi^{(S_i)}}(P_\pi^{(S_i)})}_{\text{distance to independence of } P_\pi^{(S_i)}}. \end{aligned}$$

Proof. Recall that $P_\pi^{(i)}$ is the i th marginal transition matrix of P with respect to π . For arbitrary $(S_i)_{i=1}^n$, we take $L_i = \otimes_{j \in S_i} P_\pi^{(j)}$ in Theorem 2.7. The desired result follows by recalling that with these choices,

$$D_{KL}^\pi(P \| \otimes_{i=1}^n L_i) = \mathbb{I}^\pi(P), \quad D_{KL}^{\pi^{(S_i)}}(P_\pi^{(S_i)} \| L_i) = \mathbb{I}^{\pi^{(S_i)}}(P_\pi^{(S_i)}).$$

□




Figure 2: Visualizations of the set $\mathcal{L}_{\otimes_{i=1}^n S_i}(\mathcal{X})$ and $\mathcal{L}_{\otimes_{i=1}^l V_i}(\mathcal{X})$, where $(V_i)_{i=1}^l$ and $(S_i)_{i=1}^n$ are two partitions of $\llbracket d \rrbracket$. Note that $\mathcal{L}_\otimes(\mathcal{X}) \subseteq \mathcal{L}_{\otimes_{i=1}^n S_i}(\mathcal{X}) \cap \mathcal{L}_{\otimes_{i=1}^l V_i}(\mathcal{X})$, and all the arrows are based upon the divergence D_{KL}^π . We also recall that all $P_m^{(i)}$ depend on π . The Pythagorean identity of $P \in \mathcal{L}(\mathcal{X})$ is stated in Theorem 2.7.

2.5 $(C_i)_{i=1}^n$ -factorizable transition matrices with respect to a graph

In the literature of graphical model and Markov random field, factorization of probability masses with respect to the cliques of a graph has been investigated, and culminates in the Hammersley-Clifford theorem, see for instance (Wainwright, 2019, Chapter 11). In this vein,

in this subsection we consider the problem of factorizability of a transition matrix with respect to the cliques of a graph.

Let us consider a given undirected graph $G = (V = \llbracket d \rrbracket, E)$. A set $C \subseteq \llbracket d \rrbracket$ is said to be a clique of G if any pair of two vertices in C are connected by an edge in E . Let $(C_i)_{i=1}^n$ be a set of cliques of the graph G , with possibly overlapping vertices, such that $\cup_{i=1}^n C_i = \llbracket d \rrbracket$.

We are ready to define the set of $(C_i)_{i=1}^n$ -factorizable transition matrices with respect to the graph G :

Definition 2.8 ($(C_i)_{i=1}^n$ -factorizable transition matrices with respect to a graph G). *Let $(C_i)_{i=1}^n$ be a set of cliques of a graph G on $\llbracket d \rrbracket$ such that $\cup_{i=1}^n C_i = \llbracket d \rrbracket$. A transition matrix $P \in \mathcal{L}(\mathcal{X})$ is said to be $(C_i)_{i=1}^n$ -factorizable with respect to the graph G and $(L_i)_{i=1}^n$ if there exists $L_i \in \mathcal{L}(\mathcal{X}^{(C_i)})$ such that P can be written as, for $x, y \in \mathcal{X}$,*

$$\begin{aligned} P(x, y) &\propto \prod_{i=1}^n L_i(x^{(C_i)}, y^{(C_i)}) \\ &= \frac{1}{\sum_{y \in \mathcal{X}} \prod_{i=1}^n L_i(x^{(C_i)}, y^{(C_i)})} \prod_{i=1}^n L_i(x^{(C_i)}, y^{(C_i)}) \\ &=: \frac{1}{Z(x, (L_i)_{i=1}^n)} \prod_{i=1}^n L_i(x^{(C_i)}, y^{(C_i)}), \end{aligned}$$

where $Z(x, (L_i)_{i=1}^n)$ is the normalization constant of P at x . We write $\mathcal{L}_{\otimes_{i=1}^n C_i}^G(\mathcal{X})$ to be the set of all $(C_i)_{i=1}^n$ -factorizable transition matrices with respect to G . In particular, we note that $\mathcal{L}_{\otimes}(\mathcal{X}) = \mathcal{L}_{\otimes_{i=1}^d \{i\}}^G(\mathcal{X})$ for any graph G .

Let $(S_i)_{i=1}^n$ be a mutually exclusive partition of $\llbracket d \rrbracket$ as in Section 2.4, and we choose a graph G with n disjoint blocks and the members of each S_i form a clique within. With these choices we see that $\mathcal{L}_{\otimes_{i=1}^n S_i}(\mathcal{X}) = \mathcal{L}_{\otimes_{i=1}^n C_i}^G(\mathcal{X})$, and hence Definition 2.8 is a generalization of Definition 2.6. These concepts are illustrated in Figure 2.5.




Figure 3: Illustrations of C_1, C_2, S_1, S_2 on a given 5-node graph with $d = 5$. We take $C_1 = \{1, 2, 3\}$, $C_2 = \{3, 4, 5\}$, $S_1 = \{1, 2, 3\}$ and $S_2 = \{4, 5\}$. Note that these sets are all cliques of the graph and S_1, S_2 together form a partition of $\llbracket 5 \rrbracket$.

To seek the closest $(C_i)_{i=1}^n$ -factorizable transition matrix with respect to the graph G and KL divergence, one such candidate is given by

Definition 2.9. Let $\pi \in \mathcal{P}(\mathcal{X})$ be a positive probability mass and $P \in \mathcal{L}(\mathcal{X})$. We define, for all $x, y \in \mathcal{X}$,

$$\begin{aligned} \mathbf{P}_\pi(x, y) &\propto \prod_{i=1}^n P_\pi^{(C_i)}(x^{(C_i)}, y^{(C_i)}) \\ &= \frac{1}{Z(x, (P_\pi^{(C_i)})_{i=1}^n)} \prod_{i=1}^n P_\pi^{(C_i)}(x^{(C_i)}, y^{(C_i)}), \end{aligned}$$

where we recall that $P_\pi^{(C_i)}$ is the keep- C_i -in transition matrix with respect to π and $Z(x, (P_\pi^{(C_i)})_{i=1}^n)$ is introduced in Definition 2.8. Note that \mathbf{P}_π depends on $(C_i)_{i=1}^n$, P and π , and $\mathbf{P}_\pi \in \mathcal{L}_{\otimes_{i=1}^n C_i}^G(\mathcal{X})$.

We prove a Pythagorean inequality, which allows us to conclude that \mathbf{P}_π is indeed the closest $(C_i)_{i=1}^n$ -factorizable transition matrix with normalization constants greater than or equal to that of \mathbf{P}_π , with respect to the graph G and KL divergence.

Theorem 2.8. (Pythagorean inequality) Let $(C_i)_{i=1}^n$ be a set of cliques of a graph G on $\llbracket d \rrbracket$ such that $\cup_{i=1}^n C_i = \llbracket d \rrbracket$. Let $\pi \in \mathcal{P}(\mathcal{X})$ be a positive probability mass, $P \in \mathcal{L}(\mathcal{X})$, $M \in \mathcal{L}_{\otimes_{i=1}^n C_i}^G(\mathcal{X})$, $L_i \in \mathcal{L}(\mathcal{X}^{(C_i)})$ for $i \in \llbracket n \rrbracket$ such that M is $(C_i)_{i=1}^n$ -factorizable with respect to the graph G and $(L_i)_{i=1}^n$. For all M such that $Z(x, (L_i)_{i=1}^n) \geq Z(x, (P_\pi^{(C_i)})_{i=1}^n)$ for all x , we have

$$D_{KL}^\pi(P \| M) \geq D_{KL}^\pi(P \| \mathbf{P}_\pi) + \sum_{i=1}^n D_{KL}^{\pi^{(C_i)}}(P_\pi^{(C_i)} \| L_i),$$

where $P_\pi^{(C_i)}$ is the keep- C_i -in transition matrices of P with respect to π in Definition 2.5, while $D_{KL}^{\pi^{(C_i)}}(P_\pi^{(C_i)}; L_i)$ is weighted by $\pi^{(C_i)}$, the keep- C_i -in marginal distribution of π . In other words, \mathbf{P}_π is the unique minimizer of

$$\min_{M \in \mathcal{L}_{\otimes_{i=1}^n C_i}^G(\mathcal{X}); Z(x, (L_i)_{i=1}^n) \geq Z(x, (P_\pi^{(C_i)})_{i=1}^n) \forall x} D_{KL}^\pi(P \| M) = D_{KL}^\pi(P \| \mathbf{P}_\pi).$$

Proof.

$$\begin{aligned} D_{KL}^\pi(P \| M) &= D_{KL}^\pi(P \| \mathbf{P}_\pi) + \sum_{x, y} \pi(x) P(x, y) \ln \left(\frac{Z(x, (L_i)_{i=1}^n)}{Z(x, (P_\pi^{(C_i)})_{i=1}^n)} \frac{(\otimes_{i=1}^n P_\pi^{(S_i)})(x, y)}{(\otimes_{i=1}^n L_i)(x, y)} \right) \\ &\geq D_{KL}^\pi(P \| \mathbf{P}_\pi) + \sum_{i=1}^n \sum_{x^{(C_i)}, y^{(C_i)}} \pi^{(C_i)}(x^{(C_i)}) P_\pi^{(C_i)}(x^{(C_i)}, y^{(C_i)}) \ln \left(\frac{P_\pi^{(C_i)}(x^{(C_i)}, y^{(C_i)})}{L_i(x^{(C_i)}, y^{(C_i)})} \right) \\ &= D_{KL}^\pi(P \| \mathbf{P}_\pi) + \sum_{i=1}^n D_{KL}^{\pi^{(C_i)}}(P_\pi^{(C_i)} \| L_i). \end{aligned}$$

□

2.6 Comparisons of mixing and hitting time parameters between P and its information projections

Let $S \subseteq \llbracket d \rrbracket$. In Section 2.4, we have seen that $P^{(S)} \otimes P^{(-S)}$, the tensor product of the keep- S -in and leave- S -out transition matrix of a given π -stationary $P \in \mathcal{L}(\mathcal{X})$, arises naturally as an information projection of P onto the space $\mathcal{L}_{S \otimes \llbracket d \rrbracket \setminus S}(\mathcal{X})$. The objective of this subsection is to investigate the relationship of hitting and mixing time parameters such as commute time, right spectral gap, log-Sobolev constant and Cheeger's constant between P and its information projections. These parameters play important roles in bounding the hitting or mixing time of P , see for instance [Aldous and Fill \(2002\)](#); [Levin and Peres \(2017\)](#); [Montenegro and Tetali \(2006\)](#); [Saloff-Coste \(1997\)](#) and the references therein.

To this end, let us fix the notations. Throughout this subsection, we assume that $X = (X_n)_{n \in \mathbb{N}}$ is an ergodic π -reversible Markov chain with transition matrix $P \in \mathcal{L}(\pi)$. In view of Proposition 2.3, we thus see that $X^{(S)} := (X_n^{(S)})_{n \in \mathbb{N}}$ is also an ergodic $\pi^{(S)}$ -reversible Markov chain with transition matrix $P^{(S)}$. We write, for $f : \mathcal{X} \rightarrow \mathbb{R}$,

$$\begin{aligned} \mathbb{E}_\pi(f) &:= \sum_{x \in \mathcal{X}} \pi(x) f(x), & \text{Var}_\pi(f) &:= \sum_{x \in \mathcal{X}} \pi(x) (f(x) - \mathbb{E}_\pi(f))^2, \\ \text{Ent}_\pi(f) &:= \mathbb{E}_\pi \left(f \ln \frac{f}{\mathbb{E}_\pi(f)} \right), & \mathcal{D}_\pi(f, f) &:= \frac{1}{2} \sum_{x, y \in \mathcal{X}} \pi(x) P(x, y) (f(x) - f(y))^2, \end{aligned}$$

to be, respectively, the expectation, variance, entropy of f with respect to π and the Dirichlet form of P with respect to f, π . We are interested in the following list of hitting and mixing time parameters associated with X and $X^{(S)}$:

- (Right spectral gap, relaxation time and log-Sobolev constant) The right spectral gap and the log-Sobolev constant of P are defined respectively to be

$$\gamma(P) := \inf_{f; \text{Var}_\pi(f) \neq 0} \frac{\mathcal{D}_\pi(f, f)}{\text{Var}_\pi(f)}, \quad \alpha(P) := \inf_{f; \text{Ent}_\pi(f^2) \neq 0} \frac{\mathcal{D}_\pi(f, f)}{\text{Ent}_\pi(f^2)}.$$

Note that since P is assumed to be reversible and ergodic, the right spectral gap can be written as $\gamma(P) = 1 - \lambda_2(P)$, where $\lambda_2(P) < 1$ is the second largest eigenvalue of P . For $S \subseteq \llbracket d \rrbracket$, we shall analogously consider the right spectral gap $\gamma(P^{(S)})$ and the log-Sobolev constant $\alpha(P^{(S)})$ of $P^{(S)}$ by replacing π in the definitions above with $\pi^{(S)}$ and \mathcal{X} with $\mathcal{X}^{(S)}$. The relaxation time of the continuized chain of P is defined to be $t_{rel}(P) := 1/\gamma(P)$.

- (Cheeger's constant) Let $\emptyset \neq A \subseteq \mathcal{X} = \times_{i=1}^d \mathcal{X}^{(i)}$ and $A^c := \mathcal{X} \setminus A$. We define

$$\Phi_A(P) := \frac{(\pi \boxtimes P)(A, A^c)}{\pi(A)},$$

and hence the Cheeger's constant of P (Levin and Peres, 2017, Chapter 7.2) is defined to be

$$\Phi(P) := \min_{A \subset \mathcal{X}; 0 < \pi(A) \leq 1/2} \Phi_A(P).$$

Analogously, we define the Cheeger's constant of the keep- S -in transition matrix $P^{(S)}$. Let $\emptyset \neq A^{(S)} \subseteq \mathcal{X}^{(S)}$ and $A^{(S)c} = \mathcal{X}^{(S)} \setminus A^{(S)}$. We then have

$$\begin{aligned} \Phi_{A^{(S)}}(P^{(S)}) &= \frac{(\pi^{(S)} \boxtimes P^{(S)})(A^{(S)}, A^{(S)c})}{\pi^{(S)}(A^{(S)})}, \\ \Phi(P^{(S)}) &= \min_{A^{(S)} \subset \mathcal{X}^{(S)}; 0 < \pi^{(S)}(A^{(S)}) \leq 1/2} \Phi_{A^{(S)}}(P^{(S)}). \end{aligned}$$

- (Commute time and average hitting time) Let $x, y \in \mathcal{X}$. The hitting time to the state x (resp. $x^{(S)}$) of the Markov chain X (resp. $X^{(S)}$) are defined to be

$$\tau_x(P) := \inf\{n \geq 0; X_n = x\}, \quad \tau_{x^{(S)}}(P^{(S)}) := \inf\{n \geq 0; X_n^{(S)} = x^{(S)}\},$$

where the usual convention of $\inf \emptyset := 0$ applies. The mean commute time between x and y of X are given by

$$\mathbb{E}_x(\tau_y(P)) + \mathbb{E}_y(\tau_x(P)) = \sup_{0 \leq f \leq 1; f(x)=1, f(y)=0} \frac{1}{\mathcal{D}_\pi(f, f)}. \quad (15)$$

Analogously one can define the mean commute time between $x^{(S)}$ and $y^{(S)}$ of $X^{(S)}$ to be

$$\mathbb{E}_{x^{(S)}}(\tau_{y^{(S)}}(P^{(S)})) + \mathbb{E}_{y^{(S)}}(\tau_{x^{(S)}}(P^{(S)})).$$

The maximal mean commute time t_c is defined to be

$$\begin{aligned} t_c(P) &:= \max_{x, y \in \mathcal{X}} \mathbb{E}_x(\tau_y(P)) + \mathbb{E}_y(\tau_x(P)), \\ t_c(P^{(S)}) &= \max_{x^{(S)}, y^{(S)} \in \mathcal{X}^{(S)}} \mathbb{E}_{x^{(S)}}(\tau_{y^{(S)}}(P^{(S)})) + \mathbb{E}_{y^{(S)}}(\tau_{x^{(S)}}(P^{(S)})). \end{aligned}$$

The average hitting time t_{av} is defined to be

$$\begin{aligned} t_{av}(P) &:= \sum_{x, y \in \mathcal{X}} \pi(x)\pi(y)\mathbb{E}_x(\tau_y(P)), \\ t_{av}(P^{(S)}) &= \sum_{x^{(S)}, y^{(S)} \in \mathcal{X}^{(S)}} \pi^{(S)}(x^{(S)})\pi^{(S)}(y^{(S)})\mathbb{E}_{x^{(S)}}(\tau_{y^{(S)}}(P^{(S)})), \end{aligned}$$

We proceed to develop results to compare these parameters between P and its information projections. For instance, in the case of the right spectral gap, we are interested in bounding $\gamma(P)$ with $\gamma(P^{(S)})$ or $\gamma(P^{(S)} \otimes P^{(-S)})$.

The main result of this subsection recalls a contraction principle in (Aldous and Fill, 2002, Proposition 4.44): the hitting and mixing time parameters such as $\gamma, \alpha, \Phi, t_c, t_{av}$ are at least "faster" for $P^{(S)}$ than the original chain P . We also establish new monotonicity results for the parameters $S \mapsto \mathbb{I}^{\pi^{(S)}}(P^{(S)})$ and $S \mapsto D_{KL}^{\pi^{(S)}}(P^{(S)}) \|\Pi^{(S)}\|$ via the Shearer's lemma.

Corollary 2.4 (Contraction principle and monotonicity). *Let $\emptyset \neq T \subseteq S \subseteq \llbracket d \rrbracket$ and $P \in \mathcal{L}(\pi)$ be an ergodic transition matrix. We have*

$$\begin{aligned}\gamma(P) &\leq \gamma(P^{(S)}) \leq \gamma(P^{(T)}), \\ \alpha(P) &\leq \alpha(P^{(S)}) \leq \alpha(P^{(T)}), \\ \Phi(P) &\leq \Phi(P^{(S)}) \leq \Phi(P^{(T)}), \\ t_c(P) &\geq t_c(P^{(S)}) \geq t_c(P^{(T)}), \\ t_{av}(P) &\geq t_{av}(P^{(S)}) \geq t_{av}(P^{(T)}).\end{aligned}$$

If furthermore $\pi = \otimes_{i=1}^d \pi^{(i)}$ is a product stationary distribution, then

$$\begin{aligned}\mathbb{I}^\pi(P) &\geq \mathbb{I}^{\pi^{(S)}}(P^{(S)}) \geq \mathbb{I}^{\pi^{(T)}}(P^{(T)}), \\ D_{KL}^\pi(P \parallel \Pi) &\geq D_{KL}^{\pi^{(S)}}(P^{(S)} \parallel \Pi^{(S)}) \geq D_{KL}^{\pi^{(T)}}(P^{(T)} \parallel \Pi^{(T)}),\end{aligned}$$

where $\Pi \in \mathcal{L}(\pi)$ is the transition matrix with each row being π . All the above equalities hold if $T = \llbracket d \rrbracket$.

Remark 2.7. *We note that it is perhaps possible to obtain tighter bounds in Proposition 2.4 by considering the corresponding hitting or mixing time parameters of the “restriction chains” as in Jerrum et al. (2004). In this manuscript we shall not introduce these chains.*

In Section 3.1.2, we shall show that the Glauber dynamics, being a non-tensor-product transition matrix, attain equalities for γ and α , that is, there exists $\emptyset \neq S \subset \llbracket d \rrbracket$ such that $\gamma(P) = \gamma(P^{(S)})$ and $\alpha(P) = \alpha(P^{(S)})$.

Proof of Corollary 2.4. Once we have obtained the inequalities governing P and $P^{(S)}$, we replace P by $P^{(S)}$ and note that $(P^{(S)})^{(T)} = P^{(T)}$ to reach at the inequalities governing $P^{(S)}$ and $P^{(T)}$. As a result, it suffices to prove only the inequalities between P and $P^{(S)}$.

The case of $\gamma, \alpha, \Phi, t_c, t_{av}$ have already been analyzed in (Aldous and Fill, 2002, Proposition 4.44).

Using the Shearer’s lemma for distance to independence in Corollary 2.2, we see that

$$\mathbb{I}^\pi(P) \geq \mathbb{I}^{\pi^{(S)}}(P^{(S)}) + \mathbb{I}^{\pi^{(-S)}}(P^{(-S)}) \geq \mathbb{I}^{\pi^{(S)}}(P^{(S)}).$$

Finally, using again the Shearer’s lemma for the KL divergence of Markov chains in Theorem 2.6 leads to

$$D_{KL}^\pi(P \parallel \Pi) \geq D_{KL}^{\pi^{(S)}}(P^{(S)} \parallel \Pi^{(S)}) + D_{KL}^{\pi^{(-S)}}(P^{(-S)} \parallel \Pi^{(-S)}) \geq D_{KL}^{\pi^{(S)}}(P^{(S)} \parallel \Pi^{(S)}).$$

□

The next result compares $\gamma(P)$ and $\gamma(P^{(S)} \otimes P^{(-S)})$ to obtain

Corollary 2.5. *Let $\emptyset \neq S \subseteq \llbracket d \rrbracket$. Let $P \in \mathcal{L}(\pi)$ be a lazy, ergodic and π -reversible transition matrix. We have*

$$\gamma(P) \leq \gamma(P^{(S)} \otimes P^{(-S)}).$$

In particular, this yields $t_{rel}(P) \geq t_{rel}(P^{(S)} \otimes P^{(-S)})$.

Proof. As P is lazy, both $P^{(S)}, P^{(-S)}$ are also lazy by Proposition 2.3, and hence $\lambda_2(P^{(S)} \otimes P^{(-S)}) = \max\{\lambda_2(P^{(S)}), \lambda_2(P^{(-S)})\}$. The desired result follows from Proposition 2.4. \square

3 Unified view of key MCMC algorithms via rate distortion theory

In the previous Section 2, we defined $\mathbb{I}^\pi(P)$, the distance to independence of a given transition matrix P , as a divergence from the closest product chain, and we proposed to interpret this distance as a generalization of mutual information in the context of Markov chains. We further investigated factorizability of multivariate Markov chains, and information projection problems with respect to general f -divergences between transition matrices.

We now move back to the rate distortion problem introduced in Section 1. Recall from (2) that rate distortion optimization can be expressed in terms of a KL divergence in the distribution setting. In this section, we will extend this divergence formulation to the Markovian setting. We will further demonstrate that the transition matrices of a plethora of MCMC algorithms can be viewed as the optimizer of this rate distortion problem, thus offering an unified variational perspective on the optimality of these algorithms.

First, we introduce the rate distortion problem for Markov chains:

Definition 3.1 (Rate distortion function of $M \in \mathcal{L}(\mathcal{X})$ and of $\pi \in \mathcal{P}(\mathcal{X})$). *Let $c : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ be a bounded function that we call the (distortion) cost function. The rate distortion optimization problem of a source Markov chain M with respect to the cost function c and target distribution π is defined to be*

$$\min_{P \in \mathcal{L}(\mathcal{X})} R(\pi, P, M, c) \text{ where } R(\pi, P, M, c) := D_{KL}^\pi(P \| M) + \mathbb{E}_{(X,Y) \sim \pi \boxtimes P}[c(X, Y)].$$

Note that when π is a positive probability function, we can rewrite the objective function R as

$$R(\pi, P, M, c) = \tilde{D}_{KL}(\pi \boxtimes P \| \pi \boxtimes M) + \mathbb{E}_{(X,Y) \sim \pi \boxtimes P}[c(X, Y)].$$

The objective function R can be interpreted as a relaxation of a constrained optimization problem. The objective of the constrained problem is called the rate distortion function \mathbf{R} of a given M with respect to Δ , the cost function c and π :

$$\mathbf{R}(\pi, \Delta, M, c) := \min_{P; \mathbb{E}_{(X,Y) \sim \pi \boxtimes P}[c(X,Y)] \leq \Delta} D_{KL}^\pi(P \| M).$$

The above objective function can be instructively compared with the classical rate distortion function of π with respect to Δ and c ((Cover and Thomas, 2006, Section 10.7), (Polyanskiy and Wu, 2022, Equation (5.30)), (Shafieepoorfard et al., 2016, Section 4)), which can be expressed as

$$\begin{aligned}\tilde{\mathbf{R}}(\pi, \Delta, c) &:= \min_{P; (X,Y) \sim \pi \boxtimes P, \mathbb{E}[c(X,Y)] \leq \Delta} I(X;Y) \\ &= \min_{P; \mathbb{E}_{(X,Y) \sim \pi \boxtimes P}[c(X,Y)] \leq \Delta} \tilde{D}_{KL}(\pi \boxtimes P \| \pi \otimes \pi P) \\ &= \min_{P; \mathbb{E}_{(X,Y) \sim \pi \boxtimes P}[c(X,Y)] \leq \Delta} \min_{\nu \in \mathcal{P}(\mathcal{X})} \tilde{D}_{KL}(\pi \boxtimes P \| \pi \otimes \nu),\end{aligned}$$

where the first equality expresses $\tilde{\mathbf{R}}$ in terms of the mutual information $I(X;Y)$, while the third equality follows from a well-known variational principle, see e.g. (Shafieepoorfard et al., 2016, equation (2.3)), and

We define a transformation $\Psi_c : \mathcal{L}(\mathcal{X}) \rightarrow \mathcal{L}(\mathcal{X})$ with respect to a cost function c , that we call the Boltzmann-Gibbs transformation, to be, for all $x, y \in \mathcal{X}$,

$$\Psi_c(M)(x, y) := \frac{M(x, y)e^{-c(x,y)}}{\sum_{z \in \mathcal{X}} M(x, z)e^{-c(x,z)}}. \quad (16)$$

For each fixed x , the probability measure $\Psi_c(M)(x, \cdot)$ is the Boltzmann-Gibbs transformation of $M(x, \cdot)$ with respect to cost function c , see for example Del Moral (2004).

Theorem 3.1. *Consider the rate distortion optimization problem of M with respect to c and π . The Boltzmann-Gibbs transformation of M with respect to c is a minimizer, that is,*

$$\Psi_c(M) \in \arg \min_{P \in \mathcal{L}(\mathcal{X})} R(\pi, P, M, c).$$

Note that $\Psi_c(M)$ does not depend on π . When π is a positive probability mass, $\Psi_c(M)$ is furthermore the unique minimizer.

Proof. Let $x \in \mathcal{X}$. When $\pi(x) > 0$, we compute the derivative of R with respect to $P(x, y)$ to give

$$\frac{\partial}{\partial P(x, y)} R(\pi, P, M, c) = \pi(x) \left(\ln \left(\frac{P(x, y)}{M(x, y)} \right) + c(x, y) - \ln \left(\frac{P(x, z)}{M(x, z)} \right) - c(x, z) \right),$$

where z is a reference point such that $P(x, z) = 1 - \sum_{y \neq z} P(x, y)$. Since $\pi(x) > 0$, the unique $P(x, \cdot)$ that sets the above to zero is

$$P(x, y) = \Psi_c(M)(x, y).$$

When $\pi(x) = 0$, then the function $R(\pi, P, M, c)$ does not depend on $P(x, \cdot)$. In particular, any $P(x, \cdot) \in \mathcal{P}(\mathcal{X})$ minimizes R , and hence we can take $P(x, \cdot) = \Psi_c(M)(x, \cdot)$. \square

Analogous to (Hernández-Hernández and Marcus, 1996, Lemma 3.3), we denote the Legendre transformation of $\mathcal{L}(\mathcal{X}) \ni P \mapsto \tilde{D}_{KL}(\pi \boxtimes P \| \pi \boxtimes M)$ to be

$$\begin{aligned} \tilde{D}_{KL}^*(c) &:= \max_{P \in \mathcal{L}(\mathcal{X})} \mathbb{E}_{(X,Y) \sim \pi \boxtimes P}[-c(X,Y)] - \tilde{D}_{KL}(\pi \boxtimes P \| \pi \boxtimes M) \\ &= - \min_{P \in \mathcal{L}(\mathcal{X})} R(\pi, P, M, c) \\ &= \mathbb{E}_{X \sim \pi}[\ln \mathbb{E}_{Y|X \sim M(X, \cdot)}[e^{-c(X,Y)}]]. \end{aligned}$$

Note that the dependence of \tilde{D}_{KL}^* on π, M is suppressed, and \tilde{D}_{KL}^* is also known as the convex conjugate. As such, we can understand $\tilde{D}_{KL}^*(c)$ as an expected cumulant generating function of $c(X, Y)$.

As an illustration of the rate distortion function \mathbf{R} of M with respect to Δ and $\Psi_c(M)$, we consider Markov chains on the two-point state space $\mathcal{X} = \{1, 2\}$, which is a common example in the Markov chain literature, see for instance Billera and Diaconis (2001). Let the cost function be chosen such that $c(1, 1), c(2, 2) \leq 0$, $c(1, 2), c(2, 1) > 0$ and $\pi = (\pi(1), \pi(2))$ be a positive probability mass. In Figure 4, we visualize these notions. For $P \in \mathcal{L}(\{1, 2\})$, it can be parameterized by the two coordinates $(P(1, 2), P(2, 1))$ and hence it lies in the box $[0, 1] \times [0, 1]$. We represent the source Markov chain M with $(M(1, 2), M(2, 1))$ as the black cross. With these choices, the distortion cost constraint can be written as

$$\begin{aligned} &\{P; \mathbb{E}_{(X,Y) \sim \pi \boxtimes P}[c(X, Y)] \leq \Delta\} \\ &= \{P; \pi(2)P(2, 1)(c(2, 1) - c(2, 2)) + \pi(1)P(1, 2)(c(1, 2) - c(1, 1)) \leq \Delta - \pi(1)c(1, 1) - \pi(2)c(2, 2)\}, \end{aligned}$$

which is represented by the blue triangle. As the mapping $P \mapsto D_{KL}^\pi(P \| M)$ is convex, the red ellipses are the contour lines of this map with the unique minimum at M . $\Psi_c(M)$ is visualized by the red circle dot with the coordinate $(\Psi_c(M)(1, 2), \Psi_c(M)(2, 1))$. Note that all of the central MCMC algorithms that we cover in this paper can be understood as in Figure 4, where different algorithms correspond to different M (i.e. different black crosses) and cost functions c (i.e. different blue triangles). Also, different choices of π lead to different contour lines (but still convex) and different distortion constraint regions.




Figure 4: Geometric interpretation of the rate distortion function \mathbf{R} of the source Markov chain M with respect to Δ and $\Psi_c(M)$ on a two-point state space $\mathcal{X} = \{1, 2\}$, where $c(1, 1), c(2, 2) \leq 0$, $c(1, 2), c(2, 1) > 0$ and $\pi = (\pi(1), \pi(2))$ is a positive probability mass.

Let us now illustrate Theorem 3.1 with the classical Metropolis-Hastings chain. We shall provide more examples of MCMC algorithms in Section 3.1. Let P_0 be an ergodic and reversible chain with stationary distribution being the discrete uniform on \mathcal{X} and satisfies $P(x, x) > 0$ for all $x \in \mathcal{X}$. Denote the Boltzmann-Gibbs distribution associated with energy function H at inverse temperature $\beta \geq 0$ to be

$$\pi_\beta(x) \propto e^{-\beta H(x)}.$$

To generate samples (approximately) from π_β , one possible Markov chain Monte Carlo algorithm is the Metropolis-Hastings chain with proposal chain P_0 and target distribution π_β . If we take the distortion cost function c and the source Markov chain M to be respectively, for $x, y \in \mathcal{X}$,

$$c(x, y) = \begin{cases} \beta(H(y) - H(x))_+, & \text{if } x \neq y, \\ \ln P_0(x, x) - \ln \left(1 - \sum_{z; z \neq x} P_0(x, z) e^{-\beta(H(z) - H(x))_+} \right), & \text{if } x = y, \end{cases} \quad (17)$$

$$M = P_0, \quad (18)$$

then we have

$$\Psi_c(M)(x, y) = \begin{cases} P_0(x, y) e^{-\beta(H(y) - H(x))_+}, & \text{if } x \neq y, \\ 1 - \sum_{z; z \neq x} P_0(x, z) e^{-\beta(H(z) - H(x))_+}, & \text{if } x = y, \end{cases} \quad (19)$$

where $a_+ := \max\{a, 0\}$ denotes the non-negative part of $a \in \mathbb{R}$. We first note that our choice of c in (17) is non-positive on the diagonal, since

$$P_0(x, x) + \sum_{z; z \neq x} P_0(x, z) e^{-\beta(H(z) - H(x))_+} \leq P_0(x, x) + \sum_{z; z \neq x} P_0(x, z) = 1.$$

Rearranging, we arrive at $c(x, x) \leq 0$ for all x . We refer readers to Remark 3.2 for a discussion on the diagonal entries of c . We see that with these choices $\Psi_c(M)$ in (19) is exactly the classical Metropolis-Hastings chain, and it is the unique minimizer of the rate distortion optimization problem of M with respect to c and any positive probability mass π . The smaller the temperature is, the larger the value of β , and hence the larger the value of the expected cost $\mathbb{E}_{(X, Y) \sim \pi \boxtimes P}[c(X, Y)]$. This forces the optimal P to put probability masses on the diagonals $P(x, x)$ for all x , which is consistent with our understanding on the behaviour of Metropolis-Hastings chain at low temperature. We summarize this into the following corollary:

Corollary 3.1 (Variational interpretation of the transition matrix of the Metropolis-Hastings chain). *Consider the rate distortion optimization problem of M with respect to c and a positive π , where c, M are chosen as in (17) and (18) respectively. $\Psi_c(M)$, the Boltzmann-Gibbs transformation of M with respect to c in (19), is the transition matrix of the Metropolis-Hastings chain with proposal chain P_0 and target distribution π_β . $\Psi_c(M)$ is also the unique minimizer of the rate distortion optimization problem of M .*

Remark 3.1. *The pair (M, c) chosen as in (18) and (17) is not the unique pair that generates $\Psi_c(M)$. Another such pair, $(N, c_{N, M})$, can be defined as follows, where both $N \neq M$ and $c_{N, M} \neq c$. Let $\mathcal{L}(\mathcal{X}) \ni N \neq M$ be another positive transition matrix. If we consider the cost function $c_{N, M}$ defined to be*

$$c_{N, M}(x, y) = \begin{cases} \ln \left(\frac{N(x, y)}{M(x, y)} \right) + \beta(H(y) - H(x))_+, & \text{if } x \neq y, \\ \ln N(x, x) - \ln \left(1 - \sum_{z; z \neq x} M(x, z) e^{-\beta(H(z) - H(x))_+} \right), & \text{if } x = y, \end{cases}$$

then we see that

$$\begin{aligned} \Psi_{c_{N, M}}(N)(x, y) &= \begin{cases} M(x, y) e^{-\beta(H(y) - H(x))_+}, & \text{if } x \neq y, \\ 1 - \sum_{z; z \neq x} M(x, z) e^{-\beta(H(z) - H(x))_+}, & \text{if } x = y, \end{cases} \\ &= \Psi_c(M)(x, y). \end{aligned}$$

A geometric understanding is also possible as in Figure 4. If the black cross at M and the blue triangle constraint region yields an optimal red dot $\Psi_c(M)$, then another black cross at N with another constraint region leads to the same optimal red dot $\Psi_c(M)$. We argue that our choice of (M, c) is natural in the context of MCMC: M can be naturally interpreted as the proposal chain while c is chosen to enforce π -reversibility or π -stationarity of the resulting $\Psi_c(M)$. Also, we note that our choice of c does not depend on M for the off-diagonal entries.

Remark 3.2 (On the distortion cost function along the diagonal entries). *As shown earlier our choice of the distortion cost function in (17) is non-positive along all diagonal entries. This is unconventional although cost function on \mathbb{R} rather than \mathbb{R}_+ is still studied in the literature (Polyanskiy and Wu, 2022, Page 484). If one would rather have a zero cost function along all diagonal entries, there are two ways that we suggest.*

In the first way, we define a normalized cost function \tilde{c} as in (Yeung, 2008, Definition 8.5, equation (8.9)) to be

$$\tilde{c}(x, y) := c(x, y) - c(x, x) \geq 0.$$

We can then associate a $\tilde{M} = \tilde{M}(c)$ so that $\Psi_c(M) = \Psi_{\tilde{c}}(\tilde{M})$.

In the second way, let us consider a KL-type divergence that is defined to be

$$\mathbf{D}_{KL}^\pi(P\|M) := \sum_x \pi(x) \sum_{y; y \neq x} P(x, y) \ln \left(\frac{P(x, y)}{M(x, y)} \right).$$

In other words, we omit all diagonal entries in \mathbf{D}_{KL}^π . This divergence has been studied in Billera and Diaconis (2001), and is very similar to the divergence in Diaconis and Miclo (2009) in continuous-time generators. Analogously, if we consider the rate-distortion optimization problem given by

$$\min_{P \in \mathcal{L}(\mathcal{X})} \mathbf{D}_{KL}^\pi(P\|M) + \mathbb{E}_{(X, Y) \sim \pi \boxtimes P} [c(X, Y)], \quad (20)$$

then by taking

$$c(x, y) = \begin{cases} \beta(H(y) - H(x))_+, & \text{if } x \neq y, \\ 0, & \text{if } x = y, \end{cases}$$

$$M = P_0,$$

we see that a minimizer of (20) is given by

$$\Phi_c(M)(x, y) = \begin{cases} P_0(x, y) e^{-\beta(H(y) - H(x))_+}, & \text{if } x \neq y, \\ 1 - \sum_{z; z \neq x} P_0(x, z) e^{-\beta(H(z) - H(x))_+}, & \text{if } x = y, \end{cases}$$

which is the transition matrix of Metropolis-Hastings. In order to be consistent with Section 2, in this paper we shall consider the problem in Definition 3.1 rather than (20).

3.1 Variational interpretation of the probability law of MCMC algorithms

In the previous section, we have argued in Theorem 3.1 that $\Psi_c(M)$ is an optimal transition matrix in the sense of solving the rate distortion optimization problem. In this subsection, we shall demonstrate that the probability law at time n also admits a variational interpretation.

Given $\pi, \zeta \in \mathcal{P}(\mathcal{X})$ and a sequence of possibly time-dependent source chains $(M_l)_{l \in \mathbb{N}}$ and cost functions $(c_l)_{l \in \mathbb{N}}$ with $M_l \in \mathcal{L}(\mathcal{X})$ for all l , we consider the following recursively defined information-theoretic optimization problems for $l \in \mathbb{N}$:

$$P_l = \Psi_{c_l}(M_l) \in \arg \min_{P \in \mathcal{L}(\mathcal{X})} R(\pi, P, M_l, c_l), \quad (21)$$

$$\zeta_l \in \arg \min_{\eta \in \mathcal{P}(\mathcal{X})} \tilde{D}_{KL}(\zeta_{l-1} \boxtimes P_l \| \zeta_{l-1} \otimes \eta), \quad (22)$$

with the initial condition $\zeta_0 := \zeta$. We note that (21) is the rate distortion optimization problem of M_l with respect to c_l and π at time l , while (22) is the Shannon mutual information problem (see e.g. [Shafieepoorfard et al. \(2016\)](#)). We thus see that $\zeta_l = \zeta_{l-1} \Psi_{c_l}(M_l)$ and hence

$$\zeta_n = \zeta \prod_{l=1}^n \Psi_{c_l}(M_l).$$

Starting from an initial law ζ , the law of a non-homogeneous or adaptive MCMC chain at time n , ζ_n , solves the optimization problem (22).

We summarize the above into the following result:

Theorem 3.2. *Given $n \in \mathbb{N}$, an initial law $\zeta \in \mathcal{P}(\mathcal{X})$ and a sequence of possibly time-dependent source chains $(M_l)_{l \in \mathbb{N}}$ and cost functions $(c_l)_{l \in \mathbb{N}}$ with $M_l \in \mathcal{L}(\mathcal{X})$ for all n , the law $\zeta_n = \zeta \prod_{l=1}^n \Psi_{c_l}(M_l)$ solves the optimization problem (22).*

We shall illustrate Theorem 3.2 via simulated annealing in Section 3.1.5 which is a non-homogeneous MCMC algorithm.

In the case of homogeneous MCMC algorithms, we can take $M_l = M$ and $c_l = c$ so that $\zeta_n = \zeta \Psi_c(M)^n$. This yields the following corollary on a variational interpretation of the probability law of a discrete-time Metropolis-Hastings chain at time n :

Corollary 3.2 (Variational interpretation of the law of the Metropolis-Hastings chain at time n). *Given $n \in \mathbb{N}$, an initial law $\zeta \in \mathcal{P}(\mathcal{X})$, $M \in \mathcal{L}(\mathcal{X})$ and a cost function c , where c, M are chosen as in (17) and (18) respectively. The law $\zeta_n = \zeta \Psi_c(M)^n$, which is the law of the Metropolis-Hastings chain with proposal chain P_0 and target distribution π_β at time n , solves the optimization problem (22).*

In the rest of this section, we shall illustrate Theorem 3.1 and 3.2 via a variety of MCMC algorithms.

3.1.1 Example: Metropolis-Hastings type Markov chains with general acceptance function

In the first example, we consider Metropolis-Hastings type Markov chains with general acceptance function. Let $\alpha : \mathcal{X} \times \mathcal{X} \rightarrow (0, 1]$ be a function that we call an acceptance function.

If we choose, for $x, y \in \mathcal{X}$,

$$c(x, y) = \begin{cases} -\ln \alpha(x, y), & \text{if } x \neq y, \\ \ln P_0(x, x) - \ln \left(1 - \sum_{z; z \neq x} P_0(x, z) \alpha(x, z)\right), & \text{if } x = y, \end{cases} \quad (23)$$

$$M = P_0, \quad (24)$$

so that

$$\Psi_c(M)(x, y) = \begin{cases} P_0(x, y) \alpha(x, y), & \text{if } x \neq y, \\ 1 - \sum_{z; z \neq x} P_0(x, z) \alpha(x, z), & \text{if } x = y, \end{cases} \quad (25)$$

then $\Psi_c(M)$ is a Metropolis-Hastings type Markov chain with proposal chain P_0 . Analogous to Corollary 3.1 and 3.2, we obtain that

Corollary 3.3 (Variational interpretation of the transition matrix and of the law of the Metropolis-Hastings type chain). *Consider the rate distortion optimization problem of M with respect to c and a positive π , where c, M are chosen as in (23) and (24) respectively. $\Psi_c(M)$, the Boltzmann-Gibbs transformation of M with respect to c in (25), is exactly the transition matrix of the Metropolis-Hastings type chain with proposal chain P_0 . It is also the unique minimizer of the rate distortion optimization problem of M .*

Suppose we are given a time $n \in \mathbb{N}$ and an initial law $\zeta \in \mathcal{P}(\mathcal{X})$. The law $\zeta_n = \zeta \Psi_c(M)^n$, which is the law of the Metropolis-Hastings type chain with proposal chain P_0 at time n , solves the optimization problem (22).

If we choose the acceptance function to be $\alpha(x, y) = e^{-\beta(H(y)-H(x))_+}$, then we recover exactly the classical Metropolis-Hastings chain with proposal P_0 and target being the Boltzmann-Gibbs distribution π_β with energy function H and at inverse temperature β .

Other common choices of $\alpha(x, y)$ in the context of MCMC are

- (lazy Metropolis-Hastings): for fixed $\varepsilon \in [0, 1)$, $\alpha(x, y) = (1 - \varepsilon)e^{-\beta(H(y)-H(x))_+}$.
- (Barker proposal Agrawal et al. (2023)): $\alpha(x, y) = \frac{e^{\beta(H(x)-H(y))}}{1 + e^{\beta(H(x)-H(y))}}$.
- (Generalized Barker proposal Agrawal et al. (2023)): for fixed $r \in \mathbb{N}$, $\alpha(x, y) = \frac{\sum_{l=1}^r e^{l\beta(H(x)-H(y))}}{\sum_{l=0}^r e^{l\beta(H(x)-H(y))}}$.
- (Non-reversible Metropolis-Hastings Bierkens (2016)): Fix a skew-symmetric matrix Γ that is known as the vorticity matrix. We take

$$\alpha(x, y) = \min \left\{ 1, \frac{\Gamma(x, y) + e^{-\beta H(y)} P_0(y, x)}{e^{-\beta H(x)} P_0(x, y)} \right\}.$$

Except for the non-reversible Metropolis-Hastings chain, the above Metropolis-Hastings type chains are reversible with respect to π_β .

3.1.2 Example: Glauber dynamics, its marginal transition matrices and the distance to independence

In the second example, we consider the Glauber dynamics or the Gibbs sampler on the product state space $\mathcal{X} = \times_{i=1}^d \mathcal{X}^{(i)}$. The classical Glauber dynamics amounts to choosing uniformly at random a coordinate out of the d coordinates, and then run a Metropolis-Hastings step on the chosen coordinate while fixing the remaining coordinates. For $i \in \llbracket d \rrbracket$, we denote by $P_0^{(i)}$ to be an ergodic and reversible proposal chain with discrete uniform stationary distribution on $\mathcal{X}^{(i)}$. In this example, we specialize into, for $x, y \in \mathcal{X}$,

$$M = \frac{1}{d} \sum_{l=1}^d (\otimes_{i=1}^{l-1} I) \otimes P_0^{(l)} \otimes (\otimes_{i=l+1}^d I), \quad (26)$$

$$c(x, y) = \begin{cases} \beta(H(y) - H(x))_+, & \text{if } x \neq y, \\ \ln M(x, x) - \ln \left(1 - \sum_{z; z \neq x} M(x, z) e^{-\beta(H(z) - H(x))_+} \right), & \text{if } x = y, \end{cases} \quad (27)$$

so that

$$\Psi_c(M)(x, y) = \begin{cases} M(x, y) e^{-\beta(H(y) - H(x))_+}, & \text{if } x \neq y, \\ 1 - \sum_{z; z \neq x} M(x, z) e^{-\beta(H(z) - H(x))_+}, & \text{if } x = y, \end{cases} \quad (28)$$

where the I s are the identity matrices on the appropriate spaces. We thus obtain from Theorem 3.1 and 3.2 variational interpretations of Glauber dynamics as follows:

Corollary 3.4 (Variational interpretation of the transition matrix and of the law of the Glauber dynamics). *Consider the rate distortion optimization problem of M with respect to c and a positive π , where c, M are chosen as in (27) and (26) respectively. $\Psi_c(M)$, the Boltzmann-Gibbs transformation of M with respect to c in (28), is exactly the transition matrix of Glauber dynamics. It is also the unique minimizer of the rate distortion optimization problem of M .*

Suppose we are given a time $n \in \mathbb{N}$ and an initial law $\zeta \in \mathcal{P}(\mathcal{X})$. The law $\zeta_n = \zeta \Psi_c(M)^n$, which is the law of the Glauber dynamics at time n , solves the optimization problem (22).

It is instructive to compute the marginal transition matrix of $\Psi_c(M)$ with respect to π_β , that is, $\Psi_c(M)_{\pi_\beta}^{(i)}$. Using Definition 2.3, we thus see that, for $i \in \llbracket d \rrbracket$, $x^i \neq y^i \in \mathcal{X}^{(i)}$,

$$\begin{aligned} \Psi_c(M)_{\pi_\beta}^{(i)}(x^i, y^i) &= \frac{\sum_{(x^j, y^j); j \neq i} \pi_\beta(x^1, \dots, x^d) \Psi_c(M)((x^1, \dots, x^d), (y^1, \dots, y^d))}{\sum_{x^j; j \neq i} \pi_\beta(x^1, \dots, x^d)} \\ &= \frac{\sum_{x^j; j \neq i} e^{-\beta(H(x^1, \dots, x^i, \dots, x^d) \vee H(x^1, \dots, y^i, \dots, x^d))} \frac{1}{d} P_0^{(i)}(x^i, y^i)}{\sum_{x^j; j \neq i} e^{-\beta H(x^1, \dots, x^d)}} \\ &= \frac{1}{d} P_0^{(i)}(x^i, y^i) \frac{\sum_{x^j; j \neq i} e^{-\beta(H(x^1, \dots, x^i, \dots, x^d) \vee H(x^1, \dots, y^i, \dots, x^d))}}{\sum_{x^j; j \neq i} e^{-\beta H(x^1, \dots, x^d)}}, \end{aligned} \quad (29)$$

where for $a, b \in \mathbb{R}$, $a \vee b := \max\{a, b\}$.

The above expression can be simplified when the energy function H takes on a separable form, that is, there exists functions H_i for $i \in \llbracket d \rrbracket$ such that for all $x \in \mathcal{X}$, it can be written as

$$H(x^1, \dots, x^d) = \sum_{i=1}^d H_i(x^i).$$

If we take, for $i \in \llbracket d \rrbracket$, $x^i, y^i \in \mathcal{X}^{(i)}$,

$$c_i(x^i, y^i) := \begin{cases} \beta(H_i(y^i) - H_i(x^i))_+, & \text{if } x^i \neq y^i, \\ \ln P_0^{(i)}(x^i, x^i) - \ln \left(1 - \sum_{z^i; z^i \neq x^i} P_0^{(i)}(x^i, z^i) e^{-\beta(H_i(z^i) - H_i(x^i))_+} \right), & \text{if } x^i = y^i, \end{cases}$$

then we see that $\Psi_{c_i}(P_0^{(i)})$ is the transition matrix of Metropolis-Hastings with proposal $P_0^{(i)}$ and target energy function H_i at inverse temperature β .

Using the separable form of H , we note that the right hand side of (29) can be simplified using

$$\frac{\sum_{x^j; j \neq i} e^{-\beta(H(x^1, \dots, x^i, \dots, x^d) \vee H(x^1, \dots, y^i, \dots, x^d))}}{\sum_{x^j; j \neq i} e^{-\beta H(x^1, \dots, x^d)}} = e^{-\beta H_i(x^i) \vee H_i(y^i)},$$

and hence we have

$$\Psi_c(M)_{\pi_\beta}^{(i)} = \frac{1}{d} \Psi_{c_i}(P_0^{(i)}) + \frac{d-1}{d} I.$$

As a result, the closest product chain to the Glauber dynamics under the KL divergence with respect to π_β and a separable H is

$$\bigotimes_{i=1}^d \Psi_c(M)_{\pi_\beta}^{(i)} = \bigotimes_{i=1}^d \left(\frac{1}{d} \Psi_{c_i}(P_0^{(i)}) + \frac{d-1}{d} I \right),$$

and the distance to independence of the Glauber dynamics $\Psi_c(M)$ is

$$\mathbb{I}^{\pi_\beta}(\Psi_c(M)) = D_{KL}^{\pi_\beta} \left(\Psi_c(M) \left\| \bigotimes_{i=1}^d \left(\frac{1}{d} \Psi_{c_i}(P_0^{(i)}) + \frac{d-1}{d} I \right) \right. \right).$$

The right spectral gap γ and the log-Sobolev constant α are

$$\begin{aligned} \gamma(\Psi_c(M)_{\pi_\beta}^{(i)}) &= \frac{1}{d} \gamma(\Psi_{c_i}(P_0^{(i)})), \\ \alpha(\Psi_c(M)_{\pi_\beta}^{(i)}) &= \frac{1}{d} \alpha(\Psi_{c_i}(P_0^{(i)})), \end{aligned}$$

and hence, thanks to the contraction principle in Proposition 2.4,

$$\gamma(\Psi_c(M)) \leq \frac{1}{d} \min_{i \in \llbracket d \rrbracket} \gamma(\Psi_{c_i}(P_0^{(i)})),$$

$$\alpha(\Psi_c(M)) \leq \frac{1}{d} \min_{i \in \llbracket d \rrbracket} \alpha(\Psi_{c_i}(P_0^{(i)})).$$

In fact, in the case of Glauber dynamics, its transition matrix is said to be a “product-type”, and by (Levin and Peres, 2017, Corollary 12.13) or (Saloff-Coste, 1997, Lemma 2.2.11) the equalities hold:

$$\begin{aligned} \gamma(\Psi_c(M)) &= \frac{1}{d} \min_{i \in \llbracket d \rrbracket} \gamma(\Psi_{c_i}(P_0^{(i)})), \\ \alpha(\Psi_c(M)) &= \frac{1}{d} \min_{i \in \llbracket d \rrbracket} \alpha(\Psi_{c_i}(P_0^{(i)})). \end{aligned}$$

Therefore, we provide an example of non-tensor-product transition matrix where the spectral gap and log-Sobolev bounds in the contraction principle are attained.

3.1.3 Example: swapping algorithm, its marginal transition matrices and consequences of the contraction principle

In the third example, we investigate the swapping algorithm, an algorithm that incorporates temperature swapping moves to improve the convergence to equilibrium. Much of our exposition in this example follows that in Bhatnagar and Randall (2016). Recall that π_β is the Boltzmann-Gibbs distribution at inverse temperature β . Let $0 =: \beta_1 < \dots < \beta_l := \beta$ be a sequence of inverse temperatures with $l \geq 2$, and we denote by $\mathcal{X}_{sw} := \mathcal{X}^l$, the l products of the original state space \mathcal{X} , to be the state space of the swapping chain. We also write the product distribution to be

$$\pi_{sw} := \bigotimes_{i=1}^l \pi_{\beta_i}.$$

Recall that P_0 is the proposal chain on \mathcal{X} . Let P_{sw} be the transition matrix of the swapping chain.

At each step, the swapping chain chooses an inverse temperature β_i uniformly at random then chooses uniformly at random between the following two moves:

- (Level move): In a level move, the swapping chain chooses an inverse temperature β_i uniformly at random. The swapping chain moves the i th coordinate according to a Metropolis-Hastings chain at inverse temperature β_i , while the remaining coordinates are kept fixed.
- (Swap move): In a swap move, the swapping chain chooses an index $i \in \llbracket l - 1 \rrbracket$ and swaps the coordinate x_i and x_{i+1} with a suitable acceptance probability. Precisely, we have for all $i \in \llbracket l - 1 \rrbracket$, $x = (x_1, \dots, x_l)$, $y = (x_1, \dots, x_{i+1}, x_i, \dots, x_l)$,

$$\begin{aligned} P_{sw}(x, y) &= \frac{1}{2(l-1)} \min \left\{ 1, \frac{\pi_{sw}(y)}{\pi_{sw}(x)} \right\} \\ &= \frac{1}{2(l-1)} \min \left\{ 1, \frac{\pi_{\beta_i}(x_{i+1})\pi_{\beta_{i+1}}(x_i)}{\pi_{\beta_i}(x_i)\pi_{\beta_{i+1}}(x_{i+1})} \right\} \end{aligned}$$

$$= \frac{1}{2(l-1)} e^{-(\beta_{i+1}-\beta_i)(H(x_{i+1})-H(x_i))_+}.$$

In view of the above, we take

$$M(x, y) = \begin{cases} \frac{1}{2l} P_0(x_i, x'_i), & \text{if } x = (x_1, \dots, x_l), y = (x_1, \dots, x'_i, \dots, x_l), i \in \llbracket l \rrbracket, x'_i \neq x_i, \\ \frac{1}{2(l-1)}, & \text{if } x = (x_1, \dots, x_l), y = (x_1, \dots, x_{i+1}, x_i, \dots, x_l), i \in \llbracket l-1 \rrbracket, x_{i+1} \neq x_i, \\ 1 - \sum_{z; z \neq x} M(x, z), & \text{if } x = y, \\ 0, & \text{otherwise,} \end{cases} \quad (30)$$

$$c(x, y) = \begin{cases} \beta_i(H(x'_i) - H(x_i))_+, & \text{if } x = (x_1, \dots, x_l), y = (x_1, \dots, x'_i, \dots, x_l), i \in \llbracket l \rrbracket, \\ (\beta_{i+1} - \beta_i)(H(x_{i+1}) - H(x_i))_+, & \text{if } x = (x_1, \dots, x_l), y = (x_1, \dots, x_{i+1}, x_i, \dots, x_l), i \in \llbracket l-1 \rrbracket, \\ \ln M(x, x) - \ln \left(1 - \sum_{z; z \neq x} M(x, z) e^{-c(x, z)} \right), & \text{if } x = y, \\ 0, & \text{otherwise.} \end{cases} \quad (31)$$

We see that the first (resp. second) equality of c and M correspond to the level (resp. swap) move. These lead to $P_{sw} = \Psi_c(M)$, the Boltzmann-Gibbs transformation of M with respect to the cost function c , which offers a variational interpretation on the transition matrix of the swapping algorithm.

Corollary 3.5 (Variational interpretation of the transition matrix and of the law of the swapping algorithm). *Consider the rate distortion optimization problem of M with respect to c and a positive π , where c, M are chosen as in (31) and (30) respectively. $\Psi_c(M)$, the Boltzmann-Gibbs transformation of M with respect to c , is exactly the transition matrix of the swapping algorithm P_{sw} . It is also the unique minimizer of the rate distortion optimization problem of M .*

Suppose we are given a time $n \in \mathbb{N}$ and an initial law $\zeta \in \mathcal{P}(\mathcal{X}_{sw})$. The law $\zeta_n = \zeta \Psi_c(M)^n$, which is the law of the swapping chain at time n , solves the optimization problem (22) with \mathcal{X} therein replaced by \mathcal{X}_{sw} .

Next, we calculate the marginal transition matrix of $\Psi_c(M)$ with respect to π_{sw} , that is, $\Psi_c(M)_{\pi_{sw}}^{(i)}$. Using Definition 2.3 again, we first note that, for $i = 1$, $x^1 \neq y^1 \in \mathcal{X}$,

$$\begin{aligned} \Psi_c(M)_{\pi_{sw}}^{(1)}(x^1, y^1) &= \frac{\sum_{(x^j, y^j); j \neq 1} \pi_{sw}(x^1, \dots, x^l) \Psi_c(M)((x^1, \dots, x^l), (y^1, \dots, y^l))}{\sum_{x^j; j \neq 1} \pi_{sw}(x^1, \dots, x^l)} \\ &= \frac{1}{2l} P_0(x^1, y^1) e^{-\beta_1(H(y^1) - H(x^1))_+} + \frac{\min \left\{ \pi_{\beta_1}(x^1) \pi_{\beta_2}(y^1), \pi_{\beta_1}(y^1) \pi_{\beta_2}(x^1) \right\}}{2(l-1) \pi_{\beta_1}(x^1)} \\ &= \frac{1}{2l} P_0(x^1, y^1) e^{-\beta_1(H(y^1) - H(x^1))_+} + \frac{e^{\beta_1 H(x^1) - (\beta_1 H(x^1) + \beta_2 H(y^1)) \vee (\beta_1 H(y^1) + \beta_2 H(x^1))}}{2(l-1) \sum_{x \in \mathcal{X}} e^{-\beta_2 H(x)}}, \end{aligned}$$

where the first term corresponds to a level move while the second term corresponds to a swap move. Similarly, for $i = l$, $x^l \neq y^l \in \mathcal{X}$,

$$\begin{aligned} \Psi_c(M)_{\pi_{sw}}^{(l)}(x^l, y^l) &= \frac{1}{2l} P_0(x^l, y^l) e^{-\beta_l(H(y^l) - H(x^l))_+} \\ &\quad + \frac{e^{\beta_l H(x^l) - (\beta_l H(x^l) + \beta_{l-1} H(y^l)) \vee (\beta_l H(y^l) + \beta_{l-1} H(x^l))}}{2(l-1) \sum_{x \in \mathcal{X}} e^{-\beta_{l-1} H(x)}}. \end{aligned}$$

Finally, for $i \in \{2, \dots, l-1\}$, $x^i \neq y^i \in \mathcal{X}$, we calculate that

$$\begin{aligned} \Psi_c(M)_{\pi_{sw}}^{(i)}(x^i, y^i) &= \frac{1}{2l} P_0(x^i, y^i) e^{-\beta_i(H(y^i) - H(x^i))_+} \\ &\quad + \frac{e^{\beta_i H(x^i) - (\beta_i H(x^i) + \beta_{i+1} H(y^i)) \vee (\beta_i H(y^i) + \beta_{i+1} H(x^i))}}{2(l-1) \sum_{x \in \mathcal{X}} e^{-\beta_{i+1} H(x)}} \\ &\quad + \frac{e^{\beta_i H(x^i) - (\beta_i H(x^i) + \beta_{i-1} H(y^i)) \vee (\beta_i H(y^i) + \beta_{i-1} H(x^i))}}{2(l-1) \sum_{x \in \mathcal{X}} e^{-\beta_{i-1} H(x)}}, \end{aligned}$$

where the first term is a level move, the second term is a swap move between i and $i+1$ and the third term is a swap move between i and $i-1$.

If the objective is to sample from π_β , then we can discard the first $l-1$ coordinates of the sample generated by the swapping algorithm and keep only the l th coordinate since $\beta_l = \beta$. An alternative sampler would be the keep- l -in transition matrix $\Psi_c(M)_{\pi_{sw}}^{(l)}$. From the contraction principle in Proposition 2.4, we note that

$$\begin{aligned} \gamma(\Psi_c(M)) &\leq \gamma(\Psi_c(M)_{\pi_{sw}}^{(l)}), \\ \alpha(\Psi_c(M)) &\leq \alpha(\Psi_c(M)_{\pi_{sw}}^{(l)}), \end{aligned}$$

and hence the speed of convergence of $\Psi_c(M)_{\pi_{sw}}^{(l)}$ towards $\pi_{sw}^{(l)}$ is “faster” than that of $\Psi_c(M)$ towards π_{sw} . However, $\Psi_c(M)_{\pi_{sw}}^{(l)}$ is relatively more difficult to simulate than $\Psi_c(M)$ since it involves the normalization constant of the Boltzmann-Gibbs distribution at inverse temperature β_{l-1} , while the swapping algorithm admits a natural and easy-to-simulate procedure.

3.1.4 Example: Feynman-Kac path models

In this example, we look into the Feynman-Kac path models as studied in (Del Moral, 2004, Chapter 1.3). Let \mathbb{P}_n be the joint law of the Markov chain from time 0 to n with transition matrix P , that is,

$$\mathbb{P}_n(x_0, \dots, x_n) := \mathbb{P}(X_0 = x_0, \dots, X_n = x_n).$$

Let $G_l : \mathcal{X} \rightarrow (0, \infty)$ be a sequence of positive potential functions with $l \in \llbracket n \rrbracket$ such that the normalization constant is positive and finite, that is, for any $n \in \mathbb{N}$,

$$\mathcal{Z}_n := \sum_{\mathcal{X}^{n+1}} \left(\prod_{l=0}^{n-1} G_l(x_l) \right) \mathbb{P}_n(x_0, \dots, x_n) \in (0, \infty).$$

The Feynman-Kac path measure of \mathbb{P}_n with respect to the potential functions (G_l) is given by

$$\mathbb{M}_n(x_0, \dots, x_n) := \frac{1}{\mathcal{Z}_n} \left(\prod_{l=0}^{n-1} G_l(x_l) \right) \mathbb{P}_n(x_0, \dots, x_n).$$

Note that both $\mathbb{P}_n, \mathbb{M}_n \in \mathcal{P}(\mathcal{X}^{n+1})$. Now, if we consider the rate distortion optimization problem of \mathbb{P}_n , we see that

$$\mathbb{M}_n \in \arg \min_{\mu \in \mathcal{P}(\mathcal{X}^{n+1})} \left[\tilde{D}_{KL}(\mu \| \mathbb{P}_n) + \mathbb{E}_{(X_0, \dots, X_n) \sim \mu} \left[-\ln \left(\prod_{l=0}^{n-1} G_l(X_l) \right) \right] \right].$$

As a result, we can interpret and understand that the Feynman-Kac path measure \mathbb{M}_n minimizes the rate distortion optimization problem of \mathbb{P}_n with a cost function being the sum of negative logarithm of the potential functions.

3.1.5 Example: simulated annealing

In the fifth example, we consider a discrete simulated annealing chain via Metropolis-Hastings. Unlike previous MCMC algorithms that we have studied in this manuscript thus far, simulated annealing is a non-homogeneous MCMC algorithm. Let $(\beta_l)_{l \in \mathbb{N}}$ be a non-decreasing inverse temperature schedule. Similar to (17) and (18), we set for $n \in \mathbb{N}$ and $x, y \in \mathcal{X}$,

$$c_l(x, y) = \begin{cases} \beta_l(H(y) - H(x))_+, & \text{if } x \neq y, \\ \ln P_0(x, x) - \ln \left(1 - \sum_{z; z \neq x} P_0(x, z) e^{-\beta_l(H(z) - H(x))_+} \right), & \text{if } x = y, \end{cases} \quad (32)$$

$$M_l = P_0, \quad (33)$$

so that

$$\Psi_{c_l}(M_l)(x, y) = \begin{cases} P_0(x, y) e^{-\beta_l(H(y) - H(x))_+}, & \text{if } x \neq y, \\ 1 - \sum_{z; z \neq x} P_0(x, z) e^{-\beta_l(H(z) - H(x))_+}, & \text{if } x = y, \end{cases} \quad (34)$$

Using Theorem 3.2 leads to

Corollary 3.6 (Variational interpretation of the law of the simulated annealing chain at time n). *Suppose we are given a time $n \in \mathbb{N}$ and an initial law $\zeta \in \mathcal{P}(\mathcal{X})$. The law $\zeta_n = \zeta \prod_{l=1}^n \Psi_{c_l}(M_l)$, which is the law of the simulated annealing chain at time n with proposal chain P_0 and inverse temperature schedule $(\beta_l)_{l \in \mathbb{N}}$, solves the optimization problem (22).*

4 Geometry of multivariate Markov chains induced from the information divergence rate

In this section, inspired by the geometric decomposition of [Amari and Han \(1989\)](#) in the context of distributions, we analyze the information geometric structure of multivariate Markov chains arising from the information divergence rate between Markov processes, under the framework established by [Nagaoka \(2005\)](#). For $i \in \llbracket d \rrbracket$, we let $(\mathcal{X}^{(i)}, \mathcal{E}^{(i)})$ where $\mathcal{E}^{(i)} \subset \mathcal{X}^{(i)} \times \mathcal{X}^{(i)}$ be a fully connected digraph. We write $\mathcal{X} = \times_{i=1}^d \mathcal{X}^{(i)}$, $\mathcal{E} = \times_{i=1}^d \mathcal{E}^{(i)}$, and denote $\mathcal{L}(\mathcal{X}, \mathcal{E})$ the set of transition matrices over the graph $(\mathcal{X}, \mathcal{E})$. Observe that by construction, all members of $\mathcal{L}(\mathcal{X}, \mathcal{E})$ are irreducible. For $P \in \mathcal{L}(\mathcal{X}, \mathcal{E})$, we say that P is a product chain when there exist a collection $P_i \in \mathcal{L}(\mathcal{X}^{(i)}, \mathcal{E}^{(i)})$ for $i \in \llbracket d \rrbracket$ such that for any

$$(x, x') = ((x_1, x_2, \dots, x_d), (x'_1, x'_2, \dots, x'_d)) \in \mathcal{X} \times \mathcal{X}$$

it holds that

$$P[x, x'] = \prod_{i=1}^d P_i(x_i, x'_i).$$

We write $\mathcal{L}_{\otimes}(\mathcal{X}, \mathcal{E}) \subset \mathcal{L}(\mathcal{X}, \mathcal{E})$ for the subset consisting of all product chains, and we denote $\mathcal{Q}(\mathcal{X}, \mathcal{E}) = \{\pi \boxtimes P : P \in \mathcal{L}(\mathcal{X}, \mathcal{E}), \pi P = \pi\}$ the set of all edge measures. Transition matrices will be denoted P , while the corresponding edge measure will typically be denoted $Q := \pi \boxtimes P$.

4.1 Information geometry of Markov chains

Recall that the information divergence rate between two Markov chains X_1, X_2, \dots and X'_1, X'_2, \dots sampled respectively according to $P \in \mathcal{L}(\mathcal{X}, \mathcal{E})$ and $P' \in \mathcal{L}(\mathcal{X}, \mathcal{E})$ can be expressed as

$$D(P||P') := \lim_{n \rightarrow \infty} \frac{1}{n} D(X_1, \dots, X_n || X'_1, \dots, X'_n) = \sum_{x, x' \in \mathcal{E}} \pi(x) P(x, x') \ln \frac{P(x, x')}{P'(x, x')},$$

where π is the stationary distribution of P . In this section, we consider the dually flat geometry on $\mathcal{L}(\mathcal{X}, \mathcal{E})$ induced from the information divergence rate D , introduced by [Nagaoka \(2005\)](#). We let \mathfrak{s} be the mapping $\mathcal{E}^{\mathbb{R}} \rightarrow \mathcal{L}(\mathcal{X}, \mathcal{E})$, which for a function $f \in \mathcal{E}^{\mathbb{R}}$ outputs the transition matrix

$$\mathfrak{s}(f)(x, x') = \begin{cases} f(x, x') \frac{v(x')}{\rho v(x)} & \text{when } x, x' \in \mathcal{E}, \\ 0 & \text{otherwise,} \end{cases}$$

and where v and ρ are respectively the right Perron-Frobenius eigenvector and root of the linear operator defined by $(fu)(x) = \sum_{x': (x, x') \in \mathcal{E}} f(x, x') u(x')$.

Definition 4.1 (e-family ([Nagaoka, 2005](#))). *We say that $\mathcal{V}_e = \{P_\theta : \theta \in \mathbb{R}^d\}$ is an exponential family (e-family) of Markov chains (or transition matrices), when there exist functions*

$\psi: \mathbb{R}^d \rightarrow \mathbb{R}$, $K: \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$, $g_i: \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ for $i \in \llbracket d \rrbracket$ and $V: \mathbb{R}^d \times \mathcal{X} \rightarrow \mathbb{R}$, such that for any $\theta \in \mathbb{R}^d$,

$$P_\theta(x, x') = \exp \left(K(x, x') + \sum_{i=1}^d \theta^i g_i(x, x') + V(\theta, x') - V(\theta, x) - \psi(\theta) \right),$$

when $(x, x') \in \mathcal{E}$, and $P_\theta(x, x') = 0$ otherwise.

Definition 4.2 (m-family ([Hayashi and Watanabe, 2016](#))). We say that $\mathcal{V}_m = \{P_\xi: \xi \in \Xi \subset \mathbb{R}^d\}$ is a mixture family (m-family) of Markov chains (or transition matrices) when there exist edge measures $Q_0, Q_1, \dots, Q_d \in \mathcal{Q}(\mathcal{X}, \mathcal{E})$ such that for any $\xi \in \Xi$, the edge measure Q_ξ of P_ξ can be expressed as

$$Q_\xi(x, x') = \left(1 - \sum_{i=1}^d \xi^i \right) Q_0(x, x') + \sum_{i=1}^d \xi^i Q_i(x, x'),$$

and $\Xi = \{\xi \in \mathbb{R}^d: Q_\xi \in \mathcal{Q}(\mathcal{X}, \mathcal{E}), \forall (x, x') \in \mathcal{E}\}$.

In particular, $\mathcal{L}(\mathcal{X}, \mathcal{E})$ forms both an e-family and an m-family ([Nagaoka, 2005](#)). For example, it is known that reversible Markov chains form both an m-family and e-family ([Wolfer and Watanabe, 2021](#)). For a more comprehensive introduction to information geometry of Markov chains, we refer the reader to the recent survey of [Wolfer and Watanabe \(2023\)](#).

Remark 4.1. The information divergence rate D considered in this section agrees with D_{KL} introduced in [Definition 2.1](#) on the set of π -stationary transition matrices irreducible over $(\mathcal{X}, \mathcal{E})$.

4.2 The exponential family of product chains

Remarkably, product Markov chains form an e-family in multivariate Markov chains.

Theorem 4.1. $\mathcal{L}_\otimes(\mathcal{X}, \mathcal{E})$ forms an e-family in $\mathcal{L}(\mathcal{X}, \mathcal{E})$.

Proof. We let $P_0, P_1 \in \mathcal{L}_\otimes(\mathcal{X}, \mathcal{E})$ and let $t \in \mathbb{R}$. We construct

$$P_t[x, x'] := \frac{P_0[x, x']^{1-t} P_1[x, x']^t v_t(x')}{v_t(x) \rho_t},$$

and for $i \in \llbracket d \rrbracket$,

$$P_{t,i}(x_i, x'_i) := \frac{P_{0,i}(x_i, x'_i)^{1-t} P_{1,i}(x_i, x'_i)^t v_{t,i}(x'_i)}{v_{t,i}(x_i) \rho_{t,i}},$$

where for any $x, x' \in \mathcal{X}$, by factorization,

$$P_0[x, x'] = \prod_{i=1}^d P_{0,i}(x_i, x'_i), \quad P_1[x, x'] = \prod_{i=1}^d P_{1,i}(x_i, x'_i),$$

for two collections of transition matrices $P_{0,i}, P_{1,i}, i \in \llbracket d \rrbracket$, and where for $i \in \llbracket d \rrbracket$, $(v_{t,i}, \rho_{t,i})$, and (v_t, ρ_t) are the Perron-Frobenius right eigenvector and root of respectively

$$(P_{0,i})^{\circ 1-t} (P_{1,i})^{\circ t}, \quad P_0^{\circ 1-t} P_1^{\circ t},$$

with \circ denoting here the Hadamard product and induced power. On one hand,

$$\begin{aligned} P_t[x, x'] &= \frac{P_0[x, x']^{1-t} P_1[x, x']^t v_t(x')}{v_t(x) \rho_t} \\ &= \frac{\left(\prod_{i=1}^d P_{0,i}(x_i, x'_i)^{1-t} \right) \left(\prod_{i=1}^d P_{1,i}(x_i, x'_i)^t \right) v_t(x')}{v_t(x) \rho_t} \end{aligned}$$

and on the other hand, constructing the product chain from the $P_{t,i}$,

$$\left(\bigotimes_{i=1}^d P_{t,i} \right) [x, x'] := \prod_{i=1}^d P_{t,i}(x_i, x'_i) = \frac{\left(\prod_{i=1}^d P_{0,i}(x_i, x'_i)^{1-t} \right) \left(\prod_{i=1}^d P_{1,i}(x_i, x'_i)^t \right) \left(\prod_{i=1}^d v_{t,i}(x'_i) \right)}{\left(\prod_{i=1}^d v_{t,i}(x_i) \right) \left(\prod_{i=1}^d \rho_{t,i} \right)}.$$

As a result,

$$\bigotimes_{i=1}^d P_{t,i} = \mathfrak{s} \left(\left[\bigotimes_{i=1}^d (P_{0,i})^{\circ 1-t} \right] \circ \left[\bigotimes_{i=1}^d (P_{1,i})^{\circ t} \right] \right) = P_t,$$

thus $P_t \in \mathcal{L}_{\otimes}(\mathcal{X}, \mathcal{E})$ and $\mathcal{L}_{\otimes}(\mathcal{X}, \mathcal{E})$ forms an e-family. \square

For $P \in \mathcal{L}(\mathcal{X}, \mathcal{E})$ a multivariate transition matrix, it thus makes sense to define the m-projection onto $\mathcal{L}_{\otimes}(\mathcal{X}, \mathcal{E})$ as

$$P_m := \arg \min_{\bar{P} \in \mathcal{L}_{\otimes}(\mathcal{X}, \mathcal{E})} D(P || \bar{P}),$$

which can be understood as the closest product chain to P . Furthermore, the e-family structure of \mathcal{L}_{\otimes} enables us to state readily that the following Pythagorean theorem holds ([Hayashi and Watanabe, 2016](#), Corollary 4.8), ([Nagaoka, 2005](#), (23)) for any $\bar{P} \in \mathcal{L}_{\otimes}(\mathcal{X}, \mathcal{E})$,

$$D(P || \bar{P}) = D(P || P_m) + D(P_m || \bar{P}).$$

In fact, we derive an explicit expression for the m-projection P_m in Proposition 4.1. Let $P \in \mathcal{L}_{\otimes}(\mathcal{X}, \mathcal{E})$ with factors P_i and respective stationary distributions π_i and edge measures Q_i . Then the stationary distribution and edge measure of P are respectively given by

$$\pi = \bigotimes_{i=1}^d \pi_i, \quad Q = \bigotimes_{i=1}^d Q_i.$$

Proposition 4.1. *The m -projection P_m of $P \in \mathcal{L}(\mathcal{X}, \mathcal{E})$ onto $\mathcal{L}_\otimes(\mathcal{X}, \mathcal{E})$ can be factored as*

$$P_m = \otimes_{i=1}^d P^{(i)},$$

where

$$P^{(i)}(x_i, x'_i) = P_\pi^{(i)}(x_i, x'_i) = \frac{\sum_{\substack{j=1 \\ j \neq i}}^d \sum_{x_j, x'_j \in \mathcal{X}^{(j)}} Q[(x_1, \dots, x_d), (x'_1, \dots, x'_d)]}{\sum_{\substack{j=1 \\ j \neq i}}^d \sum_{x_j \in \mathcal{X}^{(j)}} \pi[(x_1, \dots, x_d)]},$$

and where π and Q are the stationary distribution and edge measure of P .

Proof. Reorganizing the Pythagorean theorem, the claims reduces to finding a solution for P_m satisfying the following equality,

$$\sum_{x, x' \in \mathcal{X}} (Q[x, x'] - Q_m[x, x']) \ln \frac{P_m[x, x']}{\bar{P}[x, x']} = 0,$$

where Q_m is the edge measure of P_m . Since P_m and \bar{P} are product chains, we can rewrite,

$$\sum_{(x_1, \dots, x_d), (x'_1, \dots, x'_d) \in \mathcal{X}} \left(Q[(x_1, \dots, x_d), (x'_1, \dots, x'_d)] - \prod_{j=1}^d Q_{m,j}(x_j, x'_j) \right) \sum_{i=1}^d \ln \frac{P_{m,i}(x_i, x'_i)}{\bar{P}_i(x_i, x'_i)} = 0,$$

and in turn,

$$\sum_{i=1}^d \sum_{x_i, x'_i \in \mathcal{X}^{(i)}} \left[\left(\sum_{\substack{j=1 \\ j \neq i}}^d \sum_{x_j, x'_j \in \mathcal{X}^{(j)}} Q[(x_1, \dots, x_d), (x'_1, \dots, x'_d)] \right) - Q_{m,i}(x_i, x'_i) \right] \ln \frac{P_{m,i}(x_i, x'_i)}{\bar{P}_i(x_i, x'_i)} = 0.$$

The equation is verified when

$$Q_{m,i}(x_i, x'_i) = Q^{(i)}(x_i, x'_i) := \sum_{\substack{j=1 \\ j \neq i}}^d \sum_{x_j, x'_j \in \mathcal{X}^{(j)}} Q[(x_1, \dots, x_d), (x'_1, \dots, x'_d)]. \quad (35)$$

We verify that the $Q^{(i)}$ are indeed edge measures. For any $i \in \llbracket d \rrbracket$, by marginalization of $Q^{(i)}$,

$$\begin{aligned} \sum_{x'_i \in \mathcal{X}^{(i)}} Q^{(i)}(x_i, x'_i) &= \sum_{x'_i \in \mathcal{X}^{(i)}} \sum_{\substack{j=1 \\ j \neq i}}^d \sum_{x_j, x'_j \in \mathcal{X}^{(j)}} Q[(x_1, \dots, x_d), (x'_1, \dots, x'_d)] \\ &= \sum_{\substack{j=1 \\ j \neq i}}^d \sum_{x_j \in \mathcal{X}^{(j)}} \sum_{(x'_1, \dots, x'_d) \in \mathcal{X}} Q[(x_1, \dots, x_d), (x'_1, \dots, x'_d)] \\ &= \sum_{\substack{j=1 \\ j \neq i}}^d \sum_{x_j \in \mathcal{X}^{(j)}} \pi[(x_1, \dots, x_d)], \end{aligned}$$

and similarly,

$$\sum_{x_i \in \mathcal{X}^{(i)}} Q^{(i)}(x_i, x'_i) = \sum_{\substack{j=1 \\ j \neq i}}^d \sum_{x'_j \in \mathcal{X}^{(j)}} \pi[(x'_1, \dots, x'_d)] = \sum_{x'_i \in \mathcal{X}^{(i)}} Q^{(i)}(x_i, x'_i).$$

As a consequence, $Q^{(i)}$ is an edge measure with $\pi^{(i)}[(x_1, \dots, x_d)] = \sum_{\substack{j=1 \\ j \neq i}}^d \sum_{x_j \in \mathcal{X}^{(j)}} \pi[(x_1, \dots, x_d)]$, and for any $x, x' \in \mathcal{X}$,

$$P^{(i)}(x_i, x'_i) = \frac{Q^{(i)}(x_i, x'_i)}{\pi^{(i)}(x'_i)},$$

whence the claim. \square

The distance of a multivariate Markov chain from product chains is thus given by the following equality as a difference of entropy rates.

Lemma 4.1. *Let $P \in \mathcal{L}(\mathcal{X}, \mathcal{E})$ and P_m is the m -projection of P onto $\mathcal{L}_{\otimes}(\mathcal{X}, \mathcal{E})$. It holds that*

$$\mathbb{I}(P) = D(P||P_m) = \sum_{i=1}^d H(P^{(i)}) - H(P),$$

where $H(P) := -\sum_{x, x' \in \mathcal{X}} Q[x, x'] \ln P[x, x']$ is the Shannon entropy rate of P , and $\mathbb{I}(P)$ can be understood as a measure of distance from independence¹ to P .

Proof.

$$\begin{aligned} D(P||P_m) &= \sum_{x, x' \in \mathcal{X}} Q[x, x'] \ln \frac{P[x, x']}{P_m[x, x']} \\ &= \sum_{(x_1, \dots, x_d), (x'_1, \dots, x'_d) \in \mathcal{X}} Q[(x_1, \dots, x_d), (x'_1, \dots, x'_d)] \sum_{i=1}^d \ln \frac{1}{P^{(i)}(x_i, x'_i)} - H(P) \\ &= \sum_{i=1}^d \left(\sum_{x_i, x'_i \in \mathcal{X}^{(i)}} \left[\sum_{\substack{j=1 \\ j \neq i}}^d \sum_{x_j, x'_j \in \mathcal{X}^{(j)}} Q[(x_1, \dots, x_d), (x'_1, \dots, x'_d)] \right] \ln \frac{1}{P^{(i)}(x_i, x'_i)} \right) - H(P) \\ &= \sum_{i=1}^d \left(\sum_{x_i, x'_i \in \mathcal{X}^{(i)}} Q^{(i)}(x_i, x'_i) \ln \frac{1}{P^{(i)}(x_i, x'_i)} \right) - H(P) \\ &= \sum_{i=1}^d H(P^{(i)}) - H(P). \end{aligned}$$

\square

¹This distance from independence \mathbb{I} will coincide with \mathbb{I}^π given in (7) on the set of π -stationary transition matrices.

4.3 The m-family of multivariate chains with prescribed marginal edge measures

Denote $\mathfrak{q}^{(i)}: \mathcal{L}(\mathcal{X}, \mathcal{E}) \rightarrow \mathcal{Q}(\mathcal{X}^{(i)}, \mathcal{E}^{(i)})$ the map which associates to a multivariate Markov chain its marginal edge measure on $(\mathcal{X}^{(i)}, \mathcal{E}^{(i)})$ established in (35), that is

$$\mathfrak{q}^{(i)}(P)(x_i, x'_i) = \sum_{\substack{j=1 \\ j \neq i}}^d \sum_{x_j, x'_j \in \mathcal{X}^{(j)}} Q[(x_1, \dots, x_d), (x'_1, \dots, x'_d)],$$

where Q pertains to P . For $i \in \llbracket d \rrbracket$, we let $Q_{\odot, i} \in \mathcal{Q}(\mathcal{X}^{(i)}, \mathcal{E}^{(i)})$ be prescribed edge measures, and we denote

$$\mathcal{K}(Q_{\odot, 1}, \dots, Q_{\odot, d}) := \{P \in \mathcal{L}(\mathcal{X}) : \forall i \in \llbracket d \rrbracket, \mathfrak{q}^{(i)}(P) = Q_{\odot, i}\},$$

the set of all multivariate chains with prescribed marginals.

Theorem 4.2. $\mathcal{K}(Q_{\odot, 1}, \dots, Q_{\odot, d})$ forms an m-family in $\mathcal{L}(\mathcal{X}, \mathcal{E})$.

Proof. Let $P_0, P_1 \in \mathcal{K}(Q_{\odot, 1}, \dots, Q_{\odot, d})$, with respective edge measures Q_0, Q_1 . We define $Q_t = (1-t)Q_0 + tQ_1$ for any suitable t . Denote P_t the corresponding transition matrix. For $i \in \llbracket d \rrbracket$, we compute

$$\begin{aligned} \mathfrak{q}^{(i)}(P_t) &= \sum_{\substack{j=1 \\ j \neq i}}^d \sum_{x_j, x'_j \in \mathcal{X}^{(j)}} Q_t[(x_1, \dots, x_d), (x'_1, \dots, x'_d)] \\ &= (1-t) \sum_{\substack{j=1 \\ j \neq i}}^d \sum_{x_j, x'_j \in \mathcal{X}^{(j)}} Q_0[(x_1, \dots, x_d), (x'_1, \dots, x'_d)] \\ &\quad + t \sum_{\substack{j=1 \\ j \neq i}}^d \sum_{x_j, x'_j \in \mathcal{X}^{(j)}} Q_1[(x_1, \dots, x_d), (x'_1, \dots, x'_d)] \\ &= (1-t)\mathfrak{q}^{(i)}(P_0) + t\mathfrak{q}^{(i)}(P_1) = (1-t)Q_{\odot, i}[x, x'] + tQ_{\odot, i}[x, x'] = Q_{\odot, i}[x, x']. \end{aligned}$$

As a result, $P_t \in \mathcal{K}(Q_{\odot, 1}, \dots, Q_{\odot, d})$ and the latter forms an m-family. \square

For $P \in \mathcal{L}(\mathcal{X}, \mathcal{E})$ a general multivariate transition matrix, it therefore makes sense to talk about the e-projection onto $\mathcal{K}(Q_{\odot, 1}, \dots, Q_{\odot, d})$, defined as

$$P_e := \arg \min_{\bar{P} \in \mathcal{K}(Q_{\odot, 1}, \dots, Q_{\odot, d})} D(\bar{P} \| P).$$

Furthermore, the following Pythagorean theorem is known to hold (Hayashi and Watanabe, 2016, Corollary 4.7), (Nagaoka, 2005, (23)) for any $\bar{P} \in \mathcal{K}(Q_{\odot, 1}, \dots, Q_{\odot, d})$,

$$D(\bar{P} \| P) = D(\bar{P} \| P_e) + D(P_e \| P).$$

4.4 Resulting geodesic foliation

In fact, we can endow multivariate transitions matrices with a mutually dual foliation.

Theorem 4.3 (m-geodesic foliation of multivariate Markov chains).

$$\mathcal{L}(\mathcal{X}, \mathcal{E}) = \bigsqcup_{\otimes_{i=1}^d P_{\odot,i} \in \mathcal{L}_{\otimes}(\mathcal{X}, \mathcal{E})} \mathcal{K}(Q_{\odot,1}, \dots, Q_{\odot,d}),$$

where for any $i \in \llbracket d \rrbracket$, $Q_{\odot,i}$ is the edge measure pertaining to the transition matrix $P_{\odot,i}$.

Proof. The fact that the leaves are disjoint and exhaust $\mathcal{L}(\mathcal{X}, \mathcal{E})$ is easy to see. Mutual duality stems from $\mathcal{L}_{\otimes}(\mathcal{X}, \mathcal{E})$ being an e-family and $\mathcal{K}(Q_{\odot,1}, \dots, Q_{\odot,d})$ be an m-family for any $\otimes_{i=1}^d P_{\odot,i} \in \mathcal{L}_{\otimes}(\mathcal{X}, \mathcal{E})$. \square

Example 4.1. We let $A \in \mathbb{R}^{\mathcal{X}^2}$ be the adjacency matrix defined by

$$A(x, x') = \begin{cases} 1 & \text{when } (x, x') \in \mathcal{E} \\ 0 & \text{otherwise,} \end{cases}$$

from which the stochastic matrix

$$U := \frac{1}{\rho_A} \text{diag}(v_A)^{-1} A \text{diag}(v_A),$$

is induced, where v_A, ρ_A is the right Perron-Frobenius pair associated with A . It is well-known that U is max-entropic in the sense where it achieves the maximum entropy rate for the connection graph $(\mathcal{X}, \mathcal{E})$ ([Burda et al., 2009](#)). Furthermore, $H(U) = \ln \rho_A$, and the e-projection of U onto the m-family $\mathcal{K}(Q_{\odot,1}, \dots, Q_{\odot,d})$ satisfies ([Csiszár et al., 1987](#))

$$U_e := \arg \min_{P' \in \mathcal{K}(Q_{\odot,1}, \dots, Q_{\odot,d})} D(P' || U) = \arg \max_{P' \in \mathcal{K}(Q_{\odot,1}, \dots, Q_{\odot,d})} H(P').$$

In other words, the e-projection of U is obtained from the maximum entropy principle under prescribed marginal edge-measures, which amounts to optimizing a convex function over a convex domain. Observe that since

$$A(x, x') = \prod_{i=1}^d A_i(x_i, x'_i),$$

where A_i is the adjacency matrix for $(\mathcal{X}^{(i)}, \mathcal{E}^{(i)})$, it follows that $U \in \mathcal{L}_{\otimes}(\mathcal{X}, \mathcal{E})$ and we can write $U = \otimes_{i=1}^d U_i$ with $U_i = \mathfrak{s}(A_i)$ for $i \in \llbracket d \rrbracket$. As a result, it also holds that $U_e \in \mathcal{L}_{\otimes}(\mathcal{X}, \mathcal{E})$, and in fact,

$$U_e = \bigotimes_{i=1}^d P_{0,i},$$

where $P_{0,i}$ is the transition matrix associated with the edge measure $Q_{\odot,i}$. On [Figure 5](#), this is represented by projecting U along the e-family $\mathcal{L}_{\otimes}(\mathcal{X}, \mathcal{E})$ and onto the leaf induced by the prescribed edge measures.




Figure 5: Illustration of the m-foliation of $\mathcal{L}(\mathcal{X}, \mathcal{E})$ with respect to the Bregman geometry induced by D . The cylinder represents the family of transition matrices at a given distance from the independent e-family $\mathcal{L}_{\otimes}(\mathcal{X}, \mathcal{E})$. Each leaf \mathcal{K} determined by a prescribed collection of marginal edge-measures forms an m-family. In particular, P_{\perp} is the e-projection of P onto $\mathcal{L}_{\otimes}(\mathcal{X}, \mathcal{E})$, the m-projection of \bar{P} onto $\mathcal{K}(Q_{r,1}, \dots, Q_{r,d})$ and the Pythagorean identity holds between them. The e-projection of the max-entropic Markov chain U onto $\mathcal{K}(Q_{3,1}, \dots, Q_{3,d})$ is represented by U_e (refer to Example 4.1 for their definition).

Similar to the decomposition presented by [Amari and Han \(1989\)](#) in the context of distributions, which leads to the analysis of multiterminal hypothesis testing by [Watanabe \(2017\)](#), we surmise that our geometric decomposition can be leveraged to provide an analysis in the context of a multiterminal Markovian setting. We leave this application as an exciting open problem.

4.5 Partial projections and the Han–Shearer entropy gap

The observation that $\mathcal{L}_{\otimes}(\mathcal{X}, \mathcal{E})$ forms an e-family gives rise to a collection of hierarchies.

Corollary 4.1 (to Theorem 4.1). *Let $S \subset \llbracket d \rrbracket$, it holds that*

$$\mathcal{L}_{\otimes}(\mathcal{X}, \mathcal{E}) \stackrel{e\text{-family}}{\subset} \mathcal{L}_{S \otimes (\llbracket d \rrbracket \setminus S)}(\mathcal{X}, \mathcal{E}) \stackrel{e\text{-family}}{\subset} \mathcal{L}(\mathcal{X}, \mathcal{E}),$$

where

$$\mathcal{L}_{\mathcal{S} \otimes ([d] \setminus \mathcal{S})}(\mathcal{X}, \mathcal{E}) = \left\{ P \in \mathcal{L}(\mathcal{X}, \mathcal{E}) : \exists P_S \in \mathcal{L}(\mathcal{X}^{(S)}, \mathcal{E}^{(S)}), \right. \\ \left. P_{-S} \in \mathcal{L}(\mathcal{X}^{(-S)}, \mathcal{E}^{(-S)}), P = P_S \otimes P_{-S} \right\}.$$

More generally, for two partitions of $[d]$, $\mathcal{R} = \{R_1, \dots, R_q\}$ and $\mathcal{S} = \{S_1, \dots, S_n\}$, when \mathcal{S} is a refinement² of \mathcal{R} , it holds that

$$\mathcal{L}_{\otimes_{i=1}^n S_i}(\mathcal{X}, \mathcal{E}) \stackrel{e\text{-family}}{\subset} \mathcal{L}_{\otimes_{i=1}^q R_i}(\mathcal{X}, \mathcal{E}).$$

As a result, for any subset S of the coordinates, one can thus consider the (orthogonal) m-projection onto the factorizable set $\mathcal{L}_{\mathcal{S} \otimes ([d] \setminus \mathcal{S})}(\mathcal{X}, \mathcal{E})$, whose expression is given by

$$P_m = P^{(S)} \otimes P^{(-S)},$$

where

$$P^{(S)}(x^{(S)}, x'^{(S)}) = \frac{\sum_{j \notin S} \sum_{x_j, x'_j \in \mathcal{X}^{(j)}} Q[(x_1, \dots, x_d), (x'_1, \dots, x'_d)]}{\sum_{j \notin S} \sum_{x_j \in \mathcal{X}^{(j)}} \pi[(x_1, \dots, x_d)]}.$$

In particular, for two partitions of $[d]$, $\mathcal{R} = \{R_1, \dots, R_q\}$ and $\mathcal{S} = \{S_1, \dots, S_n\}$, such that \mathcal{S} is a refinement of \mathcal{R} , if we denote $P_{\mathcal{S}, m}$ and $P_{\mathcal{R}, m}$ the m-projection of $P \in \mathcal{L}(\mathcal{X}, \mathcal{E})$ onto respectively $\mathcal{L}_{\otimes_{i=1}^n S_i}(\mathcal{X}, \mathcal{E})$ and $\mathcal{L}_{\otimes_{i=1}^q R_i}(\mathcal{X}, \mathcal{E})$, then for any $\bar{P} \in \mathcal{L}_{\otimes_{i=1}^n S_i}(\mathcal{X}, \mathcal{E})$, the following hierarchical decomposition (see Figure 6) is satisfied.

$$D(P || \bar{P}) = D(P || P_{\mathcal{R}, m}) + D(P_{\mathcal{R}, m} || P_{\mathcal{S}, m}) + D(P_{\mathcal{S}, m} || \bar{P}) \\ = D\left(P \left\| \bigotimes_{R_i \subset \mathcal{R}} P_{\mathcal{R}, m}^{(R_i)}\right.\right) + \sum_{i=1}^q D\left(P_{\mathcal{R}, m}^{(R_i)} \left\| \bigotimes_{S_j \in \mathcal{S}, S_j \subset R_i} P_{\mathcal{S}, m}^{(S_j)}\right.\right) + \sum_{i=1}^n D\left(P_{\mathcal{S}, m}^{(S_i)} \left\| \bar{P}^{(S_i)}\right.\right).$$

In the remainder of this section, we explore connections between the entropy rate of a transition matrix P and the entropy rates of factors of its m-projections onto factorizable subsets. Let $\mathcal{S} = \{S_1, \dots, S_n\}$, $S_i \subset [d]$ such that every integer in $[d]$ lies in at least r of these subsets. For a probability distribution $\mu \in \mathcal{P}(\mathcal{X} = \times_{i=1}^d \mathcal{X}^{(i)})$, we define the Han-Shearer entropy gap as with respect to \mathcal{S} and r as,

$$\Gamma_{\text{HS}}^{(\mathcal{S}, r)}(\mu) := \frac{1}{r} \sum_{i=1}^n H(\mu^{(S_i)}) - H(\mu). \quad (36)$$

²A partition \mathcal{S} is a refinement of a partition \mathcal{R} if every element of \mathcal{S} is a subset of some element of \mathcal{R} .




Figure 6: Hierarchical decomposition of divergence in terms of orthogonal projections with respect a partition \mathcal{R} and its refinement \mathcal{S} .

Han’s inequality (Han, 1978) and Shearer’s lemma (Chung et al., 1986) essentially states that the entropy gap $\Gamma_{\text{HS}}^{(\mathcal{S},r)}(\mu)$ is non-negative. Similarly, we define a version of the Han–Shearer entropy gap in the context of transition matrices.

$$\Gamma_{\text{HS}}^{(\mathcal{S},r)}(P) := \frac{1}{r} \sum_{i=1}^n H(P^{(S_i)}) - H(P). \quad (37)$$

We draw a connection between the entropy gap of a transition matrix and the gaps pertaining to its edge measure and stationary distribution. In particular, we show that, at least when the stationary distribution is a product, the entropy rate of a transition matrix P is smaller than or equal to the “average” entropy rate of the leave-one-out transition matrices, thereby recovering a Han-type inequality for transition matrices.

Proposition 4.2 (Han–Shearer inequality for the entropy rate of transition matrices). *Let $P \in \mathcal{L}(\mathcal{X}, \mathcal{E})$ with respective edge-measure Q and stationary distribution π . Let $\mathcal{S} = S_1, \dots, S_n \subset \llbracket d \rrbracket$ such that every integer in $\llbracket d \rrbracket$ lies in at least r of these subsets. It holds that*

$$\Gamma_{\text{HS}}^{(\mathcal{S},r)}(P) = \Gamma_{\text{HS}}^{(\mathcal{S},r)}(Q) - \Gamma_{\text{HS}}^{(\mathcal{S},r)}(\pi),$$

where $\Gamma_{\text{HS}}^{(\mathcal{S},r)}(P)$ is defined in (37) and $\Gamma_{\text{HS}}^{(\mathcal{S},r)}(Q), \Gamma_{\text{HS}}^{(\mathcal{S},r)}(\pi)$ are defined in (36). In particular, when $\pi = \otimes_{i=1}^d \pi_i$ for $\pi_i \in \mathcal{P}(\mathcal{X}^{(i)})$,

$$\Gamma_{\text{HS}}^{(\mathcal{S},r)}(P) \geq 0.$$

Proof. It suffices to observe that from the chain rule of conditional entropy

$$H(P) = H(Q) - H(\pi).$$

As a consequence,

$$\Gamma_{\text{HS}}^{(S,r)}(Q) = \frac{1}{r} \sum_{i=1}^n (H(P^{(S_i)}) + H(\pi^{(S_i)})) - H(P) - H(\pi) = \Gamma_{\text{HS}}^{(S,r)}(P) + \Gamma_{\text{HS}}^{(S,r)}(\pi).$$

In particular, when $\pi = \otimes_{i=1}^d \pi_i$ for $\pi_i \in \mathcal{P}(\mathcal{X}^{(i)})$, $\Gamma_{\text{HS}}^{(S,r)}(\pi) = 0$, thus $\Gamma_{\text{HS}}^{(S,r)}(P) \geq 0$. \square

The rest of this subsection is devoted to prove that the mappings $S \mapsto H(P^{(S)})$ and $S \mapsto D(P \parallel P^{(S)} \otimes P^{(-S)})$ are submodular in S . These two properties are analogous to the counterpart properties of the Shannon entropy and the mutual information of random variables: they are respectively monotone submodular and submodular functions (see for example (Polyanskiy and Wu, 2022, Chapter 1.4, 1.5)). Note that in the i.i.d. case, the submodularity of Shannon entropy implies the Han's inequality via the notion of self-bounding function, see for example Sason (2022, Corollary 2, (29)).

Proposition 4.3. *Let $S \subseteq \llbracket d \rrbracket$. Let $P \in \mathcal{L}(\mathcal{X}, \mathcal{E})$ with respective edge-measure Q and a product stationary distribution $\pi = \otimes_{i=1}^d \pi^{(i)}$. We have*

1. (Submodularity of the entropy rate of P) *The mapping $S \mapsto H(P^{(S)})$ is submodular.*
2. (Submodularity of the distance to $(S, \llbracket d \rrbracket \setminus S)$ -factorizability of P) *The mapping $S \mapsto D(P \parallel P^{(S)} \otimes P^{(-S)})$ is submodular, where we recall that $D(P \parallel P^{(S)} \otimes P^{(-S)})$ is the distance to $(S, \llbracket d \rrbracket \setminus S)$ -factorizability of P in Section 2.4.*
3. (Supermodularity and monotonicity of the distance to independence) *The mapping $S \mapsto \mathbb{I}(P^{(S)})$ is monotone and supermodular.*

Proof. We first prove item (1). Let $S \subseteq T \subseteq \llbracket d \rrbracket$ and suppose $i \in \llbracket d \rrbracket \setminus T$. By the definition of submodularity, it suffices to show that

$$H(P^{(S \cup \{i\})}) - H(P^{(S)}) \geq H(P^{(T \cup \{i\})}) - H(P^{(T)}).$$

Using the chain rule of conditional entropy $H(P) = H(Q) - H(\pi)$ leads to

$$H(Q^{(S \cup \{i\})}) - H(Q^{(S)}) - H(\pi^{(S \cup \{i\})}) + H(\pi^{(S)}) \geq H(Q^{(T \cup \{i\})}) - H(Q^{(T)}) - H(\pi^{(T \cup \{i\})}) + H(\pi^{(T)}).$$

Now, we use the assumption that $\pi = \otimes_{i=1}^d \pi^{(i)}$ to see that $H(\pi^{(S \cup \{i\})}) - H(\pi^{(S)}) = H(\pi^{(T \cup \{i\})}) - H(\pi^{(T)}) = H(\pi^{(i)})$, and hence

$$H(Q^{(S \cup \{i\})}) - H(Q^{(S)}) \geq H(Q^{(T \cup \{i\})}) - H(Q^{(T)}),$$

which readily follows from the submodularity property of Shannon entropy of the edge measure Q (see e.g. (Polyanskiy and Wu, 2022, Chapter 1.4)).

Next, we prove item (2). By property of submodular function, the mapping $S \mapsto H(P^{(-S)})$ is also submodular, and hence

$$D(P \parallel P^{(S)} \otimes P^{(-S)}) = H(P^{(S)}) + H(P^{(-S)}) - H(P)$$

is submodular since the sum of two submodular functions is submodular.

Finally, we prove item (3). The monotonicity is shown in Proposition 2.4, while using Lemma 4.1 we note that

$$\mathbb{I}(P^{(S)}) = \sum_{i \in S} H(P^{(i)}) - H(P^{(S)}).$$

As the sum of two supermodular functions is supermodular, $\mathbb{I}(P^{(S)})$ is supermodular. □

Acknowledgements

Michael Choi is grateful for helpful discussions on large deviations with Pierre Del Moral, and on information theory with Cheuk Ting Li and Lei Yu. Michael Choi acknowledges the financial support of the project “MAPLE: Mechanistic Accelerated Prediction of Protein Secondary Structure via LangEvin Monte Carlo” with grant number 22-5715-P0001 under the NUS Faculty of Science Ministry of Education Tier 1 grant Data for Science and Science for Data collaborative scheme, project NUSREC-HPC-00001 and NUSREC-CLD-00001 for NUS HPC-AI Priority Projects for Research Program, as well as the startup funding of the National University of Singapore with grant number A-0000178-01-00. Youjia Wang gratefully acknowledges the financial support from National University of Singapore via the Presidential Graduate Fellowship. GW expresses gratitude to Shun Watanabe for insightful discussions and for highlighting references, specifically (Amari and Han, 1989) and (Watanabe, 2017). The work of GW was supported by the Special Postdoctoral Researcher Program (SPDR) of RIKEN and by the Japan Society for the Promotion of Science KAKENHI under Grant 23K13024.

References

- S. Agrawal, D. Vats, K. Łatuszyński, and G. O. Roberts. Optimal scaling of MCMC beyond Metropolis. *Adv. in Appl. Probab.*, 55(2):492–509, 2023.
- D. Aldous and J. A. Fill. Reversible markov chains and random walks on graphs, 2002. Unfinished monograph, recompiled 2014, available at <http://www.stat.berkeley.edu/~aldous/RWG/book.html>.
- S.-I. Amari and T. S. Han. Statistical inference under multiterminal rate restrictions: A differential geometric approach. *IEEE Transactions on Information Theory*, 35(2):217–227, 1989.
- N. Bhatnagar and D. Randall. Simulated tempering and swapping on mean-field models. *J. Stat. Phys.*, 164(3):495–530, 2016.

- J. Bierkens. Non-reversible Metropolis-Hastings. *Stat. Comput.*, 26(6):1213–1228, 2016.
- L. J. Billera and P. Diaconis. A geometric interpretation of the Metropolis-Hastings algorithm. *Statistical Science*, pages 335–339, 2001.
- V. Borkar and L. Miclo. On the fastest finite Markov processes. *J. Math. Anal. Appl.*, 481(2):123488, 43, 2020.
- S. Boucheron, G. Lugosi, and P. Massart. *Concentration inequalities*. Oxford University Press, Oxford, 2013. A nonasymptotic theory of independence, With a foreword by Michel Ledoux.
- S. Boyd, P. Diaconis, and L. Xiao. Fastest mixing Markov chain on a graph. *SIAM Rev.*, 46(4):667–689, 2004.
- S. Boyd, P. Diaconis, P. Parrilo, and L. Xiao. Fastest mixing Markov chain on graphs with symmetries. *SIAM J. Optim.*, 20(2):792–819, 2009.
- Z. Burda, J. Duda, J.-M. Luck, and B. Waclaw. Localization of the maximal entropy random walk. *Physical review letters*, 102(16):160602, 2009.
- F. Chen, L. Lovász, and I. Pak. Lifting Markov chains to speed up mixing. In *Annual ACM Symposium on Theory of Computing (Atlanta, GA, 1999)*, pages 275–281. ACM, New York, 1999.
- G.-Y. Chen and T. Kumagai. Cutoffs for product chains. *Stochastic Process. Appl.*, 128(11):3840–3879, 2018.
- T.-L. Chen, W.-K. Chen, C.-R. Hwang, and H.-M. Pai. On the optimal transition matrix for Markov chain Monte Carlo sampling. *SIAM J. Control Optim.*, 50(5):2743–2762, 2012.
- Y. Chen, A. Bušić, and S. Meyn. Ergodic theory for controlled Markov chains with stationary inputs. *Ann. Appl. Probab.*, 28(1):79–111, 2018.
- M. C. Choi and G. Wolfer. Systematic approaches to generate reversiblizations of markov chains. *IEEE Transactions on Information Theory*, 2024+. doi: 10.1109/TIT.2023.3304685.
- F. R. Chung, R. L. Graham, P. Frankl, and J. B. Shearer. Some intersection theorems for ordered sets and graphs. *Journal of Combinatorial Theory, Series A*, 43(1):23–37, 1986.
- T. M. Cover and J. A. Thomas. *Elements of information theory*. Wiley-Interscience [John Wiley & Sons], Hoboken, NJ, second edition, 2006.
- I. Csiszár, T. Cover, and B.-S. Choi. Conditional limit theorems under Markov conditioning. *IEEE Transactions on Information Theory*, 33(6):788–801, 1987.

- P. Del Moral. *Feynman-Kac formulae*. Probability and its Applications (New York). Springer-Verlag, New York, 2004. Genealogical and interacting particle systems with applications.
- A. Dembo and O. Zeitouni. *Large deviations techniques and applications*, volume 38 of *Stochastic Modelling and Applied Probability*. Springer-Verlag, Berlin, 2010. Corrected reprint of the second (1998) edition.
- P. Diaconis and L. Miclo. On characterizations of Metropolis type algorithms in continuous time. *ALEA Lat. Am. J. Probab. Math. Stat.*, 6:199–238, 2009.
- G. Fayolle and A. de La Fortelle. Entropy and the principle of large deviations for discrete-time Markov chains. *Problemy Peredachi Informatsii*, 38(4):121–135, 2002. ISSN 0555-2923.
- W. H. Fleming and D. Hernández-Hernández. Risk-sensitive control of finite state machines on an infinite horizon. I. *SIAM J. Control Optim.*, 35(5):1790–1810, 1997.
- D. Gavinsky, S. Lovett, M. Saks, and S. Srinivasan. A tail bound for read- k families of functions. *Random Structures Algorithms*, 47(1):99–108, 2015.
- A. Ghassemi and N. Kiyavash. Interaction information for causal inference: The case of directed triangle. In *2017 IEEE International Symposium on Information Theory (ISIT)*, pages 1326–1330, 2017.
- T. S. Han. Nonnegative entropy measures of multivariate symmetric correlations. *Information and Control*, 36:133–156, 1978.
- Y. Hao, A. Orlitsky, and V. Pichapati. On learning markov chains. In S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 31. Curran Associates, Inc., 2018.
- M. Hayashi and S. Watanabe. Information geometry approach to parameter estimation in Markov chains. *The Annals of Statistics*, 44(4):1495 – 1535, 2016.
- D. Hernández-Hernández and S. I. Marcus. Risk sensitive control of Markov processes in countable state space. *Systems Control Lett.*, 29(3):147–155, 1996.
- D. M. Higdon. Auxiliary variable methods for markov chain monte carlo with applications. *Journal of the American Statistical Association*, 93(442):585–595, 1998.
- N. J. Higham. Matrix nearness problems and applications. In *Applications of matrix theory (Bradford, 1988)*, volume 22 of *Inst. Math. Appl. Conf. Ser. New Ser.*, pages 1–27. Oxford Univ. Press, New York, 1989.
- T. Holliday, A. Goldsmith, and P. Glynn. Entropy and mutual information for markov channels with general inputs. In *Proceedings of the annual Allerton conference on communication control and computing*, volume 40, pages 824–833. The University; 1998, 2002.

- D. Hsu, A. Kontorovich, D. A. Levin, Y. Peres, C. Szepesvári, and G. Wolfer. Mixing time estimation in reversible Markov chains from a single sample path. *Ann. Appl. Probab.*, 29(4):2439–2480, 2019.
- L.-J. Huang, Y.-T. Liao, T.-L. Chen, and C.-R. Hwang. Optimal variance reduction for Markov chain Monte Carlo. *SIAM J. Control Optim.*, 56(4):2977–2996, 2018.
- M. Jerrum, J.-B. Son, P. Tetali, and E. Vigoda. Elementary bounds on Poincaré and log-Sobolev constants for decomposable Markov chains. *Ann. Appl. Probab.*, 14(4):1741–1765, 2004.
- J. G. Kemeny and J. L. Snell. *Finite Markov chains: with a new appendix" Generalization of a fundamental matrix"*. Springer, New York, 1983.
- D. Lacker. Independent projections of diffusions: Gradient flows for variational inference and optimal mean field approximations, 2023.
- D. A. Levin and Y. Peres. *Markov chains and mixing times*, volume 107. American Mathematical Soc., 2017.
- Z. Li and L.-H. Lim. Generalized matrix nearness problems. *SIAM J. Matrix Anal. Appl.*, 44(4):1709–1730, 2023.
- P. Mathé. Relaxation of product Markov chains on product spaces. *J. Complexity*, 14(3):319–332, 1998.
- R. Montenegro and P. Tetali. Mathematical aspects of mixing times in Markov chains. *Found. Trends Theor. Comput. Sci.*, 1(3):x+121, 2006.
- H. Nagaoka. The exponential family of Markov chains and its information geometry. In *The proceedings of the Symposium on Information Theory and Its Applications*, volume 28(2), pages 601–604, 2005.
- S. Natarajan. Large deviations, hypotheses testing, and source coding for finite Markov chains. *IEEE Trans. Inform. Theory*, 31(3):360–365, 1985.
- K. Neklyudov, M. Welling, E. Egorov, and D. Vetrov. Involutive mcmc: a unifying framework. In *Proceedings of the 37th International Conference on Machine Learning, ICML'20*, 2020.
- Y. Polyanskiy and Y. Wu. Information theory: From coding to learning. *Book draft*, 2022.
- Z. Rached, F. Alajaji, and L. L. Campbell. The Kullback-Leibler divergence rate between Markov sources. *IEEE Trans. Inform. Theory*, 50(5):917–921, 2004.
- L. Saloff-Coste. Lectures on finite Markov chains. In *Lectures on probability theory and statistics (Saint-Flour, 1996)*, volume 1665 of *Lecture Notes in Math.*, pages 301–413. Springer, Berlin, 1997.

- I. Sason. Information inequalities via submodularity and a problem in extremal graph theory. *Entropy*, 24(5):597, 2022.
- W. F. Schreiber. Cameraman image. <https://hdl.handle.net/1721.3/195767>, 1978. Accessed: 10-Apr-2024. Licensed under CC BY-NC.
- E. Shafieepoorfard, M. Raginsky, and S. P. Meyn. Rationally inattentive control of Markov processes. *SIAM J. Control Optim.*, 54(2):987–1016, 2016.
- B. van Ginkel, B. van Gisbergen, and F. Redig. Run-and-tumble motion: the role of reversibility. *J. Stat. Phys.*, 183(3):Paper No. 44, 31, 2021.
- M. Vidyasagar. An elementary derivation of the large deviation rate function for finite state Markov chains. *Asian J. Control*, 16(1):1–19, 2014.
- M. J. Wainwright. *High-dimensional statistics*, volume 48 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge, 2019. A non-asymptotic viewpoint.
- Y. Wang and M. C. H. Choi. Information divergences of markov chains and their applications, 2023.
- S. Watanabe. Neyman–Pearson test for zero-rate multiterminal hypothesis testing. *IEEE Transactions on Information Theory*, 64(7):4923–4939, 2017.
- G. Wolfer and S. Watanabe. Information geometry of reversible Markov chains. *Information Geometry*, 4(2):393–433, 2021.
- G. Wolfer and S. Watanabe. Information geometry of Markov kernels: a survey. *Frontiers in Physics*, 11, 2023.
- G. Wolfer and S. Watanabe. Geometric aspects of data-processing of Markov chains. *To appear in Transactions of Mathematics and Its Applications*, 2024+.
- S.-J. Wu and M. T. Chu. Constructing optimal transition matrix for Markov chain Monte Carlo. *Linear Algebra Appl.*, 487:184–202, 2015.
- R. W. Yeung. *Information theory and network coding*. New York, NY: Springer, 2008.