

# Error Analysis of Three-Layer Neural Network Trained with PGD for Deep Ritz Method

Yuling Jiao<sup>a</sup>, Yanming Lai<sup>\*b</sup>, and Yang Wang<sup>b</sup>

<sup>a</sup>School of Mathematics and Statistics, and Hubei Key Laboratory of Computational Science, Wuhan University, Wuhan 430072, China

<sup>b</sup>Department of Mathematics, The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong, China

May 21, 2024

## Abstract

Machine learning is a rapidly advancing field with diverse applications across various domains. One prominent area of research is the utilization of deep learning techniques for solving partial differential equations (PDEs). In this work, we specifically focus on employing a three-layer tanh neural network within the framework of the deep Ritz method (DRM) to solve second-order elliptic equations with three different types of boundary conditions. We perform projected gradient descent (PDG) to train the three-layer network and we establish its global convergence. To the best of our knowledge, we are the first to provide a comprehensive error analysis of using overparameterized networks to solve PDE problems, as our analysis simultaneously includes estimates for approximation error, generalization error, and optimization error. We present error bound in terms of the sample size  $n$  and our work provides guidance on how to set the network depth, width, step size, and number of iterations for the projected gradient descent algorithm. Importantly, our assumptions in this work are classical and we do not require any additional assumptions on the solution of the equation. This ensures the broad applicability and generality of our results.

## 1 Introduction

Machine learning is a field of artificial intelligence that focuses on developing algorithms and models capable of learning from data and making predictions

---

\*Corresponding Author (ylaiam@connect.ust.hk)

or decisions. It involves the study of statistical techniques and computational algorithms to enable computers to automatically learn and improve from experience. Machine learning finds applications in various domains, such as image and speech recognition, natural language processing, recommendation systems, and autonomous vehicles. By leveraging large datasets and powerful computational resources, machine learning algorithms can uncover patterns, extract insights, and solve complex problems, driving advancements in technology and revolutionizing numerous industries.

Neural networks play a crucial role in machine learning methods, and their approximation capability is an important topic of concern for researchers. [56] is a review of the two-layer network approximation results from the 1990s. In recent years, there have been numerous studies on the approximation results of deep neural networks [77, 78, 79, 80, 65, 69, 66, 59, 42, 60, 61, 83, 82, 35, 28, 29, 76]. The pioneering work of [77] introduced a novel approach to approximate smooth functions using neural networks. By constructing neural networks that approximate Taylor expansions and partition of unity, this work provided insights into the approximation of smooth functions. [69] focuses on approximation in Besov spaces. [42] study the approximation of smooth function classes. [28, 29] investigate approximation in Sobolev spaces. These works primarily utilize ReLU and sigmoid activation functions, which are commonly used in practice, to study the approximation properties of neural networks in common function spaces. There are also other works that explore neural networks with super-approximation capabilities. [80] demonstrates that ReLU-periodic function networks can overcome the curse of dimensionality. [79] shows that sin-arcsin networks can overcome the curse of dimensionality. [60] demonstrates that ReLU-floor neural networks can overcome the curse of dimensionality. Shen et al. [61] shows that Floor-Exponential-Step networks can overcome the curse of dimensionality. [83] leverage a triangular-wave function and the softsign function to overcome the curse of dimensionality. Jiao et al. [35] demonstrate that ReLU-sine-exponential networks can overcome the curse of dimensionality. However, these results only address the curse of dimensionality at the approximation level and do not consider issues at the training level. Recently, [82] shows that ReLU neural networks can be approximated by commonly used activation functions. Therefore, the approximation results of ReLU networks can be translated into approximation results for other activation function networks.

The theoretical study of training problems in neural networks is a vast and rapidly evolving field. To gain a comprehensive understanding of this field, the readers are referred to the review article [9]. [32] proposed the framework of Neural Tangent Kernel (NTK) analysis and proved the global convergence of infinitely wide neural networks. Since then, research based on NTK theory for studying optimization problems in shallow and deep neural networks has become a popular area of study. For further exploration, please refer to [32, 3, 2, 4, 22, 21, 5, 85, 84, 13, 12, 15, 54, 53, 52, 39, 38], and the references mentioned therein. These works demonstrate that training shallow and deep neural networks using gradient descent and stochastic gradient descent algorithms exhibits global convergence. In NTK analysis, the minimum eigenvalue

of the kernel matrix plays a crucial role. However, research on the minimum eigenvalue of the kernel matrix is currently quite limited. [54, 53, 6, 55] are outstanding works in this field. Given the importance of understanding the minimum eigenvalue’s magnitude, further research in this area is highly needed. Apart from NTK theory, there are other theoretical approaches for studying neural network training. For example, mean field theory [47, 16, 25, 51, 68] has been utilized in this context as well.

The generalization performance of neural networks is an important metric, and it is related to the complexity of the networks. [8, 7, 26, 50] have studied the VC dimension and pseudo-dimension of neural networks, which are indicators of function complexity. [75] investigates upper bounds on the VC dimension and pseudo-dimension for function classes involving derivatives. Some works have shown that trained neural networks exhibit good generalization performance [5, 2, 12, 13]. However, these works still have some unresolved issues. In section 6 of [5], the authors discuss the conditions under which the learning task can achieve good performance when the samples satisfy certain underlying functions. However, the function classes discussed in that section are highly limited and do not cover most real-world scenarios. It is mentioned that two-layer networks can generalize well, but there is no clear characterization of the parameter upper bounds for approximation in two-layer networks [48, 56]. Additionally, the function classes that two-layer networks can approximate are extremely limited. For example, it seems that there is currently no result indicating that two-layer networks can approximate functions in Sobolev spaces. The upper bounds in [12] are of a similar nature to those in [5], so they face the same issues. [2] assumes the existence of an underlying neural network that achieves low error on the data distribution, and the available data is significantly more than the minimum number of samples required to learn this underlying neural network. However, it is difficult to verify this assumption from the data. The results in [13] depend on the PL coefficient, which in turn depends on the minimum eigenvalue of the NTK. However, as mentioned above, the relationship between the minimum eigenvalue and the number of samples is still not clearly explained.

Using deep learning methods to solve PDEs is a popular field of research. Neural network models have been applied to solve various types of PDEs [30, 71, 67, 43, 40, 57]. Different loss functions have been proposed: [73] introduced the deep Ritz method (DRM), [58] proposed Physics-informed Neural Networks (PINNs), [81] introduced the weak adversarial network (WAN), and [14] proposed Friedrichs learning. At the theoretical level, works such as [72, 45, 31, 74, 41] have provided error analysis for these methods based on the assumption that the solutions of the underlying equations lie in the so-called Barron spaces. Subsequently, works such as [34, 24, 23, 33, 44, 62, 63, 36, 49] have discovered that this assumption is not necessary. Instead, relying on the classical theory of partial differential equation regularity is sufficient to establish error analysis for these methods, achieving consistency with classical numerical methods such as finite element methods. However, the aforementioned works only consider either the approximation error or the generalization error, or a combination of both. They fail to account for the optimization error that arises from training

the neural network using optimization algorithms. Consequently, their analyses are incomplete. In order to provide a solid theoretical foundation for using machine learning to solve PDEs, it is essential to incorporate factors related to the training process into theoretical research. Indeed, this is a significant challenge because, as discussed earlier, the relationship between neural network optimization and generalization is not yet fully understood.

## 1.1 Main Results

Here, we present an informal version of the main results in this work. To keep it concise, we focus on providing the convergence rate of the Robin problem. For the formal and comprehensive version, please refer to Theorem 5.

**Theorem 1** (informal version). *Let  $u_R$  be the weak solution of Robin problem (1)(4). Let  $n$  be the sample size. Let the overparametrization condition be*

$$A = n^{\frac{415d^4(d+3)5^{d+2}}{288d^3+4}}.$$

*Let the step size*

$$\eta \leq C(d, \text{coe}, \Omega) n^{-\frac{103d^3}{144d^3+2}} \frac{1}{A}$$

*and the iteration step  $T = \frac{1}{\eta}$ . Let  $f_{W_T}$  be the three-layer neural network function trained by PGD after  $T$  step. Then with probability at least  $1 - \frac{C(d, \text{coe}, \Omega)}{n}$ ,*

$$\|f_{W_T} - u_R\|_{H^1(\Omega)} \leq C(d, \text{coe}, \Omega) \max\{1, 1/\beta\} n^{-\frac{1}{288d^3+4}}.$$

## 1.2 Our Contributions

The following are the main contributions of our work:

1. To the best of our knowledge, we are the first to provide a comprehensive error analysis of using overparameterized networks to solve PDE problems. Our work provides guidance on how to set the network depth, width, step size, and number of iterations for the projected gradient descent algorithm. Our assumption (5) in this work is classical, common, and weak, and we do not require any additional assumptions on the solution of the equation. Therefore, our results have strong generality.
2. In this work, we construct three-layer neural networks to approximate functions in Sobolev spaces, extending the results from [18] that were originally limited to  $W^{s, \infty}(\Omega)$  spaces to general  $W^{s, p}(\Omega)$  spaces. This result complements the research in [29], which only considers approximation results for deep networks in Sobolev spaces.

3. When considering the generalization error of the neural network function class, it is necessary for the functions to lie within a pre-defined bounded ball in  $C(\Omega)$ . For PDE problems, truncation techniques that are useful for regression problems cannot be applied due to the consideration of functions in Sobolev spaces. To overcome this challenge, we leverage the properties of the projected gradient descent algorithm to ensure that the network parameters are initialized sufficiently close to the initialization, resulting in a bounded  $C(\Omega)$  norm for the iteratively obtained neural network function sequence. Furthermore, we even find that the neural network sequence generated by projected gradient descent lies within a bounded ball in  $C^1(\Omega)$ . This allows us to apply complexity bounds for functions in the  $C^1$  bounded ball to assist in estimating the generalization error.

In this work, we have referenced the techniques used in [20, 37, 19] to establish the global convergence of optimization algorithms. However, our work still differs in many aspects from theirs. First of all, our setups are not exactly the same. They use gradient descent to optimize deep neural networks, while we use projected gradient descent to optimize three-layer neural networks. Secondly, our problem setting is different. Their study focuses on regression problems and deals with a class of smooth functions, while our research focuses on PDE problems and deals with a class of Sobolev functions. Consequently, we need to construct neural networks that can approximate functions in Sobolev spaces. Thirdly, our approach to controlling the generalization error also differs from theirs. They utilize the smoothness of activation functions to control the covering number and, consequently, the generalization error. In contrast, we make full use of the properties of the projected gradient descent algorithm to demonstrate that the neural network sequences generated by the algorithm lie within a bounded ball in  $C^1(\Omega)$ . This enables us to apply complexity bounds for functions in the  $C^1$  bounded ball to estimate the generalization error. Finally, our approach to decomposing the error is also different.

### 1.3 Related Works

Recently, Kohler et al.[20, 37, 19] investigated the convergence of gradient descent for solving regression problems by overparameterized deep networks. In their work, they considered the approximation error, generalization error, and optimization error simultaneously, and their conclusions hold for smooth function classes, making their findings generalizable. They mitigated the negative impact of overparameterization on generalization error by exploiting the smoothness of activation functions, thereby bridging the gap between optimization error and generalization error under the overparameterized condition. More importantly, differing from the works based on the NTK theory, they presented an alternative approach to studying the global convergence of optimization algorithms. They utilized random initialization to ensure that the inner parameters of the neural network are sufficiently close to the best approximation elements

and leveraged the convexity of the neural network function with respect to the outer parameters to establish the global convergence of gradient descent. Furthermore, when proving global convergence, they required the 2-norm of the outer parameters of the best approximation elements to be sufficiently small, and they achieved this by increasing the network width, leading to a proportional decrease in the 2-norm of the outer parameters. By not relying on the NTK framework, they were able to avoid explicitly analyzing the minimum eigenvalue of the NTK matrix.

## 1.4 Organization of This Paper

The paper is structured as follows: Section 2 provides a brief introduction to the Deep Ritz method. Section 3 covers the necessary background knowledge and tools employed in this work. In Section 4, we present the error decomposition, breaking down the total error into approximation error, generalization error, and optimization error. Subsequently, sections 5 to 7 delve into the investigation of these three types of errors individually. The main results are presented in Section 8, and finally, in Section 9, we conclude the paper.

## 2 Deep Ritz Method

Let  $\Omega$  be a convex bounded open set in  $\mathbb{R}^d$  and assume that  $\partial\Omega \in C^\infty$ . Without loss of generality, we assume that  $\Omega \subset (0, 1)^d$ . We consider the following second order elliptic equation:

$$-\Delta u + wu = f \text{ in } \Omega \tag{1}$$

with three kinds of boundary conditions:

$$u = 0 \text{ on } \partial\Omega \tag{2}$$

$$\frac{\partial u}{\partial n} = g \text{ on } \partial\Omega \tag{3}$$

$$u + \beta \frac{\partial u}{\partial n} = g \text{ on } \partial\Omega, \quad \beta \in \mathbb{R}, \beta \neq 0 \tag{4}$$

which are called Dirichlet, Neumann and Robin boundary conditions, respectively. Note that for the Dirichlet problem, we only consider the homogeneous boundary condition here since the inhomogeneous case can be turned into the homogeneous case by translation.

We make the following assumption on the known terms in the equation:

$$f \in L^\infty(\Omega), \quad g \in H^{1/2}(\Omega), \quad w \in L^\infty(\Omega), \quad w \geq c_w \tag{5}$$

where  $c_w$  is some positive constant. In the following we abbreviate

$$C(\|f\|_{L^\infty(\Omega)}, \|g\|_{H^{1/2}(\Omega)}, \|w\|_{L^\infty(\Omega)}, c_w),$$

constants depending on the known terms in equation, as  $C(\text{coe})$  for simplicity.

For problem (1)(2), the variational problem is to find  $u \in H_0^1(\Omega)$  such that

$$(\nabla u, \nabla v) + (wu, v) = (f, v), \quad \forall v \in H_0^1(\Omega). \quad (6a)$$

The corresponding minimization problem is

$$\min_{u \in H_0^1(\Omega)} \frac{1}{2} \int_{\Omega} (|\nabla u|^2 + wu^2 - 2fu) dx. \quad (6b)$$

The variational problem (6a) is equivalent to the minimization problems (6b). This is a well-known result. The following (7a) and (7b), (8a) and (8b) have the same kind of relationship. For a reference on this topic, please refer to [17, Theorem 1.1.2].

For problem (1)(3), the variational problem is to find  $u \in H^1(\Omega)$  such that

$$(\nabla u, \nabla v) + (wu, v) = (f, v) + (g, T_0 v)|_{\partial\Omega}, \quad \forall v \in H^1(\Omega). \quad (7a)$$

where  $T_0$  is the zero order trace operator. The corresponding minimization problem is

$$\min_{u \in H^1(\Omega)} \int_{\Omega} \left( \frac{1}{2} |\nabla u|^2 + \frac{1}{2} wu^2 - fu \right) dx - \int_{\partial\Omega} g T_0 u ds. \quad (7b)$$

For problem (1)(4), the variational problem is to find  $u \in H^1(\Omega)$  such that

$$(\nabla u, \nabla v) + (wu, v) + \frac{1}{\beta} (T_0 u, T_0 v)|_{\partial\Omega} = (f, v) + \frac{1}{\beta} (g, T_0 v)|_{\partial\Omega}, \quad \forall v \in H^1(\Omega). \quad (8a)$$

The corresponding minimization problem is

$$\min_{u \in H^1(\Omega)} \int_{\Omega} \left( \frac{1}{2} |\nabla u|^2 + \frac{1}{2} wu^2 - fu \right) dx + \frac{1}{\beta} \int_{\partial\Omega} \left( \frac{1}{2} (T_0 u)^2 - g T_0 u \right) ds. \quad (8b)$$

The next lemma says that when  $g = 0$  and  $\beta \rightarrow 0$ , the solution of the Robin problem converges to the solution of the Dirichlet problem.

**Lemma 1** ([34], Lemma 3.4). *Let assumption (5) holds. Let  $g = 0$ . Let  $u_D$  be the solution of problem (6a) (also (6b)) and  $u_R$  the solution of problem (8a) (also (8b)). There holds*

$$\|u_R - u_D\|_{H^1(\Omega)} \leq C(\text{coe})\beta.$$

With this lemma, it suffice to consider the Robin problem since the Dirichlet problem can be handled through a limit process. Define  $L_R$  as a functional on  $H^1(\Omega)$ :

$$L_R(u) := \int_{\Omega} \left( \frac{1}{2} |\nabla u|^2 + \frac{1}{2} wu^2 - fu \right) dx + \frac{1}{\beta} \int_{\partial\Omega} \left( \frac{1}{2} (T_0 u)^2 - g T_0 u \right) ds.$$

Note that  $L_R$  can be equivalently written as

$$L_R(u) = |\Omega| \mathbb{E}_{X \sim U(\Omega)} \left( \frac{1}{2} |\nabla u(X)|^2 + \frac{1}{2} w(X) u^2(X) - f(X) u(X) \right)$$

$$+ \frac{|\partial\Omega|}{\beta} \mathbb{E}_{Y \sim U(\partial\Omega)} \left( \frac{1}{2} (T_0 u)^2(Y) - g(Y) T_0 u(Y) \right), \quad (9)$$

where  $U(\Omega)$  and  $U(\partial\Omega)$  are uniform distribution on  $\Omega$  and  $\partial\Omega$ , respectively. We then introduce a discrete version of  $L_R$ :

$$\begin{aligned} \widehat{L}_R(u) &:= \frac{|\Omega|}{n} \sum_{i=1}^n \left( \frac{1}{2} |\nabla u(X_i)|^2 + \frac{1}{2} w(X_i) u^2(X_i) - f(X_i) u(X_i) \right) \\ &\quad + \frac{|\partial\Omega|}{\beta m} \sum_{j=1}^m \left( \frac{1}{2} (T_0 u)^2(Y_j) - g(Y_j) T_0 u(Y_j) \right), \end{aligned} \quad (10)$$

where  $\{X_i\}_{i=1}^n$  and  $\{Y_j\}_{j=1}^m$  are i.i.d. random variables according to  $U(\Omega)$  and  $U(\partial\Omega)$  respectively. Similarly, for the Neumann problem we define

$$\begin{aligned} L_N(u) &:= |\Omega| \mathbb{E}_{X \sim U(\Omega)} \left( \frac{1}{2} |\nabla u(X)|^2 + \frac{1}{2} w(X) u^2(X) - f(X) u(X) \right) \\ &\quad - |\partial\Omega| \mathbb{E}_{Y \sim U(\partial\Omega)} g(Y) T_0 u(Y) \end{aligned} \quad (11)$$

and

$$\begin{aligned} \widehat{L}_N(u) &:= \frac{|\Omega|}{n} \sum_{i=1}^n \left( \frac{1}{2} |\nabla u(X_i)|^2 + \frac{1}{2} w(X_i) u^2(X_i) - f(X_i) u(X_i) \right) \\ &\quad - \frac{|\partial\Omega|}{m} \sum_{j=1}^m g(Y_j) T_0 u(Y_j). \end{aligned} \quad (12)$$

We consider the following two minimization problems:

$$\min_{u \in \mathcal{P}} \widehat{L}_R(u), \quad \min_{u \in \mathcal{P}} \widehat{L}_N(u) \quad (13)$$

The minimization is taken over some parametrized function class  $\mathcal{P}$ . In this work we choose  $\mathcal{P}$  to be a neural network function class. Now we discuss in details that the neural network function class we choose. For some  $m_1, m_2, A \in \mathbb{N}_{\geq 1}$ , let  $W := \{(W_s^l, b_s^l) : s \in [A], l = 1, 2, 3\}$  be the neural network parameters with  $W_s^1 \in \mathbb{R}^{m_1 \times d}$ ,  $W_s^2 \in \mathbb{R}^{m_2 \times m_1}$ ,  $W_s^3 \in \mathbb{R}^{1 \times m_2}$ ,  $b_s^1 \in \mathbb{R}^{m_1}$ ,  $b_s^2 \in \mathbb{R}^{m_2}$ ,  $b_s^3 \in \mathbb{R}$  for  $s \in [A]$  and define three-layer subnetworks  $\{f_s^3\}_{s=1}^A$  by

$$\begin{aligned} f_s^0 &= x; \\ f_s^1 &= \sigma(f_{s,1}^{org}) = \sigma(W_s^1 f_s^0 + b_s^1); \\ f_s^2 &= \sigma(f_{s,2}^{org}) = \sigma(W_s^2 f_s^1 + b_s^2); \\ f_s^3 &= W_s^3 f_s^2 + b_s^3, \end{aligned}$$

where  $\sigma : \mathbb{R} \rightarrow \mathbb{R}$ , and define  $f_W$  to be the sum of the subnetworks:

$$f_W = \sum_{s=1}^A f_s^3.$$

$\{W_s^l\}$  are called weights and  $\{b_s^l\}$  are called biases.  $\sigma$  is called an activation function. The width of subnetworks  $\{f_s^3\}_{s=1}^A$  is defined as  $\max\{m_1, m_2\}$ . Let  $W^l = \{(W_s^l, b_s^l), s \in [A]\}$  for  $l = 1, 2, 3$ . Define  $\|W^3\| := \left[ \sum_{s=1}^A \|W_s^3\|_F^2 + |b_s^3|^2 \right]^{1/2}$  and  $\|W^3\|_1 := \left[ \sum_{s=1}^A (\|W_s^3\|_\infty + |b_s^3|) \right]^{1/2}$ . For some  $B_{inn}, B_{out} \in \mathbb{R}$ , define the neural network function class  $\mathcal{F}_{NN}(\{m_1, m_2, A\}, B_{inn}, B_{out})$  to be

$$\begin{aligned} \mathcal{F}_{NN}(\{m_1, m_2, A\}, B_{inn}, B_{out}) := \\ \{f_W : \|W_s^l\|_F, \|b_s^l\|_2 \leq B_{inn} (l = 1, 2; s \in [A]), \|W^3\|_1 \leq B_{out}\}. \end{aligned}$$

In the following we abbreviate  $\mathcal{F}_{NN}(\{m_1, m_2, A\}, B_{inn}, B_{out})$  as  $\mathcal{F}_{NN}$  for simplicity. In this work we choose  $m_1 = 5d, m_2 = \binom{2d+1}{d+1}$  and let the activation function  $\sigma$  be  $\tanh$ . Let  $B_\sigma = \max\{\|\sigma\|_{C(\Omega)}, 1\}$  and define  $B_{\sigma'}, B_{\sigma''}$  similarly. Other parameters will be specified in Theorem 5.

**Remark 1.** *Indeed, as we will see later, the requirement of  $\sigma = \tanh$  is only necessary when studying the approximation error. For the analysis of the generalization error and optimization error, it suffices for  $\sigma$  to be  $C^2$  continuous. But as highlighted by [18], the approach we employ for handling the approximation error is also applicable to other common activation functions such as sigmoid, arctan, and more. Therefore, with slight adjustments, the analysis in this paper can be extended to derive convergence rates for neural networks activated by these functions as well.*

Many algorithms can be used to solve the minimization problem (13), such as gradient descent, stochastic gradient descent, etc. In this paper we employ projected gradient descent (PGD) to solve (13). Taking the Robin problem as an example, the update rule of PGD is as follows:

$$W_{t+1} = \text{proj}_{\mathcal{C}}(W_t - \eta \nabla_W \widehat{L}_R(W_t)), \quad t = 0, 1, \dots, T-1,$$

where  $\eta \in \mathbb{R}_{>0}$  is step size,  $T \in \mathbb{N}_{>0}$  is the number of iterations and  $\mathcal{C}$  is the set to which we project the iterates sequence. We initialize the outer layer parameters to zero:  $W_s^3 = 0, b_s^3 = 0, s \in [A]$ . The initialization of the inner layer parameters will be specified in Theorem 5.

## 3 Preliminaries

### 3.1 Sobolev Spaces

In this part we summarize some concepts and results related to Sobolev spaces that we will need for our analysis.

Let  $d \in \mathbb{N}_{\geq 1}$ . Let  $\Omega \subset \mathbb{R}^d$  be an open bounded domain with smooth boundary  $\partial\Omega$ , and without loss of generality we assume that  $\Omega \subset [0, 1]^d$ . Let  $\alpha = (\alpha_1, \dots, \alpha_n)$  be an  $n$ -dimensional index with  $|\alpha| := \sum_{i=1}^n \alpha_i$  and  $s$  be a

nonnegative integer. We use the notation  $D^\alpha = \frac{\partial^{|\alpha|}}{\partial x_1^{\alpha_1} \dots \partial x_d^{\alpha_d}}$ . The standard function spaces, including continuous function space,  $L^p$  space, Sobolev spaces, are given below.

$$\begin{aligned}
C(\Omega) &:= \{\text{all the continuous functions defined on } \Omega\}, \\
C^s(\Omega) &:= \{f : \Omega \rightarrow \mathbb{R} \mid D^\alpha f \in C(\Omega)\}, \\
C(\bar{\Omega}) &:= \{\text{all the continuous functions defined on } \bar{\Omega}\}, \quad \|f\|_{C(\bar{\Omega})} := \max_{x \in \bar{\Omega}} |f(x)|, \\
C^s(\bar{\Omega}) &:= \{f : \bar{\Omega} \rightarrow \mathbb{R} \mid D^\alpha f \in C(\bar{\Omega})\}, \quad \|f\|_{C^s(\bar{\Omega})} := \max_{x \in \bar{\Omega}, |\alpha| \leq s} |D^\alpha f(x)|, \\
L^p(\Omega) &:= \left\{ f : \Omega \rightarrow \mathbb{R} \mid \int_\Omega |f|^p dx < \infty \right\}, \quad \|f\|_{L^p(\Omega)} := \left[ \int_\Omega |f|^p(x) dx \right]^{1/p}, \quad \forall p \in [1, \infty), \\
L^\infty(\Omega) &:= \{f : \Omega \rightarrow \mathbb{R} \mid \exists C > 0 \text{ s.t. } |f| \leq C \text{ a.e.}\}, \quad \|f\|_{L^\infty(\Omega)} := \inf\{C \mid |f| \leq C \text{ a.e.}\}, \\
W^{s,p}(\Omega) &:= \{f : \Omega \rightarrow \mathbb{R} \mid D^\alpha f \in L^p(\Omega), |\alpha| \leq s\}, \quad \|f\|_{W^{s,p}(\Omega)} := \left( \sum_{|\alpha| \leq s} \|D^\alpha f\|_{L^p(\Omega)}^p \right)^{1/p}.
\end{aligned}$$

If  $s$  is a nonnegative real number, the fractional Sobolev space  $W^{s,p}(\Omega)$  can be defined as follows: setting  $\theta = s - [s]$  and

$$\begin{aligned}
W^{s,p}(\Omega) &:= \left\{ f : \Omega \rightarrow \mathbb{R} \mid \int_\Omega \int_\Omega \frac{|D^\alpha f(x) - D^\alpha f(y)|^p}{|x - y|^{\theta p + d}} dx dy < \infty, \quad \forall |\alpha| = [s] \right\}, \\
\|f\|_{W^{s,p}(\Omega)} &:= \left( \|f\|_{W^{[s],p}(\Omega)}^p + \sum_{|\alpha| = [s]} \int_\Omega \int_\Omega \frac{|D^\alpha f(x) - D^\alpha f(y)|^p}{|x - y|^{\theta p + d}} dx dy \right)^{1/p}.
\end{aligned}$$

Let  $C_0^\infty(\Omega)$  be the set of smooth functions with compact support in  $\Omega$ , and  $W_0^{s,p}(\Omega)$  is the completion space of  $C_0^\infty(\Omega)$  in  $W^{s,p}(\Omega)$ . For  $s < 0$ ,  $W^{s,p}(\Omega)$  is the dual space of  $W_0^{-s,q}(\Omega)$  with  $q$  satisfying  $\frac{1}{p} + \frac{1}{q} = 1$ . When  $p = 2$ ,  $W^{s,p}(\Omega)$  is a Hilbert space and it is also denoted by  $H^s(\Omega)$ .

**Lemma 2** ([18], Lemma A.7). *Let  $d_1, d_2, k \in \mathbb{N}_{\geq 1}$ ,  $\Omega_1 \subset \mathbb{R}^{d_1}$ ,  $\Omega_2 \subset \mathbb{R}^{d_2}$ ,  $f \in C^k(\Omega_1; \mathbb{R})$  and  $g \in C^k(\Omega_2; \mathbb{R})$ . Then it holds that*

$$\|g \circ f\|_{W^{k,\infty}(\Omega_1)} \leq 16 (e^2 k^4 d_2 d_1^2)^k \|g\|_{W^{k,\infty}(\Omega_2)} \max_{1 \leq i \leq d_2} \|(f)_i\|_{W^{k,\infty}(\Omega_1)}^k.$$

**Lemma 3** ([29], Lemma B.5). *Let  $k \in \mathbb{N}_{\geq 0}$ , and assume that  $f \in W^{k,\infty}(\Omega)$  and  $g \in W^{k,p}(\Omega)$  with  $1 \leq p \leq \infty$ . If  $k \geq 3$ , additionally assume that  $f \in C^k(\Omega)$  or  $g \in C^k(\Omega)$ . Then  $fg \in W^{k,p}(\Omega)$  and*

$$\|fg\|_{W^{k,p}(\Omega)} \leq C(k, p, d) \sum_{i=0}^k \|f\|_{W^{i,\infty}(\Omega)} \|g\|_{W^{k-i,p}(\Omega)},$$

and, consequently

$$\|fg\|_{W^{k,p}(\Omega)} \leq C(k, p, d) \|f\|_{W^{k,\infty}(\Omega)} \|g\|_{W^{k,p}(\Omega)}.$$

**Lemma 4.** *Let  $s \in \mathbb{N}, 1 \leq p \leq \infty$ . Let  $\Omega \subset \mathbb{R}^d$  be open and bounded,  $x_0 \in \Omega$  and  $r > 0$  such that  $\Omega$  is star-shaped with respect to  $B := B_{r, \|\cdot\|_2}(x_0)$ , and  $r > (1/2)r_{\max}^*$ . Then for any  $f \in W^{s,p}(\Omega)$ , there exists a polynomial  $f^{(poly)} = \sum_{\tilde{s}=0}^{s-1} \sum_{\beta \in P_{\tilde{s},d}} c_\beta x^\beta$  such that*

$$\left| f - f^{(poly)} \right|_{W^{k,p}(\Omega)} \leq C(s, d, \gamma) h^{s-k} |f|_{W^{s,p}(\Omega)} \quad \text{for } k = 0, 1, \dots, s,$$

where  $h = \text{diam}(\Omega)$  and  $\gamma = h/r_{\max}^*$ . Moreover,  $|c_\beta| \leq C(s, d, R) r^{-d/p} \|f\|_{W^{s-1,p}(\Omega)}$  for all  $\beta \in P_{\tilde{s},d}$  with  $0 \leq |\tilde{s}| \leq s-1$ .

*Proof.* The existence of  $f_{poly}$  is precisely the well-known Bramble-Hilbert lemma, and its proof can be found in [11, Lemma 4.3.8]. The upper bound estimation of  $|c_\beta|$  can be found in [28, Lemma B.9].  $\square$

**Definition 1** (trace operator). *Let  $m \in \mathbb{N}_{\geq 1}$ . Let  $\Omega$  be a  $C^m$  domain. The trace operator  $T = (T_0, T_1, \dots, T_{m-1})$  is defined by*

$$\begin{aligned} T : H^m(\Omega) &\rightarrow H^{m-1/2}(\partial\Omega) \times H^{m-3/2}(\partial\Omega) \times \dots \times H^{1/2}(\partial\Omega) \\ v &\mapsto (T_0 v, T_1 v, \dots, T_{m-1} v). \end{aligned}$$

**Lemma 5** (trace theorem). *Let  $m \in \mathbb{N}_{\geq 1}$ . Let  $\Omega$  be a  $C^m$  domain. Then the trace operator is continuous and surjective.*

*Proof.* See [1, Theorem 7.33].  $\square$

The next three lemmas are classical regularity results for the PDEs we are concerned with in this work.

**Lemma 6** ([27], Theorem 2.4.2.5). *Let assumption (5) holds. Let  $u_D$  be the weak solution of the Dirichlet problem (1)(2) but replacing the right-hand side of (2) with some  $H^{3/2}(\partial\Omega)$  function  $g$ . Then  $u_D \in H^2(\Omega)$  and*

$$\|u_D\|_{H^2(\Omega)} \leq C(\Omega, w) (\|f\|_{L^2(\Omega)} + \|g\|_{H^{3/2}(\partial\Omega)}).$$

**Lemma 7** ([27], Theorem 2.4.2.7). *Let assumption (5) holds. Let  $u_N$  be the solution of the Neumann problem (1)(3). Then  $u_N \in H^2(\Omega)$  and*

$$\|u_N\|_{H^2(\Omega)} \leq C(\Omega, w) (\|f\|_{L^2(\Omega)} + \|g\|_{H^{1/2}(\partial\Omega)}).$$

**Lemma 8.** *Let assumption (5) holds. Let  $u_R$  be the solution of the Robin problem (1)(4). Then  $u_R \in H^2(\Omega)$  and*

$$\|u_R\|_{H^2(\Omega)} \leq C(\Omega, w) (\|f\|_{L^2(\Omega)} + \|g\|_{H^{3/2}(\partial\Omega)}).$$

*Proof.* The proof is given in the appendix.  $\square$

### 3.2 Convex Optimization

**Definition 2.** Let  $n \in \mathbb{N}_{\geq 1}$ . A subset  $C$  of  $\mathbb{R}^n$  is called convex if

$$\alpha x + (1 - \alpha)y \in C, \quad \forall x, y \in C, \forall \alpha \in [0, 1].$$

**Definition 3.** Let  $n \in \mathbb{N}_{\geq 1}$ . Let  $C$  be a nonempty convex subset of  $\mathbb{R}^n$ . We say that a function  $f : C \rightarrow \mathbb{R}$  is convex if

$$f(\alpha x + (1 - \alpha)y) \leq \alpha f(x) + (1 - \alpha)f(y), \quad \forall x, y \in C, \forall \alpha \in [0, 1].$$

**Lemma 9** ([10], Proposition 1.1.7). Let  $n \in \mathbb{N}_{\geq 1}$ . Let  $C$  be a nonempty convex subset of  $\mathbb{R}^n$  and let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be differentiable over an open set that contains  $C$ . Then  $f$  is convex over  $C$  if and only if

$$f(z) \geq f(x) + \nabla f(x)^T(z - x), \quad \forall x, z \in C.$$

$f$  is strictly convex over  $C$  if and only if the above inequality is strict whenever  $x \neq z$ .

**Lemma 10** ([10], Proposition 1.1.10). Let  $n \in \mathbb{N}_{\geq 1}$ . Let  $C$  be a nonempty convex subset of  $\mathbb{R}^n$  and let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be twice continuously differentiable over an open set that contains  $C$ . Let  $\nabla^2 f(x)$  be the Hessian of  $f$  at  $x$ . If  $\nabla^2 f(x)$  is positive semi-definite for all  $x \in C$ , then  $f$  is convex over  $C$ ; if  $\nabla^2 f(x)$  is positive definite for all  $x \in C$ , then  $f$  is strictly convex over  $C$ .

**Definition 4** (projection). Let  $n \in \mathbb{N}_{\geq 1}$ . Let  $C$  be a nonempty convex subset of  $\mathbb{R}^n$  and let  $z$  be a vector in  $\mathbb{R}^n$ . The vector that minimizes  $\|z - x\|_2$  over  $x \in C$  is called the projection of  $z$  on  $C$  and denoted as  $\text{proj}_C(z)$ .

**Lemma 11** ([10], Proposition 1.1.9). Let  $n \in \mathbb{N}_{\geq 1}$ . Let  $C$  be a nonempty convex subset of  $\mathbb{R}^n$  and let  $z$  be a vector in  $\mathbb{R}^n$ .  $\text{proj}_C(z)$  is uniquely determined. Furthermore, a vector  $x^* = \text{proj}_C(z)$  if and only if

$$(z - x^*)^T(x - x^*) \leq 0, \quad \forall x \in C.$$

**Lemma 12.** Let  $n \in \mathbb{N}_{\geq 1}$ . Let  $C$  be a nonempty convex subset of  $\mathbb{R}^n$  and let  $z, z'$  be two vectors in  $\mathbb{R}^n$ . There holds

$$\begin{aligned} \|\text{proj}_C(z) - \text{proj}_C(z')\|_2^2 &\leq (z - z')^T(\text{proj}_C(z) - \text{proj}_C(z')), \\ \|\text{proj}_C(z) - \text{proj}_C(z')\|_2 &\leq \|z - z'\|_2. \end{aligned}$$

*Proof.* We have

$$\begin{aligned} \|\text{proj}_C(z) - \text{proj}_C(z')\|_2^2 &= (\text{proj}_C(z) - \text{proj}_C(z'))^T(\text{proj}_C(z) - \text{proj}_C(z')) \\ &= (\text{proj}_C(z) - z)^T(\text{proj}_C(z) - \text{proj}_C(z')) + (z - z')^T(\text{proj}_C(z) - \text{proj}_C(z')) \\ &\quad + (z' - \text{proj}_C(z'))^T(\text{proj}_C(z) - \text{proj}_C(z')). \end{aligned}$$

By Lemma 11, the first and the third term on the right-hand side are nonpositive, hence we obtain the first inequality. By Cauchy-Schwarz inequality, the right-hand side of the first inequality is not greater than  $\|z - z'\|_2 \|\text{proj}_C(z) - \text{proj}_C(z')\|_2$ . Then the second inequality follows.  $\square$

**Definition 5.** Let  $n \in \mathbb{N}_{\geq 1}, L \in \mathbb{R}_{>0}$ . Let  $C$  be a nonempty convex subset of  $\mathbb{R}^n$ . Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be differentiable over an open set that contains  $C$ .  $f$  is called  $L$ -strongly smooth over  $C$  if for any  $x, y \in C$ ,

$$f(y) \leq f(x) + \nabla f(x)^T (y - x) + \frac{L}{2} \|y - x\|_2^2.$$

**Lemma 13.** Let  $n \in \mathbb{N}_+$ . Given  $x, y \in \mathbb{R}^n$ . For a differentiable function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , if there exists some constant  $L \in \mathbb{R}$  such that for any  $a \in [0, 1]$ ,

$$\|\nabla f((1-a)x + ay) - \nabla f(x)\|_2 \leq aL\|y - x\|_2,$$

then

$$f(y) \leq f(x) + \nabla f(x)^T (y - x) + \frac{L}{2} \|y - x\|_2^2.$$

*Proof.* According to Cauchy-Schwarz inequality,

$$\begin{aligned} f(y) - f(x) &= \int_0^1 \nabla f((1-a)x + ay)^T (y - x) da \\ &= \int_0^1 [\nabla f((1-a)x + ay) - \nabla f(x)]^T (y - x) da + \int_0^1 \nabla f(x)^T (y - x) da \\ &\leq \|y - x\|_2 \int_0^1 \|\nabla f((1-a)x + ay) - \nabla f(x)\|_2 da + \nabla f(x)^T (y - x) \\ &\leq \frac{L}{2} \|y - x\|_2^2 + \nabla f(x)^T (y - x). \end{aligned}$$

□

### 3.3 Function Classes Complexity and Concentration Inequality

**Definition 6** (Rademacher complexity). Let  $n \in \mathbb{N}_{\geq 1}$ . The Rademacher complexity of a set  $A \subset \mathbb{R}^n$  is defined as

$$\mathfrak{R}_n(A) = \mathbb{E}_{\{\zeta_i\}_{i=1}^n} \left[ \sup_{a \in A} \frac{1}{n} \sum_{i=1}^n \zeta_i a_i \right],$$

where,  $\{\zeta_i\}_{i=1}^n$  are  $n$  i.i.d Rademacher variables with  $\mathbb{P}(\zeta_i = 1) = \mathbb{P}(\zeta_i = -1) = \frac{1}{2}$ . The Rademacher complexity of function class  $\mathcal{G}$  associate with random sample  $\{X_i\}_{i=1}^n$  is defined as

$$\mathfrak{R}_n(\mathcal{G}) = \mathbb{E}_{\{X_i, \zeta_i\}_{i=1}^n} \left[ \sup_{g \in \mathcal{G}} \frac{1}{n} \sum_{i=1}^n \zeta_i g(X_i) \right].$$

**Definition 7** (covering number). An  $\epsilon$ -cover of a set  $T$  in a metric space  $(S, \tau)$  is a subset  $T_c \subset S$  such that for each  $t \in T$ , there exists a  $t_c \in T_c$  such that  $\tau(t, t_c) \leq \epsilon$ . The  $\epsilon$ -covering number of  $T$ , denoted as  $\mathcal{N}(\epsilon, T, \tau)$  is defined to be the minimum cardinality among all  $\epsilon$ -cover of  $T$  with respect to the metric  $\tau$ .

The Rademacher complexity and the covering number of  $\mathcal{G}$  share the following relation.

**Lemma 14** ([34], Lemma 5.7). *Let  $n \in \mathbb{N}_{\geq 1}$ . For any function class  $\mathcal{G}$  with  $|g| \leq B_{\mathcal{G}}$  for all  $g \in \mathcal{G}$ ,*

$$\mathfrak{R}_n(\mathcal{G}) \leq \inf_{0 < \delta < B_{\mathcal{G}}/2} \left( 4\delta + \frac{12}{\sqrt{n}} \int_{\delta}^{B_{\mathcal{G}}/2} \sqrt{\ln \mathcal{N}(\epsilon, \mathcal{G}, \|\cdot\|_{\infty})} d\epsilon \right).$$

**Lemma 15** (McDiarmid's inequality). *Let  $n \in \mathbb{N}_{\geq 1}$ . Let  $g$  be a function from  $\Omega^n$  to  $\mathbb{R}$ . Suppose that function  $g$  satisfies the bounded differences property: there exists constants  $\{c_i\}_{i=1}^n$  such that for any  $x_1, \dots, x_n \in \Omega$ ,*

$$\sup_{\tilde{x}_i \in \Omega} |g(x_1, \dots, \tilde{x}_i, \dots, x_n) - g(x_1, \dots, x_i, \dots, x_n)| \leq c_i, \quad i \in [n].$$

Let  $\{X_i\}_{i=1}^n$  be independent variables, then for any  $\tau > 0$ ,

$$|g(X_1, \dots, X_n) - \mathbb{E}g(X_1, \dots, X_n)| \leq \tau$$

with probability at least  $1 - 2e^{-\frac{2\tau^2}{\sum_{i=1}^n c_i^2}}$ .

*Proof.* See [70, Theorem 2.9.1]. □

### 3.4 Miscellaneous

**Lemma 16.** *For  $0 < p < \frac{1}{e}$ ,*

$$\ln x \leq x^p, \quad x \in \left[ \left( \frac{2}{p} \ln \frac{1}{p} + \frac{1}{p} \right)^{1/p}, +\infty \right).$$

For  $p \geq \frac{1}{e}$ ,

$$\ln x \leq x^p, \quad x \in (0, +\infty).$$

*Proof.* Define  $g(x) := x^p - \ln x$ . Then  $g'(x) = px^{p-1} - \frac{1}{x}$ , from which we conclude that  $g(x)$  decreases on  $\left(0, \left(\frac{1}{p}\right)^{1/p}\right)$  and increases on  $\left(\left(\frac{1}{p}\right)^{1/p}, +\infty\right)$ .

For  $0 < p < \frac{1}{e}$ , assume  $\frac{k}{e^k} \leq p \leq \frac{k-1}{e^{k-1}}$  with some  $k \in \mathbb{N}_{\geq 2}$ . Since  $p \leq \frac{k-1}{e^{k-1}} \leq \frac{e^{(k-1)/2}}{e^{k-1}} = \frac{1}{e^{(k-1)/2}}$ , we have  $k \leq 2 \ln \frac{1}{p} + 1$ . The result follows from the facts that

$$g \left( \left( \frac{k}{p} \right)^{1/p} \right) = \frac{1}{p} \left( k - \ln \frac{k}{p} \right) \geq 0.$$

$$\text{and } \left( \frac{2}{p} \ln \frac{1}{p} + \frac{1}{p} \right)^{1/p} \geq \left( \frac{k}{p} \right)^{1/p} \geq \left( \frac{1}{p} \right)^{1/p}.$$

For  $p \geq \frac{1}{e}$ , the minimum

$$g\left(\left(\frac{1}{p}\right)^{1/p}\right) = \frac{1}{p}\left(1 - \ln\frac{1}{p}\right) \geq 0.$$

Hence  $g(x)$  is nonnegative on  $(0, +\infty)$ .  $\square$

**Lemma 17.** For  $q > e$ ,

$$y^q \leq e^y, \quad y \in [2q \ln q + q, +\infty).$$

For  $0 < q \leq e$ ,

$$y^q \leq e^y, \quad y \in (0, +\infty).$$

*Proof.* We can obtain the result directly by letting  $x = y^q$  and  $p = \frac{1}{q}$  in Lemma 16.  $\square$

**Lemma 18.** For any  $a, b \geq 0, p \geq 1$ ,

$$(a + b)^p \leq 2^{p-1}(a^p + b^p).$$

*Proof.* The function  $f(x) = x^p$  is convex on  $(0, +\infty)$  since  $f''(x) = p(p-1)x^{p-2} \geq 0$ . Then

$$\left(\frac{a+b}{2}\right)^p \leq \frac{1}{2}a^p + \frac{1}{2}b^p.$$

$\square$

## 4 Error Decomposition

In this section, we investigate some properties and relations of the continuous loss and the discrete loss. Building upon these results, we derive an error decomposition, which serves as the starting point for our subsequent work.

The continuous loss  $L_R$  and  $L_N$  have the following properties.

**Lemma 19.** For any  $u \in H^1(\Omega)$ ,

$$\begin{aligned} C(\text{coe})\|u - u_R\|_{H^1(\Omega)}^2 &\leq L_R(u) - L_R(u_R) \leq \max\{1, 1/\beta\}C(\Omega, \text{coe})\|u - u_R\|_{H^1(\Omega)}^2, \\ C(\text{coe})\|u - u_N\|_{H^1(\Omega)}^2 &\leq L_N(u) - L_N(u_N) \leq C(\Omega, \text{coe})\|u - u_R\|_{H^1(\Omega)}^2. \end{aligned}$$

*Proof.* We only present a proof for the Robin problem. The proof is adapted to the Neumann problem after minor modifications. For any  $u \in \mathcal{F}_{NN}$ , set  $v = u - u_R$ , then

$$L_R(u) = L_R(u_R + v)$$

$$\begin{aligned}
&= \frac{1}{2}(\nabla(u_R + v), \nabla(u_R + v))_{L^2(\Omega)} + \frac{1}{2}(u_R + v, u_R + v)_{L^2(\Omega; w)} - \langle u_R + v, f \rangle_{L^2(\Omega)} \\
&\quad + \frac{1}{2\beta}(T_0 u_R + T_0 v, T_0 u_R + T_0 v)_{L^2(\partial\Omega)} - \frac{1}{\beta}\langle T_0 u_R + T_0 v, g \rangle_{L^2(\partial\Omega)} \\
&= \frac{1}{2}(\nabla u_R, \nabla u_R)_{L^2(\Omega)} + \frac{1}{2}(u_R, u_R)_{L^2(\Omega; w)} - \langle u_R, f \rangle_{L^2(\Omega)} + \frac{1}{2\beta}(T_0 u_R, T_0 u_R)_{L^2(\partial\Omega)} \\
&\quad - \frac{1}{\beta}\langle T_0 u_R, g \rangle_{L^2(\partial\Omega)} + \frac{1}{2}(\nabla v, \nabla v)_{L^2(\Omega)} + \frac{1}{2}(v, v)_{L^2(\Omega; w)} + \frac{1}{2\beta}(T_0 v, T_0 v)_{L^2(\partial\Omega)} \\
&\quad + [(\nabla u_R, \nabla v)_{L^2(\Omega)} + (u^*, v)_{L^2(\Omega; w)} - \langle v, f \rangle_{L^2(\Omega)} \\
&\quad + \frac{1}{\beta}(T_0 u_R, T_0 v)_{L^2(\partial\Omega)} - \frac{1}{\beta}\langle T_0 v, g \rangle_{L^2(\partial\Omega)}] \\
&= L_R(u_R) + \frac{1}{2}(\nabla v, \nabla v)_{L^2(\Omega)} + \frac{1}{2}(v, v)_{L^2(\Omega; w)} + \frac{1}{2\beta}(T_0 v, T_0 v)_{L^2(\partial\Omega)},
\end{aligned}$$

where the last equality is due to the fact that  $u_R$  is the solution of equation (8a). Hence

$$\begin{aligned}
C(\text{coe})\|v\|_{H^1(\Omega)}^2 &\leq L_R(u) - L_R(u_R) \\
&= \frac{1}{2}(\nabla v, \nabla v)_{L^2(\Omega)} + \frac{1}{2}(v, v)_{L^2(\Omega; w)} + \frac{1}{2\beta}(T_0 v, T_0 v)_{L^2(\partial\Omega)} \\
&\leq \max\{1, 1/\beta\}C(\Omega, \text{coe})\|v\|_{H^1(\Omega)}^2,
\end{aligned}$$

where in the third step we make use of Lemma 5. □

The following result provides an estimation of the difference between the minimum value of the discrete loss and the minimum value of the continuous loss.

**Lemma 20.**

$$\begin{aligned}
&\left| \inf_{f_W \in \mathcal{F}_{NN}} \widehat{L}_R(f_W) - L_R(u_R) \right| \\
&\leq \inf_{f_W \in \mathcal{F}_{NN}} [L_R(f_W) - L_R(u_R)] \\
&\quad + \max \left\{ \sup_{f_W \in \mathcal{F}_{NN}} [\widehat{L}_R(f_W) - L_R(f_W)], \sup_{f_W \in \mathcal{F}_{NN}} [L_R(f_W) - \widehat{L}_R(f_W)] \right\}, \\
&\left| \inf_{f_W \in \mathcal{F}_{NN}} \widehat{L}_N(f_W) - L_R(u_N) \right| \\
&\leq \inf_{f_W \in \mathcal{F}_{NN}} [L_N(f_W) - L_N(u_N)] \\
&\quad + \max \left\{ \sup_{f_W \in \mathcal{F}_{NN}} [\widehat{L}_N(f_W) - L_N(u)], \sup_{f_W \in \mathcal{F}_{NN}} [L_N(f_W) - \widehat{L}_N(f_W)] \right\}.
\end{aligned}$$

*Proof.* We only give a proof for the Robin problem. Let  $f_{\bar{W}}$  be any function in  $\mathcal{F}_{NN}$ . We have

$$\begin{aligned} & \inf_{f_W \in \mathcal{F}_{NN}} \widehat{L}_R(f_W) - L_R(u_R) \leq \widehat{L}_R(f_{\bar{W}}) - L_R(u_R) \\ & = [\widehat{L}_R(f_{\bar{W}}) - L_R(f_{\bar{W}})] + [L_R(f_{\bar{W}}) - L_R(u_R)] \\ & \leq \sup_{f_W \in \mathcal{F}_{NN}} [\widehat{L}_R(f_W) - L_R(f_W)] + [L_R(f_{\bar{W}}) - L_R(u_R)], \end{aligned}$$

which implies

$$\begin{aligned} & \inf_{f_W \in \mathcal{F}_{NN}} \widehat{L}_R(f_W) - L_R(u_R) \\ & \leq \sup_{f_W \in \mathcal{F}_{NN}} [\widehat{L}_R(f_W) - L_R(f_W)] + \inf_{f_W \in \mathcal{F}_{NN}} [L_R(f_W) - L_R(u_R)]. \quad (14) \end{aligned}$$

On the other hand, let  $\{u_k\}_{k=1}^\infty \subseteq \mathcal{F}_{NN}$  be a sequence such that  $\lim_{k \rightarrow \infty} \widehat{L}_R(u_k) = \inf_{f_W \in \mathcal{F}_{NN}} \widehat{L}_R(f_W)$ . For any  $k \in \mathbb{N}_{\geq 1}$ ,

$$\begin{aligned} L_R(u_R) - \widehat{L}_R(u_k) & = [L_R(u_R) - L_R(u_k)] + [L_R(u_k) - \widehat{L}_R(u_k)] \\ & \leq L_R(u_k) - \widehat{L}_R(u_k) \leq \sup_{f_W \in \mathcal{F}_{NN}} [L_R(f_W) - \widehat{L}_R(f_W)]. \end{aligned}$$

Hence

$$L_R(u_R) - \inf_{f_W \in \mathcal{F}_{NN}} \widehat{L}_R(f_W) \leq \sup_{f_W \in \mathcal{F}_{NN}} [L_R(f_W) - \widehat{L}_R(f_W)]. \quad (15)$$

Combining (14) and (15) yields the conclusion.  $\square$

The following lemma decomposes the total error into three different types of errors, which we will handle using different tools.

**Lemma 21.** (1) Consider Robin problem (8a). Suppose  $f_{W_T} \in \mathcal{F}_{NN}$ . There holds

$$\begin{aligned} & L_R(f_{W_T}) - L_R(u_R) \\ & \leq \left[ \widehat{L}_R(f_{W_T}) - \inf_{f_W \in \mathcal{F}_{NN}} \widehat{L}_R(f_W) \right] + \inf_{f_W \in \mathcal{F}_{NN}} [L_R(f_W) - L_R(u_R)] \\ & \quad + 2 \max \left\{ \sup_{f_W \in \mathcal{F}_{NN}} [L_R(f_W) - \widehat{L}_R(f_W)], \sup_{f_W \in \mathcal{F}_{NN}} [\widehat{L}_R(f_W) - L_R(f_W)] \right\}. \end{aligned}$$

(2) Consider Neumann problem (7a). Suppose  $f_{W_T} \in \mathcal{F}_{NN}$ . There holds

$$\begin{aligned} & L_N(f_{W_T}) - L_N(u_N) \\ & \leq \left[ \widehat{L}_N(f_{W_T}) - \inf_{f_W \in \mathcal{F}_{NN}} \widehat{L}_N(f_W) \right] + \inf_{f_W \in \mathcal{F}_{NN}} [L_N(f_W) - L_N(u_N)] \\ & \quad + 2 \max \left\{ \sup_{f_W \in \mathcal{F}_{NN}} [L_N(f_W) - \widehat{L}_N(f_W)], \sup_{f_W \in \mathcal{F}_{NN}} [\widehat{L}_N(f_W) - L_N(f_W)] \right\}. \end{aligned}$$

**Remark 2.** *The first term reflects the distance between the discrete loss value at the  $T$ -th iteration and the minimum value of the discrete loss, and is therefore referred to as the optimization error. According to Lemma 19, the second term shows the distance between the set  $\mathcal{F}_{NN}$  and the target function  $u_R(u_N)$ , which is exactly the classical definition of approximation error. The third term is the generalization error, which measures the uniform difference between the continuous loss and the discrete loss over  $\mathcal{F}_{NN}$ .*

*Proof.* We only give a proof for the Robin problem. It follows from Lemma 20 that

$$\begin{aligned}
& L_R(f_{W_T}) - L_R(u_R) \\
&= [L_R(f_{W_T}) - \widehat{L}_R(f_{W_T})] + \left[ \widehat{L}_R(f_{W_T}) - \inf_{f_W \in \mathcal{F}_{NN}} \widehat{L}_R(f_W) \right] \\
&\quad + \left[ \inf_{f_W \in \mathcal{F}_{NN}} \widehat{L}_R(f_W) - L_R(u_R) \right] \\
&\leq \left[ \widehat{L}_R(f_{W_T}) - \inf_{f_W \in \mathcal{F}_{NN}} \widehat{L}_R(f_W) \right] + \inf_{f_W \in \mathcal{F}_{NN}} [L_R(f_W) - L_R(u_R)] \\
&\quad + 2 \max \left\{ \sup_{f_W \in \mathcal{F}_{NN}} [\widehat{L}_R(f_W) - L_R(f_W)], \sup_{f_W \in \mathcal{F}_{NN}} [L_R(f_W) - \widehat{L}_R(f_W)] \right\}.
\end{aligned}$$

where in the second step we use the fact that  $f_{W_T} \in \mathcal{F}_{NN}$ . □

## 5 Approximation Error

In this section, we study the error of neural network approximation for functions in Sobolev spaces. We follow the proof strategy presented in [18]. To be specific, the main process can be divided into two steps. According to the Bramble-Hilbert lemma (Lemma 4), there exists polynomials that can locally approximate functions in Sobolev spaces. The first step in [77] is to approximate those polynomials by neural network functions. Secondly, following the ideas presented in [77], [18] constructs an approximate partition of unity  $\{\Phi_j^N\}_{j \in \{1, 2, \dots, N\}^d}$  by neural network functions to achieve localization. However, the study in [18] did not cover all integer order Sobolev spaces, as they only studied the approximation of neural network functions to  $W^{s, \infty}$  functions. We generalize their result to the approximation of  $W^{s, p}$  functions with  $1 \leq p \leq \infty$ .

**Remark 3.** *Here, we consider only the tanh neural network, i.e.,  $\sigma = \tanh$ . However, as mentioned in [18], this construction method is applicable to other smooth activation functions.*

We first introduce some notations we need in this section. Let  $N \in \mathbb{N}_{\geq 1}$ .

For  $j \in \{1, \dots, N\}^d$ , define

$$I_j^N := \prod_{i=1}^d \left( \frac{j_i - 1}{N}, \frac{j_i}{N} \right), \quad J_j^N := \prod_{i=1}^d \left( \frac{j_i - 2}{N}, \frac{j_i + 1}{N} \right).$$

Define

$$\mathcal{V} := \left\{ v \in \mathbb{Z}^d : \max_{1 \leq i \leq d} |v_i| \leq 1 \right\}.$$

For some  $k \in \mathbb{N}_{\geq 1}$  and some accuracy parameter  $\epsilon > 0$ , let

$$\alpha = N \ln \left( \frac{(2k)^{k+1} (Nk)^k}{e^k \epsilon} \right). \quad (16)$$

For  $y \in \mathbb{R}$ , define

$$\begin{aligned} \phi_1^N(y) &:= \frac{1}{2} - \frac{1}{2} \sigma \left( \alpha \left( y - \frac{1}{N} \right) \right), \\ \phi_j^N(y) &:= \frac{1}{2} \sigma \left( \alpha \left( y - \frac{j-1}{N} \right) \right) - \frac{1}{2} \sigma \left( \alpha \left( y - \frac{j}{N} \right) \right) \quad \text{for } 2 \leq j \leq N-1, \\ \phi_N^N(y) &:= \frac{1}{2} \sigma \left( \alpha \left( y - \frac{N-1}{N} \right) \right) + \frac{1}{2}. \end{aligned}$$

and

$$\Phi_j^N(x) := \prod_{i=1}^d \phi_{j_i}^N(x_i), \quad x \in \mathbb{R}^d.$$

**Remark 4.** Compared to [18], we have removed  $R$  in the definition of  $\alpha$ , which does not affect the subsequent conclusions.

The following lemma shows that  $\{\Phi_j\}_{j \in \{1, 2, \dots, N\}^d}$  is an approximate partition of unity.

**Lemma 22** ([18], Lemma 4.1 and Lemma 4.2). *Let  $k \in \mathbb{N}_{\geq 0}$ . If  $0 < \epsilon < 1/4$  in (16), then*

$$\begin{aligned} \left\| \sum_{v \in \mathcal{V}} \Phi_{j+v}^N - 1 \right\|_{W^{k, \infty}(I_j^N)} &\leq 2^{dk} d \epsilon, \\ \|\Phi_{j+v}^N\|_{W^{k, \infty}(I_j^N)} &\leq \max \{1, (2k)^{2k} \alpha^k\} \epsilon, \quad \forall v \in \mathbb{Z}^d \text{ with } \|v\|_\infty \geq 2. \end{aligned}$$

The next lemma constructs neural networks  $\{f_{NN,i}^{(poly)}\}_{i \in \{1, 2, \dots, N\}^d}$  that approximate the target function in each local region  $I_i^N$  by utilizing the results of neural network approximation of polynomials ([18, Lemma 3.5]) and the Bramble-Hilbert lemma.

**Lemma 23.** Let  $N \in \mathbb{N}_{\geq 1}$ . Let  $i \in \{1, \dots, N\}^d$ . Let  $s, d \in \mathbb{N}, 1 \leq p \leq \infty$ . Let  $\epsilon > 0$ . Then for any  $f \in W^{s,p}((0,1)^d)$ , there exists a two-layer tanh neural network  $f_{NN,i}^{(poly)}$  with width no more than  $3(s-1) \lceil \frac{s}{2} \rceil \binom{s+d-2}{s-1}$  such that for  $k = 0, 1, \dots, s-1$ ,

$$\begin{aligned} \left\| f - f_{NN,i}^{(poly)} \right\|_{W^{k,p}(J_i^N)} &\leq C(s, d, p) \|f\|_{W^{s,p}(J_i^N)} \left( \left( \frac{1}{N} \right)^{s-k} + \epsilon \right), \\ \left\| f - f_{NN,i+v}^{(poly)} \right\|_{W^{k,p}(I_i^N)} &\leq \|f\|_{W^{k,p}(I_i^N)} + C(s, d, p) \|f\|_{W^{s-1,p}(J_{i+v}^N)}, \quad \forall \|v\|_\infty \geq 2. \end{aligned}$$

Furthermore, the weights of  $f_{NN,i}^{(poly)}$  are upper bounded by

$$C(s, d, p) \|f\|_{W^{s-1,p}(J_i^N)} N^{d/p} \epsilon^{-(s-1)/2}.$$

**Remark 5.** This lemma is crucial for extending the approximation of [18] from  $W^{s,\infty}$  functions to  $W^{s,p}$  functions. While [18] only utilizes the Bramble-Hilbert lemma for  $W^{s,\infty}$  functions, we employ the Bramble-Hilbert lemma for arbitrary  $W^{s,p}$  functions.

*Proof.* Let  $\Omega = J_i^N$  in Lemma 4, then there exists

$$f_i^{(poly)}(x) = \sum_{\tilde{s}=0}^{s-1} \sum_{\beta \in P_{\tilde{s},d}} c_{\beta,i} x^\beta.$$

with  $|c_{\beta,i}| \leq C(s, d) \left( \frac{3}{2N} \right)^{-d/p} \|f\|_{W^{s-1,p}(J_i^N)}$  such that for  $k = 0, 1, \dots, s-1$ ,

$$\left\| f - f_i^{(poly)} \right\|_{W^{k,p}(J_i^N)} \leq C(s, d) \left( \frac{3\sqrt{2}}{N} \right)^{s-k} \|f\|_{W^{s,p}(J_i^N)}. \quad (17)$$

By Lemma 3.5 in [18], there exists a two-layer tanh neural network  $f_{NN,\tilde{s}}^{(mono)} : (0,1)^d \rightarrow \mathbb{R}^{\binom{\tilde{s}+d-1}{\tilde{s}}}$  of width  $3 \lceil \frac{\tilde{s}+1}{2} \rceil \binom{\tilde{s}+d-1}{\tilde{s}}$  such that

$$\max_{\beta \in P_{\tilde{s},d}} \left\| x^\beta - \left( f_{NN,\tilde{s}}^{(mono)}(x) \right)_{\iota(\beta)} \right\|_{W^{k,\infty}((0,1)^d)} \leq \epsilon.$$

where  $\iota : P_{\tilde{s},d} \rightarrow \{1, \dots, \binom{\tilde{s}+d-1}{\tilde{s}}\}$  is a bijection. Furthermore, the weights of the network scale as  $O\left(\epsilon^{-\tilde{s}/2} (\tilde{s}(\tilde{s}+2))^{3(\tilde{s}+2)^2}\right)$  for small  $\epsilon$  and large  $\tilde{s}$ . Define

$$f_{NN,i}^{(poly)}(x) = c_{0,i} + \sum_{\tilde{s}=1}^{s-1} \sum_{\beta \in P_{\tilde{s},d}} c_{\beta,i} \left( f_{NN,\tilde{s}}^{(mono)}(x) \right)_{\iota(\beta)}.$$

There holds

$$\left\| f_i^{(poly)} - f_{NN,i}^{(poly)} \right\|_{W^{k,p}(J_i^N)} \leq \sum_{\tilde{s}=1}^{s-1} \sum_{\beta \in P_{\tilde{s},d}} |c_{\beta,i}| \left\| x^\beta - \left( f_{NN,\tilde{s}}^{(mono)}(x) \right)_{\iota(\beta)} \right\|_{W^{k,p}(J_i^N)}$$

$$\leq \epsilon \left( \frac{3}{N} \right)^{d/p} \sum_{\tilde{s}=1}^{s-1} \sum_{\beta \in P_{\tilde{s},d}} |c_{\beta,i}| \leq C(s,d,p) \|f\|_{W^{s-1,p}(J_i^N)} \epsilon. \quad (18)$$

Combining (17) and (18) yields the first result.

For  $\beta \in P_{\tilde{s},d}$ ,  $\|x^\beta\|_{W^{k,p}(I_j^N)} \leq k^{1/p} \tilde{s}^k N^{-d/p}$ . Thus we have for  $\|v\|_\infty \geq 2$ ,

$$\|f_{i+v}^{(poly)}\|_{W^{k,p}(I_i^N)} \leq \sum_{\tilde{s}=0}^{s-1} \sum_{\beta \in P_{\tilde{s},d}} |c_{\beta,i+v}| \|x^\beta\|_{W^{k,p}(I_i^N)} \leq C(s,d,p) \|f\|_{W^{s-1,p}(J_{i+v}^N)}.$$

By a similar derivation as obtaining (18), we have

$$\left\| f_{i+v}^{(poly)} - f_{NN,i+v}^{(poly)} \right\|_{W^{k,p}(I_i^N)} \leq C(s,d,p) \|f\|_{W^{s-1,p}(J_{i+v}^N)} \epsilon.$$

The second result then follows from the triangle inequality.  $\square$

The following lemma states that multiplication operations can be approximated by neural networks.

**Lemma 24** ([18], Corollary 3.7). *Let  $d \in \mathbb{N}, k \in \mathbb{N}_{\geq 0}$  and  $M \in \mathbb{R}_{>0}$ . Then for every  $\epsilon > 0$ , there exists a two-layer tanh neural network  $f_{NN}^{(mul)} : [-M, M]^d \rightarrow \mathbb{R}$  of width  $3 \lceil \frac{d+1}{2} \rceil \binom{2d-1}{d}$  such that*

$$\left\| f_{NN}^{(mul)}(x) - \prod_{i=1}^d x_i \right\|_{W^{k,\infty}([-M,M]^d)} \leq \epsilon.$$

Furthermore, the weights of the network are upper bounded by  $C(d,k)\epsilon^{-d/2} M^{3d^2/4+2d}$ .

**Remark 6.** *The upper bound for the weights in [18] does not explicitly depend on the range of the input variable, denoted as  $M$ . However, in the subsequent estimation of the convergence rate, we cannot simply treat  $M$  as a constant. Therefore, here we explicitly calculate the relationship between  $M$  and the impact on the weights based on the proof of Lemma 3.5 in [18].*

The following properties of tanh function is also needed.

**Lemma 25** ([18], Lemma A.4). *Let  $k \in \mathbb{N}$ . Then for all  $x \in \mathbb{R}$ ,*

$$\left| \sigma^{(k)}(x) \right| \leq (2k)^{k+1} \min\{\exp(-2x), \exp(2x)\}.$$

Now we are able to present a fundamental result on neural network function approximation to functions in  $W^{s,p}$ .

**Proposition 1.** *Let  $s, d \in \mathbb{N}_{\geq 1}, 1 \leq p \leq \infty$ . Let  $N \in \mathbb{N}_{\geq 1}$ . Then for any  $f \in W^{s,p}((0,1)^d)$ , there exists a three-layer neural network  $f_{NN}$  with the first hidden*

layer width of  $N^d \left[ 3(s-1) \left\lceil \frac{s}{2} \right\rceil \binom{s+d-2}{s-1} + 2d \right]$  and the second hidden layer width of  $3N^d \left\lceil \frac{d+2}{2} \right\rceil \binom{2d+1}{d+1}$ , such that for  $k = 1, \dots, s-1$ ,

$$\|f_{NN} - f\|_{W^{k,p}((0,1)^d)} \leq C(s, d, p, f) \ln^k \left( \frac{(2k)^{k+1} (Nk)^k N^{s+2d}}{e^k} \right) \left( \frac{1}{N} \right)^{s-k}.$$

Furthermore, the weights of  $f_{NN}$  are bounded by

$$C(s, d, p, f) N^{\frac{3}{4p}d^3 + \frac{k+p+7}{2p}d^2 + \frac{2sp+2k+2p+15}{4p}d + \frac{1}{2}(s^2+3)}$$

provided  $N \geq 2k$ .

*Proof.* Let  $\epsilon$  be  $\epsilon_1$  in (16). Let  $f_{NN,j}^{(poly)}$  be the neural network in Lemma 23 with  $\epsilon = \epsilon_2$  and  $f_{NN}^{(mul)}$  be the neural network in Lemma 24 with  $\epsilon = \epsilon_3$ . Let

$$f_{NN} = \sum_{j \in \{1, \dots, N\}^d} f_{NN}^{(mul)}(f_{NN,j}^{(poly)}(x), \phi_{j_1}^N(x_1), \dots, \phi_{j_d}^N(x_d)).$$

We decompose the total error into three terms and then bound each term in succession:

$$\begin{aligned} & \|f - f_{NN}\|_{W^{k,p}((0,1)^d)} \\ & \leq \left\| f - \sum_{j \in \{1, \dots, N\}^d} f \Phi_j^N \right\|_{W^{k,p}((0,1)^d)} + \sum_{j \in \{1, \dots, N\}^d} \left\| (f - f_{NN,j}^{(poly)}) \Phi_j^N \right\|_{W^{k,p}((0,1)^d)} \\ & \quad + \sum_{j \in \{1, \dots, N\}^d} \left\| f_{NN,j}^{(poly)}(x) \Phi_j^N(x) - f_{NN}^{(mul)}(f_{NN,j}^{(poly)}(x), \phi_{j_1}^N(x_1), \dots, \phi_{j_d}^N(x_d)) \right\|_{W^{k,p}((0,1)^d)}. \end{aligned} \tag{19}$$

For the first term, we firstly study the integral over  $I_i^N$  with  $i \in \{0, \dots, N\}^d$ . Applying Lemma 3 and Lemma 22, we have

$$\begin{aligned} & \left\| f - \sum_{j \in \{1, \dots, N\}^d} f \Phi_j^N \right\|_{W^{k,p}(I_i^N)} \\ & \leq C(k, d, p) \|f\|_{W^{k,p}(I_i^N)} \left\| 1 - \sum_{v \in \mathcal{V}} \Phi_{i+v}^N \right\|_{W^{k,\infty}(I_i^N)} \\ & \quad + C(k, d, p) \|f\|_{W^{k,p}(I_i^N)} \left\| \sum_{\substack{j \in \{1, \dots, N\}^d \\ j-i \notin \mathcal{V}}} \Phi_j^N \right\|_{W^{k,\infty}(I_i^N)} \\ & \leq C(k, d, p) \|f\|_{W^{k,p}(I_i^N)} (2^{kd} d \epsilon_1 + N^d (2k)^{2k} \alpha^k \epsilon_1) \end{aligned}$$

$$\leq C(s, d, p) \|f\|_{W^{s-1,p}(I_i^N)} \epsilon_1 N^{k+d} \ln^k \left( \frac{(2k)^{k+1} (Nk)^k}{e^k \epsilon_1} \right).$$

Then

$$\begin{aligned} & \left\| f - \sum_{j \in \{1, \dots, N\}^d} f \Phi_j^N \right\|_{W^{k,p}((0,1)^d)} \\ & \leq C(s, d, p) \|f\|_{W^{s-1,p}((0,1)^d)} \epsilon_1 N^{k+2d} \ln^k \left( \frac{(2k)^{k+1} (Nk)^k}{e^k \epsilon_1} \right). \end{aligned}$$

Similarly, for the second term of (19), we firstly handle the integration over  $I_i^N$ . Employing Lemma 3, we have

$$\begin{aligned} & \sum_{j \in \{1, \dots, N\}^d} \left\| \left( f - f_{NN,j}^{(poly)} \right) \Phi_j^N \right\|_{W^{k,p}(I_i^N)} \\ & \leq C(k, d, p) \sum_{j \in \{1, \dots, N\}^d} \sum_{\tilde{k}=0}^k \left\| f - f_{NN,j}^{(poly)} \right\|_{W^{k-\tilde{k},p}(I_i^N)} \left\| \Phi_j^N \right\|_{W^{\tilde{k},\infty}(I_i^N)} \\ & \leq C(k, d, p) \sum_{v \in \mathcal{V}} \sum_{\tilde{k}=0}^k \left\| f - f_{NN,i+v}^{(poly)} \right\|_{W^{k-\tilde{k},p}(I_i^N)} \left\| \Phi_{i+v}^N \right\|_{W^{\tilde{k},\infty}(I_i^N)} \\ & \quad + C(k, d, p) \sum_{\substack{j \in \{1, \dots, N\}^d \\ j-i \notin \mathcal{V}}} \left\| f - f_{NN,j}^{(poly)} \right\|_{W^{k,p}(I_i^N)} \left\| \Phi_j^N \right\|_{W^{k,\infty}(I_i^N)}. \end{aligned}$$

From (94) in [18] we have  $\left\| \Phi_{i+v}^N \right\|_{W^{\tilde{k},\infty}(I_i^N)} \leq N^{\tilde{k}} (2\tilde{k})^{2\tilde{k}} \ln^{\tilde{k}} \left( \frac{(2\tilde{k})^{\tilde{k}+1} (N\tilde{k})^{\tilde{k}}}{e^{\tilde{k}} \epsilon_1} \right)$ . By utilizing Lemma 23, we obtain an upper bound of

$$C(s, d, p) \|f\|_{W^{s,p}(J_i^N)} \ln^k \left( \frac{(2k)^{k+1} (Nk)^k}{e^k \epsilon_1} \right) \left( \left( \frac{1}{N} \right)^{s-k} + \epsilon_2 N^k \right)$$

for the first term. By employing Lemma 22 and Lemma 23, we obtain an upper bound for the second term as follows:

$$C(s, d, p) (N^d \|f\|_{W^{k,p}(I_i^N)} + N^{d(1-1/p)} \|f\|_{W^{s-1,p}((0,1)^d)}) N^k \ln^k \left( \frac{(2k)^{k+1} (Nk)^k}{e^k \epsilon_1} \right) \epsilon_1.$$

Combining these two bounds we get

$$\begin{aligned} & \sum_{j \in \{1, \dots, N\}^d} \left\| \left( f - f_{NN,j}^{(poly)} \right) \Phi_j^N \right\|_{W^{k,p}(I_i^N)} \\ & \leq C(s, d, p) \ln^k \left( \frac{(2k)^{k+1} (Nk)^k}{e^k \epsilon_1} \right) \\ & \quad \left( \|f\|_{W^{s,p}(J_i^N)} \left( N^{k+d} \epsilon_1 + \left( \frac{1}{N} \right)^{s-k} + \epsilon_2 N^k \right) + \|f\|_{W^{s-1,p}((0,1)^d)} N^{k+d(1-1/p)} \epsilon \right). \end{aligned}$$

By utilizing Lemma 18, we obtain the global estimation

$$\begin{aligned} & \sum_{j \in \{1, \dots, N\}^d} \left\| \left( f - f_{NN,j}^{(poly)} \right) \Phi_j^N \right\|_{W^{k,p}((0,1)^d)} \\ & \leq C(s, d, p) \|f\|_{W^{s,p}((0,1)^d)} \ln^k \left( \frac{(2k)^{k+1} (Nk)^k}{e^k \epsilon_1} \right) \left( N^{k+d} \epsilon_1 + \left( \frac{1}{N} \right)^{s-k} + N^k \epsilon_2 \right). \end{aligned}$$

Next, we bound the third term of (19). Based on a similar approach as obtaining (18), we have  $\|f_i^{(poly)} - f_{NN,i}^{(poly)}\|_{W^{k,\infty}((0,1)^d)} \leq C(s, d, p) \|f\|_{W^{s-1,p}(J_i^N)} N^{d/p} \epsilon_2$ . Furthermore, due to  $\|x^\beta\|_{W^{k,\infty}((0,1)^d)} \leq \tilde{s}^k$  for  $\beta \in P_{\tilde{s},d} := \{\alpha \in \mathbb{N}_{\geq 0}^d : |\alpha| = \tilde{s}\}$ ,

$$\begin{aligned} \|f_i^{(poly)}\|_{W^{k,\infty}((0,1)^d)} & \leq \sum_{\tilde{s}=1}^{s-1} \sum_{\beta \in P_{\tilde{s},d}} |c_{\beta,i}| \|x^\beta\|_{W^{k,\infty}((0,1)^d)} \\ & \leq C(s, d, p) \|f\|_{W^{s-1,p}(J_i^N)} N^{d/p}. \end{aligned}$$

Hence by triangle inequality we have

$$\|f_{NN,i}^{(poly)}\|_{W^{k,\infty}((0,1)^d)} \leq C(s, d, p) \|f\|_{W^{s-1,p}(J_i^N)} N^{d/p}.$$

For  $i \in [d]$ ,  $\|\phi_{j_i}^N\|_{W^{k,\infty}((0,1)^d)} \leq (2k)^{k+1} \alpha^k$  due to Lemma 25. Employing Lemma 24 with  $M = C(s, d, p) \|f\|_{W^{s-1,p}((0,1)^d)} N^{d/p}$  and Lemma 2, we have

$$\begin{aligned} & \|f_{NN}^{(mul)}(f_{NN,j}^{(poly)}(x), \phi_{j_1}^N(x_1), \dots, \phi_{j_d}^N(x_d)) - f_{NN,j}^{(poly)}(x) \Phi_j^N(x)\|_{W^{k,\infty}((0,1)^d)} \\ & \leq 16(e^2 k^4 d^2 (d+1))^k \left\| f_{NN}^{(mul)}(y) - \prod_{i=1}^{d+1} y_i \right\|_{W^{k,\infty}([-M,M]^{d+1})} \\ & \quad \max\{\|f_{NN,j}^{(poly)}\|_{W^{k,\infty}((0,1)^d)}^k, \|\phi_{j_1}^N\|_{W^{k,\infty}((0,1)^d)}^k, \dots, \|\phi_{j_d}^N\|_{W^{k,\infty}((0,1)^d)}^k\} \\ & \leq C(s, d, p) \epsilon_3 \left( \|f\|_{W^{s-1,p}(J_j^N)}^k N^{kd/p} + N^k \ln^k \left( \frac{(2k)^{k+1} (Nk)^k}{e^k \epsilon_1} \right) \right). \end{aligned}$$

Then

$$\begin{aligned} & \sum_{j \in \{1, \dots, N\}^d} \|f_{NN}^{(mul)}(f_{NN,j}^{(poly)}(x), \phi_{j_1}^N(x_1), \dots, \phi_{j_d}^N(x_d)) - f_{NN,j}^{(poly)}(x) \Phi_j^N(x)\|_{W^{k,p}((0,1)^d)} \\ & \leq C(s, d, p) \epsilon_3 \left( \|f\|_{W^{s-1,p}((0,1)^d)}^k N^{d+kd/p} + N^{k+d} \ln^k \left( \frac{(2k)^{k+1} (Nk)^k}{e^k \epsilon_1} \right) \right). \end{aligned}$$

Combining the estimates for the three terms of (19), we conclude that

$$\|f_{NN} - f\|_{W^{k,p}((0,1)^d)}$$

$$\begin{aligned} &\leq C(s, d, p) \|f\|_{W^{s,p}((0,1)^d)} \ln^k \left( \frac{(2k)^{k+1} (Nk)^k}{e^k \epsilon_1} \right) \left( N^{k+2d} \epsilon_1 + \left( \frac{1}{N} \right)^{s-k} + N^k \epsilon_2 \right) \\ &\quad + C(s, d, p) \left( \|f\|_{W^{s-1,p}((0,1)^d)}^k N^{d+kd/p} + N^{k+d} \ln^k \left( \frac{(2k)^{k+1} (Nk)^k}{e^k \epsilon_1} \right) \right) \epsilon_3. \end{aligned}$$

We finish the proof by further letting

$$\epsilon_1 = \left( \frac{1}{N} \right)^{s+2d}, \quad \epsilon_2 = \left( \frac{1}{N} \right)^s, \quad \epsilon_3 = \left( \frac{1}{N} \right)^{s+d+k \max\{d/p-1, 0\}}.$$

□

Below, we present the approximation error estimation that we need in this paper. This type of result not only demonstrates the existence of optimal approximation elements that allow neural network functions to sufficiently approximate the target function but also shows that neural network functions in the vicinity of the optimal approximation elements can sufficiently approximate the target function, too. We also need the result that the 2-norm of the outer parameters of the best approximation elements is sufficiently small. Following the ideas proposed in [37, 19], we achieve this by increasing the network width, which leads to a proportional decrease in the 2-norm of the outer parameters. We will see that such results contribute to the analysis of global convergence in the subsequent discussion of projected gradient descent.

**Theorem 2.** *Let  $d, A' \in \mathbb{N}_{\geq 1}$ . Let  $N \in \mathbb{N}_{\geq 1}$  and  $N \geq C(d)$ . Suppose that  $A, A'$  can be divided by  $N^d$ . For any  $f \in \bar{H}^2((0,1)^d)$  and any injection  $\varphi : \{1, 2, \dots, A'\} \rightarrow \{1, 2, \dots, A\}$  such that for  $k \in [A']$ ,*

$$\varphi(k) \in \{(k-1)A/A' + i' : i' \in [A/A']\},$$

*there exists  $W^* = \{(W^*)_s^l, (b^*)_s^l : l = 1, 2, 3; s \in [A]\}$  such that for  $W = \{(W_s^l, b_s^l) : l = 1, 2, 3; s \in [A]\}$  satisfying for  $s \in [A]$ ,  $W_s^3 = (W^*)_s^3, b_s^3 = (b^*)_s^3$ ; for  $s \in \{\varphi(1), \varphi(2), \dots, \varphi(A')\}, l = 1, 2$ ,*

$$\|W_s^l - (W^*)_s^l\|_F, \|b_s^l - (b^*)_s^l\|_2 \leq \left( \frac{1}{N} \right)^{\frac{3}{2}d^3 + 10d^2 + \frac{31}{2}d + \frac{29}{2}}, \quad (20)$$

*there holds*

$$\|f_W - f\|_{H^1(\Omega)} \leq C(d, f, \sigma, \Omega) \frac{1}{N^{1/2}}.$$

*Moreover,*

$$\begin{aligned} \|(W^*)^3\|_1 &\leq C(d, f) N^{\frac{3}{8}d^3 + \frac{5}{2}d^2 + \frac{37}{8}d + \frac{7}{2}}, \\ \|(W^*)^3\| &\leq C(d, f) \frac{1}{A'} N^{\frac{3}{8}d^3 + \frac{5}{2}d^2 + \frac{41}{8}d + \frac{7}{2}}. \end{aligned}$$

*Proof.* Let  $s = 2, k = 1, p = 2$  in Proposition 1 and let  $p = \frac{1}{2(2d+3)}$  in Lemma 16, we know that provided  $N \geq C(d)$ , there exists  $W^{**} = \{((W^{**})_s^l, (b^{**})_s^l) : l = 1, 2, 3; s \in [N^d]\}$  with  $(W^{**})_s^1 \in \mathbb{R}^{m_1 \times d}, (W^{**})_s^2 \in \mathbb{R}^{m_2 \times m_1}, (W^{**})_s^3 \in \mathbb{R}^{1 \times m_2}, (b^{**})_s^1 \in \mathbb{R}^{m_1}, (b^{**})_s^2 \in \mathbb{R}^{m_2}, (b^{**})_s^3 \in \mathbb{R}$  for  $s \in [N^d]$ , such that

$$\|f_{W^{**}} - f\|_{H^1((0,1)^d)} \leq C(d, f) \frac{1}{N^{1/2}}.$$

Furthermore, each component in  $(W^{**})_s^l$  and  $(b^{**})_s^l$  is bounded by  $C(d, f)N^{\frac{3}{8}d^3 + \frac{5}{2}d^2 + \frac{29}{8}d + \frac{7}{2}}$ . Define  $W^* = \{((W^*)_s^l, (b^*)_s^l) : l = 1, 2, 3; s \in [A]\}$  in the following way: for  $s \in [N^d]$ ,

$$\begin{aligned} (W^*)_j^3 &= (b^*)_j^3 = 0, \quad j \in \{1, 2, \dots, A\} \setminus \{\varphi(1), \varphi(2), \dots, \varphi(A')\}; \\ (W^*)_j^3 &= (b^*)_j^3 = 0, \quad j \in \{1, 2, \dots, A\} \setminus \{\varphi(1), \varphi(2), \dots, \varphi(A')\}; \\ (W^*)_j^3 &= (b^*)_j^3 = 0, \quad j \in \{1, 2, \dots, A\} \setminus \{\varphi(1), \varphi(2), \dots, \varphi(A')\}; \\ (W^*)_{(s-1)A'/N^d+i}^l &= (W^{**})_s^l, \quad (b^*)_{(s-1)A'/N^d+i}^l = (b^{**})_s^l, \quad l = 1, 2; i \in [A/N^d]. \end{aligned}$$

It is easy to see  $f_{W^*} \equiv f_{W^{**}}$  and

$$\begin{aligned} \|(W^*)^3\|_1 &= \|(W^{**})^3\|_1 \leq C(d, f)N^{\frac{3}{8}d^3 + \frac{5}{2}d^2 + \frac{37}{8}d + \frac{7}{2}}, \\ \|(W^*)^3\| &= \frac{N^d}{A'} \|(W^{**})^3\| \leq C(d, f) \frac{1}{A'} N^{\frac{3}{8}d^3 + \frac{5}{2}d^2 + \frac{41}{8}d + \frac{7}{2}}. \end{aligned}$$

Next we estimate  $\|f_W - f_{W^*}\|_{H^1(\Omega)}$  for  $W$  satisfying condition (20). We begin with the following two inequalities:

$$\begin{aligned} \|f_{s,1}^{org} - (f^*)_{s,1}^{org}\|_\infty &\leq \|W_s^1 - (W^*)_s^1\|_\infty + \|b_s^1 - (b^*)_s^1\|_\infty, \\ \|f_{s,2}^{org} - (f^*)_{s,2}^{org}\|_\infty &\leq \|W_s^2 - (W^*)_s^2\|_\infty \|f_s^1\|_\infty + \|(W^*)_s^2\|_\infty \|(f^*)_s^1 - (f^*)_s^1\|_\infty + \|b_s^2 - (b^*)_s^2\|_\infty \\ &\leq B_\sigma \|W_s^2 - (W^*)_s^2\|_\infty + \|b_s^2 - (b^*)_s^2\|_\infty + L_\sigma \|(W^*)_s^2\|_\infty \|W_s^1 - (W^*)_s^1\|_\infty \\ &\quad + L_\sigma \|(W^*)_s^2\|_\infty \|b_s^1 - (b^*)_s^1\|_\infty. \end{aligned}$$

With these two inequalities, we have

$$\begin{aligned} |f_W - f_{W^*}| &\leq \sum_{s=1}^A |f_s^3 - (f^*)_s^3| \leq \sum_{s=1}^A \|(W^*)_s^3\|_\infty \|f_s^2 - (f^*)_s^2\|_\infty \\ &\leq \sum_{s=1}^A \|(W^*)_s^3\|_\infty [L_\sigma B_\sigma \|W_s^2 - (W^*)_s^2\|_\infty + L_\sigma \|b_s^2 - (b^*)_s^2\|_\infty \\ &\quad + L_\sigma^2 \|(W^*)_s^2\|_\infty \|W_s^1 - (W^*)_s^1\|_\infty + L_\sigma^2 \|(W^*)_s^2\|_\infty \|b_s^1 - (b^*)_s^1\|_\infty]. \end{aligned}$$

Using the formula for calculating the gradient of  $f_W$ :

$$\nabla_x f_W = \sum_{s=1}^A (W_s^1)^T \text{diag}[\sigma'(f_{s,1}^{org})] (W_s^2)^T \text{diag}[\sigma'(f_{s,2}^{org})] (W_s^3)^T.$$

and the above two inequalities, we derive that

$$\begin{aligned}
& \|\nabla_x f_W - \nabla_x f_{W^*}\|_\infty \\
& \leq \sum_{s=1}^A [\|(W_s^1)^T - ((W^*)_s^1)^T\|_\infty \|\text{diag}[\sigma'(f_{s,1}^{org})](W_s^2)^T \text{diag}[\sigma'(f_{s,2}^{org})]((W^*)_s^3)^T\|_\infty \\
& \quad + \|((W^*)_s^1)^T\|_\infty \|\text{diag}[\sigma'(f_{s,1}^{org})] - \text{diag}[\sigma'((f^*)_{s,1}^{org})]\|_\infty \\
& \quad \quad \|(W_s^2)^T \text{diag}[\sigma'(f_{s,2}^{org})]((W^*)_s^3)^T\|_\infty \\
& \quad + \|((W^*)_s^1)^T \text{diag}[\sigma'((f^*)_{s,1}^{org})]\|_\infty \|(W_s^2)^T - ((W^*)_s^2)^T\|_\infty \\
& \quad \quad \|\text{diag}[\sigma'(f_{s,2}^{org})]((W^*)_s^3)^T\|_\infty \\
& \quad + \|((W^*)_s^1)^T \text{diag}[\sigma'((f^*)_{s,1}^{org})]((W^*)_s^2)^T\|_\infty \\
& \quad \quad \|\text{diag}[\sigma'(f_{s,2}^{org})] - \text{diag}[\sigma'((f^*)_{s,2}^{org})]\|_\infty \|((W^*)_s^3)^T\|_\infty] \\
& \leq \sum_{s=1}^A [\|((W^*)_s^3)^T\|_\infty [B_{\sigma'}^2 \| (W_s^2)^T \|_\infty \| (W_s^1)^T - ((W^*)_s^1)^T \|_\infty \\
& \quad + L_{\sigma'} B_{\sigma'} \| (W_s^2)^T \|_\infty \| ((W^*)_s^1)^T \|_\infty \| f_{s,1}^{org} - (f^*)_{s,1}^{org} \|_\infty \\
& \quad + B_{\sigma'}^2 \| ((W^*)_s^1)^T \|_\infty \| (W_s^2)^T - ((W^*)_s^2)^T \|_\infty \\
& \quad + L_{\sigma'} B_{\sigma'} \| ((W^*)_s^2)^T \|_\infty \| ((W^*)_s^1)^T \|_\infty \| f_{s,2}^{org} - (f^*)_{s,2}^{org} \|_\infty] \\
& \leq \sum_{s=1}^A [\|((W^*)_s^3)^T\|_\infty [B_{\sigma'}^2 \| ((W^*)_s^1)^T \|_\infty \| (W_s^2)^T - ((W^*)_s^2)^T \|_\infty \\
& \quad + L_{\sigma'} B_{\sigma'} B_{\sigma'} \| ((W^*)_s^2)^T \|_\infty \| ((W^*)_s^1)^T \|_\infty \| W_s^2 - (W^*)_s^2 \|_\infty \\
& \quad + L_{\sigma'} B_{\sigma'} \| ((W^*)_s^2)^T \|_\infty \| ((W^*)_s^1)^T \|_\infty \| b_s^2 - (b^*)_s^2 \|_\infty \\
& \quad + B_{\sigma'}^2 \| (W_s^2)^T \|_\infty \| (W_s^1)^T - ((W^*)_s^1)^T \|_\infty \\
& \quad + L_{\sigma'} B_{\sigma'} \| ((W^*)_s^1)^T \|_\infty \| x \|_\infty \\
& \quad \quad (L_{\sigma'} \| ((W^*)_s^2)^T \|_\infty \| (W^*)_s^2 \|_\infty + \| (W_s^2)^T \|_\infty) \| W_s^1 - (W^*)_s^1 \|_\infty \\
& \quad + L_{\sigma'} B_{\sigma'} \| ((W^*)_s^1)^T \|_\infty \\
& \quad \quad (L_{\sigma'} \| ((W^*)_s^2)^T \|_\infty \| (W^*)_s^2 \|_\infty + \| (W_s^2)^T \|_\infty) \| b_s^1 - (b^*)_s^1 \|_\infty].
\end{aligned}$$

Plugging the upper bound for the norm of weights and biases:

$$\begin{aligned}
& \|((W^*)_s^2)^T\|_\infty \leq m_2 B_2, \| (W^*)_s^2 \|_\infty \leq m_1 B_2, \\
& \|((W^*)_s^1)^T\|_\infty \leq m_1 B_1, \| (W^*)_s^1 \|_\infty \leq d B_1, \\
& \| (W^*)^3 \|_1 \leq C(d, f) N^{\frac{3}{8}} d^3 + \frac{5}{2} d^2 + \frac{37}{8} d + \frac{7}{2}, \\
& \| (W_s^2)^T \|_\infty \leq \| ((W^*)_s^2)^T \|_\infty + \| (W_s^2)^T - ((W^*)_s^2)^T \|_\infty \leq 2m_2 B_2.
\end{aligned}$$

with

$$m_1 = 3 |P_{1,d}| + 2d, m_2 = 3 \left\lceil \frac{d+2}{2} \right\rceil |P_{d+1,d+1}|,$$

$$B_{1,2} = C(d, f) N^{\frac{3}{8}d^3 + \frac{5}{2}d^2 + \frac{29}{8}d + \frac{7}{2}}$$

and simplifying the expressions, we finally obtain

$$\begin{aligned} \|f_W - f_{W^*}\|_{W^{1,\infty}(\Omega)} &= \max \left\{ \max_{x \in \Omega} |f_W - f_{W^*}|, \max_{x \in \Omega} \|\nabla_x f_W - \nabla_x f_{W^*}\|_\infty \right\} \\ &\leq C(d, f, \sigma) N^{\frac{3}{8}d^3 + \frac{5}{2}d^2 + \frac{27}{8}d + \frac{7}{2}} \max_{s=1,2,\dots,A} \\ &\quad [N^{\frac{3}{8}d^3 + \frac{5}{2}d^2 + \frac{29}{8}d + \frac{7}{2}} \|(W_s^2)^T - ((W^*)^2_s)^T\|_\infty + N^{\frac{3}{4}d^3 + 5d^2 + \frac{29}{4}d + 7} \|W_s^2 - (W^*)_s^2\|_\infty \\ &\quad + N^{\frac{3}{4}d^3 + 5d^2 + \frac{29}{4}d + 7} \|b_s^2 - (b^*)_s^2\|_\infty + N^{\frac{3}{8}d^3 + \frac{5}{2}d^2 + \frac{29}{8}d + \frac{7}{2}} \|(W_s^1)^T - ((W^*)_s^1)^T\|_\infty \\ &\quad + N^{\frac{9}{8}d^3 + \frac{15}{2}d^2 + \frac{87}{8}d + \frac{21}{2}} \|W_s^1 - (W^*)_s^1\|_\infty + N^{\frac{9}{8}d^3 + \frac{15}{2}d^2 + \frac{87}{8}d + \frac{21}{2}} \|b_s^1 - (b^*)_s^1\|_\infty] \\ &\leq C(d, f, \sigma) \frac{1}{N^{1/2}}, \end{aligned}$$

which implies the result due to triangle inequality and the fact that  $H^1$  norm can be controlled by  $W^{1,\infty}$  norm.  $\square$

## 6 Generalization Error

In this section, we utilize tools such as Rademacher complexity and covering numbers to handle generalization error

$$\max \left\{ \sup_{f_W \in \mathcal{F}_{NN}} [L_R(f_W) - \widehat{L}_R(f_W)], \sup_{f_W \in \mathcal{F}_{NN}} [\widehat{L}_R(f_W) - L_R(f_W)] \right\}$$

and

$$\max \left\{ \sup_{f_W \in \mathcal{F}_{NN}} [L_N(f_W) - \widehat{L}_N(f_W)], \sup_{f_W \in \mathcal{F}_{NN}} [\widehat{L}_N(f_W) - L_N(f_W)] \right\},$$

where  $\mathcal{F}_{NN}$  is an abbreviation for

$$\begin{aligned} \mathcal{F}_{NN}(\{m_1, m_2, A\}, B_{inn}, B_{out}) = \\ \{f_W : \|W_s^l\|_F, \|b_s^l\|_2 \leq B_{inn} (l = 1, 2; s \in [A]), \|W^3\|_1 \leq B_{out}\}. \end{aligned}$$

In this section we assume that the activation  $\sigma \in C^2(\Omega)$ .

**Lemma 26.**

$$\begin{aligned} &\mathbb{E}_{\{X_i\}_{i=1}^n, \{Y_j\}_{j=1}^m} \sup_{f_W \in \mathcal{F}_{NN}} \pm [L_R(f_W) - \widehat{L}_R(f_W)] \\ &\leq \sum_{k=1}^5 \mathbb{E}_{\{X_i\}_{i=1}^n, \{Y_j\}_{j=1}^m} \sup_{f_W \in \mathcal{F}_{NN}} \pm [L_k^{(R)}(f_W) - \widehat{L}_k^{(R)}(f_W)], \\ &\mathbb{E}_{\{X_i\}_{i=1}^n, \{Y_j\}_{j=1}^m} \sup_{f_W \in \mathcal{F}_{NN}} \pm [L_N(f_W) - \widehat{L}_N(f_W)] \end{aligned}$$

$$\leq \sum_{k=1}^5 \mathbb{E}_{\{X_i\}_{i=1}^n, \{Y_j\}_{j=1}^m} \sup_{f_W \in \mathcal{F}_{NN}} \pm \left[ L_k^{(N)}(f_W) - \widehat{L}_k^{(N)}(f_W) \right],$$

where

$$\begin{aligned} L_1^{(R)}(f_W) &= L_1^{(N)}(f_W) = \frac{|\Omega|}{2} \mathbb{E}_{X \sim U(\Omega)} \|\nabla_x f_W(X)\|_2^2, \\ L_2^{(R)}(f_W) &= L_2^{(N)}(f_W) = \frac{|\Omega|}{2} \mathbb{E}_{X \sim U(\Omega)} w(X) f_W^2(X), \\ L_3^{(R)}(f_W) &= L_3^{(N)}(f_W) = -|\Omega| \mathbb{E}_{X \sim U(\Omega)} f(X) f_W(X), \\ L_4^{(R)}(f_W) &= \frac{|\partial\Omega|}{2\beta} \mathbb{E}_{Y \sim U(\partial\Omega)} (T_0 f_W)^2(Y), \quad L_4^{(N)}(f_W) = 0, \\ L_5^{(R)}(f_W) &= -\frac{|\partial\Omega|}{\beta} \mathbb{E}_{Y \sim U(\partial\Omega)} g(Y) T_0 f_W(Y), \\ L_5^{(N)}(f_W) &= -|\partial\Omega| \mathbb{E}_{Y \sim U(\partial\Omega)} g(Y) T_0 f_W(Y), \end{aligned}$$

and  $\widehat{L}_k^{(R)}(f_W), \widehat{L}_k^{(N)}(f_W)$  is the discrete version of  $L_k^{(R)}(f_W), L_k^{(N)}(f_W)$ , for example,

$$\widehat{L}_1^{(R)}(f_W) = \frac{|\Omega|}{2n} \sum_{i=1}^n \|\nabla f_W(X_i)\|_2^2.$$

*Proof.* The result can be obtained easily from the property of supremum.  $\square$

By the symmetrization argument, we can convert the estimation of generalization error into an estimation of the Rademacher complexity of the relevant function classes.

**Lemma 27** ([34], Lemma 5.3). *Define*

$$\begin{aligned} \mathcal{F}_1 &:= \{\|\nabla_x f_W\|_2^2 : f_W \in \mathcal{F}_{NN}\}, & \mathcal{F}_2 &:= \{f_W^2 : f_W \in \mathcal{F}_{NN}\}, \\ \mathcal{F}_3 &:= \{f_W : f_W \in \mathcal{F}_{NN}\}, & \mathcal{F}_4 &:= \{f_W^2|_{\partial\Omega} : f_W \in \mathcal{F}_{NN}\}, \\ \mathcal{F}_5 &:= \{f_W|_{\partial\Omega} : f_W \in \mathcal{F}_{NN}\}. \end{aligned}$$

There holds

$$\begin{aligned} \mathbb{E}_{\{X_i\}_{i=1}^n} \sup_{f_W \in \mathcal{F}_{NN}} \pm \left[ L_1^{(R)}(f_W) - \widehat{L}_1^{(R)}(f_W) \right] &\leq C(\Omega, coe) \mathfrak{R}_n(\mathcal{F}_1), \\ \mathbb{E}_{\{X_i\}_{i=1}^n} \sup_{f_W \in \mathcal{F}_{NN}} \pm \left[ L_2^{(R)}(f_W) - \widehat{L}_2^{(R)}(f_W) \right] &\leq C(\Omega, coe) \mathfrak{R}_n(\mathcal{F}_2), \\ \mathbb{E}_{\{X_i\}_{i=1}^n} \sup_{f_W \in \mathcal{F}_{NN}} \pm \left[ L_3^{(R)}(f_W) - \widehat{L}_3^{(R)}(f_W) \right] &\leq C(\Omega, coe) \mathfrak{R}_n(\mathcal{F}_3), \\ \mathbb{E}_{\{X_i\}_{i=1}^n} \sup_{f_W \in \mathcal{F}_{NN}} \pm \left[ L_4^{(R)}(f_W) - \widehat{L}_4^{(R)}(f_W) \right] &\leq C(\Omega, coe) \max\{1, 1/\beta\} \mathfrak{R}_m(\mathcal{F}_4), \\ \mathbb{E}_{\{X_i\}_{i=1}^n} \sup_{f_W \in \mathcal{F}_{NN}} \pm \left[ L_5^{(R)}(f_W) - \widehat{L}_5^{(R)}(f_W) \right] &\leq C(\Omega, coe) \max\{1, 1/\beta\} \mathfrak{R}_m(\mathcal{F}_5) \end{aligned}$$

and

$$\begin{aligned}
\mathbb{E}_{\{X_i\}_{i=1}^n} \sup_{f_W \in \mathcal{F}_{NN}} \pm \left[ L_1^{(N)}(f_W) - \widehat{L}_1^{(N)}(f_W) \right] &\leq C(\Omega, \text{coe}) \mathfrak{R}_n(\mathcal{F}_1), \\
\mathbb{E}_{\{X_i\}_{i=1}^n} \sup_{f_W \in \mathcal{F}_{NN}} \pm \left[ L_2^{(N)}(f_W) - \widehat{L}_2^{(N)}(f_W) \right] &\leq C(\Omega, \text{coe}) \mathfrak{R}_n(\mathcal{F}_2), \\
\mathbb{E}_{\{X_i\}_{i=1}^n} \sup_{f_W \in \mathcal{F}_{NN}} \pm \left[ L_3^{(N)}(f_W) - \widehat{L}_3^{(N)}(f_W) \right] &\leq C(\Omega, \text{coe}) \mathfrak{R}_n(\mathcal{F}_3), \\
\mathbb{E}_{\{X_i\}_{i=1}^n} \sup_{f_W \in \mathcal{F}_{NN}} \pm \left[ L_4^{(N)}(f_W) - \widehat{L}_4^{(N)}(f_W) \right] &= 0, \\
\mathbb{E}_{\{X_i\}_{i=1}^n} \sup_{f_W \in \mathcal{F}_{NN}} \pm \left[ L_5^{(N)}(f_W) - \widehat{L}_5^{(N)}(f_W) \right] &\leq C(\Omega, \text{coe}) \mathfrak{R}_m(\mathcal{F}_5).
\end{aligned}$$

According to Lemma 14, we can estimate the Rademacher complexity of a given function class by derive an upper bound of the covering number of the function class. The following lemma provide upper bounds for the covering number of bounded smooth function classes.

**Lemma 28** ([64], Chapter 7, Theorem XIV). *Let  $s \in \mathbb{N}_{\geq 1}, B \in \mathbb{R}_{>0}$ . Let  $C_{\mathcal{B}(0,B)}^s(\Omega) := \{f \in C^s(\Omega) : \|f\|_{C^s(\Omega)} \leq B\}$ . We have for  $\epsilon > 0$ ,*

$$\ln \mathcal{N}(\epsilon, C_{\mathcal{B}(0,B)}^s(\Omega), \|\cdot\|_{\infty}) \leq C(s, d) \left( \frac{B}{\epsilon} \right)^{d/s}.$$

**Remark 7.** *The functions in set  $C_{\mathcal{B}(0,B)}^s(\Omega)$  have uniformly bounded derivatives, thus satisfying the conditions (64) and (67) in [64].*

**Remark 8.** *Theorem XIV in [64] does not explicitly state the dependence of the covering number on the parameter  $B$ , the upper bound for derivatives. However, this relationship can be immediately obtained using the technique of scaling.*

**Remark 9.** *In fact, Theorem XIV in [64] only provides upper bounds for the so-called minimal  $\epsilon$ -entropy and  $\epsilon$ -capacity, while what we need is an upper bound for the so-called  $\epsilon$ -entropy there. However, Theorem IV in [64] implies that for a totally bounded set, the  $\epsilon$ -entropy is always less than or equal to the  $\epsilon$ -capacity. On the other hand, the Arzela-Ascoli theorem guarantees that  $C_{\mathcal{B}(0,B)}^s(\Omega)$  is a totally bounded set.*

The following result shows that  $\{\mathcal{F}_i\}_{i=1}^5$ , the function classes we concern, are all subsets of bounded smooth function classes.

**Lemma 29.** *For function classes  $\{\mathcal{F}_i\}_{i=1}^5$  defined in Lemma 27, there holds*

$$\mathcal{F}_i \subset C_{\mathcal{B}(0, B_{\mathcal{F}_i})}^1(\Omega), \quad i = 1, 2, 3, 4, 5$$

with

$$\begin{aligned}
B_{\mathcal{F}_1} &= 2dm_2^2 m_1^{3/2} B_{\sigma'}^4 B_{\sigma''} B_{inn}^5 B_{out}^2, \\
B_{\mathcal{F}_2}, B_{\mathcal{F}_4} &= 2m_2^2 \sqrt{m_1} B_{\sigma'}^2 B_{\sigma''}^2 B_{inn} B_{out}^2, \\
B_{\mathcal{F}_3}, B_{\mathcal{F}_5} &= m_2 \sqrt{m_1} B_{\sigma'} B_{\sigma''}^2 B_{inn} B_{out}.
\end{aligned}$$

*Proof.* For any  $f_W \in \mathcal{F}_{NN}(\{m_1, m_2, A\}, B_{inn}, B_{out})$ ,

$$|f_W| \leq \sum_{s=1}^A \|W_s^3\|_F \|f_s^2\|_2 \leq m_2 B_\sigma \|W^3\|_1 \leq m_2 B_\sigma B_{out}.$$

where we employ the relation

$$\sum_{s=1}^A \|W_s^3\|_F \leq \sqrt{m_2} \sum_{s=1}^A \max_{k_2 \in [m_2]} |w_{s,k_2}^3| \leq \sqrt{m_2} \|W^3\|_1.$$

The first order derivative of  $f_W$  with respect to the spatial variable  $x$  can be calculated by

$$\frac{\partial f_W}{\partial x_j} = \sum_{s=1}^A (W_s^{1,j})^T \text{diag}[\sigma'(f_{s,1}^{org})] (W_s^2)^T \text{diag}[\sigma'(f_{s,2}^{org})] (W_s^3)^T, \quad j \in [d].$$

Hence

$$\begin{aligned} \left| \frac{\partial f_W}{\partial x_j} \right| &\leq \sum_{s=1}^A \|W_s^{1,j}\|_F \|\text{diag}[\sigma'(f_{s,1}^{org})]\|_F \|W_s^2\|_F \|\text{diag}[\sigma'(f_{s,2}^{org})]\|_F \|W_s^3\|_F \\ &\leq m_2 \sqrt{m_1} B_\sigma^2 B_{inn} B_{out}. \end{aligned}$$

By now, we have obtained upper bound estimates for the  $C^1$  norms of  $\mathcal{F}_3$  and  $\mathcal{F}_5$ . For any  $h \in \mathcal{F}_2$ , there exists  $f_W \in \mathcal{F}_{NN}$  such that  $h = f_W^2$ . Then  $\frac{\partial h}{\partial x_j} = 2f_W \frac{\partial f_W}{\partial x_j}$  and

$$\begin{aligned} |h| &= |f_W|^2 \leq m_2 B_\sigma B_{out}, \\ \left| \frac{\partial h}{\partial x_j} \right| &= 2|f_W| \left| \frac{\partial f_W}{\partial x_j} \right| \leq 2m_2 \sqrt{m_1} B_\sigma B_\sigma^2 B_{inn} B_{out}^2, \end{aligned}$$

which implies  $B_{\mathcal{F}_2}$ . We can derive  $B_{\mathcal{F}_4}$  in the same way. To estimate  $C^1$  norm of  $\mathcal{F}_1$ , we need to calculate the second order derivative of  $f_W$  with respect to the spatial variable  $x$ . For  $j, j' \in [d]$ ,

$$\begin{aligned} \frac{\partial^2 f_W}{\partial x_j \partial x_{j'}} &= \\ &\sum_{s=1}^A (W_s^{1,j'})^T [(\text{diag}[\sigma''(f_{s,1}^{org})] (W_s^2)^T \text{diag}[\sigma'(f_{s,2}^{org})] (W_s^3)^T) \odot W_s^{1,j} \\ &\quad + \text{diag}[\sigma'(f_{s,1}^{org})] (W_s^2)^T ((\text{diag}[\sigma''(f_{s,2}^{org})] W_s^2 \text{diag}[\sigma'(f_{s,1}^{org})] W_s^{1,j}) \odot (W_s^3)^T)]. \end{aligned}$$

Then

$$\left| \frac{\partial^2 f_W}{\partial x_j \partial x_{j'}} \right|$$

$$\begin{aligned}
&\leq \sum_{s=1}^A \|W_s^{1,j'}\|_F [\|\text{diag}[\sigma''(f_{s,1}^{org})]\|_F \|W_s^2\|_F \|\text{diag}[\sigma'(f_{s,2}^{org})]\|_F \|W_s^3\|_F \|W_s^{1,j}\|_F \\
&\quad + \|\text{diag}[\sigma'(f_{s,1}^{org})]\|_F \|W_s^2\|_F \|\text{diag}[\sigma''(f_{s,2}^{org})]\|_F \\
&\|W_s^2\|_F \|\text{diag}[\sigma'(f_{s,1}^{org})]\|_F \|W_s^{1,j}\|_F \|W_s^3\|_F] \\
&\leq m_2 m_1 B_{\sigma'}^2 B_{\sigma''} B_{inn}^4 B_{out}.
\end{aligned}$$

For any  $h \in \mathcal{F}_1$ , there exists  $f_W \in \mathcal{F}_{NN}$  such that  $h = \|\nabla_x f_W\|_2^2$ . It follows that

$$|h| \leq \sum_{j=1}^d \left| \frac{\partial f_W}{\partial x_j} \right|^2 \leq d m_2^2 m_1 B_{\sigma'}^4 B_{inn}^2 B_{out}^2.$$

Since  $\frac{\partial h}{\partial x_j} = \frac{\partial}{\partial x_j} \sum_{j'=1}^d \left( \frac{\partial f_W}{\partial x_{j'}} \right)^2 = 2 \sum_{j'=1}^d \frac{\partial f_W}{\partial x_{j'}} \frac{\partial^2 f_W}{\partial x_j \partial x_{j'}}$ , we have

$$\left| \frac{\partial h}{\partial x_j} \right| \leq 2 \sum_{j'=1}^d \left| \frac{\partial f_W}{\partial x_{j'}} \right| \left| \frac{\partial^2 f_W}{\partial x_j \partial x_{j'}} \right| \leq 2 d m_2^2 m_1^{3/2} B_{\sigma'}^4 B_{\sigma''} B_{inn}^5 B_{out}^2.$$

Combining the above two estimates we obtain  $B_{\mathcal{F}_1}$ .  $\square$

**Proposition 2.** *Let  $m = n$ . There holds*

$$\begin{aligned}
\mathbb{E}_{\{X_i\}_{i=1}^n, \{Y_j\}_{j=1}^m} \sup_{f_W \in \mathcal{F}_{NN}} \pm [L_R(f_W) - \widehat{L}_R(f_W)] &\leq C(d, \sigma, \Omega, coe) \max \left\{ 1, \frac{1}{\beta} \right\} \frac{B_{inn}^5 B_{out}^2}{\sqrt{n}}, \\
\mathbb{E}_{\{X_i\}_{i=1}^n, \{Y_j\}_{j=1}^m} \sup_{f_W \in \mathcal{F}_{NN}} \pm [L_N(f_W) - \widehat{L}_N(f_W)] &\leq C(d, \sigma, \Omega, coe) \frac{B_{inn}^5 B_{out}^2}{\sqrt{n}}.
\end{aligned}$$

*Proof.* We apply Lemma 14 to calculate an upper bound of  $\mathfrak{R}_n(\mathcal{F}_i)$  for  $i = 1, 2, 3, 4, 5$ :

$$\begin{aligned}
\mathfrak{R}_n(\mathcal{F}_i) &\leq \inf_{0 < \delta < B_{\mathcal{F}_i}/2} \left( 4\delta + \frac{12}{\sqrt{n}} \int_{\delta}^{B_{\mathcal{F}_i}/2} \sqrt{\ln \mathcal{N}(\epsilon, \mathcal{F}_i, \|\cdot\|_{\infty})} d\epsilon \right) \\
&\leq \inf_{0 < \delta < B_{\mathcal{F}_i}/2} \left( 4\delta + C(d) \frac{12}{\sqrt{n}} \int_{\delta}^{B_{\mathcal{F}_i}/2} \left( \frac{B_{\mathcal{F}_i}}{\epsilon} \right)^{d/2} d\epsilon \right) \\
&\leq \inf_{0 < \delta < B_{\mathcal{F}_i}/2} \left( 4\delta + C(d) \frac{B_{\mathcal{F}_i}}{\sqrt{n}} \right) = C(d) \frac{B_{\mathcal{F}_i}}{\sqrt{n}},
\end{aligned}$$

where in the second step we employ Lemma 28 with  $s = 1$ . The result is then implied by Lemma 26, 27 and 29.  $\square$

**Theorem 3.** *Let  $m = n$ . Let  $\tau \in \mathbb{R}_{>0}$ . With probability at least  $1 - 2e^{-\frac{\min\{1, \beta^2\} n \tau^2}{C(d, \sigma, coe, \Omega) B_{inn}^4 B_{out}^4}}$ ,*

$$\max \left\{ \sup_{f_W \in \mathcal{F}_{NN}} [\widehat{L}_R(f_W) - L_R(f_W)], \sup_{f_W \in \mathcal{F}_{NN}} [L_R(u) - \widehat{L}_R(f_W)] \right\}$$

$$\leq C(d, \sigma, \Omega, coe) \max\{1, 1/\beta\} \frac{B_{inn}^5 B_{out}^2}{\sqrt{n}} + \tau.$$

With probability at least  $1 - 2e^{-\frac{n\tau^2}{C(d, \sigma, coe, \Omega) B_{inn}^4 B_{out}^4}}$ ,

$$\begin{aligned} & \max \left\{ \sup_{f_W \in \mathcal{F}_{NN}} [\widehat{L}_N(f_W) - L_N(f_W)], \sup_{f_W \in \mathcal{F}_{NN}} [L_N(f_W) - \widehat{L}_N(f_W)] \right\} \\ & \leq C(d, \sigma, \Omega, coe) \frac{B_{inn}^5 B_{out}^2}{\sqrt{n}} + \tau. \end{aligned}$$

*Proof.* We only prove the inequality of the Robin problem since the inequality of the Neumann problem can be proved similarly. Define

$$h(X_1, \dots, X_n, Y_1, \dots, Y_m) := \sup_{f_W \in \mathcal{F}_{NN}} \left[ \widehat{L}_R(f_W) - L_R(f_W) \right].$$

We examine the difference of  $h$ :

$$\begin{aligned} & h(X_1, \dots, \widetilde{X}_i, \dots, X_n, Y_1, \dots, Y_m) - h(X_1, \dots, X_i, \dots, X_n, Y_1, \dots, Y_m) \\ & \leq \frac{|\Omega|}{n} \sup_{f_W \in \mathcal{F}_{NN}} \left[ \|\nabla_x f_W(\widetilde{X}_i)\|_2^2 + \frac{1}{2} w(\widetilde{X}_i) f_W^2(\widetilde{X}_i) - f(\widetilde{X}_i) f_W(\widetilde{X}_i) - \|\nabla_x f_W(X_i)\|_2^2 \right. \\ & \quad \left. - \frac{1}{2} w(X_i) f_W^2(X_i) + f(X_i) f_W(X_i) \right] \\ & \leq C(d, \sigma, coe, \Omega) \frac{B_{inn}^2 B_{out}^2}{n}. \end{aligned}$$

where in the second step we apply Lemma 29. We can bound

$$h(X_1, \dots, X_i, \dots, X_n, Y_1, \dots, Y_m) - h(X_1, \dots, \widetilde{X}_i, \dots, X_n, Y_1, \dots, Y_m)$$

similarly and hence obtain

$$\begin{aligned} & \left| h(X_1, \dots, \widetilde{X}_i, \dots, X_n, Y_1, \dots, Y_m) - h(X_1, \dots, X_i, \dots, X_n, Y_1, \dots, Y_m) \right| \\ & \leq C(d, \sigma, coe, \Omega) \frac{B_{inn}^2 B_{out}^2}{n}. \end{aligned}$$

In the same way we can show

$$\begin{aligned} & \left| h(X_1, \dots, X_n, Y_1, \dots, \widetilde{Y}_j, \dots, Y_m) - h(X_1, \dots, X_n, Y_1, \dots, Y_j, \dots, Y_m) \right| \\ & \leq C(d, \sigma, coe, \Omega) \max\{1, 1/\beta\} \frac{B_{inn}^2 B_{out}^2}{n}. \end{aligned}$$

Therefore according to Lemma 15, we have  $h - \mathbb{E}h \leq \tau$  with probability at least

$$1 - e^{-\frac{-\min\{1, \beta^2\} n \tau^2}{C(d, \sigma, coe, \Omega) B_{inn}^4 B_{out}^4}}. \text{ We can also show that } \sup_{f_W \in \mathcal{F}_{NN}} \left[ L_R(f_W) - \widehat{L}_R(f_W) \right] -$$

$\mathbb{E} \sup_{f_W \in \mathcal{F}_{NN}} [L_R(f_W) - \widehat{L}_R(f_W)] \leq \tau$  with probability at least  $1 - e^{-\frac{-\min\{1, \beta^2\} n \tau^2}{C(d, \sigma, coe, \Omega) B_{inn}^4 B_{out}^4}}$  in a similar manner. Then the estimate of

$$\max \left\{ \sup_{f_W \in \mathcal{F}_{NN}} [\widehat{L}_R(f_W) - L_R(f_W)], \sup_{f_W \in \mathcal{F}_{NN}} [L_R(f_W) - \widehat{L}_R(f_W)] \right\}$$

is implied by the union bound.  $\square$

## 7 Optimization Error

In this section we study the convergence of PGD:

$$\text{Robin problem: } W_{t+1} = \text{proj}_{\mathcal{C}}(W_t - \eta \nabla_W \widehat{L}_R(W_t)), \quad t = 0, 1, \dots, T-1;$$

$$\text{Neumann problem: } W_{t+1} = \text{proj}_{\mathcal{C}}(W_t - \eta \nabla_W \widehat{L}_N(W_t)), \quad t = 0, 1, \dots, T-1.$$

where  $\mathcal{C}$  is some projected set, which will be determined later. Our goal is to derive an upper bound of  $\widehat{L}_R(f_{W_T}) - \inf_{f_W \in \mathcal{F}_{NN}} \widehat{L}_R(f_W)$  and  $\widehat{L}_N(f_{W_T}) - \inf_{f_W \in \mathcal{F}_{NN}} \widehat{L}_N(f_W)$ , where  $\mathcal{F}_{NN}$  is an abbreviation for

$$\begin{aligned} \mathcal{F}_{NN}(\{m_1, m_2, A\}, B_{inn}, B_{out}) = \\ \{f_W : \|W_s^l\|_F, \|b_s^l\|_2 \leq B_{inn} (l = 1, 2; s \in [A]), \|W^3\|_1 \leq B_{out}\}. \end{aligned}$$

In this section we assume that the activation  $\sigma \in C^2(\Omega)$ .

**Lemma 30.** *Define*

$$\|\nabla_{W^3} \widehat{L}_R(f_W)\| := \left[ \sum_{s=1}^A (\|\nabla_{W_s^3} \widehat{L}_R(f_W)\|_F^2 + \|\nabla_{b_s^3} \widehat{L}_R(f_W)\|_2^2) \right]^{1/2}.$$

For  $f_W \in \mathcal{F}_{NN}(\{m_1, m_2, A\}, B_{inn}, B_{out})$ ,

$$\|\nabla_{W^3} \widehat{L}_R(f_W)\| \leq C(d, \sigma, coe, \Omega) \max\{1, 1/\beta\} \sqrt{A} B_{inn}^4 B_{out},$$

$$\|\nabla_{W^3} \widehat{L}_N(f_W)\| \leq C(d, \sigma, coe, \Omega) \sqrt{A} B_{inn}^4 B_{out}.$$

*Proof.* We only present a proof for  $\widehat{L}_R(f_W)$ . The inequality of  $\widehat{L}_N(f_W)$  can be proved in almost the same way.

It suffice to derive upper bounds of  $\|\nabla_{W_s^3} \widehat{L}_R(f_W)\|_F$  and  $\left| \frac{\partial}{\partial b_s^3} \widehat{L}_R(f_W) \right|$ . By definition,

$$\begin{aligned} & \nabla_{W_s^3} \widehat{L}_R(f_W) \\ &= \frac{|\Omega|}{n} \sum_{i=1}^n \left[ \sum_{j=1}^d \frac{\partial f_W(X_i)}{\partial x_j} \nabla_{W_s^3} \left( \frac{\partial f_W(X_i)}{\partial x_j} \right) + w(X_i) f_W(X_i) \nabla_{W_s^3} f_W(X_i) \right. \\ & \quad \left. - f(X_i) \nabla_{W_s^3} f_W(X_i) \right] \end{aligned}$$

$$+ \frac{|\partial\Omega|}{\beta m} \sum_{i=1}^m [f_W(Y_i) \nabla_{W_s^3} f_W(Y_i) - g(Y_j) \nabla_{W_s^3} f_W(Y_j)],$$

from which we can see that in order to obtain upper bound of  $\|\nabla_{W_s^3} \widehat{L}_R(f_W)\|_F$ , it suffices to study  $|f_W|$ ,  $\left|\frac{\partial f_W}{\partial x_j}\right|$ ,  $\|\nabla_{W_s^3} f_W\|_F$ ,  $\left\|\nabla_{W_s^3} \left(\frac{\partial f_W}{\partial x_j}\right)\right\|_F$ . But in the proof of Lemma 29 we already have  $|f_W| \leq m_2 B_\sigma B_{out}$  and  $\left|\frac{\partial f_W}{\partial x_j}\right| \leq m_2 \sqrt{m_1} B_\sigma^2 B_{inn} B_{out}$ . Since  $\nabla_{W_s^3} f_W = (f_s^2)^T$ , we have

$$\|\nabla_{W_s^3} f_W\|_F = \|f_s^2\|_F \leq \sqrt{m_2} B_\sigma.$$

The gradient of  $\frac{\partial f_W}{\partial x_j}$  with respect to  $W_s^3$  can be calculated by

$$\nabla_{W_s^3} \left(\frac{\partial f_W}{\partial x_j}\right) = (W_s^{1,j})^T \text{diag}[\sigma'(f_{s,1}^{org})] (W_s^2)^T \text{diag}[\sigma'(f_{s,2}^{org})].$$

where  $W_s^{1,j}$  is the  $j$ th column of  $W_s^1$ . Thus

$$\begin{aligned} \left\|\nabla_{W_s^3} \left(\frac{\partial f_W}{\partial x_j}\right)\right\|_F &\leq \|W_s^{1,j}\|_F \|\text{diag}[\sigma'(f_{s,1}^{org})]\|_F \|W_s^2\|_F \|\text{diag}[\sigma'(f_{s,2}^{org})]\|_F \\ &\leq \sqrt{m_2 m_1} B_\sigma^2 B_{inn}^2. \end{aligned}$$

Combining the above estimates, we derive that

$$\|\nabla_{W_s^3} \widehat{L}_R(f_W)\|_F \leq C(d, \sigma, coe, \Omega) \max\{1, 1/\beta\} B_{inn}^4 B_{out}.$$

We can derive in the same way that

$$\left|\frac{\partial}{\partial b_s^3} \widehat{L}_R(f_W)\right| \leq C(d, \sigma, coe, \Omega) \max\{1, 1/\beta\} B_{out}.$$

□

The following lemma present estimations for the strongly smooth parameter of  $\widehat{L}_R$  and  $\widehat{L}_N$ .

**Lemma 31.** *Define*

$$\begin{aligned} &\|\nabla_W \widehat{L}_R(f_W) - \nabla_W \widehat{L}_R(f_{\widetilde{W}})\| := \\ &\left[ \sum_{s=1}^A \sum_{l=1}^3 (\|\nabla_{W_s^l} \widehat{L}_R(f_W) - \nabla_{W_s^l} \widehat{L}_R(f_{\widetilde{W}})\|_F^2 + \|\nabla_{b_s^l} \widehat{L}_R(f_W) - \nabla_{b_s^l} \widehat{L}_R(f_{\widetilde{W}})\|_2^2) \right]^{1/2} \end{aligned}$$

and

$$\|W - \widetilde{W}\| := \left[ \sum_{s=1}^A \sum_{l=1}^3 \|W_s^l - (\widetilde{W})_s^l\|_F^2 + \|b_s^l - (\widetilde{b})_s^l\|_2^2 \right]^{1/2}.$$

For  $f_W \in \mathcal{F}_{NN}(\{m_1, m_2, A\}, B_{inn}, B_{out})$ , there holds

$$\begin{aligned} \|\nabla_W \widehat{L}_R(f_W) - \nabla_W \widehat{L}_R(f_{\widetilde{W}})\| &\leq C(d, \sigma, \text{coe}, \Omega) \max\{1/\beta, 1\} AB_{inn}^7 B_{out}^3 \|W - \widetilde{W}\|, \\ \|\nabla_W \widehat{L}_N(f_W) - \nabla_W \widehat{L}_N(f_{\widetilde{W}})\| &\leq C(d, \sigma, \text{coe}, \Omega) AB_{inn}^7 B_{out}^3 \|W - \widetilde{W}\|. \end{aligned}$$

*Proof.* Since the proof is too lengthy, we put it in the appendix.  $\square$

The following result shows that the discrete losses are convex with respect to the neural network parameters in the outer layer. This property helps to ensure the global convergence of optimization algorithms.

**Lemma 32.** *Given  $W^1$  and  $W^2$ ,  $\widehat{L}_R(f_W)$  and  $\widehat{L}_N(f_W)$  are convex with respect to  $W^3$ .*

*Proof.* We only present a proof for  $\widehat{L}_R(f_W)$ . The convexity of  $\widehat{L}_N(f_W)$  can be proved in almost the same way. Write  $W^3$  as  $W^3 = (W_1^3, b_1^3, W_2^3, b_2^3, \dots, W_A^3, b_A^3)^T \in \mathbb{R}^{A(m_2+1)}$ . Let  $D_s = (W_s^1)^T \text{diag}[\sigma'(f_{s,1}^{org})](W_s^2)^T \text{diag}[\sigma'(f_{s,2}^{org})] \in \mathbb{R}^{d \times m_2}$  for  $s \in [A]$  and  $D = (D_1, 0, D_2, 0, \dots, D_A, 0) \in \mathbb{R}^{d \times A(m_2+1)}$ , then  $\nabla_x f_W = \sum_{s=1}^A D_s (W_s^3)^T = DW^3$ . Let  $\overline{f}^2 = ((f_1^2)^T, 1, (f_2^2)^T, 1, \dots, (f_A^2)^T, 1) \in \mathbb{R}^{1 \times A(m_2+1)}$ , then  $f_W = \sum_{s=1}^A (W_s^3 f_s^2 + b_s^3) = \overline{f}^2 W^3$  and  $f_W^2 = (W^3)^T (\overline{f}^2)^T \overline{f}^2 W^3$ . Therefore we can express  $\widehat{L}_R(f_W)$  as

$$\widehat{L}_R(f_W) = (W^3)^T H W^3 - \left[ \frac{|\Omega|}{n} \sum_{i=1}^n f^{(source)}(X_i) \overline{f}^2(X_i) + \frac{|\partial\Omega|}{\beta m} \sum_{i=1}^m g(Y_i) \overline{f}^2(Y_i) \right] W^3,$$

where the Hessian matrix

$$\begin{aligned} H &= \frac{|\Omega|}{n} \sum_{i=1}^n D(X_i)^T D(X_i) + \frac{|\Omega|}{2n} \sum_{i=1}^n w(X_i) (\overline{f}^2(X_i))^T \overline{f}^2(X_i) \\ &\quad + \frac{|\partial\Omega|}{2\beta m} \sum_{i=1}^m (\overline{f}^2(Y_i))^T \overline{f}^2(Y_i). \end{aligned}$$

Since  $\|w\|_{L^\infty(\Omega)} \geq c_w$ ,  $H$  is positive semi-definite. According to Lemma 10, we conclude that  $\widehat{L}_R(f_W)$  is convex.  $\square$

According to [37], as long as the number of subnetworks,  $A$ , is sufficiently large, with high probability close to 1, there exist  $A'$  subnetworks whose parameters initialized randomly are sufficiently close to the parameters of the subnetwork corresponding to a given neural network function in  $\mathcal{F}_{NN}$ .

**Lemma 33.** *Let  $N, A' \in \mathbb{N}_{>0}$ . Let  $f^*$  be the target function to be estimated. Let  $f_{\widetilde{W}}$  be a given function in  $\mathcal{F}_{NN}$ . Suppose that  $A$  can be divided by  $A'$ . Let  $(W_0)_{s,i,j}^l, (b_0)_{s,i}^l (s \in [A]; l = 1, 2; i \in [m_l]; j \in [m_{l-1}])$  be uniformly distributed on*

$$\left[ -C(d, f^*) N^{10d^3}, C(d, f^*) N^{10d^3} \right].$$

Let  $E$  be the event that there exists an injection  $\varphi : \{1, 2, \dots, A'\} \rightarrow \{1, 2, \dots, A\}$  such that for  $k \in [A']$ ,

$$\varphi(k) \in \{(k-1)A/A' + i' : i' \in [A/A']\}$$

and for  $l = 1, 2; i \in [m_l]; j \in [m_{l-1}]; s \in \{\varphi(1), \dots, \varphi(A')\}$ ,

$$\begin{aligned} |(W_0)_{s,i,j}^l - (\bar{W})_{s,i,j}^l| &\leq \frac{1}{2\sqrt{m_l m_{l-1}}} \left(\frac{1}{N}\right)^{83d^3/2}, \\ |(b_0)_{s,i}^l - (\bar{b})_{s,i}^l| &\leq \frac{1}{2\sqrt{m_l}} \left(\frac{1}{N}\right)^{83d^3/2}. \end{aligned}$$

We claim that  $P(E) \geq 1 - A'e^{-C(d,f^*)} \frac{A}{A'} \left(\frac{1}{N}\right)^{206d^4(d+3)5^{d+2}}$ .

*Proof.* We follow the idea in [37]. Given  $k \in [A']$ ,  $s \in \{(k-1)A/A' + i' : i' \in [A/A']\}$ , let  $E_{k,s}$  be the event that for  $l = 1, 2; i \in [m_l]; j \in [m_{l-1}]$ ,

$$\begin{aligned} |(W_0)_{s,i,j}^l - (\bar{W})_{s,i,j}^l| &\leq \frac{1}{2\sqrt{m_l m_{l-1}}} \left(\frac{1}{N}\right)^{\frac{3}{2}d^3 + 10d^2 + \frac{31}{2}d + \frac{29}{2}}, \\ |(b_0)_{s,i}^l - (\bar{b})_{s,i}^l| &\leq \frac{1}{2\sqrt{m_l}} \left(\frac{1}{N}\right)^{\frac{3}{2}d^3 + 10d^2 + \frac{31}{2}d + \frac{29}{2}}. \end{aligned}$$

Then

$$E = \bigcap_{k \in [A']} \bigcup_{s \in \{(k-1)A/A' + i' : i' \in [A/A']\}} E_{k,s}.$$

The probability of  $E_{k,s}$  being true can be calculated directly.

$$\begin{aligned} P(E_{k,s}) &= \left(\frac{1}{\sqrt{m_1 d}}\right)^{m_1 d} \left(\frac{1}{\sqrt{m_1}}\right)^{m_1} \left(\frac{1}{\sqrt{m_2 m_1}}\right)^{m_2 m_1} \left(\frac{1}{\sqrt{m_2}}\right)^{m_2} \\ &\quad \left(\frac{1}{N}\right)^{83d^3(m_1 d + m_1 + m_2 m_1 + m_2)/2} \left(\frac{1}{C(d, f^*) N^{10d^3}}\right)^{(m_1 d + m_1 + m_2 m_1 + m_2)} \\ &\geq C(d, f^*) \left(\frac{1}{N}\right)^{206d^4(d+3)5^{d+2}}, \end{aligned}$$

where we use that  $m_1 = 5d, m_2 = 3 \lceil \frac{d+2}{2} \rceil \binom{2d+1}{d+1} \leq \frac{3}{2}(d+3)5^{d+1}$ . Denote

$$E_k = \bigcup_{s \in \{(k-1)A/A' + i' : i' \in [A/A']\}} E_{k,s}.$$

Then

$$P(E_k^c) = P\left(\bigcap_{s \in \{(k-1)A/A' + i' : i' \in [A/A']\}} E_{k,s}^c\right)$$

$$= \prod_{s \in \{(k-1)A/A' + i' : i' \in [A/A']\}} P(E_{k,s}^c) \leq \left[ 1 - C(d, f^*) \left( \frac{1}{N} \right)^{206d^4(d+3)5^{d+2}} \right]^{A/A'}.$$

By union bound we have

$$P(E^c) = P\left( \bigcup_{k \in [A']} E_k^c \right) \leq A' \left[ 1 - C(d, f^*) \left( \frac{1}{N} \right)^{206d^4(d+3)5^{d+2}} \right]^{A/A'}.$$

Applying the inequality  $e^x \geq 1 + x$  for any  $x \in \mathbb{R}$ , we finish the proof.  $\square$

The following result, with the initial idea from [19], demonstrates that the optimization error can be controlled by the sum of four terms. We can see that generalization error and approximation error occur again here. The difference from traditional approximation error lies in the fact that the approximation error here considers not only the approximation properties of a single neural network function but also the approximation properties of all neural network functions in the vicinity of a given neural network function.

**Theorem 4.** *Let  $N, A' \in \mathbb{N}_{>0}$ . Suppose that  $A$  can be divided by  $A'$ . Let  $(W_0)_{s,i,j}^l, (b_0)_{s,i}^l (s \in [A]; l = 1, 2; i \in [m_l]; j \in [m_{l-1}])$  be uniformly distributed on*

$$\left[ -C(d, \sigma, \text{coe}, \Omega)N^{10d^3}, C(d, \sigma, \text{coe}, \Omega)N^{10d^3} \right].$$

Let  $B_{inn} = C(d, \text{coe}, \Omega)N^{10d^3}$ ,  $B_{out} = C(d, \text{coe}, \Omega)N^{11d^3}$ . Let

$$\mathcal{C} = \left\{ W : \|W^3\|_1 \leq C(d, \text{coe}, \Omega)N^{11d^3}; \right. \\ \left. \|W_s^l - (W_0)_s^l\|_F, \|b_s^l - (b_0)_s^l\|_2 \leq \frac{1}{2} \left( \frac{1}{N} \right)^{83d^3/2}, l = 1, 2, s \in [A] \right\}.$$

Let  $f_{\bar{W}}$  be a given function in  $\mathcal{F}_{NN}$ . Let

$$\mathcal{B}_{\bar{W}} = \\ \left\{ W : W^3 = \bar{W}^3; \exists \varphi \text{ s.t. } \varphi(k) \in \{(k-1)A/A' + i' : i' \in [A/A']\} \text{ for } k \in [A'] \text{ and} \right. \\ \left. \|W_s^l - (\bar{W})_s^l\|_F, \|b_s^l - (\bar{b})_s^l\|_2 \leq \left( \frac{1}{N} \right)^{83d^3/2}, s \in \{\varphi(1), \dots, \varphi(A')\}, l = 1, 2 \right\}.$$

Let

$$\eta \leq C(d, \sigma, \text{coe}, \Omega) \left( \frac{1}{N} \right)^{103d^3} \frac{1}{A}.$$

With probability at least  $1 - A'e^{-C(d, f^*) \frac{A}{A'} \left( \frac{1}{N} \right)^{206d^4(d+3)5^{d+2}}}$ ,

$$\widehat{L}_R(f_{W_T}) - \inf_{f_W \in \mathcal{F}_{NN}} \widehat{L}_R(f_W)$$

$$\begin{aligned}
&\leq 2 \max \left\{ \sup_{f_W \in \mathcal{F}_{NN}} [\widehat{L}_R(f_W) - L_R(f_W)], \sup_{f_W \in \mathcal{F}_{NN}} [L_R(f_W) - \widehat{L}_R(f_W)] \right\} \\
&\quad + 2 \sup_{W \in \mathcal{B}_{\bar{W}}} [L_R(f_W) - L_R(u_R)] + \frac{1}{2\eta T} \|(W_0)^3 - (\bar{W})^3\|^2 + \frac{\eta}{2T} \sum_{t=0}^{T-1} \|\nabla_{W^3} \widehat{L}_R(f_{W_t})\|^2, \\
&\widehat{L}_N(f_{W_T}) - \inf_{f_W \in \mathcal{F}_{NN}} \widehat{L}_N(f_W) \\
&\leq 2 \max \left\{ \sup_{f_W \in \mathcal{F}_{NN}} [\widehat{L}_N(f_W) - L_N(f_W)], \sup_{f_W \in \mathcal{F}_{NN}} [L_N(f_W) - \widehat{L}_N(f_W)] \right\} \\
&\quad + 2 \sup_{W \in \mathcal{B}_W} [L_N(f_W) - L_N(u_N)] + \frac{1}{2\eta T} \|(W_0)^3 - (\bar{W})^3\|^2 + \frac{\eta}{2T} \sum_{t=0}^{T-1} \|\nabla_{W^3} \widehat{L}_N(f_{W_t})\|^2.
\end{aligned}$$

*Proof.* We only present a proof for the Robin problem. The Neumann problem can be proved in the same way.

Let  $\bar{W}_t = \{((W_t)_s^1, (b_t)_s^1), ((W_t)_s^2, (b_t)_s^2), ((\bar{W})_s^3, (\bar{b})_s^3), s \in [A]\}$ . By Lemma 9 and Lemma 32, we have

$$\begin{aligned}
\widehat{L}_R(f_{W_t}) - \widehat{L}_R(f_{\bar{W}_t}) &\leq [\nabla_{W^3} \widehat{L}_R(f_{W_t})]^T ((W_t)^3 - (\bar{W})^3) \\
&= \frac{1}{\eta} [\eta \nabla_{W^3} \widehat{L}_R(f_{W_t})]^T ((W_t)^3 - (\bar{W})^3) \\
&= \frac{1}{2\eta} (-\|(W_t)^3 - (\bar{W})^3 - \eta \nabla_{W^3} L_R(f_{W_t})\|^2 \\
&\quad + \|(W_t)^3 - (\bar{W})^3\|^2 + \|\eta \nabla_{W^3} \widehat{L}_R(f_{W_t})\|^2).
\end{aligned}$$

By Lemma 12,

$$\begin{aligned}
&\|(W_t)^3 - (\bar{W})^3 - \eta \nabla_{W^3} \widehat{L}_R(f_{W_t})\|^2 \\
&\geq \|\text{proj}_{\mathcal{C}}((W_t)^3 - \eta \nabla_{W^3} \widehat{L}_R(f_{W_t})) - \text{proj}_{\mathcal{C}}((\bar{W})^3)\|^2 = \|(W_{t+1})^3 - (\bar{W})^3\|^2.
\end{aligned}$$

Thus

$$\begin{aligned}
&\widehat{L}_R(f_{W_t}) - \widehat{L}_R(f_{\bar{W}_t}) \\
&\leq \frac{1}{2\eta} \left( -\|(W_{t+1})^3 - (\bar{W})^3\|^2 + \|(W_t)^3 - (\bar{W})^3\|^2 + \|\eta \nabla_{W^3} \widehat{L}_R(f_{W_t})\|^2 \right),
\end{aligned}$$

which implies

$$\begin{aligned}
&\frac{1}{T} \sum_{t=0}^{T-1} \widehat{L}_R(f_{W_t}) \\
&\leq \frac{1}{T} \sum_{t=0}^{T-1} \widehat{L}_R(f_{\bar{W}_t}) + \frac{1}{2\eta T} \|(W_0)^3 - (\bar{W})^3\|^2 + \frac{\eta}{2T} \sum_{t=0}^{T-1} \|\nabla_{W^3} \widehat{L}_R(f_{W_t})\|^2. \quad (21)
\end{aligned}$$

Next we prove that  $\hat{L}_R$  decreases during iteration. Let  $a \in [0, 1]$ . For  $l = 1, 2; s \in [A]$ , since  $W_{t+1}, W_t \in \mathcal{F}_{NN}(\{m_1, m_2, A\}, B_{inn}, B_{out})$ , there holds

$$\begin{aligned} \|a(W_{t+1})_s^l + (1-a)(W_t)_s^l\|_F &\leq a\|(W_{t+1})_s^l\|_F + (1-a)\|(W_t)_s^l\|_F \leq B_{inn}, \\ \|a(b_{t+1})_s^l + (1-a)(b_t)_s^l\|_2 &\leq a\|(b_{t+1})_s^l\|_F + (1-a)\|(b_t)_s^l\|_F \leq B_{inn}, \\ \|a(W_{t+1})^3 + (1-a)(W_t)^3\|_1 &\leq a\|(W_{t+1})^3\|_1 + (1-a)\|(W_t)^3\|_1 \leq B_{out}, \end{aligned}$$

which implies  $f_{aW_{t+1}+(1-a)W_t} \in \mathcal{F}_{NN}(\{m_1, m_2, A\}, B_{inn}, B_{out})$ . By Lemma 31,

$$\|\nabla_W \hat{L}_R(f_{aW_{t+1}+(1-a)W_t}) - \nabla_W \hat{L}_R(f_{W_t})\| \leq L\|W_{t+1} - W_t\|.$$

where  $L = C(d, \sigma, coe, \Omega) \max\{1/\beta, 1\} AB_{inn}^7 B_{out}^3$ . Now we can apply Lemma 13 and get

$$\begin{aligned} \hat{L}_R(f_{W_{t+1}}) &\leq \hat{L}_R(f_{W_t}) + [\nabla_W \hat{L}_R(f_{W_t})]^T (W_{t+1} - W_t) + \frac{L}{2} \|W_{t+1} - W_t\|^2 \\ &\leq \hat{L}_R(f_{W_t}) - \left(\frac{1}{\eta} - \frac{L}{2}\right) \|W_{t+1} - W_t\|^2, \end{aligned}$$

where we also make use of the relation

$$\begin{aligned} &[\nabla_W \hat{L}_R(f_{W_t})]^T (W_{t+1} - W_t) \\ &= -\frac{1}{\eta} [W_t - \eta \nabla_W \hat{L}_R(f_{W_t}) - W_t]^T [\text{proj}_{\mathcal{C}}(W_t - \eta \nabla_W \hat{L}_R(f_{W_t})) - \text{proj}_{\mathcal{C}}(W_t)] \\ &\leq -\frac{1}{\eta} \|\text{proj}_{\mathcal{C}}(W_t - \eta \nabla_W \hat{L}_R(f_{W_t})) - \text{proj}_{\mathcal{C}}(W_t)\|^2 = -\frac{1}{\eta} \|W_{t+1} - W_t\|^2. \end{aligned}$$

Here the second step is due to Lemma 12. Now we know that  $\hat{L}_R$  decreases during iteration provided  $\eta \leq \frac{2}{L}$ . So based on (21), we obtain

$$\begin{aligned} \hat{L}_R(f_{W_T}) - \inf_{f_W \in \mathcal{F}_{NN}} \hat{L}_R(f_W) &\leq \frac{1}{T} \sum_{t=0}^{T-1} \hat{L}_R(f_{W_t}) - \inf_{f_W \in \mathcal{F}_{NN}} \hat{L}_R(f_W) \\ &\leq \frac{1}{T} \sum_{t=0}^{T-1} \left[ \hat{L}_R(f_{\bar{W}_t}) - \inf_{f_W \in \mathcal{F}_{NN}} \hat{L}_R(f_W) \right] + \frac{1}{2\eta T} \|(W_0)^3 - (\bar{W})^3\|^2 \\ &\quad + \frac{\eta}{2T} \sum_{t=0}^{T-1} \|\nabla_{W^3} \hat{L}_R(f_{W_t})\|^2. \end{aligned}$$

We further divide the first term on the right-hand side into three terms:

$$\begin{aligned} &\hat{L}_R(f_{\bar{W}_t}) - \inf_{f_W \in \mathcal{F}_{NN}} \hat{L}_R(f_W) \\ &= [\hat{L}_R(f_{\bar{W}_t}) - L_R(f_{\bar{W}_t})] + [L_R(f_{\bar{W}_t}) - L_R(u_R)] + \left[ L_R(u_R) - \inf_{f_W \in \mathcal{F}_{NN}} \hat{L}_R(f_W) \right]. \end{aligned}$$

Since  $W_t \in \mathcal{C}$  and  $f_{\bar{W}} \in \mathcal{F}_{NN}$ , there holds  $f_{\bar{W}_t} \in \mathcal{F}_{NN}$  by our choice of  $B_{inn}$  and  $B_{out}$ . Thus first term can be controlled by  $\sup_{f_W \in \mathcal{F}_{NN}} [\hat{L}_R(f_W) - L_R(f_W)]$ . An

upper bound of the third term is provided by Lemma 20. So it suffice to show that with probability at least  $1 - A'e^{-C(d, \text{coe}, \Omega) \frac{A}{A'}} \left(\frac{1}{N}\right)^{206d^4(d+3)5^{d+2}}$ ,  $\bar{W}_t \in \mathcal{B}_{\bar{W}}$  and hence

$$L_R(f_{\bar{W}_t}) - L_R(u_R) \leq \sup_{W \in \mathcal{B}_{\bar{W}}} [L_R(f_W) - L_R(u_R)].$$

By Lemma 33, with that probability, there exists an injection  $\varphi : \{1, 2, \dots, A'\} \rightarrow \{1, 2, \dots, A\}$  such that for  $k \in [A']$ ,

$$\varphi(k) \in \{(k-1)A/A' + i' : i' \in [A/A']\}$$

and for  $l = 1, 2; s \in \{\varphi(1), \dots, \varphi(A')\}$ ,

$$\|(W_0)_s^l - (\bar{W})_s^l\|_F, \|(b_0)_s^l - (\bar{b})_s^l\|_2 \leq \frac{1}{2} \left(\frac{1}{N}\right)^{83d^3/2}.$$

Since  $W_t \in \mathcal{C}$ ,

$$\|(W_t)_s^l - (W_0)_s^l\|_F, \|(b_t)_s^l - (b_0)_s^l\|_2 \leq \frac{1}{2} \left(\frac{1}{N}\right)^{83d^3/2}, s \in [A], l = 1, 2.$$

From triangle inequality, we conclude that  $\bar{W}_t \in \mathcal{B}_{\bar{W}}$  with that probability.  $\square$

## 8 Convergence Rate of DRM

**Theorem 5.** *Let  $m = n$ . Let  $(W_0)_{s,i,j}^l, (b_0)_{s,i}^l (s \in [A]; l = 1, 2; i \in [m_l]; j \in [m_{l-1}])$  be uniformly distributed on*

$$\left[ -C(d, \text{coe}, \Omega) n^{\frac{10d^3}{144d^3+2}}, C(d, \text{coe}, \Omega) n^{\frac{10d^3}{144d^3+2}} \right].$$

Let  $B_{inn} = C(d, \text{coe}, \Omega) n^{\frac{10d^3}{144d^3+2}}, B_{out} = C(d, \text{coe}, \Omega) n^{\frac{11d^3}{144d^3+2}}$ . Let

$$A = n^{\frac{415d^4(d+3)5^{d+2}}{288d^3+4}}. \quad (22)$$

Let

$$\mathcal{C} = \left\{ W : \|W^3\|_1 \leq C(d, \text{coe}, \Omega) n^{\frac{11d^3}{144d^3+2}}; \right. \\ \left. \|W_s^l - (W_0)_s^l\|_F, \|b_s^l - (b_0)_s^l\|_2 \leq \frac{1}{2} n^{-\frac{83d^3}{288d^3+4}}, l = 1, 2, s \in [A] \right\}.$$

Let

$$\eta \leq C(d, \text{coe}, \Omega) n^{-\frac{103d^3}{144d^3+2}} \frac{1}{A} \quad (23)$$

and  $T = \frac{1}{\eta}$ .

(1) Consider Robin problem (1)(4). With probability at least  $1 - \frac{C(d, \text{coe}, \Omega)}{n}$ ,

$$\|f_{W_T} - u_R\|_{H^1(\Omega)} \leq C(d, \text{coe}, \Omega) \max\{1, 1/\beta\} n^{-\frac{1}{288d^3+4}}.$$

(2) Consider Neumann problem (1)(3). With probability at least  $1 - \frac{C(d, \text{coe}, \Omega)}{n}$ ,

$$\|f_{W_T} - u_N\|_{H^1(\Omega)} \leq C(d, \text{coe}, \Omega) n^{-\frac{1}{288d^3+4}}.$$

(3) Consider Dirichlet problem (1)(2). With probability at least  $1 - \frac{C(d, \text{coe}, \Omega)}{n}$ ,

$$\|f_{W_T} - u_D\|_{H^1(\Omega)} \leq C(d, \text{coe}, \Omega) n^{-\frac{1}{576d^3+8}}.$$

**Remark 10.** [44] provides a lower bound of  $\mathcal{O}(n^{-1/d})$  (when the solution of the equation is in  $H^2(\Omega)$ ) for DRM, and they found a neural network estimator with an upper error bound of  $\mathcal{O}(n^{-1/(d+2)} \log^{1/2} n)$ . The upper bound we obtained here is far from this lower bound, mainly due to the large parameter values used in our three-layer network estimator (Theorem 2, Proposition 1). However, from the perspective of constructing a three-layer network to approximate functions in Sobolev space, this is inevitable. One solution is to decrease the parameter values by increasing the depth of the neural network (refer to [29]). However, increasing the network depth also means increasing the number of parameters, leading to an increase in computational complexity during network training. Therefore, finding a neural network estimator that improves the error bound of this work while maintaining manageable computational complexity is one of the future research focuses.

*Proof.* We only provide a proof for the Robin problem. The result of the Neumann problem can be derived in the same way. The result of the Dirichlet problem can be obtained by combining the convergence rate of the Robin problem and Lemma 1.

Since  $W_t \in \mathcal{C}$  and  $\|(W_0)_s^l\|_F, \|(b_0)_s^l\|_2 \leq C(d, \text{coe}, \Omega) n^{\frac{10d^3}{144d^3+2}}$  for  $l = 1, 2; s \in [A]$ , we have  $f_{W_t} \in \mathcal{F}_{NN}$  by our choice of  $B_{inn}$  and  $B_{out}$ . Choosing  $\bar{W}$  in Theorem 4 to be  $W^*$ , the approximator in Theorem 2 when approximating  $u_R$ , and plugging it into Lemma 21, we have with probability at least  $1 - A'e^{-C(d, \text{coe}, \Omega) \frac{A}{A'} n^{-\frac{206d^4(d+3)5^{d+2}}{144d^3+2}}}$ ,

$$\begin{aligned} & L_R(f_{W_T}) - L_R(u_R) \\ & \leq 4 \max \left\{ \sup_{f_W \in \mathcal{F}_{NN}} [\widehat{L}_R(f_W) - L_R(f_W)], \sup_{f_W \in \mathcal{F}_{NN}} [L_R(f_W) - \widehat{L}_R(f_W)] \right\} \\ & \quad + 3 \sup_{W \in \mathcal{B}_{W^*}} [L_R(f_W) - L_R(u_R)] + \frac{1}{2\eta T} \|(W^*)^3\|^2 + \frac{\eta}{2T} \sum_{t=0}^{T-1} \|\nabla_{W^3} \widehat{L}_R(f_{W_t})\|^2. \end{aligned}$$

Here we set

$$N = C(d, \text{coe}, \Omega) n^{\frac{1}{144d^3+2}}$$

in Theorem 4. Our previous results provide estimates for each term on the right-hand side of the inequality. Choosing  $\tau = C(d, \Omega, coe) \max\{1, 1/\beta\} n^{-\frac{1}{144d^3+2}}$  in Theorem 3, we have with probability at least  $1 - 2e^{-C(d, coe, \Omega) n^{\frac{60d^3}{144d^3+2}}}$ ,

$$\begin{aligned} & \max \left\{ \sup_{f_W \in \mathcal{F}_{NN}} [\widehat{L}_R(f_W) - L_R(f_W)], \sup_{f_W \in \mathcal{F}_{NN}} [L_R(u) - \widehat{L}_R(f_W)] \right\} \\ & \leq C(d, \Omega, coe) \max\{1, 1/\beta\} n^{-\frac{1}{144d^3+2}}. \end{aligned}$$

Choosing

$$N = C(d, coe, \Omega) n^{\frac{1}{144d^3+2}}$$

in Theorem 2, there holds

$$\|(W^*)^3\| \leq C(d, coe, \Omega) \frac{1}{A'} N^{23d^3/2} = C(d, coe, \Omega) \frac{1}{A'} n^{\frac{23d^3}{288d^3+4}}.$$

By Lemma 30,

$$\|\nabla_{W^3} \widehat{L}_R(f_W)\| \leq C(d, \sigma, \Omega) \max\{1, 1/\beta\} \sqrt{A} n^{\frac{51d^3}{144d^3+2}}.$$

According to Lemma 8, Lemma 19 and Theorem 2,

$$\begin{aligned} \sup_{W \in \mathcal{B}_{W^*}} [L_R(f_W) - L_R(u_R)] & \leq C(\Omega, coe) \max\{1, 1/\beta\} \sup_{W \in \mathcal{B}_{W^*}} \|f_W - u_R\|_{H^1(\Omega)}^2 \\ & \leq C(d, coe, \Omega) \max\{1, 1/\beta\} n^{-\frac{1}{144d^3+2}}. \end{aligned}$$

Combining all the above upper bounds, we derive that with probability at least  $1 - A' e^{-C(d, coe, \Omega) \frac{A}{A'} n^{-\frac{206d^4(d+3)5^{d+2}}{144d^3+2}}} - 2e^{-C(d, coe, \Omega) n^{\frac{60d^3}{144d^3+2}}}$ ,

$$\begin{aligned} & L_R(f_{W_T}) - L_R(u_R) \\ & \leq C(d, coe, \Omega) \max\{1, 1/\beta^2\} \left( n^{-\frac{1}{144d^3+2}} + \frac{1}{A'^2} n^{\frac{23d^3}{144d^3+2}} + \eta A n^{\frac{51d^3}{72d^3+1}} \right). \end{aligned}$$

Let  $A$  satisfy (22) and  $\eta$  satisfy (23), and further let

$$A' = C(d, coe, \Omega) n^{\frac{23d^3+1}{288d^3+4}}.$$

After organizing and simplifying, we obtain that with probability at least  $1 - \frac{C(d, coe, \Omega)}{n}$ ,

$$L_R(f_{W_T}) - L_R(u_R) \leq C(d, coe, \Omega) \max\{1, 1/\beta^2\} n^{-\frac{1}{144d^3+2}}.$$

The convergence rate of  $\|f_W - u_R\|_{H^1(\Omega)}$  is then implied by Lemma 19.  $\square$

## 9 Conclusion

In this work, we focus on employing a three-layer tanh neural network within the framework of the deep Ritz method(DRM) to solve second-order elliptic equations with three different types of boundary conditions. To the best of our knowledge, we are the first to provide a comprehensive error analysis of using overparameterized networks to solve PDE problems. We present error bound in terms of the sample size  $n$  and our work provides guidance on how to set the network depth, width, step size, and number of iterations for the projected gradient descent algorithm. Importantly, our assumptions in this work are classical and we do not require any additional assumptions on the solution of the equation. This ensures the broad applicability and generality of our results.

As an initial study, this work focuses on a three-layer neural network. However, we believe that the framework presented in this work can also be extended to analyze deep neural networks. We speculate that deep neural network estimators may have better upper bounds, and this will be one of the topics for our future research. Additionally, investigating other solution formats for PDEs, such as PINNs and WAN, as well as considering other types of PDEs, are also promising directions for future research. On the training front, this work investigates the projected gradient descent algorithm. Analyzing the convergence properties of gradient descent and stochastic gradient descent algorithms when applied to PDE problems presents technical challenges. In the future, we aim to address these technical difficulties and explore the convergence analysis of these optimization algorithms in the context of PDE problem-solving.

## A Proof of Lemma 8

We first give an auxiliary lemma.

**Lemma 34.** *Let assumption (5) holds. Let  $u$  be the weak solution of the following Robin problem*

$$\begin{cases} -\Delta u + wu = 0 & \text{in } \Omega \\ u + \beta \frac{\partial u}{\partial n} = g & \text{on } \partial\Omega. \end{cases} \quad (24)$$

Then  $u_R \in H^2(\Omega)$  and

$$\|u\|_{H^2(\Omega)} \leq C(\Omega, w) \frac{1}{\beta} \|g\|_{H^{1/2}(\partial\Omega)}.$$

*Proof.* We follow the idea proposed in [46] in a slightly different context. We first estimate the trace  $T_0 u = u|_{\partial\Omega}$ . We define the Dirichlet-to-Neumann map

$$\tilde{T} : u|_{\partial\Omega} \mapsto \frac{\partial u}{\partial n} \Big|_{\partial\Omega},$$

where  $u$  satisfies  $-\Delta u + wu = 0$  in  $\Omega$ , then

$$T_0 u = \left( \beta \tilde{T} + I \right)^{-1} g.$$

Now we are going to show that  $\beta \tilde{T} + I$  is a positive definite operator in  $L^2(\partial\Omega)$ . We notice that the variational formulation of (24) can be read as follow:

$$\int_{\Omega} \nabla u \cdot \nabla v dx + \int_{\Omega} w u v dx + \frac{1}{\beta} \int_{\partial\Omega} u v ds = \frac{1}{\beta} \int_{\partial\Omega} g v ds, \quad \forall v \in H^1(\Omega).$$

Taking  $v = u$ , then we have

$$\|T_0 u\|_{L^2(\partial\Omega)}^2 \leq \left\langle \left( \beta \tilde{T} + I \right) T_0 u, T_0 u \right\rangle.$$

This means that  $\beta \tilde{T} + I$  is a positive definite operator in  $L^2(\partial\Omega)$ , and further,  $(\beta \tilde{T} + I)^{-1}$  is bounded. We have the estimate

$$\|T_0 u\|_{H^{1/2}(\partial\Omega)} \leq C(\Omega) \|g\|_{H^{1/2}(\partial\Omega)}. \quad (25)$$

We rewrite the Robin problem (24) as follows

$$\begin{cases} -\Delta u + wu = 0 & \text{in } \Omega \\ u + \frac{\partial u}{\partial n} = \frac{1}{\beta} (g - (1 - \beta) u) & \text{on } \partial\Omega. \end{cases}$$

By Lemma 7 we have

$$\begin{aligned} \|u\|_{H^2(\Omega)} &\leq C(\Omega, w) \frac{1}{\beta} \|g - (1 - \beta) T_0 u\|_{H^{1/2}(\partial\Omega)} \\ &\leq C(\Omega, w) \frac{1}{\beta} \left( \|g\|_{H^{1/2}(\partial\Omega)} + \|T_0 u\|_{H^{1/2}(\partial\Omega)} \right). \end{aligned} \quad (26)$$

Combining (25) and (26), we obtain the desired estimation.  $\square$

With the help of the above lemma, we now turn to prove Lemma 8.

*Proof of Lemma 8.* We decompose the Robin problem (1)(4) into two equations

$$\begin{cases} -\Delta u_0 + wu_0 = f & \text{in } \Omega \\ u_0 = g & \text{on } \partial\Omega, \end{cases}$$

$$\begin{cases} -\Delta u_1 + wu_1 = 0 & \text{in } \Omega \\ u_1 + \beta \frac{\partial u_1}{\partial n} = -\frac{\partial u_0}{\partial n} & \text{on } \partial\Omega. \end{cases}$$

and obtain the solution of (1)(4):

$$u = u_0 + \beta u_1.$$

According to Lemma 6, we have

$$\|u_0\|_{H^2(\Omega)} \leq C(\Omega, w)(\|f\|_{L^2(\Omega)} + \|g\|_{H^{3/2}(\partial\Omega)}). \quad (27)$$

Using Lemma 34, it is easy to obtain

$$\|u_1\|_{H^2(\Omega)} \leq C(\Omega, w) \frac{1}{\beta} \left\| \frac{\partial u_0}{\partial n} \right\|_{H^{1/2}(\partial\Omega)} \leq C(\Omega, w) \frac{1}{\beta} \|u_0\|_{H^2(\Omega)}, \quad (28)$$

where the last inequality follows from Lemma 5. Combining (27) and (28), the desired estimation can be derived by triangle inequality.  $\square$

## B Proof of Lemma 31

*Proof.* We only present a proof for  $\widehat{L}_R(f_W)$ . The inequality of  $\widehat{L}_N(f_W)$  can be proved in almost the same way. The whole proof is divided into five steps. In the first step, we provide upper bounds for  $|f_W|$  and  $|f_W - f_{\widetilde{W}}|$ . The former can be derived easily:

$$|f_W| \leq \sum_{s=1}^A [\|W_s^3\|_F \|f_s^2\|_2 + |b_s^3|] \leq \sum_{s=1}^A (\sqrt{m_2} B_\sigma \|W_s^3\|_F + |b_s^3|).$$

In order to estimate  $|f_W - f_{\widetilde{W}}|$ , we need the following two inequalities:

$$\begin{aligned} \|f_{s,1}^{org}(W) - f_{s,1}^{org}(\widetilde{W})\|_2 &\leq \|W_s^1 - (\widetilde{W})_s^1\|_2 \|x\|_2 + \|b_s^1 - (\widetilde{b})_s^1\|_2, \\ \|f_{s,2}^{org}(W) - f_{s,2}^{org}(\widetilde{W})\|_2 &\leq \|W_s^2 - (\widetilde{W})_s^2\|_2 \|f_s^1(W)\|_2 + \|(\widetilde{W})_s^2\|_2 \|f_s^1(W) - f_s^1(\widetilde{W})\|_2 + \|b_s^2 - (\widetilde{b})_s^2\|_2 \\ &\leq \sqrt{m_1} B_\sigma \|W_s^2 - (\widetilde{W})_s^2\|_2 + \|b_s^2 - (\widetilde{b})_s^2\|_2 + L_\sigma \|(\widetilde{W})_s^2\|_2 \|x\|_2 \|W_s^1 - (\widetilde{W})_s^1\|_2 \\ &\quad + L_\sigma \|(\widetilde{W})_s^2\|_2 \|b_s^1 - (\widetilde{b})_s^1\|_2. \end{aligned}$$

Thus

$$\begin{aligned} |f_W - f_{\widetilde{W}}| &\leq \sum_{s=1}^A |f_s^3(W) - f_s^3(\widetilde{W})| \\ &\leq \sum_{s=1}^A (\|W_s^3 - (\widetilde{W})_s^3\|_2 \|f^2(W)\|_2 + \|(\widetilde{W})_s^3\|_2 \|f_s^2(W) - f_s^2(\widetilde{W})\|_2 + |b_s^3 - (\widetilde{b})_s^3|) \\ &\leq \sum_{s=1}^A (\sqrt{m_2} B_\sigma \|W_s^3 - (\widetilde{W})_s^3\|_2 + |b_s^3 - (\widetilde{b})_s^3| + \sqrt{m_1} L_\sigma B_\sigma \|(\widetilde{W})_s^3\|_2 \|W_s^2 - (\widetilde{W})_s^2\|_2 \\ &\quad + L_\sigma \|(\widetilde{W})_s^3\|_2 \|b_s^2 - (\widetilde{b})_s^2\|_2 + L_\sigma^2 B_x \|(\widetilde{W})_s^3\|_2 \|(\widetilde{W})_s^2\|_2 \|W_s^1 - (\widetilde{W})_s^1\|_2 \\ &\quad + L_\sigma^2 \|(\widetilde{W})_s^3\|_2 \|(\widetilde{W})_s^2\|_2 \|b_s^1 - (\widetilde{b})_s^1\|_2). \end{aligned}$$

The second step is to give upper bounds of  $\|\nabla_x f_W\|_2$  and  $\|\nabla_x f_W - \nabla_x f_{\widetilde{W}}\|_2$ . The gradient of  $f_W$  with respect to the spatial variable  $x$  can be calculated by

$$\nabla_x f_W = \sum_{s=1}^A (W_s^1)^T \text{diag}[\sigma'(f_{s,1}^{org})] (W_s^2)^T \text{diag}[\sigma'(f_{s,2}^{org})] (W_s^3)^T,$$

therefore

$$\begin{aligned} \|\nabla_x f_W\|_2 &\leq \sum_{s=1}^A \|W_s^1\|_2 \|\text{diag}[\sigma'(f_{s,1}^{org})]\|_2 \|W_s^2\|_2 \|\text{diag}[\sigma'(f_{s,2}^{org})]\|_2 \|W_s^3\|_2 \\ &\leq \sum_{s=1}^A \sqrt{m_2 m_1} B_{\sigma'} \|W_s^1\|_2 \|W_s^2\|_2 \|W_s^3\|_2. \end{aligned}$$

and

$$\begin{aligned} &\|\nabla_x f_W - \nabla_x f_{\widetilde{W}}\|_2 \\ &\leq \sum_{s=1}^A [\|W_s^1 - (\widetilde{W})_s^1\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(W))]\|_F \|W_s^2\|_F \|\text{diag}[\sigma'(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \\ &\quad + \|(\widetilde{W})_s^1\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(W))] - \text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \\ &\quad \|W_s^2\|_F \|\text{diag}[\sigma'(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \\ &\quad + \|(\widetilde{W})_s^1\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|W_s^2 - (\widetilde{W})_s^2\|_F \\ &\quad \|\text{diag}[\sigma'(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \\ &\quad + \|(\widetilde{W})_s^1\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^2\|_F \\ &\quad \|\text{diag}[\sigma'(f_{s,2}^{org}(W))] - \text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F \|(W_s^3)^T\|_2 \\ &\quad + \|(\widetilde{W})_s^1\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^2\|_F \\ &\quad \|\text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F \|W_s^3 - (\widetilde{W})_s^3\|_F] \\ &\leq C(\sigma) \sum_{s=1}^A [\sqrt{m_2 m_1} \|(\widetilde{W})_s^1\|_F \|(\widetilde{W})_s^2\|_F \|W_s^3 - (\widetilde{W})_s^3\|_F \\ &\quad + \max\{\sqrt{m_2}, \sqrt{m_1}\} \sqrt{m_1} \|(\widetilde{W})_s^1\|_F \|(\widetilde{W})_s^2\|_F \|W_s^3\|_F \|W_s^2 - (\widetilde{W})_s^2\|_F \\ &\quad + \sqrt{m_1} \|(\widetilde{W})_s^1\|_F \|(\widetilde{W})_s^2\|_F \|W_s^3\|_F \|b_s^2 - (\widetilde{b})_s^2\|_2 \\ &\quad + \sqrt{m_2 m_1} \|W_s^3\|_F \|(\widetilde{W})_s^1\|_F \max\{\|W_s^2\|_F^2, \|(\widetilde{W})_s^2\|_F^2\} \|W_s^1 - (\widetilde{W})_s^1\|_F \\ &\quad + \max\{\sqrt{m_2}, \sqrt{m_1}\} \|(\widetilde{W})_s^1\|_F \|W_s^3\|_F \max\{\|W_s^2\|_F^2, \|(\widetilde{W})_s^2\|_F^2\} \|b_s^1 - (\widetilde{b})_s^1\|_2]. \end{aligned}$$

The third step is to derive the estimates of  $\|\nabla_{W_s^l} f_W\|_F$ ,  $\|\nabla_{b_s^l} f_W\|_F$  and  $\|\nabla_{W_s^l} f_W - \nabla_{W_s^l} f_{\widetilde{W}}\|_F$ ,  $\|\nabla_{b_s^l} f_W - \nabla_{b_s^l} f_{\widetilde{W}}\|_F$  ( $l = 1, 2, 3$ ). The gradient of  $f_W$  with respect to weights and biases in each layer can be calculated by

$$\nabla_{W_s^3} f_W = (f_s^2)^T,$$

$$\begin{aligned}
\nabla_{b_s^3} f_W &= 1, \\
\nabla_{W_s^2} f_W &= \text{diag}[\sigma'(f_{s,2}^{org})](W_s^3)^T (f_s^1)^T, \\
\nabla_{b_s^2} f_W &= \text{diag}[\sigma'(f_{s,2}^{org})](W_s^3)^T, \\
\nabla_{W_s^1} f_W &= \text{diag}[\sigma'(f_{s,1}^{org})](W_s^2)^T \text{diag}[\sigma'(f_{s,2}^{org})](W_s^3)^T x^T, \\
\nabla_{b_s^1} f_W &= \text{diag}[\sigma'(f_{s,1}^{org})](W_s^2)^T \text{diag}[\sigma'(f_{s,2}^{org})](W_s^3)^T,
\end{aligned}$$

hence we have

$$\begin{aligned}
\|\nabla_{W_s^3} f_W\|_F &= \|f_s^2\|_F \leq \sqrt{m_2} B_\sigma, \quad |\nabla_{b_s^3} f_W| = 1, \\
\|\nabla_{W_s^2} f_W\|_F &\leq \|\text{diag}[\sigma'(f_{s,2}^{org})]\|_F \|W_s^3\|_F \|f_s^1\|_F \leq \sqrt{m_2 m_1} B_\sigma B_{\sigma'} \|W_s^3\|_F, \\
\|\nabla_{b_s^2} f_W\|_F &\leq \|\text{diag}[\sigma'(f_{s,2}^{org})]\|_F \|W_s^3\|_F \leq \sqrt{m_2} B_{\sigma'} \|W_s^3\|_F, \\
\|\nabla_{W_s^1} f_W\|_F &= \|\text{diag}[\sigma'(f_{s,1}^{org})]\|_F \|W_s^2\|_F \|\text{diag}[\sigma'(f_{s,2}^{org})]\|_F \|W_s^3\|_F \|x\|_F \\
&\leq \sqrt{m_1 m_2} B_{\sigma'}^2 B_x \|W_s^2\|_F \|W_s^3\|_F, \\
\|\nabla_{b_s^1} f_W\|_F &= \|\text{diag}[\sigma'(f_{s,1}^{org})]\|_F \|W_s^2\|_F \|\text{diag}[\sigma'(f_{s,2}^{org})]\|_F \|W_s^3\|_F \\
&\leq \sqrt{m_1 m_2} B_{\sigma'}^2 \|W_s^2\|_F \|W_s^3\|_F.
\end{aligned}$$

and

$$\begin{aligned}
\|\nabla_{W_s^3} f_W - \nabla_{W_s^3} f_{\widetilde{W}}\|_F &= \|f_s^2(W) - f_s^2(\widetilde{W})\|_2 \\
&\leq \sqrt{m_1} L_\sigma B_\sigma \|W_s^2 - (\widetilde{W})_s^2\|_2 + L_\sigma \|b_s^2 - (\widetilde{b})_s^2\|_2 + L_\sigma^2 B_x \|(\widetilde{W})_s^2\|_2 \|W_s^1 - (\widetilde{W})_s^1\|_2 \\
&\quad + L_\sigma^2 \|(\widetilde{W})_s^2\|_2 \|b_s^1 - (\widetilde{b})_s^1\|_2, \\
\|\nabla_{b_s^3} f_W - \nabla_{b_s^3} f_{\widetilde{W}}\|_F &= 0, \\
\|\nabla_{W_s^2} f_W - \nabla_{W_s^2} f_{\widetilde{W}}\|_F &\leq \|\text{diag}[\sigma'(f_{s,2}^{org}(W))] - \text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F \|W_s^3\|_F \|f_s^1(W)\|_2 \\
&\quad + \|\text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F \|W_s^3 - (\widetilde{W})_s^3\|_F \|f_s^1(W)\|_2 \\
&\quad + \|\text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^3\|_F \|f_s^1(W) - f_s^1(\widetilde{W})\|_2 \\
&\leq \sqrt{m_2 m_1} B_\sigma B_{\sigma'} \|W_s^3 - (\widetilde{W})_s^3\|_F + \sqrt{m_1} \sqrt{m_1} L_\sigma B_\sigma^2 \|W_s^3\|_F \|W_s^2 - (\widetilde{W})_s^2\|_2 \\
&\quad + \sqrt{A m_1} L_\sigma B_\sigma \|W_s^3\|_F \|b_s^2 - (\widetilde{b})_s^2\|_2 \\
&\quad + L_\sigma B_x (\sqrt{m_1} L_\sigma B_\sigma \|W_s^3\|_F \|(\widetilde{W})_s^2\|_2 + \sqrt{m_2} B_{\sigma'} \|(\widetilde{W})_s^3\|_F) \|W_s^1 - (\widetilde{W})_s^1\|_2 \\
&\quad + L_\sigma (\sqrt{m_1} L_\sigma B_\sigma \|W_s^3\|_F \|(\widetilde{W})_s^2\|_2 + \sqrt{m_2} B_{\sigma'} \|(\widetilde{W})_s^3\|_F) \|b_s^1 - (\widetilde{b})_s^1\|_2, \\
\|\nabla_{b_s^2} f_W - \nabla_{b_s^2} f_{\widetilde{W}}\|_F &\leq \|\text{diag}[\sigma'(f_{s,2}^{org}(W))] - \text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F \|W_s^3\|_F \\
&\quad + \|\text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F \|W_s^3 - (\widetilde{W})_s^3\|_F \\
&\leq \sqrt{m_2} B_{\sigma'} \|W_s^3 - (\widetilde{W})_s^3\|_F + \sqrt{m_1} L_\sigma B_\sigma \|W_s^3\|_F \|W_s^2 - (\widetilde{W})_s^2\|_2 \\
&\quad + L_\sigma \|W_s^3\|_F \|b_s^2 - (\widetilde{b})_s^2\|_2 + L_\sigma L_\sigma B_x \|W_s^3\|_F \|(\widetilde{W})_s^2\|_2 \|W_s^1 - (\widetilde{W})_s^1\|_2 \\
&\quad + L_\sigma L_\sigma \|W_s^3\|_F \|(\widetilde{W})_s^2\|_2 \|b_s^1 - (\widetilde{b})_s^1\|_2,
\end{aligned}$$

$$\begin{aligned}
& \|\nabla_{W_s^1} f_W - \nabla_{W_s^1} f_{\widetilde{W}}\|_F \\
& \leq \|\text{diag}[\sigma'(f_{s,1}^{org}(W))] - \text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|W_s^2\|_F \\
& \quad \|\text{diag}[\sigma'(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \|x\|_2 \\
& \quad + \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|W_s^2 - (\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \|x\|_2 \\
& \quad + \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^2\|_F \\
& \quad \|\text{diag}[\sigma'(f_{s,2}^{org}(W))] - \text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F \|W_s^3\|_F \|x\|_2 \\
& \quad + \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F \|W_s^3 - (\widetilde{W})_s^3\|_F \|x\|_2 \\
& \leq \sqrt{m_2 m_1} B_{\sigma'}^2 \|(\widetilde{W})_s^2\|_F \|W_s^3 - (\widetilde{W})_s^3\|_F \\
& \quad + \sqrt{m_1} B_{\sigma'} \|W_s^3\|_F (\sqrt{m_2} B_{\sigma'} + \sqrt{m_1} L_{\sigma'} B_{\sigma'} \|(\widetilde{W})_s^2\|_F) \|W_s^2 - (\widetilde{W})_s^2\|_F \\
& \quad + \sqrt{m_1} L_{\sigma'} B_{\sigma'} \|(\widetilde{W})_s^2\|_F \|W_s^3\|_F \|b_s^2 - (\widetilde{b})_s^2\|_2 \\
& \quad + L_{\sigma'} B_{\sigma'} \|W_s^3\|_F (\sqrt{m_2} \|W_s^2\|_F + \sqrt{m_1} L_{\sigma'} \|(\widetilde{W})_s^2\|_F) \|W_s^1 - (\widetilde{W})_s^1\|_2 \\
& \quad + L_{\sigma'} B_{\sigma'} \|W_s^3\|_F (\sqrt{m_2} \|W_s^2\|_F + \sqrt{m_1} L_{\sigma'} \|(\widetilde{W})_s^2\|_F) \|b_s^1 - (\widetilde{b})_s^1\|_2, \\
& \|\nabla_{b_s^1} f_W - \nabla_{b_s^1} f_{\widetilde{W}}\|_F \\
& \leq \|\text{diag}[\sigma'(f_{s,1}^{org}(W))] - \text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|W_s^2\|_F \\
& \quad \|\text{diag}[\sigma'(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \\
& \quad + \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|W_s^2 - (\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \\
& \quad + \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^2\|_F \\
& \quad \|\text{diag}[\sigma'(f_{s,2}^{org}(W))] - \text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F \|W_s^3\|_F \\
& \quad + \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F \|W_s^3 - (\widetilde{W})_s^3\|_F \\
& \leq \sqrt{m_2 m_1} B_{\sigma'}^2 \|(\widetilde{W})_s^2\|_F \|W_s^3 - (\widetilde{W})_s^3\|_F \\
& \quad + \sqrt{m_1} B_{\sigma'} \|W_s^3\|_F (\sqrt{m_2} B_{\sigma'} + \sqrt{m_1} L_{\sigma'} B_{\sigma'} \|(\widetilde{W})_s^2\|_F) \|W_s^2 - (\widetilde{W})_s^2\|_F \\
& \quad + \sqrt{m_1} L_{\sigma'} B_{\sigma'} \|(\widetilde{W})_s^2\|_F \|W_s^3\|_F \|b_s^2 - (\widetilde{b})_s^2\|_2 \\
& \quad + L_{\sigma'} B_{\sigma'} \|W_s^3\|_F (\sqrt{m_2} \|W_s^2\|_F + \sqrt{m_1} L_{\sigma'} \|(\widetilde{W})_s^2\|_F) \|W_s^1 - (\widetilde{W})_s^1\|_2 \\
& \quad + L_{\sigma'} B_{\sigma'} \|W_s^3\|_F (\sqrt{m_2} \|W_s^2\|_F + \sqrt{m_1} L_{\sigma'} \|(\widetilde{W})_s^2\|_F) \|b_s^1 - (\widetilde{b})_s^1\|_2.
\end{aligned}$$

The fourth step is to present upper bounds of  $\left\| \nabla_{W_s^l} \left( \frac{\partial f_W}{\partial x_j} \right) \right\|_F$ ,  $\left\| \nabla_{b_s^l} \left( \frac{\partial f_W}{\partial x_j} \right) \right\|_F$  and  $\left\| \nabla_{W_s^l} \left( \frac{\partial f_W}{\partial x_j} \right) - \nabla_{W_s^l} \left( \frac{\partial f_{\widetilde{W}}}{\partial x_j} \right) \right\|_F$ ,  $\left\| \nabla_{b_s^l} \left( \frac{\partial f_W}{\partial x_j} \right) - \nabla_{b_s^l} \left( \frac{\partial f_{\widetilde{W}}}{\partial x_j} \right) \right\|_F$  ( $l = 1, 2, 3$ ). Let  $W_s^1 = (W_s^{1,1}, \dots, W_s^{1,d})$  with  $W_s^{1,j} \in \mathbb{R}^{m_1}$ ,  $j = 1, \dots, d$ . The partial derivative of  $f_W$  with respect to the spatial variables  $x_j$  ( $j = 1, \dots, d$ ) is

$$\frac{\partial f_W}{\partial x_j} = \sum_{s=1}^A (W_s^{1,j})^T \text{diag}[\sigma'(f_{s,1}^{org})] (W_s^2)^T \text{diag}[\sigma'(f_{s,2}^{org})] (W_s^3)^T.$$

and its gradient with respect to weights and biases in each layer can be calculated

by

$$\begin{aligned}
\nabla_{W_s^3} \left( \frac{\partial f_W}{\partial x_j} \right) &= (W_s^{1,j})^T \text{diag}[\sigma'(f_{s,1}^{org})] (W_s^2)^T \text{diag}[\sigma'(f_{s,2}^{org})], \\
\nabla_{W_s^2} \left( \frac{\partial f_W}{\partial x_j} \right) &= \text{diag}[\sigma'(f_{s,2}^{org})] (W_s^3)^T (W_s^{1,j})^T \text{diag}[\sigma'(f_{s,1}^{org})] \\
&\quad + (W_s^2 \text{diag}[\sigma'(f_{s,1}^{org})] W_s^{1,j}) \odot (\text{diag}[\sigma''(f_{s,2}^{org})] (W_s^3)^T) (f_s^1)^T, \\
\nabla_{b_s^2} \left( \frac{\partial f_W}{\partial x_j} \right) &= (W_s^2 \text{diag}[\sigma'(f_{s,1}^{org})] W_s^{1,j}) \odot (\text{diag}[\sigma''(f_{s,2}^{org})] (W_s^3)^T), \\
\nabla_{W_s^1} \left( \frac{\partial f_W}{\partial x_j} \right) &= (0, \dots, 0, \underbrace{\text{diag}[\sigma'(f_{s,1}^{org})] (W_s^2)^T \text{diag}[\sigma'(f_{s,2}^{org})] (W_s^3)^T}_{j\text{th}}, 0, \dots, 0) \\
&\quad + (\text{diag}[\sigma''(f_{s,1}^{org})] W_s^{1,j}) \odot ((W_s^2)^T \text{diag}[\sigma'(f_{s,2}^{org})] (W_s^3)^T) x^T \\
&\quad + \text{diag}[\sigma'(f_{s,1}^{org})] (W_s^2)^T \\
&\quad \quad ((W_s^2 \text{diag}[\sigma'(f_{s,1}^{org})] W_s^{1,j}) \odot (\text{diag}[\sigma''(f_{s,2}^{org})] (W_s^3)^T)) x^T, \\
\nabla_{b_s^1} \left( \frac{\partial f_W}{\partial x_j} \right) &= (\text{diag}[\sigma''(f_{s,1}^{org})] W_s^{1,j}) \odot ((W_s^2)^T \text{diag}[\sigma'(f_{s,2}^{org})] (W_s^3)^T) \\
&\quad + \text{diag}[\sigma'(f_{s,1}^{org})] (W_s^2)^T \\
&\quad \quad ((W_s^2 \text{diag}[\sigma'(f_{s,1}^{org})] W_s^{1,j}) \odot (\text{diag}[\sigma''(f_{s,2}^{org})] (W_s^3)^T)).
\end{aligned}$$

Employing the inequality  $\|a \odot b\|_2 \leq \|a\|_2 \|b\|_2$  for any vectors  $a$  and  $b$ , we have

$$\begin{aligned}
\left\| \nabla_{W_s^3} \left( \frac{\partial f_W}{\partial x_j} \right) \right\|_F &\leq \|W_s^{1,j}\|_F \|\text{diag}[\sigma'(f_{s,1}^{org})]\|_F \|W_s^2\|_F \|\text{diag}[\sigma'(f_{s,2}^{org})]\|_F \\
&\leq \sqrt{m_2 m_1} \|W_s^{1,j}\|_F \|W_s^2\|_F, \\
\left\| \nabla_{W_s^2} \left( \frac{\partial f_W}{\partial x_j} \right) \right\|_F &\leq \|\text{diag}[\sigma'(f_{s,2}^{org})]\|_F \|W_s^3\|_F \|W_s^{1,j}\|_F \|\text{diag}[\sigma'(f_{s,1}^{org})]\|_F \\
&\quad + \|W_s^2\|_F \|\text{diag}[\sigma'(f_{s,1}^{org})]\|_F \|W_s^{1,j}\|_F \\
&\quad \quad \|\text{diag}[\sigma''(f_{s,2}^{org})]\|_F \|W_s^3\|_F \|f_s^1\|_F \\
&\leq 2\sqrt{m_2 m_1} B_\sigma B_{\sigma'}^2 B_{\sigma''} \|W_s^{1,j}\|_F \|W_s^2\|_F \|W_s^3\|_F, \\
\left\| \nabla_{b_s^2} \left( \frac{\partial f_W}{\partial x_j} \right) \right\|_F &\leq \|W_s^2\|_F \|\text{diag}[\sigma'(f_{s,1}^{org})]\|_F \|W_s^{1,j}\|_F \|\text{diag}[\sigma''(f_{s,2}^{org})]\|_F \|W_s^3\|_F \\
&\leq \sqrt{m_2 m_1} B_{\sigma'} B_{\sigma''} \|W_s^{1,j}\|_F \|W_s^2\|_F \|W_s^3\|_F, \\
\left\| \nabla_{W_s^1} \left( \frac{\partial f_W}{\partial x_j} \right) \right\|_F &\leq \|\text{diag}[\sigma'(f_{s,1}^{org})]\|_F \|W_s^2\|_F \|\text{diag}[\sigma'(f_{s,2}^{org})]\|_F \|W_s^3\|_F \\
&\quad + \|\text{diag}[\sigma''(f_{s,1}^{org})]\|_F \|W_s^{1,j}\|_F \|W_s^2\|_F \\
&\quad \quad \|\text{diag}[\sigma'(f_{s,2}^{org})]\|_F \|W_s^3\|_F \|x\|_2 \\
&\quad + \|\text{diag}[\sigma'(f_{s,1}^{org})]\|_F \|W_s^2\|_F \|W_s^2\|_F \\
&\quad \quad \|\text{diag}[\sigma'(f_{s,1}^{org})]\|_F \|W_s^{1,j}\|_F \|\text{diag}[\sigma''(f_{s,2}^{org})]\|_F \|W_s^3\|_F \|x\|_2
\end{aligned}$$

$$\begin{aligned}
&\leq 3\sqrt{m_2}m_1B_{\sigma'}^2B_{\sigma''}B_x\|W_s^{1,j}\|_F\|W_s^2\|_F^2\|W_s^3\|_F, \\
\left\|\nabla_{b_s^1}\left(\frac{\partial f_W}{\partial x_j}\right)\right\|_F &\leq \|\text{diag}[\sigma''(f_{s,1}^{org})]\|_F\|W_s^{1,j}\|_F\|W_s^2\|_F\|\text{diag}[\sigma'(f_{s,2}^{org})]\|_F\|W_s^3\|_F \\
&\quad + \|\text{diag}[\sigma'(f_{s,1}^{org})]\|_F\|W_s^2\|_F\|W_s^2\|_F \\
&\quad \|\text{diag}[\sigma'(f_{s,1}^{org})]\|_F\|W_s^{1,j}\|_F\|\text{diag}[\sigma''(f_{s,2}^{org})]\|_F\|W_s^3\|_F \\
&\leq 2\sqrt{m_2}m_1B_{\sigma'}^2B_{\sigma''}\|W_s^{1,j}\|_F\|W_s^2\|_F^2\|W_s^3\|_F.
\end{aligned}$$

and

$$\begin{aligned}
&\left\|\nabla_{W_s^3}\left(\frac{\partial f_W}{\partial x_j}\right) - \nabla_{W_s^3}\left(\frac{\partial f_{\widetilde{W}}}{\partial x_j}\right)\right\|_F \\
&\leq \|W_s^{1,j} - (\widetilde{W})_s^{1,j}\|_F\|\text{diag}[\sigma'(f_{s,1}^{org}(W))]\|_F\|W_s^2\|_F\|\text{diag}[\sigma'(f_{s,2}^{org}(W))]\|_F \\
&\quad + \|(\widetilde{W})_s^{1,j}\|_F\|\text{diag}[\sigma'(f_{s,1}^{org}(W))]\|_F - \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \\
&\quad \|W_s^2\|_F\|\text{diag}[\sigma'(f_{s,2}^{org}(W))]\|_F \\
&\quad + \|(\widetilde{W})_s^{1,j}\|_F\|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F\|W_s^2 - (\widetilde{W})_s^2\|_F\|\text{diag}[\sigma'(f_{s,2}^{org}(W))]\|_F \\
&\quad + \|(\widetilde{W})_s^{1,j}\|_F\|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \\
&\quad \|(\widetilde{W})_s^2\|_F\|\text{diag}[\sigma'(f_{s,2}^{org}(W))]\|_F - \|\text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F \\
&\leq C(\sigma)(\max\{m_2, m_1\})\|(\widetilde{W})_s^{1,j}\|_F\|(\widetilde{W})_s^2\|_F\|W_s^2 - (\widetilde{W})_s^2\|_F \\
&\quad + \sqrt{m_1}\|(\widetilde{W})_s^{1,j}\|_F\|(\widetilde{W})_s^2\|_F\|b_s^2 - (\widetilde{b})_s^2\|_2 \\
&\quad + \max\{\sqrt{m_2}, \sqrt{m_1}\}\|(\widetilde{W})_s^{1,j}\|_F\max\{\|(\widetilde{W})_s^2\|_F^2, \|W_s^2\|_F^2\}\|W_s^1 - (\widetilde{W})_s^1\|_F \\
&\quad + \max\{\sqrt{m_2}, \sqrt{m_1}\}\|(\widetilde{W})_s^{1,j}\|_F\max\{\|(\widetilde{W})_s^2\|_F^2, \|W_s^2\|_F^2\}\|b_s^1 - (\widetilde{b})_s^1\|_2 \\
&\quad + \sqrt{m_2m_1}\|W_s^2\|_F\|W_s^{1,j} - (\widetilde{W})_s^{1,j}\|_F, \\
&\left\|\nabla_{b_s^3}\left(\frac{\partial f_W}{\partial x_j}\right) - \nabla_{b_s^3}\left(\frac{\partial f_W}{\partial x_j}\right)\right\|_F = 0, \\
&\left\|\nabla_{W_s^2}\left(\frac{\partial f_W}{\partial x_j}\right) - \nabla_{W_s^2}\left(\frac{\partial f_{\widetilde{W}}}{\partial x_j}\right)\right\|_F \\
&\leq \|\text{diag}[\sigma'(f_{s,2}^{org}(W))]\|_F - \|\text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F\|W_s^3\|_F \\
&\quad \|W_s^{1,j}\|_F\|\text{diag}[\sigma'(f_{s,1}^{org}(W))]\|_F \\
&\quad + \|\text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F\|W_s^3 - (\widetilde{W})_s^3\|_F\|W_s^{1,j}\|_F\|\text{diag}[\sigma'(f_{s,1}^{org}(W))]\|_F \\
&\quad + \|\text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F\|(\widetilde{W})_s^3\|_F\|W_s^{1,j} - (\widetilde{W})_s^{1,j}\|_F\|\text{diag}[\sigma'(f_{s,1}^{org}(W))]\|_F \\
&\quad + \|\text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F\|(\widetilde{W})_s^3\|_F \\
&\quad \|(\widetilde{W})_s^{1,j}\|_F\|\text{diag}[\sigma'(f_{s,1}^{org}(W))]\|_F - \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \\
&\quad + \|W_s^2 - (\widetilde{W})_s^2\|_F\|\text{diag}[\sigma'(f_{s,1}^{org}(W))]\|_F\|W_s^{1,j}\|_F \\
&\quad \|\text{diag}[\sigma''(f_{s,2}^{org}(W))]\|_F\|W_s^3\|_F\|f_s^1(W)\|_F \\
&\quad + \|(\widetilde{W})_s^2\|_F\|\text{diag}[\sigma'(f_{s,1}^{org}(W))]\|_F - \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F\|W_s^{1,j}\|_F
\end{aligned}$$

$$\begin{aligned}
& \|\text{diag}[\sigma''(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \|f_s^1(W)\|_F \\
& + \|(\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|W_s^{1,j} - (\widetilde{W})_s^{1,j}\|_F \\
& \quad \|\text{diag}[\sigma''(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \|f_s^1(W)\|_F \\
& + \|(\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^{1,j}\|_F \\
& \quad \|\text{diag}[\sigma''(f_{s,2}^{org}(W))] - \text{diag}[\sigma''(f_{s,2}^{org}(\widetilde{W}))]\|_F \|W_s^3\|_F \|f_s^1(W)\|_F \\
& + \|(\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^{1,j}\|_F \\
& \quad \|\text{diag}[\sigma''(f_{s,2}^{org}(\widetilde{W}))]\|_F \|W_s^3 - (\widetilde{W})_s^3\|_F \|f_s^1(W)\|_F \\
& + \|(\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^{1,j}\|_F \\
& \quad \|\text{diag}[\sigma''(f_{s,2}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^3\|_F \|f_s^1(W) - f_s^1(\widetilde{W})\|_F \\
\leq & C(\sigma)(\sqrt{m_2}m_1 \max\{\|W_s^{1,j}\|_F, \|(\widetilde{W})_s^{1,j}\|_F\}) \|(\widetilde{W})_s^2\|_F \|W_s^3 - (\widetilde{W})_s^3\|_F \\
& + \max\{\sqrt{m_2}, \sqrt{m_1}\} m_1 \|W_s^3\|_F \\
& \quad \max\{\|W_s^{1,j}\|_F, \|(\widetilde{W})_s^{1,j}\|_F\} \|(\widetilde{W})_s^2\|_F \|W_s^2 - (\widetilde{W})_s^2\|_F \\
& + m_1 \|W_s^3\|_F \max\{\|W_s^{1,j}\|_F, \|(\widetilde{W})_s^{1,j}\|_F\} \|(\widetilde{W})_s^2\|_F \|b_s^2 - (\widetilde{b})_s^2\|_2 \\
& + \sqrt{m_2}m_1 \max\{\|W_s^{1,j}\|_F, \|(\widetilde{W})_s^{1,j}\|_F\} \|(\widetilde{W})_s^2\|_F^2 \\
& \quad \max\{\|W_s^3\|_F, \|(\widetilde{W})_s^3\|_F\} \|W_s^1 - (\widetilde{W})_s^1\|_F \\
& + \sqrt{m_2}m_1 \|(\widetilde{W})_s^2\|_F \max\{\|W_s^3\|_F, \|(\widetilde{W})_s^3\|_F\} \|W_s^{1,j} - (\widetilde{W})_s^{1,j}\|_F \\
& + \sqrt{m_2}m_1 \max\{\|W_s^{1,j}\|_F, \|(\widetilde{W})_s^{1,j}\|_F\} \|(\widetilde{W})_s^2\|_F^2 \\
& \quad \max\{\|W_s^3\|_F, \|(\widetilde{W})_s^3\|_F\} \|b_s^1 - (\widetilde{b})_s^1\|_2, \\
& \left\| \nabla_{b_s^2} \left( \frac{\partial f_W}{\partial x_j} \right) - \nabla_{b_s^2} \left( \frac{\partial f_{\widetilde{W}}}{\partial x_j} \right) \right\|_F \\
\leq & \|W_s^2 - (\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(W))]\|_F \|W_s^{1,j}\|_F \|\text{diag}[\sigma''(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \\
& + \|(\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(W))] - \text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \\
& \quad \|W_s^{1,j}\|_F \|\text{diag}[\sigma''(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \\
& + \|(\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|W_s^{1,j} - (\widetilde{W})_s^{1,j}\|_F \\
& \quad \|\text{diag}[\sigma''(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \\
& + \|(\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^{1,j}\|_F \\
& \quad \|\text{diag}[\sigma''(f_{s,2}^{org}(W))] - \text{diag}[\sigma''(f_{s,2}^{org}(\widetilde{W}))]\|_F \|W_s^3\|_F \\
& + \|(\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^{1,j}\|_F \\
& \quad \|\text{diag}[\sigma''(f_{s,2}^{org}(\widetilde{W}))]\|_F \|W_s^3 - (\widetilde{W})_s^3\|_F \\
\leq & C(\sigma)(\sqrt{m_2}m_1 \|(\widetilde{W})_s^2\|_F \|(\widetilde{W})_s^{1,j}\|_F \|W_s^3 - (\widetilde{W})_s^3\|_F \\
& + \max\{\sqrt{m_2}, \sqrt{m_1}\} m_1 \|W_s^3\|_F \|(\widetilde{W})_s^2\|_F
\end{aligned}$$

$$\begin{aligned}
& \max\{\|W_s^{1,j}\|_F, \|(\widetilde{W})_s^{1,j}\|_F\} \|W_s^2 - (\widetilde{W})_s^2\|_F \\
& + m_1 \|W_s^3\|_F \|(\widetilde{W})_s^2\|_F \|(\widetilde{W})_s^{1,j}\|_F \|b_s^2 - (\widetilde{b})_s^2\|_2 \\
& + \sqrt{m_2} m_1 \|(\widetilde{W})_s^2\|_F \|W_s^3\|_F \|W_s^{1,j} - (\widetilde{W})_s^{1,j}\|_F \\
& + \max\{\sqrt{m_2}, \sqrt{m_1}\} \|(\widetilde{W})_s^2\|_F \|W_s^3\|_F \|(\widetilde{W})_s^2\|_F \\
& \quad \max\{\|W_s^{1,j}\|_F, \|(\widetilde{W})_s^{1,j}\|_F\} \|W_s^1 - (\widetilde{W})_s^1\|_F \\
& + \max\{\sqrt{m_2}, \sqrt{m_1}\} \|(\widetilde{W})_s^2\|_F \|W_s^3\|_F \|(\widetilde{W})_s^2\|_F \\
& \quad \max\{\|W_s^{1,j}\|_F, \|(\widetilde{W})_s^{1,j}\|_F\} \|b_s^1 - (\widetilde{b})_s^1\|_2, \\
& \left\| \nabla_{W_s^1} \left( \frac{\partial f_W}{\partial x_j} \right) - \nabla_{W_s^1} \left( \frac{\partial f_{\widetilde{W}}}{\partial x_j} \right) \right\|_F \\
& \leq \|\text{diag}[\sigma'(f_{s,1}^{org}(W))] - \text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|W_s^2\|_F \\
& \quad \|\text{diag}[\sigma'(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \\
& + \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|W_s^2 - (\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \\
& + \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^2\|_F \\
& \quad \|\text{diag}[\sigma'(f_{s,2}^{org}(W))] - \text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F \|W_s^3\|_F \\
& + \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F \|W_s^3 - (\widetilde{W})_s^3\|_F \\
& + \|\text{diag}[\sigma''(f_{s,1}^{org}(W))] - \text{diag}[\sigma''(f_{s,1}^{org}(\widetilde{W}))]\|_F \|W_s^{1,j}\|_F \|W_s^2\|_F \\
& \quad \|\text{diag}[\sigma'(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \|x\|_2 \\
& + \|\text{diag}[\sigma''(f_{s,1}^{org}(\widetilde{W}))]\|_F \|W_s^{1,j} - (\widetilde{W})_s^{1,j}\|_F \|W_s^2\|_F \\
& \quad \|\text{diag}[\sigma'(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \|x\|_2 \\
& + \|\text{diag}[\sigma''(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^{1,j}\|_F \|W_s^2 - (\widetilde{W})_s^2\|_F \\
& \quad \|\text{diag}[\sigma'(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \|x\|_2 \\
& + \|\text{diag}[\sigma''(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^{1,j}\|_F \|(\widetilde{W})_s^2\|_F \\
& \quad \|\text{diag}[\sigma'(f_{s,2}^{org}(W))] - \text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F \|W_s^3\|_F \|x\|_2 \\
& + \|\text{diag}[\sigma''(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^{1,j}\|_F \|(\widetilde{W})_s^2\|_F \\
& \quad \|\text{diag}[\sigma'(f_{s,2}^{org}(\widetilde{W}))]\|_F \|W_s^3 - (\widetilde{W})_s^3\|_F \|x\|_2 \\
& + \|\text{diag}[\sigma'(f_{s,1}^{org}(W))] - \text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|W_s^2\|_F \|W_s^2\|_F \\
& \quad \|\text{diag}[\sigma'(f_{s,1}^{org}(W))]\|_F \|W_s^{1,j}\|_F \|\text{diag}[\sigma''(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \|x\|_2 \\
& + \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|W_s^2 - (\widetilde{W})_s^2\|_F \|W_s^2\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(W))]\|_F \\
& \quad \|W_s^{1,j}\|_F \|\text{diag}[\sigma''(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \|x\|_2 \\
& + \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^2\|_F \|W_s^2 - (\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(W))]\|_F \\
& \quad \|W_s^{1,j}\|_F \|\text{diag}[\sigma''(f_{s,2}^{org}(W))]\|_F \|W_s^3\|_F \|x\|_2 \\
& + \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^2\|_F \|(\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(W))] - \text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F
\end{aligned}$$

$$\begin{aligned}
& \|W_s^{1,j}\|_F \|\text{diag}[\sigma''(f_2^{org}(W))]\|_F \|W_s^3\|_F \|x\|_2 \\
& + \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^2\|_F \|(\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \\
& \|W_s^{1,j} - (\widetilde{W})_s^{1,j}\|_F \|\text{diag}[\sigma''(f_2^{org}(W))]\|_F \|W_s^3\|_F \|x\|_2 \\
& + \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^2\|_F \|(\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^{1,j}\|_F \\
& \|\text{diag}[\sigma''(f_2^{org}(W)) - \text{diag}[\sigma''(f_2^{org}(\widetilde{W}))]\|_F \|W_s^3\|_F \|x\|_2 \\
& + \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^2\|_F \|(\widetilde{W})_s^2\|_F \|\text{diag}[\sigma'(f_{s,1}^{org}(\widetilde{W}))]\|_F \|(\widetilde{W})_s^{1,j}\|_F \\
& \|\text{diag}[\sigma''(f_2^{org}(\widetilde{W}))]\|_F \|W_s^3 - (\widetilde{W})_s^3\|_F \|x\|_2 \\
\leq & C(\sigma)(\sqrt{m_2}m_1\|(\widetilde{W})_s^2\|_F^2\|(\widetilde{W})_s^{1,j}\|_F\|W_s^3 - (\widetilde{W})_s^3\|_F \\
& + \sqrt{m_2}m_1^{3/2}\|W_s^3\|_F \max\{\|W_s^{1,j}\|_F, \|(\widetilde{W})_s^{1,j}\|_F\} \\
& \max\{\|W_s^2\|_F^2, \|(\widetilde{W})_s^2\|_F^2\}\|W_s^2 - (\widetilde{W})_s^2\|_F \\
& + m_1\|(\widetilde{W})_s^2\|_F^2\|W_s^3\|_F\|(\widetilde{W})_s^{1,j}\|_F\|b_s^2 - (\widetilde{b})_s^2\|_2 \\
& + \sqrt{m_2}m_1\|W_s^3\|_F \max\{\|W_s^{1,j}\|_F, \|(\widetilde{W})_s^{1,j}\|_F\} \\
& \max\{\|W_s^2\|_F^3, \|(\widetilde{W})_s^2\|_F^3\}\|W_s^1 - (\widetilde{W})_s^1\|_F \\
& + \sqrt{m_2}m_1\|W_s^3\|_F \max\{\|W_s^2\|_F^2, \|(\widetilde{W})_s^2\|_F^2\}\|W_s^{1,j} - (\widetilde{W})_s^{1,j}\|_F \\
& + \sqrt{m_2}m_1\|W_s^3\|_F \max\{\|W_s^{1,j}\|_F, \|(\widetilde{W})_s^{1,j}\|_F\} \\
& \max\{\|W_s^2\|_F^3, \|(\widetilde{W})_s^2\|_F^3\}\|b_s^1 - (\widetilde{b})_s^1\|_F, \\
& \left\| \nabla_{b_s^1} \left( \frac{\partial f_W}{\partial x_j} \right) - \nabla_{b_s^1} \left( \frac{\partial f_{\widetilde{W}}}{\partial x_j} \right) \right\|_F \\
\leq & C(\sigma)(\sqrt{m_2}m_1\|(\widetilde{W})_s^2\|_F^2\|(\widetilde{W})_s^{1,j}\|_F\|W_s^3 - (\widetilde{W})_s^3\|_F \\
& + \sqrt{m_2}m_1^{3/2}\|W_s^3\|_F \max\{\|W_s^{1,j}\|_F, \|(\widetilde{W})_s^{1,j}\|_F\} \\
& \max\{\|W_s^2\|_F^2, \|(\widetilde{W})_s^2\|_F^2\}\|W_s^2 - (\widetilde{W})_s^2\|_F \\
& + m_1\|(\widetilde{W})_s^2\|_F^2\|W_s^3\|_F\|(\widetilde{W})_s^{1,j}\|_F\|b_s^2 - (\widetilde{b})_s^2\|_2 \\
& + \sqrt{m_2}m_1\|W_s^3\|_F \max\{\|W_s^{1,j}\|_F, \|(\widetilde{W})_s^{1,j}\|_F\} \\
& \max\{\|W_s^2\|_F^3, \|(\widetilde{W})_s^2\|_F^3\}\|W_s^1 - (\widetilde{W})_s^1\|_F \\
& + \sqrt{m_2}m_1\|W_s^3\|_F \max\{\|W_s^2\|_F^2, \|(\widetilde{W})_s^2\|_F^2\}\|W_s^{1,j} - (\widetilde{W})_s^{1,j}\|_F \\
& + \sqrt{m_2}m_1\|W_s^3\|_F \max\{\|W_s^{1,j}\|_F, \|(\widetilde{W})_s^{1,j}\|_F\} \\
& \max\{\|W_s^2\|_F^3, \|(\widetilde{W})_s^2\|_F^3\}\|b_s^1 - (\widetilde{b})_s^1\|_F).
\end{aligned}$$

In the final step, we deal with  $\|\nabla_{W_s^l} \widehat{L}_R(f_W) - \nabla_{W_s^l} \widehat{L}_R(f_{\widetilde{W}})\|_F, \|\nabla_{b_s^l} \widehat{L}_R(f_W) - \nabla_{b_s^l} \widehat{L}_R(f_{\widetilde{W}})\|_2$  for  $l = 1, 2, 3$ . To avoid excessive verbosity, we only present an estimate for  $\|\nabla_{W_s^1} \widehat{L}_R(f_W) - \nabla_{W_s^1} \widehat{L}_R(f_{\widetilde{W}})\|_F$ , and similar estimates can be obtained for the other terms.  $\nabla_{W_s^1} \widehat{L}_R(f_W) - \nabla_{W_s^1} \widehat{L}_R(f_{\widetilde{W}})$  can be calculated by

$$\nabla_{W_s^1} \widehat{L}_R(f_W) - \nabla_{W_s^1} \widehat{L}_R(f_{\widetilde{W}})$$

$$\begin{aligned}
&= \frac{|\Omega|}{n} \sum_{i=1}^n \sum_{j=1}^d \left[ \frac{\partial f_W(X_i)}{\partial x_j} \nabla_{W_s^1} \left( \frac{\partial f_W(X_i)}{\partial x_j} \right) - \frac{\partial f_{\widetilde{W}}(X_i)}{\partial x_j} \nabla_{W_s^1} \left( \frac{\partial f_{\widetilde{W}}(X_i)}{\partial x_j} \right) \right] \\
&+ \frac{|\Omega|}{n} \sum_{i=1}^n w(X_i) [f_W(X_i) \nabla_{W_s^1} f_W(X_i) - f_{\widetilde{W}}(X_i) \nabla_{W_s^1} f_{\widetilde{W}}(X_i)] \\
&- \frac{|\Omega|}{n} \sum_{i=1}^n f(X_i) [\nabla_{W_s^1} f_W(X_i) - \nabla_{W_s^1} f_{\widetilde{W}}(X_i)] \\
&+ \frac{|\partial\Omega|}{\beta m} \sum_{i=1}^m [f_W(Y_i) \nabla_{W_s^1} f_W(Y_i) - f_{\widetilde{W}}(Y_i) \nabla_{W_s^1} f_{\widetilde{W}}(Y_i)] \\
&- \frac{|\partial\Omega|}{\beta m} \sum_{i=1}^m g(Y_i) [\nabla_{W_s^1} f_W(Y_i) - \nabla_{W_s^1} f_{\widetilde{W}}(Y_i)].
\end{aligned}$$

The first term on the right-hand side is bounded by

$$\begin{aligned}
&\sum_{j=1}^d \left\| \frac{\partial f_W(X_i)}{\partial x_j} \nabla_{W_s^1} \left( \frac{\partial f_W(X_i)}{\partial x_j} \right) - \frac{\partial f_{\widetilde{W}}(X_i)}{\partial x_j} \nabla_{W_s^1} \left( \frac{\partial f_{\widetilde{W}}(X_i)}{\partial x_j} \right) \right\|_F \\
&\leq \sum_{j=1}^d \left| \frac{\partial f_W(X_i)}{\partial x_j} - \frac{\partial f_{\widetilde{W}}(X_i)}{\partial x_j} \right| \left\| \nabla_{W_s^1} \left( \frac{\partial f_W(X_i)}{\partial x_j} \right) \right\|_F \\
&+ \sum_{j=1}^d \left| \frac{\partial f_{\widetilde{W}}(X_i)}{\partial x_j} \right| \left\| \nabla_{W_s^1} \left( \frac{\partial f_W(X_i)}{\partial x_j} \right) - \nabla_{W_s^1} \left( \frac{\partial f_{\widetilde{W}}(X_i)}{\partial x_j} \right) \right\|_F \\
&\leq \|\nabla_x f_W(X_i) - \nabla_x f_{\widetilde{W}}(X_i)\|_2 \left( \sum_{j=1}^d \left\| \nabla_{W_s^1} \left( \frac{\partial f_W(X_i)}{\partial x_j} \right) \right\|_F^2 \right)^{1/2} \\
&+ \|\nabla_x f_{\widetilde{W}}(X_i)\|_2 \left( \sum_{j=1}^d \left\| \nabla_{W_s^1} \left( \frac{\partial f_W(X_i)}{\partial x_j} \right) - \nabla_{W_s^1} \left( \frac{\partial f_{\widetilde{W}}(X_i)}{\partial x_j} \right) \right\|_F^2 \right)^{1/2}.
\end{aligned}$$

The fourth term on the right-hand side is bounded by (and there is a similar bound for the second term)

$$\|f_W \nabla_{W_s^1} f_W - f_{\widetilde{W}} \nabla_{W_s^1} f_{\widetilde{W}}\|_F \leq |f_W - f_{\widetilde{W}}| \|\nabla_{W_s^1} f_W\|_F + |f_{\widetilde{W}}| \|\nabla_{W_s^1} f_W - \nabla_{W_s^1} f_{\widetilde{W}}\|_F$$

Now, making use of the estimates of

$$\begin{aligned}
&|f_W|, |f_W - f_{\widetilde{W}}|, \|\nabla_x f_W\|_2, \|\nabla_x f_W - \nabla_x f_{\widetilde{W}}\|_2, \|\nabla_{W_s^l} f_W\|_F, \|\nabla_{b_s^l} f_W\|_2, \\
&\|\nabla_{W_s^l} f_W - \nabla_{W_s^l} f_{\widetilde{W}}\|_F, \|\nabla_{b_s^l} f_W - \nabla_{b_s^l} f_{\widetilde{W}}\|_2, \left\| \nabla_{W_s^l} \left( \frac{\partial f_W}{\partial x_j} \right) \right\|_F, \left\| \nabla_{b_s^l} \left( \frac{\partial f_W}{\partial x_j} \right) \right\|_2, \\
&\left\| \nabla_{W_s^l} \left( \frac{\partial f_W}{\partial x_j} \right) - \nabla_{W_s^l} \left( \frac{\partial f_{\widetilde{W}}}{\partial x_j} \right) \right\|_F, \left\| \nabla_{b_s^l} \left( \frac{\partial f_W}{\partial x_j} \right) - \nabla_{b_s^l} \left( \frac{\partial f_{\widetilde{W}}}{\partial x_j} \right) \right\|_F \quad (l = 1, 2, 3)
\end{aligned}$$

just obtained in the first four steps, we derive that

$$\begin{aligned}
& \|\nabla_{W_s^1} \widehat{L}_R(f_W) - \nabla_{W_s^1} \widehat{L}_R(f_{\widetilde{W}})\|_F \\
& \leq C(\sigma, f, w, \Omega) \sqrt{m_2 m_1} \|W_s^1\|_F \|W_s^2\|_F^2 \|W_s^3\|_F \\
& \quad \sum_{s'=1}^A [\sqrt{m_2 m_1} \|(\widetilde{W})_{s'}^1\|_F \|(\widetilde{W})_{s'}^2\|_F \|W_{s'}^3 - (\widetilde{W})_{s'}^3\|_F + |b_{s'}^3 - (\widetilde{b})_{s'}^3| \\
& \quad + \max\{\sqrt{m_2}, \sqrt{m_1}\} \sqrt{m_1} \|(\widetilde{W})_{s'}^1\|_F \|(\widetilde{W})_{s'}^2\|_F \\
& \quad \max\{\|W_{s'}^3\|_F, \|(\widetilde{W})_{s'}^3\|_F\} \|W_{s'}^2 - (\widetilde{W})_{s'}^2\|_F \\
& \quad + \sqrt{m_1} \|(\widetilde{W})_{s'}^1\|_F \|(\widetilde{W})_{s'}^2\|_F \max\{\|W_{s'}^3\|_F, \|(\widetilde{W})_{s'}^3\|_F\} \|b_{s'}^2 - (\widetilde{b})_{s'}^2\|_2 \\
& \quad + \sqrt{m_2 m_1} \max\{\|W_{s'}^3\|_F, \|(\widetilde{W})_{s'}^3\|_F\} \|(\widetilde{W})_{s'}^1\|_F \\
& \quad \max\{\|W_{s'}^2\|_F^2, \|(\widetilde{W})_{s'}^2\|_F^2\} \|W_{s'}^1 - (\widetilde{W})_{s'}^1\|_F \\
& \quad + \max\{\sqrt{m_2}, \sqrt{m_1}\} \|(\widetilde{W})_{s'}^1\|_F \max\{\|W_{s'}^3\|_F, \|(\widetilde{W})_{s'}^3\|_F\} \\
& \quad \max\{\|W_{s'}^2\|_F^2, \|(\widetilde{W})_{s'}^2\|_F^2\} \|b_{s'}^1 - (\widetilde{b})_{s'}^1\|_2] \\
& + C(\sigma, f, w, \Omega) \left( \sum_{s'=1}^A \max\{\sqrt{m_2 m_1} \|(\widetilde{W})_{s'}^1\|_2 \|(\widetilde{W})_{s'}^2\|_2 \|(\widetilde{W})_{s'}^3\|_2, |(\widetilde{b})_{s'}^3|\} \right) \\
& m_1 \left( \sum_{j=1}^d \max\{\|W_s^{1,j}\|_F^2, \|(\widetilde{W})_s^{1,j}\|_F^2\} \right)^{1/2} \\
& (\sqrt{m_2} \|(\widetilde{W})_s^2\|_F^2 \|W_s^3 - (\widetilde{W})_s^3\|_F \\
& + \sqrt{m_2} m_1^{1/2} \|W_s^3\|_F \max\{\|W_s^2\|_F^2, \|(\widetilde{W})_s^2\|_F^2\} \|W_s^2 - (\widetilde{W})_s^2\|_F \\
& + \|(\widetilde{W})_s^2\|_F^2 \|W_s^3\|_F \|b_s^2 - (\widetilde{b})_s^2\|_2 \\
& + \sqrt{m_2} \|W_s^3\|_F \max\{\|W_s^2\|_F^3, \|(\widetilde{W})_s^2\|_F^3\} \|W_s^1 - (\widetilde{W})_s^1\|_F \\
& + \sqrt{m_2} \|W_s^3\|_F \max\{\|W_s^2\|_F^3, \|(\widetilde{W})_s^2\|_F^3\} \|b_s^1 - (\widetilde{b})_s^1\|_F) \\
& + C(\sigma, g, \partial\Omega) \frac{1}{\beta} \sqrt{m_1 m_2} \|W_s^2\|_F \|W_s^3\|_F \\
& \sum_{s'=1}^A (\sqrt{m_2} \|W_{s'}^3 - (\widetilde{W})_{s'}^3\|_2 + |b_{s'}^3 - (\widetilde{b})_{s'}^3| + \sqrt{m_1} \|(\widetilde{W})_{s'}^3\|_2 \|W_{s'}^2 - (\widetilde{W})_{s'}^2\|_2 \\
& + \|(\widetilde{W})_{s'}^3\|_2 \|b_{s'}^2 - (\widetilde{b})_{s'}^2\|_2 + \|(\widetilde{W})_{s'}^3\|_2 \|(\widetilde{W})_{s'}^2\|_2 \|W_{s'}^1 - (\widetilde{W})_{s'}^1\|_2 \\
& + L_\sigma^2 \|(\widetilde{W})_{s'}^3\|_2 \|(\widetilde{W})_{s'}^2\|_2 \|b_{s'}^1 - (\widetilde{b})_{s'}^1\|_2) \\
& + C(\sigma, g, \partial\Omega) \frac{1}{\beta} \sum_{s'=1}^A (\sqrt{m_2} \|(\widetilde{W})_{s'}^3\|_F + |(\widetilde{b})_{s'}^3|) (\sqrt{m_2 m_1} \|(\widetilde{W})_{s'}^2\|_F \|W_{s'}^3 - (\widetilde{W})_{s'}^3\|_F \\
& + \max\{\sqrt{m_2}, \sqrt{m_1}\} \sqrt{m_1} \|W_{s'}^3\|_F \|(\widetilde{W})_{s'}^2\|_F \|W_{s'}^2 - (\widetilde{W})_{s'}^2\|_F \\
& + \sqrt{m_1} \|(\widetilde{W})_{s'}^2\|_F \|W_{s'}^3\|_F \|b_{s'}^2 - (\widetilde{b})_{s'}^2\|_2)
\end{aligned}$$

+ m

$$+ \max\{\sqrt{m_2}, \sqrt{m_1}\} \max\{\|W_s^2\|_F^2, \|(\widetilde{W})_s^2\|_F^2\} \\ \|W_s^3\|_F \|b_s^1 - (\widetilde{b})_s^1\|_2.$$

Combining the estimates of  $\|\nabla_{W_s^l} \widehat{L}_R(f_W) - \nabla_{W_s^l} \widehat{L}_R(f_{\widetilde{W}})\|_F, \|\nabla_{b_s^l} \widehat{L}_R(f_W) - \nabla_{b_s^l} \widehat{L}_R(f_{\widetilde{W}})\|_2$  for  $l = 1, 2, 3$ , we arrive at

$$\|\nabla_W \widehat{L}_R(f_W) - \nabla_W \widehat{L}_R(f_{\widetilde{W}})\|^2 \\ \leq C(d, \sigma, f, g, w, \Omega) A^2 \{1/\beta^2, 1\} \max_{1 \leq s \leq A} [\max\{\|W_s^1\|_F^6, \|(\widetilde{W})_s^1\|_F^6\} \\ \max\{\|W_s^2\|_F^8, \|(\widetilde{W})_s^2\|_F^8\} \max\{\|W_s^3\|_F^6, \|(\widetilde{W})_s^3\|_F^6, |(\widetilde{b})_s^3|^6\}] \|W - \widetilde{W}\|^2.$$

□

## References

- [1] Robert A Adams and John JF Fournier. *Sobolev spaces*. Elsevier, 2003.
- [2] Zeyuan Allen-Zhu, Yuanzhi Li, and Yingyu Liang. Learning and generalization in overparameterized neural networks, going beyond two layers. *Advances in neural information processing systems*, 32, 2019.
- [3] Zeyuan Allen-Zhu, Yuanzhi Li, and Zhao Song. A convergence theory for deep learning via over-parameterization. In *International conference on machine learning*, pages 242–252. PMLR, 2019.
- [4] Zeyuan Allen-Zhu, Yuanzhi Li, and Zhao Song. On the convergence rate of training recurrent neural networks. *Advances in neural information processing systems*, 32, 2019.
- [5] Sanjeev Arora, Simon Du, Wei Hu, Zhiyuan Li, and Ruosong Wang. Fine-grained analysis of optimization and generalization for overparameterized two-layer neural networks. In *International Conference on Machine Learning*, pages 322–332. PMLR, 2019.
- [6] Arindam Banerjee, Pedro Cisneros-Velarde, Libin Zhu, and Mikhail Belkin. Neural tangent kernel at initialization: linear width suffices. In *Uncertainty in Artificial Intelligence*, pages 110–118. PMLR, 2023.
- [7] Peter L Bartlett, Nick Harvey, Christopher Liaw, and Abbas Mehrabian. Nearly-tight vc-dimension and pseudodimension bounds for piecewise linear neural networks. *Journal of Machine Learning Research*, 20(63):1–17, 2019.
- [8] Peter L Bartlett and Shahar Mendelson. Rademacher and gaussian complexities: Risk bounds and structural results. *Journal of Machine Learning Research*, 3(Nov):463–482, 2002.
- [9] Peter L Bartlett, Andrea Montanari, and Alexander Rakhlin. Deep learning: a statistical viewpoint. *Acta numerica*, 30:87–201, 2021.

- [10] Dimitri Bertsekas. *Convex optimization theory*, volume 1. Athena Scientific, 2009.
- [11] Susanne C Brenner and L Ridgway Scott. The mathematical theory of finite element methods.
- [12] Yuan Cao and Quanquan Gu. Generalization bounds of stochastic gradient descent for wide and deep neural networks. *Advances in neural information processing systems*, 32, 2019.
- [13] Yuan Cao and Quanquan Gu. Generalization error bounds of gradient descent for learning over-parameterized deep relu networks. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 34, pages 3349–3356, 2020.
- [14] Fan Chen, Jianguo Huang, Chunmei Wang, and Haizhao Yang. Friedrichs learning: Weak solutions of partial differential equations via deep learning. *SIAM Journal on Scientific Computing*, 45(3):A1271–A1299, 2023.
- [15] Zixiang Chen, Yuan Cao, Difan Zou, and Quanquan Gu. How much over-parameterization is sufficient to learn deep relu networks? In *International Conference on Learning Representation (ICLR)*, 2021.
- [16] Lenaïc Chizat and Francis Bach. On the global convergence of gradient descent for over-parameterized models using optimal transport. *Advances in neural information processing systems*, 31, 2018.
- [17] Philippe G Ciarlet. *The finite element method for elliptic problems*. SIAM, 2002.
- [18] Tim De Ryck, Samuel Lanthaler, and Siddhartha Mishra. On the approximation of functions by tanh neural networks. *Neural Networks*, 143:732–750, 2021.
- [19] Selina Drews and Michael Kohler. Analysis of the expected  $l_2$  error of an over-parametrized deep neural network estimate learned by gradient descent without regularization. *arXiv preprint arXiv:2311.14609*, 2023.
- [20] Selina Drews and Michael Kohler. On the universal consistency of an over-parametrized deep neural network estimate learned by gradient descent. *Annals of the Institute of Statistical Mathematics*, pages 1–31, 2024.
- [21] Simon Du, Jason Lee, Haochuan Li, Liwei Wang, and Xiyu Zhai. Gradient descent finds global minima of deep neural networks. In *International conference on machine learning*, pages 1675–1685. PMLR, 2019.
- [22] Simon S Du, Xiyu Zhai, Barnabas Poczos, and Aarti Singh. Gradient descent provably optimizes over-parameterized neural networks. In *International Conference on Learning Representations*, 2018.

- [23] C Duan, Y Jiao, Y Lai, X Lu, Q Quan, and JZ Yang. Deep ritz methods for laplace equations with dirichlet boundary condition. *Commun. Comput. Phys.*, 31(4):1020–1048, 2022.
- [24] Chenguang Duan, Yuling Jiao, Yanming Lai, Dingwei Li, Jerry Zhijian Yang, et al. Convergence rate analysis for deep ritz method. *Communications in Computational Physics*, 31(4):1020–1048, 2022.
- [25] Cong Fang, Jason Lee, Pengkun Yang, and Tong Zhang. Modeling from features: a mean-field framework for over-parameterized deep neural networks. In *Conference on learning theory*, pages 1887–1936. PMLR, 2021.
- [26] Noah Golowich, Alexander Rakhlin, and Ohad Shamir. Size-independent sample complexity of neural networks. In *Conference On Learning Theory*, pages 297–299. PMLR, 2018.
- [27] Pierre Grisvard. *Elliptic problems in nonsmooth domains*. SIAM, 2011.
- [28] Ingo Gühring, Gitta Kutyniok, and Philipp Petersen. Error bounds for approximations with deep relu neural networks in  $w$ ,  $s$ ,  $p$  norms. *Analysis and Applications*, 18(05):803–859, 2020.
- [29] Ingo Gühring and Mones Raslan. Approximation rates for neural networks with encodable weights in smoothness spaces. *Neural Networks*, 134:107–130, 2021.
- [30] Jiequn Han, Arnulf Jentzen, and Weinan E. Solving high-dimensional partial differential equations using deep learning. *Proceedings of the National Academy of Sciences*, 115(34):8505–8510, 2018.
- [31] Qingguo Hong, Jonathan W Siegel, and Jinchao Xu. A priori analysis of stable neural network solutions to numerical pdes. *arXiv preprint arXiv:2104.02903*, 2021.
- [32] Arthur Jacot, Franck Gabriel, and Clément Hongler. Neural tangent kernel: Convergence and generalization in neural networks. *Advances in neural information processing systems*, 31, 2018.
- [33] Yuling Jiao, Yanming Lai, Dingwei Li, Xiliang Lu, Fengru Wang, Jerry Zhijian Yang, et al. A rate of convergence of physics informed neural networks for the linear second order elliptic pdes. *Communications in Computational Physics*, 31(4):1272–1295, 2022.
- [34] Yuling Jiao, Yanming Lai, Yisu Lo, Yang Wang, and Yunfei Yang. Error analysis of deep ritz methods for elliptic equations. *Analysis and Applications*, 22(01):57–87, 2024.
- [35] Yuling Jiao, Yanming Lai, Xiliang Lu, Fengru Wang, Jerry Zhijian Yang, and Yuanyuan Yang. Deep neural networks with relu-sine-exponential activations break curse of dimensionality in approximation on hölder class. *SIAM Journal on Mathematical Analysis*, 55(4):3635–3649, 2023.

- [36] Yuling Jiao, Yanming Lai, Yang Wang, Haizhao Yang, and Yunfei Yang. Convergence analysis of the deep galerkin method for weak solutions. In *From Classical Analysis to Analysis on Fractals: A Tribute to Robert Strichartz, Volume 1*, pages 53–82. Springer, 2023.
- [37] Michael Kohler and Adam Krzyzak. Analysis of the rate of convergence of an over-parametrized deep neural network estimate learned by gradient descent. *arXiv preprint arXiv:2210.01443*, 2022.
- [38] Chaoyue Liu, Libin Zhu, and Mikhail Belkin. Loss landscapes and optimization in over-parameterized non-linear systems and neural networks. *Applied and Computational Harmonic Analysis*, 59:85–116, 2022.
- [39] Chaoyue Liu, Libin Zhu, and Misha Belkin. On the linearity of large non-linear models: when and why the tangent kernel is constant. *Advances in Neural Information Processing Systems*, 33:15954–15964, 2020.
- [40] Zichao Long, Yiping Lu, Xianzhong Ma, and Bin Dong. Pde-net: Learning pdes from data. In *International conference on machine learning*, pages 3208–3216. PMLR, 2018.
- [41] Jianfeng Lu and Yulong Lu. A priori generalization error analysis of two-layer neural networks for solving high dimensional schrödinger eigenvalue problems. *Communications of the American Mathematical Society*, 2(1):1–21, 2022.
- [42] Jianfeng Lu, Zuowei Shen, Haizhao Yang, and Shijun Zhang. Deep network approximation for smooth functions. *SIAM Journal on Mathematical Analysis*, 53(5):5465–5506, 2021.
- [43] Lu Lu, Xuhui Meng, Zhiping Mao, and George Em Karniadakis. Deepxde: A deep learning library for solving differential equations. *SIAM review*, 63(1):208–228, 2021.
- [44] Yiping Lu, Haoxuan Chen, Jianfeng Lu, Lexing Ying, and Jose Blanchet. Machine learning for elliptic pdes: Fast rate generalization bound, neural scaling law and minimax optimality. In *International Conference on Learning Representations*, 2022.
- [45] Yulong Lu, Jianfeng Lu, and Min Wang. A priori generalization analysis of the deep ritz method for solving high dimensional elliptic partial differential equations. In *Conference on learning theory*, pages 3196–3241. PMLR, 2021.
- [46] Costabel Martin and Dauge Monique. A singularly mixed boundary value problem. *Communications in Partial Differential Equations*, 21(11-12):1919–1949, 1996.

- [47] Song Mei, Andrea Montanari, and Phan-Minh Nguyen. A mean field view of the landscape of two-layer neural networks. *Proceedings of the National Academy of Sciences*, 115(33):E7665–E7671, 2018.
- [48] Hrushikesh N Mhaskar. Neural networks for optimal approximation of smooth and analytic functions. *Neural computation*, 8(1):164–177, 1996.
- [49] Johannes Müller and Marius Zeinhofer. Error estimates for the deep ritz method with boundary penalty. In *Mathematical and Scientific Machine Learning*, pages 215–230. PMLR, 2022.
- [50] Behnam Neyshabur, Ryota Tomioka, and Nathan Srebro. Norm-based capacity control in neural networks. In *Conference on learning theory*, pages 1376–1401. PMLR, 2015.
- [51] Phan-Minh Nguyen and Huy Tuan Pham. A rigorous framework for the mean field limit of multilayer neural networks. *Mathematical Statistics and Learning*, 6(3):201–357, 2023.
- [52] Quynh Nguyen. On the proof of global convergence of gradient descent for deep relu networks with linear widths. In *International Conference on Machine Learning*, pages 8056–8062. PMLR, 2021.
- [53] Quynh N Nguyen and Marco Mondelli. Global convergence of deep networks with one wide layer followed by pyramidal topology. *Advances in Neural Information Processing Systems*, 33:11961–11972, 2020.
- [54] Samet Oymak and Mahdi Soltanolkotabi. Toward moderate overparameterization: Global convergence guarantees for training shallow neural networks. *IEEE Journal on Selected Areas in Information Theory*, 1(1):84–105, 2020.
- [55] Abhishek Panigrahi, Abhishek Shetty, and Navin Goyal. Effect of activation functions on the training of overparametrized neural nets. In *International Conference on Learning Representations*, 2019.
- [56] Allan Pinkus. Approximation theory of the mlp model in neural networks. *Acta numerica*, 8:143–195, 1999.
- [57] Maziar Raissi. Deep hidden physics models: Deep learning of nonlinear partial differential equations. *Journal of Machine Learning Research*, 19(25):1–24, 2018.
- [58] Maziar Raissi, Paris Perdikaris, and George E Karniadakis. Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations. *Journal of Computational physics*, 378:686–707, 2019.
- [59] Zuowei Shen, Haizhao Yang, and Shijun Zhang. Deep network approximation characterized by number of neurons. *arXiv preprint arXiv:1906.05497*, 2019.

- [60] Zuowei Shen, Haizhao Yang, and Shijun Zhang. Deep network with approximation error being reciprocal of width to power of square root of depth. *Neural Computation*, 33(4):1005–1036, 2021.
- [61] Zuowei Shen, Haizhao Yang, and Shijun Zhang. Neural network approximation: Three hidden layers are enough. *Neural Networks*, 141:160–173, 2021.
- [62] Yeonjong Shin. On the convergence of physics informed neural networks for linear second-order elliptic and parabolic type pdes. *Communications in Computational Physics*, 28(5):2042–2074, 2020.
- [63] Yeonjong Shin, Zhongqiang Zhang, and George Em Karniadakis. Error estimates of residual minimization using neural networks for linear pdes. *Journal of Machine Learning for Modeling and Computing*, 4(4), 2023.
- [64] Albert Nikolayevich Shiriyayev. *Selected Works of AN Kolmogorov: Volume III Information Theory and the Theory of Algorithms*. Springer Publishing Company, Incorporated, 2010.
- [65] Jonathan W Siegel and Jinchao Xu. Approximation rates for neural networks with general activation functions. *Neural Networks*, 128:313–321, 2020.
- [66] Jonathan W Siegel and Jinchao Xu. Sharp bounds on the approximation rates, metric entropy, and n-widths of shallow neural networks. *Foundations of Computational Mathematics*, pages 1–57, 2022.
- [67] Justin Sirignano and Konstantinos Spiliopoulos. Dgm: A deep learning algorithm for solving partial differential equations. *Journal of computational physics*, 375:1339–1364, 2018.
- [68] Justin Sirignano and Konstantinos Spiliopoulos. Mean field analysis of neural networks: A law of large numbers. *SIAM Journal on Applied Mathematics*, 80(2):725–752, 2020.
- [69] Taiji Suzuki. Adaptivity of deep relu network for learning in besov and mixed smooth besov spaces: optimal rate and curse of dimensionality. In *International Conference on Learning Representations*, 2018.
- [70] Roman Vershynin. *High-dimensional probability: An introduction with applications in data science*, volume 47. Cambridge university press, 2018.
- [71] E Weinan, Jiequn Han, and Arnulf Jentzen. Algorithms for solving high dimensional pdes: from nonlinear monte carlo to machine learning. *Nonlinearity*, 35(1):278, 2021.
- [72] E Weinan and Stephan Wojtowytsch. Some observations on high-dimensional partial differential equations with barron data. In *Mathematical and Scientific Machine Learning*, pages 253–269. PMLR, 2022.

- [73] E Weinan and Bing Yu. The deep ritz method: A deep learning-based numerical algorithm for solving variational problems. *Communications in Mathematics and Statistics*, 1(6):1–12, 2018.
- [74] Jinchao Xu. Finite neuron method and convergence analysis. *Communications in Computational Physics*, 28(5):1707–1745, 2020.
- [75] Yahong Yang, Haizhao Yang, and Yang Xiang. Nearly optimal vc-dimension and pseudo-dimension bounds for deep neural network derivatives. *Advances in Neural Information Processing Systems*, 36, 2024.
- [76] Yunfei Yang and Ding-Xuan Zhou. Optimal rates of approximation by shallow relu k neural networks and applications to nonparametric regression. *Constructive Approximation*, pages 1–32, 2024.
- [77] Dmitry Yarotsky. Error bounds for approximations with deep relu networks. *Neural Networks*, 94:103–114, 2017.
- [78] Dmitry Yarotsky. Optimal approximation of continuous functions by very deep relu networks. In *Conference on learning theory*, pages 639–649. PMLR, 2018.
- [79] Dmitry Yarotsky. Elementary superexpressive activations. In *International Conference on Machine Learning*, pages 11932–11940. PMLR, 2021.
- [80] Dmitry Yarotsky and Anton Zhevnerchuk. The phase diagram of approximation rates for deep neural networks. *Advances in neural information processing systems*, 33:13005–13015, 2020.
- [81] Yaohua Zang, Gang Bao, Xiaojing Ye, and Haomin Zhou. Weak adversarial networks for high-dimensional partial differential equations. *Journal of Computational Physics*, 411:109409, 2020.
- [82] Shijun Zhang, Jianfeng Lu, and Hongkai Zhao. Deep network approximation: Beyond relu to diverse activation functions. *Journal of Machine Learning Research*, 25(35):1–39, 2024.
- [83] Shijun Zhang, Zuowei Shen, and Haizhao Yang. Deep network approximation: Achieving arbitrary accuracy with fixed number of neurons. *Journal of Machine Learning Research*, 23(276):1–60, 2022.
- [84] Difan Zou, Yuan Cao, Dongruo Zhou, and Quanquan Gu. Gradient descent optimizes over-parameterized deep relu networks. *Machine learning*, 109:467–492, 2020.
- [85] Difan Zou and Quanquan Gu. An improved analysis of training over-parameterized deep neural networks. *Advances in neural information processing systems*, 32, 2019.