Global Behavior of Learning Dynamics in Zero-Sum Games with Memory Asymmetry

Yuma Fujimoto CyberAgent The University of Tokyo Soken University fujimoto.yuma1991@gmail.com

Kaito Ariu CyberAgent kaitoariu@gmail.com

Kenshi Abe CyberAgent The University of Electro-Communications abekenshi1224@gmail.com

Abstract

This study examines the global behavior of dynamics in learning in games between two players, X and Y. We consider the simplest situation for memory asymmetry between two players: X memorizes the other Y's previous action and uses reactive strategies, while Y has no memory. Although this memory complicates the learning dynamics, we discover two novel quantities that characterize the global behavior of such complex dynamics. One is an extended Kullback-Leibler divergence from the Nash equilibrium, a well-known conserved quantity from previous studies. The other is a family of Lyapunov functions of X 's reactive strategy. These two quantities capture the global behavior in which X's strategy becomes more exploitative, and the exploited Y's strategy converges to the Nash equilibrium. Indeed, we theoretically prove that Y's strategy globally converges to the Nash equilibrium in the simplest game equipped with an equilibrium in the interior of strategy spaces. Furthermore, our experiments also suggest that this global convergence is universal for more advanced zero-sum games than the simplest game. This study provides a novel characterization of the global behavior of learning in games through a couple of indicators.

1 Introduction

Learning in games targets how multiple agents learn their optimal strategies in the repetition of games [\[1\]](#page-8-0). The set of such players' best strategies is defined as Nash equilibrium [\[2\]](#page-8-1), where every player has no motivation to change his/her strategy. However, this equilibrium is hard to compute in general because one's best strategy depends on the others' strategies. Indeed, the behavior of multi-agent learning is complicated in zero-sum games, where players conflict in their payoffs. Even when players try to learn their optimal strategies there, their strategies often cycle around the Nash equilibrium and fail to converge to the equilibrium.

In order to understand such strange behaviors, which are unique in multi-agent learning, the dynamics of how multiple agents learn their strategies, say, learning dynamics, are frequently studied [\[3,](#page-8-2) [4,](#page-8-3) [5,](#page-8-4) [6\]](#page-8-5). The representative dynamics of interest are the replicator dynamics, which is based on the evolutionary dynamics in biology [\[7,](#page-8-6) [8,](#page-8-7) [9,](#page-8-8) [10,](#page-8-9) [11\]](#page-9-0). These dynamics are also known as the multiplicative weight updates (MWU) in its discrete-time version [\[12,](#page-9-1) [13\]](#page-9-2). Furthermore, their connection to other representative learning dynamics, such as gradient ascent [\[14,](#page-9-3) [15,](#page-9-4) [16,](#page-9-5) [17\]](#page-9-6) and

Q-learning [\[18,](#page-9-7) [19,](#page-9-8) [20\]](#page-9-9), should be noted. Such replicator dynamics are known to be characterized by Kullbuck-Leibler (KL) divergence, which is the distance from the Nash equilibrium to the players' present strategies. This KL divergence is conserved during the learning dynamics, and the distance from the Nash equilibrium is invariant. Follow the Regularized Leader (FTRL) is a class of learning algorithms including the replicator dynamics and also has its conserved quantity, which is the summation of divergences for all the players [\[21,](#page-9-10) [22\]](#page-9-11). To summarize, such complex learning dynamics have been discussed based on their conserved quantity.

In this study, we define memory as an agent's ability to change its action choice depending on the outcome of past games. By definition, this memory allows the agent to make more complex and intelligent decisions. When memory is introduced into a normal-form game, the players can achieve a wider range of strategies as the Nash equilibria (known as Folk theorem [\[23\]](#page-9-12)). Furthermore, memory is also introduced into learning algorithms, such as replicator dynamics [\[24,](#page-9-13) [25,](#page-9-14) [26,](#page-9-15) [27\]](#page-9-16) and Q-learning [\[28,](#page-9-17) [29,](#page-9-18) [30,](#page-9-19) [31,](#page-9-20) [32\]](#page-10-0). Here, since this memory causes feedback from the past, the global dynamics of such learning algorithms become more complex. Indeed, replicator dynamics diverge from the Nash equilibrium under symmetric memory lengths between players [\[26\]](#page-9-15), while converging under asymmetric memory lengths [\[27\]](#page-9-16). Here, KL divergence is no longer useful to capture the global dynamics because it increases or decreases over time. The analysis of the dynamics in with-memory games is limited to the local, linearized stability analysis in the vicinity of Nash equilibrium [\[27\]](#page-9-16). To summarize, since memory crucially complicates learning dynamics, the global behavior of the dynamics is still unexplored.

This study provides the first theoretical analysis of the global behavior of learning in with-memory games. We assume games where their memory structure is simplest and asymmetric; One side adopts a reactive strategy that can memorize the other's previous action [\[33,](#page-10-1) [34,](#page-10-2) [35,](#page-10-3) [24,](#page-9-13) [36,](#page-10-4) [25\]](#page-9-14), while the other has no memory. In order to characterize the global behavior of such with-memory games, we extend KL divergence and prove that such extended divergence increases or decreases with time depending on the reactive strategy (see Fig. [1A](#page-1-0)). Furthermore, we propose a family of Lyapunov functions that characterize the dynamics of reactive strategy (see Fig. [1B](#page-1-0)). These Lyapunov functions show that the with-memory side monotonically learns to exploit the no-memory side. As an application of these functions, we prove that the convergence from arbitrary initial strategies, i.e., global convergence, occurs in matching pennies. We also experimentally confirm that such global convergence is observed in general zero-sum games.

Figure 1: A. Illustration of the global behavior of the conditional divergence, $D(X, y)$. Three trajectories (red, black, and blue) are plotted with the Nash equilibrium (the black star marker). The horizontal and vertical axes show X's strategy (x_1^{st}) and Y's strategy (y_1) in matching pennies (formulated in Fig. [2\)](#page-2-0). This divergence decreases (red: $D < 0$), cycles (black: $D = 0$), or increases (blue: $D > 0$) with time. These three lines are plotted for the different initial strategies, i.e., X and y. B. Illustration of the global behavior of the family of Lyapunov functions, $H(X; \delta)$. The colored line shows a trajectory (from purple to red) of Lyapunov functions H_1 , H_2 , and H_3 , each of which is $H(X; \delta)$ for some specific δ . The gray broken lines are the projections of the black solid line to H_1 - H_2 , H_2 - H_3 , and H_3 - H_1 planes. All of H_1 , H_2 , and H_3 monotonically increase with time.

2 Preliminary

2.1 Settings

First, we formulate two-player normal-form games. We consider two players, denoted as X and Y. X's actions are denoted as $\{a_i\}_{1\leq i\leq m_\chi}$, while Y's are $\{b_j\}_{1\leq j\leq m_\gamma}$. When X and Y choose a_i and b_j , they obtain the payoffs of $u_{ij} \in \mathbb{R}$ and $v_{ij} \in \mathbb{R}$, respectively. Thus, all their possible payoffs are given by the matrices, $U := (u_{ij})_{ij}$ and $V := (v_{ij})_{ij}$. Here, when $V = -U$ holds, the games are called zero-sum. Although our formulation of learning algorithms can be used for general games, this study focuses on zero-sum games.

Figure 2: Illustration of games between reactive and zero-memory strategies. The area surrounded by the magenta dotted line shows the normal-form game. In each round, X chooses action $i = 1$ or 2 in the row, following its strategy, i.e., the probability distribution of $x = (x_1, x_2)$. On the other hand, Y chooses action $j = 1$ or 2 in the column, following its strategy, i.e., the probability distribution of $y = (y_1, y_2)$. Depending on their choices i and j, X receives a payoff u_{ij} , given by a matrix form of $U = (u_{ij})_{i,j} = ((u_{11}, u_{12}), (u_{21}, u_{22}))$. Furthermore, in zero-sum games, Y receives $-u_{ij}$. Especially in the matching pennies, their actions of 1 (2) correspond to the choice of "head" ("tail") of a coin. When their choices match $i = j$, X wins, i.e., $u_{11} = u_{22} = 1$ (the orange blocks). Else when their choices mismatch $i \neq j$, Y wins, i.e., $u_{12} = u_{21} = -1$ (the blue blocks). The area outside of the magenta dotted line shows the difference due to an effect of memory. The gray box shows that X memorizes Y's previous action, represented as $j = 1$ or 2. Thus, X uses a reactive strategy and can choose its action with the conditional probability of x_{1j} and x_{2j} for Y's previous action.

We assume that X can use reactive strategies, i.e., can change its action choice depending on the other's previous action. This reactive strategy is denoted as $X := (x_{i|j})_{1 \le i \le m_X, 1 \le j \le m_Y}$ $\prod_{1 \le j \le m_\Upsilon} \Delta^{m_\Upsilon-1}$, a matrix composed of m_Υ vectors each of which are an element of a $m_\Upsilon - 1$ dimensional simplex. Here, $x_{i|j}$ means the probability that X chooses a_i in the condition when Y's previous action is b_j . Thus, $\sum_i x_{i|j} = 1$ should be satisfied for all j. On the other hand, Y only can use classical mixed strategies and choose its own action without reference to the previous actions. This mixed strategy is denoted as $y = (y_j)_{1 \leq j \leq m_Y} \in \Delta^{m_Y-1}$, a vector which is an element of a $(m_Y - 1)$ -dimensional simplex. Thus, $\Sigma_j y_j = 1$ should be satisfied.

2.2 Stationary state and expected payoff

We now discuss the stationary state and expected payoff of repeated games. Since Y determines its action independent of the outcomes of previous rounds, X's stationary strategy, defined as $x^{\text{st}} := (x_i^{\text{st}})_{1 \le i \le m_\chi}$, is given by $x_i^{\text{st}}(x_i, y) = \sum_j x_{i|j} y_j$. Here, x_i^{st} means the probability that X chooses a_i in the stationary state. Furthermore, the stationary state is described as $P^{\rm st}:=x^{\rm st}\otimes y$ with use of x^{st} and y . Last, X's expected payoff is given by $u^{\text{st}}(x^{\text{st}}, y) := \sum_i \sum_j u_{ij} p_{ij}^{\text{st}} = \sum_i \sum_j u_{ij} x_i^{\text{st}} y_j$.

2.3 Nash equilibrium

We now define the Nash equilibrium in the normal-form game. Here, note that this equilibrium is based on games without memories, where X's strategy does not refer to the past games, i.e., $\bm{x}:=(x_i)_i.$

By using the expected payoff $u^{st}(\bm{x}, \bm{y})$ for games without memories, the Nash equilibrium (\bm{x}^*, \bm{y}^*) is formulated as

$$
\begin{cases}\nx^* = \operatorname{argmax}_{\mathbf{x}} u^{\operatorname{st}}(\mathbf{x}, \mathbf{y}^*) \\
\mathbf{y}^* = \operatorname{argmin}_{\mathbf{y}} u^{\operatorname{st}}(\mathbf{x}^*, \mathbf{y})\n\end{cases} \tag{1}
$$

From the definition, $u^{st}(x, y)$ is the linear function for x and y, and the Nash equilibrium condition is characterized by the gradient of such expected payoffs as

$$
\begin{cases} \partial u^{\text{st}}/\partial x_i = \Sigma_j u_{ij} y_j^* = C & (x_i^* > 0) \\ \partial u^{\text{st}}/\partial x_i = \Sigma_j u_{ij} y_j^* < C & (x_i^* = 0) \end{cases},\tag{2}
$$

$$
\begin{cases} \partial u^{\text{st}}/\partial y_j = \sum_j u_{ij} x_i^* = C & (y_i^* > 0) \\ \partial u^{\text{st}}/\partial y_i = \sum_j u_{ij} x_i^* > C & (y_i^* = 0) \end{cases} \tag{3}
$$

From these conditions, for all i and j such that $x_i^* > 0$ and $y_j^* > 0$ hold, respectively, we obtain

$$
\sum_i u_{ij} x_i^* = \sum_j u_{ij} y_j^* =: u^*.
$$
\n⁽⁴⁾

Let us interpret this equation. First, $\Sigma_i u_{ij} x_i^* = u^*$ means that when X takes its Nash equilibrium strategy, its own payoff is fixed to u^* , independent of Y's strategy. On the other hand, $\Sigma_j u_{ij}^* y_j^* = u^*$ similarly means that Y's Nash equilibrium strategy fixes X's payoff to u^* . In other words, either X or Y takes its Nash equilibrium strategy, their payoffs are fixed. This is the special property in zero-sum games.

2.4 Learning algorithm: replicator dynamics

Let us define the replicator dynamics as a representative learning algorithm. X's and Y's replicator dynamics are formulated as

$$
\dot{x}_{i|j} = x_{i|j} \left(\frac{\mathrm{d}u^{\mathrm{st}}}{\mathrm{d}x_{i|j}} - \Sigma_i x_{i|j} \frac{\mathrm{d}u^{\mathrm{st}}}{\mathrm{d}x_{i|j}} \right),\tag{5}
$$

$$
\dot{y}_j = -y_j \left(\frac{\mathrm{d}u^{\mathrm{st}}}{\mathrm{d}y_j} - \Sigma_j y_j \frac{\mathrm{d}u^{\mathrm{st}}}{\mathrm{d}y_j} \right). \tag{6}
$$

Here, following the theorems in [\[26\]](#page-9-15), X's and Y's replicator dynamics include the gradient for the expected payoff u^{st} . Thus, the update of X's strategy increases its payoff u^{st} , while that of Y's strategy decreases the other's payoff u^{st} .

3 Novel quantities characterizing learning dynamics

This section analyzes the dynamics of Eqs. [\(5\)](#page-3-0) and [\(6\)](#page-3-1). First, we compute in detail the gradient terms, which appear to be complex. Next, as a preliminary, we define positive definite matrices for some special vectors. Based on this definition, we introduce two quantities characterizing the dynamics of Eqs. [\(5\)](#page-3-0) and [\(6\)](#page-3-1): An extended KL divergence and a family of Lyapunov functions.

3.1 Polynomial expressions of learning dynamics

First, the gradient terms in Eqs. [\(5\)](#page-3-0) and [\(6\)](#page-3-1) are computed as

$$
\frac{\mathrm{d}u^{\mathrm{st}}(\boldsymbol{x}^{\mathrm{st}}(\boldsymbol{X},\boldsymbol{y}),\boldsymbol{y})}{\mathrm{d}x_{i|j}} = \frac{\partial x_i^{\mathrm{st}}(\boldsymbol{x}_i,\boldsymbol{y})}{\partial x_{i|j}} \frac{\partial u^{\mathrm{st}}(\boldsymbol{x}^{\mathrm{st}},\boldsymbol{y})}{\partial x_i^{\mathrm{st}}} \tag{7}
$$

$$
= y_j \Sigma_{j'} u_{ij'} y_{j'}, \tag{8}
$$

$$
\frac{du^{st}(\boldsymbol{x}^{st}(\boldsymbol{X},\boldsymbol{y}),\boldsymbol{y})}{dy_j} = \frac{\partial u^{st}(\boldsymbol{x}^{st},\boldsymbol{y})}{\partial y_j} + \Sigma_i \frac{\partial x_i^{st}(\boldsymbol{x}_i,\boldsymbol{y})}{\partial y_j} \frac{\partial u^{st}(\boldsymbol{x}^{st},\boldsymbol{y})}{\partial x_i^{st}}
$$
(9)

$$
= \sum_{i} u_{ij} x_i^{\text{st}} + \sum_{i} x_{i|j} \sum_{j'} u_{ij'} y_{j'}.
$$
\n(10)

Here, we remark that Eqs. [\(5\)](#page-3-0) and [\(6\)](#page-3-1) are nonlinear functions of X and y , which is a feature of learning in with-memory games. Notably, however, these equations are polynomial expressions with X and y . Such polynomial expressions cannot be seen in the games of other memory lengths [\[26,](#page-9-15) [27\]](#page-9-16) but are special in the games between reactive and no memory strategies.

3.2 Positive definiteness for zero-sum vectors

Next, let us introduce a definiteness of matrices. Here, however, this definite matrix is for vectors whose elements are summed to 0, named "zero-sum vectors". In mathematics, zero-sum vector $\delta := (\delta_k)_k$ satisfies $\Sigma_k \delta_k = 0$ but $\delta \neq 0$.

Definition 1 (Positive definiteness for zero-sum vectors). *A square matrix* M *is "positive definite for zero-sum vectors" when for all vectors* $\delta \neq 0$ *such that* $\Sigma_k \delta_k = 0$, $\delta \cdot (M \delta) < 0$ *holds.*

The positive definiteness for zero-sum vectors connects with an ordinary positive definiteness by a simple transformation of a matrix (see Appendix [B](#page-13-0) for details).

3.3 Extended Kullback-Leibler divergence

The first quantity is an extended version of divergence. Before considering the extension, we introduce the classical version of divergence D_c , which is the function of X's mixed strategies $(\boldsymbol{x} := (x_i)_{1 \leq i \leq m_\mathsf{X}} \in \Delta^{m_\mathsf{X}-1})$ and Y's mixed strategies (\boldsymbol{y}) as

$$
D_{\mathrm{c}}(\boldsymbol{x},\boldsymbol{y}) := D_{\mathrm{KL}}(\boldsymbol{x}^*\|\boldsymbol{x}) + D_{\mathrm{KL}}(\boldsymbol{y}^*\|\boldsymbol{y}),\tag{11}
$$

$$
D_{\mathrm{KL}}(\boldsymbol{p}^* \| \boldsymbol{p}) := \boldsymbol{p}^* \cdot \log \boldsymbol{p}^* - \boldsymbol{p}^* \cdot \log \boldsymbol{p}.\tag{12}
$$

We now give an intuitive interpretation of this quantity. First, $D_{KL}(p^*||p)$ is the KL divergence, meaning the distance from the reference point p^* to the target point p . Thus, $D_c(x, y)$ means the total distance from the Nash equilibrium (x^*, y^*) to the current state (x, y) .

Let us extend the classical divergence to the case of this study, where X refers to the previous action of the other and can use reactive strategies X. This extended divergence, i.e., $D(X, y)$, is named the "conditional-sum" divergence, formulated as

$$
D(\boldsymbol{X}, \boldsymbol{y}) := \Sigma_j D_{\mathrm{KL}}(\boldsymbol{x}^* \| \boldsymbol{x}_j) + D_{\mathrm{KL}}(\boldsymbol{y}^* \| \boldsymbol{y}).
$$
\n(13)

We now remark the difference between $D(X, y)$ and $D_c(x, y)$. Recall that X's reactive strategy is defined as $(x_j)_{1 \le j \le m_\gamma}$, which shows how to choose its action with the condition that Y chose b_j in the previous round. Hence, $D(X, y)$ represents the summation of KL divergence from x^* to x_j for all the conditions of j . Here, we also remark that when the reactive strategy does not use memory, i.e., $x_j = x$ for all j, this conditional-sum divergence also captures the behavior of the classical divergence (see Appendix [C](#page-14-0) for details).

This conditional-sum divergence satisfies the following theorem (see Appendix [A.1](#page-11-0) for its full proof).

Theorem 1 (Monotonic decrease of D for positive definite X^TU). *If* X^TU *is positive definite for* zero-sum vector, $D^{\dagger}(\bm{X}; {\mathrm d} \bm{y}) := \dot{D}(\bm{X}, \bm{y}) < 0$ for all ${\mathrm d} \bm{y} \neq \bm{0}$.

Proof Sketch. We calculation $D(X, y)$ in practice. In the calculation, the contribution of X's gradient (Eq. [\(8\)](#page-3-2)) cancels out the contribution of the first term of the gradient of Y (Eq. [\(10\)](#page-3-3)). (Here, we remark that the same canceling out also occurs in the calculation for the conservation of the classical divergence $D_c(x, y)$ in games without memory.) However, the contribution of the second term of Eq. [\(10\)](#page-3-3) is special in games of a reactive strategy. By using the constant payoff condition in the Nash equilibrium (Eqs. [\(4\)](#page-3-4)), we obtain

$$
\dot{D}(\boldsymbol{X}, \boldsymbol{y}) = -\mathrm{d}\boldsymbol{y}^{\mathrm{T}} \boldsymbol{X}^{\mathrm{T}} \boldsymbol{U} \mathrm{d}\boldsymbol{y} \ (=:D^{\dagger}(\boldsymbol{X}; \mathrm{d}\boldsymbol{y})),\tag{14}
$$

where we defined $dy := y - y^*$, which means the difference from Y's equilibrium strategy and is a zero-sum vector. Thus, when $\bar{X}^{\mathrm{T}}U$ is positive definite for zero-sum vectors, $\mathrm{d}y^{\mathrm{T}}X^{\mathrm{T}}U\mathrm{d}y$ is always positive, leading to $D^{\dagger}(\bm{X}; \mathrm{d}\bm{y}) < 0$ for all $\mathrm{d}\bm{y} \neq \bm{0}$. \Box

If $X^{\mathrm{T}}U$ are always positive definite for zero-sum vectors, Thm. [1](#page-4-0) gives the global behavior of learning dynamics as below.

Corollary 1 (Convergence of Y's strategy). *If* X^TU *continues to be positive definite for zero-sum* $vectors, y = y^*$ *holds after sufficiently long time passes.*

Proof. This corollary is straightforwardly proved by Thm. [1.](#page-4-0) If X^TU is positive definite and if $y \neq y^*$, $D^{\dagger}(X; dy) < 0$ holds. Here, $D(X, y)$ has its lower bound $(\min_{X, y} D(X, y) = 0)$. Thus, $D^{\dagger}(\boldsymbol{X};\mathrm{d}\boldsymbol{y})=0 \Leftrightarrow \boldsymbol{y}=\boldsymbol{y}^*$ should be satisfied eventually. \Box

3.4 Family of Lyapunov functions

Furthermore, we introduce a Lyapunov function, which characterizes the learning dynamics of X's reactive strategy. Based on an arbitrary zero-sum vector $\bm{\delta}:=(\delta_i)_{1\leq i\leq m_\textsf{X}},$ this function is defined as

$$
H(\mathbf{X};\delta) := \delta^{\mathrm{T}} U \log \mathbf{X}^{\mathrm{T}} \delta. \tag{15}
$$

The following theorem holds for this Lyapunov function (see Appendix [A.2](#page-12-0) for its full proof).

Theorem 2 (Monotonic increase of H). *For all* δ *such that* $\Sigma_i \delta_i = 0$, $H(X; \delta)$ *monotonically increases with time.*

Proof Sketch. By using Eq. [\(4\)](#page-3-4), we calculate

$$
\dot{H}(\mathbf{X};\delta) = (\delta^{\mathrm{T}} U \mathrm{d} \mathbf{y})^{2} (= :H^{\dagger}(\mathbf{y};\delta)).
$$
\n(16)

 \Box

This means that $H^{\dagger}(\boldsymbol{y};\boldsymbol{\delta}) \geq 0$ for all $\boldsymbol{\delta}$.

Let us interpret the function of $H(X; \delta)$. For simplicity, we consider a simple class of zero-sum vectors, $\delta = e_i - e_{i'}$ for some i and $i' \neq i$, where e_i denotes the unit vector for i-th element. Then, we calculate the function as $H(\bm{X}; \bm{e}_i - \bm{e}_{i'}) = \bm{u}_i \cdot \log \bm{x}_i - \bm{u}_i \cdot \log \bm{x}_{i'} - \bm{u}_{i'} \cdot \log \bm{x}_i + \bm{u}_{i'} \cdot \log \bm{x}_{i'}.$ The monotonic increase of $H(\bm{X};\bm{e}_i-\bm{e}_{i'})$ shows that $\log \bm{x}_i$ becomes correlated to \bm{u}_i , and $\log \bm{x}_{i'}$ to $u_{i'}$. Here, $x_i = (x_{i|j})_j$ represents the vector of the probabilities to choose *i*-th action in the condition that the other chose j-th action in the previous round. $u_i = (u_{ij})_j$ also represents the vector of the payoffs for action i. Thus, the correlation between $\log x_i$ and $\overline{u_i}$ means that the larger its reward u_{ij} is, the more frequently X returns its *i*-th action against the other's *j*-th action, in other words, X becomes more exploitative.

This theorem characterizes the state of learning dynamics after a sufficiently long time passes, as follows.

Corollary 2 (Convergence of Y's strategy or divergence of H). *After sufficiently long time passes,* $y = y^*$ *holds or* $H(\bar{X}; \delta)$ diverges to infinity for all $\delta \neq 0$ such that $\Sigma_i \check{\delta_i} = 0$.

Proof. Since $H(X; \delta) \ge 0$ always holds for all δ , either $H(X; \delta) = 0$ or $H(X; \delta) > 0$ holds in the final state. The former is equivalent to $y = y^*$, while the latter is equivalent to $H(\bm{X}; \bm{\delta}) \to \infty$.

4 Global behavior of learning dynamics

Now, Cors. [1](#page-4-1) and [2](#page-5-0) are utilized to capture the global behavior of the learning dynamics. One of the global behaviors seen in the zero-sum games is the global convergence to the Nash equilibrium independent of X's and Y's initial strategies. First, Cor. [1](#page-4-1) guarantees that after a sufficiently long time passes, either Y's strategy converges to the equilibrium or $H(X; \delta)$ diverges. Since the former trivially leads to such global convergence, we now consider the latter condition, i.e., $H(X; \delta) =$ $\delta^T U \log X^T \delta \to \infty$. Here, $H(X; \delta) = \delta^T U \log X^T \delta \to \infty$ naively connects to $-D^{\dagger}(X, dy) =$ $dy^T X^T U dy > 0$ (explained in the next paragraph), meaning that $X^T U$ is positive definite for zero-sum vectors. If so, Cor. [2](#page-5-0) becomes applicable, and either Y's strategy converges to the Nash equilibrium.

We explain the connection between $H(X; \delta) = \delta^{\mathrm{T}} U \log X^{\mathrm{T}}$ and $-D^{\dagger}(X, dy) = dy^{\mathrm{T}} X^{\mathrm{T}} U dy >$ 0. Here, $H(\bm{X};\bm{\delta})$ and $-D^{\dagger}(\bm{X},\mathrm{d}\bm{y})$ have two commonalities. The first one is that $\bm{\delta}$ and $\mathrm{d}\bm{y}$ are both zero-sum vectors in common. The other one is that both H and $-D^{\dagger}$ are given by the product of X's strategy X and the payoff matrix U. This means that the more X is correlated to U, the larger both H and $-D^{\dagger}$ are. Here, however, we also remark two differences of H from $-D^{\dagger}$; H considers the logarithm in X's strategy X and the reversed order in the product of X and U .

In the following, we prove such global convergence of either player's strategy in the matching pennies, the simplest game equipped with a full-support equilibrium. We also experimentally show global convergence in more complicated games than the matching pennies later.

4.1 Example: Matching-pennies

Let us define the matching-pennies game (see Fig. [2](#page-2-0) for the illustration of its payoff matrix). This game considers the action numbers of $m_X = m_Y = 2$ and the payoff matrix of $U =$

 $((u_{11}, u_{12}), (u_{21}, u_{22})) = ((+1, -1), (-1, +1))$. The Nash equilibrium of this game is only $x^* = y^* = (1/2, 1/2)$. This game is the simplest example of games that have a full-support equilibrium. In addition, it has been known that the replicator dynamics in games without memories cycle around the Nash equilibrium and cannot reach the equilibrium. Nevertheless, the learning dynamics in games of a reactive strategy achieve the convergence to the Nash equilibrium for Y's strategy. The following theorem holds (see Appendix [A.3](#page-12-1) for its full proof).

Theorem 3 (Global convergence in matching-pennies). *In the matching-pennies (*U = $((+1,-1),(-1,+1))$), Y's strategy converges to the Nash equilibrium ($y = y^*$), independent *of both the players' initial strategies.*

Proof Sketch. First, we define a special notation in two-action games; $(x_{1|j}, x_{2|j}) =: (x_j, 1 - x_j)$ for $j = 1, 2$. We also define the logit function of x_j as $q_j := \log x_j - \log(1 - x_j)$. Here, note that $q_1 > q_2 \Leftrightarrow x_1 > x_2$. By the direct calculation, we prove $H(\mathbf{X}; \boldsymbol{\delta}) \propto q_1 - q_2$. Thus, Cor. [2](#page-5-0) shows that after a sufficiently long time passes, $y = y^*$ holds or $H(X; \delta)$ diverges, in other words, $H(X; \delta) > 0 \Leftrightarrow q_1 > q_2 \Leftrightarrow x_1 > x_2$ holds. In the former case, Y obviously reaches the Nash equilibrium strategy, so we now assume the latter case $(x_1 > x_2 \text{ holds})$. We can also prove that $x_1 > x_2$ is equivalent to the positive definiteness of $X^{\mathrm{T}}U$. Because $X^{\mathrm{T}}U$ continues to hold after a sufficiently long time passes, $y = y^*$ holds after a further long time passes, meaning that Y's strategy converges to its equilibrium. \Box

5 Experimental supports for theory

This section experimentally shows the global convergence to the Nash equilibrium. First, we visualize the mechanism of global convergence in matching pennies. Following Thm. [2,](#page-5-1) $H(\boldsymbol{X};\boldsymbol{\delta})$ increases with time, and $X^T U$ becomes positive definite. After that, following Thm. [1,](#page-4-0) the distance from the Nash equilibrium $D(X, y)$ decreases with time. Next, we also demonstrate global convergence for several examples of more advanced games than matching pennies. We have observed the global convergence in all zero-sum games as far as we experimented, even though we could not prove it in general zero-sum games. All the simulations in this section are performed following the Runge-Kutta fourth-order method of Eqs. [\(5\)](#page-3-0) and [\(6\)](#page-3-1) with a sufficiently small step size of 5×10^{-2} .

Figure 3: A. Trajectories of q_1 and q_2 . The rainbow contour plot indicates the value of $q_1 - q_2$. All the trajectories monotonically increase $q_1 - q_2$ with time and converge in the area of $q_1 > q_2$ in their final states. B. Trajectories of the learning dynamics. The black broken line corresponds to the region of Nash equilibria, $x^{st} = y = (1/2, 1/2)$. Each colored line shows a trajectory of the learning dynamics. First, the circle markers show the initial states. Following the blue lines, the trajectories diverge from the Nash equilibria $(D(X, y))$ increases with time). However, the trajectories stop to diverge and switch to converge to the Nash equilibria $(D(X, y))$ decreases), following the red lines. The star markers are the final states and correspond to one of the Nash equilibria.

5.1 Visualization of global convergence mechanism

Let us see that learning dynamics in the matching pennies converge to the equilibrium following the mechanism we provided. To analyze the matching pennies in which both the players have two actions, we use the notation of $(x_{1|j}, x_{2|j}) =: (x_j, 1 - x_j), q_j := \log x_j - \log(1 - x_j)$, and $(y_1, y_2) =: (y, 1 - y).$

Fig. [3A](#page-6-0) shows the dynamics of q_1 and q_2 . Here, the colors indicate the contour plot for q_1 – q_2 , showing that $H(\bm{X}; \bm{\delta}) \propto q_1 - q_2$ monotonically increases with time and thus Thm. [2](#page-5-1) holds. Furthermore, we also see that X's strategy reaches the region of $q_1 > q_2 \Leftrightarrow x_1 > x_2$ after a sufficiently long time passes. In the region, X^TU is positive definite, and thus Thm. [1](#page-4-0) is applicable after sufficiently long time passes.

Next, Fig. [3B](#page-6-0) plots the global behavior of the learning dynamics, which is described by the three parameters of x_1, x_2 , and y . The gray line shows the region that corresponds to the Nash equilibrium, i.e., $x^{st}(X, y) = y = (1/2, 1/2)$. Furthermore, the colored lines show example trajectories of the learning dynamics. The blue part of the line shows that $D(X, y)$ increases at the beginning of the learning dynamics. This part is in the region of $x_1 < x_2$, following Thm. [1.](#page-4-0) After that, the red part shows that $D(X, y)$ decreases, and the learning dynamics converge to the equilibrium. This part is in the region of $x_1 > x_2$, following Thm. [1.](#page-4-0)

5.2 Global convergence in more advanced zero-sum games

The global convergence is observed in more advanced zero-sum games than the matching pennies. Fig. [4](#page-7-0) shows three representative examples. In all the examples, we basically assume that there is no dominant (pure-)strategy, where X's $i = \sigma(j)$ -th action is advantageous for Y's j-th action, and vise versa, for the permutation σ (i.e., $\sigma(1) = 2$, $\sigma(2) = 3$, $\sigma(3) = 4$, and $\sigma(4) = 1$). In other words, the payoff matrices satisfy $u_{i\sigma(i)} = 2$ and $u_{\sigma(j)j} = -2$. Three panels take different values in the other elements of their payoff matrix.

Figure 4: Global convergence in three advanced games. In all the games, we consider the rock-paperscissors-like game extended to the four-action case, where the second, third, fourth, and first actions win the other's first, second, third, and fourth actions, respectively. The winner receives the payoff of 2 (the orange blocks in the matrices for the winning of X), while the loser sends the payoff of 2 (the blue blocks). We now introduce three variants for the other blocks in the payoff matrix. A. We set each of the other blocks by random numbers in $[-1, 1]$ (the gray blocks). Then, Y's strategy converges to the unique Nash equilibrium (the red star marker) independent of its initial state (the blue circle markers). B. We set each of the other blocks by 0, where the payoff matrix degenerates. Y's strategy converges to one of the Nash equilibria (the line consisting of the red star markers) depending on its initial state. C. Only the block for the interaction between an action is set to -1 , and the others are 0. If so, X's strategy converges to the unique Nash equilibrium (the orange star markers) independent of its initial state (the green circle markers). Instead, Y's strategies do not converge.

First, Fig. [4A](#page-7-0) shows the case where the other elements of the payoff matrix are random numbers following the uniform distribution of $[-1, 1]$. In this case, the payoff matrix does not degenerate. In other words, X has no two different strategies that generate equal payoffs for all Y's strategies. In mathematics, there exists no $\boldsymbol{a} = (a_i)_{1 \leq i \leq m_\chi} \in \mathbb{R}^{m_\chi}$ such that $\sum_i a_i \boldsymbol{u}_i = \boldsymbol{0}$. We observe that the single Nash equilibrium exists, and Y's strategy always converges to its equilibrium independent of X's and Y's initial strategies.

Second, Fig. [4B](#page-7-0) shows the case where all the other elements take 0. In this case, the payoff matrix degenerates; Indeed, $u_i + u_{i'} = 0$ holds for $(i, i') = (1, 3)$, $(2, 4)$. Continuous Nash equilibria are

seen there, $x^* = r_X(0, 1/2, 0, 1/2) + (1 - r_X)(1/2, 0, 1/2, 0)$ and $y^* = r_Y(0, 1/2, 0, 1/2) + (1 - r_Y)(1/2, 0, 1/2)$ r_Y)(1/2, 0, 1/2, 0) for all $0 \le r_X$ and $0 \le r_Y \le 1$. We observe that Y's strategy converges to one of its equilibria depending on X's and Y's initial strategies.

Third, Fig. [4C](#page-7-0) shows the case where the other elements take 0 in principle except for $u_{11} = -1$. In this case, the payoff matrix degenerates because $u_i + u_{i'} = 0$ holds for $(i, i') = (2, 4)$. Furthermore, the only Nash equilibrium exists on the boundary of strategy spaces, $x^* = (0, 1/2, 0, 1/2)$, and $y^* = (1/2, 0, 1/2, 0)$. As far as we experiment, when the payoff matrix degenerates and only the boundary Nash equilibrium exists, Y's strategy does not converge to its equilibrium exceptionally. Nevertheless, we observe that X's strategy converges to its equilibrium instead of Y's.

6 Conclusion

This study considered the simplest situation of memory asymmetry between two players; only player X memorizes the other's previous action, while player Y cannot. We formulated their learning dynamics based on the replicator dynamics. Although the existence of memory complicates the dynamics, we captured the global behavior of the learning dynamics by introducing two new quantities. One is the conditional-sum divergence, which is an extension of the previous divergence to the case of games of a reactive strategy. The time change of this extended divergence is characterized by the definiteness of X^TU . The other is a family of Lyapunov functions, characterized by U log X^T . This is proved to monotonically increase with time. As a valid application of these two quantities, we proved the global convergence in the learning dynamics in the simplest game equipped with an interior Nash equilibrium. We further experimentally observed the global convergence in more advanced zero-sum games. It is still a conjecture whether the learning dynamics in games of a reactive strategy converge to the Nash equilibrium even in general zero-sum payoff matrices. This study provides novel and valid indicators to analyze dynamics in learning in games.

References

- [1] Drew Fudenberg and David K Levine. *The theory of learning in games*, volume 2. MIT press, 1998.
- [2] John F Nash Jr. Equilibrium points in n-person games. *Proceedings of the National Academy of Sciences*, 36(1):48–49, 1950.
- [3] Karl Tuyls, Pieter Jan'T Hoen, and Bram Vanschoenwinkel. An evolutionary dynamical analysis of multi-agent learning in iterated games. In *AAMAS*, volume 12, pages 115–153. Springer, 2006.
- [4] Lucian Busoniu, Robert Babuska, and Bart De Schutter. A comprehensive survey of multiagent reinforcement learning. *IEEE Transactions on Systems, Man, and Cybernetics, Part C (Applications and Reviews)*, 38(2):156–172, 2008.
- [5] Karl Tuyls and Gerhard Weiss. Multiagent learning: Basics, challenges, and prospects. *Ai Magazine*, 33(3):41–41, 2012.
- [6] Daan Bloembergen, Karl Tuyls, Daniel Hennes, and Michael Kaisers. Evolutionary dynamics of multi-agent learning: A survey. *Journal of Artificial Intelligence Research*, 53:659–697, 2015.
- [7] Peter D Taylor and Leo B Jonker. Evolutionary stable strategies and game dynamics. *Mathematical biosciences*, 40(1-2):145–156, 1978.
- [8] Daniel Friedman. Evolutionary games in economics. *Econometrica: Journal of the Econometric Society*, pages 637–666, 1991.
- [9] Tilman Börgers and Rajiv Sarin. Learning through reinforcement and replicator dynamics. *Journal of Economic Theory*, 77(1):1–14, 1997.
- [10] Josef Hofbauer, Karl Sigmund, et al. *Evolutionary games and population dynamics*. Cambridge university press, 1998.
- [11] Martin A Nowak and Karl Sigmund. Evolutionary dynamics of biological games. *Science*, 303(5659):793–799, 2004.
- [12] Sanjeev Arora, Elad Hazan, and Satyen Kale. The multiplicative weights update method: a meta-algorithm and applications. *Theory of computing*, 8(1):121–164, 2012.
- [13] James P Bailey and Georgios Piliouras. Multiplicative weights update in zero-sum games. In *EC*, pages 321–338, 2018.
- [14] Satinder Singh, Michael J Kearns, and Yishay Mansour. Nash convergence of gradient dynamics in general-sum games. In *UAI*, pages 541–548, 2000.
- [15] Michael Bowling and Manuela Veloso. Multiagent learning using a variable learning rate. *Artificial Intelligence*, 136(2):215–250, 2002.
- [16] Martin Zinkevich. Online convex programming and generalized infinitesimal gradient ascent. In *ICML*, pages 928–936, 2003.
- [17] Michael Bowling. Convergence and no-regret in multiagent learning. In *NeurIPS*, pages 209–216, 2004.
- [18] Christopher JCH Watkins and Peter Dayan. Q-learning. *Machine learning*, 8(3):279–292, 1992.
- [19] Aamal Abbas Hussain, Francesco Belardinelli, and Georgios Piliouras. Asymptotic convergence and performance of multi-agent q-learning dynamics. In *AAMAS*, pages 1578–1586, 2023.
- [20] Aamal Hussain, Francesco Belardinelli, and Georgios Piliouras. Beyond strict competition: approximate convergence of multi-agent q-learning dynamics. In *IJCAI*, pages 135–143, 2023.
- [21] Panayotis Mertikopoulos and William H Sandholm. Learning in games via reinforcement and regularization. *Mathematics of Operations Research*, 41(4):1297–1324, 2016.
- [22] Panayotis Mertikopoulos, Christos Papadimitriou, and Georgios Piliouras. Cycles in adversarial regularized learning. In *SODA*, pages 2703–2717, 2018.
- [23] Drew Fudenberg and Eric Maskin. The folk theorem in repeated games with discounting or with incomplete information. In *A long-run collaboration on long-run games*, pages 209–230. World Scientific, 2009.
- [24] Yuma Fujimoto and Kunihiko Kaneko. Emergence of exploitation as symmetry breaking in iterated prisoner's dilemma. *Physical Review Research*, 1(3):033077, 2019.
- [25] Yuma Fujimoto and Kunihiko Kaneko. Exploitation by asymmetry of information reference in coevolutionary learning in prisoner's dilemma game. *Journal of Physics: Complexity*, 2(4):045007, 2021.
- [26] Yuma Fujimoto, Kaito Ariu, and Kenshi Abe. Learning in multi-memory games triggers complex dynamics diverging from nash equilibrium. In *IJCAI*, 2023.
- [27] Yuma Fujimoto, Kaito Ariu, and Kenshi Abe. Memory asymmetry creates heteroclinic orbits to nash equilibrium in learning in zero-sum games. In *AAAI*, 2024.
- [28] Naoki Masuda and Hisashi Ohtsuki. A theoretical analysis of temporal difference learning in the iterated prisoner's dilemma game. *Bulletin of mathematical biology*, 71:1818–1850, 2009.
- [29] Wolfram Barfuss. Reinforcement learning dynamics in the infinite memory limit. In *AAMAS*, pages 1768–1770, 2020.
- [30] Yuki Usui and Masahiko Ueda. Symmetric equilibrium of multi-agent reinforcement learning in repeated prisoner's dilemma. *Applied Mathematics and Computation*, 409:126370, 2021.
- [31] Janusz M Meylahn, Lars Janssen, et al. Limiting dynamics for q-learning with memory one in symmetric two-player, two-action games. *Complexity*, 2022, 2022.
- [32] Masahiko Ueda. Memory-two strategies forming symmetric mutual reinforcement learning equilibrium in repeated prisoners' dilemma game. *Applied Mathematics and Computation*, 444:127819, 2023.
- [33] Martin Nowak. Stochastic strategies in the prisoner's dilemma. *Theoretical population biology*, 38(1):93–112, 1990.
- [34] Hisashi Ohtsuki. Reactive strategies in indirect reciprocity. *Journal of Theoretical Biology*, 227(3):299–314, 2004.
- [35] Seung Ki Baek, Hyeong-Chai Jeong, Christian Hilbe, and Martin A Nowak. Comparing reactive and memory-one strategies of direct reciprocity. *Scientific reports*, 6(1):25676, 2016.
- [36] Laura Schmid, Christian Hilbe, Krishnendu Chatterjee, and Martin A Nowak. Direct reciprocity between individuals that use different strategy spaces. *PLoS Computational Biology*, 18(6):e1010149, 2022.

Appendix

A Proofs

A.1 Proof of Thm. [1](#page-4-0)

Proof. The dynamics of $D(X, y)$ are calculated as

$$
\dot{D}(\boldsymbol{X}, \boldsymbol{y}) = -\Sigma_{i}\Sigma_{j}x_{i}^{*}\frac{\dot{x}_{ij}}{x_{ij}} - \Sigma_{j}y_{j}^{*}\frac{\partial y}{\partial y_{j}} \\
= -\Sigma_{i}\Sigma_{j}x_{i}^{*}\left(\frac{\partial u^{st}}{\partial x_{ij}} - \Sigma_{i}x_{ij}\frac{\partial u^{st}}{\partial x_{ij}}\right) - \Sigma_{j}y_{j}^{*}\left(\frac{\partial u^{st}}{\partial y_{j}} - \Sigma_{j}y_{j}\frac{\partial u^{st}}{\partial y_{j}}\right) \\
= -\Sigma_{i}\Sigma_{j}x_{i}^{*}y_{j}\left(\Sigma_{i}u_{ij}x_{j}y_{j} - \Sigma_{i}x_{ij}\Sigma_{j}u_{ij}xy_{j} + \Sigma_{j}y_{j}^{*}\left(\Sigma_{i}u_{ij}x_{i}^{*t} - \Sigma_{j}y_{j}\Sigma_{i}u_{ij}x_{i}^{*t}\right)\right) \\
+ \Sigma_{j}y_{j}^{*}\left(\Sigma_{i}x_{ij}\Sigma_{j}u_{ij}xy_{j} - \Sigma_{j}y_{j}\Sigma_{i}x_{ij}\Sigma_{j}u_{ij}xy_{j}\right) \\
= -\frac{\Sigma_{i}\Sigma_{j}\Sigma_{j}x_{i}^{*}y_{j}u_{ij}xy_{j} - \Sigma_{i}\Sigma_{j}\Sigma_{j}x_{ij}\Sigma_{j}u_{ij}xy_{j}y_{j}}{\Sigma_{i}\Sigma_{j}y_{ij}x_{ij}x_{ij}x_{j}} - \frac{\Sigma_{i}\Sigma_{j}y_{j}y_{ij}x_{ij}u_{ij}xy_{j}}{\Sigma_{i}\Sigma_{j}y_{ij}x_{ij}u_{ij}xy_{j}} \\
= -u^{*} + u^{st} + u^{*} - u^{st} - d y^{T} \boldsymbol{X}^{T} \boldsymbol{U} d y \\
= -dy^{T} \boldsymbol{X}^{T} \boldsymbol{U} d y \\
= -dy^{T} \boldsymbol{X}^{T} \boldsymbol{U} d y \\
= -dy^{T} \boldsymbol{X}^{T} \boldsymbol{U} d y \\
= -\frac{1}{2}y^{T} \boldsymbol{X}^{T} \boldsymbol{U} d y \\
= -\frac{1}{2}y^{T} \boldsymbol{X}^{T} \boldsymbol{U} d y \\
= -\frac{1}{2}y^{T} \boldsymbol{X}^{T} \boldsymbol{U} d y \\
= \Sigma_{j}y_{j}y_{j}y_{j}x
$$

Here, note that $dy = y - y^*$ is zero-sum vector. Thus, if $X^T U$ is positive definite for zero-sum vectors, $D^{\dagger}(\bm{X};\mathrm{d}\bm{y}) \leq 0$ holds independent of $\mathrm{d}\bm{y}$, meaning that $D(\hat{\bm{X}},\bm{y})$ monotonically decreases with time.

A.2 Proof of Thm. [2](#page-5-1)

Proof. By Eqs. [\(5\)](#page-3-0) and [\(8\)](#page-3-2), we obtain

$$
\dot{H}(\boldsymbol{X}, \boldsymbol{y}) = \sum_{i} \sum_{i'} \sum_{j} \delta_{i} u_{ij} \frac{\dot{x}_{i' \mid j}}{x_{i' \mid j}} \delta_{i'}
$$
\n
$$
= \sum_{i} \sum_{i'} \sum_{j} \delta_{i} u_{ij} \left(\frac{\partial u^{st}}{\partial x_{i \mid j}} - \sum_{i} x_{i \mid j} \frac{\partial u^{st}}{\partial x_{i \mid j}} \right) \delta_{i'}
$$
\n
$$
= \sum_{i} \sum_{i'} \sum_{j} \delta_{i} u_{ij} y_{j} \delta_{i'} (\sum_{j'} u_{i'j'} y_{j'} - \sum_{i'} x_{i'' \mid j} \sum_{j'} u_{i''j'} y_{j'})
$$
\n
$$
= \sum_{i} \sum_{i'} \sum_{j} \sum_{j'} \delta_{i} u_{ij} y_{j} \delta_{i'} u_{i'j'} y_{j'} - \sum_{i} \sum_{i'} \sum_{j'} \sum_{j'} \frac{\left(\sum_{i'} \delta_{i'} \right)}{\delta_{i} u_{i j} y_{j} x_{i'' \mid j} u_{i''j'} y_{j'}}
$$
\n
$$
= \sum_{i} \sum_{i'} \sum_{j} \sum_{j'} \delta_{i} u_{ij} y_{j} \delta_{i'} u_{i'j'} y_{j'}
$$
\n
$$
= (\sum_{i} \sum_{j} \delta_{i} u_{ij} y_{j})^{2}
$$
\n
$$
= (\sum_{i} \sum_{j} \delta_{i} u_{ij} dy_{j} + \sum_{i} \sum_{j} \delta_{i} u_{ij} y_{j'}^{*})^{2}
$$
\n
$$
= (\sigma^{\mathrm{T}} U \, \mathrm{d} \boldsymbol{y})^{2}
$$
\n
$$
(= : H^{\dagger}(\boldsymbol{y}; \boldsymbol{\delta})). \tag{23}
$$

This equation shows that $H^\dagger({\bm y};{\bm \delta})\geq 0$ always holds. Furthermore, $H^\dagger({\bm y};{\bm \delta})=0$ for all ${\bm \delta}$ is satisfied if and only if $dy = 0 \Leftrightarrow y = y^*$. \Box

A.3 Proof of Thm. [3](#page-6-1)

Proof. Now, by using the special properties of two-action games, let us simplify the notation of $\boldsymbol{X} := \{x_{i|j}\}_{i,j}, \boldsymbol{y} := \{y_j\}_j$, and $\boldsymbol{\delta} := \{\delta_i\}_i$. From the definition, $\Sigma_i x_{i|j} = 1 \Leftrightarrow x_{1|j} + x_{2|j} = 1$ for all $j \in \{1,2\}$ holds, and thus we simply denote it by $x_{1|j} =: x_j$ and $x_{2|j} = 1 - x_j$. In a similar manner, because $y_1 + y_2 = 1$ holds, we simply denote it by $y_1 := y$ and $y_2 = 1 - y$. Furthermore, since $\Sigma_i \delta_i = 0$ holds, we simply denote it by $\delta_1 =: \delta$ and $\delta_2 = -\delta$. In two-action games, we can write

$$
\delta^{\mathrm{T}} = (\delta_1 \quad \delta_2) = (\delta \quad -\delta), \tag{24}
$$

$$
\log \mathbf{X} = \begin{pmatrix} \log x_{1|1} & \log x_{1|2} \\ \log x_{2|1} & \log x_{2|2} \end{pmatrix} = \begin{pmatrix} \log x_1 & \log x_2 \\ \log(1 - x_1) & \log(1 - x_2) \end{pmatrix}.
$$
 (25)

Using these notations, we calculate $H(\mathbf{X}; \delta)$ as

$$
H(\mathbf{X}; \delta) = \delta^{\mathrm{T}} \mathbf{U} \log \mathbf{X}^{\mathrm{T}} \delta
$$

= $(\delta - \delta) \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} \log x_1 & \log(1 - x_1) \\ \log x_2 & \log(1 - x_2) \end{pmatrix} \begin{pmatrix} \delta \\ -\delta \end{pmatrix}$
= $2\delta^2 \left(\log \frac{x_1}{1 - x_1} + \log \frac{x_2}{1 - x_2} \right)$
= $2\delta^2 (q_1 - q_2).$ (26)

Here, in the final line, we used $q_j := \log x_j - \log(1 - x_j) \Leftrightarrow x_j = \exp(q_j)/(1 + \exp(q_j))$. The dynamics of $H(X; \delta)$ is also calculated as

$$
H^{\dagger}(\mathbf{y};\delta) = (\delta^{\mathrm{T}} U \mathrm{d}\mathbf{y})^2
$$

=
$$
\begin{pmatrix} (\delta - \delta) \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} y - y^* \\ -y + y^* \end{pmatrix} \end{pmatrix}^2
$$

=
$$
16\delta^2(y - y^*)^2.
$$
 (27)

This equation shows that for $\delta \neq 0, H^{\dagger}(\bm{y};\bm{\delta})=0$ holds if and only if $y=y^*$. Thus, after sufficiently long time passes, $y = y^*$ holds or $H(\tilde{X}; \delta)$ diverges to infinity. If so, $H(\tilde{X}; \delta) > 0$ is satisfied and is equivalent to $q_1 > q_2$, which is also equivalent to $x_1 > x_2$ from the definition. If $x_1 > x_2$ holds, $D^{\dagger}(\mathbf{X};\mathrm{d}\mathbf{y}) < 0$ is satisfied because we can derive

$$
D^{\dagger}(\mathbf{X};\mathrm{d}\mathbf{y}) = -\mathrm{d}\mathbf{y}^{\mathrm{T}}\mathbf{X}^{\mathrm{T}}U\mathrm{d}\mathbf{y}
$$

= $-(y - y^* - y + y^*)\begin{pmatrix} x_1 & 1 - x_1 \\ x_2 & 1 - x_2 \end{pmatrix} \begin{pmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{pmatrix} \begin{pmatrix} y - y^* \\ -y + y^* \end{pmatrix}$
= $-(y - y^*)^2(x_1 - x_2)(u_{11} - u_{12} - u_{21} + u_{22}).$ (28)

Since $D(X, y)$ has its lower bound, $D^{\dagger}(X; dy) = 0 \Leftrightarrow y = y^*$ holds after sufficiently long time passes. Thus, it was proved that Y's strategy globally converges to its Nash equilibrium strategy. \square

B Connection with ordinary definiteness

This study considered positive definiteness with some constraint, i.e., for zero-sum vectors. However, this constrained positive definiteness is related to the ordinarily positive definiteness without constraint, introduced as follows (describing "tilde" for distinction).

Definition 2 (Positive definiteness). A square matrix M is positive definite when for all vectors $\tilde{\boldsymbol{\delta}} \neq \boldsymbol{0}, \, \tilde{\boldsymbol{\delta}}\cdot(\tilde{\boldsymbol{M}}\tilde{\boldsymbol{\delta}}) > 0.$

Now, we show how the positive definiteness for zero-sum vectors is tied to the ordinary positive definiteness. First, freely choose and fix $\hat{k} \in \{1, \dots, m\}$. From any $m \times m$ matrix $\mathbf{M} = (m_{kk'})_{k,k'}$, we define a $(m-1) \times (m-1)$ matrix $\tilde{M} := (\tilde{m}_{kk'})_{k \neq \hat{k}, k' \neq \hat{k}}$ by

$$
\tilde{m}_{kk'} := m_{kk'} - m_{k\hat{k}} - m_{\hat{k}k'} + m_{\hat{k}\hat{k}}.\tag{29}
$$

Then, the positive definiteness of M for zero-sum vectors is equivalent to the positive definiteness of M in the ordinary definition as follows.

Theorem 4 (Equivalence to positive definiteness). *If and only if* $m \times m$ *matrix* M *is positive definite for zero-sum vectors,* $(m - 1) \times (m - 1)$ *matrix* \tilde{M} *is positive definite.*

Proof. For any $(m-1)$ -dimensional vector $\tilde{\delta} = (\delta_k)_{k \neq \hat{k}}$, we define a m-dimensional zero-sum vector δ by

$$
\delta_k := \begin{cases}\n-\Sigma_{k \neq \hat{k}} \tilde{\delta}_k & (k = \hat{k}) \\
\tilde{\delta}_k & (k \neq \hat{k})\n\end{cases}.
$$
\n(30)

Conversely, it is trivially possible to define δ from any δ . Thus, there is one-to-one correspondence between the m-dimensional zero-sum vector δ and the $(m - 1)$ -dimensional (general-sum) vector δ . Now, we can prove $\delta \cdot (M \delta) = \tilde{\delta} \cdot (\tilde{M} \tilde{\delta})$ as follows.

$$
\delta \cdot (M\delta) = \sum_{k} \sum_{k'} \delta_{k} m_{kk'} \delta_{k'}
$$
\n
$$
= \sum_{k \neq \hat{k}} \sum_{k' \neq \hat{k}} \delta_{k} m_{kk'} \delta_{k'} + \sum_{k \neq \hat{k}} \delta_{k} m_{k\hat{k}} \delta_{\hat{k}} + \sum_{k' \neq \hat{k}} \delta_{\hat{k}} m_{\hat{k}k'} \delta_{k'} + \delta_{\hat{k}} m_{\hat{k}\hat{k}} \delta_{\hat{k}}
$$
\n
$$
= \sum_{k \neq \hat{k}} \sum_{k' \neq \hat{k}} \delta_{k} m_{kk'} \delta_{k'} - \sum_{k \neq \hat{k}} \sum_{k' \neq \hat{k}} \delta_{k} m_{k\hat{k}} \delta_{k'} - \sum_{k \neq \hat{k}} \sum_{k' \neq \hat{k}} \delta_{k} m_{\hat{k}k'} \delta_{k'} + \sum_{k \neq \hat{k}} \sum_{k' \neq \hat{k}} \delta_{k} m_{\hat{k}\hat{k}} \delta_{k'}
$$
\n
$$
= \sum_{k \neq \hat{k}} \sum_{k' \neq \hat{k}} \delta_{k} (m_{kk'} - m_{k\hat{k}} - m_{\hat{k}k'} + m_{\hat{k}\hat{k}}) \delta_{k'}
$$
\n
$$
= \sum_{k \neq \hat{k}} \sum_{k' \neq \hat{k}} \delta_{k} \tilde{m}_{kk'} \delta_{k'}
$$
\n
$$
= \tilde{\delta} \cdot (\tilde{M}\tilde{\delta}). \tag{31}
$$

Here, in the third equal sign, we used Eq. [\(30\)](#page-13-1). Thus, if and only if M is positive definite for zero-sum vectors (i.e., $\delta \cdot (\overline{M} \delta) > 0$), \tilde{M} is positive definite (i.e., $\tilde{\delta} \cdot (\tilde{M} \tilde{\delta}) > 0$). \Box

C Connection to the classical divergence

This section is dedicated to understanding the connection between our conditional-sum divergence $D(X, y)$ and the classical divergence $D_c(x, y)$. Here, recall that X's reactive strategy consists of multiple vectors $\bm{X}=(\bm{x}_j)_{1\leq j\leq m_\mathsf{Y}}$, where \bm{x}_j is the vector of the probability distribution of its action choice when the other chose j -th action in the previous round. On the other hand, in the classical divergence, X's mixed strategy x is the single vector of the probability distribution of its action choice independent of the other's previous choice. Thus, if the reactive strategy of X satisfies $x_j = x$ for all j, it corresponds to the mixed strategy of x. Then, $\dot{D}(\bm{X}, \bm{y}) = \dot{D}_{\rm c}(\bm{x}, \bm{y})$ holds, as the following theorem shows.

Theorem 5 (Connection with the classical total divergence). When X *satisfies* $x_j = x$ *for all j*, $\dot{D}(\boldsymbol{X}, \boldsymbol{y}) = \dot{D}_{\rm c}(\boldsymbol{x}, \boldsymbol{y}) = 0$ holds.

Proof. Let us prove that $\dot{D} = \sum_j \dot{D}_{\text{KL}}(\boldsymbol{x}^* \| \boldsymbol{x}_j) + \dot{D}_{\text{KL}}(\boldsymbol{y}^* \| \boldsymbol{y})$ and $\dot{D}_{\text{c}} = \dot{D}_{\text{KL}}(\boldsymbol{x}^* \| \boldsymbol{x}) + \dot{D}_{\text{KL}}(\boldsymbol{y}^* \| \boldsymbol{y})$ are equal. First, the time evolution of Y's divergence, i.e., $\dot{D}_{\text{KL}}(\bm{y}^*||\bm{y})$, is trivially equal between these two equations. Thus, we prove $\Sigma_j \dot{D}_{\text{KL}}(\bm{x}^* || \bm{x}_j) = \dot{D}_{\text{KL}}(\bm{x}^* || \bm{x})$ below.

$$
\Sigma_j \dot{D}_{\text{KL}}(\boldsymbol{x}^* \| \boldsymbol{x}_j) = -\Sigma_j \Sigma_i x_i^* \frac{\dot{x}_{i|j}}{x_{i|j}} \n= -\Sigma_j \Sigma_i x_i^* y_j (\Sigma_{j'} u_{ij'} y_{j'} - \Sigma_i \underbrace{x_{i|j}}_{=x_i} \Sigma_{j'} u_{ij'} y_{j'}) \n= -\Sigma_i x_i^* (\Sigma_{j'} u_{ij'} y_{j'} - \Sigma_i x_i \Sigma_{j'} u_{ij'} y_{j'}) \n= \dot{D}_{\text{KL}}(\boldsymbol{x}^* \| \boldsymbol{x}).
$$
\n(32)

In the final equal sign, we used the replicator dynamics of X's strategy for games without memories as

$$
\dot{x}_i = x_i \left(\frac{\mathrm{d}u_{\mathrm{st}}(\boldsymbol{x}, \boldsymbol{y})}{\mathrm{d}x_i} - \Sigma_i x_i \frac{\mathrm{d}u_{\mathrm{st}}(\boldsymbol{x}, \boldsymbol{y})}{\mathrm{d}x_i} \right)
$$
\n
$$
= x_i (\Sigma_j u_{ij} y_j - \Sigma_i x_i \Sigma_j u_{ij} y_j)
$$
\n(33)

 \Box

Thus, we proved $\dot{D} = \dot{D}_c$.

In this proof, we proved $\dot{D}(\bm{X},\bm{y})=\dot{D}_{\rm c}(\bm{x},\bm{y})$ separately for each divergence of X's strategy and Y's strategy.

Furthermore, note that $\dot{D}(\bm{X}, y) = \dot{D}_c(\bm{x}, y)$ holds only for a moment. Because the learning dynamics give $\dot{x}_j \neq \dot{x}, x_j$ gradually differs from x . In conclusion, $\dot{D}(\bm{X}, \bm{y}) = \dot{D}_\mathrm{c}(\bm{x}, \bm{y})$ becomes unsatisfied immediately. (However, $\dot{D}_{\rm c}(\mathbf{x},\mathbf{y})=0$ continues to hold.)

D Computational environment

The simulations presented in this paper were conducted using the following computational environment.

- Operating System: macOS Monterey (version 12.4)
- Programming Language: Python 3.11.3
- Processor: Apple M1 Pro (10 cores)
- Memory: 32 GB