Optimal Rates for Vector-Valued Spectral Regularization Learning Algorithms

Dimitri Meunier^{*,†}

Zikai Shen^{*,‡}

Mattes Mollenhauer[§]

Arthur Gretton[†]

Zhu Li[†]

Abstract

We study theoretical properties of a broad class of regularized algorithms with vector-valued output. These *spectral algorithms* include kernel ridge regression, kernel principal component regression, various implementations of gradient descent and many more. Our contributions are twofold. First, we rigorously confirm the so-called *saturation effect* for ridge regression with vector-valued output by deriving a novel lower bound on learning rates; this bound is shown to be suboptimal when the smoothness of the regression function exceeds a certain level. Second, we present the upper bound for the finite sample risk general vector-valued spectral algorithms, applicable to both well-specified and misspecified scenarios (where the true regression function lies outside of the hypothesis space) which is minimax optimal in various regimes. All of our results explicitly allow the case of infinite-dimensional output variables, proving consistency of recent practical applications.

1 Introduction

We investigate a fundamental topic in modern machine learning—the behavior and efficiency of learning algorithms for regression in high-dimensional and potentially infinite-dimensional output spaces \mathcal{Y} . Given two random variables X and Y, we seek to empirically minimize the squared expected risk

$$\mathcal{E}(F) \coloneqq \mathbb{E}\left[\|Y - F(X)\|_{\mathcal{V}}^2 \right] \tag{1}$$

over functions F in a reproducing kernel Hilbert space consisting of vector-valued functions from a topological space \mathcal{X} to a Hilbert space \mathcal{Y} . The study of this setting as an ill-posed statistical inverse problem is well established: see e.g. Smale and Zhou, 2007; Caponnetto and De Vito, 2007a; Yao et al., 2007; Bauer et al., 2007; Blanchard and Mücke, 2018; Fischer and Steinwart, 2020. In this work, we study the setting when \mathcal{Y} is high- or infinite-dimensional, since it has been less well covered by the literature, yet has many applications in multitask regression (Caponnetto et al., 2008; Baldassarre et al., 2012) and infinite-dimensional learning problems, including the conditional mean embedding (Grünewälder et al., 2012a,b; Park and Muandet, 2020), structured prediction (Ciliberto et al., 2016, 2020), causal inference (Singh et al., 2023), regression with instrumental and proximal variables (Singh et al., 2019; Mastouri et al., 2021), the estimation of linear operators and dynamical systems (Song et al., 2009; Mollenhauer and Koltai, 2020; Kostic et al., 2022; Mollenhauer et al.,

^{*}Equal Contribution.

[†]Gatsby Computational Neuroscience Unit, University College London, dimitri.meunier.21@ucl.ac.uk, arthur.gretton@gmail.com,zhu.li@ucl.ac.uk

[‡]University College London, zikai.shen.22@ucl.ac.uk

[§]Merantix Momentum, mattes.mollenhauer@merantix-momentum.com

2022; Kostic et al., 2023), and functional regression (Kadri et al., 2016). Interestingly, the aforementioned infinite-dimensional applications typically use the classical *ridge regression* algorithm. Our goal is to motivate the use of alternative learning algorithms in these settings, while providing strong theoretical guarantees.

Classically, the ill-posed problem (1) is solved via regularization strategies, which are often implemented in terms of so-called *spectral filter functions* in the context of inverse problems in Hilbert spaces (Engl et al., 2000). When applied to the learning problem given by (1), these filter functions correspond to learning algorithms including ridge regression, a variety of different implementations of gradient descent, principal component regression and many more (we refer the reader to Gerfo et al., 2008b and Baldassarre et al., 2012 for overviews of the real-valued and vector-valued output variable case, respectively). Algorithms based on spectral filter functions when $\mathcal{Y} = \mathbb{R}$ are studied extensively, see e.g. Blanchard and Mücke (2018); Lin et al. (2020b). To the best of our knowledge, the detailed behavior of this general class of algorithms has remained unknown when \mathcal{Y} is a general Hilbert space, with the exception of a few results for special cases in the setting of ridge regression (Caponnetto and De Vito, 2007a; Li et al., 2023b).

Overview of our contributions. In this manuscript, we aim to theoretically understand vector-valued spectral learning algorithms. The contribution of our work is twofold: (i) we rigorously confirm the saturation effect of ridge regression for general Hilbert spaces \mathcal{Y} (see paragraph below) in the context of lower bounds on rates for the learning problem (1) and (ii) we cover a gap in the existing literature by providing upper rates for general spectral algorithms in high- and infinite-dimensional spaces. Our results explicitly allow the misspecified learning case in which the true regression function is not contained in the hypothesis space. We base our analysis on the concept of vector-valued interpolation spaces introduced by Li et al. (2022, 2023b). The interpolation space norms measure the smoothness of the true regression function, replacing typical source conditions found in the literature which only cover the well-specified case. To the best of our knowledge, these are the first bounds covering this general setting for vector-valued spectral algorithms.

Saturation effect of ridge regression. The widely-used ridge regression algorithm is known to exhibit the so-called saturation effect: it fails to exploit additional smoothness in the target function beyond a certain threshold. This effect has been thoroughly investigated in the context of Tikhonov regularization in inverse problems (Engl et al., 2000, Chapter 5), but is generally reflected only in upper rates in the learning literature, see e.g. Lin et al. (2020b); Blanchard and Mücke (2018). Interestingly, existing lower bounds (Caponnetto and De Vito, 2007a; Blanchard and Mücke, 2018; Li et al., 2023b) are usually formulated in a more general setting and do not reflect this saturation effect, leaving a gap between upper and lower rates. We leverage the bias-variance decomposition paradigm to lower bound the learning risk of kernel ridge regression with vector-valued output, in order to close this gap.

Learning rates of vector-valued spectral algorithms. Motivated by the fact that the saturation effect is technically unavoidable with vector-valued ridge regression, we proceed to study the generalization error of popular alternative learning algorithms. In particular, we provide upper rates in the vector-valued setting consistent with the known behavior of spectral algorithms in the real-valued learning setting, based on their so-called *qualification property* (Blanchard and Mücke, 2018; Lin et al., 2020b). In particular, we confirm that a saturation effect can be bypassed in high and infinite dimensions by algorithms such as principal component regression and gradient descent, allowing for a better sample complexity for high-smoothness problems. Furthermore, we study the misspecified setting and show that upper rates for spectral algorithms match the state-of-the-art upper rates for misspecified vector-valued ridge regression recently obtained by Li et al. (2023b). Those rates are optimal for a wide variety of settings. Moreover, we argue that applications of vector-valued spectral algorithms are easy to implement by making use of an extended *representer theorem* based on Baldassarre et al. (2012), allowing for the numerical evaluation based on empirical data—even in the infinite-dimensional case.

Related Work. The saturation effect of regularization techniques in deterministic inverse problems is wellknown. For example, Neubauer (1997); Mathé (2004); Herdman et al. (2011) study the saturation effect for Tikhonov regularization and general spectral algorithms. In the kernel statistical learning framework, the general phenomenon of saturation is discussed by e.g. Bauer et al. (2007); Gerfo et al. (2008a). Recent work by Li et al. (2023a) investigates saturation effect in the learning context by providing a lower bound on the learning rate. To the best of our knowledge, however, all studies in the statistical learning context focus on the case when Y is real-valued. General upper bounds of kernel ridge regression with real-valued or finite-dimensional Y have been extensively studied in the literature (see e.g., Caponnetto and De Vito (2007b); Steinwart et al. (2009); Caponnetto and Yao (2010); Fischer and Steinwart (2020)), where minimax optimal learning rates are derived. Recent work Li et al. (2022, 2023b) studies the infinite-dimensional output space setting with Tikhonov regularization and obtains analogous minimax optimal learning rates. For kernel learning with spectral algorithms, existing work (see e.g., Bauer et al. (2007); Blanchard and Mücke (2018); Lin and Cevher (2018); Lin et al. (2020b); Zhang et al. (2023b)) focuses on real-valued output space setting and obtains optimal upper learning rates depending on the qualification number of the spectral algorithms, where only Zhang et al. (2023b) considers the misspecified learning scenario where the target regression function does not lie in the hypothesis space. However, general investigations of spectral algorithms for vector-valued output spaces are absent in the literature.

Structure of this paper. This paper is structured as follows. In Section 2, we introduce mathematical preliminaries related to reproducing kernel Hilbert spaces, vector-valued regression as well as the concept of vector-valued interpolation spaces. Section 3 contains a brief review the so-called saturation effect and a corresponding novel lower bound for vector-valued kernel ridge regression. In Section 4, we investigate general spectral learning algorithms in the context of vector-valued interpolations spaces and provide our main result: upper learning rates for this setting.

2 Background and Preliminaries

Throughout the paper, we consider a random variable X (the covariate) defined on a second countable locally compact Hausdorff space¹ \mathcal{X} endowed with its Borel σ -field $\mathcal{F}_{\mathcal{X}}$, and the random variable Y (the output) defined on a potentially infinite dimensional separable real Hilbert space $(\mathcal{Y}, \langle \cdot, \cdot \rangle_{\mathcal{Y}})$ endowed with its Borel σ -field $\mathcal{F}_{\mathcal{Y}}$. We let $(\Omega, \mathcal{F}, \mathbb{P})$ be the underlying probability space with expectation operator \mathbb{E} . Let P be the push-forward of \mathbb{P} under (X, Y) and π and ν be the marginal distributions on \mathcal{X} and \mathcal{Y} , respectively; i.e., $X \sim \pi$ and $Y \sim \nu$. We use the Markov kernel $p: \mathcal{X} \times \mathcal{F}_{\mathcal{Y}} \to \mathbb{R}_+$ to express the distribution of Y conditioned on X as

$$\mathbb{P}[Y \in A | X = x] = \int_A p(x, dy),$$

for all $x \in \mathcal{X}$ and events $A \in \mathcal{F}_{\mathcal{Y}}$, see e.g. Dudley (2002). We introduce some notation related to linear operators on Hilbert spaces and vector-valued integration; formal definitions can be found in Appendix A for completeness, or we refer the reader to Weidmann (1980); Diestel and Uhl (1977). The spaces of Bochner square-integrable functions with respect to π and taking values in \mathcal{Y} are written as $L_2(\mathcal{X}, \mathcal{F}_{\mathcal{X}}, \pi; \mathcal{Y})$, abbreviated as $L_2(\pi; \mathcal{Y})$. We obtain the classical Lebesgue spaces as $L_2(\pi) \coloneqq L_2(\pi; \mathbb{R})$. Throughout the paper, we write [F] or more explicitly $[F]_{\pi}$ for the π -equivalence class of (potentially pointwise defined) measurable functions from \mathcal{X} to \mathcal{Y} , which we naturally interpret as elements in $L_2(\pi; \mathcal{Y})$ whenever they are square integrable. Let H be a separable real Hilbert space with inner product $\langle \cdot, \cdot \rangle_H$. We write $\mathcal{L}(H, H')$ as the Banach space of bounded linear operators from H to another Hilbert space H', equipped with the operator norm $\|\cdot\|_{H \to H'}$. When H = H', we simply write $\mathcal{L}(H)$ instead. We write $S_2(H, H')$ as the Hilbert space of Hilbert-Schmidt operators from H to H' and $S_1(H, H')$ as the Banach space of trace class operators (see Appendix A for a complete definition). For two Hilbert spaces H, H', we say that H is (continuously) embedded in H' and denote it as $H \to H'$ if H can be interpreted as a vector subspace of H' and the inclusion operator $i: H \to H'$ performing the change of norms with ix = x for $x \in H$ is continuous; and we say that H is isometrically

¹Under additional technical assumptions, the results in this paper can also be formulated when \mathcal{X} is a more general topological space. However, some properties of kernels defined on \mathcal{X} such as the so-called c_0 -universality (Carmeli et al., 2010) simplify the exposition when \mathcal{X} is a second countable locally compact Hausdorff space.

isomorphic to H' and denote it as $H \simeq H'$ if there is a linear isomorphism between H and H' which is an isometry.

Tensor Product of Hilbert Spaces: Denote $H \otimes H'$ the tensor product of Hilbert spaces H, H'. The element $x \otimes x' \in H \otimes H'$ is treated as the linear rank-one operator $x \otimes x' : H' \to H$ defined by $y' \to \langle y', x' \rangle_{H'} x$ for $y' \in H'$. Based on this identification, the tensor product space $H \otimes H'$ is isometrically isomorphic to the space of Hilbert-Schmidt operators from H' to H, i.e., $H \otimes H' \simeq S_2(H', H)$. We will hereafter not make the distinction between these two spaces, and treat them as being identical.

Remark 1 (Aubin, 2000, Theorem 12.6.1). Consider the Bochner space $L_2(\pi; H)$ where H is a separable Hilbert space. One can show that $L_2(\pi; H)$ is isometrically identified with the tensor product space $H \otimes L_2(\pi)$, and we denote as Ψ the isometric isomorphism between the two spaces. See Appendix A for more details on tensor product spaces and the explicit definition of Ψ .

Scalar-valued Reproducing Kernel Hilbert Space (RKHS). We let $k : \mathcal{X} \times \mathcal{X} \to \mathbb{R}$ be a symmetric and positive definite kernel function and \mathcal{H} be a vector space of functions from \mathcal{X} to \mathbb{R} , endowed with a Hilbert space structure via an inner product $\langle \cdot, \cdot \rangle_{\mathcal{H}}$. We say that k is a reproducing kernel of \mathcal{H} if and only if for all $x \in \mathcal{X}$ we have $k(\cdot, x) \in \mathcal{H}$ and for all $x \in \mathcal{X}$ and $f \in \mathcal{H}$, we have $f(x) = \langle f, k(x, \cdot) \rangle_{\mathcal{H}}$. A space \mathcal{H} which possesses a reproducing kernel is called a reproducing kernel Hilbert space (RKHS; see e.g. Berlinet and Thomas-Agnan, 2011). We denote the canonical feature map of \mathcal{H} as $\phi(x) = k(\cdot, x)$.

We require some technical assumptions on the previously defined RKHS and kernel, which we assume to be satisfied throughout the text:

- 1. \mathcal{H} is separable, this is satisfied if k is continuous, given that \mathcal{X} is separable;²
- 2. $k(\cdot, x)$ is measurable for π -almost all $x \in \mathcal{X}$;
- 3. $k(x, x) \leq \kappa^2$ for π -almost all $x \in \mathcal{X}$.

The above assumptions are not restrictive in practice, as well-known kernels such as the Gaussian, Laplace and Matérn kernels satisfy them on $\mathcal{X} \subseteq \mathbb{R}^d$ (Sriperumbudur et al., 2011). We now introduce some facts about the interplay between \mathcal{H} and $L_2(\pi)$, which has been extensively studied by Smale and Zhou (2004, 2005), De Vito et al. (2006) and Steinwart and Scovel (2012). We first define the (not necessarily injective) embedding $I_{\pi} : \mathcal{H} \to L_2(\pi)$, mapping a function $f \in \mathcal{H}$ to its π -equivalence class [f]. The embedding is a well-defined compact operator as long as its Hilbert-Schmidt norm is finite. In fact, this requirement is satisfied since its Hilbert-Schmidt norm can be computed as (Steinwart and Scovel, 2012, Lemma 2.2 & 2.3) $\|I_{\pi}\|_{S_2(\mathcal{H},L_2(\pi))} = \|k\|_{L_2(\pi)} \leq \kappa$. The adjoint operator $S_{\pi} \coloneqq I_{\pi}^* : L_2(\pi) \to \mathcal{H}$ is an integral operator with respect to the kernel k, i.e. for $f \in L_2(\pi)$ and $x \in \mathcal{X}$ we have (Steinwart and Christmann, 2008, Theorem 4.27)

$$(S_{\pi}f)(x) = \int_{\mathcal{X}} k(x, x') f(x') d\pi(x').$$

Next, we define the self-adjoint, positive semi-definite and trace class integral operators

$$L_X \coloneqq I_\pi S_\pi \colon L_2(\pi) \to L_2(\pi) \quad \text{and} \quad C_X \coloneqq S_\pi I_\pi \colon \mathcal{H} \to \mathcal{H}.$$

Vector-valued Reproducing Kernel Hilbert Space (vRKHS). Let $K : \mathcal{X} \times \mathcal{X} \to \mathcal{L}(\mathcal{Y})$ be an operator valued positive-semidefinite (psd) kernel. Fix $K, x \in \mathcal{X}$, and $h \in \mathcal{Y}$, then $(K_x h)(\cdot) := K(\cdot, x)h$ defines a function from \mathcal{X} to \mathcal{Y} . The completion of

$$\mathcal{G}_{\text{pre}} \coloneqq \text{span} \{ K_x h \mid x \in \mathcal{X}, h \in \mathcal{Y} \}$$

²This follows from Steinwart and Christmann (2008, Lemma 4.33). Note that the Lemma requires separability of \mathcal{X} , which is satisfied since we assume that \mathcal{X} is a second countable locally compact Hausdorff space.

with inner product on \mathcal{G}_{pre} defined on the elementary elements as $\langle K_x h, K_{x'} h' \rangle_{\mathcal{G}} \coloneqq \langle h, K(x, x') h' \rangle_{\mathcal{Y}}$, defines a vRKHS denoted as \mathcal{G} . For a more complete overview of the vector-valued reproducing kernel Hilbert space, we refer the reader to Carmeli et al. (2006), Carmeli et al. (2010) and Li et al. (2023b, Section 2). In the following, we will denote \mathcal{G} as the vRKHS induced by the kernel $K : \mathcal{X} \times \mathcal{X} \to \mathcal{L}(\mathcal{Y})$ with

$$K(x, x') \coloneqq k(x, x') \operatorname{Id}_{\mathcal{Y}}, \quad x, x' \in \mathcal{X}.$$
(2)

We emphasize that this family of kernels is the de-facto standard for high- and infinite-dimensional applications (Grünewälder et al., 2012a,b; Park and Muandet, 2020; Ciliberto et al., 2016, 2020; Singh et al., 2019; Mastouri et al., 2021; Singh et al., 2023; Mollenhauer and Koltai, 2020; Kostic et al., 2022; Mollenhauer et al., 2022; Kostic et al., 2023; Kadri et al., 2016) due to the crucial *representer theorem* which gives a closed form solution for the ridge regression problem based on the data. We generalize this representer theorem to cover the general spectral algorithm case in Proposition 1.

Remark 2 (General multiplicative kernel). Without loss of generality, we provide our results for the vRKHS \mathcal{G} induced by the operator-valued kernel given by $K(x,x') = k(x,x') \operatorname{Id}_{\mathcal{Y}}$. However, with suitably adjusted constants in the assumptions, our results transfer directly to the more general vRKHS $\tilde{\mathcal{G}}$ induced by the more general operator-valued kernel

$$\widetilde{K}(x,x') \coloneqq k(x,x')T$$

where $T: \mathcal{Y} \to \mathcal{Y}$ is any positive-semidefinite self-adjoint operator. The precise characterization of the adjusted constants is given by Li et al. (2023b, Section 4.1).

An important property of \mathcal{G} is that it is isometrically isomorphic to the space of Hilbert-Schmidt operators between \mathcal{H} and \mathcal{Y} (Li et al., 2023b, Corollary 1). Similarly to the scalar case we can map every element in \mathcal{G} into its π -equivalence class in $L_2(\pi; \mathcal{Y})$ and we use the shorthand notation $[F] = [F]_{\pi}$ (see Definition 6 in Appendix A for more details).

Theorem 1 (vRKHS isomorphism). For every function $F \in \mathcal{G}$ there exists a unique operator $C \in S_2(\mathcal{H}, \mathcal{Y})$ such that $F(\cdot) = C\phi(\cdot) \in \mathcal{Y}$ with $||C||_{S_2(\mathcal{H}, \mathcal{Y})} = ||F||_{\mathcal{G}}$ and vice versa. Hence $\mathcal{G} \simeq S_2(\mathcal{H}, \mathcal{Y})$ and we denote the isometric isomorphism between $S_2(\mathcal{H}, \mathcal{Y})$ and \mathcal{G} as $\overline{\Psi}$. It follows that \mathcal{G} can be written as $\mathcal{G} = \{F : \mathcal{X} \to \mathcal{Y} \mid F = C\phi(\cdot), C \in S_2(\mathcal{H}, \mathcal{Y})\}.$

2.1 Vector-valued Regression

We briefly recall the basic setup of regularized least squares regression with Hilbert space-valued random variables. The squared expected risk for vector-valued regression is

$$\mathcal{E}(F) \coloneqq \mathbb{E}\left[\|Y - F(X)\|_{\mathcal{Y}}^2\right] = \int_{\mathcal{X} \times \mathcal{Y}} \|y - F(x)\|_{\mathcal{Y}}^2 p(x, dy) \pi(dx), \tag{3}$$

for measurable functions $F : \mathcal{X} \to \mathcal{Y}$. The analytical minimizer of the risk over all those measurable functions is the regression function or the conditional mean function $F_{\star} \in L_2(\pi; \mathcal{Y})$ given by

$$F_*(x) \coloneqq \mathbb{E}[Y \mid X = x] = \int_{\mathcal{Y}} y p(x, dy), \quad x \in \mathcal{X}.$$

Throughout the paper, we assume that $\mathbb{E}[||Y||_{\mathcal{Y}}^2] < +\infty$, i.e., the random variable Y is square integrable. Note that this implies $F_* \in L_2(\pi; \mathcal{Y})$. Our focus in this work is to approximate F_* with kernel-based regularized least-squares algorithms, where we pay special attention to the case when \mathcal{Y} is of high or infinite dimension. We pick \mathcal{G} as a hypothesis space of functions in which to estimate F_* . Note that by Theorem 1, minimizing the functional \mathcal{E} on \mathcal{G} is equivalent to minimizing the following functional on $S_2(\mathcal{H}, \mathcal{Y})$,

$$\bar{\mathcal{E}}(C) \coloneqq \mathbb{E}\left[\|Y - C\phi(X)\|_{\mathcal{Y}}^2 \right]. \tag{4}$$

It is shown in Mollenhauer et al. (2022, Proposition 3.5 and Section 3.4) that the optimality condition can be written as

$$C_{YX} = C_* C_X, \qquad C_* \in S_2(\mathcal{H}, \mathcal{Y}), \tag{5}$$

where $C_{YX} := \mathbb{E}[Y \otimes \phi(X)]$ is the cross-covariance operator. As discussed in full detail by Mollenhauer et al. (2022), the problem 5 can be formulated as a potentially ill-posed inverse problem on the space of Hilbert-Schmidt operators. As such, a regularization is required; we introduce regularized solutions of this problem in Section 4 through the classical concept of spectral filter functions.

2.2Vector-valued Interpolation Space and Source Condition

We now introduce the background required in order to characterize the smoothness of the target function F_* , both in the well-specified setting $(F_* \in \mathcal{G})$ and in the misspecified setting $(F_* \notin \mathcal{G})$. We review the results of Steinwart and Scovel (2012) and Fischer and Steinwart (2020) in constructing scalar-valued interpolation spaces, and Li et al. (2022) in defining vector-valued interpolation spaces.

Real-valued Interpolation Space: By the spectral theorem for self-adjoint compact operators, there exists an at most countable index set I, a non-increasing sequence $(\mu_i)_{i \in I} > 0$, and a family $(e_i)_{i \in I} \in \mathcal{H}$, such that $([e_i])_{i \in I}$ ³ is an orthonormal basis (ONB) of $\overline{\operatorname{ran} I_{\pi}} \subseteq L_2(\pi)$ and $(\mu_i^{1/2} e_i)_{i \in I}$ is an ONB of $(\ker I_{\pi})^{\perp} \subseteq \mathcal{H}$, and we have

$$L_X = \sum_{i \in I} \mu_i \langle \cdot, [e_i] \rangle_{L_2(\pi)} [e_i], \qquad C_X = \sum_{i \in I} \mu_i \langle \cdot, \mu_i^{\frac{1}{2}} e_i \rangle_{\mathcal{H}} \mu_i^{\frac{1}{2}} e_i \tag{6}$$

For $\alpha \geq 0$, the α -interpolation space Steinwart and Scovel (2012) is defined by

$$[\mathcal{H}]^{\alpha} \coloneqq \left\{ \sum_{i \in I} a_i \mu_i^{\alpha/2} \left[e_i \right] \colon \left(a_i \right)_{i \in I} \in \ell_2(I) \right\} \subseteq L_2(\pi),$$

equipped with the inner product

$$\left\langle \sum_{i \in I} a_i(\mu_i^{\alpha/2}[e_i]), \sum_{i \in I} b_i(\mu_i^{\alpha/2}[e_i]) \right\rangle_{[\mathcal{H}]^{\alpha}} = \sum_{i \in I} a_i b_i,$$

for $(a_i)_{i \in I}$, $(b_i)_{i \in I} \in \ell_2(I)$. The α -interpolation space defines a Hilbert space. Moreover, $(\mu_i^{\alpha/2}[e_i])_{i \in I}$ forms an ONB of $[\mathcal{H}]^{\alpha}$ and consequently $[\mathcal{H}]^{\alpha}$ is a separable Hilbert space. In the following, we use the abbreviation $\|\cdot\|_{\alpha} \coloneqq \|\cdot\|_{[\mathcal{H}]^{\alpha}}.$

Vector-valued Interpolation Space: Introduced in Li et al. (2022), vector-valued interpolation spaces generalize the notion of scalar-valued interpolation spaces to vRKHS with a kernel of the form (2).

Definition 1 (Vector-valued interpolation space). Let k be a real-valued kernel with associated RKHS \mathcal{H} and let $[\mathcal{H}]^{\alpha}$ be the real-valued interpolation space associated to \mathcal{H} with some $\alpha \geq 0$. The vector-valued interpolation space $[\mathcal{G}]^{\alpha}$ is defined as (refer to Remark 1 for the definition of Ψ)

$$[\mathcal{G}]^{\alpha} \coloneqq \Psi\left(S_2([\mathcal{H}]^{\alpha}, \mathcal{Y})\right) = \{F \mid F = \Psi(C), \ C \in S_2([\mathcal{H}]^{\alpha}, \mathcal{Y})\}.$$

The space $[\mathcal{G}]^{\alpha}$ is a Hilbert space equipped with the inner product

$$\langle F, G \rangle_{\alpha} \coloneqq \langle C, L \rangle_{S_2([\mathcal{H}]^{\alpha}, \mathcal{Y})} \qquad (F, G \in [\mathcal{G}]^{\alpha}),$$

where $C = \Psi^{-1}(F)$, $L = \Psi^{-1}(G)$. For $\alpha = 0$, we retrieve $||F||_0 = ||C||_{S_2(L_2(\pi),\mathcal{Y})}$. ³We recall that the bracket [·] denotes the embedding that maps f to its equivalence class $I_{\pi}(f) \in L_2(\pi)$.

Remark 3 (Interpolation space inclusions). Note that we have $F_* \in L_2(\pi; \mathcal{Y})$ since $Y \in L_2(\mathbb{P}; \mathcal{Y})$ by assumption. Furthermore, for $0 < \beta < \alpha$, Fischer and Steinwart (2020, Eq. (7)) imply the inclusions

$$[\mathcal{G}]^{\alpha} \hookrightarrow [\mathcal{G}]^{\beta} \hookrightarrow [\mathcal{G}]^{0} \subseteq L_{2}(\pi; \mathcal{Y}).$$

Under assumptions 1 to 3 and with \mathcal{X} being a second-countable locally compact Hausdorff space, $[\mathcal{G}]^0 = L_2(\pi; \mathcal{Y})$ if and only if \mathcal{H} is dense in the space of continuous functions vanishing at infinity, equipped with the uniform norm (Li et al., 2023b, Remark 4).

Remark 4 (Well-specified versus misspecified setting). We say that we are in the well-specified setting if $F_* \in [\mathcal{G}]^1$. In this case, there exists $\overline{F} \in \mathcal{G}$ such that $F_* = \overline{F} \pi$ -almost surely and $||F_*||_1 = ||\overline{F}||_{\mathcal{G}}$, i.e. F_* admits a representer in \mathcal{G} (see Remark 5 in Appendix A). When $F_* \in [\mathcal{G}]^\beta$ for $\beta < 1$, F_* may not admit such a representation and we are in the misspecified setting, as $[\mathcal{G}]^1 \subseteq [\mathcal{G}]^\beta$.

Definition 1 and Remarks 3 and 4 motivate the use of following assumption on the smoothness of the target function: there exists $\beta > 0$ and a constant $B \ge 0$ such that $F_* \in [\mathcal{G}]^{\beta}$

$$\|F_*\|_{\beta} \le B. \tag{SRC}$$

We let $C_* := \Psi^{-1}(F_*) \in S_2([\mathcal{H}]^\beta, \mathcal{Y})$. (SRC) directly generalizes the notion of a so-called Hölder-type source condition in the learning literature (Caponnetto and De Vito, 2007b; Fischer and Steinwart, 2020; Lin and Cevher, 2018; Lin et al., 2020b) and allows to characterize the misspecified learning scenario.

2.3 Further Assumptions

In addition to (SRC), we require standard assumptions to obtain the precise learning rate for kernel learning algorithms. We list them below. For constants $D_2 > 0$ and $p \in (0, 1]$ and for all $i \in I$,

$$\mu_i \le D_2 i^{-1/p}. \tag{EVD}$$

For constants $D_1, D_2 > 0$ and $p \in (0, 1)$ and for all $i \in I$,

$$D_1 i^{-\frac{i}{p}} \le \mu_i \le D_2 i^{-1/p}. \tag{EVD+}$$

(EVD) and (EVD+) are standard assumptions on the *eigenvalue decay* of the integral operator: they describe the interplay between the marginal distribution π and the RKHS \mathcal{H} (see more details in Caponnetto and De Vito, 2007b; Fischer and Steinwart, 2020). (EVD+) is needed in order to establish lower bounds on the excess risk. Note that we have excluded the value p = 1 from (EVD+); indeed, p = 1 is incompatible with the assumption of a bounded kernel, a fact missed by previous works and of independent interest (see Appendix, Remark 7).

For $\alpha \in [p, 1]$, the inclusion $I_{\pi}^{\alpha, \infty} : [\mathcal{H}]^{\alpha} \hookrightarrow L_{\infty}(\pi)$ is continuous, and $\exists A > 0$ such that

$$\|I_{\pi}^{\alpha,\infty}\|_{[\mathcal{H}]^{\alpha}\to L_{\infty}(\pi)} \le A.$$
(EMB)

Property (EMB) is referred to as the embedding property in Fischer and Steinwart (2020). It can be shown that it holds if and only if there exists a constant $A \ge 0$ with $\sum_{i \in I} \mu_i^{\alpha} e_i^2(x) \le A^2$ for π -almost all $x \in \mathcal{X}$ (Fischer and Steinwart, 2020, Theorem 9). Since we assume k to be bounded, the embedding property always hold true when $\alpha = 1$. Furthermore, (EMB) implies a polynomial eigenvalue decay of order $1/\alpha$, which is why we take $\alpha \ge p$. (EMB) is not needed when we deal with the well-specified setting, but is crucial to bound the excess risk in the misspecified setting.

Finally, we assume that there are constants $\sigma, R > 0$ such that

$$\int_{\mathcal{Y}} \|y - F_*(x)\|_{\mathcal{Y}}^q p(x, dy) \le \frac{1}{2} q! \sigma^2 R^{q-2},$$
(MOM)

is satisfied for π -almost all $x \in \mathcal{X}$ and all $q \ge 2$. The (MOM) condition on the Markov kernel p(x, dy) is a *Bernstein moment condition* used to control the noise of the observations (see Caponnetto and De Vito, 2007b; Fischer and Steinwart, 2020 for more details). If Y is almost surely bounded, for example $||Y||_{\mathcal{Y}} \le Y_{\infty}$ almost surely, then (MOM) is satisfied with $\sigma = R = 2Y_{\infty}$. It is possible to prove that the Bernstein condition is equivalent to sub-exponentiality, see Mollenhauer et al. (2022, Remark 4.9).

3 Saturation Effect of Kernel Ridge Regression

The most established way of learning F_* is by kernel ridge regression (KRR), which can be formulated as the following optimization problem: given a dataset $D = \{(x_i, y_i)\}_{i=1}^n$ independently and identically sampled from the joint distribution of X and Y,

$$\hat{F}_{\lambda} \coloneqq \underset{F \in \mathcal{G}}{\operatorname{arg\,min}} \frac{1}{n} \sum_{i=1}^{n} \|y_i - F(x_i)\|_{\mathcal{Y}}^2 + \lambda \|F\|_{\mathcal{G}}^2, \tag{7}$$

where $\lambda > 0$ is the regularization parameter. The generalization error of vector-valued KRR is expressed as $\hat{F}_{\lambda} - F_*$, and controlled in different norms: see Li et al. (2023b) for an extensive study. We recall here a simplified special case of the key results obtained in this work. In the next Theorem, \leq, \geq are inequality up to positive multiplicative constants that are independent of n.

Theorem 2 (Upper and lower bounds for KRR in the well-specified regime). Let \hat{F}_{λ} be the KRR estimator from (7). Furthermore, let the conditions (EVD+), (SRC) and (MOM) be satisfied for some $0 and <math>\beta \ge 1$. Then, with high probability we have

$$\left\| \left[\hat{F}_{\lambda_n} \right] - F_* \right\|_{L_2(\pi)}^2 \lesssim n^{-\frac{\min\{\beta,2\}}{\min\{\beta,2\}+p}} \quad for \ a \ choice \ \lambda_n = \Theta\left(n^{-\frac{1}{\beta+p}}\right),$$

and furthermore for all learning methods (i.e., measurable maps) of the form $D \rightarrow \hat{F}_D$,

$$\|[\hat{F}_D] - F_*\|_{L_2(\pi)}^2 \gtrsim n^{-\frac{\beta}{\beta+p}}.$$

Theorem 2 shows the minimax optimal learning rate for vector-valued KRR for $\beta \in [1, 2]$. However, when $\beta > 2$, the obtained upper bound saturates at $n^{-\frac{1}{2+p}}$, creating a gap with the lower bound. This phenomenon is referred to as the saturation effect of Tikhonov regularization, and has been well investigated in deterministic inverse problems (Neubauer, 1997). In the case where \mathcal{Y} is real-valued, Li et al. (2023a) prove that the saturation effect cannot be avoided with Tikhonov regularization. Below, we give a similar but generalized bound on lower rates for the case that \mathcal{Y} is a Hilbert space. For this result only, we assume that \mathcal{X} is a compact subset of \mathbb{R}^d . We give the proof in Appendix B.

Theorem 3 (Saturation of KRR). Let \mathcal{X} be a compact subset of \mathbb{R}^d . Let $\lambda = \lambda(n)$ be an arbitrary choice of regularization parameter satisfying $\lambda(n) \to 0$ as $n \to +\infty$ and let \hat{F}_{λ} be the KRR estimator from (7). We assume that the noise is non-zero and bounded below, i.e. there exists $\sigma > 0$, such that

$$\int_{\mathcal{Y}} \|y - F_*(x)\|_{\mathcal{Y}}^2 p(x, dy) \ge \sigma^2,$$

is satisfied for π -almost all $x \in \mathcal{X}$. We assume in addition and for this result only that k is Hölder continuous (see Definition 11 in the appendix), i.e., $k \in C^{\theta}(\mathcal{X} \times \mathcal{X})$ for $\theta \in (0,1]$. Suppose that Assumptions (EVD+) and (SRC) hold with $p \in (0,1)$ and $\beta \geq 2$. For $\tau \geq 0$, for sufficiently large n > 0, where the hidden index bound depends on τ , with probability greater than $1 - e^{-\tau}$, there exists some constant $c_{\tau} > 0$ such that

$$\mathbb{E}\left[\left\|\left[\hat{F}_{\lambda}\right]-F_{\star}\right\|^{2}_{L_{2}(\pi;\mathcal{Y})}\left|x_{1},\ldots,x_{n}\right]\geq c_{\tau}n^{-\frac{2}{2+p}}.$$

The assumption that k is Hölder continuous is crucial in lower bounding the variance with a covering number argument. Kernels satisfying this assumption include Gaussian kernels, Laplace kernels and Matérn kernels. Theorem 3 clearly demonstrates that the learning rate from vector-valued KRR cannot reach the information theoretic lower rate.

As discussed above, Li et al. (2023a) propose a similar lower bound in the real-valued case, and we now highlight two fundamental differences with Li et al. (2023a) in the proof. First, while both works adopt the same biasvariance decomposition, we need to lower bound the bias and the variance term with infinite-dimensional output in our setting. Second, we adopt a different and simpler approach in proving the lower bound, since there are a number of issues with the proof of Li et al. (2023a), both in the treatment of the bias and of the variance. For a detailed comparison with the earlier work, and an explanation of the differences in our approach, please refer to Remark 6 in the Appendix.

4 Consistency and optimal rates for general spectral algorithms

Regularized population solution: Our goal is to regularize (5) in such a way that we get a unique and well-defined solution that provides a good approximation to F_* . We first recall the concept of a filter function (i.e., a function on an interval which is applied on self-adjoint operators to each individual eigenvalue via the spectral calculus, see Engl et al., 2000), that will allow to define a regularization strategy. One may think of the following definition as a class of functions approximating the inversion map $x \mapsto 1/x$ while still being defined for x = 0 in a reasonable way. We use the definition given by Lin et al. (2020a), but equivalent definitions can be found throughout the literature.

Definition 2 (Filter function). Let $\Lambda \subseteq \mathbb{R}^+$. A family of functions $g_{\lambda} : [0, \infty) \to [0, \infty)$ indexed by $\lambda \in \Lambda$ is called a filter with qualification $\rho \ge 0$ if it satisfies the following two conditions:

1. There exists a positive constant E such that, for all $\lambda \in \Lambda$

$$\sup_{\alpha \in [0,1]} \sup_{x \in [0,\kappa^2]} \lambda^{1-\alpha} x^{\alpha} g_{\lambda}(x) \le E$$
(8)

2. There exists a positive constant $\omega_{\rho} < \infty$ such that

$$\sup_{\alpha \in [0,\rho]} \sup_{\lambda \in \Lambda} \sup_{x \in [0,\kappa^2]} |r_{\lambda}(x)| x^{\alpha} \lambda^{-\alpha} \le \omega_{\rho},$$
(9)

with $r_{\lambda}(x) \coloneqq 1 - g_{\lambda}(x)x$.

Below, we give some standard examples which are discussed by e.g. Gerfo et al. (2008b); Blanchard and Mücke (2018) in the context of kernel regression with scalar output variables, and in Baldassarre et al. (2012) for the vector-valued case. A variety of additional algorithms can be expressed in terms of a filter function.

1. Ridge regression. From the Tikhonov filter function $g_{\lambda}(x) = (x+\lambda)^{-1}$, we obtain the known ridge regression algorithm. In this case, we have $E = \rho = \omega_{\rho} = 1$.

2. Gradient Descent. From the Landweber iteration filter function given by

$$g_k(x) \coloneqq \tau \sum_{i=0}^{k-1} (1-\tau x)^i \text{ for } k \coloneqq 1/\lambda, k \in \mathbb{N}$$

we obtain the gradient descent scheme with constant step size $\tau > 0$, which corresponds to the population gradient iteration given by $F_{k+1} := F_k - \tau \nabla \mathcal{E}(F_k)$ for $k \in \mathbb{N}$. In this case, we have E = 1 and arbitrary qualification with $\omega_{\rho} = 1$ whenever $0 < \rho \leq 1$ and $\omega_{\rho} = \rho^{\rho}$ otherwise. Gradient schemes with more complex update rules can be expressed in terms of filter functions as well (Mücke et al., 2019; Lin and Cevher, 2018; Lin et al., 2020b).

3. Kernel principal component regression. The truncation filter function $g_{\lambda}(x) = x^{-1}\mathbb{1}[x \ge \lambda]$ yields kernel principal component regression, corresponding to a hard thresholding of eigenvalues at a truncation level λ . In this case we have $E = \omega_{\rho} = 1$ for arbitrary qualification ρ .

Population solution: Given a filter function g_{λ} , we call $g_{\lambda}(C_X)^4$ the regularized inverse of C_X . We may think of the regularized inverse as approximating the *pseudoinverse* of C_X (see e.g. Engl et al. (2000)) when $\lambda \to 0$. We define the regularized population solution to (4) as

$$C_{\lambda} \coloneqq C_{YX} g_{\lambda}(C_X) \in S_2(\mathcal{H}, \mathcal{Y}), \qquad F_{\lambda}(\cdot) \coloneqq C_{\lambda} \phi(\cdot) \in \mathcal{G}.$$

$$(10)$$

The solution arising from standard regularization strategies leads to well-known statistical methodologies. We refer to Engl et al. (2000) for the background on filter functions in classical regularization theory.

Empirical solution: Given the dataset $D = \{(x_i, y_i)\}_{i=1}^n$, the empirical analogue of (10) is

$$\hat{C}_{\lambda} \coloneqq \hat{C}_{YX} g_{\lambda}(\hat{C}_X), \qquad \hat{F}_{\lambda}(\cdot) \coloneqq \hat{C}_{\lambda} \phi(\cdot) \in \mathcal{G}, \tag{11}$$

where \hat{C}_{YX} , \hat{C}_X are empirical covariance operators define as

$$\hat{C}_X \coloneqq \frac{1}{n} \sum_{i=1}^n \phi(x_i) \otimes \phi(x_i) \qquad \hat{C}_{YX} \coloneqq \frac{1}{n} \sum_{i=1}^n y_i \otimes \phi(x_i).$$

Note that (11) is the regularized solution of the empirical inverse problem

$$\hat{C}_{YX} = \hat{C}\hat{C}_X, \qquad \hat{C} \in S_2(\mathcal{H}, \mathcal{Y})$$

which arises as the optimality condition for minimizers on \mathcal{G} of the empirical analogue of (3), given by $\mathcal{E}_n(F) := \frac{1}{n} \sum_{i=1}^n \|y_i - F(x_i)\|_{\mathcal{Y}}^2$; see Proposition 2 in the Appendix for a proof. For the vector-valued kernel given in (2), it is well-known that \hat{F}_{λ} can be computed in closed-form for the ridge regression estimator—even in infinite dimensions Song et al. (2009). For general filter functions, an extended representer theorem is given by Baldassarre et al. (2012) in the context of finite-dimensional multitask learning: this approach works in infinite dimensions as well. We give the closed form solution based on Baldassarre et al. (2012) below (we include the proof in Appendix D.1).

Proposition 1 (Representer theorem for general spectral filter). Let $(\mathbf{K})_{ij} = k(x_i, x_j), 1 \le i, j \le n$ denote the Gram matrix associated to the scalar-valued kernel k. We have

$$\hat{F}_{\lambda}(x) = \sum_{i=1}^{n} y_i \alpha_i(x), \qquad \alpha(x) = \frac{1}{n} g_{\lambda}\left(\frac{\mathbf{K}}{n}\right) \mathbf{k}_x \in \mathbb{R}^n, \qquad (\mathbf{k}_x)_i = k(x, x_i), \quad 1 \le i \le n.$$
(12)

Example 1 (Conditional integration). Consider now a random variable Z taking values in a topological space Z on which we define a second RKHS $\mathcal{H}' \subseteq \mathbb{R}^{Z}$ with kernel $\ell : Z \times Z \to \mathbb{R}$ and canonical feature map $\psi : Z \to \mathcal{H}', z \mapsto \ell(z, \cdot)$. The conditional mean embedding (Song et al., 2009; Grünewälder et al., 2012a) is defined as

$$F_*(x) \coloneqq \mathbb{E}[\psi(Z) \mid X = x], \qquad x \in \mathcal{X}.$$

We immediately see the link with vector-valued regression with $Y = \psi(Z)$ and $\mathcal{Y} = \mathcal{H}'$. The conditional mean embedding allows us to compute the conditional expectation of any element of \mathcal{H}' . Indeed, using the reproducing property, for $f \in \mathcal{H}'$, we have for all $x \in \mathcal{X}$,

$$\mathbb{E}[f(Z) \mid X = x] = \langle f, \mathbb{E}[\psi(Z) \mid X = x] \rangle_{\mathcal{H}'}.$$

 $[\]overline{{}^{4}g_{\lambda}(C_X)}$ is defined with the rules of spectral calculus, see Definition 9 in the Appendix.

Given a dataset $\{(x_i, z_i)\}_{i=1}^{n-5}$ and an estimate of the conditional mean embedding F_* with a spectral algorithm \hat{F}_{λ} as in Eq. (11), and substituting the formula in Eq. (12), we obtain $\mathbb{E}[f(Z) \mid X = x] \approx \langle f, \hat{F}_{\lambda}(x) \rangle_{\mathcal{H}'} = \sum_{i=1}^{n} \langle f, \psi(z_i) \rangle_{\mathcal{H}'} \alpha_i(x) = \mathbf{f}_z^\top \alpha(x)$, where $(\mathbf{f}_z)_i = f(z_i), 1 \leq i \leq n$.

Learning rates: We now give our main result, the learning rates for the difference between $[\hat{F}_{\lambda}]$ and F_* in the interpolation norm, where F_{λ} and \hat{F}_{λ} are given by (10) and (11) based on a general spectral filter satisfying Definition 2. The proof is deferred to Section C in the Appendix.

Theorem 4 (Upper learning rates). Let \hat{F}_{λ} be an estimator based on a general spectral filter with qualification $\rho \ge 0$. Furthermore, let the conditions (EVD), (EMB), (MOM) be satisfied with $0 . With <math>0 \le \gamma \le 1$, if (SRC) is satisfied with $\gamma < \beta \le 2\rho$, we have

1. in the case $\beta + p \leq \alpha$, let $\lambda_n = \Theta\left(\left(n/\log^{\theta}(n)\right)^{-\frac{1}{\alpha}}\right)$ for some $\theta > 1$, for all $\tau > \log(6)$ and sufficiently large $n \geq 1$, there is a constant J > 0 independent of n and τ such that

$$\left\| \left[\hat{F}_{\lambda_n} \right] - F_* \right\|_{\gamma}^2 \le \tau^2 J \left(\frac{n}{\log^{\theta} n} \right)^{-\frac{\beta - \gamma}{\alpha}}$$

is satisfied with P^n -probability not less than $1 - 6e^{-\tau}$.

2. in the case $\beta + p > \alpha$, let $\lambda_n = \Theta\left(n^{-\frac{1}{\beta+p}}\right)$, for all $\tau > \log(6)$ and sufficiently large $n \ge 1$, there is a constant J > 0 independent of n and τ such that

$$\left\| \left[\hat{F}_{\lambda_n} \right] - F_* \right\|_{\gamma}^2 \le \tau^2 J n^{-\frac{\beta - \gamma}{\beta + p}}$$

is satisfied with P^n -probability not less than $1 - 6e^{-\tau}$.

Theorem 4 provides the upper rate for vector-valued spectral algorithms. In particular, in combination with the lower bound in Theorem 2, we see that vector-valued spectral algorithms with qualification ρ achieve an optimal learning rate when the smoothness β of the regression function is in the range $(\alpha - p, 2\rho]$. For algorithms with infinite ρ such as gradient descent and principal component regression, we confirm that they can exploit smoothness of the target function just as in the real-valued setting (Bauer et al., 2007; Blanchard and Mücke, 2018; Li et al., 2022), while not suffering from saturation. For Tikhonov regularization, where $\rho = 1$, the rates recover the state-of-the-art results from Li et al. (2023b). Finally, we point out that obtaining minimax optimal learning rates for $\beta < \alpha - p$ still remains challenging even in the real-valued output scenario. Note however that for a large variety of RKHS, α is arbitrarily close to p and we obtain optimal rates for the whole range $(0, 2\rho]$: we refer to Li et al. (2023b); Zhang et al. (2023b) for a detailed discussion.

References

Jean-Pierre Aubin. Applied Functional Analysis. John Wiley & Sons, Inc., 2nd edition, 2000.

- Luca Baldassarre, Lorenzo Rosasco, Annalisa Barla, and Alessandro Verri. Multi-output learning via spectral filtering. *Machine Learning*, 87(3):259–301, 2012.
- Frank Bauer, Sergei Pereverzev, and Lorenzo Rosasco. On regularization algorithms in learning theory. Journal of Complexity, 23(1):52–72, 2007.

⁵Note that this induces a dataset $D = \{(x_i, \psi(z_i))\}_{i=1}^n$ where we identify $y_i = \psi(z_i)$.

- Alain Berlinet and Christine Thomas-Agnan. Reproducing Kernel Hilbert Spaces in Probability and Statistics. Springer, 2011.
- Gilles Blanchard and Nicole Mücke. Optimal rates for regularization of statistical inverse learning problems. *Foundations of Computational Mathematics*, 18(4):971–1013, 2018.
- A. Caponnetto and E. De Vito. Optimal rates for the regularized least-squares algorithm. Foundations of Computational Mathematics, 7(3):331–368, 2007a.
- Andrea Caponnetto and Ernesto De Vito. Optimal rates for the regularized least-squares algorithm. Foundations of Computational Mathematics, 7(3):331–368, 2007b.
- Andrea Caponnetto and Yuan Yao. Cross-validation based adaptation for regularization operators in learning theory. *Analysis and Applications*, 8(02):161–183, 2010.
- Andrea Caponnetto, Charles A. Micchelli, Massimiliano Pontil, and Yiming Ying. Universal multi-task kernels. Journal of Machine Learning Research, 9:1615–1646, 2008. ISSN 1532-4435.
- Claudio Carmeli, Ernesto De Vito, and Alessandro Toigo. Vector valued reproducing kernel Hilbert spaces of integrable functions and Mercer theorem. *Analysis and Applications*, 4(04):377–408, 2006.
- Claudio Carmeli, Ernesto De Vito, Alessandro Toigo, and Veronica Umanitá. Vector valued reproducing kernel Hilbert spaces and universality. *Analysis and Applications*, 8(01):19–61, 2010.
- Carlo Ciliberto, Lorenzo Rosasco, and Alessandro Rudi. A consistent regularization approach for structured prediction. Advances in Neural Information Processing Systems, 29, 2016.
- Carlo Ciliberto, Lorenzo Rosasco, and Alessandro Rudi. A general framework for consistent structured prediction with implicit loss embeddings. *The Journal of Machine Learning Research*, 21(1):3852–3918, 2020.
- Ernesto De Vito, Lorenzo Rosasco, and Andrea Caponnetto. Discretization error analysis for tikhonov regularization. *Analysis and Applications*, 4(01):81–99, 2006.
- Joe Diestel and J.J. Uhl. Vector Measures. American Mathematical Society, 1977.
- R.M. Dudley. Real Analysis and Probability. Cambridge University Press, 2nd edition edition, 2002.
- Heinz Werner Engl, Martin Hanke, and A. Neubauer. Regularization of Inverse Problems. Kluwer, 2000.
- Simon Fischer and Ingo Steinwart. Sobolev norm learning rates for regularized least-squares algorithms. Journal Of Machine Learning Research, 21:205–1, 2020.
- Jun Ichi Fujii, Masatoshi Fujii, Takayuki Furuta, and Ritsuo Nakamoto. Norm inequalities equivalent to heinz inequality. Proceedings of the American Mathematical Society, 118(3):827–830, 1993.
- L Lo Gerfo, Lorenzo Rosasco, Francesca Odone, E De Vito, and Alessandro Verri. Spectral algorithms for supervised learning. *Neural Computation*, 20(7):1873–1897, 2008a.
- Lorenzo Lo Gerfo, Lorenzo Rosasco, Francesca Odone, Ernesto De Vito, and Alessandro Verri. Spectral algorithms for supervised learning. *Neural Computation*, 20:1873–1897, 2008b.
- Steffen Grünewälder, Guy Lever, Luca Baldassarre, Sam Patterson, Arthur Gretton, and Massimilano Pontil. Conditional mean embeddings as regressors. In *Proceedings of the 29th International Coference on International Conference on Machine Learning*, pages 1803—1810, 2012a.
- Steffen Grünewälder, Guy Lever, Luca Baldassarre, Massimilano Pontil, and Arthur Gretton. Modelling transition dynamics in mdps with rkhs embeddings. In *Proceedings of the 29th International Conference* on Machine Learning, pages 535–542, 2012b.

- Terry Herdman, Ruben D Spies, and Karina G Temperini. Global saturation of regularization methods for inverse ill-posed problems. *Journal of optimization theory and applications*, 148(1):164–196, 2011.
- Hachem Kadri, Emmanuel Duflos, Philippe Preux, Stéphane Canu, Alain Rakotomamonjy, and Julien Audiffren. Operator-valued kernels for learning from functional response data. *Journal of Machine Learning Research*, 17(20):1–54, 2016.
- Vladimir Kostic, Pietro Novelli, Andreas Maurer, Carlo Ciliberto, Lorenzo Rosasco, and Massimiliano Pontil. Learning dynamical systems via Koopman operator regression in reproducing kernel Hilbert spaces. Advances in Neural Information Processing Systems, 35:4017–4031, 2022.
- Vladimir Kostic, Karim Lounici, Pietro Novelli, and Massimiliano Pontil. Koopman operator learning: Sharp spectral rates and spurious eigenvalues. *arXiv preprint arXiv:2302.02004*, 2023.
- Yicheng Li, Haobo Zhang, and Qian Lin. On the saturation effect of kernel ridge regression. In *The Eleventh International Conference on Learning Representations*, 2023a.
- Zhu Li, Dimitri Meunier, Mattes Mollenhauer, and Arthur Gretton. Optimal rates for regularized conditional mean embedding learning. In Advances in Neural Information Processing Systems, volume 35, pages 4433–4445, 2022.
- Zhu Li, Dimitri Meunier, Mattes Mollenhauer, and Arthur Gretton. Towards optimal Sobolev norm rates for the vector-valued regularized least-squares algorithm. arXiv preprint arXiv:2312.07186, 2023b.
- Junhong Lin and Volkan Cevher. Optimal distributed learning with multi-pass stochastic gradient methods. In *International Conference on Machine Learning*, pages 3092–3101. PMLR, 2018.
- Junhong Lin and Volkan Cevher. Optimal convergence for distributed learning with stochastic gradient methods and spectral algorithms. *Journal of Machine Learning Research*, 21(147):1–63, 2020.
- Junhong Lin, Alessandro Rudi, Lorenzo Rosasco, and Volkan Cevher. Optimal rates for spectral algorithms with least-squares regression over Hilbert spaces. *Applied and Computational Harmonic Analysis*, 48(3):868–890, 2020a.
- Junhong Lin, Alessandro Rudi, Lorenzo Rosasco, and Volkan Cevher. Optimal rates for spectral algorithms with least-squares regression over Hilbert spaces. *Applied and Computational Harmonic Analysis*, 48(3):868–890, 2020b.
- Afsaneh Mastouri, Yuchen Zhu, Limor Gultchin, Anna Korba, Ricardo Silva, Matt Kusner, Arthur Gretton, and Krikamol Muandet. Proximal causal learning with kernels: Two-stage estimation and moment restriction. In *International Conference on Machine Mearning*, pages 7512–7523. PMLR, 2021.
- Peter Mathé. Saturation of regularization methods for linear ill-posed problems in hilbert spaces. **SIAM** *journal on numerical analysis*, 42(3):968–973, 2004.
- Mattes Mollenhauer and Péter Koltai. Nonparametric approximation of conditional expectation operators. arXiv preprint arXiv:2012.12917, 2020.
- Mattes Mollenhauer, Nicole Mücke, and T.J. Sullivan. Learning linear operators: Infinite-dimensional regression as a well-behaved non-compact inverse problem. *arXiv preprint arXiv:2211.08875*, 2022.
- Nicole Mücke, Gergely Neu, and Lorenzo Rosasco. Beating SGD saturation with tail-averaging and minibatching. In *Advances in Neural Information Processing Systems*, volume 32, 2019.
- Andreas Neubauer. On converse and saturation results for Tikhonov regularization of linear ill-posed problems. SIAM Journal On Numerical Analysis, 34(2):517–527, 1997.
- Junhyung Park and Krikamol Muandet. A measure-theoretic approach to kernel conditional mean embeddings. Advances in Neural Information Processing Systems, 33:21247–21259, 2020.

- Rahul Singh, Maneesh Sahani, and Arthur Gretton. Kernel instrumental variable regression. Advances in Neural Information Processing Systems, 32, 2019.
- Rahul Singh, Liyuan Xu, and Arthur Gretton. Kernel methods for causal functions: dose, heterogeneous and incremental response curves. *Biometrika*, page asad042, 07 2023. ISSN 1464-3510.
- Steve Smale and Ding-Xuan Zhou. Shannon sampling and function reconstruction from point values. *Bulletin* of the American Mathematical Society, 41(3):279–305, 2004.
- Steve Smale and Ding-Xuan Zhou. Shannon sampling II: Connections to learning theory. *Applied and Computational Harmonic Analysis*, 19(3):285–302, 2005.
- Steve Smale and Ding-Xuan Zhou. Learning theory estimates via integral operators and their approximations. *Constructive Approximation*, 26(2):153–172, 2007.
- Le Song, Jonathan Huang, Alex Smola, and Kenji Fukumizu. Hilbert space embeddings of conditional distributions with applications to dynamical systems. In *Proceedings of the 26th Annual International Conference on Machine Learning*, pages 961–968. ACM, 2009.
- Bharath K Sriperumbudur, Kenji Fukumizu, and Gert RG Lanckriet. Universality, characteristic kernels and RKHS embedding of measures. *Journal of Machine Learning Research*, 12(Jul):2389–2410, 2011.
- Ingo Steinwart and Andreas Christmann. Support Vector Machines. Springer, 2008.
- Ingo Steinwart and Clint Scovel. Mercer's theorem on general domains: On the interaction between measures, kernels, and RKHSs. *Constructive Approximation*, 35(3):363–417, 2012.
- Ingo Steinwart, Don R Hush, Clint Scovel, et al. Optimal rates for regularized least squares regression. In *COLT*, pages 79–93, 2009.
- Joachim Weidmann. Linear Operators in Hilbert Spaces. Springer, 1980.
- Yuan Yao, Lorenzo Rosasco, and Andrea Caponnetto. On early stopping in gradient descent learning. Constructive Approximation, 26(2):289–315, 2007.
- Haobo Zhang, Yicheng Li, Weihao Lu, and Qian Lin. On the optimality of misspecified kernel ridge regression. arXiv preprint arXiv:2305.07241, 2023a.
- Haobo Zhang, Yicheng Li, Weihao Lu, and Qian Lin. On the optimality of misspecified spectral algorithms. arXiv preprint arXiv:2303.14942, 2023b.

Appendices

The appendix is organized as follows. In Section A, we give additional mathematical background and notations. In Section B, we give the proof of Theorem 3 and provide a technical comparison of our proof with Li et al. (2023a). In Section C, we prove Theorem 4. Finally, in Section D, we provide auxiliary results used in the main proofs.

A Additional Background

A.1 Hilbert spaces and linear operators

Definition 3 (Bochner L_q -spaces, Diestel and Uhl (1977)). Let H be a separable Hilbert space and π a probability measure on \mathcal{X} . For $1 \leq q \leq \infty$, $L_q(\mathcal{X}, \mathcal{F}_{\mathcal{X}}, \pi; H)$, abbreviated $L_q(\pi; H)$, is the space of strongly $\mathcal{F}_{\mathcal{X}} - \mathcal{F}_H$ measurable and Bochner q-integrable functions from \mathcal{X} to H, with the norms

$$\|f\|_{L_q(\pi;H)}^q = \int_{\mathcal{X}} \|f\|_H^q \,\mathrm{d}\pi, \quad 1 \le q < \infty, \qquad \|f\|_{L_\infty(\pi;H)} = \inf\left\{C \ge 0 : \pi\{\|f\|_H > C\} = 0\right\}.$$

Definition 4 (*p*-Schatten class, e.g. Weidmann (1980)). Let H, H' be separable Hilbert spaces. For $1 \le q \le \infty$, $S_p(H, H')$, abbreviated $S_p(H)$ if H = H', is the Banach space of all compact operators C from H to H' such that $\|C\|_{S_p(H,H')} \coloneqq \|(\sigma_i(C))_{i\in I}\|_{\ell_p}$ is finite. Here $\|(\sigma_i(C))_{i\in I}\|_{\ell_p}$ is the ℓ_p -sequence space norm of the sequence of the strictly positive singular values of C indexed by the at most countable set I. For p = 2, we retrieve the space of Hilbert-Schmidt operators, for p = 1 we retrieve the space of Trace Class operators, and for $p = +\infty$, $\|\cdot\|_{S_\infty(H,H')}$ corresponds to the operator norm $\|\cdot\|_{H\to H'}$.

Definition 5 (Tensor Product of Hilbert Spaces, Aubin (2000)). Let H, H' be Hilbert spaces. The Hilbert space $H \otimes H'$ is the completion of the algebraic tensor product with respect to the norm induced by the inner product $\langle x_1 \otimes x'_1, x_2 \otimes x'_2 \rangle_{H \otimes H'} = \langle x_1, x_2 \rangle_H \langle x'_1, x'_2 \rangle_{H'}$ for $x_1, x_2 \in H$ and $x'_1, x'_2 \in H'$ defined on the elementary tensors of $H \otimes H'$. This definition extends to span $\{x \otimes x' | x \in H, x' \in H'\}$ and finally to its completion. The space $H \otimes H'$ is separable whenever both H and H' are separable. If $\{e_i\}_{i \in I}$ and $\{e'_j\}_{j \in J}$ are orthonormal basis in H and H', $\{e_i \otimes e'_j\}_{i \in I, j \in J}$ is an orthonormal basis in $H \otimes H'$.

Theorem 5 (Isometric Isomorphism between $L_2(\pi; \mathcal{Y})$ and $S_2(L_2(\pi), \mathcal{Y})$, Theorem 12.6.1 Aubin (2000)). Let H be a separable Hilbert space. The Bochner space $L_2(\pi; H)$ is isometrically isomorphic to $S_2(L_2(\pi), \mathcal{Y})$ and the isometric isomorphism is realized by the map $\Psi: S_2(L_2(\pi), \mathcal{Y}) \to L_2(\pi; H)$ acting on elementary tensors as $\Psi(f \otimes y) = (\omega \to f(\omega)y)$.

A.2 RKHS embbedings into L₂ and Well-specifiedness

Recall that $I_{\pi} : \mathcal{H} \to L_2(\pi)$ is the embedding that maps every function in \mathcal{H} into its π -equivalence class in $L_2(\pi)$ and that we used the shorthand notation $[f] = I_{\pi}(f)$ for all $f \in \mathcal{H}$. We define similarly $\mathcal{I}_{\pi} : \mathcal{G} \to L_2(\pi; \mathcal{Y})$ as the embedding that maps every function in \mathcal{G} into its π -equivalence class in $L_2(\pi; \mathcal{Y})$.

Definition 6 (Embedding \mathcal{G} into $L_2(\pi; \mathcal{Y})$). Let $\mathcal{I}_{\pi} \coloneqq I_{\mathcal{Y}} \otimes I_{\pi}$ be the tensor product of the operator $\mathrm{Id}_{\mathcal{Y}}$ with the operator I_{π} (see Aubin (2000, Definition 12.4.1.) for the definition of tensor product of operators). \mathcal{I}_{π} maps every function in \mathcal{G} into its π -equivalence class in $L_2(\pi; \mathcal{Y})$. We then use the shorthand notation $[F] = \mathcal{I}_{\pi}(F)$ for all $F \in \mathcal{G}$.

Remark 5. Let $\{d_j\}_{j\in J}$ be an orthonormal basis of \mathcal{Y} and recall that $\{\sqrt{\mu_i}[e_i]\}_{i\in I}$ forms an orthonormal basis of $[\mathcal{H}]^1$. Let $F \in [\mathcal{G}]^1$. Then F can be represented as the element $C \coloneqq \sum_{i\in I, j\in J} a_{ij}d_j \otimes \sqrt{\mu_i}[e_i]$ in $S_2([\mathcal{H}]^1, \mathcal{Y})$

by definition of $[\mathcal{G}]^1$ with $\|C\|_1^2 = \sum_{i,j} a_{ij}^2$. Hence defining $\overline{F} := \sum_{i \in I, j \in J} a_{ij} d_j \otimes \sqrt{\mu_i} e_i$ we have $C = \overline{C} \pi - a.e.$ and

$$\|\bar{C}\|_{\mathcal{G}}^{2} = \sum_{i \in I, j \in J} a_{i,j}^{2} = \|C\|_{1}^{2} < +\infty$$

Taking the elements identifying \overline{C} in \mathcal{G} gives a representer \overline{F} of F in \mathcal{G} .

A.3 Additional Notations

In the following, we fix $\{d_j\}_{j\in J}$ an orthonormal basis of \mathcal{Y} , where J is at most countable. Recall that $\{\mu_i^{1/2}e_i\}_{i\in I}$ is an ONB of $(\ker I_\pi)^{\perp}$ in \mathcal{H} , and $\{[e_i]\}_{i\in I}$ is an ONB of $\overline{\operatorname{ran} I_\pi}$ in $L_2(\pi)$. Let $\{\tilde{e}_i\}_{i\in I'}$ be an ONB of $\ker I_\pi$ (with $I \cap I' = \emptyset$), then $\{\mu_i^{1/2}e_i\}_{i\in I} \cup \{\tilde{e}_i\}_{i\in I'}$ forms an ONB of \mathcal{H} , and $\{d_j \otimes \mu_i^{1/2}e_i\}_{i\in I, j\in J} \cup \{d_j \otimes \tilde{e}_i\}_{i\in I', j\in J}$ forms an ONB of $\mathcal{Y} \otimes \mathcal{H} \simeq \mathcal{G}$.

For any Hilbert space H, linear operator $T: H \to H$ and scalar $\lambda > 0$, we define $T_{\lambda} := T + \lambda I_H$.

B Saturation Effect with Tikhonov Regularization

In the following proofs a quantity $h_n \ge 0$ depending on $n \ge 1$, but independent of τ the confidence level, is equal to o(1) if $h_n \to 0$ when $n \to +\infty$.

We will make extensive use of the following notation in the subsequent analysis.

Definition 7 (Empirical $L_2(\pi)$ -norm). Denoted by $\langle \cdot, \cdot \rangle_{2,n}$, the empirical $L_2(\pi)$ -norm associated to points $\{x_i\}_{i=1}^n$ independently and identically sampled from the distribution of X, is defined as, for any $f, g \in \mathcal{H}$,

$$\langle f,g\rangle_{2,n} \coloneqq \left\langle \hat{C}_X, f\otimes g \right\rangle_{S_2(\mathcal{H})} = \left\langle \hat{C}_Xf,g \right\rangle_{\mathcal{H}} = \left\langle \hat{C}_X^{\frac{1}{2}}f, \hat{C}_X^{\frac{1}{2}}g \right\rangle_{\mathcal{H}} = \frac{1}{n}\sum_{i=1}^n f(x_i)g(x_i).$$

This induces an inner product on \mathcal{H} , with associated norm,

$$||f||_{2,n}^2 = \langle f, f \rangle_{2,n} = \frac{1}{n} \sum_{i=1}^n f(x_i)^2.$$

Definition 8. Fix $x \in \mathcal{X}$. The regularized canonical feature map is defined as

$$f_x(\cdot) = C_{X,\lambda}^{-1} k(x, \cdot) : \mathcal{X} \to \mathcal{H}.$$

Recall from Eq. (11) that the ridge estimator \hat{F}_{λ} defined in Eq. (7) can be expressed as

$$\hat{C}_{\lambda} = \hat{C}_{YX} g_{\lambda}(\hat{C}_X), \qquad \hat{F}_{\lambda}(\cdot) = \hat{C}_{\lambda} \phi(\cdot) \in \mathcal{G},$$

where in Theorem 3 we focus on Tikhonov regularization where $g_{\lambda}(x) = (x + \lambda)^{-1}$. In that setting we have

$$r_{\lambda}(x) \coloneqq 1 - \frac{x}{x+\lambda} = -\frac{\lambda}{x+\lambda}.$$
(13)

Proof of Theorem 3. Since $\beta \geq 2$, $F_* \in [\mathcal{G}]^{\beta} \subseteq [\mathcal{G}]^1$, therefore F_* has a representer \overline{F} in \mathcal{G} such that $F_* = \overline{F} \pi$ -a.e., and by Theorem 1, $\overline{F}(\cdot) = \overline{C}\phi(\cdot)$, with $\overline{C} \in S_2(\mathcal{H}, \mathcal{Y})$. Define the errors $\epsilon_i \coloneqq y_i - \overline{C}\phi(x_i)$, $i = 1, \ldots, n$,

that are i.i.d samples with the same distribution as $\epsilon \coloneqq Y - \overline{C}\phi(X)$. By assumption $\mathbb{E}\left[\|\epsilon\|_{\mathcal{Y}}^2 \mid X\right] \ge \sigma^2$ and by definition $\mathbb{E}\left[\epsilon \mid X\right] = 0$. By Eq. (13), we have

$$r_{\lambda}\left(\hat{C}_{X}\right) \coloneqq I - \hat{C}_{X}\hat{C}_{X,\lambda}^{-1} = -\lambda\hat{C}_{X}.$$

The following **bias-variance decomposition** is the essence of the proof. In the following derivation we abbreviate $S_2(L_2(\pi), \mathcal{Y})$ to S_2 $L_2(\pi; \mathcal{Y})$ to L_2 and x_1, \ldots, x_n to \underline{x}_n to save space.

$$\begin{split} \mathbb{E}\left[\left\|\left[\hat{F}_{\lambda}\right]-F_{*}\right\|_{L_{2}}^{2}\mid\underline{x}_{n}\right] &= \mathbb{E}\left[\left\|\left[\left(\hat{C}_{YX}\hat{C}_{X,\lambda}^{-1}-\bar{C}\right]\right\|_{S_{2}}^{2}\mid\underline{x}_{n}\right]\right] \\ &= \mathbb{E}\left[\left\|\left[\left(\frac{1}{n}\sum_{i=1}^{n}y_{i}\otimes\phi(x_{i})\right)\hat{C}_{X,\lambda}^{-1}-\bar{C}\right]\right\|_{S_{2}}^{2}\mid\underline{x}_{n}\right] \\ &= \mathbb{E}\left[\left\|\left[\frac{1}{n}\sum_{i=1}^{n}\left(\bar{C}\phi(x_{i})+\epsilon_{i}\right)\otimes\phi(x_{i})\hat{C}_{X,\lambda}^{-1}-\bar{C}\right]\right\|_{S_{2}}^{2}\mid\underline{x}_{n}\right] \\ &= \mathbb{E}\left[\left\|\left[-\bar{C}r_{\lambda}\left(\hat{C}_{X}\right)+\frac{1}{n}\sum_{i=1}^{n}\epsilon_{i}\otimes\left(\hat{C}_{X,\lambda}^{-1}\phi(x_{i})\right)\right]\right\|_{S_{2}}^{2}\mid\underline{x}_{n}\right] \\ &= \left\|\left[\bar{C}r_{\lambda}\left(\hat{C}_{X}\right)\right]\right\|_{S_{2}}^{2}+\frac{1}{n^{2}}\sum_{i=1}^{n}\mathbb{E}\left[\left\|\epsilon_{i}\right\|_{\mathcal{V}}^{2}\midx_{i}\right]\left\|\left[\hat{C}_{X,\lambda}^{-1}\phi(x_{i})\right]\right\|_{L_{2}(\pi)}^{2} \\ &\geq \lambda^{2}\left\|\left[\bar{C}\hat{C}_{X,\lambda}^{-1}\right]\right\|_{S_{2}}^{2}+\frac{\sigma^{2}}{n^{2}}\sum_{i=1}^{n}\left\|\left[\hat{C}_{X,\lambda}^{-1}\phi(x_{i})\right]\right\|_{L_{2}(\pi)}^{2}. \end{split}$$

The second term is a lower bound on the *variance* while the first term is a lower bound on the *bias*.

Bounding the Bias term. The idea is to first show that the population analogue of $\|[\bar{C}\hat{C}_{X,\lambda}^{-1}]\|_{S_2(L_2(\pi),\mathcal{Y})}^2$ can be bounded below by a non-zero constant. We can then bound the difference between the empirical and population version of $\|[\bar{C}\hat{C}_{X,\lambda}^{-1}]\|_{S_2(L_2(\pi),\mathcal{Y})}^2$ using a concentration inequality. By Lemma 1, for $\lambda > 0$, there is a constant c > 0 (see Lemma 1 for the exact value of c) such that

$$\| \left[\bar{C} C_{X,\lambda}^{-1} \right] \|_{S_2(L_2(\pi),\mathcal{Y})}^2 \ge c > 0.$$

Furthermore by Lemma 2, there is a constant $c_0 > 0$ (see Lemma 2 for the exact value of c_0) such that for any $\tau \ge \log(4)$, with probability at least $1 - 4e^{-\tau}$, for $n \ge (c_0 \tau)^{(4+2p)}$ and $1 \ge \lambda \ge n^{-\frac{1}{2+p}}$, we have

$$\left\| \left[\bar{C}C_{X,\lambda}^{-1} \right] \right\|_{S_2(L_2(\pi),\mathcal{Y})}^2 - \left\| \left[\bar{C}\hat{C}_{X,\lambda}^{-1} \right] \right\|_{S_2(L_2(\pi),\mathcal{Y})}^2 \right\| = \tau^2 o(1).$$

Therefore, under the same high probability,

$$\| [\bar{C}\hat{C}_{X,\lambda}^{-1}] \|_{S_2(L_2(\pi),\mathcal{Y})}^2 \ge c - \tau^2 o(1).$$

It leads to our final bound on the bias term, for a constant $\rho_2 \ge 0$ and for sufficiently large $n \ge 1$, where the hidden index bound depends on τ , we have

$$\lambda^{2} \left\| \left[\bar{C} \hat{C}_{X,\lambda}^{-1} \right] \right\|_{S_{2}(L_{2}(\pi),\mathcal{Y})}^{2} \ge \rho_{1} \lambda^{2}.$$
(14)

Bounding the Variance Term. Using the norm from Definition 7, we have the following chain of identities.

$$\frac{\sigma^2}{n^2} \sum_{i=1}^n \left\| \left[\hat{C}_{X,\lambda}^{-1} \phi(x_i) \right] \right\|_{L_2(\pi)}^2 = \frac{\sigma^2}{n^2} \sum_{i=1}^n \int_{\mathcal{X}} \left\langle \phi(X), \hat{C}_{X,\lambda}^{-1} \phi(x_i) \right\rangle_{\mathcal{H}}^2 d\pi(x) = \frac{\sigma^2}{n} \int_{\mathcal{X}} \| \hat{C}_{X,\lambda}^{-1} \phi(X) \|_{2,n}^2 d\pi(x).$$
(15)

Therefore it suffices to consider $\int_{\mathcal{X}} \|\hat{C}_{X,\lambda}^{-1}k(x,\cdot)\|_{2,n}^2 d\pi(x).$

Combining the result of Lemma 4 and Lemma 5 we obtain that for $1 \ge \lambda \ge n^{-\frac{1}{2+p}}$ with probability at least $1 - 6e^{-\tau}$, for $n \ge (c_0 \tau)^{4+2p}$, the following bounds hold simultaneously for all $x \in \mathcal{X}$:

$$\begin{split} \left\| \hat{C}_{X}^{\frac{1}{2}} \left(\hat{C}_{X,\lambda}^{-1} - C_{X,\lambda}^{-1} \right) k(x,\cdot) \right\|_{\mathcal{H}} &\leq \tau o(1) \\ \| [C_{X,\lambda}^{-1} k(x,\cdot)] \|_{2,n}^{2} - \frac{1}{2} \| [C_{X,\lambda}^{-1} k(x,\cdot)] \|_{L_{2}(\pi)}^{2} &\geq -\tau o(1) \\ \| [C_{X,\lambda}^{-1} k(x,\cdot)] \|_{2,n}^{2} - \frac{3}{2} \| [C_{X,\lambda}^{-1} k(x,\cdot)] \|_{L_{2}(\pi)}^{2} &\leq \tau o(1). \end{split}$$

Fix $x \in \mathcal{X}$. Using the algebraic identity $a^2 - b^2 = (a - b)(2b + (a - b))$, and recalling that by Definition 7,

$$||f||_{2,n}^2 = \left\|\hat{C}_X^{\frac{1}{2}}f\right\|_{\mathcal{H}}^2,$$

we deduce

$$\begin{aligned} & \left\| \left\| \hat{C}_{X}^{\frac{1}{2}} \hat{C}_{X,\lambda}^{-1} k(x,\cdot) \right\|_{\mathcal{H}}^{2} - \left\| \hat{C}_{X}^{\frac{1}{2}} C_{X,\lambda}^{-1} k(x,\cdot) \right\|_{\mathcal{H}}^{2} \right\| \\ & \leq \left\| \hat{C}_{X}^{\frac{1}{2}} \left(\hat{C}_{X,\lambda}^{-1} - C_{X,\lambda}^{-1} \right) k(x,\cdot) \right\|_{\mathcal{H}} \cdot \left(\left\| \hat{C}_{X}^{\frac{1}{2}} \left(\hat{C}_{X,\lambda}^{-1} - C_{X,\lambda}^{-1} \right) k(x,\cdot) \right\|_{\mathcal{H}} + 2 \left\| C_{X,\lambda}^{-1} k(x,\cdot) \right\|_{2,n} \right) \\ & \leq \tau o(1) \left(\tau o(1) + 2 \left\| C_{X,\lambda}^{-1} k(x,\cdot) \right\|_{2,n} \right). \end{aligned}$$

Using Definition 7 again, this reads

$$\left\|\hat{C}_{X,\lambda}^{-1}k(x,\cdot)\right\|_{2,n}^{2} \ge \left\|C_{X,\lambda}^{-1}k(x,\cdot)\right\|_{2,n}^{2} - \tau o(1)\left(\tau o(1) + 2\left\|C_{X,\lambda}^{-1}k(x,\cdot)\right\|_{2,n}\right).$$

We have

$$\|C_{X,\lambda}^{-1}k(x,\cdot)\|\|_{2,n}^2 \leq \frac{3}{2} \|[C_{X,\lambda}^{-1}k(x,\cdot)]\|_{L_2(\pi)}^2 + \tau o(1) \leq \left(\sqrt{1.5} \|[C_{X,\lambda}^{-1}k(x,\cdot)]\|_{L_2(\pi)} + \sqrt{\tau}o(1)\right)^2.$$

Hence,

$$\begin{split} \left\| \hat{C}_{X,\lambda}^{-1} k(x,\cdot) \right\|_{2,n}^{2} &\geq \left\| C_{X,\lambda}^{-1} k(x,\cdot) \right\|_{2,n}^{2} - \tau o(1) \left(\left\| \left[C_{X,\lambda}^{-1} k(x,\cdot) \right] \right\|_{L_{2}(\pi)} + \sqrt{\tau} o(1) + \tau o(1) \right) \\ &\geq \frac{1}{2} \left\| \left[C_{X,\lambda}^{-1} k(x,\cdot) \right] \right\|_{L_{2}(\pi)}^{2} - \tau o(1) \\ &- \tau o(1) \left(\left\| \left[C_{X,\lambda}^{-1} k(x,\cdot) \right] \right\|_{L_{2}(\pi)} + \sqrt{\tau} o(1) + \tau o(1) \right) \\ &\geq \frac{1}{2} \left\| \left[C_{X,\lambda}^{-1} k(x,\cdot) \right] \right\|_{L_{2}(\pi)}^{2} - \tau^{2} o(1) - \tau o(1) \left\| \left[C_{X,\lambda}^{-1} k(x,\cdot) \right] \right\|_{L_{2}(\pi)}. \end{split}$$

By Lemma 17,

$$\int_{\mathcal{X}} \left\| \left[C_{X,\lambda}^{-1} k(x,\cdot) \right] \right\|_{L_2(\pi)}^2 d\pi(x) = \mathcal{N}_2(\lambda).$$

Furthermore, by Jensen's inequality,

$$\int_{\mathcal{X}} \left\| \left[C_{X,\lambda}^{-1} k(x,\cdot) \right] \right\|_{L_2(\pi)}^2 \mathrm{d}\pi(x) \ge \left(\int_{\mathcal{X}} \left\| \left[C_{X,\lambda}^{-1} k(x,\cdot) \right] \right\|_{L_2(\pi)} \mathrm{d}\pi(x) \right)^2.$$

Recall from Lemma 16 that

$$c_{1,2}\lambda^{-p} \leq \mathcal{N}_2(\lambda) \leq c_{2,2}\lambda^{-p}.$$

Therefore we have

$$\int_{\mathcal{X}} \| [C_{X,\lambda}^{-1} k(x,\cdot)] \|_{L_{2}(\pi)} \mathrm{d}\pi(x) \leq \sqrt{c_{2,2}} \lambda^{-\frac{p}{2}} \\ \int_{\mathcal{X}} \| [C_{X,\lambda}^{-1} k(x,\cdot)] \|_{L_{2}(\pi)}^{2} \mathrm{d}\pi(x) \geq c_{1,2} \lambda^{-p}.$$

Hence

$$\int_{\mathcal{X}} \|\hat{C}_{X,\lambda}^{-1}k(x,\cdot)\|_{2,n}^{2} d\pi(x) \ge \frac{c_{1,2}}{2}\lambda^{-p} - \tau^{2}o(1) - \tau o(1)\sqrt{c_{2,2}}\lambda^{-\frac{p}{2}} \\ \ge \left(\frac{c_{1,2}}{2} - \tau o(1)\sqrt{c_{2,2}}\right)\lambda^{-p} - \tau^{2}o(1)$$

Combined with Eq. (15), it leads to our final bound on the variance term, for a constant $\rho_2 \ge 0$ and for sufficiently large $n \ge 1$, where the hidden index bound depends on τ , we have

$$\frac{\sigma^2}{n^2} \sum_{i=1}^n \left\| \left[\hat{C}_{X,\lambda}^{-1} \phi(x_i) \right] \right\|_{L_2(\pi)}^2 \ge \frac{\rho_2}{n\lambda^p}.$$
(16)

Putting it together. We are now ready to assemble the lower bounds on the variance and on the bias. For a fixed confidence parameter $\tau \ge \log(10)$, for sufficiently large n > 0, where the hidden index bound depends on τ , with probability at least $1 - 10e^{-\tau}$, we have by Eq. (14) and Eq. (16), that for $\lambda = \lambda(n)$ satisfying $1 \ge \lambda > n^{-\frac{1-q}{2+p}}$ for some $q \in (0, \frac{1}{2})$,

$$\mathbb{E}\left[\left\|\left[\hat{F}_{\lambda}\right]-F_{\star}\right\|^{2}_{L_{2}(\pi;\mathcal{Y})}\mid x_{1},\ldots,x_{n}\right]\geq\rho_{1}\lambda^{2}+\rho_{2}n^{-1}\lambda^{-p}$$

where ρ_1, ρ_2 have no dependence on n. Recall Young's inequality, for r, q > 1 satisfying $r^{-1} + q^{-1} = 1$, we have for all $a, b \ge 0$,

$$a + b \ge r^{\frac{1}{r}} q^{\frac{1}{q}} a^{\frac{1}{r}} b^{\frac{1}{q}}$$

We apply Young's inequality with $r^{-1} = p/(2+p)$ and $q^{-1} = 2/(2+p)$, there exists a constant $c_1 > 0$ such that

$$\rho_1 \lambda^2 + \rho_2 n^{-1} \lambda^{-p} \ge c_1 \left(\lambda^2\right)^{\frac{p}{2+p}} \left(\lambda^{-p} n^{-1}\right)^{\frac{2}{2+p}} = c_1 n^{-\frac{2}{2+p}}.$$

To conclude the proof, let $\lambda = \lambda(n)$ be an arbitrary choice of regularization parameter satisfying $\lambda(n) \to 0$. We have just covered the case $1 \ge \lambda \ge n^{-\frac{1}{2+p}}$ and the case $0 < \lambda \le n^{-\frac{1}{2+p}}$ is covered by (Li et al., 2023a, Section B.4).

Lemma 1. For any $\lambda \leq 1$ and $C \in S_2(\mathcal{H}, \mathcal{Y})$, with $C \neq S_2(\overline{\operatorname{ran} S_{\pi}}, \mathcal{Y})^6$, we have

$$\left\| \left[CC_{X,\lambda}^{-1} \right] \right\|_{S_2(L_2(\pi),\mathcal{Y})}^2 \ge \sum_{i \in I, j \in J} a_{ij}^2 \frac{\mu_i}{(\mu_i + 1)^2} > 0,$$

with $a_{ij} \coloneqq \langle d_j, C \sqrt{\mu_i} e_i \rangle_{\mathcal{Y}}, \ i \in I, j \in J.$

Proof. Define $\{a_{ij}\}_{i \in I \cap I', j \in J}$ such that $a_{ij} \coloneqq \langle d_j, C \sqrt{\mu_i} e_i \rangle_{\mathcal{Y}}$ for $i \in I, j \in J$ and $a_{ij} \coloneqq \langle d_j, C\tilde{e}_i \rangle_{\mathcal{Y}}$ for $i \in I', j \in J$. Then, on one hand, since $C \in S_2(\mathcal{H}, \mathcal{Y})$,

$$C = \sum_{i \in I, j \in J} a_{ij} d_j \otimes (\sqrt{\mu_i} e_i) + \sum_{i \in I', j \in J} a_{ij} d_j \otimes \tilde{e}_i.$$

On the other hand,

$$C_{X,\lambda}^{-1} = \sum_{i \in I} (\mu_i + \lambda)^{-1} (\sqrt{\mu_i} e_i) \otimes (\sqrt{\mu_i} e_i) + \lambda^{-1} \sum_{i \in I'} \tilde{e}_i \otimes \tilde{e}_i$$

 $^{^{6}}$ ± is the notation for "not being orthogonal to".

Therefore, noting that $\tilde{e}_i = 0 \pi$ -a.e. for all $i \in I'$, we have,

$$\begin{bmatrix} CC_{X,\lambda}^{-1} \end{bmatrix} = \left[\sum_{i \in I, j \in J} a_{ij} (\mu_i + \lambda)^{-1} d_j \otimes (\sqrt{\mu_i} e_i) + \sum_{i \in I', j \in J} \frac{a_{ij}}{\lambda} d_j \otimes \tilde{e}_i \right]$$
$$= \sum_{i \in I, j \in J} a_{ij} \frac{\sqrt{\mu_i}}{\mu_i + \lambda} d_j \otimes [e_i].$$

Therefore the $S_2(L_2(\pi), \mathcal{Y})$ -norm can be evaluated in closed form using Parseval's identity,

$$\left\| \left[CC_{X,\lambda}^{-1} \right] \right\|_{S_2(L_2(\pi),\mathcal{Y})}^2 = \sum_{i \in I, j \in J} a_{ij}^2 \frac{\mu_i}{(\mu_i + \lambda)^2} \ge \sum_{i \in I, j \in J} a_{ij}^2 \frac{\mu_i}{(\mu_i + 1)^2},$$

where we used that $\{d_j\}_{j\in J}$ is orthonormal in \mathcal{Y} , $\{[e_i\}_{i\in I})$ is orthonormal in $L_2(\pi)$, and $\lambda \leq 1$. The right hand side has no dependence on λ or n. Furthermore, under assumption (EVD+), $\mu_i > 0$ for all $i \in I$, therefore the right hand side term equals zero if and only if $a_{ij} = 0$ for all $i \in I, j \in J$. Since by assumption $C \neq S_2(\overline{\operatorname{ran} S_{\pi}}, \mathcal{Y})$, the right hand side is strictly positive.

Lemma 2. Suppose Assumption (EVD) holds with $p \in (0,1]$. Let $C \in S_2(\mathcal{H}, \mathcal{Y})$ such that $[C] \in S_2([\mathcal{H}]^2, \mathcal{Y})$. There is a constant $c_0 > 0$ such that for any $\delta \ge \log(4)$, with probability at least $1 - 4e^{-\tau}$, for $n \ge (c_0 \tau)^{(4+2p)}$ and $1 \ge \lambda \ge n^{-\frac{1}{2+p}}$, we have

$$\left\| \left\| \left[CC_{X\lambda}^{-1} \right] \right\|_{S_2(L_2(\pi),\mathcal{Y})}^2 - \left\| \left[C\hat{C}_{X\lambda}^{-1} \right] \right\|_{S_2(L_2(\pi),\mathcal{Y})}^2 \right\| \le \tau^2 o(1)$$

We have $c_0 \coloneqq 8\kappa \max\{\sqrt{c_{2,1}}, 1\}$ where $c_{2,1}$ is defined in Lemma 16.

Proof. Using the identity $A^{-1} - B^{-1} = A^{-1}(B - A)B^{-1}$, we obtain

$$C_{X,\lambda}^{-1} - \hat{C}_{X,\lambda}^{-1} = C_{X,\lambda}^{-1} (\hat{C}_X - C_X) \hat{C}_{X,\lambda}^{-1}$$

We apply Lemma 22 with $\gamma = 0$,

$$\begin{split} \left\| \left[C \left(C_{X,\lambda}^{-1} - \hat{C}_{X,\lambda}^{-1} \right) \right] \right\|_{S_{2}(L_{2}(\pi),\mathcal{Y})} &= \left\| \left[C \hat{C}_{X,\lambda}^{-1} (\hat{C}_{X} - C_{X}) C_{X,\lambda}^{-1} \right] \right\|_{S_{2}(L_{2}(\pi),\mathcal{Y})} \\ &= \left\| C \hat{C}_{X,\lambda}^{-1} (\hat{C}_{X} - C_{X}) C_{X,\lambda}^{-1} C_{X}^{\frac{1}{2}} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} \\ &\leq \left\| C \hat{C}_{X,\lambda}^{-\frac{1}{2}} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} \left\| \hat{C}_{X,\lambda}^{-\frac{1}{2}} C_{X,\lambda}^{\frac{1}{2}} \right\|_{\mathcal{H} \to \mathcal{H}} \\ &\cdot \left\| C_{X,\lambda}^{-\frac{1}{2}} (\hat{C}_{X} - C_{X}) C_{X,\lambda}^{-\frac{1}{2}} \right\|_{\mathcal{H} \to \mathcal{H}} \left\| C_{X,\lambda}^{-\frac{1}{2}} C_{X,\lambda}^{\frac{1}{2}} \right\|_{\mathcal{H} \to \mathcal{H}} \end{split}$$
(17)

We consider each of the four terms in line (17). The last term is bounded above by 1 and the first term is bounded above by $\lambda^{-\frac{1}{2}} \|C\|_{S_2(\mathcal{H},\mathcal{Y})}$. By Lemma 20 applied with s = 1/2, we have for the second term

$$\left\| \hat{C}_{X,\lambda}^{-\frac{1}{2}} C_{X,\lambda}^{\frac{1}{2}} \right\|_{\mathcal{H} \to \mathcal{H}} \leq \left\| \hat{C}_{X,\lambda}^{-1} C_{X,\lambda} \right\|_{\mathcal{H} \to \mathcal{H}}^{\frac{1}{2}}$$

Then, by Lemma 18, for $\tau \ge \log(2)$, with probability at least $1 - 2e^{-\tau}$, for $\sqrt{n\lambda} \ge 8\tau\kappa\sqrt{\max\{\mathcal{N}(\lambda),1\}}$, we have

$$\left\|\hat{C}_{X,\lambda}^{-1}C_{X,\lambda}\right\|_{\mathcal{H}\to\mathcal{H}} \leq 2.$$

Since $\mathcal{N}(\lambda) \leq c_{2,1}\lambda^{-p}$ by Lemma 16, and $\lambda \leq 1$, it suffices to verify that λ satisfies

$$\sqrt{n\lambda} \ge 8\tau\kappa \max\{\sqrt{c_{2,1}}, 1\}\lambda^{-\frac{p}{2}}.$$

Since $\lambda \ge n^{-\frac{1}{2+p}}$ by assumption, we deduce the sufficient condition $n \ge (\tau c_0)^{2(2+p)}$, where $c_0 \coloneqq 8\kappa \max\{\sqrt{c_{2,1}}, 1\}$. We bound the third term using Lemma 16 Lin and Cevher (2020). For $\tau \ge \log(2)$, with probability at least $1 - 2e^{-\tau}$, we have

$$\lambda^{-\frac{1}{2}} \left\| C_{X,\lambda}^{-\frac{1}{2}} (C_X - \hat{C}_X) C_{X,\lambda}^{-\frac{1}{2}} \right\|_{\mathcal{H} \to \mathcal{H}} \leq \frac{4\kappa^2 \xi_{\delta}}{3n\lambda^{\frac{3}{2}}} + \sqrt{\frac{2\kappa^2 \xi_{\delta}}{n\lambda^2}}$$

where we define

$$\xi_{\delta} \coloneqq \log \frac{2\kappa^2 (\mathcal{N}_1(\lambda) + 1)}{e^{-\tau} \|C_X\|_{\mathcal{H} \to \mathcal{H}}}.$$

By assumption $\lambda \ge n^{-\frac{1}{2+p}}$. We thus have

$$n\lambda^{\frac{3}{2}} \ge n^{\frac{1+2p}{4+2p}}$$
 and $n\lambda^{2} \ge n^{\frac{p}{2+p}}$.

On the other hand, since $1 \ge \lambda \ge n^{-\frac{1}{2+p}}$, using Lemma 16, we have

$$\xi_{\delta} \le \log \frac{82(c_{2,1}\lambda^{-p}+1)}{e^{-\tau} \|C_X\|_{\mathcal{H}\to\mathcal{H}}} \le \log \frac{2(c_{2,1}+1)n^{\frac{p}{2+p}}}{e^{-\tau} \|C_X\|_{\mathcal{H}\to\mathcal{H}}} \le \log \frac{2(c_{2,l}+1)}{e^{-\tau} \|C_X\|_{\mathcal{H}\to\mathcal{H}}} + \frac{p}{2+p}\log n.$$

The first term does not depend on n, and the second term is logarithmic in n. Putting everything together with a union bound, we get a bound on (17). With probability at least $1 - 4e^{-\tau}$, for $n \ge (c_0 \tau)^{(4+2p)}$, we have

$$\left\| \left[C \left(C_{X,\lambda}^{-1} - \hat{C}_{X,\lambda}^{-1} \right) \right] \right\|_{S_2(L_2(\pi),\mathcal{Y})} \le \| C \|_{S_2(\mathcal{H},\mathcal{Y})} \sqrt{2} \left(\frac{4\xi_{\delta}}{3n^{\frac{0.5+p}{2+p}}} + \sqrt{\frac{2\xi_{\delta}}{n^{\frac{p}{2+p}}}} \right) = \tau o(1)$$

The derivations in the proof of Lemma 1 show that

$$\left[CC_{X,\lambda}^{-1}\right] = \sum_{i \in I, j \in J} a_{ij} \frac{\sqrt{\mu_i}}{\mu_i + \lambda} d_j \otimes \left[e_i\right].$$

with $a_{ij} \coloneqq \langle d_j, C \sqrt{\mu_i} e_i \rangle_{\mathcal{Y}}, i \in I, j \in J$. Note that since $[C] \in S_2([\mathcal{H}]^2, \mathcal{Y})$, we have

$$\|[C]\|_{S_2([\mathcal{H}]^2,\mathcal{Y})}^2 = \left\|\sum_{i\in I, j\in J} a_{ij}d_j \otimes (\sqrt{\mu_i}e_i)\right\|_{S_2([\mathcal{H}]^2,\mathcal{Y})}^2 = \sum_{i\in I, j\in J} \frac{a_{ij}^2}{\mu_i} < +\infty.$$

Hence,

$$\left\| \left[CC_{X,\lambda}^{-1} \right] \right\|_{S_2(L_2(\pi),\mathcal{Y})}^2 = \sum_{i \in I, j \in J} a_{ij}^2 \frac{\mu_i}{(\mu_i + \lambda)^2} \le \sum_{i \in I, j \in J} \frac{a_{ij}^2}{\mu_i} = \left\| \left[C \right] \right\|_{S_2([\mathcal{H}]^2,\mathcal{Y})}^2 < +\infty.$$
(18)

Using the equality $a^2 - b^2 = (a - b)(a + b)$ and the reverse triangular inequality, we obtain the following bound, with probability at least $1 - 4e^{-\tau}$, for $n \ge (c_0 \tau)^{(4+2p)}$,

$$\begin{aligned} \left\| \left\| \left[CC_{X,\lambda}^{-1} \right] \right\|_{S_{2}(L_{2}(\pi),\mathcal{Y})}^{2} - \left\| \left[C\hat{C}_{X,\lambda}^{-1} \right] \right\|_{S_{2}(L_{2}(\pi),\mathcal{Y})}^{2} \right\| \\ \leq \left\| \left[C\left(C_{X,\lambda}^{-1} - \hat{C}_{X,\lambda}^{-1} \right) \right] \right\|_{S_{2}(L_{2}(\pi),\mathcal{Y})} \left(\left\| \left[CC_{X,\lambda}^{-1} \right] \right\|_{S_{2}(L_{2}(\pi),\mathcal{Y})} + \left\| \left[C\hat{C}_{X,\lambda}^{-1} \right] \right\|_{S_{2}(L_{2}(\pi),\mathcal{Y})} \right) \\ \leq \tau o(1) \left(2 \left\| \left[CC_{X,\lambda}^{-1} \right] \right\|_{S_{2}(L_{2}(\pi),\mathcal{Y})} + \left\| \left[C\left(C_{X,\lambda}^{-1} - \hat{C}_{X,\lambda}^{-1} \right) \right] \right\|_{S_{2}(L_{2}(\pi),\mathcal{Y})} \right) \\ \leq \tau o(1) \left(2 \left\| \left[C \right] \right\|_{S_{2}([\mathcal{H}]^{2},\mathcal{Y})}^{2} + \tau o(1) \right) \\ = \tau^{2} o(1), \end{aligned}$$

where in the second last line we used Equation (18).

Lemma 3. Fix $x \in \mathcal{X}$ and f_x as in Definition 8. For $\tau \ge \log(2)$, with probability at least $1 - 2e^{-\tau}$ (note that this event depends on x),

$$\|f_x\|_{2,n}^2 - \|[f_x]\|_{L_2(\pi)}^2 \le \frac{1}{2} \|[f_x]\|_{L_2(\pi)}^2 + \frac{5\tau\kappa^2}{3\lambda^2 n}.$$

Proof. We start with

$$\|f_x\|_{\infty} \le \kappa \|f_x\|_{\mathcal{H}} \le \kappa^2 \lambda^{-1}.$$

We apply Proposition 3 to $f = f_x$, with $M = \kappa^2 \lambda^{-1}$. For $\tau \ge \log(2)$, with probability at least $1 - 2e^{-\tau}$,

$$\left| \|f_x\|_{2,n}^2 - \|[f_x]\|_{L_2(\pi)}^2 \right| \le \frac{1}{2} \|[f_x]\|_{L_2(\pi)}^2 + \frac{5\tau\kappa^2}{3\lambda^2 n}.$$

Lemma 4. Suppose that \mathcal{X} is a compact set in \mathbb{R}^d and that $k \in C^{\theta}(\mathcal{X} \times \mathcal{X})$ for $\theta \in (0,1]$ (Definition 11). Assume that $1 \ge \lambda \ge n^{-\frac{1}{2+p}}$. With probability at least $1 - 2e^{-\tau}$, it holds for all $x \in \mathcal{X}$ simultaneously that

$$\begin{split} \|C_{X,\lambda}^{-1}k(x,\cdot)\|_{2,n}^2 &\geq \frac{1}{2} \| [C_{X,\lambda}^{-1}k(x,\cdot)] \|_{L_2(\pi)}^2 - \tau o(1), \\ \|C_{X,\lambda}^{-1}k(x,\cdot)\|_{2,n}^2 &\leq \frac{3}{2} \| [C_{X,\lambda}^{-1}k(x,\cdot)] \|_{L_2(\pi)}^2 + \tau o(1). \end{split}$$

Proof. The proof follows Li et al. (2023a, Lemma C.11). As we use different notations and tracking of constants, we provide a similar proof in our setting for completeness. By Lemma 24, there exists an ϵ -net $\mathcal{F} \subseteq \mathcal{K}_{\lambda} \subseteq \mathcal{H}$ with respect to $\|\cdot\|_{\infty}$ such that there exists a positive constant c with

$$|\mathcal{F}| \le c(\lambda \epsilon)^{-\frac{2d}{\theta}}.$$

for ϵ to be determined later. Using Lemma 3 and a union bound over the finite set \mathcal{F} , with probability at least $1 - 2e^{-\tau}$, it holds simultaneously for all $f \in \mathcal{F}$ that

$$\left| \|f\|_{2,n}^2 - \|[f]\|_{L_2(\pi)}^2 \right| \le \frac{1}{2} \|[f]\|_{L_2(\pi)}^2 + \frac{5(\tau + \log(|\mathcal{F}|))\kappa^2}{3\lambda^2 n}.$$
(19)

We work in the event where Equation (19) holds for all $f \in \mathcal{F}$. By definition of an ϵ -net and \mathcal{K}_{λ} , for any $x \in \mathcal{X}$, there exists some $f \in \mathcal{F}$ such that

$$\|C_{X,\lambda}^{-1}k(x,\cdot) - f\|_{\infty} \le \epsilon,$$

which in particular implies that

$$\begin{aligned} \| \| [C_{X,\lambda}^{-1}k(x,\cdot)] \|_{L_{2}(\pi)} - \| [f] \|_{L_{2}(\pi)} | \leq \epsilon \\ \| \| C_{X,\lambda}^{-1}k(x,\cdot) \|_{2,n} - \| f \|_{2,n} | \leq \epsilon. \end{aligned}$$

Since $\|C_{X,\lambda}^{-1}k(x,\cdot)\|_{\infty} \leq \kappa^2 \lambda^{-1}$, using the algebraic identity $a^2 - b^2 = (a - b)(2b + (a - b))$, we obtain

$$\begin{split} \| \| [C_{X,\lambda}^{-1}k(x,\cdot)] \|_{L_{2}(\pi)}^{2} - \| [f] \|_{L_{2}(\pi)}^{2} | &\leq \epsilon (2\kappa^{2}\lambda^{-1} + \epsilon) \\ \| \| C_{X,\lambda}^{-1}k(x,\cdot) \|_{2,n}^{2} - \| f \|_{2,n}^{2} | &\leq \epsilon (2\kappa^{2}\lambda^{-1} + \epsilon). \end{split}$$

We therefore have,

$$\begin{split} \|C_{X,\lambda}^{-1}k(x,\cdot)\|_{2,n}^2 &\leq \|f\|_{2,n}^2 + \epsilon(2\kappa^2\lambda^{-1} + \epsilon) \\ &\leq \frac{3}{2}\|[f]\|_{L_2(\pi)}^2 + \frac{5(\tau + \log(|\mathcal{F}|)\kappa^2}{3\lambda^2 n} + \epsilon(2\kappa^2\lambda^{-1} + \epsilon) \\ &\leq \frac{3}{2}\|[C_{X,\lambda}^{-1}k(x,\cdot)]\|_{L_2(\pi)}^2 + \frac{5(\tau + \log(|\mathcal{F}|)\kappa^2}{3\lambda^2 n} + 2\epsilon(2\kappa^2\lambda^{-1} + \epsilon). \end{split}$$

We now choose $\epsilon = \frac{1}{n}$ and bound the error term. Recall that $1 \ge \lambda \ge n^{-\frac{1}{2+p}}$, therefore,

$$\frac{5(\tau + \log(|\mathcal{F}|)\kappa^2}{3\lambda^2 n} + 2\epsilon(2\kappa^2\lambda^{-1} + \epsilon) \leq \frac{5(\tau + \log(|\mathcal{F}|)\kappa^2}{3}n^{-\frac{p}{2+p}} + 2\left(2\kappa^2n^{\frac{-1-p}{2+p}} + \frac{1}{n^2}\right) \\ \leq \frac{5\kappa^2}{3}\left(\tau + \log(c\lambda^{-\frac{2d}{\theta}}n^{\frac{2d}{\theta}})\right)n^{-\frac{p}{2+p}} + 2\left(2\kappa^2n^{\frac{-1-p}{2+p}} + \frac{1}{n^2}\right) \\ = \tau o(1).$$

Lemma 5. For $1 \ge \lambda \ge n^{-\frac{1}{2+p}}$, with probability at least $1 - 4e^{-\tau}$, for $n \ge (c_0\tau)^{4+2p}$, we have for all $x \in \mathcal{X}$ simultaneously

$$\left\| \hat{C}_{X}^{\frac{1}{2}} \hat{C}_{X,\lambda}^{-1} (C_{X} - \hat{C}_{X}) C_{X,\lambda}^{-1} k(x, \cdot) \right\|_{\mathcal{H}} = \tau o(1),$$
(20)

where c_0 is the same constant as in Lemma 2.

Proof.

$$\begin{aligned} & \left\| \hat{C}_{X}^{\frac{1}{2}} \hat{C}_{X,\lambda}^{-1} (C_{X} - \hat{C}_{X}) C_{X,\lambda}^{-1} k(x, \cdot) \right\|_{\mathcal{H}} \\ &= \left\| \hat{C}_{X}^{\frac{1}{2}} \hat{C}_{X,\lambda}^{-\frac{1}{2}} \hat{C}_{X,\lambda}^{-\frac{1}{2}} C_{X,\lambda}^{\frac{1}{2}} C_{X,\lambda}^{-\frac{1}{2}} (C_{X} - \hat{C}_{X}) C_{X,\lambda}^{-1} k(x, \cdot) \right\|_{\mathcal{H}} \\ &\leq \left\| \hat{C}_{X}^{\frac{1}{2}} \hat{C}_{X,\lambda}^{-\frac{1}{2}} \right\|_{\mathcal{H} \to \mathcal{H}} \left\| \hat{C}_{X,\lambda}^{-\frac{1}{2}} C_{X,\lambda}^{\frac{1}{2}} \right\|_{\mathcal{H} \to \mathcal{H}} \\ &\cdot \left\| C_{X,\lambda}^{-\frac{1}{2}} (C_{X} - \hat{C}_{X}) C_{X,\lambda}^{-\frac{1}{2}} \right\|_{\mathcal{H} \to \mathcal{H}} \left\| C_{X,\lambda}^{-\frac{1}{2}} k(x, \cdot) \right\|_{\mathcal{H}} \end{aligned}$$

We already saw in the proof of Lemma 2 that the first term is bounded by 1 and there is a constant $c_0 > 0$ such that for $\tau \ge \log(2)$, with probability at least $1 - 2e^{-\tau}$, for $n \ge (c_0\tau)^{4+2p}$, the second term is bounded by $\sqrt{2}$. For the third term we also saw in the proof of Lemma 4 that for $\tau \ge \log(2)$, with probability at least $1 - 2e^{-\tau}$, we have

$$\lambda^{-\frac{1}{2}} \left\| C_{X,\lambda}^{-\frac{1}{2}} (C_X - \hat{C}_X) C_{X,\lambda}^{-\frac{1}{2}} \right\|_{\mathcal{H} \to \mathcal{H}} \le \frac{4\kappa^2 \xi_{\delta}}{3n\lambda^{\frac{3}{2}}} + \sqrt{\frac{2\kappa^2 \xi_{\delta}}{n\lambda^2}}$$

where we defined

$$\xi_{\delta} = \log \frac{2\kappa^2 (\mathcal{N}_1(\lambda) + 1)}{e^{-\tau} \|C_X\|_{\mathcal{H} \to \mathcal{H}}}.$$

Finally, the fourth term is bounded above by $\lambda^{-\frac{1}{2}}\kappa$. Note that the bound on the fourth term is independent of x, so it holds simultaneously for all $x \in \mathcal{X}$. This is in contrast with the setting of Lemma 4 where for each fixed $x \in \mathcal{X}$ corresponds an element in the ϵ -net of \mathcal{F} for which we have a high probability bound, and therefore we must use a union bound in order for the bound to hold simultaneously for all $x \in \mathcal{X}$ in the proof of Lemma 4. As in the proof of Lemma 4 since $1 \ge \lambda \ge n^{-\frac{1}{2+p}}$, we have

$$\xi_{\delta} \le \log \frac{2(c_{2,l}+1)}{e^{-\tau} \|C_X\|_{\mathcal{H} \to \mathcal{H}}} + \frac{p}{2+p} \log n$$

In the bound on ξ_{δ} above, the first term does not depend on n, and the second term is logarithmic in n. Putting everything together by union bound, with probability at least $1 - 4e^{-\tau}$, for $n \ge (c_0 \tau)^{4+2p}$, we have

$$\left\| \hat{C}_{X}^{\frac{1}{2}} \hat{C}_{X,\lambda}^{-1} (C_{X} - \hat{C}_{X}) C_{X,\lambda}^{-1} k(x, \cdot) \right\|_{\mathcal{H}} = \tau o(1).$$

Remark 6 (Comparison to Li et al. (2023a)). We explicit the differences between our proof strategy and the proof strategy of Li et al. (2023a).

- Scalar versus vector-valued: lower bounding the bias in our case require us to accommodate for the vectorvalued setting (see Lemma 1).
- New proof of the bias: we lower bound the bias through Lemma 2, while Li et al. (2023a) obtain the lower bound in Lemma C.7; however the proof of Lemma C.7 implicitly uses the equality ||A⁻¹|| = ||A||⁻¹, with || · || the operator norm, see Eq. (69) Li et al. (2023a) and the preceding equations. It holds that ||A⁻¹|| ≥ ||A||⁻¹, but ||A⁻¹|| ≤ ||A||⁻¹ may not hold in general. We therefore develop a new proof for this step, leading to Lemma 2.
- New proof of the variance: we lower bound the variance in Lemma 5, while Li et al. (2023a) lower bound the variance in Lemma C.12; to show Eq. (20), Li et al. (2023a) use a covering argument involving $\mathcal{N}(\mathcal{K}_{\lambda}, \|\cdot\|_{\mathcal{H}}, \epsilon)$ (Lemma C.10). However, a close look at the proof of Lemma C.10 (last inequality of the proof) reveals that $\frac{\lambda_i}{\lambda+\lambda_i}$ was mistaken for $\frac{\lambda}{\lambda+\lambda_i}$ and plugging the correct term in the proof would lead to a vacuous bound. As explained in the proof of Lemma 5, we therefore develop a proof that is free of a covering number argument for this step.

C Learning rates for spectral algorithms

To upper bound the excess-risk, we use a decomposition involving the *approximation error* expressed as $F_{\lambda} - F_{\star}$ and the *estimation error* expressed as $\hat{F}_{\lambda} - F_{\lambda}$.

$$\left\| \left[\hat{F}_{\lambda} \right] - F_{*} \right\|_{\gamma} \leq \left\| \left[\hat{F}_{\lambda} - F_{\lambda} \right] \right\|_{\gamma} + \left\| \left[F_{\lambda} \right] - F_{*} \right\|_{\gamma},$$

where \hat{F}_{λ} is the empirical estimator based on general spectral regularization (Eq. (11)) and F_{λ} is its counterpart in population (Eq. (10)). Note that this is a different decomposition than the **bias-variance decomposition** used in the proof of Theorem 3.

The proof structure is as follows:

- 1. Fourier expansion C.1.
- 2. Approximation Error C.2.
- 3. Estimation error C.3

C.1 Fourier expansion

Recall the notations defined in Appendix A.3. The family $\{d_j\}_{j\in J}$ is an ONB of \mathcal{Y} , the family $\{\sqrt{\mu_i}e_i\}_{i\in I}$ is an ONB of $(\ker I_{\pi})^{\perp}$ and the family $\{\tilde{e}_i\}_{i\in I'}$ is an ONB of $\ker I_{\pi}$ such that $\{\mu_i^{1/2}e_i\}_{i\in I} \cup \{\tilde{e}_i\}_{i\in I'}$ forms an ONB of \mathcal{H} . Furthermore, recall that $\{\mu_i^{\beta/2}[e_i]\}_{i\in I}$ is an ONB of $[\mathcal{H}]^{\beta}$, $\beta \geq 0$.

Lemma 6 (Fourier expansion). Suppose Assumption (SRC) holds with $\beta \ge 0$. By definition of the vector-valued interpolation space and by Theorem 5, we have

$$F_* = \sum_{i \in I, j \in J} a_{ij} d_j [e_i], \qquad a_{ij} = \langle F_*, d_j [e_i] \rangle_{L_2(\pi, \mathcal{Y})}, \qquad \|F_*\|_\beta^2 = \sum_{i \in I, j \in J} \frac{a_{ij}^2}{\mu_i^\beta}.$$
 (21)

0

Then, we have the following equalities with respect to this Fourier decomposition.

1. The Hilbert-Schmidt operator $C_{\lambda} \in S_2(\mathcal{H}, \mathcal{Y})$, Eq. (10), can be written as

$$C_{\lambda} = \sum_{i \in I, j \in J} a_{ij} g_{\lambda}(\mu_i) \sqrt{\mu_i} d_j \otimes \sqrt{\mu_i} e_i.$$
⁽²²⁾

2. The Hilbert-Schmidt operator $(C_{YX} - C_{\lambda}C_X) C_{X,\lambda}^{-\frac{1}{2}} \in S_2(\mathcal{H}, \mathcal{Y})$ can be written as

$$(C_{YX} - C_{\lambda}C_X) C_{X,\lambda}^{-\frac{1}{2}} = \sum_{ij} a_{ij}r_{\lambda}(\mu_i)(\mu_i + \lambda)^{-\frac{1}{2}}\sqrt{\mu_i} (d_j \otimes \sqrt{\mu_i}e_i)$$
(23)

3. The Hilbert-Schmidt operator $C_{YX} \in S_2(\mathcal{H}, \mathcal{Y})$ can be written as

$$C_{YX} = \left(\sum_{i \in I, j \in J} a_{ij} \mu_i^{-\frac{\beta}{2}} d_j \otimes \sqrt{\mu_i} e_i\right) C_X^{\frac{\beta+1}{2}}$$
(24)

Proof. We first derive the Fourier expansion of C_{YX} ,

where in Eq. (25) we used the tower property of conditional expectation and in Eq. (26) we used the fact that $\{[e_i]\}_{i \in I}$ forms an orthonormal system in $L_2(\pi)$. We can manipulate Eq. (26) to derive Eq. (24),

$$C_{YX} = \sum_{i \in I, j \in J} a_{ij} \mu_i^{\frac{1}{2} - \frac{\beta+1}{2}} d_j \otimes \left(C_X^{\frac{\beta+1}{2}}(\sqrt{\mu_i}e_i) \right)$$
$$= \left(\sum_{i \in I, j \in J} a_{ij} \mu_i^{-\frac{\beta}{2}} d_j \otimes \sqrt{\mu_i}e_i \right) C_X^{\frac{\beta+1}{2}}.$$

By the spectral decomposition of C_X Eq. (6) and spectral calculus (Definition 9), we have that

$$g_{\lambda}(C_X) = \sum_{i \in I} g_{\lambda}(\mu_i) \sqrt{\mu_i} e_i \otimes \sqrt{\mu_i} e_i + g_{\lambda}(0) \sum_{i \in I'} \tilde{e}_i \otimes \tilde{e}_i,$$
(27)

$$r_{\lambda}(C_X) = \sum_{i \in I} r_{\lambda}(\mu_i) \sqrt{\mu_i} e_i \otimes \sqrt{\mu_i} e_i + \sum_{i \in I'} \tilde{e}_i \otimes \tilde{e}_i.$$
(28)

where we used $r_{\lambda}(0) = 1$.

Proof of Eq. (22). Using Eq. (26) and (27), we have

$$C_{\lambda} = \left(\sum_{i \in I, j \in J} a_{ij} \sqrt{\mu_i} d_j \otimes (\sqrt{\mu_i} e_i)\right) \left(\sum_{k \in I} g_{\lambda}(\mu_k) (\sqrt{\mu_k} e_k) \otimes (\sqrt{\mu_k} e_k) + g_{\lambda}(0) \sum_{l \in I'} \tilde{e}_l \otimes \tilde{e}_l\right)$$

$$= \sum_{ijk} a_{ij} \sqrt{\mu_i} g_{\lambda}(\mu_k) \delta_{ik} d_j \otimes (\sqrt{\mu_k} e_k)$$

$$= \sum_{i \in I, j \in J} a_{ij} \sqrt{\mu_i} g_{\lambda}(\mu_i) d_j \otimes (\sqrt{\mu_i} e_i),$$

(29)

where in Eq. (29), we recall the fact that $\{\sqrt{\mu_i}e_i\}_{i\in I}$ forms an ONB of $(\ker I_{\pi})^{\perp}$ and $\{\tilde{e}_i\}_{i\in I'}$ forms an ONB of $\ker I_{\pi}$.

Proof of Eq. (23). Using Eq. (26) and (28), we have

$$\begin{split} \left(C_{YX} - C_{\lambda} C_{X}\right) C_{X,\lambda}^{-\frac{1}{2}} &= C_{YX} r_{\lambda} (C_{X}) C_{X,\lambda}^{-\frac{1}{2}} \\ &= \left(\sum_{i \in I, j \in J} a_{ij} \sqrt{\mu_{i}} d_{j} \otimes (\sqrt{\mu_{i}} e_{i})\right) \left(\sum_{k \in I} r_{\lambda} (\mu_{k}) \sqrt{\mu_{k}} e_{k} \otimes \sqrt{\mu_{k}} e_{k} + \sum_{l \in I'} \tilde{e}_{l} \otimes \tilde{e}_{l}\right) C_{X,\lambda}^{-\frac{1}{2}} \\ &= \left(\sum_{ijk} a_{ij} \sqrt{\mu_{i}} r_{\lambda} (\mu_{k}) d_{j} \otimes (\sqrt{\mu_{k}} e_{k}) \delta_{ik}\right) C_{X,\lambda}^{-\frac{1}{2}} \\ &= \sum_{i \in I, j \in J} a_{ij} \sqrt{\mu_{i}} r_{\lambda} (\mu_{i}) d_{j} \otimes (C_{X,\lambda}^{-\frac{1}{2}} (\sqrt{\mu_{i}} e_{i})) \\ &= \sum_{i \in I, j \in J} a_{ij} \sqrt{\mu_{i}} (\mu_{i} + \lambda)^{-\frac{1}{2}} r_{\lambda} (\mu_{i}) d_{j} \otimes (\sqrt{\mu_{i}} e_{i}), \end{split}$$

Lemma 7. Suppose Assumption (SRC) holds with $\beta \ge 0$, then the following bound is satisfied, for all $\lambda > 0$ and $0 \le \gamma \le 1$, we have

$$\|[F_{\lambda}]\|_{\gamma}^{2} \leq E^{2} \|F_{*}\|_{\min\{\gamma,\beta\}}^{2} \lambda^{-(\gamma-\beta)_{+}}$$

For the definition of E, see Eq. (8).

Proof. We adopt the notation of Lemma 6. By Parseval's identity and Eq. (22), we have

$$\begin{split} \|[F_{\lambda}]\|_{\gamma}^{2} &= \|C_{\lambda}\|_{S_{2}([\mathcal{H}]^{\gamma},\mathcal{Y})}^{2} \\ &= \sum_{i \in I, j \in J} a_{ij}^{2} g_{\lambda}(\mu_{i})^{2} \mu_{i}^{2-\gamma} \end{split}$$

In the case of $\gamma \leq \beta$, we bound $g_{\lambda}(\mu_i)\mu_i \leq E$ using Eq. (8). Then, by Eq. (21),

$$\|[F_{\lambda}]\|_{\gamma}^{2} \leq E^{2} \sum_{i \in I, j \in J} \frac{a_{ij}^{2}}{\mu_{i}^{\gamma}} = E^{2} \|F_{*}\|_{\gamma}^{2}.$$

In the case of $\gamma > \beta$, we apply Eq. (8) to $g_{\lambda}(\mu_i)\mu_i^{1-\frac{\gamma-\beta}{2}} \leq E\lambda^{-\frac{\gamma-\beta}{2}}$ to obtain, using Eq. (21) again,

$$\begin{split} \|[F_{\lambda}]\|_{\gamma}^{2} &= \sum_{i \in I, j \in J} g_{\lambda}(\mu_{i})^{2} \mu_{i}^{2-(\gamma-\beta)} \mu_{i}^{-\beta} a_{ij}^{2} \\ &\leq E^{2} \lambda^{-(\gamma-\beta)} \sum_{i \in I, j \in J} \mu_{i}^{-\beta} a_{ij}^{2} \\ &= E^{2} \lambda^{-(\gamma-\beta)} \|F_{*}\|_{\beta}^{2}. \end{split}$$

Lemma 8. Suppose Assumption (SRC) holds for $0 \le \beta \le 2\rho$, with ρ the qualification. Then, the following bound is satisfied, for all $\lambda > 0$, we have

$$\left\| \left(C_{YX} - C_{\lambda} C_{X} \right) C_{X,\lambda}^{-\frac{1}{2}} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} \leq \omega_{\rho} \|F_{*}\|_{\beta} \lambda^{\frac{\beta}{2}}.$$

For the definition of ω_{ρ} , see Eq. (9).

Proof. Recall that in Lemma 6 we used the decomposition

$$F_* = \sum_{i \in I, j \in J} a_{ij} d_j [e_i],$$

where Assumption (SRC) implies that $||F_*||_{\beta}^2 = \sum_{ij} \frac{a_{ij}^2}{\mu_i^{\beta}} < \infty$. Using Eq. (23) in Lemma 6 and Parseval's identity w.r.t. the ONS $\{d_j \otimes \mu_i^{1/2} e_i\}_{i \in I, j \in J}$ in $S_2(\mathcal{H}, \mathcal{Y})$, we have

$$\left\| \left(C_{YX} - C_{\lambda} C_{X} \right) C_{X,\lambda}^{-\frac{1}{2}} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} = \left(\sum_{i \in I, j \in J} a_{ij}^{2} r_{\lambda}^{2}(\mu_{i})(\mu_{i} + \lambda)^{-1} \mu_{i} \right)^{\frac{1}{2}}$$

$$\leq \left(\sum_{i \in I, j \in J} \frac{a_{ij}^{2}}{\mu_{i}^{\beta}} r_{\lambda}^{2}(\mu_{i}) \mu_{i}^{\beta} \right)^{\frac{1}{2}}$$

$$\leq \| F_{*} \|_{\beta} \sup_{i \in I} r_{\lambda}(\mu_{i}) \mu_{i}^{\frac{\beta}{2}}$$

$$\leq \| F_{*} \|_{\beta} \omega_{\rho} \lambda^{\frac{\beta}{2}}.$$

Lemma 9. Suppose Assumption (SRC) holds with $\beta \ge 0$, then for all $\lambda > 0$, we have

$$\left\| C_{\lambda} r_{\lambda} \left(\hat{C}_{X} \right) \hat{C}_{X,\lambda}^{\frac{1}{2}} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} \leq B \left\| \hat{C}_{X,\lambda}^{\frac{1}{2}} r_{\lambda} (\hat{C}_{X}) g_{\lambda} (C_{X}) C_{X}^{\frac{\beta+1}{2}} \right\|_{\mathcal{H} \to \mathcal{H}}$$

where $||F_*||_{\beta} = B < \infty$.

Proof. Recall that Lemma 6 we used the decomposition

$$F_* = \sum_{i \in I, j \in j} a_{ij} d_j [e_i],$$

where $||F_*||_{\beta}^2 = \sum_{ij} \frac{a_{ij}^2}{\mu_i^{\beta}} = B^2 < \infty$. Using Eq. (24) in Lemma 6 and $C_{\lambda} = C_{YX}g_{\lambda}(C_X)$, we have

$$\begin{aligned} \left\| C_{\lambda} r_{\lambda} \left(\hat{C}_{X} \right) \hat{C}_{X,\lambda}^{\frac{1}{2}} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} &= \left\| \left(\sum_{ij} a_{ij} \mu_{i}^{-\frac{\beta}{2}} d_{j} \otimes \sqrt{\mu_{i}} e_{i} \right) C_{X}^{\frac{\beta+1}{2}} g_{\lambda}(C_{X}) r_{\lambda}(\hat{C}_{X}) \hat{C}_{X,\lambda}^{\frac{1}{2}} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} \\ &\leq B \left\| C_{X}^{\frac{\beta+1}{2}} g_{\lambda}(C_{X}) r_{\lambda}(\hat{C}_{X}) \hat{C}_{X,\lambda}^{\frac{1}{2}} \right\|_{\mathcal{H} \to \mathcal{H}}, \end{aligned}$$

where we notice that the $S_2(\mathcal{H}, \mathcal{Y})$ norm of the first term is exactly the β norm of F_* , which is given by B. Recalling that C_X, \hat{C}_X are self adjoint, we prove the final result by taking the adjoint and using that an operator has the same operator norm as its adjoint.

C.2 Approximation Error

Lemma 10. Let F_{λ} be given by Eq. (10) based on a general spectral filter satisfying Definition 2 with qualification $\rho \ge 0$. Suppose Assumption (SRC) holds with parameter $\beta \ge 0$ and define $\beta_{\rho} = \min\{\beta, 2\rho\}$, then the following bound is satisfied, for all $\lambda > 0$ and $0 \le \gamma \le \beta_{\rho}$,

$$\left\| \left[F_{\lambda} \right] - F_{*} \right\|_{\gamma}^{2} \leq \omega_{\rho}^{2} \left\| F_{*} \right\|_{\beta_{\rho}}^{2} \lambda^{\beta_{\rho} - \gamma}.$$

		I
		I

Proof. In Eq. (10), we defined $F_{\lambda}(\cdot) = C_{\lambda}\phi(\cdot)$. On the other hand, in Lemma 6 we obtained the Fourier expansion of C_{λ} leading to Eq. (22). Thus we have for π -almost all $x \in \mathcal{X}$,

$$F_{\lambda}(x) = \sum_{i \in I, j \in J} a_{ij} \mu_i g_{\lambda}(\mu_i) d_j e_i(x).$$

Therefore,

$$[F_{\lambda}] - F_* = \sum_{i \in I, j \in J} a_{ij} (1 - \mu_i g_{\lambda}(\mu_i)) d_j [e_i] = \sum_{i \in I, j \in J} a_{ij} r_{\lambda}(\mu_i) d_j [e_i]$$

Suppose $\beta \leq 2\rho$, using Parseval's identity w.r.t. the ONB $\{d_j \mu_i^{\gamma/2}[e_i]\}_{i \in I, j \in J}$ of $[\mathcal{G}]^{\gamma}$, we have

$$\begin{split} \|[F_{\lambda}] - F_{*}\|_{\gamma}^{2} &= \left\| \sum_{i \in I, j \in J} \frac{a_{ij}}{\mu_{i}^{\gamma/2}} r_{\lambda}(\mu_{i}) d_{j} \mu_{i}^{\gamma/2} e_{i} \right\|_{\gamma}^{2} \\ &= \sum_{i \in I, j \in J} \frac{a_{ij}^{2}}{\mu_{i}^{\gamma}} r_{\lambda}^{2}(\mu_{i}) \\ &= \sum_{i \in I, j \in J} \frac{a_{ij}^{2}}{\mu_{i}^{\beta}} r_{\lambda}^{2}(\mu_{i}) \mu_{i}^{\beta - \gamma} \\ &\leq \omega_{\rho}^{2} \lambda^{\beta - \gamma} \sum_{i \in I, j \in J} \frac{a_{ij}^{2}}{\mu_{i}^{\beta}} \\ &= \|F_{*}\|_{\beta}^{2} \omega_{\rho}^{2} \lambda^{\beta - \gamma} \end{split}$$

where we used Eq. (9) in the definition of a filter function, together with $0 \leq \beta \leq 2\rho$ and $0 \leq \gamma \leq \beta$, which taken together implies that $0 \leq \frac{\beta - \gamma}{2} \leq \rho$. Finally, if $\beta \geq 2\rho$, then since $[\mathcal{G}]^{2\rho} \subseteq [\mathcal{G}]^{2\beta}$, we can perform the last derivations again with $\beta = 2\rho$ to obtain the final result.

C.3 Estimation error

Before proving the main results we recall two *embedding properties* for the vector-valued interpolation space $[\mathcal{G}]^{\beta}$ (Definition 1). The first embedding property lifts the property (EMB) defined for the scalar-valued RKHS $[\mathcal{H}]^{\alpha}$ to the vector-valued RKHS $[\mathcal{G}]^{\alpha}$.

Lemma 11 (L_{∞} -embedding property - Lemma 4 Li et al. (2023b)). Under (EMB) the inclusion operator $\mathcal{I}_{\pi}^{\alpha,\beta}: [\mathcal{G}]^{\alpha} \hookrightarrow L_{\infty}(\pi; \mathcal{Y})$ is bounded with operator norm A,

Theorem 6 (L_q -embedding property - Theorem 3 Li et al. (2023b)). Let Assumption (EMB) be satisfied with parameter $\alpha \in (0,1]$. For any $\beta \in [0, \alpha)$, the inclusion map

$$\mathcal{I}_{\pi}^{q_{\alpha,\beta}}:[\mathcal{G}]^{\beta} \hookrightarrow L_{q_{\alpha,\beta}}(\pi;\mathcal{Y})$$

is bounded, where $q_{\alpha,\beta} \coloneqq \frac{2\alpha}{\alpha-\beta}$.

The L_q -embedding property was first introduced in the scalar-valued setting in Zhang et al. (2023a) and then lifted to the vector-valued setting by Li et al. (2023b). Its role is to replace a boundedness condition on the ground truth function F_* . We now explain how the L_q -embedding property can be combined with Assumption (EMB) and a truncation technique.

Lemma 12. Recall that π is the marginal measure of X on \mathcal{X} . For $t \ge 0$, define the measurable set Ω_t as follows

$$\Omega_t \coloneqq \{x \in \mathcal{X} : \|F_*(x)\|_{\mathcal{Y}} \le t\}$$

Let q > 0. Assume that $F_* \in L_q(\pi; \mathcal{Y})$. In other words, there exists some constant $c_q > 0$ such that

$$\|F_*\|_{L_q(\pi;\mathcal{Y})} = \left(\int_{\mathcal{X}} \|F_*(x)\|_{\mathcal{Y}}^q \mathrm{d}\pi(x)\right)^{\frac{1}{q}} = c_q < +\infty,$$

Then we have the following conclusions

1. The π -measure of the complement of Ω can be bounded by

$$\pi(\{x \notin \Omega_t\}) \le \frac{c_q^q}{t^q}$$

2. Recall that $\{x_i\}_{i=1}^n$ are i.i.d. samples distributed according to π . If $t = n^{\frac{1}{q}}$ for $\tilde{q} < q$, then we can conclude as follows. For a fixed parameter $\tau > 0$, for all sufficiently large n, where the hidden index bound depends on $q\tilde{q}^{-1}$ and τ , we have

$$^{\otimes n}\left(\cap_{i=1}^{n}\left\{x_{i}\in\Omega_{t}\right\}\right)\geq1-e^{-\tau}.$$

Proof. The first claim is a straightforward application of Markov's inequality, as follows

π

$$\pi(\{x \notin \Omega_t\}) = \pi(\|F_*(x)\|_{\mathcal{Y}} > t) \le \frac{\mathbb{E}_{\pi}\left[\|F_*(X)\|_{\mathcal{Y}}^q\right]}{t^q} = \frac{c_q^q}{t^q}$$

To show the second claim, we first evaluate the probability that there exists some x_i 's that lies outside Ω_t ,

$$\pi^{\otimes n} \left(\cup_{i=1}^{n} \{ x_i \notin \Omega_t \} \right) = 1 - \pi^{\otimes n} \left(\cap_{i=1}^{n} \{ x_i \in \Omega_t \} \right)$$
$$= 1 - \pi \left(\{ x_i \in \Omega_t \} \right)^n$$
$$\leq 1 - \left(1 - \frac{c_q^q}{t^q} \right)^n$$
$$\leq \frac{c_q^q n}{t^q},$$

where in the last inequality we used Bernoulli's inequality, which states that for $r \ge 1$ and $0 \le x \le 1$,

$$(1-x)^r \ge 1 - rx.$$

By assumption $t = n^{\frac{1}{q_t}}$ for some fixed $q > q_t > 0$. We thus have

$$\pi^{\otimes n}\left(\cup_{i=1}^n \{x_i \notin \Omega_t\}\right) \le c_q^q n^{1-\frac{q}{q_t}} \le e^{-\tau},$$

for sufficiently large n, where the hidden index bound depends on $\frac{q}{q_t}$ and τ .

We adapt Li et al. (2023b, Lemma 5) to the spectral algorithms setting.

Lemma 13. Suppose Assumptions (SRC) and (EMB) hold for some $0 \le \beta \le 2\rho$, with ρ the qualification, then the following bounds are satisfied, for all $0 < \lambda \le 1$,

$$\|[F_{\lambda}] - F_{*}\|_{L_{\infty}}^{2} \le \left(\|F_{*}\|_{L_{\infty}} + A \max\{E, \omega_{\rho}\}\|F_{*}\|_{\beta}\right)^{2} \lambda^{\beta - \alpha},$$
(30)

$$\|[F_{\lambda}]\|_{L_{\infty}}^{2} \le A^{2} E^{2} \|F_{*}\|_{\min\{\alpha,\beta\}}^{2} \lambda^{-(\alpha-\beta)_{+}},$$
(31)

Proof. We use Lemma 11 and Lemma 7 to write:

$$\|[F_{\lambda}]\|_{\infty}^{2} \leq A^{2} \|[F_{\lambda}]\|_{\alpha}^{2} \leq A^{2} E^{2} \|F_{*}\|_{\min\{\alpha,\beta\}}^{2} \lambda^{-(\alpha-\beta)_{+}}.$$

This proves Eq. (31). To show Eq. (30), in the case $\beta \leq \alpha$ we use the triangle inequality, Eq. (31) and $\lambda \leq 1$ to obtain

$$\begin{aligned} \|[F_{\lambda}] - F_*\|_{\infty} &\leq \|F_*\|_{\infty} + \|[F_{\lambda}]\|_{\infty} \\ &\leq (\|F_*\|_{\infty} + AE\|F_*\|_{\beta})\lambda^{-\frac{\alpha-\beta}{2}}. \end{aligned}$$

In the case $\beta > \alpha$, Eq. (30) is a consequence of Lemma 11 and Lemma 10 with $\gamma = \alpha$ (here we use the assumption $0 \le \beta \le 2\rho$),

$$\|[F_{\lambda}] - F_{*}\|_{\infty}^{2} \leq A^{2} \|[F_{\lambda}] - F_{*}\|_{\alpha}^{2} \leq A^{2} \omega_{\rho}^{2} \|F_{*}\|_{\beta}^{2} \lambda^{\beta - \alpha} \leq (\|F_{*}\|_{\infty} + A\omega_{\rho}\|F_{*}\|_{\beta})^{2} \lambda^{\beta - \alpha}.$$

We adapt Zhang et al. (2023b, Theorem 13) to the vector-valued setting.

Theorem 7. Suppose that Assumptions (EMB), (EVD), (MOM) and (SRC) hold for $0 \le \beta \le 2\rho$, where ρ is the qualification, and $p \le \alpha \le 1$. Denote, for i = 1, ..., n,

$$\xi_i = \xi(x_i, y_i) = \left(\left(y_i - C_\lambda \phi(x_i) \right) \otimes \phi(x_i) \right) C_{X,\lambda}^{-\frac{1}{2}},$$

and for $t \ge 0$,

$$\Omega_t = \{ x \in \mathcal{X} : \|F_*(x)\|_{\mathcal{Y}} \le t \}$$

Then for all $\tau \ge 1$, with probability at least $1 - 2e^{-\tau}$, we have

$$\left\|\frac{1}{n}\sum_{i=1}^{n}\xi_{i}\mathbb{1}\left\{x_{i}\in\Omega_{t}\right\}-\mathbb{E}\left[\xi(X,Y)\mathbb{1}\left\{X\in\Omega_{t}\right\}\right]\right\|_{S_{2}(\mathcal{H},\mathcal{Y})}$$
$$\leq\tau\left(c_{1}\lambda^{\frac{\beta}{2}-\alpha}n^{-1}+c_{2}\lambda^{-\frac{\alpha}{2}}n^{-1}(t+R+A)+\frac{c_{3}\sqrt{\mathcal{N}_{1}(\lambda)}}{\sqrt{n}}+\frac{c_{4}}{\sqrt{n}\lambda^{\frac{\alpha-\beta}{2}}}\right)$$

where R is the constant from Assumption (MOM), and

$$c_{1} = 8\sqrt{2}A^{2} \max\{E, \omega_{\rho}\} \|F_{*}\|_{\beta}$$

$$c_{2} = 8\sqrt{2}A$$

$$c_{3} = 8\sqrt{2}\sigma$$

$$c_{4} = 8\sqrt{2}A \|F_{*}\|_{\beta}\omega_{\rho}$$

where A is the constant from Assumption (EMB), and E, ω_{ρ} are defined in Eq. (8) and (9) respectively.

Proof. We wish to apply vector-valued Bernstein's inequality, namely Theorem 11. We thus compute,

$$\mathbb{E}\left[\left\|\xi(X,Y)\mathbbm{1}\left\{X\in\Omega_{t}\right\}\right\|_{S_{2}(\mathcal{H},\mathcal{Y})}^{m}\right] = \mathbb{E}\left[\mathbbm{1}\left\{X\in\Omega_{t}\right\}\left\|\left(Y-C_{\lambda}\phi(X)\right)\otimes\left(C_{X,\lambda}^{-\frac{1}{2}}\phi(X)\right)\right\|_{S_{2}(\mathcal{H},\mathcal{Y})}^{m}\right]\right]$$
$$= \mathbb{E}\left[\mathbbm{1}\left\{X\in\Omega_{t}\right\}\left\|\left(Y-C_{\lambda}\phi(X)\right)\right\|_{\mathcal{Y}}^{m}\left\|C_{X,\lambda}^{-\frac{1}{2}}\phi(X)\right\|_{\mathcal{H}}^{m}\right]$$
$$= \int_{\Omega_{t}}\left\|C_{X,\lambda}^{-\frac{1}{2}}\phi(x)\right\|_{\mathcal{H}}^{m}\int_{\mathcal{Y}}\left\|y-C_{\lambda}\phi(x)\right\|_{\mathcal{Y}}^{m}dp(x,dy)d\pi(x).$$
(32)

First we consider the inner integral, by Assumption (MOM),

$$\begin{split} \int_{\mathcal{Y}} \| (y - C_{\lambda}\phi(x)) \|_{\mathcal{Y}}^{m} \, \mathrm{d}p(x, \mathrm{d}y) &\leq 2^{m-1} \left(\int_{\mathcal{Y}} \| y - F_{*}(x) \|_{\mathcal{Y}}^{m} + \| F_{\lambda}(x) - F_{*}(x) \|_{\mathcal{Y}}^{m} \right) \mathrm{d}p(x, \mathrm{d}y) \\ &= m! \sigma^{2} (2R)^{m-2} + 2^{m-1} \| F_{\lambda}(x) - F_{*}(x) \|_{\mathcal{Y}}^{m} \,. \end{split}$$

Plugging the above inequality into Eq. (32), as well as introducing the shorthand,

$$h_x \coloneqq C_{X,\lambda}^{-\frac{1}{2}}\phi(x),$$

we have

$$\mathbb{E}\left[\|\xi(X,Y)\mathbb{1}\{X\in\Omega_t\}\|_{S_2(\mathcal{H},\mathcal{Y})}^m\right] \le m!\sigma^2(2R)^{m-2} \int_{\Omega_t} \|h_x\|_{\mathcal{H}}^m \mathrm{d}\pi(x)$$

$$+ 2^{m-1} \int_{\Omega_t} \|h_x\|_{\mathcal{H}}^m \|F_\lambda(x) - F_*(x)\|_{\mathcal{Y}}^m \mathrm{d}\pi(x).$$

$$(33)$$

We bound term (33) using Lemma 15 and Lemma 17 with l = 1. We have,

$$\int_{\Omega_t} \|h_x\|_{\mathcal{H}}^m \mathrm{d}\pi(x) \le (A\lambda^{-\frac{\alpha}{2}})^{m-2} \mathcal{N}_1(\lambda).$$

Therefore we bound term (33) as follows,

$$m!\sigma^2(2R)^{m-2}\int_{\Omega_t} \|h_x\|_{\mathcal{H}}^m \mathrm{d}\pi(x) \le m!\sigma^2 \left(\frac{2AR}{\lambda^{\frac{\alpha}{2}}}\right)^{m-2} \mathcal{N}_1(\lambda)$$

If $\beta \ge \alpha$, by Assumption (EMB),

$$||F_*||_{\infty} \le A ||F_*||_{\alpha} \le A ||F_*||_{\beta}.$$

Hence by Lemma 13,

$$\|[F_{\lambda}] - F_{*}\|_{\infty} \leq (\|F_{*}\|_{\infty} + A\max\{E, \omega_{\rho}\}\|F_{*}\|_{\beta})\lambda^{\frac{\beta-\alpha}{2}} \leq A(1 + \max\{E, \omega_{\rho}\})\|F_{*}\|_{\beta}\lambda^{\frac{\beta-\alpha}{2}}.$$

If $\beta < \alpha$, by Lemma 13, we have for π -almost all $x \in \Omega_t$,

$$\|F_*(x) - F_{\lambda}(x)\|_{\mathcal{Y}} \le t + \|[F_{\lambda}]\|_{L_{\infty}(\pi;\mathcal{Y})} \le t + AE\|F_*\|_{\beta}\lambda^{\frac{\beta-\alpha}{2}}.$$

Therefore, for all $\beta \in [0, 2\rho]$,

$$\|(F_* - [F_{\lambda}])\mathbb{1}_{X \in \Omega_t}\|_{L_{\infty}(\pi;\mathcal{Y})} \le t + A(1 + \max\{E, \omega_{\rho}\}\|F_*\|_{\beta}\lambda^{\frac{\beta-\alpha}{2}}) =: \chi(t, \lambda).$$

Using Lemma 17 with l = 1, we have,

$$2^{m-1} \int_{\Omega_t} \|h_x\|_{\mathcal{H}}^m \|F_*(x) - F_{\lambda}(x)\|_{\mathcal{Y}}^m d\pi(x)$$

$$\leq 2^{m-1} \chi(t,\lambda)^{m-2} (A\lambda^{-\frac{\alpha}{2}})^m \|F_* - [F_{\lambda}]\|_{L_2(\pi;\mathcal{Y})}^2$$

$$= \left(\frac{2\chi(t,\lambda)A}{\lambda^{\frac{\alpha}{2}}}\right)^{m-2} \|F_* - [F_{\lambda}]\|_{L_2(\pi;\mathcal{Y})}^2 \frac{2A^2}{\lambda^{\alpha}}$$

$$\leq m! \left(\frac{2\chi(t,\lambda)A}{\lambda^{\frac{\alpha}{2}}}\right)^{m-2} \|F_* - [F_{\lambda}]\|_{L_2(\pi;\mathcal{Y})}^2 \frac{2A^2}{\lambda^{\alpha}}.$$

Putting everything together,

$$\mathbb{E}\Big[\|\xi(X,Y)\mathbb{1}\{X\in\Omega_t\}\|_{S_2(\mathcal{H},\mathcal{Y})}^m\Big] \le m! \left(\frac{2(R+\chi(t,\lambda))A}{\lambda^{\frac{\alpha}{2}}}\right)^{m-2} \left(\sigma^2 \mathcal{N}_1(\lambda) + \|F_*-[F_\lambda]\|_{L_2(\pi;\mathcal{Y})}^2 \frac{2A^2}{\lambda^{\alpha}}\right).$$

We now apply Theorem 11 with

$$\begin{split} L &\leftarrow \frac{2(R + \chi(t, \lambda))A}{\lambda^{\frac{\alpha}{2}}} \\ \sigma &\leftarrow 2\sigma \sqrt{\mathcal{N}_1(\lambda)} + \|F_* - [F_\lambda]\|_{L_2(\pi;\mathcal{Y})} \frac{2A}{\lambda^{\frac{\alpha}{2}}} \end{split}$$

We bound $||F_* - [F_{\lambda}]||_{L_2(\pi;\mathcal{Y})}$ using Lemma 10 with $\gamma = 0$,

$$\|F_* - [F_\lambda]\|_{L_2(\pi;\mathcal{Y})} \le \omega_\rho \|F_*\|_\beta \lambda^{\frac{\beta}{2}}.$$

The conclusion is, for all $\tau \ge 1$, with probability at least $1 - 2e^{-\tau}$, we have

$$\begin{aligned} \left\| \frac{1}{n} \sum_{i=1}^{n} \xi_{i} \mathbb{1}\left\{ x_{i} \in \Omega_{t} \right\} - \mathbb{E}[\xi(X, Y) \mathbb{1}\left\{ X \in \Omega_{t} \right\}] \right\|_{S_{2}(\mathcal{H}, \mathcal{Y})} \\ &\leq 4\sqrt{2}\tau \left(\frac{2\sigma\sqrt{\mathcal{N}_{1}(\lambda)} + \|F_{*} - [F_{\lambda}]\|_{L_{2}(\pi; \mathcal{Y})} \frac{2A}{\lambda^{\frac{\alpha}{2}}}}{\sqrt{n}} + \frac{2(R + \chi(t, \lambda))A}{n\lambda^{\frac{\alpha}{2}}} \right) \\ &\leq 4\sqrt{2}\tau \left(\frac{2\sigma}{\sqrt{n}}\sqrt{\mathcal{N}_{1}(\lambda)} + \frac{2A\|F_{*}\|_{\beta}\omega_{\rho}}{\sqrt{n}\lambda^{\frac{\alpha-\beta}{2}}} + \frac{2(R + t + A)A}{n\lambda^{\frac{\alpha}{2}}} + \frac{2A^{2}\max\{E, \omega_{\rho}\}\|F_{*}\|_{\beta}}{n\lambda^{\alpha-\frac{\beta}{2}}} \right). \end{aligned}$$

Lemma 14. Suppose that the same assumptions and notations listed in Theorem 7 hold.

1. Suppose $\beta + p > \alpha$, and $\lambda \approx n^{-\frac{1}{\beta+p}}$. For any fixed $\tau \ge 1$, with probability at least $1 - 2e^{-\tau}$, suppose that the truncation level t satisfies

$$t < n^{\frac{1}{2}\left(1 + \frac{p-\alpha}{p+\beta}\right)}.$$

then there exists a constant c > 0 such that

$$\left\|\frac{1}{n}\sum_{i=1}^{n}\xi_{i}\mathbb{1}\left\{x_{i}\in\Omega_{t}\right\}-\mathbb{E}\left[\xi(X,Y)\mathbb{1}\left\{X\in\Omega_{t}\right\}\right]\right\|_{S_{2}(\mathcal{H},\mathcal{Y})}\leq c\tau n^{-\frac{1}{2}\frac{\beta}{\beta+p}}$$

2. Suppose $\beta + p \leq \alpha$, and $\lambda \approx \left(\frac{n}{\log^{\theta}(n)}\right)^{\frac{1}{\alpha}}$ for some $\theta > 1$. For any fixed $\tau \geq 1$, with probability at least $1 - 2e^{-\tau}$, suppose that the truncation level t satisfies

$$t \le n^{\frac{1}{2}\left(1 - \frac{\beta}{\alpha}\right)},$$

then there exists a constant c > 0 such that

$$\left\|\frac{1}{n}\sum_{i=1}^{n}\xi_{i}\mathbb{1}\left\{x_{i}\in\Omega\right\}-\mathbb{E}\left[\xi(X,Y)\mathbb{1}\left\{X\in\Omega\right\}\right]\right\|_{\mathcal{G}}\leq c\tau\left(\frac{n}{\log^{\theta}(n)}\right)^{-\frac{\beta}{2\alpha}}$$

Proof. Note that Theorem 7 yields the same conclusion as in the scalar-valued case proved in Zhang et al. (2023b, Theorem 13). The Lemma then follows from the analysis for the scalar-valued case in the proof of Zhang et al. (2023b, Theorem 15). \Box

We adapt Zhang et al. (2023a, Theorem 15) to the vector-valued setting.

Theorem 8. Suppose that Assumptions (EMB), (EVD), (MOM) and (SRC) hold for $0 \le \beta \le 2\rho$, where ρ is the qualification, and $p \le \alpha \le 1$.

1. In the case of $\beta + p > \alpha$, choosing $\lambda \approx n^{-\frac{1}{\beta+p}}$, for any fixed $\tau \ge \log(4)$, when n is sufficiently large, with probability at least $1 - 4e^{-\tau}$, we have

$$\left\| \left(\left(\hat{C}_{YX} - C_{\lambda} \hat{C}_{X} \right) - \left(C_{YX} - C_{\lambda} C_{X} \right) \right) C_{X,\lambda}^{-\frac{1}{2}} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} \le c\tau n^{-\frac{1}{2}\frac{\beta}{\beta+p}}$$
(34)

where c is a constant independent of n, τ, λ .

2. In the case of $\beta + p \leq \alpha$, choosing $\lambda \asymp \left(\frac{n}{\log^{\theta}(n)}\right)^{-\frac{1}{\alpha}}$ for some $\theta > 1$. We make the additional assumption that there exists some $\alpha' < \alpha$ such that Assumption (MOM) is satisfied for $\alpha' < \alpha$. Then, for any fixed $\tau \geq \log(4)$, when n is sufficiently large, where the hidden index bound depends on $\alpha - \alpha'$, with probability at least $1 - 4e^{-\tau}$, we have

$$\left\| \left(\left(\hat{C}_{YX} - C_{\lambda} \hat{C}_{X} \right) - \left(C_{YX} - C_{\lambda} C_{X} \right) \right) C_{X,\lambda}^{-\frac{1}{2}} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} \le c\tau \left(\frac{n}{\log^{\theta}(n)} \right)^{-\frac{\omega}{2\alpha}}$$
(35)

where c is a constant independent of n, τ, λ .

Proof. By assumption (EMB) and Theorem 6, if $\beta < \alpha$, then the inclusion map

$$\mathcal{I}^{q_{\alpha,\beta}}_{\pi}:[\mathcal{G}]^{\beta} \hookrightarrow L_{q_{\alpha,\beta}}(\pi;\mathcal{Y})$$

is bounded, where $q_{\alpha,\beta} \coloneqq \frac{2\alpha}{\alpha-\beta}$. If $\beta \ge \alpha$, then by Lemma 11 the inclusion map

$$\mathcal{I}^{\infty}_{\pi} : [\mathcal{G}]^{\beta} \hookrightarrow L_{\infty}(\pi; \mathcal{Y})$$

is bounded and therefore $[\mathcal{G}]^{\beta}$ is continuously embedded into $L_q(\pi; \mathcal{Y})$ for any $q \ge 1$. In the rest of the proof, we will use q to denote $q_{\alpha,\beta}$, unless otherwise specified. Furthermore, we will use $c_q = \|F_*\|_{L_q(\pi;\mathcal{Y})}$.

We first consider the case $\beta + p > \alpha$. We can easily verify using $\beta + p > \alpha$ that the following inequality holds

$$\frac{1}{2}\left(1+\frac{p-\alpha}{p+\beta}\right) > \frac{1}{2}\left(\frac{p}{p+\beta}\right) > \frac{\alpha-\beta}{2\alpha} = \frac{1}{q_{\alpha,\beta}}.$$

Choose $t = n^{\tilde{q}^{-1}}$, where

$$\frac{1}{\tilde{q}} = \frac{1}{2} \left(\frac{1}{2} \left(1 + \frac{p - \alpha}{p + \beta} \right) + \frac{1}{q} \right).$$

We thus have

$$n^{\frac{1}{2}\left(1+\frac{p-\alpha}{p+\beta}\right)} > t = n^{\tilde{q}^{-1}} > n^{\frac{1}{q_{\alpha,\beta}}}.$$

Thus the assumptions for both Lemma 12 and Lemma 14 are satisfied.

We then consider the case $\beta + p \leq \alpha$. We now apply Assumption (EMB) and Theorem 6 to α' instead of α . We obtain that the inclusion map $I_{\pi}^{q_{\alpha',\beta}}$ is bounded, where we recall that $q_{\alpha',\beta}$ is defined to be $\frac{2\alpha'}{\alpha'-\beta}$. Since $x \mapsto \frac{2x}{x-\beta}$ is monotonically decreasing for $x > \beta$, we obtain the inequality

$$\frac{2\alpha'}{\alpha'-\beta} > \frac{2\alpha}{\alpha-\beta}$$

We choose $t = n^{\frac{1}{q_{\alpha,\beta}}}$. By construction, t satisfies the assumptions in Lemma 14. Furthermore, the assumptions of Lemma 12 are satisfied, with $F_* \in L_{q'}(\pi, \mathcal{Y})$, and $t = n^{\frac{1}{q_{\alpha,\beta}}}$.

Having established the applicability of Lemma 14 and Lemma 12, let us turn our attention to proving the results of the Theorem. Denote

$$\xi(x,y) = (y - C_{\lambda}\phi(x)) \otimes \left(C_{X,\lambda}^{-\frac{1}{2}}\phi(x)\right)$$

We compute

$$\mathbb{E}[\xi(x,y)] = (C_{YX} - C_{\lambda}C_X)C_{X,\lambda}^{-\frac{1}{2}}$$

1. The $\beta + p > \alpha$ case. Have

$$\begin{aligned} \left\| \frac{1}{n} \sum_{i=1}^{n} \xi_{i} - \mathbb{E}[\xi(x,y)] \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} &\leq \left\| \frac{1}{n} \sum_{i=1}^{n} \xi_{i} \mathbb{1}\{x_{i} \in \Omega_{t}\} - \mathbb{E}[\xi(x,y)\mathbb{1}\{x \in \Omega_{t}\}] \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} \\ &+ \left\| \frac{1}{n} \sum_{i=1}^{n} \xi_{i} \mathbb{1}\{x_{i} \in \Omega_{t}^{c}\} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} \\ &+ \left\| \mathbb{E}[\xi(x,y)\mathbb{1}\{x \in \Omega_{t}^{c}\} \|_{S_{2}(\mathcal{H},\mathcal{Y})} \end{aligned}$$

We can bound the first term with probability at least $1 - 2e^{-\tau}$ by $c\tau \left(\frac{n}{\log^{\theta}(n)}\right)^{-\frac{\beta}{2\alpha}}$, according to Lemma 14. By Lemma 12, with probability at least $1 - e^{-\tau}$, for sufficiently large $n, x_i \in \Omega_t$ for all $i \in [n]$, whereby the second term is zero. It remains to bound the third term, where our bound will be deterministic. Using Jensen's inequality, we have,

$$\begin{split} \|\mathbb{E}[\xi(x,y)\mathbb{1}\left\{x\in\Omega_{t}^{c}\right\}]\|_{S_{2}(\mathcal{H},\mathcal{Y})} &\leq \mathbb{E}\left[\|\xi(x,y)\mathbb{1}\left\{x\in\Omega_{t}^{c}\right\}\|_{S_{2}(\mathcal{H},\mathcal{Y})}\right] \\ &= \mathbb{E}\left[\|(y-C_{\lambda}\phi(x))\mathbb{1}\left\{x\notin\Omega_{t}\right\}\|_{\mathcal{Y}} \cdot \left\|C_{X,\lambda}^{-\frac{1}{2}}\phi(x)\right\|_{\mathcal{H}}\right] \\ &\leq A\lambda^{-\frac{\alpha}{2}}\mathbb{E}\left[\|(y-C_{\lambda}\phi(x))\mathbb{1}\left\{x\notin\Omega_{t}\right\}\|_{\mathcal{Y}}\right] \end{split}$$

where in the third line we used Lemma 17. We first split the second term into an approximation error and a noise term using triangle inequality.

$$\mathbb{E}\left[\left\|\left(y - C_{\lambda}\phi(x)\right)\mathbb{1}\left\{x \notin \Omega_{t}\right\}\right\|_{\mathcal{Y}}\right] \leq \mathbb{E}\left[\left\|\left(y - F_{*}(x)\right)\mathbb{1}\left\{x \notin \Omega_{t}\right\}\right\|_{\mathcal{Y}}\right] + \mathbb{E}\left[\left\|\left(F_{*}(x) - F_{\lambda}(x)\right)\mathbb{1}\left\{x \notin \Omega_{t}\right\}\right\|_{\mathcal{Y}}\right]\right]$$

We bound the first term using the tower property of conditional expectation,

$$\mathbb{E}\left[\|(y - C_{\lambda}\phi(x))\mathbb{1}\left\{x \notin \Omega_{t}\right\}\|_{\mathcal{Y}}\right] \leq \mathbb{E}_{\pi}\left[\mathbb{E}\left[\|y - C_{\lambda}\phi(x)\|_{\mathcal{Y}} \mid x\right]\mathbb{1}\left\{x \notin \Omega_{t}\right\}\right]$$
$$\leq \mathbb{E}_{\pi}\left[\mathbb{E}\left[\|y - C_{\lambda}\phi(x)\|_{\mathcal{Y}}^{2} \mid x\right]^{\frac{1}{2}}\mathbb{1}\left\{x \notin \Omega_{t}\right\}\right]$$
$$\leq \sigma\pi(x \notin \Omega_{t})$$
$$\leq \frac{\sigma c_{q}^{q}}{t^{q}}.$$

where in the third inequality we used Assumption (MOM) with q = 2, and in the fourth inequality we used Lemma 12. We bound the second term using Cauchy-Schwarz inequality and Lemma 10 with $\gamma = 0$,

$$\mathbb{E}\left[\|\left(F_{*}(x)-F_{\lambda}(x)\right)\mathbb{1}\left\{x\notin\Omega_{t}\right\}\|_{\mathcal{V}}\right] \leq \mathbb{P}(x\notin\Omega_{t})^{\frac{1}{2}}\|F_{*}\|_{\beta}\omega_{\rho}\lambda^{\frac{\beta}{2}}$$

Therefore, using Lemma 12, we have,

$$\|\mathbb{E}[\xi(x,y)\mathbb{1}\left\{x\in\Omega_{t}^{c}\right\}\|_{S_{2}(\mathcal{H},\mathcal{Y})} \leq A\lambda^{-\frac{\alpha}{2}}\left(\frac{\sigma c_{q}^{q}}{t^{q}} + \frac{c_{q}^{\frac{q}{2}}}{t^{\frac{q}{2}}}\|F_{*}\|_{\beta}\omega_{\rho}\lambda^{\frac{\beta}{2}}\right).$$
(36)

We now plug in $\lambda \approx n^{-\frac{1}{\beta+p}}$. Recall that by construction $t > n^{\frac{1}{q}}$. Thus,

$$\begin{split} \lambda^{-\frac{\alpha}{2}} t^{-q} &\lesssim n^{-1} n^{\frac{\alpha}{2(\beta+p)}} < n^{-1} n^{\frac{\beta+p}{2(\beta+p)}} = n^{-\frac{1}{2}} \le n^{-\frac{1}{2}\frac{\beta}{\beta+p}} \\ \lambda^{\frac{\beta-\alpha}{2}} t^{-\frac{q}{2}} &\lesssim n^{-\frac{1}{2}} n^{\frac{-(\beta-\alpha)}{2(\beta+p)}} < n^{-\frac{1}{2}} n^{\frac{p}{2(\beta+p)}} = n^{-\frac{1}{2}\frac{\beta}{\beta+p}} \end{split}$$

We've therefore proved inequality (34).

2. The $\beta + p \leq \alpha$ case. We proceed similarly to the $\beta + p > \alpha$ case. We have,

$$\begin{split} \left\| \frac{1}{n} \sum_{i=1}^{n} \xi_{i} - \mathbb{E}[\xi(x,y)] \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} &\leq \left\| \frac{1}{n} \sum_{i=1}^{n} \xi_{i} \mathbb{1}\{x_{i} \in \Omega_{t}\} - \mathbb{E}[\xi(x,y)\mathbb{1}\{x \in \Omega_{t}\}] \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} \\ &+ \left\| \frac{1}{n} \sum_{i=1}^{n} \xi_{i} \mathbb{1}\{x_{i} \in \Omega_{t}^{c}\} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} \\ &+ \left\| \mathbb{E}[\xi(x,y)\mathbb{1}\{x \in \Omega_{t}^{c}\}] \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} \end{split}$$

We can bound the first term with probability at least $1 - 2e^{-\tau}$ by $c\tau \left(\frac{n}{\log^{\theta}(n)}\right)^{-\frac{\beta}{2\alpha}}$, according to Lemma 14. By Lemma 12, with probability at least $1 - e^{-\tau}$, for sufficiently large n, $x_i \in \Omega_t$ for all $i \in [n]$, whereby the second term is zero. We bound the third term by Eq. (36). We now plug in $\lambda \asymp \left(\frac{n}{\log^{\theta}(n)}\right)^{-\frac{1}{\alpha}}$. Recall that by construction $t > n^{\frac{1}{q}}$. Thus,

$$\lambda^{-\frac{\alpha}{2}}t^{-q} \lesssim n^{-1} \left(\frac{n}{\log^{\theta}(n)}\right)^{\frac{1}{2}} < \left(\frac{n}{\log^{\theta}(n)}\right)^{-\frac{1}{2}} \le \left(\frac{n}{\log^{\theta}(n)}\right)^{-\frac{\beta}{2\alpha}}$$
$$\lambda^{\frac{\beta-\alpha}{2}}t^{-\frac{q}{2}} \lesssim n^{-\frac{1}{2}} \left(\frac{n}{\log^{\theta}(n)}\right)^{\frac{\alpha-\beta}{2\alpha}} < \left(\frac{n}{\log^{\theta}(n)}\right)^{\frac{-\beta}{2\alpha}}$$

We have therefore proved inequality (35).

We adapt Zhang et al. (2023b, Theorem 16) to the vector-valued setting.

Theorem 9 (Bound of estimation error). Suppose that assumptions (EMB), (EVD), (MOM) and (SRC) hold for $0 \le \beta \le 2\rho$, where ρ is the qualification, and $p \le \alpha < 1$. For $0 \le \gamma \le 1$, with $\gamma \le \beta$,

1. In the case of $\beta + p > \alpha$, by choosing $\lambda \asymp n^{-\frac{1}{\beta+p}}$, for any fixed $\tau \ge \log(4)$, when n is sufficiently large, with probability at least $1 - 4e^{-\tau}$, we have

$$\|[\hat{C}_{\lambda} - C_{\lambda}]\|_{S_2([\mathcal{H}]^{\gamma}, \mathcal{Y})}^2 \le c\tau^2 n^{-\frac{\beta-\gamma}{\beta+p}},$$

where c is a constant independent of n, τ .

2. In the case of $\beta + p \leq \alpha$, by choosing $\lambda \asymp \left(\frac{n}{\log^{\theta}(n)}\right)^{-\frac{1}{\alpha}}$, for any fixed $\tau \geq \log(4)$, when n is sufficiently large, with probability at least $1 - 4e^{-\tau}$, we have

$$\| [\hat{C}_{\lambda} - C_{\lambda}] \|_{S_{2}([\mathcal{H}]^{\gamma}, \mathcal{Y})}^{2} \leq c\tau^{2} \left(\frac{n}{\log^{\theta}(n)} \right)^{-\frac{\beta - \gamma}{\alpha}}$$

where c is a constant independent of n, τ .

Proof. Firstly, we establish the applicability of Lemma 19.

1. The $\beta + p > \alpha$ case. Have $\lambda \asymp n^{-\frac{1}{\beta+p}}$, hence

$$n\lambda^{\alpha} \gtrsim n^{\frac{\beta+p-\alpha}{\beta+p}}$$

whereas using $\lambda \leq \|C_X\|_{\mathcal{H} \to \mathcal{H}}$ for sufficiently large *n*, as well as Lemma 16,

$$8A^{2}\tau\log\left(2e\mathcal{N}(\lambda)\frac{\|C_{X}\|_{\mathcal{H}\to\mathcal{H}}+\lambda}{\|C_{X}\|_{\mathcal{H}\to\mathcal{H}}}\right) \leq 8A^{2}\tau\log(4ec_{2,1}\lambda^{-p}) \leq 8A^{2}\tau\left(\log(4ec_{2,1})+\frac{p}{\beta+p}\log(n)\right)$$

Therefore, for a fixed $\tau > 0$, for all sufficiently large n, Eq. (39) in Lemma 19 is satisfied.

2. The $\beta + p \leq \alpha$ case. Have $\lambda \asymp \left(\frac{n}{\log^{\theta}(n)}\right)^{-\frac{1}{\alpha}}$ for some $\theta > 1$, hence

$$n\lambda^{\alpha} \ge \log^{\theta}(n)$$

whereas similar to the $\beta+p>\alpha$ case, we alve

$$8A^{2}\tau\log\left(2e\mathcal{N}(\lambda)\frac{\|C_{X}\|_{\mathcal{H}\to\mathcal{H}}+\lambda}{\|C_{X}\|_{\mathcal{H}\to\mathcal{H}}}\right) \lesssim 8A^{2}\tau\left(\log(4ec_{2,1})+\frac{p}{\alpha}\log\left(\frac{n}{\log^{\theta}(n)}\right)\right)$$

Therefore, for a fixed $\tau > 0$, for all sufficiently large n, Eq. (39) in Lemma 19 is satisfied.

We thus conclude for all $\alpha \in (0,1]$, with probability $\geq 1 - 2e^{-\tau}$, Eq. (40) and (41) are satisfied simultaneously. We exploit the following decomposition

$$\begin{split} \left\| \begin{bmatrix} \hat{C}_{\lambda} - C_{\lambda} \end{bmatrix} \right\|_{S_{2}([\mathcal{H}]^{\gamma}, \mathcal{Y})} &\leq \left\| (\hat{C}_{\lambda} - C_{\lambda}) C_{X}^{\frac{1-\gamma}{2}} \right\|_{S_{2}(\mathcal{H}, \mathcal{Y})} \\ &\leq \left\| (\hat{C}_{\lambda} - C_{\lambda}) \hat{C}_{X, \lambda}^{\frac{1}{2}} \right\|_{S_{2}(\mathcal{H}, \mathcal{Y})} \cdot \left\| \hat{C}_{X, \lambda}^{-\frac{1}{2}} C_{X, \lambda}^{\frac{1}{2}} \right\|_{\mathcal{H} \to \mathcal{H}} \cdot \left\| C_{X, \lambda}^{-\frac{1}{2}} C_{X}^{\frac{1-\gamma}{2}} \right\|_{\mathcal{H} \to \mathcal{H}} \\ &\leq \left\| (\hat{C}_{\lambda} - C_{\lambda}) \hat{C}_{X, \lambda}^{\frac{1}{2}} \right\|_{S_{2}(\mathcal{H}, \mathcal{Y})} \cdot 3 \cdot \sup_{i \in \mathbb{N}} \frac{\mu_{i}^{\frac{1-\gamma}{2}}}{\sqrt{\mu_{i} + \lambda}} \\ &\leq \left\| (\hat{C}_{\lambda} - C_{\lambda}) \hat{C}_{X, \lambda}^{\frac{1}{2}} \right\|_{S_{2}(\mathcal{H}, \mathcal{Y})} \cdot 3\lambda^{-\frac{\gamma}{2}}, \end{split}$$

where in the first inequality we used Lemma 22, in the third inequality we used Eq. (41) and in the last inequality we used Lemma 21. We consider the following decomposition

$$\hat{C}_{\lambda} - C_{\lambda} = \hat{C}_{\lambda} - C_{\lambda} \left(\hat{C}_X g_{\lambda} \left(\hat{C}_X \right) + r_{\lambda} \left(\hat{C}_X \right) \right) = \left(\hat{C}_{YX} - C_{\lambda} \hat{C}_X \right) g_{\lambda} \left(\hat{C}_X \right) - C_{\lambda} r_{\lambda} \left(\hat{C}_X \right).$$

Hence

$$\left\| \left[\hat{C}_{\lambda} - C_{\lambda} \right] \right\|_{S_{2}([\mathcal{H}]^{\gamma}, \mathcal{Y})}^{2} \leq 18\lambda^{-\gamma} \left((I)^{2} + (II)^{2} \right),$$

where

$$(I) = \left\| \left(\hat{C}_{YX} - C_{\lambda} \hat{C}_{X} \right) \hat{C}_{X,\lambda}^{\frac{1}{2}} g_{\lambda} \left(\hat{C}_{X} \right) \right\|_{S_{2}(\mathcal{H},\mathcal{Y})}$$
$$(II) = \left\| C_{\lambda} r_{\lambda} \left(\hat{C}_{X} \right) \hat{C}_{X,\lambda}^{\frac{1}{2}} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})}.$$

Term (I). The high level idea is to bound (I) by exploiting the first axiom of the filter function (8), where $g_{\lambda}(\hat{C}_X)$ is intuitively a regularized inverse of \hat{C}_X , by grouping it with $\hat{C}_{X,\lambda}$.

$$(I) \leq \left\| \left(\hat{C}_{YX} - C_{\lambda} \hat{C}_{X} \right) C_{X,\lambda}^{-\frac{1}{2}} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} \cdot \left\| C_{X\lambda}^{\frac{1}{2}} \hat{C}_{X,\lambda}^{-\frac{1}{2}} \right\|_{\mathcal{H} \to \mathcal{H}} \cdot \left\| \hat{C}_{X,\lambda} g_{\lambda} \left(\hat{C}_{X} \right) \right\|_{\mathcal{H} \to \mathcal{H}}$$
$$\leq \left\| \left(\hat{C}_{YX} - C_{\lambda} \hat{C}_{X} \right) C_{X,\lambda}^{-\frac{1}{2}} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} \cdot \sqrt{3} \sup_{t \in [0,\kappa^{2}]} (t + \lambda) g_{\lambda}(t)$$
$$\leq \left\| \left(\hat{C}_{YX} - C_{\lambda} \hat{C}_{X} \right) C_{X,\lambda}^{-\frac{1}{2}} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})} \cdot 2\sqrt{3} E.$$

where the second inequality follows from Eq. (41). We consider the following decomposition

$$\begin{aligned} \left\| \left(\hat{C}_{YX} - C_{\lambda} \hat{C}_{X} \right) C_{X,\lambda}^{-\frac{1}{2}} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})}^{2} &\leq 2 \left\| \left(\left(\hat{C}_{YX} - C_{\lambda} \hat{C}_{X} \right) - \left(C_{YX} - C_{\lambda} C_{X} \right) \right) C_{X,\lambda}^{-\frac{1}{2}} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})}^{2} \\ &+ 2 \left\| \left(C_{YX} - C_{\lambda} C_{X} \right) C_{X,\lambda}^{-\frac{1}{2}} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})}^{2} \end{aligned}$$

We bound the first term by Theorem 8 and the second term by Lemma 8. This yields, for $\tau \ge \log(4)$, with probability at least $1 - 4e^{-\tau}$, for some constant c > 0 which does not depend on n, τ, λ ,

$$\begin{split} \left\| \left(\hat{C}_{YX} - C_{\lambda} \hat{C}_{X} \right) C_{X,\lambda}^{-\frac{1}{2}} \right\|_{S_{2}(\mathcal{H},\mathcal{Y})}^{2} &\leq 2\omega_{\rho}^{2} \|F_{*}\|_{\beta}^{2} \lambda^{\beta} + \begin{cases} c\tau^{2} n^{-\frac{\beta}{\beta+p}} & \beta+p \geq \alpha \\ c\tau^{2} \left(\frac{n}{\log^{\theta}(n)}\right)^{-\frac{\beta}{\alpha}} & \beta+p < \alpha \end{cases} \\ &\leq \begin{cases} \tau^{2} (c+2\|F_{*}\|_{\beta}^{2} \omega_{\rho}^{2} \lambda^{\beta}) n^{-\frac{\beta}{\beta+p}} & \beta+p \geq \alpha \\ \tau^{2} (c+2\|F_{*}\|_{\beta}^{2} \omega_{\rho}^{2} \lambda^{\beta}) \left(\frac{n}{\log^{\theta}(n)}\right)^{-\frac{\beta}{\alpha}} & \beta+p < \alpha \end{cases} \end{split}$$

where we used that $\tau > 1$. So collecting all the relevant constants together, we can write the upper bound of term (I) as follows: with probability at least $1 - 4e^{-\tau}$, for some constant c' > 0 (different from the c before) which does not depend on n, τ, λ , we have

$$(I) \le c'\tau \cdot \begin{cases} n^{-\frac{1}{2}\frac{\beta}{\beta+p}} & \beta+p \ge \alpha\\ \left(\frac{n}{\log^{\theta}(n)}\right)^{-\frac{\beta}{2\alpha}} & \beta+p < \alpha. \end{cases}$$

Term (II). Using Lemma 9, we have

$$(II) \leq B \left\| \hat{C}_{X,\lambda}^{\frac{1}{2}} r_{\lambda}(\hat{C}_{X}) g_{\lambda}(C_{X}) C_{X}^{\frac{\beta+1}{2}} \right\|_{\mathcal{H} \to \mathcal{H}}$$

The second term is the same as the scalar-valued case, which is bounded in Step 3 of the proof of Zhang et al. (2023b, Theorem 16). We define

$$\Delta_1 \coloneqq 32 \max\left\{\frac{\beta-1}{2}, 1\right\} E \omega_\rho \kappa^{\beta-1} \lambda^{\frac{1}{2}} n^{-\frac{\min(\beta,3)-1}{4}}$$

By the proof of Zhang et al. (2023b, Theorem 16), we have, with probability at least $1 - 6e^{-\tau}$

$$(II) \le 6B\omega_{\rho}E\lambda^{\frac{\beta}{2}} + \Delta_1B\tau \mathbb{1}\{\beta > 2\}.$$

1. Case $\beta + p > \alpha$. In this case $\lambda \asymp n^{-\frac{1}{\beta+p}}$. We note that for $\beta > 2$, Δ_1 as a function of n can be written as

$$\Delta_1 \asymp n^{-\frac{1}{2(\beta+p)}-\frac{\min(\beta,3)-1}{4}}$$

Note that

$$\frac{1}{2(\beta+p)} + \frac{\min(\beta,3) - 1}{4} - \frac{\beta}{2(\beta+p)} = \frac{1}{2} \left(\frac{p}{\beta+p} + \frac{\min(\beta,3) - 1}{2} \right) > 0$$

Hence

$$\Delta_1 \lesssim n^{-\frac{\beta}{2(\beta+p)}}$$

Therefore we have shown that there exists some constant c'' > 0, independent of n, λ, τ , such that with probability at least $1 - 6e^{-\tau}$, for sufficiently large n,

$$\left\| \left[\hat{C}_{\lambda} - C_{\lambda} \right] \right\|_{S_{2}([\mathcal{H}]^{\gamma}, \mathcal{Y})} \leq c'' \tau n^{-\frac{1}{2} \frac{\beta - \gamma}{\beta + p}}.$$

2. Case $\beta + p \leq \alpha$. In this case $\beta \leq \alpha \leq 1$, and $\lambda \asymp \left(\frac{n}{\log^{\theta}(n)}\right)^{-\frac{1}{\alpha}}$. We have also shown that there exists some constant c'' > 0, independent of n, λ, τ , such that with probability at least $1 - 6e^{-\tau}$, for sufficiently large n,

$$\left\| \left[\hat{C}_{\lambda} - C_{\lambda} \right] \right\|_{S_{2}([\mathcal{H}]^{\gamma}, \mathcal{Y})} \leq c'' \tau \left(\frac{n}{\log^{\theta}(n)} \right)^{-\frac{\beta - \gamma}{2\alpha}}$$

Putting together Lemma 10 and Theorem 9, we have proved Theorem 4.

D Auxiliary Results

D.1 Spectral Calculus, Proof of Proposition 1 and Empirical Solution

Definition 9 (Spectral Calculus; see Engl et al., 2000, Chapter 2.3). Let H be a Hilbert space. Consider $g: \mathbb{R} \to \mathbb{R}$ and a self-adjoint compact operator $A: H \to H$ admitting a spectral decomposition written as

$$A = \sum_{i \in I} \lambda_i h_i \otimes h_i.$$

We then define $g(A): H \to H$ as

$$g(A) \coloneqq \sum_{i \in I} g(\lambda_i) h_i \otimes h_i$$

whenever this series converges in operator norm.

Proof of Proposition 1. We define the sampling operator $S : \mathbb{R}^n \to \mathcal{H}$ and it dual $S^* : \mathcal{H} \to \mathbb{R}^n$,

$$S: \mathbb{R}^n \to \mathcal{H}, \qquad S^*: \mathcal{H} \to \mathbb{R}^n,$$

$$\alpha \mapsto \sum_{i=1}^n \alpha_i \phi(x_i) \qquad \qquad f \mapsto (f(x_i))_{i=1}^n$$

We can verify that $\hat{C}_X = n^{-1}SS^*$ and $\mathbf{K} = S^*S$. Let $\mathbf{Y} = (y_i)_{i=1}^n \in \mathbb{R}^n$. We have, for all $x \in \mathcal{X}$,

$$\hat{F}_{\lambda}(x) = \hat{C}_{\lambda}\phi(X)$$

$$= \left(\frac{1}{n}\sum_{i=1}^{n} y_{i} \otimes \phi(x_{i})\right)g_{\lambda}(\hat{C}_{X})\phi(X)$$

$$= \sum_{i=1}^{n} y_{i}\left(\phi(x_{i}), \frac{1}{n}g_{\lambda}(n^{-1}SS^{*})\phi(X)\right)_{\mathcal{H}}$$

$$= \mathbf{Y}^{T}S^{*}\left(\frac{1}{n}g_{\lambda}(n^{-1}SS^{*})\phi(X)\right)$$

$$(37)$$

$$\mathbf{Y}^{T} = \left(-\frac{1}{n}G^{*}G\right)G^{*}(X)$$

$$= \mathbf{Y}^{T} \frac{1}{n} g_{\lambda}(n^{-1}S^{*}S) S^{*} \phi(X)$$

$$= \mathbf{Y}^{T} \frac{1}{n} g_{\lambda}(n^{-1}\mathbf{K}) S^{*} \phi(X)$$
(38)

$$\mathbf{Y}^T \frac{1}{n} g_{\lambda}(n^{-1}K) \mathbf{k}_x.$$

=

To go from (37) to (38), we make the following observation. Consider the singular value decomposition of the compact operator S, there is $m \leq n$ such that

$$S = \sum_{i=1}^{m} \sqrt{\sigma_i} f_i \otimes e_i$$

where $(e_i)_i$, $(f_i)_i$ are orthonormal sequences in \mathbb{R}^n and \mathcal{H} respectively. We then have

$$SS^* = \sum_{i=1}^m \sigma_i f_i \otimes f_i, \quad S^*S = \sum_{i=1}^m \sigma_i e_i \otimes e_i.$$

Therefore, we deduce

$$S^* g_{\lambda} \left(\frac{SS^*}{n} \right) = \left(\sum_{i} \sqrt{\sigma_i} e_i \otimes f_i \right) \left(\sum_{j} g_{\lambda} \left(\frac{\sigma_j}{n} \right) f_j \otimes f_j \right)$$
$$= \sum_{i,j} g_{\lambda} \left(\frac{\sigma_j}{n} \right) \sqrt{\sigma_i} e_i \otimes f_j \langle f_i, f_j \rangle_{\mathcal{H}}$$
$$= \sum_{i} g_{\lambda} \left(\frac{\sigma_i}{n} \right) \sqrt{\sigma_i} e_i \otimes f_i$$

Similarly,

$$g_{\lambda}\left(\frac{S^{*}S}{n}\right)S^{*} = \left(\sum_{j}g_{\lambda}\left(\frac{\sigma_{j}}{n}\right)e_{j}\otimes e_{j}\right)\left(\sum_{i}\sqrt{\sigma_{i}}e_{i}\otimes f_{i}\right)$$
$$= \sum_{i,j}g_{\lambda}\left(\frac{\sigma_{j}}{n}\right)\sqrt{\sigma_{i}}e_{j}\otimes f_{i}\langle e_{i}, e_{j}\rangle_{\mathbb{R}^{n}}$$
$$= \sum_{i}g_{\lambda}\left(\frac{\sigma_{i}}{n}\right)\sqrt{\sigma_{i}}e_{i}\otimes f_{i}$$

Hence we have proved

$$S^*g_{\lambda}\left(\frac{SS^*}{n}\right) = g_{\lambda}\left(\frac{S^*S}{n}\right)S^*$$

as desired.

Proposition 2. Any minimizer $F \in \mathcal{G}$ of

$$\mathcal{E}_n(F) \coloneqq \frac{1}{n} \sum_{i=1}^n \|y_i - F(x_i)\|_{\mathcal{Y}}^2$$

on $\mathcal G$ must satisfy

$$\hat{C}_{YX} = \hat{C}\hat{C}_X, \qquad C \in S_2(\mathcal{H}, \mathcal{Y}),$$

where $F(\cdot) = C\phi(\cdot)$.

Proof. By Corollary 1, it is equivalent to solve the following optimization problem on $S_2(\mathcal{H}, \mathcal{Y})$,

$$\min_{C \in S_2(\mathcal{H},\mathcal{Y})} \frac{1}{n} \sum_{i=1}^n \|y_i - C\phi(x_i)\|_{\mathcal{Y}}^2.$$

Recall for a Hilbert-Schmidt operator $L \in S_2(\mathcal{H}, \mathcal{Y})$, we have

$$\langle L, a \otimes b \rangle_{S_2(\mathcal{H}, \mathcal{Y})} = \langle a, Lb \rangle_{\mathcal{Y}}.$$

	-	

Using this, we re-write the objective as an inner product in $S_2(\mathcal{H}, \mathcal{Y})$:

$$\frac{1}{n}\sum_{i=1}^{n} \|y_i - C\phi(x_i)\|_{\mathcal{Y}}^2 = \frac{1}{n}\sum_{i=1}^{n} -2\langle C, y_i \otimes \phi(x_i) \rangle_{S_2(\mathcal{H},\mathcal{Y})} + \langle C, (C\phi(x_i)) \otimes \phi(x_i) \rangle_{S_2(\mathcal{H},\mathcal{Y})} + \text{ constant}$$
$$= -2\langle C, \hat{C}_{YX} \rangle_{S_2(\mathcal{H},\mathcal{Y})} + \langle C, C\hat{C}_X \rangle_{S_2(\mathcal{H},\mathcal{Y})}$$

Taking the Fréchet derivative with respect to C and setting in to zero, we obtain the following first order condition

$$\hat{C}_{YX} = C\hat{C}_X$$

D.2 Properties Related to Assumptions (EMB) and (EVD)

Lemma 15 (Lemma 13 Fischer and Steinwart (2020)). Under (EMB), the following inequality is satisfied, for $\lambda > 0$ and π -almost all $x \in \mathcal{X}$,

$$\left\| \left(C_{XX} + \lambda I d_{\mathcal{H}} \right)^{-\frac{1}{2}} k(x, \cdot) \right\|_{\mathcal{H}} \leq A \lambda^{-\frac{\alpha}{2}}.$$

Definition 10 (*l*-effective dimension). For $l \ge 1$, the *l*-effective dimension $\mathcal{N}_l: (0, \infty) \to [0, \infty)$ is defined by

$$\mathcal{N}_{l}(\lambda) \coloneqq \operatorname{Tr}\left[C_{X}^{l}C_{X\lambda}^{-l}\right] = \sum_{i\geq 1} \left(\frac{\mu_{i}}{\mu_{i}+\lambda}\right)^{l}$$

The 1-effective dimension is widely considered in the statistical analysis of kernel ridge regression (see Caponnetto and De Vito (2007a), Blanchard and Mücke (2018), Lin and Cevher (2018), Lin et al. (2020a), Fischer and Steinwart (2020)). The following lemma provides upper and lower bounds for the l-effective dimension.

Lemma 16. Suppose Assumption (EVD) holds with parameter $p \in (0,1]$, for any $\lambda \in (0,1]$, there exists a constant $c_{2,l} > 0$ independent of λ such that

$$\mathcal{N}_l(\lambda) \leq c_{2,l} \lambda^{-p}.$$

If furthermore, Assumption (EVD+) holds with parameter $p \in (0,1)$, for any $\lambda \in (0,1]$, there exists a constant $c_{1,l} > 0$ independent of λ such that

$$c_{1,l}\lambda^{-p} \leq \mathcal{N}_l(\lambda) \leq c_{2,l}\lambda^{-p}.$$

The proof can be found in Li et al. (2023a) (Proposition D.1), but as the proof is incomplete we provide a full proof for completeness. This allows us to detect that the value p = 1 in Assumption (EVD+) is not compatible with the assumption of a bounded kernel (see Remark 7 below).

Proof.

$$\mathcal{N}_{l}(\lambda) \leq \sum_{i \geq 1} \left(\frac{D_{2}}{D_{2} + \lambda i^{\frac{1}{p}}} \right)^{l} \qquad (x \mapsto \frac{x}{x + \lambda} \text{ is monotonically increasing})$$

$$\leq \int_{0}^{+\infty} \left(\frac{D_{2}}{D_{2} + \lambda x^{\frac{1}{p}}} \right)^{l} dx \qquad (i \mapsto \left(\frac{D_{2}}{D_{2} + \lambda i^{1/p}} \right)^{l} \text{ is positive and decreasing})$$

$$\leq \left(\frac{D_{2}}{D_{2} + \lambda} \right)^{l} + \int_{1}^{+\infty} \left(\frac{D_{2}}{D_{2} + \lambda x^{\frac{1}{p}}} \right)^{l} dx$$

$$= \left(\frac{D_{2}}{D_{2} + \lambda} \right)^{l} + \int_{1}^{+\infty} \left(\frac{D_{2}}{D_{2} + y^{\frac{1}{p}}} \right)^{l} \frac{dy}{\lambda^{p}} \qquad (y^{1/p} = \lambda x^{1/p})$$

$$\leq 1 + \lambda^{-p} \int_{1}^{+\infty} \left(\frac{D_{2}}{D_{2} + y^{\frac{1}{p}}} \right)^{l} dy$$

Let us now consider the integral. Let us first consider $p \leq 1 < l,$

$$\int_{1}^{\infty} \left(\frac{D_2}{D_2 + y^{\frac{1}{p}}} \right)^l \mathrm{d}y \le D_2^l \int_{1}^{\infty} y^{-\frac{l}{p}} \mathrm{d}y$$
$$= D_2^l \frac{p}{l-p}$$

Therefore, using $\lambda \leq 1$, we can take $c_{2,l} = 1 + D_2^l \frac{p}{l-p}$. The remaining edge case p = 1 = l, is covered by Fischer and Steinwart (2020, Lemma 11) with $c_{2,1} = \|C_X\|_{S_1(\mathcal{H})}$. For the lower bound, we proceed similarly. For $p \in (0, 1)$ (and therefore p < l),

$$\mathcal{N}_{l}(\lambda) \geq \sum_{i\geq 1} \left(\frac{D_{1}}{D_{1}+\lambda i^{\frac{1}{p}}}\right)^{l} \qquad (x\mapsto \frac{x}{x+\lambda} \text{ is monotonically increasing})$$

$$\geq \int_{1}^{\infty} \left(\frac{D_{1}}{D_{1}+\lambda x^{\frac{1}{p}}}\right)^{l} dx \qquad (i\mapsto \left(\frac{D_{1}}{D_{1}+\lambda i^{1/p}}\right)^{l} \text{ is positive and decreasing})$$

$$= \int_{1}^{\infty} \left(\frac{D_{1}}{D_{1}+y^{\frac{1}{p}}}\right)^{l} \frac{dy}{\lambda^{p}} \qquad (y^{1/p} = \lambda x^{1/p})$$

$$\geq \lambda^{-p} \int_{1}^{\infty} \left(\frac{D_{1}}{D_{1}+1}\right)^{l} y^{-\frac{l}{p}} dy$$

$$= \lambda^{-p} \left(\frac{D_{1}}{D_{1}+1}\right)^{l} \frac{p}{l-p}.$$

Therefore, we can take $c_{1,l} = \left(\frac{D_1}{D_1+1}\right)^l \frac{p}{l-p}$.

Remark 7. We note that Assumption (EVD+) with p = 1 is not compatible with the assumption that k is bounded (Assumption 3). Indeed, suppose that $\mu_i \ge D_1 i^{-1}$, for all $i \ge 1$. Recall that $\{[e_i]\}_{i\ge 1}$ forms an orthonormal set in $L_2(\pi)$. By Mercer's theorem,

$$\kappa^2 \ge \int_{\mathcal{X}} k(x,x)\pi(\mathrm{d}x) = \sum_{i\ge 1} \mu_j \int_{\mathcal{X}} e_i(x)^2 \pi(\mathrm{d}x) = \sum_{i\ge 1} \mu_i \ge D_1 \sum_{i\ge 1} i^{-1} = +\infty,$$

which leads to a contradiction.

Lemma 17. For any $l \in [1, 2]$, the following equality holds,

$$\int_{\mathcal{X}} \left\| \left[C_{X,\lambda}^{-\frac{l}{2}} k(x,\cdot) \right] \right\|_{2-l}^{2} d\pi(x) = \mathcal{N}_{l}(\lambda).$$

In particular for l = 1,

$$\int_{\mathcal{X}} \left\| C_{X,\lambda}^{-\frac{1}{2}} k(x,\cdot) \right\|_{\mathcal{H}}^{2} d\pi(x) = \mathcal{N}_{1}(\lambda),$$
$$\int_{\mathcal{X}} \left\| \left[C_{X,\lambda}^{-1} k(x,\cdot) \right] \right\|_{L_{2}(\pi)}^{2} d\pi(x) = \mathcal{N}_{2}(\lambda).$$

and for l = 0,

Proof. Fix $x \in \mathcal{X}$. Since $k(x, \cdot) \in \mathcal{H}$, and $\left\{\mu_i^{1/2} e_i\right\}_{i \in I}$ is an ONB of $(\ker I_\pi)^{\perp}$, we have that π -almost everywhere

$$k(x,\cdot) = \sum_{i \in I} \langle k(x,\cdot), \mu_i^{1/2} e_i \rangle_{\mathcal{H}} \mu_i^{1/2} e_i = \sum_{i \in I} \mu_i e_i(x) e_i.$$

This is Mercer's Theorem (Steinwart and Scovel, 2012). On the other hand, π -almost everywhere,

$$C_{X,\lambda}^{-l/2} = \sum_{i \in I} (\mu_i + \lambda)^{-l/2} (\sqrt{\mu_i} e_i) \otimes (\sqrt{\mu_i} e_i).$$

Therefore,

$$\left[C_{X,\lambda}^{-l/2}k(x,\cdot)\right] = \sum_{i\in I} \frac{\mu_i}{(\mu_i + \lambda)^{l/2}} e_i(x)e_i,$$

and by Parseval's identity, using that $\{\mu_i^{(2-l)/2}[e_i]\}_{i\in I}$ is an ONB of $[\mathcal{H}]^{2-l}$,

$$\| [C_{X,\lambda}^{-l/2} k(x,\cdot)] \|_{2-l}^2 = \sum_{i \in I} \left(\frac{\mu_i}{\mu_i + \lambda} \right)^l e_i(x)^2$$

Therefore,

$$\int_{\mathcal{X}} \| [C_{X,\lambda}^{-l/2} k(x,\cdot)] \|_{2-l}^2 \mathrm{d}\pi(x) = \sum_{i \in I} \left(\frac{\mu_i}{\mu_i + \lambda} \right)^l \int_{\mathcal{X}} e_i(x)^2 \mathrm{d}\pi(x) = \mathcal{N}_l(\lambda),$$

where we used that $([e_i])_{i \in I}$ forms an orthonormal set in $L_2(\pi)$.

Theorem 10 (Theorem 3 by Li et al., 2023b). Let Assumption (EMB) be satisfied with parameter $\alpha \in (0, 1]$. For any $\beta \in (0, \alpha]$, the inclusion map $I_{\pi}^{\beta, q_{\alpha, \beta}} : [\mathcal{G}]^{\beta} \hookrightarrow L_{q_{\alpha, \beta}}(\pi, \mathcal{Y})$ is continuous, where $q_{\alpha, \beta} := \frac{2\alpha}{\alpha - \beta}$.

D.3 Concentration Inequalities

The following Theorem is from Fischer and Steinwart (2020, Theorem 26).

Theorem 11 (Bernstein's inequality). Let (Ω, \mathcal{B}, P) be a probability space, H be a separable Hilbert space, and $\xi : \Omega \to H$ be a random variable with

$$\mathbb{E}[\|\xi\|_H^m] \le \frac{1}{2}m!\sigma^2 L^{m-2}$$

for all $m \ge 2$. Then, for $\tau \ge 1$ and $n \ge 1$, the following concentration inequality is satisfied

$$P^{n}\left((\omega_{1},\ldots,\omega_{n})\in\Omega^{n}:\left\|\frac{1}{n}\sum_{i=1}^{n}\xi\left(\omega_{i}\right)-\mathbb{E}_{P}\xi\right\|_{H}^{2}\geq32\frac{\tau^{2}}{n}\left(\sigma^{2}+\frac{L^{2}}{n}\right)\right)\leq2e^{-\tau}.$$

In particular, for $\tau \ge 1$ and $n \ge 1$,

$$P^{n}\left((\omega_{1},\ldots,\omega_{n})\in\Omega^{n}:\left\|\frac{1}{n}\sum_{i=1}^{n}\xi\left(\omega_{i}\right)-\mathbb{E}_{P}\xi\right\|_{H}\geq4\sqrt{2}\frac{\tau}{\sqrt{n}}\left(\sigma+\frac{L}{\sqrt{n}}\right)\leq2e^{-\tau}.$$

Lemma 18. Let $\tau \ge \log(2)$, with probability at least $1 - 2e^{-\tau}$, for $\sqrt{n\lambda} \ge 8\tau\kappa\sqrt{\max\{\mathcal{N}(\lambda),1\}}$, we have

$$\left\|\hat{C}_{X,\lambda}^{-1}C_{X,\lambda}\right\|_{\mathcal{H}\to\mathcal{H}} \le 2$$

Proof. The proof is identical to Blanchard and Mücke (2018, Proposition 5.4) with the only difference that in their setting $\kappa = 1$.

Proposition 3 (Proposition C.9 Li et al., 2023a). Let π be a probability measure on $\mathcal{X}, f \in L_2(\pi)$ and $||f||_{\infty} \leq M$. Suppose we have x_1, \ldots, x_n sampled i.i.d. from π . Then, for any $\tau \geq \log(2)$, the following holds with probability at least $1 - 2e^{-\tau}$:

$$\frac{1}{2} \|f\|_{L_2(\pi)}^2 - \frac{5\tau M^2}{3n} \le \|f\|_{2,n}^2 \le \frac{3}{2} \|f\|_{L_2(\pi)}^2 + \frac{5\tau M^2}{3n},$$

where $\|\cdot\|_{2,n}$ was defined in Definition 7.

Lemma 19 (Lemma 12 Zhang et al., 2023b). Let Assumptions (EMB), (SRC) and (MOM) be satisfied. For $\tau \ge 1$, if λ and n satisfy that

$$n \ge 8A^2 \tau \lambda^{-\alpha} \log \left(2e\mathcal{N}(\lambda) \frac{\|C_X\|_{\mathcal{H} \to \mathcal{H}} + \lambda}{\|C_X\|_{\mathcal{H} \to \mathcal{H}}} \right)$$
(39)

then the following operator norm bounds are satisfied with probability not less than $1 - 2e^{-\tau}$

$$\left\|C_{X,\lambda}^{-\frac{1}{2}}\hat{C}_{X,\lambda}^{\frac{1}{2}}\right\|_{\mathcal{H}\to\mathcal{H}}^{2} \le 2,\tag{40}$$

$$\left\| C_{X,\lambda}^{\frac{1}{2}} \hat{C}_{X,\lambda}^{-\frac{1}{2}} \right\|_{\mathcal{H} \to \mathcal{H}}^{2} \le 3.$$

$$\tag{41}$$

D.4 Miscellaneous results

Lemma 20 (Cordes inequality Fujii et al. (1993)). Let A, B be two positive bounded linear operators on a separable Hilbert space H and $s \in [0, 1]$. Then

$$\|A^sB^s\|_{H\to H} \leq \|A\|^s_{H\to H} \|B\|^s_{H\to H}$$

Lemma 21 (Lemma 25 Fischer and Steinwart (2020)). For $\lambda > 0$ and $s \in [0, 1]$, we have

$$\sup_{t \ge 0} \frac{t^s}{t + \lambda} \le \lambda^{s-1}$$

We recall the following basic Lemma from Li et al. (2022, Lemma 2).

Lemma 22. For $0 \le \gamma \le 1$ and $F \in \mathcal{G}$, the inequality

$$\|[F]\|_{\gamma} \le \left\| CC_X^{\frac{1-\gamma}{2}} \right\|_{S_2(\mathcal{H},\mathcal{Y})}$$

holds, where $C = \overline{\Psi}^{-1}(F) \in S_2(\mathcal{H}, \mathcal{Y})$. If, in addition, $\gamma < 1$ or $C \perp \mathcal{Y} \otimes \ker I_{\pi}$ is satisfied, then the result is an equality.

Definition 11. Let $\mathcal{X} \subseteq \mathbb{R}^d$ be a compact set and $\theta \in (0,1]$. For a function $f : \mathcal{X} \to \mathbb{R}$, we introduce the Hölder semi-norm

$$[f]_{\theta,\mathcal{X}} \coloneqq \sup_{x,y \in \mathcal{X}, x \neq y} \frac{|f(x) - f(y)|}{\|x - y\|^{\theta}},$$

where $\|\cdot\|$ represents the usual Euclidean norm. Then, we define the Hölder space

$$C^{\theta}(\mathcal{X}) \coloneqq \{f : \mathcal{X} \to \mathbb{R} \mid [f]_{\theta, \mathcal{X}} < +\infty\},\$$

which is equipped with the norm

$$||f||_{C^{\theta}(\mathcal{X})} \coloneqq \sup_{x \in \mathcal{X}} |f(x)| + [f]_{\theta, \mathcal{X}}$$

The next lemma is used to prove Lemma 24 below. It appears in Li et al. (2023a, Lemma A.3), albeit the use of an erroneous equality in their proof: $||k(x,\cdot) - k(y,\cdot)||_{\mathcal{H}}^2 = k(x,x)k(y,y) - k(x,y)^2$. We therefore provide our own proof of this result.

Lemma 23. Assume that \mathcal{H} is an RKHS over a compact set $\mathcal{X} \subseteq \mathbb{R}^d$ associated with a kernel $k \in C^{\theta}(\mathcal{X} \times \mathcal{X})$ for $\theta \in (0,1]$. Then, we have $\mathcal{H} \subseteq C^{\frac{\theta}{2}}(\mathcal{X})$ and

$$[f]_{\frac{\theta}{2},\mathcal{X}} \le \sqrt{2[k]_{\theta,\mathcal{X}\times\mathcal{X}}} \|f\|_{\mathcal{H}}$$

Proof. For all $(x, y) \in \mathcal{X}$ and $f \in \mathcal{H}$, by the reproducing property and Cauchy–Schwarz inequality,

$$|f(x) - f(y)| = |\langle k(x, \cdot) - k(y, \cdot), f \rangle_{\mathcal{H}}| \le ||f||_{\mathcal{H}} ||k(x, \cdot) - k(y, \cdot)||_{\mathcal{H}}.$$

Then, using $k \in C^{\theta}(\mathcal{X} \times \mathcal{X})$, we obtain

$$||k(x,\cdot) - k(y,\cdot)||_{\mathcal{H}}^2 = k(x,x) + k(y,y) - 2k(x,y) \le 2[k]_{\theta,\mathcal{X}\times\mathcal{X}} ||x-y||^{\alpha},$$

which concludes the proof.

We derive as a corollary a quantitative upper bound on the ϵ -covering number of the set of (spectral) regularized kernel basis function with respect to the $\|\cdot\|_{\infty}$ norm.

Lemma 24 (Lemma C.10 Li et al., 2023a). Assume that \mathcal{H} is an RKHS over a compact set $\mathcal{X} \subseteq \mathbb{R}^d$ associated with a kernel $k \in C^{\theta}(\mathcal{X} \times \mathcal{X})$ for $\theta \in (0,1]$. Assume that $k(x,x) \leq \kappa^2$ for all $x \in \mathcal{X}$. Then, we have that for all $\epsilon > 0$,

$$\mathcal{N}(\mathcal{K}_{\lambda}, \|\cdot\|_{\infty}, \epsilon) \le c(\lambda\epsilon)^{-\frac{2d}{\theta}} \tag{42}$$

where $\mathcal{K}_{\lambda} \coloneqq \{C_{X,\lambda}^{-1}k(x,\cdot)\}_{x\in\mathcal{X}}$, and c is a positive constant which does not depend on λ, ϵ and only depends on κ and $[k]_{\theta,\mathcal{X}\times\mathcal{X}}$. $\mathcal{N}(\mathcal{K}_{\lambda}, \|\cdot\|_{\infty}, \epsilon)$ denotes the ϵ -covering number of the set \mathcal{K}_{λ} in the norm $\|\cdot\|_{\infty}$ (see Steinwart and Christmann (2008, Definition 6.19) for the definition of covering numbers).