# <span id="page-0-0"></span>The Collusion of Memory and Nonlinearity in Stochastic Approximation With Constant Stepsize

Dongyan (Lucy) Huo,§ Yixuan Zhang,† Yudong Chen,‡ Qiaomin Xie,†<sup>∗</sup>

§School of Operations Research and Information Engineering, Cornell University

†Department of Industrial and Systems Engineering, University of Wisconsin-Madison

‡Department of Computer Sciences, University of Wisconsin-Madison

#### Abstract

In this work, we investigate stochastic approximation (SA) with Markovian data and nonlinear updates under constant stepsize  $\alpha > 0$ . Existing work has primarily focused on either i.i.d. data or linear update rules. We take a new perspective and carefully examine the simultaneous presence of Markovian dependency of data and nonlinear update rules, delineating how the interplay between these two structures leads to complications that are not captured by prior techniques. By leveraging the smoothness and recurrence properties of the SA updates, we develop a fine-grained analysis of the correlation between the SA iterates  $\theta_k$  and Markovian data  $x_k$ . This enables us to overcome the obstacles in existing analysis and establish for the first time the weak convergence of the joint process  $(x_k, \theta_k)_{k>0}$ . Furthermore, we present a precise characterization of the asymptotic bias of the SA iterates, given by  $\mathbb{E}[\theta_{\infty}] - \theta^* = \alpha(b_m + b_n + b_c) + \mathcal{O}(\alpha^{3/2})$ . Here,  $b_m$  is associated with the Markovian noise,  $b_n$  is tied to the nonlinearity, and notably,  $b_c$  represents a multiplicative interaction between the Markovian noise and nonlinearity, which is absent in previous works. As a by-product of our analysis, we derive finite-time bounds on higher moment  $\mathbb{E}[\|\theta_k - \theta^*\|^{2p}]$  and present non-asymptotic geometric convergence rates for the iterates, along with a Central Limit Theorem.

### 1 Introduction

Stochastic Approximation (SA) is an iterative scheme for solving fixed-point equations using noisy observations. Its application spans various domains, including stochastic control [\[KY03,](#page-11-0) [Bor08](#page-9-0)], reinforcement learning (RL) [\[SB18,](#page-12-0) [Ber19\]](#page-9-1) and stochastic optimization [\[Lan20\]](#page-11-1). A typical SA algorithm takes the form  $\theta_{k+1} = \theta_k + \alpha g(\theta_k, x_k)$ , where  $(x_k)_{k\geq 0}$  represents the underlying noisy data sequence and  $\alpha > 0$  is the constant stepsize. The goal of SA is to approximate the target solution  $\theta^*$  that solves  $\mathbb{E}_{x \sim \pi}[g(\theta^*, x)] = 0$ , with  $\pi$  being the stationary distribution of the process  $(x_k)_{k\geq 0}$ .

SA subsumes many important algorithms. A prime example is stochastic gradient descent (SGD) for minimizing a function  $J(\theta)$  given a noisy estimate  $q(\theta, x)$  of its gradient. Linear SA schemes include SGD for quadratic objective functions, as well as various RL algorithms such as linear TD-Learning (in which g is not the gradient of any function and standard SGD results do not apply).

Of particular interest to us are SA updates given by a *nonlinear* function  $g(\theta, x)$  of  $\theta$ . One motivating example is learning a Generalized Linear Model (GLM)  $y \approx u(z^{\top} \theta)$  with a nonlinear mean function  $u : \mathbb{R} \to \mathbb{R}$ . A power approach, developed in [\[WZDD23,](#page-13-0) [KKSK11,](#page-11-2) [KS09](#page-11-3), [DGK](#page-10-0)+20], considers an appropriate surrogate loss function, for which the corresponding SGD update takes the form  $\theta_{k+1} = \theta_k + \alpha(\sigma(w_k^{\dagger} \theta_k) - y_k)w_k$ , where  $x_k = (w_k, y_k)$  is the observed covariate-response pair. Common choices of  $\sigma$  include the identity map for linear regression, the Sigmoid function for logistic regression, as well as Rectified Linear Unit (ReLU) and its various smoothed versions (e.g., ELU and SoftPlus) for ReLU regression [\[CUH16,](#page-10-1) [DGK](#page-10-0)<sup>+</sup>20, [GBB11,](#page-11-4) [BKBP22,](#page-9-2) [KKSK11\]](#page-11-2).

<sup>∗</sup>Emails: dh622@cornell.edu, yzhang2554@wisc.edu, yudong.chen@wisc.edu, qiaomin.xie@wisc.edu

<span id="page-1-1"></span>Furthermore, we are interested in the setting where the data sequence  $(x_k)_{k>0}$  forms a *Markov chain*, going beyond the common i.i.d. data setting. The Markovian model captures a wide range of SA problems in machine learning where stochastic data exhibit serial dependence  $[KOG+22, BSS+23, HCX23a, NWB+20].$  $[KOG+22, BSS+23, HCX23a, NWB+20].$ 

Classical work on SA focuses on diminishing stepsizes [\[RM51](#page-12-2), [BM00](#page-9-4)]. Constant stepsize schemes have recently gained popularity due to easy parameter tuning, fast initial convergence, and robust empirical performance. Non-asymptotic error bounds have been obtained for constant stepsize SA [\[SY19](#page-12-3), [CZD](#page-10-2)+22]. Recent work further provides fine-grained characterization of the distributional and steady-state behaviors of the iterates [\[DDB20,](#page-10-3) [YBVE21,](#page-13-1) [HCX23a](#page-11-6), [LM23,](#page-11-7) [ZX24\]](#page-13-2). Two recurring themes in these results are weak convergence of the distribution of  $\theta_t$  and the presence of an asymptotic bias  $\mathbb{E}[\theta_\infty] - \theta^* \propto \alpha$ , both having important implications for iterate averaging, bias reduction and statistical inference [\[HCX23b](#page-11-8)].

Note that most previous work studied the nonlinear update setting and Markovian data setting separately—e.g., in [\[DDB20](#page-10-3), [YBVE21\]](#page-13-1) for nonlinear SGD with i.i.d. data, and in [\[HCX23a](#page-11-6), [HCX23b\]](#page-11-8) for Markovian linear SA. The linearity or i.i.d. assumptions imposed in these prior works are restrictive, especially in the face of modern machine/reinforcement learning paradigms where nonlinear models are the norm and dependent data is common. Moreover, the absence of prior work dealing with Markovian nonlinear SA is not merely an overlook—as argued below, this setting is significantly more challenging.

**Our Contributions:** In this work, we study constant-stepsize SA with *both* Markovian data and nonlinear update. In Section [3,](#page-3-0) we elucidate the new challenges that arise from the simultaneous presence of these two structures, which break key steps in previous analysis of the i.i.d. or linear setting. Due to the interaction between these two structures, establishing weak convergence is far from obvious, and the asymptotic bias exhibits new behaviors. Consequently, analyzing the nonlinear Markovian setting requires more than simply combining previous techniques.

To address the above confounding complication, we exploit the smoothness and recurrence structures of the SA update, thereby developing a fine-grained analysis of the correlation of the parameter  $\theta_k$  and data  $x_k$ . This allows us to establish for the first time the weak convergence of the joint process  $(x_k, \theta_k)_{k>0}$  to a unique invariant distribution, represented by the limiting random variable  $(x_{\infty}, \theta_{\infty})$ . As a by-product of our analysis, we derive finite-time bounds on  $\mathbb{E}[\|\theta_k - \theta^*\|^{2p}]$ , the 2p-th moments of the errors, generalizing the results in  $[DDB20, CZD+22, SY19]$  to higher moments and to the nonlinear Markovian setting. In addition, we prove a Central Limit Theorem (CLT) for averaged iterates.

We further show that nonlinearity and Markovian structure contribute in a *multiplicative* way to the asymptotic bias of the SA iterates. In particular, we obtain the following bias characterization:  $\mathbb{E}[\theta_{\infty}^{(\alpha)}]-\theta^* =$  $\alpha(b_m+b_n+b_c)+\mathcal{O}((\alpha\tau_\alpha)^{3/2})$ . We provide explicit expressions for the vectors  $b_m, b_n, b_c$ , which are independent of  $\alpha$ . Here,  $b_m$  represents the bias component due to Markovian data (quantified by the mixing property of  $x_k$ ), and  $b_n$  the bias due to the nonlinearity of g (quantified by the second derivative g''). Importantly, we identify the additional compound term  $b<sub>c</sub>$ , which is absent in both nonlinear SA with i.i.d. data and linear SA with Markovian data. We explore the algorithmic implications of the above results on Polyak-Ruppert (PR) averaging [\[Rup88,](#page-12-4) [Pol90,](#page-12-5) [JKK](#page-11-9)<sup>+</sup>18] and Richardson-Romberg (RR) extrapolation [\[Hil87](#page-11-10)]. We show that PR averaging reduces the variance but not the bias, whereas RR extrapolation eliminates the leading bias term  $\alpha(b_m + b_n + b_c)$ , reducing the asymptotic bias to a higher order of  $\alpha$ .

Related work Postponing a detailed literature review to Section [5,](#page-8-0) here we remark on the very recent work [\[LM23](#page-11-7)], which also studies Markovian nonlinear SA using coupling. They prove weak convergence of  $(x_t, \theta_t)$  only in the linear setting, not for nonlinear SA. In the latter setting, their weak convergence analysis is thwarted by challenges similar to what we elucidate in Section [3,](#page-3-0) due to the interplay between nonlinearity and Markovian data leading to "double recursions". The coupling techniques in [\[LM23\]](#page-11-7) and ours are also different. We couple two processes by sharing data  $x_t = x'_t$ , and construct  $\theta, \theta'$  such that  $(x_t, \theta_t) \stackrel{\text{d}}{=}$  $(x'_{t+1}, \theta'_{t+1})$ . In [\[LM23\]](#page-11-7) they initialize two processes with different  $x_0$  and  $x'_0$ , and analyzes the stopping time  $\tau$  when  $\theta_{\tau} = \theta_{\tau}'$ . Moreover, the work [\[LM23](#page-11-7)] and only presents an *upper bound* for asymptotic bias, while ours presents a fine-grained characterization in Theorem [4.6](#page-6-0) necessary for justifying RR-extrapolation.

<span id="page-1-0"></span>**Notations** The Euclidean norm is denoted by  $\|\cdot\|$ . We use  $B(\beta) := \{\theta \in \mathbb{R}^d : \|\theta\| \leq \beta\}$  to denote the ball with radius  $\beta$ .  $\mathcal{L}(z)$  denotes the distribution of a random vector z and Var(z) its covariance matrix. Let  $P_2(\mathbb{R}^d)$  be the space of square-integrable distributions on  $\mathbb{R}^d$  and  $\mathcal{P}_2(\mathcal{X} \times \mathbb{R}^d)$  be the space of distributions

<span id="page-2-4"></span> $\bar{\nu}$  on  $\mathcal{X} \times \mathbb{R}^d$  with square-integrable second marginal on  $\mathbb{R}^d$ . The Wasserstein-2 between two probability measures  $\mu$  and  $\nu$  in  $\mathcal{P}_2(\mathbb{R}^d)$  is defined as  $W_2(\mu,\nu) = \inf_{\psi \in \Pi(\mu,\nu)} \left\{ \left( \mathbb{E}[\|\theta - \theta'\|^2] \right)^{\frac{1}{2}} : \mathcal{L}(\theta) = \mu, \mathcal{L}(\theta') = \nu \right\},$ where  $\Pi(\mu, \nu)$  denotes the set of all couplings between  $\mu$  and  $\nu$ . Extending to the space  $\mathcal{X} \times \mathbb{R}^d$ , we define the  $\text{metric}_{\mathcal{I}}(\vec{d}((x,\theta), (x', \theta')) := \sqrt{\mathbb{1}\{x \neq x'\} + \|\theta - \theta'\|^2}$ , and denote by  $\bar{W}_2$  the extended Wasserstein-2 distance w.r.t.  $\overline{d}$ .

The lowercase letter c and its derivatives  $c', c_0$ , etc. denote universal numerical constants, whose value may change from line to line. We use  $s \equiv s(\theta_0, \theta^*, \mu, L, R)$  and its derivatives to denote quantities (scalars, vectors, or matrices) that are independent of the stepsize  $\alpha$  and the iteration index k, but may depend on the initialization  $\theta_0$ , SA primitives  $\theta^*, \mu$  and L, and the coefficient R for the geometric mixing rate of  $(x_k)$ in Assumption [1.](#page-2-0) As we are primarily interested in dependence on  $\alpha$  and k, we adopt the following big-O notation:  $||f|| = \mathcal{O}(h(\alpha, k))$  if it holds that  $||f|| \leq s \cdot ||h(\alpha, k)||$ .

### 2 Problem Setup and Preliminaries

Let  $(x_k)_{k>0}$  be a Markov chain on a general state space X. Consider the following projected stochastic approximation (SA) iteration:

<span id="page-2-1"></span>
$$
\theta_{k+1}^{(\alpha)} = \Pi_{B(\beta)} \left[ \theta_k^{(\alpha)} + \alpha \big( g(\theta_k^{(\alpha)}, x_k) + \xi_{k+1}(\theta_k^{(\alpha)}) \big) \right],\tag{2.1}
$$

where  $g: \mathbb{R}^d \times \mathcal{X} \to \mathbb{R}^d$  is a deterministic function,  $\{\xi_k\}_{k\geq 1}$  are i.i.d. zero-mean random fields,  $\alpha > 0$  is a constant stepsize, and  $\Pi_{B(\beta)}(\theta) := \arg \min_{z: \|z\| \le \beta} \|z - \theta\|$  is the projection operator. We shall omit the superscript  $(\alpha)$  in  $\theta_k$  when the dependence on  $\alpha$  is clear from the context. In this work, we also consider the projection-free variant of the iteration  $(2.1)$  with  $\beta = \infty$ .

We denote by  $\pi$  the stationary distribution of the Markov chain  $(x_k)_{k>1}$  and define the shorthand  $\bar{g}(\theta) := \mathbb{E}_{\pi}[g(\theta, x)]$ , where  $\mathbb{E}_{\pi}[\cdot]$  denotes the expectation with respect to  $x \sim \pi$ . The algorithm [\(2.1\)](#page-2-1) computes an estimation of the target vector  $\theta^*$  that solves the steady-state equation  $\mathbb{E}_{\pi}[g(\theta,x)] = 0$ . Our general goal is to characterize the relationship between the iterate  $\theta_k$  and the target solution  $\theta^*$ .

In the following, we state the assumptions needed for our main results.

<span id="page-2-0"></span>**Assumption 1** (Uniform Ergodicity).  $(x_k)_{k>0}$  is a uniformly ergodic Markov chain on a Borel state space  $(\mathcal{X}, \mathcal{B}(\mathcal{X}))$  with transition kernel P and a unique stationary distribution  $\pi$ . That is, there exist constants  $r \in [0,1)$  and  $R > 0$  such that  $||P^k(x, \cdot) - \pi||_{TV} \leq Rr^k, \forall x \in \mathcal{X}$ .

All irreducible, aperiodic, and finite state space Markov chains are uniformly ergodic. The uniform ergodicity assumption is common in prior work on SA with Markovian noise [\[BRS21](#page-9-5), [DT22](#page-10-4), [DMNS22,](#page-10-5) [HCX23a](#page-11-6), [LLZ23\]](#page-11-11). Relaxing this uniform ergodicity assumption, in the style of [\[MPWB21](#page-12-6), [SY19](#page-12-3), [LM23\]](#page-11-7) is possible but orthogonal to our focus, and thus we do not pursue this direction in this work.

We allow the chain  $(x_k)_{k\geq 0}$  to be arbitrarily initialized rather than from the stationary distribution  $\pi$ . An important quantity is the mixing time of the Markov chain, defined as follows.

<span id="page-2-5"></span>**Definition 2.1.** For  $\epsilon \in (0,1)$ , the  $\epsilon$ -mixing time of  $(x_k)_{k\geq 0}$ , denoted by  $\tau_{\epsilon} \geq 1$ , is defined as  $\tau_{\epsilon} := \min\left\{k \geq 1\right\}$ 1 :  $\sup_{x \in X} ||P^k(x, \cdot) - \pi||_{\text{TV}} \leq \epsilon$ .

Under Assumption [1,](#page-2-0) the  $\epsilon$ -mixing time satisfies  $\tau_{\epsilon} \leq K \log \frac{1}{\epsilon}$  for all  $\epsilon \in (0,1)$ , where  $K \geq 1$  is independent of  $\epsilon$ . In the sequel, unless otherwise specified, we always choose  $\epsilon = \alpha$  and let  $\tau \equiv \tau_{\alpha}$ .

The following assumptions on the nonlinear function g in  $(2.1)$  is standard in the literature [\[DDB20,](#page-10-3) [CZD](#page-10-2)<sup>+</sup>22, [LM23](#page-11-7), [LLZ23,](#page-11-11) [HIMM19](#page-11-12). A wide family of g functions satisfies these assumptions, with the  $L_2$ -regularized logistic regression of GLM being a standard example.

<span id="page-2-2"></span>**Assumption 2** (Differentiability and Linear Growth). For each  $x \in \mathcal{X}$ , the function  $g(\theta, x)$  is three times continuously differentiable in  $\theta$  with uniformly bounded first to third derivatives, i.e.,  $\sup_{\theta \in \mathbb{R}^d} ||g^{(i)}(\theta, x)||$  $+\infty$  for  $i = 1, 2, 3, x \in \mathcal{X}$ . Moreover, there exists a constant  $L_1 > 0$  such that  $(1) \| g^{(i)}(\theta, x) - g^{(i)}(\theta', x) \| \leq L_1$ , for all  $\theta, \theta' \in \mathbb{R}^d$ ,  $i = 0, 1, 2$  and  $x \in \mathcal{X}$ , and  $(2) ||g(0, x)|| \le L_1$  for all  $x \in \mathcal{X}$ .

<span id="page-2-3"></span>Assumption 3 (Strong Monotonicity). There exists  $\mu > 0$  such that  $\langle \theta - \theta', \bar{g}(\theta) - \bar{g}(\theta') \rangle \le -\mu ||\theta - \theta'||^2$ ,  $\forall \theta, \theta' \in \mathbb{R}^d$ . Consequently, the target equation  $\bar{g}(\theta) = 0$  has a unique solution  $\theta^*$ .

<span id="page-3-5"></span>Assumption [2](#page-2-2) implies that  $g(\theta, x)$  is L<sub>1</sub>-Lipschitz w.r.t.  $\theta$  uniformly in x. When g is a linear function, i.e.,  $g(\theta, x) = A(x)\theta + b(x)$ , this assumption is satisfied with  $\sup_{x \in \mathcal{X}} \|A(x)\| < \infty$  and  $\sup_{x \in \mathcal{X}} \|b(x)\| < \infty$ , which are commonly assumed for linear SA. The above assumption immediately implies that the growth rate of  $||g||$ and  $\|\bar{g}\|$  will be at most linear in  $\theta$ , i.e.,  $\|g(\theta, x)\| \le L_1(\|\theta - \theta^*\| + 1)$  and  $\|\bar{g}(\theta)\| \le L_1(\|\theta - \theta^*\| + 1)$ . When g is a gradient field, Assumption [3](#page-2-3) is equivalent to strong convexity. For notational simplicity, we assume the strong monotonicity parameter satisfies  $\mu \leq 1-r$ , where r is the convergence factor in Assumption [1.](#page-2-0) For general  $\mu$ , our results remain valid with  $\mu$  replaced by min{ $\mu$ , 1 – r}.

We next consider the noise. Denote by  $\mathcal{F}_k$  the filtration generated by  $\{x_t, \theta_t, \xi_{t+1}\}_{t=0}^{k-1} \cup \{x_k, \theta_k\}.$ 

<span id="page-3-4"></span>**Assumption 4** (Noise Sequence). Let  $p \in \mathbb{Z}_+$  be given. The noise sequence  $(\xi_k)_{k>1}$  is a collection of i.i.d. random fields satisfying the following conditions with  $L_{2,p} > 0$ :

<span id="page-3-1"></span>
$$
\mathbb{E}[\xi_{k+1}(\theta)|\mathcal{F}_k] = 0 \quad \text{and} \quad \mathbb{E}^{1/(2p)}[\|\xi_1(\theta)\|^{2p}] \le L_{2,p}(\|\theta - \theta^*\| + 1), \quad \forall \theta \in \mathbb{R}^d. \tag{2.2}
$$

 $Define C(\theta) = \mathbb{E}[\xi_1(\theta)^{\otimes 2}]$  and assume that  $C(\theta)$  is at least twice differentiable. There also exist  $M_{\epsilon}, k_{\epsilon} \geq 0$ such that for  $\theta \in \mathbb{R}^d$ , we have  $\max_{i=1,2} ||C^{(i)}(\theta)|| \leq M_{\epsilon} \{1 + ||\theta - \theta^*||^{k_{\epsilon}}\}.$ 

In the sequel, we set  $L := L_1 + L_2$ , and without loss of generality, we assume  $L \geq 1$ .

When  $p = 1$ , the second inequality in [\(2.2\)](#page-3-1) only requires linear growth in expectation, which relaxes the almost sure linear growth condition in  $\left[\text{CZD}+22\right]$ . The constraint on the covariance matrix  $C(\theta)$  is lenient and satisfied in most regular enough settings, as shown in [\[DDB20](#page-10-3)].

### <span id="page-3-0"></span>3 Analytical Challenges and Techniques

In this section, we elaborate on the challenges and techniques used to prove the above results.

Previous work has established weak convergence of  $(x_k, \theta_k)$  separately for nonlinear SA with i.i.d. data, and for Markovian linear SA. The high-level approaches used in two representative prior works can be summarized as follows. The work **DDB20** on nonlinear SGD leverages *local linearization* of g through Taylor expansion. The work [\[HCX23a\]](#page-11-6) on Markovian linear SA exploits the mixing property of the Markovian noise to regain approximate independence, particularly between  $x_k$  and  $\theta_{k-\tau}$  for sufficiently large  $\tau$ . It is tempting to expect that nonlinear SA can be analyzed by combining these two approaches. Perhaps surprisingly, such a simple combination would not work due to the interplay between nonlinearity and Markovian structures.

To demonstrate this challenge, let us seek to establish weak convergence in the Wasserstein distance  $W_2$ via forward coupling [\[FT98](#page-11-13)], an approach employed by both [\[DDB20,](#page-10-3) [HCX23a\]](#page-11-6) as well as others [\[DMN](#page-10-6)<sup>+</sup>21]. Specifically, we consider two SA iterate sequences  $(\theta_k^{[1]}$  $\binom{[1]}{k}_{k}$ <sub>k</sub> $\geq$ <sup>0</sup> and  $(\theta_k^{[2]}$  $\binom{[2]}{k}$ <sub>k</sub> $\geq 0$  from different initializations  $\theta_{0}^{[1]}$  and  $\theta_0^{[2]}$  coupled by sharing the data sequence  $(x_k)_{k\geq 0}$ :  $\theta_{k+1}^{[1]} = \theta_k^{[1]} + \alpha g(\theta_k^{[1]})$  $\theta_k^{[1]}, x_k$ ) and  $\theta_{k+1}^{[2]} = \theta_k^{[2]} + \alpha g(\theta_k^{[2]})$  $_{k}^{[2]},x_{k}).$ To establish convergence in  $W_2$ , we consider the difference sequence

<span id="page-3-2"></span>
$$
w_{k+1} := \theta_{k+1}^{[1]} - \theta_{k+1}^{[2]} = w_k + \alpha \big(g(\theta_k^{[1]}, x_k) - g(\theta_k^{[2]}, x_k)\big),\tag{3.1}
$$

and it suffices to prove  $w_k$  converges to 0 in mean square:  $\mathbb{E}[\|w_{k+1}\|^2] \lesssim \rho^k \mathbb{E}[\|w_0\|^2]$  for  $\rho < 1$ .

With this goal in mind and following the idea from [\[DDB20](#page-10-3)], one may first linearize the right-hand side of the difference dynamic  $(3.1)$  and obtain the approximation

<span id="page-3-3"></span>
$$
w_{k+1} \approx w_k + \alpha g'(\theta_k^{[2]}, x_k) w_k. \tag{3.2}
$$

Next, to analyze the drift of the Lyapunov function  $\mathbb{E}[\|w_k\|^2]$  and handle the Markovian noise  $(x_k)$ , we use the conditioning technique from  $[HCX23a]$ . We condition on the information of  $\tau$  steps before, denoted by  $\mathcal{F}_{k-\tau} := \sigma((\theta_t^{[1]}, \theta_t^{[2]}, x_t) : t \leq k - \tau)$ . Ignoring higher-order terms and assuming a one-dimensional problem for simplicity, we obtain that

$$
\mathbb{E}[\|w_{k+1}\|^2] \approx \mathbb{E}\Big[\mathbb{E}\big[\|w_k\|^2\big(1+2\alpha g'(\theta_k^{[2]}, x_k)\big) \mid \mathcal{F}_{k-\tau}\big]\Big] \approx \mathbb{E}\Big[\|w_{k-\tau}\|^2\big(1+2\alpha \mathbb{E}\big[g'(\theta_k^{[2]}, x_k) \mid \mathcal{F}_{k-\tau}\big]\big)\Big],
$$
\n(3.3)

<span id="page-4-1"></span>where we use  $w_k \approx w_{k-\tau}$  for small  $\alpha$  (this argument, which is made precise in [\[HCX23a,](#page-11-6) [SY19\]](#page-12-3), essentially exploits the fact that  $x_k$  evolves faster than  $\theta_k$ ).

To prove dynamic [\(3.3\)](#page-3-3) converges, it boils down to showing the "gain matrix"  $\mathbb{E}[g'(\theta_k^{[2]}$  $\left[ \begin{smallmatrix} [2]_k \ k \end{smallmatrix} \right], x_k$  |  $\left[ \begin{smallmatrix} \mathcal{F}_{k-\tau} \end{smallmatrix} \right]$  is negative/Hurwitz. To further simplify, we assume k is large so that the chain  $(x_k)$  is distributed per its stationary distribution  $\pi$ , in which case the gain matrix simplifies to  $\mathbb{E}_{x_{\infty}\sim\pi}[g'(\theta_{\infty}^{[2]},x_{\infty})].$ 

Analyzing this gain matrix is where our analysis diverges from previous work. If the SA update were linear, i.e.,  $g(\theta, x) = A(x)\theta$ , then the gain  $\mathbb{E}[g'(\theta_{\infty}^{[2]}, x_{\infty})] = \mathbb{E}_{\pi}[A(x_{\infty})]$  would be independent of  $\theta_{\infty}^{[2]}$ , and its Hurwitz property is a standard and necessary condition for proving convergence of linear SA. If the data sequence  $(x_k)$  were *i.i.d.*, then  $\theta_k$  would be *independent* of  $x_k$  and hence the gain becomes  $\mathbb{E}[g'(\theta_{\infty}^{[2]}, x_{\infty})] =$  $\mathbb{E}[\mathbb{E}[g'(\theta_{\infty}^{[2]},x_{\infty})|\theta_{\infty}^{[2]}]] = \mathbb{E}[\bar{g}'(\theta_{\infty}^{[2]})]$  with  $\bar{g}(\cdot) := \mathbb{E}_{x \sim \pi}[g(\cdot,x)]$ , where the Hurwitz property again follows from standard assumptions on  $\bar{q}$ .

However, both arguments fail for the Markovian nonlinear setting. Common assumptions for nonlinear SA only ensure Hurwitz  $\mathbb{E}_{x \sim \pi}[g'(\theta, x)|\theta]$  given  $\theta$ . This does not imply the desired Hurwitz  $\mathbb{E}[g'(\theta_{\infty}^{[2]}, x_{\infty})]$ , precisely owing to the simultaneous presence of (i) the *dependence* of g' on both  $\theta_{\infty}$  and  $x_{\infty}$  (due to nonlinearity) and (ii) the *correlation* between  $\theta_{\infty}$  and  $x_{\infty}$  (due to Markovian).

**Our approaches:** We overcome this challenge by carefully analyzing the properties of the above dependence and correlation. Therefore, for sufficiently large  $\tau$ , we further decompose [\(3.3\)](#page-3-3) as

$$
\mathbb{E}[\|w_{k+1}\|^2] \approx \mathbb{E}\Big[\|w_{k-\tau}\|^2 \Big(1 + 2\alpha \mathbb{E}\big[g'(\theta_k^{[2]}, x_k) \mid \mathcal{F}_{k-\tau}\big]\Big)\Big] \n= \mathbb{E}\Big[\|w_{k-\tau}\|^2 \Big(1 + 2\alpha \mathbb{E}\big[g'(\theta_{k-\tau}^{[2]}, x_k) \mid \mathcal{F}_{k-\tau}\big] + 2\alpha \big(\mathbb{E}\big[g'(\theta_k^{[2]}, x_k) - g'(\theta_{k-\tau}^{[2]}, x_k) \mid \mathcal{F}_{k-\tau}\big]\big)\Big)\Big] \n\approx \mathbb{E}[g'(\theta_{k-\tau}^{[2]}, x_{\infty})|\mathcal{F}_{k-\tau}] \text{ Hurwitz} \n\lesssim \rho \mathbb{E}[\|w_{k-\tau}\|^2] + \alpha \mathbb{E}\Big[\mathbb{E}\Big[\underbrace{\langle w_{k-\tau}, g(\theta_k^{[1]}, x_k) - g(\theta_k^{[2]}, x_k) - g(\theta_{k-\tau}^{[1]}, x_k) + g(\theta_{k-\tau}^{[2]}, x_k) \rangle}_{\blacklozenge}\big|\mathcal{F}_{k-\tau}\Big]\Big],
$$

where we approximate  $w_k \approx w_{k-\tau}$ ,  $w_t g'(\theta_t^{[2]}, x_k) \approx (g(\theta_t^{[1]}, x_k) - g(\theta_k^{[2]})$  $(k, k, k)$ ) for  $t = k, k - \tau$  and obtain the second term in the last inequality. Next, we propose employing two different Taylor expansions to prove that  $\bullet$  is of higher orders of  $\alpha$ . We first apply the Taylor expansion to  $g(\theta_k^{[1]}, x_k)$  – k  $g(\theta_k^{[2]}$  $_{k}^{[2]},x_{k})$  and  $g(\theta_{k-}^{[1]}$  $\binom{[1]}{k-\tau}, x_k$  –  $g(\theta_{k-\tau}^{[2]})$  $\binom{[2]}{k-1}$ ,  $x_k$ ). However, this only achieves  $\blacklozenge$   $\lesssim$   $||w_k||^2 (||w_k|| + \alpha \tau T_1)$ , where  $T_1 = \min(\|\theta_k^{[1]}$  $_{k}^{[1]}$   $\|\theta_{k}^{[2]}$  $_{k}^{[2]} \|, \| \theta_{k-}^{[1]}$  $\|\theta^{[1]}_{k-\tau}\|, \|\theta^{[2]}_{k-\tau}\|$  $\binom{[2]}{k-\tau}$  | | ) + 1. When  $\theta_k^{[1]}$  $\ell_k^{[1]}$  and  $\theta_k^{[2]}$  $\kappa^{[2]}$  are not close to each order, i.e., when  $||w_k||$ is large,  $\spadesuit$  is not necessarily of higher order. Therefore, we consider a second type of Taylor expansion on  $g(\theta_k^{[1]}$  $g_k^{[1]}, x_k) - g(\theta_{k-1}^{[1]}$  ${}_{k-\tau}^{[1]}, x_k$ ) and  $g(\theta_k^{[2]})$  $g_k^{[2]}, x_k) - g(\theta_{k-}^{[2]})$  $(k_{k-\tau}^{\lfloor 2 \rfloor}, x_k)$ . The intuition for the second type of Taylor expansion is to analyze and bound  $\spadesuit$  by the small distance between  $\theta_k^{[j]}$  $\begin{array}{c} [j]_k \ [j] \end{array}$  and  $\theta_{k-\tau}^{[j]}$  for  $j \in \{1,2\}$ , even when  $||w_k||$  is large. This achieves  $\blacklozenge \lesssim ||w_k|| \alpha \tau T_1(||w_k|| + \alpha \tau T_1)$ . Simultaneously applying the two Taylor expansions will yield  $\clubsuit \lesssim \alpha \tau \|w_k\|^2 T_1$ . Finally, we overcome this challenge by carefully analyzing the boundedness of  $T_1$ ; see Theorem [4.1](#page-5-0) and its proof.

In parallel to the above coupling approach, we also explore an alternative approach by verifying the joint Markov chain  $(x_k, \theta_k)$  satisfies certain irreducibility and Lyapunov drift conditions, which in turn imply the chain is ergodic. To apply this approach, we exploit additional properties of the SA noise, namely minorization, which is satisfied in many applications where additional randomness is injected to the SA update. While the high level strategy of this approach is well developed [\[MT09](#page-12-7), [DMPS18](#page-10-7)], carrying out the analysis of each step is technically involved. In particular, we need to translate the minorization property of the noise to the irreducibility of the joint chain  $(x_k, \theta_k)$ , which is nontrivial in the presence of Markovian noise and nonlinearity.

### <span id="page-4-2"></span><span id="page-4-0"></span>4 Main Results

### 4.1 Weak Convergence of Projected SA

Our first main result proves the ergodicity of the joint process  $(x_k, \theta_k)_{k\geq 0}$  of the projected SA [\(2.1\)](#page-2-1).

<span id="page-5-3"></span><span id="page-5-0"></span>**Theorem 4.[1](#page-2-0)** (Ergodicity of Projected SA). Suppose that Assumption  $1-\lambda$  ( $p=1$ ) hold. The projected SA [\(2.1\)](#page-2-1) is applied with radius parameter  $2\|\theta^*\| \leq \beta < \infty$ . For stepsize  $\alpha > 0$  that satisfies the constraint  $\alpha \tau_{\alpha} \leq$  $\frac{\mu}{(940+96\beta)L^2}$ , the Markov chain  $(x_k, \theta_k)_{k\geq 0}$  converges to a unique stationary distribution  $\bar{\nu}_{\alpha} \in \mathcal{P}_2(\mathcal{X} \times \mathbb{R}^d)$ . Let  $\nu_{\alpha} := \mathcal{L}(\theta_{\infty})$  be the second marginal of  $\bar{\nu}_{\alpha}$ . For  $k \geq 2\tau_{\alpha}$ , it holds that

$$
W_2(\mathcal{L}(\theta_k), \nu_\alpha) \leq \bar{W}_2(\mathcal{L}(x_k, \theta_k), \bar{\nu}_\alpha) \leq (1 - \alpha \mu)^{k/2} \cdot s(\theta_0, \theta^*, \mu, L, R).
$$

Theorem [4.1](#page-5-0) generalizes prior weak convergence results for constant stepsize SA/SGD either under i.i.d. noise [\[DDB20](#page-10-3), [YBVE21\]](#page-13-1) or linear update [\[HCX23a,](#page-11-6) [LM23\]](#page-11-7). Our stepsize condition  $\alpha\tau_{\alpha} \lesssim \mu/L^2$  coincides with [\[SY19](#page-12-3), [HCX23a](#page-11-6)] on linear SA, a special case of our setting.

The proof of Theorem [4.1](#page-5-0) highlights the stabilizing effect of the projection operation in [\(2.1\)](#page-2-1). This effect, together with the smoothness of update function g, controls how the Markovian correlation propagates through the nonlinear update, allowing us to overcome the challenges discussed in Section [3.](#page-3-0) It is unclear whether our proof, which is based on Markov chain coupling, can be fully generalized to SA without projection. Nevertheless, we show that such a generalization is possible for a sub-family of nonlinear SA where g possesses the additional structure termed "asymptotic linearity", which is satisfied by, e.g., SGD applied to certain settings of logistic regression. For a formal statement of this result and proof, we refer the readers to Appendix [C.](#page-26-0)

As a by-product of our analysis, we establish the following non-asymptotic  $2p$ -th moment bound on the error  $\theta_k - \theta^*$ . Let  $\theta_{t+1/2} := \theta_t + \alpha(g(\theta_t, x_t) + \xi_{t+1}(\theta_t))$  denote the pre-projection iterate.

<span id="page-5-1"></span>**Proposition [4](#page-3-4).2.** Consider  $(\theta_k)_{k\geq 0}$  of iteration  $(2.1)$  with  $\beta \in [2||\theta^*||, \infty]$ . Let Assumption  $1-4(2p)$  hold. If stepsize  $\alpha$  satisfies  $\alpha \tau_{\alpha} L^2 \leq c_p \mu$ , with  $c_p \leq 1$ , the following holds for all  $k \geq \tau_{\alpha}$ ,

$$
\mathbb{E}[\|\theta_{k+1}-\theta^*\|^{2p}] \leq \mathbb{E}[\|\theta_{k+1/2}-\theta^*\|^{2p}] \leq c_{p,1}(1-\alpha\mu)^{k+1}\mathbb{E}[\|\theta_0-\theta^*\|^{2p}] + c_{p,2}(\alpha\tau_\alpha)^p \cdot s(\theta_0,\theta^*,L,\mu).
$$

Proposition [4.2](#page-5-1) implies that  $\mathbb{E}[\|\theta_k - \theta^*\|^{2p}] \leq (\alpha \tau)^p$  for sufficiently large k. This result generalizes those in  $[DDB20, CZD+22, SY19]$  to higher moments and the nonlinear Markovian setting. Note that Proposition [4.2](#page-5-1) is valid even without the projection operation in the SA update [\(2.1\)](#page-2-1), i.e.,  $\beta = \infty$ .

#### 4.2 Weak Convergence without Projection

Parallel to the coupling approach, we consider an alternative approach for establishing weak convergence via verifying irreducibility, positive Harris recurrence, and V -uniform ergodicity [\[MT09\]](#page-12-7) of the Markov chain  $(x_k, \theta_k)$ . This approach applies to nonlinear SA even without projection. To verify irreducibility, we exploit the following additional noise structure.

<span id="page-5-2"></span>**Assumption 5** (Noise Minorization). For each  $\theta \in \mathbb{R}^d$ , the distribution of the random variable  $\xi_1(\theta)$ , denoted by  $\zeta_{\theta}$ , can be decomposed as  $\zeta_{\theta} = \zeta_{1,\theta} + \zeta_{2,\theta}$ , where the measure  $\zeta_{1,\theta}$  has a density, denoted by  $p_{\theta}$ , which satisfies  $\inf_{\theta \in C} p_{\theta}(t) > 0$  for any bounded set C and any  $t \in \mathbb{R}^d$ .

A similar assumption is considered in [\[YBVE21,](#page-13-1) [BSS](#page-9-3)+23]. This assumption is mild and satisfied by any continuous random field supported on  $\mathbb{R}^d$ . Introducing such (small) continuous noise is often part of the algorithm design for inducing privacy [\[BW18](#page-9-6), [DRS22\]](#page-10-8) or exploration [\[PHD](#page-12-8)+18, [FAP](#page-10-9)+19]. Without Assumption [5,](#page-5-2) the chain may fail to be irreducible even when the other assumptions are satisfied; see [\[HCX23a\]](#page-11-6) for a counterexample.

Under Assumption [5,](#page-5-2) we obtain the following ergodicity result paralleling Theorem [4.1.](#page-5-0)

<span id="page-5-4"></span>**Theorem [4](#page-3-4).3** (Ergodicity of SA – Minorization). Suppose that Assumption [1](#page-2-0)[–3,](#page-2-3) Assumption  $\mathcal{A}(p = 1)$ , and Assumption [5](#page-5-2) hold. For stepsize  $\alpha > 0$  that satisfies the constraint  $\alpha \tau_{\alpha} L^2 < c_2 \mu$ , the Markov chain  $(x_k, \theta_k)_{k\geq 0}$  of [\(2.1\)](#page-2-1) with  $\beta = \infty$  is V-uniformly ergodic with Lyapunov function  $V(x, \theta) = ||\theta - \theta^*||^2 + 1$  and a unique stationary distribution  $\bar{\nu}_{\alpha} \in \mathcal{P}_2(\mathcal{X} \times \mathbb{R}^d)$ . Moreover, defining the V-norm  $\|\nu\|_V := \int |\nu(\mathrm{d}x)|V(x)$ , we have

<span id="page-5-5"></span>
$$
\left\| \mathcal{L}(x_k, \theta_k) - \bar{\nu}_{\alpha} \right\|_V \le \kappa \rho^k, \qquad \forall (x_0, \theta_0) \in \mathcal{X} \times \mathbb{R}^d, \forall k \ge 0,
$$
\n
$$
(4.1)
$$

where the constants  $\rho \in (0,1)$  and  $\kappa \in (0,\infty)$  may depend on  $\alpha$ .

### <span id="page-6-7"></span>4.3 Non-Asymptotic Convergence Rate and Central Limit Theorem

In the sequel, let  $(x_\infty, \theta_\infty^{(\alpha)})$  denote the random vector whose law is the stationary distribution  $\bar{\nu}_\alpha$  given in Theorem [4.1.](#page-5-0) As a corollary, we have geometric convergence for the first 2 moments of  $\theta_k$ .

<span id="page-6-8"></span>Corollary 4.4 (Non-Asymptotic Convergence Rate). Under the setting of Theorem [4.1,](#page-5-0) for any initialization of  $\theta_0 \in \mathbb{R}^d$ , we have

$$
\|\mathbb{E}[\theta_k] - \mathbb{E}[\theta_{\infty}^{(\alpha)}]\| \le (1 - \alpha \mu)^{k/2} \cdot s'(\theta_0, \theta^*, \mu, L, R), \text{ and}
$$
  

$$
\|\mathbb{E}[\theta_k \theta_k^{\top}] - \mathbb{E}[\theta_{\infty}^{(\alpha)}(\theta_{\infty}^{(\alpha)})^{\top}]\| \le (1 - \alpha \mu)^{k/2} \cdot s''(\theta_0, \theta^*, \mu, L, R).
$$

Moreover, the convergence rate established in Theorem [4.1](#page-5-0) is fast enough that we can use it to prove a Central Limit Theorem for the average iterates.

<span id="page-6-6"></span>**Corollary 4.5** (Central Limit Theorem). Under the setting of Theorem [4.1,](#page-5-0) as  $k \to \infty$  we have  $\frac{1}{\sqrt{k}}$  $\frac{1}{k} \sum_{t=0}^{k-1} (\theta_t \mathbb{E}[\theta_{\infty}]\big) \Rightarrow \mathcal{N}(0, \Sigma^{(a)})$ , where  $\Sigma^{(\alpha)} := \lim_{k \to \infty} \frac{1}{k} \mathbb{E}\big[\big(\sum_{t=0}^{k-1} (\theta_t - \mathbb{E}[\theta_{\infty}^{(\alpha)}])\big)^{\otimes 2}\big].$ 

Establishing the CLT sets the stage for using the SA iterates for statistical inference tasks such as confidence interval estimation. We discuss this in greater detail in Section [4.4](#page-6-1) below after characterizing the asymptotic bias, another important ingredient for using SA for inference.

#### <span id="page-6-1"></span>4.4 Bias Characterization

In this subsection, we characterize the asymptotic bias  $\mathbb{E}[\theta_{\infty}^{(\alpha)}] - \theta^*$ . Understanding the bias structure has important algorithmic implications for bias reduction, which we explore in Section [4.5,](#page-7-0) as well as for more efficient statistical inference and confidence interval estimation [\[HCX23b\]](#page-11-8).

<span id="page-6-0"></span>**Theorem 4.6** (Bias Characterization). Suppose Assumptions  $1-4(p = 3)$  $1-4(p = 3)$  hold. For each stepsize  $\alpha > 0$ satisfying  $\alpha \tau_{\alpha} L^2 < c_3 \mu$ , the following holds for some vector b independent of  $\alpha$ :

<span id="page-6-5"></span><span id="page-6-4"></span><span id="page-6-3"></span><span id="page-6-2"></span>
$$
\mathbb{E}[\theta_{\infty}^{(\alpha)}] - \theta^* = \alpha b + \mathcal{O}\big((\alpha \tau_{\alpha})^{3/2}\big). \tag{4.2}
$$

More specifically, the leading bias can be decomposed as  $b = b_m + b_n + b_c$ , where

$$
b_{\mathbf{m}} = -(\bar{g}'(\theta^*))^{-1} \mathbb{E}[g'(\theta^*, x_{\infty})h(\theta^*, x_{\infty})],
$$
\n(4.3)

$$
b_{n} = \frac{1}{2} (\bar{g}'(\theta^{*}))^{-1} \bar{g}''(\theta^{*}) A \Big( \mathbb{E}[g(\theta^{*}, x_{\infty})^{\otimes 2}] + \mathbb{E}[(\xi_{1}(\theta^{*}))^{\otimes 2}] \Big), \tag{4.4}
$$

$$
b_{\rm c} = \frac{1}{2} (\bar{g}'(\theta^*))^{-1} \bar{g}''(\theta^*) A \Big( \mathbb{E}[g(\theta^*, x_{\infty}) \otimes h(\theta^*, x_{\infty})] + \mathbb{E}[h(\theta^*, x_{\infty}) \otimes g(\theta^*, x_{\infty})] \Big), \tag{4.5}
$$

with  $A = (\bar{g}'(\theta^*)\otimes I + I\otimes \bar{g}'(\theta^*))^{-1}$  and  $h(\theta^*, x) = \int_{\mathcal{X}} (I - P^* + \Pi)^{-1} (P^* - \Pi)(x, dx')g(\theta^*, x')$ , with the kernel  $P^*$  being a regular conditional probability on X that satisfies  $\int_B \pi(\mathrm{d} x) P(x, C) = \int_C \pi(\mathrm{d} y) P^*(y, B)$ , for all  $B, C \in \mathcal{B}(\mathcal{X})$ .

We defer the detailed proof to Appendix [G.](#page-43-0) A few remarks are in order. First, we emphasize that  $(4.2)$  is essentially an equality, indicating a non-zero bias of order  $\alpha$  whenever  $b \neq 0$  (up to higher order terms). Notably, the Polyak-Ruppert averaging of the iterates cannot eliminate this bias. Note that the bias expansion in [\(4.2\)](#page-6-2) applies to both weakly converged projected and non-projected SA. Our analysis shows that compared with the non-projected SA, the projection operator induces an extra bias term of the order  $\mathcal{O}((\alpha^2 \tau_\alpha^3))$ , which is negligible relative to the main terms in in [\(4.2\)](#page-6-2).

More importantly, Theorem [4.6](#page-6-0) provides an explicit expression of the leading bias, which decomposes into three components: the Markovian part, the nonlinearity contribution, and a compound term, which is unique in nonlinear Markovian SA. Specifically,  $b<sub>m</sub>$  in  $(4.3)$  is associated with the Markovian multiplicative noise, where the matrix  $P^* - \Pi$  in the h function determines the mixing time of the data sequence  $(x_k)_{k \geq 1}$ . The term  $b_n$  in [\(4.4\)](#page-6-4) is linked to nonlinearity, as reflected by the Hessian term  $\bar{g}''(\theta^*) = \mathbb{E}[g''(\theta^*, x_{\infty})]$ , which quantifies the nonlinearity of g and is equal to zero in the case of a linear g. Lastly,  $b_c$  in [\(4.5\)](#page-6-5) is the compound term, due to its dependence on both the Markov noise (h function) and the nonlinearity measure  $\bar{g}^{\prime\prime}$ . In particular, we note the following two special cases:

- <span id="page-7-2"></span>• When g is a linear function,  $\bar{g}''(\theta^*) = 0$ . Hence,  $b_n = b_c = 0$ , and  $b_m$  recovers the result in [\[HCX23a\]](#page-11-6).
- When  $(x_k)_{k\geq 0}$  is i.i.d. sampled from the stationary distribution  $\pi$ , we have  $h(\theta^*, x) \equiv 0 \ \forall x \in \mathcal{X}$ , for  $P = P^* = \Pi$ . As such,  $b_m = b_c = 0$ , recovering the result in [\[DDB20\]](#page-10-3).

The presence of the compound term  $b_c$  suggests that as the SA structure becomes more nonlinear and the underlying Markov chain mixes more slowly, the impact on the bias is multiplicative rather than simply additive, a surprising phenomenon not unveiled in previous studies.

#### <span id="page-7-0"></span>4.5 Algorithmic Implications

We examine the practical implications of our weak convergence and bias characterization results, particularly for Polyak-Ruppert (PR) tail averaging and Richardson-Romberg (RR) extrapolation. In this subsection, we focus on the dependence on the stepsize  $\alpha$  and iteration index k, and make use of the big-O notation from Section [1.](#page-1-0) Recall that b is the bias vector defined in Theorem [4.6.](#page-6-0)

PR averaging [\[Rup88,](#page-12-4) [Pol90\]](#page-12-5) is a classical approach for reducing the variance and accelerating the convergence of SA. Here we consider the tail-averaging variant of PR averaging, defined as  $\bar{\theta}_{k_0,k} := \frac{1}{k-k_0} \sum_{t=k_0}^{k-1} \theta_t$ , for  $k \geq k_0$ , with a user-specified burn-in period  $k_0 \geq 0$  (a common choice is  $k_0 = k/2$ ). The following corollary, proved in Appendix [H,](#page-52-0) provides a non-asymptotic bound on the mean squared error (MSE) for the averaged iterates  $\theta_{k_0,k}$ .

<span id="page-7-1"></span>**Corollary 4.7** (Tail Averaging). Under the setting of Theorem  $4.6$ , the tail-averaged iterates satisfy the following bounds for all  $k > k_0 + 2\tau_\alpha$  and  $k_0 \ge \tau_\alpha + \frac{1}{\alpha \mu} \log \left( \frac{1}{\alpha \tau_\alpha} \right)$ ,

$$
\mathbb{E}\left[\|\bar{\theta}_{k_0,k}-\theta^*\|^2\right] = \underbrace{\alpha^2\|b\|^2 + \mathcal{O}\Big(\alpha\cdot(\alpha\tau)^{\frac{3}{2}}\Big)}_{T_1:\ asymptotic\ squared\ bias} + \underbrace{\mathcal{O}\Big(\frac{\tau_{\alpha}}{k-k_0}\Big)}_{T_2:\ variance} + \underbrace{\mathcal{O}\Big(\frac{(1-\alpha\mu)^{k_0/2}}{\alpha\left(k-k_0\right)^2}\Big)}_{T_3:\ optimization\ error}.
$$

Corollary [4.7](#page-7-1) shows that the MSE can be decomposed into three terms and elucidates how these terms depend on  $\alpha, k$ , and other problem parameters. In particular, the term  $T_1$  corresponds to the asymptotic squared bias  $\|\mathbb{E}[\theta_{\infty}^{(\alpha)} - \theta^*\] \|^{2}$ , which is not affected by averaging. The term  $T_2$  is associated with the variance  $Var(\bar{\theta}_{k_0,k})$ , which decays at rate  $1/k$  due to averaging. Lastly, the term  $T_3$  represents the optimization error  $\|\mathbb{E} \hat{\theta}_{k_0,k} - \theta_{\infty}\|^2$ , which decays geometrically in  $k_0$  thanks to the use of a constant stepsize  $\alpha$  and the tail-averaging procedure.

Note that averaging does not affect the bias of order  $\alpha$ . With the precise bias characterization in Theo-rem [4.6,](#page-6-0) we can order-wise reduce the bias to  $\mathcal{O}(\alpha^{1.5})$  by employing the RR extrapolation technique [\[SB02\]](#page-12-9). Let  $\bar{\theta}_{k_0,k}^{(\alpha)}$  and  $\bar{\theta}_{k_0,k}^{(2\alpha)}$  denote the tail-averaged iterates using two stepsizes  $\alpha$  and  $2\alpha$  with the same data  $(x_k)_{k\geq 0}$ . The RR extrapolated iterates are defined as  $\tilde{\theta}_{k_0,k}^{(\alpha)} = 2\bar{\theta}_{k_0,k}^{(\alpha)} - \bar{\theta}_{k_0,k}^{(2\alpha)}$ .

Corollary 4.8 (RR-Extrapolation). Under the setting of Theorem  $4.6$ , the RR-extrapolated iterates satisfy the following bounds for all  $k > k_0 + 2\tau_\alpha$  and  $k_0 \ge \tau_\alpha + \frac{1}{\alpha \mu} \log \left( \frac{1}{\alpha \tau_\alpha} \right)$ ,

$$
\mathbb{E}\Big[\|\widetilde{\theta}_{k_0,k}-\theta^*\|^2\Big] = \mathcal{O}\Big((\alpha\tau_{\alpha})^3\Big) + \mathcal{O}\Big(\frac{\tau_{\alpha}}{k-k_0}\Big) + \mathcal{O}\Big(\frac{(1-\alpha\mu)^{k_0/2}}{\alpha\left(k-k_0\right)^2}\Big).
$$

Backed by the CLT in Corollary [4.5,](#page-6-6) the iterates of constant-stepsize SA can be used to construct confidence intervals of  $\theta^*$ . For i.i.d. data or linear SA, this approach has been explored in [\[LLKC18,](#page-11-14) [YBVE21,](#page-13-1) [XZ22](#page-13-3), [HCX23b\]](#page-11-8) along with an appropriate variance estimator [\[FJ10](#page-10-10), [XZ22\]](#page-13-3). In our Markovian nonlinear setting, where the iterates are biased, it is crucial to use RR extrapolation for bias reduction. Once the bias is accounted for, the power of using constant stepsizes reveals itself as it leads to rapid mixing and low correlation of the iterates. Together, they lead to efficient confidence interval estimation schemes using nonlinear Markovian SA; see the empirical results in [\[HCX23b\]](#page-11-8) showing its efficacy. In contrast, the classical diminishing stepsize paradigm often suffers from high correlation [\[CLTZ20\]](#page-10-11) and in turn inaccurate variance estimation, resulting in unsatisfactory coverage probability with finite data [\[HCX23b\]](#page-11-8).

#### <span id="page-8-1"></span>4.6 Implications for Learning GLM

Generalized linear models (GLM) extend linear regression to the model  $\mathbb{E}[Y|W] = \sigma(W^{\top}\theta^*)$ , where W is the covariate, Y the response variable, and  $\sigma$  is called the *mean function*. For any monotone (and potentially nonlinear)  $\sigma$ , the powerful framework developed in [\[WZDD23,](#page-13-0) [KKSK11,](#page-11-2) [KS09](#page-11-3), [DGK](#page-10-0)+20] allows one to formulate the estimation of  $\theta^*$  as minimizing an appropriate *convex* (surrogate) loss function. Applying SGD to this loss leads to a nonlinear SA update, to which our results are applicable. Below we discuss their applications in two concrete examples of GLMs.

**Logistic Regression** Logistic regression uses a sigmoid mean function  $\sigma(x) = \frac{1}{1+\exp(-x)}$ . Suppose the covariate  $w_k$  is sequentially sampled from a uniformly ergodic Markov chain with a bounded state space  $W \subset \mathbb{R}^d$ , and conditioned on  $w_k$  the response  $y_k$  is Bernoulli distributed with parameter  $(1 + \exp(-w_k^{\top} \theta^*))^{-1}$ . SGD applied to the L<sub>2</sub>-regularized negative log-likelihood function takes the form of the SA update  $\theta_{k+1}$  $\theta_k + \alpha g(\theta_k, x_k)$ , where  $x_k = (w_k, y_k) \in W \times \{0, 1\}$  and  $g(\theta_k, x_k) = -w_k(\sigma(-w_k^{\top}\theta_k) - y_k) - \lambda \theta_k$ . For simplicity, we do not consider  $\xi$ -perturbation, i.e.,  $\xi_{k+1}(\theta_k) \equiv 0$ . It is easy to verify that this g is strongly monotone and sufficiently smooth with at most linear growth in  $|\theta|$ , hence satisfying Assumption [1–](#page-2-0)[3.](#page-2-3) Therefore, all the results in Sections [4.1](#page-4-0)[–4.5](#page-7-0) apply to logistic regression with constant stepsizes and Markovian data.

**Smooth ReLU Regression** The mean function  $\sigma$  can be interpreted as playing a similar role as the activation function in neural networks. Widely adopted is ReLU activation  $\sigma(x) = \max(0, x)$  as well as its various smooth approximations [\[BKBP22,](#page-9-2) [GBB11](#page-11-4)]. The problem of learning  $\theta^*$  in this setting, sometimes called ReLU Regression, has been studied in the last decade and recently regained attention [\[WZDD23](#page-13-0), [KKSK11,](#page-11-2) [KS09,](#page-11-3) [DGK](#page-10-0)<sup>+</sup>20]. Unlike linear or logistic regression, the least squares and maximum likelihood formulation associated with such nonlinear mean functions  $\sigma$  is non-convex. Nevertheless, the convex surrogate loss framework in  $[WZDD23, DGK^+20]$  $[WZDD23, DGK^+20]$  $[WZDD23, DGK^+20]$  still applies. As an example, we focus on the SoftPlus activation  $\sigma(x) = \log(1 + \exp(x))/\iota$  with a temperature parameter  $\iota > 0$  [\[GBB11\]](#page-11-4). With  $L_2$ -regularization the resulting SGD iteration is  $\theta_{k+1} = \theta_k - \alpha \big( w_k \big( \frac{1}{\iota} \log(1 + \exp(\iota w_k^{\top} \theta_k)) - y_k \big) + \lambda \theta_k \big)$ , where the covariateresponse pair  $(\theta_k, x_k)$  is as before. This problem can again be cast as nonlinear SA with a strongly monotone and smooth g, satisfying Assumptions  $1-3$  $1-3$ . All results in Sections  $4.1-4.5$  $4.1-4.5$  apply.

### <span id="page-8-0"></span>5 Related Work

General SA and SGD. SA and SGD can be traced back to the seminal work of [\[RM51](#page-12-2)]. Classical work assumes a diminishing stepsize sequence, and has shown almost sure asymptotic convergence to  $\theta^*$  [\[RM51,](#page-12-2) [Blu54](#page-9-7)]. Subsequent works propose the iterate averaging technique, now known as Polyak-Ruppert (PR) averaging, to reduce variance and accelerate convergence [\[Rup88](#page-12-4), [Pol90\]](#page-12-5), and also establish a Central Limit Theorem for the asymptotic normality of the averaged iterates [\[PJ92\]](#page-12-10). The asymptotic convergence theory of SA and SGD is well developed and extensively addressed in many exemplary textbooks, see [\[KY03,](#page-11-0) [BMP90,](#page-9-8) [WR22](#page-12-11)]. There are also recent works studying the non-asymptotic convergence with diminishing stepsizes [\[CMSS23](#page-10-12), [CBD22\]](#page-9-9). The recent work [\[CMZ23\]](#page-10-13) establishes the high probability bound on the estimation error of contractive SA with diminishing stepsize.

SA and SGD with Constant Stepsizes. There has been an increasing interest in studying SA with constant stepsize. Many works in this line provide non-asymptotic upper bounds on mean squared error (MSE)  $\mathbb{E}[\|\theta_t - \theta^*\|^2]$ . Works in [\[LS18](#page-12-12), [MLW](#page-12-13)<sup>+</sup>20, [DMN](#page-10-6)<sup>+</sup>21] study linear SA (LSA) under i.i.d. data. Recent works extend the analysis of the MSE to LSA with Markovian data, such as [\[SY19,](#page-12-3) [MPWB21,](#page-12-6) [DMNS22\]](#page-10-5). There are also works providing upper bounds of MSE for general contractive SA with Markovian noise [\[CZD](#page-10-2)<sup>+</sup>22, [CMSS23\]](#page-10-12).

In addition to obtaining non-asymptotic guarantees, there are also works focusing on understanding the asymptotic behavior of SA iterates. Recent works have shown that when using constant stepsize, one loses the almost sure convergence guarantee in the diminishing stepsize sequence regime, and at best can achieve distributional convergence, as demonstrated in [\[DDB20,](#page-10-3) [DMN](#page-10-6)<sup>+</sup>21, [YBVE21,](#page-13-1) [CMM22](#page-10-14), [XZ22,](#page-13-3) [HCX23a,](#page-11-6) [ZX24\]](#page-13-2). The presence of asymptotic bias is also a recurring theme in recent literature, with precise characterization given in [\[DDB20](#page-10-3)] for strongly-convex SGD with i.i.d. data and in [\[HCX23a\]](#page-11-6) for LSA with Markovian data. <span id="page-9-10"></span>Works in [\[MLW](#page-12-13)+20, [YBVE21,](#page-13-1) [XZ22,](#page-13-3) [HCX23b,](#page-11-8) [ZX24\]](#page-13-2) also establish Central Limit Theorems for averaged SA iterates with constant stepsizes.

# 6 Conclusion

We provide the first weak convergence and steady-state analysis for constant-stepsize SA with both nonlinear update and Markovian data. Our analysis elucidates the compound effect of nonlinearity and memory, which leads to new analytical challenges and behaviors. A limitation of our results is the use of a projection step or the noise minorization assumption. Whether they can be removed is worth investigating. Other future directions include refining the dimension dependence in our results, as well as a theoretical investigation of statistical inference.

### Acknowledgments

Y. Chen is partially supported in part by NSF grant CCF-2233152. Y. Zhang and Q. Xie are supported in part by NSF grants CNS-1955997 and ECCS-2339794.

### References

- <span id="page-9-1"></span>[Ber19] Dimitri P. Bertsekas. Reinforcement learning and Optimal Control. Athena Scientific, Belmont, Massachusetts, USA, 2019. [1](#page-0-0)
- <span id="page-9-7"></span><span id="page-9-2"></span>[BKBP22] Koushik Biswas, Sandeep Kumar, Shilpak Banerjee, and Ashish Kumar Pandey. Smooth maximum unit: Smooth activation function for deep networks using smoothing maximum technique. In 2022 IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR), pages 784–793, 2022. [1,](#page-0-0) [9](#page-8-1)
	- [Blu54] Julius R. Blum. Approximation methods which converge with probability one. The Annals of Mathematical Statistics, 25(2):382 – 386, 1954. [9](#page-8-1)
	- [BM00] Vivek S. Borkar and Sean P. Meyn. The O.D.E. method for convergence of stochastic approximation and reinforcement learning. SIAM Journal on Control and Optimization, 38(2):447–469, Jan 2000. [2](#page-1-1)
- <span id="page-9-8"></span><span id="page-9-4"></span>[BMP90] Albert Benveniste, Michel Metivier, and Pierre Priouret. Adaptive Algorithms and Stochastic Approximations. Springer Berlin Heidelberg, 1st edition, 1990. [9](#page-8-1)
- <span id="page-9-0"></span>[Bor08] Vivek S. Borkar. Stochastic Approximation: A Dynamical Systems Viewpoint. Hindustan Book Agency Gurgaon, 2008. [1](#page-0-0)
- <span id="page-9-5"></span>[BRS21] Jalaj Bhandari, Daniel Russo, and Raghav Singal. A finite time analysis of temporal difference learning with linear function approximation. Operations Research, 69(3):950–973, May 2021. [3](#page-2-4)
- <span id="page-9-3"></span>[BSS+23] Aleksandr Beznosikov, Sergey Samsonov, Marina Sheshukova, Alexander Gasnikov, Alexey Naumov, and Eric Moulines. First order methods with Markovian noise: from acceleration to variational inequalities, 2023. [2,](#page-1-1) [6](#page-5-3)
- <span id="page-9-6"></span>[BW18] Borja Balle and Yu-Xiang Wang. Improving the Gaussian mechanism for differential privacy: Analytical calibration and optimal denoising. In Jennifer Dy and Andreas Krause, editors, Proceedings of the 35th International Conference on Machine Learning, volume 80 of Proceedings of Machine Learning Research, pages 394–403. PMLR, 10–15 Jul 2018. [6](#page-5-3)
- <span id="page-9-9"></span>[CBD22] Siddharth Chandak, Vivek S. Borkar, and Parth Dodhia. Concentration of contractive stochastic approximation and reinforcement learning. Stochastic Systems, Jul 2022. [9](#page-8-1)
- <span id="page-10-11"></span>[CLTZ20] Xi Chen, Jason D Lee, Xin T Tong, and Yichen Zhang. Statistical inference for model parameters in stochastic gradient descent. Annals of Statistics, 48(1):251–273, 2020. [8](#page-7-2)
- <span id="page-10-14"></span>[CMM22] Zaiwei Chen, Shancong Mou, and Siva Theja Maguluri. Stationary behavior of constant stepsize sgd type algorithms: An asymptotic characterization. Proceedings of the ACM on Measurement and Analysis of Computing Systems, 6(1), 02 2022. [9](#page-8-1)
- <span id="page-10-12"></span>[CMSS23] Zaiwei Chen, Siva Theja Maguluri, Sanjay Shakkottai, and Karthikeyan Shanmugam. A Lyapunov theory for finite-sample guarantees of Markovian stochastic approximation. Operations Research, 2023. [9](#page-8-1)
- <span id="page-10-13"></span>[CMZ23] Zaiwei Chen, Siva Theja Maguluri, and Martin Zubeldia. Concentration of contractive stochastic approximation: Additive and multiplicative noise.  $arXiv$  preprint  $arXiv:2303.15740$ , 2023. [9](#page-8-1)
- <span id="page-10-1"></span>[CUH16] Djork-Arn´e Clevert, Thomas Unterthiner, and Sepp Hochreiter. Fast and accurate deep network learning by exponential linear units (elus), 2016. [1](#page-0-0)
- <span id="page-10-2"></span>[CZD+22] Zaiwei Chen, Sheng Zhang, Thinh T. Doan, John-Paul Clarke, and Siva Theja Maguluri. Finitesample analysis of nonlinear stochastic approximation with applications in reinforcement learning. Automatica, 146:110623, 2022. [2,](#page-1-1) [3,](#page-2-4) [4,](#page-3-5) [6,](#page-5-3) [9,](#page-8-1) [16,](#page-15-0) [32,](#page-31-0) [33](#page-32-0)
- <span id="page-10-3"></span>[DDB20] Aymeric Dieuleveut, Alain Durmus, and Francis Bach. Bridging the gap between constant step size stochastic gradient descent and Markov chains. The Annals of Statistics, 48(3):1348 – 1382, 2020. [2,](#page-1-1) [3,](#page-2-4) [4,](#page-3-5) [6,](#page-5-3) [8,](#page-7-2) [9](#page-8-1)
- <span id="page-10-0"></span>[DGK<sup>+</sup>20] Ilias Diakonikolas, Surbhi Goel, Sushrut Karmalkar, Adam R. Klivans, and Mahdi Soltanolkotabi. Approximation schemes for ReLU regression. In Jacob Abernethy and Shivani Agarwal, editors, Proceedings of Thirty Third Conference on Learning Theory, volume 125 of Proceedings of Machine Learning Research, pages 1452–1485. PMLR, 09–12 Jul 2020. [1,](#page-0-0) [9](#page-8-1)
- <span id="page-10-6"></span>[DMN<sup>+</sup>21] Alain Durmus, Eric Moulines, Alexey Naumov, Sergey Samsonov, Kevin Scaman, and Hoi-To Wai. Tight high probability bounds for linear stochastic approximation with fixed stepsize. In M. Ranzato, A. Beygelzimer, Y. Dauphin, P.S. Liang, and J. Wortman Vaughan, editors, Advances in Neural Information Processing Systems, volume 34, pages 30063–30074. Curran Associates, Inc., 2021. [4,](#page-3-5) [9](#page-8-1)
- <span id="page-10-5"></span>[DMNS22] Alain Durmus, Eric Moulines, Alexey Naumov, and Sergey Samsonov. Finite-time highprobability bounds for Polyak-Ruppert averaged iterates of linear stochastic approximation, 2022. [3,](#page-2-4) [9](#page-8-1)
- <span id="page-10-8"></span><span id="page-10-7"></span>[DMPS18] Randal Douc, Eric Moulines, Pierre Priouret, and Philippe Soulier. Markov Chains. Springer Cham, 1st edition, 2018. [5](#page-4-1)
	- [DRS22] Jinshuo Dong, Aaron Roth, and Weijie J. Su. Gaussian Differential Privacy. Journal of the Royal Statistical Society Series B: Statistical Methodology, 84(1):3–37, 02 2022. [6](#page-5-3)
	- [DT22] Jing Dong and Xin T. Tong. Stochastic gradient descent with dependent data for offline reinforcement learning, 2022. [3](#page-2-4)
- <span id="page-10-15"></span><span id="page-10-10"></span><span id="page-10-9"></span><span id="page-10-4"></span>[FAP+19] Meire Fortunato, Mohammad Gheshlaghi Azar, Bilal Piot, Jacob Menick, Ian Osband, Alex Graves, Vlad Mnih, Remi Munos, Demis Hassabis, Olivier Pietquin, Charles Blundell, and Shane Legg. Noisy networks for exploration, 2019. [6](#page-5-3)
	- [FJ10] James M. Flegal and Galin L. Jones. Batch means and spectral variance estimators in Markov chain Monte Carlo. The Annals of Statistics,  $38(2):1034 - 1070$  $38(2):1034 - 1070$  $38(2):1034 - 1070$ ,  $2010.8$
	- [Fol99] Gerald B. Folland. Real analysis: modern techniques and their applications. Wiley, New York, 2nd ed. edition, 1999. [15](#page-14-0)
- <span id="page-11-13"></span>[FT98] Sergey G. Foss and Richard L. Tweedie. Perfect simulation and backward coupling. Communications in Statistics. Stochastic Models, 14(1-2):187–203, 1998. [4](#page-3-5)
- <span id="page-11-4"></span>[GBB11] Xavier Glorot, Antoine Bordes, and Yoshua Bengio. Deep sparse rectifier neural networks. In Geoffrey Gordon, David Dunson, and Miroslav Dudík, editors, *Proceedings of the Fourteenth* International Conference on Artificial Intelligence and Statistics, volume 15 of Proceedings of Machine Learning Research, pages 315–323, Fort Lauderdale, FL, USA, 11–13 Apr 2011. PMLR. [1,](#page-0-0) [9](#page-8-1)
- <span id="page-11-6"></span>[HCX23a] Dongyan (Lucy) Huo, Yudong Chen, and Qiaomin Xie. Bias and extrapolation in Markovian linear stochastic approximation with constant stepsizes. In Abstract Proceedings of the 2023 ACM SIGMETRICS International Conference on Measurement and Modeling of Computer Systems, pages 81–82, 2023. [2,](#page-1-1) [3,](#page-2-4) [4,](#page-3-5) [5,](#page-4-1) [6,](#page-5-3) [8,](#page-7-2) [9,](#page-8-1) [44,](#page-43-1) [54,](#page-53-0) [55](#page-54-0)
- <span id="page-11-10"></span><span id="page-11-8"></span>[HCX23b] Dongyan (Lucy) Huo, Yudong Chen, and Qiaomin Xie. Effectiveness of constant stepsize in Markovian LSA and statistical inference, 2023. [2,](#page-1-1) [7,](#page-6-7) [8,](#page-7-2) [10](#page-9-10)
	- [Hil87] F B Hildebrand. Introduction to numerical analysis. Dover Books on Mathematics. Dover Publications, Mineola, NY, 2 edition, June 1987. [2](#page-1-1)
- <span id="page-11-12"></span>[HIMM19] Yu-Guan Hsieh, Franck Iutzeler, Jérôme Malick, and Panayotis Mertikopoulos. On the convergence of single-call stochastic extra-gradient methods. In H. Wallach, H. Larochelle, A. Beygelzimer, F. d'Alché-Buc, E. Fox, and R. Garnett, editors, Advances in Neural Information Processing Systems, volume 32. Curran Associates, Inc., 2019. [3](#page-2-4)
- <span id="page-11-9"></span>[JKK<sup>+</sup>18] Prateek Jain, Sham M. Kakade, Rahul Kidambi, Praneeth Netrapalli, and Aaron Sidford. Parallelizing stochastic gradient descent for least squares regression: Mini-batching, averaging, and model misspecification. Journal of Machine Learning Research, 18(223):1–42, 2018. [2](#page-1-1)
- <span id="page-11-2"></span>[KKSK11] Sham M Kakade, Varun Kanade, Ohad Shamir, and Adam Kalai. Efficient learning of generalized linear and single index models with isotonic regression. In J. Shawe-Taylor, R. Zemel, P. Bartlett, F. Pereira, and K.Q. Weinberger, editors, Advances in Neural Information Processing Systems, volume 24. Curran Associates, Inc., 2011. [1,](#page-0-0) [9](#page-8-1)
- <span id="page-11-5"></span><span id="page-11-3"></span>[KOG<sup>+</sup>22] Kyurae Kim, Jisu Oh, Jacob Gardner, Adji Bousso Dieng, and Hongseok Kim. Markov chain score ascent: A unifying framework of variational inference with markovian gradients. In S. Koyejo, S. Mohamed, A. Agarwal, D. Belgrave, K. Cho, and A. Oh, editors, Advances in Neural Information Processing Systems, volume 35, pages 34802–34816. Curran Associates, Inc., 2022. [2](#page-1-1)
	- [KS09] Adam Tauman Kalai and Ravi Sastry. The isotron algorithm: High-dimensional isotonic regression. In COLT, 2009. [1,](#page-0-0) [9](#page-8-1)
	- [KY03] Harold J. Kushner and G. George Yin. Stochastic Approximation and Recursive Algorithms and Applications. Stochastic Modelling and Applied Probability. Springer, New York, NY, USA, 2nd edition, 2003. [1,](#page-0-0) [9](#page-8-1)
	- [Lan20] Guanghui Lan. First-order and stochastic optimization methods for machine learning, volume 1. Springer, 2020. [1](#page-0-0)
- <span id="page-11-14"></span><span id="page-11-11"></span><span id="page-11-7"></span><span id="page-11-1"></span><span id="page-11-0"></span>[LLKC18] Tianyang Li, Liu Liu, Anastasios Kyrillidis, and Constantine Caramanis. Statistical inference using SGD. Proceedings of the AAAI Conference on Artificial Intelligence, 32(1), Apr. 201[8](#page-7-2). 8
	- [LLZ23] Xiang Li, Jiadong Liang, and Zhihua Zhang. Online statistical inference for nonlinear stochastic approximation with Markovian data, 2023. [3](#page-2-4)
	- [LM23] Caio Kalil Lauand and Sean Meyn. The curse of memory in stochastic approximation. In 2023 62nd IEEE Conference on Decision and Control (CDC), pages 7803–7809, 2023. [2,](#page-1-1) [3,](#page-2-4) [6](#page-5-3)
- <span id="page-12-12"></span>[LS18] Chandrashekar Lakshminarayanan and Csaba Szepesv´ari. Linear stochastic approximation: How far does constant step-size and iterate averaging go? In Amos Storkey and Fernando Perez-Cruz, editors, Proceedings of the Twenty-First International Conference on Artificial Intelligence and Statistics, volume 84 of Proceedings of Machine Learning Research, pages 1347–1355. PMLR, 09–11 Apr 2018. [9](#page-8-1)
- <span id="page-12-13"></span>[MLW+20] Wenlong Mou, Chris Junchi Li, Martin J. Wainwright, Peter L. Bartlett, and Michael I. Jordan. On linear stochastic approximation: Fine-grained Polyak-Ruppert and non-asymptotic concentration. In Jacob Abernethy and Shivani Agarwal, editors, Proceedings of Thirty Third Conference on Learning Theory, volume 125 of Proceedings of Machine Learning Research, pages 2947–2997. PMLR, 09–12 Jul 2020. [9,](#page-8-1) [10](#page-9-10)
- <span id="page-12-7"></span><span id="page-12-6"></span>[MPWB21] Wenlong Mou, Ashwin Pananjady, Martin J. Wainwright, and Peter L. Bartlett. Optimal and instance-dependent guarantees for markovian linear stochastic approximation, 2021. [3,](#page-2-4) [9](#page-8-1)
	- [MT09] Sean P. Meyn and Richard L. Tweedie. Markov Chains and Stochastic Stability. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 2nd edition, 2009. [5,](#page-4-1) [6,](#page-5-3) [39,](#page-38-0) [43,](#page-42-0) [44](#page-43-1)
	- [MW00] Michael Maxwell and Michael Woodroofe. Central limit theorems for additive functionals of Markov chains. The Annals of Probability,  $28(2)$ :713 – 724, 2000. [38](#page-37-0)
- <span id="page-12-15"></span><span id="page-12-1"></span>[NWB<sup>+</sup>20] Dheeraj Nagaraj, Xian Wu, Guy Bresler, Prateek Jain, and Praneeth Netrapalli. Least squares regression with Markovian data: Fundamental limits and algorithms. In H. Larochelle, M. Ranzato, R. Hadsell, M.F. Balcan, and H. Lin, editors, Advances in Neural Information Processing Systems, volume 33, pages 16666–16676. Curran Associates, Inc., 2020. [2](#page-1-1)
- <span id="page-12-14"></span><span id="page-12-11"></span><span id="page-12-10"></span><span id="page-12-9"></span><span id="page-12-8"></span><span id="page-12-5"></span><span id="page-12-4"></span><span id="page-12-3"></span><span id="page-12-2"></span><span id="page-12-0"></span>[PHD<sup>+</sup>18] Matthias Plappert, Rein Houthooft, Prafulla Dhariwal, Szymon Sidor, Richard Y. Chen, Xi Chen, Tamim Asfour, Pieter Abbeel, and Marcin Andrychowicz. Parameter space noise for exploration. In International Conference on Learning Representations, 2018. [6](#page-5-3)
	- [PJ92] Boris T. Polyak and Anatoli B. Juditsky. Acceleration of stochastic approximation by averaging. SIAM Journal on Control and Optimization, 30(4):838–855, Jul 1992. [9](#page-8-1)
	- [Pol90] Boris T. Polyak. New stochastic approximation type procedures. Automation and Remote Control, 51(7):98–107, Jul 1990. [2,](#page-1-1) [8,](#page-7-2) [9](#page-8-1)
	- [RM51] Herbert Robbins and Sutton Monro. A stochastic approximation method. The Annals of Mathematical Statistics, 22(3):400 – 407, 1951. [2,](#page-1-1) [9](#page-8-1)
	- [Rup88] David Ruppert. Efficient estimations from a slowly convergent Robbins-Monro process. Technical report, Cornell University, February 1988. [2,](#page-1-1) [8,](#page-7-2) [9](#page-8-1)
	- [SB02] Josef Stoer and Roland Bulirsch. Introduction to Numerical Analysis. Springer, New York, NY, USA, 3rd edition, 2002. [8](#page-7-2)
	- [SB18] Richard S. Sutton and Andrew G. Barto. Reinforcement Learning: An Introduction. A Bradford Book, Cambridge, MA, USA, 2018. [1](#page-0-0)
	- [SY19] Rayadurgam Srikant and Lei Ying. Finite-time error bounds for linear stochastic approximation and TD learning. In Proceedings of the Thirty-Second Conference on Learning Theory, volume 99 of Proceedings of Machine Learning Research, pages 2803–2830. PMLR, 25–28 Jun 2019. [2,](#page-1-1) [3,](#page-2-4) [5,](#page-4-1) [6,](#page-5-3) [9,](#page-8-1) [16](#page-15-0)
	- [Vil09] Cédric Villani. *Optimal Transport: Old and New.* Grundlehren der mathematischen Wissenschaften. Springer Berlin Heidelberg, 2009. [28,](#page-27-0) [29,](#page-28-0) [37,](#page-36-0) [39](#page-38-0)
	- [WR22] Stephen J. Wright and Benjamin Recht. Optimization for Data Analysis. Cambridge University Press, 2022. [9](#page-8-1)
- <span id="page-13-3"></span><span id="page-13-0"></span>[WZDD23] Puqian Wang, Nikos Zarifis, Ilias Diakonikolas, and Jelena Diakonikolas. Robustly learning a single neuron via sharpness. arXiv preprint arXiv:2306.07892, 2023. [1,](#page-0-0) [9](#page-8-1)
	- [XZ22] Chuhan Xie and Zhihua Zhang. A statistical online inference approach in averaged stochastic approximation. In S. Koyejo, S. Mohamed, A. Agarwal, D. Belgrave, K. Cho, and A. Oh, editors, Advances in Neural Information Processing Systems, volume 35, pages 8998–9009. Curran Associates, Inc., 2022. [8,](#page-7-2) [9,](#page-8-1) [10](#page-9-10)
- <span id="page-13-2"></span><span id="page-13-1"></span>[YBVE21] Lu Yu, Krishnakumar Balasubramanian, Stanislav Volgushev, and Murat A. Erdogdu. An analysis of constant step size SGD in the non-convex regime: Asymptotic normality and bias. In Advances in Neural Information Processing Systems, volume 34, pages 4234–4248. Curran Associates, Inc., 2021. [2,](#page-1-1) [6,](#page-5-3) [8,](#page-7-2) [9,](#page-8-1) [10](#page-9-10)
	- [ZX24] Yixuan Zhang and Qiaomin Xie. Constant stepsize Q-learning: Distributional convergence, bias and extrapolation, 2024. [2,](#page-1-1) [9,](#page-8-1) [10](#page-9-10)

### <span id="page-14-0"></span>A Additional Notations

**General Probability** We write  $z_1 \perp z_2 \mid z_3$  if random variables  $z_1$  and  $z_2$  are conditionally independent given z<sub>3</sub>. Recall that we define the metric  $\overline{d}((x,\theta),(x',\theta')):=\sqrt{\mathbb{1}\{x\neq x'\}+\|\theta-\theta'\|^2}$  for the space  $\mathcal{X}\times\mathbb{R}^d$ . Thus, for  $\bar{\mu}$  and  $\bar{\nu}$  in  $\mathcal{P}_2(\mathcal{X} \times \mathbb{R}^d)$ , the Wasserstein-2 distance w.r.t.  $\bar{d}$  is computed as

$$
\overline{W}_2(\overline{\mu}, \overline{\nu}) = \inf \left\{ \left( \mathbb{E}[\mathbb{1}\{x \neq x'\} + ||\theta - \theta'||^2] \right)^{\frac{1}{2}} : \mathcal{L}((x, \theta)) = \overline{\mu}, \mathcal{L}((x', \theta')) = \overline{\nu} \right\}.
$$

**General State Space Markov Chains** Throughout the paper, we assume that  $\mathcal{X}$  is a Borel space. Let P denote the transition kernel. We call  $\pi$  the stationary distribution of P if it satisfies  $\int_{\mathcal{X}} \pi(dx)P(x, B) = \pi(B)$ , for  $B \in \mathcal{B}(\mathcal{X})$ . Define the  $\pi$ -weighted inner product  $\langle f, g \rangle_{L^2(\pi)} = \int_{\mathcal{X}} \pi(dx) f^\top(x) g(x)$  and the induced norm  $||f||_{L^2(\pi)} = (\langle f, f \rangle_{L^2(\pi)})^{1/2}$ . Let  $L^2(\pi) = \{f : ||f||_{L^2(\pi)} < \infty\}$  denote the corresponding Hilbert space of  $\mathbb{R}^d$ . valued, square-integrable and measurable functions on  $\mathcal{X}$ . For an operator  $T: L^2(\pi) \to L^2(\pi)$ , its operator norm is defined as  $||T||_{L^2(\pi)} = \sup_{||f||_{L^2(\pi)}=1} ||Tf||_{L^2(\pi)}$ . The transition kernel is a bounded linear operator on  $L^2(\pi)$ , in particular with norm  $||P||_{L^2(\pi)=1}$ . Also, we define the kernel/operator  $\Pi = 1 \otimes \pi$  by  $\Pi(x, \cdot) = \pi$ .

Throughout the paper, we assume that X is a Borel space. Let P denote the transition kernel. We call  $\pi$ the stationary distribution of P if it satisfies  $\int_{\mathcal{X}} \pi(dx)P(x, B) = \pi(B)$ , for  $B \in \mathcal{B}(\mathcal{X})$ . There exists a kernel  $P^*$  as a regular conditional probability that satisfies  $\int_A \pi(\mathrm{d}s)P(x, B) = \int_B \pi(\mathrm{d}y)P^*(y, A)$ , for  $A, B \in \mathcal{B}(\mathcal{X})$ [\[Fol99](#page-10-15), Chapter 21.4, Theorem 19], and  $P^*$  defines the probability law for the time-reversed chain of  $(x_k)_{k\geq 0}$ .

### B Proof of Pilot Results (Proposition [4.2\)](#page-5-1)

In this section, we prove the pilot result, namely Proposition [4.2.](#page-5-1) We prove the desired moments for  $\beta = \infty$ , i.e., without any projection. It is easy to see that when the projection radius  $\beta \in [2||\theta^*||, \infty]$ ,

$$
\mathbb{E}[\|\theta_{t+1}-\theta^*\|^{2p}]\leq \mathbb{E}[\|\theta_{t+1/2}-\theta^*\|^{2p}],
$$

where  $\theta_{t+1/2}$  denotes the iterate before projection. The term on the right hand side can be further bounded by the moment bounds for iteration without projection. Therefore, it suffices for us to prove the respective moment bounds without any projection.

Given Assumption [4](#page-3-4) hold for 2p-th moment, with  $p \ge 1$ , we prove the moment bound in Proposition [4.2](#page-5-1) for *n* with  $1 \leq n \leq p$  by induction.

#### B.1 Base Case

In this section, we prove the base case of Proposition [4.2,](#page-5-1) i.e., with  $n = 1$ . The base case gives the desired mean squared error (MSE) convergence bound, which will subsequently be used in the proof of weak convergence.

We start by noting the following decomposition,

$$
\mathbb{E}[\|\theta_{k+1} - \theta^*\|^2] - \mathbb{E}[\|\theta_k - \theta^*\|^2]
$$
  
=  $2\alpha \mathbb{E}[\langle \theta_k - \theta^*, g(\theta_k, x_k) \rangle] + \alpha^2 \mathbb{E}[\|g(\theta_k, x_k)\|^2] + \alpha^2 \mathbb{E}[\|\xi_{k+1}(\theta_k)\|^2]$   
=  $2\alpha \mathbb{E}[\langle \theta_k - \theta^*, g(\theta_k, x_k) - \bar{g}(\theta_k) \rangle] + 2\alpha \mathbb{E}[\langle \theta_k - \theta^*, \bar{g}(\theta_k) \rangle]$   
+  $\alpha^2 \mathbb{E}[\|g(\theta_k, x_k)\|^2] + \alpha^2 \mathbb{E}[\|\xi_{k+1}(\theta_k)\|^2].$ 

It is easy to see that under Assumption [3,](#page-2-3) we have

<span id="page-14-1"></span>
$$
\langle \theta_k - \theta^*, \bar{g}(\theta_k) \rangle = \langle \theta_k - \theta^*, \bar{g}(\theta_k) - \bar{g}(\theta^*) \rangle \le -\mu \|\theta_k - \theta^*\|^2.
$$
 (B.1)

Additionally, under Assumption [2](#page-2-2) and [4,](#page-3-4) we have the following upper bound

$$
\alpha^2 \Big( \mathbb{E}[\|g(\theta_k, x_k)\|^2] + \mathbb{E}[\|\xi_{k+1}(\theta_k)\|^2] \Big) \leq \alpha^2 \Big( L_1^2 \mathbb{E}[(\|\theta_k - \theta^*\| + 1)^2] + L_2^2 \mathbb{E}[(\|\theta_k - \theta^*\| + 1)^2] \Big)
$$

$$
\leq 2\alpha^2 L^2 \Big( \mathbb{E}[\|\theta_k - \theta^*\|^2] + 1 \Big). \tag{B.2}
$$

<span id="page-15-0"></span>Therefore, the key to analyze the remaining inner product  $\langle \theta_k - \theta^*, g(\theta_k, x_k) - \bar{g}(\theta_k) \rangle$ .

Consider the following decomposition

$$
\langle \theta_k - \theta^*, g(\theta_k, x_k) - \bar{g}(\theta_k) \rangle = \langle \theta_k - \theta_{k-\tau}, g(\theta_k, x_k) - \bar{g}(\theta_k) \rangle \tag{B.3}
$$

<span id="page-15-8"></span><span id="page-15-3"></span><span id="page-15-2"></span>+ 
$$
\langle \theta_{k-\tau} - \theta^*, g(\theta_{k-\tau}, x_k) - \bar{g}(\theta_{k-\tau}) \rangle
$$
 (B.4)

+ 
$$
\langle \theta_{k-\tau} - \theta^*, g(\theta_k, x_k) - g(\theta_{k-\tau}, x_k) \rangle
$$
 (B.5)

<span id="page-15-11"></span><span id="page-15-10"></span><span id="page-15-7"></span><span id="page-15-6"></span><span id="page-15-5"></span><span id="page-15-4"></span><span id="page-15-1"></span>+ 
$$
\langle \theta_{k-\tau} - \theta^*, \bar{g}(\theta_k) - \bar{g}(\theta_{k-\tau}) \rangle
$$
. (B.6)

Hence, we need some upper bound on  $\|\theta_k - \theta_{k-\tau}\|$ .

We next note the following technical Lemma, which is adapted from [\[SY19](#page-12-3), [CZD](#page-10-2)+22] for the updated unbounded i.i.d. noise assumption in Assumption [4.](#page-3-4) The proof of the technical Lemma is delayed to Section [B.1.1.](#page-17-0)

<span id="page-15-9"></span>**Lemma B.1.** For  $16\alpha\tau \leq \mu/(4L^2)$ , we have

$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\| | \mathcal{F}_{k-\tau}] \le 2\alpha \tau L \|\theta_{k-\tau} - \theta^*\| + 2\alpha \tau L \tag{B.7}
$$

$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\| | \mathcal{F}_{k-\tau}] \le 4\alpha \tau L \mathbb{E}[\|\theta_k - \theta^*\| | \mathcal{F}_{k-\tau}] + 4\alpha \tau L
$$
\n(B.8)

$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\|^2 | \mathcal{F}_{k-\tau}] \le 8\alpha^2 \tau^2 L^2 \|\theta_{k-\tau} - \theta^*\|^2 + 8\alpha^2 \tau^2 L^2
$$
\n(B.9)

$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\|^2 | \mathcal{F}_{k-\tau}] \le 32\alpha^2 \tau^2 L^2 \mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 32\alpha^2 \tau^2 L^2. \tag{B.10}
$$

Given [\(B.10\)](#page-15-1), we additionally note that

$$
\|\theta_{k-\tau} - \theta^*\|^2 + 1 = \mathbb{E}[\|\theta_{k-\tau} - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 1
$$
  
\n
$$
\leq 2 \Big( \mathbb{E}[\|\theta_k - \theta_{k-\tau}\|^2 | \mathcal{F}_{k-\tau}] + \mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] \Big) + 1
$$
  
\n
$$
\leq 2 \Big( 32\alpha^2 \tau^2 L^2 (\mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 1) + \mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] \Big) + 1
$$
  
\n
$$
\leq 4 (\mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 1).
$$
\n(B.11)

We next use the above four technical inequalities to analyze the four terms in  $(B.3)$ – $(B.4)$ . To bound [\(B.3\)](#page-15-2), we first note that

$$
\|\mathbb{E}[\langle \theta_k - \theta_{k-\tau}, g(\theta_k, x_k) - \bar{g}(\theta_k) \rangle | \mathcal{F}_{k-\tau}] \| \n\leq \mathbb{E}[\|\theta_k - \theta_{k-\tau}\| \cdot 2L(\|\theta_k - \theta^*\| + 1) | \mathcal{F}_{k-\tau}] \n\stackrel{(i)}{\leq} 2L\sqrt{\mathbb{E}[\|\theta_k - \theta_{k-\tau}\|^2 | \mathcal{F}_{k-\tau}]} \sqrt{\mathbb{E}[(\|\theta_k - \theta^*\| + 1)^2 | \mathcal{F}_{k-\tau}]} \n\stackrel{(ii)}{\leq} 2L\sqrt{32\alpha^2 \tau^2 L^2(\mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 1)} \sqrt{2(\mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 1)} \n\leq 16\alpha \tau L^2(\mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 1),
$$

where (i) holds for the Cauchy-Schwarz inequality and (ii) holds for  $(B.10)$ .

To bound [\(B.4\)](#page-15-3), we next note that

$$
\|\mathbb{E}[\langle \theta_{k-\tau} - \theta^*, g(\theta_{k-\tau}, x_k) - \bar{g}(\theta_{k-\tau})\rangle | \mathcal{F}_{k-\tau}] \| \n= \| \langle \theta_{k-\tau} - \theta^*, \mathbb{E}[g(\theta_{k-\tau}, x_k) - \bar{g}(\theta_{k-\tau}) | \mathcal{F}_{k-\tau}] \rangle \| \n\le \| \theta_{k-\tau} - \theta^* \| \| \mathbb{E}[g(\theta_{k-\tau}, x_k) - \bar{g}(\theta_{k-\tau}) | \mathcal{F}_{k-\tau}] \| \n\overset{\text{(iii)}}{\le} \| \theta_{k-\tau} - \theta^* \| \cdot \left( \alpha L(\|\theta_{k-\tau} - \theta^* \| + 1) \right) \n\le 2\alpha L(\|\theta_{k-\tau} - \theta^* \|^2 + 1) \n\overset{\text{(iv)}}{\le} 8\alpha L(\mathbb{E}[\|\theta_k - \theta^* \|^2 | \mathcal{F}_{k-\tau}] + 1)
$$

$$
\leq 8\alpha\tau L^{2}(\mathbb{E}[\|\theta_{k}-\theta^{*}\|^{2}|\mathcal{F}_{k-\tau}]+1),
$$

where (iii) holds due to the mixing property of Markov chain  $(x_k)_{k\geq 0}$  and (iv) holds for [\(B.11\)](#page-15-4). To bound  $(B.5)$ , we have

$$
\|\mathbb{E}[\langle \theta_{k-\tau} - \theta^*, g(\theta_k, x_k) - g(\theta_{k-\tau}, x_k) \rangle | \mathcal{F}_{k-\tau}] \|
$$
\n
$$
= \| \langle \theta_{k-\tau} - \theta^*, \mathbb{E}[g(\theta_k, x_k) - g(\theta_{k-\tau}, x_k) | \mathcal{F}_{k-\tau}] \rangle \|
$$
\n
$$
\leq L \|\theta_{k-\tau} - \theta^*\| \cdot \mathbb{E}[\|\theta_k - \theta_{k-\tau}\| | \mathcal{F}_{k-\tau}]
$$
\n
$$
\leq L \|\theta_{k-\tau} - \theta^*\| \cdot 2\alpha \tau L (\|\theta_{k-\tau} - \theta^*\| + 1)
$$
\n
$$
\leq 4\alpha \tau L^2 (\|\theta_{k-\tau} - \theta^*\|^2 + 1)
$$
\n
$$
\leq 16\alpha \tau L^2 (\mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 1),
$$

where  $(v)$  holds for  $(B.7)$  and  $(vi)$  holds for  $(B.11)$ .

Lastly, to bound  $(B.6)$ , we apply the similar technique used in bounding the third term in  $(B.5)$  and obtain a similar result

$$
\|\mathbb{E}[\langle \theta_{k-\tau}-\theta^*, \bar{g}(\theta_k)-\bar{g}(\theta_{k-\tau})\rangle | \mathcal{F}_{k-\tau}]\| \leq 16\alpha\tau L^2(\mathbb{E}[\|\theta_k-\theta^*\|^2 | \mathcal{F}_{k-\tau}]+1).
$$

Combining all analyses above, we have

<span id="page-16-0"></span>
$$
\|2\alpha \mathbb{E}[\langle \theta_k - \theta^* \rangle, g(\theta_k, x_k) - \bar{g}(\theta_k)|\mathcal{F}_{k-\tau}]\|
$$
  
\n
$$
\leq 2\alpha (16\alpha \tau L^2 + 8\alpha \tau L^2 + 32\alpha \tau L^2) (\mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 1)
$$
  
\n
$$
\leq 112\alpha^2 \tau L^2 (\mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 1).
$$
\n(B.12)

Hence, making use of  $(B.1)$ ,  $(B.2)$ , and  $(B.12)$ , we obtain the following

$$
\mathbb{E}[\|\theta_{k+1} - \theta^*\|^2 | \mathcal{F}_{k-\tau}] - \mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] \n\leq -2\alpha\mu \mathbb{E}[\|\theta - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 112\alpha^2 \tau L^2 (\mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 1) \n+ 2\alpha^2 L^2 (\mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 1) \n\leq -2\alpha\mu \mathbb{E}[\|\theta - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 114\alpha^2 \tau L^2 \mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 114\alpha^2 \tau L^2 \n= -2\alpha(\mu - 57\alpha \tau L^2) \mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 114\alpha^2 \tau L^2.
$$

Therefore, when we have  $\alpha$  satisfying the constraint, i.e.,  $\alpha \tau L^2 < c_{2,1} \mu$ , we obtain

$$
\mathbb{E}[\|\theta_{k+1} - \theta^*\|^2 | \mathcal{F}_{k-\tau}] \le (1 - \alpha \mu) \mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 114\alpha^2 \tau L^2.
$$

Recursively, we get

$$
\mathbb{E}[\|\theta_k - \theta^*\|^2] \le (1 - \alpha \mu)^{k-\tau} \mathbb{E}[\|\theta_\tau - \theta^*\|^2] + \frac{114\alpha\tau L^2}{\mu}
$$
  
\n
$$
\le 2(1 - \alpha \mu)^{k-\tau} \Big( \mathbb{E}[\|\theta_0 - \theta^*\|^2] + \mathbb{E}[\|\theta_\tau - \theta_0\|^2] \Big) + \frac{114\alpha\tau L^2}{\mu}
$$
  
\n
$$
\le 2(1 - \alpha \mu)^{k-\tau} \Big( \mathbb{E}[\|\theta_0 - \theta^*\|^2] + 8\alpha^2\tau^2 L^2 \Big( \mathbb{E}[\|\theta_0 - \theta^*\|^2] + 1 \Big) + \frac{114\alpha\tau L^2}{\mu}
$$
  
\n
$$
\le 4(1 - \alpha \mu)^{k-\tau} \mathbb{E}[\|\theta_0 - \theta^*\|^2] + \frac{122\alpha\tau L^2}{\mu}.
$$

Lastly, we note that

<span id="page-16-1"></span>
$$
\frac{1}{(1-\alpha\mu)^{\tau}} \stackrel{\text{(i)}}{\leq} \frac{1}{1-\alpha\tau\mu} \stackrel{\text{(ii)}}{\leq} \frac{1}{1-\alpha\tau L} \stackrel{\text{(iii)}}{\leq} 2,\tag{B.13}
$$

where (i) holds by the Bernoulli inequality, that  $(1+x)^r \ge 1+rx$  for  $x \ge -1$  and  $r \ge 1$ ; (ii) holds for  $\mu \le L$ ; (iii) holds for  $\alpha \tau L < \mu/(114L) < \frac{1}{2}$ .

Hence, for  $k \geq \tau$ , we have

$$
\mathbb{E}[\|\theta_k - \theta^*\|^2] \le c_{2,1} (1 - \alpha \mu)^k \|\theta_0 - \theta^*\|^2 + c_{2,2} \alpha \tau_\alpha \frac{L^2}{\mu},
$$

for  $c_{2,1}$  and  $c_{2,2}$  some universal constants. As such, we have completed the proof of base case for Proposition [4.2.](#page-5-1)

#### <span id="page-17-0"></span>B.1.1 Proof of Lemma [B.1](#page-15-9)

In this section, we provide the proofs of the four technical inequalities in Lemma [B.1.](#page-15-9)

### Proof of [\(B.7\)](#page-15-6).

$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\| | \mathcal{F}_{k-\tau}] \leq 2\alpha \tau L \|\theta_{k-\tau} - \theta^*\| + 2\alpha \tau L.
$$

Proof. Note that

$$
\|\theta_k - \theta_{k-\tau}\| \le \sum_{t=k-\tau}^{k-1} \|\theta_{t+1} - \theta_t\|,
$$

so we start with analyzing  $\|\theta_{t+1} - \theta_t\|.$ 

$$
\|\theta_{t+1} - \theta^*\| - \|\theta_t - \theta^*\| \le \|\theta_{t+1} - \theta_t\| = \alpha \|g(\theta_t, x_t) + \xi_{t+1}(\theta_t)\|
$$
  
\n
$$
\le \alpha \|g(\theta_t, x_t)\| + \alpha \|\xi_{t+1}(\theta_t)\| \le \alpha L_1(\|\theta_t - \theta^*\| + 1) + \alpha \|\xi_{t+1}(\theta_t)\|
$$
  
\n
$$
\|\theta_{t+1} - \theta^*\| \le (1 + \alpha L_1) \|\theta_t - \theta^*\| + \alpha L_1 + \alpha \|\xi_{t+1}(\theta_t)\|.
$$

Recall that we assume

$$
\mathbb{E}^{1/2}[\|\xi_{t+1}(\theta_t)\|^2|\mathcal{F}_t] \le L_2(\|\theta_t\|+1),
$$

then we have for  $k - \tau \leq t \leq k$ ,

$$
\mathbb{E}[\|\xi_{t+1}(\theta_t)\||\mathcal{F}_{k-\tau}] = \mathbb{E}[\mathbb{E}[\|\xi_{t+1}(\theta_k)|\mathcal{F}_t]|\mathcal{F}_{k-\tau}] \leq \mathbb{E}[L_2(\|\theta_k\|+1)|\mathcal{F}_{k-\tau}]
$$
  

$$
\mathbb{E}[\|\theta_{t+1} - \theta_t\||\mathcal{F}_{k-\tau}] \leq \alpha L(\mathbb{E}[\|\theta_t - \theta^*\||\mathcal{F}_{k-\tau}] + 1)
$$
  

$$
\mathbb{E}[\|\theta_{k+1} - \theta^*\||\mathcal{F}_{k-\tau}] \leq (1 + \alpha L)\mathbb{E}[\|\theta_k - \theta^*\||\mathcal{F}_{k-\tau}] + \alpha L.
$$

Hence, for  $0\leq n\leq \tau,$ 

$$
\mathbb{E}[\|\theta_{k-\tau+n} - \theta^*\| | \mathcal{F}_{k-\tau}] \le (1+\alpha L)^n \mathbb{E}[\|\theta_{k-\tau} - \theta^*\| | \mathcal{F}_{k-\tau}] + \alpha L \sum_{l=0}^{n-1} (1+\alpha L)^l
$$
  
=  $(1+\alpha L)^n \|\theta_{k-\tau} - \theta^*\| + ((1+\alpha L)^n - 1).$ 

We next note that

$$
(1+x)^y = e^{y \log(1+x)} \le e^{xy} \le 1 + 2xy, xy \in [0, 1/2].
$$

Hence, at this stage, if we require  $\alpha \tau L < \mu/(4L) < 1/4$ , we have the following upper bound

$$
(1 + \alpha L)^n \le (1 + \alpha L)^\tau \le 1 + 2\alpha \tau L \le 2.
$$

Therefore, for  $0 \leq n \leq \tau$ ,

$$
\mathbb{E}[\|\theta_{k-\tau+n} - \theta^*\| | \mathcal{F}_{k-\tau}] \le (1 + 2\alpha \tau L) \|\theta_{k-\tau} - \theta^*\| + 2\alpha \tau L
$$
  

$$
\le 2 \|\theta_{k-\tau} - \theta^*\| + 2\alpha \tau L.
$$

As such, we have

$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\| | \mathcal{F}_{k-\tau}] \leq \sum_{t=k-\tau}^{k-1} \mathbb{E}[\|\theta_{t+1} - \theta_t\| | \mathcal{F}_{k-\tau}]
$$

$$
\leq \alpha L \sum_{t=k-\tau}^{k-1} \mathbb{E}[\|\theta_t - \theta^*\| | \mathcal{F}_{k-\tau}] + \alpha \tau L
$$
  
\n
$$
\leq \alpha \tau L(2\|\theta_{k-\tau} - \theta^*\| + 2\alpha \tau L) + \alpha \tau L
$$
  
\n
$$
\leq 2\alpha \tau L \|\theta_{k-\tau} - \theta^*\| + 2\alpha \tau L,
$$

and prove the desired inequality.

Proof of  $(B.8)$ .

$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\||\mathcal{F}_{k-\tau}] \le 4\alpha\tau L\mathbb{E}[\|\theta_k - \theta^*\||\mathcal{F}_{k-\tau}] + 4\alpha\tau L.
$$

Proof. We prove this inequality based on the claim that we have just shown,

$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\||\mathcal{F}_{k-\tau}] \le 2\alpha \tau L \|\theta_{k-\tau} - \theta^*\| + 2\alpha \tau L.
$$

We simply note that

$$
\|\theta_{k-\tau} - \theta^*\| = \mathbb{E}[\|\theta_{k-\tau} - \theta^*\| | \mathcal{F}_{k-\tau}]
$$
  
\$\leq\$  $\mathbb{E}[\|\theta_k - \theta_{k-\tau} \| | \mathcal{F}_{k-\tau}] + \mathbb{E}[\|\theta_k - \theta^*\| | \mathcal{F}_{k-\tau}].$ 

Hence,

$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\||\mathcal{F}_{k-\tau}] \le 2\alpha\tau L(\mathbb{E}[\|\theta_k - \theta_{k-\tau}\||\mathcal{F}_{k-\tau}] + \mathbb{E}[\|\theta_k - \theta^*\||\mathcal{F}_{k-\tau}] + 1)
$$
  

$$
(1 - 2\alpha\tau L)\mathbb{E}[\|\theta_k - \theta_{k-\tau}\||\mathcal{F}_{k-\tau}] \le 2\alpha\tau L\mathbb{E}[\|\theta_k - \theta^*\||\mathcal{F}_{k-\tau}] + 2\alpha\tau L.
$$

Therefore, we obtain

$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\||\mathcal{F}_{k-\tau}] \le 4\alpha\tau L \mathbb{E}[\|\theta_k - \theta^*\||\mathcal{F}_{k-\tau}] + 4\alpha\tau L.
$$

Proof of [\(B.9\)](#page-15-11).

$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\|^2 | \mathcal{F}_{k-\tau}] \leq 32\alpha^2 \tau^2 L^2 \mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 32\alpha^2 \tau^2 L^2.
$$

*Proof.* To analyze  $\mathbb{E}[\|\theta_k - \theta_{k-\tau}\|^2 | \mathcal{F}_{k-\tau}]$ , we consider the following attempt.

$$
\mathbb{E}[\|\theta_{k} - \theta_{k-\tau}\|^{2} | \mathcal{F}_{k-\tau}] \leq \tau \sum_{t=k-\tau}^{k-1} \mathbb{E}[\|\theta_{t+1} - \theta_{t}\|^{2} | \mathcal{F}_{k-\tau}]
$$
\n
$$
= \alpha^{2} \tau \sum_{t=k-\tau}^{k-1} \mathbb{E}[(\|g(\theta_{t}, x_{t})\| + \|\xi_{t+1}(\theta_{t})\|)^{2} | \mathcal{F}_{k-\tau}]
$$
\n
$$
\leq 2\alpha^{2} \tau \sum_{t=k-\tau}^{k-1} \left( \mathbb{E}[\|g(\theta_{t}, x_{t})\|^{2} | \mathcal{F}_{k-\tau}] + \mathbb{E}[\|\xi_{t+1}(\theta_{t})\|^{2} | \mathcal{F}_{k-\tau}]\right)
$$
\n
$$
\leq 2\alpha^{2} \tau L^{2} \sum_{t=k-\tau}^{k-1} \mathbb{E}[(\|\theta_{t} - \theta^{*}\| + 1)^{2} | \mathcal{F}_{k-\tau}]
$$
\n
$$
\leq 4\alpha^{2} \tau L^{2} \sum_{t=k-\tau}^{k-1} \mathbb{E}[\|\theta_{t} - \theta^{*}\|^{2} | \mathcal{F}_{k-\tau}] + 4\alpha^{2} \tau^{2} L^{2}.
$$

Next, we study  $\mathbb{E}[\|\theta_t - \theta^*\|^2 | \mathcal{F}_{k-\tau}]$ . We start with the following, for  $k-\tau \le t < k$ ,

$$
\mathbb{E}[\|\theta_{t+1}-\theta^*\|^2|\mathcal{F}_{k-\tau}]
$$
  
= 
$$
\mathbb{E}[\|\theta_t-\theta^*\|^2|\mathcal{F}_{k-\tau}] + 2\alpha \mathbb{E}[\langle\theta_t-\theta^*, g(\theta_t, x_t)\rangle|\mathcal{F}_{k-\tau}] + \alpha^2 \mathbb{E}[\|g(\theta_t, x_t) + \xi_{t+1}(\theta_t)\|^2|\mathcal{F}_{k-\tau}]
$$



 $\Box$ 

$$
\leq \mathbb{E}[\|\theta_t - \theta^*\|^2 |\mathcal{F}_{k-\tau}] + 2\alpha^2 \Big( \mathbb{E}[\|g(\theta_t, x_t)\|^2 |\mathcal{F}_{k-\tau}] + \mathbb{E}[\|\xi_{t+1}(\theta_t)\|^2 |\mathcal{F}_{k-\tau}]\Big) + 2\alpha \mathbb{E}[\|\theta_t - \theta^*\| \|g(\theta_t, x_t)\| |\mathcal{F}_{k-\tau}].
$$

We note that

$$
2\mathbb{E}[\|\theta_t - \theta^*\|\|g(\theta_t, x_t)\||\mathcal{F}_{k-\tau}] \le 2\sqrt{\mathbb{E}[\|\theta_t - \theta^*\|^2|\mathcal{F}_{k-\tau}]\mathbb{E}[\|g(\theta_t, x_t)\|^2|\mathcal{F}_{k-\tau}]}\le 2\sqrt{\mathbb{E}[\|\theta_t - \theta^*\|^2|\mathcal{F}_{k-\tau}]\mathbb{E}[(L(\|\theta_t - \theta^*\| + 1))^2|\mathcal{F}_{k-\tau}]}\le 2\sqrt{\mathbb{E}[\|\theta_t - \theta^*\|^2|\mathcal{F}_{k-\tau}]2L^2\mathbb{E}[\|\theta_t - \theta^*\|^2 + 1|\mathcal{F}_{k-\tau}]}\le 4L(\mathbb{E}[\|\theta_t - \theta^*\|^2|\mathcal{F}_{k-\tau}] + 1).
$$

Substituting the above inequality back, we obtain

$$
\mathbb{E}[\|\theta_{t+1} - \theta^*\|^2 | \mathcal{F}_{k-\tau}]
$$
  
\n
$$
\leq \mathbb{E}[\|\theta_t - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 4\alpha^2 L^2 \Big( \mathbb{E}[\|\theta_t - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 1 \Big) + 4\alpha L \mathbb{E}[\|\theta_t - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 4\alpha L
$$
  
\n
$$
\leq (1 + 4\alpha^2 L^2 + 4\alpha L) \mathbb{E}[\|\theta_t - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + (4\alpha^2 L^2 + 4\alpha L).
$$

We further recall that

$$
4\alpha^2 L^2 \le 4\alpha L(\alpha \tau L) \le \alpha L,
$$

and hence we obtain the following upper bound

$$
\mathbb{E}[\|\theta_{t+1} - \theta^*\|^2 | \mathcal{F}_{k-\tau}] \le (1 + 5\alpha L) \mathbb{E}[\|\theta_t - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 5\alpha L.
$$

Then, recursively, for  $0\leq n\leq \tau,$  we have

$$
\mathbb{E}[\|\theta_{k-\tau+n}-\theta^*\|^2|\mathcal{F}_{k-\tau}]\leq (1+5\alpha L)^n \|\theta_{k-\tau}-\theta^*\|^2+5\alpha L\sum_{l=0}^{n-1} (1+5\alpha L)^l.
$$

As such, under the assumption that  $4\alpha\tau L<\mu/(4L)<1/4,$  then for  $k-\tau\leq t\leq k,$  we have

$$
\mathbb{E}[\|\theta_t - \theta^*\|^2 | \mathcal{F}_{k-\tau}] \le (1 + 10\alpha \tau L) \|\theta_{k-\tau} - \theta^*\|^2 + 10\alpha \tau L
$$
  

$$
\le 2 \|\theta_{k-\tau} - \theta^*\|^2 + 10\alpha \tau L.
$$

Combining all the analyses above, we have

$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\|^2 | \mathcal{F}_{k-\tau}] \le 4\alpha^2 \tau L^2 \sum_{t=k-\tau}^{k-1} \mathbb{E}[\|\theta_t - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 4\alpha^2 \tau^2 L^2
$$
  

$$
\le 4\alpha^2 \tau^2 L^2 \left(2\|\theta_{k-\tau} - \theta^*\|^2 + 10\alpha \tau L\right) + 4\alpha^2 \tau^2 L^2
$$
  

$$
\le 8\alpha^2 \tau^2 L^2 \|\theta_{k-\tau} - \theta^*\|^2 + 8\alpha^2 \tau^2 L^2.
$$



Proof of  $(B.10)$ .

$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\|^2 | \mathcal{F}_{k-\tau}] \leq 32\alpha^2 \tau^2 L^2 \mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 32\alpha^2 \tau^2 L^2.
$$

*Proof.* This inequality simply extends the result from  $(B.9)$ , i.e.,

$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\|^2 | \mathcal{F}_{k-\tau}] \leq 8\alpha^2 \tau^2 L^2 \|\theta_{k-\tau} - \theta^*\|^2 + 8\alpha^2 \tau^2 L^2.
$$

We first note that

$$
\|\theta_{k-\tau} - \theta^*\|^2 = \mathbb{E}[\|\theta_{k-\tau} - \theta^*\|^2 | \mathcal{F}_{k-\tau}]
$$

$$
\leq 2\mathbb{E}[\|\theta_k-\theta_{k-\tau}\|^2|\mathcal{F}_{k-\tau}]+2\mathbb{E}[\|\theta_k-\theta^*\|^2|\mathcal{F}_{k-\tau}].
$$

Hence,

$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\|^2 | \mathcal{F}_{k-\tau}] \le 8\alpha^2 \tau^2 L^2 (2\mathbb{E}[\|\theta_k - \theta_{k-\tau}\|^2 | \mathcal{F}_{k-\tau}] + 2\mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 1) (1 - 16\alpha^2 \tau^2 L^2)\mathbb{E}[\|\theta_k - \theta_{k-\tau}\|^2 | \mathcal{F}_{k-\tau}] \le 16\alpha^2 \tau^2 L^2 \mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 8\alpha^2 \tau^2 L^2.
$$

Again, under the assumption that  $16\alpha\tau L^2 < \mu/4$ , we can conclude that

$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\|^2 | \mathcal{F}_{k-\tau}] \le 32\alpha^2 \tau^2 L^2 \mathbb{E}[\|\theta_k - \theta^*\|^2 | \mathcal{F}_{k-\tau}] + 32\alpha^2 \tau^2 L^2.
$$

 $\Box$ 

### B.2 Induction Step

In this step, assume that the moment bound in Proposition [4.2](#page-5-1) has been proven for  $k \leq n-1$ , we now proceed to show that the desired moment convergence holds for n with  $2 \le n \le p$ .

We start with the following decomposition of  $\|\theta_{k+1} - \theta^*\|^{2n}$ 

$$
\|\theta_{k+1} - \theta^*\|^{2n} = \left( \|\theta_k - \theta^*\|^2 + 2\alpha \langle \theta_k - \theta^*, g(\theta_k, x_k) + \xi_{k+1}(\theta_k) \rangle + \alpha^2 \|g(\theta_k, x_k) + \xi_{k+1}(\theta_t)\|^2 \right)^n
$$
  
= 
$$
\sum_{\substack{i,j,l \\ i+j+l=n}} {n \choose i,j,l} \|\theta_k - \theta^*\|^{2i} \Big( 2\alpha \langle \theta_k - \theta^*, g(\theta_k, x_k) + \xi_{k+1}(\theta_k) \rangle \Big)^j \Big( \alpha \|g(\theta_k, x_k) + \xi_{k+1}(\theta_k) \| \Big)^{2l}
$$

We note the following cases.

- 1.  $i = n, j = l = 0$ . In this case, the summand is simply  $\|\theta_k \theta^*\|^{2i}$ .
- 2. When  $i = n 1$ ,  $j = 1$  and  $l = 0$ . In this case, the summand is of order  $\alpha$ , i.e.,  $\alpha 2n(\theta_k \theta^*, g(\theta_k, x_k) +$  $\xi_{k+1}(\theta_k)$ <sup>j</sup> $\|\theta_k - \theta^*\|^{2(n-1)}$ . We can further compose it as

$$
2n\alpha\langle\theta_k-\theta^*,g(\theta_k,x_k)+\xi_{k+1}(\theta_k)\rangle\|\theta_k-\theta^*\|^{2(n-1)}=\underbrace{2n\alpha\langle\theta_k-\theta^*,g(\theta_k,x_k)-\bar{g}(\theta_k)+\xi_{k+1}(\theta_k)\rangle\|\theta_k-\theta^*\|^{2(n-1)}}_{T_1}+\underbrace{2n\alpha\langle\theta_k-\theta^*,\bar{g}(\theta_k)\rangle\|\theta_k-\theta^*\|^{2(n-1)}}_{T_2}.
$$

Note that, when  $(x_k)$  is i.i.d. or from a martingale noise sequence, we have

$$
\mathbb{E}[T_1|\theta_k] = 0.
$$

However, when  $(x_k)$  is Markovian, the above equality does not hold and  $T_1$  requires careful analysis. Nonetheless, under the strong monotonicity assumption, we have

$$
T_2 \le -2n\alpha\mu \|\theta_k - \theta^*\|^{2n}.
$$

3. For the remaining terms, we see that they are of higher orders of  $\alpha$ . Therefore, when  $\alpha$  is selected sufficiently small, these terms do not raise concern.

Therefore, to prove the desired moment bound, we spend the remaining section analyzing  $T_1$ . Immediately, we note that

$$
\mathbb{E}[T_1|\mathcal{F}_{k-\tau}] = \mathbb{E}\Big[2n\alpha\langle\theta_k - \theta^*, g(\theta_k, x_k) - \bar{g}(\theta_k) + \mathbb{E}[\xi_{k+1}(\theta_k)|\theta_k]\rangle\|\theta_k - \theta^*\|^{2(n-1)}|\mathcal{F}_{k-\tau}\Big]
$$
  
= 
$$
\mathbb{E}\Big[2n\alpha\langle\theta_k - \theta^*, g(\theta_k, x_k) - \bar{g}(\theta_k)\rangle\|\theta_k - \theta^*\|^{2(n-1)}|\mathcal{F}_{k-\tau}\Big].
$$

Subsequently, we focus on analyzing  $T_1'$ .

We start with the following decomposition of  $T_1'$ .

$$
2n\alpha \langle g(\theta_k, x_k) - \bar{g}(\theta_k), \theta_k - \theta^* \rangle \|\theta_k - \theta^*\|^{2(n-1)}
$$

$$
\leq 2n\alpha \|g(\theta_{k-\tau}, x_k) - \bar{g}(\theta_{k-\tau})\| \|\theta_{k-\tau} - \theta^*\|^{2n-1}
$$
\n(B.14)

$$
+2n\alpha||g(\theta_k, x_k) - g(\theta_{k-\tau}, x_k)||\|\theta_{k-\tau} - \theta^*\|^{2n-1}
$$
\n(B.15)

$$
+ 2n\alpha \|\bar{g}(\theta_{k-\tau}) - \bar{g}(\theta_k)\|\|\theta_{k-\tau} - \theta^*\|^{2n-1}
$$
\n(B.16)

$$
+ 2n\alpha \|g(\theta_k, x_k) - \bar{g}(\theta_k)\| \|\theta_k - \theta^*\|^{2(n-1)} \|\theta_k - \theta_{k-\tau}\|
$$
\n(B.17)

$$
+2n\alpha||g(\theta_k,x_k)-\bar{g}(\theta_k)||\|\theta_{k-\tau}-\theta^*\| \cdot ( \|\theta_k-\theta^*\|^{2(n-1)} - \|\theta_{k-\tau}-\theta^*\|^{2(n-1)} ). \tag{B.18}
$$

We note the following technical lemma, which will offer significant help in the analysis of  $T_1'$ . We postpone the proof of the lemma to the end of this subsection.

<span id="page-21-5"></span>**Lemma B.2.** For  $\tilde{c}_n \alpha \tau \leq \mu/(4L^2)$ , where  $\tilde{c}_n$  denotes some constant dependent of the higher-moment  $2n$ , we have

<span id="page-21-4"></span><span id="page-21-3"></span><span id="page-21-2"></span><span id="page-21-1"></span><span id="page-21-0"></span>
$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\|^{2n} |\mathcal{F}_{k-\tau}] \le c_n \alpha^{2n} \tau^{2n} L^{2n}(\|\theta_{k-\tau} - \theta^*\|^{2n} + 1).
$$

Following the lemma, we observe that a natural consequence is for any  $m \leq 2n$ , we have

$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\|^m | \mathcal{F}_{k-\tau}] \leq \left(\mathbb{E}[\|\theta_k - \theta_{k-\tau}\|^{2n} | \mathcal{F}_{k-\tau}]\right)^{\frac{m}{2n}} \leq \left(c_n \alpha^{2n} \tau^{2n} L^{2n} (\|\theta_{k-\tau} - \theta^*\|^{2n} + 1)\right)^{\frac{m}{2n}} \leq c_m \alpha^m \tau^m L^m \left(\|\theta_{k-\tau} - \theta^*\|^m + 1\right),
$$

where we use the inequality  $a^p + b^p > (a + b)^p$  for  $a, b > 0$ ,  $p \in (0, 1)$  to obtain the final inequality.

Now, we are ready to analyze [\(B.14\)](#page-21-0)–[\(B.18\)](#page-21-1). Firstly, for [\(B.14\)](#page-21-0), we make use of the mixing assumption of  $\tau$ , and have that

$$
\mathbb{E}[(|B.14)||\mathcal{F}_{k-\tau}] \le 2n\alpha \|\theta_{k-\tau} - \theta^*\|^{2n-1} \mathbb{E}[\|g(\theta_{k-\tau}, x_k) - \bar{g}(\theta_{k-\tau})\|] \mathcal{F}_{k-\tau}]
$$
  
\n
$$
\le 2n\alpha^2 L \|\theta_{k-\tau} - \theta^*\|^{2n-1} (\|\theta_{k-\tau} - \theta^*\| + 1)
$$
  
\n
$$
\le 2n\alpha^2 L \|\theta_{k-\tau} - \theta^*\|^{2n} + 2n\alpha^2 L \|\theta_{k-\tau} - \theta^*\|^{2n-1}
$$
  
\n
$$
\le 3n\alpha^2 L \|\theta_{k-\tau} - \theta^*\|^{2n} + n\alpha^2 L \|\theta_{k-\tau} - \theta^*\|^{2(n-1)},
$$

where we make use of the inequality  $2|x|^3 \leq x^2 + x^4$  to obtain the final step. Next, we proceed to analyze  $(B.15)$ . It is easy to see that

$$
\mathbb{E}[|(B.15)||\mathcal{F}_{k-\tau}] = 2n\alpha \|\theta_{k-\tau} - \theta^*\|^{2n-1} \mathbb{E}[\|g(\theta_k, x_k) - g(\theta_{k-\tau}, x_k)|\mathcal{F}_{k-\tau}]
$$
  
\n
$$
\leq 2n\alpha \|\theta_{k-\tau} - \theta^*\|^{2n-1} \mathbb{E}[\|\theta_k - \theta_{k-\tau}\||\mathcal{F}_{k-\tau}]
$$
  
\n
$$
\leq 2n\alpha \|\theta_{k-\tau} - \theta^*\|^{2n-1} \Big(2\alpha \tau L(\|\theta_{k-\tau} - \theta^*\| + 1)\Big)
$$
  
\n
$$
\leq 4n\alpha^2 \tau L \|\theta_{k-\tau} - \theta^*\|^{2n} + 4n\alpha^2 \tau L \|\theta_{k-\tau} - \theta^*\|^{2n-1}
$$
  
\n
$$
\leq 6n\alpha^2 \tau L \|\theta_{k-\tau} - \theta^*\|^{2n} + 2n\alpha^2 \tau L \|\theta_{k-\tau} - \theta^*\|^{2(n-1)}.
$$

The term in  $(B.16)$  can be analyzed in a similar fashion as the  $(B.15)$ .

For [\(B.17\)](#page-21-4), we first derive the following

$$
\mathbb{E}[|(B.17)||\mathcal{F}_{k-\tau}]
$$
\n
$$
\leq 2n\alpha \mathbb{E}\Big[2L\Big(||\theta_{k}-\theta^{*}||+1\Big)||\theta_{k}-\theta_{k-\tau}||||\theta_{k}-\theta^{*}||^{2(n-1)}|\mathcal{F}_{k-\tau}\Big]
$$
\n
$$
=\underbrace{4n\alpha L\mathbb{E}[||\theta_{k}-\theta_{k-\tau}||||\theta_{k}-\theta^{*}||^{2n-1}|\mathcal{F}_{k-\tau}]}_{T_{a}}+\underbrace{4n\alpha L\mathbb{E}[||\theta_{k}-\theta_{k-\tau}||||\theta_{k}-\theta^{*}||^{2(n-1)}|\mathcal{F}_{k-\tau}]}_{T_{b}}
$$

.

We next analyze the two terms  $T_a$  and  $T_b$  respectively. Starting with  $T_a$ , we have

$$
4n\alpha L\mathbb{E}[\|\theta_k - \theta_{k-\tau}\|\|\theta_k - \theta^*\|^{2n-1}|\mathcal{F}_{k-\tau}]
$$
\n(B.19)

$$
\leq 4n\alpha L\mathbb{E}\Big[\|\theta_k - \theta_{k-\tau}\|\Big(\|\theta_k - \theta_{k-\tau}\| + \|\theta_{k-\tau} - \theta^*\|\Big)^{2n-1}|\mathcal{F}_{k-\tau}\Big] \tag{B.20}
$$

$$
\leq 2^{2(n-1)} 4n\alpha L \mathbb{E}[\|\theta_k - \theta_{k-\tau}\| (\|\theta_k - \theta_{k-\tau}\|^{2n-1} + \|\theta_{k-\tau} - \theta^*\|^{2n-1} | \mathcal{F}_{k-\tau}]
$$
(B.21)

$$
=4^n n\alpha L\left(\underbrace{\mathbb{E}[\|\theta_k-\theta_{k-\tau}\|^{2n}|\mathcal{F}_{k-\tau}]}_{\text{by Lemma B.2}}+\|\theta_{k-\tau}-\theta^*\|^{2n-1}\mathbb{E}[\|\theta_k-\theta_{k-\tau}\||\mathcal{F}_{k-\tau}]\right) \tag{B.22}
$$

$$
\leq 4^n n \alpha L \Big( \underbrace{c_n \alpha^{2n} \tau^{2n} L^{2n}}_{\leq 2\alpha \tau L} (\|\theta_{k-\tau} - \theta^*\|^{2n} + 1) + \|\theta_{k-\tau} - \theta^*\|^{2n-1} (2\alpha \tau L(\|\theta_{k-\tau} - \theta^*\| + 1)) \Big) \tag{B.23}
$$

$$
\leq 4^n n \alpha L \Big( 4\alpha \tau L \|\theta_{k-\tau} - \theta^*\|^{2n} + 2\alpha \tau L \|\theta_{k-\tau} - \theta^*\|^{2n-1} + c_n \alpha^{2n} \tau^{2n} L^{2n} \Big) \tag{B.24}
$$

$$
\leq 4^n n \alpha L \Big( 5 \alpha \tau L \| \theta_{k-\tau} - \theta^* \|^2 + \alpha \tau L \| \theta_{k-\tau} - \theta^* \|^2 (n-1)} + c'_n \alpha^{2n-1} \tau^{2n-1} L^{2n-1} \Big). \tag{B.25}
$$

For  $T_b$ , we have

$$
4n\alpha L\mathbb{E}[\|\theta_{k} - \theta_{k-\tau}\|\|\theta_{k} - \theta^{*}\|^{2(n-1)}|\mathcal{F}_{k-\tau}]
$$
  
\n
$$
\leq 4n\alpha L\mathbb{E}\Big[\|\theta_{k} - \theta_{k-\tau}\|\Big(\|\theta_{k} - \theta_{k-\tau}\| + \|\theta_{k-\tau} - \theta^{*}\|\Big)^{2(n-1)}|\mathcal{F}_{k-\tau}\Big]
$$
  
\n
$$
\leq 2^{2n-1}n\alpha L\mathbb{E}[\|\theta_{k} - \theta_{k-\tau}\|(\|\theta_{k} - \theta_{k-\tau}\|^{2(n-1)} + \|\theta_{k-\tau} - \theta^{*}\|^{2(n-1)}|\mathcal{F}_{k-\tau}]
$$
  
\n
$$
= 2^{2n-1}n\alpha L\Big(\underbrace{\mathbb{E}[\|\theta_{k} - \theta_{k-\tau}\|^{2n-1}|\mathcal{F}_{k-\tau}]}_{\text{by Lemma B.2}} + \|\theta_{k-\tau} - \theta^{*}\|^{2(n-1)}\mathbb{E}[\|\theta_{k} - \theta_{k-\tau}\||\mathcal{F}_{k-\tau}]\Big)
$$
  
\n
$$
\leq 2^{2n-1}n\alpha L\Big(\underbrace{c_{n-1}\alpha^{2n-1}\tau^{2n-1}L^{2n-1}}_{\leq 2\alpha\tau L}(\|\theta_{k-\tau} - \theta^{*}\|^{2n-1} + 1) + \|\theta_{k-\tau} - \theta^{*}\|^{2(n-1)}(2\alpha\tau L(\|\theta_{k-\tau} - \theta^{*}\| + 1))\Big)
$$
  
\n
$$
\leq 2^{2n-1}n\alpha L\Big(4\alpha\tau L\|\theta_{k-\tau} - \theta^{*}\|^{2n-1} + 2\alpha\tau L\|\theta_{k-\tau} - \theta^{*}\|^{2(n-1)} + c_{n-1}\alpha^{2n-1}\tau^{2n-1}L^{2n-1}\Big)
$$
  
\n
$$
\leq 2^{2n-1}n\alpha L\Big(2\alpha\tau L\|\theta_{k-\tau} - \theta^{*}\|^{2n} + 4\alpha\tau L\|\theta_{k-\tau} - \theta^{*}\|^{2(n-1)} + c_{n-1}\alpha^{2n-1}\tau^{2n-1}L^{2n-1}\Big).
$$

Combining the analyses of the two terms, we get the following upper bound to [\(B.17\)](#page-21-4)

$$
\mathbb{E}[|(B.17)||\mathcal{F}_{k-\tau}]
$$
\n
$$
\leq 4^{n} n \alpha L \Big( 5 \alpha \tau L ||\theta_{k-\tau} - \theta^{*}||^{2n} + \alpha \tau L ||\theta_{k-\tau} - \theta^{*}||^{2(n-1)} + c'_{n} \alpha^{2n-1} \tau^{2n-1} L^{2n-1} \Big)
$$
\n
$$
+ 2^{2n-1} n \alpha L \Big( 2 \alpha \tau L ||\theta_{k-\tau} - \theta^{*}||^{2n} + 4 \alpha \tau L ||\theta_{k-\tau} - \theta^{*}||^{2(n-1)} + c_{n-1} \alpha^{2n-1} \tau^{2n-1} L^{2n-1} \Big)
$$
\n
$$
= 2^{2n-1} n \alpha L \Big( 12 \alpha \tau L ||\theta_{k-\tau} - \theta^{*}||^{2n} + 6 \alpha \tau L ||\theta_{k-\tau} - \theta^{*}||^{2(n-1)} + c''_{n-1} \alpha^{2n-1} \tau^{2n-1} L^{2n-1} \Big).
$$

Lastly, we analyze [\(B.18\)](#page-21-1). We first make use of the mean-value theorem, with  $a \in [0,1]$ , we have

$$
\|\theta_k - \theta^*\|^{2(n-1)} - \|\theta_{k-\tau} - \theta^*\|^{2(n-1)}
$$
  
=  $\|\theta_k - \theta_{k-\tau}\| \cdot 2(n-1) \|a(\theta_k - \theta^*) + (1-a)(\theta_{k-\tau} - \theta^*)\|^{2n-3}$   
=  $\|\theta_k - \theta_{k-\tau}\| \cdot 2(n-1) \|a(\theta_k - \theta_{k-\tau}) + \theta_{k-\tau} - \theta^*\|^{2n-3}$   
 $\leq 2^{2n-3}(n-1) \|\theta_k - \theta_{k-\tau}\| \left( \|\theta_k - \theta_{k-\tau}\|^{2n-3} + \|\theta_{k-\tau} - \theta^*\|^{2n-3} \right)$ 

Substituting the above upper bound back into  $(B.18)$ , we obtain

$$
\mathbb{E}[|(B.18)||\mathcal{F}_{k-\tau}]
$$
  
\n
$$
\leq 2^{2n-1}n(n-1)\alpha L \|\theta_{k-\tau} - \theta^*\|\mathbb{E}\Big[ (\|\theta_k - \theta^*\| + 1) \|\theta_k - \theta_{k-\tau}\| \Big( \|\theta_k - \theta_{k-\tau}\|^{2n-3} + \|\theta_{k-\tau} - \theta^*\|^{2n-3} \Big) |\mathcal{F}_{k-\tau}\Big]
$$

$$
\leq 2^{2n-1}n(n-1)\alpha L\Big(\|\theta_{k-\tau}-\theta^*\|E[\|\theta_k-\theta_{k-\tau}\|^{2n-1}|\mathcal{F}_{k-\tau}] + \|\theta_{k-\tau}-\theta^*\|^2\mathbb{E}[\|\theta_k-\theta_{k-\tau}\|^{2n-2}|\mathcal{F}_{k-\tau}] \n+ \|\theta_{k-\tau}-\theta^*\|\mathbb{E}[\|\theta_k-\theta_{k-\tau}\|^{2n-2}|\mathcal{F}_{k-\tau}] + \|\theta_{k-\tau}-\theta^*\|^{2n-2}\mathbb{E}[\|\theta_k-\theta_{k-\tau}\|^2|\mathcal{F}_{k-\tau}] \n+ \|\theta_{k-\tau}-\theta^*\|^{2n-1}\mathbb{E}[\|\theta_k-\theta_{k-\tau}\||\mathcal{F}_{k-\tau}] + \|\theta_{k-\tau}-\theta^*\|^{2n-2}\mathbb{E}[\|\theta_k-\theta_{k-\tau}\||\mathcal{F}_{k-\tau}]\Big) \n\leq 2^{2n-1}n(n-1)\alpha L\Big(c_n\alpha\tau L\|\theta_{k-\tau}-\theta^*\|^{2n}+c_{n-1}\alpha\tau L\|\theta_{k-\tau}-\theta^*\|^{2(n-1)}+c_{n-1}\alpha^{2n-1}\tau^{2n-1}L^{2n-1}\Big).
$$

Combining the analyses above, we have the following bound for  $T_1$ ,

$$
\mathbb{E}[|T_1||\mathcal{F}_{k-\tau}]
$$
\n
$$
\leq \mathbb{E}[||(B.14)||\mathcal{F}_{k-\tau}] + \mathbb{E}[||(B.15)||\mathcal{F}_{k-\tau}] + \mathbb{E}[||(B.16)||\mathcal{F}_{k-\tau}]
$$
\n
$$
+ \mathbb{E}[||(B.17)||\mathcal{F}_{k-\tau}] + \mathbb{E}[||(B.18)||\mathcal{F}_{k-\tau}]
$$
\n
$$
\leq c_{n,1}\alpha^2 \tau L^2 ||\theta_{k-\tau} - \theta^*||^{2n} + c_{n,2}\alpha^2 \tau L^2 ||\theta_{k-\tau} - \theta^*||^{2(n-1)} + c_{n,3}\alpha^{2n} \tau^{2n-1} L^{2n},
$$

where  $c_{n,1}$ ,  $c_{n,2}$  and  $c_{n,3}$  are some constants that depend on n.

Additionally, we note that

$$
\|\theta_{k-\tau} - \theta^*\|^{2n} = \mathbb{E}[\|\theta_{k-\tau} - \theta^*\|^{2n} |\mathcal{F}_{k-\tau}]
$$
  
\n
$$
\leq \mathbb{E}\left[\left(\|\theta_k - \theta_{k-\tau}\| + \|\theta_k - \theta^*\|\right)^{2n} |\mathcal{F}_{k-\tau}\right]
$$
  
\n
$$
\leq 2^{2n-1} \mathbb{E}[\|\theta_k - \theta_{k-\tau}\|^{2n} |\mathcal{F}_{k-\tau}] + 2^{2n-1} \mathbb{E}[\|\theta_k - \theta^*\|^{2n} |\mathcal{F}_{k-\tau}]
$$
  
\n
$$
\leq c_n \alpha^{2n} \tau^{2n} L^{2n} (\|\theta_{k-\tau} - \theta^*\|^{2n} + 1) + 2^{2n-1} \mathbb{E}[\|\theta_k - \theta^*\|^{2n} |\mathcal{F}_{k-\tau}].
$$

Therefore, for sufficiently small  $\alpha \tau L < \mu / (c'_{n}L)$ , we have

$$
(1 - c'_n \alpha^{2n} \tau^{2n} L^{2n}) \|\theta_{k-\tau} - \theta^*\|^{2n} \le c''_n \mathbb{E}[\|\theta_k - \theta^*\|^{2n} |\mathcal{F}_{k-\tau}] + c_n \alpha^{2n} \tau^{2n} L^{2n}
$$
  
\n
$$
\Rightarrow \|\theta_{k-\tau} - \theta^*\|^{2n} \le 2c''_n \mathbb{E}[\|\theta_k - \theta^*\|^{2n} |\mathcal{F}_{k-\tau}] + 2c_n \alpha^{2n} \tau^{2n} L^{2n}.
$$

As such, for sufficiently small  $\alpha$ , we have

$$
\mathbb{E}[|T_1||\mathcal{F}_{k-\tau}] \leq c_{n,1}\alpha^2 \tau L^2 \Big(c_n \mathbb{E}[\|\theta_k - \theta^*\|^{2n} |\mathcal{F}_{k-\tau}] + c'_n \alpha^{2n} \tau^{2n} L^{2n}\Big) + c_{n,2}\alpha^2 \tau L^2 \Big(c_{n-1} \mathbb{E}[\|\theta_k - \theta^*\|^{2(n-1)} |\mathcal{F}_{k-\tau}] + c'_{n-1}\alpha^{2(n-1)} \tau^{2(n-1)} L^{2(n-1)}\Big) + c_{n,3}\alpha^{2n} \tau^{2n-1} L^{2n} = c_{n,1}\alpha^2 \tau L^2 \mathbb{E}[\|\theta_k - \theta^*\|^{2n} |\mathcal{F}_{k-\tau}] + c_{4,2}\alpha^2 \tau L^2 \mathbb{E}[\|\theta_k - \theta^*\|^{2(n-1)} |\mathcal{F}_{k-\tau}] + c_{4,3}\alpha^{2n} \tau^{2n-1} L^{2n}.
$$

Hence, up til this point, we have obtained

$$
\mathbb{E}[\|\theta_{k+1} - \theta^*\|^{2n} |\mathcal{F}_{k-\tau}]
$$
  
\n
$$
\leq (1 - 2n\alpha\mu)\mathbb{E}[\|\theta_k - \theta^*\|^{2n} |\mathcal{F}_{k-\tau}]
$$
  
\n
$$
+ c_{n,1}\alpha^2 \tau L^2 \mathbb{E}[\|\theta_k - \theta^*\|^{2n} |\mathcal{F}_{k-\tau}] + c_{n,2}\alpha^2 \tau L^2 \mathbb{E}[\|\theta_k - \theta^*\|^{2(n-1)} |\mathcal{F}_{k-\tau}] + c_{n,3}\alpha^{2n} \tau^{2n-1} L^{2n}
$$
  
\n
$$
\leq (1 - 2n\alpha(\mu - c'_{n,1}\alpha\tau L^2))\mathbb{E}[\|\theta_k - \theta^*\|^{2n} |\mathcal{F}_{k-\tau}] + c_{n,2}\alpha^2 \tau L^2 \mathbb{E}[\|\theta_k - \theta^*\|^{2(n-1)}] |\mathcal{F}_{k-\tau}] + c_{n,3}\alpha^{2n} \tau^{2n-1} L^{2n}.
$$

Following the induction hypothesis, when  $k$  is sufficiently large, we have

$$
\mathbb{E}[\|\theta_k - \theta^*\|^{2(n-1)} | \mathcal{F}_{k-\tau}] \le c_{n-1} \alpha^{n-1} \tau^{n-1} s(\theta_0, L, \mu).
$$

Substituting the above upper bound back into our analysis of the  $2n$ -th moment bound, we obtain

$$
\mathbb{E}[\|\theta_{k+1}-\theta^*\|^{2n}|\mathcal{F}_{k-\tau}] \leq (1-2n\alpha(\mu-c'_{n,1}\alpha\tau L^2))\mathbb{E}[\|\theta_k-\theta^*\|^{2n}|\mathcal{F}_{k-\tau}] + \alpha^{n+1}\tau^n L^2 c_{n,2} \cdot c_{n-1} s(\theta_0,L,\mu) + c_{n,3}\alpha^{2n}\tau^{2n-1}L^{2n}.
$$

Subsequently, if we set  $\alpha$  sufficiently small, such that

$$
\alpha \tau L^2 < c_n \cdot \mu,
$$

we obtain

$$
\mathbb{E}[\|\theta_{k+1}-\theta^*\|^{2n}|\mathcal{F}_{k-\tau}] \le (1-\alpha\mu)\mathbb{E}[\|\theta_k-\theta^*\|^{2n}|\mathcal{F}_{k-\tau}] + \alpha^{n+1}\tau^n c'_{n,2} \cdot s(\theta_0,L,\mu),
$$

where  $s(\theta_0, L, \mu)$  is some constant that may depend on the initialization  $\theta_0$  and the problem primitives  $\mu$ and L but is independent of  $\alpha$ .

Recursively, we get

$$
\mathbb{E}[\|\theta_k - \theta^*\|^{2n}] \le (1 - \alpha \mu)^{k - \tau} \mathbb{E}[\|\theta_{\tau} - \theta^*\|^{2n}] + \alpha^n \tau^n \cdot s(\theta_0, L, \mu).
$$

Lastly, we recall that

$$
\mathbb{E}[\|\theta_{\tau} - \theta^*\|^{2n}] \le 2^{2n-1} \mathbb{E}[\|\theta_{\tau} - \theta_0\|^{2n}] + 2^{2n-1} \mathbb{E}[\|\theta_0 - \theta^*\|^{2n}]
$$
  
\n
$$
\le c_{n,1} \alpha^{2n} \tau^{2n} L^{2n} (\mathbb{E}[\|\theta_0 - \theta^*\|^{2n}] + 1) + c_{n,2} \|\theta_0 - \theta^*\|^{2n}
$$
  
\n
$$
\le c_{n,1} \mathbb{E}[\|\theta_0 - \theta^*\|^{2n}] + c_{n,2} \alpha^{2n} \tau^{2n} L^{2n}.
$$

Substituting back, we obtain for sufficiently large  $k$ ,

$$
\mathbb{E}[\|\theta_k - \theta^*\|^{2n}] \le c_{n,1}(1 - \alpha \mu)^{k-\tau} \mathbb{E}[\|\theta_0 - \theta^*\|^{2n}] + \alpha^{2n} \tau^{2n} s(\theta_0, L, \mu).
$$

As such, we have proven the desired  $n$ -th moment bound.

#### B.2.1 Proof of Lemma [B.2](#page-21-5)

We now come back to Lemma [B.2](#page-21-5) and provide the complete proof.

Proof. The proof follows a similar strategy as  $(B.9)$  and  $(B.10)$  in Section [B.1.1.](#page-17-0)

We start with the following relaxation and obtain that

$$
\mathbb{E}[\|\theta_{k} - \theta_{k-\tau}\|^{2n} |\mathcal{F}_{k-\tau}] \leq \mathbb{E}\Big[\Big(\sum_{t=k-\tau}^{k-1} \|\theta_{t+1} - \theta_{t}\|\Big)^{2n} |\mathcal{F}_{k-\tau}\Big]
$$
\n
$$
\leq \tau^{2n-1} \sum_{t=k-\tau}^{k-1} \mathbb{E}[\|\theta_{t+1} - \theta_{t}\|^{2n} |\mathcal{F}_{k-\tau}]
$$
\n
$$
= \alpha^{2n} \tau^{2n-1} \sum_{t=k-\tau}^{k-1} \mathbb{E}[\|g(\theta_{t}, x_{t}) + \xi_{t+1}(\theta_{t})\|^{2n} |\mathcal{F}_{k-\tau}]
$$
\n
$$
\leq 2^{2n-1} \alpha^{2n} \tau^{2n-1} \sum_{t=k-\tau}^{k-1} \Big(\mathbb{E}[\|g(\theta_{t}, x_{t})\|^{2n} |\mathcal{F}_{k-\tau}] + \mathbb{E}[\|\xi_{t+1}(\theta_{t})\|^{2n} |\mathcal{F}_{k-\tau}]\Big)
$$
\n
$$
\leq 2^{2n-1} \alpha^{2n} \tau^{2n-1} \sum_{t=k-\tau}^{k-1} \Big(L_{1}^{2n} \mathbb{E}[(\|\theta_{t} - \theta^{*}\| + 1)^{2n} |\mathcal{F}_{k-\tau}] + L_{2}^{2n} (\mathbb{E}[\|\theta_{t} - \theta^{*}\| |\mathcal{F}_{k-\tau}] + 1)^{2n}\Big)
$$
\n
$$
\leq 4^{2n-1} \alpha^{2n} \tau^{2n-1} L^{2n} \sum_{t=k-\tau}^{k-1} \mathbb{E}[\|\theta_{t} - \theta^{*}\|^{2n} |\mathcal{F}_{k-\tau}] + 4^{2n-1} \alpha^{2n} \tau^{2n} L^{2n}.
$$

Next, in order to obtain a bound on  $\|\theta_t - \theta^*\|^{2n}$ , we study the following term.

$$
\mathbb{E}[\|\theta_{t+1} - \theta^*\|^{2n} |\mathcal{F}_{k-\tau}] \leq \mathbb{E}\Big[\Big(\|\theta_{t+1} - \theta_t\| + \|\theta_t - \theta^*\|\Big)^{2n} |\mathcal{F}_{k-\tau}\Big]
$$
  
\n
$$
= \sum_{i=0}^{2n} {2n \choose i} \mathbb{E}[\|\theta_{t+1} - \theta_t\|^i \|\theta_t - \theta^*\|^{2n-i} |\mathcal{F}_{k-\tau}]
$$
  
\n
$$
= \mathbb{E}[\|\theta_t - \theta^*\|^{2n} |\mathcal{F}_{k-\tau}] + \sum_{i=1}^{2n} \alpha^i \mathbb{E}[\|g(\theta_t, x_t) + \xi_{t+1}(\theta_t)\|^i \|\theta_t - \theta^*\|^{2n-i} |\mathcal{F}_{k-\tau}]
$$

Note that

$$
\mathbb{E}[\|g(\theta_t, x_t) + \xi_{t+1}(\theta_t)\|^i \|\theta_t - \theta^*\|^{2n-i} |\mathcal{F}_{k-\tau}]
$$
\n
$$
\leq 2^{i-1} \mathbb{E}\Big[\Big(\|g(\theta_t, x_t)\|^i + \|\xi_{t+1}(\theta_t)\|^i\Big)\|\theta_t - \theta^*\|^{2n-i} |\mathcal{F}_{k-\tau}\Big]
$$
\n
$$
= 2^{i-1} \mathbb{E}\Big[\mathbb{E}[(\|g(\theta_t, x_t)\|^i + \|\xi_{t+1}(\theta_t)\|^i)|\theta_t]\|\theta_t - \theta^*\|^{2n-i} |\mathcal{F}_{k-\tau}\Big]
$$
\n
$$
\leq 2^{i-1} L^i \mathbb{E}\Big[(\|\theta_t - \theta^*\| + 1)^{2n} |\mathcal{F}_{k-\tau}\Big]
$$
\n
$$
\leq 2^{2(n-1)} 2^i L^i \Big(\mathbb{E}[\|\theta_t - \theta^*\| |\mathcal{F}_{k-\tau}| + 1\Big)
$$

Substituting back, we obtain

$$
\sum_{i=1}^{2n} \alpha^{i} \mathbb{E}[\|g(\theta_t, x_t) + \xi_{t+1}(\theta_t)\|^{i} \|\theta_t - \theta^*\|^{2n-i} |\mathcal{F}_{k-\tau}]
$$
\n
$$
\leq 2^{2(n-1)} \sum_{i=1}^{2n} 2^{i} \alpha^{i} L^{i} \Big( \mathbb{E}[\|\theta_t - \theta^*\| |\mathcal{F}_{k-\tau}] + 1 \Big)
$$
\n
$$
= 2^{2(n-1)} \Big( \mathbb{E}[\|\theta_t - \theta^*\| |\mathcal{F}_{k-\tau}] + 1 \Big) \cdot 2\alpha L (1 + 2\alpha L)^{2n-1}
$$
\n
$$
\leq 4^{2n-1} \alpha L \Big( \mathbb{E}[\|\theta_t - \theta^*\| |\mathcal{F}_{k-\tau}] + 1 \Big)
$$

Consolidating the terms, we have

$$
\mathbb{E}[\|\theta_{t+1} - \theta^*\|^{2n} | \mathcal{F}_{k-\tau}] \le (1 + 4^{2n-1} \alpha L) \Big( \mathbb{E}[\|\theta_t - \theta^*\| | \mathcal{F}_{k-\tau}] + 1 \Big)
$$

Recursively, for  $0\leq l\leq \tau,$  we have

$$
\mathbb{E}[\|\theta_{k-\tau+l} - \theta^*\|^{2n} |\mathcal{F}_{k-\tau}] \le (1 + 4^{2n-1} \alpha L)^l \|\theta_{k-\tau} - \theta^*\|^{2n} + 4^{2n-1} \alpha L \sum_{i=0}^{l-1} (1 + 4^{2n-1} \alpha L)^i
$$

$$
= (1 + 4^{2n-1} \alpha L)^l \|\theta_{k-\tau} - \theta^*\|^{2n} + (1 + 4^{2n-1} \alpha L)^l
$$

Then, for

$$
4^{2n-1}\alpha\tau L \le \mu/4L < 1/4,
$$

we have for  $k - \tau \leq t \leq k$ ,

$$
\mathbb{E}[\|\theta_t - \theta^*\|^{2n} |\mathcal{F}_{k-\tau}] \le (1 + 2^{4n-1} \alpha \tau L) \|\theta_{k-\tau} - \theta^*\|^{2n} + 2^{4n-1} \alpha \tau L
$$
  

$$
\le 2 \|\theta_{k-\tau} - \theta^*\|^{2n} + 2^{4n-1} \alpha \tau L
$$

Finally, we have

$$
\mathbb{E}[\|\theta_k - \theta_{k-\tau}\|^{2n} |\mathcal{F}_{k-\tau}] \le 4^{2n-1} \alpha^{2n} \tau^{2n-1} L^{2n} \sum_{t=k-\tau}^{k-1} \mathbb{E}[\|\theta_t - \theta^*\|^{2n} |\mathcal{F}_{k-\tau}] + 4^{2n-1} \alpha^{2n} \tau^{2n} L^{2n}
$$
  

$$
\le 4^{2n-1} \alpha^{2n} \tau^{2n} L^{2n} \left( 2\|\theta_{k-\tau} - \theta^*\|^{2n} + 2^{4n-1} \alpha \tau L + 1 \right)
$$
  

$$
\le 2^{4n-1} \alpha^{2n} \tau^{2n} L^{2n} \left( \|\theta_{k-\tau} - \theta^*\|^{2n} + 1 \right).
$$

As such, we have completed the proof.

 $\Box$ 

### <span id="page-26-0"></span>C Proof of Theorem [4.1](#page-5-0)

In this section, we prove the weak convergence result in Theorem [4.1.](#page-5-0) In fact, the proof of the projected SA weak convergence result can be seen as a special case of unprojected SA with the asymptotic linearity condition, which we have briefly discussed in Section [4.](#page-4-2) Therefore, the proof proceeds in the following two subsections. First, we formally define the asymptotic linearity condition and present our weak convergence result for unprojected SA under this additional assumption. Next, we relate this result for unprojected SA to projected SA and specialize the proof to obtain Theorem [4.1.](#page-5-0)

### C.1 Asymptotic Linearity

In this subsection, we formally introduce the asymptotic linearity condition, which is crucial for establishing weak convergence in the context of unprojected SA  $(\beta = \infty)$ . Additionally, we explore the implications of this condition.

<span id="page-26-1"></span>**Assumption 6** (Asymptotic Linearity). The noise sequence  $(\xi_k)_{k\geq 1}$  is a collection of i.i.d. random fields satisfying the following conditions: (1)  $\mathbb{E}[\xi_{k+1}(\theta)|\mathcal{F}_k] = 0$ , (2) there exists a constant  $L_3 > 0$  such that  $\xi_1$  is  $L_3\text{-}Lipschitz, i.e., }||\xi_1(\theta) - \xi_1(\theta')|| \leq L_3||\theta - \theta'||$ , for all  $\theta, \theta' \in \mathbb{R}^d$ , and  $(3) ||\xi_1(0)|| \leq L_3$ . Moreover, there exists a function  $G(\cdot): \mathcal{X} \to \mathbb{R}^{d \times d}$  such that given  $\epsilon > 0$ , define

$$
\delta(\epsilon) := \min \big\{ \delta : ||g'(\theta, x) - G(x)|| \le \epsilon, \forall x \in \mathcal{X} \quad and \quad \forall \theta \in \{\theta : ||\theta|| \ge \delta\} \big\},\
$$

and we have  $\lim_{\epsilon \to 0} \epsilon \delta(\epsilon) = 0$ .

The first part of Assumption [6](#page-26-1) states that the random field grows at most linearly in  $\theta$ . The second part of Assumption [6](#page-26-1) implies that  $g'(\theta, x)$  converges to a limit  $G(x)$  when  $\|\theta\| \to \infty$  for all  $x \in \mathcal{X}$ , which shows the asymptotic linearity of  $g(\theta, x)$ . Furthermore, Assumption [6](#page-26-1) also requires how fast  $g'(\theta, x)$  converges to  $G(x)$ . A sufficient condition under which the second part of Assumption [6](#page-26-1) holds is that there exists  $\omega > 0$  such that  $\|\theta\|^{1+\omega} \|g'(\theta,x)-G(x)\| < \infty$  for  $\forall \theta \in \mathbb{R}^d$  and  $x \in \mathcal{X}$ . We can verify that to ensure  $\|g'(\theta,x)-G(x)\| < \epsilon$ , we can set  $\|\theta\| \in \Theta(\epsilon^{-\frac{1}{1+\omega}})$ , which can ensure  $\epsilon\delta(\epsilon) \in \mathcal{O}(\epsilon^{\frac{\omega}{1+\omega}}) \to 0$  as  $\epsilon \to 0$ . This sufficient condition implies that  $g'(\theta, x)$  uniformly converge to  $G(x)$  with convergence rate of  $\mathcal{O}(\|\theta\|^{-(1+w)})$ . By definition, we conclude that the structure of linear SA is also asymptotic linear. Besides that, the 1-dimensional logistic regression also satisfies Assumption [6.](#page-26-1) For 1-dimensional logistic regression, we have  $g(\theta, x, y) = x\left(\frac{1}{1+e^{-\theta x}} - y\right) + \lambda \theta$ , where  $(x, y)$  presents the data. Therefore, we have  $g'(\theta, x, y) = \frac{x^2 e^{-\theta x}}{(1 + e^{-\theta x})^2}$  $\frac{x^2e^{-\sigma x}}{(1+e^{-\theta x})^2} + \lambda$  and  $g'(\theta, x, y)$  uniformly converges to  $\lambda$  with geometric convergence rate, thereby satisfies the Assumption [6.](#page-26-1)

#### C.2 Proof Under Assumption [6](#page-26-1)

With the asymptotic linearity condition now formally defined, we proceed to prove the weak convergence for unprojected SA. For convenient reference, we state the theorem below.

<span id="page-26-3"></span>**Theorem C.1** (Ergodicity of SA–Asymptotic Linearity). Suppose that Assumption [1–](#page-2-0)Assumption [4](#page-3-4) hold. Additionally, assume [6.](#page-26-1) For stepsize  $\alpha > 0$  that satisfies the constraint  $\alpha \tau_{\alpha} L^2 < \min(c_2 \mu, \kappa_{\mu})$ , with  $c_2$ formalized in Proposition [4.2](#page-5-1) and  $\kappa_{\mu}$  defined in [\(C.1\)](#page-26-2), the Markov chain  $(x_k, \theta_k)_{k\geq 0}$  converges to a unique stationary distribution  $\bar{\nu}_{\alpha} \in \mathcal{P}_2(\mathcal{X} \times \mathbb{R}^d)$ .

Moreover, there exist  $\kappa_{\mu} > 0$  and some universal constant c' such that

<span id="page-26-2"></span>
$$
\epsilon \delta(\epsilon) \le c'\mu, \quad \forall \epsilon \le \kappa_{\mu}.\tag{C.1}
$$

We let  $\nu_\alpha := \mathcal{L}(\theta_\infty)$  be the second marginal of  $\bar{\nu}_\alpha$ . For  $k \geq 2\tau_\alpha$ , it holds that

<span id="page-26-4"></span>
$$
W_2(\mathcal{L}(\theta_k), \nu_\alpha) \le \bar{W}_2(\mathcal{L}(x_k, \theta_k), \bar{\nu}_\alpha) \le (1 - \alpha \mu)^{k/2} \cdot s(\theta_0, L, \mu). \tag{C.2}
$$

The proof of Theorem [C.1](#page-26-3) consists of two major steps. Firstly, we assume that  $x_0 \sim \pi$ , and show that  $(x_k, \theta_k)_{k>0}$  converges to a unique limiting invariant distribution. Next, we relax the assumption of  $x_0 \sim \pi$ , and prove that for arbitrary initialization  $(x_0, \theta_0) \in \mathcal{X} \times \mathbb{R}^d$ , the Markov chain will converge to the same limit.

<span id="page-27-0"></span>Step 1: Initialization with  $x_0 \sim \pi$ . To prove the convergence of the Markov chain, we consider the following coupling construction. We have a pair of Markov chains  $(x_k, \theta_k^{[1]})_{k \geq 0}$  and  $(x_k, \theta_k^{[2]})_{k \geq 0}$  sharing the same underlying process and noise, i.e.,  $(x_k, \xi_{k+1})_{k>0}$ , i.e.,

<span id="page-27-3"></span>
$$
\theta_{k+1}^{[1]} = \theta_k^{[1]} + \alpha(g(\theta_k^{[1]}, x_k) + \xi_{k+1}(\theta_k^{[1]})),
$$
  
\n
$$
\theta_{k+1}^{[2]} = \theta_k^{[2]} + \alpha(g(\theta_k^{[2]}, x_k) + \xi_{k+1}(\theta_k^{[2]})).
$$
\n(C.3)

We assume that the initial iterates  $\theta_0^{[1]}$  and  $\theta_0^{[2]}$  may depend on each other and on  $x_0$ , but are independent of subsequent  $(x_k)_{k\geq 1}$  given  $x_0$ . For the iterates difference  $\theta_k^{[1]} - \theta_k^{[2]}$  $k^{[2]}$ , we have the following Proposition [C.2,](#page-27-1) whose proof is given at the end of this subsection.

<span id="page-27-1"></span>**Proposition C.2.**  $\forall k \geq \tau$  and  $\alpha \tau \leq \min(\frac{\mu}{908L^2}, \frac{\kappa_{\mu}}{L^2})$ , we have

$$
\mathbb{E}[\|\theta_k^{[1]} - \theta_k^{[2]}\|^2] \le 4(1 - \mu\alpha)^{k - \tau} \mathbb{E}[\|\theta_0^{[1]} - \theta_0^{[2]}\|^2],
$$

where  $\kappa_{\mu} > 0$  and  $\epsilon \delta(\epsilon) \leq \frac{\mu}{768}, \forall \epsilon \leq \kappa_{\mu}$ .

By Proposition [C.2](#page-27-1) and the definition of  $W_2$  and  $\bar{W}_2$ , we have

<span id="page-27-2"></span>
$$
W_2^2 \left( \mathcal{L}(\theta_k^{[1]}), \mathcal{L}(\theta_k^{[2]}) \right) \stackrel{\text{(i)}}{\leq} \bar{W}_2^2 \left( \mathcal{L}(x_k, \theta_k^{[1]}), \mathcal{L}(x_k, \theta_k^{[2]}) \right)
$$
  
\n
$$
\stackrel{\text{(ii)}}{\leq} \mathbb{E}[\|\theta_k^{[1]} - \theta_k^{[2]}\|^2]
$$
  
\n
$$
\stackrel{\text{(iii)}}{\leq} 4(1 - \mu\alpha)^{k-\tau} \mathbb{E}[\|\theta_0^{[1]} - \theta_0^{[2]}\|^2],
$$
\n(C.4)

where (i) and (ii) hold by the definition of  $W_2$  and  $\overline{W}_2$  and (iii) holds by applying Proposition [C.2.](#page-27-1)

Note that equation [\(C.4\)](#page-27-2) always holds for any joint distribution of initial iterates  $(x_0, \theta_0^{[1]}, \theta_0^{[2]})$ . Recall that  $P^*$  represents the transition kernel for the time-reversed Markov chain of  $\{x_k\}_{k\geq 0}$ , and the initial distribution of  $x_0$  is assumed to be mixed already. Given a specific  $x_0$ , we sample  $x_{-1}$  from  $P^*(\cdot | x_0)$ . Additionally, we use  $\theta_{-1}^{[2]}$  to denote the random varible that satisfies  $\theta_{-1}^{[2]} \stackrel{d}{=} \theta_0^{[1]}$  and is independent of  $\{x_k\}_{k\geq 0}$ . Finally, we set  $\theta_0^{[2]}$  as

$$
\theta_0^{[2]} = \theta_{-1}^{[2]} + \alpha(g(x_{-1}, \theta_{-1}^{[2]}) + \xi_0(\theta_{-1}^{[2]})).
$$

By the property of time-reversed Markov chain, we have  $\{x_k\}_{k\geq -1} \stackrel{d}{=} \{x_k\}_{k\geq 0}$ . Given that  $\theta_{-1}^{[2]} \stackrel{d}{=} \theta_0^{[1]}$  and  $\theta_{-1}^{[2]}$  is independent with  $\{x_k\}_{k\geq -1}$ , we can prove  $(x_k, \theta_k^{[2]}) \stackrel{d}{=} (x_{k+1}, \theta_{k+1}^{[1]})$  for  $k \geq 0$ . We thus have for all  $k \geq \tau$ :

$$
\bar{W}_2^2\left(\mathcal{L}\left(x_k,\theta_k^{[1]}\right),\mathcal{L}\left(x_{k+1},\theta_{k+1}^{[1]}\right)\right)=\bar{W}_2^2\left(\mathcal{L}\left(x_k,\theta_k^{[1]}\right),\mathcal{L}\left(x_k,\theta_k^{[2]}\right)\right)
$$
\n
$$
\stackrel{\text{(i)}}{\leq} 4(1-\mu\alpha)^{k-\tau}\mathbb{E}[\|\theta_0^{[1]}-\theta_0^{[2]}\|^2],
$$

where (i) holds by inequality  $(C.4)$ . Then, we have

$$
\sum_{k=0}^{\infty} \overline{W}_2^2 \left( \mathcal{L} \left( x_k, \theta_k^{[1]} \right), \mathcal{L} \left( x_{k+1}, \theta_{k+1}^{[1]} \right) \right)
$$
\n
$$
\leq \sum_{k=0}^{t_{\alpha}-1} \overline{W}_2^2 \left( \mathcal{L} \left( x_k, \theta_k^{[1]} \right), \mathcal{L} \left( x_{k+1}, \theta_{k+1}^{[1]} \right) \right) + 4 \mathbb{E}[\|\theta_0^{[1]} - \theta_0^{[2]}\|^2] \sum_{k=0}^{\infty} (1 - \mu \alpha)^k
$$
\n
$$
< \infty.
$$

Consequently,  $\{\mathcal{L}(x_k, \theta_k^{[1]})\}_{k\geq 0}$  forms a Cauchy sequence w.r.t. the metric  $\bar{W}_2$ . Since the space  $\mathcal{P}_2(\mathcal{X}\times\mathbb{R}^d)$ endowed with  $\overline{W}_2$  is a Polish space, every Cauchy sequence converges [\[Vil09,](#page-12-14) Theorem 6.18]. Furthermore, <span id="page-28-0"></span>convergence in Wasserstein 2-distance also implies weak convergence [\[Vil09](#page-12-14), Theorem 6.9]. Therefore, we conclude that the sequence  $(\mathcal{L}(x_k, \theta_k^{[1]}))_{k \geq 0}$  converges weakly to a limit distribution  $\bar{\mu} \in \mathcal{P}_2(\mathcal{X} \times \mathbb{R}^d)$ .

Now that we have established the existence of a limiting distribution, we next proceed to show the uniqueness. We prove this by contradiction. Note that we currently assume that  $x_0 \sim \pi$ , hence to show that the limit  $(x_{\infty}, \theta_{\infty})$  is unique, we only need to show that the limit is independent of the initial distribution of  $\theta_0$ , which can be correlated to  $x_0$ .

Consider two Markov chains  $(x_k, \theta_k)_{k\geq 0}$  and  $(x_k, \theta'_k)_{k\geq 0}$ , sharing  $(x_k, \xi_{k+1})_{k\geq 0}$  but with arbitrary initialization of  $\theta_0$  and  $\theta'_0$ . For the sake of contradiction, we assume that  $(x_0, \theta_0) \Rightarrow (x_\infty, \theta_\infty)$  and  $(x_0, \theta'_0) \Rightarrow$  $(x_{\infty}, \theta'_{\infty})$  respectively. Then, by the triangle inequality, we have that

$$
\begin{split} &\bar{W}_2\Big((x_\infty,\theta_\infty),(x_\infty,\theta'_\infty)\Big) \\ &\leq \bar{W}_2\Big((x_\infty,\theta_\infty),(x_k,\theta_k)\Big)+\bar{W}_2\Big((x_k,\theta_k),(x_k,\theta'_k)\Big)+\bar{W}_2\Big((x_k,\theta'_k),(x_\infty,\theta'_\infty)\Big) \\ &\to 0. \end{split}
$$

As such, we have shown that the limit  $\bar{\nu}$  is unique.

Lastly, we prove that  $\bar{\nu}$  is invariant. Suppose that we initialize the joint process at its limit, i.e.,  $(x_0, \theta_0) \sim$  $\bar{\nu}$ . We first apply the triangle inequality, and we obtain

$$
\bar{W}_2((x_1,\theta_1),(x_0,\theta_0)) \leq \bar{W}_2((x_1,\theta_1),(x_{k+1},\theta_{k+1})) + \bar{W}_2((x_{k+1},\theta_{k+1}),(x_0,\theta_0)).
$$

Clearly, as  $k \to \infty$ ,  $\overline{W}_2((x_{k+1},\theta_{k+1}),(x_0,\theta_0)) \to 0$ . To bound  $\overline{W}_2((x_1,\theta_1),(x_{k+1},\theta_{k+1}))$ , we need the following lemma.

<span id="page-28-1"></span>**Lemma C.3.** Consider two copies of the SA trajectory, where  $\mathcal{L}(x_0, \theta_0) = \bar{\nu}$  and  $\mathcal{L}(x'_0, \theta'_0)$  is allowed to be arbitrary.

$$
\bar{W}_2^2(\mathcal{L}(x_1,\theta_1),\mathcal{L}(x_1',\theta_1')\Big)\leq \rho_1\cdot \bar{W}_2^2\Big(\mathcal{L}(x_0,\theta_0),\mathcal{L}(x_0',\theta_0')\Big)+\rho_2\cdot \sqrt{\bar{W}_2^2\Big(\mathcal{L}(x_0,\theta_0),\mathcal{L}(x_0',\theta_0')\Big)},
$$

where

$$
\rho_1 := 1 + 2(1 + \alpha L)^2 + 16\alpha^2 L^2 < \infty \quad \text{and} \quad \rho_2 := 16\alpha^2 L^2 \sqrt{\mathbb{E}[\|\theta_0\|^4]} < \infty
$$

are independent of  $\mathcal{L}(x'_0, \theta'_0)$ .

*Proof.* Consider the following coupling between the two processes  $(x_k, \theta_0)_{k \geq 0}$  and  $(x'_k, \theta'_k)_{k \geq 0}$ 

$$
\bar{W}_2^2(\mathcal{L}(x_0, \theta_0), \mathcal{L}(x'_0, \theta'_0)) = \mathbb{E}\Big[d_0(x_0, x'_0) + ||\theta_0 - \theta'_0||^2\Big] \text{ and}
$$
  

$$
x_{k+1} = x'_{k+1} \text{ if } x_k = x'_k, \quad \forall k \ge 0.
$$

Then, it is clear that

$$
\overline{W}_2^2(\mathcal{L}(x_1,\theta_1),\mathcal{L}(x_1',\theta_1')) \leq \mathbb{E}\Big[d_0(x_1,x_1') + \|\theta_1 - \theta_1'\|^2\Big].
$$

Recall the metric  $d_0(x, x') = \mathbb{1}\{x \neq x'\}$  and hence, we have

$$
g(\theta_0, x_0) = g(\theta_0, x'_0) + d_0(x'_0, x_0)(g(\theta_0, x_0) - g(\theta_0, x'_0)).
$$

Therefore, it is easy to see that

$$
\theta_1 - \theta_1' = \theta_0 - \theta_0' + \alpha(g(\theta_0, x_0) - g(\theta_0', x_0')) + \alpha(\xi_1(\theta_0) - \xi_1(\theta_0'))
$$
  
=  $\theta_0 - \theta_0' + \alpha(g(\theta_0, x_0) \mp g(\theta_0, x_0') - g(\theta_0', x_0')) + \alpha(\xi_1(\theta_0) - \xi_1(\theta_0'))$   
=  $\theta_0 - \theta_0' + \alpha(g(\theta_0, x_0') - g(\theta_0', x_0')) + \alpha(\xi_1(\theta_0) - \xi_1(\theta_0'))$   
+  $\alpha d_0(x_0', x_0)(g(\theta_0, x_0) - g(\theta_0, x_0'))$ ,

whence

$$
\|\theta_1 - \theta'_1\| \le (1 + \alpha L) \|\theta_0 - \theta'_0\| + \alpha d_0(x'_0, x_0) \|g(\theta_0, x_0) - g(\theta_0, x'_0)\|
$$
  

$$
\le (1 + \alpha L) \|\theta_0 - \theta'_0\| + \alpha d_0(x'_0, x_0) \cdot 2L(\|\theta_0\| + 1).
$$

As such, we see that

$$
\mathbb{E}[d_0(x_1, x_1') + ||\theta_1 - \theta_1'||^2] \leq \mathbb{E}[d_0(x_0, x_0')] + 2(1 + \alpha L)^2 \cdot \mathbb{E}[\|\theta_0 - \theta_0'\|^2] + 16\alpha^2 L^2 \cdot \mathbb{E}[d_0(x_0', x_0)(\|\theta_0\|^2 + 1)].
$$

Next, we make use of Cauchy-Schwarz inequality and obtain

$$
\mathbb{E}[d_0(x'_0, x_0) \cdot ||\theta_0||^2] \le \sqrt{\mathbb{E}[d_0(x'_0, x_0)]} \sqrt{\mathbb{E}_{\theta_0 \sim \mu}[\|\theta_0\|^4]}.
$$

Because Assumption [6](#page-26-1) implies Assumption  $4(p = 2)$  $4(p = 2)$ , by Proposition [4.2](#page-5-1) and Fatou's lemma, we have

$$
\mathbb{E}[\|\theta_{\infty}-\theta^*\|^4] \leq \lim\inf_{k\to\infty}\mathbb{E}[\|\theta_k-\theta^*\|_{\infty}^4] < \infty,
$$

which implies  $\mathbb{E}[\|\theta_{\infty}\|^4]<\infty.$  Hence, the desired inequality follows through.

By Lemma [C.3,](#page-28-1) we can set  $\mathcal{L}(x'_0, \theta'_0) = \mathcal{L}(x_k, \theta_k)$ , then

$$
\bar{W}_2^2\left(\mathcal{L}\left(x_1,\theta_1\right),\mathcal{L}(x_{k+1},\theta_{k+1})\right) \leq \rho_1 \bar{W}_2^2\left(\bar{\mu},\mathcal{L}(x_k,\theta_k)\right) + \rho_2 \sqrt{\bar{W}_2^2\left(\bar{\nu},\mathcal{L}(x_k,\theta_k)\right)}.
$$

Therefore,  $\bar{W}_2^2(\mathcal{L}(x_1,\theta_1),\mathcal{L}(x_{k+1},\theta_{k+1})) \to 0$  as  $k \to 0$ , which implies  $\bar{W}_2((x_1,\theta_1),(x_0,\theta_0)) = 0$ . As such, we have proved the joint sequence  $(x_k, \theta_k)_{k\geq 0}$  converges weakly to the unique invariant distribution  $\bar{\nu} \in \mathcal{P}_2(\mathcal{X} \times \mathbb{R}^d)$ . As a result,  $\{\theta_k\}_{k \geq 0}$  converges weakly to  $\mu \in \mathcal{P}_2(\mathbb{R}^d)$ , where  $\mu$  is the second marginal of  $\bar{\mu}$ over  $\mathbb{R}^d$ .

Lastly, before proceeding to the next step, in which we remove the assumption  $x_0 \sim \pi$ , we first derive the convergence rate of  $\{\theta_k\}_{k>0}$  under  $x_0 \sim \pi$  as presented in the following lemma. This lemma will help us to establish the convergence rate without  $x_0 \sim \pi$ .

<span id="page-29-0"></span>**Lemma C.4.** Under  $x_0 \sim \pi$ , Assumption [1](#page-2-0)[–4](#page-3-4) and [6](#page-26-1) and the same setting as Proposition [C.2,](#page-27-1)

$$
W_2^2\left(\mathcal{L}(\theta_k),\nu_\alpha\right) \leq \bar{W}_2^2\left(\mathcal{L}(x_k,\theta_k),\bar{\nu}_\alpha\right) \leq 16(1-\mu\alpha)^k\cdot \left(\mathbb{E}\left[\|\theta_0^{[1]}-\theta^*\|^2\right]+c_{2,2}'\right).
$$

*Proof.* Let us consider the coupled processes defined as equation  $(C.3)$ . Suppose that the initial iterate  $(x_0, \theta_0^{[2]})$  follows the stationary distribution  $\bar{\nu}$ , thus  $\mathcal{L}(x_k, \theta_k^{[2]}) = \bar{\nu}$  and  $\mathcal{L}(\theta_k^{[2]})$  $\binom{[2]}{k} = \nu$  for all  $k \geq 0$ . By equation [\(C.4\)](#page-27-2), we have for all  $k \geq \tau$ :

$$
W_2^2 \left( \mathcal{L}(\theta_k^{[1]}), \mu \right) = W_2^2 \left( \mathcal{L}(\theta_k^{[1]}), \mathcal{L}(\theta_k^{[2]}) \right)
$$
  
\n
$$
\leq \bar{W}_2^2 \left( \mathcal{L}(x_k, \theta_k^{[1]}), \mathcal{L}(x_k, \theta_k^{[2]}) \right)
$$
  
\n
$$
\leq 4(1 - \mu \alpha)^{k - \tau} \mathbb{E}[\|\theta_0^{[1]} - \theta_0^{[2]}\|^2]
$$
  
\n
$$
\leq 8(1 - \mu \alpha)^{k - \tau} \cdot \left( \mathbb{E} \left[ \|\theta_0^{[1]} - \theta^* \|^2 \right] + \mathbb{E} \left[ \|\theta_\infty - \theta^* \|^2 \right] \right)
$$
  
\n
$$
\leq 16(1 - \mu \alpha)^k \cdot \left( \mathbb{E} \left[ \|\theta_0^{[1]} - \theta^* \|^2 \right] + \mathbb{E} \left[ \|\theta_\infty - \theta^* \|^2 \right] \right),
$$
\n(C.5)

where we make use of the derivation in  $(B.13)$  to obtain the last inequality.

We note that

$$
\mathbb{E}[\|\theta_{\infty}-\theta^*\|^2] \leq \lim \inf_{k \to \infty} \mathbb{E}[\|\theta_k-\theta^*\|_{\infty}^2] \leq c_{2,2} \cdot \alpha \tau L^2/\mu \leq c_{2,2}',
$$

the last inequality holds for  $\alpha \tau \le \min(\frac{\mu}{908L^2}, \frac{\kappa_{\mu}}{L^2})$ . Therefore, we prove the desired inequality

$$
W_2^2\left(\mathcal{L}(\theta_k^{[1]}), \mu\right) \le 16(1-\mu\alpha)^k \cdot \left(\mathbb{E}\left[\|\theta_0^{[1]} - \theta^*\|^2\right] + c'_{2,2}\right).
$$

 $\Box$ 



Step 2: Arbitrary Initialization for  $(x_0, \theta_0)$ . In this step, we remove the assumption of  $x_0 \sim \pi$  needed in the previous step. We need the following lemma to prove our result.

<span id="page-30-0"></span>**Lemma C.5.** Consider two trajectories  $(x_k, \theta_k)_{k \geq 0}$  and  $(x'_k, \theta'_k)_{k \geq 0}$ . Suppose that  $\theta_0 = \theta'_0$ ,  $x'_0 \sim \pi$  and  $x_0$ is initialized from some arbitrary distribution that satisfies  $\|\mathcal{L}(x_0) - \pi\|_{TV} = \epsilon$ . Then for  $k \geq \tau$ , we have

$$
\overline{W}_2(\mathcal{L}(x_k,\theta_k),\mathcal{L}(x'_k,\theta'_k)) \le \epsilon \Big(4c_{2,1}(1-\alpha\mu)^k \mathbb{E}[\|\theta_0-\theta^*\|^2] + 4c_{s,2}\alpha\tau \cdot \frac{L^2}{\mu} + 1\Big)^{1/2}.
$$

*Proof.* We consider the following coupling between two joint processes  $(x_k, \theta_k)_{k\geq 0}$  and  $(x'_k, \theta'_k)_{k\geq 0}$ . We first apply the maximal coupling on  $x_0$  and  $x'_0$  such that

$$
\mathbb{P}(x_0 \neq x'_0) = \|\mathcal{L}(x_0) - \mathcal{L}(x'_0)\|_{\mathrm{TV}} = \epsilon.
$$

For the case  $x_0 = x'_0$ , we can couple the two Markov chains  $\{x_k\}_{k\geq 0}$  and  $\{x'_k\}_{k\geq 0}$  such that

$$
x_k \equiv x'_k, \forall k \ge 0.
$$

Under this coupling, we have  $\theta_k \equiv \theta'_k, \forall k \geq 0$ .

For the case  $x_0 \neq x'_0$ , we let the two processes  $(x_k, \theta_k)_{k\geq 1}$  and  $(x'_k, \theta'_k)_{k\geq 1}$  evolve independently. Given the above coupling, we first observe that

$$
\begin{aligned} \bar{W}_2(\mathcal{L}(x_k, \theta_k), \mathcal{L}(x'_k, \theta'_k)) &= \mathbb{E}[\bar{W}_2(\mathcal{L}(x_k, \theta_k), \mathcal{L}(x'_k, \theta'_k)) | x_0 = x'_0] \mathbb{P}(x_0 = x'_0) \\ &+ \mathbb{E}[\bar{W}_2(\mathcal{L}(x_k, \theta_k), \mathcal{L}(x'_k, \theta'_k)) | x_0 \neq x'_0] \mathbb{P}(x_0 \neq x'_0) \\ &= \epsilon \mathbb{E}[\bar{W}_2(\mathcal{L}(x_k, \theta_k), \mathcal{L}(x'_k, \theta'_k)) | x_0 \neq x'_0]. \end{aligned}
$$

The second equality holds since  $\mathbb{E}[\bar{W}_2(\mathcal{L}(x_k, \theta_k), \mathcal{L}(x_k, \theta'_k)) | x_0 = x'_0] = 0$  and  $\mathbb{P}(x_0 \neq x'_0) = \epsilon$ .

Next, we note the following upper bound of the Wasserstein distance,

$$
\begin{split} \bar{W}_2^2(\mathcal{L}(x_k,\theta_k),\mathcal{L}(x'_k,\theta'_k))&=\inf\mathbb{E}\Big[\,\mathbbm{1}\{x_k\neq x'_k\}+\|\theta_k-\theta'_k\|^2\Big] \\ &\leq 1+2\Big(\mathbb{E}[\|\theta_k-\theta^*\|^2]+\mathbb{E}[\|\theta'_k-\theta^*\|^2]\Big). \end{split}
$$

Making use of Proposition [4.2,](#page-5-1) we have

$$
\begin{aligned} & \bar{W}_2^2(\mathcal{L}(x_k, \theta_k), \mathcal{L}(x'_k, \theta'_k)) \\ &\leq & 2c_{2,1}(1 - \alpha \mu)^k \left( \mathbb{E}[\|\theta_0 - \theta^*\|^2] + \mathbb{E}[\|\theta'_0 - \theta^*\|^2] \right) + 4c_{2,2}\alpha\tau \cdot \frac{L^2}{\mu} + 1 \\ &\leq & 4c_{2,1}(1 - \alpha \mu)^k \mathbb{E}[\|\theta_0 - \theta^*\|^2] + 4c_{2,2}\alpha\tau \cdot \frac{L^2}{\mu} + 1, \end{aligned}
$$

where the second inequality holds for  $\theta_0 = \theta'_0$  by assumption.

Note that the above upper bound to the Wasserstein distance is independent of the choice of  $(x_0, x'_0)$ . Hence, we can conclude that

$$
\overline{W}_2(\mathcal{L}(x_k, \theta_k), \mathcal{L}(x'_k, \theta'_k)) = \epsilon \mathbb{E}[\overline{W}_2(\mathcal{L}(x_k, \theta_k), \mathcal{L}(x'_k, \theta'_k)) | x_0 \neq x'_0] \leq \epsilon \Big(4c_{2,1}(1-\alpha\mu)^k \mathbb{E}[\|\theta_0 - \theta^*\|^2] + 4c_{2,2}\alpha\tau \cdot \frac{L^2}{\mu} + 1\Big)^{1/2}.
$$

We complete the proof of the lemma.

By Lemma [C.5,](#page-30-0) we see that when  $x_0$  is close to its stationary distribution  $\pi$ ,  $\theta_k$  would not deviate too much from  $\theta'_{k}$ , as if it were initialized from the stationary distribution.

Now we consider a joint process  $(x_k, \theta_k)_{k>0}$  with arbitrary initialization. By the property of uniform ergodicity of  $(x_k)_{k\geq 0}$ , we know that  $\|\mathcal{L}(x_k) - \pi\|_{TV} \leq Rr^k$ . Choose time  $t_0 \geq 0$ . We construct a second

□

<span id="page-31-0"></span>Markov chain  $(x'_k, \theta'_k)_{k \geq t_0}$  with the following properties: (1)  $x'_{t_0} \sim \pi$  and is maximally coupled to  $x_{k_0}$ , i.e.,  $\|\mathcal{L}(x_{t_0}) - \mathcal{L}(x'_{t_0})\|_{TV} = \mathbb{P}(x_{t_0} \neq x'_{t_0})$  and (2)  $\theta'_{t_0} = \theta_{k_0}$ . Under this construction, for  $k \geq t_0 + \tau$ , we have

$$
\begin{split} \bar{W}_2(\mathcal{L}(x_k, \theta_k), \bar{\nu}) \leq & \bar{W}_2(\mathcal{L}(x_k, \theta_k), \mathcal{L}(x'_k, \theta'_k)) + \bar{W}_2(\mathcal{L}(x'_k, \theta'_k), \bar{\nu}) \\ \leq & Rr^{t_0} \Big( 4c_{2,1} (1 - \alpha \mu)^{k - t_0} \mathbb{E}[\|\theta_{t_0} - \theta^*\|^2] + 4c_{2,2}\alpha \tau \cdot \frac{L^2}{\mu} + 1 \Big)^{1/2} \\ & + 16(1 - \alpha \mu)^{k - t_0} \Big( \mathbb{E} \left[ \|\theta_{t_0} - \theta^*\|^2 \right] + c'_{2,2} \Big), \end{split}
$$

where the last inequality follows from Lemma [C.4](#page-29-0) and Lemma [C.5.](#page-30-0)

For each t with  $t_0 \geq \tau$ , set  $t_0 = t/2$ . From the above inequality, we obtain that

$$
\bar{W}_2(\mathcal{L}(x_t, \theta_t), \bar{\nu})
$$
\n
$$
\leq R r^{t/2} \Big( 4c_{2,1} (1 - \alpha \mu)^{t/2} \mathbb{E}[\|\theta_{t_0} - \theta^*\|^2] + 4c_{2,2} \alpha \tau \cdot \frac{L^2}{\mu} + 1 \Big)^{1/2}
$$
\n
$$
+ 16(1 - \alpha \mu)^{t/2} \Big( \mathbb{E} \left[ \|\theta_{t_0} - \theta^*\|^2 \right] + c'_{2,2} \Big)
$$
\n
$$
\leq R r^{t/2} \Big( 4c_{2,1} (1 - \alpha \mu)^{t/2} \cdot \Big( 8(1 - \alpha \mu)^{t_0} \mathbb{E}[\|\theta_0 - \theta^*\|^2] + c_{2,2} \alpha \tau \cdot \frac{L^2}{\mu} \Big) + 4c_{2,2} \alpha \tau \cdot \frac{L^2}{\mu} + 1 \Big)^{1/2}
$$
\n
$$
+ 16(1 - \alpha \mu)^{t/2} \Big( \Big( 8(1 - \alpha \mu)^{t_0} \mathbb{E}[\|\theta_0 - \theta^*\|^2] + c_{2,2} \alpha \tau \cdot \frac{L^2}{\mu} \Big) + c'_{2,2} \Big)
$$
\n
$$
\leq \max(r, 1 - \alpha \mu)^{t/2} \cdot s(\theta_0, \theta^*, \mu, L, R)
$$
\n
$$
\leq (1 - \alpha \mu)^{t/2} \cdot s(\theta_0, \theta^*, \mu, L, R)
$$

where  $s(\theta_0, L, \mu)$  denote some constant that depends on the initialization of  $\theta_0$ , and problem primitives L,  $\mu$ , but independent of stepsize  $\alpha$  and iteration index t. Last inequality holds because  $\mu \leq 1 - r$  and  $\alpha \leq \frac{\mu}{908L^2} \leq 1.$ 

Therefore, as  $t \to \infty$ , we obtain that  $\overline{W}_2(\mathcal{L}(x_k, \theta_k), \overline{\nu}) \to 0$ , which implies that the Markov chain  $(x_k, \theta_k)_{k>0}$  with arbitrary initailization converges to the same  $\bar{\nu}$ . As such, we have proved the desired weak convergence result without the assumption on  $x_0 \sim \pi$  initialization.

Additionally, we obtain the following convergence rate. For any initialization  $(x_0, \theta_0) \in \mathcal{X} \times \mathbb{R}^d$ , we have

$$
W_2(\mathcal{L}(\theta_t), \mu) \leq \overline{W}_2(\mathcal{L}(x_t, \theta_t), \overline{\nu}) \leq (1 - \alpha \mu)^{t/2} \cdot s(\theta_0, L, \mu).
$$

#### C.2.1 Proof of Proposition [C.2](#page-27-1)

<span id="page-31-1"></span>First, we present the following lemma that is similar to  $\left[\text{CZD}^+\text{22}, \text{Lemma 2.3}\right]$ .

**Lemma C.6.** For any  $k_1 < k_2$  satisfying  $\alpha(k_2 - k_1) \leq \frac{1}{8L}$ , the following six inequalities hold:

$$
\begin{aligned}\n\|\theta_{k_2}^{[1]} - \theta_{k_2}^{[2]} - \theta_{k_1}^{[1]} + \theta_{k_1}^{[2]}\| &\leq 8\alpha L(k_2 - k_1) \|\theta_{k_2}^{[1]} - \theta_{k_2}^{[2]}\| \\
\|\theta_{k_2}^{[1]} - \theta_{k_2}^{[2]} - \theta_{k_1}^{[1]} + \theta_{k_1}^{[2]}\| &\leq 8\alpha L(k_2 - k_1) \|\theta_{k_1}^{[1]} - \theta_{k_1}^{[2]}\| \\
\|\theta_{k_2}^{[1]} - \theta_{k_1}^{[1]}\| &\leq 8\alpha L(k_2 - k_1) \left(\|\theta_{k_2}^{[1]} \| + 1\right) \\
\|\theta_{k_2}^{[1]} - \theta_{k_1}^{[1]}\| &\leq 8\alpha L(k_2 - k_1) \left(\|\theta_{k_1}^{[1]} \| + 1\right) \\
\|\theta_{k_2}^{[2]} - \theta_{k_1}^{[2]}\| &\leq 8\alpha L(k_2 - k_1) \left(\|\theta_{k_2}^{[2]} \| + 1\right) \\
\|\theta_{k_2}^{[2]} - \theta_{k_1}^{[2]}\| &\leq 8\alpha L(k_2 - k_1) \left(\|\theta_{k_2}^{[2]} \| + 1\right) \\
\|\theta_{k_2}^{[2]} - \theta_{k_1}^{[2]}\| &\leq 8\alpha L(k_2 - k_1) \left(\|\theta_{k_1}^{[2]} \| + 1\right).\n\end{aligned}
$$

*Proof.* Consider the coupling given by equation  $(C.3)$ , by Assumption [2](#page-2-2) and [6,](#page-26-1) we have

 $\|\theta_{k+1}^{[1]} - \theta_{k+1}^{[2]}\| - \|\theta_{k}^{[1]} - \theta_{k}^{[2]}\|$  $\|\hat{\theta}_{k+1}^{[2]}\| \leq \|\theta_{k+1}^{[1]} - \theta_{k+1}^{[2]} - \theta_{k}^{[1]} + \theta_{k}^{[2]}.$  $k^{\prime}$  ||

$$
= \alpha ||g(\theta_k^{[1]}, x_k) + \xi_{k+1}(\theta_k^{[1]}) - g(\theta_k^{[1]}, x_k) - \xi_{k+1}(\theta_k^{[1]})||
$$
  

$$
\leq 2\alpha L ||\theta_k^{[1]} - \theta_k^{[2]}||.
$$

<span id="page-32-0"></span>Given  $k_1 < k_2$ , for  $\forall t \in [k_1, k_2]$ , since  $1 + x \leq e^x$  for  $\forall x \in R$ , we have

$$
\begin{aligned} \|\theta_t^{[1]} - \theta_t^{[2]}\| &\leq \prod_{j=k_1}^{t-1} (1 + 2\alpha L) \|\theta_{k_1}^{[1]} - \theta_{k_1}^{[2]}\| \\ &\leq \exp(2\alpha (k_2 - k_1)L) \|\theta_{k_1}^{[1]} - \theta_{k_1}^{[2]}\| \\ &\overset{\text{(i)}}{\leq} (1 + 4\alpha (k_2 - k_1)L) \|\theta_{k_1}^{[1]} - \theta_{k_1}^{[2]}\| \\ &\leq 2 \|\theta_{k_1}^{[1]} - \theta_{k_1}^{[2]}\|, \end{aligned}
$$

where (i) holds for  $e^x \leq 1 + 2x \,\forall x \in [0, \frac{1}{2}]$  and  $\alpha(k_2 - k_1) \leq \frac{1}{8L}$ .

Then, we have

$$
\|\theta_{k_2}^{[1]} - \theta_{k_2}^{[2]} - \theta_{k_1}^{[1]} + \theta_{k_1}^{[2]}\| \le \sum_{t=k_1}^{k_2-1} \|\theta_{t+1}^{[1]} - \theta_{t+1}^{[2]} - \theta_{t}^{[1]} + \theta_{t}^{[2]}\| \le 4\alpha(k_2 - k_1)L\|\theta_{k_1}^{[1]} - \theta_{k_1}^{[2]}\|.
$$

Therefore, following  $\alpha(k_2 - k_1) \leq \frac{1}{8L}$ , we have

$$
\begin{aligned} \|\theta_{k_2}^{[1]} - \theta_{k_2}^{[2]} - \theta_{k_1}^{[1]} + \theta_{k_1}^{[2]} \| &\le 4\alpha (k_2 - k_1)L \|\theta_{k_1}^{[1]} - \theta_{k_1}^{[2]}\| \\ &\le 4\alpha (k_2 - k_1)L (\|\theta_{k_2}^{[1]} - \theta_{k_2}^{[2]} - \theta_{k_1}^{[1]} + \theta_{k_1}^{[2]} \| + \|\theta_{k_2}^{[1]} - \theta_{k_2}^{[2]} \|) \\ &\le \frac{1}{2} \|\theta_{k_2}^{[1]} - \theta_{k_2}^{[2]} - \theta_{k_1}^{[1]} + \theta_{k_1}^{[2]} \| + 4\alpha (k_2 - k_1)L \|\theta_{k_2}^{[1]} - \theta_{k_2}^{[2]} \| .\end{aligned}
$$

Then, by rearranging the terms, we have

$$
\|\theta_{k_2}^{[1]} - \theta_{k_2}^{[2]} - \theta_{k_1}^{[1]} + \theta_{k_1}^{[2]}\| \le 8\alpha(k_2 - k_1)L\|\theta_{k_2}^{[1]} - \theta_{k_2}^{[2]}\|,
$$

thereby we have proved the first two inequalities of Lemma [C.6.](#page-31-1)

By Assumption [2](#page-2-2) and  $6$  and  $[CZD+22, \text{Lemma 2.3}],$  we can prove the last four inequalities and we omit the details here.  $\Box$ 

Now, we are ready to prove Proposition [C.2.](#page-27-1) We start with the following decomposition. By equation [\(C.3\)](#page-27-3), we first have

$$
\mathbb{E}[\|\theta_{k+1}^{[1]} - \theta_{k+1}^{[2]}\|^2] = \mathbb{E}\left[\|\theta_k^{[1]} - \theta_k^{[2]} + \alpha\left(g(\theta_k^{[1]}, x_k) - g(\theta_k^{[2]}, x_k) + \xi_{k+1}(\theta_k^{[1]}) - \xi_{k+1}(\theta_k^{[2]})\right)\|^2\right]
$$
  
\n
$$
= \mathbb{E}\left[\|\theta_k^{[1]} - \theta_k^{[2]}\|^2\right] + 2\alpha \underbrace{\mathbb{E}\left[\langle\theta_k^{[1]} - \theta_k^{[2]}, g(\theta_k^{[1]}, x_k) - g(\theta_k^{[2]}, x_k)\rangle\right]}_{T_1}
$$
  
\n
$$
+ \alpha^2 \underbrace{\mathbb{E}\left[\|g(\theta_k^{[1]}, x_k) - g(\theta_k^{[2]}, x_k) + \xi_{k+1}(\theta_k^{[1]}) - \xi_{k+1}(\theta_k^{[2]})\|^2\right]}_{T_2}.
$$

For  $T_2$  $T_2$ , by Assumption 2 and [6,](#page-26-1) we have

$$
T_2\le 4L^2\mathbb{E}\left[\|\theta_k^{[1]}-\theta_k^{[2]}\|^2\right].
$$

We denote  $\varepsilon(\cdot, x_k) := g(\cdot, x_k) - \overline{g}(\cdot)$  to be the noise function. Then, by Assumption [2,](#page-2-2) we conclude that  $\varepsilon(\cdot, x_k)$  is 2L-Lipschitz continuous. Therefore,  $T_1$  can be rewritten as:

$$
T_1 = \mathbb{E}\left[\langle \theta_k^{[1]} - \theta_k^{[2]}, \varepsilon(\theta_k^{[1]}, x_k) - \varepsilon(\theta_k^{[2]}, x_k)\rangle\right] + \mathbb{E}\left[\langle \theta_k^{[1]} - \theta_k^{[2]}, \bar{g}(\theta_k^{[1]}) - \bar{g}(\theta_k^{[2]})\rangle\right].
$$

For  $\mathbb{E}\left[\langle \theta_k^{[1]}-\theta_k^{[2]}\right]$  $_{k}^{[2]},\bar{g}(\theta_{k}^{[1]}% )_{k}^{[2]}$  $[\binom{[1]}{k} - \bar{g}(\theta_k^{[2]})$  $\binom{[2]}{k}$ , by Assumption [3,](#page-2-3) we have  $\mathbb{E}\left[\langle \theta_k^{[1]}-\theta_k^{[2]}\right]$  $_{k}^{[2]},\bar{g}(\theta_{k}^{[1]}% )_{k}^{[2]}$  $[\binom{[1]}{k} - \bar{g}(\theta_k^{[2]})]$  $\left[ \begin{smallmatrix} [2] \ k \end{smallmatrix} \right] \rangle \Big] \leq - \mu \mathbb{E} \left[ \|\theta_k^{[1]} - \theta_k^{[2]}\right]$  $\binom{[2]}{k}$  $\left\| \begin{matrix} 2 \end{matrix} \right\}$ . Let  $\mathcal{F}_k := \sigma\left( (\theta_t^{[1]}, \theta_t^{[2]}, x_t) \mid t \leq k \right)$ . For  $\mathbb{E} \left[ \langle \theta_k^{[1]} - \theta_k^{[2]} \right]$  $_k^{[2]}, \varepsilon(\theta_k^{[1]}$  $\mathcal{E}_k^{[1]}, x_k) - \varepsilon(\theta_k^{[2]}$  $\left[\begin{matrix} [2], x_k \end{matrix}\right]$ , we have  $\mathbb{E}\left[\langle \theta_k^{[1]}-\theta_k^{[2]}\right]$  $_k^{[2]}, \varepsilon(\theta_k^{[1]}$  $\binom{[1]}{k}, x_k - \varepsilon(\theta_k^{[2]})$  $_{k}^{[2]},x_{k})\rangle$  $=$   $\mathbb{E}\left[\mathbb{E}\left[\langle \theta_k^{[1]}-\theta_k^{[2]}\right. \right]$  $_k^{[2]}, \varepsilon(\theta_k^{[1]}$  $\binom{[1]}{k},x_k)-\varepsilon(\theta_k^{[2]}$  $\left[ \begin{smallmatrix} [2], 1 \ k \end{smallmatrix} \right], x_k) \rangle \mid \mathcal{F}_{k-\tau}$  $=$   $\mathbb{E}\left[\mathbb{E}\left[\langle\theta_{k-\tau}^{[1]}-\theta_{k-\tau}^{[2]}\right.\right.$  ${}^{[2]}_{k-\tau}, \varepsilon(\theta^{[1]}_{k-\tau})$  ${}_{k-\tau}^{[1]}, x_k) - \varepsilon(\theta_{k-\tau}^{[2]}$  $\vert_{k=\tau}^{\lfloor 2 \rfloor}, x_k) \rangle \mid \mathcal{F}_{k-\tau}$  $\left[\begin{matrix} \end{matrix}\right]$   $(T_3)$  $+ \ \mathbb{E} \left[ \langle \theta_k^{[1]} - \theta_k^{[2]} - \theta_{k-\tau}^{[1]} - \theta_{k-\tau}^{[2]} \right.$  ${}^{[2]}_{k-\tau}, \varepsilon(\theta^{[1]}_{k-\tau})$  ${}_{k-\tau}^{[1]}, x_k) - \varepsilon(\theta_{k-\tau}^{[2]}$  $\left[\begin{smallmatrix} [2] \ k-\tau \end{smallmatrix}, x_k) \right\rangle$  $(T_4)$  $+ \ \mathbb{E}\Big[ \big\langle \theta_k^{[1]}-\theta_k^{[2]}$  $_k^{[2]}, \varepsilon(\theta_k^{[1]}$  $\binom{[1]}{k}, x_k - \varepsilon(\theta_k^{[2]})$  $\binom{[2]}{k}, x_k - \varepsilon(\theta_{k-1}^{[1]})$  ${}_{k-\tau}^{[1]}, x_k) + \varepsilon(\theta_{k-\tau}^{[2]}$  $\vert_{k-\tau}^{^{[2]}},x_k)\rangle$  $\overbrace{\hspace{2.5cm}}^{\bullet}$ i .

We assume  $\alpha \tau \leq \frac{1}{8L}$ . For  $T_3$ , by definition of mixing time  $\tau$ , we obtain

$$
T_3 = \mathbb{E} \left[ \mathbb{E} \left[ \langle \theta_{k-\tau}^{[1]} - \theta_{k-\tau}^{[2]}, \varepsilon(\theta_{k-\tau}^{[1]}, x_k) - \varepsilon(\theta_{k-\tau}^{[2]}, x_k) \rangle \mid \theta_{k-\tau}^{[1]}, \theta_{k-\tau}^{[2]}, x_{k-\tau} \right] \right]
$$
  
\n
$$
\leq 2\alpha L \mathbb{E} \left[ \| \theta_{k-\tau}^{[1]} - \theta_{k-\tau}^{[2]} \|^{2} \right]
$$
  
\n
$$
\leq 4\alpha L \mathbb{E} \left[ \| \theta_{k}^{[1]} - \theta_{k}^{[2]} \|^{2} \right] + 4\alpha L \mathbb{E} \left[ \| \theta_{k-\tau}^{[1]} - \theta_{k-\tau}^{[2]} - \theta_{k}^{[1]} + \theta_{k}^{[2]} \|^{2} \right]
$$
  
\n
$$
\leq (4\alpha L + 256\alpha^{3} \tau^{2} L^{3}) \mathbb{E} \left[ \| \theta_{k}^{[1]} - \theta_{k}^{[2]} \|^{2} \right],
$$

where the last inequality holds by Lemma [C.6.](#page-31-1)

For  $T_4$ , we obtain

<span id="page-33-2"></span><span id="page-33-1"></span><span id="page-33-0"></span>
$$
T_4 \leq \mathbb{E}\left[\|\theta_k^{[1]} - \theta_k^{[2]} - \theta_{k-\tau}^{[1]} + \theta_{k-\tau}^{[2]}\|\|\varepsilon(\theta_{k-\tau}^{[1]}, x_k) - \varepsilon(\theta_{k-\tau}^{[2]}, x_k)\|\right]
$$
  
\n
$$
\leq 16\alpha\tau L^2 \mathbb{E}\left[\|\theta_k^{[1]} - \theta_k^{[2]}\|\|\theta_{k-\tau}^{[1]} - \theta_{k-\tau}^{[2]}\|\right]
$$
  
\n
$$
\leq (16\alpha\tau L^2 + 128\alpha^2\tau^2 L^3)\mathbb{E}\left[\|\theta_k^{[1]} - \theta_k^{[2]}\|^2\right].
$$

Below, we bound term  $\spadesuit$  by two different Taylor expansions. One the one hand, there exist  $\lambda_1, \lambda_2 \in [0,1]$ such that  $h_k = \lambda_1 \theta_k^{[1]} + (1 - \lambda_1) \theta_k^{[2]}$  $\lambda_k^{[2]}, h_{k-\tau} = \lambda_2 \theta_{k-\tau}^{[1]} + (1-\lambda_2)\theta_{k-\tau}^{[2]}$  $\int_{k-\tau}^{\lfloor 2 \rfloor}$  and

$$
|\spadesuit| = |\langle \theta_k^{[1]} - \theta_k^{[2]}, \varepsilon'(h_k, x_k)(\theta_k^{[1]} - \theta_k^{[2]}) - \varepsilon'(h_{k-\tau}, x_k)(\theta_{k-\tau}^{[1]} - \theta_{k-\tau}^{[2]})\rangle|
$$
\n
$$
= |\langle \theta_k^{[1]} - \theta_k^{[2]}, \varepsilon'(h_{k-\tau}, x_k)(\theta_k^{[1]} - \theta_k^{[2]} - \theta_{k-\tau}^{[1]} + \theta_{k-\tau}^{[2]})\rangle
$$
\n
$$
+ \langle \theta_k^{[1]} - \theta_k^{[2]}, (\varepsilon'(h_k, x_k) - \varepsilon'(h_{k-\tau}, x_k))( \theta_k^{[1]} - \theta_k^{[2]})\rangle|
$$
\n
$$
\leq 16\alpha\tau L^2 ||\theta_k^{[1]} - \theta_k^{[2]} ||^2 + |\langle \theta_k^{[1]} - \theta_k^{[2]}, (\varepsilon'(h_k, x_k) - \varepsilon'(h_{k-\tau}, x_k))( \theta_k^{[1]} - \theta_k^{[2]})\rangle|
$$
\n
$$
\leq 16\alpha\tau L^2 ||\theta_k^{[1]} - \theta_k^{[2]} ||^2 + 2L ||\theta_k^{[1]} - \theta_k^{[2]} ||^2 ||h_k - h_{k-\tau}||
$$
\n
$$
\leq 16\alpha\tau L^2 ||\theta_k^{[1]} - \theta_k^{[2]} ||^2 + 2L ||\theta_k^{[1]} - \theta_k^{[2]} ||^2 (||h_k - \theta_k^{[2]}|| + ||\theta_k^{[2]} - \theta_{k-\tau}^{[2]}|| + ||\theta_{k-\tau}^{[2]} - h_{k-\tau}||)
$$
\n
$$
\leq 16\alpha\tau L^2 ||\theta_k^{[1]} - \theta_k^{[2]} ||^2 + 2L ||\theta_k^{[1]} - \theta_k^{[2]} ||^2 (||\theta_k^{[1]} - \theta_k^{[2]}|| + ||\theta_k^{[2]} - \theta_{k-\tau}^{[2]}|| + ||\theta_{k-\tau}^{[1]} - \theta_{k-\tau}^{[2]}||)
$$
\n
$$
\leq 16\alpha\tau L^2 ||\theta_k^{[1]} - \theta_k^{[2]} ||^2 + 2L ||\theta_k^{[1]} - \theta_k^{[2]} ||^2 (||\theta_k^{[1]} - \theta_k^{[2]}
$$

where we note that choosing the second iterates for triangle inequality in equation [\(C.7\)](#page-33-0) and choosing  $\theta_k^{[2]}$ k for bounding  $\theta_k^{[2]} - \theta_{k-1}^{[2]}$  $\binom{[2]}{k-\tau}$  in equation [\(C.8\)](#page-33-1) are both symmetric, which implies that we can replace the  $\theta_k^{[2]}$  $\prod_{k=1}^{\lfloor 2 \rfloor}$  in term A with arbitrary one in  $\{\theta_{k-1}^{[1]}$  $\theta_{k-\tau}^{[1]}, \theta_{k-\tau}^{[2]}, \theta_{k-\tau}^{[2]}\}.$ 

On the other hand, there exist  $\bar{\lambda}_1, \bar{\lambda}_2 \in [0, 1]$  such that  $p_k = \bar{\lambda}_1 \theta_k^{[1]} + (1 - \bar{\lambda}_1) \theta_{k-1}^{[1]}$  $\bar{\lambda}_{k-\tau}^{[1]}, q_k = \bar{\lambda}_2 \theta_k^{[2]} + (1 - \bar{\lambda}_2) \theta_{k-\tau}^{[2]}$  $k-\tau$ and

$$
|\spadesuit| = |\langle \theta_k^{[1]} - \theta_k^{[2]}, \varepsilon'(p_k, x_k)(\theta_k^{[1]} - \theta_{k-\tau}^{[1]}) - \varepsilon'(q_k, x_k)(\theta_k^{[2]} - \theta_{k-\tau}^{[2]})\rangle|
$$
  
\n
$$
= |\langle \theta_k^{[1]} - \theta_k^{[2]}, \varepsilon'(p_k, x_k)(\theta_k^{[1]} - \theta_k^{[2]} - \theta_{k-\tau}^{[1]} + \theta_{k-\tau}^{[2]})\rangle
$$
  
\n
$$
+ \langle \theta_k^{[1]} - \theta_k^{[2]}, (\varepsilon'(p_k, x_k) - \varepsilon'(q_k, x_k))(\theta_k^{[2]} - \theta_{k-\tau}^{[2]})\rangle|
$$
  
\n
$$
\leq 16\alpha\tau L^2 \|\theta_k^{[1]} - \theta_k^{[2]}\|^2 + 2L \|\theta_k^{[1]} - \theta_k^{[2]}\| \|p_k - q_k\| \|\theta_k^{[2]} - \theta_{k-\tau}^{[2]}\|
$$
  
\n
$$
\leq 16\alpha\tau L^2 \|\theta_k^{[1]} - \theta_k^{[2]}\|^2 + 16\alpha\tau L^2 \|\theta_k^{[1]} - \theta_k^{[2]}\| \|p_k - q_k\| \|(\|\theta_k^{[2]}\| + 1),
$$
\n(C.10)

where adding and subtracting the second iterates in equation [\(C.7\)](#page-33-0) and choosing  $\theta_k^{[2]}$  $\theta_k^{[2]}$  for bounding  $\theta_k^{[2]} - \theta_{k-1}^{[2]}$  $k-\tau$ in equation  $(C.8)$  are both symmetric. We have

$$
\|p_k - q_k\| = \|\bar{\lambda}_1(\theta_k^{[1]} - \theta_k^{[2]}) + (1 - \bar{\lambda}_1)(\theta_{k-\tau}^{[1]} - \theta_{k-\tau}^{[2]}) + (\bar{\lambda}_1 - \bar{\lambda}_2)(\theta_k^{[2]} - \theta_{k-\tau}^{[2]})\| \\
\leq (2 + 8\alpha\tau L)\|\theta_k^{[1]} - \theta_k^{[2]}\| + 8\alpha\tau L(\|\theta_k^{[2]}\| + 1) \\
\leq 3\|\theta_k^{[1]} - \theta_k^{[2]}\| + 8\alpha\tau L(\|\theta_k^{[2]}\| + 1).
$$

Therefore, we obtain

$$
|\spadesuit| \leq 16\alpha\tau L^2 \|\theta_k^{[1]} - \theta_k^{[2]}\|^2
$$
  
+ 2L $\|\theta_k^{[1]} - \theta_k^{[2]}\| \left(3\|\theta_k^{[1]} - \theta_k^{[2]}\| + \frac{8\alpha\tau L(\|\theta_k^{[2]}\| + 1)}{B}\right) \underbrace{8\alpha\tau L(\|\theta_k^{[2]}\| + 1)}_C := \text{(Bound2)},$ 

and we can replace the  $\theta_k^{[2]}$  $\binom{[2]}{k}$  in term B and C with arbitrary one in  $\{\theta_{k-1}^{[1]}$  ${}_{k-\tau}^{[1]}, \theta_k^{[1]}, \theta_{k-\tau}^{[2]}, \theta_k^{[2]}\}.$ By (Bound1) and (Bound2), we have

$$
|\spadesuit| \le \min(\text{Bound1}, \text{Bound2})
$$
  
\$\le 16\alpha \tau L^2 \|\theta\_k^{[1]} - \theta\_k^{[2]}\|^2 + 48\alpha \tau L^2 \|\theta\_k^{[1]} - \theta\_k^{[2]}\|^2 (\|\theta\_k^{[2]}\| + 1).

Below, we discuss the upper bound for  $|\spadesuit|$  by three cases.

**Case 1:** If one of  $\{\theta_{k-}^{[1]}$  $\{\mu^{[1]}_{k-\tau}, \theta^{[2]}_k, \theta^{[2]}_{k-\tau}, \theta^{[2]}_k\}$  has norm that is less than  $4\delta_{\alpha\tau L^2}$ , where we define  $\delta_{\alpha\tau L^2} :=$  $\delta(\alpha \tau L^2)$ , without loss of generality, we assume  $\|\theta_k^{[2]}$  $\|k^{[2]}\| \leq 4\delta_{\alpha\tau L^2}$ . Multiply  $\mathbb{1}(\|\theta_k^{[2]})$  $||\mathbf{k}^{[2]}|| \leq 4\delta_{\alpha\tau L^2}$  to both sides of the above inequality, and we have

$$
\begin{aligned} & \|\spadesuit\|1(\|\theta_k^{[2]}\| \le 4\delta_{\alpha\tau L^2}) \\ &\le \left(16\alpha\tau L^2 \|\theta_k^{[1]} - \theta_k^{[2]}\|^2 + 48\alpha\tau L^2 \|\theta_k^{[1]} - \theta_k^{[2]}\|^2 (\|\theta_k^{[2]}\| + 1)\right)1(\|\theta_k^{[2]}\| \le 4\delta_{\alpha\tau L^2}) \\ &\le (64 + 192\delta_{\alpha\tau L^2})\alpha\tau L^2 \|\theta_k^{[1]} - \theta_k^{[2]}\|^2, \end{aligned}
$$

where we can replace the  $\theta_k^{[2]}$  $\frac{[2]}{k}$  in the left hand of the above inequality with an arbitrary one in  $\{\theta_{k-1}^{[1]}$  $\theta_k^{[1]}, \theta_k^{[2]}, \theta_{k-\tau}^{[2]}, \theta_k^{[2]} \}$ by the similar argument under (Bound1) and (Bound2).

**Case 2:** If  $\|\theta_k^{[1]}$  $\|\kappa^{[1]}\| < 2 \|\theta_k^{[1]} - \theta_k^{[2]}$  $\|k^{\left(2\right)}\|$ , we obtain

$$
\|\theta_k^{[2]}\|\leq \|\theta_k^{[1]}-\theta_k^{[2]}\|+\|\theta_k^{[1]}\|\leq 3\|\theta_k^{[1]}-\theta_k^{[2]}\|.
$$

Therefore, by definition of  $\spadesuit$  and Lemma [C.6,](#page-31-1) we obtain

$$
|\spadesuit|\mathbbm{1}(\|\theta_k^{[1]}\|\leq 2\|\theta_k^{[1]}-\theta_k^{[2]}\|)\leq 2L\|\theta_k^{[1]}-\theta_k^{[2]}\|(\|\theta_k^{[1]}-\theta_{k-\tau}^{[1]}\|+\|\theta_k^{[2]}-\theta_{k-\tau}^{[2]}\|)\mathbbm{1}(\|\theta_k^{[1]}\|\leq 2\|\theta_k^{[1]}-\theta_k^{[2]}\|)
$$

$$
\leq 16\alpha\tau L^2 \|\theta_k^{[1]} - \theta_k^{[2]} \| (\|\theta_k^{[1]}\| + \|\theta_k^{[2]}\|) \mathbb{1}(\|\theta_k^{[1]}\| \leq 2 \|\theta_k^{[1]} - \theta_k^{[2]}\|)
$$
  

$$
\leq 80\alpha\tau L^2 \|\theta_k^{[1]} - \theta_k^{[2]}\|.
$$

**Case 3:** If all four variables in  $\{\theta_{k-1}^{[1]}\}$  $\begin{bmatrix} [1]_{k-\tau}, \theta_k^{[1]}, \theta_{k-\tau}^{[2]}, \theta_k^{[2]} \end{bmatrix}$  have a norm larger than  $4\delta_{\alpha\tau L^2}$ , and  $\|\theta_k^{[1]}$  $\|k\|$   $\geq$  $2\|\theta_k^{[1]}-\theta_k^{[2]}$  $\|k^2\|$ , we obtain

$$
||h_k|| = ||\lambda_1 \theta_k^{[1]} + (1 - \lambda_1) \theta_k^{[2]}|| \ge ||\theta_k^{[1]}|| - ||\theta_k^{[1]} - \theta_k^{[2]}|| \ge \frac{||\theta_k^{[1]}||}{2} \ge 2\delta_{\alpha\tau L^2}
$$
  
\n
$$
||h_{k-\tau}|| = ||\lambda_2 \theta_{k-\tau}^{[1]} + (1 - \lambda_2) \theta_{k-\tau}^{[2]}||
$$
  
\n
$$
\ge ||\theta_{k-\tau}^{[1]}|| - ||\theta_{k-\tau}^{[1]} - \theta_{k-\tau}^{[2]}||
$$
  
\n
$$
\stackrel{(i)}{\ge} \delta_{\alpha\tau L^2} - (1 + 8\alpha\tau L) ||\theta_k^{[1]} - \theta_k^{[2]}||
$$
  
\n
$$
\ge \delta_{\alpha\tau L^2},
$$

where (i) holds by choosing  $\alpha \tau \leq \frac{1}{16L}$ . Therefore, we have  $||h_k|| \geq \delta_{\alpha \tau L^2}$  and  $||h_{k-\tau}|| \geq \delta_{\alpha \tau L^2}$ . Therefore, by equation  $(C.6)$  and Assumption  $6,$  we obtain

$$
\|\Phi\| \mathbb{1}(\text{Case 3})
$$
  
\n
$$
\leq 16\alpha \tau L^2 \|\theta_k^{[1]} - \theta_k^{[2]}\|^2 + \|\theta_k^{[1]} - \theta_k^{[2]}\|^2 (\|\varepsilon'(h_k, x_k) - G(x_k)\| + \|\varepsilon'(h_{k-\tau}, x_k) - G(x_k)\|)
$$
  
\n
$$
\leq 18\alpha \tau L^2 \|\theta_k^{[1]} - \theta_k^{[2]}\|^2.
$$

Therefore, we obtain

$$
|\mathbb{E}[\spadesuit]| \leq \mathbb{E}[|\spadesuit|] \leq \mathbb{E}[|\spadesuit|1(||\theta_{k-\tau}^{[1]}|| \leq \delta_{\alpha\tau L^2})] + \mathbb{E}[\spadesuit|1(||\theta_{k}^{[1]}|| \leq \delta_{\alpha\tau L^2})]
$$
  
+  $\mathbb{E}[|\spadesuit|1(||\theta_{k-\tau}^{[2]}|| \leq \delta_{\alpha\tau L^2})] + \mathbb{E}[|\spadesuit|1(||\theta_{k}^{[2]}|| \leq \delta_{\alpha\tau L^2})]$   
+  $\mathbb{E}[\spadesuit|1(||\theta_{k}^{[1]}|| \leq 2||\theta_{k}^{[1]} - \theta_{k}^{[2]}||)] + \mathbb{E}[\spadesuit|1(\text{Case 3})]$   
 $\leq (354 + 768\delta_{\alpha\tau L^2})\alpha\tau L^2 \mathbb{E}[\|\theta_{k}^{[1]} - \theta_{k}^{[2]}\|^2].$ 

By Assumption [6,](#page-26-1) there exists  $\kappa_{\mu} > 0$  such that  $\epsilon \delta(\epsilon) \leq \frac{\mu}{3072}$  for  $\forall \epsilon \leq \kappa_{\mu}$ . By the above bounds, when  $\alpha \tau \leq \min(\frac{c\mu}{L^2}, \frac{\kappa_\mu}{L^2})$ , where we specify the constant  $c < 1$  later, we obtain

$$
\mathbb{E}[\|\theta_{k+1}^{[1]} - \theta_{k+1}^{[2]}\|^2]
$$
\n
$$
\leq (1 + 2\alpha(-\mu + 4\alpha L + 256\alpha^3 \tau^2 L^3 + 128\alpha^2 \tau^2 L^3 + 784\alpha \tau L^2 + 768\delta_{\alpha\tau L^2}\alpha \tau L^2) + 4\alpha^2 L^2) \mathbb{E}[\|\theta_k^{[1]} - \theta_k^{[2]}\|^2]
$$
\n
$$
= (1 + \alpha(-2\mu + 1568\alpha \tau L^2 + 256\alpha^2 \tau^2 L^3 + 1536\delta_{\alpha\tau L^2}\alpha \tau L^2) + \alpha^2 (8L + 512\alpha^2 \tau^2 L^3 + 4L^2) \mathbb{E}[\|\theta_k^{[1]} - \theta_k^{[2]}\|^2]
$$
\n
$$
\leq (1 + \alpha\mu(-2 + 1568c + 256c + \frac{1}{2} + 524c)) \mathbb{E}[\|\theta_k^{[1]} - \theta_k^{[2]}\|^2]
$$
\n
$$
\leq (1 - \mu\alpha) \mathbb{E}[\|\theta_k^{[1]} - \theta_k^{[2]}\|^2],
$$

where the last inequality holds by choosing  $c = \frac{1}{4696}$ . Therefore, for  $k \geq \tau$  and  $\alpha \tau \leq \min(\frac{\mu}{906L^2}, \frac{\kappa_{\mu}}{L^2})$  we have

$$
\mathbb{E}[\|\theta_k^{[1]} - \theta_k^{[2]}\|^2] \le (1 - \mu\alpha)^{k-\tau} \mathbb{E}[\|\theta_\tau^{[1]} - \theta_\tau^{[2]}\|^2]
$$
  
\n
$$
\le 2(1 - \mu\alpha)^{k-\tau} \mathbb{E}[\|\theta_0^{[1]} - \theta_0^{[2]}\|^2 + \|\theta_\tau^{[1]} - \theta_\tau^{[2]} - \theta_0^{[1]} + \theta_0^{[2]}\|^2]
$$
  
\n
$$
\le 2(1 + 8\alpha\tau L)(1 - \mu\alpha)^{k-\tau} \mathbb{E}[\|\theta_0^{[1]} - \theta_0^{[2]}\|^2]
$$
  
\n
$$
\le 4(1 - \mu\alpha)^{k-\tau} \mathbb{E}[\|\theta_0^{[1]} - \theta_0^{[2]}\|^2].
$$

### <span id="page-36-0"></span>C.3 Proof of Projected SA

Now, we specialize the proof of Theorem [C.1](#page-26-3) to the projected SA iterates.

We consider the same coupling:

$$
\begin{aligned}\n\theta_{k+\frac{1}{2}}^{[1]} &= \theta_k^{[1]} + \alpha \big( g(\theta_k^{[1]}, x_k) + \xi_{k+1}(\theta_k^{[1]}) \big), \\
\theta_{k+1}^{[1]} &= \Pi_{B(\beta)} \Big[ \theta_{k+\frac{1}{2}}^{[1]} \Big], \\
\theta_{k+\frac{1}{2}}^{[2]} &= \theta_k^{[2]} + \alpha \big( g(\theta_k^{[2]}, x_k) + \xi_{k+1}(\theta_k^{[2]}) \big), \\
\theta_{k+1}^{[2]} &= \Pi_{B(\beta)} \Big[ \theta_{k+\frac{1}{2}}^{[2]} \Big].\n\end{aligned}
$$

We first need to verify that Proposition [C.2](#page-27-1) holds for projected SA. By the non-expansion property of  $\Pi_{B(\beta)}$  with respect to  $\|\cdot\|$ , we obtain the following inequality:

$$
\|\theta_{k+1}^{[1]}-\theta_{k+1}^{[2]}\|-\|\theta_{k}^{[1]}-\theta_{k}^{[2]}\|\leq \|\theta_{k+\frac{1}{2}}^{[1]}-\theta_{k+\frac{1}{2}}^{[2]}\|-\|\theta_{k}^{[1]}-\theta_{k}^{[2]}\|,
$$

which implies Lemma  $C.6$  still holds for projected SA. For Proposition  $C.2$ , we can notice that the iterates of projected SA will always satisfy Case 1 with a finite bound  $\beta$  and we do not need to discuss Cases 2 and 3.

Therefore, when  $\alpha \tau \le \min(\frac{c\mu}{L^2}, \frac{1}{8L})$ , where we specify the constant  $c < 1$  later, we obtain

$$
\mathbb{E}[\|\theta_{k+1}^{[1]} - \theta_{k+1}^{[2]}\|^2]
$$
\n
$$
\leq (1 + \alpha(-2\mu + (160 + 96\beta))\alpha\tau L^2 + 256\alpha^2\tau^2 L^3) + \alpha^2(8L + 512\alpha^2\tau^2 L^3 + 4L^2)\Big) \mathbb{E}[\|\theta_k^{[1]} - \theta_k^{[2]}\|^2]
$$
\n
$$
\leq (1 + \alpha\mu(-2 + (160 + 96\beta)c + 256c + 524c)) \mathbb{E}[\|\theta_k^{[1]} - \theta_k^{[2]}\|^2]
$$
\n
$$
= (1 - \mu\alpha)\mathbb{E}[\|\theta_k^{[1]} - \theta_k^{[2]}\|^2],
$$

where we set  $c = \frac{1}{940+96\beta}$ .

Therefore,  $\forall k \geq \tau$  and  $\alpha \tau \leq \frac{\mu}{(940+96\beta)L^2}$ , we have

$$
\mathbb{E}[\|\theta_k^{[1]} - \theta_k^{[2]}\|^2] \le 4(1 - \mu\alpha)^{k - \tau} \mathbb{E}[\|\theta_0^{[1]} - \theta_0^{[2]}\|^2].
$$

Then, still by the non-expansion of  $\Pi_{B(\beta)}$  with respect to  $\|\cdot\|$ , the rest of the proof simply follows the same proof for non-projected SA. As such, we have proven Theorem [4.1.](#page-5-0)

# D Proof of Corollary [4.4](#page-6-8)

In this section, we present the proof of Corollary [4.4.](#page-6-8)

Recall that by Theorem [4.3,](#page-5-4) we obtain for  $k \geq 2\tau$ ,

$$
W_2^2(\mathcal{L}(\theta_k), \nu_\alpha) \le (1 - \alpha \mu)^k \cdot s(\theta_0, \theta^*, \mu, L, R).
$$

By [\[Vil09](#page-12-14), Theorem 4.1], there exists a coupling between  $\theta_k$  and  $\theta_{\infty}$  such that

$$
W_2^2(\mathcal{L}(\theta_k), \nu_\alpha) = \mathbb{E}[\|\theta_k - \theta_\infty\|^2].
$$

Applying Jensen's inequality twice, we obtain that

$$
\|\mathbb{E}[\theta_k - \theta_\infty]\|^2 \le (\mathbb{E}[\|\theta_k - \theta_\infty\|])^2 \le \mathbb{E}[\|\theta_k - \theta_\infty\|^2] \le (1 - \alpha\mu)^k \cdot s(\theta_0, \theta^*, \mu, L, R).
$$

We thus have for all  $k \geq 2\tau$ ,

$$
\|\mathbb{E}[\theta_k] - \mathbb{E}[\theta_\infty]\| \le \mathbb{E}[\|\theta_k - \theta_\infty\|] \le (1 - \alpha \mu)^{k/2} \cdot s'(\theta_0, \theta^*, \mu, L, R).
$$

<span id="page-37-0"></span>For the second moment, we first note that

$$
\|\mathbb{E}[\theta_k \theta_k^{\top}] - \mathbb{E}[\theta_{\infty} \theta_{\infty}^{\top}]\|
$$
  
\n=  $\|\mathbb{E}[(\theta_k - \theta_{\infty})(\theta_k - \theta_{\infty})^{\top}] + \mathbb{E}[\theta_{\infty}(\theta_k - \theta_{\infty})^{\top}] + \mathbb{E}[(\theta_k - \theta_{\infty})\theta_{\infty}^{\top}]\|$   
\n $\leq \|\mathbb{E}[(\theta_k - \theta_{\infty})(\theta_k - \theta_{\infty})^{\top}]\| + \|\mathbb{E}[\theta_{\infty}(\theta_k - \theta_{\infty})^{\top}]\| + \|\mathbb{E}[(\theta_k - \theta_{\infty})\theta_{\infty}^{\top}]\|$   
\n $\leq \mathbb{E}[\|\theta_k - \theta_{\infty}\|^2]\| + 2\mathbb{E}[\|\theta_{\infty}\|\|\theta_k - \theta_{\infty}\|]$   
\n $\leq \mathbb{E}[\|\theta_k - \theta_{\infty}\|^2]\| + 2\sqrt{\mathbb{E}[\|\theta_{\infty}\|^2]\mathbb{E}[\|\theta_k - \theta_{\infty}\|^2]},$  (D.1)

where we apply Cauchy-Schwarz to obtain the last inequality.

Meanwhile, we have

$$
\mathbb{E}\left[\left\|\theta_k-\theta_\infty\right\|^2\right] \le (1-\alpha\mu)^k \cdot s(\theta_0,\theta^*,\mu,L,R) \quad \text{and} \quad \mathbb{E}\left[\left\|\theta_\infty\right\|^2\right] = \mathcal{O}(1).
$$

Substituting the above bounds into the right-hand side of inequality  $(D.1)$  yields

$$
\left\| \mathbb{E} \left[ \theta_k \theta_k^{\top} \right] - \mathbb{E} \left[ \theta_\infty \theta_\infty^{\top} \right] \right\| \leq (1 - \alpha \mu)^{k/2} \cdot s''(\theta_0, \theta^*, \mu, L, R).
$$

# E Proof of Corollary [4.5](#page-6-6)

In this section, we prove the CLT result.

*Proof.* Consider the following centered test function  $\bar{h}: \mathcal{X} \times \mathbb{R}^d \to \mathbb{R}^d$  defined as

<span id="page-37-1"></span>
$$
\bar{h}(x,\theta) = \theta - \mathbb{E}[\theta_{\infty}].
$$

To prove that the CLT for function  $\bar{h}$ , we need to verify the Maxwell-Woodroofe condition [\[MW00\]](#page-12-15), i.e.,

$$
\sum_{n=1}^{\infty} n^{-3/2} \Big\| \sum_{t=0}^{n-1} Q^t h \Big\|_{L^2(\bar{\nu})} < \infty,
$$

where Q denotes the transition kernel of the joint Markov chain. If we can show the following

<span id="page-37-2"></span>
$$
\left\| \sum_{t=0}^{n-1} Q^t h \right\|_{L^2(\bar{\nu})} = \mathcal{O}(n^r)
$$
 (E.1)

with  $r \in [0, 1/2)$ , then the Maxwell-Woodroofe condition is verified, as

$$
\sum_{n=1}^{\infty} n^{-3/2} \Big\| \sum_{t=0}^{n-1} Q^t h \Big\|_{L^2(\bar{\nu})} = \sum_{n=1}^{\infty} n^{-3/2} \mathcal{O}(n^r) < \infty.
$$

We now proceed to prove the desired order in [\(E.1\)](#page-37-2). For sufficiently large  $n \geq 2\tau_{\alpha}$ , we observe

$$
\Big\|\sum_{t=0}^{n-1} Q^t h\Big\|_{L^2(\bar{\nu})} = \mathbb{E}_{\bar{\nu}} \Big\|\sum_{t=0}^{n-1} Q^t h\Big\|_2 \le \sum_{t=0}^{n-1} \mathbb{E}_{\bar{\nu}} \|Q^t h\|_2
$$
  
= 
$$
\underbrace{\sum_{t=0}^{2\tau_\alpha-1} \mathbb{E}_{\bar{\nu}} \|Q^t h\|_2}_{T_1} + \underbrace{\sum_{t=2\tau_\alpha}^{n-1} \mathbb{E}_{\bar{\nu}} \|Q^t h\|_2}_{T_2}.
$$

We now show that both terms  $T_1$  and  $T_2$  are of order  $\mathcal{O}(1)$  with respect to the parameter n.

For  $T_1$ , since Q is a transition kernel, so its  $||Q||_{L^2(\mu)}$  operator norm equals to 1. Hence,  $T_1$  can be upper bounded as

$$
T_1 \le \tau_\alpha \mathbb{E}_{\bar{\mu}}[\|h(\theta, x)\|_2^2] = \tau_\alpha \operatorname{Tr}(\operatorname{Var}(\theta_\infty)) < C_1,
$$

<span id="page-38-0"></span>where the last inequality follows from  $Tr(Var(\theta_{\infty})) \leq \alpha \tau$  established in  $(F.4)$  and  $\alpha \tau_{\alpha}^2 \to 0$  by Definition [2.1.](#page-2-5) Before proceeding to analyze the summation in  $T_2$ , we first recall [\(C.2\)](#page-26-4), that for  $t \geq 2\tau_\alpha$ ,

$$
\bar{W}_2(\mathcal{L}(x_t,\theta_t),\bar{\nu})=\mathcal{O}((1-\alpha\mu)^{t/2}),
$$

which holds for any  $(x, \theta) \in \mathcal{X} \times \mathbb{R}^d$ . Hence, by the property of Wasserstein distance [\[Vil09](#page-12-14)], there always exists a coupling that attains the optimality, i.e.,

$$
\mathbb{E}_{\Gamma((x_t,\theta_t),\bar{\nu})}\Big[\|\theta_t-\theta'\|_2^2+\delta_0(x_t\neq x')\Big] = \mathcal{O}((1-\alpha\mu)^t).
$$

Making use of this relationship, we can therefore bound  $T_2$ ,

$$
T_2 = \sum_{t=\tau_\alpha}^{n-1} \mathbb{E}_{\bar{\nu}} ||Q^t h||_2 \le \sum_{t=\tau_\alpha}^{\infty} \mathbb{E}_{\bar{\nu}} ||Q^t h||_2 = \mathcal{O}\Big(\frac{1}{1 - (1 - \alpha \mu)^{1/2}}\Big) = \mathcal{O}(1),
$$

where the last  $\mathcal{O}(\cdot)$  is asymptotic in *n*.

Combining the analysis of  $T_1$  and  $T_2$ , we have shown the desired order in  $(E.1)$ . Therefore, the Maxwell-Woodroofe condition has been verified and we establish the CLT for averaged nonlinear iterates with constant stepsize and Markovian data.  $\Box$ 

### F Proofs under Minorization Condition

When assuming the perturbed continuous noise condition in Assumption [5,](#page-5-2) one takes the alternative route to prove weak convergence. This is achieved by establishing the satisfaction of both a minorization condition and a drift condition. In this section, we prove the weak convergence result in Theorem [4.3](#page-5-4) by following this alternative approach. The subsequent corollaries of weak convergence, namely the non-asymptotic convergence rate in Corollary [4.4](#page-6-8) and the Central Limit Theorem (CLT) in Corollary [4.5,](#page-6-6) also hold, and we will provide the proofs for these results as well.

### F.1 Proof of Theorem [4.3](#page-5-4)

In this section, we prove the weak convergence under Assumption  $5(a)$  $5(a)$ . The proof consists of two major steps. Firstly, built upon the MSE convergence established in Proposition [4.2,](#page-5-1) we derive a multi-state drift condition. Subsequently, we show that under the minorization condition, the Markov chain is  $(x_k, \theta_k)_{k>0}$  is  $\varphi$ -irreducible. Then, follow [\[MT09,](#page-12-7) Theorem 19.1.3], we can conclude that the Markov chain  $(x_k, \theta_k)_{k\geq 0}$  is geometrically ergodic.

For completeness, we include the Theorem 19.1.3 from [\[MT09](#page-12-7)] below.

**Theorem F.1.** Suppose that  $\Phi$  is a  $\varphi$ -irreducible chain on X, and let  $n(x)$  be a measurable function from  $\mathcal{X} \to \mathbb{Z}_+$ . The chain is geometrically ergodic if it is aperiodic and there exists some petite set C, a nonnegative function  $V \geq 1$  and bounded on C, and positive constants  $\lambda < 1$  and b satisfying

<span id="page-38-1"></span>
$$
\int P^{n(x)}(x, dy)V(y) \leq \lambda^{n(x)}[V(x) + b \mathbb{1}_C(x)].
$$
\n(F.1)

We note that the function  $n(x)$  can be interpreted as the number of steps we must wait, starting from any x, for the drift to become negative.

Step 1: Deriving the Drift Condition Given the iteration step

$$
\theta_{k+1} = \theta_k + \alpha(g(\theta_k, x_k) + \xi_{t+1}(\theta_k)),
$$

we have already shown the following convergence rate on the MSE in Proposition [4.2,](#page-5-1) that

$$
\mathbb{E}[\|\theta_k - \theta^*\|^2] \le c_{2,1} (1 - \alpha \mu)^k \|\theta_0 - \theta^*\|^2 + c_{2,2} \alpha \tau_\alpha \frac{L^2}{\mu}.
$$

Inspired by the MSE convergence bound, we define the Lyapunov function  $V: \mathcal{X} \times \mathbb{R}^d \to [1, \infty]$ 

<span id="page-39-1"></span>
$$
V(x, \theta) = \|\theta - \theta^*\|^2 + 1.
$$
 (F.2)

Therefore, the major goal in this step is to obtain the desired drift condition as shown in [\(F.1\)](#page-38-1).

Therefore, from the above MSE convergence rate, we first obtain that

<span id="page-39-0"></span>
$$
Q^{k}V(x_0, \theta_0) \le c_{2,1}(1 - \alpha \mu)^k V(x_0, \theta_0) + c_{2,2} \frac{L^2}{\mu} \alpha \tau + 1.
$$
 (F.3)

Consider  $k = \min\{t \ge 0 : c_{2,1}(1 - \alpha \mu)^t < 1\}$ . Then, set  $\eta = 8(1 - \alpha \mu)^k$ ,  $\beta = \frac{1 - \eta}{2}$ , and  $m = c_{2,2} \frac{L^2}{\mu}$  $\frac{L}{\mu}\alpha\tau+1,$ and we consider the following bounded sublevel set,

$$
C_{\Theta} = \{ \theta : (\|\theta - \theta^*\|^2 + 1) \le m/\beta \}.
$$

From [\(F.3\)](#page-39-0), we derive that

$$
Q^{k}V(x_0, \theta_0) - V(x_0, \theta_0) \le -\beta V(x_0, \theta_0) + m \mathbb{1}_{\bar{C}}(x_0, \theta_0),
$$

where  $\overline{C} = \mathcal{X} \times C_{\Theta}$ . Rewriting the above Lyapunov drift condition, with  $b = m/(1 - \beta)$ , we obtain

$$
Q^k V(x_0, \theta_0) \le (1 - \beta) \Big( V(x_0, \theta_0) + b \mathbb{1}_{\bar{C}}(x_0, \theta_0) \Big).
$$

Setting  $\lambda$  such that  $\lambda^k = 1 - \beta$ , we have

$$
Q^k V(x_0, \theta_0) \leq \lambda^k \Big( V(x_0, \theta_0) + b \mathbb{1}_{\bar{C}}(x_0, \theta_0) \Big),
$$

which gives the desired multi-step drift condition.

Step 2: Proving the Minorization Condition Now that we have established the desired multi-step drift condition, it remains for us to show that  $\tilde{C}$  is accessible, small, and aperiodic.

Under this setup of  $\xi_t(\theta)$  in Assumption [5\(](#page-5-2)a), it is straightforward to verify the accessibility of  $\overline{C}$ . For any  $(x, \theta) \in \mathcal{X} \times \mathbb{R}^d$ , we have

$$
Q((x,\theta),\bar{C}) = \mathbb{P}((x',\theta') \in \mathcal{X} \times C_{\Theta} | (x,\theta))
$$
  
=  $\mathbb{P}(\theta' \in C_{\Theta} | (x,\theta)) \ge \int_{\theta' \in C_{\Theta}} \frac{1}{\alpha^d} p_{\theta} \left( \frac{\theta' - \theta}{\alpha} - g(x,\theta) \right) d\theta' > 0.$ 

As such, we have shown that  $\overline{C}$  is accessible.

Assuming  $\bar{C}$  is small, we can directly conclude aperiodicity following the definition of period of an accessible small set,  $d(C) = g.c.d.\Big\{n \in \mathbb{N}^* : \inf_{x \in C} P^n(x, C) > 0\Big\}.$ 

Therefore, what remains to show is that  $\overline{C}$  is small. For  $(x, \theta) \in \overline{C}$  and  $\overline{A} \in \mathcal{B}(\mathcal{X}) \times \mathcal{B}(\mathbb{R}^d)$ , we define the following projection sets,

$$
A_x = \{ \theta \in \mathbb{R}^d | (x, \theta) \in \overline{A} \} \text{ and } A^{\theta} = \{ x \in \mathcal{X} | (x, \theta) \in \overline{A} \}.
$$

Therefore,

$$
Q^{m}((x,\theta),\bar{A}) = \int_{\{(x^{(k)},\theta^{(k)})\}_{k=1}^{m-1}} \mathbb{P}((x_{m},\theta_{m}) \in \bar{A} | (x_{m-1},\theta_{m-1}) = (x^{(m-1)},\theta^{(m-1)}))
$$
  

$$
\cdots \mathbb{P}((x_{1},\theta_{1}) = d(x^{(1)},\theta^{(1)}) | (x_{0},\theta_{0}) = (x,\theta))
$$
  

$$
\geq \int_{\{(x^{(k)},\theta^{(k)})\}_{k=1}^{m-1}} \mathbb{P}((x_{m},\theta_{m}) \in \bar{A} | (x_{m-1},\theta_{m-1}) = (x^{(m-1)},\theta^{(m-1)}))
$$
  

$$
\cdots \mathbb{P}((x_{1},\theta_{1}) = d(x^{(1)},\theta^{(1)}) | (x_{0},\theta_{0}) = (x,\theta))
$$

$$
\begin{split}\n&= \int_{\{(x^{(k)},\theta^{(k)})\}_{k=1}^{m-1}} \Big( \int_{x' \in \mathcal{X}} \mathbb{P}(x_m = dx'|x_{m-1} = x^{(m-1)}) \mathbb{P}(\theta_m \in A_{x'}|(x_{m-1}, \theta_{m-1}) = (x^{(m-1)}, \theta^{(m-1)})) \Big) \\
& \mathbb{P}(x_{m-1} = dx^{(m-1)}|x_{m-2} = x^{(m-2)}) \mathbb{P}(\theta_{m-1} = d\theta^{(m-1)}|(x_{m-2}, \theta_{m-2}) = (x^{(m-2)}, \theta^{(m-2)})) \Big) \\
&\dots \\
&\mathbb{P}(x_1 = dx^{(1)}|x_0 = x) \mathbb{P}(\theta_1 = d\theta^{(1)}|(x_0, \theta_0) = (x, \theta)) \\
&= \int_{x' \in \mathcal{X}} \int_{\{x^{(k)}\}_{k=1}^{m-1}} \mathbb{P}(x_m = dx'|x_{m-1} = x^{(m-1)}) \mathbb{P}(x_{m-1} = dx^{(m-1)}|x_{m-2} = x^{(m-2)}) \cdots \mathbb{P}(x_1 = dx^{(1)}|x_0 = x) \\
&\quad \left( \int_{\{\theta^{(k)}\}_{k=1}^{m-1}} \mathbb{P}(\theta_m \in A_{x'}|(x_{m-1}, \theta_{m-1}) = (x^{(m-1)}, \theta^{(m-1)})) \right. \\
&\quad \mathbb{P}(\theta_{m-1} = d\theta^{(m-1)}|(x_{m-2}, \theta_{m-2}) = (x^{(m-2)}, \theta^{(m-2)})) \Big) \\
&\dots \mathbb{P}(\theta_1 = d\theta^{(1)}|(x_0, \theta_0) = (x, \theta))\Big). \n\end{split}
$$

Next, for  $(x, \theta) \in \overline{C}$ , we observe that

$$
\mathbb{P}(\theta_k = d\theta' | (x_{k-1}, \theta_{k-1}) = (x, \theta)) = \mathbb{P}\Big(\xi_k(\theta) = d\Big(\frac{\theta' - \theta}{\alpha} - g(x, \theta)\Big)| (x_k, \theta_k) = (x, \theta)\Big)
$$

$$
\geq \frac{1}{\alpha^d} p_\theta \Big(\frac{\theta' - \theta}{\alpha} - g(x, \theta)\Big) d\theta'
$$

$$
\geq \frac{1}{\alpha^d} \inf_{\hat{\theta} \in C_{\Theta}} p_{\hat{\theta}} \Big(\frac{\theta' - \theta}{\alpha} - g(x, \theta)\Big) d\theta'.
$$

We next recall the linear growth assumption in Assumption [2](#page-2-2) that  $||g(x, \theta)|| \le L(||\theta|| + 1)$ . Hence, given  $\theta\in C_{\Theta},\,\forall x\in\mathcal{X},$  we have

$$
\|\theta + \alpha g(x,\theta)\| \le \|\theta\| + \alpha \|g(x,\theta)\| \le (1 + \alpha L)(\|\theta\| + 1) \le (1 + \alpha L)(\sqrt{M/\beta} + \|\theta^*\| + 1) \le B,
$$

for some bounded value B.

We now define the measure  $\varsigma^{\dagger}$  on  $\mathbb{R}^{d}$  as  $\varsigma^{\dagger}(A) = \frac{1}{\alpha^{d}} \inf_{\|z\| \leq B} \int_{\theta' \in A \cap C_{\Theta}} \inf_{\hat{\theta} \in C_{\Theta}} p_{\hat{\theta}} \left( \frac{\theta' - z}{\alpha} \right)$  $\partial d\theta'$ . Hence, it is easy to see that  $\zeta^{\dagger}(C_{\Theta}^{c})=0$ . Moreover, we note the following property of measure  $\zeta^{\dagger}$ .

<span id="page-40-0"></span>**Claim 1.** For  $A \subseteq C_{\Theta}$  and  $\lambda(A) > 0$ ,  $\zeta^{\dagger}(A) > 0$ , where  $\lambda$  denotes the Lebesgue measure.

We delay the proof to the end. Taking the claim as true, we derive that

$$
\int_{\{\theta^{(k)}\}_{k=1}^{m-1}} \mathbb{P}(\theta_m \in A_{x'}|(x_{m-1}, \theta_{m-1}) = (x^{(m-1)}, \theta^{(m-1)}))
$$
\n
$$
\mathbb{P}(\theta_{m-1} = d\theta^{(m-1)}|(x_{m-2}, \theta_{m-2}) = (x^{(m-2)}, \theta^{(m-2)})) \cdots \mathbb{P}(\theta_1 = d\theta^{(1)}|(x_0, \theta_0) = (x, \theta))
$$
\n
$$
= \int_{\theta^{(m)} \in A_{x'}} \int_{\theta^{(m-1)} \in C_{\Theta}} \mathbb{P}(\theta_m d\theta^{(m)}|(x_{m-1}, \theta_{m-1}) = (x^{(m-1)}, \theta^{(m-1)}))
$$
\n
$$
\int_{\theta^{(m-2)} \in C_{\Theta}} \mathbb{P}(\theta_{m-1} = d\theta^{(m-1)}|(x_{m-2}, \theta_{m-2}) = (x^{(m-2)}, \theta^{(m-2)}))
$$
\n
$$
\cdots
$$
\n
$$
\int_{\theta^{(1)} \in C_{\Theta}} \mathbb{P}(\theta_2 = d\theta^{(2)}|(x_1, \theta_1) = (x^{(1)}, \theta^{(1)})) \mathbb{P}(\theta_1 = d\theta^{(1)}|(x_0, \theta_0) = (x, \theta))
$$

$$
\geq \int_{\theta^{(m)} \in A_{x'}} \int_{\theta^{(m-1)}} \frac{1}{cC_{\Theta}} \frac{1}{\alpha^{d}} \inf_{\hat{\theta} \in C_{\Theta}} p_{\hat{\theta}} \left( \frac{\theta^{(m)} - \theta^{(m-1)}}{\alpha} - g(x^{(m-1)}, \theta^{(m-1)}) \right) d\theta^{(m)} \n\int_{\theta^{(m-2)} \in C_{\Theta}} \mathbb{P}(\theta_{m-1} = d\theta^{(m-1)} | (x_{m-2}, \theta_{m-2}) = (x^{(m-2)}, \theta^{(m-2)})) \right) \n\cdots \n\int_{\theta^{(1)} \in C_{\Theta}} \mathbb{P}(\theta_{2} = d\theta^{(2)} | (x_{1}, \theta_{1}) = (x^{(1)}, \theta^{(1)}) \mathbb{P}(\theta_{1} = d\theta^{(1)} | (x_{0}, \theta_{0}) = (x, \theta)) \n\geq \inf_{(\tilde{x}, \tilde{\theta}) \in \tilde{C}} \int_{\theta^{(m)} \in A_{x'}} \int_{\theta^{(m-1)} \in C_{\Theta}} \frac{1}{\alpha^{d}} \inf_{\tilde{\theta} \in C_{\Theta}} p_{\hat{\theta}} \left( \frac{\theta^{(m)} - \tilde{\theta}}{\alpha} - g(\tilde{x}, \tilde{\theta}) \right) d\theta^{(m)} \n\int_{\theta^{(m-2)} \in C_{\Theta}} \mathbb{P}(\theta_{m-1} = d\theta^{(m-1)} | (x_{m-2}, \theta_{m-2}) = (x^{(m-2)}, \theta^{(m-2)})) \right) \n\cdots \n\int_{\theta^{(1)} \in C_{\Theta}} \mathbb{P}(\theta_{2} = d\theta^{(2)} | (x_{1}, \theta_{1}) = (x^{(1)}, \theta^{(1)}) \mathbb{P}(\theta_{1} = d\theta^{(1)} | (x_{0}, \theta_{0}) = (x, \theta)) \n\geq \inf_{\|z\| \leq B} \int_{\theta^{(m)} \in A_{x'}} \int_{\theta^{(m-1)} \in C_{\Theta}} \frac{1}{\alpha^{d}} \inf_{\hat{\theta} \in C_{\Theta}} p_{\hat{\theta}} \left( \frac{\theta^{(m)} - z}{\alpha} \right) d\theta^{(m)} \n\int_{\theta^{(
$$

$$
\geq \varsigma^{\dagger} (A_{x^{\prime\prime}}) \varsigma^{\dagger} (C_{\Theta})^{m-1}.
$$

Therefore, by combining all the analyses, we obtain

$$
Q^{m}((x,\theta),\bar{A})
$$
\n
$$
\geq \int_{x' \in \mathcal{X}} \int_{\{x^{(k)}\}_{k=1}^{m-1}} \mathbb{P}(x_{m} = dx'|x_{m-1} = x^{(m-1)}) \mathbb{P}(x_{m-1} = dx^{(m-1)}|x_{m-2} = x^{(m-2)})
$$
\n
$$
\cdots \mathbb{P}(x_{1} = dx^{(1)}|x_{0} = x)
$$
\n
$$
\left( \int_{\{\theta^{(k)}\}_{k=1}^{m-1}} \mathbb{P}(\theta_{m} \in A_{x'}|(x_{m-1}, \theta_{m-1}) = (x^{(m-1)}, \theta^{(m-1)}))
$$
\n
$$
\mathbb{P}(\theta_{m-1} = d\theta^{(m-1)}|(x_{m-2}, \theta_{m-2}) = (x^{(m-2)}, \theta^{(m-2)})) \right)
$$
\n
$$
\cdots \mathbb{P}(\theta_{1} = d\theta^{(1)}|(x_{0}, \theta_{0}) = (x, \theta))
$$
\n
$$
\geq \zeta^{\dagger}(C_{\Theta})^{m-1} \int_{x' \in \mathcal{X}} \zeta^{\dagger}(A_{x'})
$$
\n
$$
\int_{\{x^{(k)}\}_{k=1}^{m-1}} \mathbb{P}(x_{m} = dx'|x_{m-1} = x^{(m-1)}) \cdots \mathbb{P}(x_{1} = dx^{(1)}|x_{0} = x)
$$
\n
$$
\geq \zeta^{\dagger}(C_{\Theta})^{m-1} \int_{x' \in \mathcal{X}} \zeta^{\dagger}(A_{x'})\phi(dx')
$$

$$
= \zeta \cdot (\phi \times \zeta^{\dagger})(\bar{A}),
$$

<span id="page-42-0"></span>where  $\zeta = \zeta^{\dagger} (C_{\Theta})^{m-1}$  and  $\phi \times \zeta^{\dagger}$  being the unique induced product measure on  $\mathcal{X} \times \mathbb{R}^d$ .

As such, we have proven that  $\overline{C}$  is  $(m, \phi \times \varsigma^{\dagger})$ -small, and hence  $(\delta_m, \phi \times \varsigma^{\dagger})$ -petite, and subsequently shown that  $(x_k, \theta_k)_{k>0}$  is geometrically ergodic.

By [\[MT09](#page-12-7), Theorem 16.0.1 (iv)], we can further conclude that the geometrically ergodic  $(\theta_t, x_t)_{t>0}$  is also V-uniformly ergodic with the same V as defined in  $(F.2)$ . Therefore, we have the following convergence rate in the V-norm  $\left\|\mathcal{L}(x_k,\theta_k), -\bar{\nu}_{\alpha}\right\|_V \leq \kappa \rho^k$ , where  $\kappa$  and  $\rho$  implicitly depend on the stepsize  $\alpha$ .

Lastly, we provide the proof of Claim [1.](#page-40-0)

*Proof.* We first recall that for any set  $B \subseteq \mathbb{R}^d$  such that  $\lambda(B) > 0$ , where  $\lambda$  refers to the Lebesgue measure, then for  $\theta \in C_{\Theta}$ , we know that  $\int_{t \in B} p_{\theta}(t) dt \ge \int_{t \in B} \inf_{\theta \in C_{\Theta}} p_{\theta}(t) dt > 0$ . Moreover, for a given (translation)  $z \in \mathbb{R}^d$  and  $\theta \in C_{\Theta}$ , we have

$$
\int_{t\in B} p_{\theta}(t-z)dt = \int_{t'\in B-z} p_{\theta}(t')dt' \ge \int_{t'\in B-z} \inf_{\theta\in C} p_{\theta}(t')dt' > 0.
$$

Following the properties stated above, we define  $h(z) = \int_{t \in B} \inf_{\theta \in C} p_{\theta}(t - z) dt$ , and it is easy to verify that  $h(z)$  is a continuous function. Hence, for a bounded set  $\overline{D}$ , we have  $\inf_{z\in D}h(z) > 0$ . Subsequently, we define the measure  $\varsigma^{\dagger}$  induced by h and we verify that

$$
\varsigma^{\dagger}(A) = \inf_{\|z\| \le B} \int_{t \in A \cap C_{\Theta}} \inf_{\theta \in C} p_{\theta}(t - z) dt > 0.
$$

 $\Box$ 

### F.2 Proof of Corollary [4.4](#page-6-8)

As shown in the previous section, the joint process  $(x_k, \theta_k)_{k>0}$  is V-uniformly ergodic and hence also exhibits a geometric non-asymptotic convergence rate under the V -weighted norm. Subsequently, this corresponds to a version of Corollary [4.4](#page-6-8) resulting in a different set of convergence rate coefficients. Thus, we restate Corollary [4.4](#page-6-8) in the context of the minorization setting and provide the proof below.

**Corollary F.2** (Non-Asymptotic Convergence Rate). For any initialization of  $\theta_0 \in \mathbb{R}^d$ , under the setting of Theorem  $\angle 4.3$ , we have

$$
\left\| \mathbb{E}[\theta_k] - \mathbb{E}[\theta_{\infty}^{(\alpha)}] \right\| \leq \kappa \cdot \rho^k \cdot s'(\theta_0, L, \mu), \quad \text{and} \quad \left\| \mathbb{E}[\theta_k \theta_k^{\top}] - \mathbb{E}[\theta_{\infty}^{(\alpha)}(\theta_{\infty}^{(\alpha)})^{\top}] \right\| \leq \kappa \cdot \rho^k \cdot s''(\theta_0, L, \mu),
$$

where  $\kappa$  and  $\rho$  are defined in [\(4.1\)](#page-5-5) and implicitly depend on  $\alpha$ .

*Proof.* For V-uniformly ergodic Markov chain  $(x_k, \theta_k)_{k \geq 0}$ , when functions  $f : \mathcal{X} \times \mathbb{R}^d \to \mathbb{R}^d$  is dominated by the Lyapunov function, i.e.,  $||f|| \leq V$ , it enjoys the following convergence property,

$$
||Q^n f(x_0, \theta_0) - \pi f|| \le \kappa \rho^n V(\theta_0).
$$

Consider test function  $f(\theta, x) = \theta - \theta^*$ . It is easy to see that  $||f|| \leq V$ . Hence, we obtain

$$
\|\mathbb{E}[\theta_n] - \mathbb{E}[\theta_\infty]\| = \|Q^n f(x, \theta) - \pi f\| \le \kappa \rho^n V(\theta_0).
$$

Next, consider  $f'(x) = (\theta - \theta^*)(\theta - \theta^*)^\top$ . Clearly,  $||f'|| \leq V$ . Therefore,

$$
\|\mathbb{E}[(\theta_t - \theta^*)(\theta_t - \theta^*)^{\top}] - \mathbb{E}[(\theta_{\infty} - \theta^*)(\theta_{\infty} - \theta^*)^{\top}]\| \le \kappa \rho^n V(\theta_0).
$$

For the LHS, we have

$$
\|\mathbb{E}[(\theta_t - \theta^*)(\theta_t - \theta^*)^\top] - \mathbb{E}[(\theta_\infty - \theta^*)(\theta_\infty - \theta^*)^\top]\|
$$
  
= 
$$
\|\mathbb{E}[\theta_t \theta_t^\top] - \mathbb{E}[\theta_\infty \theta_\infty^\top] - \mathbb{E}[\theta_t - \theta_\infty](\theta^*)^\top - \theta^* \mathbb{E}[(\theta_t - \theta_\infty)^\top]\|
$$
  

$$
\geq \|\mathbb{E}[\theta_t \theta_t^\top] - \mathbb{E}[\theta_\infty \theta_\infty^\top]\| - 2\|\mathbb{E}[\theta_t - \theta_\infty]\|\|\theta^*\|.
$$

Subsequently,

$$
\|\mathbb{E}[\theta_t \theta_t^\top] - \mathbb{E}[\theta_\infty \theta_\infty^\top] \| \leq (2\|\theta^*\| + 1)\kappa \rho^n V(\theta_0).
$$



<span id="page-43-1"></span>The above results imply the convergence of the first two moments. Moreover, we conclude that

$$
\text{Var}(\theta_{\infty}) = \text{Var}(\theta_{\infty} - \theta^*) \le \mathbb{E}[\|\theta_{\infty} - \theta^*\|^2] = \lim_{t \to \infty} \mathbb{E}[\|\theta_t - \theta^*\|^2] \lesssim \alpha \tau.
$$

Additionally,

<span id="page-43-2"></span>
$$
\mathbb{E}[\|\theta_{\infty} - \theta^*\|]^2 \le \mathbb{E}[\|\theta_{\infty} - \theta^*\|^2] \lesssim \alpha \tau.
$$
 (F.4)

### F.3 Proof of Corollary [4.5](#page-6-6)

After establishing the V-uniform ergodicity of the joint process  $(x_k, \theta_k)_{k>0}$ , the central limit theorem for averaged iterates follows as a straightforward consequence.

*Proof.* For any test function  $h: \mathcal{X} \times \mathbb{R}^d \to \mathbb{R}^d$  that satisfies  $||h||^2 \leq V$ , by Theorem 17.0.1 in [\[MT09\]](#page-12-7), it has the following CLT results,

$$
\frac{1}{\sqrt{k}} \Big[ \sum_{t=0}^{k-1} \left( h_t - \mathbb{E}[h_\infty] \right) \Big] \Rightarrow \mathcal{N}(0, \Sigma^{(a)}).
$$

Therefore, consider  $h(x, \theta) = \theta - \theta^*$ , it is easy to see that  $||h||^2 \leq V$ , and hence we naturally obtain the  $\frac{1}{k} \left[ \sum_{t=0}^{k-1} (\theta_t - \mathbb{E}[\theta_\infty]) \right] \Rightarrow \mathcal{N}(0, \Sigma^{(a)}) \text{ as } k \to \infty.$ desired CLT result,  $\frac{1}{\sqrt{2}}$  $\Box$ 

# <span id="page-43-0"></span>G Proof of Theorem [4.6](#page-6-0)

We now provide the proof of Theorem [4.6](#page-6-0) on characterizing the asymptotic bias of nonlinear SA.

### G.1 BAR and Preliminaries

The proof utilizes the basic adjoint relationship (BAR) approach to study the stationary distribution

<span id="page-43-7"></span><span id="page-43-6"></span><span id="page-43-3"></span>
$$
\mathbb{E}_{\bar{\nu}}[(P-I)h(\theta,x)]=0,
$$

via carefully designed test functions h. We refer readers to  $[HCX23a]$  for the derivation of the following properties of Markovian SA at stationarity,

$$
\mathbb{E}[\mathbb{1}\{\theta_{\infty}\in S\} \mid x_{\infty}](x) = \mathbb{E}[\mathbb{1}\{\theta_{\infty+1}\in S\} \mid x_{\infty+1}](x),\tag{G.1}
$$

$$
\mathbb{E}[\theta_{\infty} \mid x_{\infty}](x) = \mathbb{E}[\theta_{\infty+1} \mid x_{\infty+1}](x), \tag{G.2}
$$

$$
\mathbb{E}[(\theta_{\infty})^{\otimes 2} \mid x_{\infty}](x) = \mathbb{E}[(\theta_{\infty+1})^{\otimes 2} \mid x_{\infty+1}](x). \tag{G.3}
$$

Following the Borel state space assumption in [1,](#page-2-0)  $\mathbb{E}[\mathbb{1}\{\theta_\infty \in \cdot\}|x_\infty = x]$  induce a regular conditional probability measure, which we denote as  $\tilde{\nu}(\cdot, x_{\infty} = x)$ , and hence [\(G.1\)](#page-43-3) can be reformulated as

 $\tilde{\nu}(\theta_{\infty} \in S, x_{\infty} = x) = \tilde{\nu}(\theta_{\infty+1} \in S, x_{\infty+1} = x).$ 

Before proceeding to the proof, we introduce the following shorthands and notations. For  $x \in \mathcal{X}$ ,

<span id="page-43-5"></span><span id="page-43-4"></span>
$$
z_i(x) := \mathbb{E}[(\theta_{\infty} - \theta^*)^{\otimes i}|x_{\infty} = x],
$$
  
\n
$$
\delta_i(x) := z_i(x) - \pi z_i,
$$

Following the differentiability assumption of g in Assumption [3,](#page-2-3) and we can apply Taylor expansion to g and we note the following notation on residuals.

$$
g(\theta, x) = g(\theta^*, x) + g'(\theta^*, x)(\theta - \theta^*) + \frac{1}{2}g''(\theta^*, x)(\theta - \theta^*)^{\otimes 2} + R_3(\theta, x)
$$
(G.4)

$$
= g(\theta^*, x) + g'(\theta^*, x)(\theta - \theta^*) + R_2(\theta, x).
$$
 (G.5)

By Assumption [3](#page-2-3) and results from Proposition [4.2](#page-5-1) and [4.2,](#page-5-1) we note that the residual  $R_n(\theta, x)$  satisfies

$$
\sup_{x \in \mathcal{X}, \theta \in \mathbb{R}^d} \left\{ \|R_n(\theta, x)\| / \|\theta - \theta^*\|^n \right\} < +\infty.
$$

Hence, we have

$$
||R_n(\theta, x_{\infty})||_{L^2(\pi)} \lesssim \mathbb{E}[||\theta_{\infty} - \theta^*||^n] = \mathcal{O}((\alpha\tau)^{n/2}), \quad n = 2, 3, 4.
$$

Lastly, we denote

$$
\begin{aligned}\n\bar{g}(\theta) &:= \mathbb{E}_{x \sim \pi}[g(\theta, x)], \quad \bar{g}_2(\theta) := \mathbb{E}_{x \sim \pi}[(g(\theta, x))^{\otimes 2}] \\
\bar{g}^{(1)}(\theta) &:= \mathbb{E}_{x \sim \pi}[g'(\theta, x)], \quad \bar{g}_2^{(1)}(\theta) := \mathbb{E}_{x \sim \pi}[(g'(\theta, x))^{\otimes 2}], \quad \bar{g}^{(2)}(\theta) := \mathbb{E}_{x \sim \pi}[g''(\theta, x)].\n\end{aligned}
$$

We are now ready to present our proof. The proof consists of two major steps. For the complexity of this problem, we first set aside the projection constraint and focus on the BAR analysis, and we shall present the asymptotic bias characterization without the projection step. The analysis in this step thus shall work for the minorization proof technique as well. Then, in the second step, we elaborate on the impact brought along by the projection analysis and conclude our proof.

#### <span id="page-44-6"></span>G.2 Step 1: Bias Characterization without Projection

### G.2.1 Step 1: First Moment Analysis

Consider test function  $h_1(x, \theta) = \theta - \theta^*$ . Therefore, we first have

$$
\mathbb{E}[\theta_{\infty+1} - \theta^*] = \mathbb{E}[\theta_{\infty} - \theta^*] + \alpha \Big( \mathbb{E}[g(\theta_{\infty}, x_{\infty})] + \mathbb{E}[\xi_{\infty+1}(\theta_{\infty})] \Big),
$$

which immediately implies that

<span id="page-44-4"></span><span id="page-44-3"></span><span id="page-44-2"></span><span id="page-44-1"></span><span id="page-44-0"></span>
$$
0 = \mathbb{E}[g(\theta_{\infty}, x_{\infty})].
$$
 (G.6)

Substituting the Taylor expansion  $(G.4)$  back into  $(G.6)$ , we have

$$
0 = \mathbb{E}[g(\theta^*, x)] + \mathbb{E}[g'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*)] + \frac{1}{2} \mathbb{E}[g''(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*)^{\otimes 2}] + \mathbb{E}[R_3(\theta_{\infty}, x_{\infty})]
$$
  
=  $\mathbb{E}[g'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*)] + \frac{1}{2} \mathbb{E}[g''(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*)^{\otimes 2}] + \mathcal{O}((\alpha \tau)^{3/2}),$  (G.7)

where we make use of  $\mathbb{E}[g(\theta^*,x)] = \bar{g}(\theta^*) = 0$  by definition and the order or  $\mathbb{E}[R_3(\theta_\infty, x_\infty)] = \mathcal{O}((\alpha\tau)^{3/2})$  to obtain the second equality.

Next, we proceed to analyze the two terms in  $(G.7)$ . For the first term, we have

$$
\mathbb{E}[g'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*)] = \mathbb{E}[g'(\theta^*, x_{\infty})z_1(x_{\infty})]
$$
  
= 
$$
\mathbb{E}[g'(\theta^*, x_{\infty})\delta_1(x_{\infty})] + \bar{g}^{(1)}(\theta^*)\mathbb{E}[\theta_{\infty} - \theta^*].
$$
 (G.8)

We now move on to analyze the second term, and obtain

$$
\mathbb{E}[g''(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*)^{\otimes 2}] = \mathbb{E}[g''(\theta^*, x_{\infty})z_2(x_{\infty})]
$$
  
= 
$$
\mathbb{E}[g''(\theta^*, x_{\infty})\delta_2(x_{\infty})] + \bar{g}^{(2)}(\theta^*)\mathbb{E}[(\theta_{\infty} - \theta^*)^{\otimes 2}].
$$
 (G.9)

Hence, substituting  $(G.8)$  and  $(G.9)$  back into  $(G.7)$ , we reorganize the terms and arrive at

$$
\mathbb{E}[\theta_{\infty} - \theta^*]
$$
  
= -(\bar{g}^{(1)}(\theta^\*))^{-1} \Big( \mathbb{E}[g'(\theta^\*, x\_{\infty})\delta\_1(x\_{\infty})] \qquad (G.10)

<span id="page-44-5"></span>+ 
$$
\frac{1}{2}
$$
 $\left( \mathbb{E}[g''(\theta^*, x_\infty)\delta_2(x_\infty)] + \bar{g}^{(2)}(\theta^*) \mathbb{E}[(\theta_\infty - \theta^*)^{\otimes 2}] \right) \right)$  (G.11)  
+  $\mathcal{O}((\alpha\tau)^{3/2}).$ 

To obtain a refined characterization of the asymptotic bias, we carefully analyze the remaining three terms in  $(G.10)$ – $(G.11)$ . We focus on each term in the next three sections respectively.

#### G.2.2 Step 2: Second Moment Analysis

We start with analyzing  $\mathbb{E}[(\theta_{\infty} - \theta^*)^{\otimes 2}]$  in this section.

Following the BAR approach, we consider the test function  $h_2(x, \theta) = (\theta - \theta^*)^{\otimes 2}$  and obtain

$$
\mathbb{E}[(\theta_{\infty+1} - \theta^*)^{\otimes 2}] = \mathbb{E}[(\theta_{\infty} - \theta^* + \alpha(g(\theta_{\infty}, x_{\infty}) + \xi_{\infty+1}(\theta_{\infty}))^{\otimes 2}] \n= \mathbb{E}[(\theta_{\infty} - \theta^*)^{\otimes 2}] + \alpha^2 (\mathbb{E}[(g(\theta_{\infty}, x_{\infty}))^{\otimes 2}] + \mathbb{E}[(\xi_{\infty+1}(\theta_{\infty}))^{\otimes 2}]) \n+ \alpha (\mathbb{E}[g(\theta_{\infty}, x_{\infty}) \otimes (\theta_{\infty} - \theta^*)] + \mathbb{E}[(\theta_{\infty} - \theta^*) \otimes g(\theta_{\infty}, x_{\infty})]).
$$

Simplifying the above expression, we have

<span id="page-45-3"></span><span id="page-45-1"></span><span id="page-45-0"></span>
$$
0 = \alpha \left( \mathbb{E}[(g(\theta_{\infty}, x_{\infty}))^{\otimes 2}] + \mathbb{E}[(\xi_{\infty+1}(\theta_{\infty}))^{\otimes 2}] \right) + \left( \mathbb{E}[(\theta_{\infty} - \theta^*) \otimes g(\theta_{\infty}, x_{\infty})] + \mathbb{E}[g(\theta_{\infty}, x_{\infty}) \otimes (\theta_{\infty} - \theta^*)] \right).
$$
 (G.12)

We adopt a similar approach in analyzing the above relationship that contains  $g(\theta_{\infty}, x_{\infty})$  as in the previous step. We make use of the Taylor expansion of g at  $\theta^*$  but at a lower order. We substitute the Taylor expansion  $(G.5)$  into  $(G.12)$  and obtain

$$
0 = \alpha \mathbb{E}[(g(\theta^*, x_{\infty}) + g'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*) + R_2(\theta_{\infty}, x_{\infty}))^{\otimes 2}] + \alpha \mathbb{E}[(\xi_{\infty+1}(\theta_{\infty}))^{\otimes 2}] + \mathbb{E}[(g(\theta^*, x_{\infty}) + g'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*) + R_2(\theta_{\infty}, x_{\infty})) \otimes (\theta_{\infty} - \theta^*)] + \mathbb{E}[(\theta_{\infty} - \theta^*) \otimes (g(\theta^*, x_{\infty}) + g'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*) + R_2(\theta_{\infty}, x_{\infty}))] = \mathbb{E}[g(\theta^*, x_{\infty}) \otimes (\theta_{\infty} - \theta^*)] + \mathbb{E}[(\theta_{\infty} - \theta^*) \otimes g(\theta^*, x_{\infty})] + \mathbb{E}[(\alpha'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*))] \otimes (\theta_{\infty} - \theta^*)] + \mathbb{E}[(\theta_{\infty} - \theta^*) \otimes (\alpha'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*))]
$$
(G.13)

+ 
$$
\mathbb{E}[(g'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*)) \otimes (\theta_{\infty} - \theta^*)] + \mathbb{E}[(\theta_{\infty} - \theta^*) \otimes (g'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*))]
$$
 (G.14)  
+  $\alpha \mathbb{E}[(g'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*))^{\otimes 2}]$  (G.15)

+ 
$$
\alpha \mathbb{E}[g(\theta^*, x_{\infty}) \otimes (g'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*))] + \alpha \mathbb{E}[(g'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*)) \otimes g(\theta^*, x_{\infty})]
$$
 (G.16)

+ 
$$
\mathbb{E}[R_2(\theta_{\infty}, x_{\infty}) \otimes (\theta_{\infty} - \theta^*)] + \mathbb{E}[(\theta_{\infty} - \theta^*) \otimes R_2(\theta_{\infty}, x_{\infty})]
$$
  
+  $\alpha \mathbb{E}[(g(\theta^*, x_{\infty}))^{\otimes 2}] + \alpha \mathbb{E}[(\xi_{\infty+1}(\theta_{\infty}))^{\otimes 2}] + \mathcal{O}(\alpha^2 \tau).$ 

Therefore, we proceed to analyze the terms in  $(G.13)$ – $(G.16)$ . Starting with the terms in [\(G.13\)](#page-45-1), we have

$$
\mathbb{E}[g(\theta^*, x_{\infty}) \otimes (\theta_{\infty} - \theta^*)] = \mathbb{E}[g(\theta^*, x_{\infty}) \otimes (\delta_1(x_{\infty}) + \pi z_1)]
$$
  
\n
$$
= \mathbb{E}[g(\theta^*, x_{\infty}) \otimes \delta_1(x_{\infty})] + \underbrace{\mathbb{E}[g(\theta^*, x_{\infty})]}_{=0} \otimes \mathbb{E}[\theta_{\infty} - \theta^*]
$$
  
\n
$$
= \mathbb{E}[g(\theta^*, x_{\infty}) \otimes \delta_1(x_{\infty})].
$$

Similarly,

<span id="page-45-4"></span><span id="page-45-2"></span>
$$
\mathbb{E}[(\theta_{\infty}-\theta^*)\otimes g(\theta^*,x_{\infty})]=\mathbb{E}[\delta_1(x_{\infty})\otimes g(\theta^*,x_{\infty})].
$$

Next, for the terms in  $(G.14)$ , we have

$$
\mathbb{E}[(g'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*)) \otimes (\theta_{\infty} - \theta^*)] = \mathbb{E}[g'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*)^{\otimes 2}]
$$
  
\n
$$
= \mathbb{E}[g'(\theta^*, x_{\infty})(\delta_2(x_{\infty}) + \pi z_2)]
$$
  
\n
$$
= \mathbb{E}[g'(\theta^*, x_{\infty})\delta_2(x_{\infty})] + \bar{g}^{(1)}(\theta^*)\mathbb{E}[(\theta_{\infty} - \theta^*)^{\otimes 2}].
$$

Similarly,

$$
\mathbb{E}[(\theta_{\infty}-\theta^*)\otimes (g'(\theta^*,x_{\infty})(\theta_{\infty}-\theta^*))]=\mathbb{E}[\delta_2(x_{\infty})g'(\theta^*,x_{\infty})]+\mathbb{E}[(\theta_{\infty}-\theta^*)^{\otimes2}]\bar{g}^{(1)}(\theta^*).
$$

Moving on to the second term in  $(G.15)$ 

$$
\mathbb{E}[(g'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*))^{\otimes 2}] = \mathbb{E}[g'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*)^{\otimes 2}g'(\theta^*, x_{\infty})]
$$
  
=  $\mathbb{E}[g'(\theta^*, x_{\infty})(\delta_2(x_{\infty}) + \pi z_2)g'(\theta^*, x_{\infty})]$ 

$$
= \mathbb{E}[g'(\theta^*, x_{\infty})\delta_2(x_{\infty})g'(\theta^*, x_{\infty})] + \bar{g}_2^{(1)}(\theta^*)\mathbb{E}[(\theta_{\infty} - \theta^*)^{\otimes 2}].
$$

Last, for the terms in  $(G.16)$ 

$$
\mathbb{E}[g(\theta^*, x_{\infty}) \otimes (g'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*))] = \mathbb{E}[g(\theta^*, x_{\infty}) \otimes (g'(\theta^*, x_{\infty})(\delta_1(x_{\infty}) + \pi z_1))]
$$
  
\n
$$
= \mathbb{E}[g(\theta^*, x_{\infty}) \otimes (g'(\theta^*, x_{\infty})\delta_1(x_{\infty}))]
$$
  
\n
$$
+ \mathbb{E}[g(\theta^*, x_{\infty}) \otimes (g'(\theta^*, x_{\infty})\mathbb{E}[\theta_{\infty} - x_{\infty}])].
$$

Similarly,

$$
\mathbb{E}[(g'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*)) \otimes g(\theta^*, x_{\infty})] = \mathbb{E}[(g'(\theta^*, x_{\infty})\delta_1(x_{\infty})) \otimes g(\theta^*, x_{\infty})] + \mathbb{E}[(g'(\theta^*, x_{\infty})\mathbb{E}[\theta_{\infty} - x_{\infty}]) \otimes g(\theta^*, x_{\infty})].
$$

Lastly, by a second order Taylor expansion around  $\theta^*$  of  $C(\theta_{\infty}) = \mathbb{E}[(\xi_{\infty+1}(\theta_{\infty}))^{\otimes 2}]$ , we have

$$
C(\theta_{\infty})=C(\theta^*)+\mathbb{C}'(\theta^*)\mathbb{E}[\theta_{\infty}-\theta^*]+\mathbb{E}[R_2'(\theta_{\infty})],
$$

where  $R'_2(\theta)$  satisfies  $\sup_{x \in \mathbb{R}^d} \left\{ ||R'_2(\theta)||/(||\theta - \theta^*||^2 + \|\theta - \theta^*\|^{k_{\epsilon}+2}) \right\} < \infty$ .

Substituting the above analyses of the terms and consolidating the terms, we obtain

<span id="page-46-0"></span>
$$
-(\bar{g}^{(1)}(\theta^*)\otimes I + I \otimes \bar{g}^{(1)}(\theta^*))\mathbb{E}[(\theta_{\infty}-\theta^*)^{\otimes 2}]
$$
  
\n
$$
= \alpha\bar{g}_2(\theta^*) + \alpha \mathbb{E}[(\xi_{\infty+1}(\theta^*))^{\otimes 2}] + \alpha\bar{g}_2^{(1)}(\theta^*)\mathbb{E}[(\theta_{\infty}-\theta^*)^{\otimes 2}]
$$
  
\n
$$
+ \mathbb{E}[g(\theta^*,x_{\infty})\otimes \delta_1(x_{\infty})] + \mathbb{E}[\delta_1(x_{\infty})\otimes g(\theta^*,x_{\infty})]
$$
  
\n
$$
+ \mathbb{E}[g'(\theta^*,x_{\infty})\delta_2(x_{\infty})] + \mathbb{E}[\delta_2(x_{\infty})g'(\theta^*,x_{\infty})]
$$
  
\n
$$
+ \alpha \mathbb{E}[g(\theta^*,x_{\infty})\otimes (g'(\theta^*,x_{\infty})\delta_1(x_{\infty}))] + \alpha \mathbb{E}[g(\theta^*,x_{\infty})\otimes (g'(\theta^*,x_{\infty})\mathbb{E}[\theta_{\infty}-x_{\infty}])]
$$
  
\n
$$
+ \alpha \mathbb{E}[g'(\theta^*,x_{\infty})\delta_1(x_{\infty}))\otimes g(\theta^*,x_{\infty})] + \alpha \mathbb{E}[(g'(\theta^*,x_{\infty})\mathbb{E}[\theta_{\infty}-x_{\infty}])\otimes g(\theta^*,x_{\infty})]
$$
  
\n
$$
+ \alpha \mathbb{E}[g'(\theta^*,x_{\infty})\delta_2(x_{\infty})g'(\theta^*,x_{\infty})]
$$
  
\n
$$
+ \mathbb{E}[R_2(\theta_{\infty},x_{\infty})\otimes (\theta_{\infty}-\theta^*)] + \mathbb{E}[(\theta_{\infty}-\theta^*)\otimes R_2(\theta_{\infty},x_{\infty})] + \alpha C'(\theta^*)\mathbb{E}[\theta_{\infty}-\theta^*]
$$
  
\n
$$
+ \mathcal{O}(\alpha^2\tau).
$$
 (G.17)

By far, we observe that the remaining terms all contain  $\delta_1$  and  $\delta_2$ . Therefore, we conclude our analysis of  $\mathbb{E}[(\theta_{\infty}-\theta^*)^{\otimes 2}]$  at this step and leave the analysis of  $\delta_1$  and  $\delta_2$  to the next section.

### G.2.3 Step 3: Analysis of the  $\delta$ -System

In this section, we analyze  $\delta_1$  and  $\delta_2$ .

Analysis of  $\delta_1$ . Starting with  $\delta_1$ , we first consider the following recursive relationship induced by [\(G.2\)](#page-43-6).

$$
\mathbb{E}[\theta_{\infty+1} - \theta^* | x_{\infty+1} = s] = \int_{\mathbb{R}} P^*(s, ds') \mathbb{E}[\theta_{\infty+1} - \theta^* | x_{\infty+1} = s, x_{\infty} = s']
$$
  
\n
$$
\stackrel{\text{(i)}}{=} \int_{\mathbb{R}} P^*(s, ds') \mathbb{E}[\theta_{\infty} - \theta^* + \alpha g(\theta_{\infty}, x_{\infty}) | x_{\infty} = s']
$$
  
\n
$$
\stackrel{\text{(ii)}}{=} \int_{\mathbb{R}} P^*(s, ds') \mathbb{E} [\theta_{\infty} - \theta^* + \alpha (g(\theta^*, x_{\infty}) + g'(\theta^*, x_{\infty}) (\theta_{\infty} - \theta^*) + R_2(\theta_{\infty}, x_{\infty}) ] | x_{\infty} = s'
$$
  
\n
$$
= \int_{\mathbb{R}} P^*(s, ds') \mathbb{E} [\theta_{\infty} - \theta^* | x_{\infty} = s']
$$
  
\n
$$
+ \alpha \int_{\mathbb{R}} P^*(s, ds') (g(\theta^*, s') + g'(\theta^*, s') \mathbb{E}[\theta_{\infty} - \theta^* | x_{\infty} = s'] + \mathbb{E} [R_2(\theta_{\infty}, x_{\infty}) | x_{\infty} = s'] ),
$$

i

where in (i) we make use of the update rule in [\(2.1\)](#page-2-1),  $\mathbb{E}[\xi_{\infty+1}(\theta_{\infty})] = 0$  and conditional independence  $x_{\infty+1} \perp \!\!\!\perp \theta_{\infty} | x_{\infty}$ . Next, we substitute the Taylor expansion [\(G.5\)](#page-43-5) to obtain (ii).

Writing with notation shorthands z and  $\delta$ , we have

<span id="page-47-0"></span>
$$
z_1(s) = \int_{\mathbb{R}} P^*(s, ds') z_1(s')
$$
  
+  $\alpha \int_R P^*(s, ds') (g(\theta^*, s') + g'(\theta^*, s') z_1(s') + \mathbb{E}[R_2(\theta_\infty, s') | x_\infty = s']).$  (G.18)

If we apply  $\pi$  to both sides of [\(G.18\)](#page-47-0), we obtain

$$
\int \pi(ds) P^*(s, ds') \Big( g(\theta^*, s') + g'(\theta^*, s') z_1(s') + \mathbb{E}[R_2(\theta_\infty, s') | x_\infty = s'] \Big) = 0.
$$

Analyzing the three terms closely, we observe that

$$
\int \pi(\mathrm{d}s) \int P^*(s, \mathrm{d}s') g(\theta^*, s') = \int g(\theta^*, s') \underbrace{\int \pi(\mathrm{d}s) P^*(s, \mathrm{d}s')}_{=\pi(\mathrm{d}s')}
$$

$$
\int \pi(\mathrm{d}s) \int P^*(s, \mathrm{d}s') g'(\theta^*, s') z_1(s') = \mathbb{E}[g'(\theta^*, x_{\infty}) z_1(x_{\infty})]
$$

$$
\int \pi(\mathrm{d}s) \int P^*(s, \mathrm{d}s') \mathbb{E}[R_2(\theta_{\infty}, s') | x_{\infty} = s'] = \mathbb{E}[R_2(\theta_{\infty}, x_{\infty})],
$$

and hence we first obtain

<span id="page-47-1"></span>
$$
\mathbb{E}[g'(\theta^*, x_{\infty})z_1(x_{\infty})] + \mathbb{E}[R_2(\theta_{\infty}, x_{\infty})] = 0.
$$
\n(G.19)

We now subtract  $\pi z_1$  on both sides of [\(G.18\)](#page-47-0), and obtain

$$
\delta_1(s) = \int_{\mathbb{R}} P^*(s, ds') \delta_1(s')
$$
  
+  $\alpha \int_{R} P^*(s, ds') (g(\theta^*, s') + g'(\theta^*, s') z_1(s') + \mathbb{E}[R_2(\theta_\infty, s')] x_\infty = s']$   
=  $\int_{\mathbb{R}} (P^*(s, ds') - \pi(ds')) \delta_1(s')$   
+  $\alpha \int_{R} (P^*(s, ds') - \pi(ds')) (g(\theta^*, s') + g'(\theta^*, s') z_1(s') + \mathbb{E}[R_2(\theta_\infty, s')] x_\infty = s']$ .

Consolidating the terms, we have

<span id="page-47-2"></span>
$$
(I - P^* + \Pi)\delta_1(s)
$$
  
=  $\alpha \int_R \left( P^*(s, ds') - \pi(ds') \right) \left( g(\theta^*, s') + g'(\theta^*, s') z_1(s') + \mathbb{E}[R_2(\theta_\infty, s') | x_\infty = s' ] \right).$  (G.20)

We next note the following properties,

<span id="page-47-3"></span>
$$
\|\mathbb{E}[R_2(\theta_{\infty}, x_{\infty})|x_{\infty} = s]\|_{L^2(\pi)}^2 \le \mathbb{E}[\|R_2(\theta_{\infty}, x_{\infty})^2\|] = \mathcal{O}((\alpha \tau)^2),
$$
  

$$
\|z_1(s)\|_{L^2(\pi)}^2 \le \mathbb{E}[\|\theta_{\infty} - \theta^*\|^2] = \mathcal{O}(\alpha \tau) = \mathcal{O}(1).
$$

Therefore, we can first conclude that  $\|\delta\|_{L^2(\pi)} = \mathcal{O}(\alpha)$ .

Subsequently, from [\(G.19\)](#page-47-1), we can derive that

$$
0 = \mathbb{E}[g'(\theta^*, x_{\infty})z_1(x_{\infty})] + \mathbb{E}[R_2(\theta_{\infty}, x_{\infty})]
$$
  
\n
$$
= \mathbb{E}[g'(\theta^*, x_{\infty})\delta_1(x_{\infty})] + \bar{g}^{(1)}(\theta^*)\mathbb{E}[\theta_{\infty} - \theta^*] + \mathbb{E}[R_2(\theta_{\infty}, x_{\infty})]
$$
  
\n
$$
\mathbb{E}[\theta_{\infty} - \theta^*] = -(\bar{g}^{(1)}(\theta^*))^{-1} (\mathbb{E}[g'(\theta^*, x_{\infty})\delta_1(x_{\infty})] + \mathbb{E}[R_2(\theta_{\infty}, x_{\infty})])
$$
. (G.21)

Hence, together with the relationship between  $\delta_1$  and  $z_1$ , we have

<span id="page-48-0"></span>
$$
z_1 = \delta_1 - (\bar{g}^{(1)}(\theta^*))^{-1} \Big( \mathbb{E}[g'(\theta^*, x_\infty)\delta_1(x_\infty)] + \mathbb{E}[R_2(\theta_\infty, x_\infty)] \Big). \tag{G.22}
$$

Lastly, we substitute  $(G.22)$  back into  $(G.20)$  and obtain

$$
(I - P^* + \Pi)\delta_1(s)
$$
  
\n
$$
= \alpha \int_R (P^*(s, ds') - \pi(ds'))g(\theta^*, s')
$$
  
\n
$$
+ \alpha \int_R (P^*(s, ds') - \pi(ds'))g'(\theta^*, s')(\delta_1(s') - (\bar{g}^{(1)}(\theta^*))^{-1}\mathbb{E}[g'(\theta^*, x_{\infty})\delta_1(x_{\infty})])
$$
  
\n
$$
- \alpha \int_R (P^*(s, ds') - \pi(ds'))g'(\theta^*, s')((\bar{g}^{(1)}(\theta^*))^{-1}\mathbb{E}[R_2(\theta_{\infty}, x_{\infty})])
$$
  
\n
$$
+ \alpha \int_R (P^*(s, ds') - \pi(ds'))\mathbb{E}[R_2(\theta_{\infty}, x_{\infty})|x_{\infty} = s'])
$$
  
\n
$$
= \alpha v(\theta^*, s) + \mathcal{O}(\alpha^2 \tau),
$$

where

$$
v(\theta^*, s) = (I - P^* + \Pi)^{-1} (P^* - \Pi) g_{\theta^*}(s) = \int_R (I - P^* + \Pi)^{-1} (P^* - \Pi)(s, ds') g(\theta^*, s').
$$

Therefore, for the terms that involve  $\delta_1,$  we can conclude that

$$
\mathbb{E}[g(\theta^*, x_{\infty}) \otimes \delta_1(x_{\infty})] = \alpha M + \mathcal{O}(\alpha^2 \tau) \text{ and } \mathbb{E}[\delta_1(x_{\infty}) \otimes g(\theta^*, x_{\infty})] = \alpha M + \mathcal{O}(\alpha^2 \tau), \mathbb{E}[g'(\theta^*, x_{\infty})\delta_1(x_{\infty})] = \alpha v' + \mathcal{O}(\alpha^2 \tau),
$$
\n(G.23)

where  $M$  and  $v$  are independent of  $\alpha$ .

Note that  $(G.23)$  together with  $(G.21)$  implies that

<span id="page-48-1"></span>
$$
\mathbb{E}[\theta_{\infty} - \theta^*] = \alpha v^{(1)} + \mathcal{O}(\alpha \tau).
$$

**Analysis of**  $\delta_2$ . For  $z_2$ , we first note that

$$
|\mathbb{E}[(\theta_{\infty}-\theta^*)^{\otimes2}|x_{\infty}]||_{L^2(\pi)}^2 \leq \mathbb{E}[\|\theta_{\infty}-\theta^*\|^4] = \mathcal{O}((\alpha\tau)^2),
$$

and hence this implies that

$$
||z_2||_{L^2(\pi)} = \mathcal{O}(\alpha \tau).
$$

Next, following  $(G.3)$ , we obtain the following recursive relationship.

$$
\mathbb{E}[(\theta_{\infty+1} - \theta^*)^{\otimes 2}|x_{\infty+1} = s'] = \int P^*(s', ds)\mathbb{E}[(\theta_{\infty+1} - \theta^*)^{\otimes 2}|x_{\infty+1} = s', x_{\infty} = s]
$$
  
\n
$$
\stackrel{\text{(i)}}{=} \int P^*(s', ds)\mathbb{E}[(\theta_{\infty} - \theta^* + \alpha(g(\theta_{\infty}, x_{\infty}) + \xi_{\infty+1}(\theta_{\infty})))^{\otimes 2}|x_{\infty} = s]
$$
  
\n
$$
\stackrel{\text{(ii)}}{=} \int P^*(s', ds)\mathbb{E}[(\theta_{\infty} - \theta^*)^{\otimes 2}|x_{\infty} = s]
$$
  
\n
$$
+ \alpha^2 \int P^*(s', ds)\mathbb{E}[(g(\theta_{\infty}, s))^{\otimes 2}|x_{\infty} = s] + \alpha^2 \int P^*(s', ds)\mathbb{E}[(\xi_{\infty+1}(\theta_{\infty}))^{\otimes 2}|x_{\infty} = s]
$$
  
\n
$$
+ \alpha \int P^*(s', ds)\left(\mathbb{E}[(\theta_{\infty} - \theta^*) \otimes g(\theta_{\infty}, s)|x_{\infty} = s] + \mathbb{E}[g(\theta_{\infty}, s) \otimes (\theta_{\infty} - \theta^*)|x_{\infty} = s]\right)
$$

where in (i) we make use of the update rule in [\(2.1\)](#page-2-1) and conditional independence  $x_{\infty+1} \perp \!\!\!\perp \theta_{\infty} | x_{\infty}$ . Next, we substitute the Taylor expansion  $(G.5)$  to obtain (ii).

,

Writing with  $z_2$  shorthand, we have

$$
z_2(s') = \int P^*(s', ds) z_2(s)
$$
  
+  $\alpha^2 \int P^*(s', ds) \mathbb{E}[(g(\theta_{\infty}, s))^{\otimes 2} | x_{\infty} = s] + \alpha^2 \int P^*(s', ds) \mathbb{E}[(\xi_{\infty+1}(\theta_{\infty}))^{\otimes 2} | x_{\infty} = s]$   
+  $\alpha \int P^*(s', ds) \Big( \mathbb{E}[(\theta_{\infty} - \theta^*) \otimes g(\theta_{\infty}, s) | x_{\infty} = s] + \mathbb{E}[g(\theta_{\infty}, s) \otimes (\theta_{\infty} - \theta^*) | x_{\infty} = s] \Big).$ 

Making use of the relationship  $(P^* - \Pi)z_2 = (P^* - \Pi)\delta_2$ , we have

$$
\delta_2(s') = \int \left( P^*(s', ds) - \pi(ds) \right) \delta_2(s)
$$
  
+  $\alpha^2 \int P^*(s', ds) \mathbb{E}[(g(\theta_{\infty}, s))^{\otimes 2} | x_{\infty} = s] + \alpha^2 \int P^*(s', ds) \mathbb{E}[(\xi_{\infty+1}(\theta_{\infty}))^{\otimes 2} | x_{\infty} = s]$   
+  $\alpha \int P^*(s', ds) \Big( \mathbb{E}[(\theta_{\infty} - \theta^*) \otimes g(\theta_{\infty}, s) | x_{\infty} = s] + \mathbb{E}[g(\theta_{\infty}, s) \otimes (\theta_{\infty} - \theta^*) | x_{\infty} = s] \Big).$ 

Hence,

$$
(I - P^* + \Pi)\delta_2(s')
$$
  
=  $\alpha \int P^*(s', ds) \Big( \mathbb{E}[(\theta_{\infty} - \theta^*) \otimes g(\theta_{\infty}, s)|x_{\infty} = s] + \mathbb{E}[g(\theta_{\infty}, s) \otimes (\theta_{\infty} - \theta^*)|x_{\infty} = s]$   
+  $\alpha^2 \int P^*(s', ds) \Big( \mathbb{E}[g((\theta_{\infty}, s))^{\otimes 2}|x_{\infty} = s] + \mathbb{E}[(\xi_{\infty+1}(\theta_{\infty}))^{\otimes 2}|x_{\infty} = s] \Big).$ 

To analyze the above system of  $\delta_2$ , we make use of the Taylor expansion of  $g(\theta_{\infty}, s)$  and  $\xi_{\infty+1}(\theta_{\infty})$ . Starting with the first term, we have

$$
\mathbb{E}[(\theta_{\infty} - \theta^*) \otimes g(\theta_{\infty}, s)|x_{\infty} = s]
$$
\n
$$
= \mathbb{E}\Big[\Big(\theta_{\infty} - \theta^*\Big) \otimes \Big(g(\theta^*, s) + g'(\theta^*, s)(\theta_{\infty} - \theta^*) + R_2(\theta, s)\Big)|x_{\infty} = s\Big]
$$
\n
$$
= z_1(s) \otimes g(\theta^*, s) + z_2(s)g'(\theta^*, s) + \mathbb{E}[(\theta_{\infty} - \theta^*) \otimes R_2(\theta, s)|x_{\infty} = s]
$$
\n
$$
= \Big(\delta_1(s) - (\bar{g}^{(1)}(\theta^*))^{-1}\Big(\mathbb{E}[g'(\theta^*, x_{\infty})\delta_1(x_{\infty})] + \mathbb{E}[R_2(\theta_{\infty}, x_{\infty})]\Big)\Big) \otimes g(\theta^*, s)
$$
\n
$$
+ z_2(s)g'(\theta^*, s) + \mathbb{E}[(\theta_{\infty} - \theta^*) \otimes R_2(\theta, s)|x_{\infty} = s].
$$

Similarly, we have the following relationship for the second term,

$$
\mathbb{E}[g(\theta_{\infty},s) \otimes (\theta_{\infty} - \theta^*) | x_{\infty} = s]
$$
  
=  $g(\theta^*,s) \otimes (\delta_1(s) - (\bar{g}^{(1)}(\theta^*))^{-1} (\mathbb{E}[g'(\theta^*, x_{\infty})\delta_1(x_{\infty})] + \mathbb{E}[R_2(\theta_{\infty}, x_{\infty})])$   
+  $g'(\theta^*, s)z_2(s) + \mathbb{E}[R_2(\theta, s) \otimes (\theta_{\infty} - \theta^*) | x_{\infty} = s].$ 

Next, we proceed to analyze the third term.

$$
\mathbb{E}[(g(\theta_{\infty},s))^{\otimes2}|x_{\infty}=s]
$$
\n
$$
= \mathbb{E}\Big[\Big(g(\theta^*,s) + g'(\theta^*,s)(\theta_{\infty}-\theta^*) + R_2(\theta_{\infty},s)\Big)^{\otimes2}|x_{\infty}=s\Big]
$$
\n
$$
= \mathbb{E}[(g(\theta^*,s))^{\otimes2}|x_{\infty}=s] + \mathbb{E}[(g'(\theta^*,s)(\theta_{\infty}-\theta^*))^{\otimes2}|x_{\infty}=s]
$$
\n
$$
+ \mathbb{E}[g(\theta^*,s) \otimes (g'(\theta^*,s)(\theta_{\infty}-\theta^*))|x_{\infty}=s] + \mathbb{E}[(g'(\theta^*,s)(\theta_{\infty}-\theta^*)) \otimes g(\theta^*,s)|x_{\infty}=s]
$$
\n
$$
+ \mathbb{E}[g(\theta^*,s) \otimes R_2(\theta_{\infty},s)|x_{\infty}=s] + \mathbb{E}[R_2(\theta_{\infty},s) \otimes g(\theta^*,s)|x_{\infty}=s]
$$
\n
$$
+ \mathbb{E}[(g'(\theta^*,s)(\theta_{\infty}-\theta^*)) \otimes R_2(\theta_{\infty},s)|x_{\infty}=s]
$$
\n
$$
+ \mathbb{E}[R_2(\theta_{\infty},s) \otimes (g'(\theta^*,s)(\theta_{\infty}-\theta^*))|x_{\infty}=s] + \mathbb{E}[(R_2(\theta_{\infty},s))^{\otimes2}|x_{\infty}=s].
$$

Lastly, for the noise term, we derive that

$$
\mathbb{E}[(\xi_{\infty+1}(\theta_{\infty}))^{\otimes 2}|x_{\infty}=s] = \mathbb{E}[(\xi_{\infty+1}(\theta^*))^{\otimes 2}] + C'(\theta^*)\mathbb{E}[\theta_{\infty}-\theta^*|x_{\infty}=s] + \mathbb{E}[R_2'(\theta_{\infty})|x_{\infty}=s].
$$

Leveraging on the respective orders, we can conclude that

<span id="page-50-0"></span>
$$
\|\delta_2\|_{L^2(\pi)} = \mathcal{O}(\alpha^2 \tau).
$$

Therefore, for second-moment cross-terms, we have the following orders

$$
\mathbb{E}[g'(\theta^*, x_{\infty})\delta_2(x_{\infty})] = \mathcal{O}(\alpha^2 \tau) \text{ and } \mathbb{E}[\delta_2(x_{\infty})g'(\theta^*, x_{\infty})] = \mathcal{O}(\alpha^2 \tau), \n\mathbb{E}[g''(\theta^*, x_{\infty})\delta_2(x_{\infty})] = \mathcal{O}(\alpha^2 \tau).
$$
\n(G.24)

#### G.2.4 Step 4: Bias Characterization

Finally, we are ready to consolidate the above analyses and conclude the characterization of the asymptotic bias.

We recall that we have already shown the following expansion of the asymptotic bias

$$
\mathbb{E}[\theta_{\infty} - \theta^*] = -(\bar{g}^{(1)}(\theta^*))^{-1} \Big( \mathbb{E}[g'(\theta^*, x_{\infty})\delta_1(x_{\infty})] + \frac{1}{2} \Big( \mathbb{E}[g''(\theta^*, x_{\infty})\delta_2(x_{\infty})] + \bar{g}^{(2)}(\theta^*) \mathbb{E}[(\theta_{\infty} - \theta^*)^{\otimes 2}] \Big) \Big) + \mathcal{O}((\alpha \tau)^{3/2}).
$$

By our analyses above, we have shown that

$$
\delta_1 = \alpha (I - P^* + \Pi)^{-1} (P^* - \Pi) g_{\theta}^* + \mathcal{O}(\alpha^2 \tau).
$$

Hence, we derive that

$$
\mathbb{E}[g'(\theta^*, x_{\infty})\delta_1(x_{\infty})] = \alpha \mathbb{E}[g'(\theta^*, x_{\infty})(I - P^* + \Pi)^{-1}(P^* - \Pi)g_{\theta}^*(x_{\infty})] + \mathcal{O}(\alpha^2 \tau).
$$

For the second term, we simply use the order shown in [\(G.24\)](#page-50-0).

Lastly, we substitute our analyses of  $\delta_1$  and  $\delta_2$  into the expansion of MSE [\(G.17\)](#page-46-0) and derive that

$$
- (\bar{g}^{(1)}(\theta^*) \otimes I + I \otimes \bar{g}^{(1)}(\theta^*)) \mathbb{E}[(\theta_{\infty} - \theta^*)^{\otimes 2}]
$$
  
\n
$$
= \alpha \bar{g}_2(\theta^*) + \alpha \mathbb{E}[(\xi_{\infty+1}(\theta^*))^{\otimes 2}]
$$
  
\n
$$
+ \alpha \mathbb{E}[g(\theta^*, x_{\infty}) \otimes (I - P^* + \Pi)^{-1}(P^* - \Pi)g_{\theta^*}(x_{\infty})]
$$
  
\n
$$
+ \alpha \mathbb{E}[(I - P^* + \Pi)^{-1}(P^* - \Pi)g_{\theta^*}(x_{\infty}) \otimes g(\theta^*, x_{\infty})] + \mathcal{O}(\alpha^2 \tau).
$$

Therefore, combining all the analyses, we have shown the bias characterization in Theorem [4.6,](#page-6-0)

$$
\mathbb{E}[\theta_{\infty} - \theta^*]
$$
\n
$$
= -\alpha \cdot (\bar{g}^{(1)}(\theta^*))^{-1} \mathbb{E}[g'(\theta^*, x_{\infty})h(\theta^*, x_{\infty})]
$$
\n
$$
+ \alpha \cdot \frac{1}{2}(\bar{g}^{(1)}(\theta^*))^{-1}(\bar{g}^{(1)}A(\bar{g}_2(\theta^*) + \alpha \mathbb{E}[(\xi_{\infty+1}(\theta^*))^{\otimes 2}])
$$
\n
$$
+ \alpha \cdot \frac{1}{2}(\bar{g}^{(1)}(\theta^*))^{-1}A(\mathbb{E}[g(\theta^*, x_{\infty}) \otimes h(\theta^*, x_{\infty})] + \mathbb{E}[h(\theta^*, x_{\infty}) \otimes g(\theta^*, x_{\infty})]) + \mathcal{O}((\alpha \tau)^{3/2}).
$$

where

$$
A = (\bar{g}^{(1)}(\theta^*) \otimes I + I \otimes \bar{g}^{(1)}(\theta^*))^{-1},
$$
  
\n
$$
h(\theta^*, s) = (I - P^* + \Pi)^{-1}(P^* - \Pi)g_{\theta^*}(s)
$$
  
\n
$$
= \int_{\mathcal{X}} (I - P^* + \Pi)^{-1}(P^* - \Pi)(s, ds')g(\theta^*, s').
$$

Therefore, we see that assuming weak convergence without projection, the bias admits a leading term of order  $\alpha$ . We emphasize that the expansion holds as equality, rather than an upper bound.

#### G.3 Step 2: Impact of Projection on Bias

Now, we proceed to analyze the impact of having the additional projection step on the asymptotic bias characterization. In the following, we use the shorthand  $\theta_{t+1/2}$  to denote the iterate we obtain before the projection step, i.e.,

$$
\theta_{t+1/2} = \theta_t + \alpha (g(\theta_t, x_t) + \xi_{t+1}(\theta_t))
$$
 and  $\theta_{t+1} = \Pi_{B(\beta)} \theta_{t+1/2}$ .

Therefore, we see that our analysis from Step 1 can be understood as the analysis for  $\theta_{\infty+1/2}$ . Starting with the first moment analysis with test function  $h_1(x, \theta) = \theta - \theta^*$ , we have

$$
\mathbb{E}[\theta_{\infty+1} - \theta^*] = \mathbb{E}[\theta_{\infty+1/2} - \theta^*] + \mathbb{E}[\theta_{\infty+1} - \theta_{\infty+1/2}]
$$
  
=  $\mathbb{E}[\theta_{\infty} - \theta^*] + \alpha \big(\mathbb{E}[g(\theta_{\infty}, x_{\infty})] + \mathbb{E}[\xi_{\infty+1}(\theta_{\infty})]\big) + \mathbb{E}[\theta_{\infty+1} - \theta_{\infty+1/2}],$ 

which implies that

<span id="page-51-0"></span>
$$
-\frac{1}{\alpha}\mathbb{E}[\theta_{\infty+1} - \theta_{\infty+1/2}] = \mathbb{E}[g'(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*)] + \frac{1}{2}\mathbb{E}[g''(\theta^*, x_{\infty})(\theta_{\infty} - \theta^*)^{\otimes 2}] + \mathcal{O}((\alpha\tau)^{3/2}).
$$
 (G.25)

Therefore, we turn our focus to analyzing  $\mathbb{E}[\theta_{\infty+1} - \theta_{\infty+1/2}]$ .

$$
\begin{split} &\mathbb{E}[\theta_{\infty+1}-\theta_{\infty+1/2}] \\ &=\mathbb{E}[(\theta_{\infty+1}-\theta_{\infty+1/2})\mathbbm{1}\{\|\theta_{t+1/2}-\theta^*\|<\beta\}]+\mathbb{E}[(\theta_{\infty+1}-\theta_{\infty+1/2})\mathbbm{1}\{\|\theta_{t+1/2}-\theta^*\|\geq\beta\}] \\ &=\mathbb{E}[(\theta_{\infty+1}-\theta_{\infty+1/2})\mathbbm{1}\{\|\theta_{t+1/2}-\theta^*\|\geq\beta\}], \end{split}
$$

where we note that when  $\|\theta_{t+1/2} - \theta^*\| \leq \beta$  implies that  $\|\theta_{t+1/2}\| \leq \beta + \|\theta^*\| \leq 2\beta$  and hence  $\theta_{\infty+1} = \theta_{t+1/2}$ in this case. To analyze the remaining term, we use Hölder's inequality and obtain

$$
\begin{aligned} &\|\mathbb{E}[(\theta_{\infty+1}-\theta_{\infty+1/2})\mathbbm{1}\{\|\theta_{t+1/2}-\theta^*\| \geq \beta\}]\| \\ &\leq \mathbb{E}^{1/p}[\|(\theta_{\infty+1}-\theta^*)-(\theta_{\infty+1/2}-\theta^*)\|^p]\mathbb{E}^{1/q}[\mathbbm{1}\{\|\theta_{t+1/2}-\theta^*\| \geq \beta\}] \\ &\leq 2\mathbb{E}^{1/p}[\|\theta_{\infty+1/2}-\theta^*\|^p]\big(\mathbb{P}(\|\theta_{t+1/2}-\theta^*\| \geq \beta)\big)^{1/q}. \end{aligned}
$$

Setting  $p = 6$  and  $q = 6/5$ , and making use of the property that  $\mathbb{E}[\|\theta_{\infty+1/2} - \theta^*\|^6] = \mathcal{O}((\alpha\tau)^3)$  from Proposition [4.2,](#page-5-1) we have

$$
\mathbb{E}^{1/6}[\|\theta_{\infty+1/2}-\theta^*\|^6] \big(\mathbb{P}(\|\theta_{t+1/2}-\theta^*\| \ge R)\big)^{5/6} \lesssim (\alpha \tau)^{3/6} \big(\mathbb{E}[\|\theta_{t+1/2}-\theta^*\|^6]/R^6)^{5/6} \lesssim (\alpha \tau)^3.
$$

Hence, we can conclude that

$$
\mathbb{E}[\theta_{\infty+1} - \theta_{\infty+1/2}] = \mathcal{O}((\alpha\tau)^3).
$$

Substituting this order information back into [\(G.25\)](#page-51-0), we can see that  $\mathbb{E}[\theta_{\infty+1} - \theta_{\infty+1/2}]/\alpha = \mathcal{O}(\alpha^2 \tau^3)$ . Recall that  $\tau = \mathcal{O}(\log(1/\alpha))$ , and hence we can assimilate this residual order from projection into the existing  $\mathcal{O}((\alpha\tau)^{3/2})$  residual term.

For the remaining terms, we follow the existing analysis in Section [G.2.1](#page-44-6) and again obtain

$$
\mathbb{E}[\theta_{\infty} - \theta^*] = -(\bar{g}^{(1)}(\theta^*))^{-1} \Big( \mathbb{E}[g'(\theta^*, x_{\infty})\delta_1(x_{\infty})] + \frac{1}{2} \Big( \mathbb{E}[g''(\theta^*, x_{\infty})\delta_2(x_{\infty})] + \bar{g}^{(2)}(\theta^*) \mathbb{E}[(\theta_{\infty} - \theta^*)^{\otimes 2}] \Big) \Big) + \mathcal{O}((\alpha \tau)^{3/2}).
$$

Now, we proceed to analyze  $\mathbb{E}[(\theta_{\infty}-\theta^*)^{\otimes 2}]$  and examine the impact of projection. Consider test function  $h_2(x, \theta) = (\theta - \theta^*)^{\otimes 2}$  and follow a similar strategy as the first moment analysis, we have

$$
\mathbb{E}[(\theta_{\infty+1} - \theta^*)^{\otimes 2}] = \mathbb{E}[(\theta_{\infty+1/2} - \theta^*)^{\otimes 2}] + \mathbb{E}[(\theta_{\infty+1} - \theta^*)^{\otimes 2} - (\theta_{\infty+1/2} - \theta^*)^{\otimes 2}]
$$

$$
= \mathbb{E}[(\theta_{\infty} - \theta^*)^{\otimes 2}] + \alpha^2 (\mathbb{E}[(g(\theta_{\infty}, x_{\infty}))^{\otimes 2}] + \mathbb{E}[(\xi_{\infty+1}(\theta_{\infty}))^{\otimes 2}]) + \alpha (\mathbb{E}[g(\theta_{\infty}, x_{\infty}) \otimes (\theta_{\infty} - \theta^*)] + \mathbb{E}[(\theta_{\infty} - \theta^*) \otimes g(\theta_{\infty}, x_{\infty})]) + \mathbb{E}[(\theta_{\infty+1} - \theta^*)^{\otimes 2} - (\theta_{\infty+1/2} - \theta^*)^{\otimes 2}].
$$

Hence, by reorganizing the terms, we have

$$
\begin{split} &-\frac{1}{\alpha}\mathbb{E}[(\theta_{\infty+1}-\theta^*)^{\otimes 2}-(\theta_{\infty+1/2}-\theta^*)^{\otimes 2}] \\ &=\alpha(\mathbb{E}[(g(\theta_{\infty},x_{\infty}))^{\otimes 2}]+\mathbb{E}[(\xi_{\infty+1}(\theta_{\infty}))^{\otimes 2}]) \\ &+(\mathbb{E}[g(\theta_{\infty},x_{\infty})\otimes(\theta_{\infty}-\theta^*)]+\mathbb{E}[(\theta_{\infty}-\theta^*)\otimes g(\theta_{\infty},x_{\infty})]). \end{split}
$$

Therefore, we turn our focus to analyzing  $\mathbb{E}[(\theta_{\infty+1}-\theta^*)^{\otimes2}-(\theta_{\infty+1/2}-\theta^*)^{\otimes2}]$ . Similar as the first moment analysis, we have

$$
\mathbb{E}[(\theta_{\infty+1} - \theta^*)^{\otimes 2} - (\theta_{\infty+1/2} - \theta^*)^{\otimes 2}]
$$
  
= 
$$
\mathbb{E}[((\theta_{\infty+1} - \theta^*)^{\otimes 2} - (\theta_{\infty+1/2} - \theta^*)^{\otimes 2}) \mathbb{1}\{ \|\theta_{t+1/2} - \theta^*\| \ge \beta \}].
$$

To analyze the term on the right hand side, we again make use of Hölder's inequality and obtain

$$
\begin{split} &\|\mathbb{E}[((\theta_{\infty+1}-\theta^*)^{\otimes 2}-(\theta_{\infty+1/2}-\theta^*)^{\otimes 2})\mathbbm{1}\{\|\theta_{t+1/2}-\theta^*\| \geq \beta\}]\| \\ &\leq \mathbb{E}^{1/p}[\|(\theta_{\infty+1}-\theta^*)^{\otimes 2}-(\theta_{\infty+1/2}-\theta^*)^{\otimes 2}\|^p]\mathbb{E}^{1/q}[\mathbbm{1}\{\|\theta_{t+1/2}-\theta^*\| \geq \beta\}] \\ &\leq 2\mathbb{E}^{1/p}[\|\theta_{\infty+1/2}-\theta^*\|^2]^p] \big(\mathbb{P}(\|\theta_{t+1/2}-\theta^*\| \geq \beta)\big)^{1/q}. \end{split}
$$

Setting  $p = 3$  and  $q = 3/2$ , and making use of the property that  $\mathbb{E}[\|\theta_{\infty+1/2} - \theta^*\|^6] = \mathcal{O}((\alpha\tau)^3)$  from Proposition [4.2,](#page-5-1) we have

$$
\mathbb{E}^{1/3}[\|\theta_{\infty+1/2} - \theta^*\|^6] \big(\mathbb{P}(\|\theta_{t+1/2} - \theta^*\| \ge R)\big)^{2/3} \lesssim (\alpha \tau) \Big(\mathbb{E}[\|\theta_{t+1/2} - \theta^*\|^6]/R^6\Big)^{2/3} \lesssim (\alpha \tau)^3.
$$

Hence, we can conclude that

$$
\mathbb{E}[(\theta_{\infty+1}-\theta^*)^{\otimes 2}-(\theta_{\infty+1/2}-\theta^*)^{\otimes 2}]=\mathcal{O}((\alpha\tau)^3).
$$

From the analyses above, we can also conclude that

$$
\|\mathbb{E}[\theta_{\infty+1} - \theta_{\infty+1/2}|x_{\infty}]\|_{L^2(\pi)} = \mathcal{O}((\alpha\tau)^{3/2})
$$
  

$$
\|\mathbb{E}[(\theta_{\infty+1} - \theta^*)^{\otimes 2} - (\theta_{\infty+1/2} - \theta^*)^{\otimes 2}|x_{\infty}]\|_{L^2(\pi)} = \mathcal{O}((\alpha\tau)^{3/2}).
$$

Therefore, combining the analyses above, we see that the projection only introduces error terms of order  $\mathcal{O}(\alpha^2 \tau^3)$ . Hence, combining the analysis from Section [G.2.1,](#page-44-6) we can conclude the same desired order that

<span id="page-52-2"></span><span id="page-52-1"></span>
$$
\mathbb{E}[\theta_{\infty}^{(\alpha)} - \theta^*] = \alpha b + \mathcal{O}((\alpha \tau)^{3/2}).
$$

## <span id="page-52-0"></span>H Additional Insights on TA and RR

In this section, we present more detailed results that characterize the first and second moment of Polayk-Ruppert (PR) tail-averaged iterates and Richardson-Romberg (RR) extrapolated iterates.

The following corollary provides non-asymptotic characterization for the first two moments of PR tailaveraged iterates  $\theta_{k_0,k}$ .

Corollary H.1 (Tail Averaging). Under the setting of Theorem [4.6,](#page-6-0) the tail-averaged iterates satisfy the following bounds for all  $k > k_0 + 2\tau$  and  $k_0 \geq \tau + \frac{1}{\alpha \mu} \log \left( \frac{1}{\alpha \tau_{\alpha}} \right)$ :

$$
\mathbb{E}[\bar{\theta}_{k_0,k}^{(\alpha)} - \theta^*] = \alpha b + \mathcal{O}\left((\alpha \tau_\alpha)^{3/2}\right) + \mathcal{O}\left(\frac{(1 - \alpha \mu)^{k_0/2}}{\alpha (k - k_0)}\right) \quad \text{and} \tag{H.1}
$$

$$
\mathbb{E}\left[(\bar{\theta}_{k_0,k}^{(\alpha)} - \theta^*)(\bar{\theta}_{k_0,k}^{(\alpha)} - \theta^*)^\top\right] = \alpha^2 b b^T + \mathcal{O}(\alpha \cdot (\alpha \tau_\alpha)^{3/2}) + \mathcal{O}\left(\frac{\tau_\alpha}{k - k_0} + \frac{(1 - \alpha \mu)^{k_0/2}}{\alpha (k - k_0)^2}\right). \tag{H.2}
$$

<span id="page-53-0"></span>With this result, taking the trace on both sides of  $(H.2)$  recovers Corollary [4.7.](#page-7-1)

Proof. First, we have

$$
\mathbb{E}[\bar{\theta}_{k_0,k}^{(\alpha)} - \theta^*] = (\mathbb{E}[\theta_{\infty}] - \theta^*) + \frac{1}{k - k_0} \sum_{t = k_0}^{k-1} \mathbb{E}[\theta_t - \theta_{\infty}].
$$

By Corollary [4.4,](#page-6-8) we obtain

$$
\|\mathbb{E}[\theta_t] - \mathbb{E}[\theta_\infty]\| \le (1 - \alpha \mu)^{\frac{t}{2}} \cdot s'(\theta_0, L, \mu).
$$

Hence, it follows that

$$
\left\| \sum_{t=k_0}^{k-1} \mathbb{E} \left[ \theta_t - \theta_\infty \right] \right\| \leq \sum_{t=k_0}^{k-1} \left\| \mathbb{E} \left[ \theta_t \right] - \mathbb{E}[\theta_\infty] \right\|
$$
  

$$
\leq s'(\theta_0, L, \mu) \cdot \left( 1 - \alpha \mu \right)^{\frac{k_0}{2}} \frac{1}{1 - \sqrt{(1 - \alpha \mu)}}
$$
  

$$
\leq s'(\theta_0, L, \mu) \cdot \left( 1 - \alpha \mu \right)^{\frac{k_0}{2}} \frac{2}{\alpha \mu}.
$$

Together with Theorem [4.6,](#page-6-0) we have

$$
\mathbb{E}\left[\bar{\theta}_{k_0,k}^{(\alpha)}\right] - \theta^* = \alpha b + \mathcal{O}\left((\alpha\tau_\alpha)^{3/2}\right) + \mathcal{O}\left(\frac{(1-\alpha\mu)^{k_0/2}}{\alpha(k-k_0)}\right),
$$

thereby finishing the proof of the first moment.

To bound the second moment of the tail-averaged iterate, we follow the proof technique in [\[HCX23a,](#page-11-6) Section A.6.2] . We notice that

$$
\mathbb{E}\left[\left(\bar{\theta}_{k_{0},k}^{(\alpha)}-\theta^{*}\right)\left(\bar{\theta}_{k_{0},k}^{(\alpha)}-\theta^{*}\right)^{\top}\right] \n= \mathbb{E}\left[\left(\bar{\theta}_{k_{0},k}^{(\alpha)}-\mathbb{E}\left[\theta_{\infty}\right]+\mathbb{E}\left[\theta_{\infty}\right]-\theta^{*}\right)\left(\bar{\theta}_{k_{0},k}^{(\alpha)}-\mathbb{E}\left[\theta_{\infty}\right]+\mathbb{E}\left[\theta_{\infty}\right]-\theta^{*}\right)^{\top}\right] \n= \mathbb{E}\left[\left(\bar{\theta}_{k_{0},k}^{(\alpha)}-\mathbb{E}\left[\theta_{\infty}\right]\right)\left(\bar{\theta}_{k_{0},k}^{(\alpha)}-\mathbb{E}\left[\theta_{\infty}\right]\right)^{\top}\right]+\mathbb{E}\left[\left(\bar{\theta}_{k_{0},k}^{(\alpha)}-\mathbb{E}\left[\theta_{\infty}\right]\right)\left(\mathbb{E}\left[\theta_{\infty}\right]-\theta^{*}\right)^{\top}\right] \n+ \mathbb{E}\left[\left(\mathbb{E}\left[\theta_{\infty}\right]-\theta^{*}\right)\left(\bar{\theta}_{k_{0},k}^{(\alpha)}-\mathbb{E}\left[\theta_{\infty}\right]\right)^{\top}\right]+\mathbb{E}\left[\left(\mathbb{E}\left[\theta_{\infty}\right]-\theta^{*}\right)\left(\mathbb{E}\left[\theta_{\infty}\right]-\theta^{*}\right)^{\top}\right].
$$

For  $T_2$ , we have

$$
T_2 = \frac{1}{k - k_0} \left( \sum_{t = k_0}^{k - 1} \mathbb{E} \left[ \theta_t - \theta_\infty \right] \right) \left( \mathbb{E}[\theta_\infty] - \theta^* \right)^\top
$$
  
=  $\mathcal{O} \left( \frac{(1 - \alpha \mu)^{k_0/2}}{\alpha (k - k_0)} \right) \cdot (\alpha b + \mathcal{O}((\alpha \tau_\alpha)^{3/2})) = \mathcal{O} \left( \frac{(1 - \alpha \mu)^{k_0/2}}{(k - k_0)} \right).$ 

The term  $\mathcal{T}_3$  is similar to  $\mathcal{T}_2$  and obeys the same bound.

For  $T_4$ , we have

$$
T_4 = (\alpha b + \mathcal{O}((\alpha \tau_\alpha)^{3/2}))(\alpha b + \mathcal{O}((\alpha \tau_\alpha)^{3/2}))^T = \alpha^2 b b^T + \mathcal{O}(\alpha \cdot (\alpha \tau_\alpha)^{3/2}).
$$

For  $T_1$ , we have

$$
T_1 = \frac{1}{(k - k_0)^2} \mathbb{E}\bigg[ \Big( \sum_{t = k_0}^{k-1} (\theta_t - \mathbb{E}[\theta_\infty]) \Big) \Big( \sum_{t = k_0}^{k-1} (\theta_t - \mathbb{E}[\theta_\infty]) \Big)^{\top} \bigg]
$$

<span id="page-54-0"></span>
$$
=\frac{1}{(k-k_0)^2}\sum_{t=k_0}^{k-1}\mathbb{E}\left[\left(\theta_t-\mathbb{E}[\theta_\infty]\right)\left(\theta_t-\mathbb{E}[\theta_\infty]\right)^{\top}\right]
$$
(H.3)

<span id="page-54-2"></span><span id="page-54-1"></span>
$$
+\frac{1}{(k-k_0)^2}\sum_{t=k_0}^{k-1}\sum_{l=t+1}^{k-1}\mathbb{E}\left[\left(\theta_t-\mathbb{E}[\theta_\infty]\right)\left(\theta_l-\mathbb{E}[\theta_\infty]\right)^{\top}\right]
$$
(H.4)

<span id="page-54-3"></span>
$$
+\frac{1}{(k-k_0)^2}\sum_{t=k_0}^{k-1}\sum_{l=t+1}^{k-1}\mathbb{E}\left[\left(\theta_l-\mathbb{E}[\theta_\infty]\right)\left(\theta_t-\mathbb{E}[\theta_\infty]\right)^{\top}\right].
$$
\n(H.5)

By Corollary [4.4](#page-6-8) and Proposition [4.2](#page-5-1) we have

$$
\mathbb{E}\Big[\big(\theta_t - \mathbb{E}[\theta_\infty]\big)\big(\theta_t - \mathbb{E}[\theta_\infty]\big)^\top\Big] \n= \Big(\mathbb{E}\big[\theta_t\theta_t^\top\big] - \mathbb{E}\big[\theta_\infty\theta_\infty^\top\big]\Big) + \Big(\mathbb{E}\big[\theta_\infty\theta_\infty^\top\big] - \mathbb{E}[\theta_\infty]\mathbb{E}\big[\theta_\infty^\top\big]\Big) \n- \Big(\mathbb{E}\big[\theta_t\big]\mathbb{E}\big[\theta_\infty^\top\big] + \mathbb{E}[\theta_\infty]\mathbb{E}\big[\theta_t^\top\big] - 2\mathbb{E}[\theta_\infty]\mathbb{E}\big[\theta_\infty^\top\big]\Big) \n= \Big(\mathbb{E}\big[\theta_t\theta_t^\top\big] - \mathbb{E}\big[\theta_\infty\theta_\infty^\top\big]\Big) + \text{Var}\big(\theta_\infty\big) - \mathbb{E}\big[\theta_t - \theta_\infty\big]\mathbb{E}\big[\theta_\infty^\top\big] - \mathbb{E}[\theta_\infty]\mathbb{E}\Big[(\theta_t - \theta_\infty)^\top\Big] \n= \mathcal{O}\Big((1 - \alpha\mu)^{\frac{t}{2}} + \alpha\tau_\alpha\Big),
$$

where we bound Var  $(\theta_{\infty})$  with  $\mathcal{O}(\alpha \tau_{\alpha})$  by Proposition [4.2](#page-5-1) and Fatou's lemma.

Then, for [\(H.3\)](#page-54-1), we have

$$
\begin{split} \text{(H.3)} &= \frac{1}{\left(k - k_0\right)^2} \sum_{t = k_0}^{k - 1} \mathcal{O}\left((1 - \alpha \mu)^{\frac{t}{2}} + \alpha \tau_\alpha\right) \\ &= \mathcal{O}\left(\frac{1}{\left(k - k_0\right)^2} \sum_{t = k_0}^{\infty} \left(1 - \alpha \mu\right)^{\frac{t}{2}}\right) + \mathcal{O}\left(\frac{\alpha \tau_\alpha}{k - k_0}\right) \\ &= \mathcal{O}\left(\frac{\left(1 - \alpha \mu\right)^{k_0/2}}{\alpha \left(k - k_0\right)^2} + \frac{\alpha \tau_\alpha}{k - k_0}\right). \end{split}
$$

We restate the following claim, whose proof closely resembles Claim 4 in [\[HCX23a](#page-11-6)].

**Claim 2.** For 
$$
t \ge \tau + \frac{1}{\alpha \mu} \log \left( \frac{1}{\alpha \tau_{\alpha}} \right)
$$
 and  $l \ge t + 2\tau_{\alpha}$ , we have  

$$
\left\| \mathbb{E} \left[ \left( \theta_t - \mathbb{E}[\theta^{(\alpha)}] \right) \left( \theta_l - \mathbb{E}[\theta^{(\alpha)}] \right)^{\top} \right] \right\| = \mathcal{O} \left( \left( \alpha \tau_{\alpha} \right) \cdot \left( 1 - \alpha \mu \right)^{\frac{(l-t)}{2}} \right).
$$

Then, by [\[HCX23a,](#page-11-6) Claim 4], we have term  $(H.4) = \mathcal{O}(\frac{\tau_{\alpha}}{k-k_0})$ . Similarly, we have term  $(H.5) = \mathcal{O}(\frac{\tau_{\alpha}}{k-k_0})$ . Therefore, we have

<span id="page-54-4"></span>
$$
T_1 = \mathcal{O}\left(\frac{(1 - \alpha\mu)^{k_0/2}}{\alpha (k - k_0)^2}\right) + \frac{\tau_\alpha}{k - k_0}.
$$
\n(H.6)

By adding  $T_1\hbox{--} T_4$  together, we obtain

$$
\mathbb{E}\left[\left(\bar{\theta}_{k_0,k}^{(\alpha)}-\theta^*\right)\left(\bar{\theta}_{k_0,k}^{(\alpha)}-\theta^*\right)^{\top}\right]=\alpha^2bb^T+\mathcal{O}(\alpha\cdot(\alpha\tau_{\alpha})^{3/2})+\mathcal{O}\left(\frac{(1-\alpha\mu)^{k_0/2}}{(k-k_0)}\right)\\+\mathcal{O}\left(\frac{(1-\alpha\mu)^{k_0/2}}{\alpha(k-k_0)^2}+\frac{\tau_{\alpha}}{k-k_0}\right)\\=\alpha^2bb^T+\mathcal{O}(\alpha\cdot(\alpha\tau_{\alpha})^{3/2})+\mathcal{O}\left(\frac{\tau_{\alpha}}{k-k_0}+\frac{(1-\alpha\mu)^{k_0/2}}{\alpha(k-k_0)^2}\right).
$$

 $\Box$ 

Next, we present the following corollary formalizes the non-asymptotic characterization for the first two moments of the RR-extrapolated iterate  $\widetilde{\theta}_{k_0,k}^{(\alpha)}$ .

Corollary H.2 (Richardson-Romberg Extrapolation). Under the setting of Theorem [4.6,](#page-6-0) the RR extrapolated iterates with stepsizes  $\alpha$  and  $2\alpha$  satisfy the following bounds for all  $k > k_0 + 2\tau_\alpha$  and  $k_0 \ge \tau_\alpha +$  $\frac{1}{\alpha\mu}\log\left(\frac{1}{\alpha\tau_{\alpha}}\right)$ :

$$
\mathbb{E}[\tilde{\theta}_{k_0,k}^{(\alpha)} - \theta^*] = \mathcal{O}\big((\alpha \tau_\alpha)^{3/2}\big) + \mathcal{O}\Big(\frac{(1 - \alpha \mu)^{k_0/2}}{\alpha(k - k_0)}\Big), \quad \text{and} \tag{H.7}
$$

$$
\mathbb{E}\Big[(\tilde{\theta}_{k_0,k}^{(\alpha)} - \theta^*)(\tilde{\theta}_{k_0,k} - \theta^*)^\top\Big] = \mathcal{O}\big((\alpha \tau_\alpha)^3\big) + \mathcal{O}\Big(\frac{\tau_\alpha}{k - k_0} + \frac{(1 - \alpha \mu)^{k_0/2}}{\alpha (k - k_0)^2}\Big). \tag{H.8}
$$

*Proof.* By equation  $(H.1)$ , we obtain

$$
\mathbb{E}\left[\tilde{\theta}_{k_0,k}^{(\alpha)}\right] - \theta^* = \mathbb{E}\left[2\bar{\theta}_{k_0,k}^{(\alpha)} - \bar{\theta}_{k_0,k}^{(2\alpha)}\right] - \theta^* = 2\mathbb{E}\left[\bar{\theta}_{k_0,k}^{(\alpha)} - \theta^*\right] - \mathbb{E}\left[\bar{\theta}_{k_0,k}^{(2\alpha)} - \theta^*\right]
$$

$$
= 2\left(\alpha b + \mathcal{O}\left((\alpha\tau_{\alpha})^{3/2}\right) + \mathcal{O}\left(\frac{(1 - \alpha\mu)^{k_0/2}}{\alpha(k - k_0)}\right)\right)
$$

$$
-\left(2\alpha b + \mathcal{O}\left((2\alpha\tau_{2\alpha})^{3/2}\right) + \mathcal{O}\left(\frac{(1 - 2\alpha\mu)^{k_0/2}}{\alpha(k - k_0)}\right)\right)
$$

$$
= \mathcal{O}\left((\alpha\tau_{\alpha})^{3/2}\right) + \mathcal{O}\left(\frac{(1 - \alpha\mu)^{k_0/2}}{\alpha(k - k_0)}\right).
$$

Let  $u_1 := \bar{\theta}_{k_0,k}^{(\alpha)} - \mathbb{E}\left[\theta_{\infty}^{(\alpha)}\right]$ ∞  $\Big], u_2 := \bar{\theta}_{k_0,k}^{(2\alpha)} - \mathbb{E}\left[\theta_{\infty}^{(2\alpha)}\right]$ ∞ and  $v := 2 \mathbb{E} \left[ \theta_{\infty}^{(\alpha)} \right]$ ∞  $\Big] - \mathbb{E} \left[ \theta_{\infty}^{(2\alpha)} \right]$ ∞  $\Big] - \theta^*$ . With these notations,  $\tilde{\theta}_{k_0,k} - \theta^* = 2u_1 - u_2 + v$ . We then have the following bound

$$
\left\| \mathbb{E}\left[ \left( \tilde{\theta}_{k_0,k}^{(\alpha)} - \theta^* \right) \left( \tilde{\theta}_{k_0,k}^{(\alpha)} - \theta^* \right)^\top \right] \right\| \leq \mathbb{E}\left[ \left\| 2u_1 - u_2 + v \right\|^2 \right] \leq 3 \mathbb{E} \left\| 2u_1 \right\|^2 + 3 \mathbb{E} \left\| u_2 \right\|^2 + 3 \left\| v \right\|^2.
$$

By equation  $(H.6)$ , we have

$$
\mathbb{E} \|u_1\|^2 = \text{Tr} \left( \mathbb{E} \left[ u_1 u_1^\top \right] \right) = \mathcal{O} \left( \frac{\left(1 - \alpha \mu \right)^{k_0/2}}{\alpha \left( k - k_0 \right)^2} + \frac{\tau_\alpha}{k - k_0} \right).
$$

Similarly, we have

$$
\mathbb{E} \|u_2\|_2^2 = \mathcal{O}\left(\frac{(1 - 2\alpha\mu)^{k_0/2}}{\alpha (k - k_0)^2} + \frac{\tau_{2\alpha}}{k - k_0}\right).
$$

By Theorem [4.6,](#page-6-0) we have  $||v||_2^2 = \mathcal{O}((\alpha \tau_\alpha)^3)$ .

Combining these bounds, we have

$$
\mathbb{E}\Big[\big(\tilde{\theta}_{k_0,k}-\theta^*\big)\big(\tilde{\theta}_{k_0,k}-\theta^*\big)^{\top}\Big] = \mathcal{O}\big((\alpha\tau_{\alpha})^3\big)+\mathcal{O}\Big(\frac{\tau_{\alpha}}{k-k_0}+\frac{(1-\alpha\mu)^{k_0/2}}{\alpha\left(k-k_0\right)^2}\Big).
$$

