

When is exponential asymptotic optimality achievable in average-reward restless bandits?

Yige Hong^{1*} Qiaomin Xie² Yudong Chen³ Weina Wang¹

¹Computer Science Department, Carnegie Mellon University

²Department of Industrial and Systems Engineering, University of Wisconsin-Madison

³Department of Computer Sciences, University of Wisconsin-Madison

{yigeh, weinaw}@cs.cmu.edu

{qiaomin.xie, yudong.chen}@wisc.edu

Abstract

We consider the discrete-time infinite-horizon average-reward restless bandit problem. We propose a novel policy that maintains two dynamic subsets of arms: one subset of arms has a nearly optimal state distribution and takes actions according to an Optimal Local Control routine; the other subset of arms is driven towards the optimal state distribution and gradually merged into the first subset. We show that our policy is asymptotically optimal with an $O(\exp(-CN))$ optimality gap for an N -armed problem, under the mild assumptions of aperiodic-unichain, non-degeneracy, and local stability. Our policy is the first to achieve exponential asymptotic optimality under the above set of easy-to-verify assumptions, whereas prior work either requires a strong Global Attractor assumption or only achieves an $O(1/\sqrt{N})$ optimality gap. We further discuss the fundamental obstacles in significantly weakening our assumptions. In particular, we prove a lower bound showing that local stability is fundamental for exponential asymptotic optimality.

1 Introduction

Restless bandits (RBs) [Whi88] is a stochastic sequential decision-making problem consisting of multiple arms coupled by a constraint. Each arm operates as a Markov decision process (MDP) with two possible actions: activating or keeping passive. At each time step, the decision maker observes the states of the arms and chooses a fixed fraction of arms to activate, aiming to maximize the expected reward collected from the arms. The RB problem has a long history and diverse applications. Readers can refer to the recent survey paper [NM23] for a comprehensive review of the literature.

In this paper, we study the infinite-horizon RBs in discrete time, with the long-run average-reward objective. Since computing the exact optimal policy for an RB problem is intractable as the number of arms N is large [PT99], we focus on developing asymptotically optimal policies that can be efficiently computed. Specifically, the optimality gap of a policy is the difference between the average reward per arm and that of an optimal policy. A policy is considered to be *asymptotically optimal* if the optimality gap is $o(1)$ in the asymptotic limit of $N \rightarrow \infty$.¹

*Corresponding author

¹In this paper, we use standard Bachmann–Landau notation and focus on the asymptotics with respect to N .

The asymptotic optimality of average-reward RB in the large N regime has been studied for decades, starting with the seminal papers on the renowned Whittle index policy [Whi88, WW90]. Since then researchers have been weakening the assumptions for asymptotic optimality [Ver16, HXCW23, HXCW24, Yan24] or improving the order of optimality gaps [GGY23a, GGY23b]. A recent milestone is the *exponential asymptotic optimality* established in [GGY23a, GGY23b], where a class of policies called LP-Priority is proved to achieve an optimality gap $O(\exp(-CN))$ for some constant C , under some assumptions. Such an exponential gap has significant practical value, since it implies that a policy has nearly-optimal performance even for RBs with small N . Moreover, an exponential gap also “beats” the Central Limit Theorem (CLT), an intriguing theoretical property that is not a priori obvious to be possible. In particular, on a high-level all asymptotic optimality results in the RB literature are based on the concentration of the empirical distribution of the arms’ states around a certain optimal distribution. Therefore, an $O(1/\sqrt{N})$ bound owing to CLT was believed to be fundamental. An important observation made in [GGY23a, GGY23b] is that if the system has a certain notion of *local linearity*, the optimality gap only depends on the distance between the optimal distribution and the *expected* empirical state distribution, allowing one to go beyond CLT.

Despite the above intuition, it remains unclear what is the fundamental mechanism leading to exponential asymptotic optimality. Although a simple assumption called non-degeneracy (or non-singularity) suffices to ensure the local linearity property described above, non-degeneracy alone is not enough to guarantee exponential optimality of the LP-Priority policies considered in [GGY23a, GGY23b]. In particular, another crucial assumption is needed, namely the Uniform Global Attractor Property (UGAP). Without UGAP, LP-Priority may even have a constant optimality gap; see [WW90, GGY20, HXCW23] for concrete examples. Unfortunately, UGAP is hard to interpret and verify, as it concerns the global convergence of a certain non-linear difference equation.

In light of the difficulty caused by UGAP, very recent work has studied new policies that are asymptotically optimal without assuming UGAP [HXCW23, HXCW24, Yan24]. Different from LP-Priority and Whittle index policies, these new policies actively control the empirical distribution of the states of the arms, driving the distribution to *globally converge* towards a certain optimal distribution, without relying on extraneous assumptions. However, while these results conclude that UGAP is not needed for asymptotic optimality, it remains unclear whether UGAP is fundamental for exponential asymptotic optimality — the best optimality gap proved in this line of work is $O(1/\sqrt{N})$.

Given these developments, it is natural to ask if removing UGAP necessarily results in a sacrifice in the optimality gap. Is it possible to efficiently find policies that achieve *global exponential asymptotic optimality*, i.e., an $O(\exp(-CN))$ optimality gap without assuming global attraction?

Our contributions. The primary contribution of the paper is to design the first policy, the *Two-Set Policy*, that achieves exponential asymptotic optimality without UGAP. The Two-Set Policy maintains two dynamic subsets of arms and applies two different subroutines to the subsets: one subset of arms has a nearly optimal state distribution and takes actions according to a certain Optimal Local Control; the other subset of arms is driven towards the optimal state distribution and gradually merged into the first subset. Intuitively, the Optimal Local Control in the first subset induces local linearity, while the second subset achieves global convergence. While the ideas in each subroutine is not completely new, the Two-Set Policy leverages them in a novel way to achieve the best of both worlds. The theoretical guarantees of the Two-Set Policy is informally summarized in the theorem below.

Theorem (Informal version of Theorem 1). *Suppose each arm is unichain, aperiodic, non-degenerate*

and locally stable. Then we can efficiently compute a N -armed policy π such that

$$R^{\text{rel}} - R(\pi, \mathbf{S}_0) \leq O(\exp(-CN)), \quad (1)$$

where R^{rel} is an (efficiently computable) upper bound on the optimal long-run average reward, and $R(\pi, \mathbf{S}_0)$ is the long-run average reward under the policy π given the vector of initial states \mathbf{S}_0 .

We will make precise the definitions of non-degeneracy and local stability. We emphasize that we only need aperiodic unichain under a particular policy rather than all policies. The assumptions of Theorem 1 are a subset of, and hence strictly weaker than, the assumptions made in prior work [GGY23a, GGY23b] for exponential asymptotic optimality. Moreover, all our assumptions pertain to a linear program defined by problem primitives and are thus much easier to verify than UGAP.

The proof of the above theorem employs a novel multivariate Lyapunov function technique, which generalizes the focus-set approach in [HXCW24]. This multivariate Lyapunov function enables us to decouple the complex dynamics under the Two-Set Policy, where the states of the arms and the two dynamic subsets are coupled and change simultaneously.

We next elucidate the roles of our assumptions for exponential optimality. The importance of unichain, aperiodicity and non-degeneracy is well recognized; we complement the discussion with concrete examples. We further establish the fundamental importance of local stability by proving the following.

Theorem (Informal version of Theorem 2). *For regular unstable RBs, every policy π satisfies*

$$R^{\text{rel}} - R(\pi, \mathbf{S}_0) \geq \Omega(1/\sqrt{N}). \quad (2)$$

We will formally define the class of regular (locally) unstable RB and provide concrete examples. Note that all known efficient policies rely on the upper bound R^{rel} . For these policies, Theorem 2 shows that local stability is crucial for achieving exponential asymptotic optimality for a natural class of RB instances. The proof of Theorem 2 uses Lyapunov analysis to reveal an interesting trade-off between two competing needs when local stability fails: to keep the states of the arms near the optimal distribution, and to take the optimal actions when the states are near the optimal distribution.

Other related work. Apart from the average-reward RBs, there are also a rich body of literature on RBs with finite-horizon total-reward [HF17, ZCJW19, BS20, GNJT23, ZF21, GGY23b, GGY24] or or the infinite-horizon discounted-reward criteria [ZF22, GNJT23]. The approaches in these setting do not directly apply to the average-reward setting, as their computational complexities scale with the (effective) time horizon. However, there are still a large intersection of ideas — in particular, the first $O(1/N)$ optimality gap result was established in the finite-horizon setting by [ZF21].

2 Problem setup

Consider the discrete-time average-reward restless bandit problem with N homogeneous arms, referred to as the N -armed problem. Index the arms by $[N] \triangleq \{1, 2, \dots, N\}$. To avoid confusion with the Whittle index, We refer to i as the *ID* of Arm i . Each arm is associated with a Markov decision process (MDP) defined by $(\mathbb{S}, \mathbb{A}, P, r)$, which is called the single-armed MDP. Here \mathbb{S} is a finite state space; $\mathbb{A} = \{0, 1\}$ is the action space, where action 1 is activating/pulling the arm; $P : \mathbb{S} \times \mathbb{A} \times \mathbb{S} \rightarrow [0, 1]$ is the transition kernel; $r : \mathbb{S} \times \mathbb{A} \rightarrow \mathbb{R}$ is the reward function. Let

$r_{\max} = \max_{s \in \mathbb{S}, a \in \mathbb{A}} |r(s, a)|$. The RB problem has a *budget constraint*, given by a constant $\alpha \in (0, 1)$, which requires that α fraction of arms must be pulled at every time step. We assume that αN is an integer. We focus on the setting where all the model parameters, $\mathbb{S}, \mathbb{A}, P, r, \alpha$, are known.

A policy π for the N -armed problem chooses the action for each of the N arms in each time step. The policy can be randomized and history-dependent. Under π , the *state vector* $\mathbf{S}_t^\pi \triangleq (S_t^\pi(i))_{i \in [N]} \in \mathbb{S}^N$ represents the states of all arms at time t , and the *action vector* $\mathbf{A}_t^\pi \triangleq (A_t^\pi(i))_{i \in [N]} \in \mathbb{A}^N$ denotes the actions applied to each arm. For each policy π , consider the following two quantities: the limsup average reward $R^+(\pi, \mathbf{S}_0) \triangleq \limsup_{T \rightarrow \infty} \frac{1}{T} \sum_{t=0}^{T-1} \frac{1}{N} \sum_{i \in [N]} \mathbb{E}[r(S_t^\pi(i), A_t^\pi(i))]$; and the liminf average reward $R^-(\pi, \mathbf{S}_0) \triangleq \liminf_{T \rightarrow \infty} \frac{1}{T} \sum_{t=0}^{T-1} \frac{1}{N} \sum_{i \in [N]} \mathbb{E}[r(S_t^\pi(i), A_t^\pi(i))]$. When the limsup and liminf coincide, the long-run average reward can be defined as follows:

$$R(\pi, \mathbf{S}_0) \triangleq \lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=0}^{T-1} \frac{1}{N} \sum_{i \in [N]} \mathbb{E}[r(S_t^\pi(i), A_t^\pi(i))].$$

Note that $R(\pi, \mathbf{S}_0)$ is well-defined when π is Markovian w.r.t. the state \mathbf{S}_t , as \mathbf{S}_t is a finite-state Markov chain under such π [Put05, Proposition 8.1.1]. Using a similar argument, one can show that $R(\pi, \mathbf{S}_0)$ is well-defined if π is Markovian w.r.t. an augmented system state with a finite state space.

Our goal is to solve the following optimization problem.

$$\begin{aligned} & \underset{\text{policy } \pi}{\text{maximize}} && R^-(\pi, \mathbf{S}_0) && \text{subject to} && \sum_{i \in [N]} A_t^\pi(i) = \alpha N, \quad \forall t \geq 0. \end{aligned} \quad (\text{RB})$$

Let $R^*(N)$ be the optimal value of (RB), referred to as the *optimal reward*. Note that $R^*(N) = \sup_{\pi'} R^-(\pi', \mathbf{S}_0) = \sup_{\pi'} R^+(\pi', \mathbf{S}_0)$ because (RB) is an MDP with finite state and action spaces [Put05, Theorem 9.1.6]. For any policy π , we define its optimality gap as $R^*(N) - R^-(\pi, \mathbf{S}_0)$. A policy is *asymptotically optimal* if its optimality gap vanishes as $N \rightarrow \infty$, i.e., $R^*(N) - R^-(\pi, \mathbf{S}_0) = o(1)$. This notion of asymptotic optimality is consistent with the literature [Ver16, Definition 4.11].

We remark that there always exists a stationary Markovian policy that achieves the optimal reward [Put05, Theorem 9.1.8]. In later parts of the paper, although we will not restrict ourselves to stationary Markovian policies, we focus on policies that induce finite-state Markov chains. Under each of these policies, π , the long-run average reward $R(\pi, \mathbf{S}_0)$ is well-defined. For simplicity, we will refer to $R(\pi, \mathbf{S}_0)$ as the objective function of (RB) and write the optimality gap as $R^*(N) - R(\pi, \mathbf{S}_0)$.

Scaled state-count vector. We introduce an alternative representation of the N -armed system state, which will be used extensively throughout the paper. For each subset $D \subseteq [N]$, we define the *scaled state-count vector on D* as $X_t^\pi(D) = (X_t^\pi(D, s))_{s \in \mathbb{S}}$, where $X_t^\pi(D, s) \triangleq \frac{1}{N} \sum_{i \in D} \mathbb{1}\{S_t^\pi(i) = s\}$ is the number of arms in D in state s scaled by $1/N$. When $D = [N]$ is the set of all arms, we simply call $X_t^\pi([N])$ the *scaled state-count vector*. Sometimes we view $X_t^\pi(D)$ as a vector-valued function of $D \subseteq [N]$. We refer to this function X_t as the *system state* at time t . Note that the system state X_t^π contains the same information as the state vector \mathbf{S}_t^π . We also define the *scaled state-action-count vector* as $Y_t^\pi = (Y_t^\pi(s, a))_{s \in \mathbb{S}, a \in \mathbb{A}}$, where $Y_t^\pi(s, a) \triangleq \frac{1}{N} \sum_{i \in [N]} \mathbb{1}\{S_t^\pi(i) = s, A_t^\pi(i) = a\}$ is the fraction of arms with state s and action a at time t .

Additional notation. Let $[0, 1]_N = \{0, 1/N, 2/N, \dots, 1\}$. For each subset $D \subseteq [N]$, $m(D) = |D|/N$ denotes the fraction of arms contained in D , thus $m(D) \in [0, 1]_N$. Let $\Delta(\mathbb{S})$ be probability simplex on \mathbb{S} . We treat each distribution $v \in \Delta(\mathbb{S})$ as a row vector. Recall that π denotes a policy for the N -armed problem. When the context is clear, we drop superscript π from \mathbf{S}_t^π , \mathbf{A}_t^π , and X_t^π .

3 Two-Set Policy

In this section, we describe our Two-Set Policy. We first discuss a linear programming (LP) relaxation of (RB) in Section 3.1 and state assumptions. The optimal solution of the LP is then used to define two subroutines, *Exact Proportional Control* in Section 3.2, and *Optimal Local Control* in Section 3.3. We then use these two subroutines to construct the Two-Set Policy in Section 3.4. The ideas in the two subroutines are not new; our novelty lies in how we build them into the Two-Set Policy.

3.1 LP relaxation

We consider the linear programming (LP) relaxation of the N -armed problem (RB), which is crucial for the design and analysis of RB policies.

$$\begin{aligned} & \underset{\{y(s,a)\}_{s \in \mathbb{S}, a \in \mathbb{A}}}{\text{maximize}} && \sum_{s \in \mathbb{S}, a \in \mathbb{A}} r(s,a)y(s,a) \end{aligned} \tag{LP}$$

$$\text{subject to} \quad \sum_{s \in \mathbb{S}} y(s,1) = \alpha, \tag{3}$$

$$\sum_{s' \in \mathbb{S}, a \in \mathbb{A}} y(s',a)P(s',a,s) = \sum_{a \in \mathbb{A}} y(s,a), \forall s; \quad \sum_{s \in \mathbb{S}, a \in \mathbb{A}} y(s,a) = 1; \quad y(s,a) \geq 0, \forall s,a. \tag{4}$$

To see why (LP) is a relaxation of (RB), for any stationary Markovian policy π , consider $y^\pi(s,a) = \lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E}[\frac{1}{N} \sum_{i \in [N]} \mathbb{1}\{S_t^\pi(i) = s, A_t^\pi(i) = a\}]$, $\forall s \in \mathbb{S}, a \in \mathbb{A}$. It is not hard to see that $R(\pi, \mathbf{S}_0) = \sum_{s \in \mathbb{S}, a \in \mathbb{A}} r(s,a)y^\pi(s,a)$, and $(y^\pi(s,a))_{s \in \mathbb{S}, a \in \mathbb{A}}$ satisfies the constraints (3)-(4). Therefore, letting R^{rel} be the optimal value of (LP), it can be shown that $R^{\text{rel}} \geq R^*(N)$ (See Appendix A.1). Note that the LP relaxation can be efficiently solved. We can bound the optimality gap of any policy π using the inequality $R^*(N) - R^-(\pi, \mathbf{S}_0) \leq R^{\text{rel}} - R^-(\pi, \mathbf{S}_0)$. This bound has been widely used in prior work [WW90, Ver16, GGY23a, GGY23b, HXCW23].

Optimal single-armed policy. For a fixed optimal solution to (LP), $y^* = \{y^*(s,a)\}_{s \in \mathbb{S}, a \in \mathbb{A}}$, we consider the conditional distribution $\bar{\pi}^* = (\bar{\pi}^*(a|s))_{s \in \mathbb{S}, a \in \mathbb{A}}$ given by:

$$\bar{\pi}^*(a|s) = \begin{cases} y^*(s,a)/(y^*(s,0) + y^*(s,1)), & \text{if } y^*(s,0) + y^*(s,1) > 0, \\ 1/2, & \text{if } y^*(s,0) + y^*(s,1) = 0. \end{cases} \quad \text{for } s \in \mathbb{S}, a \in \mathbb{A}. \tag{5}$$

We call $\bar{\pi}^*$ the *optimal single-armed policy*, due to the fact that $\bar{\pi}^*$ can be viewed as the optimal policy for the single-armed MDP $(\mathbb{S}, \mathbb{A}, P, r)$ with a relaxed budget constraint [HXCW24].

Let $P_{\bar{\pi}^*}$ be the transition matrix induced by the policy $\bar{\pi}^*$ on the single-armed MDP, i.e., $P_{\bar{\pi}^*}(s,s') = \sum_{a \in \mathbb{A}} \bar{\pi}^*(a|s)P(s,a,s')$. Since $P_{\bar{\pi}^*}$ is a stochastic matrix, its largest eigenvalue is equal to 1. For the other eigenvalues of $P_{\bar{\pi}^*}$, we consider the following assumption.

Assumption 1 (Aperiodic unichain). The moduli of all eigenvalues of $P_{\bar{\pi}^*}$ except the largest one are strictly less than 1; that is, $P_{\bar{\pi}^*}$ defines an aperiodic unichain on \mathbb{S} .

With Assumption 1, the transition matrix $P_{\bar{\pi}^*}$ induces a unique stationary distribution, $\mu^* \triangleq (\mu^*(s))_{s \in \mathbb{S}}$, where $\mu^*(s) = y^*(s,1) + y^*(s,0)$, as one can verify using the definition of $P_{\bar{\pi}^*}$. We call μ^* the *optimal stationary distribution*. We define $S^\emptyset \triangleq \{s \in \mathbb{S}: y^*(s,1) = y^*(s,0) = 0\}$, the set of transient states under policy $\bar{\pi}^*$.

Our next assumption imposes a non-degenerate condition on the optimal solution to (LP).

Subroutine 1 Exact Proportional Control

Input: A set of arms with number of arms in each state $(z(s))_{s \in \mathbb{S}}$, optimal single-armed policy $\bar{\pi}^*$

- 1: **for** $s \in \mathbb{S}$ **do**
 - 2: Compute p such that $p \lfloor \bar{\pi}^*(1|s)z(s) \rfloor + (1-p) \lceil \bar{\pi}^*(1|s)z(s) \rceil = \bar{\pi}^*(1|s)z(s)$.
 - 3: With probability p and $1-p$, activate $\lfloor \bar{\pi}^*(1|s)z(s) \rfloor$ or $\lceil \bar{\pi}^*(1|s)z(s) \rceil$ arms in state s .
-

Assumption 2 (Non-degenerate). For the fixed optimal solution to (LP), y^* , there exists a unique *neutral state*, i.e., a state $\tilde{s} \in \mathbb{S}$ s.t. $y^*(\tilde{s}, 1) > 0$ and $y^*(\tilde{s}, 0) > 0$.

Note that Assumption 2 stipulates the existence of a neutral state \tilde{s} — one can always find an optimal solution of (LP) such that there is at most one neutral state [GGY23b, Proposition 2].

Assumption 3 (Local stability). Given the non-degenerate condition, define Φ as a $|\mathbb{S}|$ -by- $|\mathbb{S}|$ matrix:

$$\Phi \triangleq P_{\bar{\pi}^*} - \mathbb{1}^\top \mu^* - (c_{\bar{\pi}^*} - \alpha \mathbb{1})^\top (P_1(\tilde{s}) - P_0(\tilde{s})), \quad (6)$$

where $c_{\bar{\pi}^*} \triangleq (\bar{\pi}^*(1|s))_{s \in \mathbb{S}}$ and $P_a(\tilde{s}) \triangleq (P(\tilde{s}, a, s))_{s \in \mathbb{S}}$ are both row vectors; $\mathbb{1}$ is the all-one row vector. We assume that the moduli of all eigenvalues of Φ are strictly less than 1.

The implication of Assumption 3 will become clear after we introduce the subroutine *Optimal Local Control* in Section 3.3, which is constructed from the optimal solution to (LP) and has dynamics given by the matrix Φ . We note that [GGY23a] also assumes a version of local stability, in addition to UGAP, to achieve exponential asymptotic optimality. Our local stability assumption is substantially weaker, and it is implied by their version of local stability together with UGAP (see Appendix C.4 of [GGY23a]). Importantly, we do not require a global assumption like UGAP.

Remark 1. Assumptions 1, 2 and 3 are easy to verify as they pertain to either the *single-armed* MDP primitives (P, r, α) or a linear program define by these problem primitives. In contrast, verifying UGAP is significantly harder due to the dependency on the *non-linear* N -armed dynamics.

Remark 2. Assumptions 1, 2 and 3 all depend on an optimal solution y^* to the (LP). When we state our results with more than one assumption among the three, we consider the same y^* .

3.2 Subroutine 1: Exact Proportional Control

We now describe the *Exact Proportional Control*, as presented in Subroutine 1. The subroutine takes the optimal single-armed policy $\bar{\pi}^*$ defined in (5) and a subset of arms D from the RB system as the input. The subroutine decides the actions for all arms in D to match with $\bar{\pi}^*$. Specifically, let the number of arms from D in state $s \in \mathbb{S}$ be $z(s)$. The subroutine aims to activate $\bar{\pi}^*(1|s)$ fraction of the $z(s)$ arms. To handle the integer effect, it activates $\lfloor \bar{\pi}^*(1|s)z(s) \rfloor$ or $\lceil \bar{\pi}^*(1|s)z(s) \rceil$ arms with properly chosen probability (computed in Line 2) such that the expected proportion is exactly $\bar{\pi}^*(1|s)$.

As shown in Lemma 4, if all N arms in the RB system follow Exact Proportional Control, then as $t \rightarrow \infty$, the expected scaled state-count vector $\mathbb{E}[X_t([N])]$ converges to μ^* ; the expected fraction of arms in state s taking action a converge to $y^*(s, a)$; the *expected budget usage* of the arms converges to αN . Consequently, the average expected reward achieves the optimum R^{rel} of (LP).

We remark that Subroutine 1 let the arms follow $\bar{\pi}^*$ in a way different from some prior work [HXCW23, HXCW24], where the actions are sampled from $\bar{\pi}^*$ independently across the arms.

Subroutine 2 Optimal Local Control

Input: A set of arms with number of arms in each state $(z(s))_{s \in \mathbb{S}}$, budget B , optimal single-armed policy $\bar{\pi}^*$, neutral state \tilde{s}

- 1: **Assert** Assumption 2 and conditions in (7)
 - 2: **for** $s \in \mathbb{S} \setminus \{\tilde{s}\}$ **do**
 - 3: Compute p such that $p \lfloor \bar{\pi}^*(1|s)z(s) \rfloor + (1-p) \lceil \bar{\pi}^*(1|s)z(s) \rceil = \bar{\pi}^*(1|s)z(s)$.
 - 4: With probability p and $1-p$, activate $\lfloor \bar{\pi}^*(1|s)z(s) \rfloor$ or $\lceil \bar{\pi}^*(1|s)z(s) \rceil$ arms with state s .
 - 5: Activate the arms with state \tilde{s} using the remaining budget out of B .
-

3.3 Subroutine 2: Optimal Local Control

Next, we describe Subroutine 2, *Optimal Local Control*, another building block for constructing our policy. The subroutine also takes the optimal single-armed policy $\bar{\pi}^*$ and a subset of arms D as the input. One key difference from Subroutine 1 is that here Subroutine 2 requires a predetermined budget B as an input, independent of the states of the arms. Specifically, for the input subset D of n arms, the budget B is set as $\lfloor \alpha n \rfloor$ or $\lceil \alpha n \rceil$ with suitable probability such that $\mathbb{E}[B] = \alpha n$, to handle αn being non-integer. When $D = [N]$, we set $B = \alpha N$ as the original budget on the N -armed system. For each non-neutral state $s \in \mathbb{S} \setminus \{\tilde{s}\}$, the policy also aims to activate $\bar{\pi}^*(1|s)$ fraction of the $z(s)$ arms from D that are in state s . To ensure this is feasible with the budget B constraint, we consider the following conditions for the input subset D :

$$\sum_{s \neq \tilde{s}} \bar{\pi}^*(1|s)z(s) \leq \alpha n - |S^\emptyset| - 1, \quad \text{and} \quad \sum_{s \neq \tilde{s}} \bar{\pi}^*(0|s)z(s) \leq (1-\alpha)n - |S^\emptyset| - 1. \quad (7)$$

Recall that $S^\emptyset = \{s \in \mathbb{S} : y^*(s, 1) = y^*(s, 0) = 0\}$ is the set of transient states under $\bar{\pi}^*$. By the definition of $\bar{\pi}^*$ in (5), it is not hard to see that under conditions in (7), there hold $\sum_{s \neq \tilde{s}} \lceil \bar{\pi}^*(1|s)z(s) \rceil \leq \lfloor \alpha n \rfloor$ and $z(\tilde{s}) + \sum_{s \neq \tilde{s}} \lfloor \bar{\pi}^*(1|s)z(s) \rfloor \geq \lceil \alpha n \rceil$. Therefore, the budget requested by non-neutral states $\mathbb{S} \setminus \{\tilde{s}\}$ does not exceed B (Line 2-4 of Subroutine 2), and the neutral state ensures exactly B unit of budget is used (Line 5).

If all arms in the RB system follow Subroutine 2, the expected scaled state-count vector $\mathbb{E}[X_t([N])]$ has a linear dynamics, as stated by following the lemma (proof provided in Appendix A).

Lemma 1. *Under Assumption 2, suppose (7) holds for the N -armed system at time step t . We have*

$$\mathbb{E}[X_{t+1}([N]) - \mu^* \mid X_t, \text{ all arms follow Subroutine 2}] = (X_t([N]) - \mu^*)\Phi, \quad (8)$$

where Φ is the $|\mathbb{S}|-by-|\mathbb{S}|$ matrix defined in (6).

Thus under Assumption 3, Subroutine 2 induces locally stable dynamics. Specifically, if the RB system satisfies (7) for a sufficiently long time and we let all N arms follow Subroutine 2, then the scaled state-count vector $X_t([N])$ would converge to μ^* . Consequently, the expected fraction of arms in state s taking action a would be close to $y^*(s, a)$ and the time-average of expected reward would be close to the upper bound R^{rel} —hence the name “Optimal Local Control” for Subroutine 2.

3.4 Constructing the Two-Set Policy

We now present our *Two-Set Policy* for the N -armed system, whose pseudo-code is given in Algorithm 1. On a high level, the Two-Set Policy works by picking out two disjoint subsets of arms

Algorithm 1 Two-Set Policy

- Input:** number of arms N , budget αN , an optimal solution of LP-relaxation y^* , threshold η , initial system state X_0 , initial state vector \mathbf{S}_0 , initial focus set $D_{-1} = \emptyset$
- 1: **for** $t = 0, 1, \dots$ **do**
 - 2: Let $D_{\text{temp}}^{\text{OL}}$ be a set with largest cardinality such that $D_{\text{temp}}^{\text{OL}} \subseteq D_{t-1}^{\text{OL}}$, $\delta(X_t, D_{\text{temp}}^{\text{OL}}) \geq 0$.
 - 3: Let D_t^{OL} be a set with largest cardinality such that $D_t^{\text{OL}} \supseteq D_{\text{temp}}^{\text{OL}}$, $\delta(X_t, D_t^{\text{OL}}) \geq 0$.
 - 4: Let $D_t^{\bar{\pi}^*}$ be s.t. $D_{t-1}^{\bar{\pi}^*} \setminus D_t^{\text{OL}} \subseteq D_t^{\bar{\pi}^*} \subseteq [N] \setminus D_t^{\text{OL}}$ and $|D_t^{\bar{\pi}^*}| = (\lfloor \beta(N - |D_t^{\text{OL}}|) \rfloor - 1)^+$.
 - 5: Set $A_t(i)$ for $i \in D_t^{\text{OL}}$ using Subroutine 2 Optimal Local Control.
 - 6: Set $A_t(i)$ for $i \in D_t^{\bar{\pi}^*}$ using Subroutine 1 exact propotional policy.
 - 7: Set $A_t(i)$ for $i \notin D_t^{\text{OL}} \cup D_t^{\bar{\pi}^*}$ such that $\sum_{i \in [N]} A_t(i) = \alpha N$.
 - 8: Apply $A_t(i)$ for $i \in [N]$ and observe the new state vector \mathbf{S}_{t+1} .
-

among the N arms in each time step, and letting them follow two subroutines, Optimal Local Control and Exact Proportional Control, respectively. When the two sets are picked properly, the policy pushes the scaled state-count vector $X_t([N])$ of the N -armed system towards μ^* . Ultimately, the system will stabilize in the region where all arms can follow Subroutine 2 Optimal Local Control.

Specifically, at time step t , the Two-Set Policy first picks a subset D_t^{OL} , where the arms will follow Optimal Local Control. The policy chooses D_t^{OL} in two steps, specified in Line 2-3 of Algorithm 1. Roughly speaking, it intends to let D_t^{OL} have a large overlap with D_{t-1}^{OL} as well as a large cardinality, subject the constraint $\delta(X_t, D_t^{\text{OL}}) \geq 0$, where $\delta(\cdot, \cdot)$ is a function called *slack*, defined as follows:

$$\delta(x, D^{\text{OL}}) = \eta m(D^{\text{OL}}) - \|x(D^{\text{OL}}) - m(D^{\text{OL}})\mu^*\|_U, \quad (9)$$

where η is an input parameter with suggested value $|\mathbb{S}|^{-1/2}(\min\{y^*(\tilde{s}, 0), y^*(\tilde{s}, 1)\} - (|S^\emptyset| + 1)/N)$, U is a matrix defined as $U \triangleq \sum_{k=0}^{\infty} \Phi^k (\Phi^\top)^k$, and $\|v\|_U = \sqrt{v^\top U v}$ for $v \in \mathbb{R}^{|\mathbb{S}|}$. In particular, the criteria that $\delta(X_t, D_t^{\text{OL}}) \geq 0$ ensures that arms in D_t^{OL} satisfy a ‘‘closedness’’ property: they can continue to follow Optimal Local Control almost infinitely while satisfying condition (7), until some rare event happens due to stochastic transitions. We show this property in Lemma 7 and Lemma 11.

After choosing D_t^{OL} , the Two-Set Policy picks a subset $D_t^{\bar{\pi}^*}$ that contains $\beta \triangleq \min(\alpha, 1 - \alpha)$ fraction of the remaining arms, rounded to a certain nearby integer. The set $D_t^{\bar{\pi}^*}$ should be such that $D_t^{\bar{\pi}^*} \cap D_t^{\text{OL}} = \emptyset$ and $D_t^{\bar{\pi}^*} \supseteq D_{t-1}^{\bar{\pi}^*} \setminus D_t^{\text{OL}}$.

With D_t^{OL} and $D_t^{\bar{\pi}^*}$ chosen, the Two-Set Policy let arms in D_t^{OL} follow Subroutine 2 Optimal Local Control and let the arms in $D_t^{\bar{\pi}^*}$ follow Subroutine 1 Exact Proportional Control, and chooses the actions of remaining arms to meet the budget constraint. We will show that this is always feasible in Lemma 7 in Appendix B.

4 Main results

In this section, we present our main results.

4.1 Optimality guarantee on the Two-Set Policy

Our first main theorem states an upper bound on the optimality gap of the Two-Set Policy.

Theorem 1. *Suppose Assumption 1, 2 and 3 hold. Let π be the Two-Set Policy in Algorithm 1 with $\eta = |\mathbb{S}|^{-1/2}(\min\{y^*(\tilde{s}, 0), y^*(\tilde{s}, 1)\} - (|S^\emptyset| + 1)/N)$. Then for some $C > 0$ independent of N :*

$$R^{\text{rel}} - R(\pi, \mathbf{S}_0) = O(\exp(-CN)). \quad (10)$$

Theorem 1 establishes the *first* exponential asymptotic optimality without UGAP. It implies that even in an RB system with a small number of arms, Two-Set Policy is nearly optimal, under three mild and easy-to-verify assumptions. In contrast, prior work either requires UGAP to achieve exponential asymptotic optimality or only has $O(1/\sqrt{N})$ optimality gap. We prove Theorem 1 in Appendix B.

We discuss the relationship between our Two-Set Policy and the set-expansion policy from prior work [HXCW24]. Set-expansion achieves an $O(1/\sqrt{N})$ optimality gap under the aperiodic unichain assumption (Assumption 1). It works by letting arms in a specific subset follow $\bar{\pi}^*$ and gradually expanding the subset. Our Two-Set Policy refines this approach by maintaining two judiciously chosen sets instead of one set. Furthermore, we argue that when Assumption 2 or 3 fails, a variant of the Two-Set Policy can achieve $O(1/\sqrt{N})$ optimality gap, since it becomes almost the same as the set-expansion policy. To see this, note that when Assumption 2 (non-degeneracy) fails, it holds that $\eta = -O(1/N)$; when Assumption 3 (local stability) fails, the matrix U becomes infinity. In either case, D_t^{OL} will become the maximal set such that $x(D_t^{\text{OL}}) = m(D_t^{\text{OL}})\mu^*$ if we allow $\delta(X_t, D_t^{\text{OL}})$ to be slightly negative. Consequently, D_t^{OL} equivalently follows the Exact Proportional Control and can be combined with $D_t^{\bar{\pi}^*}$. Although the Two-Set Policy and the set-expansion policy differ in how the arms follow $\bar{\pi}^*$ and the criteria for choosing the subset, we believe these differences are minor. Using similar arguments as [HXCW24], one can prove an $O(1/\sqrt{N})$ bound for the Two-Set Policy.

4.2 Discussion of the assumptions

In this subsection, we discuss the importance of the assumptions in Theorem 1. We will show that there are scenarios where, for every policy π and certain initial states \mathbf{S}_0 , it holds that

$$R^{\text{rel}} - R(\pi, \mathbf{S}_0) \geq \Omega\left(\frac{1}{\sqrt{N}}\right). \quad (11)$$

When Assumption 1 (aperiodic unichain) or Assumption 2 (non-degeneracy) is violated, it is not hard to construct examples where (11) holds for every policy π and certain initial states \mathbf{S}_0 ; see Appendix C.1. In the non-unichain example, $R^*(N)$ may depend on the initial states \mathbf{S}_0 , in which case the initial-state-independent upper bound R^{rel} is loose. In the periodic and degenerate examples, the upper bound R^{rel} can only be achieved by a policy that satisfies the budget constraint on average over time rather than every time step, again causing a large gap between R^{rel} and $R^*(N)$.

We emphasize that although R^{rel} is just an upper bound of $R^*(N)$, *all* existing bounds on the optimality gap use R^{rel} or its variants. Hence, the above examples reveal the limitations of existing technology. Specifically, while the unichain assumption may be circumvented by a tighter upper bound than R^{rel} , the aperiodic and non-degenerate assumptions appear more fundamental — they are closely tied to the *per-step* constraint of RB problems. Therefore, relaxing the aperiodic and non-degeneracy assumptions requires fundamentally new approaches.

4.2.1 A lower bound when Assumption 3 is violated

When local stability (Assumption 3) fails, instead of finding one counterexample, we take a step further to establish the lower bound (11) for a large class of instances. To establish this lower bound, we need a fundamental lemma that relates the gap $R^{\text{rel}} - R^*(N)$ to our Subroutine 2.

Lemma 2. *Suppose that the LP relaxation (LP) has a unique optimal solution y^* , and $y^*(s, 1) + y^*(s, 0) > 0, \forall s \in \mathbb{S}$. Then*

$$R^{\text{rel}} - R(\pi, \mathbf{S}_0) \geq \epsilon_0 \lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E}[d_{\text{IC}}(Y_t^\pi)], \quad (12)$$

where ϵ_0 is a positive constant; d_{IC} is a function of state-action distribution given by

$$d_{\text{IC}}(y) = \sum_{s \in S^+} y(s, 0) + \sum_{s \in S^-} y(s, 1),$$

where $S^+ = \{s \in \mathbb{S}: y^*(s, 1) > 0, y(s, 0) = 0\}$, $S^- = \{s \in \mathbb{S}: y^*(s, 1) = 0, y^*(s, 0) > 0\}$.

Note that $d_{\text{IC}}(Y_t)$ is zero if and only if Y_t is generated by Subroutine 2 (Optimal Local Control). Therefore, Lemma 2 shows that the gap $R^{\text{rel}} - R(\pi, \mathbf{S}_0)$ is determined by how closely a policy π follows Subroutine 2. Lemma 2 is proved in Appendix C.2 using a primal-dual argument.

Because $d_{\text{IC}}(Y_t) \geq |\sum_{s \in S^+} X_t([N], s) - \alpha|$, Lemma 2 implies that for an RB satisfying conditions of the lemma, to achieve exponential asymptotic optimality, one must keep the long-run average of $|\sum_{s \in S^+} X_t([N], s) - \alpha|$ exponentially small. Doing so could be hard when the RB is degenerate, as there is no neutral state to absorb the randomness of $X_t([N])$.

Lemma 2 can also be used to show a lower bound on $R^{\text{rel}} - R(\pi, \mathbf{S}_0)$ for a class of RB instances with an unstable Φ matrix, which we call *regular unstable RB*, as defined below.

Definition 1 (Regular unstable RBs). For an RB instance satisfying Assumption 1 and Assumption 2, we call it regular unstable if it has the following properties:

- (a) (LP) has a unique optimal solution y^* , and $y^*(s, 1) + y^*(s, 0) > 0$ for all $s \in \mathbb{S}$.
- (b) The matrix Φ defined in (6) has at least one eigenvalue whose modulus is strictly larger than 1, and the modulus of any eigenvalue of Φ does not equal 1.
- (c) For each $\xi \in \mathbb{R}^{|\mathbb{S}|}$ with $\|\xi\|_2 = 1$ and $\mu^* \xi^\top = 0$, for each large-enough N and each time step t ,

$$\mathbb{E}[(X_{t+1}([N]) - \mu^* - (X_t([N]) - \mu^*)\Phi)\xi^\top \mid X_t, \mathbf{A}_t] \geq \frac{K_{\text{noise}}}{\sqrt{N}} \quad \text{if } \|X_t([N]) - \mu^*\|_1 \leq \eta',$$

where $\eta', K_{\text{noise}} > 0$ are some constants; \mathbf{A}_t follows the Optimal Local Control (Subroutine 2), which is always feasible if $\|X_t([N]) - \mu^*\|_1 \leq \min\{y^*(\tilde{s}, 0), y^*(\tilde{s}, 1)\} - 1/N$.

Our next main theorem is a lower bound for regular unstable RBs; it is proved in Appendix C.2.

Theorem 2. For every regular unstable RB, every policy π and any initial state vector \mathbf{S}_0 , we have

$$R^{\text{rel}} - R(\pi, \mathbf{S}_0) \geq \Omega\left(\frac{1}{\sqrt{N}}\right).$$

Theorem 2 implies that an exponential optimality gap cannot be achieved for a significant subset of RB instances violating local stability (Assumption 3). Therefore, in addition to the widely accepted non-degenerate condition, local stability is also fundamental for exponential optimality.

We briefly comment on Property (c) of regular unstable RB: this property requires the state-count vector to have sufficient randomness in most directions under the Optimal Local Control when the system is near the optimal state distribution μ^* . In Appendix C.3, we provide an intuitive sufficient condition for (c), and a weaker version of the condition under which Theorem 2 remains valid.

The proof of Theorem 2 is based on the following idea: By Lemma 2, arms need to closely follow the Optimal Local Control to maintain a small optimality gap; however, if Optimal Local Control is locally unstable, the system will quickly drift away from the region of the state space where Optimal Local Control is applicable. In the proof, we use Lyapunov functions to quantify how much the system drifts away from μ^* in each step, based on how closely it follows the Optimal Local Control.

We prove the following in Appendix C.4 by giving a concrete example of regular unstable RB.

Corollary 1. There exists an RB satisfying Assumptions 1 and 2 and violating Assumption 3 such that $R^{\text{rel}} - R(\pi, \mathbf{S}_0) \geq \Omega\left(\frac{1}{\sqrt{N}}\right)$.

5 Proof sketch of Theorem 1

The Two-Set Policy maintains two dynamics subsets, each running a different subroutine. The main challenge of the proof lies in analyzing the complex dynamics of the two subsets. To tackle this challenge, we use a novel *multivariate Lyapunov function* that takes not only the system state X_t but also the subsets $D_t^{\text{OL}}, D_t^{\bar{\pi}^*}$ as variables.

Step 1. We first show that the optimality gap can be bounded as follows: $R^{\text{rel}} - R(\pi, \mathbf{S}_0) \leq O(1) \cdot \mathbb{P}[D_\infty^{\text{OL}} \neq [N]]$, where the subscript ∞ indicates the steady-state distribution. The proof of this bound is similar to those of Theorems 3 and 8 of [GGY23a] and is provided in Lemma 8.

Step 2. Next, by the definition of D_t^{OL} , it is not hard to see that if $\|X_t([N]) - \mu^*\|_U \leq \eta$, then $D_t^{\text{OL}} = [N]$. Therefore, it suffices to show that $\mathbb{P}[\|X_\infty([N]) - \mu^*\|_U > \eta]$ is exponentially small. To bound the distance $\|X_\infty([N]) - \mu^*\|_U$, an important step is to construct proper bivariate functions that capture the dynamics of the arms in the subsets D_t^{OL} and $D_t^{\bar{\pi}^*}$. We define

$$h_W(x, D) = \|x(D) - m(D)\mu^*\|_W, \quad h_U(x, D) = \|x(D) - m(D)\mu^*\|_U,$$

where W is a $|\mathcal{S}|$ -by- $|\mathcal{S}|$ weight matrix and $v_W = \sqrt{vWv^\top}$ for $v \in \mathbb{R}^{|\mathcal{S}|}$. The matrices W and U are designed such that $P_{\bar{\pi}^*}$ and Φ are pseudo-contractions on $\Delta(\mathcal{S})$ with fixed point μ^* under the U -weighted norm and the W -weighted norm, respectively (Lemma 6). The bivariate functions h_W and h_U are referred to as *subset Lyapunov functions*, because they witness the convergence of state distributions on a fixed subset of arms when that subset continues to follow either the Exact Proportional Control or the Optimal Local Control (Lemma 10).

Step 3. We use subset Lyapunov functions h_W and h_U as building block for constructing our global Lyapunov function $V(x, D^{\text{OL}}, D^{\bar{\pi}^*})$, defined as

$$V(x, D^{\text{OL}}, D^{\bar{\pi}^*}) = h_U(x, D^{\text{OL}}) + h_W(x, D^{\bar{\pi}^*}) + L(1 - m(D^{\text{OL}})),$$

where L is a constant ensuring that $V(x, D^{\text{OL}}, D^{\bar{\pi}^*})$ bounds $h_U(x, [N]) = \|X_\infty([N]) - \mu^*\|_U$. We show that $V(X_t, D_t^{\text{OL}}, D_t^{\bar{\pi}^*})$ decreases by a large amount every time step with high probability, unless $V(X_t, D_t^{\text{OL}}, D_t^{\bar{\pi}^*}) = O(1/\sqrt{N})$, which implies that the Lyapunov function is small in steady-state.

A key idea in analyzing the changes of $V(X_t, D_t^{\text{OL}}, D_t^{\bar{\pi}^*})$ is to analyze those changes caused by state transitions and those caused by set-updates separately, leading to the key inequality below:

$$\begin{aligned} & V(X_{t+1}, D_{t+1}^{\text{OL}}, D_{t+1}^{\bar{\pi}^*}) - V(X_t, D_t^{\text{OL}}, D_t^{\bar{\pi}^*}) \\ & \leq (h_U(X_{t+1}, D_t^{\text{OL}}) - h_U(X_t, D_t^{\text{OL}})) + (h_W(X_{t+1}, D_t^{\bar{\pi}^*}) - h_W(X_t, D_t^{\bar{\pi}^*})) \\ & \quad + 4(\lambda_U^{1/2} + \lambda_W^{1/2})m(D_t^{\text{OL}} \setminus D_{t+1}^{\text{OL}}) + O\left(\frac{1}{N}\right). \end{aligned}$$

To upper bound the RHS of this inequality, we prove two lemmas to show that $m(D_t^{\text{OL}} \setminus D_{t+1}^{\text{OL}})$ is small (Lemma 11) and that $(h_U(X_{t+1}, D_t^{\text{OL}}) - h_U(X_t, D_t^{\text{OL}})) + (h_W(X_{t+1}, D_t^{\bar{\pi}^*}) - h_W(X_t, D_t^{\bar{\pi}^*}))$ is sufficiently negative (Lemma 12). See Appendix B for the full details of the proof.

References

- [BS20] David B. Brown and James E. Smith. Index policies and performance bounds for dynamic selection problems. *Management Science*, 66(7):3029–3050, 2020.
- [GGY20] Nicolas Gast, Bruno Gaujal, and Chen Yan. Exponential convergence rate for the asymptotic optimality of whittle index policy. *arXiv:2012.09064 [cs.PF]*, 2020.

- [GGY23a] Nicolas Gast, Bruno Gaujal, and Chen Yan. Exponential asymptotic optimality of whittle index policy. *Queueing Systems*, 104(1):107–150, 2023.
- [GGY23b] Nicolas Gast, Bruno Gaujal, and Chen Yan. Linear program-based policies for restless bandits: Necessary and sufficient conditions for (exponentially fast) asymptotic optimality. *Math. Oper. Res.*, 2023.
- [GGY24] Nicolas Gast, Bruno Gaujal, and Chen Yan. Reoptimization nearly solves weakly coupled markov decision processes. *arXiv preprint arXiv:2211.01961*, 2024.
- [GNJT23] Abheek Ghosh, Dheeraj Nagaraj, Manish Jain, and Milind Tambe. Indexability is not enough for whittle: Improved, near-optimal algorithms for restless bandits. In *Proc. Int. Conf. Autonomous Agents and Multiagent Syst.*, page 1294–1302, Richland, SC, 2023.
- [GT56] Alan J Goldman and Albert W Tucker. Theory of linear programming. *Linear inequalities and related systems*, 38:53–97, 1956.
- [HF17] Weici Hu and Peter Frazier. An asymptotically optimal index policy for finite-horizon restless bandits. *arXiv:1707.00205 [math.OC]*, 2017.
- [HXCW23] Yige Hong, Qiaomin Xie, Yudong Chen, and Weina Wang. Restless bandits with average reward: Breaking the uniform global attractor assumption. In A. Oh, T. Neumann, A. Globerson, K. Saenko, M. Hardt, and S. Levine, editors, *Advances in Neural Information Processing Systems*, volume 36, pages 12810–12844. Curran Associates, Inc., 2023.
- [HXCW24] Yige Hong, Qiaomin Xie, Yudong Chen, and Weina Wang. Unichain and aperiodicity are sufficient for asymptotic optimality of average-reward restless bandits. *arXiv preprint arXiv:2402.05689*, 2024.
- [NM23] José Niño-Mora. Markovian restless bandits and index policies: A review. *Mathematics*, 11(7), 2023.
- [PT99] Christos H. Papadimitriou and John N. Tsitsiklis. The complexity of optimal queuing network control. *Math. Oper. Res.*, 24(2):293–305, 1999.
- [Put05] Martin L Puterman. *Markov decision processes: Discrete stochastic dynamic programming*. John Wiley & Sons, 2005.
- [Ver16] I. M. Verloop. Asymptotically optimal priority policies for indexable and nonindexable restless bandits. *Ann. Appl. Probab.*, 26(4):1947–1995, 2016.
- [Whi88] Peter Whittle. Restless bandits: activity allocation in a changing world. *J. Appl. Probab.*, 25:287 – 298, 1988.
- [WW90] Richard R. Weber and Gideon Weiss. On an index policy for restless bandits. *J. Appl. Probab.*, 27(3):637–648, 1990.
- [Yan24] Chen Yan. An optimal-control approach to infinite-horizon restless bandits: Achieving asymptotic optimality with minimal assumptions. *arXiv preprint arXiv:2403.11913*, 2024.

- [ZCJW19] Gabriel Zayas-Cabán, Stefanus Jasin, and Guihua Wang. An asymptotically optimal heuristic for general nonstationary finite-horizon restless multi-armed, multi-action bandits. *Advances in Applied Probability*, 51:745–772, 2019.
- [ZF21] Xiangyu Zhang and Peter I. Frazier. Restless bandits with many arms: Beating the central limit theorem. *arXiv:2107.11911 [math.OC]*, 2021.
- [ZF22] Xiangyu Zhang and Peter I. Frazier. Near-optimality for infinite-horizon restless bandits with many arms. *arXiv:2203.15853 [cs.LG]*, 2022.

A Preliminary lemmas and proofs

In this section, we state and prove several preliminary lemmas. In Appendix A.1, we show that the linear program (LP) is a relaxation of the N -armed problem. In Appendix A.2, we prove two lemmas about the transition dynamics under the two subroutines, Exact Proportional Control (Subroutine 1) and Optimal Local Control (Subroutine 2). In Appendix A.3, we define two weight matrices, W and U , and show the pseudo-contraction properties of P_{π^*} and Φ under the L_2 norms weighted by W and U , respectively.

A.1 Justifying the LP relaxation

In this section, we show that linear program (LP) is indeed a relaxation of the N -armed problem, i.e., $R^{\text{rel}} \geq R^*(N)$. While this fact is used throughout the literature, our paper assumes weaker assumptions than most prior work, so for the sake of rigor, we prove it in our setting for completeness.

Lemma 3 (LP relaxation). *Let R^{rel} be the optimal value of the linear program (LP), and let $R^*(N)$ be the optimal reward of the N -armed restless bandit problem (RB). Then we have*

$$R^{\text{rel}} \geq R^*(N). \quad (13)$$

Proof. For any stationary Markovian policy π , define

$$y^\pi(s, a) \triangleq \lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \left[\frac{1}{N} \sum_{i \in [N]} \mathbb{1}\{S_t^\pi(i) = s, A_t^\pi(i) = a\} \right] \quad \forall s \in \mathbb{S}, a \in \mathbb{A}.$$

We first show that $R(\pi, \mathbf{S}_0) = \sum_{s \in \mathbb{S}, a \in \mathbb{A}} r(s, a) y^\pi(s, a)$.

$$\begin{aligned} \sum_{s \in \mathbb{S}, a \in \mathbb{A}} r(s, a) y^\pi(s, a) &= \sum_{s \in \mathbb{S}, a \in \mathbb{A}} r(s, a) \lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \left[\frac{1}{N} \sum_{i \in [N]} \mathbb{1}\{S_t^\pi(i) = s, A_t^\pi(i) = a\} \right] \\ &= \lim_{T \rightarrow \infty} \frac{1}{NT} \sum_{t=0}^{T-1} \sum_{i \in [N]} \mathbb{E} \left[\sum_{s \in \mathbb{S}, a \in \mathbb{A}} r(s, a) \mathbb{1}\{S_t^\pi(i) = s, A_t^\pi(i) = a\} \right] \\ &= \lim_{T \rightarrow \infty} \frac{1}{NT} \sum_{t=0}^{T-1} \sum_{i \in [N]} \mathbb{E} [r(S_t^\pi(i), A_t^\pi(i))] \\ &= R(\pi, \mathbf{S}_0). \end{aligned}$$

Next, we show that $(y^\pi(s, a))_{s \in \mathbb{S}, a \in \mathbb{A}}$ satisfies the constraints of (LP). We first consider (3):

$$\begin{aligned}
\sum_{s \in \mathbb{S}} y^\pi(s, 1) &= \sum_{s \in \mathbb{S}} \lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \left[\frac{1}{N} \sum_{i \in [N]} \mathbb{1}\{S_t^\pi(i) = s, A_t^\pi(i) = 1\} \right] \\
&= \lim_{T \rightarrow \infty} \frac{1}{NT} \sum_{t=0}^{T-1} \sum_{i \in [N]} \mathbb{E} \left[\sum_{s \in \mathbb{S}} \mathbb{1}\{S_t^\pi(i) = s, A_t^\pi(i) = 1\} \right] \\
&= \lim_{T \rightarrow \infty} \frac{1}{NT} \sum_{t=0}^{T-1} \alpha N \\
&= \alpha.
\end{aligned}$$

Next, we look at the constraint (4):

$$\begin{aligned}
\sum_{s' \in \mathbb{S}, a \in \mathbb{A}} y^\pi(s', a) P(s', a, s) &= \lim_{T \rightarrow \infty} \frac{1}{NT} \sum_{t=0}^{T-1} \sum_{i \in [N]} \sum_{s \in \mathbb{S}, a \in \mathbb{A}} P(s', a, s) \mathbb{P}[S_t^\pi(i) = s', A_t^\pi(i) = a] \\
&= \lim_{T \rightarrow \infty} \frac{1}{NT} \sum_{t=0}^{T-1} \sum_{i \in [N]} \mathbb{P}[S_{t+1}^\pi(i) = s] \\
&= \lim_{T \rightarrow \infty} \frac{1}{NT} \sum_{t=1}^T \sum_{i \in [N]} \mathbb{P}[S_t^\pi(i) = s] \\
&= \sum_{a \in \mathbb{A}} y^\pi(s, a) \\
\sum_{s \in \mathbb{S}, a \in \mathbb{A}} y^\pi(s, a) &= \frac{1}{NT} \sum_{t=0}^{T-1} \sum_{i \in [N]} \mathbb{E} \left[\sum_{s \in \mathbb{S}, a \in \mathbb{A}} \mathbb{1}\{S_t^\pi(i) = s, A_t^\pi(i) = a\} \right] = 1,
\end{aligned}$$

and it is obvious that $\sum_{s \in \mathbb{S}, a \in \mathbb{A}} y^\pi(s, a) \geq 0$.

Combining the above argument, $(y^\pi(s, a))_{s \in \mathbb{S}, a \in \mathbb{A}}$ is a feasible solution to (LP), so $R(\pi, \mathbf{S}_0) = \sum_{s \in \mathbb{S}, a \in \mathbb{A}} r(s, a) y^\pi(s, a) \leq R^{\text{rel}}$.

By standard results for MDP with finite state and action spaces, there always exists a stationary Markovian policy whose long-run average reward achieves the optimal reward [Put05, Theorem 9.1.8]. Letting π be this optimal stationary Markovian policy, then $R^*(N) = R(\pi, \mathbf{S}_0) \leq R^{\text{rel}}$. \square

A.2 Transition dynamics under subroutines

The lemma below characterizes the dynamics of a N -armed RB system if all arms follow Exact Proportional Control (Subroutine 1).

Lemma 4. *For any time step t ,*

$$\mathbb{E}[X_{t+1}([N]) \mid X_t, \text{all arms follow Subroutine 1}] = X_t([N]) P_{\bar{\pi}^*} \quad a.s., \quad (14)$$

where expectation is taken entry-wise for the random vector $X_{t+1}([N])$; $P_{\bar{\pi}^*}$ is the transition matrix induced by $\bar{\pi}^*$ on the single-armed MDP, i.e., $P_{\bar{\pi}^*}(s, s') = \sum_{a \in \mathbb{A}} \bar{\pi}^*(a|s) P(s, a, s')$ for $s, s' \in \mathbb{S}$.

Proof. By definition, if all N arms follow Exact Proportional Control at time t , the expected number of arms in state s taking action a is equal to $X_t([N], s)\bar{\pi}^*(a|s)$. Among these arms, the fraction of arms transitioning to state s' in the next time step equals $P(s, a, s')$. Since the state transition of each arm is Markovian given the state and action, the expected number of arms going from state s to state s' and taking action a equals $X_t([N], s)\bar{\pi}^*(a|s)P(s, a, s')$. Summing over s and a , we get

$$\begin{aligned} & \mathbb{E} [X_{t+1}([N], s') | X_t, \text{all arms follow Subroutine 1}] \\ &= \sum_{s \in \mathbb{S}, a \in \mathbb{A}} X_t([N], s)\bar{\pi}^*(a|s)P(s, a, s') \quad \forall s' \in \mathbb{S}, \end{aligned}$$

which completes the proof of the lemma. \square

If all arms in the RB system follow Optimal Local Control (Subroutine 2), the expectation of the scaled state-count vector $X_t([N])$ has a linear dynamics, as stated in the lemma below. The lemma is basically a restatement of Lemma 2 in [GGY23a], although in [GGY23a], the LP-Priority policy was considered, and it was assumed that $y^*(s, 1) + y^*(s, 0) > 0$ for all $s \in \mathbb{S}$.

Lemma 1. *Under Assumption 2, suppose (7) holds for the N -armed system at time step t . We have*

$$\mathbb{E}[X_{t+1}([N]) - \mu^* | X_t, \text{all arms follow Subroutine 2}] = (X_t([N]) - \mu^*)\Phi, \quad (8)$$

where Φ is the $|\mathbb{S}|$ -by- $|\mathbb{S}|$ matrix defined in (6).

Proof. Throughout this proof, we use the shorthand x for $x([N])$, and omit “ $\mathbf{A}_t \sim$ Optimal Local Control” in the conditionings.

We fixed an arbitrary x and let $X_t = x$. For any $s \neq \tilde{s}$ and $a \in \mathbb{A}$, the expected number of arms taking action a equals $x(s)\bar{\pi}^*(a|s)$; for $s = \tilde{s}$, the expected number of arms taking action 1 equals $\alpha - \sum_{s \neq \tilde{s}} x(s)\bar{\pi}^*(1|s)$. and the expected number of arms taking action 0 equals $1 - \alpha - \sum_{s \neq \tilde{s}} x(s)\bar{\pi}^*(0|s)$. Therefore, for any $s' \in \mathbb{S}$ such that $s' \neq \tilde{s}$,

$$\begin{aligned} & \mathbb{E} [X_{t+1}(s') | X_t = x] \\ &= \sum_{s \neq \tilde{s}, a \in \mathbb{A}} x(s)\bar{\pi}^*(a|s)P(s, a, s') \\ & \quad + \left(\alpha - \sum_{s \neq \tilde{s}} x(s)\bar{\pi}^*(1|s) \right) P(\tilde{s}, 1, s') + \left(1 - \alpha - \sum_{s \neq \tilde{s}} x(s)\bar{\pi}^*(0|s) \right) P(\tilde{s}, 0, s') \\ &= \sum_{s \neq \tilde{s}} x(s)P_{\bar{\pi}^*}(s, s') + \left(\alpha - \sum_{s \neq \tilde{s}} x(s)\bar{\pi}^*(1|s) \right) P_1(\tilde{s}, s') + \left(1 - \alpha - \sum_{s \neq \tilde{s}} x(s)\bar{\pi}^*(0|s) \right) P_0(\tilde{s}, s') \\ &= \sum_{s \in \mathbb{S}} x(s)P_{\bar{\pi}^*}(s, s') - x(\tilde{s})P_{\bar{\pi}^*}(\tilde{s}, s') \\ & \quad + \left(\alpha - \sum_{s \neq \tilde{s}} x(s)\bar{\pi}^*(1|s) \right) P_1(\tilde{s}, s') + \left(1 - \alpha - \sum_{s \neq \tilde{s}} x(s)\bar{\pi}^*(0|s) \right) P_0(\tilde{s}, s'). \end{aligned} \quad (15)$$

Because $P_{\bar{\pi}^*}(\tilde{s}, s') = \bar{\pi}^*(1|\tilde{s})P_1(\tilde{s}, s') + \bar{\pi}^*(0|\tilde{s})P_0(\tilde{s}, s')$, we expand the second term in (15) and rearrange the terms to get

$$\begin{aligned} & \mathbb{E} [X_{t+1}(s') | X_t = x] \\ &= \sum_{s \in \mathbb{S}} x(s)P_{\bar{\pi}^*}(s, s') + \left(\alpha - \sum_{s \in \mathbb{S}} x(s)\bar{\pi}^*(1|s) \right) P_1(\tilde{s}, s') + \left(1 - \alpha - \sum_{s \in \mathbb{S}} x(s)\bar{\pi}^*(0|s) \right) P_0(\tilde{s}, s') \end{aligned}$$

$$= \sum_{s \in \mathbb{S}} x(s) P_{\bar{\pi}^*}(s, s') + \left(\alpha - \sum_{s \in \mathbb{S}} x(s) \bar{\pi}^*(1|s) \right) P_1(\tilde{s}, s') - \left(\alpha - \sum_{s \in \mathbb{S}} x(s) \bar{\pi}^*(1|s) \right) P_0(\tilde{s}, s') \quad (16)$$

$$= \sum_{s \in \mathbb{S}} x(s) \left(P_{\bar{\pi}^*}(s, s') - \bar{\pi}^*(1|s) (P_1(\tilde{s}, s') - P_0(\tilde{s}, s')) \right) + \alpha (P_1(\tilde{s}, s') - P_0(\tilde{s}, s')), \quad (17)$$

where (16) uses the facts that $\bar{\pi}^*(0|s) = 1 - \bar{\pi}^*(1|s)$ and $\sum_{s \in \mathbb{S}} x(s) = 1$.

Next we calculate $\mathbb{E}[X_{t+1}(\tilde{s}) | X_t = x]$. Observe that $\mathbb{E}[X_t(\tilde{s})] = 1 - \sum_{s' \neq \tilde{s}} \mathbb{E}[X_t(s')]$, so

$$\begin{aligned} & \mathbb{E}[X_{t+1}(\tilde{s}) | X_t = x] \\ &= 1 - \sum_{s' \neq \tilde{s}} \mathbb{E}[X_t(s')] \\ &= \sum_{s \in \mathbb{S}} x(s) \left(1 - \sum_{s' \neq \tilde{s}} P_{\bar{\pi}^*}(s, s') + \sum_{s' \neq \tilde{s}} \bar{\pi}^*(1|s) (P_1(\tilde{s}, s') - P_0(\tilde{s}, s')) \right) - \alpha \sum_{s' \neq \tilde{s}} (P_1(\tilde{s}, s') - P_0(\tilde{s}, s')) \\ &= \sum_{s \in \mathbb{S}} x(s) \left(P_{\bar{\pi}^*}(s, \tilde{s}) - \bar{\pi}^*(1|s) (P_1(\tilde{s}, \tilde{s}) - P_0(\tilde{s}, \tilde{s})) \right) + \alpha (P_1(\tilde{s}, \tilde{s}) - P_0(\tilde{s}, \tilde{s})). \end{aligned} \quad (18)$$

Note that (18) has the same form as (17), with s' replaced by \tilde{s} .

We rewrite (17) and (18) in matrix form to get

$$\mathbb{E}[X_{t+1} | X_t = x] = x \left(P_{\bar{\pi}^*} - c_{\bar{\pi}^*}^\top (P_1(\tilde{s}) - P_0(\tilde{s})) \right) + \alpha (P_1(\tilde{s}) - P_0(\tilde{s})). \quad (19)$$

Recall that we have let $\Phi = P_{\bar{\pi}^*} - \mathbb{1}^\top \mu^* - (c_{\bar{\pi}^*} - \alpha \mathbb{1})^\top (P_1(\tilde{s}) - P_0(\tilde{s}))$, we can verify that the RHS of (19) is the same as $(x - \mu^*)\Phi + \mu^*$. Specifically,

$$\begin{aligned} (x - \mu^*)\Phi + \mu^* &= (x - \mu^*) \left(P_{\bar{\pi}^*} - \mathbb{1}^\top \mu^* - (c_{\bar{\pi}^*} - \alpha \mathbb{1})^\top (P_1(\tilde{s}) - P_0(\tilde{s})) \right) + \mu^* \\ &= x \left(P_{\bar{\pi}^*} - c_{\bar{\pi}^*}^\top (P_1(\tilde{s}) - P_0(\tilde{s})) \right) + x \left(-\mathbb{1}^\top \mu^* - \alpha \mathbb{1}^\top (P_1(\tilde{s}) - P_0(\tilde{s})) \right) \\ &\quad - \mu^* \left(P_{\bar{\pi}^*} - c_{\bar{\pi}^*}^\top (P_1(\tilde{s}) - P_0(\tilde{s})) \right) - \mu^* \left(-\mathbb{1}^\top \mu^* - \alpha \mathbb{1}^\top (P_1(\tilde{s}) - P_0(\tilde{s})) \right) + \mu^* \\ &= x \left(P_{\bar{\pi}^*} - c_{\bar{\pi}^*}^\top (P_1(\tilde{s}) - P_0(\tilde{s})) \right) - \mu^* - \alpha (P_1(\tilde{s}) - P_0(\tilde{s})) \\ &\quad - \mu^* + \alpha (P_1(\tilde{s}) - P_0(\tilde{s})) + \mu^* + \alpha (P_1(\tilde{s}) - P_0(\tilde{s})) + \mu^* \\ &= x \left(P_{\bar{\pi}^*} - c_{\bar{\pi}^*}^\top (P_1(\tilde{s}) - P_0(\tilde{s})) \right) + \alpha (P_1(\tilde{s}) - P_0(\tilde{s})). \end{aligned} \quad (20)$$

where to get (20) we use the facts that $x \mathbb{1}^\top = \mu^* \mathbb{1}^\top = 1$ and $\mu^* c_{\bar{\pi}^*}^\top = \alpha$. Substituting the RHS of (19) with $(x - \mu^*)\Phi + \mu^*$, we have proved that under Optimal Local Control, $\mathbb{E}[X_{t+1} | X_t = x] = (x - \mu^*)\Phi + \mu^*$. This completes the proof of Lemma 1. \square

A.3 Weighted L_2 norms for quantifying the convergence of distributions

As discussed in Section 3.2 and Section 3.3, if a set of arms follow Exact Proportional Control or Optimal Local Control for a long time, the distribution of their states converge to the optimal stationary distribution μ^* . In our analysis, it is convenient to define a proper metric to witness this convergence in each time step and quantify the convergence rate. For this purpose, we introduce two matrices, W and U , and study the W -weighted (U -weighted) L_2 norm on \mathbb{S} . Note that the same matrix W and W -weighted norm have also been used in [HXCW24].

The definition of W and U are given as follows.

Definition 2. Assume Assumption 1, 2 and 3. Let W and U be $|\mathbb{S}|$ -by- $|\mathbb{S}|$ matrices given by

$$W = \sum_{k=0}^{\infty} (P_{\bar{\pi}^*} - \mathbb{1}^\top \mu^*)^k ((P_{\bar{\pi}^*} - \mathbb{1}^\top \mu^*)^\top)^k \quad (21)$$

$$U = \sum_{k=0}^{\infty} \Phi^k (\Phi^\top)^k, \quad (22)$$

Next, we show that the matrices W and U are well-defined and positive definite, and lower bound their eigenvalues by 1. Then if we let λ_W and λ_U denote maximal eigenvalues of W and U , the eigenvalues of W and U are in the range $[1, \lambda_W]$ and $[1, \lambda_U]$, respectively.

Lemma 5. Assume Assumption 1, 2 and 3, the matrices W and U given in Definition 2 are well-defined. Moreover, their eigenvalues are lower bounded by 1.

Proof. The well-definedness and eigenvalue lower bound for W has been proved in [HXCW24]. The main fact used in the proof is that all eigenvalues of $P_{\bar{\pi}^*} - \mathbb{1}^\top \mu^*$ have moduli strictly less than 1 under Assumption 1. Since we have assumed the same things for Φ in Assumption 3, the proof for U is analogous with $P_{\bar{\pi}^*} - \mathbb{1}^\top \mu^*$ replaced by Φ . \square

With matrices W and U , we can define some weighted L_2 norms. Let $\|\cdot\|_W$ denote the W -weighted L_2 norm, given by $\|u\|_W = \sqrt{uWu^\top}$ for any row vector $u \in \mathbb{R}^{|\mathbb{S}|}$; let $\|\cdot\|_U$ denote the U -weighted L_2 norm, defined similarly.

Lemma 6 below shows that $P_{\bar{\pi}^*}$ and Φ are pseudo-contractions on the probability simplex $\Delta(\mathbb{S})$, under the W -weighted L_2 norm and the U -weighted L_2 norm, respectively.

Lemma 6 (Pseudo-contraction under the weighted L_2 norms). Assume Assumption 1, 2 and 3. For any distribution $v \in \Delta(\mathbb{S})$, we have

$$\|(v - \mu^*)P_{\bar{\pi}^*}\|_W \leq \rho_w \|v - \mu^*\|_W \quad (23)$$

$$\|(v - \mu^*)\Phi\|_U \leq \rho_u \|v - \mu^*\|_U. \quad (24)$$

where $\|\cdot\|_W$ is the W -weighted L_2 norm, i.e., $\|u\|_W = \sqrt{uWu^\top}$ for any row vector $u \in \mathbb{R}^{|\mathbb{S}|}$; $\|\cdot\|_U$ is the U -weighted L_2 norm defined similarly; $\rho_w = 1 - 1/(2\lambda_W)$; $\rho_u = 1 - 1/(2\lambda_U)$.

Proof. The inequality (23) has been proved in [HXCW24]. The proof of (24) is analogous. \square

B Proving the exponential asymptotic optimality of Two-Set Policy (Theorem 1)

In this section, we prove Theorem 1, which states that Two-Set Policy achieve exponential asymptotic optimality under Assumptions 1, 2 and 3. In Appendix B.1, we state all the lemmas needed for proving Theorem 1. Then in Appendix B.2, we prove Theorem 1 using these lemmas. Finally, we prove the lemmas. In Appendix B.3, we first prove the core lemmas that are closely related to the properties of Two-Set Policy; then in Appendix B.4, we prove the rest of the lemmas, which are technical and relatively routine.




Figure 1: The flowchart for the proof of Theorem 1. We omit Lemma 7 from the flowchart for readability since it is used in the proofs of all other lemmas and the theorem.

B.1 Lemma statements

In this section, we state a sequence of lemmas to prepare for proving Theorem 1. The logical relationship between the lemmas and the theorem is illustrated in the flowchart Figure 1.

We first introduce a few notational conventions. Note that the Two-Set Policy in Algorithm 1 induces a finite-state Markov chain whose state at time t is $(X_t, D_t^{\text{OL}}, D_t^{\bar{\pi}^*})$. We use Σ_t as a shorthand for $(X_t, D_t^{\text{OL}}, D_t^{\bar{\pi}^*})$. We use $X_\infty, D_\infty^{\text{OL}}, D_\infty^{\bar{\pi}^*}, \Sigma_\infty$ to represent variables following long-run average distribution: For example, for any function f of Σ_t ,

$$\mathbb{E}[f(\Sigma_\infty)] \triangleq \lim_{t \rightarrow \infty} \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E}[f(\Sigma_t)],$$

where the limit always exists because Σ_t is a finite-state Markov chain, though it could depend on initial state $(X_0, D_0^{\text{OL}}, D_0^{\bar{\pi}^*})$.

Our first lemma, Lemma 7, shows that the arms in D^{OL} and $D^{\bar{\pi}^*}$ can indeed follow the intended subroutine when η is small enough. This lemma is proved in Appendix B.3.

Lemma 7 (Subroutine conformity). *Assume Assumptions 1, 2 and 3 hold. Consider the Two-Set Policy described in Algorithm 1 with $\eta = |\mathbb{S}|^{-1/2} (\min\{y^*(\tilde{s}, 0), y^*(\tilde{s}, 1)\} - (|S^\emptyset| + 1)/N)$: The steps in Line 5-7 are well-defined. In other words, we can always let the arms in D_t^{OL} follow Optimal Local Control, let the arms in $D_t^{\bar{\pi}^*}$ follow Exact Proportional Control, and choose actions for the arms not in $D_t^{\text{OL}} \cup D_t^{\bar{\pi}^*}$ to satisfy the budget constraint $\sum_{i \in [N]} A_t(i) = \alpha N$.*

Our next lemma, Lemma 8, relates the optimality gap with the probability of $D_\infty^{\text{OL}} \neq [N]$. We prove it in Appendix B.4 with a similar argument as the proofs of Theorem 3 and Theorem 8 in [GGY23a]; however, we circumvent the use of UGAP in our proof.

Lemma 8. *Assume Assumptions 1, 2 and 3 hold. The optimality gap under the Two-Set Policy (Algorithm 1) is bounded as*

$$R^{\text{rel}} - R(\pi, \mathbf{S}_0) \leq O(1) \cdot \mathbb{P}[D_\infty^{\text{OL}} \neq [N]]. \quad (25)$$

Lemma 8 implies that it suffices to show that $\mathbb{P}[D_\infty^{\text{OL}} \neq [N]]$ is exponentially small, which we will prove via a Lyapunov argument, with the help of Lemma 9 below. Lemma 9 is proved in Appendix B.4.

Lemma 9. *Suppose there exists a function $V(x, D^{\text{OL}}, D^{\bar{\pi}^*})$ and constants V_{\max} and $\bar{\eta} > 0$ such that for any $t \geq 0$,*

$$0 \leq V(\Sigma_t) \leq V_{\max} \quad a.s.; \quad (26)$$

if $V(\Sigma_t) > \bar{\eta}$,

$$\mathbb{E}[V(\Sigma_{t+1}) - V(\Sigma_t) \mid \Sigma_t] \leq -\gamma \quad (27)$$

$$\mathbb{E}[(V(\Sigma_{t+1}) - V(\Sigma_t))^+ \mid \Sigma_t] = O(1/\sqrt{N}); \quad (28)$$

if $\bar{\eta}/2 < V(\Sigma_t) \leq \bar{\eta}$,

$$\mathbb{E}[(V(\Sigma_{t+1}) - V(\Sigma_t))^+ \mid \Sigma_t] \leq O(\exp(-CN)); \quad (29)$$

if $V(\Sigma_t) \leq \bar{\eta}/2$,

$$\mathbb{P}[V(\Sigma_{t+1}) > \bar{\eta}/2 \mid \Sigma_t] \leq O(\exp(-C'N)), \quad (30)$$

where γ, C, C' are positive constants. Then we have

$$\mathbb{P}[V(\Sigma_\infty) > \bar{\eta}] = O(\exp(-C''N)), \quad (31)$$

where C'' is another positive constant.

The remaining lemmas prepare for constructing the Lyapunov function $V(x, D^{\text{OL}}, D^{\bar{\pi}^*})$ satisfying the conditions in Lemma 9. We first define two classes of functions, $h_W = \{h_W(x, D)\}_{D \subseteq [N]}$ and $h_U = \{h_U(x, D)\}_{D \subseteq [N]}$: For any system state x and subset $D \subseteq [N]$, define

$$h_W(x, D) = \|x(D) - m(D)\mu^*\|_W \quad (32)$$

$$h_U(x, D) = \|x(D) - m(D)\mu^*\|_U. \quad (33)$$

These two classes of functions quantify the convergence of state distributions for subsets of arms that follow Exact Proportional Control or Optimal Local Control. Such functions are referred to as *subset Lyapunov functions* and are used as building blocks in the Lyapunov analysis of the prior work [HXCW24].

Next, we show several properties of h_W and h_U . They are proved in Appendix B.4 using properties of the two weighted L_2 norms.

Lemma 10 (Properties of h_W and h_U). *Assume Assumptions 1, 2 and 3 hold. For any time step t and $D \subseteq [N]$, let X'_{t+1} be the system state at time $t+1$ if the arms in D follow Exact Proportional Control at time t , then we have*

$$\mathbb{E}[(h_W(X'_{t+1}, D) - \rho_w h_W(X_t, D))^+ \mid X_t] \leq O(1/\sqrt{N}) \quad a.s. \quad (34)$$

$$\mathbb{P}[h_W(X'_{t+1}, D) > \rho_w h_W(X_t, D) + r \mid X_t] \leq O(\exp(-C Nr^2)) \quad a.s. \quad \forall r \geq 0, \quad (35)$$

where $\rho_w = 1 - 1/(2\lambda_W)$, and C is a positive constant. In addition, for any $D, D' \subseteq [N]$,

$$h_W(x, D) \geq \frac{1}{|\mathbb{S}|^{1/2}} \|x(D) - m(D)\mu^*\|_1 \quad (36)$$

$$|h_W(x, D) - h_W(x, D')| \leq 2\lambda_W^{1/2} (m(D' \setminus D) + m(D \setminus D')), \quad (37)$$

The same inequalities hold for h_U with Exact Proportional Control and W replaced by the Optimal Local Control and U , which we omit here.

Finally, we state two important lemmas characterizing the dynamics of the set D_t^{OL} . In Lemma 11, we show that the arms in D_t^{OL} are most likely to remain in D_{t+1}^{OL} , so that D_t^{OL} is almost non-shrinking in the set-inclusive sense. In Lemma 12, we utilize the maximality of D_t^{OL} to show that the fraction of arms not in D_t^{OL} is small when $X_t(D_t^{\bar{\pi}^*})$ is close to $m(D_t^{\bar{\pi}^*})\mu^*$ or when $X_t([N])$ is close to μ^* . These two lemmas imply that D_t^{OL} tends to gradually expand and cover all arms as the two subroutines drive the states of the arms in D_t^{OL} and $D_t^{\bar{\pi}^*}$ to the optimal distribution. Both lemmas are proved in Appendix B.3.

Lemma 11 (Almost non-shrinking). *Assume Assumptions 1, 2 and 3 hold. Under the Two-Set Policy (Algorithm 1), for any $t \geq 0$,*

$$\mathbb{E}[m(D_t^{\text{OL}} \setminus D_{t+1}^{\text{OL}}) \mid \Sigma_t] \leq O(m(D_t^{\text{OL}}) \exp(-CNm(D_t^{\text{OL}})^2)) \quad a.s. \quad (38)$$

$$\mathbb{P}[m(D_t^{\text{OL}} \setminus D_{t+1}^{\text{OL}}) > 0 \mid \Sigma_t] \leq O(\exp(-CNm(D_t^{\text{OL}})^2)) \quad a.s., \quad (39)$$

where C is a positive constant.

Lemma 12 (Sufficient coverage). *Assume Assumptions 1, 2 and 3 hold. Consider the Two-Set Policy (Algorithm 1) with $\eta = |\mathcal{S}|^{-1/2}(\min\{y^*(\bar{s}, 0), y^*(\bar{s}, 1)\} - (|\mathcal{S}^\emptyset| + 1)/N)$. For any $t \geq 0$,*

$$1 - m(D_t^{\text{OL}}) \leq \frac{\lambda_U^{1/2}}{\eta\beta} h_W(X_t, D_t^{\bar{\pi}^*}) + \frac{2}{\beta N} \quad a.s. \quad (40)$$

$$m(D_t^{\text{OL}}) = 1 \quad \text{if } h_U(X_t, [N]) \leq \eta. \quad (41)$$

B.2 Proof of Theorem 1

Proof. Define the Lyapunov function $V(x, D^{\text{OL}}, D^{\bar{\pi}^*})$ as

$$V(x, D^{\text{OL}}, D^{\bar{\pi}^*}) = h_U(x, D^{\text{OL}}) + h_W(x, D^{\bar{\pi}^*}) + L(1 - m(D^{\text{OL}})), \quad (42)$$

where $L = 2\lambda_U^{1/2} + 4\lambda_W^{1/2} - 2\lambda_U^{1/2}\beta$. We claim that

$$\mathbb{P}[V(X_\infty, D_\infty^{\text{OL}}, D_\infty^{\bar{\pi}^*}) > \eta] = O(\exp(-CN)), \quad (43)$$

for some positive constant C . To see why (43) helps, note that by Lemma 10,

$$V(x, D^{\text{OL}}, D^{\bar{\pi}^*}) \geq h_U(x, D^{\text{OL}}) + 2\lambda_U^{1/2}(1 - m(D^{\text{OL}})) \geq h_U(x, [N]),$$

so $V(X_\infty, D_\infty^{\text{OL}}, D_\infty^{\bar{\pi}^*}) \leq \eta$ implies $h_U(X_\infty, [N]) \leq \eta$, which further implies $D_\infty^{\text{OL}} = [N]$ by Lemma 12. Therefore, suppose we can show (43), with Lemma 8 and the above argument, we have

$$\begin{aligned} R^{\text{rel}} - R(\pi, \mathbf{S}_0) &\leq O(1) \cdot \mathbb{P}[D_\infty^{\text{OL}} \neq [N]] \\ &\leq O(1) \cdot \mathbb{P}[h_U(X_\infty, [N]) > \eta] \\ &\leq O(1) \cdot \mathbb{P}[V(X_\infty, D_\infty^{\text{OL}}, D_\infty^{\bar{\pi}^*}) > \eta] \\ &= O(\exp(-CN)), \end{aligned}$$

which will finish the proof.

In the rest of the proof, we will show (43) by invoking Lemma 9 with $\bar{\eta} = \eta$. First, $V(x, D^{\text{OL}}, D^{\bar{\pi}^*})$ is non-negative and upper bounded by $\lambda_U^{1/2} + \lambda_W^{1/2} + L$, so the condition (26) holds. To show the other conditions, we first examine the difference $V(\Sigma_{t+1}) - V(\Sigma_t)$. Consider the decomposition

$$V(\Sigma_{t+1}) - V(\Sigma_t) = (V(X_{t+1}, D_{t+1}^{\text{OL}}, D_{t+1}^{\bar{\pi}^*}) - V(X_{t+1}, D_t^{\text{OL}}, D_t^{\bar{\pi}^*})) \quad (44)$$

$$+ (V(X_{t+1}, D_t^{\text{OL}}, D_t^{\bar{\pi}^*}) - V(X_t, D_t^{\text{OL}}, D_t^{\bar{\pi}^*})). \quad (45)$$

We call the difference term in RHS of (44) the *set-update term*, and the difference term in (45) the *state-transition term*. We bound them separately.

For the set-update term in (44), we can derive a bound fully in terms of the sizes of sets, using the Lipschitz continuity of $h_W(x, D)$ and $h_U(x, D)$ with respect to D given in (37) of Lemma 10:

$$\begin{aligned} & V(X_{t+1}, D_{t+1}^{\text{OL}}, D_{t+1}^{\bar{\pi}^*}) - V(X_{t+1}, D_t^{\text{OL}}, D_t^{\bar{\pi}^*}) \\ &= h_U(X_{t+1}, D_{t+1}^{\text{OL}}) - h_U(X_{t+1}, D_t^{\text{OL}}) + h_W(X_{t+1}, D_{t+1}^{\bar{\pi}^*}) - h_W(X_{t+1}, D_t^{\bar{\pi}^*}) \\ &\quad - L(m(D_{t+1}^{\text{OL}}) - m(D_t^{\text{OL}})) \\ &\leq 2\lambda_U^{1/2}(m(D_{t+1}^{\text{OL}} \setminus D_t^{\text{OL}}) + m(D_t^{\text{OL}} \setminus D_{t+1}^{\text{OL}})) + 2\lambda_W^{1/2}(m(D_{t+1}^{\bar{\pi}^*} \setminus D_t^{\bar{\pi}^*}) + m(D_t^{\bar{\pi}^*} \setminus D_{t+1}^{\bar{\pi}^*})) \end{aligned} \quad (46)$$

$$- L(m(D_{t+1}^{\text{OL}} \setminus D_t^{\text{OL}}) - m(D_t^{\text{OL}} \setminus D_{t+1}^{\text{OL}})). \quad (47)$$

We can bound $m(D_{t+1}^{\bar{\pi}^*} \setminus D_t^{\bar{\pi}^*})$ and $m(D_t^{\bar{\pi}^*} \setminus D_{t+1}^{\bar{\pi}^*})$ in the last expression in terms of D_{t+1}^{OL} and D_t^{OL} . By definition, $D_{t+1}^{\bar{\pi}^*} \supseteq D_t^{\bar{\pi}^*} \setminus D_{t+1}^{\text{OL}}$, and $D_t^{\bar{\pi}^*}$ is disjoint from D_t^{OL} , so it is not hard to see that

$$D_t^{\bar{\pi}^*} \setminus D_{t+1}^{\bar{\pi}^*} \subseteq D_{t+1}^{\text{OL}} \setminus D_t^{\text{OL}}, \quad m(D_t^{\bar{\pi}^*} \setminus D_{t+1}^{\bar{\pi}^*}) \leq m(D_{t+1}^{\text{OL}} \setminus D_t^{\text{OL}}). \quad (48)$$

Moreover, by definition $m(D_t^{\bar{\pi}^*}) \geq \beta(1 - m(D_t^{\text{OL}})) - \frac{2}{N}$ and $m(D_{t+1}^{\bar{\pi}^*}) \leq \beta(1 - m(D_{t+1}^{\text{OL}}))$, so

$$\begin{aligned} m(D_{t+1}^{\bar{\pi}^*} \setminus D_t^{\bar{\pi}^*}) &= m(D_t^{\bar{\pi}^*} \setminus D_{t+1}^{\bar{\pi}^*}) + m(D_{t+1}^{\bar{\pi}^*}) - m(D_t^{\bar{\pi}^*}) \\ &\leq m(D_{t+1}^{\text{OL}} \setminus D_t^{\text{OL}}) - \beta(m(D_{t+1}^{\text{OL}}) - m(D_t^{\text{OL}})) + \frac{2}{N}. \end{aligned} \quad (49)$$

Plugging (48) and (49) into (46) and rearranging the terms, we get

$$V(X_{t+1}, D_{t+1}^{\text{OL}}, D_{t+1}^{\bar{\pi}^*}) - V(X_{t+1}, D_t^{\text{OL}}, D_t^{\bar{\pi}^*}) \leq 4(\lambda_U^{1/2} + \lambda_W^{1/2})m(D_t^{\text{OL}} \setminus D_{t+1}^{\text{OL}}) + \frac{4\lambda_W^{1/2}}{N}, \quad (50)$$

where the terms related to $m(D_{t+1}^{\text{OL}} \setminus D_t^{\text{OL}})$ have been canceled out.

For the state-transition term in (45), it follows directly from the definition of V that

$$\begin{aligned} & V(X_{t+1}, D_t^{\text{OL}}, D_t^{\bar{\pi}^*}) - V(X_t, D_t^{\text{OL}}, D_t^{\bar{\pi}^*}) \\ &\leq (h_U(X_{t+1}, D_t^{\text{OL}}) - h_U(X_t, D_t^{\text{OL}})) + (h_W(X_{t+1}, D_t^{\bar{\pi}^*}) - h_W(X_t, D_t^{\bar{\pi}^*})) \end{aligned} \quad (51)$$

Combine the above calculations, we get the following key inequality:

$$\begin{aligned} & V(\Sigma_{t+1}) - V(\Sigma_t) \\ &\leq (h_U(X_{t+1}, D_t^{\text{OL}}) - h_U(X_t, D_t^{\text{OL}})) + (h_W(X_{t+1}, D_t^{\bar{\pi}^*}) - h_W(X_t, D_t^{\bar{\pi}^*})) \end{aligned} \quad (52)$$

$$+ 4(\lambda_U^{1/2} + \lambda_W^{1/2})m(D_t^{\text{OL}} \setminus D_{t+1}^{\text{OL}}) + \frac{4\lambda_W^{1/2}}{N}. \quad (53)$$

Next, we verify the drift condition (27) of Lemma 9, restated below: when $V(\Sigma_t) \geq \eta$,

$$\mathbb{E}[V(\Sigma_{t+1}) - V(\Sigma_t) \mid \Sigma_t] \leq -\gamma + O(1/\sqrt{N}) \quad a.s., \quad (54)$$

for a constant $\gamma > 0$. We take expectation in the bound (52)-(53). Because $\mathbb{E}[h_U(X_{t+1}, D_t^{\text{OL}}) \mid \Sigma_t] \leq \rho_u h_U(X_t, D_t^{\text{OL}}) + O(1/\sqrt{N})$, $\mathbb{E}[h_W(X_{t+1}, D_t^{\bar{\pi}^*}) \mid \Sigma_t] \leq \rho_w h_W(X_t, D_t^{\bar{\pi}^*}) + O(1/\sqrt{N})$ (Lemma 10), $\mathbb{E}[m(D_t^{\text{OL}} \setminus D_{t+1}^{\text{OL}}) \mid \Sigma_t] = O(1/\sqrt{N})$ (Lemma 11), we get

$$\mathbb{E}[V(\Sigma_{t+1}) - V(\Sigma_t) \mid \Sigma_t] \leq -((1 - \rho_u)h_U(X_t, D_t^{\text{OL}}) + (1 - \rho_w)h_W(X_t, D_t^{\bar{\pi}^*})) + O(1/\sqrt{N}). \quad (55)$$

To show (54), it remains to lower bound the term $(1 - \rho_u)h_U(X_t, D_t^{\text{OL}}) + (1 - \rho_w)h_W(X_t, D_t^{\bar{\pi}^*})$ when $V(\Sigma_t) > \eta$. We invoke Lemma 12 to get

$$\begin{aligned} V(\Sigma_t) &\leq h_U(X_t, D_t^{\text{OL}}) + h_W(X_t, D_t^{\bar{\pi}^*}) + L\left(\frac{\lambda_U^{1/2}}{\eta\beta}h_W(X_t, D_t^{\bar{\pi}^*}) + \frac{2}{\beta N}\right) \\ &= K_{Vh} \cdot ((1 - \rho_u)h_U(X_t, D_t^{\text{OL}}) + (1 - \rho_w)h_W(X_t, D_t^{\bar{\pi}^*})) + O(1/N), \end{aligned} \quad (56)$$

for some positive constant K_{Vh} . So $\mathbb{E}[V(\Sigma_{t+1}) - V(\Sigma_t) \mid \Sigma_t] \leq -\eta/K_{Vh} + O(1/\sqrt{N})$ when $V(\Sigma_t) > \eta$, which implies (54).

Next, we verify the condition (28) of Lemma 9, restated below: when $V(\Sigma_t) > \eta$,

$$\mathbb{E}[(V(\Sigma_{t+1}) - V(\Sigma_t))^+ \mid \Sigma_t] \leq O(1/\sqrt{N}) \quad a.s. \quad (57)$$

Again, using the bound in (52)-(53), we have

$$\begin{aligned} &\mathbb{E}[(V(\Sigma_{t+1}) - V(\Sigma_t))^+ \mid \Sigma_t] \\ &\leq \mathbb{E}[(h_U(X_{t+1}, D_t^{\text{OL}}) - h_U(X_t, D_t^{\text{OL}}))^+ \mid \Sigma_t] + \mathbb{E}[(h_W(X_{t+1}, D_t^{\bar{\pi}^*}) - h_W(X_t, D_t^{\bar{\pi}^*}))^+ \mid \Sigma_t] \\ &\quad + 4(\lambda_U^{1/2} + \lambda_W^{1/2})\mathbb{E}[m(D_t^{\text{OL}} \setminus D_{t+1}^{\text{OL}}) \mid \Sigma_t] + \frac{4\lambda_W^{1/2}}{N}. \end{aligned}$$

Each term on the RHS above is $O(1/\sqrt{N})$: In particular, by Lemma 11. $\mathbb{E}[m(D_t^{\text{OL}} \setminus D_{t+1}^{\text{OL}}) \mid \Sigma_t] = O(m(D_t^{\text{OL}})O(\exp(-CNm(D_t^{\text{OL}})))) = O(1/\sqrt{N})$.

Finally, we show that when $V(\Sigma_t) \leq \eta$,

$$\mathbb{P}[V(\Sigma_{t+1}) > \rho_1 V(\Sigma_t) + r \mid \Sigma_t] = O(\exp(-CN)) \quad a.s., \quad (58)$$

where $r = (1 - \rho_1)\eta/2$, ρ_1 is a constant in $(0, 1)$ and C is a positive constant. It is not hard to see that (58) implies the conditions (29) and (30) of Lemma 9, We rearrange the bound in (52)-(53) and apply (56) to get

$$\begin{aligned} &V(\Sigma_{t+1}) - \left(1 - \frac{1}{K_{Vh}}\right)V(\Sigma_t) \\ &\leq (h_U(X_{t+1}, D_t^{\text{OL}}) - \rho_u h_U(X_t, D_t^{\text{OL}})) + (h_W(X_{t+1}, D_t^{\bar{\pi}^*}) - \rho_w h_W(X_t, D_t^{\bar{\pi}^*})) \\ &\quad + 4(\lambda_U^{1/2} + \lambda_W^{1/2})m(D_t^{\text{OL}} \setminus D_{t+1}^{\text{OL}}) + O(1/N). \end{aligned}$$

For N larger than a certain constant threshold, the $O(1/N)$ term in the last expression is less than $r/4$. Using a union bound and invoking Lemma 10, we get

$$\begin{aligned} &\mathbb{P}\left[V(\Sigma_{t+1}) > \left(1 - \frac{1}{K_{Vh}}\right)V(\Sigma_t) + r \mid \Sigma_t\right] \\ &\leq \mathbb{P}[h_U(X_{t+1}, D_t^{\text{OL}}) > \rho_u h_U(X_t, D_t^{\text{OL}}) + r/4 \mid \Sigma_t] \\ &\quad + \mathbb{P}[h_W(X_{t+1}, D_t^{\bar{\pi}^*}) > \rho_w h_W(X_t, D_t^{\bar{\pi}^*}) + r/4 \mid \Sigma_t] \\ &\quad + \mathbb{P}[4(\lambda_U^{1/2} + \lambda_W^{1/2})m(D_t^{\text{OL}} \setminus D_{t+1}^{\text{OL}}) > r/4 \mid \Sigma_t] + \mathbb{1}\{O(1/N) > r/4\} \\ &\leq O(\exp(-CN)) + O(\exp(-CNm(D_t^{\text{OL}}))). \end{aligned}$$

By Lemma 12, $V(\Sigma_t) < \eta$ implies $m(D_t^{\text{OL}}) = 1$, so $O(\exp(-CNm(D_t^{\text{OL}}))) = O(\exp(-CN))$. Therefore, (58) holds with $\rho_1 = 1 - 1/(K_{Vh})$. \square

B.3 Proofs of the core lemmas

Lemma 7 (Subroutine conformity). *Assume Assumptions 1, 2 and 3 hold. Consider the Two-Set Policy described in Algorithm 1 with $\eta = |\mathbb{S}|^{-1/2}(\min\{y^*(\tilde{s}, 0), y^*(\tilde{s}, 1)\} - (|S^\emptyset| + 1)/N)$: The steps in Line 5-7 are well-defined. In other words, we can always let the arms in D_t^{OL} follow Optimal Local Control, let the arms in $D_t^{\bar{\pi}^*}$ follow Exact Proportional Control, and choose actions for the arms not in $D_t^{\text{OL}} \cup D_t^{\bar{\pi}^*}$ to satisfy the budget constraint $\sum_{i \in [N]} A_t(i) = \alpha N$.*

Proof. In this proof, we need to verify the condition of following the Optimal Local Control, (7), for the arms in D_t^{OL} . Then we need to show that $A_t(i)$ for $i \notin D_t^{\text{OL}} \cup D_t^{\bar{\pi}^*}$ can be chosen such that $\sum_{i \in [N]} A_t(i) = \alpha N$, if the arms in D_t^{OL} and $D_t^{\bar{\pi}^*}$ follow Optimal Local Control and Exact Proportional Control, respectively.

The condition (7) specialized to D_t^{OL} are stated as follows.

$$\sum_{s \neq \tilde{s}} \bar{\pi}^*(1|s) X_t(D_t^{\text{OL}}, s) \leq \alpha m(D_t^{\text{OL}}) - (|S^\emptyset| + 1)/N, \quad (59)$$

$$\sum_{s \neq \tilde{s}} \bar{\pi}^*(0|s) X_t(D_t^{\text{OL}}, s) \leq (1 - \alpha) m(D_t^{\text{OL}}) - (|S^\emptyset| + 1)/N. \quad (60)$$

We first show (59). Observe that $\sum_{s \in \mathbb{S}} \bar{\pi}^*(1|s) \mu^*(s) = \sum_{s \in \mathbb{S}, a \in \mathbb{A}} y^*(s, a) = \alpha$, so

$$\begin{aligned} \alpha m(D_t^{\text{OL}}) - \sum_{s \neq \tilde{s}} \bar{\pi}^*(1|s) X_t(D_t^{\text{OL}}, s) &= y^*(\tilde{s}, 1) m(D_t^{\text{OL}}) - \sum_{s \neq \tilde{s}} \bar{\pi}^*(1|s) (m(D_t^{\text{OL}}) \mu^*(s) - X_t(D_t^{\text{OL}}, s)) \\ &\geq y^*(\tilde{s}, 1) m(D_t^{\text{OL}}) - \sum_{s \neq \tilde{s}} |m(D_t^{\text{OL}}) \mu^*(s) - X_t(D_t^{\text{OL}}, s)| \\ &\geq y^*(\tilde{s}, 1) m(D_t^{\text{OL}}) - \|m(D_t^{\text{OL}}) \mu^* - X_t(D_t^{\text{OL}}, s)\|_1 \\ &\geq y^*(\tilde{s}, 1) m(D_t^{\text{OL}}) - |\mathbb{S}|^{1/2} \|m(D_t^{\text{OL}}) \mu^* - X_t(D_t^{\text{OL}}, s)\|_U, \end{aligned}$$

where to get the last expression, we use the argument that U 's eigenvalues are at least 1, so $\|v\|_1 \leq |\mathbb{S}|^{1/2} \|v\|_2 \leq |\mathbb{S}|^{1/2} \|v\|_U$ for any $|\mathbb{S}|$ -dim vector v . To bound $\|m(D_t^{\text{OL}}) \mu^* - X_t(D_t^{\text{OL}}, s)\|_U$, recall that D_t^{OL} is chosen such that $\delta(X_t, D_t^{\text{OL}}) \geq 0$, so by definition, we have

$$\|X_t(D_t^{\text{OL}}) - m(D_t^{\text{OL}}) \mu^*\|_U \leq \eta m(D_t^{\text{OL}}) \leq |\mathbb{S}|^{-1/2} (y^*(\tilde{s}, 1) - (|S^\emptyset| + 1)/N) m(D_t^{\text{OL}}).$$

Therefore, combining the above calculations,

$$\alpha m(D_t^{\text{OL}}) - \sum_{s \neq \tilde{s}} \bar{\pi}^*(1|s) X_t(D_t^{\text{OL}}, s) \geq (|S^\emptyset| + 1)/N,$$

which implies (59). (60) can be similarly proved.

It remains to show that $\sum_{i \in [N]} A_t(i) = \alpha N$ can be satisfied. If $D^{\text{OL}} = [N]$, by definition we have $\sum_{i \in D_t^{\text{OL}}} A_t(i) = \alpha N$. Thus we only need to consider the case when $D^{\text{OL}} \subsetneq [N]$. By definition, $\sum_{i \in D_t^{\text{OL}}} A_t(i) \in \{\lfloor \alpha |D_t^{\text{OL}}| \rfloor, \lceil \alpha |D_t^{\text{OL}}| \rceil\}$. Because $A_t(i) \in \{0, 1\}$ for $i \in D_t^{\bar{\pi}^*}$, we have

$$\lfloor \alpha |D_t^{\text{OL}}| \rfloor \leq \sum_{i \in D_t^{\text{OL}} \cup D_t^{\bar{\pi}^*}} A_t(i) \leq \lceil \alpha |D_t^{\text{OL}}| \rceil + |D_t^{\bar{\pi}^*}|.$$

Since $\sum_{i \notin D_t^{\text{OL}} \cup D_t^{\bar{\pi}^*}} A_t(i)$ can be anything integer between 0 and $N - |D_t^{\text{OL}}| - |D_t^{\bar{\pi}^*}|$, it suffices if

$$\lfloor \alpha |D_t^{\text{OL}}| \rfloor + (N - |D_t^{\text{OL}}| - |D_t^{\bar{\pi}^*}|) \geq \alpha N \quad a.s.$$

$$[\alpha|D_t^{\text{OL}}|] + |D_t^{\bar{\pi}^*}| \leq \alpha N \quad a.s.$$

These two inequalities can be verified using the facts that $|D_t^{\bar{\pi}^*}| = (\lfloor \beta(N - |D_t^{\text{OL}}|) - 1 \rfloor)^+$, $\alpha N \in \{\lfloor \alpha N \rfloor, \lceil \alpha N \rceil\}$ and $D_t^{\text{OL}} \subsetneq [N]$. \square

Lemma 11 (Almost non-shrinking). *Assume Assumptions 1, 2 and 3 hold. Under the Two-Set Policy (Algorithm 1), for any $t \geq 0$,*

$$\mathbb{E}[m(D_t^{\text{OL}} \setminus D_{t+1}^{\text{OL}}) \mid \Sigma_t] \leq O(m(D_t^{\text{OL}}) \exp(-CNm(D_t^{\text{OL}})^2)) \quad a.s. \quad (38)$$

$$\mathbb{P}[m(D_t^{\text{OL}} \setminus D_{t+1}^{\text{OL}}) > 0 \mid \Sigma_t] \leq O(\exp(-CNm(D_t^{\text{OL}})^2)) \quad a.s., \quad (39)$$

where C is a positive constant.

Proof. By the definition of D_{t+1}^{OL} , it suffices to construct a set $\bar{D} \subseteq D_t^{\text{OL}}$ based on the realization of X_{t+1} such that

$$\delta(X_{t+1}, \bar{D}) \geq 0 \quad a.s. \quad (61)$$

$$\mathbb{E}[(m(D_t^{\text{OL}} \setminus \bar{D}) \mid \Sigma_t] \leq O(\exp(-CNm(D_t^{\text{OL}})^2)) \quad a.s. \quad (62)$$

$$\mathbb{P}[m(D_t^{\text{OL}} \setminus \bar{D}) > 0 \mid \Sigma_t] \leq O(\exp(-CNm(D_t^{\text{OL}})^2)) \quad a.s. \quad (63)$$

Consider taking \bar{D} as:

$$\bar{D} = \begin{cases} D_t^{\text{OL}} & \text{if } \delta(X_{t+1}, D_t^{\text{OL}}) \geq 0 \\ \emptyset & \text{if } \delta(X_{t+1}, D_t^{\text{OL}}) < 0. \end{cases} \quad (64)$$

With \bar{D} defined in (64), (61) is obviously true. To show (62) and (63), observe that

$$\begin{aligned} \mathbb{E}[(m(D_t^{\text{OL}} \setminus \bar{D}))^+ \mid X_t, D_t^{\text{OL}}, D_t^{\bar{\pi}^*}] &= m(D_t^{\text{OL}}) \mathbb{P}[\delta(X_{t+1}, D_t^{\text{OL}}) < 0] \\ \mathbb{P}[m(D_t^{\text{OL}} \setminus \bar{D}) > 0 \mid \Sigma_t] &= \mathbb{P}[\delta(X_{t+1}, D_t^{\text{OL}}) < 0]. \end{aligned}$$

Therefore, it remains to bound $\mathbb{P}[\delta(X_{t+1}, D_t^{\text{OL}}) < 0]$. By definition, $\delta(x, D) = \eta m(D) - \|x(D) - m(D)\mu^*\|_U = \eta m(D) - h_U(x, D)$, so

$$\mathbb{P}[\delta(X_{t+1}, D_t^{\text{OL}}) < 0] = \mathbb{P}[h_U(X_{t+1}, D_t^{\text{OL}}) > \eta m(D_t^{\text{OL}})],$$

Because $\delta(X_t, D_t^{\text{OL}}) \geq 0$, we have $h_U(X_t, D_t^{\text{OL}}) \leq \eta m(D_t^{\text{OL}})$. Therefore,

$$\begin{aligned} h_U(X_{t+1}, D_t^{\text{OL}}) &\leq h_U(X_{t+1}, D_t^{\text{OL}}) - \rho_u h_U(X_t, D_t^{\text{OL}}) + \rho_u \eta m(D_t^{\text{OL}}) \\ \mathbb{P}[h_U(X_{t+1}, D_t^{\text{OL}}) \geq \eta m(D_t^{\text{OL}})] &\leq \mathbb{P}[h_U(X_{t+1}, D_t^{\text{OL}}) - h_U(X_t, D_t^{\text{OL}}) \geq (1 - \rho_u) \eta m(D_t^{\text{OL}})] \\ &= O(\exp(-C'N(1 - \rho_u)^2 \eta^2 m(D_t^{\text{OL}})^2)), \end{aligned}$$

where the last inequality comes from Lemma 10 and C' is a positive constant. Because $(1 - \rho_u)^2 \eta^2 > 0$ and is a constant, $\mathbb{P}[\delta(X_{t+1}, D_t^{\text{OL}}) < 0] = O(\exp(-CNm(D_t^{\text{OL}})^2))$ with $C = C'(1 - \rho_u)^2 \eta^2$. \square

Lemma 12 (Sufficient coverage). *Assume Assumptions 1, 2 and 3 hold. Consider the Two-Set Policy (Algorithm 1) with $\eta = |\mathcal{S}|^{-1/2}(\min\{y^*(\tilde{s}, 0), y^*(\tilde{s}, 1)\} - (|\mathcal{S}^\emptyset| + 1)/N)$. For any $t \geq 0$,*

$$1 - m(D_t^{\text{OL}}) \leq \frac{\lambda_U^{1/2}}{\eta \beta} h_W(X_t, D_t^{\bar{\pi}^*}) + \frac{2}{\beta N} \quad a.s. \quad (40)$$

$$m(D_t^{\text{OL}}) = 1 \quad \text{if } h_U(X_t, [N]) \leq \eta. \quad (41)$$

Proof. First, we show that

$$h_W(X_t, D_t^{\bar{\pi}^*}) \geq \lambda_U^{-1/2} \eta m(D_t^{\bar{\pi}^*}). \quad (65)$$

If we can show (65), combined with the fact that $m(D_t^{\bar{\pi}^*}) = (\lfloor \beta(N - |D_t^{\text{OL}}|) \rfloor - 1)^+ / N$, we get

$$h_W(X_t, D_t^{\bar{\pi}^*}) \geq \lambda_U^{-1/2} \eta (\beta(N - |D_t^{\text{OL}}|) - 2) / N = \lambda_U^{-1/2} \eta \beta (1 - m(D_t^{\text{OL}})) - 2\lambda_U^{-1/2} \eta / N,$$

which implies (40).

To show (65), we only need to consider the case where $m(D_t^{\bar{\pi}^*}) \neq 0$. Suppose, to get a contradiction, that $h_W(X_t, D_t^{\bar{\pi}^*}) < \lambda_U^{-1/2} \eta m(D_t^{\bar{\pi}^*})$. Because $\|v\|_U \leq \lambda_U^{1/2} \|v\|_2 \leq \lambda_U^{1/2} \|v\|_W$ for any $v \in \mathbb{R}^{|\mathbb{S}|}$,

$$\|X_t(D_t^{\bar{\pi}^*}) - m(D_t^{\bar{\pi}^*})\mu^*\|_U \leq \lambda_U^{1/2} \|X_t(D_t^{\bar{\pi}^*}) - m(D_t^{\bar{\pi}^*})\mu^*\|_W < \eta m(D_t^{\bar{\pi}^*}).$$

Combined with the triangular inequality and the definition of D_t^{OL} , we have

$$\begin{aligned} \|X_t(D_t^{\text{OL}} \cup D_t^{\bar{\pi}^*}) - m(D_t^{\text{OL}} \cup D_t^{\bar{\pi}^*})\mu^*\|_U &\leq \|X_t(D_t^{\text{OL}}) - m(D_t^{\text{OL}})\mu^*\|_U + \|X_t(D_t^{\bar{\pi}^*}) - m(D_t^{\bar{\pi}^*})\mu^*\|_U \\ &< \eta m(D_t^{\text{OL}}) + \eta m(D_t^{\bar{\pi}^*}). \end{aligned}$$

Consequently, $D_t^{\text{OL}} \cup D_t^{\bar{\pi}^*}$ is a strict superset of D_t^{OL} (thus also a superset of $D_{\text{temp}}^{\text{OL}}$) such that $\delta(X_t, D_t^{\text{OL}} \cup D_t^{\bar{\pi}^*}) \geq 0$. So the arms in $D_t^{\bar{\pi}^*}$ should have instead been taken into D_t^{OL} when the algorithm chooses D_t^{OL} to be the maximal set such that $D_t^{\text{OL}} \supseteq D_{\text{temp}}^{\text{OL}}$ and $\delta(X_t, D_t^{\text{OL}}) \geq 0$, which causes a contradiction. Therefore, (65) holds.

Finally, we show (41). When $h_U(X_t, [N]) \leq \eta$, we have $\delta(X_t, [N]) \geq 0$. Again, because D_t^{OL} is chosen as the largest-cardinality set containing $D_{\text{temp}}^{\text{OL}}$, we must have $D_t^{\text{OL}} = [N]$. \square

B.4 Proofs of the technical lemmas

Lemma 8. *Assume Assumptions 1, 2 and 3 hold. The optimality gap under the Two-Set Policy (Algorithm 1) is bounded as*

$$R^{\text{rel}} - R(\pi, \mathbf{S}_0) \leq O(1) \cdot \mathbb{P}[D_\infty^{\text{OL}} \neq [N]]. \quad (25)$$

Proof. We will first bound the norm of the long-run expectation of the $X_t([N]) - \mu^*$ using $\mathbb{P}[D_\infty^{\text{OL}} \neq [N]]$, and then derive the optimality gap bound.

Let $Q = (I - \Phi)^{-1}$, which is well-defined because all eigenvalues of Φ have moduli strictly less than 1 (Assumption 3). By direct calculation, we have that for any $v \in \Delta(\mathbb{S})$,

$$(v - \mu^*)Q - (v - \mu^*)\Phi Q = v - \mu^*. \quad (66)$$

We define the positive constant λ_Q as

$$\lambda_Q = \max \left\{ \max_{v \in \Delta(\mathbb{S})} \|(v - \mu^*)Q\|_1, \max_{v \in \Delta(\mathbb{S})} \|(v - \mu^*)\Phi Q\|_1 \right\},$$

which must be finite because $\Delta(\mathbb{S})$ is a compact set.

Letting $v = X_t([N])$ and taking the expectation (for random vectors) in (66), we get

$$\begin{aligned} \mathbb{E}[X_t([N]) - \mu^*] &= \mathbb{E}[(X_t([N]) - \mu^*)Q - (X_t([N]) - \mu^*)\Phi Q] \\ &= \mathbb{E}[(X_{t+1}([N]) - \mu^*)Q - (X_t([N]) - \mu^*)\Phi Q] \end{aligned}$$

$$- \mathbb{E}[(X_{t+1}([N]) - \mu^*)Q - (X_t([N]) - \mu^*)Q].$$

Using the fact that

$$\lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E}[(X_{t+1}([N]) - \mu^*)Q - (X_t([N]) - \mu^*)Q] = 0,$$

we get

$$\lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E}[X_t([N]) - \mu^*] = \lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E}[(X_{t+1}([N]) - \mu^*)Q - (X_t([N]) - \mu^*)\Phi Q]. \quad (67)$$

Therefore, it remains to bound $\mathbb{E}[(X_{t+1}([N]) - \mu^*)Q - (X_t([N]) - \mu^*)\Phi Q]$ for every time step t .

Let $\Sigma_t = (X_t, D_t^{\text{OL}}, D_t^{\bar{\pi}^*})$. We will bound $\mathbb{E}[(X_{t+1}([N]) - \mu^*)Q - (X_t([N]) - \mu^*)\Phi Q \mid \Sigma_t]$ by decomposing it into two terms:

$$\begin{aligned} & \mathbb{E}[(X_{t+1}([N]) - \mu^*)Q - (X_t([N]) - \mu^*)\Phi Q \mid \Sigma_t] \\ &= \mathbb{E}[(X_{t+1}([N]) - \mu^*)Q - (X_t([N]) - \mu^*)\Phi Q \mid \Sigma_t] \mathbb{1}\{D_t^{\text{OL}} = [N]\} \end{aligned} \quad (68)$$

$$+ \mathbb{E}[(X_{t+1}([N]) - \mu^*)Q - (X_t([N]) - \mu^*)\Phi Q \mid \Sigma_t] \mathbb{1}\{D_t^{\text{OL}} \neq [N]\}. \quad (69)$$

For the term in (68), note that by Lemma 1, $\mathbb{E}[X_{t+1}([N]) - \mu^* \mid \Sigma_t] = (X_t([N]) - \mu^*)\Phi$ when $D_t^{\text{OL}} = [N]$, so

$$\mathbb{E}[(X_{t+1}([N]) - \mu^*)Q \mid \Sigma_t] \mathbb{1}\{D_t^{\text{OL}} = [N]\} = (X_t([N]) - \mu^*)\Phi Q \mathbb{1}\{D_t^{\text{OL}} = [N]\},$$

which implies that the term in (68) is zero. For the term in (69), its L_1 norm can be bounded as

$$\begin{aligned} & \left\| \mathbb{E}[(X_{t+1}([N]) - \mu^*)Q - (X_t([N]) - \mu^*)\Phi Q \mid \Sigma_t] \right\|_1 \mathbb{1}\{D_t^{\text{OL}} \neq [N]\} \\ & \leq \mathbb{E} \left[\left\| (X_{t+1}([N]) - \mu^*)Q - (X_t([N]) - \mu^*)\Phi Q \right\|_1 \mid \Sigma_t \right] \mathbb{1}\{D_t^{\text{OL}} \neq [N]\} \\ & \leq \mathbb{E} \left[\left\| (X_{t+1}([N]) - \mu^*)Q \right\|_1 + \left\| (X_t([N]) - \mu^*)\Phi Q \right\|_1 \mid \Sigma_t \right] \mathbb{1}\{D_t^{\text{OL}} \neq [N]\} \\ & \leq 2\lambda_Q \mathbb{1}\{D_t^{\text{OL}} \neq [N]\}, \end{aligned} \quad (70)$$

where (70) follows from the definition of λ_Q and the fact that $X_{t+1}([N]), X_t([N]) \in \Delta(\mathbb{S})$. Combining the calculations of the two terms in (68) and (69), we have

$$\begin{aligned} & \left\| \mathbb{E}[(X_{t+1}([N]) - \mu^*)Q - (X_t([N]) - \mu^*)\Phi Q] \right\|_1 \\ &= \mathbb{E} \left[\left\| \mathbb{E}[(X_{t+1}([N]) - \mu^*)Q - (X_t([N]) - \mu^*)\Phi Q \mid \Sigma_t] \right\|_1 \mathbb{1}\{D_t^{\text{OL}} \neq [N]\} \right] \\ & \leq 2\lambda_Q \mathbb{P} [D_t^{\text{OL}} \neq [N]]. \end{aligned} \quad (71)$$

Plugging (71) into (67), we get

$$\begin{aligned} \left\| \mathbb{E}[X_\infty([N]) - \mu^*] \right\|_1 & \leq \lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=0}^{T-1} \left\| \mathbb{E}[(X_{t+1}([N]) - \mu^*)Q - (X_t([N]) - \mu^*)\Phi Q] \right\|_1 \\ & \leq 2\lambda_Q \mathbb{P} [D_\infty^{\text{OL}} \neq [N]]. \end{aligned} \quad (72)$$

where the expectations or probabilities of variables or events with subscript ∞ are shorthands for the long-run time averages.

Now we start to bound the optimality gap. We first rewrite the long-run average reward $R(\pi, \mathbf{S}_0)$ in terms of an expectation with respect to the states:

$$R(\pi, \mathbf{S}_0) = \mathbb{E}[r(\mathbf{S}_\infty, \mathbf{A}_\infty)] = \mathbb{E}\left[r^\pi(X_\infty, D_\infty^{\text{OL}}, D_\infty^{\bar{\pi}^*})\right],$$

where $r^\pi(x, D^{\text{OL}}, D^{\bar{\pi}^*})$ is the expected reward at a given state under the Two-Set Policy:

$$r^\pi(x, D^{\text{OL}}, D^{\bar{\pi}^*}) \triangleq \mathbb{E}[r(\mathbf{S}_t, \mathbf{A}_t) \mid X_t = x, D_t^{\text{OL}} = D^{\text{OL}}, D_t^{\bar{\pi}^*} = D^{\bar{\pi}^*}].$$

Note that the RHS of $r^\pi(x, D^{\text{OL}}, D^{\bar{\pi}^*})$'s definition does not depend on t because the Two-Set Policy is stationary and Markovian policy with state $(X_t, D_t^{\text{OL}}, D_t^{\bar{\pi}^*})$.

Next, we examine the value of $r^\pi(x, D^{\bar{\pi}^*}, D^{\text{OL}})$ when $D^{\text{OL}} = [N]$ and $D^{\bar{\pi}^*} = \emptyset$. Let $Y_t(s, a)$ be the random variable representing the fraction of arms in state s taking action a at time t . Then

$$r^\pi(X_t, [N], \emptyset) = \sum_{s \in \mathbb{S}, a \in \mathbb{A}} r(s, a) \mathbb{E}[Y_t(s, a) \mid X_t, D_t^{\text{OL}} = [N], D_t^{\bar{\pi}^*} = \emptyset].$$

By the definition of Optimal Local Control,

$$\mathbb{E}[Y_t(s, a) \mid X_t, D_t^{\text{OL}} = [N], D_t^{\bar{\pi}^*} = \emptyset] = \bar{\pi}^*(a|s) X_t([N], s) \quad s \neq \tilde{s} \quad (73)$$

$$\mathbb{E}[Y_t(\tilde{s}, 1) \mid X_t, D_t^{\text{OL}} = [N], D_t^{\bar{\pi}^*} = \emptyset,] = \alpha - \sum_{s \neq \tilde{s}} \bar{\pi}^*(1|s) X_t([N], s) \quad (74)$$

$$\mathbb{E}[Y_t(\tilde{s}, 0) \mid X_t, D_t^{\text{OL}} = [N], D_t^{\bar{\pi}^*} = \emptyset,] = 1 - \alpha - \sum_{s \neq \tilde{s}} \bar{\pi}^*(0|s) X_t([N], s). \quad (75)$$

Therefore,

$$\begin{aligned} r^\pi(X_t, [N], \emptyset) &= \sum_{s \neq \tilde{s}, a \in \mathbb{A}} (r(s, a) - r(\tilde{s}, a)) \bar{\pi}^*(a|s) X_t([N], s) \\ &\quad + \alpha r(\tilde{s}, 1) + (1 - \alpha) r(\tilde{s}, 0), \end{aligned} \quad (76)$$

Let $\hat{r}(v)$ be the affine function on $\Delta(\mathbb{S})$ such that $\hat{r}(X_t([N])) = r^\pi(X_t, \emptyset, [N])$:

$$\hat{r}(v) \triangleq \sum_{s \neq \tilde{s}, a \in \mathbb{A}} (r(s, a) - r(\tilde{s}, a)) \bar{\pi}^*(a|s) v(s) + \alpha r(\tilde{s}, 1) + (1 - \alpha) r(\tilde{s}, 0).$$

It is easy to see that $\hat{r}(v)$ is $4r_{\max}$ -Lipschitz continuous with respect to L_1 -norm, and $\hat{r}(\mu^*) = R^{\text{rel}}$.

With the above definitions, we can lower bound $R(\pi, \mathbf{S}_0)$ as

$$\begin{aligned} R(\pi, \mathbf{S}_0) &= \mathbb{E}[r^\pi(X_\infty, D_\infty^{\text{OL}}, D_\infty^{\bar{\pi}^*})] \\ &\geq \mathbb{E}[\hat{r}(X_\infty([N])) \mathbb{1}\{D_\infty^{\text{OL}} = [N]\}] - r_{\max} \mathbb{P}[D_\infty^{\text{OL}} \neq [N]] \\ &= \mathbb{E}[\hat{r}(X_\infty([N]))] - 2r_{\max} \mathbb{P}[D_\infty^{\text{OL}} \neq [N]] \\ &= \hat{r}(\mathbb{E}[X_\infty([N])]) - 2r_{\max} \mathbb{P}[D_\infty^{\text{OL}} \neq [N]] \\ &= \hat{r}(\mu^*) + \hat{r}(\mathbb{E}[X_\infty([N])]) - \hat{r}(\mu^*) - 2r_{\max} \mathbb{P}[D_\infty^{\text{OL}} \neq [N]] \\ &\geq R^{\text{rel}} - 4r_{\max} \|\mathbb{E}[X_\infty([N]) - \mu^*]\|_1 - 2r_{\max} \mathbb{P}[D_\infty^{\text{OL}} \neq [N]] \\ &\geq R^{\text{rel}} - 2r_{\max}(4\lambda_Q + 1) \mathbb{P}[D_t^{\text{OL}} \neq [N]]. \end{aligned}$$

Because $2r_{\max}(4\lambda_Q + 1) = O(1)$, we have $R^{\text{rel}} - R(\pi, \mathbf{S}_0) \leq O(1) \cdot \mathbb{P}[D_t^{\text{OL}} \neq [N]]$. \square

Lemma 9. *Suppose there exists a function $V(x, D^{\text{OL}}, D^{\bar{\pi}^*})$ and constants V_{\max} and $\bar{\eta} > 0$ such that for any $t \geq 0$,*

$$0 \leq V(\Sigma_t) \leq V_{\max} \quad a.s.; \quad (26)$$

if $V(\Sigma_t) > \bar{\eta}$,

$$\mathbb{E}[V(\Sigma_{t+1}) - V(\Sigma_t) \mid \Sigma_t] \leq -\gamma \quad (27)$$

$$\mathbb{E}[(V(\Sigma_{t+1}) - V(\Sigma_t))^+ \mid \Sigma_t] = O(1/\sqrt{N}); \quad (28)$$

if $\bar{\eta}/2 < V(\Sigma_t) \leq \bar{\eta}$,

$$\mathbb{E}[(V(\Sigma_{t+1}) - V(\Sigma_t))^+ \mid \Sigma_t] \leq O(\exp(-CN)); \quad (29)$$

if $V(\Sigma_t) \leq \bar{\eta}/2$,

$$\mathbb{P}[V(\Sigma_{t+1}) > \bar{\eta}/2 \mid \Sigma_t] \leq O(\exp(-C'N)), \quad (30)$$

where γ, C, C' are positive constants. Then we have

$$\mathbb{P}[V(\Sigma_\infty) > \bar{\eta}] = O(\exp(-C''N)), \quad (31)$$

where C'' is another positive constant.

Proof. Let $f(a) = (a - \bar{\eta}/2)^+$. To obtain an bound on $\mathbb{P}[V(\Sigma_\infty) > \bar{\eta}]$, we will calculate $\mathbb{E}[f(V(\Sigma_{t+1}))] - \mathbb{E}[f(V(\Sigma_t))]$ for each time step t and invoking the fact that

$$\lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=0}^{T-1} (\mathbb{E}[f(V(\Sigma_{t+1}))] - \mathbb{E}[f(V(\Sigma_t))]) = 0. \quad (77)$$

We will bound $\mathbb{E}[f(V(\Sigma_{t+1}))] - \mathbb{E}[f(V(\Sigma_t))]$ in three cases, $V(\Sigma_t) > \bar{\eta}$, $\bar{\eta}/2 < V(\Sigma_t) \leq \bar{\eta}$, and $V(\Sigma_t) \leq \bar{\eta}/2$.

First, if $V(\Sigma_t) > \bar{\eta}$,

$$\begin{aligned} & \mathbb{E}[f(V(\Sigma_{t+1})) \mid \Sigma_t] - f(V(\Sigma_t)) \\ &= \mathbb{E}[(V(\Sigma_{t+1}) - V(\Sigma_t)) \mathbb{1}\{V(\Sigma_{t+1}) > V(\Sigma_t)\} \mid \Sigma_t] \\ & \quad + \mathbb{E}[(V(\Sigma_{t+1}) - V(\Sigma_t)) \mathbb{1}\{\bar{\eta}/2 < V(\Sigma_{t+1}) \leq V(\Sigma_t)\} \mid \Sigma_t] \\ & \quad + \mathbb{E}[(\bar{\eta}/2 - V(\Sigma_t)) \mathbb{1}\{V(\Sigma_{t+1}) \leq \bar{\eta}/2\} \mid \Sigma_t] \\ & \leq \mathbb{E}[(V(\Sigma_{t+1}) - V(\Sigma_t)) \mathbb{1}\{V(\Sigma_{t+1}) > V(\Sigma_t)\} \mid \Sigma_t] \\ & \quad + \frac{1}{2} \mathbb{E}[(V(\Sigma_{t+1}) - V(\Sigma_t)) \mathbb{1}\{V(\Sigma_{t+1}) \leq V(\Sigma_t)\} \mid \Sigma_t] \end{aligned} \quad (78)$$

$$= \frac{1}{2} \mathbb{E}[(V(\Sigma_{t+1}) - V(\Sigma_t))^+ \mid \Sigma_t] + \frac{1}{2} \mathbb{E}[V(\Sigma_{t+1}) - V(\Sigma_t) \mid \Sigma_t] \quad (79)$$

$$\leq -\frac{1}{2} \gamma + O(1/\sqrt{N}). \quad (80)$$

where the inequality in (78) follows uses the fact that when $V(\Sigma_t) \geq \bar{\eta}$, $V(\Sigma_{t+1}) \geq 0$,

$$\frac{1}{2}(V(\Sigma_{t+1}) - V(\Sigma_t)) - (\bar{\eta}/2 - V(\Sigma_t)) = \frac{1}{2}(V(\Sigma_{t+1}) + V(\Sigma_t)) - \bar{\eta}/2 \geq 0;$$

the inequality in (80) applies the properties of V assumed in (27) and (28).

Next, consider the case of $\bar{\eta}/2 < V(\Sigma_t) \leq \bar{\eta}$.

$$\mathbb{E}[f(V(\Sigma_{t+1})) \mid \Sigma_t] - f(V(\Sigma_t))$$

$$\begin{aligned}
&\leq \mathbb{E}[(f(V(\Sigma_{t+1})) - f(V(\Sigma_t))) \mathbb{1}\{V(\Sigma_{t+1}) > V(\Sigma_t)\} \mid \Sigma_t] \\
&= \mathbb{E}[(V(\Sigma_{t+1}) - V(\Sigma_t))^+ \mid \Sigma_t] \\
&\leq O(\exp(-CN)),
\end{aligned} \tag{81}$$

where (81) follows from (29).

Finally, consider the case of $V(\Sigma_t) \leq \bar{\eta}/2$.

$$\begin{aligned}
&\mathbb{E}[f(V(\Sigma_{t+1})) \mid \Sigma_t] - f(V(\Sigma_t)) \\
&= \mathbb{E}[(V(\Sigma_{t+1}) - \bar{\eta}/2)^+ \mid \Sigma_t] \\
&\leq V_{\max} \mathbb{P}[V(\Sigma_{t+1}) - \bar{\eta}/2 \mid \Sigma_t] \\
&= O(\exp(-C'N)),
\end{aligned} \tag{82}$$

where (82) follows from (30).

Combining the above calculations,

$$\mathbb{E}[f(V(\Sigma_{t+1}))] - \mathbb{E}[f(V(\Sigma_t))] \leq \left(-\frac{1}{2}\gamma + O(1/\sqrt{N})\right) \mathbb{P}[V(\Sigma_t) > \bar{\eta}] + O(\exp(-\min(C, C')N)).$$

Rearranging the terms, we get

$$\mathbb{P}[V(\Sigma_t) > \bar{\eta}] \leq \frac{1}{\gamma/2 - O(1/\sqrt{N})} \left(\mathbb{E}[f(V(\Sigma_{t+1}))] - \mathbb{E}[f(V(\Sigma_t))] + O(\exp(-\min(C, C')N)) \right), \tag{83}$$

for any time step t . Plugging the bound (83) into (77), we get

$$\mathbb{P}[V(\Sigma_\infty) > \bar{\eta}] = O(\exp(-\min(C, C')N)). \quad \square$$

Lemma 10 (Properties of h_W and h_U). *Assume Assumptions 1, 2 and 3 hold. For any time step t and $D \subseteq [N]$, let X'_{t+1} be the system state at time $t+1$ if the arms in D follow Exact Proportional Control at time t , then we have*

$$\mathbb{E}[(h_W(X'_{t+1}, D) - \rho_w h_W(X_t, D))^+ \mid X_t] \leq O(1/\sqrt{N}) \quad a.s. \tag{34}$$

$$\mathbb{P}[h_W(X'_{t+1}, D) > \rho_w h_W(X_t, D) + r \mid X_t] \leq O(\exp(-CNr^2)) \quad a.s. \quad \forall r \geq 0, \tag{35}$$

where $\rho_w = 1 - 1/(2\lambda_W)$, and C is a positive constant. In addition, for any $D, D' \subseteq [N]$,

$$h_W(x, D) \geq \frac{1}{|\mathbb{S}|^{1/2}} \|x(D) - m(D)\mu^*\|_1 \tag{36}$$

$$|h_W(x, D) - h_W(x, D')| \leq 2\lambda_W^{1/2} (m(D' \setminus D) + m(D \setminus D')), \tag{37}$$

The same inequalities hold for h_U with Exact Proportional Control and W replaced by the Optimal Local Control and U , which we omit here.

Proof. Proving (34) and (35) By the pseudo-contraction properties of W -weighted norms given in Lemma 6,

$$\begin{aligned}
\rho_w h_W(X_t, D) &= \rho_w \|X_t(D) - m(D)\mu^*\|_W \\
&\geq \|(X_t(D) - m(D)\mu^*)P_{\bar{\pi}^*}\|_W \\
&= \|X_t(D)P_{\bar{\pi}^*} - m(D)\mu^*\|_W.
\end{aligned}$$

Consequently,

$$\begin{aligned}
h_W(X'_{t+1}, D) - \rho_w h_W(X_t, D) &\leq \|X'_{t+1}(D) - m(D)\mu^*\|_W - \|X_t(D)P_{\bar{\pi}^*} - m(D)\mu^*\|_W \\
&\leq \|X'_{t+1}(D) - X_t(D)P_{\bar{\pi}^*}\|_W \\
&\leq \lambda_W^{1/2} \|X'_{t+1}(D) - X_t(D)P_{\bar{\pi}^*}\|_2 \\
&\leq \lambda_W^{1/2} \|X'_{t+1}(D) - X_t(D)P_{\bar{\pi}^*}\|_1.
\end{aligned}$$

Therefore, it suffices to show that

$$\mathbb{E}[\|X'_{t+1}(D) - X_t(D)P_{\bar{\pi}^*}\|_1 \mid X_t] = O(1/\sqrt{N}) \quad a.s. \quad (84)$$

$$\mathbb{P}[\|X'_{t+1}(D) - X_t(D)P_{\bar{\pi}^*}\|_1 > r \mid X_t] = O(\exp(-C\lambda_W N r^2)) \quad a.s. \quad \forall r \geq 0. \quad (85)$$

To show (84) and (85), we examine the distribution of $\|X'_{t+1}(D) - X_t(D)P_{\bar{\pi}^*}\|_1$ given X_t . Note that by Lemma 4, $\mathbb{E}[X'_{t+1}(D) \mid X_t] = X_t(D)P_{\bar{\pi}^*}$.

$$\begin{aligned}
\|X'_{t+1}(D) - X_t(D)P_{\bar{\pi}^*}\|_1 &= \|X'_{t+1}(D) - \mathbb{E}[X'_{t+1}(D) \mid X_t]\|_1 \\
&\leq \|X'_{t+1}(D) - \mathbb{E}[X'_{t+1}(D) \mid X_t, (A_t(i))_{i \in D}]\|_1 \quad (86)
\end{aligned}$$

$$+ \|\mathbb{E}[X'_{t+1}(D) \mid X_t, (A_t(i))_{i \in D}] - \mathbb{E}[X'_{t+1}(D) \mid X_t]\|_1. \quad (87)$$

We first derive an expectation bound and a high-probability bound for (86). Note that for each $s \in \mathbb{S}$,

$$X'_{t+1}(D, s) = \frac{1}{N} \sum_{i \in D} \mathbb{1}\{S'_{t+1}(i) = s\},$$

where $S'_{t+1}(i)$ denote the state of arm i at time $t+1$ if the arms in D follow Exact Proportional Control at time t . Given X_t and $(A_t(i))_{i \in D}$, $\mathbb{1}\{S'_{t+1}(i) = s\}$ for $i \in D$ are independent Bernoulli random variables, each with mean $P(S_t(i), A_t(i), s)$. By Cauchy-Schwartz,

$$\begin{aligned}
&\mathbb{E}\left[\left|X'_{t+1}(D, s) - \mathbb{E}[X'_{t+1}(D, s) \mid X_t, (A_t(i))_{i \in D}]\right| \mid X_t, (A_t(i))_{i \in D}\right] \\
&\leq \text{Var}\left[X'_{t+1}(D, s) \mid X_t, (A_t(i))_{i \in D}\right]^{1/2} \\
&= \frac{1}{N} \left(\sum_{i \in D} \text{Var}\left[\mathbb{1}\{S'_{t+1}(i) = s\}, \mid X_t, (A_t(i))_{i \in D}\right] \right)^{1/2} \\
&\leq \frac{1}{\sqrt{N}}.
\end{aligned}$$

By Hoeffding inequality, for all $r \geq 0$,

$$\mathbb{P}\left[\left|X'_{t+1}(D, s) - \mathbb{E}[X'_{t+1}(D, s) \mid X_t, (A_t(i))_{i \in D}]\right| > r \mid X_t, (A_t(i))_{i \in D}\right] \leq 2 \exp(-2Nr^2).$$

Therefore, summing over all $s \in \mathbb{S}$ and taking expectations over $(A_t(i))_{i \in D}$, we get

$$\mathbb{E}\left[\left\|\left|X'_{t+1}(D, s) - \mathbb{E}[X'_{t+1}(D, s) \mid X_t, (A_t(i))_{i \in D}]\right|\right\|_1 \mid X_t\right] \leq \frac{|\mathbb{S}|}{\sqrt{N}} \quad (88)$$

$$\mathbb{P}\left[\left\|\left|X'_{t+1}(D, s) - \mathbb{E}[X'_{t+1}(D, s) \mid X_t, (A_t(i))_{i \in D}]\right|\right\|_1 > r \mid X_t\right] \leq 2|\mathbb{S}| \exp\left(-\frac{2Nr^2}{|\mathbb{S}|^2}\right), \quad (89)$$

where (89) comes from applying a union bound.

Next, we bound the term in (87). For any $s \in \mathbb{S}$, we have

$$\begin{aligned}\mathbb{E} [X'_{t+1}(D, s) | X_t, (A_t(i))_{i \in D}] &= \sum_{s' \in \mathbb{S}, a \in \mathbb{A}} \bar{\pi}^*(a|s) X_t(D, s') P(s', a, s) \\ \mathbb{E} [X'_{t+1}(D, s) | X_t] &= \sum_{s' \in \mathbb{S}, a \in \mathbb{A}} \Gamma_t(s', a) P(s', a, s),\end{aligned}$$

where $N \cdot \Gamma_t(s, a)$ represents the number of arms in D with state s taking action a at time t . By the definition of Exact Proportional Control (Subroutine 1),

$$\Gamma_t(s', 1) \in \{ \lceil N \bar{\pi}^*(1|s') X_t(D, s') \rceil / N, \lfloor N \bar{\pi}^*(1|s') X_t(D, s') \rfloor / N \},$$

and $\Gamma_t(s, 0) = X_t(D, s) - \Gamma_t(s, 1)$. Consequently,

$$\begin{aligned}& \left| \mathbb{E} [X'_{t+1}(D, s) | X_t, (A_t(i))_{i \in D}] - \mathbb{E} [X'_{t+1}(D, s) | X_t] \right| \\ & \leq \sum_{s' \in \mathbb{S}, a \in \mathbb{A}} |\bar{\pi}^*(a|s) X_t(D, s') - \Gamma_t(s', a)| P(s', a, s) \\ & \leq \frac{1}{N} \sum_{s' \in \mathbb{S}, a \in \mathbb{A}} P(s', a, s) \quad a.s.\end{aligned}\tag{90}$$

Summing over all $s \in \mathbb{S}$ in (90), we get

$$\left\| \mathbb{E} [X'_{t+1}(D, s) | X_t, (A_t(i))_{i \in D}] - \mathbb{E} [X'_{t+1}(D, s) | X_t] \right\|_1 \leq \frac{2|\mathbb{S}|}{N} \quad a.s.\tag{91}$$

Combining the above calculations, we get

$$\begin{aligned}& \mathbb{E} \left[\left\| X'_{t+1}(D) - X_t(D) P_{\bar{\pi}^*} \right\|_1 \mid X_t \right] \\ & \leq \mathbb{E} \left[\left\| X'_{t+1}(D) - \mathbb{E} [X'_{t+1}(D, s) | X_t, (A_t(i))_{i \in D}] \right\|_1 \mid X_t \right] \\ & \quad + \mathbb{E} \left[\left\| \mathbb{E} [X'_{t+1}(D, s) | X_t, (A_t(i))_{i \in D}] - \mathbb{E} [X'_{t+1}(D, s) | X_t] \right\|_1 \mid X_t \right] \\ & \leq \frac{1}{\sqrt{N}} + \frac{2|\mathbb{S}|}{N},\end{aligned}\tag{92}$$

which implies (84). We also have that for any $r \geq 2|\mathbb{S}|/N$,

$$\begin{aligned}& \mathbb{P} \left[\left\| X'_{t+1}(D) - X_t(D) P_{\bar{\pi}^*} \right\|_1 > r \mid X_t \right] \\ & \leq \mathbb{P} \left[\left\| X'_{t+1}(D) - \mathbb{E} [X'_{t+1}(D, s) | X_t, (A_t(i))_{i \in D}] \right\|_1 > r - \frac{2|\mathbb{S}|}{N} \mid X_t \right] \\ & \leq 2|\mathbb{S}| \exp \left(- \frac{2N(r - 2|\mathbb{S}|/N)^2}{|\mathbb{S}|^2} \right) \\ & \leq 2|\mathbb{S}| \exp \left(- \frac{Nr^2}{|\mathbb{S}|^2} + \frac{8}{N} \right)\end{aligned}\tag{93}$$

where we use the fact that $(r - 2|\mathbb{S}|/N)^2 \geq r^2/2 - (2|\mathbb{S}|/N)^2$ in the (93). Note that the above upper bound also holds for $r < 2|\mathbb{S}|/N$ because the term in (93) is greater than 1 for $r < 2|\mathbb{S}|/N$. Because $\exp(8/N) = O(1)$, we have proved (85):

$$\mathbb{P} \left[\left\| X'_{t+1}(D) - X_t(D) P_{\bar{\pi}^*} \right\|_1 > r \mid X_t \right] = O(\exp(-C\lambda_W N r^2)),$$

where $C = 1/(\lambda_W |\mathbb{S}|^2)$. This finishes the proof that

$$\mathbb{E}[(h_W(X'_{t+1}, D) - \rho_w h_W(X_t, D))^+ | X_t] \leq O(1/\sqrt{N}) \quad a.s. \quad (34)$$

$$\mathbb{P}[h_W(X'_{t+1}, D) > \rho_w h_W(X_t, D) + r | X_t] \leq O(\exp(-CNr^2)) \quad a.s. \quad \forall r \geq 0. \quad (35)$$

Proving the analogs of (34) and (35) for h_U . Now let X'_{t+1} denote the system at time $t+1$ given that the arms in D follow the Optimal Local Control at time t . The proof is almost verbatim: We first show

$$\mathbb{E} \left[\left\| X'_{t+1}(D, s) - \mathbb{E}[X'_{t+1}(D, s) | X_t, (A_t(i))_{i \in D}] \right\|_1 \middle| X_t \right] \leq \frac{|\mathbb{S}|}{\sqrt{N}} \quad (94)$$

$$\mathbb{P} \left[\left\| X'_{t+1}(D, s) - \mathbb{E}[X'_{t+1}(D, s) | X_t, (A_t(i))_{i \in D}] \right\|_1 > r \middle| X_t \right] \leq 2|\mathbb{S}| \exp \left(-\frac{2Nr^2}{|\mathbb{S}|^2} \right), \quad (95)$$

using identical arguments. Then we can show that

$$\left\| \mathbb{E} [X'_{t+1}(D, s) | X_t, (A_t(i))_{i \in D}] - \mathbb{E} [X'_{t+1}(D, s) | X_t] \right\|_1 \leq \frac{4|\mathbb{S}|}{N} \quad a.s., \quad (96)$$

using the fact that under Optimal Local Control (Subroutine 2), for all $s' \in \mathbb{S}$ and $a \in \mathbb{A}$,

$$\begin{aligned} |\bar{\pi}^*(a|s)X_t(D, s') - \Gamma_t(s', a)| &\leq \frac{1}{N} \quad a.s. \quad \forall s' \neq \tilde{s} \\ |\bar{\pi}^*(a|\tilde{s})X_t(D, s') - \Gamma_t(\tilde{s}, a)| &\leq \frac{|\mathbb{S}|}{N} \quad a.s. \end{aligned}$$

Finally, we combine (94), (95) and (96) to get that under h_U ,

$$\mathbb{E}[(h_W(X'_{t+1}, D) - \rho_w h_W(X_t, D))^+ | X_t] \leq O(1/\sqrt{N}) \quad a.s. \quad (97)$$

$$\mathbb{P}[h_W(X'_{t+1}, D) > \rho_w h_W(X_t, D) + r | X_t] \leq O(\exp(-C_U Nr^2)) \quad a.s. \quad \forall r \geq 0, \quad (98)$$

for some constant $C_U > 0$.

The rest of the inequalities in this lemma. The other two inequalities, (36) and (37), and their analogs for h_U are relatively straightforward from the definitions of the weighted L_2 norms, so we skip them here. Similar calculations have also been done in [HXCW24] for h_W , so we refer the readers to [HXCW24] for the detailed calculations. \square

C Examples and proofs of violating assumptions

C.1 Examples of violating Assumptions 1-2

An aperiodic but non-unichain example. We first give an example of RBs where $P_{\bar{\pi}^*}$ is aperiodic but not unichain. Consider the RB problem whose single-armed MDP has two states, A and B , with no transitions between these two states. The reward function is given by $r(A, 1) = r(B, 0) = 1$ and $r(A, 0) = r(B, 1) = 0$. Let $\alpha = 1/2$ in the budget constraint. Since there is no transition between these two states, the Markov chain induced by the optimal single-armed policy, $\bar{\pi}^*$, $P_{\bar{\pi}^*}$, is not unichain. However, each of the recurrent classes $\{A\}$ and $\{B\}$ is aperiodic in this Markov chain.

It is easy to verify that the optimal solution to the LP relaxation (LP) is $y^*(A, 1) = y^*(B, 0) = 1/2$ and $y^*(s, a) = 0$ for the rest of $s \in \mathbb{S}, a \in \mathbb{A}$; the optimal value of (LP) is $R^{\text{rel}} = 1$. Now consider the N -armed problem where all arms are initialized in state A . In this case, the long-run average reward of any policy is $1/2$. Therefore, $R^{\text{rel}} - R(\pi, \mathbf{S}_0) \geq 1/2 = \Omega(1/\sqrt{N})$ for any π .

A unichain but periodic example. Next, we give an example of RBs where P_{π^*} is unichain but periodic, and $R^{\text{rel}} - R(\pi, \mathbf{S}_0) = \Theta(1)$. This example was included in [HXCW24], and we paraphrase it here for completeness.

Consider the RB problem whose single-armed MDP has two states, A and B . The state of the single-armed MDP deterministically transitions to the other state in each time step, regardless of the action applied. Let the reward function be $r(A, 0) = r(B, 1) = 1$ and $r(A, 1) = r(B, 0) = 0$; let $\alpha = 1/2$ in the budget constraint. Obviously, the Markov chain induced by any single-armed policy is unichain and periodic with a period 2.

One can easily see the optimal solution to the LP relaxation (LP) is by $y^*(A, 0) = y^*(B, 1) = 1/2$ and $y^*(A, 1) = y^*(B, 0) = 0$; the optimal value of (LP) is $R^{\text{rel}} = 1$. Now consider the N -armed problem where all arms are initialized in state A . Then at any time t , either all arms are in state A or all arms are in state B . In this case, under any policy π , the long-run average $R(\pi, \mathbf{S}_0)$ is $1/2$, so $R^{\text{rel}} - R(\pi, \mathbf{S}_0) = 1/2 = \Omega(1/\sqrt{N})$.

An example satisfying Assumption 1 but not Assumption 2. Next, we give an example of RB satisfying Assumption 1 but violating Assumption 2. Suppose each arm in the N -armed problem has state $S_t(i) = \text{Ber}(1/2)$ i.i.d. across i and t . Let $\alpha = 1/2$ in the budget constraint. The reward function $r(s, 1) = s$ and $r(s, 0) = 0$ for $s = 0, 1$. Intuitively, the decision maker observes the state of each arm, and pulls $N/2$ of them. Each arm pulled by the decision maker generates an amount of reward equal to its state. The single-armed MDP is clearly aperiodic unichain under any policy because its state is i.i.d.

The optimal solution of the LP relaxation (LP) is $y^*(1, 1) = y^*(0, 0) = 1/2$, and $y^*(s, a) = 0$ for other $s \in \mathbb{S}, a \in \mathbb{A}$; the optimal value $R^{\text{rel}} = 1/2$. Intuitively, the optimal single-armed policy pulls the arm if and only if it is in state 1.

In the N -armed problem, the optimal policy is obviously pulling $N/2$ arms with the largest states. Then the optimal reward $R^*(N)$ is given by

$$R^*(N) = \frac{1}{N} \mathbb{E} \left[\min \left(\sum_{i \in [N]} S_i(t), \frac{N}{2} \right) \right],$$

for an arbitrary t . Since $S_t(i) = \text{Ber}(1/2)$ and are i.i.d. across i , it can be shown that $R^*(N) = 1/2 - \Theta(1/\sqrt{N})$. Therefore, $R^{\text{rel}} - R^*(N) = \Theta(1/\sqrt{N})$.

C.2 Proofs of Lemma 2 and Theorem 2

In this section, we prove our second main result, Theorem 2, which is a lower bound on $R^{\text{rel}} - R(\pi, \mathbf{S}_0)$ for regular unstable RBs (Definition 1); meanwhile we will also prove Lemma 2 stated in Section 4.2 as one of the preliminary steps for proving Theorem 2. This section is organized linearly, with a sequence of lemma statements and proofs; Theorem 2 will be proved at the end of this section.

We first restate and prove Lemma 2.

Lemma 2. *Suppose that the LP relaxation (LP) has a unique optimal solution y^* , and $y^*(s, 1) + y^*(s, 0) > 0, \forall s \in \mathbb{S}$. Then*

$$R^{\text{rel}} - R(\pi, \mathbf{S}_0) \geq \epsilon_0 \lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E}[d_{\text{IC}}(Y_t^\pi)], \quad (12)$$

where ϵ_0 is a positive constant; d_{IC} is a function of state-action distribution given by

$$d_{\text{IC}}(y) = \sum_{s \in S^+} y(s, 0) + \sum_{s \in S^-} y(s, 1),$$

where $S^+ = \{s \in \mathbb{S} : y^*(s, 1) > 0, y^*(s, 0) = 0\}$, $S^- = \{s \in \mathbb{S} : y^*(s, 1) = 0, y^*(s, 0) > 0\}$.

Proof of Lemma 2. We first investigate the dual of (LP), whose properties help us prove the lemma:

$$\begin{aligned}
& \underset{\lambda, \mu, \{f(s)\}_{s \in \mathbb{S}}}{\text{minimize}} \quad \mu && \text{(Dual-LP)} \\
& \text{subject to} \quad r(s, a) + (\alpha - \mathbb{1}\{a = 1\})\lambda - \mu + \sum_{s'} P(s, a, s')f(s') \leq f(s) \quad \forall s \in \mathbb{S}, a \in \mathbb{A} && (99) \\
& \quad \lambda, \mu, f(s) \in \mathbb{R} \quad \forall s \in \mathbb{S}
\end{aligned}$$

Note that the primal problem (LP) is always feasible and bounded, so (Dual-LP) is also feasible and bounded. By strong duality of linear programs, the optimal value of the dual problem is $\mu^* = R^{\text{rel}}$. Moreover, there always exists an optimal solution of the dual problem (λ^*, f^*, μ^*) that satisfies strict complementarity with the unique primal optimal solution y^* [GT56]:

$$r(s, a) - \lambda^* \mathbb{1}\{a = 1\} + \lambda^* \alpha - \mu^* + \sum_{s'} P(s, a, s')f^*(s') - f^*(s) = 0 \quad \text{iff} \quad y^*(s, a) = 0. \quad (100)$$

Let $r_{\lambda^*}(s, a) = r(s, a) - \lambda^* \mathbb{1}\{a = 1\} + \lambda^* \alpha$. By (100) and the definition of S^+ and S^- , there exists a constant $\epsilon_0 > 0$ such that

$$r_{\lambda^*}(s, 0) - R^{\text{rel}} + \sum_{s'} P(s, 0, s')f^*(s') \leq f^*(s) - \epsilon_0 \quad \forall s \in S^+ \quad (101)$$

$$r_{\lambda^*}(s, 1) - R^{\text{rel}} + \sum_{s'} P(s, 1, s')f^*(s') \leq f^*(s) - \epsilon_0 \quad \forall s \in S^-. \quad (102)$$

Define the function $Q^*: \mathbb{S} \times \mathbb{A} \rightarrow \mathbb{R}$ as

$$Q^*(s, a) = r_{\lambda^*}(s, a) - R^{\text{rel}} + \sum_{s'} P(s, a, s')f^*(s').$$

Then (99) implies that $f^*(s) \geq Q^*(s, a)$ for any $s \in \mathbb{S}, a \in \mathbb{A}$; (101) and (102) imply that $Q^*(s, 1) - Q^*(s, 0)$ is at least ϵ_0 for $s \in S^+$ and at most $-\epsilon_0$ for $s \in S^-$.

Now, we are ready to bound $R^{\text{rel}} - R(\pi, \mathbf{S}_0)$: for any policy π and initial state vector \mathbf{S}_0 ,

$$\begin{aligned}
& R^{\text{rel}} - R(\pi, \mathbf{S}_0) \\
&= \mathbb{E} \left[R^{\text{rel}} - \frac{1}{N} \sum_{i \in [N]} r_{\lambda^*}(S_\infty(i), A_\infty(i)) \right] \\
&= \frac{1}{N} \sum_{i \in [N]} \mathbb{E} \left[R^{\text{rel}} - r_{\lambda^*}(S_\infty(i), A_\infty(i)) \right] \\
&= \frac{1}{N} \sum_{i \in [N]} \mathbb{E} \left[R^{\text{rel}} - r_{\lambda^*}(S_\infty(i), A_\infty(i)) - \sum_{s' \in \mathbb{S}} P(S_\infty(i), A_\infty(i), s')f^*(s') + f^*(S_\infty(i)) \right] && (103) \\
&= \frac{1}{N} \sum_{i \in [N]} \mathbb{E} \left[-Q^*(S_\infty(i), A_\infty(i)) + f^*(S_\infty(i)) \right] \\
&\geq \frac{1}{N} \sum_{i \in [N]} \epsilon_0 \mathbb{E} \left[\mathbb{1}\{S_\infty(i) \in S^+, A_\infty(i) = 0\} + \mathbb{1}\{S_\infty(i) \in S^-, A_\infty(i) = 1\} \right] && (104) \\
&= \epsilon_0 \mathbb{E} \left[\sum_{s \in S^+} Y_\infty(s, 0) + \sum_{s \in S^-} Y_\infty(s, 1) \right]. && (105)
\end{aligned}$$

where to get (103), we use the fact that $\mathbb{E}[\sum_{s' \in \mathbb{S}} P(S_\infty(i), A_\infty(i), s')f^*(s')] = \mathbb{E}[f^*(S_\infty(i))]$ for any function $f: \mathbb{S} \rightarrow \mathbb{R}$. \square

Next, we show Lemma 13, which implies that under any actions for the N -arms, one can find a subset of arms (can be empty) such that these almost follow the Optimal Local Control; the size of the subset is determined by $d_{\text{IC}}(Y_t)$. Essentially, this lemma shows that under any policy, if $d_{\text{IC}}(Y_t)$ is small, the transition dynamics will be close to the transition dynamics under Optimal Local Control.

Lemma 13. *Consider an instance of regular unstable RBs (Definition 1). Under any policy π , at any time t , there exists a set $D_t \subseteq [N]$ depending on X_t and \mathbf{A}_t with size $m(D_t) \geq 1 - 2d_{\text{IC}}(Y_t)/\beta - 1/(\beta N)$, where $\beta = \min(\alpha, 1 - \alpha)$, such that if we resample actions for the arms in D_t using Optimal Local Control, we get almost the same actions as A_t up to at most 1 arm. For such D_t , we have*

- The expectation of $X_{t+1}(D_t)$ satisfies

$$\left\| \mathbb{E}[X_{t+1}(D_t) - m(D_t)\mu^* \mid X_t, \mathbf{A}_t] - (X_t(D_t) - m(D_t)\mu^*)\Phi \right\|_1 \leq \frac{2}{N}. \quad (106)$$

- Let $X'_{t+1}(D_t)$ be the state distribution at time $t + 1$ when the actions for the arms in D_t are resampled using Optimal Local Control, then there exists a coupling between X'_{t+1} and X_{t+1} such that

$$\|X'_{t+1}(D_t) - X_{t+1}(D_t)\|_1 \leq \frac{2}{N} \text{ a.s.} \quad (107)$$

Proof of Lemma 13. We construct the subset D_t by removing a suitable subset of arms from $[N]$. We first remove each arm i with state $S_t(i) \in S^+$ and $A_t(i) = 0$ or $S_t(i) \in S^-$ and $A_t(i) = 1$. After this step, we get a subset D' with $N(1 - d_{\text{IC}}(Y_t))$ arms. The budget usage of D' equals $\alpha N - N \sum_{s \in S^-} Y_t(s, 1)$. Then D_t is obtained by removing arms that take action 1 or action 0 from D' such that the budget usage of D_t equals $\lfloor \alpha |D_t| \rfloor$. If $\alpha N - N \sum_{s \in S^-} Y_t(s, 1) > \alpha |D'|$, it is not hard to show that the number of arms removed from D' is bounded by

$$|D'| - |D_t| \leq \frac{1}{1 - \alpha} \left(\left| \alpha N - N \sum_{s \in S^-} Y_t(s, 1) - \alpha |D'| \right| + 1 \right) \leq \frac{1}{1 - \alpha} (Nd_{\text{IC}}(Y_t) + 1);$$

If $\alpha N - N \sum_{s \in S^-} Y_t(s, 1) > \alpha |D'|$, it is not hard to show that the number of arms removed from D' is bounded by

$$|D'| - |D_t| \leq \frac{1}{\alpha} \left(\left| \alpha N - N \sum_{s \in S^-} Y_t(s, 1) - \alpha |D'| \right| + 1 \right) \leq \frac{1}{\alpha} (Nd_{\text{IC}}(Y_t) + 1).$$

In both cases, we have

$$|D'| - |D_t| \leq \frac{1}{\beta} (Nd_{\text{IC}}(Y_t) + 1), \quad (108)$$

where recall that $\beta = \min(\alpha, 1 - \alpha)$. Now we get a subset D_t such that $m(D_t) \geq 1 - d_{\text{IC}}(Y_t) - (d_{\text{IC}}(Y_t) + 1/N)/\beta \geq 1 - 2d_{\text{IC}}(Y_t)/\beta - 1/(\beta N)$. Moreover, for each $i \in D_t$, $A_t(i) = 1$ if $S_t(i) \in S^+$; $A_t(i) = 0$ if $S_t(i) \in S^-$; the total budget usage of D_t equals $\lfloor \alpha |D_t| \rfloor$.

Following similar calculations as the proof of Lemma 1, we can show that

$$\mathbb{E}[X_{t+1}(D_t) \mid X_t, \mathbf{A}_t] = X_t(D_t) \left(P_{\bar{\pi}^*} - c_{\bar{\pi}^*}^\top (P_1(\tilde{s}) - P_0(\tilde{s})) \right) + m(D_t) \frac{\lfloor \alpha N \rfloor}{N} (P_1(\tilde{s}) - P_0(\tilde{s})).$$

whereas by the definition of Φ ,

$$(X_t(D_t) - m(D_t)\mu^*)\Phi + m(D_t)\mu^* = X_t(D_t) \left(P_{\bar{\pi}^*} - c_{\bar{\pi}^*}^\top (P_1(\tilde{s}) - P_0(\tilde{s})) \right) + m(D_t)\alpha(P_1(\tilde{s}) - P_0(\tilde{s})).$$

Comparing the above two equations, we get (106).

To show (107), note that if we resample the actions using the Optimal Local Control for the arms in D_t , the actions of the arms in S^+ and S^- are unchanged; at most one arm with state \tilde{s} gets a different action after resampling. Therefore, we can couple X'_{t+1} and X_{t+1} such that for each arm that gets the same actions after resampling, the next states of the arm are the same. In this way, we get $\|X'_{t+1}(D_t) - X_{t+1}(D_t)\|_1 \leq 2/N$. \square

Next, we conduct Lyapunov analysis with the help of Lemma 13. To construct Lyapunov functions, we first define two seminorms. Consider the Jordan decomposition of Φ , $\Phi = VJV^{-1}$. We split J into $J = J_+ + J_-$, where J_+ contains the Jordan blocks of J with eigenvalues' moduli greater than 1, and J_- contains the Jordan blocks of J with eigenvalues' moduli smaller than 1. Let I_+ be the diagonal matrix with 1 on the entries where J_+ is non-zero; let I_- be the identity matrix subtracting I_+ . We define the matrices U_+ and U_- via the infinite series:

$$U_+ = \sum_{k=1}^{\infty} (VJ_+^\dagger V^{-1})^k (VJ_+^\dagger V^{-1})^{\top k} \quad (109)$$

$$U_- = VI_- V^{-1} (VI_- V^{-1})^\top + \sum_{k=1}^{\infty} (VJ_- V^{-1})^k (VJ_- V^{-1})^{\top k} \quad (110)$$

where the superscript \dagger denotes pseudo-inverse.

In the next lemma, we show that U_+ and U_- are well-defined, and prove several useful facts about the semi-norms weighted by U_+ and U_- .

Lemma 14. *For any regular unstable RBs (Definition 1), the matrices U_+ and U_- given in (109) and (110) are well-defined and positive-semi-definite. Further, if we let $\|v\|_{U_+} = \sqrt{vU_+v^\top}$ and $\|v\|_{U_-} = \sqrt{vU_-v^\top}$ for any $v \in \mathbb{R}^{|\mathcal{S}|}$, then*

$$\|v\Phi\|_{U_+} \geq (1 + \epsilon_1) \|v\|_{U_+} \quad (111)$$

$$\|v\Phi\|_{U_+} \leq (1 + \epsilon_2) \|v\|_{U_+} \quad (112)$$

$$\|v\Phi\|_{U_-} \leq (1 - \epsilon_3) \|v\|_{U_-}, \quad (113)$$

where $\epsilon_1, \epsilon_2, \epsilon_3$ are positive constants. Moreover, there exists a positive constant K_{norm} such that for any $v \in \mathbb{R}^{|\mathcal{S}|}$,

$$\|v\|_{U_+} + \|v\|_{U_-} \geq K_{\text{norm}} \|v\|_1. \quad (114)$$

Proof of Lemma 14. To show the well-definedness U_+ and U_- , note that the moduli of all eigenvalues of $VJ_+^\dagger V^{-1}$ and $VJ_- V^{-1}$ are strictly less than 1, so their k -th powers must converge exponentially as $k \rightarrow \infty$, implying the convergence of the series in (109) and (110). Moreover, since each term in the two series is positive semi-definite, both series converge to positive semi-definite matrices.

Next, we show (111). Observe that

$$\begin{aligned} \Phi U_+ \Phi^\top &= \sum_{k=1}^{\infty} (VJ_+^\dagger V^{-1})^k (VJ_+^\dagger V^{-1})^{\top k} \\ &= (VJ_+^\dagger V^{-1}) (VJ_+^\dagger V^{-1})^\top + \sum_{k=2}^{\infty} (VJ_+^\dagger V^{-1})^k (VJ_+^\dagger V^{-1})^{\top k} \\ &= (VI_+ V^{-1}) (VI_+ V^{-1})^\top + \sum_{k=2}^{\infty} (VJ_+^\dagger V^{-1})^k (VJ_+^\dagger V^{-1})^{\top k} \end{aligned}$$

$$= (VI_+V^{-1})(VI_+V^{-1})^\top + U_+.$$

We claim that there exists a positive constant ϵ'_1 such that for any $v \in \mathbb{R}^{|\mathbb{S}|}$,

$$v(VI_+V^{-1})(VI_+V^{-1})^\top v^\top \geq \epsilon'_1 vU_+v^\top, \quad (115)$$

Without loss of generality, we only need to show (115) for v not being the zero vector. By normalization, it suffices to show that the function $v \mapsto vU_+v^\top$ is bounded on the set $\{v: \|v(VI_+V^{-1})\|_2 = 1\}$. Let \mathcal{V}_+ be the linear subspace spanned by rows of V corresponding to the non-zero entries on I_+ 's diagonal, and let \mathcal{V}_- be the linear subspace spanned by rows of V corresponding to the zero entries on I_+ 's diagonal. Observe that $\mathbb{R}^{|\mathbb{S}|}$ can be represented as the direct sum $\mathcal{V}_+ \oplus \mathcal{V}_-$, which implies that any $v \in \mathbb{R}^{|\mathbb{S}|}$ can be decomposed as $v = v^\parallel + v^\perp$, where $v^\parallel \in \mathcal{V}_+$ and $v^\perp \in \mathcal{V}_-$. Moreover, \mathcal{V}_- is the kernel of VI_+V^{-1} , and VI_+V^{-1} is the identity mapping when restricted to \mathcal{V}_+ . Therefore, if $\|v(VI_+V^{-1})\|_2 = 1$, then $\|v^\parallel\|_2 = 1$. Observe that \mathcal{V}_- is also the kernel of U_- , so

$$vU_+v^\top = (v^\parallel + v^\perp)U_+(v^\parallel + v^\perp)^\top = v^\parallel U_+(v^\parallel)^\top.$$

Because $\{v \in \mathcal{V}_+: \|v^\parallel\|_2 = 1\}$ is a compact set, vU_+v^\top is bounded when $\|v(VI_+V^{-1})\|_2 = 1$. After getting (115), we have that

$$\begin{aligned} \|v\Phi\|_{U_+} &= \sqrt{v\Phi U_+ \Phi^\top v^\top} \\ &\geq \sqrt{(1 + \epsilon'_1)vU_+v^\top} \\ &= \sqrt{1 + \epsilon'_1} \|v\|_{U_+}. \end{aligned}$$

Letting $\epsilon_1 = \sqrt{1 + \epsilon'_1} - 1$, we obtain (111).

Next, we prove (112). For any $v \in \mathbb{R}^{|\mathbb{S}|}$, we have $vU_+v^\top \geq \|vVJ_+^\dagger V^{-1}\|_2^2$. We claim that there exists a positive constant ϵ'_2 such that

$$\|vVI_+V^{-1}\|_2 \leq \epsilon'_2 \|vVJ_+^\dagger V^{-1}\|_2. \quad (116)$$

To show (116), without loss of generality, it suffices to show that the function $v \mapsto \|vVI_+V^{-1}\|_2$ is bounded on the set $\{v: \|vVJ_+^\dagger V^{-1}\|_2 = 1\}$. Consider the decomposition $\mathbb{R}^{|\mathbb{S}|} = \mathcal{V}_+ \oplus \mathcal{V}_-$. Because the kernel of $VJ_+^\dagger V^{-1}$ is \mathcal{V}_- , and $VJ_+^\dagger V^{-1}$ is invertible when restricted to \mathcal{V}_+ , the set $\{v: \|vVJ_+^\dagger V^{-1}\|_2 = 1\}$ is the sum of \mathcal{V}_- and a compact subset of \mathcal{V}_+ . Because \mathcal{V}_- is also the kernel of vVI_+V^{-1} , $\|vVI_+V^{-1}\|_2$ is bounded. This proves (116). Because $\Phi U_+ \Phi^\top = (VI_+V^{-1})(VI_+V^{-1})^\top + U_+$, (116) implies that

$$\begin{aligned} \|v\Phi\|_{U_+} &\leq \sqrt{v\Phi U_+ \Phi^\top v^\top} \\ &= \sqrt{v(VI_+V^{-1})(VI_+V^{-1})^\top v^\top + vU_+v^\top} \\ &\leq \sqrt{(\epsilon'_2)^2 + 1} \|v\|_{U_+}. \end{aligned}$$

Taking $\epsilon_2 = \sqrt{(\epsilon'_2)^2 + 1} - 1$ finishes the proof.

Next, we show (113). Observe that

$$\Phi U_- \Phi^\top = VJ I_- V^{-1} (VJ I_- V^{-1})^\top + \sum_{k=1}^{\infty} (VJ J_-^k V^{-1}) (VJ J_-^k V^{-1})^\top$$

$$\begin{aligned}
&= \sum_{k=1}^{\infty} (VJ_-^k V^{-1})(VJ_-^k V^{-1})^\top \\
&= U_- - VI_- V^{-1}(VI_- V^{-1})^\top.
\end{aligned}$$

We claim that there exists a positive constant ϵ'_3 such that for any $v \in \mathbb{R}^{|\mathcal{S}|}$,

$$v(VI_- V^{-1})(VI_- V^{-1})^\top v^\top \geq \epsilon'_3 vU_- v^\top. \quad (117)$$

The argument for (117) is similar to the argument for (115), so we omit it here. Therefore,

$$\begin{aligned}
\|v\Phi\|_{U_-} &= \sqrt{v\Phi U_- \Phi^\top v^\top} \\
&\leq \sqrt{(1 - \epsilon'_3)vU_- v^\top} \\
&= \sqrt{1 - \epsilon'_3} \|v\|_{U_-}.
\end{aligned}$$

Taking $\epsilon_3 = 1 - \sqrt{1 - \epsilon'_3}$, we obtain (113).

To show (114), note that

$$\begin{aligned}
\|v\|_{U_+} + \|v\|_{U_-} &= \left\| vU_+ v^\top \right\| + \left\| vU_- v^\top \right\| \\
&\geq \left\| v(U_+ + U_-)v^\top \right\| \\
&= \|v\|_{U_+ + U_-}.
\end{aligned}$$

Since all norms in a finite-dimensional Euclidean space are equivalent, it suffices to show $\|\cdot\|_{U_+ + U_-}$ is a norm, or equivalently, $U_+ + U_-$ is positive definite. For any $v \in \mathbb{R}^{|\mathcal{S}|}$, we have

$$\begin{aligned}
v(U_+ + U_-)v^\top &\geq v \left((VJ_+^\dagger V^{-1})(VJ_+^\dagger V^{-1})^\top + VI_- V^{-1}(VI_- V^{-1})^\top \right) v^\top \\
&= vV(J_+^\dagger V^{-1}V^{-\top}J_+^\dagger + I_- V^{-1}V^{-\top}I_-)V^\top v^\top.
\end{aligned}$$

By the definitions of J_+ and I_- , it is not hard to see that we always have $v(U_+ + U_-)v^\top \geq 0$, and v is the zero vector whenever $v(U_+ + U_-)v^\top = 0$. Therefore, $U_+ + U_-$ is positive definite, which implies (114). \square

Now we are ready to conduct the Lyapunov analysis. We define the Lyapunov functions h_- and h_+ : for any system state x and $D \subseteq [N]$, let

$$h_-(x, D) = \|x(D) - m(D)\mu^*\|_{U_-} \quad (118)$$

$$h_+(x, D) = \|x(D) - m(D)\mu^*\|_{U_+}. \quad (119)$$

Our next lemma gives a lower bound for the drift of $h_-(x, D)$ under *any policy*.

Lemma 15. *Consider an instance of regular unstable RBs (Definition 1). Let K_1 be any positive constant. Under any policy π , for any time step t , we have*

$$\begin{aligned}
&\mathbb{E}[h_+(X_{t+1}, [N]) \mid X_t, \mathbf{A}_t] - h_+(X_t, [N]) \\
&\geq \frac{\epsilon_1 K_1 - (2 + \epsilon_1 + \epsilon_2)K_1 \mathbb{1}\{\mathcal{E}_t^c\}}{\sqrt{N}} - \frac{6\lambda_+^{1/2}}{N} - 8\lambda_+^{1/2} d_{\text{IC}}(Y_t) \quad a.s.,
\end{aligned} \quad (120)$$

where the good event \mathcal{E}_t is given by

$$\mathcal{E}_t = \left\{ h_+(X_t, [N]) > \frac{K_1}{\sqrt{N}} + \frac{2\lambda_+^{1/2}d_{\text{IC}}(Y_t)}{\beta} + \frac{\lambda_+^{1/2}}{\beta N}, \|X_t([N]) - \mu^*\|_1 \leq \eta' - \frac{6d_{\text{IC}}(Y_t)}{\beta} - \frac{3}{\beta N} \right\},$$

and $K_1 = \lambda_+^{1/2}K_{\text{noise}}/(2 + \epsilon_1 + \epsilon_2)$.

Proof of Lemma 15. Let D_t be the set given in Lemma 13. By the definition of h_+ and the fact that $m(D_t) \geq 1 - 2d_{\text{IC}}(Y_t) - 1/N$, we have

$$\begin{aligned} & \mathbb{E}[h_+(X_{t+1}, [N]) \mid X_t, \mathbf{A}_t] - h_+(X_t, [N]) \\ & \geq \mathbb{E}[h_+(X_{t+1}, D_t) \mid X_t, \mathbf{A}_t] - h_+(X_t, D_t) - 4\lambda_+^{1/2}(1 - m(D_t)) \\ & \geq \mathbb{E}[h_+(X_{t+1}, D_t) \mid X_t, \mathbf{A}_t] - h_+(X_t, D_t) - 8\lambda_+^{1/2}d_{\text{IC}}(Y_t)/\beta - 4\lambda_+^{1/2}/(\beta N). \end{aligned} \quad (121)$$

Note that (121) uses the fact that $\|v\|_{U_+} \leq \lambda_+^{1/2} \|v\|_1$.

We first consider the case with $h_+(X_t, D_t) > K_1/\sqrt{N}$. We have

$$\begin{aligned} \mathbb{E}[h_+(X_{t+1}, D_t) \mid X_t, \mathbf{A}_t] &= \mathbb{E}[\|X_{t+1}(D_t) - m(D_t)\mu^*\|_{U_+} \mid X_t, \mathbf{A}_t] \\ &\geq \|\mathbb{E}[X_{t+1}(D_t) - m(D_t)\mu^* \mid X_t, \mathbf{A}_t]\|_{U_+} \\ &\geq \|(X_t(D_t) - m(D_t)\mu^*)\Phi\|_{U_+} - 2\lambda_+^{1/2}/(\beta N), \end{aligned} \quad (122)$$

$$\geq \|(X_t(D_t) - m(D_t)\mu^*)\Phi\|_{U_+} - 2\lambda_+^{1/2}/(\beta N), \quad (123)$$

where (122) uses Jensen's inequality; (123) invokes Lemma 13 and the fact that $\|v\|_{U_+} \leq \lambda_+^{1/2} \|v\|_1$ for any $v \in \mathbb{R}^{\mathbb{S}}$. By Lemma 14, we have

$$\|v\Phi\|_{U_+} \geq (1 + \epsilon_1)\|v\|_{U_+}.$$

Letting $v = X_t(D_t) - m(D_t)\mu^*$, combined with (123), we get

$$\begin{aligned} \mathbb{E}[h_+(X_{t+1}, D_t) \mid X_t, \mathbf{A}_t] &\geq (1 + \epsilon_1)h_+(X_t, D_t) - \frac{2\lambda_+^{1/2}}{\beta N} \\ &\geq h_+(X_t, D_t) + \frac{\epsilon_1 K_1}{\sqrt{N}} - \frac{2\lambda_+^{1/2}}{\beta N}, \end{aligned}$$

which implies that

$$\mathbb{E}[h_+(X_{t+1}, [N]) \mid X_t, \mathbf{A}_t] - h_+(X_t, [N]) \geq \frac{\epsilon_1 K_1}{\sqrt{N}} - \frac{6\lambda_+^{1/2}}{\beta N} - \frac{8\lambda_+^{1/2}d_{\text{IC}}(Y_t)}{\beta}. \quad (124)$$

Next, we consider the case of $h_+(X_t, D_t) \leq K_1/\sqrt{N}$:

$$\begin{aligned} & \mathbb{E}[h_+(X_{t+1}, D_t) \mid X_t, \mathbf{A}_t] \\ &= \mathbb{E}[\|X_{t+1}(D_t) - m(D_t)\mu^*\|_{U_+} \mid X_t, \mathbf{A}_t] \\ &\geq \mathbb{E}[\|X_{t+1}(D_t) - m(D_t)\mu^* - (X_t(D_t) - m(D_t)\mu^*)\Phi\|_{U_+} \mid X_t, \mathbf{A}_t] - \|(X_t(D_t) - m(D_t)\mu^*)\Phi\|_{U_+} \\ &\geq \mathbb{E}[\|X_{t+1}(D_t) - m(D_t)\mu^* - (X_t(D_t) - m(D_t)\mu^*)\Phi\|_{U_+} \mid X_t, \mathbf{A}_t] - (1 + \epsilon_2)h_+(X_t, D_t), \end{aligned} \quad (125)$$

where (125) uses the fact that $\|v\|_{\Phi U_+ \Phi^\top} \leq (1 + \epsilon_2)\|v\|_{U_+}$ for all $v \in \mathbb{R}^{\mathbb{S}}$, proved in Lemma 14. We plan to invoke Property (c) of regular unstable RBs to bound the first term in (125). Let $X'_{t+1}(D_t)$

be the state distribution at time $t + 1$ when the arms in D_t follow Optimal Local Control. By Lemma 13, there exists a coupling such that $\|X'_{t+1}(D_t) - X_{t+1}(D_t)\|_1 \leq 2/N$ almost surely, so

$$\begin{aligned} & \mathbb{E}[\|X_{t+1}(D_t) - m(D_t)\mu^* - (X_t(D_t) - m(D_t)\mu^*)\Phi\|_{U_+} \mid X_t, \mathbf{A}_t] \\ & \geq \mathbb{E}[\|X'_{t+1}(D_t) - m(D_t)\mu^* - (X_t(D_t) - m(D_t)\mu^*)\Phi\|_{U_+} \mid X_t, \mathbf{A}_t] - \frac{2\lambda_+^{1/2}}{N}. \end{aligned} \quad (126)$$

Let ξ^\top be the right eigenvector of U_+ corresponding to the eigenvalue λ_+ , with norm $\|\xi\|_2 = 1$. Because U_+ is positive semidefinite, $U_+ \geq \lambda_+ \xi^\top \xi$, so

$$\|v\|_{U_+} \geq \sqrt{\lambda_+ v \xi^\top \xi v^\top} = \lambda_+^{1/2} |v \xi^\top|,$$

for any $v \in \mathbb{R}^{|\mathcal{S}|}$. Moreover, we argue that ξ satisfies the conditions in Property (c) of regular unstable RBs: the column space of U_+ is a subspace of the column space of Φ . Because $\mu^* \Phi = 0$, we have $\mu^* U_+ = 0$. Therefore, $\mu^* \xi^\top = \lambda_+^{-1} \mu^* U_+ \xi^\top = 0$. Therefore, by Property (c) of regular unstable RBs,

$$\begin{aligned} & \mathbb{E}[\|X'_{t+1}(D_t) - m(D_t)\mu^* - (X_t(D_t) - m(D_t)\mu^*)\Phi\|_{U_+} \mid X_t, \mathbf{A}_t] \\ & \geq \lambda_+^{1/2} \mathbb{E}[\|(X'_{t+1}(D_t) - m(D_t)\mu^* - (X_t(D_t) - m(D_t)\mu^*)\Phi)\xi^\top\| \mid X_t, \mathbf{A}_t] \\ & \geq \frac{\lambda_+^{1/2} K_{\text{noise}}}{\sqrt{N}} \mathbb{1}\{\|X_t(D_t) - m(D_t)\mu^*\|_1 \leq m(D_t)\eta'\}. \end{aligned} \quad (127)$$

Combining (125), (126) and (127), we get

$$\begin{aligned} & \mathbb{E}[h_+(X_{t+1}, D_t) \mid X_t, \mathbf{A}_t] - h_+(X_t, D_t) \\ & \geq \frac{\lambda_+^{1/2} K_{\text{noise}}}{\sqrt{N}} \mathbb{1}\{\|X_t(D_t) - m(D_t)\mu^*\|_1 \leq m(D_t)\eta'\} - (2 + \epsilon_2)h_+(X_t, D_t) - \frac{2\lambda_+^{1/2}}{N} \\ & \geq \frac{\lambda_+^{1/2} K_{\text{noise}}}{\sqrt{N}} \mathbb{1}\{\|X_t(D_t) - m(D_t)\mu^*\|_1 \leq m(D_t)\eta'\} - \frac{(2 + \epsilon_2)K_1}{\sqrt{N}} - \frac{2\lambda_+^{1/2}}{N}. \end{aligned}$$

Therefore, when $h_+(X_t, D_t) \leq K_1/\sqrt{N}$,

$$\begin{aligned} & \mathbb{E}[h_+(X_{t+1}, [N]) \mid X_t, \mathbf{A}_t] - h_+(X_t, [N]) \\ & \geq \frac{\lambda_+^{1/2} K_{\text{noise}}}{\sqrt{N}} \mathbb{1}\{\|X_t(D_t) - m(D_t)\mu^*\|_1 \leq m(D_t)\eta'\} - \frac{(2 + \epsilon_2)K_1}{\sqrt{N}} - \frac{6\lambda_+^{1/2}}{\beta N} - \frac{8\lambda_+^{1/2} d_{\text{IC}}(Y_t)}{\beta}. \end{aligned} \quad (128)$$

Summarizing the bounds (124) and (128) and substituting $K_1 = \lambda_+^{1/2} K_{\text{noise}}/(2 + \epsilon_1 + \epsilon_2)$ in the bounds, we get

- When $h_+(X_t, D_t) > K_1/\sqrt{N}$ or $\|X_t(D_t) - m(D_t)\mu^*\|_1 \leq m(D_t)\eta'$, we have

$$\mathbb{E}[h_+(X_{t+1}, [N]) \mid X_t, \mathbf{A}_t] - h_+(X_t, [N]) \geq \frac{\epsilon_1 K_1}{\sqrt{N}} - \frac{6\lambda_+^{1/2}}{\beta N} - \frac{8\lambda_+^{1/2} d_{\text{IC}}(Y_t)}{\beta}. \quad (129)$$

- When $h_+(X_t, D_t) \leq K_1/\sqrt{N}$ and $\|X_t(D_t) - m(D_t)\mu^*\|_1 > m(D_t)\eta'$, we have

$$\mathbb{E}[h_+(X_{t+1}, [N]) \mid X_t, \mathbf{A}_t] - h_+(X_t, [N]) \geq -\frac{(2 + \epsilon_2)K_1}{\sqrt{N}} - \frac{6\lambda_+^{1/2}}{\beta N} - \frac{8\lambda_+^{1/2} d_{\text{IC}}(Y_t)}{\beta}. \quad (130)$$

Note that the conditions separating the two cases above are in terms of $X_t(D_t)$. To complete the proof, we need to state the conditions in terms of $X_t([N])$ instead, observe that

$$\begin{aligned} h_+(X_t, D_t) &\geq h_+(X_t, [N]) - \lambda_+^{1/2}(1 - m(D_t)) \\ &\geq h_+(X_t, [N]) - 2\lambda_+^{1/2}d_{\text{IC}}(Y_t)/\beta - \lambda_+^{1/2}/(\beta N) \\ \|X_t(D_t) - m(D_t)\mu^*\|_1 &\leq \|X_t([N]) - \mu^*\|_1 + 2(1 - m(D_t)) \\ &\leq \|X_t([N]) - \mu^*\|_1 + 4d_{\text{IC}}(Y_t)/\beta + 2/(\beta N). \end{aligned}$$

Therefore, $h_+(X_t, [N]) > K_1/\sqrt{N} + 2\lambda_+^{1/2}d_{\text{IC}}(Y_t)/\beta + \lambda_+^{1/2}/(\beta N)$ implies $h_+(X_t, D_t) > K_1/\sqrt{N}$; $\|X_t([N]) - \mu^*\|_1 \leq \eta' - 6d_{\text{IC}}(Y_t)/\beta - 3/(\beta N)$ implies that

$$\|X_t(D_t) - m(D_t)\mu^*\|_1 \leq \eta' - \frac{2d_{\text{IC}}(Y_t)}{\beta} - \frac{1}{\beta N} \leq \eta' - (1 - m(D_t)) \leq m(D_t)\eta',$$

which implies the bounds in the lemma statement. \square

In the next lemma, we upper bound the drift of $h_-(x, D)$ under *any policy*.

Lemma 16. *Consider an instance of regular unstable RBs (Definition 1). Under any policy π , for any time step t , we have*

$$\mathbb{E}[h_-(X_{t+1}, [N]) \mid X_t, \mathbf{A}_t] \leq (1 - \epsilon_3)h_-(X_t, [N]) + \frac{\lambda_-^{1/2}|\mathbb{S}|}{\sqrt{N}} + \frac{10\lambda_-^{1/2}|\mathbb{S}|}{\beta N} + \frac{8\lambda_-^{1/2}d_{\text{IC}}(Y_t)}{\beta} \quad a.s., \quad (131)$$

where λ_- is the largest eigenvalue of U_- .

Proof of Lemma 16. Let D_t be the set given in Lemma 13. Let $X'_{t+1}(D_t)$ be the state distribution at time $t+1$ when the actions for the arms in D_t are resampled using Optimal Local Control. By Lemma 13, there exists a coupling such that $\|X'_{t+1}(D_t) - X_{t+1}(D_t)\|_1 \leq 2/N$ almost surely. By the definition of h_- and the fact that $m(D_t) \geq 1 - 2d_{\text{IC}}(Y_t)/\beta - 1/(\beta N)$, we have

$$\begin{aligned} &\mathbb{E}[h_-(X_{t+1}, [N]) \mid X_t, \mathbf{A}_t] - (1 - \epsilon_3)h_-(X_t, [N]) \\ &\leq \mathbb{E}[h_-(X_{t+1}, D_t) \mid X_t, \mathbf{A}_t] - (1 - \epsilon_3)h_-(X_t, D_t) + 4\lambda_-^{1/2}(1 - m(D_t)) \\ &\leq \mathbb{E}[h_-(X_{t+1}, D_t) \mid X_t, \mathbf{A}_t] - (1 - \epsilon_3)h_-(X_t, D_t) + 8\lambda_-^{1/2}d_{\text{IC}}(Y_t)/\beta + 4\lambda_-^{1/2}/(\beta N) \\ &\leq \mathbb{E}[h_-(X'_{t+1}, D_t) \mid X_t, \mathbf{A}_t] - (1 - \epsilon_3)h_-(X_t, D_t) + 8\lambda_-^{1/2}d_{\text{IC}}(Y_t)/\beta + 6\lambda_-^{1/2}/(\beta N). \end{aligned} \quad (132)$$

Because X'_{t+1} is obtained by letting arms in D_t follows the Optimal Local Control, using similar arguments as the proof of Lemma 10, we get

$$\mathbb{E}[h_-(X'_{t+1}, D_t) \mid X_t, \mathbf{A}_t] - (1 - \epsilon_3)h_-(X_t, D_t) \leq \frac{\lambda_-^{1/2}|\mathbb{S}|}{\sqrt{N}} + \frac{4\lambda_-^{1/2}|\mathbb{S}|}{N}. \quad (133)$$

Plugging (133) into (132), we finish the proof. \square

Now we are ready to prove Theorem 2.

Proof of Theorem 2. For any system state x and subset $D \subseteq [N]$, define

$$h_{\text{lb}}(x, D) = L_{\text{lb}}h_+(x, D) - h_-(x, D),$$

where $L_{\text{lb}} = 2\lambda_-^{1/2}|\mathbb{S}|/\epsilon_1$. We will show that with an appropriate L_{lb} and for N larger than a constant threshold, we have

$$\mathbb{E}[h_{\text{lb}}(X_{t+1}, [N])] - \mathbb{E}[h_{\text{lb}}(X_t, [N])] \geq \Omega\left(\frac{1}{\sqrt{N}}\right) - \Theta(1) \cdot \mathbb{E}[d_{\text{IC}}(Y_t)] \quad \forall t \geq 0. \quad (134)$$

If we can prove (134), then taking the long-run average on two sides of the inequality implies

$$0 = \mathbb{E}[h_{\text{lb}}(X_\infty, [N])] - \mathbb{E}[h_{\text{lb}}(X_\infty, [N])] \geq \Omega\left(\frac{1}{\sqrt{N}}\right) - \Theta(1) \cdot \mathbb{E}[d_{\text{IC}}(Y_\infty)],$$

so $\mathbb{E}[d_{\text{IC}}(Y_\infty)] = \Omega(1/\sqrt{N})$. Combined with Lemma 2, we get $R^{\text{rel}} - R(\pi, \mathbf{S}_0) = \Omega(1/\sqrt{N})$, which proves Theorem 2.

Now we prove (134) using the formulas of $h_+(X_t, [N])$ and $h_-(X_t, [N])$'s drifts given in Lemma 15 and Lemma 16.

$$\begin{aligned} & \mathbb{E}[h_{\text{lb}}(X_{t+1}, [N]) \mid X_t, \mathbf{A}_t] - h_{\text{lb}}(X_t, [N]) \\ &= L_{\text{lb}} \left(\mathbb{E}[h_+(X_{t+1}, [N]) \mid X_t, \mathbf{A}_t] - h_+(X_t, [N]) \right) - \left(\mathbb{E}[h_-(X_{t+1}, [N]) \mid X_t, \mathbf{A}_t] - h_-(X_t, [N]) \right) \\ &\geq L_{\text{lb}} \left(\frac{\epsilon_1 K_1 - (2 + \epsilon_1 + \epsilon_2) K_1 \mathbb{1}\{\mathcal{E}_t^c\}}{\sqrt{N}} - \frac{6\lambda_+^{1/2}}{\beta N} - 8\lambda_+^{1/2} \beta^{-1} d_{\text{IC}}(Y_t) \right) \\ &\quad + \epsilon_3 h_-(X_t, [N]) - \frac{\lambda_-^{1/2} |\mathbb{S}|}{\sqrt{N}} - \frac{10\lambda_-^{1/2} |\mathbb{S}|}{\beta N} - 8\lambda_-^{1/2} \beta^{-1} d_{\text{IC}}(Y_t) \\ &= \left(L_{\text{lb}} \epsilon_1 K_1 - \lambda_-^{1/2} |\mathbb{S}| \right) \frac{1}{\sqrt{N}} + \left(\epsilon_3 h_-(X_t, [N]) - \frac{L_{\text{lb}}(2 + \epsilon_1 + \epsilon_2) K_1}{\sqrt{N}} \mathbb{1}\{\mathcal{E}_t^c\} \right) \end{aligned} \quad (135)$$

$$- (8L_{\text{lb}} \lambda_+^{1/2} + 8\lambda_-^{1/2}) \beta^{-1} d_{\text{IC}}(Y_t) - O\left(\frac{1}{N}\right). \quad (136)$$

Note that the coefficient of the $1/\sqrt{N}$ in (135) equals $\lambda_-^{1/2} |\mathbb{S}|$, which is positive.

Comparing (135)-(136) with our goal, (134), it remains to lower bound the second term in (135). First, observe that given the event \mathcal{E}_t , the second term of (135) is non-negative. So we only need to consider the event \mathcal{E}_t^c . By definition, \mathcal{E}_t^c implies that

$$\begin{aligned} h_+(X_t, [N]) &\leq \frac{K_1}{\sqrt{N}} + \frac{2\lambda_+^{1/2} d_{\text{IC}}(Y_t)}{\beta} + \frac{\lambda_+^{1/2}}{\beta N}, \\ \|X_t([N]) - \mu^*\|_1 &> \eta' - \frac{6d_{\text{IC}}(Y_t)}{\beta} - \frac{3}{\beta N}. \end{aligned}$$

By (114) in Lemma 14,

$$h_+(X_t, [N]) + h_-(X_t, [N]) \geq K_{\text{norm}} \|X_t([N]) - \mu^*\|_1,$$

so given \mathcal{E}_t^c ,

$$\begin{aligned} h_-(X_t, [N]) &\geq K_{\text{norm}} \|X_t([N]) - \mu^*\|_1 - h_+(X_t, [N]) \\ &\geq K_{\text{norm}} \left(\eta' - \frac{6d_{\text{IC}}(Y_t)}{\beta} - \frac{3}{\beta N} \right) - \frac{K_1}{\sqrt{N}} - \frac{2\lambda_+^{1/2} d_{\text{IC}}(Y_t)}{\beta} - \frac{\lambda_+^{1/2}}{\beta N} \\ &= K_{\text{norm}} \eta' - \frac{K_1}{\sqrt{N}} - (6K_{\text{norm}} + 2\lambda_+^{1/2}) \beta^{-1} d_{\text{IC}}(Y_t) - O\left(\frac{1}{N}\right). \end{aligned}$$

Therefore, given \mathcal{E}_t^c , combining the above lower bound, if $N \geq K_1^2(\epsilon_3 + L_{\text{lb}}(2 + \epsilon_1 + \epsilon_2))^2 / (K_{\text{norm}}\eta'\epsilon_3)^2$,

$$\begin{aligned} & \epsilon_3 h_-(X_t, [N]) - \frac{L_{\text{lb}}(2 + \epsilon_1 + \epsilon_2)K_1}{\sqrt{N}} \mathbb{1}\{\mathcal{E}_t^c\} \\ & \geq \epsilon_3 K_{\text{norm}}\eta' - \frac{K_1(\epsilon_3 + L_{\text{lb}}(2 + \epsilon_1 + \epsilon_2))}{\sqrt{N}} - \epsilon_3(6K_{\text{norm}} + 2\lambda_+^{1/2})\beta^{-1}d_{\text{IC}}(Y_t) - O\left(\frac{1}{N}\right), \\ & \geq -\epsilon_3(6K_{\text{norm}} + 2\lambda_+^{1/2})\beta^{-1}d_{\text{IC}}(Y_t) - O\left(\frac{1}{N}\right). \end{aligned} \quad (137)$$

Substituting (137) back to the second term of (135), we get

$$\begin{aligned} & \mathbb{E} [h_{\text{lb}}(X_{t+1}, [N]) \mid X_t, \mathbf{A}_t] - h_{\text{lb}}(X_t, [N]) \\ & \geq \frac{\lambda_-^{1/2}|\mathbb{S}|}{\sqrt{N}} - (6\epsilon_3 K_{\text{norm}} + 2\epsilon_3\lambda_+^{1/2} + 8L_{\text{lb}}\lambda_+^{1/2} + 8\lambda_-^{1/2})\beta^{-1}d_{\text{IC}}(Y_t) - O\left(\frac{1}{N}\right), \end{aligned}$$

which implies (134) and finishes the proof. \square

C.3 Remarks on Property (c) of regular unstable RBs (Definition 1)

In this section, we prove an intuitive sufficient condition for Property (c) of regular unstable RBs and also provide a weaker version of this property.

We first state and prove the intuitive sufficient condition for Property (c):

Lemma 17. *Consider an RB instance with the single-armed MDP $(\mathbb{S}, \mathbb{A}, P, r)$, where $P(s, a, s') > 0$ for all $s \in \mathbb{S}, a \in \mathbb{A}, s' \in \mathbb{S}$. Then this RB instance satisfies Property (c) in Definition 1.*

Proof. Throughout this proof, we fix X_t and \mathbf{A}_t and omit them in the conditioning when writing expectations or probabilities.

Recalling the statement of Property (c), it suffices to show that for any $\xi \in \mathbb{R}^{|\mathbb{S}|}$ with $\|\xi\|_2 = 1$ and $\mu^*\xi^\top = 0$, there exists a constant $K_{\text{noise}} > 0$ such that for any N larger than a constant and any time step t , if the local optimal policy is applicable, we have

$$\mathbb{E} [(X_{t+1}([N]) - \mu^* - (X_t([N]) - \mu^*)\Phi)\xi^\top \mid X_t, \mathbf{A}_t] \geq \frac{K_{\text{noise}}}{\sqrt{N}}, \quad (138)$$

where \mathbf{A}_t follows the Optimal Local Control. Note that in this case, the parameter η' in Property (c) of Definition 1 could be any positive number such that $\|X_t([N]) - \mu^*\|_1 \leq \eta'$ guarantees the condition (7) for applying the Optimal Local Control.

We first show that for any $\xi \in \mathbb{R}^{|\mathbb{S}|}$ such that $\|\xi\|_2 = 1$ and $\mu^*\xi^\top = 0$,

$$\xi_{\max} - \xi_{\min} \geq 1/\sqrt{|\mathbb{S}|}. \quad (139)$$

where $\xi_{\max} = \max_{s \in \mathbb{S}} \xi(s)$ and $\xi_{\min} = \min_{s \in \mathbb{S}} \xi(s)$. Observe that we must have either $\xi_{\max} \geq 1/\sqrt{|\mathbb{S}|}$ or $\xi_{\min} \geq -1/\sqrt{|\mathbb{S}|}$, because otherwise $\|\xi\|_2 < 1$. We first show (139) for the case with $\xi_{\max} \geq 1/\sqrt{|\mathbb{S}|}$. In this case, we must have $\xi_{\min} \leq 1/\sqrt{|\mathbb{S}|}$, because otherwise $\|\xi\|_2 > 1$. Let $\mathbb{1}_{\text{scaled}} \in \mathbb{R}^{|\mathbb{S}|}$ be the whose entries are all $1/\sqrt{|\mathbb{S}|}$. Then

$$\|\xi - \mathbb{1}_{\text{scaled}}\|_\infty \leq \max(\xi_{\max} - 1/\sqrt{|\mathbb{S}|}, 1/\sqrt{|\mathbb{S}|} - \xi_{\min}) \leq \xi_{\max} - \xi_{\min}.$$

On the other hand, because $\|\mu^*\|_1 = 1$ and $\mu^*\mathbb{1}_{\text{scaled}}^\top = 1/\sqrt{|\mathbb{S}|}$,

$$\|\xi - \mathbb{1}_{\text{scaled}}\|_\infty \geq \mu^*\mathbb{1}_{\text{scaled}}^\top - \mu^*\xi^\top = 1/\sqrt{|\mathbb{S}|}.$$

Combining the last two inequalities yields (139). For the case with $\xi_{\min} \leq -1/\sqrt{|\mathbb{S}|}$, (139) can be proved in an analogous way.

To show (138), we invoke the Paley-Zygmund inequality to get

$$\begin{aligned} & \mathbb{P} \left[(X_{t+1}([N])\xi^\top - \mathbb{E}[X_{t+1}([N])\xi^\top])^2 > \frac{1}{2} \text{Var} [X_{t+1}([N])\xi^\top] \right] \\ & \geq \frac{1}{4} \frac{\text{Var} [X_{t+1}([N])\xi^\top]^2}{\mathbb{E} \left[(X_{t+1}([N])\xi^\top - \mathbb{E}[X_{t+1}([N])\xi^\top])^4 \right]}. \end{aligned} \quad (140)$$

Next, we lower bound the second moment of $X_{t+1}([N])\xi^\top - \mathbb{E}[X_{t+1}([N])\xi^\top]$ in (140). For any timestep t , given X_t, \mathbf{A}_t , we have

$$X_{t+1}([N])\xi^\top = \sum_{s \in \mathbb{S}} X_{t+1}([N], s) \xi(s) \quad (141)$$

$$= \frac{1}{N} \sum_{s \in \mathbb{S}} \sum_{i \in [N]} \mathbb{1}\{\mathbf{S}_{t+1}(i) = s\} \xi(s) \quad (142)$$

$$= \frac{1}{N} \sum_{i \in [N]} \xi(\mathbf{S}_{t+1}(i)). \quad (143)$$

Note that conditioned on X_t, \mathbf{A}_t , $\xi(\mathbf{S}_{t+1}(i))$ are independent across i . Therefore, the variance of $X_{t+1}([N])\xi^\top$ is given by

$$\text{Var} [X_{t+1}([N])\xi^\top] = \frac{1}{N^2} \sum_{i \in [N]} \text{Var} [\xi(\mathbf{S}_{t+1}(i))].$$

Because $P(s, a, s') > 0$ for all $s, s' \in \mathbb{S}, a \in \mathbb{A}$, the support of $\mathbf{S}_{t+1}(i)$ is the whole state space \mathbb{S} . Moreover, because $\xi_{\max} - \xi_{\min} > 1/\sqrt{|\mathbb{S}|}$, we have

$$\begin{aligned} \text{Var} [\xi(\mathbf{S}_{t+1}(i))] & \geq \min_{s, a, s'} P(s, a, s') \left((\xi_{\max} - \mathbb{E} [\xi(\mathbf{S}_{t+1}(i))])^2 + (\xi_{\min} - \mathbb{E} [\xi(\mathbf{S}_{t+1}(i))])^2 \right) \\ & \geq \frac{1}{2} \min_{s, a, s'} P(s, a, s') (\xi_{\max} - \xi_{\min})^2 \\ & \geq \frac{\min_{s, a, s'} P(s, a, s')}{2|\mathbb{S}|}, \end{aligned}$$

where $\min_{s, a, s'}$ is a shorthand for $\min_{s, s' \in \mathbb{S}, a \in \mathbb{A}}$. Therefore,

$$\text{Var} [X_{t+1}([N])\xi^\top] \geq \frac{1}{N^2} \sum_{i \in [N]} \text{Var} [\xi(\mathbf{S}_{t+1}(i))] \geq \frac{\min_{s, a, s'} P(s, a, s')}{2|\mathbb{S}|N}. \quad (144)$$

Next, we upper bound the fourth moment of $X_{t+1}([N])\xi^\top - \mathbb{E}[X_{t+1}([N])\xi^\top]$. Using the fact that $X_{t+1}([N])\xi^\top = \sum_{i \in [N]} \xi(\mathbf{S}_{t+1}(i))/N$, we get

$$\begin{aligned} & \mathbb{E} \left[(X_{t+1}([N])\xi^\top - \mathbb{E}[X_{t+1}([N])\xi^\top])^4 \right] \\ & = \frac{1}{N^4} \left(\sum_{i \in [N]} \mathbb{E} \left[(\xi(\mathbf{S}_{t+1}(i)) - \mathbb{E} [\xi(\mathbf{S}_{t+1}(i))])^4 \right] + 6 \sum_{1 \leq i < j \leq N} \text{Var} [\xi(\mathbf{S}_{t+1}(i))] \text{Var} [\xi(\mathbf{S}_{t+1}(j))] \right) \end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{N^4} \left(N(\xi_{\max} - \xi_{\min})^4 + 6 \frac{N(N-1)}{2} (\xi_{\max} - \xi_{\min})^4 \right) \\
&\leq \frac{3(\xi_{\max} - \xi_{\min})^4}{N^2}.
\end{aligned} \tag{145}$$

Substituting the calculations in (144) and (145) back to (140), we get

$$\begin{aligned}
&\mathbb{P} \left[(X_{t+1}([N])\xi^\top - \mathbb{E}[X_{t+1}([N])\xi^\top])^2 > \frac{\min_{s,a,s'} P(s, a, s')}{4|\mathbb{S}|N} \right] \\
&\geq \left(\frac{\min_{s,a,s'} P(s, a, s')}{2|\mathbb{S}|N} \right)^2 \frac{N^2}{12(\xi_{\max} - \xi_{\min})^4} \\
&= \frac{\min_{s,a,s'} P(s, a, s')^2}{47|\mathbb{S}|^2(\xi_{\max} - \xi_{\min})^4}.
\end{aligned} \tag{146}$$

Because under the Optimal Local Control, we have $\|\mathbb{E}[X_{t+1}([N]) | X_t, \mathbf{A}_t] - \mathbb{E}[X_{t+1}([N])\xi^\top | X_t]\| \leq 2/N$ and $\mathbb{E}[X_{t+1}([N]) | X_t] = X_t([N])\Phi$, we can rewrite the bound (146) to get

$$\mathbb{P} \left[|(X_{t+1}([N])\xi^\top - X_t([N])\Phi)\xi^\top| > \frac{\min_{s,a,s'} P(s, a, s')^{1/2}}{2|\mathbb{S}|^{1/2}\sqrt{N}} - \frac{2}{N} \right] \geq \frac{\min_{s,a,s'} P(s, a, s')^2}{47|\mathbb{S}|^2(\xi_{\max} - \xi_{\min})^4}.$$

Therefore,

$$\mathbb{E} \left[|(X_{t+1}([N])\xi^\top - X_t([N])\Phi)\xi^\top| \right] \geq \left(\frac{\min_{s,a,s'} P(s, a, s')^{1/2}}{2|\mathbb{S}|^{1/2}\sqrt{N}} - \frac{2}{N} \right) \cdot \frac{\min_{s,a,s'} P(s, a, s')^2}{47|\mathbb{S}|^2(\xi_{\max} - \xi_{\min})^4},$$

which implies (138) with K_{noise} being the coefficient of $1/\sqrt{N}$ in the last expression. \square

Next, we state the weaker version of Property (c) of regular unstable RBs.

Property 1 (Weaker version of Property (c) in Definition 1). There exists positive constants η' and K_{noise} with $\eta' \leq \min\{y^*(\tilde{s}, 0), y^*(\tilde{s}, 1)\} - 1/N$, such that for any large-enough N and any time step t , we have

$$\mathbb{E} \left[\|X_{t+1}([N]) - \mu^* - (X_t([N]) - \mu^*)\Phi\|_{U_+} | X_t, \mathbf{A}_t \right] \geq \frac{\lambda_+^{1/2} K_{\text{noise}}}{\sqrt{N}} \quad \text{if } \|X_t([N]) - \mu^*\|_1 \leq \eta',$$

where \mathbf{A}_t follow the Optimal Local Control (Subroutine 2), which is always feasible if $\|X_t([N]) - \mu^*\|_1 \leq \min\{y^*(\tilde{s}, 0), y^*(\tilde{s}, 1)\} - 1/N$.

Intuitively, Property 1 assumes that when $X_t([N])$ is close to μ^* , the state-count vector in the next step, $X_{t+1}([N])$, has enough randomness; or precisely, $X_{t+1}([N]) - \mu^* - \mathbb{E}[X_{t+1}([N]) - \mu^* | X_t, \mathbf{A}_t]$ has enough variances in the directions orthogonal to the kernel of U_+ .

If Property (c) of regular unstable RBs is replaced by Property 1, the whole proof of Theorem 2 goes through almost verbatim, except one change: In the proof of Lemma 15, in the argument below (126), we apply Property 1 rather than Property (c).

C.4 Proof of Corollary 1

Corollary 1. *There exists an RB satisfying Assumptions 1 and 2 and violating Assumption 3 such that $R^{\text{rel}} - R(\pi, \mathbf{S}_0) \geq \Omega\left(\frac{1}{\sqrt{N}}\right)$.*

Proof. We will verify that the example in Figure 1 of [HXCW23] is regular unstable.

In this example, the single-armed MDP $(\mathbb{S}, \mathbb{A}, P, r)$ is defined as below: $\mathbb{S} = \{1, 2, 3\}$; $\mathbb{A} = \{0, 1\}$; the transition kernel P is given by

$$P(\cdot, 0, \cdot) = \begin{bmatrix} 0.02232142 & 0.10229283 & 0.87538575 \\ 0.03426605 & 0.17175704 & 0.79397691 \\ 0.52324756 & 0.45523298 & 0.02151947 \end{bmatrix},$$

$$P(\cdot, 1, \cdot) = \begin{bmatrix} 0.14874601 & 0.30435809 & 0.54689589 \\ 0.56845754 & 0.41117331 & 0.02036915 \\ 0.25265570 & 0.27310439 & 0.4742399 \end{bmatrix};$$

The reward function r is defined as

$$r(\cdot, 0) = [0 \ 0 \ 0],$$

$$r(\cdot, 1) = [0.37401552 \ 0.11740814 \ 0.07866135].$$

Let $\alpha = 0.4$ in the budget constraint. The optimal solution of the LP-relaxation (LP) is:

$$y^*(\cdot, \cdot) = \begin{bmatrix} 0 & 0.29943 \\ 0.23768 & 0.10057 \\ 0.36232 & 0 \end{bmatrix},$$

where the first column corresponds to $y^*(s, 0)$ and the second column corresponds to $y^*(s, 1)$.

Since $P(s, a, s') > 0$ for all s, a, s' , $P_{\bar{\pi}^*}$ is an aperiodic unichain, so Assumption 1 holds. Also, from y^* , we can see that the state 2 is the unique neutral state, implying Assumption 2. Moreover, one can numerically verify that y^* is the unique solution to (LP), so Property (a) of Definition 1 is satisfied; the moduli of the eigenvalues of Φ are $\{1.133, 0, 0.059\}$, so Property (b) of Definition 1 is satisfied.

Finally, observe that in this example, $P(s, a, s') > 0$ for any $s, s' \in \mathbb{S}$ and $a \in \mathbb{A}$. Therefore, we invoke the sufficient condition Lemma 17 proved in Appendix C.3 to conclude that this example satisfies Property (c) of Definition 1. \square