Repeater-Based Quantum Communication Protocol: Maximizing Teleportation Fidelity with Minimal Entanglement

Arkaprabha Ghosal,^{1,2} Jatin Ghai,^{1,2} Tanmay Saha,^{1,2} Sibasish Ghosh,^{1,2} and Mir Alimuddin³

¹Optics & Quantum Information Group, The Institute of Mathematical Sciences, CIT Campus, Taramani, Chennai 600113, India

²Homi Bhabha National Institute, Training School Complex, Anushakti Nagar, Mumbai 400085, India

³Department of Physics of complex Systems, S. N. Bose National Center for Basic Sciences,

Block JD, Sector III, Salt Lake, Kolkata 700106, India.

Transmitting unknown quantum states to distant locations is crucial for distributed quantum information protocols. The seminal quantum teleportation scheme achieves this feat while requiring prior maximal entanglement between the sender and receiver. In scenarios with noisy entangled states, optimal teleportation fidelity characterizes the efficacy of transmitting the state, demanding the proper selection of local operations at the sender's and receiver's ends. The complexity escalates further in long-range communication setups, prompting the consideration of a repeater-based approach, which incorporates arrays of nodes with multiple segments to facilitate the efficient transmission of quantum information. The fidelity of the communication line gets degraded even if a single segment is affected by noise. In such cases, the general wisdom employs the standard entanglement swapping protocol involving maximally entangled states across the noiseless segments and applying maximally entangled basis measurement at the corresponding nodes to achieve optimal fidelity. In this Letter, we propose a more efficient protocol for a certain class of noisy states in any intermediary segment, achieving the same fidelity as the standard protocol while consuming less amount of entanglement. Our approach ensures enhanced teleportation fidelity even when the end-to-end state gets noisier, and thus promises efficient utility of quantum resources in repeater-based distributed quantum protocols.

Introduction.- In distributed communication scenarios, the successful implementation of most information processing tasks critically depends on the relentless transmission of signals across distant points [1-3]. However, in the domain of long-range communication, signal quality often deteriorates as the separation between sender and receiver increases. To address this challenge, one potential remedy involves a repeater-based approach, wherein the communication line is divided into multiple segments or short-range communication lines [4, 5]. Signals received from each segment are then replicated, amplified, and relayed to neighbouring stations (nodes) until they reach the intended endpoints. However, conventional signal replication is rendered unfeasible in quantum information processing due to the *no-cloning* principle [6]. Nonetheless, thanks to the advent of quantum teleportation protocols [7], the way has been paved for the development of a viable quantum repeater-based protocol through entanglement swapping [8–11]. Noteworthy, the protocol requires equipping each segment with maximally entangled states (MES) and performing maximally entangled measurements (MEM) at intermediary nodes to facilitate seamless teleportation from sender to receiver.

However, establishing maximal entanglement poses significant challenges, and due to the environmental noise introduction often results in noisy entangled states [12]. Addressing this challenge involves various strategies, one of kind which entails distilling maximally entangled state from many copies of the identical noisy entangled states through local operations and classical communication (LOCC) [4, 5, 13–18]. However, we are interested in a single-shot approach that involves directly utilizing a single copy of the noisy entangled state as closely as possible to the input one [19–28]. The degree to which a state is teleported close to the unknown input is usually char-

acterized by the optimal teleportation fidelity (OTF). While the optimal teleportation fidelity value 'one' quantifies perfect teleportation, for noisy states, the value remains strictly below that. Notably, the complexity of finding the OTF is substantial and it escalates further in the quantum repeater scenario when more parties are involved [4, 5, 12]. For given noisy states, OTF explicitly relies on the appropriate choice of LOCC among sender, receiver, as well as multiple nodes. Here, we are interested in a scenario where one of the segments is faulty (called noisy-segment) and shares a noisy state ρ , while the other segments (*free-segments*) can freely choose arbitrary state to achieve the optimal end-to-end teleportation fidelity aimed at the best possible quantum information processing task. In this regard, standard entanglement swapping protocol (E-SWAP) can be implemented by sharing MES in free-segments and performing MEM in nodes to teleport the state ρ between sender and receiver, thereby achieving OTF between them. However, this prompts an intriguing question: can we go beyond the standard protocol to minimize the resources in free-segments and nodes to get an equal advantage?

This Letter precisely addresses the questions raised above. We demonstrate that, for a certain class of two-qubit noisy states, an equivalent optimal teleportation fidelity (OTF) can be achieved without physically teleporting the noisy state itself. The protocol utilizes non-maximal resources in freesegments and nodes, resulting in lower resource consumption compared to the standard E-SWAP protocol. Importantly, this advantage persists even if the number of nodes approaches infinity. Furthermore, in a given quantum repeater scenario, as the noisy segment shares a noisier state, the resource consumption decreases accordingly. These features not only expand our existing knowledge but also reveal exciting possibilities for innovative approaches and might ensure the advancement of several distributed tasks, from cryptography to the quantum internet [29–40], paving the way for more efficient

distributed quantum communication protocols.

Preliminaries.– The pure state of an individual quantum system (X) is represented by $|\psi\rangle_X \in \mathcal{H}_X$, where \mathcal{H}_X stands for a Hilbert space with dimension d_X . A mixed state is defined by $\rho_X \in \mathcal{D}(\mathcal{H}_X)$, here $\mathcal{D}(\mathcal{H}_X)$ denotes the set of trace one positive operator $(\mathcal{P}(\mathcal{H}_X))$, both acting on \mathcal{H}_X . A measurement on system X is denoted by $M_X \equiv \{\Pi_X^i\}_{i=1}^N$, where $\Pi_X^i \in \mathcal{P}(\mathcal{H}_X)$ and $\sum_{i=1}^N \Pi_X^i = \mathbb{I}_X$. Any deterministic physical transformation $X \to Y$ is represented by a completely positive trace-preserving (CPTP) map $\Lambda_{X \to Y} : \mathcal{D}(\mathcal{H}_X) \mapsto \mathcal{D}(\mathcal{H}_Y)$ [41].

Let's consider a scenario where Sender (Alice) and Receiver (Bob) share an arbitrary state $\rho_{AB} \in \mathcal{H}_{AB}$. The joint Hilbert space of A and B, denoted as \mathcal{H}_{AB} , is the tensor product of their individual Hilbert spaces, $\mathcal{H}_A \otimes \mathcal{H}_B$. Alice's goal is to teleport an unknown quantum state $|\psi\rangle_X$ to Bob. The most general teleportation protocol can be delineated as a CPTP map, denoted by $\mathcal{T} := \{M_{XA}, \Lambda_{B \mapsto X'}\},\$ where X' represents the output state with $d_{X'} = d_X$. In detail, a general teleportation protocol entails an input measurement $M_{XA} \equiv \{\Pi_{XA}^i\}_{i=1}^N$ on the XA part of the composite system, followed by communication of the measurement outcomes through a noiseless classical channel. Based on this received information, Bob applies some suitable CPTP map $\Lambda_{B\to X'} \equiv {\Lambda^i}_{i=1}^N$ on his part B. This process yields an output state $\rho_{X'} = \mathcal{T}(\psi_X, \rho_{AB})$. The utility of a given teleportation protocol \mathcal{T} for a pre-shared state ρ_{AB} is characterized by calculating the average fidelity over the entire input space, defined as $f_{\mathcal{T}}(\rho_{AB}) = \int d\psi_X \,_X \langle \psi | \rho_{X'} | \psi \rangle_X$. Note that for a given state the teleportation fidelity $f_{\mathcal{T}}$ explicitly depends on the teleportation protocol \mathcal{T} . The optimal teleportation fidelity (OTF) is then defined as [19]:

$$f^{\star}(\rho_{AB}) = \max_{\mathcal{T}} f_{\mathcal{T}}(\rho_{AB}). \tag{1}$$

Another significant quantity of interest is the fully entangled fraction (FEF), which measures the overlap of an arbitrary $d \times d$ bipartite state ρ_{AB} with the nearest maximally entangled state [19, 21]. The FEF of ρ_{AB} is expressed as $F(\rho_{AB}) = \max_{|\phi\rangle} \langle \phi | \rho_{AB} | \phi \rangle$, where $|\phi\rangle$ represents an arbitrary $d \times d$ maximally entangled state. It's important to note that $F(\rho_{AB})$ does not serve as a direct measure of entanglement, as its value can increase under the set of LOCC operation (Λ_{LOCC}) [23, 42]. In contrast, a revised quantity, the optimal fully entangled fraction (OFEF), defined as:

$$F^{\star}(\rho_{AB}) = \max_{\Lambda \in \Lambda_{LOCC}} F(\Lambda(\rho_{AB})), \qquad (2)$$

which clearly retains its monotonicity under LOCC and qualifies as a *measure of entanglement* [12, 43]. Verstraete *et al* proposed a systematic method for determining $F^*(\rho_{AB})$ in a two-qubit system [43]. However, for an arbitrary state the challenge in determining the above quantity lies in finding a suitable LOCC operation, which parallels the hurdles one must encounter in finding the optimal teleportation fidelity. More explicitly, it has been established that for an arbitrary $d \times d$ state, these two optimal quantities are linked by the relation $f^*(\rho_{AB}) = (d F^*(\rho_{AB}) + 1)/(d + 1)$ [19]. In our subsequent studies, we often refer to both terms interchangeably, as they are equivalent by the above relation. Notably, in the case of pre-shared separable states or PPT entangled states [44] OTF and OFEF reach 2/(d + 1) and 1/d, respectively [13, 19, 27]. These bounds are commonly referred to as the classical bound, and for a certain state, if it crosses these lower bounds, the state proves to be useful for teleportation and can offer nontrivial contributions in quantum information processing. It's noteworthy that for an arbitrary two-qubit state ρ_{AB} , the OFEF is always upper-bounded by [43]

$$F^*(\rho_{AB}) \le \frac{1 + \mathcal{C}(\rho_{AB})}{2},\tag{3}$$

where $C(\rho_{AB})$ is the concurrence of the state ρ_{AB} [45]. Although, in the case of pure entangled states and some classes of mixed entangled states, the OFEF hits the above upper bound [43, 46], generally it is not true for an arbitrary state. This distinction underscores F^* and C as independent measures of entanglement [47, 48].



FIG. 1: (Color online) Quantum Repeater-Based Teleportation Scheme from Alice to Bob.- (a) Considering n + 1 segments (n nodes) between the sender and receiver, where each segment S_i shares a two-qubit entangled state ρ_i . The general strategy is to prepare a two-qubit state between Alice and Bob in such a way that it achieves the highest possible fully entangled fraction (F_{AB}^*) , optimized over LOCC. (b) In this scenario, one of the segments (S_i) shares a noisy state ρ . The objective is to establish $F_{AB}^* = F^*(\rho)$, assuming the other segments can share arbitrary entangled states. We aim to find a LOCC protocol that ultimately results in a two-qubit state σ between Alice and Bob with $F^*(\sigma) = F^*(\rho)$. This protocol leads to the optimal fully entangled fraction and saturates the upper bound.

Repeater-based teleportation protocol.– In the realm of long-range quantum communication, quantum repeater-based protocols stand out for their effectiveness in mitigating uncontrollable noise [5]. This approach entails dividing the communication line into multiple short-range segments, denoted as S_i , where instead of direct communication from the sender (Alice) to a receiver (Bob), quantum information is relayed through intermediary nodes N_i along these segments. Let's assume there are (n + 1) segments, where $\{S_1, S_i, S_{n+1}\}_{i=2}^n$ are directly controlled by $A - N_1$, $\{N_{i-1} - N_i\}_{i=2}^n$, and

3

 $N_n - B$ parties, respectively, as depicted in Figure 1(a). If individual segments S_i share some arbitrary state ρ_i , what is the optimal fidelity achievable between the end-to-end parties (AB)? While general studies are quite nontrivial, for two-qubit noises in the given repeater scenario, the upper bound of optimal fidelity can be expressed as follows [49]:

$$F_{AB}^{\star} \le \min\{F_S, F_C\}. \tag{4}$$

The bound $F_S = \min_i \{F^*(\rho_i)\}$ arises from F^* being an entanglement measure, where $F^{\star}(\rho_i)$ denotes the OFEF of the state ρ_i . Meanwhile, the bound $F_C = (1 + \prod_{i=1}^{n+1} C_i)/2$ is obtained using Eq. (3), where C_i is the concurrence of the state ρ_i . Interestingly, these two bounds are typically independent of each other, however, in some special cases, they boil down to being equal [49]. It's crucial to note that having separable states in any segment makes $F_{AB}^{\star} = 1/2$, while $F_{AB}^{\star} = 1$ requires maximal entanglement across every segment. However, when a single segment hosts a fixed noisy state ρ , the end-to-end OFEF becomes $F_{AB}^{\star} \leq F^{\star}(\rho)$. The upper bound can always be achievable by the E-SWAP protocol utilizing maximal resources in free-segments and nodes. Contrarily, our investigation aims to develop a more efficient LOCC protocol that consumes less entanglement compared to E-SWAP while delivering an equivalent advantage, *i.e.*, $F_{AB}^{\star} = F^{\star}(\rho)$ [see Figure 1(b)]. In this regard, earlier studies indicate that the entanglement swapping scheme emerges as the indispensable protocol to establish $C_{AB} = C(\rho)$ between end-to-end parties [50]. Here we focus on OFEF, an independent quantifier of entanglement with significant operational implications in quantum information processing. We begin with the following lemma which exhibits no-go in resource consumption, detailed calculations deferred to Appendix A.

Lemma 1. In a single-node scenario if segment S_1 shares a two-qubit fixed noisy state ρ_1 with the constraint $F^*(\rho_1) = (1+C_1)/2$, then entanglement swapping is considered to be a necessary protocol to establish $F^*_{AB} = F^*(\rho_1)$.

The above relation between OFEF and Concurrence typically arises from the property $\rho_1^{\Gamma} |\phi\rangle = -\frac{C_1}{2} |\phi\rangle$, where Γ denotes partial transposition and $|\phi\rangle$ represents a maximally entangled state [43]. Fixed noise, whether as a pure entangled state or a mixed state with a mixture of Bell states, satisfies this property and necessarily consumes one-*ebit* of entanglement in free segment S_2 . However, in our subsequent theorem, we present a nontrivial example where resource consumption is indeed lower than that of the E-SWAP protocol while achieving the same final goal.

Consider a two-dimensional subspace S in $\mathcal{H}_A \otimes \mathcal{H}_B$ for a two-qubit composite system, spanned by a product state $|P\rangle$ and its orthogonal entangled state $|\zeta(\delta)\rangle := \{\sqrt{\delta}, \sqrt{1-\delta}\}$, where $\delta \in [1/2, 1)$ is its Schmidt coefficient. We define a noisy state $\rho(p, \delta) = pP + (1 - p)\zeta(\delta) \in S$, where $P := |P\rangle\langle P|, \zeta(\delta) := |\zeta(\delta)\rangle\langle\zeta(\delta)|$, and $p \in$ [0, 1]. Now, let's introduce another pure entangled state $|\psi(\alpha)\rangle := \{\sqrt{\alpha}, \sqrt{1-\alpha}\}$, which will be utilized in the free segments. Additionally, we define a fine-grained two-qubit projective entangled measurement $M(\beta) \equiv \{|\phi_i\rangle\langle\phi_i|\}_{i=1}^4$, where $\{\sqrt{\beta}, \sqrt{1-\beta}\}$ are the Schmidt coefficients of all $|\phi_i\rangle$. Without loss of generality consider that $\alpha, \beta \in [1/2, 1)$.

Theorem 1. In a single-node scenario, suppose segment S_1 is considered to be a noisy-segment characterized by a mixed state $\rho(p, \delta)$. Meanwhile, the free segment S_2 shares an arbitrary pure state $|\psi(\alpha)\rangle$, and node N_1 performs a measurement $M(\beta)$. We observe that within a certain range of the parameters $\{p, \delta, \alpha, \beta\}$, it is indeed possible to establish $F_{AB}^* = F^*[\rho(p, \delta)]$ between the sender and receiver.

A detailed proof with calculations is deferred to Appendix B. It's worth highlighting that when $\alpha = \beta = 1/2$, this leads to the standard E-SWAP protocol, and then any arbitrary state shared in S_1 can be readily prepared between the sender and receiver. However, Theorem 1 suggests that for different choices of α and β , the exact state may not be preparable and would inherently be noisier than $\rho(p, \delta)$. Nevertheless, it remains feasible to achieve equal fidelity for some specific range of $\{p, \delta, \alpha, \beta\}$. For instance, if we choose non-maximal resources ($\alpha = 0.75, \beta = 0.60$) for $\delta = 0.5$ ($\delta = 0.7$), we get $F_{AB}^{\star} > 1/2$ for $0.515 \leq p < 1(0.493 \leq p < 1)$. This eventually implies that the proposed protocol consumes a lesser amount of resources than the standard protocol.

Given our understanding of the necessity of quantum repeater scenarios, it's vital to establish the equivalent result in an *n*-node scenario. In the subsequent corollary, we will demonstrate that the above studies can be readily extended when one of the end segments $(S_1 \text{ or } S_{n+1})$ is associated with the noisy state, which effectively boils down to a *single*-node scenario.

Corollary 1. Consider the free segments $\{S_i\}_{i=2}^{n+1}$, where S_i shares the state $|\psi_i\rangle \equiv \{\sqrt{\alpha_i}, \sqrt{1-\alpha_i}\}$ with $\alpha_i \ge 1/2$, $\forall i$. It is always possible to find noisy states $\rho(p, \delta) = pP + (1-p)\zeta(\delta)$ shared in the first segment S_1 , with non-vanishing region of $\{p, \delta\}$, such that $F_{AB}^* = F^*[\rho(p, \delta)]$ can be achieved between the sender and receiver.

Here, we make this study by considering maximal entanglement basis measurement ($\beta = 1/2$) at each node, as discussed in detail in Appendix C. A careful observation reveals that for a constant non-maximally entangled state ($\alpha_i = \alpha > 1/2$) shared in free segments, as the number of nodes increases, the selective region $\{p, \delta\}$ for which $F_{AB}^{\star} = F^{\star}[\rho(p, \delta)]$, makes ρ tending towards a separable state. However, in an arbitrary n-node scenario, the goal is often to transmit optimal quantum information through a fixed noisy state. In other words, regardless of the number of nodes, can one establish end-toend teleportation fidelity identical to the noisy state $\rho(p, \delta)$ for some constant noise parameters $\{p, \delta\}$? This requirement can be certainly met by the E-SWAP protocol for an arbitrary state. However, our study demonstrates that for a certain range of $\{p, \delta\}$, the E-SWAP protocol isn't necessary to achieve this goal; rather, it can be attained using strictly non-maximally entangled states in free segments. We evaluate the variance in resource consumption between the E-SWAP and our proposed protocol defined by $R_v = n(1-C)$. Our findings illustrate that variance in resource consumption (R_v) remains strictly positive for all values of n; it escalates at first and then finally saturates in some positive number with the number of nodes, as



FIG. 2: (Color online) (a) Variance in resource consumption R_v with the number of nodes n for given noisy state $\rho(p, \delta)$ with the parameters values p = 0.8 and $\delta = 0.5, 0.9$. (b) Variance in resource consumption with the noise parameter p for a given number of nodes n = 10 and $\delta = 0.5, 0.9$.

depicted in Figure 2a. In this context, the following observations hold considerable significance (for details see Appendix D).

Observation 1. As the number of nodes $n \to \infty$, the variance in resource consumption R_v tends toward a constant value for a given noisy state $\rho(p, \delta)$.

Observation 2. For a given number of nodes n, as the noisy state $\rho(p, \delta)$ tends towards a separable state, the variance in resource consumption R_v increases accordingly, as illustrated in Figure 2b.

So far, we have explored scenarios where noise is localized at one of the end segments. However, consider the segments $\{S_i\}_{i=1}^{n+1}$ share a composition of states $\{\{\psi(\alpha_i)\}_{i=1}^{i=m-1}, \rho_m, \{\psi(\alpha_i)\}_{i=m+1}^{i=n+1}\}$, where the m^{th} segment is noisy and shares a fixed noisy state ρ_m , while the other segments are considered free, each with an arbitrary pure entangled state $\psi(\alpha_i)$, with $\alpha_i \geq 1/2$. We demonstrate that this scenario effectively reduces to a *two*-node scenario for any value of $m \in \{2, \dots, n\}$ (see Appendix E). By setting the m^{th} segment as a reference, the left segments $\{S_i\}_{i=1}^{m-1}$ and the right segments $\{S_i\}_{i=m+1}^{i=n+1}$ can be composed into new pair of segments S_1 and S_3 , respectively, while segment S_2 will be the noisy one in the *two*-node scenario. Precisely, the segments $\{S_i\}_{i=1}^3$ will have a composition of $\{\psi(\alpha'_l), \rho_2, \psi(\alpha'_r)\}$, where $\psi(\alpha'_{l(r)})$ represents the effective pure entangled state at segment $S_1(S_3)$ and $\rho_2 = \rho_m$. In the subsequent theorem, we will derive the condition $F_{AB}^* = F^*(\rho_2)$, with a detailed proof deferred to Appendix E.

Theorem 2. Let's assume a two-node scenario, where the intermediary segment S_2 shares a noisy state $\rho(p, \delta) = p P + (1-p)\zeta(\delta) \in S$, while the free segments $(i \in \{1,3\})$ share pure entangled states $\psi(\alpha'_i)$. We establish a nontrivial LOCC protocol that achieves $F_{AB}^* = F^*[\rho(p, \delta)]$ when the condition $\alpha'_l \alpha'_r / (1-\alpha'_l)(1-\alpha'_r) \leq p^2 / \delta(1-\delta)(1-p)^2$ is satisfied.

In an *n*-node quantum repeater scenario, determining the position of a noisy segment that exhibits the narrowest (or broadest) range of $\{p, \delta\}$ according to Theorem 2 for a given noisy state explicitly depends on the distribution of $\{\alpha_i\}$ in the free segments. For instance, when all the free segments share

identical states (*i.e.*, $\alpha_i = \alpha$), the lowest value of variance in resource consumption R_v is achieved at m = n/2 for even n, and at $m \in \{(n-1)/2, (n+1)/2\}$ for odd n. Furthermore, as the noisy segment gradually transitions towards either end, R_v inevitably expands. In light of that, an intriguing observation emerges:

Observation 3. In an n-node scenario, assuming free segments share identical non-maximal pure entangled states the variance in resource consumption R_v experiences a nondecreasing trend as the noisy segment migrates from the midpoint towards either end of the repeater scenario.

A graphical plot is illustrated in Figure 3, with detailed calculations provided in Appendix F. Importantly, this observation underscores the intricate dynamics of noise that govern resource optimization in quantum repeater architectures.



FIG. 3: (Color online) Variance in resource consumption R_v with the position of the noisy node (m) for a given number of nodes n = 99 and for the noisy state $\rho(p, \delta)$ with the parameters values p = 0.8 and $\delta = 0.5, 0.9$.

Discussions.- In a quantum repeater scenario where multiple segments share noisy entangled states, effective longrange quantum communication depends on the optimal distribution of entanglement between end-to-end parties. However, finding the optimal protocol for sharing a useful entangled state is quite a complex task and depends majorly on the LOCC protocols among the parties. These protocols also vary depending on the specific tasks. For instance, while the distribution of concurrence has been investigated for multisegment scenarios [50], our work demonstrates that achieving optimal teleportation fidelity—a measure of average teleportation success-requires a different protocol. Effective entanglement distribution and its utility in teleportation are crucial for various quantum communication protocols [29-40]. If noise exists in a single segment, establishing optimal concurrence necessitates maximal entanglement in other segments and maximally entangled basis measurements at intermediary nodes. This forms an effective entanglement swapping protocol that prepares an identical noisy state between the distant sender and receiver, corresponding to the noisy segment. In contrast, for optimal teleportation fidelity, we observe that non-maximally entangled states and non-maximal basis measurements can readily establish the same amount of teleportation fidelity, even if the end-to-end state becomes

noisier. Thus, our protocol establishes the optimal fully entangled fraction and saturates the upper bound while consuming lesser entanglement resources in the given quantum repeater scenario. However, in the presence of noise across multiple segments, it remains uncertain whether the optimal teleportation fidelity can saturate the fidelity of any one of the single noisy states ρ_i shared across different segments, *i.e.*, $F_{AB}^{\star} = \min_{i} \{F^{\star}(\rho_{i})\}$. Interestingly, we found an example where a noisy state ρ , when shared in a single segment, allows $F_{AB}^{\star} = F^{\star}(\rho)$. However, if two segments are affected by this same noisy state, $F_{AB}^{\star} < F^{\star}(\rho)$ (see Appendix G). For multiple noisy segments, distillation-based repeater protocols have been extensively explored [4, 5]. Nonetheless, our focus is on evaluating average teleportation fidelity using a single copy of noisy states for unknown quantum inputs, unlike distillation protocols that first distill to a maximally entangled state

- M. M. Wilde, *Quantum information theory* (Cambridge university press, 2013).
- [2] D. C. Marinescu and G. M. Marinescu, Chapter 3 classical and quantum information theory, in *Classical and Quantum Information*, edited by D. C. Marinescu and G. M. Marinescu (Academic Press, Boston, 2012) pp. 221–344.
- [3] S. Khatri and M. M. Wilde, Principles of quantum communication theory: A modern approach (2024), arXiv:2011.04672.
- [4] W. Dür, H.-J. Briegel, J. I. Cirac, and P. Zoller, Quantum repeaters based on entanglement purification, Phys. Rev. A 59, 169 (1999).
- [5] K. Azuma, S. E. Economou, D. Elkouss, P. Hilaire, L. Jiang, H.-K. Lo, and I. Tzitrin, Quantum repeaters: From quantum networks to the quantum internet, Rev. Mod. Phys. 95, 045006 (2023).
- [6] W. K. Wootters and W. H. Zurek, A single quantum cannot be cloned, Nature 299, 802 (1982).
- [7] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, Teleporting an unknown quantum state via dual classical and einstein-podolsky-rosen channels, Phys. Rev. Lett. 70, 1895 (1993).
- [8] M. Żukowski, A. Zeilinger, M. A. Horne, and A. K. Ekert, "event-ready-detectors" bell experiment via entanglement swapping, Phys. Rev. Lett. 71, 4287 (1993).
- [9] J.-W. Pan, D. Bouwmeester, H. Weinfurter, and A. Zeilinger, Experimental entanglement swapping: Entangling photons that never interacted, Phys. Rev. Lett. 80, 3891 (1998).
- [10] J. Bouda and V. Buzek, Entanglement swapping between multiqudit systems, Journal of Physics A: Mathematical and General 34, 4301 (2001).
- [11] V. Bayrakci and F. Ozaydin, Quantum zeno repeaters, Scientific Reports 12, 10.1038/s41598-022-19170-z (2022).
- [12] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Quantum entanglement, Rev. Mod. Phys. 81, 865 (2009).
- [13] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, Mixed-state entanglement and quantum error correction, Phys. Rev. A 54, 3824 (1996).
- [14] C. H. Bennett, G. Brassard, S. Popescu, B. Schumacher, J. A. Smolin, and W. K. Wootters, Purification of noisy entanglement and faithful teleportation via noisy channels, Phys. Rev. Lett. 76, 722 (1996).
- [15] D. Deutsch, A. Ekert, R. Jozsa, C. Macchiavello, S. Popescu,

5

before teleportation. Interestingly, our findings suggest that in scenarios where multiple segments share noisy states and distillation is technically impossible in some segments, producing non-maximally entangled states in the remaining segments can be more efficient. This approach effectively requires less entanglement consumption in the quantum repeaters.

ACKNOWLEDGMENTS

We acknowledge Dr. Manik Banik for many fruitful discussions and suggestions. MA acknowledges funding from the National Mission in Interdisciplinary Cyber-Physical systems from the Department of Science and Technology through the I-HUB Quantum Technology Foundation (Grant no: I-HUB/PDF/2021-22/008).

and A. Sanpera, Quantum privacy amplification and the security of quantum cryptography over noisy channels, Phys. Rev. Lett. **77**, 2818 (1996).

- [16] H.-J. Briegel, W. Dür, J. I. Cirac, and P. Zoller, Quantum repeaters: The role of imperfect local operations in quantum communication, Phys. Rev. Lett. 81, 5932 (1998).
- [17] M. Razavi, M. Piani, and N. Lütkenhaus, Quantum repeaters with imperfect memories: Cost and scalability, Phys. Rev. A 80, 032301 (2009).
- [18] S. Broadfoot, U. Dorner, and D. Jaksch, Singlet generation in mixed-state quantum networks, Phys. Rev. A 81, 042316 (2010).
- [19] M. Horodecki, P. Horodecki, and R. Horodecki, General teleportation channel, singlet fraction, and quasidistillation, Phys. Rev. A 60, 1888 (1999).
- [20] S. Bose, V. Vedral, and P. L. Knight, Purification via entanglement swapping and conserved entanglement, Phys. Rev. A 60, 194 (1999).
- [21] P. Badziag, M. Horodecki, P. Horodecki, and R. Horodecki, Local environment can enhance fidelity of quantum teleportation, Phys. Rev. A 62, 012311 (2000).
- [22] F. Verstraete, J. Dehaene, and B. DeMoor, Local filtering operations on two qubits, Phys. Rev. A 64, 010101 (2001).
- [23] F. Verstraete and H. Verschelde, Optimal teleportation with a mixed state of two qubits, Phys. Rev. Lett. 90, 097901 (2003).
- [24] R. Pal, S. Bandyopadhyay, and S. Ghosh, Entanglement sharing through noisy qubit channels: One-shot optimal singlet fraction, Phys. Rev. A 90, 052304 (2014).
- [25] Sudha, H. S. Karthik, R. Pal, K. S. Akhilesh, S. Ghosh, K. S. Mallesh, and A. R. Usha Devi, Canonical forms of two-qubit states under local operations, Phys. Rev. A 102, 052419 (2020).
- [26] V. Siddhu and J. Smolin, Optimal one-shot entanglement sharing, Phys. Rev. A 108, 032617 (2023).
- [27] E. Chitambar and F. Leditzky, On the duality of teleportation and dense coding, IEEE Transactions on Information Theory 70, 3529 (2024).
- [28] S. Hermans, M. Pompili, H. Beukers, S. Baier, J. Borregaard, and R. Hanson, Qubit teleportation between non-neighbouring nodes in a quantum network, Nature 605, 663 (2022).
- [29] C. H. Bennett and G. Brassard, Quantum cryptography: Public key distribution and coin tossing, Theoretical Computer Science 560, 7 (2014), theoretical Aspects of Quantum Cryptogra-

phy-celebrating 30 years of BB84.

- [30] A. K. Ekert, Quantum cryptography based on bell's theorem, Phys. Rev. Lett. 67, 661 (1991).
- [31] F. Xu, X. Ma, Q. Zhang, H.-K. Lo, and J.-W. Pan, Secure quantum key distribution with realistic devices, Rev. Mod. Phys. 92, 025002 (2020).
- [32] M. Curty, K. Azuma, and H.-K. Lo, Simple security proof of twin-field type quantum key distribution protocol, npj Quantum Information 5, 10.1038/s41534-019-0175-6 (2019).
- [33] J.-Y. Li, X.-X. Fang, T. Zhang, G. N. M. Tabia, H. Lu, and Y.-C. Liang, Activating hidden teleportation power: Theory and experiment, Phys. Rev. Res. 3, 023045 (2021).
- [34] V. Scarani, H. Bechmann-Pasquinucci, N. J. Cerf, M. Dušek, N. Lütkenhaus, and M. Peev, The security of practical quantum key distribution, Rev. Mod. Phys. 81, 1301 (2009).
- [35] W. Kłobus, W. Laskowski, M. Markiewicz, and A. Grudka, Nonlocality activation in entanglement-swapping chains, Phys. Rev. A 86, 020302 (2012).
- [36] X. Meng, J. Gao, and S. Havlin, Concurrence percolation in quantum networks, Phys. Rev. Lett. **126**, 170501 (2021).
- [37] A. Acín, J. I. Cirac, and M. Lewenstein, Entanglement percolation in quantum networks, Nature Physics 3, 256 (2007).
- [38] M. Cuquet and J. Calsamiglia, Entanglement percolation in quantum complex networks, Phys. Rev. Lett. 103, 240503 (2009).
- [39] K. Azuma, H. Takeda, M. Koashi, and N. Imoto, Quantum repeaters and computation by a single module: Remote nondestructive parity measurement, Phys. Rev. A 85, 062309 (2012).
- [40] S. Wehner, D. Elkouss, and R. Hanson, Quantum internet: A vision for the road ahead, Science 362, eaam9288 (2018), https://www.science.org/doi/pdf/10.1126/science.aam9288.
- [41] M. A. Nielsen and I. L. Chuang, *Quantum computation and quantum information* (Cambridge university press, 2010).
- [42] J. Řeháček, Z. Hradil, J. Fiurášek, and i. c. v. Brukner, Design-

ing optimum completely positive maps for quantum teleportation, Phys. Rev. A **64**, 060301 (2001).

- [43] F. Verstraete and H. Verschelde, Fidelity of mixed states of two qubits, Phys. Rev. A 66, 022307 (2002).
- [44] M. Horodecki, P. Horodecki, and R. Horodecki, Separability of mixed states: necessary and sufficient conditions, Physics Letters A 223, 1 (1996).
- [45] W. K. Wootters, Entanglement of formation of an arbitrary state of two qubits, Phys. Rev. Lett. 80, 2245 (1998).
- [46] A. Ghosal, D. Das, S. Roy, and S. Bandyopadhyay, Optimal two-qubit states for quantum teleportation vis-à-vis state properties, Phys. Rev. A 101, 012304 (2020).
- [47] S. Bandyopadhyay and A. Ghosh, Optimal fidelity for a quantum channel may be attained by nonmaximally entangled states, Phys. Rev. A 86, 020304 (2012).
- [48] R. Pal and S. Bandyopadhyay, Entanglement sharing via qudit channels: Nonmaximally entangled states may be necessary for one-shot optimal singlet fraction and negativity, Phys. Rev. A 97, 032322 (2018).
- [49] A. Ghosal, J. Ghai, T. Saha, M. Alimuddin, and S. Ghosh, Optimal quantum teleportation of collaboration (2024), arXiv:2401.17201 [quant-ph].
- [50] G. Gour and B. C. Sanders, Remote preparation and distribution of bipartite entangled states, Phys. Rev. Lett. 93, 260501 (2004).
- [51] R. Horodecki, P. Horodecki, and M. Horodecki, Violating bell inequality by mixed spin-12 states: necessary and sufficient condition, Physics Letters A 200, 340 (1995).
- [52] R. Horodecki and M. Horodecki, Information-theoretic aspects of inseparability of mixed states, Phys. Rev. A 54, 1838 (1996).
- [53] R. Horodecki, M. Horodecki, and P. Horodecki, Teleportation, bell's inequalities and inseparability, Physics Letters A 222, 21 (1996).
- [54] J. Modławska and A. Grudka, Increasing singlet fraction with entanglement swapping, Phys. Rev. A 78, 032321 (2008).

Appendix A: Proof of Lemma 1

Proof. Let the segment S_1 shares a noisy two-qubit state ρ_1 with concurrence C_1 and *FEF*

$$F(\rho_1) = \frac{1 + \mathcal{C}_1}{2},\tag{A1}$$

respectively. Clearly, Eq. (A1) implies that the optimal FEF value of ρ_1 , *i.e.*, $F^*(\rho_1)$ is equal with $F(\rho_1)$. Any two-qubit entangled state satisfying Eq. (A1) must hold the property [43]

$$\rho_1^{\Gamma} \left| \phi \right\rangle = \lambda_{min} \left| \phi \right\rangle,\tag{A2}$$

where Γ is the partial transposition in computational basis, and $|\phi\rangle$ is a maximally entangled state which is also the eigenvector of the smallest eigenvalue λ_{min} . Note that if ρ_1 is entangled then λ_{min} must be negative and can also be expressed as $\lambda_{min} = -\frac{N(\rho_1)}{2}$, where $N(\rho_1)$ is the negativity of ρ_1 [43]. However, the negativity of any two-qubit state ρ_1 satisfying Eq. (A2) is equal with its concurrence $C(\rho_1) = C_1$ [43, 46]. Hence, without any loss of generality one can express $\lambda_{min} = -\frac{C_1}{2}$. Under local unitary operation $U \otimes V$, any two-qubit state satisfying Eq. (A1) can be transformed to an unique canonical state [21, 46] with its Hilbert-Schmidt form,

$$\tilde{\rho}_{1} = (U \otimes V) \ \rho_{1} \ (U^{\dagger} \otimes V^{\dagger})$$

$$= \frac{1}{4} \left[\mathbb{I} \otimes \mathbb{I} + (\mathbf{r} \cdot \sigma) \otimes \mathbb{I} - \mathbb{I} \otimes (\mathbf{r} \cdot \sigma) - \sum_{i=1}^{3} |\tau_{i}| \ (\sigma_{i} \otimes \sigma_{i}) \right], \tag{A3}$$

where \mathbb{I} is identity operation on a qubit state and $\{\sigma_i\}$ are the Pauli matrices. The matrix representation in Eq. (A3) is the Hilbert-Schmidt representation [51–53], where $\mathbf{r} = \{r_1, r_2, r_3\}$ is the local vector of $\tilde{\rho}_1$ such that $r_i = Tr[\tilde{\rho}_1(\sigma_i \otimes \mathbb{I})] = Tr[\tilde{\rho}_1(\mathbb{I} \otimes \sigma_i)]$ for i = 1, 2, 3. The parameter τ_i in Eq. (A3) is the correlation matrix element such that $\tau_i = [\tilde{\rho}_1(\sigma_i \otimes \sigma_i)]$ for i = 1, 2, 3.

It is now trivial to show that $F(\rho_1) = F(\tilde{\rho}_1) = F^*(\rho_1)$ holds because of the local unitary equivalence between $\tilde{\rho}_1$ and ρ_1 . One can always show that Eq. (A1) holds for any two-qubit state *iff*

$$\tilde{\rho}_{1}^{\Gamma} \left| \Phi_{0} \right\rangle = -\frac{\mathcal{C}_{1}}{2} \left| \Phi_{0} \right\rangle \tag{A4}$$

is satisfied (see Lemma 4 in Ref. [46]), where $\sqrt{2} |\Phi_0\rangle = |00\rangle + |11\rangle$.

Now let us suppose the free segment S_2 shares a two-qubit state ρ_2 with concurrence value C_2 . From Ref. [50] one can always argue that no three-party trace-preserving LOCC between $A - N_1 - B$ can prepare a two-qubit state between Alice-Bob with concurrence greater than C_1C_2 . Hence, from Ref. [50] and [43] one can conclude that the optimal FEF between Alice-Bob is upper bounded as

$$F_{AB}^{\star} \le \frac{1 + \mathcal{C}_1 \mathcal{C}_2}{2} \le \frac{1 + \mathcal{C}_1}{2}.$$
(A5)

Hence, from Eq. (A5) it is clear that $F_{AB}^{\star} < F^{\star}(\rho_1)$ if $C_2 < 1$. Therefore, the equality in Eq. (A5) will hold *iff* we consider $C_2 = 1$ which is a maximally entangled state shared in S_2 . This completes the proof of Lemma 1.

Appendix B: Proof of Theorem 1

Proof. Let segment S_1 shares a rank two mixed state $\rho_1 \in S$, where S is a two-dimensional subspace in two-qubit space spanned by a product state $|P\rangle$ and its orthogonal entangled state $|\zeta(\delta)\rangle$. Without loss of generality we could take $\rho_1 = \rho(p, \delta) = pP + (1-p)\zeta(\delta)$, where $\zeta(\delta) := |\zeta(\delta)\rangle \langle \zeta(\delta)|$ with $|\zeta(\delta)\rangle = \sqrt{\delta} |00\rangle + \sqrt{1-\delta} |11\rangle$, and $P := |P\rangle \langle P| = |01\rangle \langle 01|$. The optimal fully entangled fraction (OFEF) of the preshared state ρ_1 can be computed by solving a semi-definite program [23, 47, 54] as

$$F^{\star}(\rho_{1}) = \frac{1}{2}(1 + \mathcal{C}(\rho_{1}) - p), \quad \text{if } \mathcal{C}(\rho_{1}) > 2p,$$

$$= \frac{1}{2}\left(1 + \frac{C(\rho_{1})^{2}}{4p}\right), \quad \text{if } \mathcal{C}(\rho_{1}) \le 2p,$$
(B1)

where $C(\rho_1) = 2\sqrt{\delta(1-\delta)}(1-p)$ is the concurrence of ρ_1 . Furthermore from Eq. (B1) one can easily conclude that

$$F^{\star}(\rho_1) \le \frac{1 + \mathcal{C}(\rho_1)}{2}.$$
(B2)

We now assume a scenario, where the free segment S_2 has a preshared pure two-qubit entangled state, say, $|\psi_2\rangle = \sqrt{\alpha} |00\rangle + \sqrt{1-\alpha} |11\rangle$, such that $\frac{1}{2} \le \alpha < 1$. Since we want to establish end-to-end OFEF $F_{AB}^{\star} = F^{\star}(\rho_1)$, prerequisite entanglement in the free segment S_2 has to maintain the following constraint

$$F^{\star}(\rho_1) \le F^{\star}(\psi_2) = \frac{1 + \mathcal{C}(\psi_2)}{2}.$$

The joint state $(A - N_1 - B)$ shared by two segments S_1 and S_2 can thus be written as:

$$\eta_{AN_{1}B} = \eta_{12} = \rho_{1} \otimes |\psi_{2}\rangle \langle \psi_{2}|$$

= $\left(p |01\rangle \langle 01| + (1-p) |\zeta(\delta)\rangle \langle \zeta(\delta)|\right) \otimes |\psi_{2}\rangle \langle \psi_{2}|.$ (B3)

Now we will show that under some specific range of state parameters of ρ_1 and $|\psi_2\rangle$, there exist deterministic LOCC protocols for which one obtains

$$F_{AB}^{\star} = F^{\star}(\rho_1). \tag{B4}$$

$$\begin{aligned} |\phi_0\rangle &= \sqrt{\beta} |00\rangle + \sqrt{(1-\beta)} |11\rangle, \quad |\phi_1\rangle &= \sqrt{(1-\beta)} |00\rangle - \sqrt{\beta} |11\rangle, \\ |\phi_2\rangle &= \sqrt{\beta} |01\rangle + \sqrt{(1-\beta)} |10\rangle, \quad |\phi_3\rangle &= \sqrt{(1-\beta)} |01\rangle - \sqrt{\beta} |10\rangle, \end{aligned}$$
(B5)

where $\frac{1}{2} \leq \beta < 1$. Post measurement state between Alice and Bob, corresponding to the projector $\{ |\phi_i\rangle \langle \phi_i | \}$, is defined as,

$$\rho_{AB,\phi_i} = \frac{1}{p_i} Tr_{N_1} \left(\left(\left| \phi_i \right\rangle_{N_1} \left\langle \phi_i \right| \otimes \mathbb{I}_{AB} \right) \eta_{12} \right), \tag{B6}$$

where $p_i = Tr(\rho_{AB,\phi_i})$ is the probability of outcome *i*. The explicit expressions of the post-measurement states for each of the measurement outcomes can thus be evaluated as

$$\rho_{AB,\phi_0} = \frac{\left\{\alpha\beta\delta + (1-\alpha)(1-\beta)(1-\delta)\right\}(1-p)}{(1-\alpha)(1-\beta) - (1-\alpha-\beta)\delta(1-p)} \left|\chi_0\right\rangle \left\langle\chi_0\right| + \frac{(1-\alpha)(1-\beta)p}{(1-\alpha)(1-\beta) - (1-\alpha-\beta)\delta(1-p)} \left|01\right\rangle \left\langle01\right|, \quad (B7)$$

$$\rho_{AB,\phi_1} = \frac{\left\{\alpha(1-\beta)\delta + (1-\alpha)\beta(1-\delta)\right\}(1-p)}{(1-\alpha)\beta + (\alpha-\beta)\delta(1-p)} \left|\chi_1\right\rangle \left\langle\chi_1\right| + \frac{(1-\alpha)\beta p}{(1-\alpha)\beta + (\alpha-\beta)\delta(1-p)} \left|01\right\rangle \left\langle01\right|,\tag{B8}$$

$$\rho_{AB,\phi_2} = \frac{\left\{ (1-\alpha)\beta\delta + \alpha(1-\beta)(1-\delta) \right\} (1-p)}{\alpha(1-\beta) - (\alpha-\beta)\delta(1-p)} \left| \chi_2 \right\rangle \left\langle \chi_2 \right| + \frac{\alpha(1-\beta)p}{\alpha(1-\beta) - (\alpha-\beta)\delta(1-p)} \left| 00 \right\rangle \left\langle 00 \right|, \tag{B9}$$

$$\rho_{AB,\phi_3} = \frac{\left\{ (1-\alpha)(1-\beta)\delta + \alpha\beta(1-\delta) \right\}(1-p)}{\alpha\beta + (1-\alpha-\beta)\delta(1-p)} \left| \chi_3 \right\rangle \left\langle \chi_3 \right| + \frac{\alpha\beta p}{\alpha\beta + (1-\alpha-\beta)\delta(1-p)} \left| 00 \right\rangle \left\langle 00 \right|, \tag{B10}$$

where $|\chi_i\rangle$ s are given by

$$|\chi_0\rangle = \frac{\sqrt{\alpha\beta\delta} |00\rangle + \sqrt{(1-\alpha)(1-\beta)(1-\delta)} |11\rangle}{\sqrt{\alpha\beta\delta + (1-\alpha)(1-\beta)(1-\delta)}},\tag{B11}$$

$$|\chi_1\rangle = \frac{\sqrt{\alpha(1-\beta)\delta}|00\rangle - \sqrt{(1-\alpha)\beta(1-\delta)}|11\rangle}{\sqrt{\alpha(1-\beta)\delta + (1-\alpha)\beta(1-\delta)}},$$
(B12)

$$|\chi_2\rangle = \frac{\sqrt{(1-\alpha)\beta\delta}|01\rangle + \sqrt{\alpha(1-\beta)(1-\delta)}|10\rangle}{\sqrt{(1-\alpha)\beta\delta + \alpha(1-\beta)(1-\delta)}},\tag{B13}$$

$$|\chi_3\rangle = \frac{\sqrt{(1-\alpha)(1-\beta)\delta}|01\rangle - \sqrt{\alpha\beta(1-\delta)}|10\rangle}{\sqrt{(1-\alpha)(1-\beta)\delta + \alpha\beta(1-\delta)}}.$$
(B14)

Now from the knowledge of $\{|\chi_i\rangle\}$, one can figure out OFEF of every ρ_{AB,ϕ_i} as

$$F^{\star}(\rho_{AB,\phi_{0}}) = \frac{(1-\alpha)\alpha\delta p(1-p)}{2\{1-\alpha-\delta(1-p)\}\{(1-\alpha)(1-\beta)-(1-\alpha-\beta)\delta(1-p)\}} + \sqrt{\frac{(1-\alpha)\alpha(1-\beta)\beta(1-\delta)\delta(1-p)^{2}}{\{(1-\alpha)(1-\beta)-(1-\alpha-\beta)\delta(1-p)\}^{2}}} + \frac{(1-\alpha-\delta)(1-p)}{2\{1-\alpha-\delta(1-p)\}},$$
if, $\alpha\beta\delta(1-\delta)(1-p)^{2} > (1-\alpha)(1-\beta)p^{2},$

$$F^{\star}(\rho_{AB,\phi_{1}}) = \frac{\{(1-\alpha)\beta + (\alpha-\beta)\delta\}(1-p)}{2\{(1-\alpha)\beta + (\alpha-\beta)\delta(1-p)\}} + \sqrt{\frac{(1-\alpha)\alpha(1-\beta)\beta(1-\delta)\delta(1-p)^{2}}{\{(1-\alpha)\beta + (\alpha-\beta)\delta(1-p)\}^{2}}},$$

if,
$$\alpha (1-\beta)\delta(1-\delta)(1-p)^2 > (1-\alpha)\beta p^2$$
,
(B15)

and,

$$F^{\star}(\rho_{AB,\phi_0}) = \frac{\left\{1 - \delta(1-p)\right\} \left\{(1-\alpha)(1-\beta)p + \alpha\beta\delta(1-p)\right\}}{2p\left\{(1-\alpha)(1-\beta) - (1-\alpha-\beta)\delta(1-p)\right\}}, \quad \text{if,} \quad \alpha\beta\delta(1-\delta)(1-p)^2 \le (1-\alpha)(1-\beta)p^2,$$

$$F^{\star}(\rho_{AB,\phi_{1}}) = \frac{\left\{1 - \delta(1-p)\right\}\left\{(1-\alpha)\beta p + \alpha(1-\beta)\delta(1-p)\right\}}{2p\left\{(1-\alpha)\beta + (\alpha-\beta)\delta(1-p)\right\}}, \quad \text{if,} \quad \alpha(1-\beta)\delta(1-\delta)(1-p)^{2} \le (1-\alpha)\beta p^{2}$$
(B16)

whereas,

$$F^{\star}(\rho_{AB,\phi_{2}}) = \frac{1}{2} - \frac{(1-\beta)p}{2\{(1-\beta) - (1-2\beta)\delta(1-p)\}} + \sqrt{\frac{(1-\beta)\beta(1-\delta)\delta(1-p)^{2}}{\{(1-\beta) - (1-2\beta)\delta(1-p)\}^{2}}},$$

if, $(1-\alpha)\beta\delta(1-\delta)(1-p)^{2} > \alpha(1-\beta)p^{2},$

$$F^{\star}(\rho_{AB,\phi_{3}}) = \sqrt{\frac{(1-\alpha)\alpha(1-\beta)\beta(1-\delta)\delta(1-p)^{2}}{\left\{\alpha\beta + (1-\alpha-\beta)\delta(1-p)\right\}^{2}}} + \frac{\left\{\alpha\beta - (1-\alpha-\beta)\delta\right\}(1-p)}{2\left\{\alpha\beta + (1-\alpha-\beta)\delta(1-p)\right\}},$$

if, $(1-\alpha)(1-\beta)\delta(1-\delta)(1-p)^{2} > \alpha\beta p^{2},$
and, (B17)

$$F^{\star}(\rho_{AB,\phi_2}) = \frac{\{1 - \delta(1-p)\}\{\alpha(1-\beta)p + (1-\alpha)\beta\delta(1-p)\}}{2p\{\alpha(1-\beta) - (\alpha-\beta)\delta(1-p)\}}, \quad \text{if,} \quad (1-\alpha)\beta\delta(1-\delta)(1-p)^2 \le \alpha(1-\beta)p^2,$$

$$F^{*}(\rho_{AB,\phi_{3}}) = \frac{\left\{1 - \delta(1-p)\right\} \left\{\alpha\beta p + (1-\alpha)(1-\beta)\delta(1-p)\right\}}{2p\left\{\alpha\beta + (1-\alpha-\beta)\delta(1-p)\right\}}, \quad \text{if,} \quad (1-\alpha)(1-\beta)\delta(1-\delta)(1-p)^{2} \le \alpha\beta p^{2}.$$
(B18)

Now let us consider a particular domain where all four inequalities,

$$\alpha\beta\delta(1-\delta)(1-p)^{2} \le (1-\alpha)(1-\beta)p^{2},$$
(B19)

$$\alpha(1-\beta)\delta(1-\delta)(1-p)^2 \le (1-\alpha)\beta p^2,\tag{B20}$$

$$(1-\alpha)\beta\delta(1-\delta)(1-p)^2 \le \alpha(1-\beta)p^2,$$
(B21)

$$(1-\alpha)(1-\beta)\delta(1-\delta)(1-p)^2 \le \alpha\beta p^2,$$
(B22)

are simultaneously satisfied. From these conditions, one can argue that the following inequality conditions should be satisfied,

$$0
or,
$$\frac{1}{3}
(B23)$$$$

It can be checked easily that this set of inequalities arises from Eqs. (B19 - B22). Hence Eqs. (B19 - B22) are the minimal constraints required for OFEF. So while all these conditions hold, from Eqs. (B16) and (B18) one can easily find out that the average OFEF, *i.e.*,

$$F_{AB}^{\star} = \sum_{i=0}^{3} p_i F^{\star}(\rho_{AB,\phi_i})$$

$$= \frac{1}{2} \left(1 + \delta(1-\delta) \frac{(1-p)^2}{p} \right) = F^*(\rho_1) = F^*(\rho(p,\delta)),$$
(B24)

which ends the proof.

A major point to be noted here is that the established OFEF between Alice and Bob in Eq. (B24) is hitting the LOCC upper bound 'min{ $F^*(\rho_1), F^*(\psi_2)$ }' which means Eq. (B23) must obey the inequality condition, $F^*(\rho_1) \leq F^*(\psi_2)$. This means that the following inequality must hold, *i.e.*,

$$\frac{1}{2}\left(1+\delta(1-\delta)\frac{(1-p)^2}{p}\right) \le F^{\star}(\psi_2) = \frac{1+2\sqrt{\alpha(1-\alpha)}}{2},\tag{B25}$$

if Eq. (B23) is satisfied. From Eq. (B25) one can find a domain of α as

$$\frac{1}{2} \le \alpha < \frac{p^2}{p^2 + \delta(1-\delta)(1-p)^2} < \frac{1}{2} \left[1 + \sqrt{1 - \delta^2(1-\delta)^2 \frac{(1-p)^4}{p^2}} \right],$$

which is consistent with Eq. (B23).

Appendix C: Proof of Corollary 1

Proof. Let's consider a quantum repeater set-up, composed of n nodes with n+1 segments between Alice and Bob. The segments $\{S_i\}_{i=2}^{n+1}$, each shares the state $|\psi_i\rangle = \sqrt{\alpha_i} |00\rangle + \sqrt{1-\alpha_i} |11\rangle$, having concurrence $C(\psi_i) = 2\sqrt{\alpha_i(1-\alpha_i)}$ and the segment S_1 shares a noisy state $\rho_1 = \rho(p, \delta) = pP + (1-p)\zeta(\delta)$, where $\zeta(\delta) := |\zeta(\delta)\rangle \langle \zeta(\delta)|$ with $|\zeta(\delta)\rangle = \sqrt{\delta} |00\rangle + \sqrt{1-\delta} |11\rangle$ and $P = |01\rangle \langle 01|$. The joint state shared by n+1 segments can thus be written as:

$$\eta_{AN_{1}\cdots N_{n}B} = \eta = \rho_{1} \otimes |\psi_{2}\rangle \langle \psi_{2}| \otimes |\psi_{3}\rangle \langle \psi_{3}| \otimes \dots \otimes |\psi_{n+1}\rangle \langle \psi_{n+1}|$$

$$= \left(p \left|01\right\rangle \langle 01| + (1-p) \left|\zeta(\delta)\right\rangle \langle \zeta(\delta)|\right) \otimes \left(\bigotimes_{i=2}^{n+1} |\psi_{i}\rangle \langle \psi_{i}|\right). \tag{C1}$$

It is known that given two non-maximally entangled states with concurrences, say C_1 and C_2 , we can deterministically prepare another non-maximally entangled state with concurrence C_1C_2 through the RPBES protocol [50]. The state prepared will have the form

$$|\tau\rangle = \sqrt{\tau} |00\rangle + \sqrt{1 - \tau} |11\rangle, \qquad (C2)$$

where $\tau = \frac{1 + \sqrt{1 - C_1^2 C_2^2}}{2}$ is the Schmidt coefficients such that $\tau \ge \frac{1}{2}$ and the concurrence of $|\tau\rangle$ is $C(\tau) = C_1 C_2 \le 1$.

First, we apply this RPBES protocol at nodes N_2, N_3, \dots, N_n for our given chain. The resultant state between node N_1 and Bob is

$$|\Psi_n\rangle = \sqrt{\alpha'_n} |00\rangle + \sqrt{1 - \alpha'_n} |11\rangle, \qquad (C3)$$

where $\alpha'_n = \frac{1 + \sqrt{1 - C^2(\Psi_n)}}{2}$, and $C(\Psi_n) = \prod_{i=2}^{n+1} C(\psi_i)$. Now we are in a single node scenario where we have ρ_1 and $|\psi\rangle_{N_1B} \langle \psi| = |\Psi_n\rangle \langle \Psi_n|$. If we do only a Bell measurement at node N_1 ($\beta = \frac{1}{2}$) then from Appendix B we know that to obtain optimal fidelity the noise parameters p and δ of the noisy state ρ_1 , and the Schmidt coefficient of the state $|\Psi_n\rangle$ have to satisfy the following inequalities:

$$0
or,
$$\frac{1}{3}
(C4)$$$$

Thus from the third inequality we have

$$\frac{1+\sqrt{1-\mathcal{C}^2(\Psi_n)}}{2} \le \frac{p^2}{p^2+\delta(1-\delta)(1-p)^2}.$$
(C5)

Hence we see that for non-maximally entangled states with given concurrences distributed among n segments $\{S_i\}_{i=2}^{n+1}$, there exists a certain range of the noise parameters p and δ of the noisy state ρ_1 in segment S_1 for which $F_{AB}^{\star} = F^{\star}(\rho_1) = F^{\star}(\rho(p, \delta))$.

Appendix D: Resource consumption in the proposed LOCC protocol and comparison with E-SWAP

If all the segments $\{S_i\}_{i=2}^{n+1}$ share the same non-maximally entangled state, *i.e.*, $\alpha_i = \alpha$ for all $i \in [2, n+1]$ where *i* is a natural number then $\mathcal{C}(\Psi_n) = \mathcal{C}^n$, where $\mathcal{C} = 2\sqrt{\alpha(1-\alpha)}$. Substituting it in inequality (C5) we have

$$\frac{1+\sqrt{1-\mathcal{C}^{2n}}}{2} \le \frac{p^2}{p^2+\delta(1-\delta)(1-p)^2}.$$
(D1)

With some basic manipulations, we arrive at the upper bound on the number of nodes n for the given values of the noise parameters $\{p, \delta\}$ and the concurrence C of the states in segments $\{S_i\}_{i=2}^{n+1}$ as

$$n \leq \frac{\left|\log\left(1 - \left(\frac{p^2 - \delta(1 - \delta)(1 - p)^2}{p^2 + \delta(1 - \delta)(1 - p)^2}\right)^2\right)\right|}{2|\log(\mathcal{C})|}.$$
(D2)

Let us define a quantity variance in resource consumption:

$$R_v = n(1 - \mathcal{C}). \tag{D3}$$

We know that one-*ebit* resource (a maximally entangled state with concurrence C = 1) is required per free segment in the E-SWAP protocol. Our proposed protocol, which utilizes non-maximally entangled state with concurrence C < 1, will suffice to satisfy the condition $F_{AB}^{\star} = F^{\star}(\rho(p, \delta))$. Thus, R_v quantifies how much less resource is consumed in our proposed protocol compared to the E-SWAP protocol to achieve optimal fidelity in an *n*-node repeater scenario. In terms of the noise parameter and number of nodes, it is upper-bounded as

$$R_{v} \le n - n \left(1 - \left(\frac{p^{2} - \delta(1 - \delta)(1 - p)^{2}}{p^{2} + \delta(1 - \delta)(1 - p)^{2}} \right)^{2} \right)^{\frac{1}{2n}}.$$
 (D4)

As the number of nodes increases the variance in resource consumption (R_v) saturates. In fact, for $n \to \infty$, we have

$$R_v \to -\frac{1}{2} \log \left(-\frac{\delta(1-\delta)(1-p)^2 p^2}{\{\delta(1-\delta)(1-p)^2 + p^2\}^2} \right) - \log(2).$$
(D5)

Appendix E: Proof of Theorem 2

Proof. We consider a repeater scenario of n + 1 number of segments (*n* number of nodes) between Alice and Bob such that an intermediate segment, S_m , shares a noisy state of the form $\rho_m = \rho(p, \delta) = pP + (1 - p)\zeta(\delta)$, where $\zeta(\delta) := |\zeta(\delta)\rangle \langle \zeta(\delta)|$ with $|\zeta(\delta)\rangle = \sqrt{\delta} |00\rangle + \sqrt{1 - \delta} |11\rangle$, and $P = |01\rangle \langle 01|$. All the other segments $\{S_i\}_{i=1}^{n+1}$, except for i = m, each shares a pure state $|\psi_i\rangle = \sqrt{\alpha_i} |00\rangle + \sqrt{1 - \alpha_i} |11\rangle$ with concurrence $C(\psi_i)$. So the initial state of the composite network can be expressed as

$$\rho = |\eta\rangle_{1(m-1)} \langle \eta| \otimes \rho_m \otimes |\eta'\rangle_{(m+1)(n+1)} \langle \eta'|,$$

where we define

$$\left|\eta\right\rangle_{1(m-1)} = \bigotimes_{i=1}^{m-1} \left|\psi_i\right\rangle,$$

and,
$$|\eta'\rangle_{(m+1)(n+1)} = \bigotimes_{i=m+1}^{n+1} |\psi_i\rangle$$
.

The strategy is simple. We first apply the RPBES protocol [50] to the first m-1 segments $\{S_i\}_{i=1}^{m-1}$. The resultant state will become

$$|\Psi_l\rangle = \sqrt{\alpha_l'} |00\rangle + \sqrt{1 - \alpha_l'} |11\rangle, \qquad (E1)$$

where $\alpha'_l = \frac{1 + \sqrt{1 - C^2(\Psi_l)}}{2}$ and $C(\Psi_l) = \prod_{i=1}^{m-1} C(\psi_i)$. Similarly, we apply the RPBES protocol on the remaining segments ${S_i}_{i=m+1}^{n+1}$ resulting in the state

$$|\Psi_r\rangle = \sqrt{\alpha'_r} |00\rangle + \sqrt{1 - \alpha'_r} |11\rangle, \qquad (E2)$$

where $\alpha'_r = \frac{1 + \sqrt{1 - C^2(\Psi_r)}}{2}$ and $C(\Psi_r) = \prod_{i=m+1}^{n+1} C(\psi_i)$. So, we are now effectively reduced to a two-node scenario $\{N_{m-1}, N_m\}$ with three segments $S_1(A - N_{m-1}), S_2(N_{m-1} - W_m)$ N_m), and $S_3(N_m - B)$, where the total state can be written as

$$\rho = |\Psi_l\rangle \langle \Psi_l | \otimes \rho_2 \otimes |\Psi_r\rangle \langle \Psi_r |$$

where segment S_1 shares a non maximally entangled state $|\Psi_l\rangle$, segment S_2 shares the noisy state $\rho_2 = \rho_m$, and segment S_3 shares another non maximally entangled state $|\Psi_r\rangle$. Let us first consider node N_m . We implement a measurement in the Bell basis there. Following the calculations in Appendix B, we will end up with the four post-measurement states $\rho_{N_{m-1}B,\phi_i}$, where

 $i \in \{1, 2, 3, 4\}$, as in Eqs. (B7) to (B10) with $\beta = \frac{1}{2}$. Now, implement a Bell measurement at node N_{m-1} . Corresponding to each outcome, *i.e.*, $\rho_{N_{m-1}B,\phi_i}$, of the earlier Bell measurement at node N_m , there will be four more outcomes. So, in total, there will be sixteen states. The detailed calculations are as follows:

For the state $\rho_{N_{m-1}B,\phi_0}$ (as in Eq. (B7) with $\beta = \frac{1}{2}$), the outcomes corresponding to the second Bell measurement are given by

$$\rho_{AB,\phi_{0},\phi_{0}} = \frac{\left\{\alpha_{r}^{\prime}\alpha_{l}^{\prime}\delta + (1-\alpha_{r}^{\prime})(1-\alpha_{l}^{\prime})(1-\beta)\right\}(1-p)}{(1-\alpha_{r}^{\prime})(1-\alpha_{l}^{\prime}) - (1-\alpha_{r}^{\prime})(1-2\alpha_{l}^{\prime})p - (1-\alpha_{r}^{\prime}-\alpha_{l}^{\prime})\delta(1-p)} \left|\zeta_{0,0}\right\rangle\left\langle\zeta_{0,0}\right| \\ + \frac{(1-\alpha_{r}^{\prime})\alpha_{l}^{\prime}p}{(1-\alpha_{r}^{\prime})(1-\alpha_{l}^{\prime}) - (1-\alpha_{r}^{\prime})(1-2\alpha_{l}^{\prime})p - (1-\alpha_{r}^{\prime}-\alpha_{l}^{\prime})\delta(1-p)} \left|01\right\rangle\left\langle01\right|, \tag{E3}$$

$$\rho_{AB,\phi_{0},\phi_{1}} = \frac{\left\{\alpha_{r}^{\prime}\alpha_{l}^{\prime}\delta + (1-\alpha_{r}^{\prime})(1-\alpha_{l}^{\prime})(1-\delta)\right\}(1-p)}{(1-\alpha_{r}^{\prime})(1-\alpha_{l}^{\prime}) - (1-\alpha_{r}^{\prime})(1-2\alpha_{l}^{\prime})p - (1-\alpha_{r}^{\prime}-\alpha_{l}^{\prime})\delta(1-p)} \left|\zeta_{0,1}\right\rangle\left\langle\zeta_{0,1}\right| + \frac{(1-\alpha_{r}^{\prime})\alpha_{l}^{\prime}p}{(1-\alpha_{r}^{\prime})(1-\alpha_{l}^{\prime}) - (1-\alpha_{r}^{\prime})(1-2\alpha_{l}^{\prime})p - (1-\alpha_{r}^{\prime}-\alpha_{l}^{\prime})\delta(1-p)} \left|01\right\rangle\left\langle01\right|, \quad (E4)$$

$$\rho_{AB,\phi_{0},\phi_{2}} = \frac{\left\{ \alpha_{r}'(1-\alpha_{l}')\delta + (1-\alpha_{r}')\alpha_{l}'(1-\delta) \right\}(1-p)}{(1-\alpha_{r}')\alpha_{l}' + (1-\alpha_{r}')(1-2\alpha_{l}')p - (\alpha_{l}'-\alpha_{r}')\delta(1-p)} \left| \zeta_{0,2} \right\rangle \left\langle \zeta_{0,2} \right| \\ + \frac{(1-\alpha_{r}')(1-\alpha_{r}')(1-\alpha_{l}')p}{(1-\alpha_{r}')\alpha_{l}' + (1-\alpha_{r}')(1-2\alpha_{l}')p - (\alpha_{l}'-\alpha_{r}')\delta(1-p)} \left| 11 \right\rangle \left\langle 11 \right|,$$
(E5)

$$\rho_{AB,\phi_{0},\phi_{3}} = \frac{\left\{ \alpha_{r}'(1-\alpha_{l}')\delta + (1-\alpha_{r}')\alpha_{l}'(1-\delta) \right\}(1-p)}{(1-\alpha_{r}')\alpha_{l}' + (1-\alpha_{r}')(1-2\alpha_{l}')p - (\alpha_{l}'-\alpha_{r}')\delta(1-p)} \left| \zeta_{0,3} \right\rangle \left\langle \zeta_{0,3} \right| \\ + \frac{(1-\alpha_{r}')(1-\alpha_{r}')(1-\alpha_{l}')p}{(1-\alpha_{r}')\alpha_{l}' + (1-\alpha_{r}')(1-2\alpha_{l}')p - (\alpha_{l}'-\alpha_{r}')\delta(1-p)} \left| 11 \right\rangle \left\langle 11 \right|,$$
(E6)

where,

$$|\zeta_{0,0}\rangle = \frac{\sqrt{\alpha'_r \alpha'_l \delta} |00\rangle + \sqrt{(1 - \alpha'_r)(1 - \alpha'_l)(1 - \delta)} |11\rangle}{\sqrt{\alpha'_r \alpha'_l \delta + (1 - \alpha'_r)(1 - \alpha'_l)(1 - \delta)}},$$
(E7)

$$|\zeta_{0,1}\rangle = \frac{\sqrt{\alpha'_r \alpha'_l \delta} |00\rangle - \sqrt{(1 - \alpha'_r)(1 - \alpha'_l)(1 - \delta)} |11\rangle}{\sqrt{\alpha'_r \alpha'_l \delta + (1 - \alpha'_r)(1 - \alpha'_l)(1 - \delta)}},$$
(E8)

$$|\zeta_{0,2}\rangle = \frac{\sqrt{(1-\alpha_r')\alpha_l'(1-\delta)}|01\rangle + \sqrt{\alpha_r'(1-\alpha_l')\delta}|10\rangle}{\sqrt{(1-\alpha_r')\alpha_l'(1-\delta) + \alpha_r'(1-\alpha_l')\delta}},\tag{E9}$$

$$|\zeta_{0,3}\rangle = \frac{\sqrt{(1-\alpha_r')\alpha_l'(1-\delta)}|01\rangle - \sqrt{\alpha_r'(1-\alpha_l')\delta}|10\rangle}{\sqrt{(1-\alpha_r')\alpha_l'(1-\delta) + \alpha_r'(1-\alpha_l')\delta}}.$$
(E10)

Note that the states in Eqs. (E3 - E6) are of the same form as in Eqs. (B7 - B10). Following the similar calculations as in Appendix B, for finite resource consumption we get

$$\alpha'_{r}(1-\alpha'_{l})\delta(1-\delta)(1-p)^{2} \le (1-\alpha'_{r})\alpha'_{l}p^{2},$$
(E11)

and,

$$\alpha'_r \alpha'_l \delta(1-\delta)(1-p)^2 \le (1-\alpha'_r)(1-\alpha'_l)p^2.$$
 (E12)

For the state $\rho_{N_{m-1}B,\phi_1}$ (as in Eq. (B8) with $\beta = \frac{1}{2}$), the outcomes corresponding to the second Bell measurement are given by

$$\rho_{AB,\phi_{1},\phi_{0}} = \frac{\left\{\alpha_{r}^{\prime}\alpha_{l}^{\prime}\delta + (1-\alpha_{r}^{\prime})(1-\alpha_{l}^{\prime})(1-\delta)\right\}(1-p)}{(1-\alpha_{r}^{\prime})(1-\alpha_{l}^{\prime}) - (1-\alpha_{r}^{\prime})(1-2\alpha_{l}^{\prime})p - (1-\alpha_{r}^{\prime}-\alpha_{l}^{\prime})\delta(1-p)} \left|\zeta_{1,0}\right\rangle \left\langle\zeta_{1,0}\right| \\
+ \frac{(1-\alpha_{r}^{\prime})\alpha_{l}^{\prime}p}{(1-\alpha_{r}^{\prime})(1-\alpha_{l}^{\prime}) - (1-\alpha_{r}^{\prime})(1-2\alpha_{l}^{\prime})p - (1-\alpha_{r}^{\prime}-\alpha_{l}^{\prime})\delta(1-p)} \left|01\right\rangle \left\langle01\right|,$$
(E13)

$$\rho_{AB,\phi_{1},\phi_{1}} = \frac{\left\{\alpha_{r}'\alpha_{l}'\delta + (1-\alpha_{r}')(1-\alpha_{l}')(1-\delta)\right\}(1-p)}{(1-\alpha_{r}')(1-\alpha_{l}') - (1-\alpha_{r}')(1-2\alpha_{l}')p - (1-\alpha_{r}'-\alpha_{l}')\delta(1-p)} \left|\zeta_{1,1}\right\rangle\left\langle\zeta_{1,1}\right| + \frac{(1-\alpha_{r}')\alpha_{l}'p}{(1-\alpha_{r}')(1-\alpha_{l}') - (1-\alpha_{r}')(1-2\alpha_{l}')p - (1-\alpha_{r}'-\alpha_{l}')\delta(1-p)} \left|01\right\rangle\left\langle01\right|, \quad (E14)$$

$$\rho_{AB,\phi_{1},\phi_{2}} = \frac{\left\{ \alpha_{r}'(1-\alpha_{l}')\delta + (1-\alpha_{r}')\alpha_{l}'(1-\delta) \right\}(1-p)}{(1-\alpha_{r}')\alpha_{l}' + (1-\alpha_{r}')(1-2\alpha_{l}')p - (\alpha_{l}'-\alpha_{r}')\delta(1-p)} \left| \zeta_{1,2} \right\rangle \left\langle \zeta_{1,2} \right| \\ + \frac{(1-\alpha_{r}')(1-\alpha_{l}')p}{(1-\alpha_{r}')\alpha_{l}' + (1-\alpha_{r}')(1-2\alpha_{l}')p - (\alpha_{l}'-\alpha_{r}')\delta(1-p)} \left| 11 \right\rangle \left\langle 11 \right|,$$
(E15)

$$\rho_{AB,\phi_{1},\phi_{3}} = \frac{\left\{ \alpha_{r}'(1-\alpha_{l}')\delta + (1-\alpha_{r}')\alpha_{l}'(1-\delta) \right\}(1-p)}{(1-\alpha_{r}')\alpha_{l}' + (1-\alpha_{r}')(1-2\alpha_{l}')p - (\alpha_{l}'-\alpha_{r}')\delta(1-p)} \left| \zeta_{1,3} \right\rangle \left\langle \zeta_{1,3} \right| \\
+ \frac{(1-\alpha_{r}')(1-\alpha_{l}')p}{(1-\alpha_{r}')\alpha_{l}' + (1-\alpha_{r}')(1-2\alpha_{l}')p - (\alpha_{l}'-\alpha_{r}')\delta(1-p)} \left| 11 \right\rangle \left\langle 11 \right|,$$
(E16)

where,

$$|\zeta_{1,0}\rangle = \frac{\sqrt{\alpha'_r \alpha'_l \delta} |00\rangle - \sqrt{(1 - \alpha'_r)(1 - \alpha'_l)(1 - \delta)} |11\rangle}{\sqrt{\alpha'_r \alpha'_l \delta + (1 - \alpha'_r)(1 - \alpha'_l)(1 - \delta)}},$$
(E17)

$$|\zeta_{1,1}\rangle = \frac{\sqrt{\alpha'_r \alpha'_l \delta} |00\rangle + \sqrt{(1 - \alpha'_r)(1 - \alpha'_l)(1 - \delta)} |11\rangle}{\sqrt{\alpha'_r \alpha'_l \delta + (1 - \alpha'_r)(1 - \alpha'_l)(1 - \delta)}},$$
(E18)

14

$$|\zeta_{1,2}\rangle = \frac{\sqrt{(1-\alpha'_r)\alpha'_l(1-\delta)}|01\rangle - \sqrt{\alpha'_r(1-\alpha'_l)\delta}|10\rangle}{\sqrt{(1-\alpha'_r)\alpha'_l(1-\delta) + \alpha'_r(1-\alpha'_l)\delta}},$$
(E19)

$$|\zeta_{1,3}\rangle = \frac{\sqrt{(1-\alpha'_r)\alpha'_l(1-\delta)}\,|01\rangle + \sqrt{\alpha'_r(1-\alpha'_l)\delta}\,|10\rangle}{\sqrt{(1-\alpha'_r)\alpha'_l(1-\delta) + \alpha'_r(1-\alpha'_l)\delta}}.$$
(E20)

We retrieve Eqs. (E11) and (E12) for finite resource consumption. For the state $\rho_{N_{m-1}B,\phi_2}$ (as in Eq. (B9) with $\beta = \frac{1}{2}$), the outcomes corresponding to the second Bell measurement are given by

$$\rho_{AB,\phi_{2},\phi_{0}} = \frac{\left\{ (1 - \alpha_{r}')\alpha_{l}'\delta + \alpha_{r}'(1 - \alpha_{l}')(1 - \delta) \right\} (1 - p)}{\alpha_{r}'(1 - \alpha_{l}') - \alpha_{r}'(1 - 2\alpha_{l}')p - (\alpha_{r}' - \alpha_{l}')\delta(1 - p)} \left| \zeta_{2,0} \right\rangle \left\langle \zeta_{2,0} \right| \\ + \frac{\alpha_{r}'\alpha_{l}'p}{\alpha_{r}'(1 - \alpha_{l}') - \alpha_{r}'(1 - 2\alpha_{l}')p - (\alpha_{r}' - \alpha_{l}')\delta(1 - p)} \left| 00 \right\rangle \left\langle 00 \right|,$$
(E21)

$$\rho_{AB,\phi_{2},\phi_{1}} = \frac{\left\{ (1 - \alpha_{r}')\alpha_{l}'\delta + \alpha_{r}'(1 - \alpha_{l}')(1 - \delta) \right\} (1 - p)}{\alpha_{r}'(1 - \alpha_{l}') - \alpha_{r}'(1 - 2\alpha_{l}')p - (\alpha_{r}' - \alpha_{l}')\delta(1 - p)} \left| \zeta_{2,1} \right\rangle \left\langle \zeta_{2,1} \right| \\ + \frac{\alpha_{r}'\alpha_{l}'p}{\alpha_{r}'(1 - \alpha_{l}') - \alpha_{r}'(1 - 2\alpha_{l}')p - (\alpha_{r}' - \alpha_{l}')\delta(1 - p)} \left| 00 \right\rangle \left\langle 00 \right|,$$
(E22)

$$\rho_{AB,\phi_{2},\phi_{2}} = \frac{\left\{ (1 - \alpha_{r}')(1 - \alpha_{l}')\delta + \alpha_{r}'\alpha_{l}'(1 - \delta) \right\}(1 - p)}{\alpha_{r}'\alpha_{l}' + \alpha_{r}'(1 - 2\alpha_{l}')p + (1 - \alpha_{r}' - \alpha_{l}')\delta(1 - p)} \left| \zeta_{2,2} \right\rangle \left\langle \zeta_{2,2} \right| \\ + \frac{\alpha_{r}'(1 - \alpha_{l}')p}{\alpha_{r}'\alpha_{l}' + \alpha_{r}'(1 - 2\alpha_{l}')p + (1 - \alpha_{r}' - \alpha_{l}')\delta(1 - p)} \left| 10 \right\rangle \left\langle 10 \right|,$$
(E23)

$$\rho_{AB,\phi_{2},\phi_{3}} = \frac{\left\{ (1 - \alpha_{r}')(1 - \alpha_{l}')\delta + \alpha_{r}'\alpha_{l}'(1 - \delta) \right\}(1 - p)}{\alpha_{r}'\alpha_{l}' + \alpha_{r}'(1 - 2\alpha_{l}')p + (1 - \alpha_{r}' - \alpha_{l}')\delta(1 - p)} \left| \zeta_{2,3} \right\rangle \left\langle \zeta_{2,3} \right| \\ + \frac{\alpha_{r}'(1 - \alpha_{r}')p}{\alpha_{r}'\alpha_{l}' + \alpha_{r}'(1 - 2\alpha_{l}')p + (1 - \alpha_{r}' - \alpha_{l}')\delta(1 - p)} \left| 10 \right\rangle \left\langle 10 \right|,$$
(E24)

where,

$$|\zeta_{2,0}\rangle = \frac{\sqrt{(1 - \alpha'_r)\alpha'_l \delta} |01\rangle + \sqrt{\alpha'_r (1 - \alpha'_l)(1 - \delta)} |10\rangle}{\sqrt{(1 - \alpha'_r)\alpha'_l \delta + \alpha'_r (1 - \alpha'_l)(1 - \delta)}},$$
(E25)

$$|\zeta_{2,1}\rangle = \frac{\sqrt{(1-\alpha_r')\alpha_l'\delta}|01\rangle - \sqrt{\alpha_r'(1-\alpha_l')(1-\delta)}|10\rangle}{\sqrt{(1-\alpha_r')\alpha_l'\delta + \alpha_r'(1-\alpha_l')(1-\delta)}},$$
(E26)

$$|\zeta_{2,2}\rangle = \frac{\sqrt{\alpha_r'\alpha_l'(1-\delta)}\,|00\rangle + \sqrt{(1-\alpha_r')(1-\alpha_l')\delta}\,|11\rangle}{\sqrt{\alpha_r'\alpha_l'(1-\delta) + (1-\alpha_r')(1-\alpha_l')\delta}},\tag{E27}$$

$$|\zeta_{2,3}\rangle = \frac{\sqrt{\alpha'_r \alpha'_l (1-\delta)} |00\rangle - \sqrt{(1-\alpha'_r)(1-\alpha'_l)\delta} |11\rangle}{\sqrt{\alpha'_r \alpha'_l (1-\delta) + (1-\alpha'_r)(1-\alpha'_l)\delta}}.$$
(E28)

Following the same steps as before, for finite resource consumption, we get

$$(1 - \alpha'_r)(1 - \alpha'_l)\delta(1 - \delta)(1 - p)^2 \le \alpha'_r \alpha'_l p^2,$$
 (E29)

and,

$$(1 - \alpha'_r)\alpha'_l\delta(1 - \delta)(1 - p)^2 \le \alpha'_r(1 - \alpha'_l)p^2.$$
(E30)

Finally, for the state $\rho_{N_{m-1}B,\phi_3}$ (as in Eq. (B10) with $\beta = \frac{1}{2}$), the outcomes corresponding to the second Bell measurement are given by

$$\rho_{AB,\phi_{3},\phi_{0}} = \frac{\left\{ (1 - \alpha_{r}')\alpha_{l}'\delta + \alpha_{r}'(1 - \alpha_{l}')(1 - \delta) \right\} (1 - p)}{\alpha_{r}'(1 - \alpha_{l}') - \alpha_{r}'(1 - 2\alpha_{l}')p - (\alpha_{r}' - \alpha_{l}')\delta(1 - p)} \left| \zeta_{3,0} \right\rangle \left\langle \zeta_{3,0} \right| \\
+ \frac{\alpha_{r}'\alpha_{l}'p}{\alpha_{r}'(1 - \alpha_{l}') - \alpha_{r}'(1 - 2\alpha_{l}')p - (\alpha_{r}' - \alpha_{l}')\delta(1 - p)} \left| 00 \right\rangle \left\langle 00 \right|,$$
(E31)

$$\rho_{AB,\phi_{3},\phi_{1}} = \frac{\left\{ (1 - \alpha_{r}')\alpha_{l}'\delta + \alpha_{r}'(1 - \alpha_{l}')(1 - \delta) \right\}(1 - p)}{\alpha_{r}'(1 - \alpha_{l}') - \alpha_{r}'(1 - 2\alpha_{l}')p - (\alpha_{r}' - \alpha_{l}')\delta(1 - p)} \left| \zeta_{3,1} \right\rangle \left\langle \zeta_{3,1} \right| \\ + \frac{\alpha_{r}'\alpha_{l}'p}{\alpha_{r}'(1 - \alpha_{l}') - \alpha_{r}'(1 - 2\alpha_{l}')p - (\alpha_{r}' - \alpha_{l}')\delta(1 - p)} \left| 00 \right\rangle \left\langle 00 \right|,$$
(E32)

$$\rho_{AB,\phi_{3},\phi_{2}} = \frac{\left\{ (1 - \alpha_{r}')(1 - \alpha_{l}')\delta + \alpha_{r}'\alpha_{l}'(1 - \delta) \right\}(1 - p)}{\alpha_{r}'\alpha_{l}' + \alpha_{r}'(1 - 2\alpha_{l}')p + (1 - \alpha_{r}' - \alpha_{l}')\delta(1 - p)} \left| \zeta_{3,2} \right\rangle \left\langle \zeta_{3,2} \right| \\ + \frac{\alpha_{r}'(1 - \alpha_{l}')p}{\alpha_{r}'\alpha_{l}' + \alpha_{r}'(1 - 2\alpha_{l}')p + (1 - \alpha_{r}' - \alpha_{l}')\delta(1 - p)} \left| 10 \right\rangle \left\langle 10 \right|,$$
(E33)

$$\rho_{AB,\phi_{3},\phi_{3}} = \frac{\left\{ (1 - \alpha_{r}')(1 - \alpha_{l}')\delta + \alpha_{r}'\alpha_{l}'(1 - \delta) \right\}(1 - p)}{\alpha_{r}'\alpha_{l}' + \alpha_{r}'(1 - 2\alpha_{l}')p + (1 - \alpha_{r}' - \alpha_{l}')\delta(1 - p)} \left| \zeta_{3,3} \right\rangle \left\langle \zeta_{3,3} \right| \\ + \frac{\alpha_{r}'(1 - \alpha_{l}')p}{\alpha_{r}'\alpha_{l}' + \alpha_{r}'(1 - 2\alpha_{l}')p + (1 - \alpha_{r}' - \alpha_{l}')\delta(1 - p)} \left| 10 \right\rangle \left\langle 10 \right|,$$
(E34)

where,

$$|\zeta_{3,0}\rangle = \frac{\sqrt{(1-\alpha_r')\alpha_l'\delta}|01\rangle - \sqrt{\alpha_r'(1-\alpha_l')(1-\delta)}|10\rangle}{\sqrt{(1-\alpha_r')\alpha_l'\delta + \alpha_r'(1-\alpha_l')(1-\delta)}},$$
(E35)

$$|\zeta_{3,1}\rangle = \frac{\sqrt{(1-\alpha_r')\alpha_l'\delta} |01\rangle + \sqrt{\alpha_r'(1-\alpha_l')(1-\delta)} |10\rangle}{\sqrt{(1-\alpha_r')\alpha_l'\delta + \alpha_r'(1-\alpha_l')(1-\delta)}},$$
(E36)

$$|\zeta_{3,2}\rangle = \frac{\sqrt{\alpha_r'\alpha_l'(1-\delta)}\,|00\rangle - \sqrt{(1-\alpha_r')(1-\alpha_l')\delta}\,|11\rangle}{\sqrt{\alpha_r'\alpha_l'(1-\delta) + (1-\alpha_r')(1-\alpha_l')\delta}},\tag{E37}$$

$$|\zeta_{3,3}\rangle = \frac{\sqrt{\alpha_r'\alpha_l'(1-\delta)}|00\rangle + \sqrt{(1-\alpha_r')(1-\alpha_l')\delta}|11\rangle}{\sqrt{\alpha_r'\alpha_l'(1-\delta) + (1-\alpha_r')(1-\alpha_l')\delta}}.$$
(E38)

Again, for finite resource consumption, we retrieve Eqs. (E29) and (E30).

These four (*i.e.*, Eqs. (E11 - E12) and Eqs. (E29 - E30)) are exactly the constraints obtained in Eqs. (B19 - B22). So from Appendix B, it follows that the minimal condition that has to be satisfied for finite resource consumption is

$$\alpha_{r}' \alpha_{l}' \delta(1-\delta)(1-p)^{2} \leq (1-\alpha_{l}')(1-\alpha_{r}')p^{2}$$

$$\Rightarrow \qquad \frac{\alpha_{l}' \alpha_{r}'}{(1-\alpha_{l}')(1-\alpha_{r}')} \leq \frac{p^{2}}{\delta(1-\delta)(1-p)^{2}},$$
(E39)

which concludes the proof.

Appendix F: Observation 3

Proof. For better visualisation we rewrite Eq. (E39) in terms of concurrence as

$$\frac{\left(1 + \sqrt{1 - \mathcal{C}^{2}(\Psi_{l})}\right) \left(1 + \sqrt{1 - \mathcal{C}^{2}(\Psi_{r})}\right)}{\left(1 - \sqrt{1 - \mathcal{C}^{2}(\Psi_{r})}\right) \left(1 - \sqrt{1 - \mathcal{C}^{2}(\Psi_{r})}\right)} \le \frac{p^{2}}{\delta(1 - \delta)(1 - p)^{2}}.$$
(F1)

If all the consecutive $\{S_i\}_{i=1}^{n+1}$, except for i = m, share the same pure state $|\psi_i\rangle = |\psi\rangle = \sqrt{\alpha} |00\rangle + \sqrt{1-\alpha} |11\rangle$, *i.e.*, $\alpha_i = \alpha$ for $i \in [1, n+1] \setminus \{m\}$, where i is a natural number, with concurrence $\mathcal{C} = 2\sqrt{\alpha(1-\alpha)}$ then Eq. (F1) can be written as

$$\frac{\left(1+\sqrt{1-\mathcal{C}^{2(m-1)}}\right)\left(1+\sqrt{1-\mathcal{C}^{2(n-m+1)}}\right)}{\left(1-\sqrt{1-\mathcal{C}^{2(n-m+1)}}\right)} \le \frac{p^2}{\delta(1-\delta)(1-p)^2}.$$
(F2)

Thus for any given number of nodes n and noisy state parameters p and δ , we can find the value of the lower bound on the concurrence required for finite resource consumption depending on the position of the noisy state m in the given chain. We calculate the lower bound on concurrence for every value of m from 1 to n + 1 using Eq. (F2) for the given values of the state parameters. Then, using Eq. (D3), we calculate the maximum possible variance in resource consumption. The results are plotted in Figure 3.

Appendix G: Two Noisy segments

Proof. Let us consider a single node scenario $(A - N_1 - B)$ with two segments S_1 and S_2 . Segment S_1 shares a rank two mixed state $\rho_1 = \rho(p, \delta) = pP + (1 - p)\zeta(\delta)$, where $\zeta(\delta) := |\zeta(\delta)\rangle \langle \zeta(\delta)|$ with $|\zeta(\delta)\rangle = \sqrt{\delta} |00\rangle + \sqrt{1 - \delta} |11\rangle$, and $P = |01\rangle \langle 01|$. Let us assume that segment S_2 also has the same state shared, *i.e.*, $\rho_2 = \rho(p, \delta) = \rho_1$. Using Eq. (4) we have

$$F_{AB}^{\star} \le \min\{F^{\star}(\rho(p,\delta)), F_C\},\tag{G1}$$

where, $F_C = \frac{1 + C^2(\rho(p, \delta))}{2}$.

If $F_C < F^*(\rho(p, \delta))$, then according to Eq. (G1), it is impossible to maintain the same fidelity between A and B as achieved in segment S_1 between A and N_1 . Let us now consider Bell measurement ($\beta = \frac{1}{2}$) at node N_1 . Eq. (G1) determines the maximum fidelity of the final state between Alice and Bob. Using Eq. (B1) with $\delta = 0.95$, we can determine the range of the noise parameter where it is impossible to maintain the same fidelity between Alice and Bob as in segment S_1 (or segment S_2) and it can be calculated from the previously mentioned inequality as

$$0 \le p < 0.25.$$
 (G2)

In the single node scenario where $\rho_1 = \rho(p, \delta)$, but segment S_2 has a pure two-qubit entangled state $|\psi_2\rangle = \sqrt{\alpha} |00\rangle + \sqrt{1-\alpha} |11\rangle$, using Eq. (B19) with $\delta = 0.95$ and $\alpha = 0.55$, we find that for

$$0.19 \le p < 1,\tag{G3}$$

we can establish $F_{AB}^{\star} = F^{\star}(\rho_1)$.

Thus, from Eqs. (G2) and (G3), we can see that for

$$0.19 \le p < 0.25,$$
 (G4)

it is impossible to establish the same fidelity between Alice and Bob as in the initial state if both nodes are noisy. This is in contrast to the case where one of the nodes is not noisy, allowing us to maintain the same fidelity. \Box