

INVARIANT RINGS OF THE SPECIAL ORTHOGONAL GROUP HAVE NONUNIMODAL h -VECTORS

ALDO CONCA, ANURAG K. SINGH, AND MATTEO VARBARO

To Sudhir Ghorpade, in celebration of his sixtieth birthday.

ABSTRACT. For K an infinite field of characteristic other than two, consider the action of the special orthogonal group $\mathrm{SO}_t(K)$ on a polynomial ring via copies of the regular representation. When K has characteristic zero, Boutot's theorem implies that the invariant ring has rational singularities; when K has positive characteristic, the invariant ring is F -regular, as proven by Hashimoto using good filtrations. We give a new proof of this, viewing the invariant ring for $\mathrm{SO}_t(K)$ as a cyclic cover of the invariant ring for the corresponding orthogonal group; this point of view has a number of useful consequences, for example it readily yields the a -invariant and information on the Hilbert series. Indeed, we use this to show that the h -vector of the invariant ring for $\mathrm{SO}_t(K)$ need not be unimodal.

1. INTRODUCTION

Let X be an $n \times n$ symmetric matrix of indeterminates over a field K , and let $I_{t+1}(X)$ denote the ideal of the polynomial ring $K[X]$ generated by the size $t+1$ minors of X . For t a positive integer with $t+1 \leq n$, we refer to $K[X]/I_{t+1}(X)$ as a *symmetric determinantal ring*. The ring $K[X]/I_{t+1}(X)$ is a Cohen-Macaulay normal domain of dimension

$$\binom{n+1}{2} - \binom{n+1-t}{2},$$

as proven in [Ku]. These rings have been studied extensively, in part because they arise as invariant rings for the natural action of the orthogonal group

$$(1.0.1) \quad \mathrm{O}_t(K) := \{M \in \mathrm{GL}_t(K) \mid M^{\mathrm{tr}}M = \mathrm{id}\}$$

as follows: for Y a $t \times n$ matrix of indeterminates, $\mathrm{O}_t(K)$ acts K -linearly on $K[Y]$ via

$$M: Y \longmapsto MY \quad \text{for } M \in \mathrm{O}_t(K).$$

This is a right action of $\mathrm{O}_t(K)$ on the polynomial ring $K[Y]$, corresponding to a left action of $\mathrm{O}_t(K)$ on the affine space $\mathbb{A}_K^{t \times n}$. Note that $Y^{\mathrm{tr}}Y \longmapsto Y^{\mathrm{tr}}M^{\mathrm{tr}}MY = Y^{\mathrm{tr}}Y$ for $M \in \mathrm{O}_t(K)$, so the entries of $Y^{\mathrm{tr}}Y$ are invariant under the action; when the field K is infinite of characteristic other than two, the invariant ring is precisely the K -algebra generated by the entries of $Y^{\mathrm{tr}}Y$, see [DP, Theorem 5.6], and is isomorphic to the symmetric determinantal ring $K[X]/I_{t+1}(X)$ via the entrywise map $X \longmapsto Y^{\mathrm{tr}}Y$. We use this to identify the rings $K[X]/I_{t+1}(X)$ and $K[Y^{\mathrm{tr}}Y]$.

A.C. and M.V. are supported by PRIN 2020355B8Y ‘‘Squarefree Gröbner degenerations, special varieties and related topics,’’ by MIUR Excellence Department Project awarded to the Dept. of Mathematics, Univ. of Genova, CUP D33C23001110001 and by INdAM-GNSAGA; A.K.S. is supported by NSF grants DMS-2101671 and DMS-2349623; all authors were supported by NSF grant DMS-1928930 and by Alfred P. Sloan Foundation grant G-2021-16778, while in residence at SLMath/MSRI, Berkeley, during the Spring 2024 Commutative Algebra program. We are grateful to Mitsuyasu Hashimoto for valuable discussions, and to the referee for several useful comments.

By [Go1, Go2], the ring $R := K[Y^{\text{tr}}Y]$ has class group $\mathbb{Z}/2$, and is Gorenstein precisely when $n \equiv t + 1 \pmod{2}$. Taking \mathfrak{p} to be a prime ideal that serves as a generator for the class group, it follows that the symbolic power $\mathfrak{p}^{(2)}$ is isomorphic to R . We choose an explicit isomorphism $\mathfrak{p}^{(2)} \cong R$ so that the cyclic cover of R with respect to \mathfrak{p} is precisely the invariant ring for the action of the special orthogonal group $\text{SO}_t(K)$. This gives a straightforward approach towards studying the invariant ring $K[Y]^{\text{SO}_t(K)}$, for example towards determining its a -invariant and information regarding the Hilbert series.

When K is an infinite field of characteristic two, the groups $\text{O}_t(K)$ and $\text{SO}_t(K)$ coincide when taking $\text{O}_t(K)$ to be the group as defined in (1.0.1); the invariant ring in this case is

$$K[Y^{\text{tr}}Y, \sum_{i=1}^t y_{ij} \mid 1 \leq j \leq n],$$

see [Ri, Proposition 17], and a presentation is provided by [Ri, Proposition 23]. The reader is warned that there are varying definitions used for the orthogonal group in characteristic two, see for example [PS, page 10].

Section 2 includes some generalities on cyclic covers; these are used in Section 3 where we compute the a -invariant of $K[Y]^{\text{SO}_t(K)}$ and also record a proof that this ring is F -regular. Section 4 is devoted to the h -vector of $K[Y]^{\text{SO}_t(K)}$, i.e., the coefficients of the numerator of its Hilbert series: the key result here is that this invariant ring is a semistandard graded Gorenstein normal domain, for which the h -vector need not be unimodal; the context for this is discussed as well in Section 4.

2. CYCLIC COVERS AND F -REGULARITY

Let R be a normal domain. By a *divisorial ideal* of R , we mean a nonzero intersection of fractional principal ideals. Let \mathfrak{a} be a divisorial ideal that has finite order m when viewed as an element of the divisor class group of R . Then $\mathfrak{a}^{(m)} = \alpha R$, for an element α in the fraction field of R . Set

$$(2.0.1) \quad T := 1/\alpha^{1/m},$$

which is an element in an algebraic closure of the fraction field of R ; the choice of α or the m -th root is not unique. The *cyclic cover* of R with respect to \mathfrak{a} is the ring

$$\tilde{R} := R[\mathfrak{a}T, \mathfrak{a}^{(2)}T^2, \mathfrak{a}^{(3)}T^3, \dots],$$

viewed as a subring of $R[T]$. Since

$$\mathfrak{a}^{(m+k)}T^{m+k} = \alpha\mathfrak{a}^{(k)}T^{m+k} = \mathfrak{a}^{(k)}T^k$$

for each $k \geq 0$, the ring \tilde{R} is a finitely generated reflexive R -module; specifically, one has an R -module isomorphism

$$\tilde{R} \cong R \oplus \mathfrak{a} \oplus \mathfrak{a}^{(2)} \oplus \dots \oplus \mathfrak{a}^{(m-1)}.$$

When the ring R is \mathbb{N} -graded and \mathfrak{a} is a homogeneous divisorial ideal of finite order m , there exists a homogeneous element α with $\mathfrak{a}^{(m)} = \alpha R$, and the \mathbb{N} -grading on R extends to a \mathbb{Q} -grading on \tilde{R} obtained by setting

$$\deg T := -(\deg \alpha)/m.$$

It turns out that this is a $\mathbb{Q}_{\geq 0}$ -grading on \tilde{R} , and that $[\tilde{R}]_0 = R_0$, see [Si, Proposition 4.2].

Suppose that the characteristic of R is zero or relatively prime to m , and that \mathfrak{p} is a height one prime ideal of R . Then the ideal $\alpha R_{\mathfrak{p}}$ is principal; take r to be a generator. Since $r^m = \alpha u$, for u a unit in $R_{\mathfrak{p}}$, it follows that

$$\tilde{R}_{\mathfrak{p}} = R_{\mathfrak{p}}[rT] \cong R_{\mathfrak{p}}[u^{1/m}],$$

so $R_{\mathfrak{p}} \rightarrow \tilde{R}_{\mathfrak{p}}$ is étale. In particular, under this assumption on the characteristic, the ring $\tilde{R}_{\mathfrak{p}}$ is regular for each height one prime of R ; since each $\alpha^{(k)}$ is reflexive, the ring \tilde{R} also satisfies the Serre condition S_2 , and is hence a normal domain. By [Wa, Theorem 2.7], F -regularity is preserved under finite extensions that are étale at height one primes, so one has:

Theorem 2.1 (Watanabe). *Let R be an \mathbb{N} -graded ring that is finitely generated over a field R_0 of characteristic $p > 0$, and let \tilde{R} be the cyclic cover of R with respect to a homogeneous ideal of finite order relatively prime to p . Then, if R is F -regular, so is \tilde{R} .*

The restriction on the characteristic is removed in [CR, Theorem C]. For the theory of F -regularity in the graded setting, we point the reader towards [HH]. When R is an \mathbb{N} -graded ring finitely generated over a field R_0 of positive characteristic, the notions of weak F -regularity, F -regularity, and strong F -regularity all coincide as proven in [LS], so we do not make a distinction between these in the present paper.

The F -regularity of generic determinantal rings and of Plücker coordinate rings of Grassmannians is proven as [HH, Theorem 7.14]; the proof therein is readily adapted to symmetric determinantal rings, as we show next. For a different approach, see [Ló, §4.1].

Theorem 2.2. *Let X be an $n \times n$ symmetric matrix of indeterminates over a field K of positive prime characteristic. Then the ring $K[X]/I_{t+1}(X)$ is F -regular.*

Proof. If $n \equiv t + 1 \pmod{2}$, then $K[X]/I_{t+1}(X)$ is Gorenstein; otherwise, enlarge X to a symmetric matrix \tilde{X} of size $n + 1$, in which case the ring $K[\tilde{X}]/I_{t+1}(\tilde{X})$ is Gorenstein, and contains $K[X]/I_{t+1}(X)$ as a pure subring. Since F -regularity is inherited by pure subrings, it suffices to prove the desired result when $R := K[X]/I_{t+1}(X)$ is Gorenstein.

The a -invariant of R is computed in [Ba] and [Co2], and recorded in the following section; in particular, $a(R) < 0$. We next claim that R is F -injective, equivalently F -pure, since the notions coincide in the Gorenstein case. This follows by [CH2, Theorem 2.1] in combination with the main result of [Co1] asserting that the “diagonal” initial ideal of $I_{t+1}(X)$ is square-free and defines a Cohen-Macaulay ring.

The F -regularity of R now follows from [HH, Corollary 7.13], once we verify that the localization $R_{x_{ij}}$ is F -regular for each x_{ij} . Using the lemma below and induction on t , the localizations $R_{x_{11}}$ and R_{Δ} are F -regular; but then $R_{\mathfrak{p}}$ is F -regular if \mathfrak{p} is a prime ideal such that $x_{11} \notin \mathfrak{p}$ or $\Delta \notin \mathfrak{p}$. It follows that $R_{\mathfrak{p}}$ is also F -regular if $x_{12} \notin \mathfrak{p}$. Since we have accounted for the diagonal variable x_{11} and the off-diagonal variable x_{12} , the symmetry implies that $R_{x_{ij}}$ is F -regular for each x_{ij} . \square

Lemma 2.3. *Let $R := K[X]/I_{t+1}(X)$, where X is a symmetric $n \times n$ matrix of indeterminates. Then:*

- (1) *The ring $R_{x_{11}}$ is isomorphic to a localization of a polynomial ring over $K[X']/I_t(X')$, where X' is a symmetric $(n-1) \times (n-1)$ matrix of indeterminates.*
- (2) *For $\Delta := x_{11}x_{22} - x_{12}^2$, the ring R_{Δ} is isomorphic to a localization of a polynomial ring over $K[X']/I_{t-1}(X')$, for X' a symmetric $(n-2) \times (n-2)$ matrix of indeterminates.*

For a proof, see [Jo, Lemma 1.1]; the argument also appears implicitly in [MV].

3. THE a -INVARIANT

Let Y be a $t \times n$ matrix of indeterminates over a field K . In this section, we work with the grading on the subring $R := K[Y^t Y]$ that is induced by the standard grading on the polynomial ring $K[Y]$. Note that under the identification of $K[X]/I_{t+1}(X)$ with $K[Y^t Y]$, this corresponds to taking $\deg x_{ij} = 2$ for each i, j . With this grading, [Ba, Theorem 4.4] or [Co2, Theorem 2.4] imply that the a -invariant of R is

$$a(R) = \begin{cases} -t(n+1) & \text{if } n \equiv t \pmod{2}, \\ -tn & \text{if } n \not\equiv t \pmod{2}; \end{cases}$$

more generally, the graded canonical module of R is

$$\omega_R = \begin{cases} \mathfrak{p}(-tn+t) & \text{if } n \equiv t \pmod{2}, \\ R(-tn) & \text{if } n \not\equiv t \pmod{2}, \end{cases}$$

where \mathfrak{p} is the ideal of $K[Y^t Y]$ generated by the maximal minors of the first t rows of $Y^t Y$, i.e., by the maximal minors of the product matrix

$$\begin{pmatrix} y_{11} & y_{21} & \cdots & y_{t1} \\ y_{12} & y_{22} & \cdots & y_{t2} \\ \vdots & \vdots & \ddots & \vdots \\ y_{1t} & y_{2t} & \cdots & y_{tt} \end{pmatrix} \begin{pmatrix} y_{11} & y_{12} & y_{13} & \cdots & \cdots & y_{1n} \\ y_{21} & y_{22} & y_{23} & \cdots & \cdots & y_{2n} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ y_{t1} & y_{t2} & y_{t3} & \cdots & \cdots & y_{tn} \end{pmatrix}.$$

Using the identification of $K[X]/I_{t+1}(X)$ with $K[Y^t Y]$, the ideal \mathfrak{p} is prime of height one by [Ku, Theorem 1], and generates the class group of R by [Go1]. The symbolic power $\mathfrak{p}^{(2)}$ is the principal ideal of R generated by the determinant of the first t columns of the product matrix displayed above, i.e., $\mathfrak{p}^{(2)}$ is generated by the square of

$$\Delta := \det \begin{pmatrix} y_{11} & y_{21} & \cdots & y_{t1} \\ y_{12} & y_{22} & \cdots & y_{t2} \\ \vdots & \vdots & \ddots & \vdots \\ y_{1t} & y_{2t} & \cdots & y_{tt} \end{pmatrix}.$$

Choosing a unit as in (2.0.1), set

$$T := 1/\Delta.$$

The generators of $\mathfrak{p}T$ are then identified with the maximal minors of the matrix Y , so that the cyclic cover \tilde{R} of R with respect to \mathfrak{p} is the subring of the polynomial ring $K[Y]$ generated by the entries of the product matrix $Y^t Y$ along with the maximal minors of Y . It is clear that these generators are fixed under the action of the special orthogonal group

$$M: Y \mapsto MY \quad \text{for } M \in \mathrm{SO}_t(K).$$

When the field K is infinite of characteristic other than two, the invariant ring is precisely the K -algebra generated by these elements, [DP, Theorem 5.6].

We determine the graded canonical module of \tilde{R} ; while the semisimplicity of $\mathrm{SO}_t(K)$ may be used to verify that \tilde{R} is Gorenstein, [HR, page 123], our goal is to additionally obtain the a -invariant of \tilde{R} . Since $\deg T = -t$, one has

$$\tilde{R} = R \oplus \mathfrak{p}(t).$$

Let \mathfrak{m} denote the homogeneous maximal ideal of R . For an \mathbb{N} -graded R -module M , we use $\underline{\mathrm{Hom}}(M, R/\mathfrak{m})$ to denote its graded dual as in [GW, page 184]. Setting $d := \dim R$, the

graded canonical module of \tilde{R} may be computed as

$$\omega_{\tilde{R}} = \underline{\text{Hom}}(H_m^d(\tilde{R}), R/\mathfrak{m}) = \underline{\text{Hom}}(H_m^d(R), R/\mathfrak{m}) \oplus \underline{\text{Hom}}(H_m^d(\mathfrak{p}(t)), R/\mathfrak{m}).$$

The first term in this direct sum is ω_R , while the second is

$$\begin{aligned} \underline{\text{Hom}}(H_m^d(\mathfrak{p}(t)), R/\mathfrak{m}) &= \underline{\text{Hom}}(H_m^d(\omega_R) \otimes_R \omega_R^{(-1)} \otimes_R \mathfrak{p}(t), R/\mathfrak{m}) \\ &= \text{Hom}_R(\omega_R^{(-1)} \otimes_R \mathfrak{p}(t), \underline{\text{Hom}}(H_m^d(\omega_R), R/\mathfrak{m})) \\ &= \text{Hom}_R(\omega_R^{(-1)} \otimes_R \mathfrak{p}(t), R) \\ &= (\omega_R \otimes_R \mathfrak{p}^{(-1)}(-t))^{**}, \end{aligned}$$

where $(-)^{**}$ is the reflexive hull. Since $\mathfrak{p}^{(2)} = R(-2t)$, one has $\mathfrak{p}^{(-1)} = \mathfrak{p}(2t)$, so

$$(\omega_R \otimes_R \mathfrak{p}^{(-1)}(-t))^{**} = \begin{cases} R(-tn) & \text{if } n \equiv t \pmod{2}, \\ \mathfrak{p}(-tn+t) & \text{if } n \not\equiv t \pmod{2}. \end{cases}$$

Putting it all together, one gets

$$\omega_{\tilde{R}} = \begin{cases} \mathfrak{p}(-tn+t) \oplus R(-tn) & \text{if } n \equiv t \pmod{2}, \\ R(-tn) \oplus \mathfrak{p}(-tn+t) & \text{if } n \not\equiv t \pmod{2}, \end{cases}$$

so that

$$\omega_{\tilde{R}} = \tilde{R}(-tn),$$

i.e., \tilde{R} is Gorenstein with $a(\tilde{R}) = -tn$. To summarize what we have at this stage:

Theorem 3.1. *Let Y be a $t \times n$ matrix of indeterminates over a field K of characteristic other than two. Let \tilde{R} denote the K -subalgebra of $K[Y]$ generated by the entries of the product matrix $Y^t Y$ along with the maximal minors of Y . Then \tilde{R} is a Gorenstein normal domain. When K has characteristic zero, the ring \tilde{R} has rational singularities; when K has positive characteristic, \tilde{R} is F -regular.*

With the \mathbb{N} -grading on \tilde{R} inherited from the standard grading on $K[Y]$, one has

$$a(\tilde{R}) = -tn.$$

The fact that \tilde{R} has rational singularities in characteristic zero follows from Boutot's theorem [Bo]; the F -regularity in characteristic $p \geq 3$ follows by combining Theorem 2.1 and Theorem 2.2. For a different approach using good filtrations, see [Ha, Corollary 2].

Remark 3.2. The ring \tilde{R} in Theorem 3.1 has K -algebra generators in degree 2 and degree t ; it admits a standard grading in the following two cases:

(i) When $t = 1$, index the entries of Y as y_1, \dots, y_n . The ring $R := K[Y^t Y]$ is then the second Veronese subring of the polynomial ring $K[Y]$, i.e., the subring generated by the monomials $y_i y_j$. One has

$$\mathfrak{p} = (y_1^2, y_1 y_2, \dots, y_1 y_n)R \quad \text{and} \quad \mathfrak{p}^{(2)} = (y_1^2)R.$$

Taking $T := 1/y_1$, the cyclic cover \tilde{R} coincides with $K[Y]$ under the standard grading.

(ii) When $t = 2$, the K -algebra generators of \tilde{R} are the entries of $Y^t Y$, and the size two minors of Y ; these generators all have degree two, so the grading on \tilde{R} may be rescaled to a standard grading.

Remark 3.3. When t is even, the ring \tilde{R} in Theorem 3.1 has generators of even degree; rescaling by a factor of two, one obtains generators in degree one (the entries of $Y^u Y$) and generators in degree $t/2$ (the maximal minors of Y); this is the grading considered in the following section. This is a *semistandard* grading on \tilde{R} , i.e., an \mathbb{N} -grading under which the ring is integral over the K -subalgebra generated by its elements of degree one.

4. NONUNIMODAL h -VECTORS

A description for the Hilbert function of a generic determinantal ring may be found in [Ab], while an expression for its Hilbert series is presented in [CH1]. In particular, for the numerator of the Hilbert series, known as the *h-polynomial*, one has both a combinatorial description (in terms on non-intersection paths with given number of turns) and an explicit compact (and determinantal!) formula. For pfaffian rings, the corresponding results are in [DN, GK]. For symmetric determinantal rings one finds in [Co2] a combinatorial description of the *h-polynomial*, but no compact determinantal expression for it is known in general. However, for X a symmetric $n \times n$ matrix of indeterminates and $t + 1 = n - 1$, the expression of the *h-polynomial* of $K[X]/I_{t+1}(X)$ is easily obtained to be

$$(4.0.1) \quad \binom{2}{2} + \binom{3}{2}z + \cdots + \binom{n}{2}z^{n-2},$$

see for example [Co2, Example 2.3(c)].

As in Remark 3.3, an \mathbb{N} -grading on a ring A is *semistandard* if A is a finitely generated algebra over a field $K := A_0$, and A is integral over the K -subalgebra generated by its elements of degree one. This condition ensures that the Hilbert series of A may be written as a rational function

$$\frac{h_0 + h_1 z + h_2 z^2 + \cdots + h_k z^k}{(1 - z)^{\dim A}}, \quad \text{where } h_i \in \mathbb{Z} \text{ and } h_k \neq 0.$$

The coefficients of the numerator, i.e., of the *h-polynomial*, form the *h-vector* (h_0, \dots, h_k) of the ring A . When A is Cohen-Macaulay, it is readily seen that each h_i is nonnegative; when A is Gorenstein, the *h-vector* is a palindrome, i.e., $h_i = h_{k-i}$ for each $0 \leq i \leq k$. In this case, the *h-vector* is said to be *unimodal* if

$$h_0 \leq h_1 \leq \dots \leq h_{\lfloor k/2 \rfloor}.$$

Unimodality results reflect interesting geometric and combinatorial properties; they figure prominently in Ehrhart theory. Following his proof of the Anand-Dumir-Gupta conjectures regarding the enumeration of magic squares [St1, St3], Stanley asked if the *h-vector* of the corresponding affine semigroup ring is unimodal. This was indeed proven to be the case by Athanasiadis [At], see also [BR]. While Mustața and Payne [MP] have constructed examples of Gorenstein normal affine semigroup rings for which the *h-vector* is not unimodal, these are not standard graded, and the following remains unresolved:

Conjecture 4.1. The *h-vector* of a standard graded Gorenstein domain is unimodal.

This is due to Stanley [St2, Conjecture 4(a)], see also [Bra, Conjecture 1], [Bre, Conjecture 5.1], [Bru, page 36], and [Hi, Conjecture 1.5]. We show that invariant rings for the action of $SO_t(K)$ yield examples of “naturally occurring” semistandard graded Gorenstein normal domains, for which the *h-vector* is not unimodal:

Theorem 4.2. Consider a $2m \times (2m + 2)$ matrix of indeterminates Y over a field K of characteristic other than two. Let \tilde{R} denote the K -subalgebra of $K[Y]$ generated by the

entries of the product matrix $Y^{\text{tr}}Y$ and the maximal minors of Y , where the generators are assigned degree 1 and degree m respectively. If $m \geq 2$, the h -vector of \tilde{R} is not unimodal.

Proof. Viewing the subring $R := K[Y^{\text{tr}}Y]$ as a symmetric determinantal ring and using the expression (4.0.1), one see that R has Hilbert series

$$\frac{\binom{2}{2} + \binom{3}{2}z + \cdots + \binom{2m+2}{2}z^{2m}}{(1-z)^{2m^2+5m}}.$$

The ring R is not Gorenstein; the Hilbert series of R yields that of ω_R , from which it follows that the cyclic cover \tilde{R} has Hilbert series

$$\frac{\left[\binom{2}{2} + \binom{3}{2}z + \cdots + \binom{2m+2}{2}z^{2m}\right] + \left[\binom{2m+2}{2}z^m + \binom{2m+1}{2}z^{m+1} + \cdots + \binom{2}{2}z^{3m}\right]}{(1-z)^{2m^2+5m}}.$$

Hence

$$h_m - h_{m+1} = \left[\binom{m+2}{2} + \binom{2m+2}{2}\right] - \left[\binom{m+3}{2} + \binom{2m+1}{2}\right] = m - 1,$$

so the h -vector of \tilde{R} is not unimodal; for a specific example, the case $m = 2$ yields the nonunimodal h -vector

$$(1, 3, 6, 10, 15, 0, 0) + (0, 0, 15, 10, 6, 3, 1) = (1, 3, 21, 20, 21, 3, 1). \quad \square$$

REFERENCES

- [Ab] S. S. Abhyankar, *Enumerative combinatorics of Young tableaux*, Monogr. Textbooks Pure Appl. Math. **115**, Marcel Dekker, Inc., New York, 1988. 6
- [At] C. A. Athanasiadis, *Ehrhart polynomials, simplicial polytopes, magic squares and a conjecture of Stanley*, J. Reine Angew. Math. **583** (2005), 163–174. 6
- [Ba] M. Barile, *The Cohen-Macaulayness and the a -invariant of an algebra with straightening laws on a doset*, Comm. Algebra **22** (1994), 413–430. 3, 4
- [Bo] J.-F. Boutot, *Singularités rationnelles et quotients par les groupes réductifs*, Invent. Math. **88** (1987), 65–68. 5
- [Bra] B. Braun, *Unimodality problems in Ehrhart theory*, in: Recent trends in combinatorics, IMA Vol. Math. Appl. **159**, pp. 687–711, Springer, Cham, 2016. 6
- [Bre] F. Brenti, *Log-concave and unimodal sequences in algebra, combinatorics, and geometry: an update*, Contemp. Math. **178** (1994), 71–89. 6
- [Bru] W. Bruns, *Commutative algebra arising from the Anand-Dumir-Gupta conjectures*, in: Commutative algebra and combinatorics, Ramanujan Math. Soc. Lect. Notes Ser. **4**, pp. 1–38, Ramanujan Math. Soc., Mysore, 2007. 6
- [BR] W. Bruns and T. Römer, *h -vectors of Gorenstein polytopes*, J. Combin. Theory Ser. A **114** (2007), 65–76. 6
- [CR] J. A. Carvajal-Rojas, *Finite torsors over strongly F -regular singularities*, Épijournal Géom. Algébrique **6** (2022), Art. 1, 30 pp. 3
- [Co1] A. Conca, *Gröbner bases of ideals of minors of a symmetric matrix*, J. Algebra **166** (1994), 406–421. 3
- [Co2] A. Conca, *Symmetric ladders*, Nagoya Math. J. **136** (1994), 35–56. 3, 4, 6
- [CH1] A. Conca and J. Herzog, *On the Hilbert function of determinantal rings and their canonical module*, Proc. Amer. Math. Soc. **122** (1994), 677–681. 6
- [CH2] A. Conca and J. Herzog, *Ladder determinantal rings have rational singularities*, Adv. Math. **132** (1997), 120–147. 3
- [DP] C. De Concini and C. Procesi, *A characteristic free approach to invariant theory*, Adv. Math. **21** (1976), 330–354. 1, 4
- [DN] E. De Negri, *Some results on Hilbert series and a -invariant of Pfaffian ideals*, Math. J. Toyama Univ. **24** (2001), 93–106. 6
- [GK] S. R. Ghorpade and C. Krattenthaler, *The Hilbert series of Pfaffian rings*, in: Algebra, arithmetic and geometry with applications (West Lafayette, IN, 2000), pp. 337–356, Springer, Berlin, 2004. 6
- [Go1] S. Goto, *The divisor class group of a certain Krull domain*, J. Math. Kyoto Univ. **17** (1977), 47–50. 2, 4

- [Go2] S. Goto, *On the Gorensteinness of determinantal loci*, J. Math. Kyoto Univ. **19** (1979), 371–374. [2](#)
- [GW] S. Goto and K.-i. Watanabe, *On graded rings, I*, J. Math. Soc. Japan **30** (1978), 179–213. [4](#)
- [Ha] M. Hashimoto, *Good filtrations of symmetric algebras and strong F -regularity of invariant subrings*, Math. Z. **236** (2001), 605–623. [5](#)
- [Hi] T. Hibi, *Flawless O -sequences and Hilbert functions of Cohen-Macaulay integral domains*, J. Pure Appl. Algebra **60** (1989), 245–251. [6](#)
- [HH] M. Hochster and C. Huneke, *Tight closure of parameter ideals and splitting in module-finite extensions*, J. Algebraic Geom. **3** (1994), 599–670. [3](#)
- [HR] M. Hochster and J. L. Roberts, *Rings of invariants of reductive groups acting on regular rings are Cohen-Macaulay*, Adv. Math. **13** (1974), 115–175. [4](#)
- [Jo] T. Józefiak, *Ideals generated by minors of a symmetric matrix*, Comment. Math. Helv. **53** (1978), 595–607. [3](#)
- [Ku] R. Kutz, *Cohen-Macaulay rings and ideal theory in rings of invariants of algebraic groups*, Trans. Amer. Math. Soc. **194** (1974), 115–129. [1](#), [4](#)
- [Ló] A. C. Lórinçz, *On the collapsing of homogeneous bundles in arbitrary characteristic*, Ann. Sci. Éc. Norm. Supér. (4) **56** (2023), 1313–1337. [3](#)
- [LS] G. Lyubeznik and K. E. Smith, *Strong and weak F -regularity are equivalent for graded rings*, Amer. J. Math. **121** (1999), 1279–1290. [3](#)
- [MV] A. Micali and O. E. Villamayor, *Sur les algèbres de Clifford*, Ann. Sci. École Norm. Sup. (4) **1** (1968), 271–304. [3](#)
- [MP] M. Mustață and S. Payne, *Ehrhart polynomials and stringy Betti numbers*, Math. Ann. **333** (2005), 787–795. [6](#)
- [PS] A. N. Parshin and I. R. Shafarevich, editors, *Algebraic geometry IV, Linear algebraic groups. Invariant theory*, Encyclopaedia of Mathematical Sciences **55**, Springer-Verlag, Berlin, 1994. [2](#)
- [Ri] D. R. Richman, *The fundamental theorems of vector invariants*, Adv. Math. **73** (1989), 43–78. [2](#)
- [Si] A. K. Singh, *Cyclic covers of rings with rational singularities*, Trans. Amer. Math. Soc. **355** (2003), 1009–1024. [2](#)
- [St1] R. P. Stanley, *Linear homogeneous Diophantine equations and magic labelings of graphs*, Duke Math. J. **40** (1973), 607–632. [6](#)
- [St2] R. P. Stanley, *Log-concave and unimodal sequences in algebra, combinatorics, and geometry*, in: Graph theory and its applications: East and West (Jinan, 1986), Ann. New York Acad. Sci. **576**, pp. 500–535, New York Acad. Sci., New York, 1989. [6](#)
- [St3] R. P. Stanley, *Combinatorics and commutative algebra*, Second edition, Progress in Mathematics **41**, Birkhäuser, Boston, MA, 1996. [6](#)
- [Wa] K.-i. Watanabe, *F -regular and F -pure normal graded rings*, J. Pure Appl. Algebra **71** (1991), 341–350. [3](#)

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI GENOVA, DIPARTIMENTO DI ECCELLENZA 2023-2027, VIA DODECANESO 35, I-16146 GENOVA, ITALY
 Email address: conca@dimma.unige.it

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, 155 SOUTH 1400 EAST, SALT LAKE CITY, UT 84112, USA
 Email address: singh@math.utah.edu

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI GENOVA, DIPARTIMENTO DI ECCELLENZA 2023-2027, VIA DODECANESO 35, I-16146 GENOVA, ITALY
 Email address: varbaro@dimma.unige.it