

Broad band spectral modeling of M87 nucleus

ANDRZEJ NIEDŹWIECKI,¹ MICHAŁ SZANECKI,¹ AND AGNIESZKA JANIUK²

¹*Faculty of Physics and Applied Informatics, Łódź University, Pomorska 149/153, 90-236 Łódź, Poland*

²*Center for Theoretical Physics, Polish Academy of Sciences, Al. Lotnikow 32/46, 02-668 Warsaw, Poland*

ABSTRACT

We study spectra produced by weakly accreting black hole (BH) systems using the semi-analytic advection dominated accretion flow (ADAF) model and the general-relativistic magnetohydrodynamic (GRMHD) simulation. We find significant differences between these two approaches related to a wider spread of the flow parameters as well as a much steeper radial distribution of the magnetic field in the latter. We apply these spectral models to the broad-band spectral energy distribution (SED) of the nucleus of M87 galaxy. The standard (in particular, one-dimensional) formulation of the ADAF model does not allow to explain it; previous claims that this model reproduces the observed SED suffer from an inaccurate treatment of the Compton process. The spectra based on GRMHD simulation are in a much better agreement with the observed data. In our GRMHD model, in which we assumed the BH spin $a = 0.9$, bulk of radiation observed between the millimeter and the X-ray range is produced in the disk area between 2 and 4 gravitational radii from the BH. In this solution, the synchrotron component easily reproduces the spectral data between the millimeter and the UV range, and the Compton component does not violate the X-rays constraints, for $\dot{M} \lesssim 0.01 M_{\odot}/\text{year}$ and a relatively strong magnetic field, with the plasma $\beta \sim 1$ in the region where radiation is produced. However, the Compton component cannot explain the observed X-ray spectrum. Instead, the X-ray spectrum can be reproduced by a high energy tail of the synchrotron spectrum if electrons have a hybrid energy distribution with a $\sim 5\%$ nonthermal component.

Keywords: Active galactic nuclei – Radio cores – Low-luminosity active galactic nuclei – High energy astrophysics – Astrophysical black holes – Accretion

1. INTRODUCTION

Accretion flows powering the black hole (BH) X-ray binaries and active galactic nuclei (AGNs) occur in two main forms. At high luminosities, $\gtrsim 1\%$ of the Eddington luminosity, they form optically thick disks, radiating away $\sim 10\%$ of the rest-mass energy of accreting matter. At lower luminosities, the characteristic spectral features of an optically thick material disappear and the radiative efficiency drops significantly. The prevailing interpretation of these low-luminosity BH systems invokes a tenuous, two-temperature mode of accretion (Shapiro et al. 1976; Narayan & Yi 1995; Abramowicz et al. 1996), commonly referred to as an advection-dominated accretion flow (ADAF). The theory of these optically thin flows predicts that a central black hole should be surrounded by a hot, magnetized

plasma emitting considerable amounts of synchrotron radiation. These predictions are markedly consistent with the Event Horizon Telescope (EHT) images of M87* and Sgr A* (Event Horizon Telescope Collaboration et al. 2019a, 2022), in which the observed brightness at the EHT frequency, 230 GHz, can be naturally explained by synchrotron emission. Also, the morphology seen in these images indicates that the near-horizon region is directly observed without obscuration, thus confirming the optical thinness of the radiating material. Despite these theoretical and observational advances, we still rather poorly understand the details of the physical mechanisms involved, or even whether the observed radiation is produced in the accretion disk or in the inner jet. On the theoretical side, a major source of uncertainty in the quantitative predictions of the expected radiation is the lack of understanding of the physical processes involved in the transfer of the dissipated energy to the plasma particles, and hence how the plasma properties determine the heating efficiency of the radiating particles and how many of them are accelerated into a non-thermal population.

Much work has gone into developing semi-analytical ADAF models (e.g. Manmoto et al. 1997; Manmoto 2000; Xie & Yuan 2012; Niedźwiecki et al. 2012; Nemmen et al. 2014), following the original formulation by Shapiro et al. (1976) and Ichimaru (1977). These models typically involve a detailed treatment of the energy balance of electrons, including the relevant radiative cooling processes as well as advection of energy, compressive heating, and the phenomenological description of the heating efficiency (given by a parameter commonly denoted by δ). The ADAF model has been applied to many low luminosity BH systems, including M87* and Sgr A* which, being one of the best-studied AGNs emitting many orders of magnitude below the Eddington limit, have been one of the primary targets for the application of this model. In this work we revisit these models and we note that some simplifications underlying the analytical descriptions (in particular, the neglect of the multidimensional structure) are a major hindrance to the accurate estimation of spectral formation in the studied objects.

Magnetohydrodynamic (MHD) simulations of accretion driven by the magnetorotational instability (MRI; Balbus & Hawley 1991, 1998) enable a more sophisticated approach to studying flow dynamics and the involved MHD processes. Magnetic fields play an important role in transporting the angular momentum outwards, while the accretion flow drags the magnetic field towards the BH horizon. One of the key predictions of general-relativistic (GR) MHD simulations is that the black hole spin must play a key role in powering relativistic jets, while the poloidal magnetic field loops are brought twisted by the frame-dragging effect (McKinney et al. 2012; Blandford et al. 2019). These jets are directly observed in about 10% of active galaxies, and one of the most prominent cases is that in M87* galaxy (Lu et al. 2023). A powerful jet is obtained when the magnetic flux on horizon overcomes the gravitational expulsion (Komissarov & McKinney 2007). However, when the magnetic flux exceeds some threshold value, being a function of the mass accretion rate, the backreaction of the BH magnetosphere acts in the opposite way. The term magnetically-arrested disk (MAD) has been introduced by Tchekhovskoy et al. (2011) and is frequently used to describe episodic modes of accretion, connected with flaring eruptions of magnetic flux and azimuthally distributed channels of gas entering the black hole (cf. James et al. 2022). The MAD accretion mode, observed in GRMHD simulations, was in fact predicted already in the calculations based on pseudo-Newtonian framework (Igumenshchev et al. 2003; Igumenshchev 2008) which shown that the magnetic fields are dynamically important and change the effectiveness of magneto-rotational instability. In this mode, the gas is pushed outward by magnetic tension, and the interchange

instability is the one which ultimately allows it to accrete through the BH horizon.

While the GRMHD models provide a thorough description of the flow structure, they typically neglect radiative cooling processes, and the treatment of the thermodynamics of the electrons is very cursory. A few radiative GRMHD simulation were carried out with self-consistent evolution of two-temperature plasma (e.g. Ryan et al. 2018; Chael et al. 2019) and the results were found to be strongly dependent on the assumptions concerning the electron heating. In most cases, however, GRMHD simulations evolve a single fluid whose internal energy is dominated by ions, while the temperature of the emitting electrons remains unconstrained and is then typically set manually for radiation calculations.

In this work we compare spectra predicted by the ADAF and the GRMHD models for accretion rates relevant for M87*, using the radiative-transfer code of Niedźwiecki (2005) and Niedźwiecki et al. (2012). Such a comparison has never been made before. We then attempt to reproduce the broadband spectral energy distribution (SED) observed in M87* using these spectral predictions. Specifically, we apply them to the radio to X-ray range data compiled by Prieto et al. (2016), and to the results of the quasi-simultaneous multi-wavelength campaign in 2017, reported by EHT MWL Science Working Group et al. (2021).

2. NOTATION

We use the usual dimensionless parameters, $r = R/R_g$, $a = J/(cR_gM)$ and $\dot{m} = \dot{M}/\dot{M}_{\text{Edd}}$, where $R_g = GM/c^2$ is the gravitational radius, J is the black hole angular momentum, $\dot{M}_{\text{Edd}} = L_{\text{Edd}}/(\epsilon c^2)$, $L_{\text{Edd}} \equiv 4\pi GMm_p c/\sigma_T$, and to define the Eddington rate we take the radiative efficiency to be $\epsilon = 0.1$.

For M87*, we assume the distance of $d = 16.7$ Mpc (Mei et al. 2007; Blakeslee et al. 2009), the source inclination, i.e. the viewing angle of the emitting region with respect to the line of sight, of $i = 17^\circ$ (corresponding to the angle between the approaching jet and the line of sight; Walker et al. 2018), and the black-hole mass of $M = 6.5 \times 10^9 M_\odot$, as adopted by the EHT Collaboration. This corresponds to $\dot{M}_{\text{Edd}} = 148 M_\odot \text{ year}^{-1}$.

The plasma β parameter is defined as the ratio of the gas to magnetic pressure, $\beta = P_{\text{gas}}/P_{\text{mag}}$, and the fraction of accretion power used to directly heat electrons is denoted by δ . The number density is denoted by n and the electron and ion temperature by T_e and T_p .

3. ADAF MODELS FOR M87*

We first apply a semi-analytic ADAF model `kerrflow` (Niedźwiecki et al. 2022). It strictly follows the standard ADAF formalism, which is based on transonic solutions of the hydrodynamical equations for the conservation of mass,

radial momentum, angular momentum, and separate energy equations for ions and electrons, using the phenomenological α -prescription for viscosity (e.g. Narayan et al. 1997; Manmoto et al. 1997; Gammie & Popham 1998; Manmoto 2000). The hydrodynamic description underlying these solutions is essentially one-dimensional, giving only the radial distribution of the flow parameters (which turns out to be a major oversimplification, as we find below). As in all ADAF solutions, the plasma is assumed to be described by radius-independent values of α and β parameters. Despite these simplifications, the analytic ADAF models have an important advantage of including a detailed description of the energy balance of the electrons, including the relevant radiative cooling processes (the synchrotron and bremsstrahlung emission and their Comptonization), and are therefore a good starting point for estimating the expected values of the electron temperature. These models have been used extensively for the description of black-hole systems and their comparison with models based on MHD simulations can be valuable.

The `kerrflow` model improves on some computational inaccuracies present in other similar models. The most important one concerns the treatment of Comptonization, for which `kerrflow` uses an accurate, GR Monte Carlo simulations, whereas in other published models it is computed using the approximate formulae of Coppi & Blandford (1990). The Coppi & Blandford (1990) approximation, however, is only accurate at nonrelativistic temperatures (as clearly stated in that paper) and underestimates the energy increase in a scattering at large T_e . Then, using it in ADAF models, which are typically characterised by relativistic T_e , results in inaccurate spectral predictions, to the extent dependent on the model parameters. At $\dot{m} \sim 10^{-3}$, when the Comptonization spectrum has a power-law shape, the approximation gives a far too soft spectrum. At $\dot{m} \lesssim 10^{-4}$, when individual scattering orders are seen in the spectrum, it strongly underestimates the energy at which the scattering bumps are seen. Furthermore, all previous models (with a notable exception of Li et al. 2009, referred to below) used a non-GR description (in particular, pseudo-Newtonian potential for hydrodynamics and neglect of GR transfer of radiation), which fails in the region where most of the accretion power is released and radiation is produced. In contrast, `kerrflow` is fully GR.

The effect of these two shortcomings on the spectrum is illustrated in Figure 1(a), where the `kerrflow` spectrum is compared with the one computed for the same parameters with a non-GR model `riaf-sed` (Nemmen et al. 2014). Both models predict very similar values of density and electron temperature (in `riaf-sed`, T_e is slightly larger) and then the large difference in the position of the scattering bumps seen in these spectra is entirely due to deficiencies of the Coppi & Blandford (1990) approach applied

in `riaf-sed`. The difference in the shape and amplitude of the synchrotron peak is due to the neglect of both the photon advection by the BH and the GR redshift in `riaf-sed`. We note, in particular, that all previous attempts to explain the M87* spectrum with ADAF models relied on fitting the first scattering bump to the optical/UV data and the second scattering bump to the X-ray data. The strong underprediction of the energies of these bumps by the applied Coppi & Blandford (1990) approximation makes these fitting attempts incorrect.

In its most general version, the ADAF model includes a number of free parameters and exploring the full parameter space is not feasible. We are most interested in estimating the suitable values of β , δ and \dot{m} , therefore, we allowed these parameters to vary, while keeping the others fixed. Specifically, we assumed the usual value of $\alpha = 0.3$ (as discussed by Niedźwiecki et al. 2022, there is a degeneracy between α , \dot{m} and δ). Furthermore, it has become customary to assume a local accretion rate decreasing inward as a power-law, $\dot{m} = \dot{m}_{\text{out}}(r/r_{\text{out}})^s$, at all radii down to the black-hole horizon, typically with $s \sim 0.5$. However, assuming such a $\dot{m}(r)$ profile in the inner flow may strongly affect the spectral properties (cf. Niedźwiecki et al. 2022), whereas there is no clear indication from MHD simulations that such a strong mass loss should occur in the innermost region (and actually it was not found in the simulations considered in Section 4) as well as the observational evidence of winds in systems with very low accretion rates concerns the radial scales of $r \sim 10^4 - 10^5$, e.g. in M87* (Park et al. 2019), but not at lower radii. Therefore, we assume that \dot{m} is constant in the region where the radiation is produced, i.e. we set $s = 0$. Then, we also assume $a = 0.95$, which value may be relevant for a system with a powerful jet.

We found that it is not possible to reproduce the full SED with the `kerrflow` model, therefore, we focused on modeling the millimeter part. We considered $1 \leq \beta \leq 9$ and $0.01 \leq \delta \leq 0.5$ and we found that the model best-matches the millimeter data for $\dot{m} \simeq 10^{-5}$ and $\delta \simeq 0.1$, and this result does not depend on the value of β . These solutions are represented by models A1 and A9 defined in Table 1, and the solid red line in Figure 1(a) shows the spectrum for model A1 with $\beta = 1$. We now compare this result with previous ADAF models for M87*. The early models for this object (Reynolds et al. 1996; Di Matteo et al. 2003) used the simplified self-similar description of Narayan & Yi (1995) and then more accurate ones, based on transonic hydrodynamic solutions, were presented by Li et al. (2009), Nemmen et al. (2014) and Bandyopadhyay et al. (2019). Of these, only the model of Li et al. (2009) is GR and then it is closest to `kerrflow`. Indeed, we note a very good agreement between parameters and synchrotron spectra of these two models, with the main difference concerning ~ 3 times larger




Figure 1. Comparison of the observed SED of M87 core with spectra predicted by (a) the analytic ADAF model, and (b,c) the model based on the GRMHD simulation. In all panels the green points show the highest spatial resolution SED in the quiescent phase of M87* given in table 4 of Prieto et al. (2016), and the blue points show results of the quasi-simultaneous multi-wavelength campaign in 2017, reported by EHT MWL Science Working Group et al. (2021). See Table 1 for definition of the models. All models assume $M = 6.5 \times 10^9 M_\odot$ and $i = 17^\circ$; all spectra except for the dotted magenta line in (c) assume a purely thermal distribution of electron energy. In panel (a) the solid red line shows the spectrum of the *kerrflow* model A1; the dashed black line shows the spectrum of *riaf-sed* (Nemmen et al. 2014) for the same $\dot{m} = 10^{-5}$, $\delta = 0.1$ and $\beta = 1$. The difference between the model spectra is primarily due to the neglect of GR effects and the inaccurate treatment of Comptonization in *riaf-sed*. Panel (b) illustrates how the GR transfer affects the spectrum in the GRMHD model B; the synchrotron and Compton components are shown separately. The dashed black line shows the spectrum in the flat space-time (i.e. neglecting the GR transfer) and the solid red line shows the fully-GR spectrum. Panel (c) shows how the Compton component is affected when the synchrotron component is matched to the data by increasing either the magnetic field (the solid red line, model D) or the density and the electron temperature (the dashed black line, model C). The synchrotron and Compton components are shown separately. The dotted magenta line shows the synchrotron spectrum in model D with a hybrid electron energy distribution, including a 5% energy content of non-thermal electrons in the form of a high-energy power-law tail with an index equal to 3.2.

ADAF model	$\dot{m} [\times 10^{-5}]$	β	δ
A1	1	1	0.1
A9	1	9	0.1
GRMHD model	$\dot{m} [\times 10^{-5}]$	b_{mag}	T_e/T_p
A	3.8	1	0.04
B	25	0.4	0.04
C	35	0.34	0.1
D	7.6	3.9	0.04

Table 1. Parameters of the models. All models assume $M = 6.5 \times 10^9 M_\odot$; the spin parameter is $a = 0.9$ in the GRMHD and $a = 0.95$ in the ADAF model.

\dot{m} and correspondingly stronger synchrotron peak in Li et al. (2009); this larger \dot{m} likely results from the (incorrect, see above) attempt to reproduce the data above 10^{14} Hz with the Compton component, and the scarcity of radio data available at that time. Nemmen et al. (2014) and Bandyopadhyay et al. (2019) found parameters strongly differing from ours, in particular a large $\dot{m} \simeq 2 \times 10^{-4}$ in the inner part, which may reflect large computational differences between the models and again their results relied on matching the incorrectly determined Compton spectrum to the data.

4. MODEL BASED ON GRMHD SIMULATIONS

4.1. Simulations and post-processing

In this Section we analyze the results of 3D general-relativistic ideal GRMHD numerical simulations of accretion flows onto a magnetically saturated Kerr BH. For the analysis, we choose the AGN-HS100 simulation presented in Janiuk & James (2022). Our run was performed with the code *HARM_COOL* (presented in e.g. Sapountzis & Janiuk 2019, for a 2D case). The MHD equations for the continuity and energy momentum are brought in the code in conservative form in general relativity, and account for the dynamical evolution of magnetized plasma in the gravitational potential of a spinning BH.

Our simulation is performed on a fixed Kerr background metric, with proper coordinate transformation between spherical (Boyer-Lindquist) and code (Modified Kerr-Schild) coordinates, to avoid the flow penetrate the black hole horizon. The inner radius of the grid is located at $r_{\text{in}} = 1.01 r_{\text{hor}}$, and the outer radius is fixed at $10^5 R_g$. The number of points in r - θ - ϕ directions is $288 \times 256 \times 256$, respectively. Thanks to adopting a progressively sparser grid above $r = 400$, our simulations are well resolved close the black hole horizon.

The initial conditions are imposed by the equilibrium torus solution Fishbone & Moncrief (1976), located initially at $r_{\text{in}} = 12$ where the cusp forms, and parameterized with the radius of pressure maximum $r_{\text{max}} = 25$. BH spin $a = 0.9$ is adopted. To start accretion, the initial torus is embedded

in a poloidal magnetic field, prescribed by the vector potential of the form $A_\phi = r^5(\rho - \rho_{\text{max}})/0.2$, therefore, the poloidal loops are centered at the radius r_{max} . The field is initially set to zero outside the torus. Factor r^5 enables bringing more magnetic flux from large distances and formation of the MAD state. In addition, the initial distribution of internal energy is perturbed azimuthally, by a random fluctuation, of the form $u = u_0(0.95 + 0.1\text{RAND})$ where RAND is a random number uniformly distributed between 0 and 1.

The GRMHD code works in geometrized units and has no intrinsic mass scale and then to apply it to M87* we need to introduce specific scalings to convert the code variables to cgs units. This is done after adopting the BH mass (i.e. $M = 6.5 \times 10^9 M_\odot$ for M87*) and the resulting scalings for gravitational length and time, $R_g = GM/c^2$ and $\mathcal{T} = R_g/c$. The factor scaling the density to the cgs units, \mathcal{M} , is then related to the assumed accretion rate by $\mathcal{M} = \dot{M}\mathcal{T}/R_g^3$. We have found that it is not possible to reproduce the observed SED by adjusting only the \dot{m} parameter, and therefore we allow small variations in the scaling of the magnetic field by introducing a free parameter (close to unity), b_{mag} , so that the scaling factor for the magnetic field is $\mathcal{B} = b_{\text{mag}}(8\pi\mathcal{M}c^2)^{1/2}$.

Furthermore, the MHD simulation gives only the total internal energy density, u_{gas} , and not the distinct electron and ion temperatures. A common practice in radiation calculations in such models, the so-called post-processing, is to set T_e after running the simulation, as a function of local plasma parameters. As we discuss below, the popular parameterization used for this, introduced by Mościbrodzka et al. (2016) and referred to as the R - β model, may yield unrealistically high values of T_e , as it ignores limitations due to cooling efficiency. Therefore, we do not follow it. Instead, we set the electron temperature based on the results obtained in our ADAF model, which includes a detailed treatment of the electron energy balance and then gives a reasonable indication of the expected magnitude of T_e under relevant physical conditions.

Figure 2(a-c) shows the radial profiles of B , T_p and n in the *kerrflow* models A1 and A9. The lower β model A1 is characterized by a higher B (which is a trivial consequence of the assumed higher magnetization), lower T_p (because a larger fraction of the accretion power is used to build up the magnetic field instead of heating the flow), and higher n (because a more magnetized flow is geometrically thinner). Also shown in Figure 2(a-c) are the radial profiles of quantities representing the GRMHD solution, for which the average $\bar{B}(r)$ and $\bar{T}_p(r)$ are found by density-weighted averaging ($\bar{B} = \int BndV / \int ndV$ and analogously for T_p) over narrow, spherical shells, and $\bar{n}(r)$ by averaging ($\bar{n} = \int ndV / \int dV$) over narrow rings representing the mid-plane ($z \leq 0.1r$, where z is the distance from the mid-plane). The major dif-

ference between the GRMHD simulation and the analytic ADAF calculation concerns B , which for the former has a much steeper profile corresponding to a gradual build-up of the magnetic field, in contrast to comparatively flatter profiles resulting from the assumption of constant β in the ADAF model. The GRMHD solution also has a slightly flatter profile of n , due to the weak loss of accreting matter to the wind, and T_p decreasing from a large- β to a small- β value toward smaller r , but with a relative increase close to the event horizon due to the additional heating by rotational energy extraction from the BH (neglected in the ADAF model). Nevertheless, except for the difference in the gradient of B , we see a very good agreement of the parameters characterising the ADAF and MHD solutions. We can therefore expect that both solutions are also characterised by similar values of T_e .

Figure 2d shows the T_e profiles in our models. The temperature ratio in the `kerrflow` solutions A1 and A9 is in the range $T_e/T_p \simeq (0.02 - 0.07)$ and, more importantly, in the innermost part $T_e \simeq 10^{11}$ K (within a factor $\lesssim 2$). This innermost T_e is rather weakly affected by the value of δ , see Figure 3c. Guided by these results, in most of our GRMHD radiative models we set $T_e = 0.04T_p$ (we checked how the increase of T_e affects the spectrum and we found that this leads to a too strong Compton emission, as illustrated in model C defined below). For the proton temperature we assume that electrons give a negligible contribution to u_{gas} , and then the simulated values of u_{gas} are used to determine T_p . With our adopted prescription for the local T_e (i.e. $= 0.04T_p$), the average $\bar{T}_e(r)$ is close to the value corresponding to the energy balance in our ADAF calculations, as shown in Figure 2d.

We now briefly comment on the R - β model, in which the ratio between the ion and electron temperatures is given by the local plasma β ,

$$R \equiv \frac{T_p}{T_e} = R_{\text{high}} \frac{\beta^2}{1 + \beta^2} + R_{\text{low}} \frac{1}{1 + \beta^2}, \quad (1)$$

which then allows to relate T_e to u_{gas} by

$$T_e = \frac{2m_p u_{\text{gas}}}{3\rho k(2 + R)}, \quad (2)$$

see e.g. [Event Horizon Telescope Collaboration et al. \(2019b\)](#). In the above R_{high} and R_{low} are the free parameters of the model and give the temperature ratios at high and low β , respectively. The R - β prescription tentatively relates the temperature distribution to the efficiency of the electron heating, which increases with plasma magnetisation, as often seen in simulations of electron heating in turbulent or reconnecting plasmas (e.g. [Howes 2010](#); [Rowan et al. 2017](#)).

Figure 3a shows the radial profile of the density-weighted average \bar{T}_e in our GRMHD solution for $R_{\text{high}} = 80$ and $R_{\text{low}} = 1$, which are typical values used in applications of the R - β model. We see that the average \bar{T}_e approaches

$\sim 10^{12}$ K at small r (furthermore, in some low- β regions it approaches $\sim 10^{13}$ K, see the shaded region). This value appears much too high, in particular, an order of magnitude above the level of $\sim 10^{11}$ K set by the energy balance of electrons in our ADAF solutions with the same values of n and B . We note that it would be impossible to increase this balance temperature to $\sim 10^{12}$ K, because the synchrotron cooling rate rapidly increases with temperature, approximately $\propto T_e^5$, and then keeping electrons at the average $T_e \sim 10^{12}$ K would require a heating rate orders of magnitude larger than the available accretion power. The problem of too high T_e in the R - β model possibly occurs only in MAD solutions, where the plasma magnetization of the inner part of the flow is large, implying a small β . The T_e in that region can be reduced by increasing R_{low} . In our GRMHD solution, using $R_{\text{low}} = 10$ we get $\bar{T}_e \simeq 4 \times 10^{11}$ K, which appears still too high.

The above issue is also apparent in figure 17 of [Event Horizon Telescope Collaboration et al. \(2021\)](#), comparing the temperature-ratio distribution from the radiative MHD simulations and from the R - β postprocessing. In the jet region, both approaches give similar values of $T_e \simeq T_p$. However, in the innermost few R_g of the disk (which is the region where the production of radiation observed at millimeter and shorter wavelengths takes place in our GRMHD model), the R - β also gives a very high $T_e \sim T_p$, whereas the radiative simulations yield a much lower $T_e \simeq 0.04T_p$, which notably matches the temperature ratio given by the energy balance in our ADAF calculations.

4.2. SED modeling

We now apply the GRMHD model to the SED observed in M87*. We use the density, the electron temperature, and the components of the velocity and magnetic fields from the GRMHD simulation to compute the model spectrum, using the same GR Monte Carlo code as in the `kerrflow` model. In it, we generate a large number of synchrotron photons according to the local values of plasma parameters and we track their motion in the Kerr metric, subject to Compton scattering competing with synchrotron self-absorption, until they cross the event horizon, escape to a distant observer, or get absorbed.

It is instructive to compare the formation of spectra in the GRMHD and ADAF models. The strongest radiation production occurs in the innermost part of the flow. It might therefore seem that the ADAF and GRMHD models which have similar values of B , n and T_e in this region, also possess similar radiative properties. However, it turns out that the GRMHD model predicts a much weaker emission than ADAF. For example the GRMHD model A produces the synchrotron component almost by two orders of magnitude weaker than the ADAF models A1 and A9, despite having




Figure 2. Comparison of the radial profiles of (a) T_p , (b) B , (c) n , and (d) T_e for the `kerrflow` models A1 (the dashed green lines) and A9 (the dotted blue lines) and the GRMHD model A (the red solid lines), see Table 1. In the GRMHD model, T_p , B , and T_e are found by spherical, density-weighted averaging (over narrow shells with $\Delta r = 0.1r$), and n by averaging over narrow rings (again with $\Delta r = 0.1r$) with $z \leq 0.1r$.

very similar parameters of the innermost emission region (see Figure 2). The reason for this is the much steeper B profile in the GRMHD solution, which implies that the synchrotron radiation emissivity, $\propto B^2$, is much more concentrated near the event horizon and thus subjected to much stronger reduction by GR effects. In our GRMHD calculation, the *observed* radiation is reduced by over a factor of 10 (see Figure 1b and discussion below), as compared to a factor of ~ 3 in the ADADF model. Furthermore, the much faster decline of B means that the amount of *produced* radiation is significantly smaller in the GRMHD model, because the area of the disk where the synchrotron radiation is efficiently produced is smaller than in the ADADF model. Specifically, in our MHD models almost all radiation observed above the turnover frequency, ν_t (below which the emission is strongly self-absorbed), is produced at $r \lesssim 4$, whereas in the ADADF models a significant contribution comes from the region extending out to $r \sim 20$.

Another difference is related to the flow structure, which in the GRMHD solution is characterized by a range of parameter values at a given radius, related to the fluctuating nature of this solution. Such fluctuations are completely neglected in the analytical ADADF solutions, where at each r the synchrotron emission is produced at a single value of B and T_e and the resulting synchrotron spectrum has the shape of a relatively narrow bump peaking in IR, as seen in Figure 1a. Figure 3b shows the range of T_e at which the synchrotron emission is effectively produced (as estimated based on the local emissivity, $\propto T_e^2 B^2 n$) in the GRMHD solution. We see that at small r some regions have T_e an order of magnitude larger than the average T_e . Such a broad temperature spread gives rise to a broad synchrotron spectrum, extending from the millimeter to optical/UV band at an approximately constant level in νF_ν units, as seen in Figures 1bc.

Figure 1b illustrates how the GR transfer of radiation affects the observed spectrum by comparing the spectra com-




Figure 3. In panels (a) and (b), the solid lines show the radial profile of electron temperature obtained by density-weighted spherical averaging in our GRMHD solution, for two prescriptions relating T_e to T_p . (a) The $R - \beta$ relation, equations (1) and (2), with $R_{\text{high}} = 80$, $R_{\text{low}} = 1$, and β computed for model A. (b) $T_e = 0.04T_p$, as assumed in most of our spectral models. The shaded regions show the range of T_e at which a significant contribution to the synchrotron emission occurs. In panel (c) the red solid line shows the T_e profile for the GRMHD model with $T_e = 0.04T_p$, the same as in panel (b), and the black lines show the T_e profiles in `kerflow`; the dashed line is for model A1 and the dotted line is for the same parameters except for $\delta = 0.3$.

puted in model B neglecting (dashed line) and including (solid line) these transfer effects. As noted above, the synchrotron emission is strongly concentrated close to the BH and hence the observed spectrum is strongly reduced due to the bending of photon trajectories toward the BH and the capture of photons below the event horizon. Furthermore, a significant effect of the GR redshift can be seen, shifting the turnover frequency below the EHT frequency, whereas in the flat space-time it would be at $\simeq 400$ Hz.

In model A the synchrotron component is weak and then it strongly underpredicts the observed SED, in particular at the EHT frequency. The synchrotron emission can be enhanced by increasing the density, the electron temperature, and/or the magnetic field. However, a significant increase in the first two leads to an over-prediction of the X-ray data by the Compton component, whose amplitude relative to the synchrotron component increases with both density and temperature. The Thomson depth in the inner parts of the flow is small, $\sim 10^{-4}$, but in the Compton scattering at $\sim 10^{11}$ K the average energy increase is larger than a factor of 10^3 , and then the luminosity of the Compton component is comparable to that of the synchrotron. If the increase in either density or temperature is too large, the Compton component will be too strong. We then note that the model can be matched to the data, in the sense that the synchrotron spectrum agrees with the millimeter to UV data and the Compton spectrum does not exceed the X-ray data, only for a relatively strong B , corresponding to $b_{\text{mag}} > 1$. These properties are illustrated in Figure 1c, which presents spectra in models C (dashed line) and D (solid line). Model C shows the combined effect of a large increase in n , to $\sim 10^6 \text{ cm}^{-3}$ at $r \sim 2$, and an increase in T_e to $0.1T_p$. We see that whereas the synchrotron spectrum reproduces very well the lower energy data, the Compton emission exceeds the X-ray data by almost two or-




Figure 4. The dots represent the contribution from a given radial bin (for logarithmic radial grid) to the flux observed at different frequencies at $i = 17^\circ$ in model D, normalized by the flux measured by VLBA at 4.3×10^{10} Hz, GMVA at 8.6×10^{10} Hz, EHT at 2.3×10^{11} Hz, and HST at 5.2×10^{14} Hz, as reported by EHT MWL Science Working Group et al. (2021).

ders of magnitude. Instead, the increase of B in model D by $b_{\text{mag}} \simeq 4$ (yielding $\beta \sim 1$ in the emitting region) allows to fit the data between $\sim 10^{11}$ and 10^{15} Hz without exceeding the X-ray limit.

The bulk of radiation observed above $\nu_t \simeq 2 \times 10^{11}$ Hz is produced in the disk region (i.e. near the equatorial plane) between $\simeq 2$ and $4R_g$ (see Figure 4). Radiation produced within $2R_g$ is strongly reduced by GR transfer; we note that advection of photons into the BH is enhanced by the radial

inflow of the radiating material. A small amount of radiation is produced beyond $4R_g$ due to the weakness of the magnetic field. At $\nu < \nu_t$, the intrinsic absorption shifts the strongest contributing region to larger radii.

Finally, all the spectra discussed above were computed for the thermal distribution of electrons. In this version of the model, the high-energy component can be reduced below the observed X-ray flux, as in model D, but it cannot reproduce the X-ray spectrum. If the energy distribution of electrons is hybrid, the observed X-rays may be produced by nonthermal synchrotron emission. The dotted line in Figure 1c shows the synchrotron spectrum in model D assuming such a hybrid distribution of electrons, with the nonthermal component in the form of a high-energy power-law tail containing 5% of energy of the total distribution. For the power-law index of the tail we assume $p = 3.2$, for which the photon index $\Gamma = 2.1$ is roughly consistent with that observed. We only note here an approximate agreement of such a model with the observed X-ray spectrum, which was derived in [EHT MWL Science Working Group et al. \(2021\)](#) assuming that the *XMM* and *NuSTAR* measurement can be represented by a single power-law. Formal spectral fitting of the X-ray data, performed in the detector space, is not feasible as our computational model is too time-consuming.

5. SUMMARY AND DISCUSSION

We compared radiative properties of two commonly used models of low-luminosity BH systems, the one based on the GRMHD simulation and the one using a semi-analytic ADAF description of the two-temperature accretion. The latter gives a similar distribution of the parameters of the flow to the distribution of the averaged parameters in the former, except for the magnetic field, which in the GRMHD solution has a much steeper radial profile corresponding to the amplification of B by the MRI, whereas the constant- β structure typically assumed for ADAFs gives a much slower outward gradient of B . The rapid decline of B with increasing r in the GRMHD solution results in a much lower radiative efficiency, since it significantly limits the extent of the area where synchrotron radiation is produced and places this area in a region of strong reduction by GR effects.

Another difference between the two models is related to the neglect of the multidimensional structure of the flow in the ADAF description. This significantly affects the spectral shape, which in the ADAF model is represented by a relatively narrow synchrotron bump followed by similarly narrow scattering bumps. In contrast, the spread of parameters corresponding to the 3D GRMHD solution gives broad synchrotron as well as Compton components extending over many decades of photon energy.

We computed spectra for accretion rates and BH mass relevant for M 87*, assuming also a high value of the BH spin,

$a \sim 0.9$, which may favor the production of the jet observed in this object. The ADAF model can explain the millimeter part of the spectrum observed in M87, but does not agree with the spectral data at other wavelengths. We note that previous ADAF models of this object are significantly affected by order-of-magnitude inaccuracies related to the neglect of the GR transfer of radiation as well as an oversimplified description of the Compton component.

The spectra based on the GRMHD solution are in a much better agreement with the observed SED. The low radiative efficiency of this model can be compensated by setting a relatively large B , corresponding to $\beta \sim 1$ at small r . An alternative way to increase the luminosity by increasing either the accretion rate (and hence the density) or the electron temperature results in a significant overprediction of the X-ray flux by the Compton component.

The obvious caveat to our spectral modelling is that the SED data are obtained by observations with different angular resolution, which in the case of M87 translates to the spatial scale of $\sim 10R_g$ at the EHT frequency (i.e. 230 GHz) and $\gtrsim 10^5 R_g$ at higher frequencies, and therefore the optical, UV and X-ray radiation may be produced in a spatially distinct region from the emission region detected by EHT. Nevertheless, the synchrotron component produced by predominantly thermal electrons in our GRMHD model is a good match for the spectral data between the millimeter and the UV range.

At $\nu \lesssim 100$ GHz the synchrotron emission is strongly self-absorbed, which shifts the location of the effective flux maximum to larger radii, increasing with decreasing ν (Figure 4). This is consistent with the difference of sizes of the ring-like structures in the radio core of M87* resolved at 86 GHz and 230 GHz ([Lu et al. 2023](#)). Due to the strong self-absorption, the synchrotron emission from the inner accretion flow cannot explain the total flux measured at the low frequencies. This low-frequency radiation must then be generated, at least in part, on larger spatial scales, as indeed suggested by the core shifts observed in M87 at 86 GHz and lower frequencies (e.g. [EHT MWL Science Working Group et al. 2021](#)). The contribution from the jet region is negligible in our calculations (this is also consistent with the assessment in [Lu et al. 2023](#), that the core emission comes from the accretion flow). Some relevant physical processes, in particular internal shocks which can dissipate the bulk kinetic energy and channel some of it into particle acceleration (e.g. [Rees 1978](#); [Spada et al. 2001](#)), are not handled accurately in current GRMHD codes, and therefore we have not attempted to model the jet emission that could explain the radio data.

The Compton component predicted in the GRMHD model is too hard to fit the data in the X-ray range. Nevertheless, comparison of this component with X-ray data imposes important constraints on the model and indicates that the emitting region must be characterized by a relatively large $\beta \sim 1$.

The observed X-ray emission may be produced within the inner few R_g by nonthermal synchrotron emission of a hybrid electron plasma with a high energy tail with the power-law index $\simeq 3.2$, which implies the acceleration index of $\simeq 2.2$ (taking into account the steepening of the electron distribution by synchrotron cooling).

The maximum contribution to radiation observed at 230 Hz in our best-matching solution (model D) comes from $\simeq 3R_g$, which region is characterised by the average $B \simeq 25$ G, $T_e \simeq 6 \times 10^{10}$ K and $n \simeq 1.5 \times 10^5$ cm $^{-3}$. These values are consistent with parameters of the emission region estimated in [Event Horizon Telescope Collaboration et al. \(2021\)](#) using a one-zone model, although we note that such one-zone estimations are significantly affected by the neglect of the GR transfer effects. On the other hand, the accretion rate of our best-matching model, $\dot{M} \simeq 0.011 M_\odot$ year $^{-1}$, is larger than $\dot{M} \lesssim 0.002 M_\odot$ year $^{-1}$ assessed in the GRMHD models applied by [Event Horizon Telescope Collaboration et al. \(2021\)](#) to explain the EHT observation. These models, however, assumed the $R - \beta$ prescription for the electron temperature, which strongly affects the comparison with our results. We note that it may be risky to base radiative computations on the $R - \beta$ relation without checking that T_e is within physical limits.

Our preferred value of $\dot{M} \simeq 0.01 M_\odot$ year $^{-1}$ is a factor of 10 lower than the Bondi accretion rate ([Di Matteo et al. 2003](#)). This is sufficient to produce jet with a power on the order of $\sim 3 \times 10^{43}$ erg s $^{-1}$, however exceeds the recent estimates by eg. [Punsly \(2023\)](#).

In our GRMHD solution the synchrotron radiation at $\nu \gtrsim 10^{14}$ Hz is produced by electrons with $T_e \sim 10^{12}$ K. The re-

gions with such a large T_e represent a very small fraction of the total flow volume, but their high emissivity makes them dominate the high-energy spectrum. This high emissivity also implies a high cooling rate, which then requires enhanced heating to maintain the high temperature. The density in these high- T regions is an order of magnitude lower, and the plasma magnetization $\sigma \propto B^2/n$ is similarly larger, than average in the disk. Then, electrons in these regions may be indeed more efficiently heated if the heating is due to magnetic reconnection (e.g. [Rowan et al. 2017](#)).

Finally, we note that radiative GRMHD simulations of [Chael et al. \(2019\)](#) using prescriptions for electron heating based on either a Landau-damped turbulent cascade ([Howes 2010](#)) or magnetic reconnection ([Rowan et al. 2017](#)) give similar to our model parameters in the disk, in particular $T_e \sim 10^{11}$ K. However, those simulations also give a much higher $T_e \sim 10^{12}$ K in the strongly magnetized regions in the jet, and their spectra are fully dominated by these regions. Then, these spectra depend on an arbitrary cut in σ more than on physical parameters (as shown in figure 17 in [Chael et al. 2019](#)).

ACKNOWLEDGEMENTS

We acknowledge support from the Polish National Science Center under grants 2023/48/Q/ST9/00138 and 2023/50/A/ST9/00527. We gratefully acknowledge Polish high-performance computing infrastructure PLGrid (HPC Center: ACK Cyfronet AGH) for providing computer facilities and support within computational grant no. PLG/2024/017013.

REFERENCES

- Abramowicz, M. A., Chen, X. M., Granath, M., & Lasota, J. P. 1996, *ApJ*, 471, 762, doi: [10.1086/178004](#)
- Balbus, S. A., & Hawley, J. F. 1991, *ApJ*, 376, 214, doi: [10.1086/170270](#)
- . 1998, *Reviews of Modern Physics*, 70, 1, doi: [10.1103/RevModPhys.70.1](#)
- Bandyopadhyay, B., Xie, F.-G., Nagar, N. M., et al. 2019, *MNRAS*, 490, 4606, doi: [10.1093/mnras/stz2874](#)
- Blakeslee, J. P., Jordán, A., Mei, S., et al. 2009, *ApJ*, 694, 556, doi: [10.1088/0004-637X/694/1/556](#)
- Blandford, R., Meier, D., & Readhead, A. 2019, *ARA&A*, 57, 467, doi: [10.1146/annurev-astro-081817-051948](#)
- Chael, A., Narayan, R., & Johnson, M. D. 2019, *MNRAS*, 486, 2873, doi: [10.1093/mnras/stz988](#)
- Coppi, P. S., & Blandford, R. D. 1990, *MNRAS*, 245, 453, doi: [10.1093/mnras/245.3.453](#)
- Di Matteo, T., Allen, S. W., Fabian, A. C., Wilson, A. S., & Young, A. J. 2003, *ApJ*, 582, 133, doi: [10.1086/344504](#)
- EHT MWL Science Working Group, Algaba, J. C., Anczarski, J., et al. 2021, *ApJL*, 911, L11, doi: [10.3847/2041-8213/abef71](#)
- Event Horizon Telescope Collaboration, Akiyama, K., Alberdi, A., et al. 2019a, *ApJL*, 875, L1, doi: [10.3847/2041-8213/ab0ec7](#)
- . 2019b, *ApJL*, 875, L5, doi: [10.3847/2041-8213/ab0f43](#)
- Event Horizon Telescope Collaboration, Akiyama, K., Algaba, J. C., et al. 2021, *ApJL*, 910, L13, doi: [10.3847/2041-8213/abe4de](#)
- Event Horizon Telescope Collaboration, Akiyama, K., Alberdi, A., et al. 2022, *ApJL*, 930, L12, doi: [10.3847/2041-8213/ac6674](#)
- Fishbone, L. G., & Moncrief, V. 1976, *ApJ*, 207, 962, doi: [10.1086/154565](#)
- Gammie, C. F., & Popham, R. 1998, *ApJ*, 498, 313, doi: [10.1086/305521](#)

- Howes, G. G. 2010, MNRAS, 409, L104, doi: [10.1111/j.1745-3933.2010.00958.x](https://doi.org/10.1111/j.1745-3933.2010.00958.x)
- Ichimaru, S. 1977, ApJ, 214, 840, doi: [10.1086/155314](https://doi.org/10.1086/155314)
- Igumenshchev, I. V. 2008, ApJ, 677, 317, doi: [10.1086/529025](https://doi.org/10.1086/529025)
- Igumenshchev, I. V., Narayan, R., & Abramowicz, M. A. 2003, ApJ, 592, 1042, doi: [10.1086/375769](https://doi.org/10.1086/375769)
- James, B., Janiuk, A., & Nouri, F. H. 2022, ApJ, 935, 176, doi: [10.3847/1538-4357/ac81b7](https://doi.org/10.3847/1538-4357/ac81b7)
- Janiuk, A., & James, B. 2022, A&A, 668, A66, doi: [10.1051/0004-6361/202244196](https://doi.org/10.1051/0004-6361/202244196)
- Komissarov, S. S., & McKinney, J. C. 2007, MNRAS, 377, L49, doi: [10.1111/j.1745-3933.2007.00301.x](https://doi.org/10.1111/j.1745-3933.2007.00301.x)
- Li, Y.-R., Yuan, Y.-F., Wang, J.-M., Wang, J.-C., & Zhang, S. 2009, ApJ, 699, 513, doi: [10.1088/0004-637X/699/1/513](https://doi.org/10.1088/0004-637X/699/1/513)
- Lu, R.-S., Asada, K., Krichbaum, T. P., et al. 2023, Nature, 616, 686, doi: [10.1038/s41586-023-05843-w](https://doi.org/10.1038/s41586-023-05843-w)
- Manmoto, T. 2000, ApJ, 534, 734, doi: [10.1086/308768](https://doi.org/10.1086/308768)
- Manmoto, T., Mineshige, S., & Kusunose, M. 1997, ApJ, 489, 791, doi: [10.1086/304817](https://doi.org/10.1086/304817)
- McKinney, J. C., Tchekhovskoy, A., & Blandford, R. D. 2012, MNRAS, 423, 3083, doi: [10.1111/j.1365-2966.2012.21074.x](https://doi.org/10.1111/j.1365-2966.2012.21074.x)
- Mei, S., Blakeslee, J. P., Côté, P., et al. 2007, ApJ, 655, 144, doi: [10.1086/509598](https://doi.org/10.1086/509598)
- Mościbrodzka, M., Falcke, H., & Shiokawa, H. 2016, A&A, 586, A38, doi: [10.1051/0004-6361/201526630](https://doi.org/10.1051/0004-6361/201526630)
- Narayan, R., Kato, S., & Honma, F. 1997, ApJ, 476, 49, doi: [10.1086/303591](https://doi.org/10.1086/303591)
- Narayan, R., & Yi, I. 1995, ApJ, 452, 710, doi: [10.1086/176343](https://doi.org/10.1086/176343)
- Nemmen, R. S., Storchi-Bergmann, T., & Eracleous, M. 2014, MNRAS, 438, 2804, doi: [10.1093/mnras/stt2388](https://doi.org/10.1093/mnras/stt2388)
- Niedźwiecki, A. 2005, MNRAS, 356, 913, doi: [10.1111/j.1365-2966.2004.08557.x](https://doi.org/10.1111/j.1365-2966.2004.08557.x)
- Niedźwiecki, A., Szanecki, M., Zdziarski, A. A., & Xie, F.-G. 2022, ApJ, 931, 167, doi: [10.3847/1538-4357/ac6c8b](https://doi.org/10.3847/1538-4357/ac6c8b)
- Niedźwiecki, A., Xie, F.-G., & Zdziarski, A. A. 2012, MNRAS, 420, 1195, doi: [10.1111/j.1365-2966.2011.20106.x](https://doi.org/10.1111/j.1365-2966.2011.20106.x)
- Park, J., Hada, K., Kino, M., et al. 2019, ApJ, 871, 257, doi: [10.3847/1538-4357/aaf9a9](https://doi.org/10.3847/1538-4357/aaf9a9)
- Prieto, M. A., Fernández-Ontiveros, J. A., Markoff, S., Espada, D., & González-Martín, O. 2016, MNRAS, 457, 3801, doi: [10.1093/mnras/stw166](https://doi.org/10.1093/mnras/stw166)
- Punsly, B. 2023, A&A, 677, A180, doi: [10.1051/0004-6361/202346831](https://doi.org/10.1051/0004-6361/202346831)
- Rees, M. J. 1978, MNRAS, 184, 61P, doi: [10.1093/mnras/184.1.61P](https://doi.org/10.1093/mnras/184.1.61P)
- Reynolds, C. S., Di Matteo, T., Fabian, A. C., Hwang, U., & Canizares, C. R. 1996, MNRAS, 283, L111, doi: [10.1093/mnras/283.4.L111](https://doi.org/10.1093/mnras/283.4.L111)
- Rowan, M. E., Sironi, L., & Narayan, R. 2017, ApJ, 850, 29, doi: [10.3847/1538-4357/aa9380](https://doi.org/10.3847/1538-4357/aa9380)
- Ryan, B. R., Ressler, S. M., Dolence, J. C., Gammie, C., & Quataert, E. 2018, ApJ, 864, 126, doi: [10.3847/1538-4357/aad73a](https://doi.org/10.3847/1538-4357/aad73a)
- Sapountzis, K., & Janiuk, A. 2019, ApJ, 873, 12, doi: [10.3847/1538-4357/ab0107](https://doi.org/10.3847/1538-4357/ab0107)
- Shapiro, S. L., Lightman, A. P., & Eardley, D. M. 1976, ApJ, 204, 187, doi: [10.1086/154162](https://doi.org/10.1086/154162)
- Spada, M., Ghisellini, G., Lazzati, D., & Celotti, A. 2001, MNRAS, 325, 1559, doi: [10.1046/j.1365-8711.2001.04557.x](https://doi.org/10.1046/j.1365-8711.2001.04557.x)
- Tchekhovskoy, A., Narayan, R., & McKinney, J. C. 2011, MNRAS, 418, L79, doi: [10.1111/j.1745-3933.2011.01147.x](https://doi.org/10.1111/j.1745-3933.2011.01147.x)
- Walker, R. C., Hardee, P. E., Davies, F. B., Ly, C., & Junor, W. 2018, ApJ, 855, 128, doi: [10.3847/1538-4357/aaafcc](https://doi.org/10.3847/1538-4357/aaafcc)
- Xie, F.-G., & Yuan, F. 2012, MNRAS, 427, 1580, doi: [10.1111/j.1365-2966.2012.22030.x](https://doi.org/10.1111/j.1365-2966.2012.22030.x)