

Stochastic First-Order Methods with Non-smooth and Non-Euclidean Proximal Terms for Nonconvex High-Dimensional Stochastic Optimization

Yue Xie · Jiawen Bi · Hongcheng Liu

Received: date / Accepted: date

Abstract When the nonconvex problem is complicated by stochasticity, the sample complexity of stochastic first-order methods may depend linearly on the problem dimension, which is undesirable for large-scale problems. In this work, we propose dimension-insensitive stochastic first-order methods (DISFOMs) to address nonconvex optimization with expected-valued objective function. Our algorithms allow for non-Euclidean and non-smooth distance functions as the proximal terms. Under mild assumptions, we show that DISFOM using minibatches to estimate the gradient enjoys sample complexity of $\mathcal{O}((\log d)/\epsilon^4)$ to obtain an ϵ -stationary point. Furthermore, we prove that DISFOM employing variance reduction can sharpen this bound to $\mathcal{O}((\log d)^{2/3}/\epsilon^{10/3})$, which perhaps leads to the best-known sample complexity result in terms of d . We provide two choices of the non-smooth distance functions, both of which allow for closed-form solutions to the proximal step. Numerical experiments are conducted to illustrate the dimension insensitive property of the proposed frameworks.

Keywords Nonconvex Large-scale Optimization · Stochastic First-order Methods · Non-Euclidean distances · Sample Complexity · Variance Reduction

Mathematics Subject Classification (2010) 90C06 · 90C15 · 90C26 · 90C30

Research is supported from HKU-IDS start-up fund; and Guangdong Province Fundamental and Applied Fundamental Research Regional Joint Fund, 2022B1515130009.

Yue Xie

Department of Mathematics and Musketeers Foundation Institute of Data Science, The University of Hong Kong, Pokfulam, Hong Kong.

E-mail: yxie21@hku.hk

Jiawen Bi

Department of Mathematics, The University of Hong Kong, Pokfulam, Hong Kong.

E-mail: jwbi@hku.hk

Hongcheng Liu

Department of Industrial and Systems Engineering, University of Florida

E-mail: hliu@ise.ufl.edu

1 Introduction

In this paper, we consider a stochastic optimization (SO) problem formulated as below:

$$\begin{aligned} \min_{\mathbf{x} \in \mathbb{R}^d} \quad & f(\mathbf{x}) = \mathbb{E}[F(\mathbf{x}, \zeta)] \\ \text{s.t.} \quad & \mathbf{x} \in X, \end{aligned} \quad (1)$$

where $f : \mathbb{R}^d \rightarrow \mathbb{R}$ and $F : \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}$. $(\Omega, \mathcal{F}, \mathbb{P})$ is the probability space. X is a closed and convex set. Denote by $\mathbf{x}^* \in \mathbb{R}^d$ an optimal solution to (1). We do not have explicit assumption on problem dimension d , but we are particularly interested in a large d . We assume that the expectation $\mathbb{E}[F(\mathbf{x}, \zeta)] = \int_{\Omega} F(\mathbf{x}, \zeta) d\mathbb{P}(\zeta)$ is well-defined and finite-valued for every $\mathbf{x} \in \mathbb{R}^d$ and that it is possible to generate independent and identically distributed realizations, ζ_1, ζ_2, \dots , of the random vector ζ . For this problem, we are particularly interested in solution algorithms when $f(\mathbf{x})$ is smooth but potentially nonconvex.

In view of stochasticity and potential nonconvexity of the problem, we aim to compute an stationary point with small expected value of first-order optimality residual.

Definition 1 For any $\epsilon \geq 0$ and nonnegative residual function $r(\cdot)$, a feasible random solution $\mathbf{x}_\epsilon^* \in X$ is said to be an ϵ -stationary solution/point if and only if

$$\mathbb{E}[r(\mathbf{x}_\epsilon^*)] \leq \epsilon. \quad (2)$$

Many stochastic first-order methods (S-FOMs) apply to smooth and nonconvex optimization problems in the form of (1). These methods date back to [26, 6, 28], and have been much studied (e.g., by [24, 25, 20, 16, 12, 27, 9, 11, 4, 5, 2, 19, 30, 22, 8, 18, 21, 29, 10, 23, 14], among others). In terms of the stochastic first-order oracle or sample, most existing S-FOMs incur analytical complexities that grow polynomially in dimensionality.

Indeed, for S-FOMs in smooth and nonconvex optimization, (See [12, 13, 23] and references therein.) Here σ^2 is defined as an upper bound on $\mathbb{E}[\|\nabla F(\mathbf{x}, \zeta) - \nabla f(\mathbf{x})\|^2]$. In the worst case, σ^2 is linear in d because

$$\sigma^2 \geq \max_{x \in X} \mathbb{E}[\|\nabla F(x, \zeta) - \nabla f(x)\|^2] = \max_{x \in X} \sum_{i=1}^d \mathbb{E}[(\nabla_i F(x, \zeta) - \nabla_i f(x))^2],$$

where the right-hand side is often $\mathcal{O}(d)$. This then leads to an analytical complexity in terms of SFO as following:

$$\mathcal{O}\left(\frac{d}{\epsilon^4}\right). \quad (3)$$

More recent results such as by [10, 29] further enhances (3) into:

$$\mathcal{O}\left(\frac{d}{\epsilon^2} + \frac{\sqrt{d}}{\epsilon^3}\right). \quad (4)$$

However, with the ever-increasing demand for more sophisticated and comprehensive models, (SO) problems of excessively high dimensionality are rapidly emerging. Based on the existing theories, the SGD and many other S-FOM alternatives

may have to perform at least hundreds of millions of iterations or stochastic first-order oracles to achieve the desired solution quality. Without fundamental innovations vis-à-vis high dimensionality, the SO problem (1) will become extremely difficult to solve in the near future.

Dimension-insensitive S-FOMs have been discussed by [20,17,32], especially about mirror descent algorithm (MDA) and its analysis. The authors of [1] base their algorithm on successively solving a series of ℓ_1 -regularized optimization problems using Nesterov's dual averaging algorithm and obtain the dimension insensitive result. However, [1,20,17] focused on convex SO problems. [32] extended the MDA to nonconvex settings and showed a complexity of

$$\mathcal{O}\left(\frac{\max_{\mathbf{x} \in X} \mathbb{E} [\|\nabla F(\mathbf{x}, \zeta)\|_\infty^2]}{\epsilon^4}\right). \quad (5)$$

This bound is verifiable to be much better than (3) and (4) in terms of the dependence on d ($\mathcal{O}(\log d)$ instead of $\mathcal{O}(d)$ because of the infinity norm) under certain mild assumptions on ζ (e.g. sub-Gaussian). To our knowledge, this is the only dimension-insensitive algorithm available in the nonconvex regime from the literature. However, the result is lacking in two key respects: (i). The authors assume that the distance generating function should be continuously differentiable. This is somewhat restrictive and does not allow non-smooth alternatives to enhance the algorithm. (ii). The term ϵ^{-4} in (5) can be further improved.

In this work, we propose dimension-insensitive stochastic first-order methods (DISFOMs) that allow non-Euclidean and non-smooth distances to construct the proximal term. Throughout this work, the residual function in Def. 1 is naturally defined via the following:

$$r(\bar{\mathbf{x}}) \triangleq \text{dist}_{\|\cdot\|_\infty}(0, \partial(f + \delta_X)(\bar{\mathbf{x}})) \triangleq \inf\{\|\mathbf{v}\|_\infty \mid \mathbf{v} \in \partial(f + \delta_X)(\bar{\mathbf{x}})\}. \quad (6)$$

We point out that although the residual function is given in $\|\cdot\|_\infty$, the dimension-insensitive result is not trivial. In fact, the residual function is stronger than a residual function discussed in literature ([32]) and defined in $\|\cdot\|_1$:

$$r_1(\mathbf{x}) \triangleq \frac{1}{\lambda} \|\mathbf{x} - \text{prox}_{\lambda f}(\mathbf{x})\|_1,$$

where $\text{prox}_{\lambda f}(\mathbf{x})$ is the proximal operator w.r.t. Bregman divergence. Detailed discussions are in Section 3.

Under mild assumptions, we show that DISFOM using minibatches to estimate the gradient enjoys sample complexity of $\mathcal{O}((\log d)/\epsilon^4)$ to obtain an ϵ -stationary point. Furthermore, we prove that DISFOM employing variance reduction can sharpen this bound to $\mathcal{O}((\log d)^{2/3}/\epsilon^{10/3})$, which further improves over (5) in terms of dependence on both d and ϵ , and perhaps leads to the best-known sample complexity result in terms of d in the nonconvex regime. We discuss two choices of the non-smooth distance functions, both of which allow closed-form solutions to the proximal step. Numerical experiments are conducted to illustrate the dimension-insensitive property of the proposed frameworks and show comparison with other popular algorithms.

In Section 2 we clarify some mild assumptions used in this article. We discuss the residual function defined in (6) and how it compares with others in Section 3. In

Section 4, we formally introduce the DISFOMs. Theoretical analysis are provided in Section 5. Section 6 includes numerical experiments. A final conclusion can be found in Section 7.

2 Preliminary

We consider problem 1 where the following assumptions are satisfied:

Assumption 1 *There is a stochastic first-order oracle (SFO) that returns the gradient $\nabla F(\mathbf{x}, \zeta)$ for any given input point $(\mathbf{x}, \zeta) \in \mathbb{R}^d \times \Omega$.*

Assumption 2 *Let $w(\mathbf{x}) \triangleq \nabla F(\mathbf{x}, \zeta) - \nabla f(\mathbf{x})$. $w_i(\mathbf{x})$ denotes its i th component. Then*

1. $\mathbb{E}[w(\mathbf{x})] = 0, \quad \forall \mathbf{x} \in X,$
 2. $\mathbb{E}[\exp(tw_i(\mathbf{x}))] \leq \exp(\sigma_\infty^2 t^2 / 2), \quad \forall t \in \mathbb{R}, \forall i = 1, \dots, d, \forall \mathbf{x} \in X,$
- i.e., $w_i(\mathbf{x})$ is a random variable that has σ_∞^2 -sub-Gaussian distribution.*

By Assumption 2, we immediately have

$$\mathbb{E}[\|w(\mathbf{x})\|^2] \leq d\sigma_\infty^2. \quad (7)$$

Assumption 3

$$\|\nabla F(\mathbf{x}, \zeta) - \nabla F(\mathbf{y}, \zeta)\| \leq L\|\mathbf{x} - \mathbf{y}\|, \quad \forall \zeta \in \Omega, \quad \forall \mathbf{x}, \mathbf{y} \in X. \quad (8)$$

By Assumption 3, we have that $\nabla f(\mathbf{x})$ is also Lipschitz continuous with constant L . Therefore, the following inequality holds:

$$f(\mathbf{y}) \leq f(\mathbf{x}) + \nabla f(\mathbf{x})^T (\mathbf{y} - \mathbf{x}) + \frac{L}{2} \|\mathbf{y} - \mathbf{x}\|^2. \quad (9)$$

Assumption 4

$$f(\mathbf{x}) \geq \underline{f}, \quad \forall \mathbf{x} \in X. \quad (10)$$

Assumption 1 stipulates that the oracle accessible here is the tractable computation of a stochastic gradient. Assumption 2 assumes that each component of the stochastic gradient is unbiased and its error is a sub-Gaussian random variable with parameter σ_∞ . Assumption 3 states that function $\nabla F(\cdot, \zeta)$ is Lipschitz smooth. All these assumptions above are standard or equivalent to the common conditions in the literature of stochastic approximation or S-FOMs. For example, Assumptions 1 and 2 are imposed by [20, 12, 17] (though some important results therein do not rely on Assumption 2). Note that if a random variable has a bounded support, it is sub-Gaussian. Assumption 3 or its slightly weaker version (inequality holds after taking expectation on the left) is necessary for the discussions by [29, 21, 23].

Notation. Given a sequence $\{\mathbf{v}^k\}$, $\Delta \mathbf{v}^{k+1} \triangleq \mathbf{v}^{k+1} - \mathbf{v}^k$. Given a vector $\mathbf{v} \in \mathbb{R}^d$, \mathbf{v}_i means its i th component unless specified otherwise in the context. For a positive integer K , $[K]$ denotes the set $\{1, 2, \dots, K\}$. $\|\cdot\|$ denotes the 2-norm $\|\cdot\|_2$. Denote $g^k \triangleq \nabla f(\mathbf{x}^k)$ for any k .

3 Discussion of residual function

It is true that the dimension-insensitive complexity result is dependent on the criterion used to measure optimality. If the criterion is weak, then the dimension-insensitive result could be meaningless. We demonstrate that the dimension-insensitive result associated with the residual function (6) is nontrivial by comparing $r(\mathbf{x})$ in (6) with the following two discussed in [32]:

$$\begin{aligned} r_1(\mathbf{x}) &\triangleq \frac{1}{\lambda} \|\mathbf{x} - \text{prox}_{\lambda f}(\mathbf{x})\|_1, \\ r_2(\mathbf{x}) &\triangleq \frac{1}{\lambda} \sqrt{D_h(\mathbf{x}, \text{prox}(\mathbf{x})) + D_h(\text{prox}(\mathbf{x}), \mathbf{x})}, \end{aligned}$$

where

$$\text{prox}_{\lambda f}(\mathbf{x}) \triangleq \arg \min_{\mathbf{z} \in X} f(\mathbf{z}) + \frac{1}{\lambda} D_h(\mathbf{z}, \mathbf{x}) \quad (11)$$

and $D_h(\cdot, \cdot)$ is the Bregman divergence induced by 1-strongly convex continuously differentiable distance generating function $h(\cdot)$ w.r.t $\|\cdot\|_1$, i.e.,

$$D_h(\mathbf{u}, \mathbf{v}) \triangleq h(\mathbf{u}) - h(\mathbf{v}) - \langle \nabla h(\mathbf{v}), \mathbf{u} - \mathbf{v} \rangle \geq \frac{1}{2} \|\mathbf{u} - \mathbf{v}\|_1^2.$$

The residual function $r_1(\mathbf{x})$ is a natural generalization from the one used in [7] to ℓ_1 -norm. In fact, $r_2(\mathbf{x})$ is stronger than $r_1(\mathbf{x})$ since

$$\begin{aligned} D_h(\mathbf{x}, \text{prox}(\mathbf{x})) + D_h(\text{prox}(\mathbf{x}), \mathbf{x}) &= \langle \nabla h(\mathbf{x}) - \nabla h(\text{prox}(\mathbf{x})), \mathbf{x} - \text{prox}(\mathbf{x}) \rangle \\ &\geq \|\mathbf{x} - \text{prox}(\mathbf{x})\|_1^2. \end{aligned}$$

We point out that $r(\mathbf{x})$ is stronger than $r_2(\mathbf{x})$. In fact, if f is ρ -relatively weakly convex ($f(\mathbf{x}) + \rho h(\mathbf{x})$ is convex for such $h(\cdot)$ and some $\rho > 0$) and $\lambda \in (0, \rho^{-1})$ so that (11) is well-defined, we have the following statement.

Lemma 1 *Let $\mathbf{x} \in X$, then*

$$r(\mathbf{x}) \geq (1 - \lambda\rho)r_2(\mathbf{x}). \quad (12)$$

Proof Denote $\hat{\mathbf{x}} \triangleq \text{prox}_{\lambda f}(\mathbf{x})$. Then by definition (11),

$$\begin{aligned} 0 &\in \nabla f(\hat{\mathbf{x}}) + \frac{1}{\lambda} (\nabla h(\hat{\mathbf{x}}) - \nabla h(\mathbf{x})) + \mathcal{N}_X(\hat{\mathbf{x}}), \\ &\iff \left\langle \left(\nabla f(\hat{\mathbf{x}}) + \frac{1}{\lambda} (\nabla h(\hat{\mathbf{x}}) - \nabla h(\mathbf{x})) \right), \mathbf{y} - \hat{\mathbf{x}} \right\rangle \geq 0, \quad \forall \mathbf{y} \in X, \\ &\implies \langle \nabla f(\mathbf{x}), \mathbf{x} - \hat{\mathbf{x}} \rangle \geq \left\langle \nabla f(\hat{\mathbf{x}}) + \frac{1}{\lambda} \nabla h(\hat{\mathbf{x}}) - \nabla f(\mathbf{x}) - \frac{1}{\lambda} \nabla h(\mathbf{x}), \hat{\mathbf{x}} - \mathbf{x} \right\rangle \\ &\quad \geq \left(\frac{1}{\lambda} - \rho \right) \langle \nabla h(\hat{\mathbf{x}}) - \nabla h(\mathbf{x}), \hat{\mathbf{x}} - \mathbf{x} \rangle. \end{aligned} \quad (13)$$

Choose arbitrary $\mathbf{v} \in \partial(f + \delta_X)(\mathbf{x})$, then we have

$$\langle \mathbf{v} - \nabla f(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle \leq 0, \quad \forall \mathbf{y} \in X. \quad (14)$$

By (13) and (14), we have

$$(1/\lambda - \rho) \langle \nabla h(\hat{\mathbf{x}}) - \nabla h(\mathbf{x}), \hat{\mathbf{x}} - \mathbf{x} \rangle \leq \langle \mathbf{v}, \mathbf{x} - \hat{\mathbf{x}} \rangle. \quad (15)$$

WLOG, suppose that $\hat{\mathbf{x}} \neq \mathbf{x}$. Then by (15),

$$(1/\lambda - \rho) \frac{\langle \nabla h(\hat{\mathbf{x}}) - \nabla h(\mathbf{x}), \hat{\mathbf{x}} - \mathbf{x} \rangle}{\|\hat{\mathbf{x}} - \mathbf{x}\|_1} \leq \left\langle \mathbf{v}, \frac{\mathbf{x} - \hat{\mathbf{x}}}{\|\mathbf{x} - \hat{\mathbf{x}}\|_1} \right\rangle \leq \|\mathbf{v}\|_\infty. \quad (16)$$

However,

$$\begin{aligned} \langle \nabla h(\hat{\mathbf{x}}) - \nabla h(\mathbf{x}), \hat{\mathbf{x}} - \mathbf{x} \rangle &\geq \|\hat{\mathbf{x}} - \mathbf{x}\|_1^2 \\ \implies \frac{\langle \nabla h(\hat{\mathbf{x}}) - \nabla h(\mathbf{x}), \hat{\mathbf{x}} - \mathbf{x} \rangle}{\|\hat{\mathbf{x}} - \mathbf{x}\|_1} &\geq \sqrt{\langle \nabla h(\hat{\mathbf{x}}) - \nabla h(\mathbf{x}), \hat{\mathbf{x}} - \mathbf{x} \rangle}. \end{aligned} \quad (17)$$

By combining (16)(17) and the fact that \mathbf{v} is arbitrary, we have that (12) holds. \square

4 Dimension-insensitive stochastic first-order methods

We start from a general framework of a stochastic first-order method as follows.

Algorithm 1 Stochastic first-order method

Step 1. Initialize \mathbf{x}^1 and hyper-parameters $K, \eta > 0$.

Step 2. Invoke the following operations for $k = 1, \dots, K$:

Step 2.1 Generate the gradient estimate G^k .

Step 2.2 Solve the proximal projection problem:

$$\mathbf{x}^{k+1} = P_X^k(\mathbf{x}^k - \eta G^k) \quad (18)$$

Step 3. Output \mathbf{x}^{Y+1} for a random Y , which has a discrete distribution on $[K]$ with a probability mass function $\mathbb{P}[Y = k] = 1/K$.

Now we discuss the details of the proximal projection operator $P_X^k(\cdot)$ and gradient estimate G^k that can achieve the dimension-insensitive property. In particular, the proximal projection operator $P_X^k(\cdot)$ is defined as

$$P_X^k(\mathbf{v}) = \arg \min_{\mathbf{x} \in X} \left\{ \frac{1}{2} \|\mathbf{x} - \mathbf{v}\|^2 + \phi(\mathbf{x} - \mathbf{x}^k) \right\}, \quad (19)$$

where $\phi : \mathbb{R}^d \rightarrow \mathbb{R}$ is a convex function and we suppose that calculation of (19) is easy and exact when $X = \mathbb{R}^d$ (we will discuss the specific form of ϕ later) or when $\phi \equiv 0$. Moreover, suppose that $\phi(0) = 0$. We could employ ADMM (see Appendix B) to resolve (18)(19) to accuracy level $\hat{\epsilon} > 0$ so that the following conditions hold: $\exists \xi^{k+1}, \Gamma^{k+1}, \mathbf{y}^{k+1}$, s.t.,

$$(\Delta \mathbf{x}^{k+1} + \eta G^k + \xi^{k+1} + \Gamma^{k+1})^T (\mathbf{x} - \mathbf{x}^{k+1}) \geq 0, \quad \forall \mathbf{x} \in X, \quad (20a)$$

$$\xi^{k+1} \in \partial \phi(\mathbf{y}^{k+1} - \mathbf{x}^k), \quad (20b)$$

$$\|\Gamma^{k+1}\|_\infty \leq \hat{\epsilon}, \quad \|\mathbf{x}^{k+1} - \mathbf{y}^{k+1}\|_1 \leq \hat{\epsilon}. \quad (20c)$$

In particular, when $X = \mathbb{R}^d$, by assumption we have that (19) can be solved easily and exactly, and (20a)-(20c) hold true with $\hat{\epsilon} = 0$. In fact, when $X = \mathbb{R}^d$ we have

$$-\Delta \mathbf{x}^{k+1} - \eta G^k = \xi^{k+1} \in \partial \phi(\mathbf{x}^{k+1} - \mathbf{x}^k). \quad (21)$$

If $X \neq \mathbb{R}^d$, we point out that ADMM is also fast in resolving (19) under the assumption that X is a polyhedron and $\partial \phi(\mathbf{x})$ is a piecewise linear multifunctions ($Gr(\partial \phi(\mathbf{x})) \triangleq \{(\mathbf{x}, \mathbf{y}) \mid \mathbf{y} \in \partial \phi(\mathbf{x})\}$ is the union of finitely many polyhedra). Then by [31], ADMM converges in linear rate in 2-norm. That means to obtain (20a)-(20c), the iterations of ADMM is at most $\mathcal{O}(\log(\sqrt{d}/\hat{\epsilon}))$. Considering that each step of ADMM is cheap, additional computation introduced by $P_X^k(\cdot)$ is incremental.

4.1 Choices of ϕ

Next we provide two possible choices of ϕ .

Case 1: Let

$$\phi(\mathbf{x}) = \frac{\hat{\rho}}{2} \|\mathbf{x}\|_1^2, \quad \hat{\rho} > 0. \quad (22)$$

For such a $\phi(\mathbf{x})$, the following lemma holds by application of Danskin's Theorem.

Lemma 2 *Let $\phi(\mathbf{x})$ be in (22). Then*

$$\partial \phi(\mathbf{x}) = \hat{\rho} \|\mathbf{x}\|_1 \partial \|\mathbf{x}\|_1 = \left\{ \hat{\rho} \|\mathbf{x}\|_1 \mathbf{v} : \mathbf{v} \in \mathbb{R}^d, \begin{array}{ll} \mathbf{v}_i = 1 & \text{if } \mathbf{x}_i > 0 \\ \mathbf{v}_i = -1 & \text{if } \mathbf{x}_i < 0 \\ \mathbf{v}_i \in [-1, 1] & \text{if } \mathbf{x}_i = 0 \end{array} \right\}. \quad (23)$$

Proof Let $\text{conv}(S)$ denote the convex hull of a set S . Note that $\|\mathbf{x}\|_1^2 = \max_{\alpha \in [-1, 1]^d} (\alpha^T \mathbf{x})^2$. Then by Danskin's Theorem,

$$\begin{aligned} & \partial \|\mathbf{x}\|_1^2 \\ &= \text{conv} \left\{ 2(\alpha^T \mathbf{x}) \alpha : \begin{array}{l} \left\{ \begin{array}{ll} \alpha_i = 1 & \text{if } \mathbf{x}_i > 0, \\ \alpha_i = -1 & \text{if } \mathbf{x}_i < 0, \\ \alpha_i \in [-1, 1] & \text{if } \mathbf{x}_i = 0, \end{array} \right\} \text{ or } \left\{ \begin{array}{ll} \alpha_i = -1 & \text{if } \mathbf{x}_i > 0, \\ \alpha_i = 1 & \text{if } \mathbf{x}_i < 0, \\ \alpha_i \in [-1, 1] & \text{if } \mathbf{x}_i = 0, \end{array} \right\} \end{array} \right\} \\ &= \text{conv} \left\{ 2\|\mathbf{x}\|_1 \alpha : \begin{array}{l} \left\{ \begin{array}{ll} \alpha_i = 1 & \text{if } \mathbf{x}_i > 0, \\ \alpha_i = -1 & \text{if } \mathbf{x}_i < 0, \\ \alpha_i \in [-1, 1] & \text{if } \mathbf{x}_i = 0, \end{array} \right\} \end{array} \right\} \\ &= 2\|\mathbf{x}\|_1 \partial \|\mathbf{x}\|_1. \end{aligned}$$

Therefore the result follows. \square

Now we discuss the formula to calculate the following proximal problem (one of the subproblems in ADMM iterations) for such a choice of ϕ :

$$\min_{\mathbf{z}} \frac{1}{2} \|\mathbf{z} - \mathbf{v}\|^2 + \frac{\hat{\rho}}{2} \|\mathbf{z}\|_1^2. \quad (24)$$

The next lemma demonstrates that it is indeed tractable to find the solution of (24).

Lemma 3 Suppose that \mathbf{z}^* denotes the optimal solution of (24). Then, if $\mathbf{v} = 0$, $\mathbf{z}^* = 0$. Otherwise, suppose that $\mathbf{v} \neq 0$. Let i_1, i_2, \dots, i_d be the permutation of $\{1, \dots, d\}$ such that

$$|\mathbf{v}_{i_1}| \leq |\mathbf{v}_{i_2}| \leq \dots \leq |\mathbf{v}_{i_d}|.$$

Denote a dummy index $i_0 = 0$ and dummy scalar $\mathbf{v}_{i_0} = 0$. Let $s_k \triangleq \sum_{t=0}^{k-1} |\mathbf{v}_{i_t}| + \frac{(d-k+1)\hat{\rho}+1}{\hat{\rho}} |\mathbf{v}_{i_k}|$, $\forall k = 0, \dots, d$. (By convention, $\sum_{t=0}^{-1} |\mathbf{v}_{i_t}| \triangleq 0$). Then s_k is non-decreasing and $\|\mathbf{v}\|_1 < s_d$. Suppose that \bar{k} satisfies $s_{\bar{k}} \leq \|\mathbf{v}\|_1 < s_{\bar{k}+1}$. Then we have that

$$\mathbf{z}_{i_t}^* = \begin{cases} 0 & \text{if } 1 \leq t \leq \bar{k} \\ \mathbf{v}_{i_t} - \frac{\text{sgn}(\mathbf{v}_{i_t})\hat{\rho}}{\hat{\rho}(d-k)+1} \sum_{t=k+1}^d |\mathbf{v}_{i_t}| & \text{if } t > \bar{k}. \end{cases} \quad (25)$$

Proof When $\mathbf{v} = 0$, it is trivial to see that $\mathbf{z}^* = 0$. Suppose that $\mathbf{v} \neq 0$. Note that we have

$$\begin{aligned} 0 &\in \mathbf{z}^* - \mathbf{v} + \frac{\hat{\rho}}{2} \partial \|\mathbf{z}\|_1^2 \Big|_{\mathbf{z}=\mathbf{z}^*} \stackrel{(23)}{=} \mathbf{z}^* - \mathbf{v} + \hat{\rho} \|\mathbf{z}^*\|_1 \partial \|\mathbf{z}\|_1 \Big|_{\mathbf{z}=\mathbf{z}^*} \\ \iff \mathbf{z}^* &= \arg \min_{\mathbf{z}} \frac{1}{2} \|\mathbf{z} - \mathbf{v}\|^2 + \hat{\rho} \|\mathbf{z}^*\|_1 \|\mathbf{z}\|_1 \\ \iff \mathbf{z}_i^* &= \begin{cases} 0 & \text{if } |\mathbf{v}_i| \leq \hat{\rho} \|\mathbf{z}^*\|_1 \\ \mathbf{v}_i - \hat{\rho} \|\mathbf{z}^*\|_1 & \text{if } \mathbf{v}_i > \hat{\rho} \|\mathbf{z}^*\|_1 \\ \mathbf{v}_i + \hat{\rho} \|\mathbf{z}^*\|_1 & \text{if } \mathbf{v}_i < -\hat{\rho} \|\mathbf{z}^*\|_1 \end{cases} \end{aligned} \quad (26)$$

It is easy to see that $\hat{\rho} \|\mathbf{z}^*\|_1 \leq |\mathbf{v}_{i_d}|$, otherwise by (26) $\mathbf{z}^* = 0$, a contradiction. If $\hat{\rho} \|\mathbf{z}^*\|_1 = |\mathbf{v}_{i_d}|$, then by (26), $\|\mathbf{z}^*\|_1 = |\mathbf{v}_{i_d}| = 0 \implies \mathbf{v} = 0$, a contradiction to our assumption. Therefore, $\hat{\rho} \|\mathbf{z}^*\|_1 < |\mathbf{v}_{i_d}|$. Suppose that $|\mathbf{v}_{i_k}| \leq \hat{\rho} \|\mathbf{z}^*\|_1 < |\mathbf{v}_{i_{k+1}}|$ for some k s.t. $0 \leq k \leq d-1$. Note that such k is unique for a fixed \mathbf{v} . Then by (26) we have

$$\mathbf{z}_{i_t}^* = \begin{cases} 0 & \text{if } 1 \leq t \leq k \\ \mathbf{v}_{i_t} - \text{sgn}(\mathbf{v}_{i_t}) \hat{\rho} \|\mathbf{z}^*\|_1 & \text{if } t > k \end{cases}$$

This indicates that

$$\|\mathbf{z}^*\|_1 = \sum_{t=k+1}^d |\mathbf{v}_{i_t}| - \hat{\rho}(d-k) \|\mathbf{z}^*\|_1 \implies \|\mathbf{z}^*\|_1 = \frac{1}{\hat{\rho}(d-k)+1} \sum_{t=k+1}^d |\mathbf{v}_{i_t}|.$$

Therefore,

$$\mathbf{z}_{i_t}^* = \begin{cases} 0 & \text{if } 1 \leq t \leq k \\ \mathbf{v}_{i_t} - \frac{\text{sgn}(\mathbf{v}_{i_t})\hat{\rho}}{\hat{\rho}(d-k)+1} \sum_{t=k+1}^d |\mathbf{v}_{i_t}| & \text{if } t > k \end{cases} \quad (27)$$

By the fact that $|\mathbf{v}_{i_k}| \leq \hat{\rho} \|\mathbf{z}^*\|_1 < |\mathbf{v}_{i_{k+1}}|$, this solution leads to $s_k \leq \|\mathbf{v}\|_1 < s_{k+1}$. By the fact that $s_{t+1} - s_t = \frac{(d-t)+1}{\hat{\rho}} (|\mathbf{v}_{i_{t+1}}| - |\mathbf{v}_{i_t}|)$ and $s_d = \|\mathbf{v}\|_1 + \frac{1}{\hat{\rho}} |\mathbf{v}_{i_d}|$, s_t is non-decreasing and $\|\mathbf{v}\|_1 < s_d$. \square

An alternative to case 1 is an indicator function defined as follows.

Case 2: Let

$$\phi(\mathbf{z}) = \delta_{\bar{X}}(\mathbf{z}) = \begin{cases} 0 & \text{if } \mathbf{z} \in \bar{X} \\ +\infty & \text{if } \mathbf{z} \notin \bar{X}, \end{cases} \quad (28)$$

where $\bar{X} \triangleq \{\mathbf{z} \mid \|\mathbf{z}\|_1 \leq \psi\}$ and $\psi > 0$.

In this case, we need to solve the following problem in the proximal projection step.

$$\begin{aligned} \min_{\mathbf{z}} \quad & \frac{1}{2} \|\mathbf{z} - \mathbf{v}\|_2^2 \\ \text{s.t.} \quad & \|\mathbf{z}\|_1 \leq \psi \end{aligned} \quad (29)$$

The next lemma reveals the closed-form solution to (29). Since this result utilizes similar technique as in Lemma 3, we relegate its proof to Appendix A.

Lemma 4 *Suppose that \mathbf{z}^* denotes the optimal solution of (29). If $\|\mathbf{v}\|_1 \leq \psi$, then $\mathbf{z}^* = \mathbf{v}$. Otherwise, suppose that i_1, \dots, i_d is a permutation of $\{1, 2, \dots, d\}$ such that $|\mathbf{v}_{i_1}| \geq |\mathbf{v}_{i_2}| \geq \dots \geq |\mathbf{v}_{i_d}|$. Denote $s_0 \triangleq 0$, $s_d \triangleq \|\mathbf{v}\|_1$ and $s_m \triangleq \sum_{i=1}^m |\mathbf{v}_{i_m}| - m|\mathbf{v}_{i_{m+1}}|$, $m = 1, 2, \dots, d-1$. Then s_m is a non-decreasing sequence. Suppose that $s_{\bar{m}-1} < \psi \leq s_{\bar{m}}$ for some $\bar{m} \in [d]$. Then we have*

$$\mathbf{z}_{i_t}^* = \begin{cases} 0 & \text{if } \bar{m} < t \leq d \\ \mathbf{v}_{(i_t)} - \frac{\text{sgn}(\mathbf{v}_{(i_t)})}{\bar{m}} (\sum_{k=1}^{\bar{m}} |\mathbf{v}_{(i_k)}| - \psi) & \text{if } 1 \leq t \leq \bar{m} \end{cases}$$

Remark 1 According to the formulae given by Lemma 3 and 4, computing \mathbf{z}^* of (24) or (29) only requires $O(d)$ fundamental operations.

4.2 Choices of G^k

Next we discuss choices of the gradient estimate G^k and its variance in infinity norm.

Minibatch. First we consider using a finite batch of samples to estimate the gradient $g^k \triangleq \nabla f(\mathbf{x}^k)$. Although the batchsize is not necessarily small, we borrow the name *minibatch* considering that accurate estimation of $\nabla f(\mathbf{x}^k)$ will take probably huge or even infinite number of samples. More specifically, let

$$G^k \triangleq \frac{1}{m} \sum_{i=1}^m \nabla F(\mathbf{x}^k, \zeta_i^k), \quad (30)$$

where ζ_i^k are i.i.d realizations of ζ . Then the following results regarding the sub-Gaussian assumption of the bias hold. Since the results are relatively basic, we relegate the proof to Appendix A.

Lemma 5 *Assumption 2 holds and consider the minibatch sampling (30). Let $\bar{w}^k \triangleq \frac{1}{m} \sum_{i=1}^m (\nabla F(\mathbf{x}^k, \zeta_i^k) - \nabla f(\mathbf{x}^k))$, then for any k ,*

1. $\mathbb{E}[\bar{w}^k \mid \mathbf{x}^k] = 0$,
2. $\mathbb{E}[\exp(t\bar{w}_i^k) \mid \mathbf{x}^k] \leq \exp(t^2 \cdot \sigma_\infty^2 / (2m))$, for any $t \in \mathbb{R}$, $i = 1, \dots, d$,

$$3. \mathbb{E}[\|\bar{w}^k\|_\infty^2 \mid \mathbf{x}^k] \leq 6 \log(2d)\sigma_\infty^2/m.$$

Remark 2 By (30) and Lemma 5, we have that $\exists c > 0$, s.t., $\forall k$,

$$\mathbb{E}[\|g^k - G^k\|_\infty^2] \leq \frac{c(\log d)\sigma_\infty^2}{m}. \quad (31)$$

Variance Reduction. Second, we can also apply the variance reduction technique to construct the sampled gradient [15]. The technique requires using relatively large number of samples intermittently when estimating g^k , and reuse this relatively more accurate estimation in other iterations to improve sampling efficiency.

Given interval length q , define $n_k \triangleq \lfloor \frac{k-1}{q} \rfloor q + 1$. Then G^k is computed as follows:

$$G^k \triangleq \begin{cases} \frac{1}{m_k} \sum_{i=1}^{m_k} \nabla f(\mathbf{x}^k, \zeta_i^k), & \text{if } \text{mod}(k, q) = 1, \\ \frac{1}{m_k} (\sum_{i=1}^{m_k} \nabla F(\mathbf{x}^k, \zeta_i^k) - \sum_{i=1}^{m_k} \nabla F(\mathbf{x}^{n_k}, \zeta_i^k)) + G^{n_k}, & \text{otherwise,} \end{cases} \quad (32)$$

where ζ_i^k are i.i.d realizations of ζ . Similar to (31), for any n_k ,

$$\mathbb{E}[\|g^{n_k} - G^{n_k}\|_\infty^2] \leq \frac{c(\log d)\sigma_\infty^2}{m_{n_k}}. \quad (33)$$

Suppose that

$$\begin{cases} m_k \equiv m & \text{if } k \neq n_k \\ m_{n_k} \equiv m_1 & \text{otherwise.} \end{cases} \quad (34)$$

Then we have the following result regarding the upper bound for the averaged variance across all the iterates in infinity norm.

Lemma 6 *Algorithm 1 and Assumption 3 hold. Consider using variance reduction to estimate G^k ((32)(34)). Then*

$$\mathbb{E}[\|\nabla f(\mathbf{x}^Y) - G^Y\|_\infty^2] \leq \frac{8L^2q^2}{m} \mathbb{E}[\|\mathbf{x}^{Y+1} - \mathbf{x}^Y\|^2] + \frac{2c(\log d)\sigma_\infty^2}{m_1}. \quad (35)$$

Remark 3 We defer the proof to Appendix A since the proof in $\|\cdot\|_\infty$ is similar to the one in $\|\cdot\|$, which is much discussed in literature. Lemma 6 indicates that we can bound the average of gradient variance in $\|\cdot\|_\infty$ by summation of two terms, the average square of difference of consecutive iterates and a constant depending only on $\log d$. Note that Lemma 6 holds regardless of how we define the proximal projection step (Step 2.2 in Algorithm 1) as long as the iterates are within X . To control the variance to a low level, we only need to pick a large m_1 . Requirement on the size of m is less restrictive since the average $\mathbb{E}[\|x^{Y+1} - x^Y\|^2]$ can be controlled at a low level when iteration count K is large. Therefore the sampling efficiency is improved.

5 Convergence and complexity analysis

In this section we analyze Algorithm 1 in the various settings discussed in the last section and derive dimension-insensitive sample complexity guarantees.

First we provide two key lemmas useful for analyzing all the cases.

5.1 Key lemmas

Throughout this section, suppose that Assumption 1-4 hold and let $\Delta \triangleq \mathbb{E}[f(x^1)] - f$. We start with the following lemma which holds without specifying the choices of ϕ or G^k .

Lemma 7 Consider Algorithm 1 with (20). Then for any positive scalars γ, τ, t , we have

$$\begin{aligned} & \left(\frac{1}{\eta} - \frac{L}{2} - \frac{\gamma}{2\eta} \right) \mathbb{E}[\|\Delta \mathbf{x}^{Y+1}\|^2] + \frac{1}{\eta} \mathbb{E}[(\xi^{Y+1})^T (\mathbf{y}^{Y+1} - \mathbf{x}^Y)] \\ & \leq \frac{\Delta}{K} + \frac{\tau+t}{2} \mathbb{E}[\|g^Y - G^Y\|_\infty^2] + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} \right) \hat{\epsilon}^2 + \frac{1}{2t} \mathbb{E}[\|\mathbf{y}^{Y+1} - \mathbf{x}^Y\|_1^2] \\ & \quad + \frac{\hat{\epsilon}}{\eta} \mathbb{E}[\|\xi^{Y+1}\|_\infty]. \end{aligned} \quad (36)$$

Proof We have that

$$\begin{aligned} & f(\mathbf{x}^{k+1}) \\ & \stackrel{(9)}{\leq} f(\mathbf{x}^k) + (g^k)^T \Delta \mathbf{x}^{k+1} + \frac{L}{2} \|\Delta \mathbf{x}^{k+1}\|^2 \\ & \leq f(\mathbf{x}^k) + (g^k - G^k)^T \Delta \mathbf{x}^{k+1} + (G^k)^T \Delta \mathbf{x}^{k+1} + \frac{L}{2} \|\Delta \mathbf{x}^{k+1}\|^2 \\ & \stackrel{(20a)}{\leq} f(\mathbf{x}^k) + (g^k - G^k)^T (\mathbf{x}^{k+1} - \mathbf{y}^{k+1}) + (g^k - G^k)^T (\mathbf{y}^{k+1} - \mathbf{x}^k) \\ & \quad + \frac{1}{\eta} (\Delta \mathbf{x}^{k+1} + \xi^{k+1} + \Gamma^{k+1})^T (\mathbf{x}^k - \mathbf{x}^{k+1}) + \frac{L}{2} \|\Delta \mathbf{x}^{k+1}\|^2 \\ & \leq f(\mathbf{x}^k) + \frac{\tau}{2} \|g^k - G^k\|_\infty^2 + \frac{1}{2\tau} \|\mathbf{x}^{k+1} - \mathbf{y}^{k+1}\|_1^2 + \frac{t}{2} \|g^k - G^k\|_\infty^2 \\ & \quad + \frac{1}{2t} \|\mathbf{y}^{k+1} - \mathbf{x}^k\|_1^2 - \left(\frac{1}{\eta} - \frac{L}{2} \right) \|\Delta \mathbf{x}^{k+1}\|^2 - \frac{1}{\eta} (\Gamma^{k+1})^T \Delta \mathbf{x}^{k+1} \\ & \quad + \frac{1}{\eta} (\xi^{k+1})^T (\mathbf{x}^k - \mathbf{y}^{k+1}) + \frac{1}{\eta} (\xi^{k+1})^T (\mathbf{y}^{k+1} - \mathbf{x}^{k+1}) \\ & \stackrel{(20c)}{\leq} f(\mathbf{x}^k) + \frac{\tau}{2} \|g^k - G^k\|_\infty^2 + \frac{\hat{\epsilon}^2}{2\tau} + \frac{t}{2} \|g^k - G^k\|_\infty^2 + \frac{1}{2t} \|\mathbf{y}^{k+1} - \mathbf{x}^k\|_1^2 \\ & \quad - \left(\frac{1}{\eta} - \frac{L}{2} - \frac{\gamma}{2\eta} \right) \|\Delta \mathbf{x}^{k+1}\|^2 + \frac{\hat{\epsilon}^2}{2\eta\gamma} + \frac{1}{\eta} (\xi^{k+1})^T (\mathbf{x}^k - \mathbf{y}^{k+1}) \\ & \quad + \frac{1}{\eta} \|\xi^{k+1}\|_\infty \|\mathbf{y}^{k+1} - \mathbf{x}^{k+1}\|_1 \\ & \leq f(\mathbf{x}^k) + \frac{\tau}{2} \|g^k - G^k\|_\infty^2 + \frac{\hat{\epsilon}^2}{2\tau} + \frac{t}{2} \|g^k - G^k\|_\infty^2 + \frac{1}{2t} \|\mathbf{y}^{k+1} - \mathbf{x}^k\|_1^2 \\ & \quad - \left(\frac{1}{\eta} - \frac{L}{2} - \frac{\gamma}{2\eta} \right) \|\Delta \mathbf{x}^{k+1}\|^2 + \frac{\hat{\epsilon}^2}{2\eta\gamma} + \frac{1}{\eta} (\xi^{k+1})^T (\mathbf{x}^k - \mathbf{y}^{k+1}) + \frac{\hat{\epsilon}}{\eta} \|\xi^{k+1}\|_\infty \end{aligned}$$

By rearranging terms, we have that

$$\left(\frac{1}{\eta} - \frac{L}{2} - \frac{\gamma}{2\eta} \right) \|\Delta \mathbf{x}^{k+1}\|^2 + \frac{1}{\eta} (\xi^{k+1})^T (\mathbf{y}^{k+1} - \mathbf{x}^k)$$

$$\begin{aligned} &\leq f(\mathbf{x}^k) - f(\mathbf{x}^{k+1}) + \frac{\tau+t}{2} \|g^k - G^k\|_\infty^2 + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma}\right) \hat{\epsilon}^2 + \frac{1}{2t} \|\mathbf{y}^{k+1} - \mathbf{x}^k\|_1^2 \\ &\quad + \frac{\hat{\epsilon}}{\eta} \|\xi^{k+1}\|_\infty. \end{aligned}$$

Sum up the above inequality for $k = 1, \dots, K$ and take expectation on both sides, we have

$$\begin{aligned} &\left(\frac{1}{\eta} - \frac{L}{2} - \frac{\gamma}{2\eta}\right) \sum_{k=1}^K \mathbb{E}[\|\Delta\mathbf{x}^{k+1}\|^2] + \frac{1}{\eta} \sum_{k=1}^K \mathbb{E}[(\xi^{k+1})^T (\mathbf{y}^{k+1} - \mathbf{x}^k)] \\ &\leq \mathbb{E}[f(\mathbf{x}^1) - f(\mathbf{x}^{K+1})] + \frac{\tau+t}{2} \sum_{k=1}^K \mathbb{E}[\|g^k - G^k\|_\infty^2] + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma}\right) K \hat{\epsilon}^2 \\ &\quad + \frac{1}{2t} \sum_{k=1}^K \mathbb{E}[\|\mathbf{y}^{k+1} - \mathbf{x}^k\|_1^2] + \frac{\hat{\epsilon}}{\eta} \sum_{k=1}^K \mathbb{E}[\|\xi^{k+1}\|_\infty]. \end{aligned}$$

Recall $\Delta \triangleq \mathbb{E}[f(\mathbf{x}^1)] - \underline{f} \geq \mathbb{E}[f(\mathbf{x}^1) - f(\mathbf{x}^{K+1})]$ and divide both sides by K . Then the result follows. \square

The residual of interest is $\text{dist}_{\|\cdot\|_\infty}(0, \partial(f + \delta_X)(\mathbf{x}^{Y+1}))$. The following lemma provides an upper bound for its expected value.

Lemma 8 Consider Algorithm 1 with (20). We have

$$\begin{aligned} &\mathbb{E}[\text{dist}_{\|\cdot\|_\infty}(0, \partial(f + \delta_X)(\mathbf{x}^{Y+1}))] \\ &\leq \left(L + \frac{1}{\eta}\right) \sqrt{\mathbb{E}[\|\Delta\mathbf{x}^{Y+1}\|^2]} + \sqrt{\mathbb{E}[\|g^Y - G^Y\|_\infty^2]} + \frac{1}{\eta} \mathbb{E}[\|\xi^{Y+1}\|_\infty] + \frac{\hat{\epsilon}}{\eta}. \quad (37) \end{aligned}$$

Proof Note that due to (20a), we have that

$$\nabla f(\mathbf{x}^{Y+1}) - \frac{1}{\eta} (\Delta\mathbf{x}^{Y+1} + \eta G^Y + \xi^{Y+1} + \Gamma^{Y+1}) \in \partial(f + \delta_X)(\mathbf{x}^{Y+1}).$$

Therefore,

$$\begin{aligned} &\mathbb{E}[\text{dist}_{\|\cdot\|_\infty}(0, \partial(f + \delta_X)(\mathbf{x}^{Y+1}))] \\ &\leq \mathbb{E}[\|\nabla f(\mathbf{x}^{Y+1}) - \frac{1}{\eta} (\Delta\mathbf{x}^{Y+1} + \eta G^Y + \xi^{Y+1} + \Gamma^{Y+1})\|_\infty] \\ &\leq \mathbb{E}[\|\nabla f(\mathbf{x}^{Y+1}) - g^Y\|_\infty] + \mathbb{E}[\|g^Y - G^Y\|_\infty] \\ &\quad + \frac{1}{\eta} \mathbb{E}[\|\Delta\mathbf{x}^{Y+1} + \xi^{Y+1} + \Gamma^{Y+1}\|_\infty] \\ &\leq L \mathbb{E}[\|\Delta\mathbf{x}^{Y+1}\|] + \mathbb{E}[\|g^Y - G^Y\|_\infty] \\ &\quad + \frac{1}{\eta} (\mathbb{E}[\|\Delta\mathbf{x}^{Y+1}\|] + \mathbb{E}[\|\xi^{Y+1}\|_\infty] + \mathbb{E}[\|\Gamma^{Y+1}\|_\infty]) \\ &\leq \left(L + \frac{1}{\eta}\right) \mathbb{E}[\|\Delta\mathbf{x}^{Y+1}\|] + \mathbb{E}[\|g^Y - G^Y\|_\infty] + \frac{1}{\eta} \mathbb{E}[\|\xi^{Y+1}\|_\infty] + \frac{\hat{\epsilon}}{\eta} \\ &\leq \left(L + \frac{1}{\eta}\right) \sqrt{\mathbb{E}[\|\Delta\mathbf{x}^{Y+1}\|^2]} + \sqrt{\mathbb{E}[\|g^Y - G^Y\|_\infty^2]} + \frac{1}{\eta} \mathbb{E}[\|\xi^{Y+1}\|_\infty] + \frac{\hat{\epsilon}}{\eta}. \end{aligned}$$

\square

5.2 Analysis for Case 1

The main result for minibatch sampling is as follows.

Theorem 1 For Algorithm 1 with (20), consider case 1 (22) and estimating G^k using minibatch (30). Then

$$\begin{aligned} & \mathbb{E}[\text{dist}_{\|\cdot\|_\infty}(0, \partial(f + \delta_X)(\mathbf{x}^{Y+1}))] \\ & \leq \left(L + \frac{1}{\eta}\right) \sqrt{\frac{\frac{\Delta}{K} + \frac{(\tau+t)c(\log d)\sigma_\infty^2}{2m} + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} + \frac{2\hat{\rho}}{\eta}\right)\hat{\epsilon}^2}{1/\eta - L/2 - \gamma/(2\eta)}} + \sqrt{\frac{c(\log d)\sigma_\infty^2}{m}} \\ & \quad + \frac{\hat{\rho}}{\eta} \sqrt{\frac{\frac{\Delta}{K} + \frac{(\tau+t)c(\log d)\sigma_\infty^2}{2m} + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} + \frac{2\hat{\rho}}{\eta}\right)\hat{\epsilon}^2}{\hat{\rho}/(8\eta)}} + \frac{\hat{\epsilon}}{\eta}, \end{aligned} \quad (38)$$

holds when $1/\eta - L/2 - \gamma/(2\eta) > 0$, $t = 2\eta/\hat{\rho}$, $\tau > 0$, $\gamma > 0$.

Proof By Lemma 2 we have that

$$\partial\phi(\mathbf{z} - \mathbf{x}^k) \subseteq \{\hat{\rho}\|\mathbf{z} - \mathbf{x}^k\|_1 \mathbf{v} \mid v_i \in [-1, 1]\}.$$

Therefore,

$$\|\xi^{k+1}\|_\infty \leq \hat{\rho}\|\mathbf{y}^{k+1} - \mathbf{x}^k\|_1 \quad (39)$$

Note that $\phi(0) = 0$, then by (20b) and convexity of ϕ ,

$$(\xi^{k+1})^T(\mathbf{x}^k - \mathbf{y}^{k+1}) + \phi(\mathbf{y}^{k+1} - \mathbf{x}^k) \leq \phi(\mathbf{x}^k - \mathbf{x}^k) = \phi(0) = 0. \quad (40)$$

Since we consider using a minibatch of samples to estimate the gradient (30). Then (36)(22)(39)(40) lead to

$$\begin{aligned} & \left(\frac{1}{\eta} - \frac{L}{2} - \frac{\gamma}{2\eta}\right) \mathbb{E}[\|\Delta\mathbf{x}^{Y+1}\|^2] + \frac{\hat{\rho}}{2\eta} \mathbb{E}[\|\mathbf{y}^{Y+1} - \mathbf{x}^Y\|_1^2] \\ & \leq \left(\frac{1}{\eta} - \frac{L}{2} - \frac{\gamma}{2\eta}\right) \mathbb{E}[\|\Delta\mathbf{x}^{Y+1}\|^2] + \frac{1}{\eta} \mathbb{E}[(\xi^{Y+1})^T(\mathbf{y}^{Y+1} - \mathbf{x}^Y)] \\ & \leq \frac{\Delta}{K} + \frac{\tau+t}{2} \mathbb{E}[\|g^Y - G^Y\|_\infty^2] + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma}\right) \hat{\epsilon}^2 + \frac{1}{2t} \mathbb{E}[\|\mathbf{y}^{Y+1} - \mathbf{x}^Y\|_1^2] \\ & \quad + \frac{\hat{\epsilon}}{\eta} \mathbb{E}[\|\xi^{Y+1}\|_\infty] \\ & \leq \frac{\Delta}{K} + \frac{\tau+t}{2} \mathbb{E}[\|g^Y - G^Y\|_\infty^2] + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} + \frac{2\hat{\rho}}{\eta}\right) \hat{\epsilon}^2 + \frac{1}{2t} \mathbb{E}[\|\mathbf{y}^{Y+1} - \mathbf{x}^Y\|_1^2] \\ & \quad + \frac{1}{8\eta\hat{\rho}} \mathbb{E}[\|\xi^{Y+1}\|_\infty^2] \\ & \stackrel{(t=\frac{2\eta}{\hat{\rho}})}{=} \frac{\Delta}{K} + \frac{\tau+t}{2} \mathbb{E}[\|g^Y - G^Y\|_\infty^2] + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} + \frac{2\hat{\rho}}{\eta}\right) \hat{\epsilon}^2 \\ & \quad + \frac{\hat{\rho}}{4\eta} \mathbb{E}[\|\mathbf{y}^{Y+1} - \mathbf{x}^Y\|_1^2] + \frac{\hat{\rho}}{8\eta} \mathbb{E}[\|\mathbf{y}^{Y+1} - \mathbf{x}^Y\|_1^2] \end{aligned}$$

$$\begin{aligned}
&\implies \left(\frac{1}{\eta} - \frac{L}{2} - \frac{\gamma}{2\eta} \right) \mathbb{E}[\|\Delta \mathbf{x}^{Y+1}\|^2] + \frac{\hat{\rho}}{8\eta} \mathbb{E}[\|\mathbf{y}^{Y+1} - \mathbf{x}^Y\|_1^2] \\
&\leq \frac{\Delta}{K} + \frac{\tau+t}{2} \mathbb{E}[\|g^Y - G^Y\|_\infty^2] + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} + \frac{2\hat{\rho}}{\eta} \right) \hat{\epsilon}^2 \\
&\stackrel{(31)}{\leq} \frac{\Delta}{K} + \frac{(\tau+t)c(\log d)\sigma_\infty^2}{2m} + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} + \frac{2\hat{\rho}}{\eta} \right) \hat{\epsilon}^2. \tag{41}
\end{aligned}$$

By (31)(37)(39)(41), the result follows. \square

Corollary 1 *Under the setting of Theorem 1, let*

$$\eta = \frac{1}{L}, K = \left\lceil \frac{\Delta L}{\epsilon^2} \right\rceil, m = \left\lceil \frac{c(\log d)\sigma_\infty^2}{\epsilon^2} \right\rceil, \gamma = \frac{1}{2}, \tau = \frac{1}{L}, \hat{\epsilon} = \frac{\epsilon}{L}, t = \frac{2\eta}{\hat{\rho}}.$$

Then by (38) we have

$$\mathbb{E}[\text{dist}_{\|\cdot\|_\infty}(0, \partial(f + \delta_X)(\mathbf{x}^{Y+1}))] \leq (4\sqrt{3 + 1/\hat{\rho} + 2\hat{\rho}} + 2\sqrt{4\hat{\rho}^2 + 6\hat{\rho} + 2} + 2)\epsilon.$$

The sample complexity of the algorithm is

$$Km = \left\lceil \frac{\Delta L}{\epsilon^2} \right\rceil \left\lceil \frac{c(\log d)\sigma_\infty^2}{\epsilon^2} \right\rceil = \mathcal{O}\left(\frac{\Delta L(\log d)\sigma_\infty^2}{\epsilon^4}\right).$$

The next theorem reveals the convergence rate by applying variance reduction in Case 1. The proof is similar to Theorem 1 and relegated to Appendix A.

Theorem 2 *For Algorithm 1 with (20), consider case 1 (22) and estimating G^k using variance reduction (30)(34). Then the following holds for $1/\eta - L/2 - \gamma/(2\eta) - \frac{4(\tau+t)L^2q^2}{m} > 0$, $t = 2\eta/\hat{\rho}$, $\tau > 0$, $\gamma > 0$.*

$$\begin{aligned}
&\mathbb{E}[\text{dist}_{\|\cdot\|_\infty}(0, \partial(f + \delta_X)(\mathbf{x}^{Y+1}))] \\
&\leq \left(L + \frac{1}{\eta} + \sqrt{\frac{8L^2q^2}{m}} \right) \sqrt{\frac{\frac{\Delta}{K} + \frac{(\tau+t)c(\log d)\sigma_\infty^2}{m_1} + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} + \frac{2\hat{\rho}}{\eta} \right) \hat{\epsilon}^2}{\frac{1}{\eta} - \frac{L}{2} - \frac{\gamma}{2\eta} - \frac{4(\tau+t)L^2q^2}{m}}} \\
&+ \sqrt{\frac{2c(\log d)\sigma_\infty^2}{m_1}} + \frac{\hat{\rho}}{\eta} \sqrt{\frac{\frac{\Delta}{K} + \frac{(\tau+t)c(\log d)\sigma_\infty^2}{m_1} + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} + \frac{2\hat{\rho}}{\eta} \right) \hat{\epsilon}^2}{\hat{\rho}/(8\eta)}} + \frac{\hat{\epsilon}}{\eta}. \tag{42}
\end{aligned}$$

Corollary 2 *Under the setting of Theorem 2, let*

$$\begin{aligned}
\eta &= \frac{1}{L}, K = \left\lceil \frac{\Delta L}{\epsilon^2} \right\rceil, m_1 = \left\lceil \frac{c(\log d)\sigma_\infty^2}{\epsilon^2} \right\rceil, m = q^2 = \left\lceil \left(\frac{c(\log d)\sigma_\infty^2}{\epsilon^2} \right)^{1/3} \right\rceil^2, \\
\gamma &= \frac{1}{2}, \tau = t = \frac{1}{64L}, \hat{\epsilon} = \frac{\epsilon}{L}, \hat{\rho} = 128.
\end{aligned}$$

Then by (42) we have

$$\mathbb{E}[\text{dist}_{\|\cdot\|_\infty}(0, \partial(f + \delta_X)(\mathbf{x}^{Y+1}))] = C\epsilon,$$

where C is an absolute constant independent of any parameter of the algorithm or the problem data. The sample complexity is upper bounded by

$$\begin{aligned} Km + \left\lceil \frac{K}{q} \right\rceil m_1 &\leq 2K(m + q^2) = 4 \left\lceil \frac{\Delta L}{\epsilon^2} \right\rceil \left[\left(\frac{c(\log d)\sigma_\infty^2}{\epsilon^2} \right)^{1/3} \right]^2 \\ &= \mathcal{O} \left(\frac{\Delta L (c(\log d)\sigma_\infty^2)^{2/3}}{\epsilon^{10/3}} \right). \end{aligned}$$

5.3 Analysis for Case 2

The main result is given as follows.

Theorem 3 For Algorithm 1 with (20), consider case 2 (45) and estimating G^k using minibatch (30). Then

$$\begin{aligned} &\mathbb{E}[\text{dist}_{\|\cdot\|_\infty}(0, \partial(f + \delta_X)(\mathbf{x}^{Y+1}))] \\ &\leq \left(L + \frac{1}{\eta} \right) \sqrt{\frac{\frac{\Delta}{K} + \frac{(\tau+t)c(\log d)\sigma_\infty^2}{2m} + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} \right) \hat{\epsilon}^2 + \frac{\psi^2}{2t}}{1/\eta - L/2 - \gamma/(2\eta)}} + \sqrt{\frac{c(\log d)\sigma_\infty^2}{m}} \\ &+ \frac{1}{\eta} \cdot \frac{\frac{\Delta}{K} + \frac{(\tau+t)c(\log d)\sigma_\infty^2}{2m} + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} \right) \hat{\epsilon}^2 + \frac{\psi^2}{2t}}{(\psi - \hat{\epsilon})/\eta} + \frac{\hat{\epsilon}}{\eta}. \end{aligned} \quad (43)$$

holds when $1/\eta - L/2 - \gamma/(2\eta) > 0$, $t > 0$, $\tau > 0$, $\gamma > 0$, $\psi > \hat{\epsilon}$.

Proof Note that (20b) is equivalent to

$$\mathbf{y}^{k+1} \in \arg \min_{\mathbf{y}} -(\xi^{k+1})^T \mathbf{y} \quad \text{s.t.} \quad \|\mathbf{y} - \mathbf{x}^k\|_1 \leq \psi. \quad (44)$$

Therefore, $\exists \hat{\rho}_k$ such that

$$\xi^{k+1} \in \hat{\rho}_k \partial(\|\cdot - \mathbf{x}^k\|_1)(\mathbf{y}^{k+1}), \quad 0 \leq \hat{\rho}_k \perp \|\mathbf{y}^{k+1} - \mathbf{x}^k\|_1 - \psi \leq 0. \quad (45)$$

Then

$$\xi^{k+1} \in \{\hat{\rho}_k v \mid v_i \in [-1, 1]\}.$$

Therefore,

$$\|\xi^{k+1}\|_\infty \leq \hat{\rho}_k. \quad (46)$$

Moreover, by (45),

$$(\xi^{k+1})^T (\mathbf{y}^{k+1} - \mathbf{x}^k) \geq \hat{\rho}_k \|\mathbf{y}^{k+1} - \mathbf{x}^k\|_1 - \hat{\rho}_k \|\mathbf{x}^k - \mathbf{x}^k\|_1 = \hat{\rho}_k \|\mathbf{y}^{k+1} - \mathbf{x}^k\|_1. \quad (47)$$

Consider the minibatch sampling (30). Then (36)(45)(46)(47) lead to

$$\left(\frac{1}{\eta} - \frac{L}{2} - \frac{\gamma}{2\eta} \right) \mathbb{E}[\|\Delta \mathbf{x}^{Y+1}\|^2] + \frac{1}{\eta} \mathbb{E}[\hat{\rho}_Y \|\mathbf{y}^{Y+1} - \mathbf{x}^Y\|_1]$$

$$\begin{aligned}
&\leq \frac{\Delta}{K} + \frac{\tau+t}{2} \mathbb{E}[\|g^Y - G^Y\|_\infty^2] + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma}\right) \hat{\epsilon}^2 + \frac{1}{2t} \mathbb{E}[\|\mathbf{y}^{Y+1} - \mathbf{x}^Y\|_1^2] \\
&+ \frac{\hat{\epsilon}}{\eta} \mathbb{E}[\|\xi^{Y+1}\|_\infty] \\
&\leq \frac{\Delta}{K} + \frac{\tau+t}{2} \mathbb{E}[\|g^Y - G^Y\|_\infty^2] + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma}\right) \hat{\epsilon}^2 + \frac{\psi^2}{2t} + \frac{\hat{\epsilon}}{\eta} \mathbb{E}[\hat{\rho}_Y] \\
&\implies \left(\frac{1}{\eta} - \frac{L}{2} - \frac{\gamma}{2\eta}\right) \mathbb{E}[\|\Delta \mathbf{x}^{Y+1}\|^2] + \frac{\psi - \hat{\epsilon}}{\eta} \mathbb{E}[\hat{\rho}_Y] \\
&\leq \frac{\Delta}{K} + \frac{\tau+t}{2} \mathbb{E}[\|g^Y - G^Y\|_\infty^2] + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma}\right) \hat{\epsilon}^2 + \frac{\psi^2}{2t} \\
&\stackrel{(31)}{\leq} \frac{\Delta}{K} + \frac{(\tau+t)c(\log d)\sigma_\infty^2}{2m} + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma}\right) \hat{\epsilon}^2 + \frac{\psi^2}{2t}. \tag{48}
\end{aligned}$$

By (31)(37)(46)(48), the result follows. \square

Corollary 3 *Under the settings of Theorem 3, let*

$$\begin{aligned}
\eta &= \frac{1}{L}, \quad K = \left\lceil \frac{\Delta L}{\epsilon^2} \right\rceil, \quad m = \left\lceil \frac{c(\log d)\sigma_\infty^2}{\epsilon^2} \right\rceil, \quad \gamma = \frac{1}{2}, \quad \tau = \frac{1}{L}, \quad \hat{\epsilon} = \frac{\epsilon}{L}, \quad \psi = 2\hat{\epsilon}, \\
t &= \frac{1}{L}.
\end{aligned}$$

Then by (43) we have

$$\mathbb{E}[\text{dist}_{\|\cdot\|_\infty}(0, \partial(f + \delta_X)(\mathbf{x}^{Y+1}))] = C\epsilon,$$

where C is an absolute constant. The sample complexity is

$$Km = \mathcal{O}\left(\frac{\Delta L c(\log d)\sigma_\infty^2}{\epsilon^4}\right).$$

Apply variance reduction to case 2, and the following statements hold.

Theorem 4 *For Algorithm 1 with (20), consider case 2 (45) and estimating G^k using variance reduction (30)(34). Then the following holds when $1/\eta - L/2 - \gamma/(2\eta) - \frac{4(\tau+t)L^2q^2}{m} > 0$, $t > 0$, $\tau > 0$, $\gamma > 0$, $\psi > \hat{\epsilon}$.*

$$\begin{aligned}
&\mathbb{E}[\text{dist}_{\|\cdot\|_\infty}(0, \partial(f + \delta_X)(\mathbf{x}^{Y+1}))] \\
&\leq \left(L + \frac{1}{\eta} + \sqrt{\frac{8L^2q^2}{m}}\right) \sqrt{\frac{\frac{\Delta}{K} + \frac{(\tau+t)c(\log d)\sigma_\infty^2}{m_1} + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma}\right) \hat{\epsilon}^2 + \frac{\psi^2}{2t}}{\frac{1}{\eta} - \frac{L}{2} - \frac{\gamma}{2\eta} - \frac{4(\tau+t)L^2q^2}{m}}} \\
&+ \frac{1}{\psi - \hat{\epsilon}} \left(\frac{\Delta}{K} + \frac{(\tau+t)c(\log d)\sigma_\infty^2}{m_1} + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma}\right) \hat{\epsilon}^2 + \frac{\psi^2}{2t}\right) + \frac{\hat{\epsilon}}{\eta} \tag{49} \\
&+ \sqrt{\frac{2c(\log d)\sigma_\infty^2}{m_1}}.
\end{aligned}$$

Proof See Appendix A.

Corollary 4 *Under the settings of Theorem 4, let*

$$\eta = \frac{1}{L}, K = \left\lceil \frac{\Delta L}{\epsilon^2} \right\rceil, m_1 = \left\lceil \frac{c(\log d)\sigma_\infty^2}{\epsilon^2} \right\rceil, m = q^2 = m_1^{2/3}, \gamma = \frac{1}{2},$$

$$\tau = t = \frac{1}{64L}, \hat{\epsilon} = \frac{\epsilon}{L}, \psi = 2\hat{\epsilon},$$

Then by (49) we have

$$\mathbb{E}[\text{dist}_{\|\cdot\|_\infty}(0, \partial(f + \delta_X)(\mathbf{x}^{Y+1}))] = C\epsilon,$$

where C is an absolute constant independent of any parameter of the algorithm or the problem data. The sample complexity is upper bounded by

$$\mathcal{O}\left(\frac{\Delta L(c(\log d)\sigma_\infty^2)^{2/3}}{\epsilon^{10/3}}\right).$$

6 Numerical Experiment

In this section, we present some numerical experiments to illustrate the dimension insensitive property of our proposed methods.¹

Problem setting We consider a nonconvex stochastic quadratic programming problem as follows:

$$\min_{\mathbf{x} \in X} f(\mathbf{x}) \triangleq \frac{1}{2} \mathbb{E}[(\boldsymbol{\alpha}^T \mathbf{x} - b)^2] + \lambda \sum_{i=1}^d \frac{\mathbf{x}_i^2}{1 + \mathbf{x}_i^2}, \quad (50)$$

where the feasible region $X = [-R, R]^d$, $(\boldsymbol{\alpha}, b) \in \mathbb{R}^d \times \mathbb{R}$ is a random pair that satisfies linear relationship $b = \boldsymbol{\alpha}^T \mathbf{x}_{\text{true}} + w$. We generate the scaled covariance matrix for $\boldsymbol{\alpha}$ by the following procedure. First we generate a $d \times d$ identity matrix, and then replace its top-left $\frac{d}{16} \times \frac{d}{16}$ principal sub-matrix by $\Sigma_{\text{sub}} := \mathbf{Q}\mathbf{D}\mathbf{Q}^T \in \mathbb{R}^{\frac{d}{16} \times \frac{d}{16}}$, where \mathbf{Q} consists of the orthonormal basis of a $\frac{d}{16} \times \frac{d}{16}$ matrix whose entries are i.i.d. uniformly distributed on $(0, 1)$ and \mathbf{D} is a diagonal matrix with each diagonal entry i.i.d. uniformly distributed on $(1, 2)$. Let $\boldsymbol{\alpha} = \Sigma^{\frac{1}{2}} \mathbf{s}$, where $s_i, i = 1, \dots, d$ are i.i.d and obey truncated standard normal distribution over $[-u, u]$. w also obeys a truncated standard normal distribution over $[-u, u]$ and is independent of $\boldsymbol{\alpha}$. Therefore, the variance σ^2 of s_i and w has the following formula

$$\sigma^2 = 1 - \frac{\frac{2u}{\sqrt{2\pi}} \exp\left(-\frac{u^2}{2}\right)}{\Phi(u) - \Phi(-u)}, \quad (51)$$

where Φ is the CDF of standard normal distribution. By construction, function f has the following closed form:

$$f(\mathbf{x}) = \frac{\sigma^2}{2} (\mathbf{x} - \mathbf{x}_{\text{true}})^T \Sigma (\mathbf{x} - \mathbf{x}_{\text{true}}) + \lambda \sum_{i=1}^d \frac{\mathbf{x}_i^2}{1 + \mathbf{x}_i^2} + \frac{\sigma^2}{2}.$$

¹ Data and codes have been uploaded to <https://github.com/gxybrh/DISFOM>

Moreover, the error of sampled gradient $\nabla_{\mathbf{x}}F(\mathbf{x}, \alpha, w) - \nabla f(\mathbf{x})$ is sub-Gaussian since it has bounded support. The Lipschitz constant L of ∇f is $\lambda_{\max} + 2\lambda$, where λ_{\max} is the largest eigenvalue of $\sigma^2\Sigma$, ranging from σ^2 to $2\sigma^2$ by construction. Therefore, L has a fixed range independent of d . f is nonconvex when the minimal eigenvalue of $\nabla^2 f$, $\lambda_{\min} - \lambda/2$ is negative, where λ_{\min} denotes the minimal eigenvalue of $\sigma^2\Sigma$ and equals to σ^2 .

The experiments present the quality of solutions, measured by the averaged gap $(f - f^*)/\Delta$ and residual out of 3 replications. The value of f^* is given by using the projected gradient method with backtracking (see Algorithm 3 in Appendix B), to solve the correlated closed-form problem, and $\Delta \triangleq f(\mathbf{x}^1) - f^*$. We carry on our experiments with dimension $d \in \{2^7, 2^8, \dots, 2^{14}\}$, and start with $\mathbf{x}_1 = \mathbf{0}$. $R = 3$, $u = 3$, $\lambda = 2.5$.

Gradient generation According to the two estimating methods of gradient mentioned above, mini-batch and variance reduction, we compare three algorithms with both methods of gradient estimation. In the mini-batch setting, the batch size $m_k \equiv m = 1000$ and algorithms stop with $K = 300$. In the variance reduction setting, we set $m = 1000, q = 9$. The batch size is $m_k = m = 1000$ when $\text{mod}(k, q) = 1$ and $m_k = m^{\frac{2}{3}} = 100$ otherwise, and the algorithms stop with $K = 1350$.

Methods Here we present the detailed implementation of DISFOMs, proximal stochastic gradient descent (SGD), proximal stochastic variance reduced gradient (SVRG) and stochastic mirror descent (SMDs).

1. **DISFOM_minibatch** Algorithm 1 with $\phi(\mathbf{z}) = \frac{\hat{\rho}}{2}\|\mathbf{z}\|_1^2$ (Case 1) and mini-batch sampling (30). We choose $\hat{\rho} = 2$ in minibatch. The stepsize is $\eta_k \equiv \eta = \frac{1}{L}$.
2. **DISFOM_svr** Algorithm 1 with $\phi(\mathbf{z}) = \frac{\hat{\rho}}{2}\|\mathbf{z}\|_1^2$ (Case 1) and variance reduction (30). We choose $\hat{\rho} = 128$. The stepsize is $\eta_k \equiv \eta = \frac{1}{L}$.
3. (Proximal) **SGD** Algorithm 1 with $P_X^k \equiv P_X$ (Euclidean projection on X). G^k is estimated using minibatch (30). The step size is $\eta_k \equiv \eta = \frac{1}{L}$.
4. (Proximal) **SVRG** Algorithm 1 with $P_X^k \equiv P_X$ (Euclidean projection on X). G^k is estimated using variance reduction (32). The step size is $\eta_k \equiv \eta = \frac{1}{10L}$.²
5. **SMD_minibatch** Algorithm 1. Solve the proximal projection problem in Step 2.2 as follows:

$$\mathbf{x}^{k+1} = \arg \min_{\mathbf{z} \in X} \left\{ \langle G^k, \mathbf{z} \rangle + \frac{1}{\alpha_k} D_\omega(\mathbf{z}, \mathbf{x}^k) \right\}$$

where $D_\omega(x, y) = \omega(x) - \omega(y) - \langle \nabla \omega(y), x - y \rangle$ Let the distance generating function $\omega(x) = \frac{C}{2}\|x\|_p^2$ where $p = 1 + \frac{1}{\ln d}$ and $C = e^2 \ln d$ (so that $\omega(x)$ is 1-strongly convex w.r.t. $\|\cdot\|_1$), see [3] and the reference therein. The step size is $\alpha_k \equiv \alpha = \frac{c}{\sqrt{K}}$, where $c = \sqrt{\frac{f(x^1)}{\rho L^2}}$, $\rho = \frac{\lambda}{2} - \lambda_{\min}$, given by [32]. G^k is estimated via minibatch.

² Here we need to decrease the stepsize of SVRG to stabilize its performance.

6. **SMD_svrg** Same settings as in SMD_minibatch, except that G^k is estimated via variance reduction.

Results and interpretation




Fig. 1 averaged gap $(f - f^*)/\Delta$ vs $\log(d)$ and residual vs $\log(d)$ for mini-batch




Fig. 2 averaged gap $(f - f^*)/\Delta$ vs $\log(d)$ and residual vs $\log(d)$ for variance reduction

From Figure 1 and 2 we can see that the performance of vanilla SGD and SVRG would deteriorate rapidly with the increase of d . On the contrary, the performance of DISFOMs and SMDs are comparable and less sensitive to d , in terms of both the averaged relative function value gap $(f - f^*)/\Delta$ and the averaged residual $r(\mathbf{x})$ defined in (6). To our best knowledge, there is not a proper theory for the dimension-insensitive property of SMD_svrg in literature, and the numerical experiments imply this empirically. The results for Case 2 are similar to Case 1 thus suppressed.

7 Conclusion

In this study, we introduce dimension-insensitive stochastic first-order methods (DISFOMs) as a solution for nonconvex optimization with an expected-valued objective function. Our algorithms are designed to accommodate non-Euclidean and

non-smooth distance functions as proximal terms. When dealing with a stochastic nonconvex optimization problem, the sample complexity of stochastic first-order methods can become linearly dependent on the problem dimension. This causes trouble for solving large-scale problems. We demonstrate that DISFOM achieves a sample complexity of $O((\log d)/\epsilon^4)$ in order to obtain an ϵ -stationary point via minibatch sampling. Furthermore, we show that DISFOM with variance reduction can enhance this bound to $O((\log d)^{2/3}/\epsilon^{10/3})$. We present two options for the non-smooth distance functions, both of which allow for closed-form solutions in the proximal steps. Preliminary numerical experiments illustrate the dimension-insensitive property of our proposed methods. In our future work, we intend to extend our results to the case when the objective function f is non-smooth.

References

1. Agarwal, A., Negahban, S., Wainwright, M.J.: Stochastic optimization and sparse statistical recovery: Optimal algorithms for high dimensions. *Advances in Neural Information Processing Systems* **25** (2012)
2. Bach, F., Moulines, E.: Non-strongly-convex smooth stochastic approximation with convergence rate $o(1/n)$. In: *Advances in neural information processing systems*, pp. 773–781 (2013)
3. Beck, A., Teboulle, M.: Mirror descent and nonlinear projected subgradient methods for convex optimization. *Operations Research Letters* **31**(3), 167–175 (2003)
4. Chambolle, A., Ehrhardt, M.J., Richtárik, P., Schonlieb, C.B.: Stochastic primal-dual hybrid gradient algorithm with arbitrary sampling and imaging applications. *SIAM Journal on Optimization* **28**(4), 2783–2808 (2018)
5. Chambolle, A., Pock, T.: A first-order primal-dual algorithm for convex problems with applications to imaging. *Journal of mathematical imaging and vision* **40**(1), 120–145 (2011)
6. Chung, K.L.: On a stochastic approximation method. *The Annals of Mathematical Statistics* pp. 463–483 (1954)
7. Davis, D., Drusvyatskiy, D.: Stochastic subgradient method converges at the rate $o(k^{-1/4})$ on weakly convex functions. *arXiv preprint arXiv:1802.02988* (2018)
8. Davis, D., Drusvyatskiy, D.: Stochastic model-based minimization of weakly convex functions. *SIAM Journal on Optimization* **29**(1), 207–239 (2019)
9. Devolder, O., et al.: Stochastic first order methods in smooth convex optimization. Tech. rep., CORE (2011)
10. Fang, C., Li, C.J., Lin, Z., Zhang, T.: Spider: Near-optimal non-convex optimization via stochastic path-integrated differential estimator. *Advances in neural information processing systems* **31** (2018)
11. Ghadimi, S., Lan, G.: Optimal stochastic approximation algorithms for strongly convex stochastic composite optimization i: A generic algorithmic framework. *SIAM Journal on Optimization* **22**(4), 1469–1492 (2012)
12. Ghadimi, S., Lan, G.: Stochastic first-and zeroth-order methods for nonconvex stochastic programming. *SIAM Journal on Optimization* **23**(4), 2341–2368 (2013)
13. Ghadimi, S., Lan, G., Zhang, H.: Mini-batch stochastic approximation methods for non-convex stochastic composite optimization. *Mathematical Programming* **155**(1-2), 267–305 (2016)
14. Horváth, S., Lei, L., Richtárik, P., Jordan, M.I.: Adaptivity of stochastic gradient methods for nonconvex optimization. *SIAM Journal on Mathematics of Data Science* **4**(2), 634–648 (2022)
15. Johnson, R., Zhang, T.: Accelerating stochastic gradient descent using predictive variance reduction. *Advances in neural information processing systems* **26** (2013)
16. Lan, G.: An optimal method for stochastic composite optimization. *Mathematical Programming* **133**(1-2), 365–397 (2012)
17. Lan, G.: *First-order and Stochastic Optimization Methods for Machine Learning*. Springer Nature (2020)

18. Li, Z.: Ssrgd: Simple stochastic recursive gradient descent for escaping saddle points. In: Advances in Neural Information Processing Systems, pp. 1523–1533 (2019)
19. Metel, M., Takeda, A.: Simple stochastic gradient methods for non-smooth non-convex regularized optimization. In: International Conference on Machine Learning, pp. 4537–4545. PMLR (2019)
20. Nemirovski, A., Juditsky, A., Lan, G., Shapiro, A.: Robust stochastic approximation approach to stochastic programming. SIAM Journal on optimization **19**(4), 1574–1609 (2009)
21. Nguyen, L.M., Liu, J., Scheinberg, K., Takáč, M.: Sarah: A novel method for machine learning problems using stochastic recursive gradient. In: International Conference on Machine Learning, pp. 2613–2621. PMLR (2017)
22. Nitanda, A., Suzuki, T.: Stochastic difference of convex algorithm and its application to training deep boltzmann machines. In: Artificial intelligence and statistics, pp. 470–478 (2017)
23. Pham, N.H., Nguyen, L.M., Phan, D.T., Tran-Dinh, Q.: Proxsarah: An efficient algorithmic framework for stochastic composite nonconvex optimization. Journal of Machine Learning Research **21**(110), 1–48 (2020)
24. Polyak, B.T.: New stochastic approximation type procedures. Automat. i Telemekh **7**(98–107), 2 (1990)
25. Polyak, B.T., Juditsky, A.B.: Acceleration of stochastic approximation by averaging. SIAM journal on control and optimization **30**(4), 838–855 (1992)
26. Robbins, H., Monro, S.: A stochastic approximation method. The annals of mathematical statistics pp. 400–407 (1951)
27. Rosasco, L., Villa, S., Vũ, B.C.: Convergence of stochastic proximal gradient algorithm. Applied Mathematics & Optimization pp. 1–27 (2019)
28. Sacks, J.: Asymptotic distribution of stochastic approximation procedures. The Annals of Mathematical Statistics **29**(2), 373–405 (1958)
29. Wang, Z., Ji, K., Zhou, Y., Liang, Y., Tarokh, V.: Spiderboost and momentum: Faster variance reduction algorithms. Advances in Neural Information Processing Systems **32**, 2406–2416 (2019)
30. Xu, Y., Qi, Q., Lin, Q., Jin, R., Yang, T.: Stochastic optimization for dc functions and non-smooth non-convex regularizers with non-asymptotic convergence. In: International Conference on Machine Learning, pp. 6942–6951. PMLR (2019)
31. Yang, W.H., Han, D.: Linear convergence of the alternating direction method of multipliers for a class of convex optimization problems. SIAM journal on Numerical Analysis **54**(2), 625–640 (2016)
32. Zhang, S., He, N.: On the convergence rate of stochastic mirror descent for nonsmooth nonconvex optimization. arXiv preprint arXiv:1806.04781 (2018)

A Proofs

Proof of Lemma 4.

Proof The corresponding KKT-condition is as follows

$$\nabla_{\mathbf{z}} L(\mathbf{z}, \lambda) = (\mathbf{z} - \mathbf{v}) + \lambda \partial \|\mathbf{z}\|_1 = 0 \quad (52)$$

$$g(x) = \|\mathbf{z}\|_1 - \psi \leq 0 \quad (53)$$

$$\lambda \geq 0 \quad (54)$$

$$\lambda g(x) = 0 \quad (55)$$

From (52) we have

$$\begin{aligned} \mathbf{z}_i - \mathbf{v}_i + \lambda &= 0 & \text{if } \mathbf{z}_i > 0 \\ \mathbf{z}_i - \mathbf{v}_i - \lambda &= 0 & \text{if } \mathbf{z}_i < 0 \\ -\lambda \leq \mathbf{z}_i - \mathbf{v}_i \leq \lambda & & \text{if } \mathbf{z}_i = 0 \end{aligned}$$

Thus we have

$$\mathbf{z}_i = \mathbf{v}_i - \lambda \quad \text{if } \mathbf{v}_i - \lambda > 0 \quad (56)$$

$$\mathbf{z}_i = \mathbf{v}_i + \lambda \quad \text{if } \mathbf{v}_i + \lambda < 0 \quad (57)$$

$$\mathbf{z}_i = 0 \quad \text{if } |\mathbf{v}_i| \leq \lambda \quad (58)$$

Suppose $\lambda = 0$, then $\mathbf{z} = \mathbf{v}$. This is the solution if $\|\mathbf{v}\|_1 \leq \psi$. Otherwise suppose that $\lambda > 0$. Then we have

$$\|\mathbf{z}\|_1 = \sum_i^{|\mathbf{v}_i| > \lambda} |\mathbf{z}_i| = \sum_i^{\mathbf{v}_i > \lambda} (\mathbf{v}_i - \lambda) + \sum_i^{\mathbf{v}_i < -\lambda} (-\mathbf{v}_i - \lambda) = \sum_i^{|\mathbf{v}_i| > \lambda} (|\mathbf{v}_i| - \lambda) = \psi \quad (59)$$

Suppose there exists m such that $s_{m-1} < \psi \leq s_m$, then we can deduce that $|\mathbf{v}_{i_m}| > \lambda \geq |\mathbf{v}_{i_{m+1}}|$, therefore,

$$\psi = \sum_{t=1}^m |\mathbf{v}_{i_t}| - m\lambda \implies \lambda = \frac{1}{m} \left(\sum_{t=1}^m |\mathbf{v}_{i_t}| - \psi \right). \quad (60)$$

Finally we could derive \mathbf{z} from (56), (57), (58), (60) as follows.

$$\mathbf{z}_{i_t} = \begin{cases} 0 & \text{if } m+1 \leq t \leq d \\ \mathbf{v}_{i_t} - \frac{\text{sgn}(\mathbf{v}_{i_t})}{m} (\sum_{k=1}^m |\mathbf{v}_{i_k}| - \psi) & \text{if } 1 \leq t \leq m \end{cases} \quad (61)$$

□

Proof of Lemma 5.

Proof (1) is obvious from Assumption 2. (2) is based on the fact that summation of independent mean-zero σ_i^2 -sub-Gaussian ($i = 1, \dots, m$) is $\sum_{i=1}^m \sigma_i^2$ -sub-Gaussian. (1) and (2) indicate that given \mathbf{x}^k , \bar{w}_j^k is σ_∞^2/m -sub-Gaussian, for any $j = 1, \dots, d$. Therefore, according to the property of sub-Gaussian, we have that $\mathbb{E}[\exp((\bar{w}_j^k)^2/(6\sigma_\infty^2/m)) | \mathbf{x}^k] \leq 2$. Then, for any $t > 0$,

$$\begin{aligned} & \exp(t\mathbb{E}[\|\bar{w}^k\|_\infty^2 | \mathbf{x}^k]) \\ & \leq \mathbb{E}[\exp(t \max_{1 \leq j \leq d} (\bar{w}_j^k)^2) | \mathbf{x}^k] \\ & = \mathbb{E}[\max_{1 \leq j \leq d} \exp(t(\bar{w}_j^k)^2) | \mathbf{x}^k] \\ & \leq \mathbb{E}[\sum_{j=1}^d \exp(t(\bar{w}_j^k)^2) | \mathbf{x}^k] \\ & = \sum_{j=1}^d \mathbb{E}[\exp(t(\bar{w}_j^k)^2) | \mathbf{x}^k], \end{aligned}$$

where the first inequality is from Jensen's inequality. If we let $t = m/(6\sigma_\infty^2)$, then

$$\exp\left(\frac{m}{6\sigma_\infty^2} \mathbb{E}[\|\bar{w}^k\|_\infty^2 | \mathbf{x}^k]\right) \leq 2d \implies \mathbb{E}[\|\bar{w}^k\|_\infty^2 | \mathbf{x}^k] \leq 6 \log(2d)\sigma_\infty^2/m,$$

and we proved (3). □

Proof of Lemma 6.

Proof If we let

$$B^k(\cdot) := \frac{1}{m_k} \sum_{i=1}^{m_k} \nabla f(\cdot, \zeta_i^k),$$

we then have for any k ,

$$\begin{aligned} & \mathbb{E}[\|\nabla f(\mathbf{x}^k) - B^k(\mathbf{x}^k) + B^k(\mathbf{x}^{n_k}) - \nabla f(\mathbf{x}^{n_k})\|^2 | \mathbf{x}^k, \mathbf{x}^{n_k}] \\ & = \mathbb{E}[\|\nabla f(\mathbf{x}^k) - \nabla f(\mathbf{x}^k, \zeta_1^k) + \nabla f(\mathbf{x}^{n_k}, \zeta_1^k) - \nabla f(\mathbf{x}^{n_k})\|^2 | \mathbf{x}^k, \mathbf{x}^{n_k}]/m_k, \end{aligned} \quad (62)$$

since $\nabla f(\mathbf{x}^k, \zeta_i^k) - \nabla f(\mathbf{x}^{n_k}, \zeta_i^k)$, $i = 1, \dots, m_k$ are i.i.d. with mean $\nabla f(\mathbf{x}^k) - \nabla f(\mathbf{x}^{n_k})$ given \mathbf{x}^k and \mathbf{x}^{n_k} . Therefore,

$$\begin{aligned}
& \mathbb{E}[\|\nabla f(\mathbf{x}^k) - G^k\|_\infty^2] \\
&= \mathbb{E}[\|\nabla f(\mathbf{x}^k) - B^k(\mathbf{x}^k) + B^k(\mathbf{x}^{n_k}) - G^{n_k}\|_\infty^2] \\
&= \mathbb{E}[\|\nabla f(\mathbf{x}^k) - B^k(\mathbf{x}^k) + B^k(\mathbf{x}^{n_k}) - G^{n_k} - \nabla f(\mathbf{x}^{n_k}) + \nabla f(\mathbf{x}^{n_k})\|_\infty^2] \\
&\leq 2\mathbb{E}[\|\nabla f(\mathbf{x}^k) - B^k(\mathbf{x}^k) + B^k(\mathbf{x}^{n_k}) - \nabla f(\mathbf{x}^{n_k})\|^2] + 2\mathbb{E}[\| -G^{n_k} + \nabla f(\mathbf{x}^{n_k})\|_\infty^2] \\
&\stackrel{(33), (62)}{\leq} 2\mathbb{E}[\mathbb{E}[\|\nabla f(\mathbf{x}^k) - \nabla f(\mathbf{x}^{n_k}) - \nabla f(\mathbf{x}^k, \zeta_1^k) + \nabla f(\mathbf{x}^{n_k}, \zeta_1^k)\|^2 \mid \mathbf{x}^k, \mathbf{x}^{n_k}]/m_k] \\
&+ \frac{2c(\log d)\sigma_\infty^2}{m_{n_k}} \\
&\leq \frac{4}{m_k} \mathbb{E}[\mathbb{E}[\|\nabla f(\mathbf{x}^k) - \nabla f(\mathbf{x}^{n_k})\|^2 + \| -\nabla f(\mathbf{x}^k, \zeta_1^k) + \nabla f(\mathbf{x}^{n_k}, \zeta_1^k)\|^2 \mid \mathbf{x}^k, \mathbf{x}^{n_k}]] \\
&+ \frac{2c(\log d)\sigma_\infty^2}{m_{n_k}} \\
&= \frac{4}{m_k} \mathbb{E}[\|\nabla f(\mathbf{x}^k) - \nabla f(\mathbf{x}^{n_k})\|^2] + 4\mathbb{E}[\mathbb{E}[\| -\nabla f(\mathbf{x}^k, \zeta_1^k) + \nabla f(\mathbf{x}^{n_k}, \zeta_1^k)\|^2 \mid \mathbf{x}^k, \mathbf{x}^{n_k}]/m_k] \\
&+ \frac{2c(\log d)\sigma_\infty^2}{m_{n_k}} \\
&\leq \frac{4L^2}{m_k} \mathbb{E}[\|\mathbf{x}^k - \mathbf{x}^{n_k}\|^2] + 4\mathbb{E}[\mathbb{E}[L^2\|\mathbf{x}^k - \mathbf{x}^{n_k}\|^2 \mid \mathbf{x}^k, \mathbf{x}^{n_k}]/m_k] + \frac{2c(\log d)\sigma_\infty^2}{m_{n_k}} \\
&= \frac{8L^2}{m_k} \mathbb{E}[\|\mathbf{x}^k - \mathbf{x}^{n_k}\|^2] + \frac{2c(\log d)\sigma_\infty^2}{m_{n_k}}. \tag{63}
\end{aligned}$$

Note that for $t \geq n_k$,

$$\begin{aligned}
\|\mathbf{x}^t - \mathbf{x}^{n_k}\|^2 &= \left\| \sum_{i=n_k}^{t-1} (\mathbf{x}^{i+1} - \mathbf{x}^i) \right\|^2 = (t - n_k)^2 \left\| \frac{1}{t - n_k} \sum_{i=n_k}^{t-1} (\mathbf{x}^{i+1} - \mathbf{x}^i) \right\|^2 \\
&\leq (t - n_k)^2 \cdot \frac{1}{t - n_k} \sum_{i=n_k}^{t-1} \|\mathbf{x}^{i+1} - \mathbf{x}^i\|^2 = (t - n_k) \sum_{i=n_k}^{t-1} \|\mathbf{x}^{i+1} - \mathbf{x}^i\|^2. \tag{64}
\end{aligned}$$

Therefore, for any k ,

$$\begin{aligned}
\sum_{t=n_k}^k \mathbb{E}[\|\mathbf{x}^t - \mathbf{x}^{n_k}\|^2] &\stackrel{(64)}{\leq} \sum_{t=n_k}^k (t - n_k) \sum_{i=n_k}^{t-1} \mathbb{E}[\|\mathbf{x}^{i+1} - \mathbf{x}^i\|^2] \\
&\leq \sum_{t=n_k}^{k-1} \frac{(1+k-n_k)(k-n_k)}{2} \mathbb{E}[\|\mathbf{x}^{t+1} - \mathbf{x}^t\|^2] \\
&\leq q^2 \sum_{t=n_k}^{k-1} \mathbb{E}[\|\mathbf{x}^{t+1} - \mathbf{x}^t\|^2] \\
&\leq q^2 \sum_{t=n_k}^k \mathbb{E}[\|\mathbf{x}^{t+1} - \mathbf{x}^t\|^2]. \tag{65}
\end{aligned}$$

Note that we assume $m_k \equiv m$ if $k \neq n_k$ and $m_{n_k} \equiv m_1$. Therefore,

$$\begin{aligned}
& \mathbb{E}[\|\nabla f(\mathbf{x}^Y) - G^Y\|_\infty^2] \\
&= \frac{1}{K} \sum_{k=1}^K \mathbb{E}[\|\nabla f(\mathbf{x}^k) - G^k\|_\infty^2]
\end{aligned}$$

$$\begin{aligned}
&\stackrel{(63)}{\leq} \frac{8L^2}{Km} \sum_{k=1}^K \mathbb{E}[\|\mathbf{x}^k - \mathbf{x}^{n_k}\|^2] + \frac{2c(\log d)\sigma_\infty^2}{m_1} \\
&\stackrel{(65)}{\leq} \frac{8L^2q^2}{Km} \sum_{k=1}^K \mathbb{E}[\|\mathbf{x}^{k+1} - \mathbf{x}^k\|^2] + \frac{2c(\log d)\sigma_\infty^2}{m_1} \\
&= \frac{8L^2q^2}{m} \mathbb{E}[\|\mathbf{x}^{Y+1} - \mathbf{x}^Y\|^2] + \frac{2c(\log d)\sigma_\infty^2}{m_1}.
\end{aligned}$$

□

Proof of Theorem 2.

Proof Note that similar to deriving (41), (36)(22)(39)(40) lead to

$$\begin{aligned}
&\left(\frac{1}{\eta} - \frac{L}{2} - \frac{\gamma}{2\eta}\right) \mathbb{E}[\|\Delta\mathbf{x}^{Y+1}\|^2] + \frac{\hat{\rho}}{8\eta} \mathbb{E}[\|\mathbf{y}^{Y+1} - \mathbf{x}^Y\|_1^2] \\
&\leq \frac{\Delta}{K} + \frac{\tau+t}{2} \mathbb{E}[\|g^Y - G^Y\|_\infty^2] + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} + \frac{2\hat{\rho}}{\eta}\right) \hat{\epsilon}^2 \\
&\stackrel{(35)}{\leq} \frac{\Delta}{K} + \frac{\tau+t}{2} \left(\frac{8L^2q^2}{m} \mathbb{E}[\|\mathbf{x}^{Y+1} - \mathbf{x}^Y\|^2] + \frac{2c(\log d)\sigma_\infty^2}{m_1}\right) + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} + \frac{2\hat{\rho}}{\eta}\right) \hat{\epsilon}^2 \\
&\implies \left(\frac{1}{\eta} - \frac{L}{2} - \frac{\gamma}{2\eta} - \frac{4(\tau+t)L^2q^2}{m}\right) \mathbb{E}[\|\Delta\mathbf{x}^{Y+1}\|^2] + \frac{\hat{\rho}}{8\eta} \mathbb{E}[\|\mathbf{y}^{Y+1} - \mathbf{x}^Y\|_1^2] \\
&\leq \frac{\Delta}{K} + \frac{(\tau+t)c(\log d)\sigma_\infty^2}{m_1} + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} + \frac{2\hat{\rho}}{\eta}\right) \hat{\epsilon}^2. \tag{66}
\end{aligned}$$

By (35)(37)(39)(66),

$$\begin{aligned}
&\mathbb{E}[\text{dist}_{\|\cdot\|_\infty}(0, \partial(f + \delta_X)(\mathbf{x}^{Y+1}))] \\
&\leq \left(L + \frac{1}{\eta}\right) \sqrt{\mathbb{E}[\|\Delta\mathbf{x}^{Y+1}\|^2]} + \sqrt{\mathbb{E}[\|g^Y - G^Y\|_\infty^2]} + \frac{1}{\eta} \mathbb{E}[\|\xi^{Y+1}\|_\infty] + \frac{\hat{\epsilon}}{\eta} \\
&\leq \left(L + \frac{1}{\eta}\right) \sqrt{\mathbb{E}[\|\Delta\mathbf{x}^{Y+1}\|^2]} + \sqrt{\frac{8L^2q^2}{m} \mathbb{E}[\|\mathbf{x}^{Y+1} - \mathbf{x}^Y\|^2] + \frac{2c(\log d)\sigma_\infty^2}{m_1}} \\
&+ \frac{\hat{\rho}}{\eta} \mathbb{E}[\|\mathbf{y}^{Y+1} - \mathbf{x}^Y\|_1] + \frac{\hat{\epsilon}}{\eta} \\
&\leq \left(L + \frac{1}{\eta} + \sqrt{\frac{8L^2q^2}{m}}\right) \sqrt{\mathbb{E}[\|\Delta\mathbf{x}^{Y+1}\|^2]} + \sqrt{\frac{2c(\log d)\sigma_\infty^2}{m_1}} + \frac{\hat{\rho}}{\eta} \mathbb{E}[\|\mathbf{y}^{Y+1} - \mathbf{x}^Y\|_1] + \frac{\hat{\epsilon}}{\eta} \\
&\leq \left(L + \frac{1}{\eta} + R\sqrt{\frac{8L^2q^2}{m}}\right) \sqrt{\frac{\frac{\Delta}{K} + \frac{(\tau+t)c(\log d)\sigma_\infty^2}{m_1} + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} + \frac{2\hat{\rho}}{\eta}\right) \hat{\epsilon}^2}{\frac{1}{\eta} - \frac{L}{2} - \frac{\gamma}{2\eta} - \frac{4(\tau+t)L^2q^2}{m}}} \\
&+ \sqrt{\frac{2c(\log d)\sigma_\infty^2}{m_1}} + \frac{\hat{\rho}}{\eta} \sqrt{\frac{\frac{\Delta}{K} + \frac{(\tau+t)c(\log d)\sigma_\infty^2}{m_1} + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} + \frac{2\hat{\rho}}{\eta}\right) \hat{\epsilon}^2}{\hat{\rho}/(8\eta)}} + \frac{\hat{\epsilon}}{\eta}.
\end{aligned}$$

□

Proof of Theorem 4.

Proof Similar to the derivation of (48), we have

$$\begin{aligned}
&\left(\frac{1}{\eta} - \frac{L}{2} - \frac{\gamma}{2\eta}\right) \mathbb{E}[\|\Delta\mathbf{x}^{Y+1}\|^2] + \frac{\psi - \hat{\epsilon}}{\eta} \mathbb{E}[\hat{\rho}_Y] \\
&\leq \frac{\Delta}{K} + \frac{\tau+t}{2} \mathbb{E}[\|g^Y - G^Y\|_\infty^2] + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma}\right) \hat{\epsilon}^2 + \frac{\psi^2}{2t}
\end{aligned}$$

$$\begin{aligned}
&\stackrel{(35)}{\leq} \frac{\Delta}{K} + \frac{\tau+t}{2} \left(\frac{8L^2q^2}{m} \mathbb{E}[\|\mathbf{x}^{Y+1} - \mathbf{x}^Y\|^2] + \frac{2c(\log d)\sigma_\infty^2}{m_1} \right) + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} \right) \hat{\epsilon}^2 + \frac{\psi^2}{2t} \\
\implies &\left(\frac{1}{\eta} - \frac{L}{2} - \frac{\gamma}{2\eta} - \frac{4(\tau+t)L^2q^2}{m} \right) \mathbb{E}[\|\Delta\mathbf{x}^{Y+1}\|^2] + \frac{\psi - \hat{\epsilon}}{\eta} \mathbb{E}[\hat{\rho}_Y] \\
&\leq \frac{\Delta}{K} + \frac{(\tau+t)c(\log d)\sigma_\infty^2}{m_1} + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} \right) \hat{\epsilon}^2 + \frac{\psi^2}{2t} \tag{67}
\end{aligned}$$

Then by (35)(37),

$$\begin{aligned}
&\mathbb{E}[\text{dist}_{\|\cdot\|_\infty}(0, \partial(f + \delta_X)(\mathbf{x}^{Y+1}))] \\
&\leq \left(L + \frac{1}{\eta} \right) \sqrt{\mathbb{E}[\|\Delta\mathbf{x}^{Y+1}\|^2]} + \sqrt{\mathbb{E}[\|g^Y - G^Y\|_\infty^2]} + \frac{1}{\eta} \mathbb{E}[\|\xi^{Y+1}\|_\infty] + \frac{\hat{\epsilon}}{\eta} \\
&\leq \left(L + \frac{1}{\eta} \right) \sqrt{\mathbb{E}[\|\Delta\mathbf{x}^{Y+1}\|^2]} + \sqrt{\frac{8L^2q^2}{m} \mathbb{E}[\|\mathbf{x}^{Y+1} - \mathbf{x}^Y\|^2] + \frac{2c(\log d)\sigma_\infty^2}{m_1}} + \frac{1}{\eta} \mathbb{E}[\|\xi^{Y+1}\|_\infty] \\
&\quad + \frac{\hat{\epsilon}}{\eta} \\
&\stackrel{(46)}{\leq} \left(L + \frac{1}{\eta} + \sqrt{\frac{8L^2q^2}{m}} \right) \sqrt{\mathbb{E}[\|\Delta\mathbf{x}^{Y+1}\|^2]} + \sqrt{\frac{2c(\log d)\sigma_\infty^2}{m_1} + \frac{1}{\eta} \mathbb{E}[\hat{\rho}_Y]} + \frac{\hat{\epsilon}}{\eta} \\
&\stackrel{(67)}{\leq} \left(L + \frac{1}{\eta} + \sqrt{\frac{8L^2q^2}{m}} \right) \sqrt{\frac{\frac{\Delta}{K} + \frac{(\tau+t)c(\log d)\sigma_\infty^2}{m_1} + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} \right) \hat{\epsilon}^2 + \frac{\psi^2}{2t}}{\frac{1}{\eta} - \frac{L}{2} - \frac{\gamma}{2\eta} - \frac{4(\tau+t)L^2q^2}{m}}} \\
&\quad + \sqrt{\frac{2c(\log d)\sigma_\infty^2}{m_1} + \frac{1}{\psi - \hat{\epsilon}}} \left(\frac{\Delta}{K} + \frac{(\tau+t)c(\log d)\sigma_\infty^2}{m_1} + \left(\frac{1}{2\tau} + \frac{1}{2\eta\gamma} \right) \hat{\epsilon}^2 + \frac{\psi^2}{2t} \right) + \frac{\hat{\epsilon}}{\eta}.
\end{aligned}$$

□

B Algorithm

Here we provide the details of the related algorithms: Alternating direction method of multiplier (ADMM) to solve (18)(19), gradient descent with backtracking (Algorithm 3) used for computing f^* in the numerical experiment.

Algorithm 2 Alternating direction method of multiplier (ADMM) to solve (18)(19)

Step 1. Initialize primal variable \mathbf{x}^1 , dual variable λ^1 , penalty parameter $\rho > 0$ and accuracy parameter $\hat{\epsilon} > 0$.

Step 2. Invoke the following operations for $k = 1, 2, \dots$

Step 2.1 $\mathbf{x}^{k+1} = \arg \min_{\mathbf{x} \in X} \frac{1}{2} \|\mathbf{x} - (\mathbf{x}^k - \eta G^k)\|^2 + (\lambda^k)^T (\mathbf{x} - \mathbf{y}^k) + \frac{\rho}{2} \|\mathbf{x} - \mathbf{y}^k\|^2$

Step 2.2 $\mathbf{y}^{k+1} = \arg \min_{\mathbf{y}} \phi(\mathbf{y} - \mathbf{x}^k) + (\lambda^k)^T (\mathbf{x}^{k+1} - \mathbf{y}) + \frac{\rho}{2} \|\mathbf{x}^{k+1} - \mathbf{y}\|^2$

Step 2.3 $\lambda^{k+1} = \lambda^k + \rho(\mathbf{x}^{k+1} - \mathbf{y}^{k+1})$.

Step 2.4 Stop when $\rho \|\mathbf{y}^{k+1} - \mathbf{y}^k\|_\infty \leq \hat{\epsilon}$ and $\|\mathbf{x}^{k+1} - \mathbf{y}^{k+1}\|_1 \leq \hat{\epsilon}$.

Step 3. Output the triple $(\mathbf{x}^{k+1}, \mathbf{y}^{k+1}, \lambda^{k+1})$.

Algorithm 3 Projected gradient method with backtracking

Initialization Step Set $c_1 = \frac{1}{4}, \beta = \frac{1}{2}, \epsilon = 1e - 10$. Initialize $\mathbf{x}^0 = \mathbf{0}$.

Main Step Set $\alpha_k \equiv \alpha = 1$;

While $f(P_X(\mathbf{x}^k - \alpha_k \nabla f(\mathbf{x}^k))) > f(\mathbf{x}^k) + c_1 (\nabla f(\mathbf{x}^k))^T (P_X(\mathbf{x}^k - \alpha_k \nabla f(\mathbf{x}^k)) - \mathbf{x}^k)$

$\alpha_k = \beta \alpha_k$

Stopping Stop if $\|\mathbf{x}^{k+1} - \mathbf{x}^k\|_1 \leq \epsilon$ and output \mathbf{x}^k ; otherwise replace k by $k + 1$ and repeat the Main Step.
