

Subgroup Identification with Latent Factor Structure

Yong He^a, Dong Liu^b, Fuxin Wang^c, Mingjuan Zhang^d, Wen-Xin Zhou^e

^a*Institute for Financial Studies, Shandong University, Jinan, 250100, China*

^b*Shanghai University of Finance and Economics, Shanghai, 200433, China*

^c*University of Wisconsin-Madison, Madison, WI 53706, USA*

^d*School of Statistics and Mathematics, Shanghai Lixin University of Accounting and Finance, Shanghai 201209, China*

^e*Department of Information and Decision Sciences, University of Illinois at Chicago, Chicago, IL 60607, USA*

Abstract

Subgroup analysis has garnered increasing attention for its ability to identify meaningful subgroups within heterogeneous populations, thereby enhancing predictive power. However, in many fields such as social science and biology, covariates are often highly correlated due to common factors. This correlation poses significant challenges for subgroup identification, an issue that is often overlooked in existing literature. In this paper, we aim to address this gap in the “diverging dimension” regime by proposing a center-augmented subgroup identification method within the Factor Augmented (sparse) Linear Model framework. This method bridges dimension reduction and sparse regression. Our proposed approach is adaptable to the high cross-sectional dependence among covariates and offers computational advantages with a complexity of $O(nK)$, compared to the $O(n^2)$ complexity of the conventional pairwise fusion penalty method in the literature, where n is the sample size and K is the number of subgroups. We also investigate the asymptotic properties of the oracle estimators under conditions on the minimal distance between group centroids. To implement the proposed approach, we introduce a Difference of Convex functions-based Alternating Direction Method of Multipliers (DC-ADMM) algorithm and demonstrate its convergence to a local minimizer in a finite number of steps. We illustrate the superiority of the proposed method through extensive numerical experiments and a real macroeconomic data example. An R package, SILFS, implementing the method is also available on CRAN¹.

Keywords: Subgroup Analysis; Factor Augmented Sparse Linear Model; Oracle Property; Center-augmented Regularization

1. Introduction

With advancements in data collection and information technology, the dimensionality of data has exponentially increased across various research and application fields. Concurrently, it is believed that meaningful subgroups exist within heterogeneous populations in many real datasets. In macroeconomics, [Phillips and](#)

¹<https://cran.r-project.org/web/packages/SILFS/index.html>

Sul (2007) analyzes the convergence in the cost of living indices among 19 U.S. metropolitan cities, demonstrating that cities within different groups behave quite differently. In portfolio allocations, stocks within the same industries exhibit similar characteristics (Livingston, 1977). In precision medicine, patients may belong to various latent groups, and those within the same group can be considered together for making treatment decisions; see, for example, Ma and Huang (2017) and Chen et al. (2021a). Wang and Su (2021) also points out that geographic adjacency is a natural criterion for grouping when analyzing international trade and economic geography datasets. These examples underline the importance of identifying latent group structures prior to conducting statistical inference, as this approach significantly enhances statistical efficiency.

Statisticians and econometricians typically characterize group structures using group-specific parameters in statistical models. This topic is closely related to concepts such as integrative analysis, transfer learning, and multi-task representation learning. For integrative analysis, Gertheiss and Tutz (2012) and Ollier and Viallon (2017) assume that the coefficients of the k -th group, denoted as β_k , can be decomposed into ω and ζ_k , where ω is a common parameter shared by all groups, and ζ_k is the group-specific parameter representing the heterogeneity of group structures. Similarly, in the context of transfer learning or multi-task representation learning, Li et al. (2022) and Tian et al. (2023) also use group-specific parameters to characterize the heterogeneity of group structures. However, it is worth noting that the aforementioned works presume the group membership is known a priori, which is not the case in many real applications. In the context of subgroup analysis, Ma and Huang (2017), Zhang et al. (2019) and He et al. (2022c) employ group-specific regression coefficients while assuming unknown group membership. They aim to cluster the samples and conduct statistical inference simultaneously. In this paper, we also adopt the unknown group membership framework and consider the following linear model with group-specific intercepts:

$$Y_i = \alpha_i + \mathbf{x}_i^\top \boldsymbol{\beta} + \epsilon_i, \quad i = 1, 2, \dots, n, \quad \text{or equivalently, } \mathbf{Y} = \boldsymbol{\alpha} + \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}, \quad (1.1)$$

where $\mathbf{Y} = (Y_1, Y_2, \dots, Y_n)^\top$ is the response vector, $\mathbf{X} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)^\top$ denotes the design matrix, $\boldsymbol{\epsilon} = (\epsilon_1, \epsilon_2, \dots, \epsilon_n)^\top$ is the noise vector, and the unknown vector of regression coefficients $\boldsymbol{\beta} \in \mathbb{R}^p$ is sparse. The group structure is characterized by the intercept parameters $\boldsymbol{\alpha} = (\alpha_1, \alpha_2, \dots, \alpha_n)^\top$, meaning that individuals within the same group share the same group-specific intercept parameter. Specifically, let the number of subgroups be K , and define $\mathcal{G} = \{\mathcal{G}_k\}_{k=1}^K$ as a partition of $\{1, 2, \dots, n\}$, satisfying $\mathcal{G}_k \cap \mathcal{G}_{k'} = \emptyset$ for $k \neq k'$. It is assumed that $\alpha_i = \gamma_k$ for i in \mathcal{G}_k , and $\boldsymbol{\gamma} = (\gamma_1, \gamma_2, \dots, \gamma_K)^\top$ represents the centroid of the group-specific parameters. It is worth noting that α_i may reflect heterogeneity driven by latent variables \mathbf{z}_i , that is, $\alpha_i = \alpha + \mathbf{z}_i^\top \boldsymbol{\theta}$. For example, in microeconomics, the impact of education on wages is a widely studied topic. However, individuals may have different backgrounds, such as family background and work experience, which can be modeled as \mathbf{z}_i . See, for example, Ashenfelter and Krueger (1994), Michele and Francesco (2018)

and [Connolly and Gottschalk \(2006\)](#). More generally, the coefficients for \mathbf{z}_i can be subject-specific, as the same family background and work experience may have different impacts for different individuals. In this case, the model would become

$$Y_i = \alpha + \mathbf{z}_i^\top \boldsymbol{\theta}_i + \mathbf{x}_i^\top \boldsymbol{\beta} + \epsilon_i, \quad i = 1, 2, \dots, n, \quad (1.2)$$

Throughout this article, we focus on the model [\(1.1\)](#), assuming that the heterogeneity arises from unobserved covariates \mathbf{z}_i . In fact, model [\(1.2\)](#) is similar to model [\(1.1\)](#), and our proposed algorithms and the associated theoretical properties for model [\(1.1\)](#) can be extended to model [\(1.2\)](#) with slight modifications.

In datasets from social science and biology, high-dimensional covariates are often highly correlated, possibly due to the existence of common factors. See, for example, [Stock and Watson \(2002\)](#), [Bai and Ng \(2002\)](#), [Fan et al. \(2020\)](#), [Vatcheva et al. \(2016\)](#), [Porcu et al. \(2019\)](#) and [He et al. \(2022a\)](#). The high cross-sectional dependence among covariates can lead to unsatisfactory results in various statistical inference problems, including the subgroup identification problem discussed in this paper. In the following, we present a toy example to illustrate the impacts of high collinearity on the subgroup identification problem. For model [\(1.1\)](#), each group-specific α_i takes value of either 1 or -1 with equal probability. We generate the residuals $\boldsymbol{\epsilon}$ from $\mathcal{N}(0, 0.01\mathbf{I}_p)$, and let $n = p = 100$. Set $\boldsymbol{\beta} = (\beta_1, \dots, \beta_{10}, \mathbf{0}_{p-10}^\top)^\top$, where the nonzero coefficients β_i ($1 \leq i \leq 10$) are drawn from $\mathcal{U}(2, 5)$. Let $\mathbf{X} = (\mathbf{x}_1, \dots, \mathbf{x}_n)^\top$ be n independent copies from multivariate normal distribution $\mathcal{N}(\mathbf{0}, \boldsymbol{\Xi})$, where $\boldsymbol{\Xi} \in \mathbb{R}^{p \times p}$ is a covariance matrix with off-diagonal elements being ρ and diagonal elements being 1. In this toy example, the parameter ρ is varied over the interval $[0, 0.95]$ with a step size of 0.05 to illustrate different strengths of cross-sectional dependency. For each given ρ , we consider the Center-augmented Regularizer (CAR) method proposed by [He et al. \(2022c\)](#), referred to as Standard CAR (S-CAR). We report the averages of model size, estimation errors (MSE), and Rand Index over 100 replications in [Figure 1](#). As illustrated in the [Figure 1](#) (the blue lines), both the clustering performance and the estimation accuracy of S-CAR deteriorate as the cross-sectional dependency strength, ρ , increases. The CAR method shows limited power in detecting group structures under high cross-sectional dependency, especially when $\rho > 0.7$.

To the best of our knowledge, there exists no literature addressing subgroup analysis under conditions of high collinearity among covariates, due to both computational and statistical challenges. In this work, we propose to address the high cross-sectional dependency among covariates using a factor model approach, inspired by the Factor Augmented (sparse linear) Regression Model (abbreviated as FARM) introduced in [Fan et al. \(2020\)](#) and [Fan et al. \(2023\)](#). Factor model is widely used as an effective tool for dimension reduction and can capture various levels of cross-sectional dependence. It is well recognized that many macroeconomic and financial datasets exhibit high cross-sectional dependence. For instance, [Fama and French \(1992\)](#) and [Fama and French \(2015\)](#) provide evidence that average stock returns, characterized by high cross-sectional




Figure 1: Comparison of line charts of estimation results for S-CAR and SILFS models based on 100 replications. The model size is defined as the number of non-zero components in $\hat{\beta}$.

dependence, are driven by Fama-French factors. Additionally, [McCracken and Ng \(2016\)](#) confirm that the well-studied microeconomic dataset FRED-MD also exhibits latent factor structures. [Johnstone and Paul \(2018\)](#) investigate numerous datasets showing spiked structures in covariance matrices across various fields, including microarrays, satellite images, medical shapes, climate data, and signal detection. Therefore, it is reasonable to mitigate high collinearity and achieve dimension reduction simultaneously by assuming a latent factor model. In this paper, we adopt the FARM framework to address the challenges posed by high collinearity and high-dimensionality in the subgroup analysis problem. More specifically, we assume the vector of regression coefficients β is sparse with a support set \mathcal{S} , and the observed \mathbf{x}_i 's satisfy the following factor structure:

$$\mathbf{x}_i = \mathbf{B}\mathbf{f}_i + \mathbf{u}_i, \quad i = 1, 2, \dots, n, \quad \text{or in matrix form, } \mathbf{X} = \mathbf{F}\mathbf{B}^\top + \mathbf{U}, \quad (1.3)$$

where $\mathbf{f}_i \in \mathbb{R}^r$, $\mathbf{F} = (\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_n)^\top$ is the $n \times r$ factor score matrix, \mathbf{B} is the $p \times r$ factor loading matrix, and $\mathbf{U} = (\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n)^\top$ is the matrix of idiosyncratic errors of dimension $n \times p$. To illustrate the advantage of the FARM framework in the presence of high collinearity, we substitute (1.3) into (1.1) to form the following sparse regression model

$$Y_i = \alpha_i + \mathbf{f}_i^\top \boldsymbol{\theta} + \mathbf{u}_i^\top \boldsymbol{\beta} + \epsilon_i, \quad i = 1, 2, \dots, n, \quad \text{or in matrix form, } \mathbf{Y} = \boldsymbol{\alpha} + \mathbf{F}\boldsymbol{\theta} + \mathbf{U}\boldsymbol{\beta} + \boldsymbol{\epsilon}, \quad (1.4)$$

where $\boldsymbol{\theta} = \mathbf{B}^\top \boldsymbol{\beta}$ and $\mathbf{F} \in \mathbb{R}^{n \times r}$ extract essential information from \mathbf{X} , and $\mathbf{U} \in \mathbb{R}^{n \times p}$ contributes additional sparse information to the response. The identifiability condition on \mathbf{B} and \mathbf{F} will be discussed in Section 2.1. This construction augments the highly cross-dependent covariates $\mathbf{X} \in \mathbb{R}^{n \times p}$ to weakly dependent covariates $\mathbf{F}, \mathbf{U} \in \mathbb{R}^{n \times (p+r)}$. Subsequently, we adopt the FARM framework to capture the cross-sectional dependence of covariates in subgroup analysis. Our goal is to identify latent groups characterized by the

intercept parameters $\boldsymbol{\alpha} = (\alpha_1, \alpha_2, \dots, \alpha_n)^\top$ and achieve variable selection simultaneously under model (1.4). Hereafter, we treat the group number K and factor number r as fixed, while allowing the dimension p to increase with the sample size n .

1.1. Literature Review

A closely related line of research focuses on sparse linear regression and variable selection. Over the last two decades, numerous regularization-based methods have been proposed, including well-known approaches such as LASSO (Tibshirani, 1996), SCAD (Fan and Li, 2001), elastic net (Zou and Hastie, 2005) and adaptive LASSO (Zou, 2006). The effectiveness of regularized regression methods often relies on constraints imposed on the covariance matrix of the covariates and limitations on collinearity levels. However, these constraints may be inappropriate in real macroeconomic or financial applications.

Another closely related line of research focuses on large-dimensional approximate factor models. Bai (2003) explored the theory of large-dimensional approximate factor models, and theoretical guarantees for determining the number of factors can be found in works such as Bai and Ng (2002) and Ahn and Horenstein (2013). Robust factor analysis has also gained increasing attention in recent years, as evidenced by studies like Chen et al. (2021b), He et al. (2022b) and He et al. (2023). Furthermore, Stock and Watson (2002) demonstrates the consistency of Ordinary Least Squares (OLS) estimates in factor-augmented regressions, treating latent factors as predictors. Ando and Tsay (2011) extends this approach to quantile factor-augmented regressions, with further extensions discussed in Wang et al. (2019) and Fan et al. (2021). In high-dimensional regression settings, latent factors often cannot fully explain the response, particularly when their effect is weak. Fan et al. (2020) proposes the Factor Augmented (sparse linear) Regression Model, which assumes sparsity in the coefficient vector and integrates both latent factors and idiosyncratic components into the linear regression framework. Fan et al. (2023) introduces hypothesis tests for assessing the adequacy of latent factor regression models. Additionally, Tu and Wang (2024) present a new class of information criteria aimed at achieving consistent factor and covariate selection jointly in factor-augmented regression.

The last closely related line of research focuses on subgroup analysis. Typical approaches for subgroup identification can generally be categorized into two distinct types. The first type is the finite mixture model (FMM), which employs model-based clustering techniques. Representative works include but are not limited to Banfield and Raftery (1993), Hastie and Tibshirani (1996), and Wei and Kosorok (2013) for mixture models under Gaussian distributions; Shen and Huang (2010) for logistic-normal regression; Yao et al. (2014) for the Student t distribution; and Song et al. (2014) for the Laplace distribution. For finite mixture models, Khalili and Chen (2007) propose penalized likelihood methods for variable selection and establish their consistency. In high-dimensional cases, references include Zhang et al. (2020) and Wang et al. (2023). Although the EM algorithm is effective and powerful in finding maximum likelihood estimators with latent group

variables in FMM, it suffers from significant computational burdens in high dimensions. The second type is regularization methods, which use fusion penalties to enforce sparsity in pairwise differences among group-specific parameters to achieve clustering. Examples include [Hocking et al. \(2011\)](#), [Pan et al. \(2013\)](#), [Chi and Lange \(2015\)](#), and [Wu et al. \(2016\)](#). For subgroup analysis of linear models, [Ma and Huang \(2017\)](#) propose a concave Pairwise Fusion Penalty (PFP) capable of recovering latent groups without prior information on the number of groups. Many subsequent studies adopt this penalty for its favorable statistical properties, including extensions to quantile regressions ([Zhang et al., 2019](#)) and functional regressions ([Zhang et al., 2022](#)) for group pursuits. From a computational perspective, the PFP, which pairs n samples, has a computational complexity of $O(n^2)$. To mitigate this, [Wang and Zhu \(2024\)](#) introduce a threshold PFP method to reduce computational time and enhance group detection accuracy. Moreover, [He et al. \(2022c\)](#) propose a more efficient Center-augmented Regularization (CAR) method, reducing computational complexity to $O(nK)$ by incorporating group centroids as parameters. However, due to the concavity of PFP and CAR, their performance is sensitive to the selection of initial values. Therefore, choosing appropriate initial values remains a challenging task.

1.2. Contributions and Structure of the Paper

In this work, we propose a clustering method within the FARM framework, coined Subgroup Identification with Latent Factor Structures (SILFS). SILFS demonstrates superior performance compared to existing methods in two critical aspects: identifying group memberships and simultaneously recovering the support of regression coefficients in the presence of high collinearity among covariates. Returning to Figure 1 (the red lines), we observe that SILFS exhibits robust performance across varying levels of cross-sectional dependence parameterized by ρ . Notably, for $\rho > 0.5$, SILFS significantly outperforms the S-CAR method, highlighting the necessity of accounting for high cross-sectional dependency among covariates in subgroup identification problems and the novelty of our approach. Additionally, the SILFS method draws inspiration from Center-augmented Regularization, offering computational advantages over pairwise fusion penalization methods. Theoretically, we establish properties of the local minima for the optimization problem under an ℓ_1 -type CAR penalty. We provide, to the best of our knowledge, the first rigorous theoretical guarantee that CAR can effectively identify subgroups under the ℓ_1 distance. To achieve these local minima, we propose a DC-ADMM algorithm that fully exploits the mathematical structure of the CAR penalty. In implementing the DC decomposition of the CAR penalty under inequality constraints, we convert these inequalities into equivalent equalities by introducing slack variables and incorporating indicator functions into the objective function. We derive explicit iteration formulas for the equivalent optimization problem and demonstrate that our algorithm converges to a Karush-Kuhn-Tucker (KKT) point of the objective function after a finite number of iterations.

The remainder of this paper is organized as follows. Section 2 introduces the optimization problem of

SILFS and presents a two-step algorithm to solve it. In Section 3, we establish the consistency of the oracle estimators and demonstrate that they serve as local minimizers of the proposed optimization problem with probability approaching one. Section 4 presents extensive simulation studies that illustrate the superior performance of the SILFS method in both clustering and variable selection. Furthermore, computational speed comparisons with various methods highlight the computational advantages of our proposed approach. In Section 5, we apply the SILFS method to a large panel of China’s commodity trade data involving all countries worldwide from 2019. We conclude with a discussion on potential future research directions and summarize our findings in Section 6. The supplementary material includes proofs for all theorems and additional details on algorithm implementations.

To conclude this section, we introduce the notations that will be used throughout the remainder of this paper. For a matrix $\mathbf{M} = (M_{ij})_{i \in p, j \in q}$, denote $\|\mathbf{M}\|_{\mathbb{F}}$ and $\|\mathbf{M}\|_2$ as its Frobenius norm and operator norm, respectively. Write $\|\mathbf{M}\|_1 = \max_{j \in \{1, \dots, q\}} \sum_{i=1}^p |M_{ij}|$, $\|\mathbf{M}\|_{\infty} = \max_{i \in \{1, \dots, p\}} \sum_{j=1}^q |M_{ij}|$ and $\|\mathbf{M}\|_{\max} = \max_{i,j} |M_{ij}|$. Let \mathcal{S}_1 and \mathcal{S}_2 be two subsets of $\{1, 2, \dots, p\}$. We denote $\mathbf{M}_{\mathcal{S}_1 \mathcal{S}_2}$ as the submatrix of \mathbf{M} with row indices in \mathcal{S}_1 and column indices in \mathcal{S}_2 . The matrix $\mathbf{M}_{\mathcal{S}}$ denotes the submatrix of \mathbf{M} consisting of columns whose indices belong to the set \mathcal{S} . Here, $|\mathcal{S}_1|$ denotes the cardinality of \mathcal{S}_1 . Moreover, \mathbf{I}_n represents the n -dimensional identity matrix, and $\mathbf{0}_{(p \times q)}$ denotes the zero-matrix of size of $p \times q$. $\mathbf{1}_p$ and $\mathbf{0}_p$ denote the p -dimensional vectors with all elements being 1 and 0, respectively. The function $\text{sgn}(a)$ is defined such that $\text{sgn}(a) = 1$ if $a > 0$, $\text{sgn}(a) = -1$ if $a < 0$, and $\text{sgn}(0) = 0$. For a set of indices \mathcal{S} and a function $f(\mathbf{x})$ where $\mathbf{x} \in \mathbb{R}^n$, $\nabla_{\mathcal{S}} f(\mathbf{x})$ denotes the sub-gradient of $f(\mathbf{x})$ with respect to $\mathbf{x}_{\mathcal{S}}$, where $\mathbf{x}_{\mathcal{S}}$ consists of elements of \mathbf{x} indexed by \mathcal{S} . For two sequences of real numbers $\{a_n\}_{n=1}^{\infty}$ and $\{b_n\}_{n=1}^{\infty}$, we use the notation $a_n = o(b_n)$ or $b_n \gg a_n$ to indicate that $a_n/b_n \rightarrow 0$ as $n \rightarrow \infty$, and $a_n = O(b_n)$ if there exist a positive integer N and a positive constant C such that $|a_n| \leq C|b_n|$ for all $n \geq N$.

2. Methodology

In this section, we introduce a two-step procedure to estimate the parameters β_i and α_i in model (1.4). In Section 2.1, we first estimate the unobserved \mathbf{F} and \mathbf{U} . Subsequently, in Section 2.2, we formulate a center-augmented regularization optimization problem using plug-in estimators to achieve simultaneous clustering and variable selection. Section 2.3 discusses implementation details of the algorithm, including the selection of initial values, the number of subgroups K , and the shrinkage parameters λ_1 and λ_2 .

2.1. Estimation of Factor Model

Given the presence of an unknown factor structure in model (1.1), our first step is to estimate these factors from the observed predictors \mathbf{X} . To achieve this, we employ Principal Component Analysis (PCA) to estimate the latent factors \mathbf{F} and the factor loading matrix \mathbf{B} . Specifically, Fan et al. (2013) demonstrate

that the PCA method is equivalent to the following constrained least squares:

$$\begin{aligned} (\widehat{\mathbf{F}}, \widehat{\mathbf{B}}) &= \underset{\mathbf{F} \in \mathbb{R}^{n \times r}, \mathbf{B} \in \mathbb{R}^{p \times r}}{\operatorname{argmin}} \|\mathbf{X} - \mathbf{F}\mathbf{B}^\top\|_{\mathbb{F}}^2, \\ \text{s.t.} \quad &\frac{1}{n}\mathbf{F}^\top\mathbf{F} = \mathbf{I}_r, \quad \mathbf{B}^\top\mathbf{B} \text{ is diagonal,} \end{aligned}$$

where the normalization constraints address identifiability concerns (Bai, 2003). Through simple algebra, the estimated factors $\widehat{\mathbf{F}}/\sqrt{n}$ correspond to the eigenvectors associated with the largest r eigenvalues of $\mathbf{X}\mathbf{X}^\top$, and $\widehat{\mathbf{B}} = \widehat{\mathbf{F}}^\top\mathbf{X}/n$. The idiosyncratic components can be computed straightforwardly as $\widehat{\mathbf{U}} = \mathbf{X} - \widehat{\mathbf{F}}\widehat{\mathbf{B}}^\top$.

However, the number of latent factors r is unknown in practice. There are various methods in the literature to derive consistent estimators of r , such as the Information Criteria (IC) proposed by Bai and Ng (2002) and the Eigenvalue-Ratio (ER) criterion proposed by Lam and Yao (2012) and Ahn and Horenstein (2013). Since the estimation of r is often conducted separately, we assume that the number of factors r is given.

2.2. Clustering Procedure

To recovery the group membership and estimate the regression coefficients $\boldsymbol{\beta}$ simultaneously, we plug in the PCA estimators $\widehat{\mathbf{F}}$ and $\widehat{\mathbf{U}}$ and formulate the following optimization problem with a given number of subgroups K :

$$\underset{\boldsymbol{\Theta}}{\operatorname{argmin}} Z(\boldsymbol{\Theta}) = \frac{1}{2n} \|\mathbf{Y} - \boldsymbol{\alpha} - \widehat{\mathbf{F}}\boldsymbol{\theta} - \widehat{\mathbf{U}}\boldsymbol{\beta}\|_2^2 + \lambda_1 g(\boldsymbol{\alpha}, \boldsymbol{\gamma}) + \lambda_2 \|\boldsymbol{\beta}\|_1, \quad (2.1)$$

where $\boldsymbol{\Theta} = (\boldsymbol{\alpha}^\top, \boldsymbol{\beta}^\top, \boldsymbol{\theta}^\top, \boldsymbol{\gamma}^\top)^\top$, λ_1 and λ_2 are tuning parameters, and $g(\boldsymbol{\alpha}, \boldsymbol{\gamma})$ is a penalty for group pursuits. The ℓ_1 penalty, $\|\boldsymbol{\beta}\|_1$, encourages sparsity for variable selection.

In this paper, we choose $g(\boldsymbol{\alpha}, \boldsymbol{\gamma})$ as the CAR penalty for group pursuit. The CAR method, initially proposed by He et al. (2022c), aims to estimate both the group-specific parameter $\boldsymbol{\alpha}$ and the group centroids $\boldsymbol{\gamma}$ simultaneously. This is achieved by clustering each subject i into group k based on the nearest distance. Specifically,

$$g(\boldsymbol{\alpha}, \boldsymbol{\gamma}) = \sum_{i=1}^n \min\{d(\alpha_i, \gamma_1), \dots, d(\alpha_i, \gamma_K)\},$$

where $d(x, y)$ is a distance function between x and y . This approach is analogous to the K -means method when using ℓ_2 -type distance ($d(x, y) = (x - y)^2$) or the K -median method when employing ℓ_1 -type distance ($d(x, y) = |x - y|$). From an optimization perspective, CAR exhibits a computational complexity of $O(nK)$, whereas pairwise penalties used in Ma and Huang (2017) and Zhang et al. (2019), such as $\sum_{i < j} p_\lambda(|\alpha_i - \alpha_j|)$ with penalty function $p_\lambda(\cdot)$, require $O(n^2)$ computational complexity. Hence, CAR significantly reduces computational burdens by incorporating centroid parameters. Furthermore, as CAR is a non-convex penalty, it presents challenges for both theoretical analysis and optimization. It is worth noting that CAR does not

directly offer clustering results. If $\hat{\gamma}_k$ represents the estimated group centroids, our clustering rule for the i -th subject is defined as $i \in \mathcal{G}_{k^*}$ if and only if $k^* = \operatorname{argmin}_k d(\hat{\alpha}_i, \hat{\gamma}_k)$ for $k \leq K$.

Recall that the optimization problem outlined in (2.1) depends on the choice of the distance function. In the following, we introduce the DC-ADMM and Cyclic Coordinate Descent (CCD) algorithm to solve the problem for ℓ_1 -type and ℓ_2 -type distances, respectively.

2.2.1. DC-ADMM Algorithm

In this section, we propose the DC-ADMM algorithm to solve (2.1) under the ℓ_1 -type distance. The corresponding theoretical results are elaborated in Theorem 2.1. The optimization problem is as follows:

$$\operatorname{argmin}_{\Theta} Z(\Theta) = \frac{1}{2n} \|\mathbf{Y} - \boldsymbol{\alpha} - \widehat{\mathbf{F}}\boldsymbol{\theta} - \widehat{\mathbf{U}}\boldsymbol{\beta}\|_2^2 + \lambda_1 \sum_{i=1}^n \min\{d(\alpha_i, \gamma_1), \dots, d(\alpha_i, \gamma_K)\} + \lambda_2 \|\boldsymbol{\beta}\|_1. \quad (2.2)$$

This optimization problem is challenging due to the identifiability issue with the parameter $\boldsymbol{\gamma}$ and its non-convex nature. For example, when $K = 2$ and $(\tilde{\gamma}_1, \tilde{\gamma}_2)$ minimizes the loss function in (2.1), it is evident that $(\tilde{\gamma}_2, \tilde{\gamma}_1)$ also serves as a minimizer. To address the identifiability issue, we impose the condition that $\gamma_1 \leq \gamma_2 \leq \dots \leq \gamma_K$, which is also crucial for the following optimization. Motivated by the works of [An and Tao \(2005\)](#) and [Wu et al. \(2016\)](#) on Difference of Convex (DC) programming to address non-convex optimization problems, we reformulate the CAR penalty $g(\boldsymbol{\alpha}, \boldsymbol{\gamma})$ as a difference of convex functions under the identifiability condition of $\boldsymbol{\gamma}$. Specifically, we have $g(\boldsymbol{\alpha}, \boldsymbol{\gamma}) = g_1(\boldsymbol{\alpha}, \boldsymbol{\gamma}) - g_2(\boldsymbol{\alpha}, \boldsymbol{\gamma})$ with

$$g_1(\boldsymbol{\alpha}, \boldsymbol{\gamma}) = \sum_{i=1}^n \sum_{k=1}^K d(\alpha_i, \gamma_k), \quad g_2(\boldsymbol{\alpha}, \boldsymbol{\gamma}) = \sum_{i=1}^n \sum_{k=2}^K \max\{d(\alpha_i, \gamma_{k-1}), d(\alpha_i, \gamma_k)\}.$$

We reparameterize the variables by letting $\delta_{ik} = \alpha_i - \gamma_k$ for $i = 1, \dots, n$ and $k = 1, \dots, K$. Now, both $g_1(\boldsymbol{\delta})$ and $g_2(\boldsymbol{\delta})$ are convex functions with respect to $\boldsymbol{\delta}_i$, where $\boldsymbol{\delta} = (\boldsymbol{\delta}_1^\top, \dots, \boldsymbol{\delta}_n^\top)^\top$ with $\boldsymbol{\delta}_i = (\delta_{i1}, \dots, \delta_{iK})^\top$.

Therefore, the original problem is equivalent to

$$\begin{aligned} \operatorname{argmin}_{\Theta} Z(\Theta, \boldsymbol{\delta}) &= \frac{1}{2n} \|\mathbf{Y} - \boldsymbol{\alpha} - \widehat{\mathbf{F}}\boldsymbol{\theta} - \widehat{\mathbf{U}}\boldsymbol{\beta}\|_2^2 + \lambda_1 g_1(\boldsymbol{\delta}) - \lambda_1 g_2(\boldsymbol{\delta}) + \lambda_1 \|\boldsymbol{\beta}\|_1 \\ &\text{subject to } \delta_{ik} = \alpha_i - \gamma_k, \quad i = 1, \dots, n, \quad k = 1, \dots, K, \quad \text{and } \gamma_1 \leq \gamma_2 \leq \dots \leq \gamma_K. \end{aligned}$$

We can then define a sequence of lower approximations of $g_2(\boldsymbol{\delta})$ as

$$g_2^{(m)}(\boldsymbol{\delta}) = g_2(\widehat{\boldsymbol{\delta}}^{(m-1)}) + (\nabla g_2(\widehat{\boldsymbol{\delta}}^{(m-1)}))^\top (\boldsymbol{\delta} - \widehat{\boldsymbol{\delta}}^{(m-1)}),$$

where $\nabla g_2(\boldsymbol{\delta})$ is the sub-gradient of $g_2(\boldsymbol{\delta})$ and $\widehat{\boldsymbol{\delta}}^{(m-1)}$ is the estimator of $\boldsymbol{\delta}$ from the $(m-1)$ -th iteration. More specifically, let $\nabla_{ik} g_2(\boldsymbol{\delta})$ be the sub-gradient of $g_2(\boldsymbol{\delta})$ with respect to δ_{ik} . Under the ℓ_1 -type distance

for $1 < k < K$, we have

$$\nabla_{i_k} g_2(\boldsymbol{\delta}) = \text{sgn}(\delta_{ik}) \mathbb{I}(|\delta_{ik}| > |\delta_{i(k-1)}|) + \text{sgn}(\delta_{ik}) \mathbb{I}(|\delta_{ik}| > |\delta_{i(k+1)}|),$$

and otherwise we have

$$\nabla_{i_1} g_2(\boldsymbol{\delta}) = \text{sgn}(\delta_{i_1}) \mathbb{I}(|\delta_{i_1}| > |\delta_{i_2}|), \quad \nabla_{i_K} g_2(\boldsymbol{\delta}) = \text{sgn}(\delta_{i_K}) \mathbb{I}(|\delta_{i_K}| > |\delta_{i(K-1)}|).$$

The DC programming approach encourages us to optimize an upper approximation in the $(m+1)$ -th iteration, which is

$$\begin{aligned} \underset{\boldsymbol{\Theta}, \boldsymbol{\delta}}{\text{argmin}} \quad & Z^{(m+1)}(\boldsymbol{\Theta}, \boldsymbol{\delta}) = \frac{1}{2n} \|\mathbf{Y} - \boldsymbol{\alpha} - \widehat{\mathbf{F}}\boldsymbol{\theta} - \widehat{\mathbf{U}}\boldsymbol{\beta}\|_2^2 + \lambda_1 \sum_{i=1}^n \sum_{k=1}^K d(\delta_{ik}) - \lambda_1 (\nabla g_2(\widehat{\boldsymbol{\delta}}^{(m)}))^\top (\boldsymbol{\delta} - \widehat{\boldsymbol{\delta}}^{(m)}) + \lambda_2 \|\boldsymbol{\beta}\|_1 \\ \text{subject to} \quad & \delta_{ik} = \alpha_i - \gamma_k \quad i = 1, \dots, n, \quad k = 1, \dots, K, \quad \text{and} \quad \gamma_1 \leq \gamma_2 \leq \dots \leq \gamma_K. \end{aligned} \quad (2.3)$$

Clearly, the optimization problem (2.3) is convex with equality and inequality constraints. We denote the global minimizer in the m -th iteration as $\{\widehat{\boldsymbol{\Theta}}^{(m+1)}, \widehat{\boldsymbol{\delta}}^{(m+1)}\}$. Inspired by [Giesen and Laue \(2019\)](#), who employed slack variables for inequality constraints in the standard ADMM, we construct the slack vector $\boldsymbol{y} \in \mathbb{R}^{K-1}$ with $y_k = \gamma_k - \gamma_{k+1}$ for $k < K-1$. To ensure $y_k < 0$, we define a loss function $\mathbb{I}_\infty(y_k > 0)$, where $\mathbb{I}_\infty(y_k > 0) = \infty$ when $y_k > 0$ and 0 otherwise. By incorporating this loss function, $\mathbb{I}_\infty(y_k > 0)$, (2.3) transforms into the following equivalent optimization problem without inequality constraints:

$$\begin{aligned} \underset{\boldsymbol{\Theta}, \boldsymbol{\delta}}{\text{argmin}} \quad & \frac{1}{2n} \|\mathbf{Y} - \boldsymbol{\alpha} - \widehat{\mathbf{F}}\boldsymbol{\theta} - \widehat{\mathbf{U}}\boldsymbol{\beta}\|_2^2 + \lambda_1 \sum_{i=1}^n \sum_{k=1}^K d(\delta_{ik}) - \lambda_1 (\nabla g_2(\widehat{\boldsymbol{\delta}}^{(m)}))^\top \boldsymbol{\delta} + \lambda_2 \|\boldsymbol{\eta}\|_1 + \sum_{k=1}^{K-1} \mathbb{I}_\infty(y_k > 0) \\ \text{subject to} \quad & \boldsymbol{\delta} = \mathbf{C}_1 \boldsymbol{\alpha} - \mathbf{C}_2 \boldsymbol{\gamma}, \quad \mathbf{D} \boldsymbol{\gamma} = \boldsymbol{y} \quad \text{and} \quad \boldsymbol{\beta} = \boldsymbol{\eta}, \end{aligned} \quad (2.4)$$

where

$$\mathbf{C}_1 = \begin{pmatrix} \mathbf{1}_K & \mathbf{0}_K & \cdots & \mathbf{0}_K \\ \mathbf{0}_K & \mathbf{1}_K & \cdots & \mathbf{0}_K \\ \vdots & \vdots & \cdots & \vdots \\ \mathbf{0}_K & \mathbf{0}_K & \cdots & \mathbf{1}_K \end{pmatrix}_{((n \times K) \times n)}, \quad \mathbf{C}_2 = \begin{pmatrix} \mathbf{I}_K \\ \mathbf{I}_K \\ \vdots \\ \mathbf{I}_K \end{pmatrix}_{((n \times K) \times K)}, \quad \mathbf{D} = \begin{pmatrix} 1 & -1 & 0 & \cdots & 0 \\ 0 & 1 & -1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & \cdots & 0 & 1 & -1 \end{pmatrix}_{((K-1) \times K)}.$$

The constraint $\boldsymbol{\beta} = \boldsymbol{\eta}$ is imposed to separate the ℓ_1 -norm of $\boldsymbol{\beta}$ from the quadratic loss. Finally, we form the

scaled Lagrangian problem as

$$\begin{aligned} \operatorname{argmin}_{\Theta, \delta, \mathbf{y}, \mathbf{u}, \mathbf{v}, \mathbf{w}} \frac{1}{2n} \|\mathbf{Y} - \boldsymbol{\alpha} - \widehat{\mathbf{F}}\boldsymbol{\theta} - \widehat{\mathbf{U}}\boldsymbol{\beta}\|_2^2 + \lambda_1 \sum_{i=1}^n \sum_{k=1}^K d(\delta_{ik}) - \lambda_1 (\nabla g_2(\widehat{\boldsymbol{\delta}}^{(m)}))^\top \boldsymbol{\delta} + \lambda_2 \|\boldsymbol{\eta}\|_1 \\ + \sum_{k=1}^{K-1} \mathbb{I}_\infty(y_k > 0) + \frac{\rho_1}{2} (\|\boldsymbol{\delta} - \mathbf{C}_1 \boldsymbol{\alpha} + \mathbf{C}_2 \boldsymbol{\gamma} + \mathbf{u}\|_2^2) + \frac{\rho_2}{2} (\|\mathbf{D}\boldsymbol{\gamma} - \mathbf{y} + \mathbf{v}\|_2^2) + \frac{\rho_3}{2} (\|\boldsymbol{\eta} - \boldsymbol{\beta} + \mathbf{w}\|_2^2), \end{aligned}$$

where $\mathbf{u} = (\mathbf{u}_1^\top, \dots, \mathbf{u}_n^\top)^\top$ with $\mathbf{u}_i = (u_{i1}, \dots, u_{iK})^\top$, $\mathbf{v} = (v_1, \dots, v_{K-1})^\top$ and $\mathbf{w} = (w_1, \dots, w_p)^\top$ are the Lagrangian multipliers with corresponding augmented parameters denoted as ρ_1 , ρ_2 and ρ_3 . The standard ADMM procedures can be expressed as

$$\begin{aligned} \widehat{\Theta}^{s+1} &= \operatorname{argmin}_{\Theta} \frac{1}{2n} \|\mathbf{Y} - \boldsymbol{\alpha} - \widehat{\mathbf{F}}\boldsymbol{\theta} - \widehat{\mathbf{U}}\boldsymbol{\beta}\|_2^2 + \frac{\rho_1}{2} \|\widehat{\boldsymbol{\delta}}^s - \mathbf{C}_1 \boldsymbol{\alpha} + \mathbf{C}_2 \boldsymbol{\gamma} + \widehat{\mathbf{u}}^s\|_2^2 + \frac{\rho_2}{2} \|\mathbf{D}\boldsymbol{\gamma} - \widehat{\mathbf{y}}^s + \widehat{\mathbf{v}}^s\|_2^2 \\ &\quad + \frac{\rho_3}{2} (\|\widehat{\boldsymbol{\eta}}^s - \boldsymbol{\beta} + \widehat{\mathbf{w}}^s\|_2^2), \\ \widehat{\boldsymbol{\delta}}^{s+1} &= \operatorname{argmin}_{\boldsymbol{\delta}} \frac{\rho_1}{2} \|\boldsymbol{\delta} - \mathbf{C}_1 \widehat{\boldsymbol{\alpha}}^{s+1} + \mathbf{C}_2 \widehat{\boldsymbol{\gamma}}^{s+1} + \widehat{\mathbf{u}}^s\|_2^2 - \lambda_1 (\nabla g_2(\widehat{\boldsymbol{\delta}}^{(m-1)}))^\top (\boldsymbol{\delta} - \widehat{\boldsymbol{\delta}}^{(m-1)}) + \lambda_1 \|\boldsymbol{\delta}\|_1, \\ \widehat{\mathbf{y}}^{s+1} &= \operatorname{argmin}_{\mathbf{y}} \frac{\rho_2}{2} \|\mathbf{D}\widehat{\boldsymbol{\gamma}}^{s+1} - \mathbf{y} + \widehat{\mathbf{v}}^s\|_2^2 + \sum_{k=1}^{K-1} \mathbb{I}_\infty(y_k > 0), \\ \widehat{\boldsymbol{\eta}}^{s+1} &= \operatorname{argmin}_{\boldsymbol{\eta}} \frac{\rho_3}{2} (\|\boldsymbol{\eta} - \widehat{\boldsymbol{\beta}}^{s+1} + \widehat{\mathbf{w}}^s\|_2^2) + \lambda_2 \|\boldsymbol{\eta}\|_1, \\ \widehat{\mathbf{u}}^{s+1} &= \widehat{\mathbf{u}}^s + \widehat{\boldsymbol{\delta}}^{s+1} - \mathbf{C}_1 \widehat{\boldsymbol{\alpha}}^{s+1} + \mathbf{C}_2 \widehat{\boldsymbol{\gamma}}^{s+1}, \\ \widehat{\mathbf{v}}^{s+1} &= \widehat{\mathbf{v}}^s + \mathbf{D}\widehat{\boldsymbol{\gamma}}^{s+1} - \widehat{\mathbf{y}}^{s+1}, \\ \widehat{\mathbf{w}}^{s+1} &= \widehat{\mathbf{w}}^s + \widehat{\boldsymbol{\eta}}^{s+1} - \widehat{\boldsymbol{\beta}}^{s+1}, \end{aligned} \tag{2.5}$$

where the subscript s denotes the s -th iteration in the standard ADMM algorithm.

Note that the first optimization in (2.5) is a quadratic form, and we can find the global minimizer by setting its derivative to zero. To update $\boldsymbol{\delta}$ and $\boldsymbol{\beta}$, we apply the soft-thresholding operator to obtain the explicit forms:

$$\widehat{\boldsymbol{\delta}}^{s+1} = \operatorname{ST}\left(\left(\mathbf{C}_1 \widehat{\boldsymbol{\alpha}}^{s+1} - \mathbf{C}_2 \widehat{\boldsymbol{\gamma}}^{s+1} - \widehat{\mathbf{u}}^{s+1} + \lambda_1 \nabla g_2(\widehat{\boldsymbol{\delta}}^{(m)})\right)/\rho_1, \lambda_1/\rho_1\right), \quad \widehat{\boldsymbol{\eta}}^{s+1} = \operatorname{ST}\left(\left(\widehat{\boldsymbol{\beta}}^{s+1} - \widehat{\mathbf{w}}^s\right), \lambda_2/\rho_3\right),$$

where $\operatorname{ST}_t(\mathbf{x})$ is the soft-thresholding operator applied to each element of \mathbf{x} , defined as $\operatorname{ST}_t(u) = \operatorname{sgn}(u)(|u| - t)$. By some simple algebra, the updating formula for the slack vector \mathbf{y} is given by

$$\widehat{y}_k^{s+1} = \min\{0, \mathbf{D}\widehat{\boldsymbol{\gamma}}^{s+1} + \widehat{v}_k^s\}. \tag{2.6}$$

In the above DC-ADMM algorithm, the parameters ρ_1 , ρ_2 and ρ_3 influence the convergence speed (Boyd et al., 2011; Wu et al., 2016). For practical implementation, we typically set $\rho_1 = \rho_2 = \rho_3 = 1/2$. The

complete DC-ADMM algorithm is summarized in Algorithm 1.

Algorithm 1 DC-ADMM for the Plug-in Optimization

- 1: **Input:** Datasets \mathbf{Y} and \mathbf{X} ; PCA estimators $\widehat{\mathbf{F}}$ and $\widehat{\mathbf{U}}$; tuning parameters λ_1, λ_2 and tolerance $\epsilon_1, \epsilon_2 > 0$ for inner and outer layer algorithms.
 - 2: **Global Initialize:** Set $m = 0$ and DCA-initial value $\widehat{\boldsymbol{\alpha}}^{(m)}, \widehat{\boldsymbol{\gamma}}^{(m)}$. For inner ADMM, the initializations of the dual variables $\mathbf{u}^0, \mathbf{v}^0, \mathbf{w}^0$ are chosen to be zero vectors and $\widehat{\boldsymbol{\eta}}^0 = \mathbf{1}_p$.
 - 3: **while** $m = 0$ or $|Z(\widehat{\boldsymbol{\Theta}}^{(m)}, \widehat{\boldsymbol{\delta}}^{(m)}) - Z(\widehat{\boldsymbol{\Theta}}^{(m-1)}, \widehat{\boldsymbol{\delta}}^{(m-1)})| > \epsilon_1$ **do**
 - 4: **ADMM Initialize:** Find the upper bound function $Z^{(m+1)}(\boldsymbol{\Theta}, \boldsymbol{\delta})$ in the form of (2.3) at $(\widehat{\boldsymbol{\Theta}}^{(m)}, \widehat{\boldsymbol{\delta}}^{(m)})$. Initialize $\widehat{\boldsymbol{\alpha}}^0 = \widehat{\boldsymbol{\alpha}}^{(m)}$ and $\widehat{\boldsymbol{\gamma}}^0 = \widehat{\boldsymbol{\gamma}}^{(m)}$. $\widehat{\boldsymbol{y}}^0$ is calculated by the equation (2.6) with $\widehat{\boldsymbol{\gamma}}^{(m)}$ and \mathbf{v}^0 .
 - 5: **while** $s = 0$ or $|Z^{(m+1)}(\widehat{\boldsymbol{\Theta}}^s, \widehat{\boldsymbol{\delta}}^s) - Z^{(m+1)}(\widehat{\boldsymbol{\Theta}}^{s-1}, \widehat{\boldsymbol{\delta}}^{s-1})| > \epsilon_2$ **do**
 - 6: Update $\widehat{\boldsymbol{\Theta}}^{s+1}, \widehat{\boldsymbol{\delta}}^{s+1}, \widehat{\boldsymbol{y}}^{s+1}, \widehat{\mathbf{u}}^{s+1}, \widehat{\mathbf{v}}^{s+1}$ and $\widehat{\mathbf{w}}^{s+1}$ by (2.5).
 - 7: $s \leftarrow s + 1$.
 - 8: **end while**
 - 9: **ADMM Output:** Updated variables $(\widehat{\boldsymbol{\Theta}}^{(m+1)}, \widehat{\boldsymbol{\delta}}^{(m+1)})$.
 - 10: $m \leftarrow m + 1$.
 - 11: **end while**
 - 12: **Global Output:** Estimations of intercept $\widehat{\boldsymbol{\alpha}}$, group centroids $\widehat{\boldsymbol{\gamma}}$ and regression coefficients $\widehat{\boldsymbol{\theta}}$ and $\widehat{\boldsymbol{\beta}}$.
-

In the following theorem, we establish that the proposed DC-ADMM algorithm guarantees convergence to a local minimizer in finite steps.

Theorem 2.1. In the DC-ADMM algorithm with ℓ_1 -type distance, the sequence $\{Z(\widehat{\boldsymbol{\Theta}}^{(m)})\}$ decreases with m and converges in finite steps. Specifically, there exists an $m^* < \infty$ such that $Z(\widehat{\boldsymbol{\Theta}}^{(m)}) = Z(\widehat{\boldsymbol{\Theta}}^{(m^*)})$ for all $m \geq m^*$. Furthermore, $\widehat{\boldsymbol{\Theta}}^{(m^*)}$ is a KKT point of $Z(\boldsymbol{\Theta})$.

Although the ADMM algorithm typically ensures convergence to a global minimizer, the DC-ADMM guarantees only a KKT point due to the nonconvex nature of the CAR optimization problem. A variant DC algorithm proposed by Breiman and Cutler (1993) may achieve a global minimizer, but it is often criticized for its slow convergence speed. In Section 3, we also discuss the statistical properties of the local minimizer.

It is worth noting that the DC-ADMM can also be extended to handle ℓ_2 -type distance by modifying the updating formula of $\boldsymbol{\delta}$ in (2.5). However, the computational burden of DC-ADMM under ℓ_2 -type distance can be prohibitive. Therefore, for efficiency, we propose an alternative approach using the Cyclic Coordinate Descent (CCD) algorithm to optimize under ℓ_2 -type distance.

2.2.2. Cyclic Coordinate Decent Algorithm

In this section, we introduce the CCD algorithm tailored to optimize the problem (2.1) with ℓ_2 -type distance. Without loss of generality, the superscript m denotes the m -th step estimator in the coordinate descent algorithm. For instance, $\widehat{\boldsymbol{\alpha}}^{(m)}$ and $\widehat{\boldsymbol{\gamma}}^{(m)}$ represent the m -th step estimators of $\boldsymbol{\alpha}$ and $\boldsymbol{\gamma}$, respectively.

Given the estimation process, it is clear that $\widehat{\mathbf{F}}^\top \widehat{\mathbf{U}} = \mathbf{0}$. Consequently, we update $\boldsymbol{\theta}$ and $\boldsymbol{\beta}$ as follows:

$$\widehat{\boldsymbol{\theta}}^{(m+1)} = \widehat{\mathbf{F}}^\top (\mathbf{Y} - \widehat{\boldsymbol{\alpha}}^{(m)})/n, \quad (2.7)$$

$$\widehat{\boldsymbol{\beta}}^{(m+1)} = \underset{\boldsymbol{\beta}}{\operatorname{argmin}} \frac{1}{2n} \|\mathbf{Y} - \widehat{\boldsymbol{\alpha}}^{(m)} - \widehat{\mathbf{F}}\widehat{\boldsymbol{\theta}}^{(m+1)} - \widehat{\mathbf{U}}\boldsymbol{\beta}\|_2^2 + \lambda_2 \|\boldsymbol{\beta}\|_1. \quad (2.8)$$

Clearly, the optimization problem in (2.8) is a standard LASSO problem, and many existing packages, such as the R package `glmnet`, can be used to solve it. Given $\{\widehat{\boldsymbol{\theta}}^{(m+1)}, \widehat{\boldsymbol{\beta}}^{(m+1)}, \widehat{\boldsymbol{\gamma}}^{(m)}\}$, updating $\boldsymbol{\alpha}$ is equivalent to solve the following optimization problem:

$$\widehat{\boldsymbol{\alpha}}^{(m+1)} = \underset{\boldsymbol{\alpha}}{\operatorname{argmin}} \frac{1}{2n} \|\mathbf{Y} - \boldsymbol{\alpha} - \widehat{\mathbf{F}}\widehat{\boldsymbol{\theta}}^{(m+1)} - \widehat{\mathbf{U}}\widehat{\boldsymbol{\beta}}^{(m+1)}\|_2^2 + \lambda_1 g(\boldsymbol{\alpha}, \widehat{\boldsymbol{\gamma}}^{(m)}). \quad (2.9)$$

Following procedures similar to the DC algorithm introduced in Section 2.2.1, it is straightforward to verify that both $g_1(\boldsymbol{\alpha}, \boldsymbol{\gamma})$ and $g_2(\boldsymbol{\alpha}, \boldsymbol{\gamma})$ are convex functions with respect to $\boldsymbol{\alpha}$. Therefore, the corresponding optimization in (2.9), based on the lower approximation of $g(\boldsymbol{\alpha}, \boldsymbol{\gamma})$, yields

$$\begin{aligned} \widehat{\boldsymbol{\alpha}}^{(m+1)} = \underset{\boldsymbol{\alpha}}{\operatorname{argmin}} & \frac{1}{2n} \|\mathbf{Y} - \boldsymbol{\alpha} - \widehat{\mathbf{F}}\widehat{\boldsymbol{\theta}}^{(m+1)} - \widehat{\mathbf{U}}\widehat{\boldsymbol{\beta}}^{(m+1)}\|_2^2 \\ & + \lambda_1 \sum_{i=1}^n \sum_{k=1}^K (\alpha_i - \widehat{\gamma}_k^{(m)})^2 - \lambda_1 \left(\nabla g(\widehat{\boldsymbol{\alpha}}^{(m)}, \widehat{\boldsymbol{\gamma}}^{(m)}) \right)^\top (\boldsymbol{\alpha} - \boldsymbol{\alpha}^{(m)}), \end{aligned} \quad (2.10)$$

where $\nabla g_2(\boldsymbol{\alpha}^{(m)}, \boldsymbol{\gamma}^{(m)})$ is the sub-gradient of $g_2(\boldsymbol{\alpha}^{(m)}, \boldsymbol{\gamma}^{(m)})$ with respect to $\boldsymbol{\alpha}^{(m)}$,

$$\nabla g_2(\boldsymbol{\alpha}^{(m)}, \boldsymbol{\gamma}^{(m)}) = \left\{ 2 \sum_{i=1}^n \sum_{k=2}^K \max\{|\alpha_i^{(m)} - \gamma_{k-1}^{(m)}|, |\alpha_i^{(m)} - \gamma_k^{(m)}|\} \operatorname{sgn}(\alpha_i^{(m)} - (\gamma_{k-1}^{(m)} + \gamma_k^{(m)})/2) \right\}.$$

Note that the objective function in (2.10) is quadratic with respect to $\boldsymbol{\alpha}$, and the explicit form of $\widehat{\boldsymbol{\alpha}}^{(m+1)}$ can be derived by setting its first derivative equal to zero:

$$\widehat{\boldsymbol{\alpha}}^{(m+1)} = \frac{n}{1 + 2\lambda_1 K n} \left[\frac{1}{n} (\mathbf{Y} - \widehat{\mathbf{F}}\widehat{\boldsymbol{\theta}}^{(m+1)} - \widehat{\mathbf{U}}\widehat{\boldsymbol{\beta}}^{(m+1)}) + 2\lambda_1 \left(\sum_{k=1}^K \widehat{\gamma}_k^{(m)} \right) \mathbf{1}_n + \lambda_1 \left(\nabla g(\widehat{\boldsymbol{\alpha}}^{(m)}, \widehat{\boldsymbol{\gamma}}^{(m)}) \right) \right]. \quad (2.11)$$

The updating formula for $\boldsymbol{\gamma}$ is defined as

$$\widehat{\boldsymbol{\gamma}}^{(m+1)} = \underset{\boldsymbol{\gamma}}{\operatorname{argmin}} \sum_{i=1}^n \min\{(\widehat{\alpha}_i^{(m+1)} - \gamma_1)^2, \dots, (\widehat{\alpha}_i^{(m+1)} - \gamma_K)^2\}. \quad (2.12)$$

This degenerates into the standard K -means clustering procedure and can be implemented using existing R packages such as `Ckmeans.1d.dp`.

Once the main updating rules are determined, compute $Z(\widehat{\boldsymbol{\Theta}})$ and update the main parameters at each

iteration, repeating until the stopping criterion is met. The stopping criterion is defined such that either $Z(\hat{\Theta}^{(m)})$ is sufficiently close to $Z(\hat{\Theta}^{(m-1)})$ or the maximum number of iterations is reached. The summarized CCD algorithm for optimizing the problem with ℓ_2 -type distance is outlined in Algorithm 2.

Algorithm 2 Cyclic Coordinate Descent Algorithm for the Optimization Problem with ℓ_2 -type Distance

- 1: **Input:** Datasets \mathbf{Y} and \mathbf{X} ; PCA estimators $\hat{\mathbf{F}}$ and $\hat{\mathbf{U}}$; tuning parameters λ_1 , λ_2 and tolerance $\epsilon > 0$.
 - 2: **Initialize:** Set $m = 0$ and initial value $\hat{\alpha}^{(0)}$, $\hat{\gamma}^{(0)}$.
 - 3: **while** $m = 0$ or $|Z(\hat{\Theta}^{(m)}) - Z(\hat{\Theta}^{(m-1)})| > \epsilon$ **do**
 - 4: Update $\hat{\theta}^{(m+1)}$ by (2.7) and update $\hat{\beta}^{(m+1)}$ in (2.8) by the standard LASSO optimization.
 - 5: Update $\hat{\alpha}^{(m+1)}$ by (2.11).
 - 6: Update $\hat{\gamma}^{(m+1)}$ in (2.12) by K-means algorithms.
 - 7: $m \leftarrow m + 1$.
 - 8: **end while**
 - 9: **Output:** Estimations of intercept $\hat{\alpha}$, group centroids $\hat{\gamma}$ and regression coefficients $\hat{\theta}$ and $\hat{\beta}$.
-

Note that if the ℓ_2 -type distance in Algorithm 2 is replaced with the ℓ_1 -type distance, the optimization problem in (2.12) becomes a standard K -median algorithm, implementable using the R package `Ckmeans.1d.dp`. However, updating α in (2.10) would lack an explicit form, necessitating complex algorithms such as ADMM to obtain iterative solutions. Moreover, it remains an open question whether the cyclic coordinate descent algorithm converges to a local minimizer for the non-convex optimization problem. Simulation results show no significant difference in performances between Algorithm 1 and Algorithm 2, despite the latter bearing greater computational burden. Hence, we recommend using Algorithm 2 for ℓ_2 -type distance due to its computational efficiency, reserving DC-ADMM for solving (2.1) with ℓ_1 -type distance.

2.3. Practical Implementation Details

We first investigate the method for selecting the number of subgroups K and the shrinkage parameters λ_1 and λ_2 . To alleviate computational burdens, we use the Bayesian Information Criterion (BIC) to determine the group number K . Specifically, let

$$\text{BIC}(K) = \log \left[\sum_{i=1}^n (Y_i - \hat{Y}_i)^2 / n \right] + a_n (\hat{S} + K) \frac{\log(n)}{n},$$

where $\hat{Y}_i = \sum_{k=1}^K \mathbb{I}(i \in \hat{\mathcal{G}}_k) \hat{\gamma}_k + \hat{\mathbf{f}}_i^\top \hat{\theta} + \hat{\mathbf{u}}_i^\top \hat{\beta}$ and $\hat{S} = \sum_{j=1}^p \mathbb{I}(\hat{\beta}_j \neq 0)$. For better practical performance, we set $a_n = 2 \log(nK + p)$. As λ_1 tends to infinity, it results in $\hat{\alpha}_i = \hat{\gamma}_k$ for $i \in \hat{\mathcal{G}}_k$. Similarly, the sparsity of $\hat{\beta}$ increases as λ_2 grows. To minimize differences $|\hat{\alpha}_i - \hat{\gamma}_k|$ within each group, we initially set λ_1 to a relatively large value and λ_2 to a relatively small value to control the bias. Subsequently, we determine the optimal group number by minimizing the BIC. Similar strategies are also implemented in Ma and Huang (2017) and He et al. (2022c). With the estimated group number K , we further adopt the generalized cross-validation (GCV) method by Tang et al. (2021), which is an approximate version of the method by Pan et al. (2013),

to select λ_1 and λ_2 . Specifically, the GCV is defined as

$$\text{GCV}(\lambda_1, \lambda_2) = \frac{\sum_{i=1}^n (Y_i - \hat{Y}_i)^2}{(n - \text{df})^2},$$

where $\text{df} = \sum_{j=1}^p \mathbb{I}(\hat{\beta}_j \neq 0)$. Alternatively, one can simultaneously select the optimal K , λ_1 and λ_2 by minimizing the BIC. However, this approach is computationally intensive due to the large size of the grid search.

Next, we focus on the selection of initial values $(\hat{\boldsymbol{\alpha}}^{(0)}, \hat{\boldsymbol{\gamma}}^{(0)})$ in Algorithm 1. This step is more challenging because the algorithm may converge to a local minimizer due to the non-convexity of the objective function. Inspired by the ridge regression estimators used for initializing values by He et al. (2022c), we incorporate the primary factor structure and fit the following ridge regression to obtain $\hat{\boldsymbol{\alpha}}^{(0)}$:

$$\hat{\boldsymbol{\alpha}}^{(0)} = \underset{\boldsymbol{\alpha}}{\text{argmin}} \frac{1}{2n} \|\mathbf{Y} - \mathbf{X}^* \boldsymbol{\beta}^*\|_2^2 + \lambda^* \|\boldsymbol{\beta}^*\|_2^2, \quad (2.13)$$

where $\mathbf{X}^* = (\hat{\mathbf{F}}, \mathbf{I}_n)$ and $\boldsymbol{\beta}^* = (\boldsymbol{\theta}^\top, \boldsymbol{\alpha}^\top)^\top$. The tuning parameter λ^* is determined by cross-validation. Given $\hat{\boldsymbol{\alpha}}^{(0)}$, $\hat{\boldsymbol{\gamma}}^{(0)}$ can be obtained using the K -means (or K -median) algorithm. The rationale behind using ridge regression is that the pseudo \mathbf{X}^* has a dimension of $r + n$, which is larger than the sample size n . We exclude $\hat{\mathbf{U}}$ from the regression for two reasons: firstly, due to the latent factor structure, the common factors $\mathbf{f}_i \in \mathbb{R}^r$ are considered to contain the most crucial information; secondly, incorporating the idiosyncratic errors $\mathbf{u}_i \in \mathbb{R}^p$ into ridge regression would introduce significant computational challenges due to its high dimensionality. Indeed, our simulation results indicate that the proposed strategy performs well and offers computational advantages.

3. Theoretical Properties

In this section, we investigate the statistical properties of our proposed estimators under mild conditions. First, we study the convergence rate of the ‘‘oracle’’ estimators, assuming prior knowledge of group memberships. Subsequently, we establish the asymptotic relationship between the oracle estimator and the local minimizer of the optimization problem (2.1).

Throughout the statement of the following theorems, we denote the true values of $\boldsymbol{\alpha}$, $\boldsymbol{\gamma}$ and $\boldsymbol{\beta}$ as $\boldsymbol{\alpha}_0$, $\boldsymbol{\gamma}_0$ and $\boldsymbol{\beta}_0$, respectively, and define $\boldsymbol{\theta}_0 = \mathbf{B}^\top \boldsymbol{\beta}_0$. Additionally, we use $|\mathcal{G}_{\min}|$ and $|\mathcal{G}_{\max}|$ to represent the minimum and maximum cardinalities of \mathcal{G}_k , respectively. Let $\boldsymbol{\Omega} = (\Omega_{ik})$ be the matrix indicating the true group membership, where $\Omega_{ik} = 1$ implies that subject i belongs to group \mathcal{G}_k and $\Omega_{ik} = 0$ otherwise. Hence, it is straightforward to verify that $\boldsymbol{\alpha}_0 = \boldsymbol{\Omega} \boldsymbol{\gamma}_0$. In the ideal scenario where $\boldsymbol{\Omega}$ is known in advance, the oracle

estimator can be defined as

$$(\hat{\boldsymbol{\gamma}}^{or}, \hat{\boldsymbol{\theta}}^{or}, \hat{\boldsymbol{\beta}}^{or}) = \underset{\boldsymbol{\gamma}, \boldsymbol{\theta}, \boldsymbol{\beta}}{\operatorname{argmin}} \frac{1}{2n} \|\mathbf{Y} - \boldsymbol{\Omega}\boldsymbol{\gamma} - \hat{\mathbf{F}}\boldsymbol{\theta} - \hat{\mathbf{U}}\boldsymbol{\beta}\|_2^2 + \lambda_2 \|\boldsymbol{\beta}\|_1. \quad (3.1)$$

Write $\hat{\boldsymbol{\alpha}}^{or} = \boldsymbol{\Omega}\hat{\boldsymbol{\gamma}}^{or}$. Note that the oracle estimator only incorporates prior grouping information $\boldsymbol{\Omega}$, without prior knowledge of the sparsity of $\boldsymbol{\beta}_0$.

In the following, we first introduce some assumptions related to the estimation of factor models. These technical assumptions are commonly used in the literature on large-dimensional factor models, such as Fan et al. (2013), Fan et al. (2020) and Fan et al. (2023).

Assumption 1. (Tail Probability) The sequence $\{\mathbf{f}_i, \mathbf{u}_i\}_{-\infty}^{\infty}$ is strictly stationary with $\mathbb{E}(f_{ij}) = \mathbb{E}(u_{is}) = \mathbb{E}(f_{ij}u_{is}) = 0$ for all $s \leq p$, $j \leq r$ and $i \leq n$. Furthermore, we assume \mathbf{f}_i and \mathbf{u}_i satisfy exponential-type tail probability:

$$\mathbb{P}\left(\frac{|\mathbf{a}^\top \mathbf{f}_i|}{\|\mathbf{a}\|_2} \geq t\right) \leq \exp\left(-\left(\frac{t}{V_1}\right)^{l_1}\right) \quad \text{and} \quad \mathbb{P}\left(\frac{|\mathbf{b}^\top \mathbf{u}_i|}{\|\mathbf{b}\|_2} \geq t\right) \leq \exp\left(-\left(\frac{t}{V_2}\right)^{l_2}\right),$$

where $\mathbf{a} \in \mathbb{R}^r$, $\mathbf{b} \in \mathbb{R}^p$. V_1, V_2, l_1 and l_2 are positive constants. Denote $\operatorname{Cov}(\mathbf{u}_i)$ as $\boldsymbol{\Sigma}$ for all $i = 1, \dots, n$. There exists two positive constants M_1 and M_2 such that $\lambda_{\min}(\boldsymbol{\Sigma}) \geq M_1$, $\min_{s,t \in \{1, \dots, p\}} \operatorname{Var}(u_{is}u_{it}) \geq M_1$ and $\|\boldsymbol{\Sigma}\|_{\infty} \leq M_2$.

Assumption 2. (α -Mixing Condition) Suppose $\mathcal{A}_{-\infty}^0$ and \mathcal{A}_n^{∞} are σ -algebras generated by $\{\mathbf{f}_i, \mathbf{u}_i\}_{-\infty}^0$ and $\{\mathbf{f}_i, \mathbf{u}_i\}_n^{\infty}$ respectively. We assume that there exists a constant l_3 such that $3/l_1 + 3/(2l_2) + 1/l_3 > 1$ and the following inequality holds:

$$\sup_{E_1 \in \mathcal{A}_{-\infty}^0, E_2 \in \mathcal{A}_n^{\infty}} |\mathbb{P}(E_1)\mathbb{P}(E_2) - \mathbb{P}(E_1E_2)| \leq \exp(-an^{l_3}),$$

where a is a positive constant.

Assumption 3. There exists some constant M_0 such that $\|\mathbf{B}\|_{\max} \leq M_0$, $\mathbb{E}[p^{-2}[\mathbf{u}_i^\top \mathbf{u}_j - \mathbb{E}(\mathbf{u}_i^\top \mathbf{u}_j)]]^4 \leq M_0$ and all the eigenvalues of $p^{-1}\mathbf{B}^\top \mathbf{B}$ are bounded away from 0 and ∞ . Furthermore, $\mathbb{E}\|p^{-1/2}\mathbf{B}^\top \mathbf{u}_i\|_2^4 \leq M_0$.

Assumption 1 requires exponential-type tail probability for \mathbf{f}_i and \mathbf{u}_i , assuming they are uncorrelated. Moreover, $\|\boldsymbol{\Sigma}\|_{\infty} \leq M_2$ is imposed to ensure consistent estimation of the number of factors. Assumption 2 relaxes the independence of $\{\mathbf{f}_i, \mathbf{u}_i\}_{-\infty}^{\infty}$ to weak dependence. Combined with the exponential-type tail probability in Assumption 1, these conditions allow us to establish non-asymptotic bounds related to \mathbf{f}_i and \mathbf{u}_i . Assumption 3 is a standard condition for deriving the consistency of factor loadings and factor scores with a fixed r , adapted from conditions in Bai (2003). Under these regular conditions, we derive the convergence rates of $\hat{\mathbf{F}}, \hat{\mathbf{B}}$ and $\hat{\mathbf{U}}$, ensuring the efficiency of the plug-in estimations. In the following, we

proceed to introduce some technical assumptions for subgroup identification.

Assumption 4. Suppose ϵ has tail probability of sub-Gaussian form, that is, $\mathbb{P}(|\mathbf{c}^\top \epsilon| \geq \|\mathbf{c}\|_2 t) \leq 2 \exp(-C_0 t^2)$ for any $\mathbf{c} \in \mathbb{R}^n$, where C_0 is a positive constant.

Assumption 5. There exist two positive constants C_1 and C_2 such that

$$C_1 \leq |\mathcal{G}_{\min}|^{-1} \lambda_{\min} [\text{diag}(\boldsymbol{\Omega}^\top \boldsymbol{\Omega}, n\mathbf{I}_r, n\boldsymbol{\Sigma}_{\mathcal{S}\mathcal{S}})] \leq |\mathcal{G}_{\min}|^{-1} \lambda_{\max} [\text{diag}(\boldsymbol{\Omega}^\top \boldsymbol{\Omega}, n\mathbf{I}_r, n\boldsymbol{\Sigma}_{\mathcal{S}\mathcal{S}})] \leq C_2,$$

where \mathcal{S} stands for the support set of $\boldsymbol{\beta}_0$. Additionally, we assume the irrepresentable condition holds for $\boldsymbol{\Sigma}$, i.e., there exist a positive value ρ such that $\|\boldsymbol{\Sigma}_{\mathcal{S}^c\mathcal{S}}\boldsymbol{\Sigma}_{\mathcal{S}\mathcal{S}}^{-1}\|_\infty \leq 1 - \rho$. Moreover, we assume $\|\boldsymbol{\Sigma}^{-1}\|_\infty$ is bounded by some positive constant C_4 .

Assumption 4 requires that the random noise in (1.4) follows a sub-Gaussian distribution. By the definition of $\boldsymbol{\Omega}$, we have $\boldsymbol{\Omega}^\top \boldsymbol{\Omega} = \text{diag}(|\mathcal{G}_1|, \dots, |\mathcal{G}_K|)$. As $\lambda_{\max} [\text{diag}(\boldsymbol{\Omega}^\top \boldsymbol{\Omega}, n\mathbf{I}_r, n\boldsymbol{\Sigma}_{\mathcal{S}\mathcal{S}})] \geq \lambda_{\max}(\boldsymbol{\Omega}^\top \boldsymbol{\Omega})$, Assumption 5 implies a balanced group size: $C_2 |\mathcal{G}_{\min}| \geq |\mathcal{G}_{\max}|$. The irrepresentable condition in Assumption 5, akin to that of Zhao and Yu (2006), ensures the sign consistency of $\widehat{\boldsymbol{\beta}}^{or}$. Moreover, the requirement that $\|\boldsymbol{\Sigma}^{-1}\|_\infty$ is bounded from above is equivalent to a weak correlation requirement of the idiosyncratic errors.

With these assumptions, we then present the theoretical properties of the oracle estimators in the following theorem.

Theorem 3.1. Let $\boldsymbol{\Theta}_0 := (\boldsymbol{\alpha}_0^\top, (\mathbf{H}\boldsymbol{\theta}_0)^\top, \boldsymbol{\beta}_0^\top)^\top$, where $\mathbf{H} = 1/n\mathbf{V}^{-1}\widehat{\mathbf{F}}^\top \mathbf{F}\mathbf{B}^\top \mathbf{B}$ and \mathbf{V} denotes the diagonal matrix of the first r largest eigenvalues of the sample covariance matrix, arranged in decreasing order. Suppose Assumptions 1–5 hold, $\lambda_2 \geq (1/c)\|\nabla Z_1(\boldsymbol{\Theta}_0)\|_\infty$, where $c \leq \rho/(2 - \rho)$ and $Z_1(\boldsymbol{\Theta}) = 1/2n\|\mathbf{Y} - \boldsymbol{\Omega}\boldsymbol{\gamma} - \widehat{\mathbf{F}}\boldsymbol{\theta} - \widehat{\mathbf{U}}\boldsymbol{\beta}\|_2^2$. If $p_{\mathcal{S}}(\log p/n + 1/p)^{1/2} = o(1)$ with $p_{\mathcal{S}} = |\mathcal{S}|$, it holds

$$\|((\widehat{\boldsymbol{\gamma}}^{or} - \boldsymbol{\gamma}_0)^\top, (\widehat{\boldsymbol{\theta}}^{or} - \mathbf{H}\boldsymbol{\theta}_0)^\top, (\widehat{\boldsymbol{\beta}}^{or} - \boldsymbol{\beta}_0)^\top)^\top\|_\infty = O_{\mathbb{P}}\left(p_{\mathcal{S}}(\sqrt{\log p/n} + 1/\sqrt{p})\right).$$

Theorem 3.1 establishes the consistency and convergence rate of the oracle estimator when the number of subgroups is known in advance. The first term $O_{\mathbb{P}}\left(p_{\mathcal{S}}\sqrt{\log p/n}\right)$ corresponds to the rate of penalized regression, similar to the LASSO estimator. The second term $O_{\mathbb{P}}(1/\sqrt{p})$ arises from the additional bias incurred due to the use of a plug-in estimate for the factor model. With further assumption, we can also obtain sign consistency, as stated in the following proposition.

Proposition 3.2. Under the same conditions as in Theorem 3.1, and further assuming $\min(|(\boldsymbol{\Theta}_0)_{\mathcal{S}}|) \geq \kappa\lambda_2$, where κ is a sufficiently large positive value, then $\widehat{\boldsymbol{\beta}}^{or}$ achieves sign consistency:

$$\mathbb{P}(\text{sign}(\widehat{\boldsymbol{\beta}}^{or}) = \text{sign}(\boldsymbol{\beta}_0)) \rightarrow 1.$$

Next, we establish the asymptotic relationship between the oracle estimator and the local minimizer of the optimization problem (2.1) in the following theorem.

Theorem 3.3. Let $\mathcal{M}(\lambda_1, \lambda_2)$ be the set of all local minima of the optimization problem (2.1) with ℓ_1 -type distance. Suppose the assumptions required in Theorem 3.1 hold. If $r_n \gg p_S(\sqrt{\log p/n} + 1/\sqrt{p})$ and $\lambda_1 \gg n^{-1} \max\{\sqrt{\log n}, p_S^2(\sqrt{\log p} + \sqrt{n}/\sqrt{p})\}$, where $r_n = \min_{i,j \in \{1, \dots, K\}} |\gamma_{0i} - \gamma_{0j}|$ is the minimum gap between different group centroids with γ_{0i} denoting the i -th element of γ_0 . Then, there exist a point $\hat{\Theta}_{\mathcal{M}} \in \mathcal{M}(\lambda_1, \lambda_2)$ such that $\mathbb{P}(\hat{\Theta}_{\mathcal{M}} = \hat{\Theta}^{or}) \rightarrow 1$, where $\hat{\Theta}^{or} = (\hat{\alpha}^{or}, \hat{\gamma}^{or}, \hat{\theta}^{or}, \hat{\beta}^{or})$.

Here is the refined text with some adjustments for clarity:

Theorem 3.3 asserts that the oracle estimator is a local minimizer of the optimization problem for the SILFS method with probability tending to 1. Combined with Theorem 3.1 and Proposition 3.2, we conclude that there exists a local minimizer which achieves both estimation and sign consistency. Therefore, it is crucial to identify such a (local) minimizer for the proposed SILFS method. Fortunately, the proposed DC-ADMM algorithm ensures finite-step convergence to a local minimizer. In other words, with high probability, we can obtain consistent estimators provided we start with a “good” initial value.

4. Simulation Study

In this section, we conduct thorough simulation experiments to assess the finite sample behaviors of the proposed SILFS method. In Section 4.1, we first compare the SILFS method with existing methods to evaluate empirical performance in terms of subgroup identification and variable selection. We also examine the computational time required for different methods. In Section 4.2, we investigate the sensitivity of the SILFS method to various levels of collinearity.

In the simulation studies, we use the eigenvalue-ratio method to determine the number of factors r . To elaborate, we denote $\lambda_i(\mathbf{X}\mathbf{X}^\top)$ as the i -th largest eigenvalue of $\mathbf{X}\mathbf{X}^\top$, and the estimator of the factor number is determined by the modified ER method proposed by Chang et al. (2015):

$$\hat{r} = \operatorname{argmin}_{1 \leq i \leq r^*} \frac{\lambda_i(\mathbf{X}\mathbf{X}^\top) + C_{n,p}}{\lambda_{i+1}(\mathbf{X}\mathbf{X}^\top) + C_{n,p}},$$

where r^* is a positive integer larger than r , and $C_{n,p}$ is a constant that depends only on n and p . When \mathbf{X} itself is weakly correlated, one can directly set r as 0.

4.1. Subgroup Identification and Variable Selection

In this section, we resort to Algorithm 1 to solve the optimization problem (2.1) with the ℓ_2 -type distance. To emphasize the advantage of CAR, we construct FA-PFP for comparison. FA-PFP initially considers the FARM and then replaces CAR with $\sum_{i < j} p(a_i - a_j)$ in the optimization problem (2.1), where $p(\cdot)$ is the SCAD

penalty. Tuning parameters involved in FA-PFP is selected with the same strategy as that of the SILFS. Additionally, we compare SILFS with the standard center-augmented regularization model (S-CAR), which ignores the factor structure, to assess the impact of factor structure on subgroup identification and variable selection. Furthermore, we include the oracle estimator with known factors as a comparative benchmark for all models.

We adopt a data generation process similar to that of [Zhang et al. \(2019\)](#) and [Fan et al. \(2020\)](#). Specifically, the model setting is defined as:

$$Y_i = \alpha_i + \mathbf{X}_i^\top \boldsymbol{\beta} + \epsilon_i, \quad \mathbf{X}_i = \mathbf{B} \mathbf{f}_i + \mathbf{u}_i, \quad i = 1, \dots, n,$$

where $\mathbf{f}_i \in \mathbb{R}^r$ and $\boldsymbol{\beta} \in \mathbb{R}^p$. We set ϵ_i are i.i.d. drawn from $\mathcal{N}(0, 0.1)$, and $\boldsymbol{\beta} = (\beta_1, \dots, \beta_5, \mathbf{0}_{p-5}^\top)^\top$ with $\beta_j \sim \mathcal{U}(0.8, 1)$ for $j \leq 5$. For the factor model of \mathbf{X} , let B_{ij} are i.i.d. drawn from $\mathcal{U}(0, 1)$. Take the dependence of factor score into consideration, we employ a vector AR(1) model for \mathbf{f}_i : $\mathbf{f}_i = \boldsymbol{\Phi} \mathbf{f}_{i-1} + \boldsymbol{\xi}_i$, where the noise $\boldsymbol{\xi}_i$ are i.i.d. drawn from $\mathcal{N}(\mathbf{0}_r, 0.1 \mathbf{I}_r)$ and $\Phi_{st} = (0.5)^{\mathbb{I}(s=t)} (0.3)^{|s-t|}$. The idiosyncratic errors \mathbf{u}_i are i.i.d. generated from $\mathcal{N}(\mathbf{0}_p, 0.1 \mathbf{I}_p)$. As for group-specific parameters α_i , the following two scenarios are considered.

Scenario A Set $K = 2$ and $\mathbb{P}(\alpha_i = -a) = \mathbb{P}(\alpha_i = a) = 1/2$, where $a \in \{3, 5\}$.

Scenario B Set $K = 3$ and $\mathbb{P}(\alpha_i = -a) = \mathbb{P}(\alpha_i = 0) = \mathbb{P}(\alpha_i = a) = 1/3$, where $a \in \{3, 5\}$.

In the simulation studies, we fix the number of factors as $r = 4$, set the sample size as $n = 100$, and vary the dimensionality p across 50, 100, and 150. All simulation results in this paper are based on 100 replications. To evaluate the empirical performance of the estimators, we provide the following three criteria:

(i) Estimation error indices, used by both [Zhang et al. \(2019\)](#) and [He et al. \(2022c\)](#), involve the Root Mean Squared Error (RMSE). Specifically, the RMSE for $\boldsymbol{\alpha}$ and $\boldsymbol{\beta}$ is denoted as $\text{RMSE}_{\boldsymbol{\alpha}}$ and $\text{RMSE}_{\boldsymbol{\beta}}$ respectively. The $\text{RMSE}_{\boldsymbol{\beta}}$ is computed as $(\sum_{i=1}^N \|\widehat{\boldsymbol{\beta}}^{(i)} - \boldsymbol{\beta}\|_2^2 / (Np))^{1/2}$, where $\widehat{\boldsymbol{\beta}}^{(i)}$ represents the estimate of $\boldsymbol{\beta}$ in the i -th replication and N is the number of replications. These indices evaluate the estimation accuracy, and a smaller $\text{RMSE}_{\boldsymbol{\alpha}}$ implies better subgroup identification.

(ii) Variable selection indices include the mean of Sensitivity and Specificity over 100 replications. These metrics are common and crucial for assessing feature selections, as illustrated in [Chen et al. \(2021a\)](#).

(iii) Subgroup identification indices include the Rand index (RI), a quantity between 0 and 1. It is commonly used to measure the performance of subgroup recovery, as discussed in [Rand \(1971\)](#) and [Zhang et al. \(2019\)](#). A higher Rand index value indicates better clustering performance. Additionally, we calculate the mean value of the estimated clusters, denoted as $\widehat{K}_{\text{mean}}$. The frequency of overestimation/underestimation of the group number is presented in the form $a|b$, where a and b represent the frequencies of overestimation and underestimation, respectively. These clustering-related indices are also adopted in [Liu et al. \(2023\)](#) to comprehensively evaluate clustering performance.

Table 1: Simulation results for Scenario A. The values in the parentheses denote standard deviation.

Case (a, n, p, K)	Method	Estimation Error Indices		Subgroup Identification Indices			Variable Selection Indices	
		RMSE $_{\alpha}$	RMSE $_{\beta}$	\hat{K}_{mean}	Freq	RI	Sensitivity	Specificity
(3, 100, 50, 2)	FA-PFP	0.347	0.130	2.250(0.672)	17 0	0.997(0.007)	0.992(0.039)	0.979(0.025)
	SILFS	0.128	0.109	2.100(0.302)	10 0	0.985(0.045)	1.000(0.000)	0.986(0.023)
	Oracle	0.068	0.125	NA	NA	NA	0.996(0.040)	0.996(0.009)
	S-CAR	0.313	0.212	2.430(0.820)	27 0	0.951(0.091)	0.678(0.233)	0.937(0.043)
(3, 100, 100, 2)	FA-PFP	0.363	0.094	2.210(0.498)	17 0	0.997(0.006)	0.990(0.044)	0.978(0.019)
	SILFS	0.148	0.084	2.090(0.288)	9 0	0.987(0.042)	0.982(0.076)	0.990(0.013)
	Oracle	0.067	0.084	NA	NA	NA	1.000(0.000)	0.997(0.005)
	S-CAR	0.343	0.158	2.590(0.954)	33 0	0.932(0.104)	0.626(0.285)	0.950(0.028)
(3, 100, 150, 2)	FA-PFP	0.400	0.084	2.270(0.633)	19 0	0.996(0.013)	0.962(0.120)	0.976(0.021)
	SILFS	0.241	0.097	2.320(0.618)	13 0	0.960(0.076)	0.907(0.215)	0.999(0.003)
	Oracle	0.072	0.067	NA	NA	NA	1.000(0.000)	0.997(0.004)
	S-CAR	0.334	0.126	2.550(0.936)	31 0	0.936(0.102)	0.656(0.280)	0.965(0.021)
(5, 100, 50, 2)	FA-PFP	0.558	0.157	2.160(0.507)	11 0	0.948(0.116)	0.942(0.042)	0.998(0.005)
	SILFS	0.168	0.156	2.090(0.288)	9 0	0.989(0.037)	0.984(0.073)	0.997(0.015)
	Oracle	0.081	0.120	NA	NA	NA	1.000(0.000)	0.996(0.009)
	S-CAR	0.334	0.203	2.520(0.904)	31 0	0.942(0.097)	0.702(0.270)	0.937(0.043)
(5, 100, 100, 2)	FA-PFP	0.608	0.118	2.270(0.709)	16 0	0.920(0.136)	0.949(0.038)	0.997(0.010)
	SILFS	0.137	0.105	2.070(0.326)	5 0	0.991(0.041)	0.980(0.115)	0.999(0.003)
	Oracle	0.079	0.085	NA	NA	NA	1.000(0.000)	0.996(0.007)
	S-CAR	0.330	0.158	2.560(0.903)	34 0	0.935(0.100)	0.612(0.278)	0.955(0.024)
(5, 100, 150, 2)	FA-PFP	0.560	0.098	2.170(0.551)	12 0	0.936(0.124)	0.957(0.025)	0.998(0.005)
	SILFS	0.241	0.095	2.230(0.468)	21 0	0.968(0.065)	0.934(0.173)	0.998(0.012)
	Oracle	0.073	0.071	NA	NA	NA	0.998(0.020)	0.997(0.005)
	S-CAR	0.341	0.128	2.580(0.934)	35 0	0.935(0.098)	0.618(0.296)	0.968(0.018)

The simulation results for Scenario A and B are shown in Table 1 and 2 respectively. There are four main takeaways from Table 1 for Scenario A. Firstly, though the S-CAR method proficiently identifies the number of subgroups, its Rand index is relatively lower compared with the SILFS which incorporates the factor structure. Secondly, SILFS outperforms both the FA-PFP and S-CAR methods in terms of variable selection and estimation accuracy. It also demonstrates comparable performance to the oracle method, highlighting the importance of overcoming feature dependence. Moreover, despite both FA-PFP and SILFS incorporating factor structures, FA-PFP exhibits poor performance in terms of RMSE. This can be attributed to its tendency to generate isolated and small-sized subgroups, leading to an overestimation of K . Lastly, as the group parameter a increases, the performance of SILFS improves correspondingly with the increased distance between groups. Since the FA-PFP and SILFS is equipped with factor structures, the increase in

Table 2: Simulation results for Scenario B. The values in the parentheses denote standard deviation.

Case (a, n, p, K)	Method	Estimation Error Indices		Subgroup Identification Indices			Variable Selection Indices	
		RMSE $_{\alpha}$	RMSE $_{\beta}$	\hat{K}_{mean}	Freq	RI	Sensitivity	Specificity
(3, 100, 50, 3)	FA-PFP	0.689	0.218	4.140(1.181)	67 1	0.932(0.048)	0.778(0.244)	0.914(0.058)
	SILFS	0.455	0.184	3.000(0.000)	0 0	0.971(0.036)	0.860(0.234)	0.972(0.037)
	Oracle	0.080	0.113	NA	NA	NA	1.000(0.000)	0.994(0.011)
	S-CAR	0.733	0.244	3.100(0.560)	21 11	0.917(0.079)	0.550(0.252)	0.894(0.060)
(3, 100, 100, 3)	FA-PFP	0.712	0.162	4.270(1.370)	67 1	0.930(0.051)	0.752(0.248)	0.920(0.042)
	SILFS	0.526	0.146	3.000(0.000)	0 0	0.961(0.042)	0.770(0.285)	0.978(0.025)
	Oracle	0.090	0.081	NA	NA	NA	1.000(0.000)	0.995(0.007)
	S-CAR	0.738	0.183	3.170(0.551)	25 8	0.915(0.074)	0.456(0.272)	0.931(0.033)
(3, 100, 150, 3)	FA-PFP	0.742	0.142	4.130(1.292)	62 2	0.925(0.053)	0.674(0.275)	0.926(0.042)
	SILFS	0.599	0.122	3.000(0.000)	0 0	0.949(0.042)	0.812(0.236)	0.946(0.037)
	Oracle	0.076	0.063	NA	NA	NA	0.998(0.020)	0.996(0.005)
	S-CAR	0.720	0.152	3.070(0.432)	13 6	0.923(0.060)	0.428(0.248)	0.942(0.027)
(5, 100, 50, 3)	FA-PFP	0.684	0.192	6.580(3.075)	86 0	0.956(0.063)	0.826(0.289)	0.931(0.048)
	SILFS	0.254	0.130	3.000(0.141)	0 0	0.997(0.014)	0.976(0.104)	0.986(0.044)
	Oracle	0.081	0.112	NA	NA	NA	1.000(0.000)	0.995(0.011)
	S-CAR	0.423	0.216	3.310(0.581)	25 0	0.976(0.041)	0.678(0.224)	0.928(0.050)
(5, 100, 100, 3)	FA-PFP	0.644	0.136	6.520(2.761)	95 0	0.966(0.044)	0.864(0.225)	0.938(0.033)
	SILFS	0.346	0.101	3.010(0.100)	1 0	0.993(0.021)	0.962(0.141)	0.982(0.041)
	Oracle	0.090	0.078	NA	NA	NA	1.000(0.000)	0.995(0.008)
	S-CAR	0.480	0.164	3.310(0.506)	31 1	0.972(0.045)	0.592(0.224)	0.952(0.034)
(5, 100, 150, 3)	FA-PFP	0.713	0.118	6.750(3.010)	93 0	0.954(0.062)	0.808(0.295)	0.948(0.033)
	SILFS	0.423	0.088	3.000(0.000)	0 0	0.991(0.024)	0.934(0.184)	0.976(0.048)
	Oracle	0.083	0.063	NA	NA	NA	0.998(0.020)	0.996(0.005)
	S-CAR	0.531	0.140	3.330(0.533)	28 1	0.970(0.044)	0.596(0.248)	0.957(0.030)

dimensionality has no significant impact on the estimation errors. The results in Table 2 indicate that in the case of three groups, S-CAR performs worse in terms of subgroup identification and completely loses power for variable selection. FA-PFP also exhibits a significant decline in terms of variable selection. In contrast, SILFS remains stable and continues to perform closely well with the oracle estimator. Overall, SILFS performs satisfactorily in various cases.

Next, we show the computing time of the SILFS and FA-PFP methods for Scenario A with $(\gamma_1, \gamma_2) = (-3, 3)$ and Scenario B with $(\gamma_1, \gamma_2, \gamma_3) = (-3, 0, 3)$. For fair comparison, we use the same stopping criteria for SILFS and FA-PFP methods and only report the average time for 10 replications after selecting the optimal tuning parameters. The line charts of computing time are shown in Figure 2. As the sample size increases, the computing time for FA-PFP increases rapidly, with a quadratic curve trend. In contrast, the

computational time required for SILFS increases steadily, almost linearly with respect to sample size.




Figure 2: Comparison of Computing Time between FAPFP and SILFS Algorithms.

In summary, based on the simulation results, we draw the following conclusions: (a) Existing methods in the literature show significantly reduced accuracy in variable selection and subgroup identification in the presence of covariate dependence. In contrast, the SILFS method performs well in both clustering and variable selection tasks. This is supported by high Rand Index and simplicity measures, as well as regression parameter estimators closely aligning with their true values; (b) Compared to the FA-PFP method, SILFS substantially reduces computational burdens, especially with large sample sizes and dimensions.

4.2. Sensitivity to the Level of Collinearity

In this section, we investigate the performance of SILFS in scenarios where the covariates exhibit high collinearity but do not have a factor structure. We also consider the uncorrelated case. Specifically, we generate the data as follows:

$$Y_i = \alpha_i + \mathbf{x}_i^\top \boldsymbol{\beta} + \epsilon_i, \quad i = 1, \dots, n.$$

where $\boldsymbol{\beta} = (\beta_1, \dots, \beta_{10}, \mathbf{0}_{p-10}^\top)^\top$ with $\beta_j \sim \mathcal{U}(1, 2)$ for $j \leq 10$ and $\mathbb{P}(\alpha_i = -3) = \mathbb{P}(\alpha_i = 3) = 1/2$. For the generation of covariate matrices, we consider the following cases:

- **Collinearity Case:** We draw \mathbf{x}_i i.i.d. from $\mathcal{N}(\mathbf{0}_p, \boldsymbol{\Lambda})$, where $\boldsymbol{\Lambda} = \boldsymbol{\Gamma}\boldsymbol{\Gamma}^\top + \mathbf{I}_p$. As for $\boldsymbol{\Gamma}$, we first generate a p -dimensional random square matrix \mathbf{A} , where A_{ij} are i.i.d. from $\mathcal{U}(0, 1)$. Then, by QR decomposition, we have $\mathbf{A} = \mathbf{Q}\mathbf{R}$, where $\mathbf{Q} = (\mathbf{q}_1, \dots, \mathbf{q}_p)$ is an orthogonal matrix. Finally, we set $\boldsymbol{\Gamma} = 5(\mathbf{q}_1, \dots, \mathbf{q}_s)$. When $s = p$, $\boldsymbol{\Gamma}\boldsymbol{\Gamma}^\top = 25\mathbf{I}_p$. In the simulation, we vary s as 3, 4, and 5, respectively.
- **Uncorrelated Case:** We draw \mathbf{x}_i i.i.d. from $\mathcal{N}(\mathbf{0}_p, \mathbf{I}_p)$.

Table 3: SILFS estimation compared with S-CAR under the collinearity and uncorrelated cases. The values in parentheses denote the standard deviation.

	SILFS					S-CAR				
	RMSE $_{\alpha}$	RMSE $_{\beta}$	RI	Sensitivity	Specificity	RMSE $_{\alpha}$	RMSE $_{\beta}$	RI	Sensitivity	Specificity
Collinearity case with $s = 3$										
$p = 50$	0.432	0.146	0.990(0.030)	0.994(0.045)	0.999(0.004)	0.723	0.074	0.970(0.062)	1.000(0.000)	0.949(0.102)
$p = 100$	0.530	0.104	0.985(0.033)	0.996(0.028)	0.998(0.007)	0.605	0.055	0.979(0.055)	0.998(0.020)	0.984(0.042)
$p = 150$	0.726	0.092	0.972(0.056)	0.984(0.061)	0.999(0.008)	0.667	0.053	0.975(0.060)	1.000(0.000)	0.987(0.029)
Collinearity case with $s = 4$										
$p = 50$	0.396	0.145	0.992(0.024)	0.998(0.020)	0.999(0.007)	0.770	0.144	0.972(0.054)	0.994(0.034)	0.963(0.031)
$p = 100$	0.675	0.114	0.976(0.050)	0.990(0.052)	0.997(0.011)	0.849	0.110	0.965(0.055)	0.994(0.045)	0.987(0.015)
$p = 150$	0.796	0.097	0.965(0.055)	0.990(0.044)	0.996(0.009)	0.997	0.103	0.951(0.069)	0.982(0.064)	0.994(0.012)
Collinearity case with $s = 5$										
$p = 50$	0.531	0.162	0.985(0.032)	0.996(0.028)	0.998(0.008)	0.888	0.161	0.964(0.060)	0.992(0.039)	0.960(0.034)
$p = 100$	0.707	0.115	0.973(0.045)	0.994(0.034)	0.996(0.010)	1.011	0.120	0.950(0.079)	0.984(0.061)	0.985(0.017)
$p = 150$	0.909	0.101	0.956(0.069)	0.978(0.069)	0.995(0.011)	1.005	0.103	0.952(0.080)	0.984(0.068)	0.991(0.013)
Uncorrelated case										
$p = 50$	0.220	0.117	0.998(0.008)	1.000(0.000)	1.000(0.002)	0.226	0.115	0.998(0.007)	1.000(0.000)	1.000(0.000)
$p = 100$	0.551	0.097	0.984(0.044)	0.992(0.049)	0.999(0.006)	0.471	0.093	0.988(0.033)	0.998(0.020)	0.998(0.008)
$p = 150$	0.478	0.077	0.988(0.033)	0.998(0.020)	0.999(0.005)	0.426	0.074	0.991(0.027)	1.000(0.000)	0.999(0.002)

We set the sample size as 100, while the dimensionality varies across 50, 100, and 150. All simulation results are based on 100 replications, and the evaluation metrics are consistent with those used in the last section. We present the simulation results in Table 3. In the collinearity case, the covariance matrix exhibits a spike structure. Therefore, although there is no factor structure, the proposed SILFS can still achieve better performance in terms of subgroup identification and variable selection simultaneously. Additionally, as Table 3 illustrates, for the cases where the covariates are uncorrelated, we find that the estimated \hat{r} is quite large such that we believe there is no factor structures and directly set $\hat{r} = 0$. Therefore, the performance of SILFS is comparable with that of SCAR. This indicates that SILFS can be used as a safe replacement of the existing clustering methods, regardless of the level of collinearity among covariates.

5. A Real Data Example

In this section, we employ the proposed SILFS method to explore the relationship between China's export value of commodities and exchange rates. The explanatory variables are sourced from the General Administration of Customs People's Republic of China and are available for download from <http://stats.customs.gov.cn/>. The raw datasets comprises panel data involving 219 trading partner countries (regions) of China and 61 non-industrial commodities. We focus on the top 50 countries (regions) with the highest total trade volume in 2019, resulting in a covariate dimension of 50×61 . The response variable is the Chinese exchange rates corresponding to these 50 countries (regions) in 2019. Our objective is to conduct subgroup analysis for this response variable while accounting for the effects of the covariates.

Given the possibility for high collinearity among covariates due to commodity substitutability, we first conduct PCA on the covariates. The scree plot of the top 20 principal components, depicted in the left panel

of Figure 3, reveals that the first principal component explains 45% of the total variance, while the first five components collectively account for 80% of the variance. This underscores significant cross-sectional dependence among covariates, emphasizing the necessity of applying a factor model to mitigate the impact of collinearity.




Figure 3: Scree plot (left panel) of eigenvalues (dotted line) and proportions of variance explained (bar) by the top 20 principal components. Heatmap (right panel) of the absolute values of the factor loading matrix after orthogonal rotation.

To begin with, we determine the number of factors as $\hat{r} = 3$ using the eigenvalue-ratio method. To aid in interpreting these factors, we employ varimax orthogonal rotation through the `varimax` function in R. The heatmap in the right panel of Figure 3 displays the absolute values of the factor loading matrix after orthogonal rotation, where darker colors indicate larger values. From Figure 3, we conclude that the first factor is associated with variables such as “Plastics,” “KnitApparel” (knitted or crocheted garments and clothing accessories), “OrgChem” (Organic Chemicals), and “Leather Goods”, representing categories related to textiles, apparel, and accessories. This aligns with China’s prominent role as an exporter of garments and textile products, as highlighted in studies such as [Altenburg et al. \(2020\)](#) and [Hussain et al. \(2020\)](#). The second factor primarily relates to “MineralFuels” (mineral fuels, mineral oils, products of their distillation; bituminous substances; mineral waxes), indicating its association with fuel and crude oil exports. In contrast, the third factor is influenced by variables such as “OrgChem,” “Plastics,” “ManmadeFilaments” (synthetic filament; flat strips and similar forms of synthetic textile materials), and “ManmadeFibres” (chemical fiber staple), suggesting a connection to light industry and manufacturing. Therefore, we deduce that China’s export structure is primarily driven by three latent factors, which we summarize as the “Textiles and Apparel Category”, “Oil and Fuel Category”, and “Light Industry Category”.

Next, we apply the FARM model proposed by [Fan et al. \(2020, 2023\)](#) to estimate $\hat{\beta}$. We further plot the

kernel density estimate (KDE) of $Y_i - \mathbf{X}_i^\top \hat{\boldsymbol{\beta}}$ to assess its performance. According to Figure 4, it is evident that even after adjusting for the influence of covariates, the distribution still exhibits multiple modes. This heterogeneity may stem from unobserved latent factors, such as subgroups. Therefore, employing SILFS for subgroup analysis appears to be a more appropriate approach.




Figure 4: Density plot of the response variable after adjusting for the effects of covariates in the empirical example.

We present the results of subgroup identification in Table 4. Based on these findings, we cluster fifty countries (regions) into four groups. From the perspective of developmental status, the first two groups primarily comprise developed countries, including the United States, European countries, Australia and Singapore. Specifically, the first group exhibits greater concentration, while the second group displays variations in geographical location and industrial structure. The third group primarily consists of developing countries with significant growth potential. The last group mainly comprises countries with lower economic levels. Additionally, Hong Kong, China, and Taiwan, China, representing regions with smaller scales, are included in this group. From a geopolitical standpoint, the first group mainly comprises early-developed capitalist countries in Europe and America. The second group consists of countries from the Commonwealth. The third and fourth groups, which are less developed, are primarily situated in Asia and the Middle East.

6. Discussion

Subgroup identification is crucial for characterizing heterogeneity in datasets from various fields such as biology, economics and finance. However, existing methods often rely on the computationally intensive pairwise fusion penalty (PFP) for group pursuit (Ma and Huang, 2017; Zhang et al., 2019). Moreover, high collinearity among covariates can lead to poor performance. To address these challenges, we propose SILFS

Table 4: Subgroup identification of the top 50 countries (regions) from SILFS .

Cluster	Countries (Regions)
C1	United States; Germany; United Kingdom; Netherlands; Italy; Spain; France; Belgium; Poland; Panama; Greece.
C2	Australia; Canada; Singapore; New Zealand.
C3	Japan; Malaysia; Brazil; United Arab Emirates; Pakistan; Saudi Arabia; Israel; Peru.
C4	Hong Kong, China; Vietnam; South Korea; India; Indonesia; Philippines; Thailand; Russia; Taiwan, China; Bangladesh; Mexico; Nigeria; Turkey; Myanmar; South Africa; Cambodia; Egypt; Chile; Iran; Kazakhstan; Colombia; Iraq; Kyrgyzstan; Sri Lanka; Algeria; North Korea; Argentina.

to conduct variable selection and subgroup detection simultaneously. Essentially, we adopt a factor structure to represent collinearity and use the FARM framework (Fan et al., 2021) in high dimensions for subgroup analysis. Furthermore, we introduce a novel Center-Augmented Regularization (CAR) method from He et al. (2022c) for clustering, significantly reducing computational complexity from $O(n^2)$ to $O(nK)$, where n is the sample size and K is the number of subgroups. We also investigate the corresponding algorithms and statistical properties of our proposed method. As a byproduct, an R package called SILFS, implementing the proposed method, is available on CRAN. Extensive simulations and real applications demonstrate the superiority of our proposed method.

Our study provides valuable insights into subgroup analysis in high-dimensional cases with collinearity among covariates. An important future research direction would be an in-depth exploration of the statistical properties of global minima and the theory of clustering consistency. Addressing these challenges will be essential for achieving a comprehensive understanding of the proposed approach. We leave this as future work.

Acknowledgements

This work was supported in part by the National Natural Science Foundation of China (grant numbers: 12171282 and 12101412) and Qilu Young Scholar program of Shandong University.

References

Ahn, S.C., Horenstein, A.R., 2013. Eigenvalue ratio test for the number of factors. *Econometrica* 81, 1203–1227.

- Altenburg, T., Chen, X., Lütkenhorst, W., Staritz, C., Whitfield, L., 2020. Exporting out of China or out of Africa? Automation versus relocation in the global clothing industry. 1/2020, Discussion Paper.
- An, L.T.H., Tao, P.D., 2005. The dc (difference of convex functions) programming and dca revisited with dc models of real world nonconvex optimization problems. *Annals of operations research* 133, 23–46.
- Ando, T., Tsay, R.S., 2011. Quantile regression models with factor-augmented predictors and information criterion. *The Econometrics Journal* 14, 1–24.
- Ashenfelter, O., Krueger, A., 1994. Estimates of the economic return to schooling from a new sample of twins. *The American Economic Review* 84, 1157–1173.
- Bai, J., 2003. Inferential theory for factor models of large dimensions. *Econometrica* 71, 135–171.
- Bai, J., Ng, S., 2002. Determining the number of factors in approximate factor models. *Econometrica* 70, 191–221.
- Banfield, J.D., Raftery, A.E., 1993. Model-based gaussian and non-gaussian clustering. *Biometrics* , 803–821.
- Boyd, S., Parikh, N., Chu, E., Peleato, B., Eckstein, J., et al., 2011. Distributed optimization and statistical learning via the alternating direction method of multipliers. *Foundations and Trends® in Machine learning* 3, 1–122.
- Breiman, L., Cutler, A., 1993. A deterministic algorithm for global optimization. *Mathematical Programming* 58, 179–199.
- Chang, J., Guo, B., Yao, Q., 2015. High dimensional stochastic regression with latent factors, endogeneity and nonlinearity. *Journal of Econometrics* 189, 297–312.
- Chen, H., Guo, Y., He, Y., Ji, J., Liu, L., Shi, Y., et al., 2021a. Simultaneous differential network analysis and classification for matrix-variate data with application to brain connectivity. *Biostatistics* 23, 967–989.
- Chen, L., Dolado, J.J., Gonzalo, J., 2021b. Quantile factor models. *Econometrica* 89, 875–910.
- Chi, E.C., Lange, K., 2015. Splitting methods for convex clustering. *Journal of Computational and Graphical Statistics* 24, 994–1013.
- Connolly, H.C., Gottschalk, P., 2006. Differences in wage growth by education level: do less-educated workers gain less from work experience? .
- Fama, E.F., French, K.R., 1992. The cross-section of expected stock returns. *the Journal of Finance* 47, 427–465.

- Fama, E.F., French, K.R., 2015. A five-factor asset pricing model. *Journal of financial economics* 116, 1–22.
- Fan, J., Ke, Y., Liao, Y., 2021. Augmented factor models with applications to validating market risk factors and forecasting bond risk premia. *Journal of Econometrics* 222, 269–294.
- Fan, J., Ke, Y., Wang, K., 2020. Factor-adjusted regularized model selection. *Journal of Econometrics* 216, 71–85.
- Fan, J., Li, R., 2001. Variable selection via nonconcave penalized likelihood and its oracle properties. *Journal of the American statistical Association* 96, 1348–1360.
- Fan, J., Liao, Y., Mincheva, M., 2013. Large covariance estimation by thresholding principal orthogonal complements. *Journal of the Royal Statistical Society Series B: Statistical Methodology* 75, 603–680.
- Fan, J., Lou, Z., Yu, M., 2023. Are latent factor regression and sparse regression adequate? *Journal of the American Statistical Association* , 1–13.
- Gertheiss, J., Tutz, G., 2012. Regularization and model selection with categorical effect modifiers. *Statistica Sinica* , 957–982.
- Giesen, J., Laue, S., 2019. Combining admm and the augmented lagrangian method for efficiently handling many constraints., in: *IJCAI*, pp. 4525–4531.
- Hastie, T., Tibshirani, R., 1996. Discriminant analysis by gaussian mixtures. *Journal of the Royal Statistical Society Series B: Statistical Methodology* 58, 155–176.
- He, R., Xue, H., Pan, W., Initiative, A.D.N., 2022a. Statistical power of transcriptome-wide association studies. *Genetic epidemiology* 46, 572–588.
- He, Y., Kong, X., Yu, L., Zhang, X., 2022b. Large-dimensional factor analysis without moment constraints. *Journal of Business & Economic Statistics* 40, 302–312.
- He, Y., Li, L., Liu, D., Zhou, W.X., 2023. Huber principal component analysis for large-dimensional factor models. *arXiv preprint arXiv:2303.02817* .
- He, Y., Zhou, L., Xia, Y., Lin, H., 2022c. Center-augmented ℓ_2 -type regularization for subgroup learning. *Biometrics* .
- Hocking, T.D., Joulin, A., Bach, F., Vert, J.P., 2011. Clusterpath an algorithm for clustering using convex fusion penalties, in: *28th international conference on machine learning*, p. 1.

- Hussain, H.I., Haseeb, M., Kot, S., Jermisittiparsert, K., 2020. Non-linear impact of textile and clothing manufacturing on economic growth: The case of top-asian economies. *Fibres & Textiles in Eastern Europe* , 27–36.
- Johnstone, I.M., Paul, D., 2018. Pca in high dimensions: An orientation. *Proceedings of the IEEE* 106, 1277–1292.
- Khalili, A., Chen, J., 2007. Variable selection in finite mixture of regression models. *Journal of the American Statistical Association* 102, 1025–1038.
- Lam, C., Yao, Q., 2012. Factor modeling for high-dimensional time series: inference for the number of factors. *The Annals of Statistics* , 694–726.
- Li, S., Cai, T.T., Li, H., 2022. Transfer learning for high-dimensional linear regression: Prediction, estimation and minimax optimality. *Journal of the Royal Statistical Society Series B: Statistical Methodology* 84, 149–173.
- Liu, D., Zhao, C., He, Y., Liu, L., Guo, Y., Zhang, X., 2023. Simultaneous cluster structure learning and estimation of heterogeneous graphs for matrix-variate fmri data. *Biometrics* 79, 2246–2259.
- Livingston, M., 1977. Industry movements of common stocks. *Journal of Finance* 32, 861–874.
- Ma, S., Huang, J., 2017. A concave pairwise fusion approach to subgroup analysis. *Journal of the American Statistical Association* 112, 410–423.
- McCracken, M.W., Ng, S., 2016. Fred-md: A monthly database for macroeconomic research. *Journal of Business & Economic Statistics* 34, 574–589.
- Merlevède, F., Peligrad, M., Rio, E., 2011. A bernstein type inequality and moderate deviations for weakly dependent sequences. *Probability Theory and Related Fields* 151, 435–474.
- Michele, R., Francesco, V., 2018. From the cradle to the grave: The influence of family background on the career path of italian men. *Oxford Bulletin of Economics and Statistics* 80, 1062–1088.
- Ollier, E., Viallon, V., 2017. Regression modelling on stratified data with the lasso. *Biometrika* 104, 83–96.
- Pan, W., Shen, X., Liu, B., 2013. Cluster analysis: Unsupervised learning via supervised learning with a non-convex penalty. *Journal of Machine Learning Research* 14.
- Phillips, P.C., Sul, D., 2007. Transition modeling and econometric convergence tests. *Econometrica* 75, 1771–1855.

- Porcu, E., Rieger, S., Lepik, K., Santoni, F.A., Reymond, A., Kutalik, Z., 2019. Mendelian randomization integrating gwas and eqtl data reveals genetic determinants of complex and clinical traits. *Nature communications* 10, 3300.
- Rand, W.M., 1971. Objective criteria for the evaluation of clustering methods. *Journal of the American Statistical association* 66, 846–850.
- Shen, X., Huang, H.C., 2010. Grouping pursuit through a regularization solution surface. *Journal of the American Statistical Association* 105, 727–739.
- Song, W., Yao, W., Xing, Y., 2014. Robust mixture regression model fitting by laplace distribution. *Computational Statistics & Data Analysis* 71, 128–137.
- Stock, J.H., Watson, M.W., 2002. Forecasting using principal components from a large number of predictors. *Journal of the American statistical association* 97, 1167–1179.
- Tang, X., Xue, F., Qu, A., 2021. Individualized multidirectional variable selection. *Journal of the American Statistical Association* 116, 1280–1296.
- Tian, Y., Gu, Y., Feng, Y., 2023. Learning from similar linear representations: Adaptivity, minimaxity, and robustness. *arXiv preprint arXiv:2303.17765* .
- Tibshirani, R., 1996. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society Series B: Statistical Methodology* 58, 267–288.
- Tu, Y., Wang, S., 2024. Consistent model selection for factor-augmented regressions. *manuscript* .
- Vatcheva, K.P., Lee, M., McCormick, J.B., Rahbar, M.H., 2016. Multicollinearity in regression analyses conducted in epidemiologic studies. *Epidemiology (Sunnyvale, Calif.)* 6.
- Wang, N., Zhang, X., Mai, Q., 2023. Statistical analysis for a penalized em algorithm in high-dimensional mixture linear regression model. *arXiv preprint arXiv:2307.11405* .
- Wang, S., Wang, X., Zhao, Q., Yang, X., 2019. Quantile factor-augmented prediction model and its applications to alpha-arbitrage strategy in china’s stock market, in: *2019 International Conference on Big Data, Electronics and Communication Engineering (BDECE 2019)*, Atlantis Press. pp. 73–78.
- Wang, W., Su, L., 2021. Identifying latent group structures in nonlinear panels. *Journal of Econometrics* 220, 272–295.
- Wang, W., Zhu, Z., 2024. Homogeneity and sparsity analysis for high-dimensional panel data models. *Journal of Business & Economic Statistics* 42, 26–35.

- Wei, S., Kosorok, M.R., 2013. Latent supervised learning. *Journal of the American Statistical Association* 108, 957–970.
- Wu, C., Kwon, S., Shen, X., Pan, W., 2016. A new algorithm and theory for penalized regression-based clustering. *The Journal of Machine Learning Research* 17, 6479–6503.
- Yao, W., Wei, Y., Yu, C., 2014. Robust mixture regression using the t-distribution. *Computational Statistics & Data Analysis* 71, 116–127.
- Zhang, L., Ma, R., Cai, T.T., Li, H., 2020. Estimation, confidence intervals, and large-scale hypotheses testing for high-dimensional mixed linear regression. *arXiv preprint arXiv:2011.03598* .
- Zhang, X., Zhang, Q., Ma, S., Fang, K., 2022. Subgroup analysis for high-dimensional functional regression. *Journal of Multivariate Analysis* 192, 105100.
- Zhang, Y., Wang, H.J., Zhu, Z., 2019. Robust subgroup identification. *Statistica Sinica* 29, 1873–1889.
- Zhao, P., Yu, B., 2006. On model selection consistency of lasso. *The Journal of Machine Learning Research* 7, 2541–2563.
- Zou, H., 2006. The adaptive lasso and its oracle properties. *Journal of the American statistical association* 101, 1418–1429.
- Zou, H., Hastie, T., 2005. Regularization and variable selection via the elastic net. *Journal of the Royal Statistical Society Series B: Statistical Methodology* 67, 301–320.

APPENDIX

A. Technical Lemmas

In this section, we introduce some useful technical lemmas. The asymptotic properties of factor model estimation are frequently used in the following proof process. Hence, we summarize some useful results from [Fan et al. \(2013\)](#) in the following lemma. Besides, we also deliver some lemmas related to the subgroup identification.

Lemma A.1. [[Fan et al. \(2013\)](#)] Let $l = 3/l_1 + 3/(2l_2) + 1/l_3 + 1$. Suppose $\log(p) = o(n^{l/6})$, $n = o(p^2)$ and Assumptions 1, 2 and 3 referred in the paper hold, we have:

- (i) $1/n \sum_{i=1}^n \|\hat{\mathbf{f}}_i - \mathbf{H}\mathbf{f}_i\|_2^2 = O_{\mathbb{P}}(1/n + 1/p)$.
- (ii) $\mathbf{H}\mathbf{H}^\top = \mathbf{I}_r + O_{\mathbb{P}}(1/\sqrt{n} + 1/\sqrt{p})$.
- (iii) For any subset of $\{1, 2, \dots, p\}$, denote as \mathcal{D} , we have $\max_{j \in \mathcal{D}} \sum_{i=1}^n (\hat{u}_{ij} - u_{ij})^2 = O_{\mathbb{P}}(\log |\mathcal{D}| + n/p)$.
- (iv) $\max_{i \in \{1, 2, \dots, p\}} \|\hat{\mathbf{b}}_i - \mathbf{H}\mathbf{b}_i\|_2^2 = O_{\mathbb{P}}(\log p/n + 1/p)$.
- (v) $\max_{i \in \{1, \dots, r\}; j \in \{1, \dots, p\}} 1/n |\sum_{t=1}^n f_{ti} u_{tj}| = O_{\mathbb{P}}(\sqrt{\log p/n})$.
- (vi) $\max_{i \in \{1, \dots, p\}; j \in \{1, \dots, p\}} 1/n |\sum_{t=1}^n u_{ti} u_{tj} - \mathbb{E}(u_{ti} u_{tj})| = O_{\mathbb{P}}(\sqrt{\log p/n})$.

The next lemma reveals the relationship of global optimum to the constrained optimum under some mild conditions.

Lemma A.2. [[Fan et al. \(2020\)](#)] Suppose $Z_1(\Theta)$ is a convex, twice continuously differentiable function defined over \mathbb{R}^n . Let $g(\Theta) \geq 0$ be a convex and continuous penalty function satisfying the following properties:

- (i) $g(\alpha + \beta) = g(\alpha) + g(\beta)$, for any $\alpha \in \mathcal{U}$ and $\beta \in \mathcal{U}^\perp$, where \mathcal{U} is a subspace of \mathbb{R}^n with orthogonal complement \mathcal{U}^\perp .
- (ii) There exists a continuous function $\tilde{g}(\Theta)$, such that $|\alpha^\top \beta| \leq g(\alpha) \tilde{g}(\beta)$, for any $\alpha \in \mathcal{U}^\perp$ and $\beta \in \mathcal{R}^n$.

Let $Z_\lambda(\Theta) = Z_1(\Theta) + \lambda g(\Theta)$ where $\lambda \geq 0$ and denote the constrained minimization as

$$\hat{\Theta} \in \underset{\Theta \in \mathcal{U}}{\operatorname{argmin}} Z_\lambda(\Theta). \tag{A.1}$$

If $\tilde{g}(\nabla Z_1(\hat{\Theta})) < \lambda$ and $\Theta^\top \nabla^2 Z_1(\hat{\Theta}) \Theta > 0$ for all $\Theta \in \mathcal{U}$, then the constrained minimization defined in (A.1) is the global minimizer of $Z_\lambda(\Theta)$.

The following two lemmas provide the upper bound for invertible matrices.

Lemma A.3. For two invertible $n \times n$ matrices \mathbf{A} and \mathbf{B} , suppose $\|\mathbf{A}^{-1}\|\|\mathbf{A} - \mathbf{B}\| < 1$ where $\|\cdot\|$ is an induced norm, we have

$$\|\mathbf{B}^{-1}\| \leq \frac{\|\mathbf{A}^{-1}\|}{1 - \|\mathbf{A}^{-1}\|\|\mathbf{A} - \mathbf{B}\|}.$$

Lemma A.4. [Fan et al. (2020)] Let \mathbf{C} be a $p \times q$ real matrix, and let \mathbf{A} and \mathbf{B} are two $q \times q$ symmetric matrices. Suppose \mathbf{A} is nonsingular and $\|\mathbf{B}\mathbf{A}^{-1}\| < 1$ where $\|\cdot\|$ is an induced norm, then

$$\left\| \mathbf{C}[(\mathbf{A} + \mathbf{B})^{-1} - \mathbf{A}^{-1}] \right\| \leq \frac{\|\mathbf{C}\mathbf{A}^{-1}\|\|\mathbf{B}\mathbf{A}^{-1}\|}{1 - \|\mathbf{B}\mathbf{A}^{-1}\|}.$$

Lemma A.5. Suppose Assumptions 1, 2 and 3 hold, for any vector $\mathbf{a} \in \mathbb{R}^r$ with $\|\mathbf{a}\|_2 = 1$, we have

$$\left\| \widehat{\mathbf{U}}^\top \mathbf{F} \mathbf{a} \right\|_\infty = O_{\mathbb{P}}(\sqrt{n} + n/\sqrt{p}).$$

Lemma A.6. Under the same assumptions as Lemma A.5, we have

$$\left\| \widehat{\mathbf{U}}^\top \widehat{\mathbf{U}} - \mathbf{U}^\top \mathbf{U} \right\|_{\max} = O_{\mathbb{P}}(\log p + \sqrt{n} + n/\sqrt{p}).$$

The next lemma guarantees the separability of the oracle estimators.

Lemma A.7. [The separability of the oracle estimators.] Under the same conditions required in Theorem 3.1, for all i and k , if the i -th sample is not from the k -th group, then

$$\mathbb{P}(|\widehat{\alpha}_i^{or} - \widehat{\gamma}_k^{or}| \geq r_n) \rightarrow 1,$$

where $\widehat{\alpha}_i^{or}$ and $\widehat{\gamma}_k^{or}$ are the oracle estimators defined in (3.1). r_n is the minimum gap between different groups defined in Theorem 3.3.

B. Proof of Main Theorems

B.1. Proof of Theorem 2.1

Proof. Without loss of generality, we consider the m -th iteration in the DC procedure. Recall that the optimization problem in the m -th iteration under ℓ_1 -type distance is:

$$(\widehat{\Theta}^{(m)}, \widehat{\delta}^{(m)}) = \underset{\Theta, \delta}{\operatorname{argmin}} Z^{(m)}(\Theta, \delta) \tag{B.1}$$

subject to $\delta_{ik} = \alpha_i - \gamma_k \quad i = 1, \dots, n, k = 1, \dots, K$, and $\gamma_1 \leq \gamma_2 \leq \dots \leq \gamma_K$.

where

$$\begin{aligned} Z^{(m)}(\boldsymbol{\Theta}, \boldsymbol{\delta}) &= \frac{1}{2n} \|\mathbf{Y} - \boldsymbol{\alpha} - \widehat{\mathbf{F}}\boldsymbol{\theta} - \widehat{\mathbf{U}}\boldsymbol{\beta}\|_2^2 + \lambda_1 \sum_{i=1}^n \sum_{k=1}^K |\delta_{ik}| - \lambda_1 \sum_{i=1}^n \sum_{k=2}^K \max(|\widehat{\delta}_{i(k-1)}^{(m-1)}|, |\widehat{\delta}_{ik}^{(m-1)}|) \\ &\quad - \lambda_1 g_2(\widehat{\boldsymbol{\delta}}^{(m-1)}) - \lambda_1 (\nabla g_2(\widehat{\boldsymbol{\delta}}^{(m-1)}))^\top (\boldsymbol{\delta} - \widehat{\boldsymbol{\delta}}^{(m-1)}) + \lambda_2 \|\boldsymbol{\beta}\|_1. \end{aligned}$$

Since $Z^{(m)}(\boldsymbol{\Theta}, \boldsymbol{\delta})$ is a closed and proper convex function and the Lagrangian function of (B.1) has saddle point by the saddle point theorem, then by the argument on convergence in the [Boyd et al. \(2011\)](#) and [Giesen and Laue \(2019\)](#), the standard ADMM converges to a global minimizer. By some primary calculations, the following equalities hold:

$$Z\left(\widehat{\boldsymbol{\Theta}}^{(m-1)}, \widehat{\boldsymbol{\delta}}^{(m-1)}\right) = Z^{(m)}\left(\widehat{\boldsymbol{\Theta}}^{(m-1)}, \widehat{\boldsymbol{\delta}}^{(m-1)}\right), \quad Z\left(\widehat{\boldsymbol{\Theta}}^{(m)}, \widehat{\boldsymbol{\delta}}^{(m)}\right) = Z^{(m+1)}\left(\widehat{\boldsymbol{\Theta}}^{(m)}, \widehat{\boldsymbol{\delta}}^{(m)}\right).$$

By the construction of $Z^{(m)}(\boldsymbol{\Theta}, \boldsymbol{\delta})$, for each $m \in N$,

$$Z^{(m)}\left(\widehat{\boldsymbol{\Theta}}^{(m-1)}, \widehat{\boldsymbol{\delta}}^{(m-1)}\right) > Z^{(m)}\left(\widehat{\boldsymbol{\Theta}}^{(m)}, \widehat{\boldsymbol{\delta}}^{(m)}\right) > Z^{(m+1)}\left(\widehat{\boldsymbol{\Theta}}^{(m)}, \widehat{\boldsymbol{\delta}}^{(m)}\right), \quad (\text{B.2})$$

implying that $Z\left(\widehat{\boldsymbol{\Theta}}^{(m-1)}, \widehat{\boldsymbol{\delta}}^{(m-1)}\right) > Z\left(\widehat{\boldsymbol{\Theta}}^{(m)}, \widehat{\boldsymbol{\delta}}^{(m)}\right) \geq 0$. Here the strict inequality always holds, otherwise the algorithm will be terminated when the equality is satisfied. By the monotone convergence theorem, we claim that $Z\left(\widehat{\boldsymbol{\Theta}}^{(m)}, \widehat{\boldsymbol{\delta}}^{(m)}\right)$ converges as m get larger.

Next, we illustrate the finite step convergence. Note that in each $Z^{(m)}(\boldsymbol{\Theta}, \boldsymbol{\delta})$, $\boldsymbol{\delta}^{(m-1)}$ determines the upper bound function $Z^{(m)}(\boldsymbol{\Theta}, \boldsymbol{\delta})$ only through sign function and indicator function. Thus $Z^{(m)}(\boldsymbol{\Theta}, \boldsymbol{\delta})$ has only a finite set of possible options across all integer m . Therefore there exist a m^* such that for any $m \geq m^*$, $\widehat{\boldsymbol{\Theta}}^{(m)} = \widehat{\boldsymbol{\Theta}}^{(m^*)}$. Note that $(\widehat{\boldsymbol{\Theta}}^{(m^*)}, \widehat{\boldsymbol{\delta}}^{(m^*)})$ is the global minimizer of $Z^{(m^*)}(\boldsymbol{\Theta}, \boldsymbol{\delta})$ and $\nabla Z^{(m^*)}(\widehat{\boldsymbol{\Theta}}^{(m^*)}, \widehat{\boldsymbol{\delta}}^{(m^*)}) = \nabla Z(\widehat{\boldsymbol{\Theta}}^{(m^*)}, \widehat{\boldsymbol{\delta}}^{(m^*)})$, thus we obtain that $\widehat{\boldsymbol{\Theta}}^{(m^*)}$ is a local minimizer of $Z(\widehat{\boldsymbol{\Theta}})$. \square

B.2. Proof of Theorem 3.1 and Proposition 3.2

Proof. Recall that the oracle estimator is defined as:

$$(\widehat{\boldsymbol{\gamma}}^{or}, \widehat{\boldsymbol{\theta}}^{or}, \widehat{\boldsymbol{\beta}}^{or}) = \underset{\boldsymbol{\Theta}}{\operatorname{argmin}} Z_1(\boldsymbol{\Theta}) + \lambda_2 \|\boldsymbol{\beta}\|_1.$$

The proof proceeds in **three** steps. The first step is to show that the constrained optimization estimator converges to the oracle solution. In addition, the convergence rate is also provided. In the second step, we prove that the constrained optimal is equivalent to the global optimal solution. The third step proves the sign consistency of $\widehat{\boldsymbol{\beta}}$.

Step 1. In this step, we consider an optimization problem that limited to the subspace $\mathcal{U} := \{(\boldsymbol{\beta}^\top, \boldsymbol{\theta}^\top, \boldsymbol{\gamma}^\top)^\top \in$

$\mathbb{R}^{p+r+K} : (\boldsymbol{\beta}_S^\top, \mathbf{0}_{S^c}^\top, \boldsymbol{\theta}^\top, \boldsymbol{\gamma}^\top)^\top$. We define

$$\widehat{\boldsymbol{\Theta}}_1 := \left((\widehat{\boldsymbol{\beta}}_1)_S^\top, \mathbf{0}_{S^c}^\top, \widehat{\boldsymbol{\theta}}_1^\top, \widehat{\boldsymbol{\gamma}}_1^\top \right)^\top = \underset{\boldsymbol{\Theta} \in \mathcal{U}}{\operatorname{argmin}} \frac{1}{2n} \left\| \mathbf{Y} - \boldsymbol{\Omega} \boldsymbol{\gamma} - \widehat{\mathbf{F}} \boldsymbol{\theta} - \widehat{\mathbf{U}} \boldsymbol{\beta} \right\|_2^2 + \lambda_2 \|\boldsymbol{\beta}\|_1.$$

Note that this constrained optimization problem is equivalent to the following optimization problem,

$$(\widehat{\boldsymbol{\Theta}}_1)_S := \left((\widehat{\boldsymbol{\beta}}_1)_S^\top, \widehat{\boldsymbol{\theta}}_1^\top, \widehat{\boldsymbol{\gamma}}_1^\top \right)^\top = \underset{\boldsymbol{\beta}_S, \boldsymbol{\theta}, \boldsymbol{\gamma}}{\operatorname{argmin}} \frac{1}{2n} \left\| \mathbf{Y} - \boldsymbol{\Omega} \boldsymbol{\gamma} - \widehat{\mathbf{F}} \boldsymbol{\theta} - \widehat{\mathbf{U}}_S \boldsymbol{\beta}_S \right\|_2^2 + \lambda_2 \|\boldsymbol{\beta}_S\|_1. \quad (\text{B.3})$$

The first order KKT conditions of problem (B.3) are

$$\begin{aligned} \partial Z_1[(\widehat{\boldsymbol{\Theta}}_1)_S] / \partial \boldsymbol{\beta}_S &= -\frac{1}{n} \widehat{\mathbf{U}}_S^\top \left(\mathbf{Y} - \boldsymbol{\Omega} \widehat{\boldsymbol{\gamma}}_1 - \widehat{\mathbf{F}} \widehat{\boldsymbol{\theta}}_1 - \widehat{\mathbf{U}}_S (\widehat{\boldsymbol{\beta}}_1)_S \right) \in -\lambda_2 \partial \|(\widehat{\boldsymbol{\beta}}_1)_S\|_1, \\ \partial Z_1[(\widehat{\boldsymbol{\Theta}}_1)_S] / \partial \boldsymbol{\theta} &= -\frac{1}{n} \widehat{\mathbf{F}}^\top \left(\mathbf{Y} - \boldsymbol{\Omega} \widehat{\boldsymbol{\gamma}}_1 - \widehat{\mathbf{F}} \widehat{\boldsymbol{\theta}}_1 - \widehat{\mathbf{U}}_S (\widehat{\boldsymbol{\beta}}_1)_S \right) = \mathbf{0}, \\ \partial Z_1[(\widehat{\boldsymbol{\Theta}}_1)_S] / \partial \boldsymbol{\gamma} &= -\frac{1}{n} \boldsymbol{\Omega}^\top \left(\mathbf{Y} - \boldsymbol{\Omega} \widehat{\boldsymbol{\gamma}}_1 - \widehat{\mathbf{F}} \widehat{\boldsymbol{\theta}}_1 - \widehat{\mathbf{U}}_S (\widehat{\boldsymbol{\beta}}_1)_S \right) = \mathbf{0}, \end{aligned} \quad (\text{B.4})$$

where $\partial \|\boldsymbol{\Theta}\|_1$ is the sub-gradient of $\|\boldsymbol{\Theta}\|_1$. The above KKT condition implies $\|\nabla_S Z_1(\widehat{\boldsymbol{\Theta}}_1)\|_\infty \leq \lambda_2$. We consider the difference between the first order differential of the $Z_1(\boldsymbol{\Theta})$ at $\boldsymbol{\Theta}_0$ and $\widehat{\boldsymbol{\Theta}}_1$.

$$\nabla_S Z_1(\widehat{\boldsymbol{\Theta}}_1) - \nabla_S Z_1(\boldsymbol{\Theta}_0) = \frac{1}{n} \left(\boldsymbol{\Omega}, \widehat{\mathbf{F}}, \widehat{\mathbf{U}}_S \right)^\top \left(\boldsymbol{\Omega}, \widehat{\mathbf{F}}, \widehat{\mathbf{U}}_S \right) \begin{pmatrix} \widehat{\boldsymbol{\gamma}}_1 - \boldsymbol{\gamma}_0 \\ \widehat{\boldsymbol{\theta}}_1 - \mathbf{H} \boldsymbol{\theta}_0 \\ (\widehat{\boldsymbol{\beta}}_1)_S - (\boldsymbol{\beta}_0)_S \end{pmatrix}. \quad (\text{B.5})$$

Denote

$$\mathbf{A} := \frac{1}{n} \left(\boldsymbol{\Omega}, \widehat{\mathbf{F}}, \widehat{\mathbf{U}}_S \right)^\top \left(\boldsymbol{\Omega}, \widehat{\mathbf{F}}, \widehat{\mathbf{U}}_S \right). \quad (\text{B.6})$$

If \mathbf{A} is invertible, then left multiply both sides by matrix \mathbf{A}^{-1} and take the infinite norm on both sides, we have

$$\left\| (\widehat{\boldsymbol{\Theta}}_1)_S - (\boldsymbol{\Theta}_0)_S \right\|_\infty \leq \left\| \left[\frac{1}{n} \left(\boldsymbol{\Omega}, \widehat{\mathbf{F}}, \widehat{\mathbf{U}}_S \right)^\top \left(\boldsymbol{\Omega}, \widehat{\mathbf{F}}, \widehat{\mathbf{U}}_S \right) \right]^{-1} \right\|_\infty \left\| \nabla_S Z_1(\widehat{\boldsymbol{\Theta}}_1) - \nabla_S Z_1(\boldsymbol{\Theta}_0) \right\|_\infty. \quad (\text{B.7})$$

In the following, we need to illustrate the \mathbf{A} is invertible with probability tending to 1 and focus on the upper bounds for the two infinity norms on the right-hand side of the inequality (B.7).

To show \mathbf{A} is invertible, we set $\mathbf{D} = \operatorname{diag}(\boldsymbol{\Omega}^\top \boldsymbol{\Omega} / n, \mathbf{I}_r, \boldsymbol{\Sigma}_{SS})$ and $\boldsymbol{\Delta} = \mathbf{A} - \mathbf{D}$. By Weyl theorem, we have $|\lambda_{\min}(\mathbf{A}) - \lambda_{\min}(\mathbf{D})| \leq \|\boldsymbol{\Delta}\|_2 \leq \|\boldsymbol{\Delta}\|_\infty$. Thus, it is sufficient to show $\|\boldsymbol{\Delta}\|_\infty$ is $o_{\mathbb{P}}(1)$. On the other hand,

$\|\mathbf{A}^{-1}\|_\infty$ can be bounded by Lemma A.3

$$\|\mathbf{A}^{-1}\|_\infty \leq \frac{\|\mathbf{D}^{-1}\|_\infty}{1 - \|\mathbf{D}^{-1}\|_\infty \|\mathbf{\Delta}\|_\infty}. \quad (\text{B.8})$$

It is not hard to verify that $\|\mathbf{D}^{-1}\|_\infty$ is bounded under Assumption 5 and we assume it is smaller than C . In conclusion, we should find the convergence rate of $\|\mathbf{\Delta}\|_\infty$ to show \mathbf{A} is invertible and then bound \mathbf{A}^{-1} with infinite norm.

To analyze the gap between \mathbf{A} and \mathbf{D} , an ‘‘intermediary matrix’’ of them are introduced as:

$$\mathbf{W} := \frac{1}{n} \begin{pmatrix} \mathbf{\Omega}^\top \mathbf{\Omega} & \mathbf{\Omega}^\top \mathbf{F} \mathbf{H}^\top & \mathbf{\Omega}^\top \mathbf{U}_S \\ \mathbf{H} \mathbf{F}^\top \mathbf{\Omega} & n \mathbf{I}_r & \mathbf{0} \\ \mathbf{U}_S^\top \mathbf{\Omega} & \mathbf{0} & \mathbf{U}_S^\top \mathbf{U}_S \end{pmatrix}.$$

By triangle inequality, we have

$$\|\mathbf{\Delta}\|_\infty = \|\mathbf{D} - \mathbf{W} + \mathbf{W} - \mathbf{A}\|_\infty \leq \|\mathbf{A} - \mathbf{W}\|_\infty + \|\mathbf{D} - \mathbf{W}\|_\infty =: \|\mathbf{\Delta}_1\|_\infty + \|\mathbf{\Delta}_2\|_\infty.$$

By the property of infinite norm, we have

$$\begin{aligned} \|\mathbf{\Delta}_1\|_\infty &\leq \frac{1}{n} \max \left(\left\| \left(\widehat{\mathbf{F}} - \mathbf{F} \mathbf{H}^\top \right)^\top \mathbf{\Omega} \right\|_\infty, \left\| \left(\widehat{\mathbf{U}}_S - \mathbf{U}_S \right)^\top \mathbf{\Omega} \right\|_\infty \right) + \frac{1}{n} \left\| \mathbf{\Omega}^\top \left(\widehat{\mathbf{F}} - \mathbf{F} \mathbf{H}^\top \right) \right\|_\infty \\ &\quad + \frac{1}{n} \max \left(\left\| \mathbf{\Omega}^\top \left(\widehat{\mathbf{U}}_S - \mathbf{U}_S \right) \right\|_\infty, \left\| \widehat{\mathbf{U}}_S^\top \widehat{\mathbf{U}}_S - \mathbf{U}_S^\top \mathbf{U}_S \right\|_\infty \right). \end{aligned} \quad (\text{B.9})$$

Recalling the properties of the factor model estimators listed in Lemma A.1, we obtain that

$$\begin{aligned} \left\| \left(\widehat{\mathbf{F}} - \mathbf{F} \mathbf{H}^\top \right)^\top \mathbf{\Omega} \right\|_\infty &\leq \sqrt{K} \left\| \mathbf{\Omega} \right\|_{\mathbb{F}} \left\| \widehat{\mathbf{F}} - \mathbf{F} \mathbf{H}^\top \right\|_{\mathbb{F}} = O_{\mathbb{P}} \left(\sqrt{n} + n/\sqrt{p} \right). \\ \left\| \mathbf{\Omega}^\top \left(\widehat{\mathbf{F}} - \mathbf{F} \mathbf{H}^\top \right) \right\|_\infty &\leq \sqrt{r} \left\| \mathbf{\Omega} \right\|_{\mathbb{F}} \left\| \widehat{\mathbf{F}} - \mathbf{F} \mathbf{H}^\top \right\|_{\mathbb{F}} = O_{\mathbb{P}} \left(\sqrt{n} + n/\sqrt{p} \right). \end{aligned}$$

Similarly, we also have

$$\begin{aligned} \left\| \left(\widehat{\mathbf{U}}_S - \mathbf{U}_S \right)^\top \mathbf{\Omega} \right\|_\infty &= O_{\mathbb{P}} \left((np_S \log p_S + p_S n^2/p)^{1/2} \right). \\ \left\| \mathbf{\Omega}^\top \left(\widehat{\mathbf{U}}_S - \mathbf{U}_S \right) \right\|_\infty &= O_{\mathbb{P}} \left(p_S (n \log p_S + n^2/p)^{1/2} \right). \end{aligned}$$

According to the result of Lemma A.6, it can be obtained that

$$\left\| \widehat{\mathbf{U}}_S^\top \widehat{\mathbf{U}}_S - \mathbf{U}_S^\top \mathbf{U}_S \right\|_\infty \leq p_S \left\| \widehat{\mathbf{U}}_S^\top \widehat{\mathbf{U}}_S - \mathbf{U}_S^\top \mathbf{U}_S \right\|_{\max} = O_{\mathbb{P}} \left(p_S \left(\log p_S + \sqrt{n} + n/\sqrt{p} \right) \right).$$

Combining all these results and the decomposition in (B.9), it's obvious that

$$\|\mathbf{\Delta}_1\|_\infty = O_{\mathbb{P}}(p_S(\sqrt{\log p_S/n} + 1/\sqrt{p})). \quad (\text{B.10})$$

With similar arguments, we have

$$\|\mathbf{\Delta}_2\|_\infty \leq \frac{1}{n} \max\left(\|\mathbf{H}\mathbf{F}^\top \mathbf{\Omega}\|_\infty, \|\mathbf{U}_S^\top \mathbf{\Omega}\|_\infty\right) + \frac{1}{n} \|\mathbf{\Omega}^\top \mathbf{F}\mathbf{H}^\top\|_\infty + \frac{1}{n} \max\left(\|\mathbf{\Omega}^\top \mathbf{U}_S\|_\infty, \|\mathbf{U}_S^\top \mathbf{U}_S - n\mathbf{\Sigma}_S\|_\infty\right).$$

According to the concentration inequality in Merlevède et al. (2011), we have

$$\mathbb{P}\left(\frac{1}{|\mathcal{G}_k|} \left| \sum_{\mathcal{G}_k} f_{ij} \right| \geq t\right) \leq |\mathcal{G}_k| \exp\left(-\frac{(|\mathcal{G}_k|t)^l}{V_1}\right) + \exp\left(-\frac{(|\mathcal{G}_k|t)^2}{|\mathcal{G}_k|V_2}\right) + \exp\left(-\frac{(|\mathcal{G}_k|t)^2}{|\mathcal{G}_k|V_3} \exp\left(\frac{(|\mathcal{G}_k|t)^{l(1-l)}}{V_4 \log^l(nt)}\right)\right),$$

which implies $|\sum_{\mathcal{G}_k} f_{ij}| = O_{\mathbb{P}}(\sqrt{|\mathcal{G}_k|})$. Hence, we have

$$\|\mathbf{F}^\top \mathbf{\Omega}\|_\infty = O_{\mathbb{P}}(\sqrt{|\mathcal{G}_{\max}|}) \quad \text{and} \quad \|\mathbf{\Omega}^\top \mathbf{F}\|_\infty = O_{\mathbb{P}}(\sqrt{|\mathcal{G}_{\max}|}).$$

With similar arguments, we have

$$\|\mathbf{U}_S^\top \mathbf{\Omega}\|_\infty = O_{\mathbb{P}}(\sqrt{|\mathcal{G}_{\max}| \log p_S}) \quad \text{and} \quad \|\mathbf{\Omega}^\top \mathbf{U}_S\|_\infty = O_{\mathbb{P}}(p_S \sqrt{|\mathcal{G}_{\max}| \log p_S}).$$

Further, by Lemma A.1,

$$\left\| \mathbf{U}_S^\top \mathbf{U}_S - n\mathbf{\Sigma}_S \right\|_\infty \leq p_S \left\| \mathbf{U}_S^\top \mathbf{U}_S - n\mathbf{\Sigma}_S \right\|_{\max} = O_{\mathbb{P}}\left(p_S \sqrt{n \log p_S}\right).$$

Thus, we obtain that $\|\mathbf{\Delta}_2\|_\infty = O_{\mathbb{P}}(p_S \sqrt{\log p_S/n})$. Combining the upper bound of $\|\mathbf{\Delta}_1\|_\infty$ in equation (B.10), we obtain

$$\|\mathbf{\Delta}\|_\infty = O_{\mathbb{P}}(p_S(\sqrt{\log p_S/n} + 1/\sqrt{p})). \quad (\text{B.11})$$

It implies that $\|\mathbf{\Delta}\|_\infty \leq 1/(2C)$ holds with probability tending to 1. Combined with the assumptions in Theorem 3.1, we get $p_S(\sqrt{\log p_S/n} + 1/\sqrt{p}) \rightarrow 0$. Thus, we obtain that \mathbf{A} is invertible with probability tending to 1. Plugging this results in the inequality (B.8) we have

$$\left\| \left[\frac{1}{n} \left(\mathbf{\Omega}, \widehat{\mathbf{F}}, \widehat{\mathbf{U}}_S \right)^\top \left(\mathbf{\Omega}, \widehat{\mathbf{F}}, \widehat{\mathbf{U}}_S \right) \right]^{-1} \right\|_\infty \leq 2C \quad (\text{B.12})$$

holds with probability approaching to 1.

According to (B.7), in order to bound $\left\| (\widehat{\mathbf{\Theta}}_1)_S - (\mathbf{\Theta}_0)_S \right\|_\infty$, we still need to find the upper bound of

$\|\nabla_{\mathcal{S}} Z_1(\Theta_0)\|_{\infty}$. Recall that $\mathbf{X} = \mathbf{F}\mathbf{B}^{\top} + \mathbf{U} = \widehat{\mathbf{F}}\widehat{\mathbf{B}}^{\top} + \widehat{\mathbf{U}}$, we have

$$\mathbf{Y} - \boldsymbol{\Omega}\gamma_0 - \widehat{\mathbf{F}}\mathbf{H}\boldsymbol{\theta}_0 - \widehat{\mathbf{U}}_S(\boldsymbol{\beta}_0)_S = \boldsymbol{\epsilon} + \widehat{\mathbf{F}}\left(\widehat{\mathbf{B}}^{\top} - \mathbf{H}\mathbf{B}^{\top}\right)\boldsymbol{\beta}_0.$$

Then, we can write $\nabla Z_1(\Theta_0)$ as

$$\nabla Z_1(\Theta_0) = -\frac{1}{n}\left(\boldsymbol{\Omega}, \widehat{\mathbf{F}}, \widehat{\mathbf{U}}\right)^{\top}\left(\boldsymbol{\epsilon} + \widehat{\mathbf{F}}\left(\widehat{\mathbf{B}}^{\top} - \mathbf{H}\mathbf{B}^{\top}\right)\boldsymbol{\beta}_0\right). \quad (\text{B.13})$$

We control the two parts of the right hand side of (B.13) separately. The first part is

$$\|(\boldsymbol{\Omega}, \widehat{\mathbf{F}}, \widehat{\mathbf{U}})^{\top}\boldsymbol{\epsilon}\|_{\infty} \leq \max(\|\boldsymbol{\Omega}^{\top}\boldsymbol{\epsilon}\|_{\infty}, \|\widehat{\mathbf{F}}^{\top}\boldsymbol{\epsilon}\|_{\infty}, \|\widehat{\mathbf{U}}^{\top}\boldsymbol{\epsilon}\|_{\infty}). \quad (\text{B.14})$$

Note that $\boldsymbol{\Omega}^{\top}\boldsymbol{\epsilon} = (\sum_{\mathcal{G}_1}\epsilon_i, \dots, \sum_{\mathcal{G}_K}\epsilon_i)$, the concentration inequality of sub-Gaussian random variable leads to $\|\boldsymbol{\Omega}^{\top}\boldsymbol{\epsilon}\|_{\infty} = O_{\mathbb{P}}\left(\sqrt{|\mathcal{G}_{\max}|}\right)$. For $\|\widehat{\mathbf{F}}^{\top}\boldsymbol{\epsilon}\|_{\infty}$, we have

$$\|\widehat{\mathbf{F}}^{\top}\boldsymbol{\epsilon}\|_{\infty} \leq \|\mathbf{H}\mathbf{F}^{\top}\boldsymbol{\epsilon}\|_{\infty} + \|(\widehat{\mathbf{F}}^{\top} - \mathbf{H}\mathbf{F}^{\top})\boldsymbol{\epsilon}\|_2 \leq \|\mathbf{H}\|_{\infty}\|\mathbf{F}^{\top}\boldsymbol{\epsilon}\|_{\infty} + \|(\widehat{\mathbf{F}}^{\top} - \mathbf{H}\mathbf{F}^{\top})\boldsymbol{\epsilon}\|_2.$$

Using the concentration inequality in Merlevède et al. (2011) again, we have

$$\mathbb{P}\left(\frac{1}{n}\left|\sum_{i=1}^n f_{ij}\epsilon_i\right| \geq t\right) \leq n \exp\left(-\frac{(nt)^l}{V_1}\right) + \exp\left(-\frac{(nt)^2}{nV_2}\right) + \exp\left(-\frac{(nt)^2}{nV_3} \exp\left(\frac{(nt)^{l(1-l)}}{V_4 \log^l(nt)}\right)\right).$$

Thus, we know $\|\mathbf{F}^{\top}\boldsymbol{\epsilon}\|_{\infty} \leq \|\mathbf{F}^{\top}\boldsymbol{\epsilon}\|_2 = O_{\mathbb{P}}(\sqrt{n})$. By Cauchy-Schwarz inequality,

$$\begin{aligned} \|\widehat{\mathbf{U}}^{\top}\boldsymbol{\epsilon}\|_{\infty} &\leq \|\mathbf{U}^{\top}\boldsymbol{\epsilon}\|_{\infty} + \max_{j \in \{1 \dots p\}} \left| \sum_{i=1}^n (\widehat{u}_{ij} - u_{ij})\epsilon_i \right| \\ &\leq \|\mathbf{U}^{\top}\boldsymbol{\epsilon}\|_{\infty} + \max_{j \in \{1 \dots p\}} \left(\sum_{i=1}^n (\widehat{u}_{ij} - u_{ij})^2 \right)^{1/2} \left(\sum_{i=1}^n \epsilon_i^2 \right)^{1/2}. \end{aligned} \quad (\text{B.15})$$

With similar arguments of $\|\mathbf{F}^{\top}\boldsymbol{\epsilon}\|_{\infty}$, one can obtain $\|\mathbf{U}^{\top}\boldsymbol{\epsilon}\|_{\infty} = O_{\mathbb{P}}(\sqrt{n \log p})$. Further, combining the convergence rate of factor models in the Lemma A.1, we obtain that

$$\|\widehat{\mathbf{F}}^{\top}\boldsymbol{\epsilon}\|_{\infty} = O_{\mathbb{P}}(\sqrt{n} + n/\sqrt{p}), \quad \text{and} \quad \|\widehat{\mathbf{U}}^{\top}\boldsymbol{\epsilon}\|_{\infty} = O_{\mathbb{P}}\left((n \log p + n^2/p)^{1/2}\right). \quad (\text{B.16})$$

Recall the decomposition in (B.14), we have

$$\|(\boldsymbol{\Omega}, \widehat{\mathbf{F}}, \widehat{\mathbf{U}})^{\top}\boldsymbol{\epsilon}\|_{\infty} = O_{\mathbb{P}}\left((n \log p + n^2/p)^{1/2}\right). \quad (\text{B.17})$$

Next, we bound the second part. According to the maximum inequality we have

$$\begin{aligned} \|(\boldsymbol{\Omega}, \widehat{\mathbf{F}}, \widehat{\mathbf{U}})^\top \widehat{\mathbf{F}}(\widehat{\mathbf{B}}^\top - \mathbf{H}\mathbf{B}^\top)\boldsymbol{\beta}_0\|_\infty &\leq \max\left(\|\boldsymbol{\Omega}^\top \widehat{\mathbf{F}}(\widehat{\mathbf{B}}^\top - \mathbf{H}\mathbf{B}^\top)\boldsymbol{\beta}_0\|_\infty, \right. \\ &\quad \left.\|\widehat{\mathbf{F}}^\top \widehat{\mathbf{F}}(\widehat{\mathbf{B}}^\top - \mathbf{H}\mathbf{B}^\top)\boldsymbol{\beta}_0\|_\infty, \|\widehat{\mathbf{U}}^\top \widehat{\mathbf{F}}(\widehat{\mathbf{B}}^\top - \mathbf{H}\mathbf{B}^\top)\boldsymbol{\beta}_0\|_\infty\right). \end{aligned} \quad (\text{B.18})$$

Note that $\|\boldsymbol{\Omega}^\top \widehat{\mathbf{F}}\|_\infty \leq \|\boldsymbol{\Omega}^\top \widehat{\mathbf{F}}\|_2 \leq \|\boldsymbol{\Omega}\|_2 \|\widehat{\mathbf{F}}\|_2 = O_{\mathbb{P}}(n)$. Then we conclude

$$\|\boldsymbol{\Omega}^\top \widehat{\mathbf{F}}(\widehat{\mathbf{B}}^\top - \mathbf{H}\mathbf{B}^\top)\boldsymbol{\beta}_0\|_\infty \leq n\|(\widehat{\mathbf{B}}^\top - \mathbf{H}\mathbf{B}^\top)\boldsymbol{\beta}_0\|_\infty \leq n\|(\widehat{\mathbf{B}}^\top - \mathbf{H}\mathbf{B}^\top)\boldsymbol{\beta}_0\|_2.$$

According to the Lemma A.1, the following inequality holds

$$\|(\widehat{\mathbf{B}}^\top - \mathbf{H}\mathbf{B}^\top)\boldsymbol{\beta}_0\|_2 \leq \max_{j \in \mathcal{S}} \|\widehat{\mathbf{b}}_j - \mathbf{H}\mathbf{b}_j\|_2 \|\boldsymbol{\beta}_0\|_1 = O_{\mathbb{P}}(p_S(\log p_S/n + 1/p)^{1/2}).$$

Therefore,

$$\|\boldsymbol{\Omega}^\top \widehat{\mathbf{F}}(\widehat{\mathbf{B}}^\top - \mathbf{H}\mathbf{B}^\top)\boldsymbol{\beta}_0\|_\infty = O_{\mathbb{P}}(p_S(n \log p_S + n^2/p)^{1/2}).$$

Note that $\widehat{\mathbf{F}}^\top \widehat{\mathbf{F}} = n\mathbf{I}_r$, we have

$$\|\widehat{\mathbf{F}}^\top \widehat{\mathbf{F}}(\widehat{\mathbf{B}}^\top - \mathbf{H}\mathbf{B}^\top)\boldsymbol{\beta}_0\|_\infty \leq n\|(\widehat{\mathbf{B}}^\top - \mathbf{H}\mathbf{B}^\top)\boldsymbol{\beta}_0\|_\infty = O_{\mathbb{P}}(p_S(n \log p_S + n^2/p)^{1/2}).$$

Recall that estimation of factor model produce $\widehat{\mathbf{U}}^\top \widehat{\mathbf{F}} = \mathbf{0}$. We have

$$\|(\boldsymbol{\Omega}, \widehat{\mathbf{F}}, \widehat{\mathbf{U}})^\top \widehat{\mathbf{F}}(\widehat{\mathbf{B}}^\top - \mathbf{H}\mathbf{B}^\top)\boldsymbol{\beta}_0\|_\infty = O_{\mathbb{P}}(p_S(n \log p + n^2/p)^{1/2}). \quad (\text{B.19})$$

Therefore, combine the results for (B.13), (B.17) and (B.19), we get

$$\|\nabla Z_1(\boldsymbol{\Theta}_0)\|_\infty = O_{\mathbb{P}}(p_S(\log p/n + 1/p)^{1/2}). \quad (\text{B.20})$$

WLOG, we let $c\lambda_2 = \|\nabla Z_1(\boldsymbol{\Theta}_0)\|_\infty$. Combining the inequality in (B.7) and the convergence rate in (B.12), we obtain

$$\begin{aligned} \|(\widehat{\boldsymbol{\Theta}}_1)_S - (\boldsymbol{\Theta}_0)_S\|_\infty &\leq 2C\|\nabla_S Z_1(\widehat{\boldsymbol{\Theta}}_1) - \nabla_S Z_1(\boldsymbol{\Theta}_0)\|_\infty \\ &\leq 2C\left(\|\nabla_S Z_1(\widehat{\boldsymbol{\Theta}}_1)\|_\infty + \|\nabla_S Z_1(\boldsymbol{\Theta}_0)\|_\infty\right) \leq 2C(1+c)\lambda_2. \end{aligned} \quad (\text{B.21})$$

Step 2. In this step, we show that the constrained minima $\widehat{\boldsymbol{\Theta}}_1$ is the global minimum of (3.1). According

to Lemma A.2, we only need to show $\|\nabla_{S^c} Z_1(\widehat{\Theta}_1)\|_\infty \leq \lambda_2$. By Lagrange mean value theorem, we have

$$\nabla_{S^c} Z_1(\widehat{\Theta}_1) - \nabla_{S^c} Z_1(\Theta_0) = \mathbf{L}((\widehat{\Theta}_1)_S - (\Theta_0)_S),$$

where $\mathbf{L} = 1/n(\widehat{\mathbf{U}}_{S^c}^\top \boldsymbol{\Omega}, \mathbf{0}, \widehat{\mathbf{U}}_{S^c}^\top \widehat{\mathbf{U}}_S)$ is the sub-matrix of Hessian matrix of $Z_1(\Theta)$. By norm inequality we get

$$\left\| \nabla_{S^c} Z_1(\widehat{\Theta}_1) \right\|_\infty \leq \left\| \nabla_{S^c} Z_1(\Theta_0) \right\|_\infty + \left\| \mathbf{L} \mathbf{A}^{-1} \right\|_\infty \left\| \mathbf{A}((\widehat{\Theta}_1)_S - (\Theta_0)_S) \right\|_\infty,$$

where \mathbf{A} is defined in (B.6) and $\mathbf{A} = \boldsymbol{\Delta} + \mathbf{D}$. According to Lemma A.4,

$$\left\| \mathbf{L} [\mathbf{A}^{-1} - \mathbf{D}^{-1}] \right\|_\infty \leq \frac{\left\| \mathbf{L} \mathbf{D}^{-1} \right\|_\infty \left\| \boldsymbol{\Delta} \mathbf{D}^{-1} \right\|_\infty}{1 - \left\| \boldsymbol{\Delta} \mathbf{D}^{-1} \right\|_\infty}.$$

Combining the norm inequality and the irrepresentable condition, we have

$$\left\| \mathbf{L} \mathbf{A}^{-1} - (\mathbf{0}_{|S^c| \times K}, \mathbf{0}_{|S^c| \times r}, \boldsymbol{\Sigma}_{S^c S}) \mathbf{D}^{-1} \right\|_\infty = O_{\mathbb{P}}(p_S(\log p/n + 1/p)^{1/2}).$$

Hence, $\left\| \mathbf{L} \mathbf{A}^{-1} \right\|_\infty \leq 1 - \rho$ holds with probability approaching to 1. Combined with equation (B.5) and the definition of \mathbf{A} , we have

$$\left\| \mathbf{A}((\widehat{\Theta}_1)_S - (\Theta_0)_S) \right\|_\infty = \left\| \nabla_S Z_1(\widehat{\Theta}_1) - \nabla_S Z_1(\Theta_0) \right\|_\infty \leq \left\| \nabla_S Z_1(\widehat{\Theta}_1) \right\|_\infty + \left\| \nabla_S Z_1(\Theta_0) \right\|_\infty \leq \lambda_2 + c\lambda_2.$$

Thus, recall $c < \rho/(2 - \rho)$, we obtain

$$\left\| \nabla_{S^c} Z_1(\widehat{\Theta}_1) \right\|_\infty \leq c\lambda_2 + (1 - \rho)(\lambda_2 + c\lambda_2) < \lambda_2.$$

This implies $\widehat{\Theta}_1$ is the unique global minima of the objective function, and

$$\left\| (\widehat{\Theta}_1)_S - (\Theta_0)_S \right\|_\infty = O_{\mathbb{P}}(p_S(\sqrt{\log p/n} + 1/\sqrt{p})) = o_{\mathbb{P}}(1).$$

Thus Theorem 3.1 is proved.

Step 3. In this step, we proof the proposition 3.2. Note that

$$\mathbb{P}(\text{sign}(\widehat{\boldsymbol{\beta}}^{or}) = \text{sign}(\boldsymbol{\beta}_0)) \geq \mathbb{P}(\mathcal{A}_n \cap \mathcal{B}_n),$$

where

$$\mathcal{A}_n = \left\{ \left| \mathbf{A}^{-1} \frac{1}{\sqrt{n}} \left(\boldsymbol{\Omega}, \widehat{\mathbf{F}}, \widehat{\mathbf{U}}_S \right)^\top \boldsymbol{\epsilon} \right| \leq \sqrt{n} \left(|(\Theta_0)_S| - \frac{\lambda_2}{2} \left| \mathbf{A}^{-1} (\mathbf{0}_{|S^c| \times K}, \mathbf{0}_{|S^c| \times r}, \text{sign}[(\boldsymbol{\beta}_0)_S])^\top \right| \right) \right\}$$

and

$$\mathcal{B}_n = \left\{ \left| \left(\widehat{\mathbf{U}}_{\mathcal{S}^c}^\top \boldsymbol{\Omega}, \mathbf{0}_{|\mathcal{S}^c| \times r}, \widehat{\mathbf{U}}_{\mathcal{S}^c}^\top \widehat{\mathbf{U}}_{\mathcal{S}} \right) \mathbf{A}^{-1} \frac{1}{\sqrt{n}} (\boldsymbol{\Omega}, \widehat{\mathbf{F}}, \widehat{\mathbf{U}}_{\mathcal{S}})^\top \boldsymbol{\epsilon} - \frac{1}{\sqrt{n}} \widehat{\mathbf{U}}_{\mathcal{S}^c}^\top \boldsymbol{\epsilon} \right| \leq \sqrt{n} \lambda_2 \rho \mathbf{1} \right\},$$

where the inequality holds element-wise. We consider the complement of \mathcal{A}_n and \mathcal{B}_n respectively.

$$\mathbb{P}(\mathcal{A}_n^c) \leq \mathbb{P} \left(\bigcup_{i=1}^{p_S+r+K} \left\{ |\mathbf{s}_i^\top \boldsymbol{\epsilon}| \geq \sqrt{n} \left(\min(|(\boldsymbol{\Theta}_0)_{\mathcal{S}}|) - \frac{\lambda_2}{2} \left\| \mathbf{A}^{-1} (\mathbf{0}_{|\mathcal{S}^c| \times K}, \mathbf{0}_{|\mathcal{S}^c| \times r}, \text{sign}[(\boldsymbol{\beta}_0)_{\mathcal{S}}])^\top \right\|_\infty \right) \right\} \right),$$

where \mathbf{s}_j is the j -th row of $\mathbf{A}^{-1}(\boldsymbol{\Omega}, \widehat{\mathbf{F}}, \widehat{\mathbf{U}}_{\mathcal{S}})^\top / \sqrt{n}$. Hence, it's easy to show that $\mathbf{s}_i^\top \mathbf{s}_j = \mathbf{A}^{-1}$. According to (B.12), $\|\mathbf{A}^{-1}\|_2 \leq 2C$. Thus, $\|(\mathbf{s}_i^\top \mathbf{s}_j)\|_2$ is bounded with high probability. There exists a constant M such that

$$\|\mathbf{s}_i\|_2 \leq M \quad \text{for all } i = 1, \dots, k+r+p_S.$$

Similarly, we can prove $\|(\widehat{\mathbf{U}}_{\mathcal{S}}^\top \widehat{\mathbf{U}}_{\mathcal{S}}/n)^{-1}\|_\infty$ has finite upper bound κ .

By concentration inequality of sub-Gaussian random variable, we obtain

$$\begin{aligned} \mathbb{P}(\mathcal{A}_n^c) &\leq \mathbb{P} \left(\bigcup_{i=1}^{p_S+r+K} \left\{ \frac{|\mathbf{s}_i^\top \boldsymbol{\epsilon}|}{\|\mathbf{s}_i\|} \geq \frac{1}{M} \sqrt{n} \left(\min(|(\boldsymbol{\Theta}_0)_{\mathcal{S}}|) - \frac{\lambda_2}{2} \left\| \mathbf{A}^{-1} (\mathbf{0}_{|\mathcal{S}^c| \times K}, \mathbf{0}_{|\mathcal{S}^c| \times r}, \text{sign}[(\boldsymbol{\beta}_0)_{\mathcal{S}}])^\top \right\|_\infty \right) \right\} \right) \\ &\leq \sum_{i=1}^{p_S+r+K} \mathbb{P} \left(\frac{|\mathbf{s}_i^\top \boldsymbol{\epsilon}|}{\|\mathbf{s}_i\|} \geq \frac{1}{M} \sqrt{n} \frac{\lambda_2 \kappa}{2} \right) \leq 2p_S \exp(-B_1 p_S^2 \log p_S) \leq \frac{\widetilde{B}_1}{p_S^{p_S^2-1}}, \end{aligned}$$

where B_1 and \widetilde{B}_1 are two positive values. Denote $\widetilde{\mathbf{s}}_j$ as the j -th row of

$$\widetilde{\mathbf{S}} := (\widehat{\mathbf{U}}_{\mathcal{S}^c}^\top \boldsymbol{\Omega}, \mathbf{0}, \widehat{\mathbf{U}}_{\mathcal{S}^c}^\top \widehat{\mathbf{U}}_{\mathcal{S}}) \mathbf{A}^{-1} (\boldsymbol{\Omega}, \widehat{\mathbf{F}}, \widehat{\mathbf{U}}_{\mathcal{S}})^\top / \sqrt{n} - \widehat{\mathbf{U}}_{\mathcal{S}^c}^\top / \sqrt{n}.$$

Note that $\widetilde{\mathbf{S}} \widetilde{\mathbf{S}}^\top = \widehat{\mathbf{U}}_{\mathcal{S}^c}^\top \mathbf{P} \widehat{\mathbf{U}}_{\mathcal{S}^c} / n$, where \mathbf{P} is a projection matrix. Hence, we obtain that $\|\widetilde{\mathbf{s}}_j\|_2 \leq \widetilde{M}$ for all $j \in \mathcal{S}^c$. Further, we have

$$\mathbb{P}(\mathcal{B}_n^c) \leq \sum_{i=1}^{p-p_S} \mathbb{P} \left(\frac{|\widetilde{\mathbf{s}}_i^\top \boldsymbol{\epsilon}|}{\|\widetilde{\mathbf{s}}_i\|} \geq \frac{1}{\widetilde{M}} \sqrt{n} \lambda_2 \rho \right) \leq 2(p-p_S) \exp(-B_2 p_S^2 \log p_S) \leq \frac{\widetilde{B}_2 (p-p_S)}{p_S^{p_S^2}},$$

where B_2 and \widetilde{B}_2 are two positive values. Therefore, the estimators has sign consistency, i.e. $\mathbb{P}(\text{sign}(\widehat{\boldsymbol{\beta}}) = \text{sign}(\boldsymbol{\beta}_0)) \rightarrow 1$. □

B.3. Proof of Theorem 3.3

Proof. Before starting the proof, we define a function to facilitate the subsequent proof process. For any vector $\alpha \in \mathcal{V}_{\mathcal{G}}$, where $\mathcal{V}_{\mathcal{G}} = \{\alpha, \alpha = \mathbf{\Omega}\gamma, \gamma \in \mathbb{R}^K\}$, $T(\alpha)$ is a K -dimensional vector, i.e.,

$$T : \mathcal{V}_{\mathcal{G}} \rightarrow \mathbb{R}^K.$$

Specifically, the k -th coordinate of $T(\alpha)$ is the common value of α_i for $i \in \mathcal{G}_k$. According to this definition of T , $T^{-1}(\gamma)$ represents the n -dimensional vector obtained by restoring γ to the corresponding the true grouping structure.

With a slight abuse of notations, we define $\Theta = (\alpha^\top, \gamma^\top, \theta^\top, \beta^\top)^\top$, the corresponding oracle estimator can be defined as $\widehat{\Theta}^{or} = ((\widehat{\alpha}^{or})^\top, (\widehat{\gamma}^{or})^\top, (\widehat{\theta}^{or})^\top, (\widehat{\beta}^{or})^\top)^\top$ and

$$Z(\Theta) = 1/2n\|\mathbf{Y} - \alpha - \widehat{\mathbf{F}}\theta - \widehat{\mathbf{U}}\beta\|_2^2 + \lambda_1 \sum_{i=1}^n \min\{d(\alpha_i, \gamma_1), \dots, d(\alpha_i, \gamma_K)\} + \lambda_2 \|\beta\|_1. \quad (\text{B.22})$$

We choose an open set of Θ related to t_n , denoted as

$$\mathcal{N}_n = \left\{ \Theta : \|\Theta - \widehat{\Theta}^{or}\|_\infty < t_n \right\},$$

where t_n is a real value sequence satisfying $t_n = o(r_n)$ and hence we have $t_n = o(p_S(\sqrt{\log p/n} + 1/\sqrt{p}))$.

In the neighborhood of $\widehat{\Theta}^{or}$, we show that $\widehat{\Theta}^{or}$ is the local minima of Z with high probability in the following **two steps**. The first step is to prove that for any parameter Θ belonging to \mathcal{N}_n satisfy $Z(\widehat{\alpha}^{or}, \widehat{\gamma}^{or}, \widehat{\theta}^{or}, \widehat{\beta}^{or}) < Z(T^{-1}(\gamma), \gamma, \theta, \beta)$. In the second step, we show that $Z(T^{-1}(\gamma), \gamma, \theta, \beta) \leq Z(\alpha, \gamma, \theta, \beta)$ holds on \mathcal{N}_n . Hence, $\widehat{\Theta}^{or}$ is the local minima of the objective function.

Step 1. In this step, we focus on show that $Z(\widehat{\alpha}^{or}, \widehat{\gamma}^{or}, \widehat{\theta}^{or}, \widehat{\beta}^{or}) < Z(T^{-1}(\gamma), \gamma, \theta, \beta)$ holds for any parameter $\Theta \in \mathcal{N}_n$.

As $\widehat{\Theta}^{or}$ is the oracle estimator, taking it into (B.22), we have

$$\sum_{i=1}^n \min\{d(\widehat{\alpha}_i^{or}, \widehat{\gamma}_1^{or}), \dots, d(\widehat{\alpha}_i^{or}, \widehat{\gamma}_K^{or})\} = 0.$$

Hence, the oracle estimator of (B.22) is,

$$Z(\widehat{\alpha}^{or}, \widehat{\gamma}^{or}, \widehat{\theta}^{or}, \widehat{\beta}^{or}) = 1/2n\|\mathbf{Y} - \widehat{\alpha}^{or} - \widehat{\mathbf{F}}\widehat{\theta}^{or} - \widehat{\mathbf{U}}\widehat{\beta}^{or}\|_2^2 + \lambda_2 \|\widehat{\beta}^{or}\|_1.$$

Similarly, by taking $(T^{-1}(\gamma), \gamma, \theta, \beta)^\top$ into the objective function and using the property of $T^{-1}(\gamma)$, we have

$$Z(T^{-1}(\gamma), \gamma, \theta, \beta) = 1/2n\|\mathbf{Y} - T^{-1}(\gamma) - \widehat{\mathbf{F}}\theta - \widehat{\mathbf{U}}\beta\|_2^2 + \lambda_2 \|\beta\|_1.$$

As $\widehat{\boldsymbol{\alpha}}^{or}$ and $T^{-1}(\boldsymbol{\gamma})$ has the same grouping structure as the $\boldsymbol{\alpha}_0$, we denote them as $\boldsymbol{\Omega}\widehat{\boldsymbol{\gamma}}^{or}$ and $\boldsymbol{\Omega}\boldsymbol{\gamma}$. The objective function can be rewritten as

$$Z(\widehat{\boldsymbol{\alpha}}^{or}, \widehat{\boldsymbol{\gamma}}^{or}, \widehat{\boldsymbol{\theta}}^{or}, \widehat{\boldsymbol{\beta}}^{or}) = \frac{1}{2n} \|\mathbf{Y} - \boldsymbol{\Omega}\widehat{\boldsymbol{\gamma}}^{or} - \widehat{\mathbf{F}}\widehat{\boldsymbol{\theta}}^{or} - \widehat{\mathbf{U}}\widehat{\boldsymbol{\beta}}^{or}\|_2^2 + \lambda_2 \|\widehat{\boldsymbol{\beta}}^{or}\|_1$$

and

$$Z(T^{-1}(\boldsymbol{\gamma}), \boldsymbol{\gamma}, \boldsymbol{\theta}, \boldsymbol{\beta}) = \frac{1}{2n} \|\mathbf{Y} - \boldsymbol{\Omega}\boldsymbol{\gamma} - \widehat{\mathbf{F}}\boldsymbol{\theta} - \widehat{\mathbf{U}}\boldsymbol{\beta}\|_2^2 + \lambda_2 \|\boldsymbol{\beta}\|_1.$$

By the definition of the oracle estimator, we get

$$Z(\widehat{\boldsymbol{\alpha}}^{or}, \widehat{\boldsymbol{\gamma}}^{or}, \widehat{\boldsymbol{\theta}}^{or}, \widehat{\boldsymbol{\beta}}^{or}) < Z(T^{-1}(\boldsymbol{\gamma}), \boldsymbol{\gamma}, \boldsymbol{\theta}, \boldsymbol{\beta}).$$

Step 2. In this step, we prove that $Z(T^{-1}(\boldsymbol{\gamma}), \boldsymbol{\gamma}, \boldsymbol{\theta}, \boldsymbol{\beta}) \leq Z(\boldsymbol{\alpha}, \boldsymbol{\gamma}, \boldsymbol{\theta}, \boldsymbol{\beta})$ holds for any $(\boldsymbol{\alpha}, \boldsymbol{\gamma}, \boldsymbol{\theta}, \boldsymbol{\beta})^\top \in \mathcal{N}_n$.

If sample i belongs to the k -th subgroup, we have

$$|\alpha_i - \gamma_k| \leq |\alpha_i - \widehat{\alpha}_i^{or}| + |\widehat{\gamma}_k^{or} - \gamma_k| \leq 2t_n.$$

If sample i does not belong to the k -th subgroup, by the Lemma A.7, we have

$$|\alpha_i - \gamma_k| = |\widehat{\alpha}_i^{or} - \widehat{\gamma}_k^{or}| - |(\widehat{\alpha}_i^{or} - \widehat{\gamma}_k^{or}) - (\alpha_i - \gamma_k)| \geq r_n/2 - o(r_n) \geq 2t_n$$

holds with probability approaching to 1. Therefore, for a given α_i , we define $\gamma_{K(i)}$ as

$$\min_{k=1, \dots, K} |\alpha_i - \gamma_k| = |\alpha_i - \gamma_{K(i)}|,$$

and we account the i -th sample belongs to $K(i)$ -th group. Further, the objective function in the (B.22) can be written as

$$Z(\boldsymbol{\alpha}, \boldsymbol{\gamma}, \boldsymbol{\theta}, \boldsymbol{\beta}) = \frac{1}{2n} \|\mathbf{Y} - \boldsymbol{\alpha} - \widehat{\mathbf{F}}\boldsymbol{\theta} - \widehat{\mathbf{U}}\boldsymbol{\beta}\|_2^2 + \lambda_1 \sum_{i=1}^n |\alpha_i - \gamma_{K(i)}| + \lambda_2 \|\boldsymbol{\beta}\|_1.$$

By Lagrange mean value theorem, we have:

$$\begin{aligned} & Z(\boldsymbol{\alpha}, \boldsymbol{\gamma}, \boldsymbol{\theta}, \boldsymbol{\beta}) - Z(T^{-1}(\boldsymbol{\gamma}), \boldsymbol{\gamma}, \boldsymbol{\theta}, \boldsymbol{\beta}) \\ &= \lambda_1 \sum_{i=1}^n |\alpha_i - \gamma_{K(i)}| - \frac{1}{n} (\mathbf{Y} - \boldsymbol{\alpha}^m - \widehat{\mathbf{F}}\boldsymbol{\theta} - \widehat{\mathbf{U}}\boldsymbol{\beta})^\top (\boldsymbol{\alpha} - T^{-1}(\boldsymbol{\gamma})) \\ &= \sum_{i=1}^n \left[\lambda_1 |\alpha_i - \gamma_{K(i)}| - 1/n (Y_i - \alpha_i^m - \widehat{\mathbf{f}}_i^\top \boldsymbol{\theta} - \widehat{\mathbf{u}}_i^\top \boldsymbol{\beta}) (\alpha_i - \gamma_{K(i)}) \right]. \end{aligned} \tag{B.23}$$

where $\boldsymbol{\alpha}^m = \varsigma \boldsymbol{\alpha} + (1 - \varsigma)T^{-1}(\boldsymbol{\gamma})$, $\varsigma \in [-1, 1]$. Thus, it is sufficient to show

$$\sum_{i=1}^n \left(\lambda_1 - \frac{1}{n} |Y_i - \alpha_i^m - \hat{\mathbf{f}}_i^\top \boldsymbol{\theta} - \hat{\mathbf{u}}_i^\top \boldsymbol{\beta}| \right) |\alpha_i - \gamma_{K(i)}| \geq 0.$$

Next, we aim to show that each term of the summation is positive. By triangular inequality and the property of $\hat{\boldsymbol{\beta}}^{or}$, we have:

$$\begin{aligned} \|\mathbf{Y} - \boldsymbol{\alpha}^m - \hat{\mathbf{F}}\boldsymbol{\theta} - \hat{\mathbf{U}}\boldsymbol{\beta}\|_\infty &\leq \|\boldsymbol{\epsilon}\|_\infty + \|\boldsymbol{\alpha}^m - \boldsymbol{\alpha}_0\|_\infty + \|\hat{\mathbf{F}}\|_\infty \|\boldsymbol{\theta} - \hat{\mathbf{B}}^\top \boldsymbol{\beta}_0\|_\infty + \|\hat{\mathbf{U}}(\boldsymbol{\beta} - \boldsymbol{\beta}_0)\|_\infty \\ &\leq \|\boldsymbol{\epsilon}\|_\infty + \|\boldsymbol{\alpha}^m - \boldsymbol{\alpha}_0\|_\infty + \|\hat{\mathbf{F}}\|_\infty \|\boldsymbol{\theta} - \hat{\mathbf{B}}^\top \boldsymbol{\beta}_0\|_\infty \\ &\quad + \|\hat{\mathbf{U}}\|_\infty \|(\boldsymbol{\beta} - \hat{\boldsymbol{\beta}}^{or})\|_\infty + \|\hat{\mathbf{U}}_S\|_\infty \|(\hat{\boldsymbol{\beta}}_S^{or} - (\boldsymbol{\beta}_0)_S)\|_\infty. \end{aligned}$$

Note that there exist a sufficient large C_1 such that

$$\mathbb{P}(\|\boldsymbol{\epsilon}\|_\infty \geq C_1 \sqrt{\log n}) \leq 2 \sum_{i=1}^n \exp(-C_2 \log n) \leq 2/n^{C_2-1},$$

which implies that $\|\boldsymbol{\epsilon}\|_\infty = O_{\mathbb{P}}(\sqrt{\log n})$. By triangle inequality and Theorem 3.1, we have

$$\begin{aligned} \|\boldsymbol{\alpha}^m - \boldsymbol{\alpha}_0\|_\infty &\leq \|\boldsymbol{\alpha}^m - \hat{\boldsymbol{\alpha}}^{or}\|_\infty + \|\hat{\boldsymbol{\alpha}}^{or} - \boldsymbol{\alpha}_0\|_\infty \\ &\leq t_n + O_{\mathbb{P}}(p_S(\sqrt{\log p/n} + 1/\sqrt{p})) = O_{\mathbb{P}}(p_S(\sqrt{\log p/n} + 1/\sqrt{p})). \end{aligned}$$

Similarly, we have

$$\|\boldsymbol{\theta} - \hat{\mathbf{B}}^\top \boldsymbol{\beta}_0\|_\infty \leq \|\boldsymbol{\theta} - \hat{\boldsymbol{\theta}}^{or}\|_\infty + \|\hat{\boldsymbol{\theta}}^{or} - \hat{\mathbf{B}}^\top \boldsymbol{\beta}_0\|_\infty.$$

Note that

$$\|\hat{\boldsymbol{\theta}}_1^{or} - \hat{\mathbf{B}}^\top \boldsymbol{\beta}_0\|_\infty \leq \|\hat{\boldsymbol{\theta}}_1^{or} - \mathbf{H}\boldsymbol{\theta}_0\|_\infty + \|(\mathbf{H}\mathbf{B}^\top - \hat{\mathbf{B}}^\top)\boldsymbol{\beta}_0\|_\infty = O_{\mathbb{P}}(p_S(\sqrt{\log p/n} + 1/\sqrt{p})),$$

thus $\|\boldsymbol{\theta} - \hat{\mathbf{B}}^\top \boldsymbol{\beta}_0\|_\infty = O_{\mathbb{P}}(p_S(\sqrt{\log p/n} + 1/\sqrt{p}))$. By the concentration inequality and Lemma A.1, we obtain $\|\hat{\mathbf{F}}\|_\infty = O_{\mathbb{P}}(\sqrt{n})$. By norm inequality, we have,

$$\begin{aligned} \|\hat{\mathbf{U}}\|_\infty &\leq \|\mathbf{U}\|_\infty + \|\hat{\mathbf{U}} - \mathbf{U}\|_\infty \\ &\leq \sqrt{p}\|\mathbf{U}\|_2 + \sqrt{p}\|\hat{\mathbf{U}} - \mathbf{U}\|_2 \\ &\leq p(\|\mathbf{U}^\top \mathbf{U}\|_{\max})^{1/2} + p(\|(\hat{\mathbf{U}} - \mathbf{U})^\top (\hat{\mathbf{U}} - \mathbf{U})\|_{\max})^{1/2}. \end{aligned} \tag{B.24}$$

By the proposed assumption and Lemma A.6, we have

$$\|\mathbf{U}^\top \mathbf{U}\|_{\max} \leq n\|\boldsymbol{\Sigma}\|_{\max} + \|\mathbf{U}^\top \mathbf{U} - n\boldsymbol{\Sigma}\|_{\max} = O_{\mathbb{P}}(n) + O_{\mathbb{P}}(\sqrt{n \log p}) = O_{\mathbb{P}}(n).$$

Thus, $\|\widehat{\mathbf{U}}\|_\infty = O_{\mathbb{P}}(p\sqrt{n})$ and similarly $\|\widehat{\mathbf{U}}_{\mathcal{S}}\|_\infty = O_{\mathbb{P}}(p_{\mathcal{S}}\sqrt{n})$. Recall that $t_n = o(p_{\mathcal{S}}(\sqrt{\log p/n} + 1/\sqrt{p}))$, we obtain that

$$\|\mathbf{Y} - \boldsymbol{\alpha}^m - \widehat{\mathbf{F}}\boldsymbol{\theta} - \widehat{\mathbf{U}}\boldsymbol{\beta}\|_\infty = O_{\mathbb{P}}(\sqrt{\log n} + p_{\mathcal{S}}^2(\sqrt{\log p} + \sqrt{n}/\sqrt{p}) + p\sqrt{nt_n}).$$

By setting t_n to satisfy $pt_n = o(p_{\mathcal{S}}^2(\sqrt{\log p/n} + 1/\sqrt{p}))$, we have

$$\|\mathbf{Y} - \boldsymbol{\alpha}^m - \widehat{\mathbf{F}}\boldsymbol{\theta} - \widehat{\mathbf{U}}\boldsymbol{\beta}\|_\infty = O_{\mathbb{P}}(\sqrt{\log n} + p_{\mathcal{S}}^2(\sqrt{\log p} + \sqrt{n}/\sqrt{p})).$$

If λ_1 satisfy $\lambda_1 \gg n^{-1}\max(\sqrt{\log n}, p_{\mathcal{S}}^2(\sqrt{\log p} + \sqrt{n}/\sqrt{p}))$, then

$$\sum_{i=1}^n (\lambda_1 - \frac{1}{n}(Y_i - \alpha_i^m - \widehat{\mathbf{f}}_i^\top \boldsymbol{\theta} - \widehat{\mathbf{u}}_i^\top \boldsymbol{\beta})) |\alpha_i - \gamma_{K(i)}| \geq 0,$$

with high probability, i.e., $Z(\boldsymbol{\alpha}, \boldsymbol{\gamma}, \boldsymbol{\theta}, \boldsymbol{\beta}) - Z(T^{-1}(\boldsymbol{\gamma}), \boldsymbol{\gamma}, \boldsymbol{\theta}, \boldsymbol{\beta}) \geq 0$ is proved.

Combining with the inequality from the first step, we get

$$Z(\boldsymbol{\alpha}, \boldsymbol{\gamma}, \boldsymbol{\theta}, \boldsymbol{\beta}) - Z(\widehat{\boldsymbol{\alpha}}^{or}, \widehat{\boldsymbol{\gamma}}^{or}, \widehat{\boldsymbol{\theta}}^{or}, \widehat{\boldsymbol{\beta}}^{or}) \geq 0 \quad \text{for all } \boldsymbol{\Theta} \in \mathcal{N}_n,$$

which implies the $\widehat{\boldsymbol{\Theta}}^{or}$ is a local minima of $Z(\boldsymbol{\Theta})$.

□

C. Proof of Technical Lemmas

C.1. Proof of Lemma A.5

Proof. From Lemma A.1, we know \mathbf{H} is invertible with the probability approaching to 1 and $\|\mathbf{H}^{-1}\|_2 = O_{\mathbb{P}}(1)$, where the inverse matrix of \mathbf{H} in sense of probability denoted as \mathbf{H}^{-1} . Let \mathbf{e}_j be the unit vector whose j -th element is 1. According to the estimation process, it is clear that $\widehat{\mathbf{F}}^\top \widehat{\mathbf{U}} = 0$. Combined with the triangle inequality, we have

$$\begin{aligned} \|\widehat{\mathbf{U}}^\top \mathbf{F}\mathbf{a}\|_\infty &= \|\widehat{\mathbf{U}}^\top (\mathbf{F} - \widehat{\mathbf{F}}\mathbf{H}^{-1})\mathbf{a}\|_\infty \\ &\leq \|(\widehat{\mathbf{U}}^\top - \mathbf{U}^\top)(\mathbf{F} - \widehat{\mathbf{F}}\mathbf{H}^{-1})\mathbf{a}\|_\infty + \|\mathbf{U}^\top (\mathbf{F} - \widehat{\mathbf{F}}\mathbf{H}^{-1})\mathbf{a}\|_\infty. \end{aligned} \tag{C.1}$$

By the Cauchy-Schwarz inequality and Lemma A.1, we have

$$\begin{aligned}
\|(\widehat{\mathbf{U}}^\top - \mathbf{U}^\top)(\mathbf{F} - \widehat{\mathbf{F}}\mathbf{H}^{-1})\mathbf{a}\|_\infty &= \max_{j \in \{1, \dots, p\}} |e_j^\top (\widehat{\mathbf{U}}^\top - \mathbf{U}^\top)(\mathbf{F} - \widehat{\mathbf{F}}\mathbf{H}^{-1})\mathbf{a}| \\
&\leq \left(\max_{j \in \{1, \dots, p\}} \sum_{i=1}^n (\widehat{u}_{ij} - u_{ij})^2 \right)^{1/2} \|\mathbf{F}\mathbf{H}^\top - \widehat{\mathbf{F}}\|_{\mathbb{F}} \|\mathbf{H}^{-1}\mathbf{a}\|_2 \\
&= O_{\mathbb{P}}\left(\left[(\log p + \frac{n}{p}) \left(1 + \frac{n}{p}\right) \right]^{1/2} \right).
\end{aligned} \tag{C.2}$$

As \mathbf{H} is invertible with probability tending to 1,

$$\|\mathbf{U}^\top(\mathbf{F} - \widehat{\mathbf{F}}\mathbf{H}^{-1})\mathbf{a}\|_\infty = \|\mathbf{U}^\top(\mathbf{F}\mathbf{H}^\top - \widehat{\mathbf{F}})\mathbf{H}^{-1}\mathbf{a}\|_\infty. \tag{C.3}$$

Recall the definition of \mathbf{H} in Theorem 3.1, we have

$$\widehat{\mathbf{F}} - \mathbf{F}\mathbf{H}^\top = \frac{1}{n}\mathbf{U}\mathbf{U}^\top\widehat{\mathbf{F}}\mathbf{V}^{-1} + \frac{1}{n}\mathbf{U}\mathbf{B}\mathbf{F}^\top\widehat{\mathbf{F}}\mathbf{V}^{-1} + \frac{1}{n}\mathbf{F}\mathbf{B}^\top\mathbf{U}^\top\widehat{\mathbf{F}}\mathbf{V}^{-1}.$$

Plug it back into (C.3), we have

$$\begin{aligned}
\|\mathbf{U}^\top(\mathbf{F} - \widehat{\mathbf{F}}\mathbf{H}^{-1})\mathbf{a}\|_\infty &\leq \underbrace{\frac{1}{n}\|\mathbf{U}^\top\mathbf{U}\mathbf{U}^\top\widehat{\mathbf{F}}\mathbf{V}^{-1}\mathbf{H}^{-1}\mathbf{a}\|_\infty}_{S_1} + \underbrace{\frac{1}{n}\|\mathbf{U}^\top\mathbf{U}\mathbf{B}\mathbf{F}^\top\widehat{\mathbf{F}}\mathbf{V}^{-1}\mathbf{H}^{-1}\mathbf{a}\|_\infty}_{S_2} \\
&\quad + \underbrace{\frac{1}{n}\|\mathbf{U}^\top\mathbf{F}\mathbf{B}^\top\mathbf{U}^\top\widehat{\mathbf{F}}\mathbf{V}^{-1}\mathbf{H}^{-1}\mathbf{a}\|_\infty}_{S_3}.
\end{aligned} \tag{C.4}$$

Next, we will bound each term separately. For S_1 , according to the triangle inequality, we have

$$S_1 \leq \underbrace{\frac{1}{n}\|\mathbf{U}^\top\mathbf{U}\mathbf{U}^\top\mathbf{F}\mathbf{H}^\top\mathbf{V}^{-1}\mathbf{H}^{-1}\mathbf{a}\|_\infty}_{S_{11}} + \underbrace{\frac{1}{n}\|\mathbf{U}^\top\mathbf{U}\mathbf{U}^\top(\widehat{\mathbf{F}} - \mathbf{F}\mathbf{H}^\top)\mathbf{V}^{-1}\mathbf{H}^{-1}\mathbf{a}\|_\infty}_{S_{12}}. \tag{C.5}$$

For S_{11} , according to the definition of the infinity norm of the matrix, and using the Cauchy-Schwarz inequality, we have

$$S_{11} = \frac{1}{n} \max_{j \in \{1, \dots, p\}} |e_j^\top \mathbf{U}^\top \mathbf{U} \mathbf{U}^\top \mathbf{F} \mathbf{H}^\top \mathbf{V}^{-1} \mathbf{H}^{-1} \mathbf{a}| \leq \frac{1}{n} \max_{j \in \{1, \dots, p\}} \|\mathbf{e}_j^\top \mathbf{U}^\top \mathbf{U} \mathbf{U}^\top \mathbf{F}\|_2 \|\mathbf{V}^{-1} \mathbf{H}^{-1}\|_2.$$

Note that

$$e_j^\top \mathbf{U}^\top \mathbf{U} \mathbf{U}^\top \mathbf{F} = \left(\sum_{t=1}^n u_{tj} \sum_{s=1}^n \mathbf{u}_t^\top \mathbf{u}_s f_{s1}, \dots, \sum_{t=1}^n u_{tj} \sum_{s=1}^n \mathbf{u}_t^\top \mathbf{u}_s f_{sr} \right) = \sum_{t=1}^n u_{tj} \sum_{s=1}^n \mathbf{u}_t^\top \mathbf{u}_s \mathbf{f}_s^\top.$$

After centralization, according to the triangle inequality and the Cauchy-Schwarz inequality, we have,

$$S_{11} \leq \frac{1}{n} \max_{j \in \{1, \dots, p\}} \left(\sum_{t=1}^n u_{tj}^2 \right)^{1/2} \left[\left(\sum_{t=1}^n \left\| \sum_{s=1}^n (\mathbf{u}_t^\top \mathbf{u}_s - \mathbb{E}(\mathbf{u}_t^\top \mathbf{u}_s)) \mathbf{f}_s \right\|_2^2 \right)^{1/2} + \left(\sum_{t=1}^n \left\| \sum_{s=1}^n \mathbb{E}(\mathbf{u}_t^\top \mathbf{u}_s) \mathbf{f}_s \right\|_2^2 \right)^{1/2} \right] \|\mathbf{V}^{-1}\|_2 \|\mathbf{H}^{-1}\|_2. \quad (\text{C.6})$$

By the similar technique used in the proof of Theorem 3.1, we have

$$\mathbb{P} \left(\frac{1}{n} \left| \sum_{i=1}^n u_{ij}^2 - \sum_{i=1}^n \mathbb{E} u_{ij}^2 \right| \geq t \right) \leq n \exp \left(-\frac{(nt)^l}{V_1} \right) + \exp \left(-\frac{(nt)^2}{nV_2} \right) + \exp \left(-\frac{(nt)^2}{nV_3} \exp \left(-\frac{(nt)^{l(1-l)}}{V_4 (\log nt)^l} \right) \right),$$

which implies $\max_{j \in \{1, \dots, p\}} (\sum_{t=1}^n u_{tj}^2) = O_{\mathbb{P}}(n + \sqrt{n \log p}) = O_{\mathbb{P}}(n)$. Combined with similar arguments of Lemma 8 in Fan et al. (2013), we have

$$\sum_{t=1}^n \left\| \sum_{s=1}^n (\mathbf{u}_t^\top \mathbf{u}_s - \mathbb{E}(\mathbf{u}_t^\top \mathbf{u}_s)) \mathbf{f}_s \right\|_2^2 = O_{\mathbb{P}}(n^3 p) \quad \sum_{t=1}^n \left\| \sum_{s=1}^n \mathbb{E}(\mathbf{u}_t^\top \mathbf{u}_s) \mathbf{f}_s \right\|_2^2 = O_{\mathbb{P}}(n^2 p^2).$$

According to the Lemma 5 in the Fan et al. (2013), it's obvious that $\|\mathbf{V}^{-1}\|_2 = O_{\mathbb{P}}(1/p)$. According to the decomposition of S_{11} in equation (C.6) and above results, we obtain

$$S_{11} = O_{\mathbb{P}} \left(\sqrt{n} + \frac{n}{\sqrt{p}} \right).$$

With similar arguments, we have $S_{12} = O_{\mathbb{P}}(n/p + \sqrt{n}/\sqrt{p})$. Therefore, reference to the decomposition of S_1 in (C.5), we get

$$S_1 = O_{\mathbb{P}} \left(\sqrt{n} + \frac{n}{\sqrt{p}} \right). \quad (\text{C.7})$$

We apply Cauchy-Schwarz inequality to bound S_2 and consider the definition of the infinite norm, the following inequality holds

$$\begin{aligned} S_2 &= \frac{1}{n} \max_{j \in \{1, \dots, p\}} \left| \mathbf{e}_j^\top \mathbf{U}^\top \mathbf{U} \mathbf{B} \mathbf{F}^\top \widehat{\mathbf{F}} \mathbf{V}^{-1} \widetilde{\mathbf{H}}^\top \mathbf{a} \right| \\ &= \frac{1}{n} \max_{j \in \{1, \dots, p\}} \left| \sum_{i=1}^n u_{ij} \mathbf{u}_i^\top \mathbf{B} \sum_{s=1}^n \mathbf{f}_s \widehat{\mathbf{f}}_s^\top \mathbf{V}^{-1} \widetilde{\mathbf{H}}^\top \mathbf{a} \right| \\ &\leq \frac{1}{n} \max_{j \in \{1, \dots, p\}} \left\| \sum_{i=1}^n u_{ij} \mathbf{u}_i^\top \mathbf{B} \right\|_2 \left\| \sum_{s=1}^n \mathbf{f}_s \widehat{\mathbf{f}}_s^\top \right\|_2 \left\| \mathbf{V}^{-1} \widetilde{\mathbf{H}}^\top \right\|_2. \end{aligned}$$

By simple triangle inequality, we have

$$\max_{j \in \{1, \dots, p\}} \left\| \sum_{i=1}^n u_{ij} \mathbf{u}_i^\top \mathbf{B} \right\|_2 \leq \max_{j \in \{1, \dots, p\}} \left[\left\| \sum_{i=1}^n \mathbb{E} (u_{ij} \mathbf{u}_i^\top \mathbf{B}) \right\|_2 + \left\| \sum_{i=1}^n (u_{ij} \mathbf{u}_i^\top \mathbf{B} - \mathbb{E} (u_{ij} \mathbf{u}_i^\top \mathbf{B})) \right\|_2 \right].$$

By the distribution assumption of \mathbf{u}_i , we know the $\mathbf{b}_k^\top \mathbf{u}_i u_{ij} / \|\mathbf{b}_k\|_2$ also has the exponential-type tail, where \mathbf{b}_s is the s -th column of the matrix \mathbf{B} . Applying the concentration inequality in Merlevède et al. (2011) again, we derive

$$\max_{j \in \{1, \dots, p\}} \left\| \sum_{i=1}^n (\mathbf{b}_k^\top \mathbf{u}_i u_{ij} - \mathbb{E} \mathbf{b}_k^\top \mathbf{u}_i u_{ij}) \right\|_2 = O_{\mathbb{P}} \left(\|\mathbf{b}_k\|_2 \sqrt{n \log p} \right) = O_{\mathbb{P}} \left(\sqrt{np \log p} \right).$$

As $\mathbf{B} \in \mathbb{R}^{p \times r}$ and r is finite, thus

$$\max_{j \in \{1, \dots, p\}} \left\| \sum_{i=1}^n (u_{ij} \mathbf{u}_i^\top \mathbf{B} - \mathbb{E} (u_{ij} \mathbf{u}_i^\top \mathbf{B})) \right\|_2 = O_{\mathbb{P}} \left(\sqrt{np \log p} \right)$$

and

$$\max_{j \in \{1, \dots, p\}} \left\| \sum_{i=1}^n u_{ij} \mathbf{u}_i^\top \mathbf{B} \right\|_2 = O_{\mathbb{P}} \left(n + \sqrt{np \log p} \right). \quad (\text{C.8})$$

Combining these results, we finally obtain that

$$S_2 = O_{\mathbb{P}} \left(1 + \sqrt{p \log p/n} \right). \quad (\text{C.9})$$

Recall the definition of S_3 in (C.4), an upper bound of it is given by

$$S_3 \leq \max_{j \in \{1, \dots, p\}} 1/n \|\mathbf{e}_j^\top \mathbf{U}^\top \mathbf{F}\|_2 \|\mathbf{B}^\top \mathbf{U}^\top\|_{\mathbb{F}} \|\widehat{\mathbf{F}}\|_2 \|\mathbf{V}^{-1} \mathbf{H}^{-1}\|_2.$$

According to the results in Fan et al. (2013), we have $\max_{j \in \{1, \dots, p\}} \|\mathbf{e}_j^\top \mathbf{U}^\top \mathbf{F}\|_2 = O_{\mathbb{P}}(\sqrt{n \log p})$ and $\|\mathbf{V}^{-1} \mathbf{H}^{-1}\|_2 = O(1/\sqrt{p})$.

Since $\mathbb{E} \|\mathbf{B}^\top \mathbf{U}^\top\|_{\mathbb{F}}^2 = n \text{tr}(\mathbf{B}^\top \boldsymbol{\Sigma} \mathbf{B}) = O_{\mathbb{P}}(np)$, we conclude that $S_3 = O_{\mathbb{P}}(\sqrt{n \log p/p})$. Combining the equation (C.4), (C.5) and (C.9), we have $\|\mathbf{U}^\top (\mathbf{F} - \widehat{\mathbf{F}} \mathbf{H}^{-1}) \mathbf{a}\|_{\infty} = O_{\mathbb{P}}(\sqrt{n} + n/\sqrt{p})$. Recall the decomposition in (C.1) and results in (C.2), the final result is as follow

$$\|\widehat{\mathbf{U}}^\top \mathbf{F} \mathbf{a}\|_{\infty} = O_{\mathbb{P}}(\sqrt{n} + n/\sqrt{p}).$$

□

C.2. Proof of Lemma A.6

Proof. By the triangle inequality, we have

$$\begin{aligned} \|\widehat{\mathbf{U}}^\top \widehat{\mathbf{U}} - \mathbf{U}^\top \mathbf{U}\|_{\max} &= \left\| \widehat{\mathbf{U}}^\top (\widehat{\mathbf{U}} - \mathbf{U}) + (\widehat{\mathbf{U}} - \mathbf{U})^\top \widehat{\mathbf{U}} - (\widehat{\mathbf{U}} - \mathbf{U})^\top (\widehat{\mathbf{U}} - \mathbf{U}) \right\|_{\max} \\ &\leq 2 \left\| \widehat{\mathbf{U}}^\top (\widehat{\mathbf{U}} - \mathbf{U}) \right\|_{\max} + \left\| (\widehat{\mathbf{U}} - \mathbf{U})^\top (\widehat{\mathbf{U}} - \mathbf{U}) \right\|_{\max}. \end{aligned}$$

According to the definition of factor models and the estimation process, it is clear that

$$\mathbf{X} = \mathbf{F}\mathbf{B}^\top + \mathbf{U} = \widehat{\mathbf{F}}\widehat{\mathbf{B}}^\top + \widehat{\mathbf{U}} \quad \text{and} \quad \widehat{\mathbf{U}}^\top \widehat{\mathbf{F}} = \mathbf{0}.$$

Combining the result of Lemma A.6, it is not hard to verify that

$$\begin{aligned} \|\widehat{\mathbf{U}}^\top (\widehat{\mathbf{U}} - \mathbf{U})\|_{\max} &= \|\widehat{\mathbf{U}}^\top (\mathbf{F}\mathbf{B}^\top - \widehat{\mathbf{F}}\widehat{\mathbf{B}}^\top)\|_{\max} = \|\widehat{\mathbf{U}}^\top \mathbf{F}\mathbf{B}^\top\|_{\max} \\ &= \max_{j \in \{1, \dots, p\}} \|\widehat{\mathbf{U}}^\top \mathbf{F}\tilde{\mathbf{b}}_j\|_{\infty} = O_{\mathbb{P}}(\sqrt{n} + n/\sqrt{p}), \end{aligned} \tag{C.10}$$

where $\tilde{\mathbf{b}}_j^\top$ is the j -th row of the matrix \mathbf{B} . Applying the Lemma A.1, we have

$$\|(\widehat{\mathbf{U}} - \mathbf{U})^\top (\widehat{\mathbf{U}} - \mathbf{U})\|_{\max} \leq \max_{j \in \{1, \dots, p\}} \sum_{i=1}^n (\hat{u}_{ij} - u_{ij})^2 = O_{\mathbb{P}}(\log p + n/p). \tag{C.11}$$

Hence, combining the results of (C.10) and (C.11),

$$\|\widehat{\mathbf{U}}^\top \widehat{\mathbf{U}} - \mathbf{U}^\top \mathbf{U}\|_{\max} = O_{\mathbb{P}}\left(\log p + \sqrt{n} + \frac{n}{\sqrt{p}}\right).$$

□

C.3. Proof of Lemma A.7

Proof. The gap of the oracle estimators can be bounded from below by

$$\min_{i,k} |\hat{\alpha}_i^{or} - \hat{\gamma}_k^{or}| \geq \min_{i,k} |\alpha_{0i} - \gamma_{0k}| - \max_{i,k} |(\hat{\alpha}_i^{or} - \hat{\gamma}_k^{or}) - (\alpha_{0i} - \gamma_{0k})|. \tag{C.12}$$

Recall that $r_n = \min_{i,j \in 1, \dots, K} |\gamma_{0i} - \gamma_{0j}|$, we have

$$\min_{i,k} |\alpha_{0i} - \gamma_{0k}| \geq r_n. \tag{C.13}$$

Combined with the Theorem 3.1 and the condition $r_n \gg p_S(\sqrt{\log p/n} + 1/\sqrt{p})$, we have

$$\begin{aligned} \max_{i,k} |(\widehat{\alpha}_i^{or} - \widehat{\gamma}_k^{or}) - (\alpha_{0i} - \gamma_{0k})| &\leq \max_i |\widehat{\alpha}_i^{or} - \alpha_{0i}| + \max_k |\widehat{\gamma}_k^{or} - \gamma_{0k}| \\ &\leq 2 \|\widehat{\gamma}^{or} - \gamma_0\|_\infty = o_{\mathbb{P}}(r_n). \end{aligned}$$

Hence, we get

$$\mathbb{P}(|\widehat{\alpha}_i^{or} - \widehat{\gamma}_k^{or}| \geq r_n) \rightarrow 1.$$

□