

# Information Density Bounds for Privacy

Sara Saeidian, *Member, IEEE*, Leonhard Grosse, *Student Member, IEEE*,

Parastoo Sadeghi, *Senior Member, IEEE*, Mikael Skoglund, *Fellow, IEEE*,

Tobias J. Oechtering, *Senior Member, IEEE*

## Abstract

This paper explores the implications of guaranteeing privacy by imposing a lower bound on the information density between the private and the public data. We introduce an operationally meaningful privacy measure called *pointwise maximal cost* (PMC) and demonstrate that imposing an upper bound on PMC is equivalent to enforcing a lower bound on the information density. PMC quantifies the information leakage about a secret to adversaries who aim to minimize non-negative cost functions after observing the outcome of a privacy mechanism. When restricted to finite alphabets, PMC can equivalently be defined as the information leakage to adversaries aiming to minimize the probability of incorrectly guessing randomized functions of the secret. We study the properties of PMC and apply it to standard privacy mechanisms to demonstrate its practical relevance. Through a detailed examination, we connect PMC with other privacy measures that impose upper or lower bounds on the information density. Our results highlight that lower bounding the information density is a more stringent requirement than upper bounding it. Overall, our work significantly bridges the gaps in understanding the relationships between various privacy frameworks and provides insights for selecting a suitable framework for a given application.

## Index Terms

Privacy, information density, pointwise maximal leakage, pointwise maximal cost, local differential privacy, information privacy, gain function, cost function.

## I. INTRODUCTION

Recent advancements in machine learning and artificial intelligence have driven progress across numerous domains of human life such as healthcare, finance, education, and transportation. However, as these technologies become omnipresent in our daily lives, the need for reliable and robust privacy protection has become more urgent than ever. Research suggests that as machine learning models become more sophisticated, the potential for privacy breaches escalates [1]. This, in turn, underscores the need for new protective measures for safeguarding sensitive information and ensuring the ethical deployment of AI technologies.

Traditionally, privacy research has focused on developing techniques to enable the release of tabular data immune to specific types of attacks, such as re-identification or attribute disclosure. These methods, which fall under the umbrella term *syntactic privacy*, include  $k$ -anonymity [2],  $l$ -diversity [3], and  $t$ -closeness [4]. While these techniques are still widely used both in the public sector and by private companies, recent research has shifted towards methods

that define and study privacy as an intrinsic property of the data release mechanism. This shift was synthesized in the framework of *differential privacy* (DP) [5]. DP is a mathematical framework with provable privacy guarantees. Its goal is to enable the theoretical analysis and implementation of statistical methods on aggregated datasets while protecting the privacy of each individual data entry.

The main principle behind DP is that two datasets differing in a single entry should be hard to distinguish based on the statistics released from each. This is achieved by controlling a key quantity known as the *privacy loss random variable* [6]. Let  $x_1$  and  $x_2$  be two databases differing in a single entry, i.e., *neighboring* databases. The privacy loss random variable associated with a data release mechanism  $M$  compares the probability of producing an outcome  $Y$  from database  $x_1$  versus database  $x_2$ , that is,

$$\mathcal{L}_{x_1, x_2}(Y) := \log \frac{\mathbb{P}[M(x_1) = Y]}{\mathbb{P}[M(x_2) = Y]}. \quad (1)$$

Various definitions of DP emerge by imposing different restrictions on the privacy loss random variable. For example, *pure DP* requires the privacy loss to be bounded for all pairs of neighboring databases [5]. Later, the concept of DP was generalized from databases to data pertaining to a single individual, leading to the development of *local differential privacy* (LDP) [7, 8]. LDP can be also formulated as a constraint on a privacy loss similar to (1); the key difference is that the loss is restricted for all possible pairs of inputs.

It is natural to question whether the definition of privacy loss, and more broadly the principles underlying DP, can be justified beyond intuition. One commonly accepted explanation is that DP balances Type I and Type II error probabilities in a hypothesis test between two neighboring datasets [9, 10]. In general, the approach of defining a mathematical measure for privacy and justifying it retrospectively has inherent limitations since it is difficult to ensure that the resulting definition is neither too weak nor excessively pessimistic. In the context of DP, for instance, opinions are notably divided. Some argue that DP is overly conservative, while others believe it may fail to provide sufficient protection, especially in scenarios involving correlated datasets [11].

The limitations of ex-post justified privacy measures suggest the need for alternatives that are *operationally meaningful*. Such measures need not rely on natural language descriptions or intuition to be interpreted. This motivation has led to the development of a novel privacy measure called *pointwise maximal leakage* (PML) [12, 13]. Unlike most measures, PML arises from a broad and encompassing threat model, first introduced in [14]. In this model, adversaries aim to maximize arbitrary non-negative gain functions encoding their objectives. This flexibility allows PML to address a wide array of attack scenarios, including membership inference, reconstruction, and attribute disclosure attacks [12].

Similar to the role of the privacy loss random variable in DP, PML is a random variable that quantifies the information leakage about a private quantity  $X$  (modeling any type of sensitive information) to a public quantity  $Y$ . PML can be expressed as

$$\ell_{P_{XY}}(X \rightarrow Y) = \operatorname{ess\,sup}_{P_X} i_{P_{XY}}(X; Y),$$

where

$$i_{P_{XY}} = \log \frac{dP_{XY}}{d(P_X \times P_Y)},$$

is the *information density* between  $X$  and  $Y$ . Thus, to restrict the information leakage between  $X$  and  $Y$ , a natural approach is to impose an upper bound on PML. This, in turn, imposes an upper bound on the information density between  $X$  and  $Y$ .

PML is not the only privacy measure related to information density. In fact, information theory has a long tradition of addressing privacy and security problems by enforcing various constraints on the information density and related quantities, especially mutual information. Other notable examples of privacy measures that restrict the information density are *local information privacy* (LIP) [15–18] and *asymmetric local information privacy* (ALIP) [19, 20]. These measures are defined by axiomatically imposing lower and upper bounds on  $i_{P_{XY}}$ . While LIP imposes bounds symmetrically, ALIP allows for different upper and lower bounds.

Now, imposing an upper bound on the information density is operationally justified through PML, and its implications have been rigorously examined in previous works [12, 13, 21]. On the other hand, the lower bound on the information density is not as well understood. Definitions such as LIP and ALIP in effect use the lower bound to remain consistent with and provide (local) DP guarantees. However, beyond compliance with DP-based definitions, the consequences of imposing a lower bound on the information density or its operational significance have remained largely unexplored in the literature.

#### A. Motivations

There are two key motivations behind this paper. First, we seek to provide an in-depth understanding of the implications of defining privacy through a lower bound on the information density. To this end, we introduce an operationally meaningful privacy measure, which we call *pointwise maximal cost* (PMC). We demonstrate that imposing an upper bound on PMC is equivalent to imposing a lower bound on the information density. PMC may be thought of as a complementary definition to PML. More precisely, while PML considers *opportunistic* adversaries aiming to maximize gain functions, PMC focuses on *risk-averse* adversaries aiming to minimize arbitrary non-negative cost functions.

Our second goal is to explore the connections and the interplay between the upper and lower bounds of information density. Specifically, we investigate whether a PML guarantee inherently provides a PMC guarantee, and vice versa. We address these questions both assuming finite alphabets and more general probability spaces. This exploration leads to insightful results connecting PML, PMC, LIP, ALIP, and LDP. In doing so, our work significantly closes the gap in the known relations between various privacy frameworks.

#### B. Contributions and Summary of the Main Results

Section III is dedicated to defining pointwise maximal cost and drawing connections to similar definitions. We begin by assuming finite alphabets and define PMC as the multiplicative decrease in the probability of incorrectly guessing the value of an arbitrary randomized function of  $X$  upon observing an outcome  $y$  of the privacy mechanism. We call this threat model the *randomized function model* of PMC, and prove in Theorem 1 that PMC, denoted by  $\Lambda_{P_{XY}}(X \rightarrow y)$ , can be expressed as

$$\Lambda_{P_{XY}}(X \rightarrow y) = \log \max_{x \in \mathcal{X}} \frac{P_X(x)}{P_{X|Y=y}(x)}.$$

Next, we argue that PMC can alternatively be defined as the multiplicative decrease in the expected value of an arbitrary non-negative *cost* function upon observing an outcome  $y$  of the privacy mechanism. This setup is referred to as the *cost function model* of PMC. Essentially, we prove an equivalence between the randomized function model and the cost function model of PMC (Theorem 2). This equivalence signifies a parallelism with PML since a similar equivalence between the gain function model and the randomized function model of PML was previously established in [12, Thm. 2].

In Sections III-C and III-D, we explore the connections between PMC and other information measures designed with risk-averse adversaries in mind. These are *maximal cost leakage* [22, Def. 11], *maximal realizable cost* [22, Def. 12], and *maximal guesswork leakage* [23]. By carefully examining these definitions and their formulations, we observe that they are often more cumbersome and complex without strengthening the privacy model. For instance, both maximal cost leakage and maximal realizable cost involve a supremum over a Markov chain, which we demonstrate can be omitted. By contrast, we show that PMC offers a streamlined yet powerful and flexible formulation that effectively encompasses these alternative definitions. It is important to note that our discussions here are restricted to finite random variables, as the other measures have only been defined in this context.

In Section III-E, we extend the cost function model to random variables on standard Borel spaces. Here, PMC is defined as the multiplicative decrease in the expected value of an arbitrary non-negative measurable cost function upon observing an outcome  $y$  of the privacy mechanism. By doing so, we obtain a general form of PMC which can be written as

$$\Lambda_{P_{XY}}(X \rightarrow y) = \operatorname{ess\,sup}_{P_X} \left( -i_{P_{XY}}(X; y) \right).$$

The above expression illustrates that imposing an upper bound on PMC is equivalent to imposing a lower bound on the information density.

Moving beyond definitions, in Section IV we explore the properties of PMC and examine its application to several well-known examples. We demonstrate that PMC meets the usual criteria expected of an information measure, such as non-negativity and satisfying data-processing inequalities (Lemma 1). We then use PMC to examine some central examples in privacy, including the randomized response mechanism [24, 25], the Laplace mechanism [5] used to release the sample mean of a dataset, and the Gaussian mechanism perturbing the value of a bounded random variable, among others.

In Section V, we explore the interactions between PMC and PML guarantees and their relationships with other definitions, specifically ALIP, LIP, and LDP. We begin with finite random variables in Section V-A and demonstrate that if a mechanism satisfies a PML guarantee with a sufficiently small parameter, then it also satisfies a PMC guarantee. Conversely, we show that a PMC guarantee always implies a PML guarantee (Theorem 4). This result is then leveraged to establish connections between PML, PMC, LIP, ALIP, and LDP and derive bounds between these measures (Corollary 3). Here, we also highlight the practical implications of our results for designing mechanisms with optimal utility.

Next, in Section V-B, we extend our discussions to random variables on standard Borel spaces. First, we show that PMC is fundamentally more restrictive than PML in the sense that bounded PMC implies bounded PML

(Theorem 5). We then prove an even stronger result stating that bounded PMC is equivalent to some form of an LDP guarantee (Theorem 6). This result demonstrates that PMC effectively serves as a *context-aware*<sup>1</sup> version of LDP and its application is constrained to mechanisms compatible with LDP and DP. Overall, our findings in this section provide a clear and unified understanding of the relationships among privacy measures that bound the information density.

### C. Other Related Works

Operationally meaningful privacy measures first emerged within the area of *quantitative information flow* [26] through the concept of *min-entropy leakage* [27, 28], also known as *multiplicative Bayes leakage*. Min-entropy leakage assumes that an adversary attempts to guess the value of the secret  $X$  in one try. Various extensions of this adversarial model have since been explored [14, 29, 30, 26]. Among these works, the *g-leakage* framework of Alvim et al. [14] has been particularly influential. This framework generalizes min-entropy leakage by allowing adversaries to construct guesses of  $X$  that maximize arbitrary non-negative gain functions.

Issa et al. [22] introduced the threat model approach to privacy into the domain of information theory through the development of *maximal leakage*. This measure focuses on adversaries aiming to guess the values of arbitrary (randomized) functions of  $X$ . Issa et al. [22] showed that maximal leakage is a highly robust measure since many variations of their original threat model still yield the same quantity as maximal leakage. Liao et al. [31] further extended maximal leakage by introducing a tunable loss function called  $\alpha$ -loss. Then, they considered the class of adversaries attempting to minimize the  $\alpha$ -loss, which gave rise to the notions of  $\alpha$ -leakage and *maximal  $\alpha$ -leakage*. More recent developments in this area include pointwise maximal leakage, defined as the pointwise adaptation of *g-leakage* and maximal leakage [12], *maximal  $(\alpha, \beta)$ -leakage* [32], and *maximal  $g$ -leakage under multiple guesses* [33].

The broader area of information theory has also contributed significantly to the study of privacy measures. Here, mutual information is notably the most extensively examined privacy measure [34–40]. Beyond mutual information, generalized information measures such as *f-information* [41] have also been used to assess privacy. Specific instances of *f-information* utilized as privacy measures include  $\chi^2$ -information [42, 43] (associated with  $\chi^2$ -divergence), and total variation privacy [44] (associated with total variation distance). Additionally, some works have used per-letter *f-divergences* (as opposed to the average-case divergence in *f-information*) as privacy measures [45–47]. Other research has examined the probability of correctly guessing [48] or guesswork [49, 50] for guaranteeing privacy.

Finally, several works have studied the design of privacy mechanisms subject to various privacy guarantees. These include local differential privacy [51–53], maximal leakage [54, 55], total variation privacy [44], and LIP [56, 57, 18], among others.

<sup>1</sup>A privacy measure is said to be *context-aware* if it depends on both the privacy mechanism and the distribution of the sensitive data. Conversely, it is said to be *context-free* if it depends on the privacy mechanism alone.

## II. PRELIMINARIES

### A. Notation and Assumptions

The following notations are used throughout the paper:  $\mathbb{R}_+ = [0, \infty)$  and  $\bar{\mathbb{R}}_+ = [0, \infty]$  denote the non-negative real numbers and the extended non-negative real numbers. The set  $[n] := \{1, \dots, n\}$  represents the first  $n$  positive integers,  $\log(\cdot)$  refers to the natural logarithm, and  $\mathbf{1}_A$  denotes the indicator function of the set  $A$ . We also adopt the convention that  $0/0 = 1$ .

Let  $(\Omega, \mathcal{H}, \mathbb{P})$  be an abstract probability space fixed in the background, where  $\Omega$  is the sample space,  $\mathcal{H}$  is the event space (i.e., a  $\sigma$ -algebra on  $\Omega$ ), and  $\mathbb{P}$  is a probability measure on the measurable space  $(\Omega, \mathcal{H})$ . We use  $\mathcal{H}_+$  to denote the set of all functions that are measurable relative to  $\mathcal{H}$  and  $\mathcal{B}_{\bar{\mathbb{R}}_+}$ , where  $\mathcal{B}_{\bar{\mathbb{R}}_+}$  denotes the Borel  $\sigma$ -algebra on  $\bar{\mathbb{R}}_+$ . Given a function  $h \in \mathcal{H}_+$ , the essential supremum of  $h$  with respect to  $\mathbb{P}$  is  $\text{ess sup } h = \sup\{c \in \bar{\mathbb{R}}_+ : \mathbb{P}(\{h > c\}) > 0\}$  and its essential infimum is  $\text{ess inf } h = \inf\{c \in \bar{\mathbb{R}}_+ : \mathbb{P}(\{h < c\}) > 0\}$ . Suppose  $\lambda$  and  $\rho$  are measures on  $(\Omega, \mathcal{H})$  and assume that  $\lambda$  is  $\sigma$ -finite. If  $\rho$  is absolutely continuous with respect to  $\lambda$ , denoted by  $\rho \ll \lambda$ , then we write  $p = \frac{d\rho}{d\lambda}$  to imply that

$$\int_{\Omega} h(\omega) \rho(d\omega) = \int_{\Omega} h(\omega) p(\omega) \lambda(d\omega),$$

for all  $h \in \mathcal{H}_+$ , where  $p \in \mathcal{H}_+$  is the Radon-Nikodym derivative of  $\rho$  with respect to  $\lambda$ .

Let  $\mathcal{X}$  be a set and  $\mathcal{S}_X$  a  $\sigma$ -algebra on  $\mathcal{X}$ . A mapping  $X : \Omega \rightarrow \mathcal{X}$  is called a random variable taking values in  $(\mathcal{X}, \mathcal{S}_X)$  if  $X$  is measurable relative to  $\mathcal{H}$  and  $\mathcal{S}_X$ . In this paper, we exclusively use  $X$  to denote some data containing sensitive information, i.e., the *secret*. We use  $P_X$  to denote the distribution of  $X$ . Often, we assume that  $\mathcal{X}$  is a discrete set and use some notations specific to this case. In particular, (with a slight abuse of notation) we also use  $P_X$  to denote the probability mass function (pmf) of  $X$  and write  $P_X(x) := P_X(\{x\})$  for  $x \in \mathcal{X}$ . Moreover, we assume that  $P_X$  has full support on  $\mathcal{X}$ , that is,  $P_X(x) > 0$  for all  $x \in \mathcal{X}$ , and define  $p_{\min} := \min_{x \in \mathcal{X}} P_X(x)$ .

Let  $(\mathcal{Y}, \mathcal{S}_Y)$  be a measurable space. A mapping  $P_{Y|X} : \mathcal{X} \times \mathcal{S}_Y \rightarrow [0, 1]$  is called a transition probability kernel (or simply kernel) from  $(\mathcal{X}, \mathcal{S}_X)$  into  $(\mathcal{Y}, \mathcal{S}_Y)$  if the mapping  $x \mapsto P_{Y|X=x}(\mathcal{B})$  is in  $\mathcal{S}_{X+}$  for all  $\mathcal{B} \in \mathcal{S}_Y$ , and  $P_{Y|X=x}(\cdot)$  is a probability measure on  $(\mathcal{Y}, \mathcal{S}_Y)$  for all  $x \in \mathcal{X}$ . We sometimes use  $P_{Y|X}(\mathcal{B} | x)$  instead of  $P_{Y|X=x}(\mathcal{B})$ . This notation is less awkward when we do not want to specify the outcome of  $X$  but leave it as a random variable. The kernel  $P_{Y|X}$  induces a random variable  $Y$  taking values in  $(\mathcal{Y}, \mathcal{S}_Y)$  with distribution  $P_Y$ , where

$$P_Y(\mathcal{B}) = \int_{\mathcal{X}} P_{Y|X=x}(\mathcal{B}) P_X(dx), \quad \mathcal{B} \in \mathcal{S}_Y. \quad (2)$$

We write  $P_Y = P_{Y|X} \circ P_X$  to represent *marginalization* over  $X$  described by (2). In this paper, we exclusively use  $Y$  to denote some publicly available data that contains information about  $X$ , and refer to  $P_{Y|X}$  as the *privacy mechanism*. When  $\mathcal{Y}$  is a discrete set, we use a similar notation described above in the case of discrete  $\mathcal{X}$ . For example, we write  $P_{Y|X=x}(y) := P_{Y|X=x}(\{y\})$  with  $y \in \mathcal{Y}$ .

Let  $P_{XY}$  be a probability measure on the product space  $(\mathcal{X} \times \mathcal{Y}, \mathcal{S}_X \otimes \mathcal{S}_Y)$  with marginals  $P_X$  and  $P_Y$ . Then, we write  $P_{XY} = P_X \times P_{Y|X}$  to denote *disintegration*, i.e., to imply that

$$\mathbb{E}[h] = \int_{\mathcal{X} \times \mathcal{Y}} h(x, y) P_{XY}(dx, dy) = \int_{\mathcal{X}} P_X(dx) \int_{\mathcal{Y}} h(x, y) P_{Y|X=x}(dy),$$

for all  $h \in (\mathcal{S}_X \otimes \mathcal{S}_Y)_+$ . Let  $P_X \times P_Y$  denote the product measure of  $P_X$  and  $P_Y$ . Throughout the paper it is assumed that  $P_{XY} \ll P_X \times P_Y$  and the logarithm of the Radon-Nikodym derivative

$$i_{P_{XY}} := \log \frac{dP_{XY}}{d(P_X \times P_Y)},$$

is referred to as *information density* of  $X$  and  $Y$ . We assume that all probability measures on  $(\mathcal{X}, \mathcal{S}_X)$  are absolutely continuous with respect to a common  $\sigma$ -finite measure  $\mu$  and all probability measures on  $(\mathcal{Y}, \mathcal{S}_Y)$  are absolutely continuous with respect to a common  $\sigma$ -finite measure  $\nu$ . Additionally, we assume that  $P_{XY} \ll \mu \times \nu$ .

Finally, all measurable spaces are assumed to be *standard Borel* [58, Def. 8.35]. Standard Borel spaces have several convenient properties. Most prominently, joint distributions on standard Borel spaces can always be disintegrated into a kernel and a marginal distribution [59, Thm. IV.2.18]. That is, if  $P_{XY}$  is a distribution on  $(\mathcal{X} \times \mathcal{Y}, \mathcal{S}_X \otimes \mathcal{S}_Y)$  with marginal  $P_X$  on  $(\mathcal{X}, \mathcal{S}_X)$ , then there exists a transition probability kernel from  $(\mathcal{X}, \mathcal{S}_X)$  into  $(\mathcal{Y}, \mathcal{S}_Y)$  such that  $P_{XY} = P_X \times P_{Y|X}$ . Another advantage is that if  $P_{Y|X}$  and  $Q_{Y|X}$  are both kernels and  $P_{Y|X=x} \ll Q_{Y|X=x}$  for all  $x \in \mathcal{X}$ , then we may invoke Doob's version of the Radon-Nikodym theorem to obtain a Radon-Nikodym derivative  $\frac{dP_{Y|X}}{dQ_{Y|X}}$  which is jointly measurable in  $(x, y)$  [59, Thm. V.4.44]. This has the effect that we may alternatively use  $\log \frac{dP_{Y|X}}{dP_Y}$  or  $\log \frac{dP_{X|Y}}{dP_X}$  as the information density. Note that all measurable spaces of practical interest are standard Borel, e.g., countable sets,  $\mathbb{R}^n$ , or complete separable metric spaces endowed with Borel  $\sigma$ -algebras.

### B. Pointwise Maximal Leakage

Pointwise maximal leakage (PML) is a recently introduced privacy measure that is operationally meaningful, robust, and flexible. For secrets with finite alphabets, PML is defined by quantifying risks in two powerful adversarial threat models: the *gain function model*, first introduced by Alvim et al. [14], and the *randomized function model*, put forward by Issa et al. [22]. It has been shown that when  $X$  is a finite random variable, these two models are equivalent [12, Thm. 2]. Furthermore, the gain function model can be extended to secrets taking values in standard Borel spaces [13].

Before we define PML, let us first recall the definition of Rényi divergence of order infinity [60, 61], which is used to provide simplified expressions for PML, and later, PMC.

**Definition 1** (Rényi divergence of order  $\infty$  [61, Thm. 6]). Let  $\mathbb{P}$  and  $\mathbb{Q}$  be probability measures on the measurable space  $(\Omega, \mathcal{H})$ . Let  $\lambda$  be a  $\sigma$ -finite measure satisfying  $\mathbb{P} \ll \lambda$  and  $\mathbb{Q} \ll \lambda$ . The Rényi divergence of order  $\infty$  of  $\mathbb{P}$  from  $\mathbb{Q}$  is defined as

$$D_\infty(\mathbb{P}||\mathbb{Q}) = \log \sup_{\mathcal{A} \in \mathcal{H}} \frac{\mathbb{P}(\mathcal{A})}{\mathbb{Q}(\mathcal{A})} = \log \left( \operatorname{ess\,sup}_{\mathbb{P}} \frac{p}{q} \right), \quad (3)$$

where  $p = \frac{d\mathbb{P}}{d\lambda}$  and  $q = \frac{d\mathbb{Q}}{d\lambda}$ .

If  $\mathbb{P} \ll \mathbb{Q}$ , then the divergence can also be expressed as

$$D_\infty(\mathbb{P}||\mathbb{Q}) = \log \left( \operatorname{ess\,sup}_{\mathbb{P}} \frac{d\mathbb{P}}{d\mathbb{Q}} \right) = \log \left( \operatorname{ess\,sup}_{\mathbb{Q}} \frac{d\mathbb{P}}{d\mathbb{Q}} \right).$$

On the other hand, if  $\mathbb{P} \not\ll \mathbb{Q}$ , then  $D_\infty(\mathbb{P} \parallel \mathbb{Q}) = \infty$ . When the sample space  $\Omega$  is countable we may write (3) in the form

$$D_\infty(\mathbb{P} \parallel \mathbb{Q}) = \log \left( \sup_{\omega \in \Omega} \frac{\mathbb{P}(\omega)}{\mathbb{Q}(\omega)} \right).$$

Below, we first define PML in its most general form using the gain function model. Then, assuming that  $X$  is a finite random variable, we also define PML using the randomized function model.

**Definition 2** (Gain function view of PML [13, Def. 3]). Given a joint distribution  $P_{XY}$  on the product measurable space  $(\mathcal{X} \times \mathcal{Y}, \mathcal{S}_X \otimes \mathcal{S}_Y)$ , we define the pointwise maximal leakage from  $X$  to  $y \in \mathcal{Y}$  as

$$\ell_{P_{XY}}(X \rightarrow y) := \log \sup_{\substack{(\mathcal{W}, \mathcal{S}_W), \\ g \in \mathcal{G}}} \frac{\sup_{P_{W|Y}} \mathbb{E}[g(X, W) \mid Y = y]}{\sup_{w \in \mathcal{W}} \mathbb{E}[g(X, w)]}, \quad (4)$$

where the supremum in the numerator is over all transition probability kernels  $P_{W|Y}$  from  $(\mathcal{Y}, \mathcal{S}_Y)$  into  $(\mathcal{W}, \mathcal{S}_W)$ , and  $\mathcal{G}$  denotes the set of all gain functions defined as

$$\mathcal{G} := \left\{ g \in (\mathcal{S}_X \otimes \mathcal{S}_W)_+ \mid \sup_{w \in \mathcal{W}} \mathbb{E}[g(X, w)] < \infty \right\}.$$

Definition 2 can be explained in the following way. Consider an adversary who aims to construct an estimate of  $X$ , denoted by  $W$ , in order to maximize a non-negative gain function  $g$ . The gain function  $g$  encapsulates the adversary's goal and can be adjusted to represent a variety of privacy attacks. For example, if  $X$  is a dataset, then  $g$  might model membership inference attacks [12]. To measure the information leakage associated with a single released outcome  $y \in \mathcal{Y}$ , we compare the expected value of  $g$  after observing  $y$  (in the numerator of (4)) with the expected value of  $g$  before observing  $y$  (in the denominator). The posterior expected gain is calculated using the optimal estimation kernel  $P_{W|Y}$ , while the prior expected gain is  $\sup_{w \in \mathcal{W}} \mathbb{E}[g(X, w)]$ . Then, to ensure that the privacy measure resulting from this definition encompasses various attacks, the ratio of the posterior-to-prior expected gain is maximized over all possible measurable spaces  $(\mathcal{W}, \mathcal{S}_W)$  and all gain functions in  $\mathcal{G}$ .

Next, we consider finite  $X$ 's and define PML through the randomized function model.

**Definition 3** (Randomized function view of PML [12, Def. 1]). Suppose  $(X, Y) \sim P_{XY}$ , where  $X$  is a finite random variable. The pointwise maximal leakage from  $X$  to  $y \in \mathcal{Y}$  is defined as

$$\ell_{P_{XY}}(X \rightarrow y) = \log \sup_{U: U-X-Y} \frac{\sup_{P_{U|Y}} \mathbb{P}[U = \hat{U} \mid Y = y]}{\max_{u \in \mathcal{U}} P_U(u)},$$

where  $U$  and  $\hat{U}$  are random variables on a finite set  $\mathcal{U}$ ,  $P_U = P_{U|X} \circ P_X$ , and the Markov chain  $U - X - Y - \hat{U}$  holds.

Definition 3 can be understood similarly to Definition 2. Let  $U$  be a randomized function of  $X$ . To quantify the information leakage associated with a single outcome of the mechanism  $y \in \mathcal{Y}$ , we compare the probability of correctly guessing  $U$  after observing  $y$  with the *a priori* probability of correctly guessing  $U$  in a ratio. Then, to obtain a robust privacy measure, this posterior-to-prior ratio is maximized over all possible randomized functions of  $X$ , represented by the supremum over  $U$ 's satisfying the Markov chain  $U - X - Y$ .



It has been shown that both Definitions 2 and 3 simplify to the following expression for PML [12, 13]:

$$\ell_{P_{XY}}(X \rightarrow y) = D_\infty(P_{X|Y=y} \| P_X),$$

where  $P_{X|Y=y}$  denotes the posterior distribution of  $X$  given  $y \in \mathcal{Y}$ . Observe that PML is a context-aware privacy measure since it depends on the prior distribution  $P_X$ .

Viewing  $\ell_{P_{XY}}(X \rightarrow Y)$  as a function of the random variable  $Y$ , PML itself becomes a random variable whose various statistical properties can be examined and constrained to provide privacy guarantees. A stringent privacy guarantee can thus be defined by requiring  $\ell_{P_{XY}}(X \rightarrow Y)$  to be essentially bounded. Formally, we say that a privacy mechanism  $P_{Y|X}$  satisfies  $\varepsilon$ -PML if  $\ell_{P_{XY}}(X \rightarrow Y) \leq \varepsilon$  (almost surely) for some  $\varepsilon \geq 0$ . Building on this, [62] investigates the design of privacy mechanisms under the  $\varepsilon$ -PML constraint for secrets with finite alphabets. Specifically, the authors formulate a linear program for maximizing a class of convex utility functions called *sub-convex* functions [62, Def. 5]. The optimal solutions to this linear program are referred to as *PML extremal mechanisms*. The paper also introduces the concept of *PML privacy regions* as a partitioning of the space of the privacy parameter  $\varepsilon \in [0, \infty)$  into disjoint intervals. The privacy regions specify an upper bound on the number  $(x, y) \in \mathcal{X} \times \mathcal{Y}$  pairs with  $P_{Y|X=x}(y) = 0$ . The first region, termed the *high-privacy regime*, corresponds to the interval  $0 \leq \varepsilon < \log \frac{1}{1-p_{\min}}$ . In this region, no zero probability assignments are allowed in the mechanism. It is then proved that the PML extremal mechanism  $P_{Y|X}^*$  in the high-privacy regime has the following form:

$$P_{Y|X=i}^*(j) = \begin{cases} 1 - e^\varepsilon(1 - P_X(i)) & \text{if } i = j, \\ e^\varepsilon P_X(j) & \text{if } i \neq j, \end{cases}$$

where  $i, j \in [|\mathcal{X}|]$  [62, Thm. 3].

### C. Local Information Privacy and Asymmetric Local Information Privacy

Besides PML, several other context-aware privacy measures have been defined in the literature. Two notable examples are *local information privacy* (LIP) and *asymmetric local information privacy* (ALIP). These measures are defined by axiomatically imposing lower and upper bounds on the information density between  $X$  and  $Y$ . First, we define LIP which imposes symmetric lower and upper bounds.

**Definition 4** (Local information privacy [16, 15]). Given  $\varepsilon \geq 0$ , a privacy mechanism  $P_{Y|X}$  is said to satisfy  $\varepsilon$ -LIP if  $|i_{P_{XY}}(X; Y)| \leq \varepsilon$  almost surely.

It was observed in [19] that the distribution of  $i(X; Y)$  exhibits an asymmetry in the sense that it has a skewed long tail for negative values, and a much shorter range and a sharp fall-off for positive values. This asymmetry, which the authors call the *lift-asymmetry*, motivated their definition of ALIP.

**Definition 5** (Asymmetric local information privacy [19]). Given  $\varepsilon_l, \varepsilon_u \geq 0$ , a privacy mechanism  $P_{Y|X}$  is said to satisfy  $(\varepsilon_l, \varepsilon_u)$ -ALIP if  $-\varepsilon_l \leq i_{P_{XY}}(X; Y) \leq \varepsilon_u$  almost surely.

It follows directly from the above definitions that  $\varepsilon_u$ -LIP and  $(\varepsilon_l, \varepsilon_u)$ -ALIP both imply  $\varepsilon_u$ -PML. However, it is not immediately clear whether PML also provides LIP or ALIP guarantees, as PML does not directly impose a

lower bound on the information density. As we will see, in some cases PML does implicitly impose a lower bound on the information density, thereby providing LIP and ALIP guarantees. We discuss this subject in Section V.

#### D. Local Differential Privacy

Differential privacy in its original form assumes that all data is collected in a database managed by a trusted data curator. Here, we focus on the *local* model of privacy offering an alternative that removes the need for a central database. In this model, each user's data is perturbed before collection to ensure that only the user has access to their original, unaltered information. This approach gives rise to the concept of *local differential privacy* (LDP) [7, 8].

**Definition 6** (Local differential privacy). Let  $X$  and  $X'$  be independent random variables distributed according to  $P_X$ . Given  $\varepsilon \geq 0$ , a privacy mechanism  $P_{Y|X}$  satisfies  $\varepsilon$ -LDP if

$$\mathbb{P} \left( \left\{ \sup_{\mathcal{E} \in \mathcal{S}_Y} \log \frac{P_{Y|X}(\mathcal{E} | X)}{P_{Y|X}(\mathcal{E} | X')} \leq \varepsilon \right\} \right) = 1,$$

where the probability is over the draws of  $X$  and  $X'$ . Alternatively,  $P_{Y|X}$  satisfies  $\varepsilon$ -LDP if

$$\operatorname{ess\,sup}_{P_X \times P_X} D_\infty \left( P_{Y|X}(\cdot | X) \parallel P_{Y|X}(\cdot | X') \right) \leq \varepsilon.$$

**Remark 1.** It is usually said that local differential privacy is a context-free definition independent of the distribution  $P_X$ . That is, it describes a property of the privacy mechanism  $P_{Y|X}$ . However, note that technically Definition 6 depends on  $P_X$  through its support.

LDP has close connections to context-aware privacy measures. The following result from [18] demonstrates how LDP guarantees translate into PML, LIP, and ALIP guarantees assuming that  $\mathcal{X}$  and  $\mathcal{Y}$  are finite. Note that while [18, Thm. 1] articulates the LIP result alone, its proof also provides guarantees in terms of ALIP and PML. Recall that  $p_{\min} := \min_{x \in \mathcal{X}} P_X(x)$ .

**Proposition 1** ([18, Thm. 1]). Let  $X$  and  $Y$  be finite random variables and suppose  $X$  is distributed according to  $P_X$ . Given  $\varepsilon \geq 0$ , define

$$\begin{aligned} \varepsilon_1 &= \log \left( p_{\min} + e^\varepsilon (1 - p_{\min}) \right), \\ \varepsilon_2 &= -\log \left( p_{\min} + e^{-\varepsilon} (1 - p_{\min}) \right). \end{aligned}$$

If  $P_{Y|X}$  satisfies  $\varepsilon$ -LDP, then it also satisfies  $\varepsilon_1$ -LIP,  $(\varepsilon_1, \varepsilon_2)$ -ALIP and  $\varepsilon_2$ -PML.

The previous result is straightforward and intuitive as it illustrates the transition from a context-free property of a mechanism to guarantees that hold for a specific prior distribution. However, the reverse direction is more intricate. It involves moving from a property tied to both a mechanism and a specific prior distribution to a universal property of the mechanism that holds irrespective of the prior. Interestingly, [18, Prop. 1] showed that  $\varepsilon$ -LIP implies  $2\varepsilon$ -LDP, and [19, Prop. 1] established that  $(\varepsilon_l, \varepsilon_u)$ -ALIP yields  $\varepsilon_l + \varepsilon_u$ -LDP. In general,  $\varepsilon$ -PML may not provide any LDP guarantee since a mechanism satisfying  $\varepsilon$ -PML may contain  $P_{Y|X=x}(y) = 0$  for some  $(x, y) \in \mathcal{X} \times \mathcal{Y}$ . Nonetheless, as we will explore in Section V, in the high-privacy regime PML guarantees indeed translate into valid LDP guarantees.

### III. QUANTIFYING INFORMATION LEAKED TO RISK-AVERSE ADVERSARIES

The purpose of this section is to explore the implications of defining privacy by imposing a lower bound on the information density  $i_{P_{XY}}$ . We begin by assuming that  $X$  is a finite random variable and demonstrate that given  $y \in \mathcal{Y}$ ,  $-\min_x i_{P_{XY}}(x; y)$  describes the information leaking to *risk-averse* adversaries, as opposed to the *opportunistic* adversaries of PML. In particular, we show that  $-\min_x i_{P_{XY}}(x; y)$  is operationally meaningful in two adversarial threat models. The first setup called the *randomized function* model, considers adversaries who wish to minimize the probability of incorrectly guessing the value of an arbitrary (randomized) function of  $X$ . The second setup, called the *cost function* model, considers adversaries who wish to minimize arbitrary non-negative cost functions. We refer to the information leakage measure obtained in both threat models as *pointwise maximal cost* (PMC).

In Section III-E, we extend the cost function definition of PMC to secrets on standard Borel spaces. When  $\mathcal{X}$  is finite, the cost function definition of PMC is closely related to three other informational leakage measures, namely *maximal cost leakage*, *maximal realizable cost*, and *maximal guesswork leakage* defined in [22] and [23]. We discuss their connections in Sections III-C and III-D.

#### A. Randomized Function Model

Suppose  $\mathcal{X}$  is a finite set. Consider an adversary who attempts to guess the value of a (finite) randomized function of  $X$ , denoted by  $U$ . Upon observing  $y \in \mathcal{Y}$ , the adversary formulates a guess of  $U$ , denoted by  $\hat{U}$  using a kernel  $P_{\hat{U}|Y}$  that minimizes the probability of an incorrect guess. Essentially, this threat model is an adaptation of [22] and [12, Section II-A] to the risk-averse case, where the adversary seeks to minimize the probability of incorrectly guessing  $U$  instead of maximizing the probability of correctly guessing it. To quantify the risk incurred by such an adversary, we compare the adversary's probability of incorrectly guessing  $U$  having access to  $y$  and the probability of incorrectly guessing  $U$  without access. Consequently, we define

$$\Lambda_U(X \rightarrow y) := \log \frac{\inf_{P_{\tilde{U}}} \mathbb{P}[U \neq \tilde{U}]}{\inf_{P_{\hat{U}|Y}} \mathbb{P}[U \neq \hat{U} | Y = y]}. \quad (5)$$

In general, we may not know which function of  $X$  the adversary is interested in, or different adversaries may be interested in different functions of  $X$ . Therefore, to achieve robustness in our model, we define the pointwise maximal cost (PMC) by maximizing  $\Lambda_U(X \rightarrow y)$  over all possible  $U$ 's satisfying the Markov chain  $U - X - Y$ .

**Definition 7** (Randomized function view of PMC). Suppose  $X$  is a finite random variable and let  $P_{XY}$  be the joint distribution of  $X$  and  $Y$ . Given  $y \in \mathcal{Y}$ , the pointwise maximal cost from  $X$  to  $y$  is

$$\begin{aligned} \Lambda_{P_{XY}}(X \rightarrow y) &:= \sup_{U:U-X-Y} \Lambda_U(X \rightarrow y) \\ &= \sup_{U:U-X-Y} \log \frac{\inf_{P_{\tilde{U}}} \mathbb{P}[U \neq \tilde{U}]}{\inf_{P_{\hat{U}|Y}} \mathbb{P}[U \neq \hat{U} | Y = y]}, \end{aligned} \quad (6)$$

where  $U, \tilde{U}$ , and  $\hat{U}$  are random variables defined on a finite set  $\mathcal{U}$ .

Below, we show how  $\Lambda_{P_{XY}}(X \rightarrow y)$  relates to  $\min_x i(x; y)$ .

**Theorem 1.** Let  $P_{XY}$  denote the joint distribution of  $X$  and  $Y$ . Given  $y \in \mathcal{Y}$ , PMC can be expressed as

$$\Lambda_{P_{XY}}(X \rightarrow y) = D_\infty(P_X \| P_{X|Y=y}),$$

where  $P_{X|Y=y}$  denotes the posterior distribution of  $X$ . Thus, for all  $y \in \mathcal{Y}$ , lower bounding  $\min_{x \in \mathcal{X}} i(x; y)$  is equivalent to upper bounding  $\Lambda_{P_{XY}}(X \rightarrow y)$ . That is, given  $\varepsilon \geq 0$  it holds that

$$-\varepsilon \leq \min_{x \in \mathcal{X}} i(x; y) \iff \Lambda_{P_{XY}}(X \rightarrow y) \leq \varepsilon,$$

for all  $y \in \mathcal{Y}$ .

*Proof:* Fix an arbitrary  $y \in \mathcal{Y}$ . First, we show that  $\Lambda_{P_{XY}}(X \rightarrow y) \leq D_\infty(P_X \| P_{X|Y=y})$ . Let  $u_1 \in \arg \max_{u \in \mathcal{U}} P_U(u)$  and  $u_2 \in \arg \max_{u \in \mathcal{U}} P_{U|Y=y}(u)$ . Using the fact that

$$\begin{aligned} \inf_{P_{\tilde{U}}} \mathbb{P}[U \neq \tilde{U}] &= 1 - \sup_{P_{\tilde{U}}} \mathbb{P}[U = \tilde{U}] = 1 - \max_{u \in \mathcal{U}} P_U(u), \\ \inf_{P_{\hat{U}|Y}} \mathbb{P}[U \neq \hat{U} | Y = y] &= 1 - \sup_{P_{\hat{U}|Y}} \mathbb{P}[U = \hat{U} | Y = y] = 1 - \max_{u \in \mathcal{U}} P_{U|Y=y}(u), \end{aligned}$$

we write

$$\begin{aligned} \exp(\Lambda_{P_{XY}}(X \rightarrow y)) &= \sup_{U:U-X-Y} \frac{1 - \max_u P_U(u)}{1 - \max_u P_{U|Y=y}(u)} \\ &= \sup_{U:U-X-Y} \frac{\sum_{u \neq u_1} P_U(u)}{\sum_{u \neq u_2} P_{U|Y=y}(u)}. \end{aligned}$$

Since  $P_U(u_1) = \max_u P_U(u) \geq P_U(u_2)$ , we have

$$\begin{aligned} \frac{\sum_{u \neq u_1} P_U(u)}{\sum_{u \neq u_2} P_{U|Y=y}(u)} &\leq \frac{\sum_{u \neq u_2} P_U(u)}{\sum_{u \neq u_2} P_{U|Y=y}(u)} \\ &= \frac{\sum_{u \neq u_2} P_{U|Y=y}(u) \left( \frac{P_U(u)}{P_{U|Y=y}(u)} \right)}{\sum_{u \neq u_2} P_{U|Y=y}(u)} \\ &\leq \max_{u \neq u_2} \frac{P_U(u)}{P_{U|Y=y}(u)} \\ &= \max_{u \neq u_2} \frac{\sum_{x \in \mathcal{X}} P_{U|X=x}(u) P_X(x)}{\sum_{x \in \mathcal{X}} P_{U|X=x}(u) P_{X|Y=y}(x)} \\ &= \max_{u \neq u_2} \frac{\sum_{x \in \mathcal{X}} P_{U|X=x}(u) P_{X|Y=y}(x) \left( \frac{P_X(x)}{P_{X|Y=y}(x)} \right)}{\sum_{x \in \mathcal{X}} P_{U|X=x}(u) P_{X|Y=y}(x)} \\ &\leq \max_{x \in \mathcal{X}} \frac{P_X(x)}{P_{X|Y=y}(x)} \\ &= \exp\left(D_\infty(P_X \| P_{X|Y=y})\right). \end{aligned}$$

We conclude that  $\Lambda_{P_{XY}}(X \rightarrow y) \leq D_\infty(P_X \| P_{X|Y=y})$ .

Next, we show that  $\Lambda_{P_{XY}}(X \rightarrow y) \geq D_\infty(P_X \| P_{X|Y=y})$ . Let  $x^* \in \arg \max_{x \in \mathcal{X}} \frac{P_X(x)}{P_{X|Y=y}(x)}$ , and define  $V := \mathbf{1}_{X \setminus \{x^*\}}$ . Then,  $V$  is a binary random variable with distribution  $P_V(0) = 1 - P_V(1) = P_X(x^*)$ . Let  $k$  be a large integer. We define a random variable  $W$  with alphabet  $\mathcal{W} = \{1, \dots, k+1\}$  and induced by the kernel

$$P_{W|V=0}(w) = \begin{cases} \frac{1}{k} & \text{if } w \in \{1, \dots, k\}, \\ 0 & \text{if } w = k+1, \end{cases}$$

and

$$P_{W|V=1}(w) = \begin{cases} 0 & \text{if } w \in \{1, \dots, k\}, \\ 1 & \text{if } w = k+1. \end{cases}$$

By inspection,  $P_W(w) = \frac{P_V(0)}{k} = \frac{P_X(x^*)}{k}$  for  $w \in [k]$  and  $P_W(k+1) = P_V(1)$ . Thus, taking  $k$  to be sufficiently large, we can ensure that  $P_W(k+1) = \max_w P_W(w)$ . Moreover, since the Markov chain  $W - V - X - Y$  holds  $P_{W|Y=y}(w) = \frac{P_{V|Y=y}(0)}{k} = \frac{P_{X|Y=y}(x^*)}{k}$  for  $w \in [k]$  and  $P_{W|Y=y}(k+1) = P_{V|Y=y}(1) = P_{X|Y=y}(X \setminus \{x^*\})$ . Once again, taking  $k$  to be sufficiently large we can ensure that  $P_{W|Y=y}(k+1) = \max_w P_{W|Y=y}(w)$ .

Since  $W - X - Y$  forms a Markov chain, we get

$$\begin{aligned} \exp\left(\Lambda_{P_{XY}}(X \rightarrow y)\right) &= \sup_{U:U-X-Y} \frac{1 - \max_u P_U(u)}{1 - \max_u P_{U|Y=y}(u)} \\ &\geq \frac{1 - \max_w P_W(w)}{1 - \max_w P_{W|Y=y}(w)} \\ &= \frac{\sum_{w \neq k+1} P_W(w)}{\sum_{w \neq k+1} P_{W|Y=y}(w)} \\ &= \frac{P_X(x^*)}{P_{X|Y=y}(x^*)} \\ &= \max_{x \in \mathcal{X}} \frac{P_X(x)}{P_{X|Y=y}(x)} \\ &= \exp\left(D_\infty\left(P_X \| P_{X|Y=y}\right)\right), \end{aligned}$$

as desired. ■

**Remark 2** (Closedness under pre-processing). Importantly, it follows directly from (6) that PMC is *closed under pre-processing*, that is,

$$\Lambda_{P_{ZY}}(Z \rightarrow y) \leq \Lambda_{P_{XY}}(X \rightarrow y),$$

for all  $Z$ 's satisfying the Markov chain  $Z - X - Y$  and all  $y \in \mathcal{Y}$ . Closedness under pre-processing implies that the amount of information leaking about (random) functions of  $X$  can never exceed the amount of information leaking about  $X$  itself.

**Remark 3.** It is worth emphasizing that even when  $X$  and  $Y$  are both finite random variables, PMC may be infinite. This occurs if  $P_{Y|X=x}(y) = 0$  for some  $(x, y)$  pair with  $P_X(x)P_Y(y) > 0$ , in other words, if  $P_X \times P_Y \not\ll P_{XY}$ . This characteristic also distinguishes PMC from PML since PML is always bounded when  $X$  is finite.

### B. Cost Function Model

Once again, let  $\mathcal{X}$  be a finite set. Now, we consider adversaries aiming to minimize the expected value of non-negative *cost functions*  $c : \mathcal{X} \times \mathcal{W} \rightarrow \mathbb{R}_+$ , where  $\mathcal{W}$  is an arbitrary finite set representing the adversary's guessing space. In this scenario, the adversary observes  $y \in \mathcal{Y}$  and selects  $w \in \mathcal{W}$  to minimize  $\mathbb{E}[c(X, w) \mid Y = y]$ . Essentially, this threat model is an adaptation of [14] and [12, Section II-B] to the risk-averse case, where the adversary seeks to minimize a cost function instead of maximizing a gain function. To quantify the risk associated with such an adversary, we compare the smallest posterior expected cost with the smallest prior expected cost in a ratio. Accordingly, we define

$$\Lambda_c(X \rightarrow y) := \log \frac{\inf_{P_W} \mathbb{E}[c(X, W)]}{\inf_{P_{\hat{W}|Y}} \mathbb{E}[c(X, \hat{W}) \mid Y = y]}. \quad (7)$$

The expectation in the numerator of (7) is taken with respect to  $P_{XW} = P_X \times P_W$  since  $X$  and  $W$  are statistically independent. On the other hand, in the denominator of (7), the expectation is taken with respect to  $P_{X\hat{W}|Y} = P_{X|Y} \times P_{\hat{W}|Y}$  since  $X - Y - \hat{W}$  forms a Markov chain.

Similar to the equivalence between the randomized function model and the gain function model established in [12, Thm. 2] for PML, below we demonstrate that (5) and (7) provide equivalent characterizations of PMC. This result is proved in Appendix A.

**Theorem 2.** Suppose  $X$  is a finite random variable and let  $y \in \mathcal{Y}$ . For every randomized function of  $X$ , denoted by  $U$ , there exists a set  $\mathcal{W}_U$  and a cost function  $c_U : \mathcal{X} \times \mathcal{W}_U \rightarrow \mathbb{R}_+$  such that  $\Lambda_U(X \rightarrow y) = \Lambda_{c_U}(X \rightarrow y)$ . Conversely, for every cost function  $c : \mathcal{X} \times \mathcal{W} \rightarrow \mathbb{R}_+$ , there exists a randomized function of  $X$ , denoted by  $U_c$ , such that  $\Lambda_c(X \rightarrow y) = \Lambda_{U_c}(X \rightarrow y)$ .

**Corollary 1** (Cost function view of PMC). By Theorem 2, PMC can alternatively be defined as

$$\Lambda_{P_{XY}}(X \rightarrow y) := \sup_c \Lambda_c(X \rightarrow y),$$

where the supremum is over all non-negative cost functions with finite range.

### C. Relationship to Maximal Cost Leakage and Maximal Realizable Cost

In [22], Issa et al. defined two notions of information leakage that also correspond to risk-averse adversaries. The first notion, termed *maximal cost leakage*, quantifies the average information leaking through a mechanism. The second notion, termed *maximal realizable cost*, describes the largest amount of information leakage across all outcomes of a mechanism.

**Definition 8** (Maximal cost leakage [22, Def. 11]). Given a joint distribution  $P_{XY}$  on the finite set  $\mathcal{X} \times \mathcal{Y}$ , the maximal cost leakage from  $X$  to  $Y$  is defined as

$$\mathcal{L}^c(X \rightarrow Y) := \sup_{\substack{U: U-X-Y, \\ \mathcal{W}, c: U \times \mathcal{W} \rightarrow \mathbb{R}_+}} \log \frac{\inf_{w \in \mathcal{W}} \mathbb{E}[c(U, w)]}{\inf_{\hat{w}(\cdot)} \mathbb{E}[c(U, \hat{w}(Y))]}, \quad (8)$$

where  $U$  takes values in an arbitrary finite set  $\mathcal{U}$ , and the infimum in the denominator is over all functions  $\hat{w} : \mathcal{Y} \rightarrow \mathcal{W}$ .

Issa et al. [22, Thm. 15] then showed that maximal cost leakage takes the simple form

$$\mathcal{L}^c(X \rightarrow Y) = -\log \sum_{y \in \mathcal{Y}} \min_{x \in \mathcal{X}} P_{Y|X=x}(y).$$

The general framework of maximal cost leakage closely resembles the cost function perspective of PMC outlined in Section III-B. Similarly to PMC, Definition 8 compares the adversary's minimal prior cost with her minimal posterior cost. However, there are also two distinctions: Maximal cost leakage is defined for the average outcome of  $Y$  and evaluates the costs of guessing randomized functions of  $X$ . In contrast, PMC is defined for each specific outcome  $y \in \mathcal{Y}$  and evaluates the costs associated with guessing  $X$  itself.

Earlier in Remark 2 we argued that PMC is closed under pre-processing; thus, it suffices to consider the costs of guessing  $X$  itself. In a similar vein, below we demonstrate that when evaluating the information leaking to the average outcome of  $Y$  as in maximal cost leakage, it suffices to analyze the costs associated with guessing  $X$  alone. Specifically, we show that the supremum over  $U$ 's satisfying the Markov chain  $U - X - Y$  in (8) is superfluous.

**Proposition 2.** Given a joint distribution  $P_{XY}$  on the finite set  $\mathcal{X} \times \mathcal{Y}$ , the maximal cost leakage can alternatively be defined as

$$\mathcal{L}^c(X \rightarrow Y) := \sup_{\mathcal{W}, c: \mathcal{X} \times \mathcal{W} \rightarrow \mathbb{R}_+} \log \frac{\inf_{w \in \mathcal{W}} \mathbb{E}[c(X, w)]}{\inf_{\hat{w}(\cdot)} \mathbb{E}[c(X, \hat{w}(Y))]}.$$

In other words, the supremum over  $U$ 's satisfying the Markov chain  $U - X - Y$  in (8) is superfluous.

*Proof:* Let

$$L_1 := \sup_{\substack{U: U-X-Y, \\ \mathcal{W}, c: \mathcal{U} \times \mathcal{W} \rightarrow \mathbb{R}_+}} \log \frac{\inf_{w \in \mathcal{W}} \mathbb{E}[c(U, w)]}{\inf_{\hat{w}(\cdot)} \mathbb{E}[c(U, \hat{w}(Y))]},$$

and

$$L_2 := \sup_{\mathcal{W}, c: \mathcal{X} \times \mathcal{W} \rightarrow \mathbb{R}_+} \log \frac{\inf_{w \in \mathcal{W}} \mathbb{E}[c(X, w)]}{\inf_{\hat{w}(\cdot)} \mathbb{E}[c(X, \hat{w}(Y))]}.$$

Our goal is to show that  $L_1 = L_2$ . Clearly,  $L_1 \geq L_2$  so we show that  $L_2 \geq L_1$ .

Fix a random variable  $U$  satisfying the Markov chain  $U - X - Y$  and let  $\mathcal{U}$  denote the alphabet of  $U$ . Let  $c_1 : \mathcal{U} \times \mathcal{W} \rightarrow \mathbb{R}_+$  be a cost function, where  $\mathcal{W}$  is a finite set. Consider the cost function  $c_2 : \mathcal{X} \times \mathcal{W} \rightarrow \mathbb{R}_+$  defined as

$$c_2(x, w) = \sum_{u \in \mathcal{U}} P_{U|X=x}(u) c_1(u, w) = \mathbb{E}[c_1(U, w) | X = x].$$

For all  $w \in \mathcal{W}$ , we have

$$\mathbb{E}[c_1(U, w)] = \mathbb{E}_X[\mathbb{E}[c_1(U, w) | X]] = \mathbb{E}_X[c_2(X, w)].$$

Similarly, given a function  $\hat{w} : \mathcal{Y} \rightarrow \mathcal{W}$  it holds that

$$\mathbb{E}[c_1(U, \hat{w}(Y))] = \sum_{y \in \mathcal{Y}} P_Y(y) \mathbb{E}[c_1(U, \hat{w}(y)) | Y = y]$$

$$\begin{aligned}
&= \sum_{y \in \mathcal{Y}} P_Y(y) \sum_{x \in \mathcal{X}} P_{X|Y=y}(x) \sum_{u \in \mathcal{U}} P_{U|X=x, Y=y}(u) c_1(u, \hat{w}(y)) \\
&= \sum_{y \in \mathcal{Y}} P_Y(y) \sum_{x \in \mathcal{X}} P_{X|Y=y}(x) \sum_{u \in \mathcal{U}} P_{U|X=x}(u) c_1(u, \hat{w}(y)) \\
&= \sum_{y \in \mathcal{Y}} P_Y(y) \sum_{x \in \mathcal{X}} P_{X|Y=y}(x) \mathbb{E} \left[ c_1(U, \hat{w}(y)) \mid X = x \right] \\
&= \sum_{y \in \mathcal{Y}} P_Y(y) \sum_{x \in \mathcal{X}} P_{X|Y=y}(x) c_2(x, \hat{w}(y)) \\
&= \mathbb{E} \left[ c_2(X, \hat{w}(Y)) \right],
\end{aligned}$$

where the third equality is due to the Markov chain  $U - X - Y$ . Hence, for each random variable  $U$  satisfying the Markov chain  $U - X - Y$  and cost function  $c_1 : \mathcal{U} \times \mathcal{W} \rightarrow \mathbb{R}_+$ , there exists a cost function  $c_2 : \mathcal{X} \times \mathcal{W} \rightarrow \mathbb{R}_+$  such that

$$\frac{\inf_{w \in \mathcal{W}} \mathbb{E} \left[ c_1(U, w) \right]}{\inf_{\hat{w}(\cdot)} \mathbb{E} \left[ c_1(U, \hat{w}(Y)) \right]} = \frac{\inf_{w \in \mathcal{W}} \mathbb{E} \left[ c_2(X, w) \right]}{\inf_{\hat{w}(\cdot)} \mathbb{E} \left[ c_2(X, \hat{w}(Y)) \right]}.$$

This implies that  $L_1 \leq L_2$ , as desired.  $\blacksquare$

It is also worth noting that maximal cost leakage does *not* correspond to the expected value of the PMC. Using Corollary 1 we have

$$\begin{aligned}
\mathcal{L}^c(X \rightarrow Y) &= \sup_{U:U-X-Y} \left( -\log \mathbb{E}_{Y \sim P_Y} \left[ \exp \left( -\Lambda_{P_{UY}}(U \rightarrow Y) \right) \right] \right) \\
&\leq \sup_{U:U-X-Y} \mathbb{E}_{Y \sim P_Y} \left[ \Lambda_{P_{UY}}(U \rightarrow Y) \right] \tag{9a}
\end{aligned}$$

$$\begin{aligned}
&\leq \mathbb{E}_{Y \sim P_Y} \left[ \sup_{U:U-X-Y} \Lambda_{P_{UY}}(U \rightarrow Y) \right] \\
&= \mathbb{E}_{Y \sim P_Y} \left[ \Lambda_{P_{XY}}(X \rightarrow Y) \right], \tag{9b}
\end{aligned}$$

where (9a) is due to Jensen's inequality, and (9b) follows from the closedness of PMC under pre-processing. Note that the strict concavity of  $\log(\cdot)$  implies that Jensen's inequality is strict. Consequently, maximal cost leakage does not describe the expected value of PMC but underestimates it.

**Definition 9** (Maximal realizable cost [22, Def. 12]). Given a joint distribution  $P_{XY}$  on the finite set  $\mathcal{X} \times \mathcal{Y}$ , the maximal realizable cost from  $X$  to  $Y$  is defined as

$$\mathcal{L}^{rc}(X \rightarrow Y) := \sup_{\substack{U:U-X-Y, \\ \mathcal{W}, c: \mathcal{U} \times \mathcal{W} \rightarrow \mathbb{R}_+}} \log \frac{\inf_{w \in \mathcal{W}} \mathbb{E} \left[ c(U, w) \right]}{\min_{y \in \mathcal{Y}} \inf_{\hat{w} \in \mathcal{W}} \mathbb{E} \left[ c(U, \hat{w}) \mid Y = y \right]}, \tag{10}$$

where  $U$  takes values in an arbitrary finite set  $\mathcal{U}$ .

Once again, the threat model of maximal realizable cost is similar to that of PMC. The difference is that maximal realizable cost is defined for the worst outcome of  $Y$  and evaluates the costs of guessing randomized functions of  $X$ .



Issa et al. [22, Thm. 16] showed that maximal realizable cost can be expressed as

$$\mathcal{L}^{\text{rc}}(X \rightarrow Y) = D_\infty(P_X \times P_Y \| P_{XY}).$$

Therefore, the maximal realizable cost is related to PMC by

$$\begin{aligned} \mathcal{L}^{\text{rc}}(X \rightarrow Y) &= \max_{y \in \mathcal{Y}} \sup_{U: U-X-Y} \Lambda_{P_{UY}}(U \rightarrow y) \\ &= \max_{y \in \mathcal{Y}} \Lambda_{P_{XY}}(X \rightarrow y), \end{aligned}$$

where the second equality is due to the closedness of PMC under pre-processing. This again implies that the supremum over  $U$ 's in (10) is not required (cf. Proposition 2).

In general, defining an information measure for each outcome  $y \in \mathcal{Y}$  as done by PMC has the advantage that  $\Lambda_{P_{XY}}(X \rightarrow Y)$  may be viewed as a random variable that can be restricted in various ways. Hence, the resulting privacy analysis is not restricted to computing the average or maximum leakages but can be tailored to application-specific requirements, such as restricting the tail of  $\Lambda_{P_{XY}}(X \rightarrow Y)$  or its different moments. Consequently, PMC is rendered a flexible tool that is applicable across a wider spectrum of scenarios. The flexibility of PMC will become more apparent when we discuss concrete examples in Section IV-B.

#### D. Relationship to Maximal Guesswork Leakage

A central example of a cost function is the expected number of attempts (or moments thereof) required to correctly guess a secret [49, 50]. In this context, a guessing function  $G$  is a one-to-one mapping from  $\mathcal{X}$  to  $[[\mathcal{X}]]$ , and the *guesswork* is the minimum expected number of attempts required to correctly identify  $X$ , i.e.,  $\min_G \mathbb{E}[G(X)]$ . Recently, guesswork has been used to define two information measures called *maximal guesswork leakage* and *pointwise maximal guesswork leakage* [23]. Both measures describe adversaries aiming to minimize the guesswork associated with randomized functions of  $X$  and compare the prior guesswork with the posterior guesswork. Per usual terminology, maximal guesswork leakage describes the average amount of information leaking through a mechanism, while pointwise maximal guesswork leakage is defined for each outcome  $y \in \mathcal{Y}$ .

Here, we focus on pointwise maximal guesswork leakage and its connections to PMC. Similar discussions apply to maximal guesswork leakage.

**Definition 10** (Pointwise maximal guesswork leakage [23, Def. 6]). Given a joint distribution  $P_{XY}$  on the finite set  $\mathcal{X} \times \mathcal{Y}$ , the pointwise maximal guesswork leakage from  $X$  to  $y \in \mathcal{Y}$  is defined as

$$\mathcal{L}^{\text{G-pw}}(X \rightarrow y) := \sup_{U: U-X-Y} \log \frac{\min_G \mathbb{E}[G(U)]}{\min_{\hat{G}} \mathbb{E}[\hat{G}(U) | Y = y]},$$

where  $U$  takes values in an arbitrary finite alphabet.

Pointwise maximal guesswork leakage is a special case of the cost function model of PMC. In particular, given a random variable  $U$  satisfying the Markov chain  $U - X - Y$  and a guessing function  $G: \mathcal{U} \rightarrow [[\mathcal{U}]]$ , there exists a cost function  $d: \mathcal{U} \times \mathcal{W} \rightarrow [[\mathcal{U}]]$  such that

$$\min_G \mathbb{E}[G(U)] = \min_{w \in \mathcal{W}} \mathbb{E}[d(U, w)],$$

$$\min_{\hat{G}} \mathbb{E}[\hat{G}(U) | Y = y] = \min_{\hat{w} \in \mathcal{W}} \mathbb{E}[d(U, \hat{w}) | Y = y],$$

(see [23, Sec. III] for the exact construction). Furthermore, we showed in (the proof of) Proposition 2 that for each  $U$  satisfying the Markov chain  $U - X - Y$  and each cost function  $d : \mathcal{U} \times \mathcal{W} \rightarrow \mathbb{R}_+$ , there exists a cost function  $c : \mathcal{X} \times \mathcal{W} \rightarrow \mathbb{R}_+$  such that

$$\begin{aligned} \mathbb{E}[d(U, w)] &= \mathbb{E}[c(X, w)], \\ \mathbb{E}[d(U, w) | Y = y] &= \mathbb{E}[c(X, w) | Y = y], \end{aligned}$$

for all  $w \in \mathcal{W}$ . This argument shows that  $\mathcal{L}^{\text{G-pw}}(X \rightarrow y) \leq \Lambda(X \rightarrow y)$ . Interestingly, Kurri et al. [23] proved that the inequality is in fact an equality, i.e.,

$$\mathcal{L}^{\text{G-pw}}(X \rightarrow y) = \Lambda(X \rightarrow y) = D_\infty(P_X \| P_{X|Y=y}), \quad y \in \mathcal{Y}.$$

### E. PMC on General Alphabets

Now, we extend the cost function model described earlier to a much broader class of secrets. Suppose  $X$  is defined on a measurable space  $(\mathcal{X}, \mathcal{S}_X)$ , and  $Y$  is defined on a measurable space  $(\mathcal{Y}, \mathcal{S}_Y)$ . Let  $P_{XY}$ , denote the joint distribution of  $X$  and  $Y$ .

**Definition 11.** Given a joint distribution  $P_{XY}$  on the product space  $(\mathcal{X} \times \mathcal{Y}, \mathcal{S}_X \otimes \mathcal{S}_Y)$  the pointwise maximal cost from  $X$  to  $y \in \mathcal{Y}$  is defined as

$$\Lambda_{P_{XY}}(X \rightarrow y) := \log \sup_{\substack{(\mathcal{W}, \mathcal{S}_W), \\ c \in \mathcal{C}}} \frac{\inf_{P_W} \mathbb{E}[c(X, W)]}{\inf_{P_{W|Y}} \mathbb{E}[c(X, \hat{W}) | Y = y]}, \quad (11)$$

where the infimum in the denominator is over all transition probability kernels from  $(\mathcal{Y}, \mathcal{S}_Y)$  into  $(\mathcal{W}, \mathcal{S}_W)$ , and  $\mathcal{C}$  denotes the set of all cost functions defined as

$$\mathcal{C} := \left\{ c \in (\mathcal{S}_X \otimes \mathcal{S}_W)_+ \mid \inf_{P_W} \mathbb{E}[c(X, W)] < \infty \right\}.$$

Definition 11 describes a threat model similar to the cost function model in Section III-B. It represents an adversary who aims to construct an estimate of  $X$ , denoted by  $W$ , minimizing the expected value of a cost function. Here,  $W$  is a random variable on a measurable space  $(\mathcal{W}, \mathcal{S}_W)$ , and cost functions are selected from the collection  $\mathcal{C}$ . To quantify the information revealed about  $X$  by disclosing an outcome  $y \in \mathcal{Y}$ , we compare the minimum expected cost before observing  $y$  in the numerator of (11) with the minimum expected cost after observing  $y$  in the denominator. We then take the supremum of this ratio across all measurable spaces  $(\mathcal{W}, \mathcal{S}_W)$  and all  $c \in \mathcal{C}$  to obtain a general and robust definition that does not depend on specific instances of  $c$ . Note that the condition  $\inf_{P_W} \mathbb{E}[c(X, W)] < \infty$  ensures that (11) is well-defined since we exclude cost functions that result in infinite minimum costs both in the numerator and the denominator.

Before simplifying (11), let us give two concrete examples of privacy attacks encompassed by Definition 11.

**Example 1** (Minimizing  $L_p$ -distance). A key objective in statistics is to produce an estimate of a random quantity with the smallest mean squared error. More generally, suppose  $X$  is a real-valued random variable with distribution  $P_X$ , let  $p \geq 1$ , and suppose  $\mathbb{E}[|X|^p] < \infty$ . The  $L_p$ -norm of  $X$  is

$$\|X\|_p = \left( \int_{\mathbb{R}} |x|^p P_X(dx) \right)^{\frac{1}{p}}.$$

The  $L_p$ -norm defines a metric space through the metric  $d(X, W) := \|X - W\|_p$ , where  $W$  is also a real-valued random variable.

In this context, an adversary may aim to construct an estimate of  $X$ , denoted by  $W$ , minimizing the  $L_p$ -distance to  $X$ . Using the notation of Definition 11, this attack can be modeled by taking  $\mathcal{W} = \bar{\mathbb{R}}$  and  $\mathcal{S}_{\mathcal{W}} = \mathcal{B}_{\bar{\mathbb{R}}}$ . Then, the cost function  $c_d(X, W) = |X - W|^p$ ,  $c_d \in \mathcal{C}$  can be employed to capture the adversary's objective.

**Example 2** (Minimizing  $\alpha$ -loss). Let  $\mathcal{X}$  be a finite set with cardinality  $k$ , and let  $\Delta^{k-1}$  denote the set of all distributions on  $\mathcal{X}$ , i.e., the  $(k-1)$ -dimensional probability simplex. Liao et al. [31] introduced a class of loss functions called  $\alpha$ -loss with  $\alpha \in [1, \infty]$  that interpolate between the log-loss at  $\alpha = 1$  and the 0-1 loss at  $\alpha = \infty$ . More precisely, they defined the  $\alpha$ -loss  $c_\alpha : \mathcal{X} \times \Delta^{k-1} \rightarrow \bar{\mathbb{R}}_+$  by

$$c_\alpha(x, Q) := \begin{cases} \frac{\alpha}{\alpha-1} \left( 1 - Q(x)^{\frac{\alpha-1}{\alpha}} \right) & \text{if } \alpha \in (1, \infty), \\ \log \frac{1}{Q(x)} & \text{if } \alpha = 1, \\ 1 - Q(x) & \text{if } \alpha = \infty, \end{cases}$$

where  $Q \in \Delta^{k-1}$  is a distribution on  $\mathcal{X}$ . Then, they used the  $\alpha$ -loss to define a class of tunable information leakage measure called (*maximal*)  $\alpha$ -leakage [31, Def. 5 and 6].<sup>2</sup>

Notably, adversaries minimizing the expected value of an  $\alpha$ -loss are incorporated into the threat model of Definition 11. To illustrate this, let  $\mathcal{W} = \Delta^{k-1}$ , and let  $\mathcal{S}_{\mathcal{W}}$  be the Borel  $\sigma$ -algebra on  $\Delta^{k-1}$ . Given  $\alpha \in [1, \infty]$ , we may use the cost function  $c_\alpha(X, W) \in \mathcal{C}$  to capture the adversary's objective. More generally, we can extend this model to encompass adversaries who minimize the  $\alpha$ -loss associated with randomized functions of  $X$ .

Now, we show that PMC on general alphabets also reduces to  $D_\infty(P_X \| P_{X|Y=y})$ . This result is proved in Appendix B.

**Theorem 3.** Let  $P_{XY}$  be the joint distribution of  $X$  and  $Y$ . Given  $y \in \mathcal{Y}$ , PMC as described by Definition 11 simplifies to

$$\Lambda_{P_{XY}}(X \rightarrow y) = D_\infty(P_X \| P_{X|Y=y}).$$

Henceforth, we drop the subscript  $P_{XY}$  in  $\Lambda_{P_{XY}}(X \rightarrow y)$  and  $i_{P_{XY}}$  when the joint distribution is clear from the context.

<sup>2</sup>Note that (*maximal*)  $\alpha$ -leakage assumes that the adversary maximizes an objective function such that the optimal action corresponds to minimizing the  $\alpha$ -loss. In contrast, here we assume that the adversary directly minimizes the  $\alpha$ -loss.

**Remark 4** (Measurability of PMC). We discussed earlier that one of the advantages of PMC over similar existing notions is its flexibility, as  $\Lambda(X \rightarrow Y)$  is a random variable whose various statistical properties can be examined and controlled. For this to work, the mapping  $y \mapsto \Lambda_{P_{XY}}(X \rightarrow y)$  must be measurable. Note that the assumption that  $(\mathcal{X}, \mathcal{S}_X)$  and  $(\mathcal{Y}, \mathcal{S}_Y)$  are standard Borel spaces is sufficient for this purpose (see Section II-A). Consequently, assuming that  $P_X \times P_Y \ll P_{XY}$ , we can express PMC by

$$\Lambda(X \rightarrow y) = \operatorname{ess\,sup}_{P_X} \left( -i(X; y) \right).$$

Let  $\mu$  and  $\nu$  be dominating  $\sigma$ -finite measures on  $(\mathcal{X}, \mathcal{S}_X)$  and  $(\mathcal{Y}, \mathcal{S}_Y)$ , respectively. We can also express PMC using densities:

$$\begin{aligned} \Lambda(X \rightarrow y) &= \log \left( \operatorname{ess\,sup}_{P_X} \frac{f_X(X)}{f_{X|Y}(X|y)} \right) \\ &= \log \left( \operatorname{ess\,sup}_{P_X} \frac{f_Y(y)}{f_{Y|X}(y|X)} \right), \end{aligned}$$

where  $f_{X|Y} \in (\mathcal{S}_X \otimes \mathcal{S}_Y)_+$  is the density of  $P_{X|Y}$  with respect to  $\mu$ , and  $f_X \in \mathcal{S}_{X+}$  is the density of  $P_X$  with respect to  $\mu$ . The densities  $f_{Y|X}$  and  $f_Y$  are defined similarly.

Having settled the measurability of PMC, we say that a privacy mechanism  $P_{Y|X}$  satisfies  $\varepsilon$ -PMC if

$$\mathbb{P}\{\Lambda(X \rightarrow Y) \leq \varepsilon\} = 1,$$

where  $\varepsilon \geq 0$ . This represents the strongest type of guarantee that can be defined based on PMC, as it requires  $\Lambda(X \rightarrow Y)$  to be essentially bounded.

#### IV. PROPERTIES OF PMC AND EXAMPLES

This section focuses on two main topics. First, we explore the key properties of PMC, demonstrating that it satisfies the usual criteria expected from an information measure, such as non-negativity and data-processing inequalities. Next, we examine some prototypical examples in privacy, including the randomized response mechanism [24, 25] applied to finite secrets, and the Laplace mechanism [5] for releasing the sample mean of a dataset. In all examples, we calculate, analyze, and discuss the PMC.

##### A. Properties of Pointwise Maximal Cost

Before exploring the properties of PMC, we introduce a conditional version of PMC. This definition measures the amount of information leaked to an adversary who possesses some side information about the secret already before observing the mechanism's outcome. The side information is modeled as the outcome of a random variable  $Z$  that is correlated with both  $X$  and  $Y$ . Conditional PMC is particularly useful for deriving an upper bound on the leakage when multiple observations are available.

**Definition 12** (Conditional PMC). Let  $Z$  be a random variable on a measurable space  $(\mathcal{Z}, \mathcal{S}_Z)$ . Given a transition probability kernel  $P_{XY|Z}$  from  $(\mathcal{Z}, \mathcal{S}_Z)$  into  $(\mathcal{X} \times \mathcal{Y}, \mathcal{S}_X \otimes \mathcal{S}_Y)$ , the conditional pointwise maximal cost from  $X$  to  $y \in \mathcal{Y}$  given  $z \in \mathcal{Z}$  is defined as

$$\Lambda_{P_{XY|Z}}(X \rightarrow y | z) := \log \sup_{\substack{(W, \mathcal{S}_W), \\ c \in \mathcal{C}_z}} \frac{\inf_{P_{W|Z}} \mathbb{E} [c(X, W) | Z = z]}{\inf_{P_{\hat{W}|Y, Z}} \mathbb{E} [c(X, \hat{W}) | Y = y, Z = z]},$$

where  $\mathcal{C}_z$  denotes the set of all cost functions defined as

$$\mathcal{C}_z := \left\{ c \in (\mathcal{S}_X \otimes \mathcal{S}_W)_+ \mid \inf_{P_{W|Z}} \mathbb{E} [c(X, W) | Z = z] < \infty \right\}.$$

Technically, Definition 12 does not introduce a new concept; rather it is an adaptation of Definition 11 where all distributions are conditioned on  $Z = z$ . Thus, conditional PMC describes a similar threat model to PMC, with the key difference being the additional side information  $Z = z$ . Writing  $P_{XY|Z} = P_{Y|X, Z} \times P_{X|Z}$ , the outcome  $z \in \mathcal{Z}$  essentially parameterizes the distribution of  $X$ , denoted by  $P_{X|Z}$ , and the information release mechanism  $P_{Y|X, Z}$ . As such, by applying Theorem 3, we directly obtain a simplified form for the conditional PMC.

**Corollary 2.** Let  $P_{XY|Z}$  be a transition probability kernel from  $(\mathcal{Z}, \mathcal{S}_Z)$  into  $(\mathcal{X} \times \mathcal{Y}, \mathcal{S}_X \otimes \mathcal{S}_Y)$ . The conditional pointwise maximal cost from  $X$  to  $y \in \mathcal{Y}$  given  $z \in \mathcal{Z}$  can be expressed as

$$\Lambda_{P_{XY|Z}}(X \rightarrow y | z) = D_\infty(P_{X|Z=z} \| P_{X|Y=y, Z=z}).$$

As a special case, if the Markov chain  $Z - X - Y$  holds, then  $\Lambda_{P_{XY|Z}}(X \rightarrow y | z) = D_\infty(P_{X|Z=z} \| P_{X|Y=y})$ . This scenario corresponds to the important case where the side information determines the adversary's prior knowledge about  $X$  but does not influence the mechanism through which information is released.

Now, we discuss the properties of PMC. Most properties are consequences of the fact that PMC is a Rényi divergence between two distributions. The following lemma confirms that PMC satisfies all the usual properties expected from an information measure, e.g., non-negativity, data-processing inequalities, and additivity.

**Lemma 1** (Properties of PMC). The pointwise maximal cost satisfies the following properties:

1. (Non-negativity).  $\Lambda(X \rightarrow y) \geq 0$  for all  $y \in \mathcal{Y}$ .
2. (Independence). If  $X$  and  $Y$  are independent, then  $\Lambda(X \rightarrow Y) = 0$  almost surely.
3. (Additivity). If  $P_{Y_1, \dots, Y_k | X_1, \dots, X_k} = \prod_{i=1}^k P_{Y_i | X_i}$  and  $P_{X_1, \dots, X_k} = \prod_{i=1}^k P_{X_i}$ , then

$$\Lambda(X_1, \dots, X_k \rightarrow Y_1, \dots, Y_k) = \sum_{i=1}^k \Lambda(X_i \rightarrow Y_i).$$

4. (Concavity). Given  $y \in \mathcal{Y}$ ,  $\Lambda(X \rightarrow y)$  is concave in  $P_X$ .
5. (Pre-processing). Suppose the Markov chain  $Z - X - Y$  holds. Then, for all  $y \in \mathcal{Y}$  it holds that

$$\Lambda(Z \rightarrow y) \leq \Lambda(X \rightarrow y).$$

6. (Post-processing). Suppose the Markov chain  $X - Y - Z$  holds. Then, we have

$$\operatorname{ess\,sup}_{P_Z} \Lambda(X \rightarrow Z) \leq \operatorname{ess\,sup}_{P_Y} \Lambda(X \rightarrow Y).$$

7. (Composition). Given a joint distribution  $P_{XY_1Y_2}$ , it holds that

$$\Lambda_{P_{XY_1Y_2}}(X \rightarrow y_1, y_2) \leq \Lambda_{P_{XY_1}}(X \rightarrow y_1) + \Lambda_{P_{XY_2|Y_1}}(X \rightarrow y_2 | y_1),$$

for all  $y_1 \in \mathcal{Y}_1$  and  $y_2 \in \mathcal{Y}_2$ .

Lemma 1 is proved in Appendix D.

### B. Examples

In this section, we study PMC in several examples. By calculating PMC for commonly used mechanisms, such as the randomized response mechanism [24, 25] and the Laplace mechanism [5, 6], we demonstrate how PMC can be effectively used to quantify privacy.

**Example 3** (Generalized Randomized Response [24, 25]). A common mechanism used to achieve local differential privacy is the (generalized) randomized response mechanism. It is applied when the secret  $X$  is a finite random variable and produces a privatized version of  $X$  while retaining its original alphabet.

Suppose  $\mathcal{X} = \mathcal{Y} = [n]$ . The randomized response mechanism with parameter  $\varepsilon_r \geq 0$  can be expressed as follows:

$$P_{Y|X=i}(j) = \begin{cases} \frac{e^{\varepsilon_r}}{n-1+e^{\varepsilon_r}} & \text{if } i = j, \\ \frac{1}{n-1+e^{\varepsilon_r}} & \text{if } i \neq j. \end{cases}$$

It is easy to verify that the randomized response mechanism satisfies  $\varepsilon_r$ -LDP.

Now, we calculate the PMC of the randomized response mechanism. For all  $j \in [n]$ , we have  $\min_{i \in [n]} P_{Y|X=i}(j) = \frac{1}{n-1+e^{\varepsilon_r}}$  and

$$P_Y(j) = \sum_{i \in [n]} P_{Y|X=i}(j) P_X(i) = \frac{1 + P_X(j)(e^{\varepsilon_r} - 1)}{n - 1 + e^{\varepsilon_r}}.$$

Thus, the PMC associated with outcome  $j \in [n]$  of the mechanism is

$$\Lambda(X \rightarrow j) = \log \max_{i \in [n]} \frac{P_Y(j)}{P_{Y|X=i}(j)} = \log \left( 1 + P_X(j)(e^{\varepsilon_r} - 1) \right),$$

implying that the randomized response mechanism satisfies  $\varepsilon$ -PMC with

$$\varepsilon = \max_{j \in [n]} \Lambda(X \rightarrow j) = \log \left( 1 + \max_{j \in [n]} P_X(j)(e^{\varepsilon_r} - 1) \right).$$

Note that  $\varepsilon \leq \varepsilon_r$  for all distributions  $P_X$ . Interestingly,  $\varepsilon \rightarrow \varepsilon_r$  as  $\max_{j \in [n]} P_X(j) \rightarrow 1$ .

**Example 4** (PML-extremal mechanism in the high-privacy regime [62]). Suppose  $X$  is a finite random variable with alphabet  $\mathcal{X} = [n]$ . Recall from Section II-B that PML-extremal mechanisms constitute a class of optimal mechanisms maximizing sub-convex utility functions [62, Def. 5]. Let  $0 \leq \varepsilon_u < \log \frac{1}{1-p_{\min}}$ . This range of values for  $\varepsilon_u$  corresponds to the *high-privacy regime*. Extremal mechanisms in the high-privacy regime can be expressed as

$$P_{Y|X=i}(j) = \begin{cases} 1 - e^{\varepsilon_u}(1 - P_X(j)) & \text{if } i = j, \\ e^{\varepsilon_u} P_X(j) & \text{if } i \neq j, \end{cases}$$

where  $i, j \in [n]$ . It can be verified that these mechanisms satisfy  $\varepsilon_u$ -PML.

Now, we calculate the PMC of the extremal mechanisms. We have  $\min_{i \in [n]} P_{Y|X=i}(j) = 1 - e^{\varepsilon_u}(1 - P_X(j))$  and it is easy to check that  $P_Y(j) = P_X(j)$  for all  $j \in [n]$ . Thus, the PMC associated with outcome  $j \in [n]$  of the mechanism is

$$\Lambda(X \rightarrow j) = \log \max_{i \in [n]} \frac{P_Y(j)}{P_{Y|X=i}(j)} = \log \frac{P_X(j)}{1 - e^{\varepsilon_u}(1 - P_X(j))}.$$

Since the mapping  $p \mapsto \log \frac{p}{1 - e^{\varepsilon_u}(1 - p)}$  is decreasing in  $p$ , the extremal mechanism satisfies  $\varepsilon$ -PMC with

$$\varepsilon = \max_{j \in [n]} \Lambda(X \rightarrow j) = \log \frac{p_{\min}}{1 - e^{\varepsilon_u}(1 - p_{\min})}.$$

Next, we examine the Laplace mechanism [5] which is a central component of many (central) differential privacy algorithms.

**Example 5** (Laplace mechanism [5, 6]). A central problem in statistics is to estimate the expected value of a random variable based on multiple independently drawn realizations. This problem is also extensively explored under the framework of differential privacy. In particular, when aiming to satisfy pure differential privacy, the Laplace mechanism is often employed which adds noise with Laplace distribution to the sample mean. Importantly, to release the sample mean via the Laplace mechanism with DP guarantees the data domain must be bounded, e.g., an interval. This is because the global (and local) sensitivity of the sample mean is infinite when the data domain is unbounded, e.g.,  $\mathbb{R}$  [5].

Consider an i.i.d database  $X^n = (X_1, \dots, X_n)$ , where each  $X_i$  is a bounded random variable on the interval  $[c, d]$ . Suppose  $\mathbb{E}[X_i] = \mu \in [c, d]$ , and define  $Z_1 := X_1 - \mu$ . Let  $K(t) := \log \mathbb{E}[e^{tZ_1}]$  denote the cumulant generating function of  $Z_1$ , where  $t \in \mathbb{R}$ . Suppose our objective is to safely disclose the sample mean of  $X^n$  by perturbing it with Laplace noise. Let  $\text{Lap}(m, b)$  denote Laplace distribution with mean  $m \in \mathbb{R}$  and scale parameter  $b > 0$ . The Laplace mechanism releases an outcome with the conditional distribution  $Y | X^n = x^n \sim \text{Lap}\left(\frac{\sum_{i=1}^n x_i}{n}, b\right)$ .

Below, we evaluate the PMC from each data point to the released outcome. Due to the symmetry of the problem, the Laplace mechanism leaks the same amount of information about all data points, and for notational convenience, we calculate  $\Lambda(X_n \rightarrow y)$  with  $y \in \mathbb{R}$ . Our analysis in this example complements a similar analysis performed with the Laplace mechanism and PML in [21, Prop. 4.5].

The largest privacy cost across all possible released outcomes is

$$\begin{aligned} \sup_{y \in \mathbb{R}} \Lambda(X_n \rightarrow y) &= \log \sup_{y \in \mathbb{R}} \frac{f_Y(y)}{\inf_{x_n \in [c, d]} f_{Y|X_n=x_n}(y)} \\ &= \log \sup_{y \in \mathbb{R}} \frac{\frac{1}{2b} \mathbb{E}_{X^n} \left[ \exp \left( -\frac{|y - \frac{\sum_{i=1}^n X_i}{n}|}{b} \right) \right]}{\inf_{x_n \in [c, d]} \frac{1}{2b} \mathbb{E}_{X^{n-1}} \left[ \exp \left( -\frac{|y - \frac{x_n}{n} - \frac{\sum_{i=1}^{n-1} X_i}{n}|}{b} \right) \right]}. \end{aligned} \quad (12)$$

First, we argue that to find the largest PMC across the  $y$ 's, it suffices to consider  $y \geq d$  and  $y \leq c$ . To see why, note that in the numerator, using the fact that  $-|a| \leq \min\{-a, a\}$  for all  $a \in \mathbb{R}$  we have

$$\mathbb{E}_{X^n} \left[ \exp \left( -\frac{|y - \frac{\sum_{i=1}^n X_i|}{n}|}{b} \right) \right] \leq \min \left\{ \mathbb{E}_{X^n} \left[ \exp \left( -\frac{y - \frac{\sum_{i=1}^n X_i}{n}}{b} \right) \right], \mathbb{E}_{X^n} \left[ \exp \left( \frac{y - \frac{\sum_{i=1}^n X_i}{n}}{b} \right) \right] \right\},$$

for all  $y \in \mathbb{R}$ . Moreover, in the denominator, the mapping  $y \mapsto \mathbb{E}_{X^{n-1}} \left[ \exp \left( -\frac{|y - \frac{x_n}{n} - \frac{\sum_{i=1}^{n-1} X_i|}{n}}{b} \right) \right]$  is increasing in  $(-\infty, c]$  and decreasing in  $[d, \infty)$  for all  $x_n \in [c, d]$ .

Suppose  $y \geq d$ . The expectation in the numerator of (12) is

$$\begin{aligned} \mathbb{E}_{X^n} \left[ \exp \left( -\frac{y - \frac{\sum_{i=1}^n X_i}{n}}{b} \right) \right] &= \exp \left( -\frac{y}{b} \right) \mathbb{E}_{X^n} \left[ \exp \left( \frac{\sum_{i=1}^n X_i}{nb} \right) \right] \\ &= \exp \left( -\frac{y}{b} \right) \prod_{i=1}^n \mathbb{E} \left[ \exp \left( \frac{X_i}{nb} \right) \right] \\ &= \exp \left( -\frac{y}{b} \right) \left( \mathbb{E} \left[ \exp \left( \frac{X_1}{nb} \right) \right] \right)^n, \end{aligned}$$

and the expectation in the denominator is

$$\begin{aligned} \mathbb{E}_{X^{n-1}} \left[ \exp \left( -\frac{y - \frac{x_n}{n} - \frac{\sum_{i=1}^{n-1} X_i}{n}}{b} \right) \right] &= \exp \left( -\frac{y}{b} \right) \cdot \exp \left( \frac{x_n}{nb} \right) \mathbb{E}_{X^{n-1}} \left[ \exp \left( \frac{\sum_{i=1}^{n-1} X_i}{nb} \right) \right] \\ &= \exp \left( -\frac{y}{b} \right) \cdot \exp \left( \frac{x_n}{nb} \right) \left( \mathbb{E} \left[ \exp \left( \frac{X_1}{nb} \right) \right] \right)^{n-1}. \end{aligned}$$

Thus, for  $y \geq d$  the PMC is obtained as<sup>3</sup>

$$\begin{aligned} \Lambda(X_n \rightarrow y) &= \log \frac{f_Y(y)}{\inf_{x_n \in [c, d]} f_{Y|X_n=x_n}(y)} \\ &= \log \left( \exp \left( -\frac{c}{nb} \right) \cdot \mathbb{E} \left[ \exp \left( \frac{X_1}{nb} \right) \right] \right) \\ &= -\frac{c}{nb} + \log \mathbb{E} \left[ \exp \left( \frac{X_1}{nb} \right) \right] \\ &= \frac{\mu - c}{nb} + K \left( \frac{1}{nb} \right). \end{aligned} \tag{13}$$

Similarly, when  $y \leq c$  the PMC is

$$\Lambda(X_n \rightarrow y) = \frac{d - \mu}{nb} + K \left( -\frac{1}{nb} \right). \tag{14}$$

Combining (13) and (14), we conclude that

$$\sup_{y \in \mathbb{R}} \Lambda(X_n \rightarrow y) = \max \left\{ \frac{\mu - c}{nb} + K \left( \frac{1}{nb} \right), \frac{d - \mu}{nb} + K \left( -\frac{1}{nb} \right) \right\}.$$

<sup>3</sup>We assume that  $\mathbb{P}\{X_n \in [c, c + \delta]\} > 0$  for arbitrarily small  $\delta > 0$  so that  $\text{ess inf}_{X_n} \exp(\frac{X_n}{nb}) = \exp(\frac{c}{nb})$ . Otherwise, the derived expression is an upper bound on the PMC. Similar assumptions are made about intervals of the form  $(d - \delta, d]$ .



In the absence of more assumptions about the distribution of  $X_i$ 's, we may use  $c - \mu \leq Z_1 \leq d - \mu$  to bound  $K(\frac{1}{nb})$  and  $K(-\frac{1}{nb})$  by

$$\begin{aligned} K\left(\frac{1}{nb}\right) &= \log \mathbb{E} \left[ e^{\frac{Z_1}{nb}} \right] \leq \frac{d - \mu}{nb}, \\ K\left(-\frac{1}{nb}\right) &= \log \mathbb{E} \left[ e^{-\frac{Z_1}{nb}} \right] \leq \frac{\mu - c}{nb}. \end{aligned}$$

This yields

$$\sup_{y \in \mathbb{R}} \Lambda(X_n \rightarrow y) \leq \frac{d - c}{nb}.$$

Note that the right-hand side of the above bound corresponds to the differential privacy parameter of the Laplace mechanism for a query with global sensitivity  $\frac{d-c}{n}$  [5].

However, with additional assumptions about the distribution of  $X_i$ 's, we can derive tighter bounds on the PMC. For example, if  $X_i \sim \mathcal{U}[c, d]$  then  $\mu = \frac{c+d}{2}$  and we have

$$\begin{aligned} K\left(\frac{1}{nb}\right) &= K\left(-\frac{1}{nb}\right) = \log \int_{\frac{c-d}{2}}^{\frac{d-c}{2}} \frac{1}{d-c} \exp\left(\frac{x}{nb}\right) dx \\ &= \log \left( \frac{nb}{d-c} \left[ \exp\left(\frac{d-c}{2nb}\right) - \exp\left(\frac{c-d}{2nb}\right) \right] \right) \\ &= \log \left( \frac{nb}{d-c} \cdot \exp\left(\frac{c-d}{2nb}\right) \left[ \exp\left(\frac{d-c}{nb}\right) - 1 \right] \right). \end{aligned}$$

In this case, the Laplace mechanism results in the PMC

$$\sup_{y \in \mathbb{R}} \Lambda(X_n \rightarrow y) = \log \left( \frac{nb}{d-c} \left[ \exp\left(\frac{d-c}{nb}\right) - 1 \right] \right),$$

which is smaller than  $\frac{d-c}{nb}$ .

**Example 6** (Gaussian noise). Let  $X$  be a zero mean and bounded random variable with  $|X| \leq A$ , and let  $K(t) := \log \mathbb{E} [e^{tX}]$  with  $t \in \mathbb{R}$  denote the cumulant generating function of  $X$ . Suppose our goal is to safely release the realization of  $X$  by adding Gaussian noise to it. Thus, we release an outcome of  $Y$  with conditional distribution  $Y | X = x \sim \mathcal{N}(x, \sigma^2)$ .

To calculate the PMC  $\Lambda(X \rightarrow y)$ , first suppose  $y \geq 0$ . We write

$$\begin{aligned} \Lambda(X \rightarrow y) &= \log \frac{f_Y(y)}{\inf_{x \in [-A, A]} f_{Y|X=x}(y)} \\ &= \log \frac{\mathbb{E}_X \left[ \exp\left(-\frac{(y-X)^2}{2\sigma^2}\right) \right]}{\exp\left(-\frac{(y+A)^2}{2\sigma^2}\right)} \\ &= \log \mathbb{E}_X \left[ \exp\left(\frac{(y+A)^2 - (y-X)^2}{2\sigma^2}\right) \right] \\ &= \log \mathbb{E}_X \left[ \exp\left(\frac{2y(A+X) + A^2 - X^2}{2\sigma^2}\right) \right]. \end{aligned}$$

Since  $0 \leq X^2 \leq A^2$ , we obtain the following upper and lower bounds on  $\Lambda(X \rightarrow y)$ :<sup>4</sup>

$$\log \mathbb{E}_X \left[ \exp \left( \frac{2y(A+X)}{2\sigma^2} \right) \right] \leq \Lambda(X \rightarrow y) \leq \log \mathbb{E}_X \left[ \exp \left( \frac{2y(A+X) + A^2}{2\sigma^2} \right) \right],$$

which can be expressed as

$$\frac{Ay}{\sigma^2} + K\left(\frac{y}{\sigma^2}\right) \leq \Lambda(X \rightarrow y) \leq \frac{A(A+2y)}{2\sigma^2} + K\left(\frac{y}{\sigma^2}\right).$$

By Jensen's inequality and using the fact that  $\mathbb{E}[X] = 0$ , we have  $K\left(\frac{y}{\sigma^2}\right) \geq 0$ . Furthermore, since  $X \leq A$ , we have the upper bound

$$K\left(\frac{y}{\sigma^2}\right) = \log \mathbb{E} \left[ \exp \left( \frac{yX}{\sigma^2} \right) \right] \leq \frac{yA}{\sigma^2}.$$

Thus, we get

$$\frac{Ay}{\sigma^2} \leq \Lambda(X \rightarrow y) \leq \frac{A(A+4y)}{2\sigma^2}. \quad (15)$$

Similarly, when  $y \leq 0$  we have

$$\Lambda(X \rightarrow y) = \log \mathbb{E}_X \left[ \exp \left( \frac{-2y(A-X) + A^2 - X^2}{2\sigma^2} \right) \right].$$

This yields the bounds

$$-\frac{Ay}{\sigma^2} + K\left(\frac{y}{\sigma^2}\right) \leq \Lambda(X \rightarrow y) \leq \frac{A(A-2y)}{2\sigma^2} + K\left(\frac{y}{\sigma^2}\right),$$

and

$$-\frac{Ay}{\sigma^2} \leq \Lambda(X \rightarrow y) \leq \frac{A(A-4y)}{2\sigma^2}. \quad (16)$$

Combining (15) and (16) we get the PMC bounds

$$\frac{A|y|}{\sigma^2} \leq \Lambda(X \rightarrow y) \leq \frac{A(A+|4y|)}{2\sigma^2}, \quad y \in \mathbb{R}. \quad (17)$$

The lower bound in (17) implies that  $\Lambda(X \rightarrow Y)$  is unbounded. Thus, the Gaussian noise mechanism does not satisfy  $\varepsilon$ -PMC for any finite value of  $\varepsilon$ . Nevertheless, we can use the upper bound in (17) to characterize the tail of  $\Lambda(X \rightarrow Y)$ .

Let  $\beta \geq 0$  and suppose  $A^2 = r\sigma^2$  where  $r \geq 0$ . Since  $X$  is sub-Gaussian with variance proxy  $A^2$  and the noise has Gaussian distribution with variance  $\sigma^2$ , then  $Y$  is sub-Gaussian with variance proxy  $A^2 + \sigma^2 = (r+1)\sigma^2$ .

Hence, we can write

$$\begin{aligned} \mathbb{P} \left\{ \Lambda(X \rightarrow Y) \geq \beta + \frac{A^2}{2\sigma^2} \right\} &\leq \mathbb{P} \left\{ \frac{A(A+4|Y|)}{2\sigma^2} \geq \beta + \frac{A^2}{2\sigma^2} \right\} \\ &= \mathbb{P} \left\{ |Y| \geq \frac{\sigma^2 \beta}{2A} \right\} \\ &\leq 2 \exp \left( -\frac{\beta^2 \sigma^4}{8A^2(A^2 + \sigma^2)} \right) \\ &= 2 \exp \left( -\frac{\beta^2}{8(r^2 + r)} \right), \end{aligned} \quad (18)$$

<sup>4</sup>When  $y$  is small, using  $(y-X)^2 \geq 0$  yields a tighter upper bound on the PMC. Here, we are mostly interested in large  $y$  since our goal is to characterize the tail of  $\Lambda(X \rightarrow Y)$ .

where (18) follows from the Chernoff bound on the tail of a sub-Gaussian random variable. Thus,  $\Lambda(X \rightarrow Y)$  is dominated by a Gaussian random variable with mean  $\frac{r}{2}$  and variance  $4r(r+1)$ .

## V. PMC, PML, AND CONNECTIONS TO OTHER PRIVACY MEASURES

In general, the upper and lower bounds on the information density are not independent quantities. These bounds influence each other so that imposing a lower bound on the information density may imply a certain upper bound, and vice versa. In the context of privacy, this naturally raises the question of whether a PMC guarantee inherently provides a certain PML guarantee, or whether a PML guarantee implies a certain PMC guarantee.

In this section, we explore how PMC and PML guarantees interact with each other and with other privacy definitions, namely, ALIP, LIP, and LDP. First, in Section V-A, we assume that  $X$  is a finite random variable and discuss how various privacy guarantees interrelate. We also highlight the implications of our results for designing mechanisms with optimal utility. Then, in Section V-B, we extend our discussions to secrets on standard Borel spaces.

### A. Secrets with Finite Alphabets

Suppose  $X \sim P_X$  is a finite random variable, and recall that  $p_{\min} = \min_{x \in \mathcal{X}} P_X(x)$ . We begin by illustrating how an LDP privacy guarantee translates into a PMC guarantee. This result combined with Proposition 1 paints a complete picture depicting how LDP restricts the information density. Proposition 3 is proved in Appendix E.

**Proposition 3.** Suppose  $X$  is a finite random variable distributed according to  $P_X$ . If the mechanism  $P_{Y|X}$  satisfies  $\varepsilon$ -LDP, then it satisfies  $\varepsilon'$ -PMC with

$$\varepsilon' = \log \left( e^\varepsilon - p_{\min}(e^\varepsilon - 1) \right).$$

Note that similarly to Proposition 1,  $\varepsilon' \rightarrow \varepsilon$  as  $p_{\min} \rightarrow 0$ .

We now present the main result of this section. This result demonstrates how PMC and PML guarantees interact with each other, thereby allowing us to relate both measures to other privacy definitions.

**Theorem 4.** Suppose  $X$  is a finite random variable distributed according to  $P_X$ .

1. Let  $0 \leq \varepsilon_u < \log \frac{1}{1-p_{\min}}$ . If the privacy mechanism  $P_{Y|X}$  satisfies  $\varepsilon_u$ -PML, then it also satisfies  $\varepsilon_l^*(\varepsilon_u)$ -PMC, where

$$\varepsilon_l^*(\varepsilon_u) := \log \frac{p_{\min}}{1 - e^{\varepsilon_u}(1 - p_{\min})}.$$

2. Let  $\varepsilon_l \geq 0$ . If the privacy mechanism  $P_{Y|X}$  satisfies  $\varepsilon_l$ -PMC, then it also satisfies  $\varepsilon_u^*(\varepsilon_l)$ -PML, where

$$\varepsilon_u^*(\varepsilon_l) := \log \frac{1 - e^{-\varepsilon_l}(1 - p_{\min})}{p_{\min}}.$$

Theorem 4 is proved in Appendix F.

For notational convenience, we let  $\varepsilon_u^* := \varepsilon_u^*(\varepsilon_l)$  and  $\varepsilon_l^* := \varepsilon_l^*(\varepsilon_u)$  for fixed values of  $\varepsilon_l$  and  $\varepsilon_u$ .

**Remark 5.** The condition  $\varepsilon_u < \log \frac{1}{1-p_{\min}}$  restricts us to the high-privacy regime of PML and is necessary for  $\varepsilon_l^*$  to be well-defined. This is because a mechanism  $P_{Y|X}$  satisfying  $\varepsilon_u$ -PML with  $\varepsilon_u \geq \log \frac{1}{1-p_{\min}}$  may assign




Fig. 1: Relationship between the upper and lower bounds on the information density. The blue curve corresponds to  $\varepsilon_l^*(\varepsilon_u)$  and the orange curve corresponds to  $\varepsilon_u^*(\varepsilon_l)$ . When  $P_X$  is binary and uniform, the two curves coincide.

$P_{Y|X=x}(y) = 0$  for some  $(x, y)$  pair with positive probability. Such a mechanism does not satisfy  $\varepsilon_l$ -PMC for any finite value of  $\varepsilon_l$ . This is also why an  $\varepsilon_u$ -PML guarantee yields useful LIP, ALIP and LDP guarantees only if  $\varepsilon_u < \log \frac{1}{1-p_{\min}}$  (see Corollary 3).

**Remark 6.** The translation from PML to PMC in Theorem 4 in general cannot be improved. In particular, as we saw in Example 4, the high-privacy extremal mechanism satisfying  $\varepsilon_u$ -PML also satisfies  $\varepsilon_l^*$ -PMC.

Figure 1 illustrates the bounds presented in Theorem 4. Consider a pair  $(\varepsilon_u, \varepsilon_l) \geq (0, 0)$  with  $\varepsilon_u < \log \frac{1}{1-p_{\min}}$ . Theorem 4 states that if  $\varepsilon_l > \varepsilon_l^*$ , then  $\varepsilon_u$ -PML is a strictly stronger privacy guarantee compared to  $\varepsilon_l$ -PMC. Such pairs constitute the gray area above the blue curve in Figure 1a. Similarly, if  $\varepsilon_u > \varepsilon_u^*$  then  $\varepsilon_l$ -PMC is a strictly stronger privacy guarantee compared to  $\varepsilon_u$ -PML. These pairs correspond to the gray area below the orange curve in Figure 1a. In contrast, the blue region in Figure 1a represents  $(\varepsilon_u, \varepsilon_l)$  pairs that satisfy  $\varepsilon_u^* \geq \varepsilon_u$  and  $\varepsilon_l^* \geq \varepsilon_l$ . In this region, neither the  $\varepsilon_u$ -PML nor the  $\varepsilon_l$ -PMC privacy guarantee is stronger than the other. Interestingly, when  $X$  is an equiprobable Bernoulli random variable, the gap between the  $\varepsilon_u$ -PML and  $\varepsilon_l$ -PMC curves (i.e., the blue region) disappears, as illustrated in Figure 1b. We formalize this observation in Corollary 4.

Let us now discuss some of the implications of Theorem 4. The primary consequence of this theorem is that both PML and PMC privacy guarantees translate into ALIP, LIP, and LDP guarantees. The ALIP and LIP guarantees simply follow from their definitions and the LDP guarantees follow from [19, Prop. 1], where it is shown that  $(\varepsilon_l, \varepsilon_u)$ -ALIP implies  $(\varepsilon_l + \varepsilon_u)$ -LDP.

**Corollary 3.** Suppose  $X$  is a finite random variable distributed according to  $P_X$ .

1. Let  $0 \leq \varepsilon_u < \log \frac{1}{1-p_{\min}}$ . If the mechanism  $P_{Y|X}$  satisfies  $\varepsilon_u$ -PML, then it also satisfies

$$(\varepsilon_l^*, \varepsilon_u)\text{-ALIP}, \quad \max\{\varepsilon_l^*, \varepsilon_u\}\text{-LIP}, \quad (\varepsilon_l^* + \varepsilon_u)\text{-LDP}.$$

2. Let  $\varepsilon_l \geq 0$ . If the mechanism  $P_{Y|X}$  satisfies  $\varepsilon_l$ -PMC, then it also satisfies

$$(\varepsilon_l, \varepsilon_u^*)\text{-ALIP}, \quad \max\{\varepsilon_l, \varepsilon_u^*\}\text{-LIP}, \quad (\varepsilon_l + \varepsilon_u^*)\text{-LDP}.$$

Corollary 3 describes implications in one direction: from PML or PMC to ALIP, LIP, and LDP. Interestingly, when  $X$  is an equiprobable Bernoulli random variable the reverse implications also hold. This is because, in this case, we have  $(\varepsilon_l^* \circ \varepsilon_u^*)(\varepsilon_l) = \varepsilon_l$  and  $(\varepsilon_u^* \circ \varepsilon_l^*)(\varepsilon_u) = \varepsilon_u$ . In the following corollary, the LIP statement arises from the fact that when  $X$  is an equiprobable Bernoulli random variable, then  $\varepsilon_l^*(\varepsilon_u) \geq \varepsilon_u$ . The LDP statement follows from Corollary 3 and Proposition 1.

**Corollary 4.** Suppose  $\mathcal{X} = \{0, 1\}$  and  $P_X(0) = P_X(1) = \frac{1}{2}$ . Given  $0 \leq \varepsilon_u < \log 2$ , a privacy mechanism satisfies  $\varepsilon_u$ -PML if and only if it satisfies

1.  $\varepsilon_l^*$ -PMC, or
2.  $(\varepsilon_l^*, \varepsilon_u)$ -ALIP, or
3.  $\varepsilon_l^*$ -LIP, or
4.  $\varepsilon_l^* + \varepsilon_u$ -LDP.

Alternatively, we could have fixed  $\varepsilon_l \geq 0$  and stated the result for  $\varepsilon_l$ -PMC and  $\varepsilon_u^*$ -PML.

The equivalence between PML and LDP expressed in Corollary 4 also explains why the binary PML-extremal mechanism (Example 4) for the uniform prior distribution coincides with the (binary) randomized response mechanism (Example 3).<sup>5</sup> Recall that PML-extremal mechanisms are optimal for maximizing sub-convex utility functions. Similarly, the randomized response mechanism has been shown to be optimal for LDP and the class of sub-convex utility functions [25].

Finally, we discuss the application of Theorem 4 to the privacy mechanism design problem. In the following, we use the fact that  $(\varepsilon_l, \varepsilon_u)$ -ALIP is equivalent to  $\varepsilon_u$ -PML if  $\varepsilon_l \geq \varepsilon_l^*$ .

**Corollary 5.** Suppose  $X$  is a finite random variable with alphabet  $\mathcal{X} = [n]$ . Then, for all  $0 \leq \varepsilon_u < \log \frac{1}{1-p_{\min}}$ , all  $\varepsilon_l \geq \varepsilon_l^*$ , and all utility functions, the optimal  $(\varepsilon_l, \varepsilon_u)$ -ALIP mechanism coincides with the optimal  $\varepsilon_u$ -PML mechanism. In particular, the optimal  $(\varepsilon_l, \varepsilon_u)$ -ALIP mechanism maximizing sub-convex utility functions with  $\varepsilon_l \geq \varepsilon_l^*$  is

$$P_{Y|X=i}(j) = \begin{cases} 1 - e^{\varepsilon_u}(1 - P_X(i)) & \text{if } i = j, \\ e^{\varepsilon_u} P_X(j) & \text{if } i \neq j, \end{cases}$$

where  $i, j \in [n]$ .

A similar statement can be made using  $\varepsilon_l$ -PMC and  $\varepsilon_u \geq \varepsilon_u^*$ .

<sup>5</sup>More precisely, given  $\varepsilon \geq 0$ , the binary randomized response mechanism with parameter  $\varepsilon_r = \varepsilon + \log \frac{2}{2-\varepsilon}$  coincides with the binary  $\varepsilon$ -PML extremal mechanism. See [62] for more details.

### B. Secrets with General Alphabets

Now, we consider the more general case where  $(X, Y)$  takes values in the product space  $(\mathcal{X} \times \mathcal{Y}, \mathcal{S}_X \otimes \mathcal{S}_Y)$ . Note that the first statement in Theorem 4 has no useful parallel for secrets with infinite alphabets due to its dependence on  $p_{\min}$ . Nonetheless, below we provide a more general version of the second statement in Theorem 4. Its proof is provided in Appendix G.

**Theorem 5.** Suppose  $P_{XY} \ll P_X \times P_Y$ . If  $\Lambda(X \rightarrow Y)$  is essentially bounded, then  $\ell(X \rightarrow Y)$  is also essentially bounded.

Note that the opposite direction in the above result does not hold. For example, while PML is always bounded for finite random variables, PMC can be infinite even in these cases. Hence, Theorem 5 demonstrates that PMC is a stricter measure compared to PML. Interestingly, below we show that an even stronger statement is possible.

**Theorem 6.**  $\Lambda(X \rightarrow Y)$  is essentially bounded if and only if the privacy mechanism  $P_{Y|X}$  satisfies  $\epsilon$ -LDP for some  $0 \leq \epsilon < \infty$ .

The above result is proved in Appendix H. Note that Theorem 5 may be viewed as a consequence of Theorem 6 since a mechanism satisfying LDP has bounded PML.

## VI. DISCUSSION: PMC AS AN INFORMATION LEAKAGE MEASURE

While PMC and PML are both operationally meaningful and context-aware privacy measures, they are conceptually different. PML is defined by modeling opportunistic adversaries aiming to maximize gain functions, whereas PMC is obtained by modeling risk-averse adversaries interested in minimizing cost functions. This fundamental difference in the threat models results in substantial qualitative and quantitative differences between the two measures. Here, we discuss their differences along two themes.

First, in one of the earliest papers on quantitative information flow, Smith [27] posited that a measure of information leakage should capture the difference between the initial uncertainty about a random variable and the remaining uncertainty after observing a correlated quantity. Specifically, he suggested that leakage measures should satisfy

$$\text{information leakage} = \text{initial uncertainty} - \text{remaining uncertainty}. \quad (19)$$

When  $X$  is a finite random variable, the initial uncertainty in  $X$  is finite according to all the standard measures of uncertainty such as the Rényi entropies. Therefore, a direct consequence of (19) is that information leaked about finite random variables should invariably remain finite, regardless of the mechanism used. This finiteness criterion may be viewed as an axiomatic requirement for information leakage measures and is satisfied by most measures such as PML, mutual information, and  $f$ -information, among others.

However, as noted in Remark 3, PMC can become infinite even when  $X$  is finite. As an example, consider a binary random variable  $X$  with  $P_X(0) = q$  with  $q \in (0, 1)$  and suppose the posterior distribution satisfies  $P_{X|Y=y}(0) = 1$  for some  $y \in \mathcal{Y}$ . Here, according to PMC the information leaked is  $\Lambda_{P_{XY}}(X \rightarrow y) = \infty$ . Consequently, PMC

satisfies neither (19) nor the weaker finiteness criterion. This discrepancy becomes even more counter-intuitive as  $q$  approaches 1, where the seemingly incremental knowledge gained about  $X$  still yields  $\Lambda_{P_{XY}}(X \rightarrow y) = \infty$ .

A second and more practical limitation of PMC is that while it is a context-aware privacy measure, its application is effectively restricted to the class of mechanisms compatible with LDP and DP. In finite settings, this is evident from Theorem 4 where PMC bounds can be derived only in the high-privacy regime. This trend also extends to the general case as indicated by Theorem 6. These results suggest that PMC has a more limited application compared to PML and is more restrictive in terms of utility.

As a concrete example, consider the task of estimating the mean of a Gaussian random variable by adding Gaussian noise to the sample mean. Such a problem is well-defined under PML since PML allows the addition of Gaussian noise to a Gaussian random variable (see [13, Example 4]). However, this problem cannot be studied with PMC. As shown in Example 6 and specifically (17), the lower bound on PMC becomes infinite when Gaussian noise is added to an unbounded random variable. This observation aligns with the sensitivity-based analysis in DP, where the sample mean’s sensitivity becomes infinite when the data domain is unbounded.

That being said, while using DP and LDP mechanisms, allowing higher values of PMC compared to PML can result in improved utility. In this context, employing a framework such as ALIP offers better flexibility and more favorable privacy-utility tradeoffs compared to LIP.

## VII. CONCLUSIONS

In this paper, we explored the operational meaning and the implications of guaranteeing privacy by enforcing a lower bound on the information density. We introduced a novel privacy measure, pointwise maximal cost (PMC), and demonstrated that imposing an upper bound on PMC is equivalent to enforcing a lower bound on the information density. PMC quantifies the information leakage about a secret  $X$  to risk-averse adversaries who aim to minimize non-negative cost functions after observing an outcome  $y$  of a privacy mechanism. When  $X$  is a finite random variable, PMC can also be defined by considering adversaries aiming to minimize the probability of incorrectly guessing randomized functions of  $X$ . By investigating the operational meaning of the lower bound on the information density, our results have strengthened and increased the relevance of definitions such as LIP and ALIP that previously imposed the lower bound axiomatically.

This work further explored the relationship between the upper and lower bounds on the information density by examining how PMC and PML guarantees relate. Our findings reveal that PMC is a more stringent privacy measure and its application is limited to mechanisms satisfying differential privacy-type guarantees. Notably, PMC can be infinite even when applied to finite random variables. This characteristic, which sets PMC apart from other measures such as PML and  $f$ -information, arises from the definition of PMC: If the posterior expected value of a cost function is zero then PMC is infinite.

Overall, our work significantly bridges the gaps in understanding the relationships between various privacy frameworks. Looking ahead, the findings in this paper offer crucial insights for selecting the appropriate framework to ensure privacy in different applications. If the objective is to adopt a conservative approach that prevents any definitive inference about the secret, then LDP, LIP, and ALIP would be suitable choices. When information about

the prior distribution is available, LIP and ALIP are preferred as they enable higher utility compared to LDP. ALIP provides additional flexibility by allowing distinct upper and lower bounds, while LIP simplifies the design by imposing symmetric bounds. Conversely, when the prior is not known, then LDP would be a suitable choice since it offers equally strong protection regardless of the prior. In addition, PML emerges as a particularly fitting choice when working with mechanisms that do not satisfy privacy in the traditional differential privacy sense, for instance, mechanisms with unbounded domains. PML provides the flexibility to tune the privacy parameter to achieve different types of protection, from preventing any definitive inference about the secret to high-utility mechanisms that hide only the least likely and atypical features of the secret [21].

Finally, our work has practical implications for designing privacy mechanisms since a mechanism initially designed to meet a specific privacy guarantee may also satisfy other types of guarantees. Using the relations between various measures presented here, it is possible to examine if a mechanism provides other guarantees beyond its original design.

#### APPENDIX A PROOF OF THEOREM 2

Fix  $y \in \mathcal{Y}$ . First, we argue that given a  $U$  satisfying the Markov chain  $U - X - Y$  there exists a non-negative cost function  $c_U$  such that  $\Lambda_U(X \rightarrow y) = \Lambda_c(X \rightarrow y)$ . Given  $U$ , let the cost function  $c_U : \mathcal{X} \times \mathcal{U} \rightarrow \mathbb{R}_+$  be defined as  $c_U(x, u) = 1 - P_{U|X=x}(u)$  with  $x \in \mathcal{X}$  and  $u \in \mathcal{U}$ . Then, we have

$$\begin{aligned} \exp\left(\Lambda_{c_U}(X \rightarrow y)\right) &= \frac{\inf_{P_{\hat{U}}} \mathbb{E}\left[c_U(X, \hat{U})\right]}{\inf_{P_{\tilde{U}|Y}} \mathbb{E}\left[c_U(X, \tilde{U}) \mid Y = y\right]} \\ &= \frac{\min_{u \in \mathcal{U}} \mathbb{E}[c_U(X, u)]}{\min_{u \in \mathcal{U}} \mathbb{E}[c_U(X, u) \mid Y = y]} \\ &= \frac{\min_{u \in \mathcal{U}} \sum_x c_U(x, u) P_X(x)}{\min_{u \in \mathcal{U}} \sum_x c_U(x, u) P_{X|Y=y}(x)} \\ &= \frac{\min_{u \in \mathcal{U}} \sum_x (1 - P_{U|X=x}(u)) P_X(x)}{\min_{u \in \mathcal{U}} \sum_x (1 - P_{U|X=x}(u)) P_{X|Y=y}(x)} \\ &= \frac{1 - \max_{u \in \mathcal{U}} \sum_x P_{U|X=x}(u) P_X(x)}{1 - \max_{u \in \mathcal{U}} \sum_x P_{U|X=x}(u) P_{X|Y=y}(x)} \\ &= \exp\left(\Lambda_U(X \rightarrow y)\right). \end{aligned}$$

Observe that the above construction works even when  $\Lambda_U(X \rightarrow y) = \infty$  in which case we also get  $\Lambda_{c_U}(X \rightarrow y) = \infty$ .

Now, we show that for each non-negative cost function  $c$ , there exists  $U$  satisfying the Markov chain  $U - X - Y$  such that  $\Lambda_c(X \rightarrow y) = \Lambda_U(X \rightarrow y)$ . Fix a cost function  $c$ . Without loss of generality, we assume that  $0 \leq c(x, w) \leq 1$  for all  $x \in \mathcal{X}$  and  $w \in \mathcal{W}$ . This can be achieved through normalizing  $c$  by  $\max_{x,w} c(x, w)$ .



First, suppose  $\Lambda_c(X \rightarrow y) < \infty$ . Let  $k$  be a large integer. We construct two randomized functions of  $X$  denoted by  $S$  and  $T$  both on the same alphabet  $[k+1]$ . Let

$$w_S \in \arg \min_w \sum_x c(x, w) P_X(x),$$

$$w_T \in \arg \min_w \sum_x c(x, w) P_{X|Y=y}(x),$$

where  $w_S$  denotes the adversary's optimal choice prior to observing  $y$  and  $w_T$  denotes the adversary's optimal choice after observing  $y$ . For all  $x \in \mathcal{X}$ , let

$$P_{S|X=x}(i) = \frac{c(x, w_S)}{k}, \quad i \in [k],$$

$$P_{S|X=x}(k+1) = 1 - c(x, w_S),$$

$$P_{T|X=x}(i) = \frac{c(x, w_T)}{k}, \quad i \in [k],$$

$$P_{T|X=x}(k+1) = 1 - c(x, w_T).$$

Let  $0 \leq \delta \leq 1$ . We define  $U_\delta$  as the mixture of  $S$  and  $T$  with parameter  $\delta$ , that is,

$$U_\delta = \begin{cases} S & \text{with probability } \delta, \\ T & \text{with probability } 1 - \delta. \end{cases}$$

Then,  $P_{U_\delta|X=x}(i) = \delta P_{S|X=x}(i) + (1 - \delta) P_{T|X=x}(i)$  for  $i \in [k+1]$  and we get

$$\begin{aligned} \inf_{\hat{U}} \mathbb{P}[U_\delta \neq \hat{U}] &= \min_{u \in [k+1]} \sum_x (1 - P_{U_\delta|X=x}(u)) P_X(x) \\ &= \min \left\{ \sum_x \left( 1 - \delta \frac{c(x, w_S)}{k} - (1 - \delta) \frac{c(x, w_T)}{k} \right) P_X(x), \sum_x \left( \delta c(x, w_S) + (1 - \delta) c(x, w_T) \right) P_X(x) \right\} \\ &= \sum_x \left( \delta c(x, w_S) + (1 - \delta) c(x, w_T) \right) P_X(x), \end{aligned}$$

where the last equality follows by letting  $k \rightarrow \infty$ . Similarly, we have

$$\begin{aligned} \inf_{\hat{U}|Y} \mathbb{P}[U_\delta \neq \hat{U} | Y = y] &= \min_{u \in [k+1]} \sum_x (1 - P_{U_\delta|X=x}(u)) P_{X|Y=y}(x) \\ &= \min \left\{ \sum_x \left( 1 - \delta \frac{c(x, w_S)}{k} - (1 - \delta) \frac{c(x, w_T)}{k} \right) P_{X|Y=y}(x), \sum_x \left( \delta c(x, w_S) + (1 - \delta) c(x, w_T) \right) P_{X|Y=y}(x) \right\} \\ &= \sum_x \left( \delta c(x, w_S) + (1 - \delta) c(x, w_T) \right) P_{X|Y=y}(x), \end{aligned}$$

and, once again, the last equality follows by letting  $k \rightarrow \infty$ . Therefore, we have

$$\exp \left( \Lambda_{U_\delta}(X \rightarrow y) \right) = \frac{\sum_x \left( \delta c(x, w_S) + (1 - \delta) c(x, w_T) \right) P_X(x)}{\sum_x \left( \delta c(x, w_S) + (1 - \delta) c(x, w_T) \right) P_{X|Y=y}(x)}.$$

Next, note that when  $\delta = 1$  we have

$$\exp \left( \Lambda_{U_1}(X \rightarrow y) \right) = \exp \left( \Lambda_S(X \rightarrow y) \right)$$

$$\begin{aligned}
&= \frac{\sum_x c(x, w_S) P_X(x)}{\sum_x c(x, w_S) P_{X|Y=y}(x)} \\
&\leq \frac{\sum_x c(x, w_S) P_X(x)}{\sum_x c(x, w_T) P_{X|Y=y}(x)} \\
&= \frac{\min_w \sum_x c(x, w) P_X(x)}{\min_w \sum_x c(x, w) P_{X|Y=y}(x)} \\
&= \exp\left(\Lambda_c(X \rightarrow y)\right),
\end{aligned}$$

and when  $\delta = 0$  we have

$$\begin{aligned}
\exp\left(\Lambda_{U_0}(X \rightarrow y)\right) &= \exp\left(\Lambda_T(X \rightarrow y)\right) \\
&= \frac{\sum_x c(x, w_T) P_X(x)}{\sum_x c(x, w_T) P_{X|Y=y}(x)} \\
&\geq \frac{\sum_x c(x, w_S) P_X(x)}{\sum_x c(x, w_T) P_{X|Y=y}(x)} \\
&= \frac{\min_w \sum_x c(x, w) P_X(x)}{\min_w \sum_x c(x, w) P_{X|Y=y}(x)} \\
&= \exp\left(\Lambda_c(X \rightarrow y)\right).
\end{aligned}$$

In other words, we have  $\Lambda_{U_1}(X \rightarrow y) \leq \Lambda_c(X \rightarrow y) \leq \Lambda_{U_0}(X \rightarrow y)$ . Then, due to the continuity of the mapping  $\delta \mapsto \Lambda_{U_\delta}(X \rightarrow y)$ , there exists  $\delta^* \in [0, 1]$  such that  $\Lambda_{U_{\delta^*}}(X \rightarrow y) = \Lambda_c(X \rightarrow y)$ .

Finally, we consider the case  $\Lambda_c(X \rightarrow y) = \infty$ . In this case, there exists a proper subset of  $\mathcal{X}$  denoted by  $\mathcal{E}$  such that  $P_{X|Y=y}(\mathcal{E}) = 1$  and  $c(x, w_T) = 0$  for all  $x \in \mathcal{E}$ . Let  $U$  be a binary random variable described by  $P_{U|X=x}(0) = 1$  for all  $x \in \mathcal{E}$  and  $P_{U|X=x}(0) = \frac{1}{2}$  for  $x \in \mathcal{X} \setminus \mathcal{E}$ . Then, we have

$$\begin{aligned}
\Lambda_U(X \rightarrow y) &= \log \frac{1 - \max\{P_U(0), P_U(1)\}}{1 - \max_u \sum_{x \in \mathcal{X}} P_{U|X=x}(u) P_{X|Y=y}(x)} \\
&= \log \frac{1 - \max\{\frac{1}{2} + \frac{P_X(\mathcal{E})}{2}, \frac{P_X(\mathcal{X} \setminus \mathcal{E})}{2}\}}{1 - P_{X|Y=y}(\mathcal{E})} \\
&= \infty,
\end{aligned}$$

since  $0 < P_X(\mathcal{E}) < 1$ .

## APPENDIX B

### PROOF OF THEOREM 3

Fix  $y \in \mathcal{Y}$ . The proof follows along the same lines of the proof of [13, Thm. 3]. We begin by noting that since  $(\mathcal{X}, \mathcal{S}_X)$  is standard Borel, then  $P_{XY}$  can be disintegrated into the marginal  $P_Y$  and a conditional distribution  $P_{X|Y}$  [59, Thm. IV.2.18]. We use this fact to simplify the denominator of (11). The following lemma is proved in Appendix C.

**Lemma 2.** Let  $W$  be a random variable induced by a transition probability kernel  $P_{W|Y}$  from  $(\mathcal{Y}, \mathcal{S}_Y)$  into  $(\mathcal{W}, \mathcal{S}_W)$ . Suppose  $P_{XY}(dx, dy) = P_Y(dy)P_{X|Y=y}(dx)$ . For all  $c \in \mathcal{C}$ , it holds that

$$\inf_{P_{\hat{W}|Y}} \mathbb{E} \left[ c(X, \hat{W}) \mid Y = y \right] = \inf_{\hat{w} \in \mathcal{W}} \int_{\mathcal{X}} c(x, \hat{w}) P_{X|Y=y}(dx).$$

It also follows from Lemma 2 that

$$\inf_{P_W} \mathbb{E} \left[ c(X, W) \right] = \inf_{w \in \mathcal{W}} \int_{\mathcal{X}} c(x, w) P_X(dx).$$

Suppose  $P_X \ll P_{X|Y=y}$  and let  $f := \frac{dP_X}{dP_{X|Y=y}}$  denote the Radon-Nikodym derivative of  $P_X$  with respect to  $P_{X|Y=y}$ . First, we show that  $\Lambda_{P_{XY}}(X \rightarrow y) \leq D_\infty(P_X \| P_{X|Y=y})$ . Fix a measurable space  $(\mathcal{W}, \mathcal{S}_W)$  and a cost function  $c \in \mathcal{C}$ . We can write

$$\begin{aligned} \frac{\inf_{P_W} \mathbb{E} \left[ c(X, W) \right]}{\inf_{P_{\hat{W}|Y}} \mathbb{E} \left[ c(X, \hat{W}) \mid Y = y \right]} &= \frac{\inf_{w \in \mathcal{W}} \int_{\mathcal{X}} c(x, w) P_X(dx)}{\inf_{\hat{w} \in \mathcal{W}} \int_{\mathcal{X}} c(x, \hat{w}) P_{X|Y=y}(dx)} \\ &= \sup_{\hat{w} \in \mathcal{W}} \frac{\inf_{w \in \mathcal{W}} \int_{\mathcal{X}} c(x, w) P_X(dx)}{\int_{\mathcal{X}} c(x, \hat{w}) P_{X|Y=y}(dx)} \\ &\leq \sup_{\hat{w} \in \mathcal{W}} \frac{\int_{\mathcal{X}} c(x, \hat{w}) P_X(dx)}{\int_{\mathcal{X}} c(x, \hat{w}) P_{X|Y=y}(dx)} \\ &= \sup_{\hat{w} \in \mathcal{W}} \frac{\int_{\mathcal{X}} c(x, \hat{w}) f(x) P_{X|Y=y}(dx)}{\int_{\mathcal{X}} c(x, \hat{w}) P_{X|Y=y}(dx)} \\ &\leq \text{ess sup}_{P_{X|Y=y}} f \left( \sup_{\hat{w} \in \mathcal{W}} \frac{\int_{\mathcal{X}} c(x, \hat{w}) P_{X|Y=y}(dx)}{\int_{\mathcal{X}} c(x, \hat{w}) P_{X|Y=y}(dx)} \right) \\ &= \text{ess sup}_{P_X} f \\ &= \exp \left( D_\infty(P_X \| P_{X|Y=y}) \right), \end{aligned} \tag{20a}$$

where (20a) is due to the fact that  $f$  is zero on all sets  $\mathcal{A} \in \mathcal{S}_X$  such that  $P_{X|Y=y}(\mathcal{A}) > 0$  but  $P_X(\mathcal{A}) = 0$ . Thus,  $\Lambda_{P_{XY}}(X \rightarrow y) \leq D_\infty(P_X \| P_{X|Y=y})$ .

Next, we show that  $\Lambda_{P_{XY}}(X \rightarrow y) \geq D_\infty(P_X \| P_{X|Y=y})$ . Let  $\mathcal{W} = \mathbb{Z} \cup \{-\infty\}$  and  $\mathcal{S}_W$  be the discrete  $\sigma$ -algebra on  $\mathcal{W}$ . Fix  $\varepsilon > 0$  and consider the following countable partition of  $\mathcal{X}$ :

$$\mathcal{D}_w^\varepsilon = \{x \in \mathcal{X} : e^{w\varepsilon} \leq f(x) < e^{(w+1)\varepsilon}\}, \quad w \in \mathcal{W}. \tag{21}$$

Of course,  $\mathcal{D}_w^\varepsilon \in \mathcal{S}_X$  for all  $w \in \mathcal{W}$ . Fix a large constant  $M > 0$  and consider the cost function  $c^* : \mathcal{X} \times \mathcal{W} \rightarrow \mathbb{R}_+$  defined by

$$c^*(x, w) = \begin{cases} \frac{1}{P_X(\mathcal{D}_w^\varepsilon)} \mathbf{1}_{\mathcal{D}_w^\varepsilon}(x) & \text{if } P_X(\mathcal{D}_w^\varepsilon) > 0, \\ M & \text{if } P_X(\mathcal{D}_w^\varepsilon) = 0. \end{cases}$$

It is easy to see that  $c^* \in (\mathcal{S}_X \otimes \mathcal{S}_W)_+$ . Moreover, the absolute continuity  $P_X \ll P_{X|Y=y}$  ensures that if  $P_X(\mathcal{D}_w^\varepsilon) > 0$  then  $P_{X|Y=y}(\mathcal{D}_w^\varepsilon) > 0$ . Therefore, we can write

$$\exp \left( \Lambda_{P_{XY}}(X \rightarrow y) \right) \geq \frac{\inf_{w \in \mathcal{W}} \int_{\mathcal{X}} c^*(x, w) P_X(dx)}{\inf_{w \in \mathcal{W}} \int_{\mathcal{X}} c^*(x, w) P_{X|Y=y}(dx)}$$

$$\begin{aligned}
&= \frac{\min \left\{ M, \inf_{w \in \mathcal{W}: P_X(\mathcal{D}_w^\varepsilon) > 0} \int_{\mathcal{X}} \frac{1}{P_X(\mathcal{D}_w^\varepsilon)} \mathbf{1}_{\mathcal{D}_w^\varepsilon}(x) P_X(dx) \right\}}{\min \left\{ M, \inf_{w \in \mathcal{W}: P_X(\mathcal{D}_w^\varepsilon) > 0} \int_{\mathcal{X}} \frac{1}{P_X(\mathcal{D}_w^\varepsilon)} \mathbf{1}_{\mathcal{D}_w^\varepsilon}(x) P_{X|Y=y}(dx) \right\}} \\
&= \frac{\min \left\{ M, \inf_{w \in \mathcal{W}: P_X(\mathcal{D}_w^\varepsilon) > 0} \frac{P_X(\mathcal{D}_w^\varepsilon)}{P_X(\mathcal{D}_w^\varepsilon)} \right\}}{\min \left\{ M, \inf_{w \in \mathcal{W}: P_X(\mathcal{D}_w^\varepsilon) > 0} \frac{P_{X|Y=y}(\mathcal{D}_w^\varepsilon)}{P_X(\mathcal{D}_w^\varepsilon)} \right\}} \\
&= \frac{\inf_{w \in \mathcal{W}: P_X(\mathcal{D}_w^\varepsilon) > 0} \frac{P_X(\mathcal{D}_w^\varepsilon)}{P_X(\mathcal{D}_w^\varepsilon)}}{\inf_{w \in \mathcal{W}: P_X(\mathcal{D}_w^\varepsilon) > 0} \frac{P_{X|Y=y}(\mathcal{D}_w^\varepsilon)}{P_X(\mathcal{D}_w^\varepsilon)}} \tag{22a} \\
&= \sup_{w \in \mathcal{W}: P_X(\mathcal{D}_w^\varepsilon) > 0} \frac{P_X(\mathcal{D}_w^\varepsilon)}{P_{X|Y=y}(\mathcal{D}_w^\varepsilon)} \tag{22b} \\
&= \operatorname{ess\,sup}_{P_X} \bar{f}, \tag{22c}
\end{aligned}$$

where (22a) follows by taking  $M$  to be sufficiently large and

$$\bar{f}(x) := \sum_{w \in \mathcal{W}} \frac{P_X(\mathcal{D}_w^\varepsilon)}{P_{X|Y=y}(\mathcal{D}_w^\varepsilon)} \mathbf{1}_{\mathcal{D}_w^\varepsilon}(x), \quad x \in \mathcal{X}.$$

Observe that we have replaced the supremum over  $w$  in (22b) with the essential supremum over  $x$  in (22c) because  $\bar{f}$  is constant on each set  $\mathcal{D}_w^\varepsilon$ .<sup>6</sup> Finally, we have

$$\begin{aligned}
\Lambda_{P_{XY}}(X \rightarrow y) &\geq \log \operatorname{ess\,sup}_{P_X} \bar{f} \\
&\geq \log \left( \left( \operatorname{ess\,sup}_{P_X} f \right) e^{-\varepsilon} \right) \tag{23a} \\
&= \log \operatorname{ess\,sup}_{P_X} f - \varepsilon \\
&= D_\infty(P_X \| P_{X|Y=y}) - \varepsilon,
\end{aligned}$$

where (23a) is because by (21),  $\bar{f}$  never differs from  $f$  by more than a factor of  $e^\varepsilon$ . Then, by letting  $\varepsilon \rightarrow 0$ , we get  $\Lambda_{P_{XY}}(X \rightarrow y) \geq D_\infty(P_{X|Y=y} \| P_X)$ . This completes the proof for the case  $P_X \ll P_{X|Y=y}$ .

All that is left to discuss is the case  $P_X \not\ll P_{X|Y=y}$ . In this case, there exists a set  $\mathcal{A}_0 \in \mathcal{S}_X$  such that  $P_{X|Y=y}(\mathcal{A}_0) = 0$  but  $P_X(\mathcal{A}_0) > 0$ . Let  $(\mathcal{W}, \mathcal{S}_\mathcal{W})$  be an arbitrary measurable space and consider  $c(x, w) = \mathbf{1}_{\mathcal{A}_0}(x)$  for all  $w \in \mathcal{W}$ . Then, we have  $\mathbb{E}[c(X, w) | Y = y] = P_{X|Y=y}(\mathcal{A}_0) = 0$  whereas  $\mathbb{E}[c(X, w)] = P_X(\mathcal{A}_0) > 0$  for all  $w \in \mathcal{W}$ . Thus,  $\Lambda_{P_{XY}}(X \rightarrow y) = D_\infty(P_X \| P_{X|Y=y}) = \infty$ , as desired.

## APPENDIX C

### PROOF OF LEMMA 2

Fix an arbitrary kernel  $P_{W|Y}$  and  $y \in \mathcal{Y}$ . We write

$$\mathbb{E}[c(X, W) | Y = y] = \int_{\mathcal{X} \times \mathcal{W}} c(x, w) P_{XW|Y=y}(dx, dw)$$

<sup>6</sup>In fact,  $\bar{f}$  is the conditional expectation of  $f$  given  $\mathcal{F} := \sigma\{\mathcal{D}_w^\varepsilon\}$ , where  $\sigma\{\mathcal{D}_w^\varepsilon\}$  denotes the  $\sigma$ -algebra on  $\mathcal{X}$  generated by the collection of sets  $\{\mathcal{D}_w^\varepsilon\}$ .

$$\begin{aligned}
&= \int_{\mathcal{W}} P_{W|Y=y}(dw) \int_{\mathcal{X}} c(x, w) P_{X|Y=y}(dx) \\
&\geq \int_{\mathcal{W}} P_{W|Y=y}(dw) \left( \inf_{w \in \mathcal{W}} \int_{\mathcal{X}} c(x, w) P_{X|Y=y}(dx) \right) \\
&= \inf_{w \in \mathcal{W}} \int_{\mathcal{X}} c(x, w) P_{X|Y=y}(dx).
\end{aligned}$$

Taking the infimum over all kernels  $P_{W|Y}$  we get

$$\inf_{P_{W|Y}} \mathbb{E}[c(X, W) | Y = y] \geq \inf_{w \in \mathcal{W}} \int_{\mathcal{X}} c(x, w) P_{X|Y=y}(dx).$$

To prove the reverse inequality, fix  $a > \inf_{w \in \mathcal{W}} \int_{\mathcal{X}} c(x, w) P_{X|Y=y}(dx)$ . By definition, there exists  $w' \in \mathcal{W}$  such that  $\int_{\mathcal{X}} c(x, w') P_{X|Y=y}(dx) \leq a$ . Given  $w \in \mathcal{W}$ , let  $\delta_w$  denote the Dirac measure on  $(\mathcal{W}, \mathcal{S}_{\mathcal{W}})$  defined by

$$\delta_w(\mathcal{A}) = \begin{cases} 1 & \text{if } w \in \mathcal{A}, \\ 0 & \text{if } w \notin \mathcal{A}, \end{cases}$$

for each  $\mathcal{A} \in \mathcal{S}_{\mathcal{W}}$ . Then, we have

$$\begin{aligned}
\inf_{P_{W|Y}} \mathbb{E}[c(X, W) | Y = y] &\leq \int_{\mathcal{X}} P_{X|Y=y}(dx) \int_{\mathcal{W}} c(x, w) \delta_{w'}(dw) \\
&= \int_{\mathcal{X}} c(x, w') P_{X|Y=y}(dx) \\
&\leq a.
\end{aligned}$$

Then, letting  $a \rightarrow \inf_{w \in \mathcal{W}} \int_{\mathcal{X}} c(x, w) P_{X|Y=y}(dx)$  we obtain

$$\inf_{P_{W|Y}} \mathbb{E}[c(X, W) | Y = y] \leq \inf_{w \in \mathcal{W}} \int_{\mathcal{X}} c(x, w) P_{X|Y=y}(dx).$$

#### APPENDIX D

##### PROOF OF LEMMA 1

1. Non-negativity of PMC follows from the non-negativity of the Rényi divergence.
2. If  $X$  and  $Y$  are independent then  $P_{X|Y=y} = P_X$  almost surely. Thus,  $\Lambda(X \rightarrow y) = D_{\infty}(P_X \| P_{X|Y=y}) = D_{\infty}(P_X \| P_X) = 0$  almost surely.
3. The assumption implies that  $P_{X_1, \dots, X_k | Y_1, \dots, Y_k} = \prod_{i=1}^k P_{X_i | Y_i}$ . Therefore, we have

$$\begin{aligned}
\Lambda(X_1, \dots, X_k \rightarrow y_1, \dots, y_k) &= D_{\infty}(P_{X_1, \dots, X_k} \| P_{X_1, \dots, X_k | Y_1=y_1, \dots, Y_k=y_k}) \\
&= D_{\infty} \left( \prod_{i=1}^k P_{X_i} \left\| \prod_{i=1}^k P_{X_i | Y_i=y_i} \right. \right) \\
&= \sum_{i=1}^k D_{\infty}(P_{X_i} \| P_{X_i | Y_i=y_i}) \\
&= \sum_{i=1}^k \Lambda(X_i \rightarrow y_i),
\end{aligned} \tag{24}$$

almost surely, where (24) is due to the additivity property of the Rényi divergence [61, Thm. 28].

4. Fix  $y \in \mathcal{Y}$ . Let  $P_X$  and  $Q_X$  denote probability measures on  $(\mathcal{X}, \mathcal{S}_X)$ . Define  $P_\theta = \theta P_X + (1 - \theta)Q_X$ , where  $\theta \in [0, 1]$ . Let  $f_{Y|X}$  denote the conditional density (defined with respect to  $\mu$ ) of  $Y$  given  $X$ , and let

$$\begin{aligned} f_\theta &= \frac{dP_\theta}{d\mu}, \quad f_X = \frac{dP_X}{d\mu}, \quad g_X = \frac{dQ_X}{d\mu}, \\ f_Y(y) &= \int_{\mathcal{X}} f_{Y|X=x}(y) f_X(x) \mu(dx), \\ g_Y(y) &= \int_{\mathcal{X}} f_{Y|X=x}(y) g_X(x) \mu(dx), \end{aligned}$$

where  $\mu$  is a dominating  $\sigma$ -finite measure on  $\mathcal{X}$ . We have

$$\begin{aligned} \Lambda_{P_{Y|X} \times P_\theta}(X \rightarrow y) &= \log \frac{\int_{\mathcal{X}} f_{Y|X=x}(y) f_\theta(x) \mu(dx)}{\operatorname{ess\,inf}_{P_\theta} f_{Y|X}(y | X)} \\ &= \log \frac{\theta \int_{\mathcal{X}} f_{Y|X=x}(y) f_X(x) \mu(dx) + (1 - \theta) \int_{\mathcal{X}} f_{Y|X=x}(y) g_X(x) \mu(dx)}{\operatorname{ess\,inf}_{P_\theta} f_{Y|X}(y | X)} \\ &= \log \frac{\theta f_Y(y) + (1 - \theta) g_Y(y)}{\operatorname{ess\,inf}_{P_\theta} f_{Y|X}(y | X)} \\ &\geq \log \left( \theta \frac{f_Y(y)}{\operatorname{ess\,inf}_{P_X} f_{Y|X}(y | X)} + (1 - \theta) \frac{g_Y(y)}{\operatorname{ess\,inf}_{Q_X} f_{Y|X}(y | X)} \right) \tag{25} \\ &\geq \theta \log \frac{f_Y(y)}{\operatorname{ess\,inf}_{P_X} f_{Y|X}(y | X)} + (1 - \theta) \log \frac{g_Y(y)}{\operatorname{ess\,inf}_{Q_X} f_{Y|X}(y | X)} \\ &= \theta \Lambda_{P_{Y|X} \times P_X}(X \rightarrow y) + (1 - \theta) \Lambda_{P_{Y|X} \times Q_X}(X \rightarrow y), \end{aligned}$$

where (25) follows because  $P_X, Q_X \ll P_\theta$ , implying that  $\operatorname{ess\,inf}_{P_\theta} f_{Y|X}(y | X) \leq \operatorname{ess\,inf}_{P_X} f_{Y|X}(y | X)$  and  $\operatorname{ess\,inf}_{P_\theta} f_{Y|X}(y | X) \leq \operatorname{ess\,inf}_{Q_X} f_{Y|X}(y | X)$ .

5. Fix  $y \in \mathcal{Y}$ . Consider the Markov chain  $Z - X - Y$ . The kernel  $P_{Z|X}$  induces distributions  $P_Z = P_{Z|X} \circ P_X$  and  $P_{Z|Y=y} = P_{Z|X} \circ P_{X|Y=y}$ . Therefore, we can write

$$\begin{aligned} \Lambda(Z \rightarrow y) &= D_\infty(P_Z \| P_{Z|Y=y}) = D_\infty(P_{Z|X} \circ P_X \| P_{Z|X} \circ P_{X|Y=y}) \\ &\leq D_\infty(P_X \| P_{X|Y=y}) = \Lambda(X \rightarrow y), \end{aligned}$$

where the inequality is due to the data-processing inequality for the Rényi divergence [61, Thm. 9].

6. Consider the Markov chain  $X - Y - Z$ . The kernel  $P_{Z|Y}$  induces distributions  $P_{XZ} = P_{Z|Y} \circ P_{XY}$  and  $P_Z = P_{Z|Y} \circ P_Y$ . Hence, we can write

$$\begin{aligned} \operatorname{ess\,sup}_{P_Z} \Lambda(X \rightarrow Z) &= D_\infty(P_X \times P_Z \| P_{XZ}) = D_\infty(P_X \times (P_{Z|Y} \circ P_Y) \| P_{Z|Y} \circ P_{XY}) \\ &= D_\infty(P_{Z|Y} \circ (P_X \times P_Y) \| P_{Z|Y} \circ P_{XY}) \leq D_\infty(P_X \times P_Y \| P_{XY}) \\ &= \operatorname{ess\,sup}_{P_Y} \Lambda(X \rightarrow Y), \end{aligned}$$

where the inequality is due to the data-processing inequality for the Rényi divergence [61, Thm. 9].

7. Fix  $y_1 \in \mathcal{Y}_1$  and  $y_2 \in \mathcal{Y}_2$ . First, suppose  $P_X \ll P_{X|Y_1=y_1}$ . We can express the PMC using densities:

$$\Lambda(X \rightarrow y_1, y_2) = \log \operatorname{ess\,sup}_{P_X} \frac{f_{Y_1, Y_2}(y_1, y_2)}{f_{Y_1, Y_2|X}(y_1, y_2 | X)}$$

$$\begin{aligned}
&= \log \operatorname{ess\,sup}_{P_X} \frac{f_{Y_1}(y_1) f_{Y_2|Y_1}(y_2 | y_1)}{f_{Y_1|X}(y_1 | X) f_{Y_2|Y_1,X}(y_2 | y_1, X)} \\
&= \operatorname{ess\,sup}_{P_X} \left( \log \frac{f_{Y_1}(y_1)}{f_{Y_1|X}(y_1 | X)} + \log \frac{f_{Y_2|Y_1}(y_2 | y_1)}{f_{Y_2|Y_1,X}(y_2 | y_1, X)} \right) \\
&\leq \log \operatorname{ess\,sup}_{P_X} \frac{f_{Y_1}(y_1)}{f_{Y_1|X}(y_1 | X)} + \log \operatorname{ess\,sup}_{P_X} \frac{f_{Y_2|Y_1}(y_2 | y_1)}{f_{Y_2|Y_1,X}(y_2 | y_1, X)} \\
&\leq \log \operatorname{ess\,sup}_{P_X} \frac{f_{Y_1}(y_1)}{f_{Y_1|X}(y_1 | X)} + \log \operatorname{ess\,sup}_{P_{X|Y_1=y_1}} \frac{f_{Y_2|Y_1}(y_2 | y_1)}{f_{Y_2|Y_1,X}(y_2 | y_1, X)}, \quad (26) \\
&= \Lambda(X \rightarrow y_1) + \Lambda(X \rightarrow y_2 | y_1).
\end{aligned}$$

where (26) is due to the assumption that  $P_X \ll P_{X|Y_1=y_1}$ . Next, suppose  $P_X \not\ll P_{X|Y_1=y_1}$ . Then,  $\Lambda(X \rightarrow y_1) = D_\infty(P_X \| P_{X|Y_1=y_1}) = \infty$ , which results in the trivial inequality  $\Lambda(X \rightarrow y_1, y_2) \leq \infty$ .

#### APPENDIX E

##### PROOF OF PROPOSITION 3

If  $P_{Y|X}$  satisfies  $\varepsilon$ -LDP, then it holds that

$$\frac{P_{Y|X=x}(y)}{P_{Y|X=x'}(y)} \leq e^\varepsilon, \quad x, x' \in \mathcal{X}, y \in \mathcal{Y}.$$

Fix  $x \in \mathcal{X}$  and  $y \in \mathcal{Y}$ . We can write

$$\begin{aligned}
\frac{P_Y(y)}{P_{Y|X=x}(y)} &= \frac{\sum_{x' \in \mathcal{X}} P_{Y|X=x'}(y) P_X(x')}{P_{Y|X=x}(y)} \\
&\leq \frac{P_{Y|X=x}(y) P_X(x) + e^\varepsilon P_{Y|X=x}(y) (1 - P_X(x))}{P_{Y|X=x}(y)} \\
&= e^\varepsilon - P_X(x) (e^\varepsilon - 1).
\end{aligned}$$

Maximizing over  $x \in \mathcal{X}$  on both sides and taking the logarithm, we get

$$\Lambda(X \rightarrow y) = \log \max_{x \in \mathcal{X}} \frac{P_Y(y)}{P_{Y|X=x}(y)} \leq \log \left( e^\varepsilon - p_{\min} (e^\varepsilon - 1) \right).$$

#### APPENDIX F

##### PROOF OF THEOREM 4

1. Fix some  $y \in \mathcal{Y}$  and  $x^* \in \mathcal{X}$ . Since  $P_{Y|X}$  satisfies  $\varepsilon_u$ -PML we have  $i(x; y) \leq \varepsilon_u$  for all  $x \in \mathcal{X}$ . Thus, we can write

$$\begin{aligned}
1 &= \sum_{x \in \mathcal{X}} P_{X|Y=y}(x) = P_{X|Y=y}(x^*) + \sum_{x \in \mathcal{X} \setminus \{x^*\}} P_{X|Y=y}(x) \\
&\leq P_{X|Y=y}(x^*) + \sum_{x \in \mathcal{X} \setminus \{x^*\}} e^{\varepsilon_u} P_X(x) \\
&= P_{X|Y=y}(x^*) + e^{\varepsilon_u} (1 - P_X(x^*)).
\end{aligned}$$

Rearranging, we get

$$\frac{P_X(x^*)}{P_{X|Y=y}(x^*)} \leq \frac{P_X(x^*)}{1 - e^{\varepsilon_u} (1 - P_X(x^*))}.$$

This yields the following upper bound on the PMC:

$$\Lambda(X \rightarrow y) = \log \max_{x \in \mathcal{X}} \frac{P_X(x)}{P_{X|Y=y}(x)} \leq \log \max_{x \in \mathcal{X}} \frac{P_X(x)}{1 - e^{\varepsilon_u}(1 - P_X(x))} = \log \frac{p_{\min}}{1 - e^{\varepsilon_u}(1 - p_{\min})},$$

where the last equality is due to the fact that the mapping  $p \mapsto \frac{p}{1 - e^{\varepsilon_u}(1 - p)}$  is decreasing for  $0 \leq \varepsilon_u < \log \frac{1}{1 - p_{\min}}$ .

- Fix some  $y \in \mathcal{Y}$  and  $x^* \in \mathcal{X}$ . Since  $P_{Y|X}$  satisfies  $\varepsilon_l$ -PMC we have  $i(x; y) \geq -\varepsilon_l$  for all  $x \in \mathcal{X}$ . Thus, we can write

$$\begin{aligned} 1 &= \sum_{x \in \mathcal{X}} P_{X|Y=y}(x) = P_{X|Y=y}(x^*) + \sum_{x \in \mathcal{X} \setminus \{x^*\}} P_{X|Y=y}(x) \\ &\geq P_{X|Y=y}(x^*) + \sum_{x \in \mathcal{X} \setminus \{x^*\}} e^{-\varepsilon_l} P_X(x) \\ &= P_{X|Y=y}(x^*) + e^{-\varepsilon_l} (1 - P_X(x^*)). \end{aligned}$$

Rearranging, we get

$$\frac{P_{X|Y=y}(x^*)}{P_X(x^*)} \leq \frac{1 - e^{-\varepsilon_l}(1 - P_X(x^*))}{P_X(x^*)}.$$

This yields the following upper bound on the PML:

$$\ell(X \rightarrow y) = \log \max_{x \in \mathcal{X}} \frac{P_{X|Y=y}(x)}{P_X(x)} \leq \log \max_{x \in \mathcal{X}} \frac{1 - e^{-\varepsilon_l}(1 - P_X(x))}{P_X(x)} = \log \frac{1 - e^{-\varepsilon_l}(1 - p_{\min})}{p_{\min}},$$

where the last equality is due to the fact that the mapping  $p \mapsto \frac{1 - e^{-\varepsilon_l}(1 - p)}{p}$  is decreasing for  $\varepsilon_l \geq 0$ .

## APPENDIX G

### PROOF OF THEOREM 5

To prove the theorem, we show that if  $\text{ess sup}_{P_Y} \ell(X \rightarrow Y) = \infty$ , then  $\text{ess sup}_{P_Y} \Lambda(X \rightarrow Y) = \infty$  as well. To do this, first, we describe the implications of having an unbounded PML. Note that we can write

$$\text{ess sup}_{P_Y} \ell(X \rightarrow Y) = D_{\infty}(P_{XY} \| P_X \times P_Y) = \log \sup_{\mathcal{E} \in \mathcal{S}_Y} \frac{\text{ess sup}_{P_X} P_{Y|X}(\mathcal{E} | X)}{P_Y(\mathcal{E})}.$$

Hence, assuming that  $P_{XY} \ll P_X \times P_Y$ ,  $\ell(X \rightarrow Y)$  is unbounded if and only if one of the following two conditions hold: either

- there exists a decreasing sequence of events  $\mathcal{A}_1, \mathcal{A}_2, \dots \in \mathcal{S}_Y$  and a set  $\mathcal{B} \in \mathcal{S}_X$  with  $P_X(\mathcal{B}) > 0$  such that  $P_Y(\mathcal{A}_i) \rightarrow 0$  but  $P_{Y|X=x}(\mathcal{A}_i) \not\rightarrow 0$  for all  $x \in \mathcal{B}$ , or
- there exists a decreasing sequence of events  $\mathcal{A}_1, \mathcal{A}_2, \dots \in \mathcal{S}_Y$  and a set  $\mathcal{B} \in \mathcal{S}_X$  with  $P_X(\mathcal{B}) > 0$  such that  $P_Y(\mathcal{A}_i) \rightarrow 0$ , and  $P_{Y|X=x}(\mathcal{A}_i) \rightarrow 0$ , but  $P_{Y|X=x}(\mathcal{A}_i)$  converges at a strictly slower rate compared to  $P_Y(\mathcal{A}_i)$  for all  $x \in \mathcal{B}$ .

Next, we argue that the first case can never happen when  $P_{XY} \ll P_X \times P_Y$ . Recall that  $P_{XY} \ll P_X \times P_Y$  implies that  $P_{Y|X=x} \ll P_Y$  almost surely. Fix  $x \in \mathcal{X}$  and let  $f = \frac{dP_{Y|X=x}}{dP_Y}$ . Let  $(\mathcal{A}_i)_{i \in \mathbb{N}}$  be a sequence of events with  $\lim_{i \rightarrow \infty} P_Y(\mathcal{A}_i) = 0$ . Observe that

$$\lim_{i \rightarrow \infty} P_{Y|X=x}(\mathcal{A}_i) = \lim_{i \rightarrow \infty} \int_{\mathcal{Y}} \mathbf{1}_{\mathcal{A}_i}(y) P_{Y|X=x}(dy) = \lim_{i \rightarrow \infty} \int_{\mathcal{Y}} \mathbf{1}_{\mathcal{A}_i}(y) f(y) P_Y(dy).$$



When  $f$  is bounded  $\lim_{i \rightarrow \infty} P_Y(\mathcal{A}_i) = 0$  clearly implies that  $\lim_{i \rightarrow \infty} P_{Y|X=x}(\mathcal{A}_i) = 0$ . Furthermore, the argument carries over to arbitrary  $P_Y$ -integrable  $f$  by first writing  $f$  as the limit of a sequence of bounded functions, and then using the dominated convergence theorem.

We conclude that  $\ell(X \rightarrow Y)$  is unbounded if and only if the second condition above holds. Observe that if there exists a decreasing sequence of events  $(\mathcal{A}_i)_{i \in \mathbb{N}}$  and a non-negligible set  $\mathcal{B} \in \mathcal{S}_X$  such that  $P_{Y|X=x}(\mathcal{A}_i)$  converges to zero more slowly than  $P_Y(\mathcal{A}_i)$  for all  $x \in \mathcal{B}$ , then there exists a non-negligible set  $\mathcal{C} \in \mathcal{S}_X$  such that  $P_{Y|X=x}(\mathcal{A}_i)$  converges to zero faster than  $P_Y(\mathcal{A}_i)$  for all  $x \in \mathcal{C}$ . Thus, we can write

$$\begin{aligned} \operatorname{ess\,sup}_{P_Y} \Lambda(X \rightarrow Y) &= D_\infty(P_X \times P_Y \| P_{XY}) \\ &= \log \sup_{\mathcal{E} \in \mathcal{S}_Y} \frac{P_Y(\mathcal{E})}{\operatorname{ess\,inf}_{P_X} P_{Y|X} \mathcal{E} | X} \\ &\geq \log \lim_{i \rightarrow \infty} \frac{P_Y(\mathcal{A}_i)}{\operatorname{ess\,inf}_{P_X} P_{Y|X}(\mathcal{A}_i | X)} \\ &\geq \log \lim_{i \rightarrow \infty} \frac{P_Y(\mathcal{A}_i)}{\operatorname{ess\,sup}_{P_X} \left( \mathbf{1}_{\mathcal{C}}(X) P_{Y|X}(\mathcal{A}_i | X) \right)} \end{aligned} \quad (27a)$$

$$= \infty, \quad (27b)$$

where (27a) is because the essential infimum of a function is at least as small as its essential supremum over a non-negligible set, and (27b) follows from the definition of set  $\mathcal{C}$ .

## APPENDIX H PROOF OF THEOREM 6

We show that

$$\operatorname{ess\,sup}_{P_Y} \Lambda(X \rightarrow Y) = \log \sup_{\mathcal{E} \in \mathcal{S}_Y} \frac{P_Y(\mathcal{E})}{\operatorname{ess\,inf}_{P_X} P_{Y|X}(\mathcal{E} | X)} \quad (28)$$

is finite if and only if

$$\log \operatorname{ess\,sup}_{P_X \times P_X} \sup_{\mathcal{E} \in \mathcal{S}_Y} \frac{P_{Y|X}(\mathcal{E} | X)}{P_{Y|X}(\mathcal{E} | X')} = \log \sup_{\mathcal{E} \in \mathcal{S}_Y} \frac{\operatorname{ess\,sup}_{P_X} P_{Y|X}(\mathcal{E} | X)}{\operatorname{ess\,inf}_{P_X} P_{Y|X}(\mathcal{E} | X')}, \quad (29)$$

is finite. Since  $P_Y(\mathcal{E}) = \mathbb{E}_{X \sim P_X} [P_{Y|X}(\mathcal{E} | X)] \leq \operatorname{ess\,sup}_{P_X} P_{Y|X}(\mathcal{E} | X)$  for all  $\mathcal{E} \in \mathcal{S}_Y$ , then the boundedness of (29) implies the boundedness of (28). To prove the other direction, note that if

$$\operatorname{ess\,sup}_{P_X \times P_X} \sup_{\mathcal{E} \in \mathcal{S}_Y} \frac{P_{Y|X}(\mathcal{E} | X)}{P_{Y|X}(\mathcal{E} | X')} = \infty,$$

then

$$(P_X \times P_X) \left( \left\{ \sup_{\mathcal{E} \in \mathcal{S}_Y} \frac{P_{Y|X}(\mathcal{E} | X)}{P_{Y|X}(\mathcal{E} | X')} > M \right\} \right) > 0, \quad (30)$$

for arbitrarily large  $M \geq 0$ . Put differently, it holds that<sup>7</sup>

$$(P_X \times P_X) \left( \left\{ \sup_{\mathcal{E} \in \mathcal{S}_Y} \frac{P_{Y|X}(\mathcal{E} | X)}{P_{Y|X}(\mathcal{E} | X')} = \infty \right\} \right) > 0.$$

This, in turn, implies that there exist sets  $\mathcal{B}, \mathcal{C} \in \mathcal{S}_X$  with  $P_X(\mathcal{B})P_X(\mathcal{C}) > 0$  such that

$$\sup_{\mathcal{E} \in \mathcal{S}_Y} \frac{P_{Y|X=x}(\mathcal{E})}{P_{Y|X=x'}(\mathcal{E})} = \infty,$$

for all  $x \in \mathcal{B}$  and  $x' \in \mathcal{C}$ . Hence, (29) is infinite if and only if one of the following two conditions hold: either

1. there exists an event  $\mathcal{A} \in \mathcal{S}_Y$  such that  $P_{Y|X=x}(\mathcal{A}) > 0$  for all  $x \in \mathcal{B}$  and  $P_{Y|X=x'}(\mathcal{A}) = 0$  for all  $x' \in \mathcal{C}$ , or
2. there exists a decreasing sequence of events  $\mathcal{A}_1, \mathcal{A}_2, \dots \in \mathcal{S}_Y$  such that  $P_{Y|X=x}(\mathcal{A}_i) \rightarrow 0$  for all  $x \in \mathcal{B}$  and  $P_{Y|X=x'}(\mathcal{A}_i) \rightarrow 0$  for all  $x' \in \mathcal{C}$  as  $i \rightarrow \infty$ , but the convergence on  $\mathcal{B}$  is at a strictly slower rate compared to the convergence on  $\mathcal{C}$ .

Next, we argue that PMC is unbounded under both conditions. Under the first condition, we have

$$P_Y(\mathcal{A}) = \mathbb{E} \left[ P_{Y|X}(\mathcal{A} | X) \right] \geq \mathbb{E} \left[ \mathbf{1}_{\mathcal{B}}(X) P_{Y|X}(\mathcal{A} | X) \right] > 0,$$

and

$$\operatorname{ess\,inf}_{P_X} P_{Y|X}(\mathcal{A} | X') = 0.$$

Thus, we get

$$\operatorname{ess\,sup}_{P_Y} \Lambda(X \rightarrow Y) \geq \log \frac{P_Y(\mathcal{A})}{\operatorname{ess\,inf}_{P_X} P_{Y|X}(\mathcal{A} | X')} = \infty.$$

Next, let  $\mathcal{S}_{\mathcal{B}} := \{\mathcal{F} \cap \mathcal{B} : \mathcal{F} \in \mathcal{S}_X\}$  denote the trace of  $\mathcal{S}_X$  on  $\mathcal{B}$ . Let the measure  $Q(\mathcal{F}) := P_X(\mathcal{F})$  for  $\mathcal{F} \in \mathcal{S}_{\mathcal{B}}$  denote the restriction of  $P_X$  to  $\mathcal{B}$ . If the second condition is satisfied, then we can write

$$\begin{aligned} \operatorname{ess\,sup}_{P_Y} \Lambda(X \rightarrow Y) &= \log \sup_{\mathcal{E} \in \mathcal{S}_Y} \frac{P_Y(\mathcal{E})}{\operatorname{ess\,inf}_{P_X} P_{Y|X}(\mathcal{E} | X')} \\ &\geq \log \lim_{i \rightarrow \infty} \frac{P_Y(\mathcal{A}_i)}{\operatorname{ess\,inf}_{P_X} P_{Y|X}(\mathcal{A}_i | X')} \\ &= \log \lim_{i \rightarrow \infty} \frac{\int_{\mathcal{X}} P_{Y|X=x}(\mathcal{A}_i) P_X(dx)}{\operatorname{ess\,inf}_{P_X} P_{Y|X}(\mathcal{A}_i | X')} \\ &\geq \log \lim_{i \rightarrow \infty} \frac{\int_{\mathcal{B}} P_{Y|X=x}(\mathcal{A}_i) P_X(dx)}{\operatorname{ess\,inf}_{P_X} P_{Y|X}(\mathcal{A}_i | X')} \\ &\geq \log \lim_{i \rightarrow \infty} \frac{P_X(\mathcal{B}) \left( \operatorname{ess\,inf}_Q P_{Y|X}(\mathcal{A}_i | X) \right)}{\operatorname{ess\,inf}_{P_X} P_{Y|X}(\mathcal{A}_i | X')} \\ &\geq \log \lim_{i \rightarrow \infty} \frac{P_X(\mathcal{B}) \left( \operatorname{ess\,inf}_Q P_{Y|X}(\mathcal{A}_i | X) \right)}{\operatorname{ess\,sup}_{P_X} \left( \mathbf{1}_{\mathcal{C}}(X') P_{Y|X}(\mathcal{A}_i | X') \right)} \end{aligned} \tag{31a}$$

<sup>7</sup>If  $(P_X \times P_X) \left( \left\{ \sup_{\mathcal{E} \in \mathcal{S}_Y} \frac{P_{Y|X}(\mathcal{E} | X)}{P_{Y|X}(\mathcal{E} | X')} = \infty \right\} \right) = 0$ , then there exists  $A \geq 0$  such that  $\frac{P_{Y|X}(\mathcal{E} | X)}{P_{Y|X}(\mathcal{E} | X')} \leq A$  almost surely for all  $\mathcal{E} \in \mathcal{S}_Y$ . This contradicts (30).

$$= \infty, \tag{31b}$$

where (31a) is because the essential infimum of a function is at least as small as its essential supremum over a non-negligible set, and (31b) follows by assumption. We conclude that PMC is essentially bounded if and only if it satisfies  $\varepsilon$ -LDP for a finite value of  $\varepsilon$ .

## REFERENCES

- [1] M. Rigaki and S. Garcia, "A survey of privacy attacks in machine learning," *ACM Computing Surveys*, vol. 56, no. 4, pp. 1–34, 2023.
- [2] L. Sweeney, " $k$ -anonymity: A model for protecting privacy," *International journal of uncertainty, fuzziness and knowledge-based systems*, vol. 10, no. 05, pp. 557–570, 2002.
- [3] A. Machanavajjhala, D. Kifer, J. Gehrke, and M. Venkatasubramanian, " $l$ -diversity: Privacy beyond  $k$ -anonymity," *ACM Transactions on Knowledge Discovery from Data (TKDD)*, vol. 1, no. 1, pp. 3–es, 2007.
- [4] N. Li, T. Li, and S. Venkatasubramanian, " $t$ -closeness: Privacy beyond  $k$ -anonymity and  $l$ -diversity," in *IEEE 23rd international conference on data engineering*, 2007, pp. 106–115.
- [5] C. Dwork, F. McSherry, K. Nissim, and A. Smith, "Calibrating noise to sensitivity in private data analysis," in *Theory of Cryptography Conference (TCC)*. Springer Berlin Heidelberg, 2006, pp. 265–284.
- [6] C. Dwork, A. Roth *et al.*, "The algorithmic foundations of differential privacy," *Foundations and Trends® in Theoretical Computer Science*, vol. 9, no. 3–4, pp. 211–407, 2014.
- [7] S. P. Kasiviswanathan, H. K. Lee, K. Nissim, S. Raskhodnikova, and A. Smith, "What can we learn privately?" in *49th Annual IEEE Symposium on Foundations of Computer Science*, 2008, pp. 531–540.
- [8] J. C. Duchi, M. I. Jordan, and M. J. Wainwright, "Local privacy and statistical minimax rates," in *IEEE 54th Annual Symposium on Foundations of Computer Science*, 2013, pp. 429–438.
- [9] L. Wasserman and S. Zhou, "A Statistical Framework for Differential Privacy," *Journal of the American Statistical Association*, vol. 105, no. 489, pp. 375–389, Mar. 2010.
- [10] P. Kairouz, S. Oh, and P. Viswanath, "The composition theorem for differential privacy," in *International conference on machine learning*. PMLR, 2015, pp. 1376–1385.
- [11] M. C. Tschantz, S. Sen, and A. Datta, "Sok: Differential privacy as a causal property," in *IEEE Symposium on Security and Privacy (SP)*, 2020, pp. 354–371.
- [12] S. Saeidian, G. Cervia, T. J. Oechtering, and M. Skoglund, "Pointwise maximal leakage," *IEEE Transactions on Information Theory*, vol. 69, no. 12, pp. 8054–8080, 2023.
- [13] —, "Pointwise maximal leakage on general alphabets," in *IEEE International Symposium on Information Theory (ISIT)*, 2023, pp. 388–393.
- [14] M. S. Alvim, K. Chatzikokolakis, C. Palamidessi, and G. Smith, "Measuring information leakage using generalized gain functions," in *IEEE 25th Computer Security Foundations Symposium*, 2012, pp. 265–279.
- [15] F. du Pin Calmon and N. Fawaz, "Privacy against statistical inference," in *50th Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, 2012, pp. 1401–1408.
- [16] B. Jiang, M. Li, and R. Tandon, "Context-aware data aggregation with localized information privacy," in *IEEE Conference on Communications and Network Security (CNS)*, 2018, pp. 1–9.
- [17] —, "Local information privacy and its application to privacy-preserving data aggregation," *IEEE Transactions on Dependable and Secure Computing*, vol. 19, no. 3, pp. 1918–1935, 2020.
- [18] B. Jiang, M. Seif, R. Tandon, and M. Li, "Context-aware local information privacy," *IEEE Transactions on Information Forensics and Security*, vol. 16, pp. 3694–3708, 2021.
- [19] M. A. Zarrabian, N. Ding, and P. Sadeghi, "On the lift, related privacy measures, and applications to privacy–utility trade-offs," *Entropy*, vol. 25, no. 4, p. 679, 2023.
- [20] —, "Asymmetric local information privacy and the watchdog mechanism," in *IEEE Information Theory Workshop (ITW)*, 2022, pp. 7–12.
- [21] S. Saeidian, G. Cervia, T. J. Oechtering, and M. Skoglund, "Rethinking disclosure prevention with pointwise maximal leakage," *Submitted to: Journal of Privacy and Confidentiality*, 2023.

- [22] I. Issa, A. B. Wagner, and S. Kamath, “An operational approach to information leakage,” *IEEE Transactions on Information Theory*, vol. 66, no. 3, pp. 1625–1657, 2020.
- [23] G. R. Kurri, M. Managoli, and V. M. Prabhakaran, “Maximal guesswork leakage,” *arXiv preprint arXiv:2405.02585*, 2024.
- [24] S. L. Warner, “Randomized response: A survey technique for eliminating evasive answer bias,” *Journal of the American Statistical Association*, vol. 60, no. 309, pp. 63–69, 1965.
- [25] P. Kairouz, S. Oh, and P. Viswanath, “Extremal mechanisms for local differential privacy,” *The Journal of Machine Learning Research*, vol. 17, no. 1, pp. 492–542, 2016.
- [26] M. S. Alvim, K. Chatzikokolakis, A. McIver, C. Morgan, C. Palamidessi, and G. Smith, *The Science of Quantitative Information Flow*. Springer Cham, 2020.
- [27] G. Smith, “On the foundations of quantitative information flow,” in *International Conference on Foundations of Software Science and Computational Structures*. Springer, 2009, pp. 288–302.
- [28] C. Braun, K. Chatzikokolakis, and C. Palamidessi, “Quantitative notions of leakage for one-try attacks,” *Electronic Notes in Theoretical Computer Science*, vol. 249, pp. 75–91, 2009.
- [29] B. Espinoza and G. Smith, “Min-entropy as a resource,” *Information and Computation*, vol. 226, pp. 57–75, 2013.
- [30] M. S. Alvim, K. Chatzikokolakis, A. McIver, C. Morgan, C. Palamidessi, and G. Smith, “Additive and multiplicative notions of leakage, and their capacities,” in *IEEE 27th Computer Security Foundations Symposium*, 2014, pp. 308–322.
- [31] J. Liao, O. Kosut, L. Sankar, and F. du Pin Calmon, “Tunable measures for information leakage and applications to privacy-utility tradeoffs,” *IEEE Transactions on Information Theory*, vol. 65, no. 12, pp. 8043–8066, 2019.
- [32] A. Gilani, G. R. Kurri, O. Kosut, and L. Sankar, “Unifying privacy measures via maximal  $(\alpha, \beta)$ -leakage (Möbel),” *IEEE Transactions on Information Theory*, vol. 70, no. 6, pp. 4368–4395, 2024.
- [33] G. R. Kurri, L. Sankar, and O. Kosut, “An operational approach to information leakage via generalized gain functions,” *IEEE Transactions on Information Theory*, pp. 1–1, 2023.
- [34] S. Asoodeh, F. Alajaji, and T. Linder, “Notes on information-theoretic privacy,” in *52nd Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, 2014, pp. 1272–1278.
- [35] S. Asoodeh, M. Diaz, F. Alajaji, and T. Linder, “Information extraction under privacy constraints,” *Information*, vol. 7, no. 1, p. 15, 2016.
- [36] S. Asoodeh, F. Alajaji, and T. Linder, “On maximal correlation, mutual information and data privacy,” in *IEEE 14th Canadian workshop on information theory (CWIT)*, 2015, pp. 27–31.
- [37] W. Wang, L. Ying, and J. Zhang, “On the relation between identifiability, differential privacy, and mutual-information privacy,” *IEEE Transactions on Information Theory*, vol. 62, no. 9, pp. 5018–5029, 2016.
- [38] A. Makhdoumi, S. Salamatian, N. Fawaz, and M. Médard, “From the information bottleneck to the privacy funnel,” in *IEEE Information Theory Workshop (ITW)*, 2014, pp. 501–505.
- [39] J. Liao, L. Sankar, V. Y. Tan, and F. du Pin Calmon, “Hypothesis testing under mutual information privacy constraints in the high privacy regime,” *IEEE Transactions on Information Forensics and Security*, vol. 13, no. 4, pp. 1058–1071, 2017.
- [40] B. Rassouli and D. Gündüz, “On perfect privacy,” *IEEE Journal on Selected Areas in Information Theory*, vol. 2, no. 1, pp. 177–191, 2021.
- [41] M. Diaz, H. Wang, F. P. Calmon, and L. Sankar, “On the robustness of information-theoretic privacy measures and mechanisms,” *IEEE Transactions on Information Theory*, vol. 66, no. 4, pp. 1949–1978, 2020.
- [42] F. du Pin Calmon, A. Makhdoumi, M. Médard, M. Varia, M. Christiansen, and K. R. Duffy, “Principal inertia components and applications,” *IEEE Transactions on Information Theory*, vol. 63, no. 8, pp. 5011–5038, 2017.
- [43] H. Wang, L. Vo, F. P. Calmon, M. Médard, K. R. Duffy, and M. Varia, “Privacy with estimation guarantees,” *IEEE Transactions on Information Theory*, vol. 65, no. 12, pp. 8025–8042, 2019.
- [44] B. Rassouli and D. Gündüz, “Optimal utility-privacy trade-off with total variation distance as a privacy measure,” *IEEE Transactions on Information Forensics and Security*, vol. 15, pp. 594–603, 2019.
- [45] A. Zamani, T. J. Oechtering, and M. Skoglund, “Data disclosure with non-zero leakage and non-invertible leakage matrix,” *IEEE Transactions on Information Forensics and Security*, vol. 17, pp. 165–179, 2021.
- [46] —, “A design framework for strongly  $\chi^2$ -private data disclosure,” *IEEE Transactions on Information Forensics and Security*, vol. 16, pp. 2312–2325, 2021.
- [47] —, “On the privacy-utility trade-off with and without direct access to the private data,” *IEEE Transactions on Information Theory*, 2023.
- [48] S. Asoodeh, M. Diaz, F. Alajaji, and T. Linder, “Estimation efficiency under privacy constraints,” *IEEE Transactions on Information*

- Theory*, vol. 65, no. 3, pp. 1512–1534, 2018.
- [49] J. L. Massey, “Guessing and entropy,” in *IEEE International Symposium on Information Theory*, 1994, p. 204.
- [50] D. Malone and W. G. Sullivan, “Guesswork and entropy,” *IEEE Transactions on Information Theory*, vol. 50, no. 3, pp. 525–526, 2004.
- [51] K. Kalantari, L. Sankar, and A. D. Sarwate, “Robust privacy-utility tradeoffs under differential privacy and hamming distortion,” *IEEE Transactions on Information Forensics and Security*, vol. 13, no. 11, pp. 2816–2830, 2018.
- [52] J. Goseling and M. Lopuhaä-Zwakenberg, “Robust optimization for local differential privacy,” in *IEEE International Symposium on Information Theory (ISIT)*, 2022, pp. 1629–1634.
- [53] M. Lopuhaä-Zwakenberg and J. Goseling, “Mechanisms for robust local differential privacy,” *Entropy*, vol. 26, no. 3, 2024.
- [54] B. Wu, A. B. Wagner, and G. E. Suh, “Optimal mechanisms under maximal leakage,” in *IEEE Conference on Communications and Network Security (CNS)*, 2020, pp. 1–6.
- [55] S. Saeidian, G. Cervia, T. J. Oechtering, and M. Skoglund, “Optimal maximal leakage-distortion tradeoff,” in *IEEE Information Theory Workshop (ITW)*, 2021, pp. 1–6.
- [56] H. Hsu, S. Asodeh, and F. P. Calmon, “Information-theoretic privacy watchdogs,” in *IEEE International Symposium on Information Theory (ISIT)*, 2019, pp. 552–556.
- [57] P. Sadeghi, N. Ding, and T. Rakotoarivelo, “On properties and optimization of information-theoretic privacy watchdog,” in *IEEE Information Theory Workshop (ITW)*, 2021, pp. 1–5.
- [58] A. Klenke, *Probability theory: A comprehensive course*. Springer Science & Business Media, 2013.
- [59] E. Çinlar, *Probability and stochastics*. Springer New York, NY, 2011, vol. 261.
- [60] A. Rényi, “On measures of entropy and information,” in *Fourth Berkeley Symposium on Mathematical Statistics and Probability, Volume 1: Contributions to the Theory of Statistics*, vol. 4. University of California Press, 1961, pp. 547–562.
- [61] T. van Erven and P. Harremoës, “Rényi divergence and Kullback-Leibler divergence,” *IEEE Transactions on Information Theory*, vol. 60, no. 7, pp. 3797–3820, 2014.
- [62] L. Grosse, S. Saeidian, and T. J. Oechtering, “Extremal mechanisms for pointwise maximal leakage,” *arXiv preprint arXiv:2310.07381*, 2023.