Kyoto University

Asuka Takatsu asuka Takatsu asuka mwaka wa 1989, asuka wa mshindi wa mshindi wa mshindi wa mshindi wa mshindi w Tokyo Metropolitan University

Han Bao bao@i.kyoto-u.ac.jp

# Abstract

A fundamental challenge in machine learning is the choice of a loss as it characterizes our learning task, is minimized in the training phase, and serves as an evaluation criterion for estimators. Proper losses are commonly chosen, ensuring minimizers of the full risk match the true probability vector. Estimators induced from a proper loss are widely used to construct forecasters for downstream tasks such as classification and ranking. In this procedure, how does the forecaster based on the obtained estimator perform well under a given downstream task? This question is substantially relevant to the behavior of the p-norm between the estimated and true probability vectors when the estimator is updated. In the proper loss framework, the suboptimality of the estimated probability vector from the true probability vector is measured by a surrogate regret. First, we analyze a surrogate regret and show that the *strict* properness of a loss is necessary and sufficient to establish a non-vacuous surrogate regret bound. Second, we solve an important open question that the order of convergence in  $p$ -norm cannot be faster than the  $1/2$ -order of surrogate regrets for a broad class of strictly proper losses. This implies that strongly proper losses entail the optimal convergence rate.

Keywords: loss functions, proper scoring rules, supervised learning, surrogate regret bounds, convex analysis

# 1 Introduction

Proper losses, also known as proper scoring rules, are measurements of the quality of a probabilistic prediction given a true probability vector [\[BSS05,](#page-23-0) [GR07,](#page-23-1) [RW10\]](#page-25-0). Intuitively, we say a loss is proper if the target probability vector is its minimizer, and strictly proper if the minimizer is unique, which is a basic property for a reasonable loss. Proper losses are prevailing in modern machine learning: for example, the cross-entropy loss popular in deep learning essentially corresponds to the log loss (or logarithmic score), and the Brier score is used for assessing model uncertainties  $[OFR<sup>+</sup>19]$  $[OFR<sup>+</sup>19]$ . As such, probabilistic estimators are obtained via proper loss minimization. It is common to post-process a minimizer of a proper loss for downstream tasks, such as classification (by choosing the most likely label), ranking (by giving ranking scores to each label [\[NA13\]](#page-24-1)), F-measure optimization (by thresholding the estimated probability [\[KNRD14\]](#page-24-2)), and probability calibration [\[KSFF17,](#page-24-3) [BGHN23\]](#page-23-2). Here, we are interested in the predictive performance of post-processed estimators in downstream tasks. Given a true and estimated probability vectors q and  $\hat{q}$ , respectively, the *surrogate* regret  $R(\mathbf{q}, \hat{\mathbf{q}})$  (introduced in [§3\)](#page-4-0) measures the suboptimality of  $\hat{\mathbf{q}}$  from  $\mathbf{q}$  in terms of a proper loss. Can we relate the suboptimality of a forecaster for a downstream task to the surrogate regret?

<sup>©2024</sup> Han Bao and Asuka Takatsu.

License: CC-BY 4.0, see <https://creativecommons.org/licenses/by/4.0/>.

Surrogate regret bounds relate the surrogate regret to the performance for downstream tasks, and have been derived for binary classification [\[Zha04,](#page-25-1) [RW09\]](#page-25-2), bipartite ranking [\[Aga14\]](#page-21-0), property elicitation [\[AA15\]](#page-21-1), F-measure optimization [\[KD16,](#page-23-3) [ZRA20\]](#page-25-3), and learning with noisy labels [\[ZLA21\]](#page-25-4), independently. Recently, a unified surrogate regret bound across different downstream tasks has been established [\[Bao23\]](#page-22-0), where surrogate regret bounds are unified in terms of the 1-norm. This is based on the observation that the suboptimality of the aforementioned downstream tasks can be controlled by the 1-norm. However, the derived bound has been limited to the binary classification case, and it remains unclear when the surrogate regret bound is non-vacuous. A reasonable loss should entail a non-vacuous regret bound, which is crucial to tackling numerous downstream tasks simultaneously. Moreover, an important conjecture that the convergence rate of surrogate regret bounds cannot be faster than the 1{2-order has yet to be solved. This conjecture has a significant role in the choice of losses because the lower bound of the order of convergence contributes to delineating the optimality of a given proper loss.

In this article, we aim to study when the surrogate regret bounds are non-vacuous and how fast the order of convergence in the p-norm can be. To this end, we analyze the p-norm bounds by the surrogate regret  $R(\mathbf{q}, \hat{\mathbf{q}})$  jointly with a rate function  $\psi$  in the following form by extending from the binary classification case [\[Bao23\]](#page-22-0) to the multiclass classification case:

<span id="page-1-0"></span>
$$
\|\mathbf{q} - \widehat{\mathbf{q}}\|_{p} \leq \psi(R(\mathbf{q}, \widehat{\mathbf{q}})).\tag{1}
$$

After formalizing these notions in [§3,](#page-4-0) we derive the surrogate regret bounds in [§4.](#page-9-0) To derive bounds of the form  $(1)$ , we introduce the *moduli of convexity* [\[Fig76\]](#page-23-4), which describe the information of its second derivative of convex functions, for convex functions defined on the probability simplex. The rate  $\psi$  in [\(1\)](#page-1-0) can be characterized by the modulus of a Bregman generator function associated with a proper loss  $\ell$  [\(Theorem 11\)](#page-11-0). To obtain a non-vacuous bound, we first show that the strict properness of a loss is necessary and sufficient to obtain a non-vacuous surrogate regret bound, or strictly increasing  $\psi$ , to put it differently [\(Theorem 9\)](#page-10-0). Whereas it has been known that non-strictly proper losses can achieve non-vacuous bounds for classification [\[RW11,](#page-25-5) Corollary 27], our sufficiency result argues that the strict properness is a minimal requirement for an estimate to be non-vacuous in terms of the p-norm. As our second main result, we provide an affirmative answer to the above conjecture: the optimal rate  $\psi(\rho)$  as  $\rho \downarrow 0$  is  $O(\rho^{1/2})$ , for a broad class of proper losses [\(Theorem 17\)](#page-17-0). This convergence rate has already been known for a restricted class of proper losses, known as strongly proper losses [\[Aga14\]](#page-21-0). Hence, our result ensures the asymptotic optimality of strongly proper losses. This gives an answer to the question, "Do we have an interesting loss that is strictly proper but not strongly proper?" [\[Bao22,](#page-22-1) §6.2.9]: there is no better proper loss outside of strongly proper losses, as long as we are concerned with the asymptotic rate of  $\psi$  in [Eq. \(1\).](#page-1-0)

#### 1.1 Organization and contributions of this article

The organization of this article and our contributions are summarized as follows.

• [§2:](#page-2-0) Notation and necessary backgrounds on convex analysis are summarized.

- [§3:](#page-4-0) Proper losses for multiclass classification are introduced. [Lemma 2](#page-5-0) characterizes the existence of minimizers for general losses, and [Proposition 5](#page-7-0) gives a self-contained and rigorous proof of the well-known representation of proper losses [\[Sav71\]](#page-25-6).
- [§4:](#page-9-0) [Theorem 11](#page-11-0) extends surrogate regret bounds for binary classification [\[Bao23\]](#page-22-0) to multiclass classification. This is achieved by extending the moduli of convexity to multivariate functions [\(Definition 8\)](#page-10-1). [Theorem 9](#page-10-0) is our first result, proving that the strict properness of a loss is a necessary and sufficient condition for an associated surrogate regret bound to be non-vacuous.

Then, the benefits of [Theorem 11](#page-11-0) are discussed in [§4.3.](#page-12-0) In particular, we can obtain the p-norm bound in the form of  $(1)$ , which can be used to control the performance of plug-in forecasters for downstream tasks such as multiclass classification, learning with noisy labels, and bipartite ranking.

- [§5:](#page-14-0) We evaluate the rate  $\psi(\rho)$  by power functions such as  $\rho^{1/s} \leq \psi(\rho) \leq \rho^{1/S}$  for some constants  $s, S > 0$ ,<sup>[1](#page-2-1)</sup> which is based on the Simonenko order function previously adopted [\[Bao23\]](#page-22-0). Our second main result roughly shows that  $s \geq 2$  [\(Theorem 17\)](#page-17-0), establishing the asymptotic optimality  $\psi(\rho) \gtrsim \rho^{1/2}$  of strongly proper losses.
- [§6:](#page-20-0) Several examples of convex functions to generate proper losses are discussed.

# <span id="page-2-0"></span>2 Background

In this section, we summarize the notation and basic properties of convex functions.

### 2.1 Notation

Throughout this article, fix  $N \in \mathbb{N}$  and  $p \in [1, \infty]$ . The Kronecker delta is denoted by  $\delta_{ij}$ . For  $k \in \mathbb{N}$ , we set  $[k] := \{1, 2, ..., k\}$ . A vector is denoted by bold-face such as  $\xi \in \mathbb{R}^N$ , and its n-th (scalar) component is written as non-bold  $\xi_n$  for each  $n \in [N]$ . The p-norm of  $\boldsymbol{\xi} \in \mathbb{R}^N$  is denoted by  $\|\boldsymbol{\xi}\|_p$ . For a topology on  $\mathbb{R}^N$ , we refer to one induced from the 2-norm, but it makes no difference whichever norm we choose. Similarly, a convexity of a function on  $(\mathbb{R}^N, \|\cdot\|_p)$  is determined independently of the choice of p. The standard inner product on  $\mathbb{R}^N$  is denoted by  $\langle \xi, \xi' \rangle := \sum_{n \in [N]} \xi_n \xi'_n$ . We introduce the notation

$$
\triangle^N := \left\{ \mathbf{q} \in \mathbb{R}^N \mid q_n \geq 0 \ (n \in [N]), \ \langle \mathbf{q}, \mathbf{1} \rangle = 1 \right\}, \quad \triangle^N_+ := \left\{ \mathbf{q} \in \triangle^N \mid q_n > 0 \ (n \in [N]) \right\},\
$$

where  $\mathbf{1} \in \mathbb{R}^N$  is the vector with each component being one. For  $\mathbf{q} \in \Delta^N$ , we denote by supp(q) the *support* of q, that is,

$$
supp(\mathbf{q}) \coloneqq \{ n \in [N] \mid q_n > 0 \}.
$$

We adhere to the convention that

 $\pm\infty \leq \pm\infty$ ,  $a \pm \infty = \pm\infty$ ,  $b \cdot (\pm\infty) = \pm\infty$ ,  $-b \cdot (\pm\infty) = \mp\infty$ ,  $0 \cdot (\pm\infty) = 0$ ,

<span id="page-2-1"></span><sup>1.</sup> In our notation,  $\psi_1 \lesssim \psi_2$  indicates the existence of an absolute constant  $C > 0$  such that  $C\psi_1 \leq \psi_2$ .

for  $a \in \mathbb{R}$  and  $b > 0$ . We use O and  $\Omega$  to always denote the *infinitesimal* asymptotic order. To be precise, for two functions  $\phi, \psi$  defined around  $0, \phi(\varepsilon) = \Omega(\psi(\varepsilon))$  as  $\varepsilon \downarrow 0$  should be understood as

$$
\liminf_{\varepsilon \downarrow 0} \left| \frac{\phi(\varepsilon)}{\psi(\varepsilon)} \right| > 0.
$$

### 2.2 Convex analysis

In this subsection, let  $f : \mathbb{R}^N \to (-\infty, \infty]$  denote a proper convex function on  $\mathbb{R}^N$ . Its effective domain is defined by

$$
\mathrm{dom}\, f \coloneqq \left\{ \boldsymbol{\xi} \in \mathbb{R}^N \mid f(\boldsymbol{\xi}) < \infty \right\}.
$$

A vector  $\mathbf{v} \in \mathbb{R}^N$  is called a *subgradient* of f at  $\boldsymbol{\xi}^0 \in \mathbb{R}^N$  if

<span id="page-3-1"></span>
$$
f(\boldsymbol{\xi}) \ge f(\boldsymbol{\xi}^0) + \langle \mathbf{v}, \boldsymbol{\xi} - \boldsymbol{\xi}^0 \rangle \quad \text{for all } \boldsymbol{\xi} \in \mathbb{R}^N. \tag{2}
$$

We say that f is *subdifferentiable* at  $\xi^0$  if there exists a subgradient of f at  $\xi^0$ . For  $\xi^0 \in \text{dom } f$ ,

$$
f'(\xi^0; \xi) := \lim_{\varepsilon \downarrow 0} \frac{f((1-\varepsilon)\xi^0 + \varepsilon\xi) - f(\xi^0)}{\varepsilon} \quad \text{for all } \xi \in \mathbb{R}^N
$$

always exists in  $[-\infty, \infty)$  and  $\mathbf{v} \in \mathbb{R}^N$  is a subgradient of f at  $\xi^0$  if and only if

<span id="page-3-0"></span>
$$
f'(\boldsymbol{\xi}^0; \boldsymbol{\xi}) \ge \langle \mathbf{v}, \boldsymbol{\xi} \rangle \quad \text{for all } \boldsymbol{\xi} \in \mathbb{R}^N
$$
 (3)

holds [\[Roc70,](#page-25-7) Theorem 23.2].

Assume  $\triangle^N \subseteq$  dom f. Then, f is subdifferentiable at  $\mathbf{q}^0 \in \triangle^N_+$  [\[Roc70,](#page-25-7) Theorem 23.4]. Moreover, if there is no subgradient of  $f$  at  $\mathbf{q}^0 \in \triangle^N \backslash \triangle^N_+$ , then

$$
f'(\mathbf{q}^0; \mathbf{q}) = -\infty \quad \text{for all } \mathbf{q} \in \triangle^N_+
$$

holds [\[Roc70,](#page-25-7) Theorem 23.3]. From the observation, we define  $\partial f(\mathbf{q}^0)$  for  $\mathbf{q}^0 \in \Delta^N$  as the set consisting of all subgradients of f at  $q^0$  if f is subdifferentiable at  $q^0$ , and otherwise

$$
\partial f(\mathbf{q}^0) := \left\{ \mathbf{v} \in [-\infty, \infty)^N \middle| \begin{array}{c} \mathbf{v}_n \in \mathbb{R} \text{ for } n \in \text{supp}(\mathbf{q}^0) \text{ and } \mathbf{v}_n = -\infty \text{ for } n \notin \text{supp}(\mathbf{q}^0) \\ \text{such that } f(\mathbf{q}) \geqslant f(\mathbf{q}^0) + \langle \mathbf{v}, \mathbf{q} - \mathbf{q}^0 \rangle \text{ holds for all } \mathbf{q} \in \triangle^N \end{array} \right\}.
$$

Note that the notion of  $\partial f(\mathbf{q}^0)$  differs from the usual one. We adopt this definition to accommodate regular losses [\(Definition 4\)](#page-7-1) later. We call a map  $\partial f : \triangle^N \to 2^{[-\infty,\infty)^N}$  the subdifferential of f. Let us denote an arbitrary selector of  $\partial f(\mathbf{q})$  by  $\nabla f$ , that is, a map assigning to each point  $\mathbf{q} \in \Delta^N$  an element in  $\partial f(\mathbf{q})$ . Since  $\partial f(\mathbf{q}^0)$  consists of the gradient of f at  $q^0$  if f is differentiable at  $q^0$ , it is consistent to use the notation  $\nabla f$  for a selector. For  $\mathbf{q}, \mathbf{q}^0 \in \triangle^N$ , the associated *Bregman divergence* of  $\mathbf{q}$  given  $\mathbf{q}^0$  is defined by

$$
B_{(f,\nabla f)}(\mathbf{q}\|\mathbf{q}^0) := f(\mathbf{q}) - f(\mathbf{q}^0) - \left\langle \nabla f(\mathbf{q}^0), \mathbf{q} - \mathbf{q}^0 \right\rangle \in [0,\infty].
$$

We verify that the inequality [\(3\)](#page-3-0) holds for  $\mathbf{q}^0 \in \triangle^N$  and  $\mathbf{v} \in \partial f(\mathbf{q}^0)$ .

**Lemma 1** Let  $f : \mathbb{R}^N \to (-\infty, \infty]$  be a proper convex function such that  $\triangle^N \subseteq \text{dom } f$ . For  $\mathbf{q}^0 \in \triangle^N$ , the set  $\partial f(\mathbf{q}^0)$  is nonempty and  $\mathbf{v} \in \partial f(\mathbf{q}^0)$  satisfies  $f'(\mathbf{q}^0; \mathbf{q}) \geq \langle \mathbf{v}, \mathbf{q} \rangle$ for  $\mathbf{q} \in \triangle^N$ .

**Proof** Fix  $\mathbf{q}^0 \in \Delta^N$ . Since the claim trivially holds true if f is subdifferentiable at  $\mathbf{q}^0$ , we assume that f is not subdifferentiable at  $q^0$ . Since f is subdifferentiable at  $q^0 \in \triangle^N_{\frac{1}{\lambda}}$  [\[Roc70,](#page-25-7) Theorem 23.4,  $I := |\text{supp}(\mathbf{q}^0)|$  satisfies  $1 \leqslant I \leqslant N-1$ . For  $\eta \in \mathbb{R}^I$ , define  $\boldsymbol{\xi}^{\eta} \in \mathbb{R}^N$  by

$$
\xi_n^{\boldsymbol{\eta}} \coloneqq \begin{cases} \eta_n & \text{if } n \in \text{supp}(\mathbf{q}^0), \\ 0 & \text{if } n \notin \text{supp}(\mathbf{q}^0), \end{cases}
$$

#

and define a function  $f_I : \mathbb{R}^I \to (-\infty, \infty]^N$  by

$$
f_I(\boldsymbol{\eta}) := f(\boldsymbol{\xi}^{\boldsymbol{\eta}}) \quad \text{for } \boldsymbol{\eta} \in \mathbb{R}^I.
$$

Then,  $f_I$  is a proper convex function on  $\mathbb{R}^I$  such that  $\triangle^I \subseteq \text{dom } f_I$ , consequently,  $f_I$ is subdifferentiable at  $\hat{\eta} \in \Delta^I_+$  [\[Roc70,](#page-25-7) Theorem 23.4]. For  $\mathbf{q} \in \Delta^N$ , define  $\eta^{\mathbf{q}} \in \mathbb{R}^I$ by  $\eta_n^{\mathbf{q}} = q_n$  for  $n \in \text{supp}(\mathbf{q}^0)$ . Then, for  $\mathbf{q} \in \triangle^N$  with  $\text{supp}(\mathbf{q}) \subseteq \text{supp}(\mathbf{q}^0)$ , we have  $\eta^{\mathbf{q}} \in \triangle^I$ and  $f_I(\eta^q) = f(q)$ . Moreover, if  $q \in \triangle^N$  satisfies supp $(q) = \text{supp}(q^0)$ , then  $\eta^q \in \triangle^I_+$  and hence  $\partial f_I(\eta^q) \neq \emptyset$ . Choose  $\mathbf{w} \in \partial f_I(\eta^{q^0})$  and define  $\mathbf{v} \in [-\infty, \infty)^N$  by #

$$
v_n := \begin{cases} w_n & \text{if } n \in \text{supp}(\mathbf{q}^0), \\ -\infty & \text{if } n \notin \text{supp}(\mathbf{q}^0). \end{cases}
$$

From now on, we show that  $\mathbf{v} \in \partial f(\mathbf{q}^0)$ . For  $\mathbf{q} \in \Delta^N$ , if  $q_n > 0$  holds for some  $n \notin \text{supp}(\mathbf{q}^0)$ , From now on, we show that  $\mathbf{v} \in \partial f(\mathbf{q}^0)$ . For  $\mathbf{q} \in \triangle^N$ , if  $q_n > 0$  holds for some  $n \notin \text{supp}(\mathbf{q}^0)$ ,<br>then  $\langle \mathbf{v}, \mathbf{q} - \mathbf{q}^0 \rangle = -\infty$  and [\(2\)](#page-3-1) holds. On the other hand, if  $q_n = 0$  for all  $n \notin \text{supp}(\math$ then  $\mathbf{q} \in \triangle^N$  with  $\text{supp}(\mathbf{q}) \subseteq \text{supp}(\mathbf{q}^0)$  and

$$
f(\mathbf{q}) = f_I(\boldsymbol{\eta}^{\mathbf{q}}) \geqslant f_I\left(\boldsymbol{\eta}^{\mathbf{q}^0}\right) + \left\langle \mathbf{w}, \boldsymbol{\eta}^{\mathbf{q}} - \boldsymbol{\eta}^{\mathbf{q}^0} \right\rangle = f(\mathbf{q}^0) + \left\langle \mathbf{v}, \mathbf{q} - \mathbf{q}^0 \right\rangle,
$$

that is, [\(2\)](#page-3-1) holds for  $\mathbf{q} \in \Delta^N$ . Thus,  $\mathbf{v} \in \partial f(\mathbf{q}^0)$  follows. Moreover, the above inequality immediately yields  $f'(\mathbf{q}^0; \mathbf{q}) \geq \langle \mathbf{v}, \mathbf{q} \rangle$ . This completes the proof of the lemma.  $\Box$ 

### <span id="page-4-0"></span>3 Classification, proper losses, and Savage representation

After introducing the learning problem of multiclass classification, we discuss losses and their properties. We review the notion of proper losses and its connection to Bregman divergences. Although this connection is already known, we formalize it rigorously. In particular, we verify the existence of minimizers of the conditional risk and its measurable selection [\(Lemma 2\)](#page-5-0), which have been implicitly used in previous literature without any proof.

### 3.1 Multiclass classification

We regard a Radon space  $\mathcal X$  as an input space, that is, the set of possible observations, and  $\mathcal{Y} := [N]$  as a set of labels. The set  $\triangle^N$  is identified as the set of all probability measures on *y*. We fix a probability measure  $\nu$  on  $\mathcal{X} \times \mathcal{Y}$  and denote by  $\nu_{\mathcal{X}}$  the marginal of  $\nu$  on  $\mathcal{X}$ , that is,  $\nu(\mathcal{B} \times \mathcal{Y}) = \nu_{\mathcal{X}}(\mathcal{B})$  holds for any measurable set  $\mathcal{B} \subseteq \mathcal{X}$ . Then, by the disintegration

theorem [\[DM78,](#page-23-5) Chapter III-70 and 72], there exists a Borel map  $\mathbf{q}^{\bullet} : \mathcal{X} \to \triangle^{N}$ , uniquely defined  $\nu_{\mathcal{X}}$ -a.e., such that

$$
\nu(\mathcal{B} \times \{y\}) = \int_{\mathcal{B}} q_y^{\mathbf{x}} d\nu_{\mathcal{X}}(\mathbf{x}) \quad \text{for a measurable set } \mathcal{B} \subseteq \mathcal{X} \text{ and } y \in \mathcal{Y}.
$$

We deem  $\mathbf{q}^{\mathbf{x}} \in \Delta^{N}$  a true probability vector at the input **x** induced from  $\nu$ . Multiclass classification is a task to learn a forecaster to predict the most likely label

$$
y^{\mathbf{x}} \in \argmax_{y \in \mathcal{Y}} q_y^{\mathbf{x}} \quad \text{for each } \mathbf{x} \in \mathcal{X}.
$$

#### 3.2 Proper losses

We continue to use the notation in the previous subsection. To elicit  $\mathbf{q}^{\bullet}$ , we use a loss  $\ell$ , which is a Borel map from  $\triangle^N$  to  $\mathbb{R}^N$ . Define the associated *full risk* by

$$
\mathbb{L}[\hat{\mathbf{q}}^{\bullet}] := \int_{\mathcal{X}\times\mathcal{Y}} \ell_y(\hat{\mathbf{q}}^{\mathbf{x}}) \mathrm{d}\nu(\mathbf{x}, y) \quad \text{for a Borel map } \hat{\mathbf{q}}^{\bullet} : \mathcal{X} \to \triangle^N.
$$

A minimizer of  $\mathbb{L}$  among Borel maps  $\hat{\mathbf{q}}^{\bullet} : \mathcal{X} \to \triangle^{N}$  is called an *estimator* of  $\mathbf{q}^{\bullet}$ . The choice of  $\ell$  directly affects the quality of an estimator. It is more intuitive to work on the conditional counterpart of the full risk instead. Let  $\mathbf{q}, \hat{\mathbf{q}} \in \triangle^N$ . For a loss  $\ell$ , the associated *conditional* risk of  $\hat{q}$  given q and *conditional Bayes risk* of q are defined by

$$
L(\mathbf{q},\widehat{\mathbf{q}}) \coloneqq \sum_{y \in \mathcal{Y}} q_y \ell_y(\widehat{\mathbf{q}}) \quad \text{and} \quad \underline{L}(\mathbf{q}) \coloneqq \inf_{\widehat{\mathbf{q}} \in \triangle^N} L(\mathbf{q},\widehat{\mathbf{q}}),
$$

respectively. Here, we regard  $q$  as a *true* probability vector and  $\hat{q}$  as an *estimate*. The full risk is rewritten as

$$
\mathbb{L}[\hat{\mathbf{q}}^{\bullet}] = \int_{\mathcal{X}} L(\mathbf{q}^{\mathbf{x}}, \hat{\mathbf{q}}^{\mathbf{x}}) d\nu_{\mathcal{X}}(\mathbf{x}).
$$

Since the infimum of a family of linear functions is concave, L is concave on  $\triangle^N$ , consequently  $\underline{L} \circ \hat{q}^{\bullet} : \mathcal{X} \to [-\infty, \infty)$  is measurable on  $\mathcal{X}$ , which in turn shows

$$
\mathbb{L}[\hat{\mathbf{q}}^{\bullet}] \geqslant \int_{\mathcal{X}} \underline{L}(\mathbf{q}^{\mathbf{x}}) d\nu_{\mathcal{X}}(\mathbf{x}).
$$

Thus, the minimization problem of the full risk is reduced to that of the conditional risk if the map  $\mathcal{M}: \triangle^N \to 2^{\triangle^{\bar{N}}}$  defined by

<span id="page-5-0"></span>
$$
\mathcal{M}(\mathbf{q}) \coloneqq \argmin_{\widehat{\mathbf{q}} \in \triangle^N} L(\mathbf{q}, \widehat{\mathbf{q}}) \quad \text{for } \mathbf{q} \in \triangle^N.
$$

has a Borel selector. Note that  $\mathcal{M}(\mathbf{q}) = \emptyset$  may happen.

**Lemma 2** Suppose  $\ell : \Delta^N \to \mathbb{R}^N$  is lower semi-continuous. Then, we have that  $\underline{L}(\mathbf{q}) > -\infty$ and  $\mathcal{M}(\mathbf{q})$  is nonempty and closed for  $\mathbf{q} \in \Delta^N$ . Moreover, if  $\ell$  is continuous, then there exists a Borel selector of M.

**Proof** Since the first assertion,  $\mathcal{M}(q)$  is nonempty and closed for  $q \in \Delta^N$ , follows from the standard argument, we omit its proof.

Assume the continuity of  $\ell$  and we show the existence of a Borel selector of  $\mathcal M$ . Note that the continuity of  $\ell$  guarantees the continuity of  $L(\mathbf{q}, \cdot)$  on  $\Delta^N$  for each  $\mathbf{q} \in \Delta^N$ . By the Kuratowski and Ryll-Nardzewski measurable selection theorem [\[KRN65,](#page-24-4) Main Theorem & Corollary 1], it is enough to show that

$$
\mathcal{B}_{\mathcal{K}} \coloneqq \left\{ \mathbf{q} \in \triangle^{N} \bigm| \mathcal{M}(\mathbf{q}) \cap \mathcal{K} \neq \varnothing \right\}
$$

is Borel for any compact set  $\mathcal{K}$  in  $\mathbb{R}^N$  with  $\triangle^N \cap \mathcal{K} \neq \emptyset$ .

Fix a compact set  $\mathcal{K}$  in  $\mathbb{R}^N$  with  $\triangle^N \cap \mathcal{K} \neq \emptyset$ . Since  $\triangle^N \cap \mathcal{K}$  is separable, there exists Fix a compact set K in  $\mathbb{R}^N$  with  $\triangle^N \cap K \neq \emptyset$ . Since  $\triangle^N \cap K$  is separable, then dense countable set  $\{q^j\}_{j\in\mathbb{N}}$  in  $\triangle^N \cap K$ . For each  $j \in \mathbb{N}$ , define  $d_j : \triangle^N \to \mathbb{R}$  by

$$
d_j(\mathbf{q}) \coloneqq L(\mathbf{q}, \mathbf{q}^j) - \inf_{\widehat{\mathbf{q}} \in \triangle^N} L(\mathbf{q}, \widehat{\mathbf{q}}) \quad \text{for } \mathbf{q} \in \triangle^N,
$$

which is lower semi-continuous, in particular, Borel on  $\triangle^N$ . Then, č

$$
\mathcal{B} := \bigcap_{m \in \mathbb{N}} \bigcup_{j \in \mathbb{N}} d_j^{-1}([0, m^{-1}))
$$

is Borel. We will show  $\mathcal{B} = \mathcal{B}_{\mathcal{K}}$ . For  $\mathbf{q} \in \mathcal{B}_{\mathcal{K}}$ , there exists  $\hat{\mathbf{q}} \in \mathcal{M}(\mathbf{q}) \cap \mathcal{K}$ . By the continuity of  $L(\mathbf{q}, \cdot)$  on  $\triangle^N$ , for each  $m \in \mathbb{N}$ , there exists  $\delta_m > 0$  such that if  $\mathbf{q}' \in \triangle^N$ continuity of  $L(\mathbf{q}, \cdot)$  on  $\triangle^N$ , for each  $m \in \mathbb{N}$ , there exists  $\delta_m > 0$  such that if  $\mathbf{q}' \in \triangle^N$ <br>satisfies  $\|\mathbf{q}' - \widehat{\mathbf{q}}\|_2 < \delta_m$ , then  $0 \le L(\mathbf{q}, \mathbf{q}') - L(\mathbf{q}, \widehat{\mathbf{q}}) < m^{-1}$ . By the density of  $\{\mathbf{$ exists  $j_m \in \mathbb{N}$  such that  $\|\mathbf{q}^{j_m} - \hat{\mathbf{q}}\|_2 < \delta_m$  and hence

$$
d_{j_m}(\mathbf{q})=L(\mathbf{q},\mathbf{q}^{j_m})-L(\mathbf{q},\widehat{\mathbf{q}})\in[0,m^{-1}),
$$

which in turn implies  $q \in \mathcal{B}$ . Conversely, for  $q \in \mathcal{B}$  and  $m \in \mathbb{N}$ , there exists  $j_m \in \mathbb{N}$  such that

$$
L(\mathbf{q},\mathbf{q}^{j_m}) < \inf_{\widehat{\mathbf{q}} \in \triangle^N} L(\mathbf{q},\widehat{\mathbf{q}}) + m^{-1}.
$$

We extract a convergent subsequence of  $(q^{j_m})_{m\in\mathbb{N}}$  (not relabeled) with limit  $\hat{\mathbf{q}} \in \Delta^N \cap \mathcal{K}$ . The continuity of  $L(\mathbf{q}, \cdot)$  gives

$$
L(\mathbf{q},\widehat{\mathbf{q}})=\lim_{m\to\infty}L(\mathbf{q},\mathbf{q}^{j_m})\leqslant\inf_{\widehat{\mathbf{q}}'\in\triangle^N}L(\mathbf{q},\widehat{\mathbf{q}}'),
$$

 $\Box$ 

proving  $\hat{\mathbf{q}} \in \mathcal{M}(\mathbf{q})$  hence  $\mathbf{q} \in \mathcal{B}_\mathcal{K}$ . This completes the proof of the lemma.

Although this fact is not directly relevant to our main topic, we complement it in this article because we are unaware of any previous literature formalizing it for losses defined on  $\Delta^N$ . For a different type of (margin-based) losses, the existence of a measurable full risk minimizer has been studied [\[Ste07,](#page-25-8) Theorem 3.2 (ii)].

Since  $\mathcal{M}(\mathbf{q})$  is ideally a singleton of **q**, we consider such a class of losses.

Definition 3 (Proper losses) A loss  $\ell : \triangle^N \to \mathbb{R}^N$  is proper if  $q \in \mathcal{M}(q)$  holds for each  $\mathbf{q} \in \Delta^N$ . We say  $\ell$  is strictly proper if  $\mathcal{M}(\mathbf{q}) = {\mathbf{q}}$  holds for each  $\mathbf{q} \in \Delta^N$ .

For a proper loss  $\ell$ , the conditional risk is minimized at the true probability vector, and the identity map on  $\triangle^N$  becomes a Borel selector of M. In this case, it follows that  $L(\mathbf{q}) = L(\mathbf{q}, \mathbf{q})$ for  $\mathbf{q} \in \triangle^N$ .

### 3.3 Savage representation

The range of a loss can be extended from  $\mathbb{R}^N$  to  $(-\infty, \infty]^N$ . This extension is necessary to accommodate some losses, including the log loss  $\ell_y(\mathbf{q}) := -\ln q_y$  (for each  $y \in \mathcal{Y}$ ) into proper losses. We will see that a proper loss induces a Bregman divergence under the regularity.

Definition 4 (Regular losses [\[GR07,](#page-23-1) Definition 1]) A loss  $\ell : \triangle^N \to (-\infty, \infty]^N$  is said to be regular if  $\ell_y(\mathbf{q}) = \infty$  happens only for  $y \notin \text{supp}(\mathbf{q})$ .

In what follows, we consider a regular loss  $\ell$ , where its range is  $(-\infty, \infty]^N$ .

Although the following property has been well known in literature [\[Sav71,](#page-25-6) §4] [\[GR07,](#page-23-1) Theorem 2] [\[WVR16,](#page-25-9) Proposition 7], we provide its rigorous proof to handle the regularity and subdifferentials carefully.

**Proposition 5 (Savage representation [\[Sav71,](#page-25-6) §4])** Let  $\ell$  be regular. Then,  $\ell$  is proper (resp. strictly proper) if and only if there exists a proper convex (resp. strictly convex) function f on  $\mathbb{R}^N$  such that dom  $f = \Delta^N$  and, for all  $\hat{\mathbf{q}} \in \Delta^N$ , there exists a subgradient  $\hat{\mathbf{v}} \in \partial f(\hat{\mathbf{q}})$ satisfying

<span id="page-7-2"></span><span id="page-7-1"></span><span id="page-7-0"></span>
$$
L(\mathbf{q}, \hat{\mathbf{q}}) = -f(\hat{\mathbf{q}}) - \langle \hat{\mathbf{v}}, \mathbf{q} - \hat{\mathbf{q}} \rangle \quad \text{for } \mathbf{q} \in \triangle^N. \tag{4}
$$

**Proof** First, assume that  $\ell$  is proper. Then,  $\underline{L}(\mathbf{q}) = L(\mathbf{q}, \mathbf{q}) \in \mathbb{R}$ . Define  $f : \mathbb{R}^N \to (-\infty, \infty]$ by #

<span id="page-7-3"></span>
$$
f(\boldsymbol{\xi}) := \begin{cases} -\underline{L}(\boldsymbol{\xi}) & \text{if } \boldsymbol{\xi} \in \triangle^N, \\ \infty & \text{otherwise,} \end{cases}
$$
 (5)

then f is a proper convex function on  $\mathbb{R}^N$  such that dom  $f \subseteq \triangle^N$ . For  $\mathbf{q}, \hat{\mathbf{q}} \in \triangle^N$ , we have

$$
f(\mathbf{q})=-L(\mathbf{q},\mathbf{q})\geq -L(\mathbf{q},\widehat{\mathbf{q}})=-L(\widehat{\mathbf{q}},\widehat{\mathbf{q}})+\langle -\ell(\widehat{\mathbf{q}}),\mathbf{q}-\widehat{\mathbf{q}}\rangle=f(\widehat{\mathbf{q}})+\langle -\ell(\widehat{\mathbf{q}}),\mathbf{q}-\widehat{\mathbf{q}}\rangle.
$$

In particular, if f is subdifferentiable at  $\hat{q}$ , then this inequality also holds by replacing  $q \in \Delta^N$ with  $\boldsymbol{\xi} \in \mathbb{R}^N$ . Thus,  $-\ell(\widehat{\mathbf{q}}) \in \partial f(\widehat{\mathbf{q}})$  and [Eq. \(4\)](#page-7-2) hold for any  $\widehat{\mathbf{q}} \in \triangle^N$ .

Conversely, suppose that there is a proper convex function f on  $\mathbb{R}^N$  such that dom  $f = \triangle^N$ and, for all  $\hat{\mathbf{q}} \in \Delta^N$ , there exists  $\hat{\mathbf{v}} \in \partial f(\hat{\mathbf{q}})$  satisfying [Eq. \(4\).](#page-7-2) Then, for  $\mathbf{q} \in \Delta^N$ , we have  $L(\mathbf{q}, \mathbf{q}) = -f(\mathbf{q})$  and

$$
L(\mathbf{q},\mathbf{q}) = -f(\mathbf{q}) \leqslant -f(\widehat{\mathbf{q}}) - \langle \widehat{\mathbf{v}}, \mathbf{q} - \widehat{\mathbf{q}} \rangle = L(\mathbf{q},\widehat{\mathbf{q}}) \quad \text{for } \widehat{\mathbf{q}} \in \Delta^N,
$$

in turn,  $\ell$  is proper.

Next, we show the equivalence of the strict properness of  $\ell$  and the strict convexity of f on  $\triangle^N$ . Let  $\ell$  be a proper loss, f a convex function on  $\triangle^N$  such that [\(4\)](#page-7-2) holds. On the one hand, if  $\ell$  is strictly proper, then we have

$$
f((1-t)\mathbf{q} + t\mathbf{q}') = L((1-t)\mathbf{q} + t\mathbf{q}', (1-t)\mathbf{q} + t\mathbf{q}')
$$
  
=  $(1-t)L(\mathbf{q}, (1-t)\mathbf{q} + t\mathbf{q}') + tL(\mathbf{q}', (1-t)\mathbf{q} + t\mathbf{q}')$   
>  $(1-t)L(\mathbf{q}, \mathbf{q}) + tL(\mathbf{q}', \mathbf{q}')$   
=  $(1-t)f(\mathbf{q}, \mathbf{q}) + tf(\mathbf{q}', \mathbf{q}'),$ 

for  $\mathbf{q}, \mathbf{q}' \in \Delta^N$  and  $t \in (0, 1)$ , proving the strict convexity of f on  $\Delta^N$ . On the other hand, if  $\ell$  is not strict proper, then there exist distinct  $\mathbf{q}, \mathbf{q}' \in \triangle^N$  such that

$$
f(\mathbf{q}) = -L(\mathbf{q}, \mathbf{q}) = -L(\mathbf{q}, \mathbf{q}') = f(\mathbf{q}') + \langle \mathbf{v}', \mathbf{q} - \mathbf{q}' \rangle,
$$

where  $\mathbf{v}' \in \partial f(\mathbf{q}')$  satisfies [Eq. \(4\)](#page-7-2) such that  $L(\mathbf{q}, \mathbf{q}') = -f(\mathbf{q}') - \langle \mathbf{v}', \mathbf{q} - \mathbf{q}' \rangle$ . For any  $t \in (0, 1)$ , we have

$$
-L((1-t)\mathbf{q} + t\mathbf{q}', (1-t)\mathbf{q} + t\mathbf{q}') = f((1-t)\mathbf{q} + t\mathbf{q}')
$$
  
\n
$$
\leq (1-t)f(\mathbf{q}) + tf(\mathbf{q}')
$$
  
\n
$$
= f(\mathbf{q}') + (1-t)\langle \mathbf{v}', \mathbf{q} - \mathbf{q}' \rangle
$$
  
\n
$$
= f(\mathbf{q}') + \langle \mathbf{v}', (1-t)\mathbf{q} + t\mathbf{q}' - \mathbf{q}' \rangle
$$
  
\n
$$
= -L((1-t)\mathbf{q} + t\mathbf{q}', (1-t)\mathbf{q} + t\mathbf{q}').
$$

This yields  $f((1-t)\mathbf{q} + t\mathbf{q}') = (1-t)f(\mathbf{q}) + f(\mathbf{q}'),$  that is, f is not strictly convex on  $\triangle^N$ . This completes the proof of the proposition.

As a by-product of the proof of [Proposition 5,](#page-7-0) we obtain the following property thanks to the construction [\(5\)](#page-7-3).

Corollary 6 (Subgradient of conditional Bayes risk) For a regular loss  $\ell$ , define a proper convex function f on  $\mathbb{R}^N$  by [\(5\)](#page-7-3). Then, f is a proper convex function on  $\mathbb{R}^N$  such that  $-\ell(\widehat{\mathbf{q}}) \in \partial f(\widehat{\mathbf{q}})$  for  $\widehat{\mathbf{q}} \in \triangle^N$ .

[Equation \(5\)](#page-7-3) gives a closed form of a subgradient of  $-\underline{L}$  and is of interest per se.

For a regular loss  $\ell$ , define the *surrogate regret*  $R : \overline{\triangle}^N \times \overline{\triangle}^N \to (-\infty, \infty]$  by

<span id="page-8-2"></span> $R(\mathbf{q}, \mathbf{\hat{q}}) := L(\mathbf{q}, \mathbf{\hat{q}}) - L(\mathbf{q}) \quad \text{for } \mathbf{q}, \mathbf{\hat{q}} \in \triangle^N,$ 

which measures the suboptimality of an estimate  $\hat{q}$  given a true q. If  $\ell$  is regular and proper, then, by [Proposition 5,](#page-7-0) the Bregman divergence associated with  $f$  defined in [\(5\)](#page-7-3) with the measurable selector  $\nabla f = -\ell$  satisfies

<span id="page-8-1"></span>
$$
R(\mathbf{q}, \hat{\mathbf{q}}) = B_{(f, -\ell)}(\mathbf{q} \| \hat{\mathbf{q}}) \quad \text{for } \mathbf{q}, \hat{\mathbf{q}} \in \triangle^{N}.
$$
 (6)

#### 3.4 Strongly proper losses

For  $\kappa > 0$ , a loss  $\ell$  is called  $\kappa$ -strongly proper if

<span id="page-8-0"></span>
$$
R(\mathbf{q},\hat{\mathbf{q}}) = L(\mathbf{q},\hat{\mathbf{q}}) - \underline{L}(\mathbf{q}) \geqslant \frac{\kappa}{2} \|\mathbf{q} - \hat{\mathbf{q}}\|_2^2 \quad \text{for } \mathbf{q}, \hat{\mathbf{q}} \in \triangle^N. \tag{7}
$$

Strongly proper losses have been introduced for  $N = 2$  [\[Aga14\]](#page-21-0) and for general  $N \ge 3$  [\[ZLA21\]](#page-25-4) to derive a surrogate regret bound in the form of [\(1\)](#page-1-0). For example, the log loss  $\ell_y(\mathbf{q}) = -\ln q_y$ is 1-strongly proper [\[ZLA21,](#page-25-4) Lemma 3]. Interestingly, for  $N = 2$ ,  $\ell$  is regular and strongly proper if and only if its conditional Bayes risk  $-L$  is strongly convex [\[Aga14,](#page-21-0) Theorem 10]. As an immediate consequence of Eq.  $(7)$ , we have the 1/2-order surrogate regret bounds for strongly proper losses:

<span id="page-8-3"></span>
$$
\|\mathbf{q} - \hat{\mathbf{q}}\|_2 \leqslant \sqrt{\frac{2}{\kappa} R(\mathbf{q}, \hat{\mathbf{q}})}.
$$
\n(8)

Though several binary losses are shown to be strongly proper [\[Aga14,](#page-21-0) Table 1], it remains challenging to determine whether a given multiclass loss is strongly proper. In the next section, we derive surrogate regret bounds for general multiclass proper losses.

### <span id="page-9-0"></span>4 Regret bounds: Necessity of strict properness

In this section, we first study the moduli of convexity in [§4.1.](#page-9-1) Therein, the equivalence of the strict convexity of a function and the strict monotonicity of its modulus [\(Theorem 9\)](#page-10-0) is the first main result, which ensures that its surrogate regret bound is non-vacuous. Then, [§4.2](#page-11-1) shows surrogate regret bounds for general multiclass proper losses beyond strongly proper losses. Finally, [§4.3](#page-12-0) relates surrogate regret bounds to several downstream tasks.

### <span id="page-9-1"></span>4.1 Moduli of convexity

Before introducing the moduli of convexity, we study the *midpoint Jensen gap* of a convex function  $f: \triangle^N \to \mathbb{R}$ , which is defined by

$$
J(\mathbf{q}, \check{\mathbf{q}}) \coloneqq \frac{f(\mathbf{q}) + f(\check{\mathbf{q}})}{2} - f\left(\frac{\mathbf{q} + \check{\mathbf{q}}}{2}\right) \quad \text{for } \mathbf{q}, \check{\mathbf{q}} \in \triangle^N.
$$

The midpoint Jensen gap is nonnegative by the convexity of f on  $\triangle^N$ . The midpoint Jensen gap is invariant under adding an affine function, and so is the modulus of convexity. That is, the midpoint Jensen gaps of two convex functions  $f : \triangle^N \to \mathbb{R}$  and  $f_{\lambda,\mathbf{u}} : \triangle^N \to \mathbb{R}$  defined by

$$
f_{\lambda,\mathbf{u}}(\mathbf{q}) \coloneqq f(\mathbf{q}) + \langle \mathbf{u}, \mathbf{q} \rangle + \lambda \quad \text{for } \mathbf{q} \in \triangle^N
$$

are the same, for any  $\mathbf{u} \in \mathbb{R}^N$  and  $\lambda \in \mathbb{R}$ . Moreover, we will show that for continuous convex functions  $f, g : \triangle^N \to \mathbb{R}$ , their midpoint Jensen gaps are the same if and only if  $f - g$  is affine. This property is reminiscent of the condition for the universal equivalence of surrogate losses [\[NWJ09,](#page-24-5) Theorem 3] [\[DKR18,](#page-23-6) Theorem 1].

**Proposition 7 (Uniqueness up to affine functions)** Let  $f, g : \triangle^N \to \mathbb{R}$  be continuous convex functions. Then, their midpoint Jensen gaps are the same if and only if  $f - g$  is affine.

**Proof** We only show that  $f - g$  is affine under the assumption that the midpoint Jensen gaps of  $f$  and  $g$  are the same since the converse implication is trivial. Hereafter, let us write the midpoint Jensen gaps of f and g by  $J_f$  and  $J_q$ , respectively.

Without loss of generality, we pick  $\mathbf{q}^0 \in \Delta^N_+$  such that f and g are differentiable at  $\mathbf{q}^0$ because a convex function is differentiable almost everywhere in the interior of its domain. Define

$$
\mathbf{v}^0 := \nabla f(\mathbf{q}^0) - \nabla g(\mathbf{q}^0) \text{ and } \lambda := f(\mathbf{q}^0) - g(\mathbf{q}^0).
$$

Fix any  $\mathbf{q} \in \triangle^N$  and set

$$
h(t) := f(\mathbf{q}^0 + t(\mathbf{q} - \mathbf{q}^0)) - g(\mathbf{q}^0 + t(\mathbf{q} - \mathbf{q}^0)) - \langle \mathbf{v}^0, t(\mathbf{q} - \mathbf{q}^0) \rangle - \lambda \quad \text{for } t \in [0, 1].
$$

Then,  $h: [0, 1] \to \mathbb{R}$  is continuous with  $h(0) = 0$  and  $h'(0) = 0$ . With elementary algebra, we have ˆ

$$
0 = J_f(\mathbf{q}^0, \mathbf{q}^0 + t(\mathbf{q} - \mathbf{q}^0)) - J_g(\mathbf{q}^0, \mathbf{q}^0 + t(\mathbf{q} - \mathbf{q}^0)) = \frac{1}{2}h(t) - h\left(\frac{t}{2}\right) \quad \text{for all } t \in [0, 1],
$$

which implies

$$
h\left(\frac{1}{2}t\right) = \frac{1}{2}h(t) \quad \text{for all } t \in [0, 1].
$$

By invoking this relation recursively, we have  $h(t) = 2<sup>k</sup> h(2<sup>-k</sup> t)$  for any  $k \in \mathbb{N}$ , which yields

$$
h(1) = \lim_{k \to \infty} \frac{h(2^{-k}) - h(0)}{2^{-k}} = h'(0) = 0.
$$

Consequently, we have

$$
f(\mathbf{q}) = g(\mathbf{q}) + \langle \mathbf{v}^0, \mathbf{q} - \mathbf{q}^0 \rangle + \lambda.
$$

Thus, we have shown that  $f - g$  is affine.

We extend the moduli of convexity defined on  $(\Delta^2, \|\cdot\|_1)$  [\[Bao23,](#page-22-0) Definition 4] to  $(\Delta^N, \|\cdot\|_p)$ for multiclass classification. Note that the diameter of  $(\triangle^N, \|\cdot\|_p)$  is  $2^{1/p}$ .

<span id="page-10-1"></span> $\Box$ 

**Definition 8 (Modulus of convexity)** For a convex function  $f : \triangle^N \to \mathbb{R}$ , its modulus of convexity of f with respect to the p-norm is the function  $\omega : [0, 2^{1/p}] \to [0, \infty)$  defined by

$$
\omega(r) \coloneqq \inf \left\{ J(\mathbf{q}, \check{\mathbf{q}}) \mid \mathbf{q}, \check{\mathbf{q}} \in \triangle^N \ \text{with} \ \|\mathbf{q} - \check{\mathbf{q}}\|_p \geq r \right\} \quad \text{for } r \in [0, 2^{1/p}].
$$

The modulus of convexity quantifies the convexity of a function. We will show that the convexity and strict convexity of a function are translated to the monotonicity and strict monotonicity of its modulus, respectively. This is an important result throughout this article because the moduli of convexity characterize surrogate regret bounds, as we will see in [§4.2](#page-11-1) soon.

<span id="page-10-0"></span>**Theorem 9 (Monotonicity of modulus)** For a convex function  $f : \triangle^N \to \mathbb{R}$ , the modulus  $\omega$  is non-decreasing on  $[0, 2^{1/p}]$  and  $\omega(0) = 0$ . Moreover, the strict convexity of f on  $\triangle^N$ is equivalent to the strict monotonicity of  $\omega$  on  $[0, 2^{1/p}]$ .

Before proving [Theorem 9,](#page-10-0) we show a lemma used repeatedly.

**Lemma 10** Let  $f: \triangle^N \to \mathbb{R}$  be a convex function. For  $r \in [0, 2^{1/p}]$ , there exist  $\mathbf{q}^r, \check{\mathbf{q}}^r \in \triangle^N$ such that  $\omega(r) = J(\mathbf{q}^r, \check{\mathbf{q}}^r)$  and  $\|\mathbf{q}^r - \check{\mathbf{q}}^r\|_p = r$ .

**Proof** Let  $r \in [0, 2^{1/p}]$ . Define

<span id="page-10-2"></span>
$$
\mathcal{D}^N(r) \coloneqq \left\{ (\mathbf{q},\breve{\mathbf{q}}) \in \triangle^N \times \triangle^N \; \big| \; \|\mathbf{q}-\widehat{\mathbf{q}}\|_p \geqslant r \right\}.
$$

Since  $\mathcal{D}^N(r)$  is compact and J is continuous on  $\mathcal{D}^N(r)$ , there is  $(\mathbf{q}, \check{\mathbf{q}}) \in \mathcal{D}^N(r)$  such that  $\omega(r) = J(\mathbf{q}, \check{\mathbf{q}})$ . Define  $c : [0, 1] \to \triangle^N$  by

$$
c(t) := (1 - t)\mathbf{q} + t\check{\mathbf{q}} \quad \text{for } t \in [0, 1].
$$

In the case of  $\|\mathbf{q} - \check{\mathbf{q}}\|_p = r$ , we can take  $(\mathbf{q}^r, \check{\mathbf{q}}^r) = (\mathbf{q}, \check{\mathbf{q}})$ , and the assertion follows. Assume  $\|\mathbf{q} - \check{\mathbf{q}}\|_p > r$ . Then, there exists  $\tau \in (0, 1/2]$  such that

$$
||c(\tau) - c(1 - \tau)||_p = (1 - 2\tau)r = r.
$$

Since  $f \circ c : [0, 1] \to \mathbb{R}$  is convex, we have

$$
\frac{f(c(\tau)) - f(c(0))}{\tau} \leq \frac{f(c(1)) - f(c(1 - \tau))}{\tau},\tag{9}
$$

which is equivalent to

$$
J(c(\tau), c(1-\tau)) = \frac{f(c(\tau)) + f(c(1-\tau))}{2} - f(c(1/2))
$$
  
\$\leq \frac{f(c(0)) + f(c(1))}{2} - f(c(1/2)) = J(\mathbf{q}, \mathbf{\check{q}}).

This yields  $J(c(\tau), c(1-\tau)) = \omega(r)$ , and hence, we can take  $(\mathbf{q}^r, \check{\mathbf{q}}^r) = (c(\tau), c(1-\tau))$ . Thus, we have confirmed the statement.  $\Box$ 

Proof of [Theorem 9](#page-10-0) Define

$$
\mathcal{D}^N(r) \coloneqq \left\{ (\mathbf{q}, \widehat{\mathbf{q}}) \in \triangle^N \times \triangle^N \mid \|\mathbf{q} - \widehat{\mathbf{q}}\|_p \geqslant r \right\}.
$$

For  $r', r \in [0, 2^{1/p}]$  with  $r' \leq r$ , we observe from the monotonicity  $\mathcal{D}^{N}(r) \subseteq \mathcal{D}^{N}(r')$ that  $\omega(r') \leq \omega(r)$ . It is easily seen that  $J(\mathbf{q}, \mathbf{q}) = 0$  holds for any  $\mathbf{q} \in \Delta^N$  hence  $\omega(0) = 0$ . Thus, the first assertion follows.

Assume that the strict convexity of f on  $\triangle^N$ . Let  $r \in (0, 2^{1/p}]$ . By [Lemma 10,](#page-10-2) there exist  $\mathbf{q}, \check{\mathbf{q}} \in \triangle^N$  such that  $\omega(r) = J(\mathbf{q}, \check{\mathbf{q}})$  and  $\|\mathbf{q} - \check{\mathbf{q}}\|_p = r$ . Define  $c : [0, 1] \to \triangle^N$  by

 $c(t) := (1 - t)\mathbf{q} + t\breve{\mathbf{q}}$  for  $t \in [0, 1]$ .

Since  $f \circ c$  is strictly convex on [0, 1], we have the strict inequality

$$
\frac{f(c(\tau)) - f(c(0))}{\tau} < \frac{f(c(1)) - f(c(1 - \tau))}{\tau} \quad \text{for } \tau \in (0, 1/2].
$$

Consequently, we conclude

$$
\omega((1-2\tau)r) \le J(c(\tau), c(1-\tau)) < J(c(0), c(1)) = \omega(r) \text{ for } \tau \in (0, 1/2],
$$

that is, the strict monotonicity of  $\omega$  on  $[0, 2^{1/p}]$ .

Conversely, if f is not strictly convex on  $\Delta^N$ , there exist distinct  $\mathbf{q}, \check{\mathbf{q}} \in \Delta^N$  such that

$$
f((1-t)\mathbf{q} + t\breve{\mathbf{q}}) = (1-t)f(\mathbf{q}) + tf(\breve{\mathbf{q}}) \text{ for } t \in [0,1].
$$

This leads to  $J(\mathbf{q}, \check{\mathbf{q}}) = 0$ . Consequently,  $\omega$  is not strictly increasing on  $[0, ||\mathbf{q} - \check{\mathbf{q}}||_p]$ .

Thus, the proof of the theorem is achieved.

Despite the simple proof, this will lead to the necessity and sufficiency for a surrogate regret bound being non-vacuous in [§4.2,](#page-11-1) together with [Theorem 11.](#page-11-0)

#### <span id="page-11-1"></span>4.2 Surrogate regret bounds

Now, we give surrogate regret bounds with respect to the  $p$ -norm. The order of a surrogate regret bound for a proper loss  $\ell$  is essentially governed by the modulus of convexity of (the negative of) its conditional Bayes risk  $-L$ . This is an extension of surrogate regret bounds for binary classification [\[Bao23,](#page-22-0) Theorem 6] to multiclass classification.

**Theorem 11 (Surrogate regret bounds)** Let  $\ell : \triangle^N \to (-\infty, \infty]^N$  be a regular proper loss and  $f: \mathbb{R}^N \to (-\infty, \infty]$  a proper convex function defined by [\(5\)](#page-7-3). For  $\mathbf{q}, \hat{\mathbf{q}} \in \triangle^N$ , it holds

<span id="page-11-2"></span>
$$
\omega(\|\mathbf{q} - \hat{\mathbf{q}}\|_{p}) \leq \frac{1}{2}R(\mathbf{q}, \hat{\mathbf{q}})
$$
\n(10)

<span id="page-11-0"></span> $\Box$ 

with equality if  $q = \hat{q}$ .

**Proof** By the definition of  $\omega$  together with [\(6\)](#page-8-1), it is sufficient to show

$$
J(\mathbf{q},\widehat{\mathbf{q}}) \leqslant \frac{1}{2}B_{(f,-\ell)}(\mathbf{q}\|\widehat{\mathbf{q}}) \quad \text{for } \mathbf{q}, \widehat{\mathbf{q}} \in \triangle^N.
$$

By [Corollary 6,](#page-8-2) we have ˆ ˙

$$
f\left(\frac{\mathbf{q}+\hat{\mathbf{q}}}{2}\right) \geqslant f(\hat{\mathbf{q}})+\left\langle -\boldsymbol{\ell}(\hat{\mathbf{q}}),\frac{\mathbf{q}+\hat{\mathbf{q}}}{2}-\hat{\mathbf{q}}\right\rangle = f(\hat{\mathbf{q}})+\frac{1}{2}\left\langle -\boldsymbol{\ell}(\hat{\mathbf{q}}),\mathbf{q}-\hat{\mathbf{q}}\right\rangle,
$$

which implies

$$
J(\mathbf{q},\widehat{\mathbf{q}})\leqslant \frac{1}{2}\left[f(\mathbf{q})-f(\widehat{\mathbf{q}})-\langle -\boldsymbol{\ell}(\widehat{\mathbf{q}}),\mathbf{q}-\widehat{\mathbf{q}}\rangle\right]=\frac{1}{2}B_{(f,-\boldsymbol{\ell})}(\mathbf{q}\|\widehat{\mathbf{q}}).
$$

The equality can be seen immediately by choosing  $\hat{\mathbf{q}} = \mathbf{q}$ .

Let us discuss when a proper loss entails a non-vacuous bound. If  $\ell$  is strictly proper, then  $\omega$  is strictly increasing, which in turn has an inverse function  $\omega^{-1}$  and leads [\(10\)](#page-11-2) to

<span id="page-12-1"></span>
$$
\|\mathbf{q} - \hat{\mathbf{q}}\|_{p} \leqslant \begin{cases} \omega^{-1}\left(\frac{1}{2}R(\mathbf{q}, \hat{\mathbf{q}})\right) & \text{if } \frac{1}{2}R(\mathbf{q}, \hat{\mathbf{q}}) \leqslant \omega\left(2^{\frac{1}{p}}\right), \\ 2^{\frac{1}{p}} & \text{otherwise.} \end{cases}
$$
(11)

 $\Box$ 

The strict monotonicity of  $\omega^{-1}$  is essential because otherwise we cannot always expect that the estimate  $\hat{\mathbf{q}}$  approaches  $\mathbf{q}$  even if the suboptimality  $R(\mathbf{q}, \hat{\mathbf{q}})$  is minimized. By [Proposition 5](#page-7-0) and [Theorem 9,](#page-10-0) the *strict* properness of  $\ell$  is necessary and sufficient for the surrogate regret bound [\(10\)](#page-11-2) being non-vacuous. This is why strict properness matters.

Let us leave a remark on the existing surrogate regret bounds for proper *composite* losses [\[ML21,](#page-24-6) Corollary 3]. They derived a surrogate regret bound similar to [\(11\)](#page-12-1), with the moduli of continuity of the conditional risk  $L(q, \cdot)$ . While the relationship between the moduli of convexity of  $-L$  and the moduli of continuity of  $L(q, \cdot)$  has not been clear,  $-L$  suffices because a surrogate regret is solely determined by  $L$  due to  $(6)$  and [Corollary 6.](#page-8-2) Moreover, the existing surrogate regret bounds [\[ML21,](#page-24-6) Corollary 3] has been limited to the binary case  $N = 2$ . Our [Theorem 11](#page-11-0) is more general therein.

#### <span id="page-12-0"></span>4.3 Relating surrogate regret to downstream tasks

The upper bound for the  $p$ -norm [\(11\)](#page-12-1) is useful for many scenarios to assess the predictive performance of plug-in forecasters, i.e., post-processed forecasters based on the estimator  $\hat{\mathbf{q}}^{\bullet}$ . Thus, we can regard the p-norm bound as a *versatile* surrogate regret bound across different downstream tasks. Subsequently, we provide several examples of downstream tasks to support this idea.

Task 1: multiclass classification. Let us consider multiclass classification based on the post-process approach. Given true and estimated probability vectors  $\mathbf{q}, \hat{\mathbf{q}} \in \triangle^N$ , respectively, the plug-in forecaster based on the estimate  $\hat{\mathbf{q}}$  is given by  $\hat{y} \in \arg \max_{y \in \mathcal{Y}} \hat{q}_y$ , where the tie is broken arbitrarily. Here, the forecaster's suboptimality in multiclass classification is measured by the (conditional)  $0-1$  regret

$$
\operatorname{Reg}_{01}(\mathbf{q},\widehat{\mathbf{q}}) \coloneqq \sum_{n\in\mathcal{Y}} q_n(1-\delta_{n\widehat{y}}) - \min_{y\in\mathcal{Y}} \sum_{n\in\mathcal{Y}} q_n(1-\delta_{ny}) = \max_{y\in\mathcal{Y}} \left\langle \mathbf{q}, \mathbf{L}_{\widehat{y}} - \mathbf{L}_y \right\rangle,
$$

where  $\mathbf{L} \in \mathbb{R}^{N \times N}$  is the 0-1 loss matrix with each component being  $L_{ij} := 1 - \delta_{ij}$ , and  $\mathbf{L}_{y}$ denotes the y-th column vector of **L**. Let  $p^*$  denote the Hölder conjugate of p. The 0-1 regret can be bounded as

$$
\mathrm{Reg}_{01}(\mathbf{q},\widehat{\mathbf{q}})\leqslant\max_{y\in\mathcal{Y}}\left\langle \mathbf{q}-\widehat{\mathbf{q}},\mathbf{L}_{\widehat{y}}-\mathbf{L}_y\right\rangle \leqslant\|\mathbf{q}-\widehat{\mathbf{q}}\|_{p}\max_{y\in\mathcal{Y}}\|\mathbf{L}_{\widehat{y}}-\mathbf{L}_y\|_{p^*}\leqslant 2^{1-\frac{1}{p}}\|\mathbf{q}-\widehat{\mathbf{q}}\|_{p},
$$

where the first inequality holds because  $\langle \hat{\mathbf{q}}, \mathbf{L}_{\hat{y}} - \mathbf{L}_y \rangle \le 0$  for any  $y \in \mathcal{Y}$  attributed to the construction of  $\hat{y}$ , and the second inequality owes to Hölder's inequality. Eventually, the 0-1 regret is controlled by the surrogate regret  $R(\mathbf{q}, \hat{\mathbf{q}})$  via [\(11\)](#page-12-1) if  $\ell$  is strictly proper, which relates the estimation quality of  $\hat{q}$  to the predictive performance of the post-processed forecaster via the p-norm.

Task 2: learning with noisy labels. Let us consider multiclass classification with class-conditional label noises: a true label y is observed as  $\tilde{y}$  with probability  $C_{y,\tilde{y}}$  with a row-stochastic noise matrix  $C \in [0, 1]^{N \times N}$ . In this scenario, our access is limited to the noisy target probability vector  $\tilde{\mathbf{q}} = \mathbf{C}^\top \mathbf{q}$ , through which a noisy estimate  $\hat{\mathbf{q}}$  is obtained. By following the noise-correction strategy [\[ZLA21\]](#page-25-4), the plug-in forecaster based on the noisy estimate  $\hat{\mathbf{q}}$  is given by  $\check{y} \in \arg \max_{y \in \mathcal{Y}} \check{q}_y$ , where  $\check{\mathbf{q}} := (\mathbf{C}^\top)^{-1} \hat{\mathbf{q}}$  (provided that  $\mathbf{C}$  is invertible). Under this setup, the 0-1 regret of  $\check{q}$  given q is bounded as follows:

$$
\begin{aligned} \mathrm{Reg}_{01}(\mathbf{q},\widecheck{\mathbf{q}}) &= \max_{y\in\mathcal{Y}} \left\langle \mathbf{q}, \mathbf{L}_{\widecheck{y}} - \mathbf{L}_y \right\rangle \\ &\leqslant \max_{y\in\mathcal{Y}} \left\langle \mathbf{q} - \widecheck{\mathbf{q}}, \mathbf{L}_{\widecheck{y}} - \mathbf{L}_y \right\rangle = \max_{y\in\mathcal{Y}} \left\langle \widehat{\mathbf{q}} - \widehat{\mathbf{q}}, \mathbf{C}^{-1} (\mathbf{L}_{\widecheck{y}} - \mathbf{L}_y) \right\rangle \\ &\leqslant \|\widetilde{\mathbf{q}} - \widehat{\mathbf{q}}\|_p \max_{y\in\mathcal{Y}} \|\mathbf{C}^{-1} (\mathbf{L}_{\widecheck{y}} - \mathbf{L}_y)\|_{p^*} \end{aligned}
$$

where the first inequality holds because  $\langle \check{\mathbf{q}}, \mathbf{L}_{\check{y}} - \mathbf{L}_y \rangle \leq 0$  for any  $y \in \mathcal{Y}$  attributed to the construction of  $\check{y}$ , and the second inequality owes to Hölder's inequality. The p-norm  $\|\tilde{\mathbf{q}}-\hat{\mathbf{q}}\|_p$ can be minimized even with access to the noisy observations only, and the  $p$ -norm bound  $(11)$ controls this by the surrogate regret of a strictly proper loss. This is also an extension of the previous surrogate regret transfer bounds [\[ZLA21,](#page-25-4) Theorem 4] beyond strongly proper losses.

**Task 3: bipartite ranking.** Consider  $N = 2$  and identify  $\mathbf{q} = [q_1 \ q_2]^\top \in \Delta^2$  with the instance  $q_1 \in [0, 1]$ . Given two instances  $q, q' \in [0, 1]$ , we are interested in giving estimates  $\hat{q}, \hat{q}' \in [0, 1]$  that yield a consistent ranking with  $(q, q')$ . In bipartite ranking, we use the estimates  $(\hat{q}, \hat{q}')$  directly without any post process. The (conditional) ranking regret [\[CLV08\]](#page-23-7) is measured by

$$
\mathrm{Reg}_{\mathrm{rank}}(q,q',\hat{q},\hat{q}') := |q-q'| \left[ 1\!\!1_{\{(\hat{q}-\hat{q}')(q-q')<0\}} + \frac{1}{2} 1\!\!1_{\{\hat{q}=\hat{q}'\}} \right],
$$

where  $\mathbb{1}_{\{A\}} = 1$  when the predicate A holds and 0 otherwise, and the first and second terms penalize an inconsistent ranking and tie, respectively. This can be immediately related to the 1-norm [\[Aga14\]](#page-21-0):

$$
\text{Reg}_{\text{rank}}(q,q',\hat{q},\hat{q}')\leqslant |q-\hat{q}|+|q'-\hat{q}'|,
$$

where the bound [\(11\)](#page-12-1) can be further applied. Thus, the ranking regret is controlled by the surrogate regret.

**Other benefits.** In addition to the above examples, one can easily relate the *p*-norm and downstream tasks such as binary classification with generalized performance criteria [\[KD16,](#page-23-3) Eq.  $(9)$ , which we omit here. Another benefit of the p-norm bound  $(11)$  is that it relates a possibly non-metric  $R(\mathbf{q}, \hat{\mathbf{q}})$  to the metric  $\|\mathbf{q} - \hat{\mathbf{q}}\|_p$ .

To conclude this section, we raise attention to the kinship between moduli of convexity and the known devices such as calibration functions [\[BJM06\]](#page-23-8) [\[Ste07\]](#page-25-8) [\[OBLJ17\]](#page-24-7) [\[BSS20\]](#page-23-9) [\[BSX](#page-23-10)<sup>+</sup>22], comparison inequalities [\[MPRS12\]](#page-24-8) [\[CRR20\]](#page-23-11), and Fisher consistency bounds [\[AMMZ22a\]](#page-22-2) [\[AMMZ22b\]](#page-22-3) [\[MMZ23\]](#page-24-9). In spite of the relevance, moduli of convexity are different in that these devices have been tailored for a specific target loss of each downstream task, whereas moduli are concerned with the p-norm.

### <span id="page-14-0"></span>5 Lower bounds of surrogate regret order

We move on to the next main result: the surrogate regret order cannot go beyond the square root. To this end, we first review the Simonenko order function and the strong convexity used to establish the main result, and then show the main result. In this section, let  $f: \triangle^N \to \mathbb{R}$ be a convex function unless otherwise stated.

### 5.1 Power evaluation of moduli

To interpret the surrogate regret bound [\(10\)](#page-11-2), we evaluate the order of the modulus  $\omega$  by power functions. To this end, we introduce the order of  $\omega$ , which is well-defined since  $\omega(r) > 0$ for  $r \in (0, 2^{1/p}]$  from [Theorem 9.](#page-10-0)

Definition 12 (Simonenko order function [\[Sim64\]](#page-25-10)) Let  $f : \triangle^N \to \mathbb{R}$  be a strictly convex function. The Simonenko order function  $\sigma : (0, 2^{1/p}] \to [0, \infty]$  is defined by

$$
\sigma(r) \coloneqq \frac{rD^-\omega(r)}{\omega(r)} \quad \textit{for $r\in \left(0,2^{\frac{1}{p}}\right]$, \quad \textit{where} \quad D^-\omega(r) \coloneqq \limsup_{\varepsilon\downarrow 0} \frac{\omega(r)-\omega(r-\varepsilon)}{\varepsilon}.
$$

The quantity  $D^-\omega$  is called the *upper left Dini derivative* of  $\omega$  at r. If  $\omega$  is differentiable at r, then  $D^{-}\omega(r) = \omega'(r)$  holds. The Simonenko order function  $\sigma$  evaluates the order of  $\omega$ .

**Proposition 13 (Power evaluations of moduli)** Let  $f : \triangle^N \to \mathbb{R}$  be a strictly convex function. For a fixed  $r_0 \in (0, 2^{1/p}]$ , we define  $s, S \in [0, \infty]$  by

<span id="page-14-1"></span>
$$
s := \inf_{r \in (0,r_0]} \sigma(r), \quad S := \sup_{r \in (0,r_0]} \sigma(r),
$$

and assume  $S < \infty$ . Then, the function  $r \mapsto \omega(r)r^{-s}$  is non-decreasing on  $(0, r_0)$  and the function  $r \mapsto \omega(r) r^{-S}$  is non-increasing on  $(0, r_0)$ . Consequently, the following inequalities hold for any  $r \in [0, r_0]$ : " ȷ ȷ

$$
\left[\frac{\omega(r_0)}{r_0^S}\right]r^S\leqslant\omega(r)\leqslant\left[\frac{\omega(r_0)}{r_0^s}\right]r^s.
$$

Here, we show the proof by assuming the differentiability of  $\omega$  because the resulting proof is instructive. The complete proof without the differentiability is slightly convoluted and deferred to [Appendix A.](#page-25-11)

**Proof** By the definition of S, for any  $t \in (0, r_0]$ ,

$$
\frac{S}{t} = \frac{1}{t} \sup_{r \in (0,r_0]} \sigma(r) \geqslant \frac{\omega'(t)}{\omega(t)}
$$

.

By integrating both sides, for any  $r' \in [r, r_0]$ ,

$$
S \ln \frac{r'}{r} = S \int_r^{r'} \frac{dt}{t} \geqslant \int_r^{r'} \frac{\omega'(t)}{\omega(t)} dt = \ln \frac{\omega(r')}{\omega(r)},
$$

which gives the one side of the desiderata. The other side can be proven similarly.  $\Box$ 

Thus, the p-norm upper bound [\(11\)](#page-12-1) is controlled by the rate  $\omega^{-1}(\rho) = O(\rho^{1/S})$ . Since we are interested in the behavior of  $\|\mathbf{q} - \hat{\mathbf{q}}\|_p$  when  $\hat{\mathbf{q}}$  is close to the minimizer of  $R(\mathbf{q}, \cdot)$ , we focus on the asymptotic behavior of the Simonenko order function as  $r \downarrow 0$ .

#### 5.2 Strong convexity and its relation to moduli

For the asymptotic analysis of  $\sigma$ , we leverage *strong* convexity. Herein, we define the strong convexity parameter for a convex function  $f : \triangle^N \to \mathbb{R}$  and  $t \in (0, 1)$  by

$$
\begin{aligned} \kappa_p^{f,t} &:= \inf \left\{ \frac{2\left[(1-t)f(\mathbf{q}) + tf(\check{\mathbf{q}}) - f((1-t)\mathbf{q} + t\check{\mathbf{q}})\right]}{t(1-t)\|\mathbf{q} - \check{\mathbf{q}}\|_p^2} \; \middle| \; \text{distinct } \mathbf{q}, \check{\mathbf{q}} \in \Delta^N \right\}, \\ \kappa_p^f &:= \inf_{t \in (0,1)} \kappa_p^{f,t}. \end{aligned}
$$

We observe from the convexity of f that  $\kappa_p^f \in [0, \infty)$  and

$$
f((1-t)\mathbf{q} + t\breve{\mathbf{q}}) \le (1-t)f(\mathbf{q}) + tf(\breve{\mathbf{q}}) - \frac{\kappa_p^f}{2}t(1-t)\|\mathbf{q} - \breve{\mathbf{q}}\|_p^2 \quad \text{for } \mathbf{q}, \breve{\mathbf{q}} \in \triangle^N \text{ and } t \in (0,1).
$$

Remark that the strong convexity parameter depends on the underlying set where we take the infimum. Let us define the strong convexity parameter on  $\mathbb{R}^N$  by replacing  $\triangle^N$ with  $\mathbb{R}^N$  and write  $\bar{\kappa}_p^{f,t}$  instead of  $\kappa_p^{f,t}$  for each  $t \in (0,1)$ . By  $\triangle^N \subseteq \mathbb{R}^N$ , we observe that

$$
\bar{\kappa}_p^{f,t} \leqslant \kappa_p^{f,t} \ \text{ for a function } f: \mathbb{R}^N \to (-\infty, \infty],
$$

where the equality does not necessarily hold.

To calculate  $\kappa_p^f$ , we only need to know  $\kappa_p^{f,1/2}$ .

Proposition 14 (Strong convexity parameter at midpoint) For a continuous convex function  $f: \triangle^N \to \mathbb{R}$ , we have  $\kappa_p^f = \kappa_p^{f,1/2}$ .

The proof is deferred to [Appendix B.](#page-27-0) By [Proposition 14,](#page-15-0) the strong convexity parameter and the modulus of convexity are connected.

Since we have

$$
\kappa_p^{f,\frac{1}{2}} = \inf \left\{ \frac{8J(\mathbf{q}, \check{\mathbf{q}})}{\|\mathbf{q} - \check{\mathbf{q}}\|_p^2} \middle| \text{ distinct } \mathbf{q}, \check{\mathbf{q}} \in \triangle^N \right\},\
$$

<span id="page-15-0"></span>\*

the following bound holds:

$$
\inf\left\{\frac{\|\mathbf{q}\|_p^2}{\|\mathbf{q}\|_2^2}\;\middle|\;\mathbf{q}\in\triangle^N\right\}\leqslant\frac{\kappa_2^{f,\frac{1}{2}}}{\kappa_p^{f,\frac{1}{2}}}\leqslant\sup\left\{\frac{\|\mathbf{q}\|_p^2}{\|\mathbf{q}\|_2^2}\;\middle|\;\mathbf{q}\in\triangle^N\right\}.
$$

In particular, if  $N = 2$ , we have

$$
\|\mathbf{q} - \check{\mathbf{q}}\|_{p} = ((q_1 - \check{q}_1)^p + (q_2 - \check{q}_2)^p)^{\frac{1}{p}} = ((q_1 - \check{q}_1)^p + (q_1 - \check{q}_1)^p)^{\frac{1}{p}} = 2^{\frac{1}{p}}|q_1 - \check{q}_1|
$$

for  $\mathbf{q}, \check{\mathbf{q}} \in \Delta^2$ , and hence,  $2\kappa_2^{f,1/2} = 2^{2/p} \kappa_p^{f,1/2}$ . Therefore,  $\kappa_p^f$  remains the same up to constant regardless of the choice of  $p \geq 1$ .

We will use another representation of  $\kappa_p^{f,1/2} = \kappa_p^f$  later:

<span id="page-16-0"></span>
$$
\kappa_p^f = \inf \left\{ \frac{8\omega(r)}{r^2} \middle| r \in (0, 2^{\frac{1}{p}}] \right\}.
$$
 (12)

### 5.3 Asymptotic lower bound

The asymptotic behavior of the Simonenko order  $\sigma(r)$  as  $r \downarrow 0$  is controlled by the strong convexity parameter  $\kappa_p^f$ . We introduce a "local" version of the strong convexity parameter.

Definition 15 (Local strong convexity modulus) For a convex function  $f: \triangle^N \to \mathbb{R}$ , define  $K_p^f : (0, 2^{1/p}] \to \mathbb{R}$  by

<span id="page-16-1"></span>
$$
K_p^f(r) := \frac{8\omega(r)}{r^2}.
$$

This quantity is defined based on the alternative expression of the strong convexity parameter  $\kappa_p^f$  in [\(12\)](#page-16-0). From the relationship (12),  $K_p^f(r) \geqslant \kappa_p^f$  always holds on  $r \in (0, 2^{1/p}]$ . In the following lemma, we verify the continuity properties that  $K_p^f$  naturally entails.

**Lemma 16** If  $f: \triangle^N \to \mathbb{R}$  is continuous convex, then  $K_p^f: (0, 2^{1/p}] \to \mathbb{R}$  is lower semicontinuous and left-continuous.

**Proof** Fix  $r \in (0, 2^{1/p}]$ . Let  $(r_j)_{j \in \mathbb{N}} \subseteq (0, 2^{1/p}]$  be a sequence converging to r. For each  $j \in \mathbb{N}$ , there exist  $\mathbf{q}^j, \check{\mathbf{q}}^j \in \triangle^N$  satisfying

$$
\|\mathbf{q}^j - \breve{\mathbf{q}}^j\|_p = r_j \quad \text{and} \quad \omega(r_j) = J(\mathbf{q}^j, \breve{\mathbf{q}}^j)
$$

from [Lemma 10.](#page-10-2) Define  $c_j : [0, 1] \to \triangle^N$  by

$$
c_j(t) := (1-t)\mathbf{q}^j + t\breve{\mathbf{q}}^j \quad \text{for } t \in [0,1].
$$

By the Arzelá–Ascoli theorem, we can extract a subsequence  $(c_{j_m})_{m\in\mathbb{N}}$  converging uniformly to some  $c : [0, 1] \to \triangle^N$  uniformly, where

 $c(t) = (1 - t)c(0) + tc(1)$  for  $t \in [0, 1]$  and  $||c(0) - c(1)||_p = r$ 

hold. Since  $f$  is continuous, we have

$$
K_p^f(r) = \frac{8}{r^2} \omega(r) \le \frac{8}{r^2} J(c(0), c(1))
$$
  
= 
$$
\lim_{m \to \infty} \frac{8}{r_{j_m}^2} J(c_{j_m}(0), c_{j_m}(1)) = \lim_{m \to \infty} \frac{8}{r_{j_m}^2} \omega(r_{j_m}) = \lim_{m \to \infty} K_p^f(r_{j_m}).
$$

Thus,  $K_p^f : (0, 2^{1/p}] \to \mathbb{R}$  is lower semi-continuous.

Next, we choose distinct  $\mathbf{q}, \check{\mathbf{q}} \in \triangle^N$  satisfying

$$
\|\mathbf{q} - \breve{\mathbf{q}}\|_p = r
$$
 and  $\omega(r) = J(\mathbf{q}, \breve{\mathbf{q}}),$ 

which exist thanks to [Lemma 10.](#page-10-2) For these  $q, \check{q}$ , define

$$
c(t) := (1 - t)\mathbf{q} + t\check{\mathbf{q}} \quad \text{for } t \in [0, 1].
$$

For  $\tau \in (0, 1/2)$ , we have

<span id="page-17-3"></span>
$$
\frac{K_p^f((1-2\tau)r)}{8}[(1-2\tau)r]^2 = \omega((1-2\tau)r)
$$
  
\n
$$
\leq J(c(\tau), c(1-\tau))
$$
  
\n
$$
\leq J(c(0), c(1)) - \frac{\kappa_p^f}{2}\tau(1-\tau)r^2
$$
  
\n
$$
= \omega(r) - \frac{\kappa_p^f}{2}\tau(1-\tau)r^2
$$
  
\n
$$
\leq \frac{K_p^f(r)}{8}r^2 - \frac{\kappa_p^f}{2}\tau(1-\tau)r^2.
$$
\n(13)

 $\Box$ 

Dividing by  $[(1 - 2\tau)r]^2/8 \neq 0$  and then taking the limit yields

<span id="page-17-0"></span>
$$
\limsup_{\tau \downarrow 0} K_p^f((1 - 2\tau)r) \leqslant K_p^f(r).
$$

Together with the lower semi-continuity  $K_p^f$ , the left-continuity  $K_p^f$  is ensured.

This completes the proof of the lemma.

Now, we analyze the asymptotic behavior of  $\sigma(r)$  at  $r \downarrow 0$  when f is continuous on  $\triangle^N$ , which is our second main result. We need a slightly stronger continuity of  $K_p^f$  than what [Lemma 16](#page-16-1) provides. Moreover, we assume the continuity of  $f$  to prevent  $f$  from being discontinuous on the  $\triangle^N \backslash \triangle^N_+$ .

**Theorem 17 (Lower bound of order)** Let  $f : \triangle^N \to \mathbb{R}$  be a continuous strictly convex function. Assume one of the following two conditions.

- <span id="page-17-1"></span>(C1)  $\kappa_p^f > 0$ .
- <span id="page-17-2"></span>(C2)  $K_p^f$  is continuous on  $(0, r_0]$  for some  $r_0 \in (0, 2^{1/p}]$  and  $K_p^f$  converges as  $r \downarrow 0$ .

Then,

<span id="page-18-1"></span>
$$
\limsup_{r \downarrow 0} \sigma(r) \geqslant 2. \tag{14}
$$

Moreover, if we assume both conditions, then

<span id="page-18-2"></span><span id="page-18-0"></span>
$$
\liminf_{r \downarrow 0} \sigma(r) \geqslant 2. \tag{15}
$$

To prove [Theorem 17,](#page-17-0) we leverage the following lemma to locally control  $\sigma(r)$ , which is proven in [Appendix B.](#page-27-0)

**Lemma 18** Let  $f : \triangle^N \to \mathbb{R}$  be a continuous convex function. Then, for any  $\mathbf{q}, \check{\mathbf{q}} \in \triangle^N$ and  $r \in (0, 2^{1/p}],$ 

$$
\liminf_{r \downarrow 0} K_p^f(r) = \kappa_p^f, \quad J(\mathbf{q}, \check{\mathbf{q}}) \ge \frac{\kappa_p^f}{8} \|\mathbf{q} - \check{\mathbf{q}}\|_p^2, \quad \text{and} \quad D^- \omega(r) \ge \frac{\kappa_p^f}{4}r.
$$

**Proof of [Theorem 17](#page-17-0)** We observe from the strict convexity of f and [Theorem 9](#page-10-0) that  $K_p^f > 0$ on  $r \in (0, 2^{1/p}]$ . By assuming [\(C1\)](#page-17-1) only, it follows from [Lemma 18](#page-18-0) that

$$
\limsup_{r\downarrow 0} \sigma(r) = \limsup_{r\downarrow 0} \frac{rD^{-}\omega(r)}{\omega(r)} \ge \limsup_{r\downarrow 0} \frac{r \cdot \frac{\kappa_p^f}{4}r}{\frac{K_p^f(r)}{8}r^2} = \limsup_{r\downarrow 0} \frac{2\kappa_p^f}{K_p^f(r)} = 2.
$$

In addition, assume [\(C2\)](#page-17-2) together. Then, the above inequality provides

$$
\liminf_{r \downarrow 0} \sigma(r) \geqslant \liminf_{r \downarrow 0} \frac{2\kappa_p^f}{K_p^f(r)} = \lim_{r \downarrow 0} \frac{2\kappa_p^f}{K_p^f(r)} = 2.
$$

Next, assume [\(C2\)](#page-17-2) only, and  $\kappa_p^f > 0$  does not hold. In this case, [Lemma 18](#page-18-0) indicates that  $K_p^f(r) \downarrow 0$  as  $r \downarrow 0$ , from which we can inductively define  $(r_j)_{j\in\mathbb{N}} \subseteq (0, r_0]$  by

$$
r_j := \inf \left\{ r \in (0, 2^{1/p}] \; \middle| \; K_p^f(r) \geq \frac{1}{2} K_p^f(r_{j-1}) \right\}.
$$

Then,  $(r_j)_{j\in\mathbb{N}}$  converges to 0 because  $K_p^f > 0$  always holds on  $r \in (0, 2^{1/p}]$  and hence  $K_p^f(r) = 0$ if and only if  $r = 0$ . We observe that

$$
K_p^f(r) < \frac{1}{2} K_p^f(r_{j-1}) = K_p^f(r_j) \quad \text{for } r \in (0, r_j].
$$

This yields

$$
D^{-}\omega(r_j) = \limsup_{\varepsilon \downarrow 0} \frac{\omega(r_j) - \omega(r_j - \varepsilon)}{\varepsilon} = \limsup_{\varepsilon \downarrow 0} \frac{\frac{K_p^f(r_j)}{8}r_j^2 - \frac{K_p^f(r_j - \varepsilon)}{8}(r_j - \varepsilon)^2}{\varepsilon}
$$

$$
\geq \limsup_{\varepsilon \downarrow 0} \frac{\frac{K_p^f(r_j)}{8}r_j^2 - \frac{K_p^f(r_j)}{8}(r_j - \varepsilon)^2}{\varepsilon} = \frac{K_p^f(r_j)}{4}r_j,
$$

which implies

$$
\limsup_{r \downarrow 0} \sigma(r) \ge \limsup_{j \to \infty} \sigma(r_j) = \limsup_{j \to \infty} \frac{r_j D^{-} \omega(r_j)}{\omega(r_j)} \ge \limsup_{j \to \infty} \frac{r_j \cdot \frac{K_p^f(r_j)}{4} r_j}{\frac{K_p^f(r_j)}{8} r_j^2} = 2.
$$

This completes the proof of [Theorem 17.](#page-17-0)

It follows from [Proposition 13](#page-14-1) and [Theorem 17](#page-17-0) that the p-norm bound [\(11\)](#page-12-1) is controlled by the rate of  $\omega^{-1}(\rho)$  cannot be faster than  $O(\rho^{1/2})$  for a very large class of strictly proper  $\ell$ . To see this, we invoke [Proposition 13](#page-14-1) to observe that for some  $r_0 \in (0, 2^{1/p}]$ ,

 $\Box$ 

$$
\omega^{-1}(\rho) \leqslant [r_0 \omega(r_0)^{-\frac{1}{S}}] \cdot \rho^{\frac{1}{S}} \quad \text{for } r \in [0, r_0] \text{ such that } \rho = \omega(r).
$$

If the bound [\(14\)](#page-18-1) holds, then we have

$$
2 \leq \limsup_{r \downarrow 0} \sigma(r) \leq \sup_{r \in (0,r_0]} \sigma(r) = S,
$$

which implies  $\rho^{1/S} \ge \rho^{1/2}$  (for  $\rho < 1$ ). Thus, we discern the optimal rate  $\omega^{-1}(\rho) = O(\rho^{1/2})$ for  $\rho \in [0, 1]$ . This assures that strongly proper losses asymptotically achieve the optimal rate  $O(\rho^{1/2})$  as seen in [\(8\)](#page-8-3). Indeed, the condition [\(C1\)](#page-17-1) is tightly connected to the strong properness.

Lower bound of  $\omega^{-1}(\rho)$ . [Proposition 13](#page-14-1) also implies that for some  $r_0 \in (0, 2^{1/p}]$ ,

$$
\omega^{-1}(\rho) \geqslant [r_0 \omega(r_0)^{-\frac{1}{s}}] \cdot \rho^{\frac{1}{s}} \quad \text{for } r \in [0, r_0] \text{ such that } \rho = \omega(r).
$$

If the bound  $(15)$  holds with the *strict* inequality, then we can choose  $r_0$  to satisfy

$$
\frac{2+\varsigma}{2} \leq \inf_{r \in (0,r_0]} \sigma(r) = s \quad \text{for } \varsigma := \liminf_{r \downarrow 0} \sigma(r) > 2,
$$

which implies  $\rho^{1/s} \ge \rho^{2/(2+s)} > \rho^{1/2}$  (for  $\rho < 1$ ). Thus,  $\omega^{-1}$  admits the lower bound  $\Omega(\rho^{1/2})$ for  $\rho \in [0, 1]$ .

**Comparison with the known lower bound.** A relevant lower bound  $\omega^{-1}(\rho) = \Omega(\rho^{1/2})$ has been shown previously for a slightly different type of losses [\[FW21,](#page-23-12) Theorem 4]. To derive the lower bound, they assume that a loss is strongly convex and has a locally Lipschitz gradient [\[FW21,](#page-23-12) Assumption 1]. The latter condition is assumed under the loss differentiability, whereas both our [\(C1\)](#page-17-1) and [\(C2\)](#page-17-2) do not need the differentiability of  $\ell$ . Ergo, the differentiability assumption is lifted to show the optimality of  $\omega^{-1}(\rho) = O(\rho^{1/2})$ . In [§6,](#page-20-0) we will see a non-differentiable example, the max-power function. To show  $\omega^{-1}(\rho) = \Omega(\rho^{1/2})$ in our case, [\(C1\)](#page-17-1) and [\(C2\)](#page-17-2) coupled with the existence of the limit of  $K_p^f(r)$  as  $r \downarrow 0$  suffice.

**Remark 1** Our analysis with the Simonenko order evaluates the order of  $\omega$  by a power function in the form of  $r^S \leq \omega(r) \leq r^s$  for  $r \in [0, r_0]$ . This is an evaluation for a finite

<span id="page-20-1"></span>

**Figure 1:** Illustration of  $\omega(r) = r \sin\left(\frac{1}{r}\right) - C\mathrm{i}\left(\frac{1}{r}\right) + r$ .

range, which is more than an asymptotic evaluation. Despite its subtlety, it often matters, as seen in the following example:

$$
\omega(r) = r \sin\left(\frac{1}{r}\right) - Ci\left(\frac{1}{r}\right) + r \quad \text{for } r > 0, \quad \text{where} \quad \text{Ci}(z) := -\int_z^{\infty} t^{-1} \cos(t) \, \text{d}t.
$$

This  $\omega$  is monotonically increasing and satisfies  $\omega(r) = O(r)$  in the asymptotic evaluation. However, when it comes to the finite evaluation, we cannot go faster than  $\omega(r) \geq r^2$  just because this  $\omega$  satisfies [\(C2\)](#page-17-2) and [Theorem 17](#page-17-0) implies  $\limsup_{r\to 0} \sigma(r) \geq 2$ . Thus, the finite evaluation gives a better characterization when we assess the convergence rate of finitely large surrogate regret. See [Fig. 1](#page-20-1) to better understand the above example.

### <span id="page-20-0"></span>6 Examples

We overview a couple of proper losses. Since there is the one-to-one correspondence between a regular proper loss  $\ell$  and a convex function  $f = -\underline{L}$  on  $\Delta^N$  (see [Eq. \(5\)\)](#page-7-3), we show examples in terms of the corresponding convex functions. To facilitate closed-form solutions of  $\omega$ , we restrict ourselves to  $N = 2$  and  $p = 1$ . In this case, we overload  $f(q_1, q_2) = f(q, 1 - q)$ for  $q \in [0, 1]$ .

[Table 1](#page-21-2) lists several convex functions with their moduli, whose derivations are given previously [\[Bao23\]](#page-22-0). All examples satisfy at least [\(C2\)](#page-17-2) of [Theorem 17.](#page-17-0) Indeed, there exists the limit of  $K_p^f(r)$  as  $r \downarrow 0$ , as seen in [Fig. 2.](#page-22-4) Thus, we have ensured that the optimal order of the inverse modulus  $\omega^{-1}(\rho)$  for many common losses is  $O(\rho^{1/2})$ . Note that the Shannon entropy, squared  $\alpha$ -norms (for  $\alpha > 1$ ), and Tsallis entropies (for  $\alpha > 1$ ) satisfy [\(C1\)](#page-17-1), but the  $\alpha$ -norms satisfy [\(C1\)](#page-17-1) only for  $\alpha \in (1, 2]$  but violate for  $\alpha > 2$ .

	$f(\mathbf{q})$	Range of $\alpha$	Modulus $\omega(r)$	Loss $\ell$
Shannon ent.	$\langle \mathbf{q}, \ln \mathbf{q} \rangle$		$\frac{1+r}{2} \ln \frac{1+r}{2} + \frac{1-r}{2} \ln \frac{1-r}{2} + \ln 2$	Log
Sq. $\alpha$ -norms	$\ \mathbf{q}\ _{\alpha}^2$	$1 < \alpha < 2$	$\frac{1}{4}$ $\left\  \begin{bmatrix} 1+r \\ 1-r \end{bmatrix} \right\ _{\alpha}^{2} - 4^{1/\alpha - 1}$	<b>Brier</b>
		$2 \leqslant \alpha$	$\frac{1}{2}$ $\left\  \begin{bmatrix} r \\ 1-r \end{bmatrix} \right\ _{\alpha}^{2} - \frac{1}{4} \left\  \begin{bmatrix} r \\ 2-r \end{bmatrix} \right\ _{\alpha}^{2} + \frac{1}{2}$	$(\alpha = 2)$
$\alpha$ -norms	$\ {\bf q}\ _\alpha$	$1 < \alpha < 2$	(No closed-form in general)	Pseudo- spherical
		$2 \leqslant \alpha$	$\frac{1}{2}$ $\left\  \begin{bmatrix} 1 \\ 1-r \end{bmatrix} \right\ _{\alpha} - \frac{1}{2} \left\  \begin{bmatrix} 2-r \\ 2-r \end{bmatrix} \right\ _{\alpha} + \frac{1}{2}$	
Tsallis ent.	$\ \mathbf{q}\ _{\alpha}^{\alpha}$	$\alpha \in (1,2) \cup (3,\infty)$	$\frac{1}{2\alpha} \left\  \begin{bmatrix} 1+r \\ 1-r \end{bmatrix} \right\ _{\alpha}^{\alpha} - 2^{1-\alpha}$	$\alpha$ -log
		$2 \leq \alpha \leq 3$	$\frac{1}{2} \left\  \begin{bmatrix} 1 \\ 1-r \end{bmatrix} \right\ _{\alpha}^{\alpha} - \frac{1}{2^{\alpha}} \left\  \begin{bmatrix} r \\ 2-r \end{bmatrix} \right\ _{\alpha}^{\alpha} + \frac{1}{2}$	
Max-power	$\max_{n\in[N]} (q_n - \frac{1}{N})^{\alpha}$	$1 < \alpha < 2$	$\frac{1}{2} r-\frac{1}{2} ^{\alpha}-\frac{1}{2^{\alpha}} r-1 ^{\alpha}+\frac{1}{2^{1+\alpha}}$	
		$2 \leq \alpha$	$(\frac{r}{2})^{\alpha}$	

<span id="page-21-2"></span>**Table 1:** Examples of a convex function f. For  $\omega$ , we show the expressions with  $(N, p) = (2, 1)$ .

In these examples, f is differentiable entirely on  $\triangle^N_+$  except for the max-power function. For  $N = 2$ , the max-power function is  $f(q) :=$ ely on $\left| q-\frac{1}{2}\right|$ 2  $\triangle^{N}_{+}$  except for the max-power function.<br> $\Big|^{\alpha}$ , which is not differentiable at  $q = 1/2$ . [Theorem 17](#page-17-0) is applicable even for such non-differentiable functions.

**Example for**  $N \ge 3$ . Though deriving a closed form of  $\omega$  for general  $N > 2$  is challenging, we can delineate  $\omega$  for the Shannon entropy with  $p = 2$ : for  $r \in (0, 2^{1/2})$ , define  $\mathbf{q}, \check{\mathbf{q}} \in \triangle^N$  by " ȷ " ȷ

$$
\mathbf{q} = \left[ \frac{1 + 2^{-1/2}r}{2} \quad \frac{1 - 2^{-1/2}r}{2} \quad 0 \quad \dots \quad 0 \right] \quad \text{and} \quad \mathbf{\check{q}} = \left[ \frac{1 - 2^{-1/2}r}{2} \quad \frac{1 + 2^{-1/2}r}{2} \quad 0 \quad \dots \quad 0 \right].
$$

Then,  $\|\mathbf{q} - \check{\mathbf{q}}\|_2 = r$  and  $\omega(r) = J(\mathbf{q}, \check{\mathbf{q}})$ . At this minimizer,  $\omega$  can be written as

$$
\omega(r) = \frac{1 + 2^{-1/2}r}{2} \ln \frac{1 + 2^{-1/2}r}{2} + \frac{1 - 2^{-1/2}r}{2} \ln \frac{1 - 2^{-1/2}r}{2} + \ln 2.
$$

This  $\omega$  (for  $p = 2$ ) is akin to the form of  $\omega$  shown in [Table 1,](#page-21-2) which is for  $(N, p) = (2, 1)$ , with a slight difference in the scale. Its derivation is based on the method of Lagrange multipliers and deferred to [Proposition 24.](#page-33-0)

### Acknowledgments

HB and AT are supported by JSPS Grant-in-Aid for Transformative Research Areas(A) (22A201). AT is supported by JSPS Grant-in-Aid for Scientific Research(C) (19K03494).

# References

- <span id="page-21-1"></span>[AA15] Arpit Agarwal and Shivani Agarwal. On consistent surrogate risk minimization and property elicitation. In Proceedings of the 28th Conference on Learning Theory, pages 4–22, 2015.
- <span id="page-21-0"></span>[Aga14] Shivani Agarwal. Surrogate regret bounds for bipartite ranking via strongly proper losses. Journal of Machine Learning Research, 15(1):1653–1674, 2014.

<span id="page-22-4"></span>

**Figure 2:** Numerical plots of  $K_p^f(r) = 8\omega(r)/r^2$  for each f in [Table 1.](#page-21-2)

- <span id="page-22-2"></span>[AMMZ22a] Pranjal Awasthi, Anqi Mao, Mehryar Mohri, and Yutao Zhong. H-consistency bounds for surrogate loss minimizers. In Proceedings of the 39th International Conference on Machine Learning, pages 1117–1174, 2022.
- <span id="page-22-3"></span>[AMMZ22b] Pranjal Awasthi, Anqi Mao, Mehryar Mohri, and Yutao Zhong. Multi-class H-consistency bounds. Advances in Neural Information Processing Systems, 35:782–795, 2022.
- <span id="page-22-1"></span>[Bao22] Han Bao. Excess Risk Transfer and Learning Problem Reduction towards Reliable Machine Learning. PhD thesis, University of Tokyo, 2022.
- <span id="page-22-0"></span>[Bao23] Han Bao. Proper losses, moduli of convexity, and surrogate regret bounds. In Proceedings of the 36th Conference on Learning Theory, pages 525–547, 2023.

<span id="page-23-2"></span>

- <span id="page-23-8"></span>[BJM06] Peter L Bartlett, Michael I Jordan, and Jon D McAuliffe. Convexity, classification, and risk bounds. Journal of the American Statistical Association, 101(473):138–156, 2006.
- <span id="page-23-0"></span>[BSS05] Andreas Buja, Werner Stuetzle, and Yi Shen. Loss functions for binary class probability estimation and classification: Structure and applications. Technical Report, 2005.
- <span id="page-23-9"></span>[BSS20] Han Bao, Clay Scott, and Masashi Sugiyama. Calibrated surrogate losses for adversarially robust classification. In Proceedings of the 30th Conference on Learning Theory, pages 408–451, 2020.
- <span id="page-23-10"></span>[BSX`22] Han Bao, Takuya Shimada, Liyuan Xu, Issei Sato, and Masashi Sugiyama. Pairwise supervision can provably elicit a decision boundary. In Proceedings of the 25th International Conference on Artificial Intelligence and Statistics, pages 2618–2640, 2022.
- <span id="page-23-7"></span>[CLV08] Stéphan Clémençon, Gábor Lugosi, and Nicolas Vayatis. Ranking and empirical minimization of U-statistics. The Annals of Statistics, 36(2):844–874, 2008.
- <span id="page-23-11"></span>[CRR20] Carlo Ciliberto, Lorenzo Rosasco, and Alessandro Rudi. A general framework for consistent structured prediction with implicit loss embeddings. Journal of Machine Learning Research, 21(1):3852–3918, 2020.
- <span id="page-23-6"></span>[DKR18] John Duchi, Khashayar Khosravi, and Feng Ruan. Multiclass classification, information, divergence and surrogate risk. The Annals of Statistics, 46(6B):3246– 3275, 2018.
- <span id="page-23-5"></span>[DM78] Claude Dellacherie and Paul-André Meyer. Probabilities and Potential, volume 29 of North-Holland Mathematics Studies. North-Holland Publishing Co., Amsterdam-New York, 1978.
- <span id="page-23-4"></span>[Fig76] Tadeusz Figiel. On the moduli of convexity and smoothness. Studia Mathematica, 56(2):121–155, 1976.
- <span id="page-23-12"></span>[FW21] Rafael Frongillo and Bo Waggoner. Surrogate regret bounds for polyhedral losses. Advances in Neural Information Processing Systems, 34:21569–21580, 2021.
- <span id="page-23-1"></span>[GR07] Tilmann Gneiting and Adrian E Raftery. Strictly proper scoring rules, prediction, and estimation. Journal of the American statistical Association, 102(477):359– 378, 2007.
- <span id="page-23-3"></span>[KD16] Wojciech Kotłowski and Krzysztof Dembczyński. Surrogate regret bounds for generalized classification performance metrics. In Proceedings of the 8th Asian Conference on Machine Learning, pages 301–316, 2016.
- <span id="page-24-11"></span>[KK96] R. Kannan and Carole King Krueger. Advanced Analysis on the Real Line. Universitext. Springer-Verlag, New York, 1996.
- <span id="page-24-2"></span>[KNRD14] Oluwasanmi O Koyejo, Nagarajan Natarajan, Pradeep K Ravikumar, and Inderjit S Dhillon. Consistent binary classification with generalized performance metrics. Advances in Neural Information Processing Systems, 27:2744–2752, 2014.
- <span id="page-24-4"></span>[KRN65] K. Kuratowski and C. Ryll-Nardzewski. A general theorem on selectors. Bulletin de l'Académie Polonaise des Sciences. Série des Sciences Mathématiques, Astronomiques et Physiques, 13:397–403, 1965.
- <span id="page-24-3"></span>[KSFF17] Meelis Kull, Telmo Silva Filho, and Peter Flach. Beta calibration: a wellfounded and easily implemented improvement on logistic calibration for binary classifiers. In Proceedings of the 20th International Conference on Artificial Intelligence and Statistics, pages 623–631, 2017.
- <span id="page-24-6"></span>[ML21] Alexander Mey and Marco Loog. Consistency and finite sample behavior of binary class probability estimation. In Proceedings of the AAAI Conference on Artificial Intelligence, volume 35, pages 8967–8974, 2021.
- <span id="page-24-9"></span>[MMZ23] Anqi Mao, Mehryar Mohri, and Yutao Zhong. Structured prediction with stronger consistency guarantees. Advances in Neural Information Processing Systems, 36:46903–46937, 2023.
- <span id="page-24-8"></span>[MPRS12] Youssef Mroueh, Tomaso Poggio, Lorenzo Rosasco, and Jean-Jeacques Slotine. Multiclass learning with simplex coding. Advances in Neural Information Processing Systems, 25:2789–2797, 2012.
- <span id="page-24-1"></span>[NA13] Harikrishna Narasimhan and Shivani Agarwal. On the relationship between binary classification, bipartite ranking, and binary class probability estimation. Advances in Neural Information Processing Systems, 26:2913–2921, 2013.
- <span id="page-24-5"></span>[NWJ09] XuanLong Nguyen, Martin J Wainwright, and Michael I Jordan. On surrogate loss functions and f-divergences. The Annals of Statistics, 37(2):876–904, 2009.
- <span id="page-24-7"></span>[OBLJ17] Anton Osokin, Francis Bach, and Simon Lacoste-Julien. On structured prediction theory with calibrated convex surrogate losses. Advances in Neural Information Processing Systems, 31:302–313, 2017.
- <span id="page-24-0"></span>[OFR`19] Yaniv Ovadia, Emily Fertig, Jie Ren, Zachary Nado, David Sculley, Sebastian Nowozin, Joshua Dillon, Balaji Lakshminarayanan, and Jasper Snoek. Can you trust your model's uncertainty? evaluating predictive uncertainty under dataset shift. Advances in Neural Information Processing Systems, 32:13991–14002, 2019.
- <span id="page-24-10"></span>[OT13] Shin-Ichi Ohta and Asuka Takatsu. Displacement convexity of generalized relative entropies. II. Communications in Analysis and Geometry, 21(4):687– 785, 2013.

<span id="page-25-7"></span>

- <span id="page-25-2"></span>[RW09] Mark D Reid and Robert C Williamson. Surrogate regret bounds for proper losses. In Proceedings of the 26th International Conference on Machine Learning, pages 897–904, 2009.
- <span id="page-25-0"></span>[RW10] Mark D Reid and Robert C Williamson. Composite binary losses. Journal of Machine Learning Research, 11:2387–2422, 2010.
- <span id="page-25-5"></span>[RW11] Mark D Reid and Robert C Williamson. Information, divergence and risk for binary experiments. Journal of Machine Learning Research, 12(22):731–817, 2011.
- <span id="page-25-6"></span>[Sav71] Leonard J Savage. Elicitation of personal probabilities and expectations. Journal of the American Statistical Association, 66(336):783–801, 1971.
- <span id="page-25-10"></span>[Sim64] Igor Borisovich Simonenko. Interpolation and extrapolation of linear operators in Orlicz spaces. Matematicheskii Sbornik, 105(4):536–553, 1964.
- <span id="page-25-8"></span>[Ste07] Ingo Steinwart. How to compare different loss functions and their risks. Constructive Approximation, 26(2):225–287, 2007.
- <span id="page-25-9"></span>[WVR16] Robert C Williamson, Elodie Vernet, and Mark D Reid. Composite multiclass losses. Journal of Machine Learning Research, 17:1–52, 2016.
- <span id="page-25-1"></span>[Zha04] Tong Zhang. Statistical behavior and consistency of classification methods based on convex risk minimization. The Annals of Statistics, 32(1):56–85, 2004.
- <span id="page-25-4"></span>[ZLA21] Mingyuan Zhang, Jane Lee, and Shivani Agarwal. Learning from noisy labels with no change to the training process. In Proceedings of the 38th International Conference on Machine Learning, pages 12468–12478. PMLR, 2021.
- <span id="page-25-3"></span>[ZRA20] Mingyuan Zhang, Harish Guruprasad Ramaswamy, and Shivani Agarwal. Convex calibrated surrogates for the multi-label F-measure. In Proceedings of the 37th International Conference on Machine Learning, pages 11246–11255. PMLR, 2020.

# <span id="page-25-11"></span>Appendix A. Proof of power evaluations without differentiability

In [§5,](#page-14-0) we show power evaluations of the moduli with the differentiability of  $\omega$ .

**Proposition 13 (Power evaluations of moduli)** Let  $f : \triangle^N \to \mathbb{R}$  be a strictly convex function. For a fixed  $r_0 \in (0, 2^{1/p}]$ , we define  $s, S \in [0, \infty]$  by

$$
s := \inf_{r \in (0,r_0]} \sigma(r), \quad S := \sup_{r \in (0,r_0]} \sigma(r),
$$

and assume  $S < \infty$ . Then, the function  $r \mapsto \omega(r)r^{-s}$  is non-decreasing on  $(0, r_0)$  and the function  $r \mapsto \omega(r) r^{-S}$  is non-increasing on  $(0, r_0)$ . Consequently, the following inequalities hold for any  $r \in [0, r_0]$ : " ȷ ȷ

$$
\left[\frac{\omega(r_0)}{r_0^S}\right]r^S\leqslant\omega(r)\leqslant\left[\frac{\omega(r_0)}{r_0^s}\right]r^s.
$$

**Proof** We first show that  $r \mapsto \omega(r)r^{-S}$  is non-increasing on  $(0, r_0)$  in a similar way to the existing argument [\[OT13,](#page-24-10) Lemma 2.9]. For  $r \in (0, r_0]$  and  $\delta > 0$ , there exists  $\varepsilon_{r,\delta} > 0$  such that

<span id="page-26-0"></span>
$$
\frac{r}{\omega(r)} \cdot \sup_{\varepsilon \in (0,\varepsilon_{r,\delta})} \frac{\omega(r) - \omega(r - \varepsilon)}{\varepsilon} \le \sigma(r) + \frac{1}{2}\delta \tag{16}
$$

by the definition of  $D^-\omega$ . Define

$$
g(t) := S + \frac{1}{2}\delta + \frac{1}{t}[(1-t)^{S+\delta} - 1] \text{ for } t \in (0,1).
$$

Then,  $q$  is continuous on  $(0, 1)$  and

$$
\lim_{t \downarrow 0} g(t) = S + \frac{1}{2}\delta - (S + \delta) = -\frac{1}{2}\delta < 0,
$$

which implies the existence of  $\tau \in (0, 1)$  such that  $g(t) < 0$  for  $t \in (0, \tau)$ . Then, for any  $u \in (0, \varepsilon_{r,\delta}) \cap (0, r\tau),$ 

$$
-\frac{r^{S+\delta}}{\omega(r)} + \frac{(r-u)^{S+\delta}}{\omega(r-u)} = \frac{ur^{S+\delta-1}}{\omega(r-u)} \left[ \frac{r}{\omega(r)} \frac{\omega(r) - \omega(r-u)}{u} \right] - \frac{r^{S+\delta}}{\omega(r-u)} + \frac{r^{S+\delta}(1-\frac{u}{r})^{S+\delta}}{\omega(r-u)} \le \frac{ur^{S+\delta-1}}{\omega(r-u)} \left[ \sigma(r) + \frac{1}{2}\delta \right] + \frac{ur^{S+\delta-1}}{\omega(r-u)} \cdot \frac{1}{\frac{u}{r}} \left[ \left( 1 - \frac{u}{r} \right)^{S+\delta} - 1 \right] \le \frac{ur^{S+\delta-1}}{\omega(r-u)} \left[ S + \frac{1}{2}\delta + g\left(\frac{u}{r}\right) - \left( S + \frac{1}{2}\delta \right) \right] = \frac{ur^{S+\delta-1}}{\omega(r-u)} \cdot g\left(\frac{u}{r}\right) < 0,
$$

where the inequality [\(16\)](#page-26-0) is used at the second line and the third line follows from the definition of S. Hence, we have

$$
\frac{(r-u)^{S+\delta}}{\omega(r-u)} < \frac{r^{S+\delta}}{\omega(r)} \quad \text{for } r \in (0, r_0) \text{ and } u \in (0, \varepsilon_{r,\delta}) \cap (0, r\tau).
$$

Letting  $\delta \downarrow 0$ , we conclude that  $r \mapsto r^S/\omega(r)$  is non-decreasing on  $(0, r_0)$ . This is equivalent to that  $r \mapsto \omega(r)r^{-S}$  is non-increasing on  $(0, r_0)$ .

Next, we show that  $r \mapsto \omega(r)r^{-s}$  is non-decreasing on  $(0, r_0)$ . For  $r \in (0, r_0]$ , let  $u \in (0, r)$ . Since  $\ln \omega$  is non-decreasing on  $[r - u, r]$ , it follows from the fundamental theorem of calculus [\[KK96,](#page-24-11) Theorem 1.3.1] that

$$
\int_{r-u}^{r} D^{-} \ln \omega(r') dr' \leqslant \ln \frac{\omega(r)}{\omega(r-u)}.
$$

By  $D^-\omega(r') \leqslant S < \infty$ , we have

$$
\lim_{\varepsilon \downarrow 0} \omega(r' - \varepsilon) = \omega(r') \quad \text{for } r' \in (0, r_0),
$$

which yields

$$
\lim_{\varepsilon \downarrow 0} \frac{\ln \omega(r') - \ln \omega(r' - \varepsilon)}{\omega(r') - \omega(r' - \varepsilon)} = \frac{1}{\omega(r')}
$$

and

$$
D^{-} \ln \omega(r') = \limsup_{\varepsilon \downarrow 0} \left[ \frac{\ln \omega(r') - \ln \omega(r' - \varepsilon)}{\omega(r') - \omega(r' - \varepsilon)} \cdot \frac{\omega(r') - \omega(r' - \varepsilon)}{\varepsilon} \right] = \frac{1}{\omega(r')} D^{-} \omega(r') \geq \frac{s}{r'}.
$$

Thus, we have

$$
\int_{r-u}^{r} D^{-} \ln \omega(r') dr' \geqslant \int_{r-u}^{r} \frac{s}{r'} dr' = s \ln \frac{r}{r-u}.
$$

These imply

 $\omega(r-u) \cdot (r-u)^{-s} \leqslant \omega(r) \cdot r^{-s},$ 

that is,  $r \mapsto \omega(r)r^{-s}$  is non-decreasing on  $(0, r_0)$ .

# <span id="page-27-0"></span>Appendix B. Proofs

Proposition 14 (Strong convexity parameter at midpoint) For a continuous convex function  $f: \triangle^N \to \mathbb{R}$ , we have  $\kappa_p^f = \kappa_p^{f,1/2}$ .

**Proof** By definition,  $\kappa_p^f \leq \kappa_p^{f,1/2}$  trivially holds. We shall prove the converse inequality. Observe from the definition that

$$
f\left(\frac{\mathbf{q}+\check{\mathbf{q}}}{2}\right) \leq \frac{1}{2}f(\mathbf{q}) + \frac{1}{2}f(\check{\mathbf{q}}) - \frac{\kappa_p^{f,\frac{1}{2}}}{8}\|\mathbf{q} - \check{\mathbf{q}}\|_p^2 \quad \text{for all } \mathbf{q}, \check{\mathbf{q}} \in \triangle^N.
$$

For  $i, j \in \mathbb{N}$ , set

$$
t_{i,j} \coloneqq 2^{-i}j.
$$

Fix distinct  $\mathbf{q}, \check{\mathbf{q}} \in \triangle^N$  and define  $c : [0, 1] \to \triangle^N$  by

$$
c(t) := (1 - t)\mathbf{q} + t\check{\mathbf{q}} \quad \text{for } t \in [0, 1].
$$

We will show that

<span id="page-27-1"></span>
$$
f(c(t_{i,j})) \le (1 - t_{i,j})f(c(0)) + t_{i,j}f(c(1)) - \frac{\kappa_p^{f, \frac{1}{2}}}{2}t_{i,j}(1 - t_{i,j})\|\mathbf{q} - \check{\mathbf{q}}\|_p^2
$$
(17)

for  $(i, j) \in \mathbb{N} \times \mathbb{Z}_{\geq 0}$  with  $t_{i,j} \in [0, 1]$  (namely, for  $i \in \mathbb{N}$  and  $0 \leq j \leq 2^i$ ) by induction on i. We immediately observe that [\(17\)](#page-27-1) always holds for  $t_{i,0} = 0$  and  $t_{i,2^{i}} = 1$  regardless of i. Note that  $t_{i,j} \in (0, 1)$  leads to  $t_{i,j\pm 1} \in [0, 1]$ .

 $\Box$ 

The inequality [\(17\)](#page-27-1) trivially holds for  $i = 1$ . Assume that (17) holds for some  $i \in \mathbb{N}$ and all j with  $0 \leq j \leq 2^i$ . Then, [\(17\)](#page-27-1) also holds for  $t_{i+1,2j} = t_{i,j}$  with  $0 \leq j \leq 2^i$ . For  $0 \leqslant j \leqslant 2^i - 1$ , we have

$$
t_{i+1,2j+1} = \frac{t_{i+1,2j} + t_{i+1,2j+2}}{2} = \frac{t_{i,j} + t_{i,j+1}}{2}.
$$

Define  $c_{i,j} : [0, 1] \to \triangle^N$  by

$$
c_{i,j}(t) := c((1-t) \cdot t_{i,j} + t \cdot t_{i,j+1}) \text{ for } t \in [0,1].
$$

This implies

$$
f(c(t_{i+1,2j+1}))
$$
  
\n
$$
\leq \frac{1}{2}f(c_{i,j}(0)) + \frac{1}{2}f(c_{i,j}(1)) - \frac{\kappa^{f,\frac{1}{2}}}{8}||c_{i,j}(0) - c_{i,j}(1)||_p^2
$$
  
\n
$$
= \frac{1}{2}f(c_{i,j}(0)) + \frac{1}{2}f(c_{i,j}(1)) - \frac{\kappa_p^{f,\frac{1}{2}}}{8} \cdot 2^{-2i} \cdot ||\mathbf{q} - \breve{\mathbf{q}}||_p^2
$$
  
\n
$$
= \frac{1}{2}f(c(t_{i,j})) + \frac{1}{2}f(c(t_{i,j+1})) - \frac{\kappa_p^{f,\frac{1}{2}}}{8} \cdot 2^{-2i} \cdot ||\mathbf{q} - \breve{\mathbf{q}}||_p^2
$$
  
\n
$$
\leq \frac{1}{2} \Biggl[ (1 - t_{i,j})f(c(0)) + t_{i,j}f(c(1)) - \frac{\kappa_p^{f,\frac{1}{2}}}{2}t_{i,j}(1 - t_{i,j}) ||\mathbf{q} - \breve{\mathbf{q}}||_p^2 \Biggr]
$$
  
\n
$$
+ \frac{1}{2} \Biggl[ (1 - t_{i,j+1})f(c(0)) + t_{i,j+1}f(c(1)) - \frac{\kappa_p^{f,\frac{1}{2}}}{2}t_{i,j+1}(1 - t_{i,j+1}) ||\mathbf{q} - \breve{\mathbf{q}}||_p^2 \Biggr]
$$
  
\n
$$
- \frac{\kappa_p^{f,\frac{1}{2}}}{8} \cdot 2^{-2i} \cdot ||\mathbf{q} - \breve{\mathbf{q}}||_p^2
$$
  
\n
$$
= (1 - t_{i+1,2j+1})f(c(0)) + t_{i+1,2j+1}f(c(1)) - \frac{\kappa_p^{f,\frac{1}{2}}}{2}t_{i+1,2j+1}(1 - t_{i+1,2j+1}) ||\mathbf{q} - \breve{\mathbf{q}}||_p^2
$$

as desired. Noting again that the inequality [\(17\)](#page-27-1) holds for  $t_{i+1,2j+1}$  with  $j = 2^i$ , we have shown the induction case for  $0 \leq j \leq 2^{i+1}$ .

Because f is continuous on  $\triangle^N$  and  $\{t_{i,j} \in [0, 1] \mid (i, j) \in \mathbb{N} \times \mathbb{Z}_{\geq 0}\}$  is dense in  $[0, 1]$ , we find

$$
f((1-t)\mathbf{q}+t\breve{\mathbf{q}}) \leq (1-t)f(\mathbf{q})+tf(\breve{\mathbf{q}})-\frac{\kappa_p^{f,\frac{1}{2}}}{2}t(1-t)\|\mathbf{q}-\breve{\mathbf{q}}\|_p^2 \quad \text{for } t \in (0,1),
$$

which leads to

$$
\kappa_p^{f,\frac{1}{2}} \leq \frac{2\big[(1-t)f(\mathbf{q}) + tf(\check{\mathbf{q}}) - f((1-t)\mathbf{q} + t\check{\mathbf{q}})\big]}{t(1-t)\|\mathbf{q} - \check{\mathbf{q}}\|_p^2} \quad \text{for } t \in (0,1).
$$

Since  $\mathbf{q}, \tilde{\mathbf{q}} \in \Delta^N$  are arbitrary, this ensures that  $\kappa_p^{f,t} \geqslant \kappa_p^{f,1/2}$  for  $t \in (0,1)$  and completes the proof of the proposition.  $\Box$  **Lemma 18** Let  $f : \triangle^N \to \mathbb{R}$  be a continuous convex function. Then, for any  $\mathbf{q}, \check{\mathbf{q}} \in \triangle^N$ and  $r \in (0, 2^{1/p}],$ 

$$
\liminf_{r\downarrow 0} K_p^f(r) = \kappa_p^f, \quad J(\mathbf{q}, \breve{\mathbf{q}}) \geqslant \frac{\kappa_p^f}{8} \|\mathbf{q} - \breve{\mathbf{q}}\|_p^2, \quad \text{and} \quad D^- \omega(r) \geqslant \frac{\kappa_p^f}{4} r.
$$

**Proof** Assume that there exists  $r_* \in (0, 2^{1/p}]$  such that  $K_p^f(r_*) = \kappa_p^f$ . We can see from the bound [\(13\)](#page-17-3) that

$$
\frac{K_p^f((1-2\tau)r_*)}{8}[(1-2\tau)r_*]^2 \leq \frac{K_p^f(r_*)}{8}r^2 - \frac{\kappa_p^f}{2}\tau(1-\tau)r_*^2 = \frac{K_p^f(r_*)}{8}[(1-2\tau)r_*]^2
$$

for  $\tau \in (0, 1/2)$ . Since  $K_p^f(r) \geq \kappa_p^f$  for  $r \in (0, 2^{1/p}]$ , this implies  $K_p^f(r) = \kappa_p^f$  for  $r \in (0, r_*]$ hence

$$
\liminf_{r \downarrow 0} K_p^f(r) = \kappa_p^f.
$$

Assume that there is no  $r \in (0, 2^{1/p}]$  so that  $K_p^f(r) = \kappa_p^f$ . Then, there exists  $(r_j)_{j \in \mathbb{N}} \subseteq (0, 2^{1/p}]$ converging to 0 such that

$$
\lim_{j \to \infty} K_p^f(r_j) = \inf \left\{ K_p^f(r) \mid r \in (0, 2^{\frac{1}{p}}] \right\} = \kappa_p^{f, \frac{1}{2}} \leq \liminf_{r \downarrow 0} K_p^f(r) \leq \lim_{j \to \infty} K_p^f(r_j),
$$

where the second equality follows from  $(12)$ . This with [Proposition 14](#page-15-0) proves the first assertion.

For  $\mathbf{q}, \check{\mathbf{q}} \in \triangle$ , we calculate

$$
J(\mathbf{q}, \check{\mathbf{q}}) \geq \omega(\|\mathbf{q} - \check{\mathbf{q}}\|_p) = \frac{K_p^f(\|\mathbf{q} - \check{\mathbf{q}}\|_p)}{8}\|\mathbf{q} - \check{\mathbf{q}}\|_p^2 \geqslant \frac{\kappa_p^f}{8}\|\mathbf{q} - \check{\mathbf{q}}\|_p^2.
$$

This is the second assertion.

For  $r \in (0, 2^{1/p}]$ , we can pick  $\mathbf{q}, \check{\mathbf{q}} \in \triangle^N$  such that  $\omega(r) = J(\mathbf{q}, \check{\mathbf{q}})$  and  $\|\mathbf{q} - \check{\mathbf{q}}\|_p = r$  from [Lemma 10.](#page-10-2) For  $\tau \in (0, 1/2)$ , we have

$$
\omega((1-2\tau)r) \leq \omega(r) - \frac{\kappa_p^f}{2}\tau(1-\tau)r^2
$$

from [\(13\)](#page-17-3), which yields

$$
D^{-}\omega(r) = \limsup_{\tau \downarrow 0} \frac{\omega(r) - \omega((1 - 2\tau)r)}{r - (1 - 2\tau)r} \geq \limsup_{\tau \downarrow 0} \frac{\frac{\kappa_p^f}{2}\tau(1 - \tau)r^2}{2\tau r} = \frac{\kappa_p^f}{4}r.
$$

This completes the proof of the lemma.

# Appendix C. Derivation of modulus for general  $N$

In [§6,](#page-20-0) we mainly consider examples of  $\omega$  only for  $(N, p) = (2, 1)$ . Since we have extended the moduli on general  $(\Delta^N, \|\cdot\|_p)$  in [Definition 8,](#page-10-1) it is nice to have an example beyond the binary case. To this end, we calculate  $\omega$  for the Shannon entropy  $f(\mathbf{q}) = \langle \mathbf{q}, \ln \mathbf{q} \rangle$  with

 $\Box$ 

general  $N \ge 2$  and  $p = 2$ . In what follows, we focus on  $f(\mathbf{q}) = \langle \mathbf{q}, \ln \mathbf{q} \rangle$  with  $p = 2$ , and the modulus  $\omega$  and midpoint Jensen gap J is defined based on this particular f throughout this section.

Let

$$
\mathcal{U}^{N-1} := \left\{ \mathbf{u} \in (0,1)^{N-1} \, \middle| \, \sum_{n \in [N-1]} u_n < 1 \right\}
$$

and define  $\psi: \mathcal{U}^{N-1} \times \mathcal{U}^{N-1} \to \mathbb{R}$  and  $\phi: \mathcal{U}^{N-1} \to \mathbb{R}$  by  $\sqrt{ }$  $\mathsf{r}$   $\mathsf{r}$ 

$$
\psi(\mathbf{u}, \mathbf{w}) := \frac{1}{2} \Bigg\{ \sum_{n \in [N-1]} (u_n - w_n)^2 + \Bigg[ \sum_{n \in [N-1]} (u_n - w_n) \Bigg]^2 \Bigg\} \quad \text{for } \mathbf{u}, \mathbf{w} \in \mathcal{U}^{N-1},
$$

$$
\phi(\mathbf{u}) := \sum_{n \in [N-1]} u_n \ln u_n + \left( 1 - \sum_{n \in [N-1]} u_n \right) \ln \left( 1 - \sum_{n \in [N-1]} u_n \right) \quad \text{for } \mathbf{u} \in \mathcal{U}^{N-1},
$$

respectively. Here,  $\phi$  is the Shannon entropy but defined on the constrained set  $\mathcal{U}^{N-1}$ , where we drop the marginal constraint in the original  $\triangle^N$ . Indeed, we have

$$
\begin{bmatrix} u_1 \\ \vdots \\ u_{N-1} \\ 1 - \langle \mathbf{u}, \mathbf{1} \rangle \end{bmatrix} \in \Delta_+^N \text{ and } \psi(\mathbf{u}, \mathbf{w}) = \frac{1}{2} \left\| \begin{bmatrix} u_1 \\ \vdots \\ u_{N-1} \\ 1 - \langle \mathbf{u}, \mathbf{1} \rangle \end{bmatrix} - \begin{bmatrix} w_1 \\ \vdots \\ w_{N-1} \\ 1 - \langle \mathbf{w}, \mathbf{1} \rangle \end{bmatrix} \right\|_2^2,
$$

<span id="page-30-0"></span>which yields  $\psi(\mathbf{u}, \mathbf{w}) \in [0, 1]$ . First, we present a couple of necessary lemmas.

**Lemma 19** For  $r \in (0, 2^{1/2})$ , there exist  $\mathbf{q}, \check{\mathbf{q}} \in \triangle^N$  such that

$$
\|\mathbf{q} - \check{\mathbf{q}}\|_2 = r
$$
,  $\text{supp}(\mathbf{q}) \cap \text{supp}(\check{\mathbf{q}}) \neq \varnothing$ ,  $J(\mathbf{q}, \check{\mathbf{q}}) < \ln 2$ .

**Proof** Fix  $r \in (0, 2^{1/2})$  and set  $a := 2^{-1/2}r \in (0, 1)$ . Define  $\mathbf{q}, \check{\mathbf{q}} \in \triangle^N$  by

$$
q_n := \delta_{1n}, \quad \check{q}_n := (1-a)\delta_{1n} + a\delta_{2n} \quad \text{for } n \in [N].
$$

Then, we have  $\|\mathbf{q} - \check{\mathbf{q}}\|_2 = r$ ,  $\text{supp}(\dot{\mathbf{q}}) \wedge \text{supp}(\check{\mathbf{q}}) \neq \emptyset$ , and

$$
\overline{J}(a) := J(\mathbf{q}, \check{\mathbf{q}}) = \frac{1-a}{2} \ln(1-a) - \left(1 - \frac{a}{2}\right) \ln\left(1 - \frac{a}{2}\right) + \frac{a}{2} \ln 2.
$$

Since we have

$$
\overline{J}(1) = \ln 2, \quad \overline{J}'(a) = \frac{1}{2} \ln \frac{2-a}{1-a} > 0,
$$

we conclude  $J(\mathbf{q}, \check{\mathbf{q}}) < \ln 2$  as desired.

<span id="page-30-1"></span>Lemma 20 For  $\mathbf{q}, \check{\mathbf{q}} \in \triangle^N$ , if

$$
\mathrm{supp}(\mathbf{q})\cap\mathrm{supp}(\check{\mathbf{q}})\neq\varnothing,\quad\mathrm{supp}(\mathbf{q})\neq\mathrm{supp}(\check{\mathbf{q}}),
$$

then there exist  $\mathbf{q}', \check{\mathbf{q}}' \in \triangle^N$  such that

$$
\|\mathbf{q}-\mathbf{q}\|_2 = \|\mathbf{q}'-\breve{\mathbf{q}}'\|_2, \quad \mathrm{supp}(\mathbf{q}') = \mathrm{supp}(\breve{\mathbf{q}}'), \quad J(\mathbf{q}',\breve{\mathbf{q}}') < J(\mathbf{q},\breve{\mathbf{q}}).
$$

 $\Box$ 

**Proof** Take  $i \in \text{supp}(q) \cap \text{supp}(\check{q})$  and write

$$
S_1 \coloneqq [\text{supp}(\mathbf{q}) \cap \text{supp}(\breve{\mathbf{q}})] \setminus \{i\}, \quad S_2 \coloneqq \text{supp}(\mathbf{q}) \setminus \text{supp}(\breve{\mathbf{q}}), \quad S_3 \coloneqq \text{supp}(\breve{\mathbf{q}}) \setminus \text{supp}(\mathbf{q}),
$$

and  $m := |S_2 \cup S_3|$ . For sufficiently small  $\varepsilon > 0$ , define  $\mathbf{q}^{\varepsilon}, \check{\mathbf{q}}^{\varepsilon} \in \triangle^N$  by

$$
q_n^{\varepsilon} := \begin{cases} q_i - m\varepsilon & \text{for } n = i, \\ q_n & \text{for } n \in S_1, \\ q_n + \varepsilon & \text{for } n \in S_2 \cup S_3, \\ 0 & \text{otherwise,} \end{cases} \qquad \widetilde{q}_n^{\varepsilon} := \begin{cases} \widetilde{q}_i - m\varepsilon & \text{for } n = i, \\ \widetilde{q}_n & \text{for } n \in S_1, \\ \widetilde{q}_n + \varepsilon & \text{for } n \in S_2 \cup S_3, \\ 0 & \text{otherwise.} \end{cases}
$$

We find that  $\|\mathbf{q}^{\varepsilon}-\check{\mathbf{q}}^{\varepsilon}\|_2 = \|\mathbf{q}-\check{\mathbf{q}}\|_2$ ,  $\text{supp}(\mathbf{q}^{\varepsilon}) = \text{supp}(\check{\mathbf{q}}^{\varepsilon})$ , and

$$
\lim_{\varepsilon\downarrow 0}J({\bf q}^\varepsilon,\widecheck{\bf q}^\varepsilon)=J({\bf q},\widecheck{\bf q}),
$$

thanks to the continuity of J. Since we have

$$
\frac{\partial}{\partial \varepsilon} J(\mathbf{q}^{\varepsilon}, \breve{\mathbf{q}}^{\varepsilon}) = -\frac{m}{2} \ln \frac{(q_i - m\varepsilon)(\breve{q}_i - m\varepsilon)}{\left(\frac{q_i + \breve{q}_i}{2} - m\varepsilon\right)^2} + \frac{1}{2} \sum_{n \in S_2} \ln \frac{(q_n + \varepsilon)\varepsilon}{\left(\frac{q_n}{2} + \varepsilon\right)^2} + \frac{1}{2} \sum_{n \in S_3} \ln \frac{(\breve{q}_n + \varepsilon)\varepsilon}{\left(\frac{\breve{q}_n}{2} + \varepsilon\right)^2},
$$

which diverges to  $-\infty$  as  $\varepsilon \downarrow 0$ , we have  $J(q^{\varepsilon}, \check{q}^{\varepsilon}) < J(q, \check{q})$  for sufficiently small  $\varepsilon > 0$ . This completes the proof of the lemma.  $\Box$ 

Corollary 21 For  $\mathbf{q}, \check{\mathbf{q}} \in \triangle^N$  with  $\|\mathbf{q} - \check{\mathbf{q}}\|_2 < 2^{1/2}$ ,  $\omega(\|\mathbf{q} - \check{\mathbf{q}}\|_2) = J(\mathbf{q}, \check{\mathbf{q}})$  holds if and only  $if \text{supp}(\mathbf{q}) = \text{supp}(\breve{\mathbf{q}}).$ 

**Proof** Let  $\mathbf{q}, \check{\mathbf{q}} \in \triangle^N$  satisfy  $\|\mathbf{q} - \check{\mathbf{q}}\|_2 < 2^{1/2}$ . We observe from [Lemma 19](#page-30-0) that  $J(\mathbf{q}, \check{\mathbf{q}}) < \ln 2$ . On the other hand,  $J(\mathbf{q}', \check{\mathbf{q}}') = \ln 2$  follows for  $\mathbf{q}', \check{\mathbf{q}}' \in \triangle^N$  with  $\text{supp}(\mathbf{q}') \cap \text{supp}(\check{\mathbf{q}}') =$  $\emptyset$ . Thus,  $\omega(\|\mathbf{q} - \check{\mathbf{q}}\|_2) = J(\mathbf{q}, \check{\mathbf{q}})$  implies supp $(\mathbf{q}) = \text{supp}(\check{\mathbf{q}})$ . The converse implication immediately follows from [Lemma 20.](#page-30-1)

Thus, the proof of the corollary is complete.

<span id="page-31-1"></span>
$$
\Box
$$

Subsequently, we show [Lemmas 22](#page-31-0) and [23,](#page-32-0) which are needed to invoke the method of Lagrangian multipliers later.

<span id="page-31-0"></span>**Lemma 22** For  $\mathbf{u}, \mathbf{w} \in \mathcal{U}^{N-1}$ , the rank of the Jacobian of  $\psi$  at  $(\mathbf{u}, \mathbf{w})$  is zero if and only if  $u$  and  $w$  are linearly dependent.

Proof Since we have

$$
\frac{\partial \psi}{\partial u_i}(\mathbf{u}, \mathbf{w}) = u_i - w_i + \sum_{n \in [N-1]} (u_n - w_n) = -\frac{\partial \psi}{\partial w_i} \quad \text{for } i \in [N-1],
$$

the rank of the Jacobian of  $\psi$  at  $(\mathbf{u}, \mathbf{w})$  is zero if and only if

$$
u_i - w_i + \sum_{n \in [N-1]} (u_n - w_n) = 0
$$
 for  $i \in [N-1]$ .

Summing the above equation up gives

$$
N \sum_{n \in [N-1]} (u_n - w_n) = 0,
$$

and hence  $u_i - w_i = 0$  for all  $i \in [N - 1]$ , which shows the linear dependence of **u** and **w**. The converse implication is trivial.  $\Box$ 

<span id="page-32-0"></span>**Lemma 23** For  $r \in (0, 2^{1/2})$ , let  $\mathbf{u}, \mathbf{w} \in \mathcal{U}^{N-1}$  satisfy  $\psi(\mathbf{u}, \mathbf{w}) = r^2/2$ . If there exists  $\lambda \in \mathbb{R}$ such that

<span id="page-32-1"></span>
$$
\frac{1}{2} \frac{\partial \phi}{\partial u_n}(\mathbf{u}) - \frac{\partial \phi}{\partial u_n} \left( \frac{\mathbf{u} + \mathbf{w}}{2} \right) + \lambda \frac{\partial \psi}{\partial u_n}(\mathbf{u}, \mathbf{w}) = 0 \quad and
$$
\n
$$
\frac{1}{2} \frac{\partial \phi}{\partial u_n}(\mathbf{u}) - \frac{\partial \phi}{\partial u_n} \left( \frac{\mathbf{u} + \mathbf{w}}{2} \right) + \lambda \frac{\partial \psi}{\partial w_n}(\mathbf{u}, \mathbf{w}) = 0
$$
\n(18)

for  $n \in [N - 1]$ . Then,

$$
u_n = \frac{1}{(N-1)(1+\mu)}, \qquad w_n = \frac{\mu}{(N-1)(1+\mu)} \qquad \text{for } n \in [N-1],
$$

where  $\mu \in (0, 1)$  satisfies

$$
\frac{N(1-\mu)^2}{(N-1)(1+\mu)^2} = r^2.
$$

Proof For simplicity, set

$$
\mathbf{v}:=\frac{\mathbf{u}+\mathbf{w}}{2},\quad u_N:=1-\sum_{n\in[N-1]}u_n,\quad w_N:=1-\sum_{n\in[N-1]}w_n,\quad\text{and}\quad v_N:=1-\sum_{n\in[N-1]}v_n.
$$

Assuming [Eq. \(18\),](#page-32-1) we calculate

$$
0 = \frac{1}{2} \frac{\partial \phi}{\partial u_i}(\mathbf{u}) - \frac{\partial \phi}{\partial u_i}(\mathbf{v}) + \lambda \frac{\partial \psi}{\partial u_i}(\mathbf{u}, \mathbf{w}) = \frac{1}{2} \left( \ln \frac{u_i}{u_N} - \ln \frac{v_i}{v_N} \right) + \lambda [u_i - w_i - (u_N - w_N)],
$$
  
\n
$$
0 = \frac{1}{2} \frac{\partial \phi}{\partial w_i}(\mathbf{w}) - \frac{\partial \phi}{\partial w_i}(\mathbf{v}) + \lambda \frac{\partial \psi}{\partial w_i}(\mathbf{u}, \mathbf{w}) = \frac{1}{2} \left( \ln \frac{w_i}{w_N} - \ln \frac{v_i}{v_N} \right) - \lambda [u_i - w_i - (u_N - w_N)],
$$

for  $i \in [N - 1]$ , which yields

$$
\ln \frac{u_i}{u_N} - \ln \frac{v_i}{v_N} = -2\lambda [u_i - w_i - (u_N - w_N)] = \ln \frac{u_j}{u_N} - \ln \frac{v_j}{v_N}
$$

$$
= -\ln \frac{w_i}{w_N} + \ln \frac{v_i}{v_N}
$$

for  $i, j \in [N - 1]$ . Thus, we have

$$
\frac{u_i}{u_j} = \frac{v_i}{v_j} = \frac{w_i}{w_j}, \quad \frac{u_i}{u_N} \cdot \frac{w_i}{w_N} = \frac{v_i^2}{v_N^2} \quad \text{for } i, j \in [N-1].
$$

Then, there exists  $\mu > 0$  such that

$$
w_i = \mu u_i \quad \text{for } i \in [N-1].
$$

It follows from  $\psi(\mathbf{u}, \mathbf{w}) \neq 0$  that  $\mu \neq 1$ . This gives the relation

$$
v_i = \frac{1+\mu}{2}u_i
$$
 for  $i \in [N-1]$ ,  $v_N = \frac{(1+\mu)u_N + 1 - \mu}{2}$ ,  $w_N = \mu u_N + 1 - \mu$ ,

and we have

$$
\frac{\mu u_i^2}{u_N(\mu u_N + 1 - \mu)} = \frac{u_i}{u_N} \cdot \frac{w_i}{w_N} = \frac{v_i^2}{v_N^2} = \frac{(1 + \mu)^2 u_i^2}{[(1 + \mu)u_N + 1 - \mu]^2},
$$

which is equivalent to

$$
u_N = \frac{\mu}{1+\mu}.
$$

these imply

$$
0 = \ln \frac{u_i}{u_N} - \ln \frac{v_i}{v_N} + 2\lambda [u_i - w_i - (u_N - w_N)] = \ln \frac{1}{\mu} + 2\lambda (1 - \mu) \left( u_i + \frac{1}{1 + \mu} \right),
$$

and hence

$$
u_i = \frac{1}{2\lambda(1-\mu)} \ln \mu - \frac{1}{1+\mu} \quad \text{for } i \in [N-1].
$$

This with the relation

$$
1 = \sum_{n \in [N]} u_n = \frac{N-1}{2\lambda(1-\mu)} \ln \mu - \frac{N-1}{1+\mu} + \frac{\mu}{1+\mu}
$$

provides

$$
\lambda = \frac{(N-1)(1+\mu)}{2N(1-\mu)} \ln \mu \quad \text{and} \quad u_i = \frac{1}{(N-1)(1+\mu)} \quad \text{for } i \in [N-1].
$$

Thus, we have

$$
r^{2} = 2\psi(\mathbf{u}, \mathbf{w}) = (1 - \mu)^{2} \sum_{n \in [N-1]} u_{n}^{2} + (1 - \mu)^{2} \frac{1}{(1 + \mu)^{2}} = \frac{N(1 - \mu)^{2}}{(N-1)(1 + \mu)^{2}},
$$

as desired.

By combining these lemmas, we have the following claim, which is the minimizers  $(q, \check{q})$ of the Shannon entropy we show in [§6.](#page-20-0)

 $\Box$ 

**Proposition 24** For  $r \in (0, 2^{1/2})$ ,  $\mathbf{q}, \check{\mathbf{q}} \in \triangle^N$  satisfy  $\|\mathbf{q} - \check{\mathbf{q}}\|_2 = r$  and  $\omega(r) = J(\mathbf{q}, \check{\mathbf{q}})$  if and only if there exist distinct  $i, j \in [N]$  such that

<span id="page-33-0"></span>
$$
q_n := \frac{1 + 2^{-1/2}r}{2} \delta_{ni} + \frac{1 - 2^{-1/2}r}{2} \delta_{nj}, \quad \check{q}_n := \frac{1 - 2^{-1/2}r}{2} \delta_{ni} + \frac{1 + 2^{-1/2}r}{2} \delta_{nj}.
$$

**Proof** There exist  $\mathbf{q}'$ ,  $\breve{\mathbf{q}}' \in \triangle^N$  such that

$$
\|\mathbf{q}' - \breve{\mathbf{q}}'\|_2 = r
$$
,  $\omega(r) = J(\mathbf{q}, \breve{\mathbf{q}})$ , and  $\text{supp}(\mathbf{q}') = \text{supp}(\breve{\mathbf{q}}')$ 

from [Lemma 10](#page-10-2) and [Corollary 21.](#page-31-1) By relabeling the indices  $n \in [N]$  and switching  $\mathbf{q}'$  and  $\check{\mathbf{q}}'$ if necessary, we may assume that  $q'_1 > \tilde{q}'_1$  and there exists  $N' \in [N]$  such that  $N' \geq 2$  and

$$
\mathrm{supp}(\mathbf{q}') = \mathrm{supp}(\breve{\mathbf{q}}') = [N'].
$$

Define  $\mathbf{u}', \mathbf{w}' \in \mathcal{U}^{N-1}$  by

$$
u'_n = q'_n, \quad w'_n = \check{q}'_n \quad \text{for } n \in [N'-1].
$$

Then, the pair  $(\mathbf{u}', \mathbf{w}')$  minimizes  $\phi$  subject to  $\phi(\mathbf{u}, \mathbf{w}) = r^2/2$  on  $\mathcal{U}^{N-1} \times \mathcal{U}^{N-1}$ . We observe from the method of Lagrangian multipliers together with [Lemmas 22](#page-31-0) and [23](#page-32-0) that

$$
u'_{n} = \frac{1}{(N'-1)(1+\mu_{N'})}, \quad w'_{n} = \frac{\mu_{N'}}{(N'-1)(1+\mu_{N'})} \quad \text{for } n \in [N'-1],
$$

where  $\mu_{N'} \in (0, 1)$  satisfies

$$
\frac{N'(1-\mu_{N'})^2}{(N'-1)(1+\mu_{N'})^2} = r^2.
$$

This implies

$$
q_{N'} = \frac{\mu_{N'}}{1 + \mu_{N'}}
$$
,  $\breve{q}_{N'} = \frac{1}{1 + \mu_{N'}}$ , and  $J(\mathbf{q}', \breve{\mathbf{q}}') = -\ln(1 + \mu_{N'}) + \frac{\mu_{N'}}{1 + \mu_{N'}}$   $\ln \mu_{N'} + \ln 2$ .

By setting

$$
\bar{J}(\mu) := -\ln(1+\mu) + \frac{\mu}{1+\mu}\ln\mu + \ln 2 \quad \text{for } \mu \in (0,1),
$$

we see that

$$
\bar{J}'(\mu) = \frac{1}{(1+\mu)^2} \ln \mu < 0
$$

and  $\mu_{N'} \leq \mu_{n'}$  holds for  $N' \leq n'$ . Thus, we conclude that  $N' = 2$ . Since we have

$$
\mu_2 = \frac{1 - 2^{-1/2}r}{1 + 2^{-1/2}r},
$$

this in turn shows

$$
q_n' = \frac{1 + 2^{-1/2}r}{2} \delta_{n1} + \frac{1 - 2^{-1/2}r}{2} \delta_{n2}, \qquad \check{q}_n' = \frac{1 - 2^{-1/2}r}{2} \delta_{n1} + \frac{1 + 2^{-1/2}r}{2} \delta_{n2}.
$$

By the invariance of J under relabeling the indices  $n \in [N]$ , the proof of the proposition is completed.  $\Box$