

ON THE SPECTRUM OF ELECTRIC QUANTUM WALK AND RELATED CMV MATRICES

FAN YANG

ABSTRACT. In this note, we show that for a family of quantum walk models with electric fields, the spectrum is the unit circle for any irrational field. The result also holds for the associated CMV matrices defined by skew-shifts. Generalizations to CMV matrices with skew-shifts on higher dimensional torus are also obtained.

1. INTRODUCTION

The quantum walk model is a quantum mechanical analogue of the classical random walk. It can also be used to describe quantum simulation of a particle on a lattice or graph. This model has demonstrated a wide range of applications across various fields, including mathematical physics and quantum information, attracting significant attention over the past few decades, see e.g. [5, 34, 39, 44, 45, 47].

It was discovered by Cantero, Moral, Grünbaum and Velázquez [16] that quantum walk models can be reduced to CMV matrices, a family of extensively studied unitary operators that have further connections to orthogonal polynomials on the unit circle (OPUC) [41, 42]. This connection brings the tools that were largely developed in the spectral theory community into the study of quantum walks.

The study of spectrum structure is an important topic in spectral theory and mathematical physics. One class of operator that was studied widely is quasi-periodic Schrödinger operators. For example, the famous Ten Martini problem, named after Kac and Simon, conjectured Cantor spectrum for the almost Mathieu operators (quasi-periodic operator with cosine potential) for any irrational magnetic flux. The complete proof of this conjecture was given by Avila-Jitomirskaya [2], with earlier important advances by Bellissard-Simon [9], Choi-Elliott-Yui [20], Last [37], Puig [38] and Avila-Krikorian [3]. Goldstein and Schlag proved Cantor spectrum for quasi-periodic operators with general analytic potentials defined on \mathbb{T} in the positive Lyapunov exponent regime for a.e. magnetic flux [24]. See [6, 7, 17, 22, 26, 32, 43] for Cantor spectrum results for some other quasi-periodic models. Though Cantor spectrum is generally expected for many 1D single-frequency quasi-periodic operators, there are some exceptions. One such known example is the Maryland model, with a tan potential, for which the spectrum was proved to be the whole real line [40]. The monotonicity of the tan potential also leads to other unique features, e.g. the lack of phase resonance, see [28, 30, 31]. For multi-frequency or higher dimensional operators, interval spectrum is conjectured. A significant advancement in the multi-frequency case was made by Goldstein-Schlag-Voda in [25], where they solved this problem up to the spectral edges for large coupling constants. For higher dimensional continuum quasi-periodic operators, a recent breakthrough of Karpeshina-Parnowski-Shterenberg [33] established the Bethe-Sommerfeld conjecture for generic potentials, hence proving finiteness of the number of gaps.

In this short note, we show that a quantum walk model in electric field, studied actively by physicists in recent years [4, 15, 18, 19, 23], does not have spectral gaps. Our proof is elementary but the result provides a new example of non-Cantor spectrum for a 1D single-frequency quasi-periodic operators. Another interesting feature of the model is that it can be reduced to a CMV matrix defined by skew-shift. The skew-shift operators, see the dynamics in (1.2), is a different family of ergodic operators, hosting a number of major open problems. One of the conjectures is that skew-shift models do not have spectral gaps, see Chapter 15 of [11]. The results in this note, Theorem 1.3 and 1.5, provide examples for which this skew-shift conjecture is true. See also [12, 13, 27, 29, 35, 36] for some related works concerning the skew-shift dynamics.

Next let us introduce the model and our results in details. We study the discrete time quantum walk of one particle on \mathbb{Z} with two dimensional internal degree of freedom with external electric fields. Compare to quantum walk with external magnetic field, for which there is a vast literature of mathematical studies recently, see e.g. [1, 17, 46], quantum walk with electric field is less explored.

The quantum walk model acts on $\ell^2(\mathbb{Z}) \otimes \mathbb{C}^2 =: \mathcal{H}$ with the following basis:

$$\delta_n^\pm = \delta_n \otimes e_\pm, \quad n \in \mathbb{Z},$$

where $\{\delta_n\}$ is the standard basis of $\ell^2(\mathbb{Z})$ and $\{e_+ = (1, 0)^T, e_- = (0, 1)^T\}$ is the standard basis of \mathbb{C}^2 . Let $C \in U(2)$ be a coin matrix, defined by

$$C = e^{2\pi i \eta} \begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix},$$

in which $|a|^2 + |b|^2 = 1$ and $\eta \in \mathbb{T}$.

Let S be a conditional shift operator, defined as

$$S\delta_n^\pm = \delta_{n\pm 1}^\pm, \quad \text{or } (S\psi)_n^\pm = \psi_{n\mp 1}^\pm, \quad \text{for } \psi \in \ell^2(\mathbb{Z}) \otimes \mathbb{C}^2,$$

The evolution of the quantum walk for each step is described by a unitary operator, which can be represented as a product CS , i.e. a shift-coin quantum walk. Here we consider the following electric walk W acting on \mathcal{H} ,

$$W_{\omega, \theta, \eta, a, b} := Q_{\omega, \theta, \eta, a, b} S,$$

where $\omega \in \mathbb{T}$ (the discrete electric field), $\theta \in \mathbb{T}$ and $Q_{\omega, \theta, \eta, a, b}$ acts coordinate-wise via a matrix multiplication $Q_{\omega, \theta, \eta, a, b, n}$ defined by

$$Q_{\omega, \theta, \eta, a, b, n}(\psi_n^-, \psi_n^+)^T = e^{2\pi i(\theta + n\omega)} C(\psi_n^-, \psi_n^+)^T.$$

In the following, we shall identify $\ell^2(\mathbb{Z}) \otimes \mathbb{C}^2 \rightarrow \ell^2(\mathbb{Z})$ via

$$\delta_n^+ \mapsto \delta_{2n+1}, \quad \delta_n^- \mapsto \delta_{2n}.$$

Under such identification, $W_{\omega, \theta, \eta, a, b}$ can be written as a five diagonal matrix, see below.

Let $a, b \in [0, 1]$ satisfying $a^2 + b^2 = 1$. For each $n \in \mathbb{Z}$, $\tilde{\alpha}_n$ is defined as

$$\alpha_{2n}(\vec{x}) := be^{2\pi i \mathcal{P}_d(T_{d,\omega}^n(\vec{x}))}, \rho_{2n} \equiv a, \text{ and } \alpha_{2n+1} \equiv 0, \rho_{2n+1} \equiv 1.$$

Let $\mathcal{E}_{\omega,a,b}^d(\vec{x})$ be the associated CMV matrix determined by $\tilde{\alpha}_n$ and ρ_n as its Verblunsky coefficients. It is known that for irrational ω , $T_{d,\omega}$ is minimal [14], and hence the spectrum $\sigma(\mathcal{E}_{\omega,a,b}^d(\vec{x})) =: \sigma(\mathcal{E}_{\omega,a,b}^d)$ is independent of \vec{x} , see e.g. [8, 10, 21]. For this model, we have:

Theorem 1.5. *For any irrational ω , and $a, b \in [0, 1]$ such that $a^2 + b^2 = 1$. we have*

$$\sigma(\mathcal{E}_{\omega,a,b}^d) = \partial\mathbb{D}.$$

The rest of the paper is organized as follows: Section 2 provides preliminaries for Theorem 1.5. We present the short proof of Theorem 1.3 in Sec. 3 and the proof of Theorem 1.5 in Sec. 4.

2. PRELIMINARIES

2.1. Some facts about combinatorial numbers.

For $k \geq 0$, the following is the combinatorial number, with the convention that $0! = 1$,

$$\binom{n}{k} = \begin{cases} \frac{n!}{k!(n-k)!}, & \text{if } n \geq k \geq 0, \\ 0, & \text{if } 0 \leq n < k. \end{cases}$$

For $n \leq -1$ and $k \geq 1$, the following definition is standard:

$$(2.3) \quad \binom{n}{k} = \frac{n \cdots (n-k+1)}{k!} = (-1)^k \frac{(-n) \cdots (-n+k-1)}{k!} = (-1)^k \binom{-n+k-1}{k}.$$

The definition also applies when $n \leq -1$ and $k = 0$, hence

$$\binom{n}{0} = (-1)^0 \binom{-n-1}{0} = 1, \text{ if } n \leq -1.$$

With these notations, it is easy to check that for arbitrary $n \in \mathbb{Z}$ (including $n \leq -1$),

$$T_{d,\omega}^n(x_1, \dots, x_d) = (x_1 + \binom{n}{1}\omega, x_2 + \binom{n}{1}x_1 + \binom{n}{2}\omega, \dots, x_d + \sum_{k=1}^{d-1} \binom{n}{k}x_{d-k} + \binom{n}{d}\omega)$$

Recall that $\mathcal{P}_d(x_1, \dots, x_d) = x_d$ is the projection onto the d -th component. Clearly,

$$\mathcal{P}_d(T_{d,\omega}^n(x_1, \dots, x_d)) = x_d + \sum_{k=1}^{d-1} \binom{n}{k}x_{d-k} + \binom{n}{d}\omega.$$

The following is the well-known Pascal's triangle formula for $n \geq 0$ and $k \geq 1$.

$$(2.4) \quad \binom{n}{k} + \binom{n}{k-1} = \binom{n+1}{k}.$$

This formula also holds for $n \leq -1$, and is probably well-known too. We verify below for completeness.

For $n = -1$ and $k \geq 1$, by (2.3)

$$\binom{-1}{k} + \binom{-1}{k-1} = (-1)^k \binom{k}{k} + (-1)^{k-1} \binom{k-1}{k-1} = 0 = \binom{0}{k},$$

where

$$(4.9) \quad \beta_j(\vec{x}_-, \omega) := -\frac{1}{2} \left(x_{d-1} + \sum_{k=2}^{d-1} \binom{j}{k-1} x_{d-k} + \binom{j}{d-1} \omega \right),$$

in which $\vec{x}_- := (x_1, \dots, x_{d-1})$. In particular (4.8) implies

$$(4.10) \quad \sigma(\mathcal{E}_{\omega, a, b}^d(\vec{x})) = \sigma(W_{\beta(\vec{x}_-, \omega), a, b}).$$

Proof. With $D = \text{diag}(e^{2\pi i \lambda_j})$, we have

$$\begin{aligned} (D\mathcal{E}_{\omega, a, b}^d(\vec{x})D^{-1})_{2j, 2j-1} &= b e^{-2\pi i \psi_j} \cdot e^{2\pi i \lambda_{2j}} \cdot e^{-2\pi i \lambda_{2j-1}} = b e^{2\pi i(-\psi_j - \lambda_{2j-1} + \lambda_{2j})}, \\ (D\mathcal{E}_{\omega, a, b}^d(\vec{x})D^{-1})_{2j+1, 2j-1} &= a^* e^{2\pi i \lambda_{2j+1}} \cdot e^{-2\pi i \lambda_{2j-1}} = a^* e^{2\pi i(\lambda_{2j+1} - \lambda_{2j-1})} \\ (D\mathcal{E}_{\omega, a, b}^d(\vec{x})D^{-1})_{2j, 2j+2} &= a e^{2\pi i \lambda_{2j}} \cdot e^{-2\pi i \lambda_{2j+2}} = a e^{2\pi i(\lambda_{2j} - \lambda_{2j+2})} \\ (D\mathcal{E}_{\omega, a, b}^d(\vec{x})D^{-1})_{2j+1, 2j+2} &= -b^* e^{2\pi i \psi_j} \cdot e^{2\pi i \lambda_{2j+1}} \cdot e^{-2\pi i \lambda_{2j+2}} = -b^* e^{2\pi i(\psi_j + \lambda_{2j+1} - \lambda_{2j+2})}. \end{aligned}$$

Taking

$$\lambda_{2j} = \psi_j/2, \text{ and } \lambda_{2j+1} = -\psi_j/2,$$

we have

$$(4.11) \quad \begin{cases} -\psi_j - \lambda_{2j-1} + \lambda_{2j} = -\frac{1}{2}(\psi_j - \psi_{j-1}) \\ \lambda_{2j+1} - \lambda_{2j-1} = -\frac{1}{2}(\psi_j - \psi_{j-1}) \\ \lambda_{2j} - \lambda_{2j+2} = -\frac{1}{2}(\psi_{j+1} - \psi_j) \\ \psi_j + \lambda_{2j+1} - \lambda_{2j+2} = -\frac{1}{2}(\psi_{j+1} - \psi_j). \end{cases}$$

Direct computations using the Pascal's triangle (2.4) yields:

$$\begin{aligned} \psi_j - \psi_{j-1} &= \sum_{k=1}^{d-1} \left(\binom{j}{k} - \binom{j-1}{k} \right) x_{d-k} + \left(\binom{j}{d} - \binom{j-1}{d} \right) \omega \\ &= \sum_{k=1}^{d-1} \binom{j-1}{k-1} x_{d-k} + \binom{j-1}{d-1} \omega \\ &= x_{d-1} + \sum_{k=2}^{d-1} \binom{j-1}{k-1} x_{d-k} + \binom{j-1}{d-1} \omega = -2\beta_{j-1}(\vec{x}_-, \omega), \end{aligned}$$

in which we used $\binom{m}{0} \equiv 1$ for any $m \in \mathbb{Z}$. Combining the above with (4.11), we have

$$D\mathcal{E}_{\omega, a, b}^d(\vec{x})D^{-1} = W_{\beta(\vec{x}_-, \omega), a, b}.$$

This is as claimed. \square

Now we are in place to prove Theorem 1.5. Note that for any $\tau \in \mathbb{R}$, by the minimality of the skew-shift $T_{d, \omega}$, the spectrum set is independent of $\vec{x} \in \mathbb{T}^d$, in particular,

$$(4.12) \quad \sigma(\mathcal{E}_{\omega, a, b}^d(\vec{x})) = \sigma(\mathcal{E}_{\omega, a, b}^d(\vec{x} + (0, \dots, 0, \tau, 0))),$$

note τ is added onto x_{d-1} . It is easy to check by (4.9), for any $j \in \mathbb{Z}$, denoting $\vec{x}_- + (0, \dots, 0, \tau) =: \vec{x}_-^{(\tau)}$, that

$$(4.13) \quad \beta_j(\vec{x}_-^{(\tau)}), \omega) = \beta_j(\vec{x}_-, \omega) - \frac{\tau}{2}.$$

Hence, by (4.7) and (4.13),

$$W_{\beta(\vec{x}_-^{(\tau)}, \omega), a, b} = e^{-\pi i \tau} W_{\beta(\vec{x}_-, \omega), a, b},$$

which implies

$$\sigma(W_{\beta(\vec{x}_-^{(\tau)}, \omega), a, b}) = e^{-\pi i \tau} \cdot \sigma(W_{\beta(\vec{x}_-, \omega), a, b}).$$

Combining the above with (4.10), we have for any $\vec{x} \in \mathbb{T}^d$ that

$$\sigma(\mathcal{E}_{\omega, a, b}^d(\vec{x} + (0, \dots, 0, \tau, 0))) = e^{-\pi i \tau} \cdot \sigma(\mathcal{E}_{\omega, a, b}^d(\vec{x})).$$

This further combined with (4.12) implies

$$(4.14) \quad \sigma(\mathcal{E}_{\omega, a, b}^d(\vec{x})) = e^{-\pi i \tau} \cdot \sigma(\mathcal{E}_{\omega, a, b}^d(\vec{x})), \text{ for any } \tau \in \mathbb{R}.$$

Since $|a|^2 + |b|^2 = 1$, $\mathcal{E}_{\omega, a, b}(\vec{x})$ is unitary, which implies $\sigma(\mathcal{E}_{\omega, a, b}(\vec{x})) \subset \partial\mathbb{D}$. Combining this with (4.14) yields

$$\sigma(\mathcal{E}_{\omega, a, b}^d(\vec{x})) = \partial\mathbb{D}.$$

□

REFERENCES

- [1] Arnault, P., & Debbasch, F. (2016). Landau levels for discrete-time quantum walks in artificial magnetic fields. *Physica A: Statistical Mechanics and its Applications*, 443, 179-191.
- [2] Avila, A., & Jitomirskaya, S. (2009). The ten martini problem. *Annals of mathematics*, 303-342.
- [3] Avila, A., & Krikorian, R. (2006). Reducibility or nonuniform hyperbolicity for quasiperiodic Schrödinger cocycles. *Annals of Mathematics*, 911-940.
- [4] Arnault, P., Pepper, B., & Pérez, A. (2020). Quantum walks in weak electric fields and Bloch oscillations. *Physical Review A*, 101(6), 062324.
- [5] Abdul-Rahman, H., & Stolz, G. (2023). Exponentially decaying velocity bounds of quantum walks in periodic fields. *Communications in Mathematical Physics*, 403(3), 1297-1327.
- [6] Band, R., Beckus, S. and Loewy, R., (2024). The Dry Ten Martini Problem for Sturmian Hamiltonians. arXiv preprint arXiv:2402.16703.
- [7] Becker, S., Han, R. and Jitomirskaya, S., (2019). Cantor spectrum of graphene in magnetic fields. *Inventiones mathematicae*, 218, pp.979-1041.
- [8] Beckus, S., Lenz, D., Lindner, M. and Seifert, C., (2017). On the spectrum of operator families on discrete groups over minimal dynamical systems. *Mathematische Zeitschrift*, 287, pp.993-1007.
- [9] Bellissard, J. and Simon, B., (1982). Cantor spectrum for the almost Mathieu equation. *Journal of functional analysis*, 48(3), pp.408-419.
- [10] Bellissard, J., Iochum, B., Scoppola, E. and Testard, D., (1989). Spectral properties of one-dimensional quasi-crystals.
- [11] Bourgain, J., (2004). *Green's Function Estimates for Lattice Schrödinger Operators and Applications*.(AM-158). Princeton University Press.
- [12] Bourgain, J., (2007). Positive Lyapounov exponents for most energies. In *Geometric Aspects of Functional Analysis: Israel Seminar 1996–2000* (pp. 37-66). Berlin, Heidelberg: Springer Berlin Heidelberg.
- [13] Bourgain, J., Goldstein, M. and Schlag, W., (2001). Anderson Localization for Schrödinger Operators on \mathbb{Z} with Potentials Given by the Skew-Shift. *Communications in Mathematical Physics*, 220, pp.583-621.
- [14] Brin, M. and Stuck, G., (2002). *Introduction to dynamical systems*. Cambridge university press.
- [15] Bru, L. A., Hinarejos, M., Silva, F., de Valcárcel, G. J., & Roldán, E. (2016). Electric quantum walks in two dimensions. *Physical Review A*, 93(3), 032333.
- [16] Cantero, M. J., Moral, L., Grünbaum, F. A., & Velázquez, L. (2010). Matrix-valued Szegő polynomials and quantum random walks. *Communications on Pure and Applied Mathematics: A Journal Issued by the Courant Institute of Mathematical Sciences*, 63(4), 464-507.

- [17] Cedzich, C., Fillman, J., & Ong, D. C. (2023). Almost everything about the unitary almost Mathieu operator. *Communications in Mathematical Physics*, 403(2), 745-794.
- [18] Cedzich, C., Rybár, T., Werner, A. H., Alberti, A., Genske, M., & Werner, R. F. (2013). Propagation of quantum walks in electric fields. *Physical review letters*, 111(16), 160601.
- [19] Cedzich, C., & Werner, A. H. (2021). Anderson localization for electric quantum walks and skew-shift CMV matrices. *Communications in Mathematical Physics*, 387, 1257-1279.
- [20] Choi, M. D., Elliott, G. A., & Yui, N. (1990). Gauss polynomials and the rotation algebra. *Inventiones mathematicae*, 99(1), 225-246.
- [21] Cycon, H.L., Froese, R.G., Kirsch, W. and Simon, B., (2009). *Schrödinger operators: With application to quantum mechanics and global geometry*. Springer.
- [22] Damanik, D., Gorodetski, A. and Yessen, W., (2016). The fibonacci hamiltonian. *Inventiones mathematicae*, 206(3), pp.629-692.
- [23] Genske, M., Alt, W., Steffen, A., Werner, A. H., Werner, R. F., Meschede, D., & Alberti, A. (2013). Electric quantum walks with individual atoms. *Physical review letters*, 110(19), 190601.
- [24] Goldstein, M. and Schlag, W., (2011). On resonances and the formation of gaps in the spectrum of quasi-periodic Schrödinger equations. *Annals of mathematics*, pp.337-475.
- [25] Goldstein, M., Schlag, W. and Voda, M., (2019). On the spectrum of multi-frequency quasiperiodic Schrödinger operators with large coupling. *Inventiones mathematicae*, 217(2), pp.603-701.
- [26] Han, R. (2018). Dry Ten Martini problem for the non-self-dual extended Harper's model. *Transactions of the American Mathematical Society*, 370(1), 197-217.
- [27] Han, R., Continuity of measure of the spectrum for Schrödinger operators with potentials driven by shifts and skew-shifts on tori. Preprint.
- [28] Han, R., Jitomirskaya, S. and Yang, F., (2022). Anti-resonances and sharp analysis of Maryland localization for all parameters. arXiv preprint arXiv:2205.04021.
- [29] Han, R., Lemm, M. and Schlag, W., (2020). Effective multi-scale approach to the Schrödinger cocycle over a skew-shift base. *Ergodic Theory and Dynamical Systems*, 40(10), pp.2788-2853.
- [30] Jitomirskaya, S. and Liu, W., (2017). Arithmetic spectral transitions for the Maryland model. *Communications on Pure and Applied Mathematics*, 70(6), pp.1025-1051.
- [31] Jitomirskaya, S. and Yang, F., (2021). Pure point spectrum for the Maryland model: a constructive proof. *Ergodic Theory and Dynamical Systems*, 41(1), pp.283-294.
- [32] Kachkovskiy, I., Parnovski, L. and Shterenberg, R., (2024). On gaps in the spectra of quasiperiodic Schrödinger operators with discontinuous monotone potentials. arXiv preprint arXiv:2407.00705.
- [33] Karpeshina, Y., Parnovski, L. and Shterenberg, R., (2020). Bethe-Sommerfeld Conjecture and Absolutely Continuous Spectrum of Multi-Dimensional Quasi-Periodic Schrödinger Operators. arXiv preprint arXiv:2010.05881.
- [34] Konno, N., (2008). Quantum walks. In *Quantum potential theory* (pp. 309-452). Springer, Berlin, Heidelberg.
- [35] Krüger, H., (2012). The spectrum of skew-shift Schrödinger operators contains intervals. *Journal of Functional Analysis*, 262(3), pp.773-810.
- [36] Krüger, H., (2013). Orthogonal polynomials on the unit circle with Verblunsky coefficients defined by the skew-shift. *International Mathematics Research Notices*, 2013(18), pp.4135-4169.
- [37] Last, Y., (1994). Zero measure spectrum for the almost Mathieu operator. *Communications in mathematical physics*, 164, pp.421-432.
- [38] Puig, J., (2004). Cantor spectrum for the almost Mathieu operator. *Communications in mathematical physics*, 244(2), pp.297-309.
- [39] Shukla, K., & Chandrashekar, C. M. (2024). Quantum magnetometry using discrete-time quantum walk. *Physical Review A*, 109(3), 032608.
- [40] Simon, B. (1985). Almost periodic Schrödinger operators IV. The maryland model. *Annals of Physics*, 159(1), 157-183.
- [41] Simon, B. (2007). CMV matrices: Five years after. *Journal of Computational and Applied Mathematics*, 208(1), 120-154.
- [42] Simon, B. (2005). Orthogonal polynomials on the unit circle. American Mathematical Soc.
- [43] Solis, M. G., Spedale, D., & Yang, F. (2024). On the spectrum of magnetic Laplacian on the Lieb lattice. arXiv preprint arXiv:2401.11405.

- [44] Wang, J., & Manouchehri, K. (2013). Physical implementation of quantum walks. Heidelberg, Springer Berlin, 10, 978-3.
- [45] Venegas-Andraca, S. E. (2012). Quantum walks: a comprehensive review. Quantum Information Processing, 11(5), 1015-1106.
- [46] Yang, F. (2024). Anderson localization for the unitary almost Mathieu operator. Nonlinearity, 37(8), 085010.
- [47] Yalçınkaya, İ., & Gedik, Z. (2015). Two-dimensional quantum walk under artificial magnetic field. Physical Review A, 92(4), 042324.

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LA 70803, USA
Email address: yangf@lsu.edu